

Inference on Average Treatment Effect under Minimization and Other Covariate-Adaptive Randomization Methods

Ting Ye*

Yanyao Yi†

Jun Shao‡

April 9, 2024

Abstract

Covariate-adaptive randomization schemes such as the minimization and stratified permuted blocks are often applied in clinical trials to balance treatment assignments across prognostic factors. The existing theoretical developments on inference after covariate-adaptive randomization are mostly limited to situations where a correct model between the response and covariates can be specified or the randomization method has well-understood properties. Based on stratification with covariate levels utilized in randomization and a further adjusting for covariates not used in randomization, in this article we propose several estimators for model free inference on average treatment effect defined as the difference between response means under two treatments. We establish asymptotic normality of the proposed estimators under all popular covariate-adaptive randomization schemes including the minimization whose theoretical property is unclear, and we show that the asymptotic distributions are invariant with respect to covariate-adaptive randomization methods. Consistent variance estimators are constructed for asymptotic inference. Asymptotic relative efficiencies and finite sample properties of estimators are also studied. We recommend using one of our proposed estimators for valid and model free inference after covariate-adaptive randomization.

Keywords: Adjusting for covariates; Balancedness of treatment assignments; Efficiency; Generalized regression; Model free inference; Multiple treatment arms; Stratification; Variance estimation.

*Department of Statistics, University of Pennsylvania

†Global Statistical Sciences, Eli Lilly and Company

‡School of Statistics, East China Normal University

1 Introduction

Consider a clinical trial to compare k treatments with given treatment assignment proportions π_1, \dots, π_k , where $k \geq 2$ is a fixed positive integer, $\sum_{t=1}^k \pi_t = 1$, and π_t can be any known number strictly between 0 and 1. In many trials patients are not all available for simultaneous assignment of treatments but rather arrive sequentially and must be treated immediately. Thus, simple randomization, which assigns patients to treatments completely at random, may yield sample sizes not following the assignment proportions across prognostic factors or covariates, e.g., institution, disease stage, prior treatment, gender, and age, which are thought to have significant influence on the responses of interest. A remedy is to apply covariate-adaptive randomization, i.e., treatment assignment of the i th patient depends on the observed covariate value of this patient and the assignments and covariate values of all $i - 1$ previously assigned patients. In this article, we focus on enforcing assignment allocation across levels of a covariate vector Z whose components are discrete or discretized continuous covariates. There are model-based approaches of balancing discrete or continuous covariates for estimation efficiency (Atkinson 1982, 1999, 2002; Rosenberger and Sverdlov 2008; Senn et al. 2010; Baldi Antognini and Zagoraiou 2011), which are not further considered in this article. The oldest method of balancing covariates is the minimization (Taves 1974) intended to balance treatment assignments across marginal levels of Z : it assigns the i th patient by minimizing a weighted sum of squared or absolute differences between the numbers of patients, up to the i th, assigned to treatments over marginal levels of Z . Pocock and Simon (1975) extended Taves’ procedure to achieving minimization with a given probability, which is still referred to as the minimization method. Other popular covariate-adaptive randomization methods include the stratified permuted block randomization (Zelen 1974), the stratified biased coin (Shao et al. 2010; Kuznetsova and Johnson 2017), and the stratified urn design (Wei 1977; Zhao and Ramakrishnan 2016). See Schulz and Grimes (2002) and Rosenberger and Sverdlov (2008) for nice summaries. As pointed out in Taves (2010), from 1989 to 2008, over 500 clinical trials implemented the minimization method to balance important covariates, despite some criticisms by Smith (1984) and Senn et al. (2010). According to a recent review of nearly 300 clinical trials published in 2009 and 2014 (Ciolino et al. 2019), 237 of them used covariate-adaptive randomization.

Although data are collected under covariate-adaptive randomization, conventional inference procedures constructed based on simple randomization are often applied in practice. This has raised concerns because statistical inference on treatment effects should be made using procedures valid under the particular randomization scheme used in data collection. Applying conventional inference after covariate-adaptive randomization may lead to invalid results, especially for the minimization because its theoretical property remains largely unclear. In its 2015 guideline, European Medicines Agency (2015) raised concerns and specifically pointed out that “possible implications of dynamic allocation methods [minimization] on the analysis e.g. with regard to bias and Type I error control should be carefully considered, . . . conventional statistical methods do not always control the Type I error”.

Starting with Shao et al. (2010), there has been significant progress in understanding the theoretical properties of statistical tests under covariate-adaptive randomization, e.g., Hu and Hu (2012), Shao and Yu (2013), Ma et al. (2015), Bugni et al. (2018, 2019), Ye (2018), and Ye and Shao (2020). Another important stream of statistical inference methods is based on permutation tests or re-randomization inference, e.g., Simon and Simon (2011), Kaiser (2012), and Bugni et al. (2018). However, except for Bugni et al. (2019), all theoretical results are established under the assumption that either a correct model between the responses of interest and covariates is available or the covariate-adaptive randomization procedure has a well-understood property, described as type 1 or type 2 later in §2 of the current paper. It should be noted that model misspecification often occurs, especially when there are many covariates, and the minimization method is neither type 1 nor type 2. The minimization is applied in practice very often (Pocock and Simon 1975), mainly because it aims to minimize the imbalance across marginal levels of Z , not every joint level of Z , which is sufficient in many applications. Enforcing treatment balance in every joint level of Z may cause sparsity of data when the dimension of Z is not small.

To fill the gap, in this paper we propose asymptotically valid inference on the average treatment effect defined as the difference between population response means of every treatment pair, under covariate-adaptive randomization including minimization. Our main idea is to apply stratification according to the levels of discrete Z and to adjust for covariates not used in treatment random-

ization through generalized regression for improving efficiency. Our estimator without adjusting for covariates, which is not the most efficient one, coincides with the estimator derived under a different approach in the recent publication (Bugni et al. 2019). Asymptotic normality of the proposed treatment effect estimators is established with explicit limiting variance formulas that can be used for inference as well as comparing relative efficiencies. Our results are not only model-free, i.e., only the existence of second-order moments of the responses and covariates are required, but also invariant with respect to covariate-adaptive randomization schemes, i.e., the same inference procedure can be applied under any covariate-adaptive randomization. We also study and compare inference procedures by simulations and illustrate our method in a real data example. R codes for the methods proposed in this paper can be found in the R package RobinCar posed at <https://github.com/tye27/RobinCar>.

2 Preliminaries

Let I be the treatment indicator vector, which equals e_t if the patient is assigned to treatment t , $t = 1, \dots, k$, where e_t denotes a vector with only the t th component being 1 while the rest being 0. Let $Y^{(t)}$ be the potential response under treatment t , W be a vector of all observed covariates, and Z be a discrete function of W utilized in covariate-adaptive randomization. For patient i , let I_i , W_i , and $Y_i^{(t)}$, $t = 1, \dots, k$, be realizations of I , W , and $Y^{(t)}$, $t = 1, \dots, k$, respectively, where $i = 1, \dots, n$, and n is the total number of patients in all treatment arms. For every patient i , I_i is generated after Z_i is observed, and only the potential response from the treatment indicated by I_i is observed, i.e., we observe $Y_i = Y_i^{(t)}$ if and only if $I_i = e_t$.

After all treatment assignments are made and responses are collected, we would like to make inference based on the observed data $\{W_i, I_i, Y_i, i = 1, \dots, n\}$. In this paper, we consider inference on the average treatment effect between treatments t and s ,

$$\theta = E(Y^{(t)} - Y^{(s)}), \quad (1)$$

where E denotes the population expectation and $s \neq t$ are given integers between 1 and k . Note that θ in (1) depends on t and s , but a subscript is omitted for simplicity, as we mainly consider the average treatment effect for two fixed treatment arms.

For our inference procedure studied in §3, we describe some minimal conditions. The first one is about the population for potential responses and covariates.

- (C1) $(Y_i^{(1)}, \dots, Y_i^{(k)}, W_i)$, $i = 1, \dots, n$, are independent and identically distributed as $(Y^{(1)}, \dots, Y^{(k)}, W)$ and $Y^{(t)}$ has finite second-order moments, $t = 1, \dots, k$.

This condition is model-free: there is no assumption on the relationship between W and the potential response $Y^{(t)}$ that may be continuous or discrete.

Under simple randomization, I_i 's are independent of $(Y_i^{(1)}, \dots, Y_i^{(k)}, W_i)$'s and are independent and identically distributed with $\text{pr}(I_i = e_t) = \pi_t$. To enforce assignment proportions at each joint level of Z treated as stratum, three popular covariate-adaptive randomization schemes are the stratified permuted block randomization method (Zelen 1974), the stratified biased coin method (Shao et al. 2010; Kuznetsova and Johnson 2017), and the stratified urn design (Wei 1977; Zhao and Ramakrishnan 2016).

The minimization (Taves 1974; Pocock and Simon 1975; Han et al. 2009) is the same as the stratified biased coin method if Z is one-dimensional, but is very different from the above three stratification methods with a multivariate Z . It aims to enforce assignment ratio across marginal levels of Z , not every stratum defined by the joint level of Z . Assignments are made by minimizing a weighted sum of squared or absolute differences between the numbers of patients assigned to treatment arms across marginal levels of Z . Because only marginal levels of Z are considered in minimization, this method is also called the marginal method in Ma et al. (2015) and Ye and Shao (2020).

We assume the following minimal conditions for covariate-adaptive randomization.

- (C2) $(I_i, i = 1, \dots, n)$ and $(Y_i^{(1)}, \dots, Y_i^{(k)}, W_i, i = 1, \dots, n)$ are conditionally independent given Z_1, \dots, Z_n .
- (C3) Z is discrete with finitely many levels given in a set \mathcal{Z} . For each $t = 1, \dots, k$, $\text{pr}(I_i = e_t \mid Z_1, \dots, Z_n) = \pi_t$ for $i = 1, \dots, n$, and $\{n(z)\}^{-1}D_t(z)$ converges to 0 in probability as $n \rightarrow \infty$.

for every $z \in \mathcal{Z}$, where $n(z)$ is the number of patients with $Z_i = z$, and $D_t(z) = n_t(z) - \pi_t n(z)$ with $n_t(z)$ being the number of patients with $Z_i = z$ assigned to treatment t .

Condition (C2) is reasonable because (i) given Z_i 's, W_i 's contain covariates not used in randomization, and (ii) treatment assignments do not affect the potential responses, although they do affect the observed responses Y_i 's. Condition (C3) holds for most covariate-adaptive randomization schemes (Baldi Antognini and Zagoraiou 2015), certainly for all schemes considered in this paper, the minimization and three stratified designs, the permuted block, biased coin, and urn design.

We classify all covariate-adaptive randomization methods into the following three types in terms of $D_t(z)$ defined in (C3).

Type 1. For every t and z , $\{n(z)\}^{-1/2} D_t(z) \rightarrow 0$ in probability as $n \rightarrow \infty$.

Type 2. For every t , $D_t(z)$, $z \in \mathcal{Z}$, are independent and, for every t and z , $\{n(z)\}^{-1/2} D_t(z) \xrightarrow{d} N(0, v_t)$ with a known $v_t > 0$, where \xrightarrow{d} denotes convergence in distribution as $n \rightarrow \infty$.

Type 3. Methods not in type 1 or 2.

The three types are defined based on their degree in enforcing the balancedness according to the given assignment proportions within every joint level of Z . Type 1 is the strongest, since $D_t(z)$ measures the imbalance of assignments within stratum z . The property $\{n(z)\}^{-1/2} D_t(z) \rightarrow 0$ in probability is stronger than $\{n(z)\}^{-1} D_t(z) \rightarrow 0$ in probability in (C3). Type 2 is weaker than type 1 in enforcing the balancedness, as it requires $\{n(z)\}^{-1/2} D_t(z)$ converging in distribution, not in probability to 0, although it is still stronger than $\{n(z)\}^{-1} D_t(z) \rightarrow 0$ in probability.

Representatives of type 1 methods are stratified permuted block randomization and stratified biased coin methods. Specifically, under stratified permuted block randomization, $D_t(z)$ is bounded by the maximum block size. For the stratified biased coin method, it follows from a result in Efron (1971) that $D_t(z)$ is bounded in probability. The stratified urn design is type 2 with $v_t = 1/12$ when $k = 2$ and $\pi_1 = \pi_2 = 1/2$ (Wei 1978). Simple randomization treated as a special case of covariate-adaptive randomization is also type 2. Finally, the minimization is type 3, since it is neither type 1 nor type 2 (Ye and Shao 2020). Specifically, under minimization, $D_t(z)$ and $D_t(z')$ with $z \neq z'$ are not independent, and their relationship is complicated, because assignments are made according to marginal levels of Z .

For type 1 methods, some theoretical results in statistical testing have been established; see, for example, Shao et al. (2010), Shao and Yu (2013), Ye (2018), Bugni et al. (2018, 2019), and Ye and Shao (2020). Bugni et al. (2018, 2019) and Ye and Shao (2020) also considered type 2 methods. In the next section, we propose inference procedures and establish their asymptotic validity under general covariate-adaptive randomization including minimization.

3 Inference on Average Treatment Effect

To make asymptotically valid inference on θ defined in (1), the key is to construct an estimator of θ and derive its asymptotic distribution. Under simple randomization, the simplest estimator is the response mean difference $\bar{Y}_t - \bar{Y}_s$, where \bar{Y}_t is the sample mean of responses under treatment t . Although $\bar{Y}_t - \bar{Y}_s$ is asymptotically normal under type 1 or 2 covariate-adaptive randomization, it is generally not efficient as covariate information is not utilized in estimation. More seriously, the asymptotic distribution of $\bar{Y}_t - \bar{Y}_s$ is not known under type 3 covariate-adaptive randomization such as the minimization. Bugni et al. (2018) derived a different estimator of θ , called the strata fixed effect estimator in their §4.2, but its asymptotic normality is established only for type 1 or 2 covariate-adaptive randomization and, thus, it cannot be used under type 3 randomization such as the minimization.

Let $\bar{Y}_t(z)$ be the sample mean of Y_i 's with $Z_i = z$ under treatment t . The following stratified response mean differences with strata being all joint levels of Z is proposed in expression (8) of Bugni et al. (2019),

$$\hat{\theta} = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{\bar{Y}_t(z) - \bar{Y}_s(z)\}, \quad (2)$$

although Bugni et al. (2019) provided $\hat{\theta}$ in a different form derived under a fully saturated linear

regression. If the weight $n(z)/n$ in (2) is replaced by the population weight $\text{pr}(Z = z)$, then $\hat{\theta}$ is exactly the stratified estimator in survey sampling. We use $n(z)/n$ in (2) as $\text{pr}(Z = z)$ is unknown.

Although $\hat{\theta}$ in (2) utilizes information from Z by stratification and is asymptotically more efficient than the simple $\bar{Y}_t - \bar{Y}_s$ or the strata fixed effect estimator in Bugni et al. (2018), it does not make use of covariate information in W but not in Z . Note that W may contain components that are not in Z but are related with the potential responses $Y^{(t)}, t = 1, \dots, k$, or some components of Z are discretized components of W and the remaining information after discretization is still predictive of $Y^{(t)}, t = 1, \dots, k$.

Let X be a function of W that we want to further adjust for. We now consider improving $\hat{\theta}$ in (2) by utilizing X . To maintain model free estimation, we do not impose any model between $Y^{(t)}$ and X , but adjust for covariate X within each $Z = z$ by applying the generalized regression approach in survey sampling, first discussed in Cassel et al. (1976) and studied extensively in the literature, for example, Särndal et al. (2003), Lin (2013), Shao and Wang (2014), and Ta et al. (2020). Since this approach is model-assisted but not model-based, i.e., a model is used to derive efficient estimators that are still asymptotically valid even if the model is incorrect, it suits our purpose of utilizing covariates without modeling.

Let X_i be the value of covariate X for patient i , $\bar{X}_t(z)$ be the sample mean of X_i 's with $Z_i = z$ under treatment t , and

$$\hat{\beta}_t(z) = \left[\sum_{i: I_i = e_t, Z_i = z} \{X_i - \bar{X}_t(z)\} \{X_i - \bar{X}_t(z)\}^T \right]^{-1} \sum_{i: I_i = e_t, Z_i = z} \{X_i - \bar{X}_t(z)\} Y_i,$$

where a^T is the transpose of vector a . Within treatment t and $Z = z$, $\hat{\beta}_t(z)$ is the least squares estimator of the coefficient vector in front of X under a linear model between $Y^{(t)}$ and X , but the model is not required to be correct. Then, our first proposed estimator of θ after adjusting for covariates is

$$\hat{\theta}_A = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{\bar{Y}_{t,A}(z) - \bar{Y}_{s,A}(z)\}, \quad \bar{Y}_{t,A}(z) = \bar{Y}_t(z) - \{\bar{X}_t(z) - \bar{X}(z)\}^T \hat{\beta}_t(z), \quad (3)$$

where $\bar{X}(z)$ is the sample mean of X_i 's of all patients with $Z_i = z$.

Within $Z_i = z$, if we assume that the linear models under all treatments have the same coefficient vector for X , such as a homogeneous ANCOVA model with covariate vector X , then we can replace $\hat{\beta}_t(z)$ by

$$\hat{\beta}(z) = \left[\sum_{t=1}^k \sum_{i: I_i = e_t, Z_i = z} \{X_i - \bar{X}_t(z)\} \{X_i - \bar{X}_t(z)\}^T \right]^{-1} \sum_{t=1}^k \sum_{i: I_i = e_t, Z_i = z} \{X_i - \bar{X}_t(z)\} Y_i.$$

Again, the model is not required to be correct in order to use $\hat{\beta}(z)$. This leads to an alternative estimator of θ after adjusting for covariates,

$$\hat{\theta}_B = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{\bar{Y}_{t,B}(z) - \bar{Y}_{s,B}(z)\}, \quad \bar{Y}_{t,B}(z) = \bar{Y}_t(z) - \{\bar{X}_t(z) - \bar{X}(z)\}^T \hat{\beta}(z). \quad (4)$$

When $k > 2$, both $\bar{X}(z)$ and $\hat{\beta}(z)$ involve data from patients in treatment arms other than t and s .

The following theorem proved in the Supplementary Material derives the asymptotic distributions of $\hat{\theta}$ in (2), $\hat{\theta}_A$ in (3), and $\hat{\theta}_B$ in (4), under covariate-adaptive randomization including minimization.

Theorem 1. Assume (C1)-(C3), the existence of second-order moments of X , $XY^{(t)}, t = 1, \dots, k$, and that $\text{var}(X | Z = z)$ is positive definite for every $z \in \mathcal{Z}$. As $n \rightarrow \infty$,

$$\begin{aligned} \sqrt{n}(\hat{\theta} - \theta) &\xrightarrow{d} N(0, \sigma_U^2 + \sigma_V^2), \\ \sqrt{n}(\hat{\theta}_A - \theta) &\xrightarrow{d} N(0, \sigma_A^2 + \sigma_V^2), \\ \sqrt{n}(\hat{\theta}_B - \theta) &\xrightarrow{d} N(0, \sigma_B^2 + \sigma_V^2), \end{aligned}$$

where

$$\begin{aligned}\sigma_U^2 &= E\{\text{var}(Y^{(t)} | Z)/\pi_t + \text{var}(Y^{(s)} | Z)/\pi_s\}, \\ \sigma_A^2 &= E[\text{var}\{Y^{(t)} - X^T \beta_t(Z) | Z\}/\pi_t + \text{var}\{Y^{(s)} - X^T \beta_s(Z) | Z\}/\pi_s] \\ &\quad + E[\{\beta_t(Z) - \beta_s(Z)\}^T \text{var}(X | Z) \{\beta_t(Z) - \beta_s(Z)\}], \\ \sigma_B^2 &= E[\text{var}\{Y^{(t)} - X^T \beta(Z) | Z\}/\pi_t + \text{var}\{Y^{(s)} - X^T \beta(Z) | Z\}/\pi_s], \\ \sigma_V^2 &= \text{var}\{E(Y^{(t)} - Y^{(s)} | Z)\},\end{aligned}$$

$t, s = 1, \dots, k$, $\beta_t(z) = \{\text{var}(X | Z = z)\}^{-1} \text{cov}(X, Y^{(t)} | Z = z)$, and $\beta(z) = \sum_{t=1}^k \pi_t \beta_t(z)$.

Theorem 1 is model free and is applicable to any covariate-adaptive randomization method satisfying (C2)-(C3), most noticeably the minimization for which very little is known about its theoretical property, as the minimization is neither type 1 nor type 2 described in §2. This provides a solid foundation for valid and model free inference after minimization.

The asymptotic result in Theorem 1 is invariant with respect to randomization methods, i.e., σ_U^2 , σ_A^2 , σ_B^2 , and σ_V^2 do not depend on the randomization scheme. In other words, each estimator of θ in (2)-(4) has the same asymptotic distribution and efficiency regardless of which randomization scheme is used for treatment assignments, including simple randomization. This is intrinsically different from many existing results that are dependent on randomization methods (Shao and Yu 2013; Ma et al. 2015; Bugni et al. 2018). The only result invariant with respect to randomization methods can be found in the literature is in Bugni et al. (2019) for $\hat{\theta}$ in (2), although it does not explicitly state this invariance property.

Due to the use of covariate-adaptive randomization, the sample mean \bar{Y}_t is not an average of independent random variables and, thus, the asymptotic distributions of estimators in (2)-(4) cannot be obtained by directly applying the central limit theorem for sum of independent random variables. We overcome this difficulty by decomposing $\hat{\theta} - \theta$ as the sum of the following two uncorrelated terms,

$$\begin{aligned}U &= \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left[\{\bar{Y}_t(z) - \bar{Y}_s(z)\} - \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} \right], \\ V &= \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} - \theta.\end{aligned}$$

Conditioned on $(I_1, \dots, I_n, Z_1, \dots, Z_n)$, U is an average of independent terms so its limiting distribution can be derived by applying the central limit theorem, which consequently provides the unconditional asymptotic distribution of U . For V , the only random part is $n(z)$ whose limiting distribution can be easily derived. For $\hat{\theta}_A$ or $\hat{\theta}_B$, a similar decomposition can be obtained with the same V and a different U incorporating the covariate adjustment term. Details can be found in the Supplementary Material.

This decomposition is not only the key to establishing the asymptotic result, but also identifies two sources of variation. The variation of potential responses $Y^{(t)}$ and $Y^{(s)}$ explained by Z is represented by σ_U^2 . The variation from treatment effect heterogeneity is measured by σ_V^2 . Note that we allow arbitrary treatment effect heterogeneity, i.e., different subgroups according to levels of Z may benefit differently from the treatment. If there is no treatment effect heterogeneity, then $\sigma_V^2 = 0$.

The asymptotic relative efficiencies among $\hat{\theta}$, $\hat{\theta}_A$ and $\hat{\theta}_B$ are summarized in the following result.

Theorem 2. *Under the same assumptions in Theorem 1, for σ_U^2 , σ_A^2 , σ_B^2 , $\beta_t(z)$ and $\beta(z)$ defined in Theorem 1, we have*

$$\begin{aligned}\sigma_U^2 - \sigma_A^2 &= E \left[\{\pi_s \beta_t(Z) + \pi_t \beta_s(Z)\}^T \text{var}(X | Z) \{\pi_s \beta_t(Z) + \pi_t \beta_s(Z)\} \right] \{\pi_t \pi_s (\pi_t + \pi_s)\}^{-1} \\ &\quad + E \left[\{\beta_t(Z) - \beta_s(Z)\}^T \text{var}(X | Z) \{\beta_t(Z) - \beta_s(Z)\} \right] \{(\pi_t + \pi_s)^{-1} - 1\}, \\ \sigma_B^2 - \sigma_A^2 &= E \left[\{\beta_t(Z) - \beta(Z)\}^T \text{var}(X | Z) \{\beta_t(Z) - \beta(Z)\} \right] \pi_t^{-1} \\ &\quad + E \left[\{\beta_s(Z) - \beta(Z)\}^T \text{var}(X | Z) \{\beta_s(Z) - \beta(Z)\} \right] \pi_s^{-1} \\ &\quad - E \left[\{\beta_t(Z) - \beta_s(Z)\}^T \text{var}(X | Z) \{\beta_t(Z) - \beta_s(Z)\} \right].\end{aligned}$$

Consequently, $\sigma_A^2 \leq \sigma_U^2$, where the equality holds if and only if for every $z \in \mathcal{Z}$,

$$\pi_s \beta_t(z) + \pi_t \beta_s(z) = 0 \quad \text{and} \quad \{\beta_t(z) - \beta_s(z)\}(1 - \pi_t - \pi_s) = 0; \quad (5)$$

and $\sigma_A^2 \leq \sigma_B^2$, where the equality holds if and only if for every $z \in \mathcal{Z}$,

$$\beta(z) = \{\pi_s \beta_t(z) + \pi_t \beta_s(z)\} / (\pi_s + \pi_t) \quad \text{and} \quad \{\beta_t(z) - \beta_s(z)\}(1 - \pi_t - \pi_s) = 0. \quad (6)$$

Theorem 2 indicates that $\hat{\theta}_A$ is always asymptotically more efficient than $\hat{\theta}$ unless (5) holds, in which case $\hat{\theta}$ and $\hat{\theta}_A$ have the same asymptotic efficiency. This theoretically corroborates the perception that covariate adjustment with a full set of treatment-covariate interactions can not hurt efficiency. When there are more than two treatments, $1 - \pi_t - \pi_s > 0$ and, consequently, (5) holds only when $\beta_t(z) = \beta_s(z) = 0$ for every z , i.e., X is uncorrelated with the potential responses $Y^{(t)}$ and $Y^{(s)}$ after conditioning on Z so that adjusting for X is unnecessary. When there are only two treatments, (5) also holds if $\pi_t = \pi_s = 1/2$ and $\beta_t(z) = -\beta_s(z)$ for every z . An example is given in §4.

Note that $\hat{\beta}(z)$ used in $\hat{\theta}_B$ ignores the fact that $\text{cov}(X, Y^{(t)} \mid Z = z)$ may depend on treatment t . That is why $\hat{\theta}_B$ is asymptotically not as efficient as $\hat{\theta}_A$ in general, and $\sigma_B^2 = \sigma_A^2$ when these covariances are the same for every t and every z , i.e., $\beta_1(z) = \dots = \beta_k(z)$. An exceptional case is that $\sigma_A^2 = \sigma_B^2$ when there are only two treatments and $\pi_t = \pi_s = 1/2$. In fact, $\hat{\theta}_B$ may be asymptotically less efficient than $\hat{\theta}$, i.e., covariate adjustment with only the main effects may hurt efficiency, a perspective in Freedman (2008) and Lin (2013). For example, there are scenarios in which (5) holds but (6) does not. Simulation examples are given in §4, where comparisons of $\hat{\theta}$, $\hat{\theta}_A$, and $\hat{\theta}_B$ are made.

To make model free inference on the average treatment effect θ defined in (1), we only need to apply Theorem 1 and construct consistent estimators of limiting variances σ_U^2 , σ_A^2 , σ_B^2 , and σ_V^2 . Let $S_t^2(z)$ be the sample variance of Y_i 's in the group of patients under treatment t with $Z_i = z$, $S_{t,A}^2(z)$ be $S_t^2(z)$ with Y_i replaced by $Y_i - X_i^T \hat{\beta}_t(z)$, $S_{t,B}^2(z)$ be $S_t^2(z)$ with Y_i replaced by $Y_i - X_i^T \hat{\beta}(z)$, and $\hat{\Sigma}(z)$ be the sample covariance matrix of X_i 's within $Z_i = z$. It is shown in the Supplementary Material that, under (C1)-(C3), the following estimators are consistent for σ_U^2 , σ_V^2 , σ_A^2 , and σ_B^2 , respectively,

$$\begin{aligned} \hat{\sigma}_U^2 &= \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left\{ \frac{S_t^2(z)}{\pi_t} + \frac{S_s^2(z)}{\pi_s} \right\}, \quad \hat{\sigma}_V^2 = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{ \bar{Y}_t(z) - \bar{Y}_s(z) \}^2 - \hat{\theta}^2, \\ \hat{\sigma}_A^2 &= \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left[\frac{S_{t,A}^2(z)}{\pi_t} + \frac{S_{s,A}^2(z)}{\pi_s} + \{ \hat{\beta}_t(z) - \hat{\beta}_s(z) \}^T \hat{\Sigma}(z) \{ \hat{\beta}_t(z) - \hat{\beta}_s(z) \} \right], \\ \hat{\sigma}_B^2 &= \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left\{ \frac{S_{t,B}^2(z)}{\pi_t} + \frac{S_{s,B}^2(z)}{\pi_s} \right\}, \end{aligned}$$

regardless of which type of covariate-adaptive randomization method is used. Note that $\hat{\sigma}_U^2$ for $\hat{\theta}$ is different from the estimator defined in (36) of Bugni et al. (2019).

4 Simulation Results

There are many publications on empirical studies under covariate-adaptive randomization in the last four decades. Some recent results are in Senn et al. (2010), Kahan and Morris (2012), and Xu et al. (2016).

To evaluate and compare our proposed estimators $\hat{\theta}$, $\hat{\theta}_A$, and $\hat{\theta}_B$ in terms of estimation bias and standard deviation, and to examine variance estimators and the related asymptotic confidence intervals based on Theorem 1, we present some simulation results in this section. We consider two covariates, i.e., $W = (X_1, X_2)$, where X_1 is binary with $\text{pr}(X_1 = 1) = 1/2$ and, conditioned on X_1 , $X_2 \sim N(X_1 - 0.5, 1)$. For the potential responses, we consider two treatments in cases I-III and three treatments in case IV.

Case I: $Y^{(1)} \mid W \sim N(4X_1 + 2X_2, 1)$, $Y^{(2)} \mid W \sim N(\varphi + 4X_1 + 2X_2, 1)$.

Case II: $Y^{(1)} \mid W \sim N(4X_1 - 2X_2, 1)$, $Y^{(2)} \mid W \sim N(\varphi + 4X_1 + 2X_2, 1)$.

Case III: $Y^{(1)} | W \sim N(0.25 + 3X_1 + 0.2X_2^2, X_1 + 0.5)$, $Y^{(2)} | W \sim N(\varphi + 4X_1 + 2X_2, 1)$.

Case IV: $Y^{(3)} | W \sim N(\psi + 1 + 2X_1 - X_2, 1)$, and $Y^{(1)}$ and $Y^{(2)}$ are the same as those in case III. We use $\varphi = \psi = 1$ in the simulation, which does not affect the relative performance of estimators and coverage probability of related confidence intervals.

Case I has homogeneous treatment effects; case II has treatment effect heterogeneity since the effects of X_2 on $Y^{(1)}$ and $Y^{(2)}$ have different signs; case III has the most severe treatment effect heterogeneity as $Y^{(1)} | W$ and $Y^{(2)} | W$ have very different distributions; case IV considers multiple treatments.

We consider 3 different Z 's in covariate-adaptive randomization. The first one is $Z = X_1$ with 2 levels, in which case the function of W not used in randomization but still related with the potential responses is $h(W) = X_2$. The second one is $Z = (X_1, d_2)$ with 4 levels, where d_2 is the discretized X_2 with 2 categories $(-\infty, 0)$ and $[0, \infty)$, and $h(W)$ is the continuous value of X_2 in $(-\infty, 0)$ or $(0, \infty)$. The third one is $Z = (X_1, d_4)$ with 8 levels, where d_4 is the discretized X_2 with 4 categories $(-\infty, -0.8)$, $[-0.8, 0)$, $[0, 0.8)$, and $[0.8, \infty)$, and $h(W)$ is the continuous value of X_2 in $(-\infty, -0.8)$, $(-0.8, 0)$, $(0, 0.8)$, or $(0.8, \infty)$. In any case, $X = X_2$ is equivalent to $h(W)$ and is used covariate adjustment.

For the randomization method, we consider minimization with treatment allocation 1:1 or 1:2 for cases I-III, and 1:2:2 for case IV. Simulation results for two other randomization methods, the stratified permuted block randomization and the stratified urn design can be found in the Supplementary Material.

We consider the total sample size $n = 100$ or 500 . For these sample sizes, the smallest possible expected numbers of patients within a stratum and treatment according to number of Z levels are given in Table 3. It can be seen that when $n = 100$ and $Z = (X_1, d_4)$ has 8 levels, with non-negligible probability, the number of patients in some stratum-treatment combination is fewer than 2 and thus calculation of estimators in (2)-(4) and their variance estimators are not possible. Therefore, for cases I-III, we omit the scenario with $n = 100$ and $Z = (X_1, d_4)$. For case IV, we focus on $n = 500$ and $Z = (X_1, d_2)$.

Tables 1-2 report the bias, standard deviation (SD), average estimated SD (SE), and coverage probability (CP) of asymptotic 95% confidence interval, estimate ± 1.96 SE, of $\hat{\theta}$, $\hat{\theta}_A$, and $\hat{\theta}_B$ for cases I-IV. Every scenario is evaluated with 2,000 simulation runs. The following is a summary of the results in Tables 1-2.

1. All estimators have negligible biases that are smaller than 1% in most cases.
2. The variance estimators or SE's are very accurate so that the coverage probabilities of confidence intervals are adequate, except for a few cases with $n = 100$ and four Z categories.
3. With homogeneous treatment effects in case I, a more informative Z leads to a more efficient $\hat{\theta}$. However, the same phenomenon may not exist when treatment effect heterogeneity exists, though a more informative Z does not lead to a less efficient $\hat{\theta}$.
4. Adjusting for covariates, i.e., using $\hat{\theta}_A$ or $\hat{\theta}_B$, may lead to substantial improvements over $\hat{\theta}$ in terms of SD, which again agrees with our theory. The improvement is larger when a less informative Z is utilized in randomization, such as $Z = X_1$. Another interesting observation is that different Z used in randomization does not affect very much the SD of $\hat{\theta}_A$ or $\hat{\theta}_B$.
5. The comparison of $\hat{\theta}_A$ and $\hat{\theta}_B$ is also consistent with our theory in §3. Under 1:1 treatment allocation or homogeneous treatment effects, $\hat{\theta}_B$ is as good as $\hat{\theta}_A$ and sometimes slightly better in finite sample performance. When treatment allocation is 1:2 and treatment effect heterogeneity exists, $\hat{\theta}_B$ is not as good as $\hat{\theta}_A$ and could be even worse than $\hat{\theta}$. The same is observed when treatment allocation is 1:2:2.
6. In case II with 1:1 treatment allocation, $\text{cov}(X, Y^{(2)} | Z = z) = -\text{cov}(X, Y^{(1)} | Z = z)$, i.e., (5) holds and, thus, $\hat{\theta}$ and $\hat{\theta}_A$ have very similar SD's, as predicted by Theorem 2. In this particular case, $\hat{\theta}_B$ is also as good as $\hat{\theta}_A$.
7. $\hat{\theta}_A$ has large SD and low CP when some stratum-treatment combinations have small number of patients, such as the case of $n = 100$ and $Z = (X_1, d_2)$ with 4 levels when treatment allocation is 1:2. However, even if the smallest expected number of patients is as small as 7.7 in the case of $n = 100$, Z having 4 levels, and 1:1 treatment allocation, $\hat{\theta}_A$ performs well.

5 A Real Data Example

In this section, we illustrate our methods by a real data example from [Chong et al. \(2016\)](#) whose goal is to evaluate the contribution of low dietary iron intake to human capital attainment by measuring the causal effect of reducing adolescent anemia on school attainment. The dataset is publicly available at <https://www.openicpsr.org/openicpsr/project/113624/version/V1/view>. In brief, [Chong et al. \(2016\)](#) conducted an experiment on adolescents aged from 11 to 19 in rural Peru (Cajamarca) where the burden of iron deficiency is high, to study whether a low-cost intervention can encourage students to increase their iron intake and hence improve their school performance. A stratified permuted block randomization design was applied to assign $n = 219$ students to one of the following three promotional videos, considered as three treatments, with treatment allocation 1:1:1. The first video shows a popular soccer player encouraging iron supplements to maximize energy; the second video shows a physician encouraging iron supplements for overall health; and the third “placebo” video shows a dentist encouraging oral hygiene without mentioning iron at all. The strata are student’s school grades $Z \in \mathcal{Z} = \{1, 2, 3, 4, 5\}$. [Chong et al. \(2016\)](#) studied a variety of outcomes regarding cognitive function, school performance, and aspirations. As an example, we focus on the outcome of academic achievement, which is a standardized average of a students academic grades from the fall semester in subjects of math, foreign language, social sciences, science, and communications. The same outcome is also used by [Bugni et al. \(2019\)](#) in an example.

Estimates $\hat{\theta}$, $\hat{\theta}_A$, and $\hat{\theta}_B$ and their SE’s are reported in Table 4 for the average treatment effect between the soccer player and placebo videos, or physician and placebo videos, together with the p-values associated with two-sided tests of no treatment effect. The estimates from $\hat{\theta}$ are the same as those in [Bugni et al. \(2019\)](#). The covariate X used in $\hat{\theta}_A$ and $\hat{\theta}_B$ is the baseline anemia status thought to have interactive effect with treatment on the outcome, as mentioned in [Chong et al. \(2016\)](#). It can be seen that the SE of $\hat{\theta}_A$ is the smallest and, in terms of p-values, the effect between physician and placebo videos is only marginally significant when $\hat{\theta}$ is used, but very significant based on $\hat{\theta}_A$.

6 Recommendations and Discussions

To improve asymptotic efficiency, we recommend $\hat{\theta}_A$ in (3) since it is asymptotically better than $\hat{\theta}$ in (2) or $\hat{\theta}_B$ in (4). In the special case of two treatment arms with equal allocation, we recommend $\hat{\theta}_B$, since it is asymptotically equivalent to $\hat{\theta}_A$ and has better empirical performance.

As full stratification according to Z is required, one limitation of estimators in (2)-(4) is that all strata need to have large enough sizes, at least 10 per stratum and treatment combination ([Ye and Shao 2020](#)). Note that both covariate-adaptive randomization in treatment assignment and adjustment for covariates in estimation can gain efficiency, and covariate-adaptive randomization has an additional practically important advantage of balancing assignments across prognostic factors. Thus, how to choose Z and X is an important future research. It is also interesting to study estimators by combining strata of small sizes.

Supplementary Material

Supplementary material contains all technical proofs and more simulation results.

Table 1: Bias, standard deviation (SD), average estimated SD (SE), and coverage probability (CP) of 95% asymptotic confidence interval under minimization for cases I-III

n	case	Z	estimator	treatment allocation 1:1				treatment allocation 1:2			
				bias	SD	SE	CP	bias	SD	SE	CP
500	I	X_1	$\hat{\theta}$	-0.0038	0.1980	0.1999	0.9590	0.0070	0.2159	0.2124	0.9465
			$\hat{\theta}_B$	-0.0016	0.0909	0.0893	0.9445	0.0016	0.0954	0.0949	0.9510
			$\hat{\theta}_A$	-0.0016	0.0908	0.0893	0.9445	0.0017	0.0954	0.0948	0.9490
		X_1, d_2	$\hat{\theta}$	-0.0029	0.1492	0.1466	0.9450	-0.0013	0.1537	0.1553	0.9560
			$\hat{\theta}_B$	-0.0017	0.0900	0.0893	0.9455	-0.0011	0.0962	0.0947	0.9440
			$\hat{\theta}_A$	-0.0018	0.0901	0.0894	0.9455	-0.0009	0.0965	0.0945	0.9435
		X_1, d_4	$\hat{\theta}$	0.0005	0.1143	0.1150	0.9560	0.0003	0.1237	0.1219	0.9485
			$\hat{\theta}_B$	0.0010	0.0903	0.0893	0.9505	0.0013	0.0967	0.0946	0.9425
			$\hat{\theta}_A$	0.0007	0.0908	0.0893	0.9500	0.0010	0.0990	0.0944	0.9415
	II	X_1	$\hat{\theta}$	0.0085	0.2212	0.2191	0.9480	0.0067	0.2320	0.2303	0.9505
			$\hat{\theta}_B$	0.0086	0.2222	0.2185	0.9495	0.0063	0.2563	0.2541	0.9430
			$\hat{\theta}_A$	0.0086	0.2214	0.2191	0.9500	0.0078	0.2255	0.2212	0.9470
		X_1, d_2	$\hat{\theta}$	0.0084	0.2201	0.2191	0.9480	0.0076	0.2284	0.2251	0.9465
			$\hat{\theta}_B$	0.0076	0.2214	0.2178	0.9440	0.0078	0.2407	0.2344	0.9350
			$\hat{\theta}_A$	0.0085	0.2204	0.2190	0.9475	0.0077	0.2242	0.2212	0.9450
		X_1, d_4	$\hat{\theta}$	0.0057	0.2222	0.2192	0.9440	0.0104	0.2256	0.2230	0.9405
			$\hat{\theta}_B$	0.0061	0.2233	0.2177	0.9425	0.0108	0.2289	0.2254	0.9420
			$\hat{\theta}_A$	0.0057	0.2221	0.2190	0.9430	0.0101	0.2252	0.2211	0.9420
	III	X_1	$\hat{\theta}$	0.0003	0.1716	0.1731	0.9475	0.0072	0.1691	0.1667	0.9425
			$\hat{\theta}_B$	0.0029	0.1495	0.1477	0.9500	0.0048	0.1675	0.1656	0.9475
			$\hat{\theta}_A$	0.0031	0.1496	0.1479	0.9480	0.0081	0.1546	0.1533	0.9465
		X_1, d_2	$\hat{\theta}$	0.0016	0.1580	0.1593	0.9490	0.0032	0.1603	0.1595	0.9465
			$\hat{\theta}_B$	0.0050	0.1468	0.1470	0.9460	0.0061	0.1621	0.1576	0.9420
			$\hat{\theta}_A$	0.0032	0.1464	0.1474	0.9480	0.0052	0.1559	0.1523	0.9440
		X_1, d_4	$\hat{\theta}$	0.0042	0.1528	0.1525	0.9465	0.0049	0.1601	0.1557	0.9405
			$\hat{\theta}_B$	0.0080	0.1495	0.1468	0.9425	0.0094	0.1577	0.1539	0.9395
			$\hat{\theta}_A$	0.0048	0.1493	0.1474	0.9450	0.0069	0.1573	0.1521	0.9410
100	I	X_1	$\hat{\theta}$	0.0111	0.4441	0.4486	0.9545	0.0066	0.4609	0.4747	0.9580
			$\hat{\theta}_B$	0.0038	0.2035	0.1982	0.9425	-0.0017	0.2114	0.2092	0.9490
			$\hat{\theta}_A$	0.0038	0.2035	0.1983	0.9425	-0.0014	0.2126	0.2083	0.9485
		X_1, d_2	$\hat{\theta}$	0.0108	0.3316	0.3305	0.9425	0.0034	0.3475	0.3501	0.9490
			$\hat{\theta}_B$	0.0029	0.2021	0.1981	0.9465	0.0016	0.2181	0.2092	0.9365
			$\hat{\theta}_A$	0.0013	0.2148	0.2013	0.9440	-0.0035	0.4016	0.2190	0.9140
	II	X_1	$\hat{\theta}$	-0.0057	0.4857	0.4902	0.9515	-0.0143	0.5131	0.5157	0.9445
			$\hat{\theta}_B$	-0.0044	0.4958	0.4828	0.9420	-0.0152	0.5710	0.5612	0.9405
			$\hat{\theta}_A$	-0.0049	0.4878	0.4894	0.9500	-0.0101	0.4965	0.4943	0.9455
		X_1, d_2	$\hat{\theta}$	-0.0062	0.4874	0.4904	0.9495	-0.0102	0.5013	0.5031	0.9545
			$\hat{\theta}_B$	-0.0077	0.4981	0.4774	0.9355	-0.0096	0.5400	0.5114	0.9305
			$\hat{\theta}_A$	-0.0076	0.4906	0.4904	0.9455	-0.0147	0.5993	0.4999	0.9390
	III	X_1	$\hat{\theta}$	0.0003	0.3855	0.3869	0.9475	-0.0074	0.3729	0.3717	0.9470
			$\hat{\theta}_B$	0.0050	0.3349	0.3266	0.9410	-0.0052	0.3744	0.3649	0.9365
			$\hat{\theta}_A$	0.0047	0.3314	0.3291	0.9465	0.0045	0.3486	0.3390	0.9400
		X_1, d_2	$\hat{\theta}$	-0.0004	0.3541	0.3566	0.9445	-0.0054	0.3572	0.3569	0.9510
			$\hat{\theta}_B$	0.0104	0.3313	0.3236	0.9425	0.0084	0.3663	0.3445	0.9305
			$\hat{\theta}_A$	0.0015	0.3315	0.3298	0.9435	0.0003	0.5643	0.3456	0.9310

Table 2: Bias, standard deviation (SD), average estimated SD (SE), and coverage probability (CP) of 95% asymptotic confidence interval under minimization for case IV with $n = 500$

t	s	θ	estimator	bias	SD	SE	CP
2	1	1	$\hat{\theta}$	-0.0007	0.1907	0.1901	0.9515
			$\hat{\theta}_B$	0.0040	0.1840	0.1821	0.9470
			$\hat{\theta}_A$	0.0058	0.1777	0.1726	0.9375
3	1	1	$\hat{\theta}$	-0.0004	0.1541	0.1546	0.9445
			$\hat{\theta}_B$	0.0037	0.1616	0.1615	0.9445
			$\hat{\theta}_A$	0.0052	0.1479	0.1460	0.9395
3	2	0	$\hat{\theta}$	0.0004	0.2094	0.2082	0.9505
			$\hat{\theta}_B$	-0.0002	0.2077	0.2048	0.9445
			$\hat{\theta}_A$	-0.0006	0.2019	0.2007	0.9495

Table 3: Smallest expected number of patients among all stratum-treatment combinations

n	Z	number of levels	1:1 allocation	1:2 allocation	1:2:2 allocation
100	X_1	2	25.0	16.7	10.0
	X_1, d_2	4	7.7	5.1	3.1
	X_1, d_4	8	2.4	1.6	1.0
500	X_1	2	125	83.3	50
	X_1, d_2	4	38.6	25.7	15.4
	X_1, d_4	8	12.1	8.1	4.8

Table 4: Results from real data analysis

estimator	soccer versus placebo			physician versus placebo		
	estimate	SE	p-value	estimate	SE	p-value
$\hat{\theta}$	-0.051	0.205	0.803	0.409	0.207	0.048
$\hat{\theta}_B$	-0.089	0.203	0.661	0.444	0.202	0.028
$\hat{\theta}_A$	-0.048	0.199	0.807	0.480	0.198	0.015

References

- Atkinson, A. C. (1982). Optimum biased coin designs for sequential clinical trials with prognostic factors. *Biometrika*, 69(1):61–67.
- Atkinson, A. C. (1999). Optimum biased-coin designs for sequential treatment allocation with covariate information. *Statistics in Medicine*, 18(14):1741–1752.
- Atkinson, A. C. (2002). The comparison of designs for sequential clinical trials with covariate information. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 165(2):349–373.
- Baldi Antognini, A. and Zagoraiou, M. (2011). The covariate-adaptive biased coin design for balancing clinical trials in the presence of prognostic factors. *Biometrika*, 98(3):519–535.
- Baldi Antognini, A. and Zagoraiou, M. (2015). On the almost sure convergence of adaptive allocation procedures. *Bernoulli*, 21(2):881–908.
- Bugni, F. A., Canay, I. A., and Shaikh, A. M. (2018). Inference under covariate-adaptive randomization. *Journal of the American Statistical Association*, 113(524):1784–1796.
- Bugni, F. A., Canay, I. A., and Shaikh, A. M. (2019). Inference under covariate-adaptive randomization with multiple treatments. *Quantitative Economics*, 10(4):1747–1785.
- Cassel, C. M., Särndal, C. E., and Wretman, J. H. (1976). Some results on generalized difference estimation and generalized regression estimation for finite populations. *Biometrika*, 63(3):615–620.
- Chong, A., Cohen, I., Field, E., Nakasone, E., and Torero, M. (2016). Iron deficiency and schooling attainment in peru. *American Economic Journal: Applied Economics*, 8(4):222–55.
- Ciolino, J. D., Palac, H. L., Yang, A., Vaca, M., and Belli, H. M. (2019). Ideal vs. real: a systematic review on handling covariates in randomized controlled trials. *BMC Medical Research Methodology*, 19(1):136.
- Efron, B. (1971). Forcing a sequential experiment to be balanced. *Biometrika*, 58(3):403–417.
- European Medicines Agency (2015). *Guideline on adjustment for baseline covariates in clinical trials*.
- Freedman, D. A. (2008). On regression adjustments in experiments with several treatments. *Ann. Appl. Stat.*, 2(1):176–196.
- Han, B., Enas, N. H., and McEntegart, D. (2009). Randomization by minimization for unbalanced treatment allocation. *Statistics in Medicine*, 28(27):3329–3346.
- Hu, Y. and Hu, F. (2012). Asymptotic properties of covariate-adaptive randomization. *Annals of Statistics*, 40(3):1794–1815.
- Kahan, B. C. and Morris, T. P. (2012). Improper analysis of trials randomised using stratified blocks or minimisation. *Statistics in Medicine*, 31(4):328–340.
- Kaiser, L. D. (2012). Dynamic randomization and a randomization model for clinical trials data. *Statistics in Medicine*, 31(29):3858–3873.
- Kuznetsova, O. M. and Johnson, V. P. (2017). Approaches to expanding the two-arm biased coin randomization to unequal allocation while preserving the unconditional allocation ratio. *Statistics in Medicine*, 36(16):2483–2498.
- Lin, W. (2013). Agnostic notes on regression adjustments to experimental data: Reexamining freedman’s critique. *Ann. Appl. Stat.*, 7(1):295–318.

- Ma, W., Hu, F., and Zhang, L. (2015). Testing hypotheses of covariate-adaptive randomized clinical trials. *Journal of the American Statistical Association*, 110(510):669–680.
- Pocock, S. J. and Simon, R. (1975). Sequential treatment assignment with balancing for prognostic factors in the controlled clinical trial. *Biometrics*, 31(1):103–115.
- Rosenberger, W. F. and Sverdlov, O. (2008). Handling covariates in the design of clinical trials. *Statist. Sci.*, 23(3):404–419.
- Särndal, C.-E., Swensson, B., and Wretman, J. (2003). *Model assisted survey sampling*. Springer Science & Business Media.
- Schulz, K. F. and Grimes, D. A. (2002). Generation of allocation sequences in randomised trials: chance, not choice. *The Lancet*, 359(9305):515–519.
- Senn, S., Anisimov, V. V., and Fedorov, V. V. (2010). Comparisons of minimization and atkinson’s algorithm. *Statistics in Medicine*, 29(78):721–730.
- Shao, J. and Wang, S. (2014). Efficiency of model-assisted regression estimators in sample surveys. *Statistica Sinica*, 24(1):395–414.
- Shao, J. and Yu, X. (2013). Validity of tests under covariate-adaptive biased coin randomization and generalized linear models. *Biometrics*, 69(4):960–969.
- Shao, J., Yu, X., and Zhong, B. (2010). A theory for testing hypotheses under covariate-adaptive randomization. *Biometrika*, 97(2):347–360.
- Simon, R. and Simon, N. R. (2011). Using randomization tests to preserve type I error with response-adaptive and covariate-adaptive randomization. *Statistics & Probability Letters*, 81(7):767–772.
- Smith, R. L. (1984). Sequential treatment allocation using biased coin designs. *Journal of the Royal Statistical Society. Series B (Methodological)*, 46(3):519–543.
- Ta, T., Shao, J., Li, Q., and Wang, L. (2020). Generalized regression estimators with high-dimensional covariates. *Statistica Sinica*, 30(3):1135–1154.
- Taves, D. R. (1974). Minimization: A new method of assigning patients to treatment and control groups. *Clinical Pharmacology & Therapeutics*, 15(5):443–453.
- Taves, D. R. (2010). The use of minimization in clinical trials. *Contemporary Clinical Trials*, 31(2):180–184.
- Wei, L.-J. (1977). A class of designs for sequential clinical trials. *Journal of the American Statistical Association*, 72(358):382–386.
- Wei, L. J. (1978). The adaptive biased coin design for sequential experiments. *Annals of Statistics*, 6(1):92–100.
- Xu, Z., Proshan, M., and Lee, S. (2016). Validity and power considerations on hypothesis testing under minimization. *Statistics in Medicine*, 35(14):2315–2327.
- Ye, T. (2018). Testing hypotheses under covariate-adaptive randomisation and additive models. *Statistical Theory and Related Fields*, 2:96–101.
- Ye, T. and Shao, J. (2020). Robust tests for treatment effect in survival analysis under covariate-adaptive randomization. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. (To appear).
- Zelen, M. (1974). The randomization and stratification of patients to clinical trials. *Journal of Clinical Epidemiology*, 27(7):365–375.
- Zhao, W. and Ramakrishnan, V. (2016). Generalization of Wei’s urn design to unequal allocations in sequential clinical trials. *Contemporary Clinical Trials Communications*, 2:75–79.

Supplementary Material

Technical Proofs

Proof of Theorem 1

Asymptotics for $\hat{\theta}$: We start with deriving the asymptotic distribution of $\hat{\theta}$. As mentioned in the main article, the key in this proof is decomposing $\hat{\theta} - \theta$ as the sum of U and V , where

$$U = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left[\{\bar{Y}_t(z) - \bar{Y}_s(z)\} - \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} \right], \quad (\text{S1})$$

$$V = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} - \theta. \quad (\text{S2})$$

Let $\mathcal{D} = (Z_1, \dots, Z_n, I_1, \dots, I_n)$, $\mathcal{I}(A)$ be the indicator function of event A , and $n_t(z) = \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z)$ be the number of patients within stratum $Z_i = z$ and treatment t . Under (C1)-(C3), $n(z)$, $n_t(z)$, and $n_s(z)$ are functions of \mathcal{D} and the conditional expectation $E\{\bar{Y}_t(z) | \mathcal{D}\}$ is equal to $E(Y^{(t)} | Z = z)$, which implies that $E(U | \mathcal{D}) = 0$ a.s. and $E(U) = 0$. Note that

$$\begin{aligned} V &= \frac{1}{n} \sum_{z \in \mathcal{Z}} \sum_{i=1}^n \left[\mathcal{I}(Z_i = z) \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} \right] - \theta \\ &= \frac{1}{n} \sum_{i=1}^n \left[\sum_{z \in \mathcal{Z}} \mathcal{I}(Z_i = z) \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} \right] - \theta \\ &= \frac{1}{n} \sum_{i=1}^n E(Y_i^{(t)} - Y_i^{(s)} | Z_i) - \theta \end{aligned}$$

Hence, $E(V) = 0$ and consequently $E(\hat{\theta}) = \theta$, which establishes the unbiasedness of $\hat{\theta}$.

Next, we establish the asymptotic normality of $\hat{\theta}$. From the previous derivations of V , V is simply an average of independent and identically distributed terms and from the Central Limit Theorem,

$$\sqrt{n}V \xrightarrow{d} N(0, \sigma_V^2).$$

We now turn to U . Notice that I_i 's are involved in U , which results in complicated dependence and is the major difficulty in deriving the asymptotic distribution of U . A useful technique that can largely simplify the problem is to derive the distribution of U conditional on $\mathcal{D} = (Z_1, \dots, Z_n, I_1, \dots, I_n)$. We define

$$U(z) = \frac{n(z)}{n} \left[\{\bar{Y}_t(z) - \bar{Y}_s(z)\} - \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} \right]$$

so that $U = \sum_{z \in \mathcal{Z}} U(z)$. From the fact that $E\{U(z) | \mathcal{D}\} = 0$ a.s. and Lindeberg's Central Limit Theorem, we conclude that, for every z , as $n \rightarrow \infty$,

$$\frac{U(z)}{\sqrt{\text{var}\{U(z) | \mathcal{D}\}}} \mid \mathcal{D} \xrightarrow{d} N(0, 1).$$

The Lindeberg's condition will be verified at the end.

Next, we prove that $\text{cov}\{\bar{Y}_t(z) - \bar{Y}_s(z), \bar{Y}_t(z') - \bar{Y}_s(z') | \mathcal{D}\} = 0$ a.s. for $z \neq z'$. By definition this conditional covariance is equal to

$$\begin{aligned}
& \text{cov} \left[\sum_{i=1}^n \mathcal{I}(Z_i = z) \left\{ \frac{\mathcal{I}(I_i = e_t)Y_i^{(t)}}{n_t(z)} - \frac{\mathcal{I}(I_i = e_s)Y_i^{(s)}}{n_s(z)} \right\}, \right. \\
& \quad \left. \sum_{j=1}^n \mathcal{I}(Z_j = z') \left\{ \frac{\mathcal{I}(I_j = e_t)Y_j^{(t)}}{n_t(z')} - \frac{\mathcal{I}(I_j = e_s)Y_j^{(s)}}{n_s(z')} \right\} \mid \mathcal{D} \right] \\
&= \sum_{i=1}^n \sum_{j=1}^n \mathcal{I}(Z_i = z) \mathcal{I}(Z_j = z') \text{cov} \left\{ \frac{\mathcal{I}(I_i = e_t)Y_i^{(t)}}{n_t(z)} - \frac{\mathcal{I}(I_i = e_s)Y_i^{(s)}}{n_s(z)}, \right. \\
& \quad \left. \frac{\mathcal{I}(I_j = e_t)Y_j^{(t)}}{n_t(z')} - \frac{\mathcal{I}(I_j = e_s)Y_j^{(s)}}{n_s(z')} \mid \mathcal{D} \right\}
\end{aligned}$$

because $\{Z_i, i = 1, \dots, n\} \subset \mathcal{D}$. When $i = j$, $\mathcal{I}(Z_i = z)\mathcal{I}(Z_i = z') = 0$ for $z \neq z'$. When $i \neq j$, the terms are also equal to zero because $\text{cov}(Y_i^{(\ell)}, Y_j^{(m)} \mid \mathcal{D}) = 0$ a.s. for $\ell, m = 1, \dots, k$ from (C1)-(C2). From the definition of $U(z)$, this also proves that $\text{cov}\{U(z), U(z') \mid \mathcal{D}\} = 0$ a.s. for $z \neq z'$. Then, it follows from the delta method that

$$\frac{U}{\sqrt{\sum_{z \in \mathcal{Z}} \text{var}\{U(z) \mid \mathcal{D}\}}} \mid \mathcal{D} \xrightarrow{d} N(0, 1).$$

From the bounded convergence theorem, this result still holds unconditionally, i.e.,

$$\frac{U}{\sqrt{\sum_{z \in \mathcal{Z}} \text{var}\{U(z) \mid \mathcal{D}\}}} \xrightarrow{d} N(0, 1).$$

Note that

$$\begin{aligned}
& n \sum_{z \in \mathcal{Z}} \text{var}\{U(z) \mid \mathcal{D}\} \\
&= \sum_{z \in \mathcal{Z}} \frac{n^2(z)}{n} \text{var} \left\{ \frac{1}{n_t(z)} \sum_{i: Z_i = z} \mathcal{I}(I_i = e_t)Y_i^{(t)} - \frac{1}{n_s(z)} \sum_{i: Z_i = z} \mathcal{I}(I_i = e_s)Y_i^{(s)} \mid \mathcal{D} \right\} \\
&= \sum_{z \in \mathcal{Z}} \frac{n^2(z)}{n} \text{var} \left\{ \sum_{i: Z_i = z} \frac{\mathcal{I}(I_i = e_t)Y_i^{(t)}}{n_t(z)} - \frac{\mathcal{I}(I_i = e_s)Y_i^{(s)}}{n_s(z)} \mid \mathcal{D} \right\} \\
&= \sum_{z \in \mathcal{Z}} \frac{n^2(z)}{n} \sum_{i: Z_i = z} \text{var} \left\{ \frac{\mathcal{I}(I_i = e_t)Y_i^{(t)}}{n_t(z)} - \frac{\mathcal{I}(I_i = e_s)Y_i^{(s)}}{n_s(z)} \mid \mathcal{D} \right\} \\
&= \sum_{z \in \mathcal{Z}} \frac{n^2(z)}{n} \sum_{i: Z_i = z} \left\{ \frac{\mathcal{I}(I_i = e_t)\text{var}(Y_i^{(t)} \mid Z_i)}{n_t^2(z)} + \frac{\mathcal{I}(I_i = e_s)\text{var}(Y_i^{(s)} \mid Z_i)}{n_s^2(z)} \right\} \\
&= \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left\{ \frac{n(z)}{n_t(z)} \text{var}(Y^{(t)} \mid Z = z) + \frac{n(z)}{n_s(z)} \text{var}(Y^{(s)} \mid Z = z) \right\} \\
&= \sum_{z \in \mathcal{Z}} \text{pr}(Z = z) \left\{ \frac{1}{\pi_t} \text{var}(Y^{(t)} \mid Z = z) + \frac{1}{\pi_s} \text{var}(Y^{(s)} \mid Z = z) \right\} + o_p(1) \\
&= E \left\{ \frac{1}{\pi_t} \text{var}(Y^{(t)} \mid Z) + \frac{1}{\pi_s} \text{var}(Y^{(s)} \mid Z) \right\} + o_p(1) \\
&= \sigma_U^2 + o_p(1)
\end{aligned}$$

where the first equality is because $n(z)$ is a function of \mathcal{D} , the third equality is because the summands are mutually independent conditional on \mathcal{D} , the fourth equality is from (C1)-(C2), the sixth equality is because $n(z)/n = \text{pr}(Z = z) + o_p(1)$, and from (C2)-(C3), $n_t(z)/n(z) = n(z)^{-1}D_t(z) + \pi_t =$

$\pi_t + o_p(1)$ for every t . From Slutsky's theorem, this proves that as $n \rightarrow \infty$,

$$\sqrt{n}U/\sigma_U \xrightarrow{d} N(0, 1).$$

Combining the results for U and V and using the fact that U and V are uncorrelated because $E(UV) = E\{VE(U | \mathcal{D})\} = 0$, we conclude that the results for $\hat{\theta}$ hold.

For completeness, we verify the Lindeberg's condition, by rearranging $U(z)$ as

$$U(z) = \sum_{i=1}^n (K_i^{(t)} - K_i^{(s)}), \quad K_i^{(t)} = \frac{n(z)}{n} \frac{\mathcal{I}(I_i = e_t, Z_i = z)}{n_t(z)} \{Y_i^{(t)} - E(Y^{(t)} | Z_i = z)\}$$

We have already shown that $E(K_i^{(t)} - K_i^{(s)} | \mathcal{D}) = 0$ and

$$\begin{aligned} & \text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D}) \\ &= \frac{n^2(z)}{n^2 n_t^2(z)} \mathcal{I}(I_i = e_t, Z_i = z) \text{var}(Y^{(t)} | Z = z) + \frac{n^2(z)}{n^2 n_s^2(z)} \mathcal{I}(I_i = e_s, Z_i = z) \text{var}(Y^{(s)} | Z = z). \end{aligned}$$

Then, the Lindeberg's condition holds because for any $\epsilon > 0$,

$$\begin{aligned} & \sum_{i=1}^n E \left[\frac{(K_i^{(t)} - K_i^{(s)})^2}{\text{var}(U(z) | \mathcal{D})} \mathcal{I} \left\{ \frac{(K_i^{(t)} - K_i^{(s)})^2}{\text{var}(U(z) | \mathcal{D})} > \epsilon \right\} | \mathcal{D} \right] \\ &= \sum_{i=1}^n \frac{\text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D})}{\text{var}(U(z) | \mathcal{D})} E \left[\frac{(K_i^{(t)} - K_i^{(s)})^2}{\text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D})} \mathcal{I} \left\{ \frac{(K_i^{(t)} - K_i^{(s)})^2}{\text{var}(U(z) | \mathcal{D})} > \epsilon \right\} | \mathcal{D} \right] \\ &\leq \max_i E \left[\frac{(K_i^{(t)} - K_i^{(s)})^2}{\text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D})} \mathcal{I} \left\{ \frac{(K_i^{(t)} - K_i^{(s)})^2}{\text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D})} > \epsilon \frac{\text{var}(U(z) | \mathcal{D})}{\text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D})} \right\} | \mathcal{D} \right] \\ &= o(1) \end{aligned}$$

where the third line is because $\sum_{i=1}^n \text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D}) = \text{var}(U(z) | \mathcal{D})$, and the last line is because $K_i^{(t)} - K_i^{(s)} / \sqrt{\text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D})}$ has zero expectation and unit variance, and that $\max_i \text{var}(K_i^{(t)} - K_i^{(s)} | \mathcal{D}) / \text{var}(U(z) | \mathcal{D}) \leq \max(\{n_t(z)\}^{-1}, \{n_s(z)\}^{-1}) = o(1)$.

Asymptotics for $\hat{\theta}_A, \hat{\theta}_B$: To establish the results for $\hat{\theta}_A$ and $\hat{\theta}_B$, we need the following lemma about the asymptotic limits of $\hat{\beta}_t(z)$ used in $\hat{\theta}_A$ and $\hat{\beta}(z)$ used in $\hat{\theta}_B$.

Lemma 1. *Under the conditions of Theorem 1, $\hat{\beta}_t(z) = \beta_t(z) + o_p(1)$ and $\hat{\beta}(z) = \beta(z) + o_p(1)$, for every $z \in \mathcal{Z}$, where $\beta_t(z)$ and $\beta(z)$ are defined in Theorem 1.*

of Lemma 1. We prove the result for $\hat{\beta}_t(z)$. The proof for $\hat{\beta}(z)$ is analogous and omitted. The numerator of $\hat{\beta}_t(z)$ equals

$$\sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) X_i Y_i - \frac{1}{n_t(z)} \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) X_i \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) Y_i$$

Conditional on \mathcal{D} , the first term is an average of independent random variables. Assuming (C2) and existence of the second moment of $XY^{(t)}$, by the weak law of large numbers for independent random variables, we conclude that, for any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \text{pr} \left\{ \frac{1}{n} \left| \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) X_i Y_i - n_t(z) E(XY^{(t)} | Z = z) \right| \geq \epsilon \mid \mathcal{D} \right\} = 0,$$

which together with the bounded convergence theorem and $n_t(z)/n = \pi_t \text{pr}(Z = z) + o_p(1)$ by (C3)

implies that

$$\frac{1}{n_t(z)} \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) X_i Y_i = E(XY^{(t)} | Z = z) + o_p(1).$$

Similarly, we can show the result with $X_i Y_i$ replaced by X_i or Y_i and, therefore,

$$\begin{aligned} & \frac{1}{n_t(z)} \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) X_i Y_i - \frac{1}{n_t^2(z)} \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) X_i \sum_{i=1}^n \mathcal{I}(I_i = e_t, Z_i = z) Y_i \\ &= \{E(XY^{(t)} | Z = z) + o_p(1)\} - \{E(X | Z = z) + o_p(1)\} \{E(Y^{(t)} | Z = z) + o_p(1)\} \\ &= \text{cov}(X, Y^{(t)} | Z = z) + o_p(1). \end{aligned}$$

The denominator of $\hat{\beta}_t(z)$ can be treated similarly, which leads to

$$\frac{1}{n_t(z)} \sum_{i: I_i = e_t, Z_i = z} \{X_i - \bar{X}_t(z)\} \{X_i - \bar{X}_t(z)\}^T = \text{var}(X | Z = z) + o_p(1).$$

The proof is completed by using the definition of $\beta_t(z)$. □

Next, consider $\hat{\theta}_B$. Let

$$U_B = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left[\bar{Y}_t(z) - \bar{Y}_s(z) - \{\bar{X}_t(z) - \bar{X}_s(z)\}^T \beta(z) - \{E(Y^{(t)} | Z = z) - E(Y^{(s)} | Z = z)\} \right]$$

Similar to the decomposition for $\hat{\theta} - \theta$ in (S1)-(S2), we have the following decomposition for $\hat{\theta}_B - \theta$:

$$\hat{\theta}_B - \theta = U_B + V - \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{\bar{X}_1(z) - \bar{X}_0(z)\}^T \{\hat{\beta}(z) - \beta(z)\}$$

The last term is $o_p(n^{-1/2})$ because $\hat{\beta}(z) - \beta(z) = o_p(1)$ by Lemma 1 and $\bar{X}_1(z) - \bar{X}_0(z) = O_p(n^{-1/2})$ that can be shown using a similar technique in the proof of $\hat{\theta}$. To derive the asymptotic distribution of U_B , we apply the same techniques used to treat U in the proof of $\hat{\theta}$, i.e., conditioned on \mathcal{D} , U_B can be shown to be an average of independent terms so that conditioned on \mathcal{D} , U_B is asymptotically normal and, therefore, unconditionally it is also asymptotically normal. Since $E(U_B | \mathcal{D}) = 0$, it remains to find the conditional variance of U_B given \mathcal{D} , which is

$$\begin{aligned} \text{var}(\sqrt{n}U_B | \mathcal{D}) &= \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \left[\frac{n(z)}{n_t(z)} \text{var}\{Y^{(t)} - X^T \beta(z) | Z = z\} + \frac{n(z)}{n_s(z)} \text{var}\{Y^{(s)} - X^T \beta(z) | Z = z\} \right] \\ &= E \left[\frac{1}{\pi_t} \text{var}\{Y^{(t)} - X^T \beta(Z) | Z\} + \frac{1}{\pi_s} \text{var}\{Y^{(s)} - X^T \beta(Z) | Z\} \right] + o_p(1) \\ &= \sigma_B^2 + o_p(1) \end{aligned}$$

This completes the proof of $\hat{\theta}_B$.

Next, we consider $\hat{\theta}_A$. Define

$$U_A = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \sum_{\ell=t,s} (-1)^{\ell=s} \left[\bar{Y}_\ell(z) - \{\bar{X}_\ell(z) - \bar{X}(z)\}^T \beta_\ell(z) - E(Y^{(\ell)} | Z = z) \right]$$

Then,

$$\hat{\theta}_A - \theta = U_A + V - \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \sum_{\ell=t,s} (-1)^{\ell=s} \{(\bar{X}_\ell(z) - \bar{X}(z))\}^T \{\hat{\beta}_\ell(z) - \beta_\ell(z)\}$$

where the last term is $o_p(n^{-1/2})$ because $\hat{\beta}_\ell(z) - \beta_\ell(z) = o_p(1)$ by Lemma 1 and $\bar{X}_j(z) - \bar{X}(z) = O_p(n^{-1/2})$. It remains to derive the asymptotic distribution of U_A . Consider a further decomposi-

tion

$$U_A = U_{A1} + U_{A2}$$

where

$$U_{A1} = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \sum_{\ell=t,s} (-1)^{\ell=s} [\bar{Y}_\ell(z) - E(Y^{(\ell)} | Z = z) - \{\bar{X}_\ell(z) - E(X | Z = z)\}^T \beta_\ell(z)]$$

$$U_{A2} = \sum_{z \in \mathcal{Z}} \frac{n(z)}{n} \{\bar{X}(z) - E(X | Z = z)\}^T \{\beta_t(z) - \beta_s(z)\}$$

and $\text{cov}(U_{A1}, U_{A2} | \mathcal{D}) = E(U_{A1}U_{A2} | \mathcal{D}) = 0$, which can be seen from $\text{cov}\{Y_i^{(t)} - X_i\beta_t(z), X_j | Z_i = z, Z_j = z\} = 0$ a.s. for any $j, i = 1, \dots, n$, and $t = 1, \dots, k$. The asymptotic normality of U_{A1} can be derived in the same way as that for U_B with $E(U_{A1} | \mathcal{D}) = 0$ and

$$\text{var}(\sqrt{n}U_{A1} | \mathcal{D}) = E \left[\frac{1}{\pi_t} \text{var}\{Y^{(t)} - X^T \beta_t(z) | Z\} + \frac{1}{\pi_s} \text{var}\{Y^{(s)} - X^T \beta_s(z) | Z\} \right] + o_p(1).$$

Note that U_{A2} is an average of independent and identically distributed terms and its asymptotic normality follows directly from the Central Limit Theorem. Specifically, $\sqrt{n}U_{A2}$ converges in distribution to normal with mean 0 and variance $E[\{\beta_t(Z) - \beta_s(Z)\}^T \text{var}(X | Z) \{\beta_t(Z) - \beta_s(Z)\}]$. This proves the first result for $\hat{\theta}_A$ because the sum of the asymptotic variances of U_{A1} and U_{A2} is exactly σ_A^2 . The proof is completed.

Proof of Theorem 2

Define $\Sigma(z) = \text{var}(X | Z = z)$,

$$\sigma_A^2(z) = \frac{\text{var}(Y^{(t)} - X^T \beta_t(z) | Z = z)}{\pi_t} + \frac{\text{var}(Y^{(s)} - X^T \beta_s(z) | Z = z)}{\pi_s}$$

$$+ \{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\}$$

$$\sigma_U^2(z) = \frac{\text{var}(Y^{(t)} | Z = z)}{\pi_t} + \frac{\text{var}(Y^{(s)} | Z = z)}{\pi_s}$$

Then $\sigma_A^2 = \sum_{z \in \mathcal{Z}} \text{pr}(Z = z) \sigma_A^2(z)$ and $\sigma_U^2 = \sum_{z \in \mathcal{Z}} \text{pr}(Z = z) \sigma_U^2(z)$. Hence, the result for $\sigma_U^2 - \sigma_A^2$ follows from

$$\begin{aligned} & \sigma_A^2(z) - \sigma_U^2(z) \\ &= \frac{\{\beta_t(z)\}^T \Sigma(z) \beta_t(z) - 2\text{cov}(X^T \beta_t(z), Y^{(t)} | Z = z)}{\pi_t} \\ &+ \frac{\{\beta_s(z)\}^T \Sigma(z) \beta_s(z) - 2\text{cov}(X^T \beta_s(z), Y^{(s)} | Z = z)}{\pi_s} + \{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \\ &= \frac{\{\beta_t(z)\}^T \Sigma(z) \beta_t(z) - 2\{\beta_t(z)\}^T \Sigma(z) \beta_t(z)}{\pi_t} \\ &+ \frac{\{\beta_s(z)\}^T \Sigma(z) \beta_s(z) - 2\{\beta_s(z)\}^T \Sigma(z) \beta_s(z)}{\pi_s} + \{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \\ &= -\frac{\{\beta_t(z)\}^T \Sigma(z) \beta_t(z)}{\pi_t} - \frac{\{\beta_s(z)\}^T \Sigma(z) \beta_s(z)}{\pi_s} + \{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \\ &= -\frac{\{\pi_s \beta_t(z) + \pi_t \beta_s(z)\}^T \Sigma(z) \{\pi_s \beta_t(z) + \pi_t \beta_s(z)\}}{\pi_t \pi_s (\pi_t + \pi_s)} \\ &- \{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \left(\frac{1 - \pi_t - \pi_s}{\pi_t + \pi_s} \right) \end{aligned}$$

where the second equality follows from $\beta_t(z) = \Sigma(z)^{-1} \text{cov}(X, Y^{(t)} | Z = z)$. This also proves that $\sigma_A^2 \leq \sigma_U^2$, because $\Sigma(z)$ is positive definite for every z and $\pi_t + \pi_s \leq 1$. If $\sigma_A^2 = \sigma_U^2$, then we must have $\pi_s \beta_t(z) + \pi_t \beta_s(z) = 0$ and $(1 - \pi_t - \pi_s) \{\beta_t(z) - \beta_s(z)\} = 0$ for every z , which is the same as

(5).

To show the result for $\sigma_B^2 - \sigma_A^2$, note that $\sigma_B^2 = \sum_{z \in \mathcal{Z}} \text{pr}(Z = z) \sigma_B^2(z)$, where

$$\begin{aligned}
\sigma_B^2(z) &= \frac{\text{var}\{Y^{(t)} - X^T \beta(z) \mid Z = z\}}{\pi_t} + \frac{\text{var}\{Y^{(s)} - X^T \beta(z) \mid Z = z\}}{\pi_s} \\
&= \frac{\text{var}\{Y^{(t)} - X^T \beta_t(z) + X^T \beta_t(z) - X^T \beta(z) \mid Z = z\}}{\pi_t} \\
&\quad + \frac{\text{var}\{Y^{(s)} - X^T \beta_s(z) + X^T \beta_s(z) - X^T \beta(z) \mid Z = z\}}{\pi_s} \\
&= \frac{\text{var}\{Y^{(t)} - X^T \beta_t(z) \mid Z = z\} + \text{var}\{X^T \beta_t(z) - X^T \beta(z) \mid Z = z\}}{\pi_t} \\
&\quad + \frac{\text{var}\{Y^{(s)} - X^T \beta_s(z) \mid Z = z\} + \text{var}\{X^T \beta_s(z) - X^T \beta(z) \mid Z = z\}}{\pi_s}
\end{aligned}$$

where the second equality is because

$$\begin{aligned}
&\text{cov}\{Y^{(t)} - \beta_t(z)^T X, \beta_t(z)^T X - \beta(z)^T X \mid Z = z\} \\
&= \text{cov}\{Y^{(t)} - \beta_t(z)^T X, X \mid Z = z\} \{\beta_t(z) - \beta(z)\} \\
&= \{\text{cov}(Y^{(t)}, X \mid Z = z) - \beta_t(z)^T \text{var}(X \mid Z = z)\} \{\beta_t(z) - \beta(z)\} = 0
\end{aligned}$$

Then,

$$\begin{aligned}
\sigma_A^2(z) - \sigma_B^2(z) &= \{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \\
&\quad - \frac{\{\beta_t(z) - \beta(z)\}^T \Sigma(z) \{\beta_t(z) - \beta(z)\}}{\pi_t} - \frac{\{\beta_s(z) - \beta(z)\}^T \Sigma(z) \{\beta_s(z) - \beta(z)\}}{\pi_s}
\end{aligned}$$

In order to show that $\sigma_A^2(z) - \sigma_B^2(z) \leq 0$ for every z , we prove a stronger statement: for each given z , it is true that for any $\tilde{\beta}(z)$,

$$\begin{aligned}
&\{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \\
&\quad - \frac{\{\beta_t(z) - \tilde{\beta}(z)\}^T \Sigma(z) \{\beta_t(z) - \tilde{\beta}(z)\}}{\pi_t} - \frac{\{\beta_s(z) - \tilde{\beta}(z)\}^T \Sigma(z) \{\beta_s(z) - \tilde{\beta}(z)\}}{\pi_s} \leq 0. \tag{S3}
\end{aligned}$$

As a consequence, setting $\tilde{\beta}(z)$ as $\beta(z) = \sum_t \pi_t \beta_t(z)$, the statement in (S3) also holds. This proves $\sigma_A^2(z) - \sigma_B^2(z) \leq 0$.

In what follows, we prove the claim in (S3). For each given z , the gradient of the left hand side of (S3) is

$$-2 \left[\frac{\{\tilde{\beta}(z) - \beta_t(z)\}^T \Sigma(z)}{\pi_t} + \frac{\{\tilde{\beta}(z) - \beta_s(z)\}^T \Sigma(z)}{\pi_s} \right],$$

which equals zero when $\tilde{\beta}(z) = \{\pi_s \beta_t(z) + \pi_t \beta_s(z)\} / (\pi_t + \pi_s)$. This is also the unique solution from the positive definiteness of $\Sigma(z)$. It is also easy to see that the Hessian of the left hand side of (S3) is negative definite, which means that $\tilde{\beta}(z) = \{\pi_s \beta_t(z) + \pi_t \beta_s(z)\} / (\pi_t + \pi_s)$ is the global and unique maximizer of the left hand side of (S3). The statement in (S3) is true because when evaluated at

$\tilde{\beta}(z) = \{\pi_s \beta_t(z) + \pi_t \beta_s(z)\} / (\pi_t + \pi_s)$, the left hand side of (S3) equals

$$\begin{aligned} & \{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \\ & - \left\{ \beta_t(z) - \frac{\pi_s \beta_t(z) + \pi_t \beta_s(z)}{\pi_t + \pi_s} \right\}^T \Sigma(z) \left\{ \beta_t(z) - \frac{\pi_s \beta_t(z) + \pi_t \beta_s(z)}{\pi_t + \pi_s} \right\} \frac{1}{\pi_t} \\ & - \left\{ \beta_s(z) - \frac{\pi_s \beta_t(z) + \pi_t \beta_s(z)}{\pi_t + \pi_s} \right\}^T \Sigma(z) \left\{ \beta_s(z) - \frac{\pi_s \beta_t(z) + \pi_t \beta_s(z)}{\pi_t + \pi_s} \right\} \frac{1}{\pi_s} \\ & = -\{\beta_t(z) - \beta_s(z)\}^T \Sigma(z) \{\beta_t(z) - \beta_s(z)\} \left(\frac{1 - \pi_t - \pi_s}{\pi_t + \pi_s} \right) \leq 0 \end{aligned}$$

This completes the proof for $\sigma_A^2 \leq \sigma_B^2$, where the equality holds if and only if $\{\beta_t(z) - \beta_s(z)\}(1 - \pi_t - \pi_s) = 0$ and $\sum_{\ell=1}^k \pi_\ell \beta_\ell(z) = \{\pi_s \beta_t(z) + \pi_t \beta_s(z)\} / (\pi_t + \pi_s)$ for every z , which is the same as (6).

Proof of the Consistency of Variance Estimators

First, we prove the consistency of estimators $\hat{\sigma}_U^2, \hat{\sigma}_V^2, \hat{\sigma}_A^2$ and $\hat{\sigma}_B^2$. For $\hat{\sigma}_U^2$, following the same arguments as in the proof of Lemma 1, we can show that

$$S_t^2(z) = \text{var}(Y^{(t)} \mid Z = z) + o_p(1)$$

From $n(z)/n = \text{pr}(Z = z) + o_p(1)$, we conclude that $\hat{\sigma}_U^2 = \sigma_U^2 + o_p(1)$ by continuous mapping theorem. For $\hat{\sigma}_V^2$, following the same arguments as in the proof of Lemma 1, we conclude that

$$\bar{Y}_t(z) = E(Y^{(t)} \mid Z = z) + o_p(1),$$

which together with the fact that $\hat{\theta} = \theta + o_p(1)$ implied by Theorem 1, we obtain that

$$\begin{aligned} \hat{\sigma}_V^2 &= \sum_{z \in \mathcal{Z}} \text{pr}(Z = z) \left\{ E(Y^{(t)} - Y^{(s)} \mid Z = z) \right\}^2 - \theta^2 + o_p(1) \\ &= E \left[\left\{ E(Y^{(t)} - Y^{(s)} \mid Z) \right\}^2 \right] - \{E(Y^{(t)} - Y^{(s)})\}^2 + o_p(1) \\ &= \sigma_V^2 + o_p(1). \end{aligned}$$

The proof for the consistency of $\hat{\sigma}_A^2$ and $\hat{\sigma}_B^2$ with $\hat{\beta}_t(z), \hat{\beta}(z)$ respectively replaced with $\beta_t(z), \beta(z)$ is analogous and is omitted. From this, the consistency of $\hat{\sigma}_A^2$ and $\hat{\sigma}_B^2$ can be established using Lemma 1.

Additional Simulation Results

Tables S1 and S2 report the bias, standard deviation (SD), average estimated SD (SE), and coverage probability (CP) of asymptotic 95% confidence interval, estimate $\pm 1.96 \text{ SE}$, of $\hat{\theta}$, $\hat{\theta}_A$, and $\hat{\theta}_B$ for cases I-III under stratified permuted block randomization and urn design, respectively, based on 2,000 simulation runs.

Table S1: Bias, standard deviation (SD), average estimated SD (SE), and coverage probability (CP) of 95% asymptotic confidence interval under Stratified Permuted Block Randomization for cases I-III (block size = 4 for 1:1 allocation and block size = 6 for 1:2 allocation)

n	case	Z	estimator	treatment allocation 1:1				treatment allocation 1:2			
				bias	SD	SE	CP	bias	SD	SE	CP
500	I	X_1	$\hat{\theta}$	0.0042	0.2005	0.2000	0.9495	0.0011	0.2161	0.2124	0.9500
			$\hat{\theta}_B$	0.0016	0.0887	0.0894	0.9515	-0.0001	0.0967	0.0948	0.9480
			$\hat{\theta}_A$	0.0016	0.0887	0.0894	0.9520	0.0000	0.0969	0.0947	0.9495
		X_1, D_2	$\hat{\theta}$	-0.0052	0.1473	0.1467	0.9455	0.0020	0.1552	0.1556	0.9530
			$\hat{\theta}_B$	-0.0004	0.0907	0.0894	0.9430	0.0014	0.0966	0.0948	0.9445
			$\hat{\theta}_A$	-0.0003	0.0908	0.0894	0.9430	0.0015	0.0964	0.0946	0.9440
		X_1, D_4	$\hat{\theta}$	0.0024	0.1121	0.1152	0.9520	0.0009	0.1256	0.1220	0.9420
			$\hat{\theta}_B$	0.0009	0.0884	0.0895	0.9430	-0.0026	0.0992	0.0947	0.9360
			$\hat{\theta}_A$	0.0009	0.0885	0.0895	0.9465	-0.0025	0.1002	0.0944	0.9320
	II	X_1	$\hat{\theta}$	0.0004	0.2168	0.2191	0.9505	0.0030	0.2329	0.2304	0.9400
			$\hat{\theta}_B$	0.0002	0.2173	0.2184	0.9485	0.0029	0.2600	0.2543	0.9415
			$\hat{\theta}_A$	0.0005	0.2169	0.2190	0.9510	0.0034	0.2247	0.2214	0.9405
		X_1, D_2	$\hat{\theta}$	0.0039	0.2199	0.2190	0.9500	0.0006	0.2275	0.2255	0.9405
			$\hat{\theta}_B$	0.0037	0.2207	0.2177	0.9500	0.0004	0.2393	0.2350	0.9410
			$\hat{\theta}_A$	0.0038	0.2203	0.2189	0.9490	0.0007	0.2241	0.2216	0.9440
		X_1, D_4	$\hat{\theta}$	0.0020	0.2131	0.2193	0.9570	-0.0046	0.2246	0.2230	0.9465
			$\hat{\theta}_B$	0.0021	0.2140	0.2176	0.9515	-0.0058	0.2314	0.2253	0.9415
			$\hat{\theta}_A$	0.0017	0.2130	0.2190	0.9560	-0.0034	0.2239	0.2211	0.9405
	III	X_1	$\hat{\theta}$	0.0046	0.1737	0.1732	0.9500	0.0028	0.1674	0.1666	0.9465
			$\hat{\theta}_B$	0.0048	0.1485	0.1477	0.9490	0.0037	0.1688	0.1656	0.9395
			$\hat{\theta}_A$	0.0049	0.1484	0.1479	0.9500	0.0057	0.1550	0.1532	0.9455
		X_1, D_2	$\hat{\theta}$	0.0028	0.1615	0.1593	0.9450	0.0016	0.1620	0.1597	0.9455
			$\hat{\theta}_B$	0.0083	0.1501	0.1471	0.9425	0.0044	0.1626	0.1579	0.9380
			$\hat{\theta}_A$	0.0064	0.1499	0.1475	0.9440	0.0039	0.1562	0.1526	0.9380
		X_1, D_4	$\hat{\theta}$	0.0033	0.1500	0.1527	0.9540	-0.0012	0.1565	0.1560	0.9500
			$\hat{\theta}_B$	0.0062	0.1450	0.1469	0.9515	0.0001	0.1586	0.1540	0.9410
			$\hat{\theta}_A$	0.0031	0.1448	0.1475	0.9520	-0.0006	0.1558	0.1521	0.9400
100	I	X_1	$\hat{\theta}$	-0.0019	0.4505	0.4484	0.9460	-0.0087	0.4690	0.4741	0.9500
			$\hat{\theta}_B$	-0.0004	0.2004	0.1977	0.9350	0.0013	0.2132	0.2094	0.9420
			$\hat{\theta}_A$	-0.0005	0.2004	0.1978	0.9365	0.0013	0.2147	0.2084	0.9395
		X_1, D_2	$\hat{\theta}$	-0.0072	0.3305	0.3314	0.9500	-0.0043	0.3414	0.3498	0.9500
			$\hat{\theta}_B$	-0.0003	0.2036	0.1983	0.9415	0.0018	0.2184	0.2094	0.9340
			$\hat{\theta}_A$	-0.0017	0.2059	0.1997	0.9380	0.0054	0.2387	0.2113	0.9165
	II	X_1	$\hat{\theta}$	0.0151	0.4930	0.4901	0.9500	0.0167	0.5103	0.5140	0.9445
			$\hat{\theta}_B$	0.0136	0.4992	0.4827	0.9430	0.0230	0.5728	0.5607	0.9405
			$\hat{\theta}_A$	0.0149	0.4945	0.4892	0.9470	0.0152	0.4938	0.4929	0.9455
		X_1, D_2	$\hat{\theta}$	0.0112	0.4953	0.4916	0.9475	0.0209	0.5070	0.5035	0.9435
			$\hat{\theta}_B$	0.0085	0.5059	0.4781	0.9320	0.0223	0.5428	0.5130	0.9310
			$\hat{\theta}_A$	0.0102	0.4976	0.4907	0.9415	0.0226	0.5105	0.4951	0.9410
	III	X_1	$\hat{\theta}$	0.0033	0.3877	0.3869	0.9485	0.0016	0.3668	0.3714	0.9460
			$\hat{\theta}_B$	0.0110	0.3325	0.3262	0.9465	0.0175	0.3692	0.3651	0.9365
			$\hat{\theta}_A$	0.0116	0.3306	0.3287	0.9490	0.0212	0.3413	0.3390	0.9380
		X_1, D_2	$\hat{\theta}$	-0.0006	0.3642	0.3578	0.9430	0.0071	0.3547	0.3575	0.9485
			$\hat{\theta}_B$	0.0163	0.3404	0.3241	0.9285	0.0240	0.3653	0.3456	0.9375
			$\hat{\theta}_A$	0.0079	0.3394	0.3296	0.9365	0.0256	0.3701	0.3400	0.9330

Table S2: Bias, standard deviation (SD), average estimated SD (SE), and coverage probability (CP) of 95% asymptotic confidence interval under Stratified Urn Design for cases I-III

n	case	Z	estimator	treatment allocation 1:1				treatment allocation 1:2			
				bias	SD	SE	CP	bias	SD	SE	CP
500	I	X_1	$\hat{\theta}$	-0.0095	0.1984	0.2000	0.9465	0.0012	0.2119	0.2121	0.9470
			$\hat{\theta}_B$	-0.0052	0.0866	0.0893	0.9585	-0.0012	0.0957	0.0948	0.9455
			$\hat{\theta}_A$	-0.0052	0.0867	0.0893	0.9580	-0.0013	0.0957	0.0947	0.9445
		X_1, D_2	$\hat{\theta}$	0.0001	0.1486	0.1466	0.9430	0.0049	0.1560	0.1554	0.9510
			$\hat{\theta}_B$	-0.0004	0.0902	0.0894	0.9405	0.0020	0.0948	0.0947	0.9440
			$\hat{\theta}_A$	-0.0004	0.0902	0.0894	0.9420	0.0022	0.0953	0.0945	0.9435
		X_1, D_4	$\hat{\theta}$	-0.0001	0.1160	0.1150	0.9470	0.0008	0.1216	0.1220	0.9520
			$\hat{\theta}_B$	-0.0019	0.0927	0.0893	0.9390	-0.0013	0.0966	0.0947	0.9445
			$\hat{\theta}_A$	-0.0018	0.0930	0.0894	0.9390	-0.0017	0.0993	0.0947	0.9405
	II	X_1	$\hat{\theta}$	0.0037	0.2196	0.2192	0.9510	-0.0040	0.2331	0.2305	0.9480
			$\hat{\theta}_B$	0.0033	0.2203	0.2186	0.9495	-0.0046	0.2596	0.2544	0.9465
			$\hat{\theta}_A$	0.0034	0.2196	0.2191	0.9495	-0.0034	0.2230	0.2214	0.9475
		X_1, D_2	$\hat{\theta}$	0.0083	0.2200	0.2191	0.9530	-0.0012	0.2261	0.2253	0.9450
			$\hat{\theta}_B$	0.0085	0.2216	0.2180	0.9465	-0.0015	0.2382	0.2346	0.9450
			$\hat{\theta}_A$	0.0082	0.2205	0.2190	0.9480	0.0001	0.2227	0.2213	0.9455
		X_1, D_4	$\hat{\theta}$	0.0071	0.2214	0.2193	0.9475	0.0068	0.2246	0.2230	0.9460
			$\hat{\theta}_B$	0.0080	0.2226	0.2176	0.9450	0.0068	0.2306	0.2254	0.9415
			$\hat{\theta}_A$	0.0068	0.2223	0.2190	0.9465	0.0066	0.2243	0.2213	0.9440
	III	X_1	$\hat{\theta}$	-0.0037	0.1724	0.1731	0.9470	-0.0015	0.1659	0.1666	0.9530
			$\hat{\theta}_B$	-0.0003	0.1480	0.1477	0.9495	-0.0014	0.1694	0.1657	0.9385
			$\hat{\theta}_A$	-0.0002	0.1478	0.1479	0.9525	0.0008	0.1548	0.1533	0.9475
		X_1, D_2	$\hat{\theta}$	0.0029	0.1594	0.1593	0.9510	0.0013	0.1596	0.1594	0.9465
			$\hat{\theta}_B$	0.0052	0.1492	0.1470	0.9495	0.0029	0.1588	0.1576	0.9420
			$\hat{\theta}_A$	0.0031	0.1487	0.1474	0.9510	0.0028	0.1530	0.1523	0.9420
		X_1, D_4	$\hat{\theta}$	0.0012	0.1534	0.1525	0.9455	0.0032	0.1569	0.1558	0.9515
			$\hat{\theta}_B$	0.0049	0.1508	0.1468	0.9445	0.0060	0.1572	0.1539	0.9460
			$\hat{\theta}_A$	0.0014	0.1510	0.1473	0.9455	0.0039	0.1554	0.1521	0.9520
100	I	X_1	$\hat{\theta}$	0.0216	0.4468	0.4487	0.9470	-0.0132	0.4801	0.4757	0.9460
			$\hat{\theta}_B$	0.0012	0.2033	0.1981	0.9365	-0.0070	0.2115	0.2103	0.9495
			$\hat{\theta}_A$	0.0014	0.2034	0.1982	0.9390	-0.0073	0.2145	0.2093	0.9420
		X_1, D_2	$\hat{\theta}$	-0.0102	0.3299	0.3306	0.9465	-0.0070	0.3562	0.3497	0.9425
			$\hat{\theta}_B$	-0.0023	0.2073	0.1980	0.9380	-0.0051	0.2200	0.2096	0.9295
			$\hat{\theta}_A$	-0.0020	0.2137	0.1999	0.9340	-0.0352	0.7085	0.2280	0.9015
	II	X_1	$\hat{\theta}$	-0.0076	0.4882	0.4903	0.9460	-0.0131	0.5116	0.5141	0.9430
			$\hat{\theta}_B$	-0.0073	0.4993	0.4841	0.9380	-0.0143	0.5838	0.5617	0.9400
			$\hat{\theta}_A$	-0.0072	0.4890	0.4895	0.9475	-0.0159	0.4929	0.4938	0.9440
		X_1, D_2	$\hat{\theta}$	-0.0103	0.4869	0.4903	0.9485	-0.0113	0.5099	0.5029	0.9420
			$\hat{\theta}_B$	-0.0121	0.4990	0.4776	0.9375	-0.0090	0.5470	0.5125	0.9270
			$\hat{\theta}_A$	-0.0108	0.4908	0.4895	0.9450	-0.0435	0.8393	0.5090	0.9325
	III	X_1	$\hat{\theta}$	0.0057	0.3864	0.3869	0.9445	-0.0174	0.3722	0.3726	0.9475
			$\hat{\theta}_B$	0.0038	0.3348	0.3270	0.9380	-0.0071	0.3754	0.3658	0.9425
			$\hat{\theta}_A$	0.0040	0.3314	0.3291	0.9420	0.0006	0.3435	0.3396	0.9420
		X_1, D_2	$\hat{\theta}$	-0.0104	0.3566	0.3569	0.9505	-0.0078	0.3625	0.3569	0.9445
			$\hat{\theta}_B$	0.0067	0.3369	0.3234	0.9310	0.0082	0.3652	0.3451	0.9310
			$\hat{\theta}_A$	-0.0012	0.3369	0.3289	0.9400	-0.0388	1.0589	0.3615	0.9225