SHIMURA CURVES IN THE PRYM LOCI OF RAMIFIED DOUBLE COVERS

PAOLA FREDIANI AND GIAN PAOLO GROSSELLI

ABSTRACT. We study Shimura curves of PEL type in the space of polarised abelian varieties A_p^{δ} generically contained in the ramified Prym locus. We generalise to ramified double covers, the construction done in [10] in the unramified case and in the case of two ramification points. Namely, we construct families of double covers which are compatible with a fixed group action on the base curve. We only consider the case of one-dimensional families and where the quotient of the base curve by the group is \mathbb{P}^1 . Using computer algebra we obtain 184 Shimura curves contained in the (ramified) Prym loci.

1. INTRODUCTION

Let $\mathsf{R}_{g,b}$ be the moduli space parametrising isomorphism classes of triples $[(C, \eta, B)]$ where C is a smooth complex projective curve of genus g, B is a reduced effective divisor of degree b on C and η is a line bundle on C such that $\eta^2 = \mathcal{O}_C(B)$. To such data it is associated a double cover of $C, f : \tilde{C} \to C$ branched on B.

The Prym variety associated to $[(C, \eta, B)]$ is the connected component containing the origin of the kernel of the norm map $\operatorname{Nm}_f : J\tilde{C} \to JC$. For b > 0, ker Nm_f is connected. It is a polarised abelian variety of dimension $g - 1 + \frac{b}{2}$, denoted by $P(C, \eta, B)$ or equivalently $P(\tilde{C}, C)$.

Let Ξ be the restriction on $P(\tilde{C}, C)$ of the principal polarisation on $J\tilde{C}$. For b > 0 the polarisation Ξ is of type $\delta = (1, \ldots, 1, \underbrace{2, \ldots, 2}_{g \text{ times}})$. If b = 0 or $2, \Xi$ is twice a principal polarisation and we endow $P(\tilde{C}, C)$

with this principal polarisation.

This defines the Prym map

$$\mathsf{P}_{g,b}:\mathsf{R}_{g,b}\longrightarrow\mathsf{A}_{g-1+\frac{b}{2}}^{\delta},\quad [(C,\eta,B)]\longmapsto [(P(C,\eta,B),\Xi)],$$

where $\mathsf{A}_{g-1+\frac{b}{2}}^{\delta}$ is the moduli space of abelian varieties of dimension $g-1+\frac{b}{2}$ with a polarization of type δ . We define the Prym locus as the closure in $\mathsf{A}_{g-1+\frac{b}{2}}^{\delta}$ of the image of the map $\mathsf{P}_{g,b}$.

The Prym map $\mathsf{P}_{g,b}$ is generically finite, if and only if $\dim \mathsf{R}_{g,b} \leq \dim \mathsf{A}_{g-1+\frac{b}{2}}^{\delta}$, (see [21]). This holds if: either $b \geq 6$ and $g \geq 1$, or b = 4 and $g \geq 3$, b = 2 and $g \geq 5$, b = 0 and $g \geq 6$. If b = 0 the Prym map is generically injective for $g \geq 7$ ([15], [20]). If r > 0, Marcucci and Pirola [23], and, for the missing cases, Marcucci and Naranjo [24] and Naranjo and Ortega [30] have proved the generic injectivity in all the cases except for b = 4, g = 3, which was studied by Nagaraj and Ramanan, and Bardelli, Ciliberto and Verra ([29], [2]) and for which the degree of the Prym map is 3. Recently, a global Torelli theorem was proved for all g and $b \geq 6$ ([19] for g = 1 and [31] for all g).

In [10] an analogous question to the Coleman-Oort conjecture was formulated for the classical Prym maps and for $P_{g,2}$. Namely, the authors asked whether there exist Shimura subvarieties of A_g that are generically contained in the Prym loci $P_{g+1,0}(R_{g+1,0})$ and $P_{g,2}(R_{g,2})$ for g sufficiently high. A subvariety $Z \subset A_g$ is said to be generically contained in the Prym locus $P_{g+1,0}(R_{g+1,0})$ if $Z \subset \overline{P_{g+1,0}(R_{g+1,0})}$, $Z \cap P_{g+1,0}(R_{g+1,0}) \neq \emptyset$ and Z intersects the locus of irreducible principally polarized abelian varieties. The same applies for $P_{g,2}(R_{g,2})$. In [10] many examples of Shimura curves generically contained in these Prym loci were found using families of Galois covers of \mathbb{P}^1 and computations made with MAGMA. All examples found are contained in A_g for $g \leq 12$.

In this paper we address a similar question for all ramified Prym maps $P_{g,b}$. In fact in [7] and [8] the second fundamental form of the Prym map $P_{g,b}$ for b = 0, and for b > 0 was studied to give an upper bound for the dimension of a germ of a totally geodesic submanifold, and hence of a Shimura subvariety of $A_{g-1+\frac{b}{2}}^{\delta}$ contained in the Prym locus.

The authors were partially supported MIUR PRIN 2017 "Moduli spaces and Lie Theory", by MIUR, Programma Dipartimenti di Eccellenza (2018-2022) - Dipartimento di Matematica "F. Casorati", Università degli Studi di Pavia and by INdAM (GNSAGA).

The expression of the second fundamental form of the Prym maps is similar to the one of the Torelli map studied in [11], [9], [17], [14], [16] and we expect that for high values of p, there should not exist totally geodesic subvarieties of A_p^{δ} generically contained in the (ramified) Prym loci.

In this paper we use a method which is similar to the one used in [10] to find examples of Shimura curves contained in the (ramified) Prym loci. The idea is to find families of Galois covers $\psi_t : \tilde{C}_t \to \tilde{C}_t/\tilde{G} \cong \mathbb{P}^1$, with 4 branch points and such that the group \tilde{G} admits a central involution $\sigma \in \tilde{G}$. If we denote by G the quotient $\tilde{G}/\langle \sigma \rangle$, there exists a factorisation of the map ψ_t as the composition $\pi_t \circ f_t$, where $f_t : \tilde{C}_t \to C_t = \tilde{C}_t/\langle \sigma \rangle$ is a double cover and $\pi_t : C_t \to C_t/G \cong \tilde{C}_t/\tilde{G} \cong \mathbb{P}^1$. Then we give a sufficient condition that ensures that the Prym varieties $P(\tilde{C}_t, C_t)$ of the double covers $f_t : \tilde{C}_t \to C_t$ yield a Shimura curve in $A^{\delta}_{\tilde{g}-g}$, where \tilde{g} (resp. g) denotes the genus of \tilde{C}_t (resp. C_t). This is done in 3.2. The condition is that the codifferential of $\mathsf{P}_{g,b}$ at a generic point $[(\tilde{C}_t, C_t)]$ of the family is an isomorphism. If the double cover $\tilde{C}_t \to C_t$ corresponds to the point $[(C_t, \eta_t, B_t)] \in \mathsf{R}_{g,b}$, the codifferential of $\mathsf{P}_{g,b}$ is the multiplication map $m : (S^2 H^0(C_t, K_{C_t} \otimes \eta_t))^{\tilde{G}} \longrightarrow H^0(C_t, K^2_{C_t}(B_t))^{\tilde{G}}$.

Since the cover ψ_t is branched at 4 points, we first require that $(S^2 H^0(C_t, K_{C_t} \otimes \eta_t))^{\tilde{G}}$ is one dimensional (condition (A) in section 3). Since the Prym map may have positive dimensional fibres, we also need to require that the map m is non zero (condition (B) in section 3).

In [31] and [19] it is proven that $\mathsf{P}_{g,b}$ is an embedding for all $b \ge 6$ and for all g > 0, hence if $b \ge 6$, and g > 0 condition (A) implies condition (B).

Once condition (A) is true, a sufficient condition ensuring (B) is that $(S^2 H^0(C_t, K_{C_t} \otimes \eta_t))^{\tilde{G}}$ is generated by decomposable tensors (condition (B1) in section 3). This happens for example if the group \tilde{G} is abelian.

If we have (A), another condition that implies (B) is the following (condition (B2)): we require that the Prym variety decomposes up to isogeny as $P(\tilde{C}_t, C_t) \sim A \times JC'_t$, where A is a fixed abelian variety and JC'_t is the jacobian of a curve $C'_t = \tilde{C}_t/N$ defined as a quotient of \tilde{C}_t by a normal subgroup $N \triangleleft \tilde{G}$, and the family of Galois covers $C'_t \rightarrow \mathbb{P}^1 = C'_t/(G/N)$ satisfies condition (*) of [12] (hence it yields a Shimura curve).

Using a MAGMA ([22]), that is explained in the Appendix and is available at:

http://www-dimat.unipv.it/grosselli/publ/

we find many examples of such families satisfying conditions (A) and (B), hence yielding Shimura curves generically contained in the Prym loci $\mathsf{P}_{g,b}(\mathsf{R}_{g,b}) \subset \mathsf{A}_p^\delta$, for $0 \le g \le 13, 0 \le b \le 16$ and $2 \le \tilde{g} \le 31$ $(p = \tilde{g} - g)$. These computations are summarised in Table 1 in the Appendix. The highest value of $p = \tilde{g} - g$ that we find in Table 1 is 18. We recovered all the examples found in [10] and also, for g = 0, some of the examples of [12] (the ones with an action of a group containing a central involution).

Notice that in the unramified case, Mohajer [25] showed that families of Galois covers $\tilde{C}_t \to \tilde{C}_t/\tilde{G} \cong \mathbb{P}^1$, when \tilde{G} is abelian, do not give rise to high dimensional Shimura subvarieties of A_g contained in the Prym locus.

The main result is summarised in the following

Theorem 1.1. There are 184 families of (ramified) Pryms yielding Shimura curves of $A_{\tilde{g}-g}^{\delta}$ for $\tilde{g}-g \leq 18$, where $0 \leq g \leq 13$, $0 \leq b \leq 16$ and $2 \leq \tilde{g} \leq 31$.

In Table 2 we also collected all the families that we found satisfying condition (A). We performed calculations for \tilde{g} up to 7. Note that after genus 49 there are no other data satisfying condition (A).

In section 4 we described some of the examples and we explicitly showed that they satisfy condition (B).

The paper is organised as follows: in section 2 we recall some basic facts on Prym varieties and on special (or Shimura) subvarieties of A_g . In section 3 we explain our construction of special subvarieties contained in the Prym loci. In section 4 we explicitly describe some of the examples that we found and we show that they satisfy conditions (A) and (B). In the Appendix we explain the MAGMA script and we give Tables 1 and 2 that summarise all the examples that we found.

Acknowledgements

We would like to thank Diego Conti, Alessandro Ghigi and Roberto Pignatelli for interesting discussions on the MAGMA script. We thank Bert van Geemen for a helpful correspondence on one of the examples.

2. Preliminaries on Prym varieties and on special subvarieties of A_q

Denote by $\mathsf{R}_{g,b}$ the moduli space parametrising isomorphism classes of triples $[(C, \eta, B)]$ where C is a smooth complex projective curve of genus g, B is a reduced effective divisor of degree b on C and η is a line bundle on C such that $\eta^2 = \mathcal{O}_C(B)$. To such data it is associated a double cover of $C, f : \tilde{C} \to C$ branched on B, with $\tilde{C} = \operatorname{Spec}(\mathcal{O}_C \oplus \eta^{-1})$.

The Prym variety associated to $[(C, \eta, B)]$ is the connected component containing the origin of the kernel of the norm map $\operatorname{Nm}_f : J\tilde{C} \to JC$. For b > 0, ker Nm_f is connected. It is a polarised abelian variety of dimension $g - 1 + \frac{b}{2}$, denoted by $P(C, \eta, B)$ or equivalently $P(\tilde{C}, C)$.

Let Ξ be the restriction on $P(\tilde{C}, C)$ of the principal polarisation on $J\tilde{C}$. For b > 0 polarisation Ξ is of type $\delta = (1, \ldots, 1, \underbrace{2, \ldots, 2}_{a \text{ times}})$. If b = 0 or 2 then Ξ is twice a principal polarisation and we endow

 $P(\tilde{C}, C)$ with this principal polarisation.

Consider the Prym map

$$\mathsf{P}_{g,b}:\mathsf{R}_{g,b}\longrightarrow\mathsf{A}_{g-1+\frac{b}{2}}^{\delta},\ [(C,\eta,B)]\longmapsto[(P(C,\eta,B),\Xi)],$$

where $\mathsf{A}_{g-1+\frac{b}{2}}^{\delta}$ is the moduli space of abelian varieties of dimension $g-1+\frac{b}{2}$ with a polarization of type δ . We define the Prym locus as the closure in $\mathsf{A}_{g-1+\frac{b}{2}}^{\delta}$ of the image of the map $\mathsf{P}_{g,b}$.

The codifferential of $\mathsf{P}_{q,b}$ at a generic point $[(C,\eta,B)]$ is given by the multiplication map

(2.1)
$$(d\mathsf{P}_{g,b})^* : S^2 H^0(C, K_C \otimes \eta) \to H^0(C, K_C^2(B))$$

which is known to be surjective ([21]), therefore $P_{g,b}$ is generically finite, if and only if

$$\dim \mathsf{R}_{g,b} \leq \dim \mathsf{A}_{g-1+\frac{b}{2}}^{\delta}$$

This holds if: either $b \ge 6$ and $g \ge 1$, or b = 4 and $g \ge 3$, b = 2 and $g \ge 5$, b = 0 and $g \ge 6$.

If b = 0 the Prym map is generically injective for $g \ge 7$ ([15], [20]). If b > 0, in [23], [24], [30], it is proved the generic injectivity in all the cases except for b = 4, g = 3, which was previously studied in [29], [2] and for which the degree of the Prym map is 3. Recently, a global Torelli theorem was proved for all g and $b \ge 6$ ([19] for g = 1 and [31] for all g).

2.1. Consider a rank 2g lattice $\Lambda \cong \mathbb{Z}^{2g}$ and an alternating form $Q : \Lambda \times \Lambda \to \mathbb{Z}$ of type $\delta = (1 \dots, 1, 2, \dots, 2)$. There exists a basis of Λ such that the corresponding matrix is

$$\begin{pmatrix} 0 & \Delta_g \\ -\Delta_g & 0 \end{pmatrix},$$

where Δ_g is the diagonal matrix whose entries are $(1, \ldots, 1, 2, \ldots, 2)$. Set $U := \Lambda \otimes \mathbb{R}$. The Siegel space is defined as follows

$$\mathfrak{S}(U,Q) := \{ J \in \mathrm{GL}(U) : J^2 = -I, J^*Q = Q, Q(x,Jx) > 0, \ \forall x \neq 0 \}.$$

The symplectic group $\operatorname{Sp}(\Lambda, Q)$ of the form Q acts on the Siegel space $\mathfrak{S}(U, Q)$ by conjugation and this action is properly discontinuous. The moduli space of abelian varieties of dimension g with a polarisation of type δ is the quotient $\operatorname{A}_g^{\delta} = \operatorname{Sp}(\Lambda, Q) \setminus \mathfrak{S}(U, Q)$. This is a complex analytic orbifold and also a smooth algebraic stack. We will consider $\operatorname{A}_g^{\delta}$ with the orbifold structure. To an element $J \in \mathfrak{S}(U,Q)$ we associate the real torus $U/\Lambda \cong \mathbb{R}^{2g}/\mathbb{Z}^{2g}$ endowed with the complex structure J and the polarization Q. This gives a polarised abelian variety that we denote by A_J . On $\mathfrak{S}(U,Q)$ there is a natural variation of rational Hodge structure, whose local system is $\mathfrak{S}(U,Q) \times \mathbb{Q}^{2g}$, and that corresponds to the Hodge decomposition of \mathbb{C}^{2g} in $\pm i$ eigenspaces for J. This gives a variation of Hodge structure on $\operatorname{A}_g^{\delta}$ in the orbifold sense.

2.2. A special or Shimura subvariety $Z \subseteq A_g^{\delta}$ is by definition a Hodge locus of the natural variation of Hodge structure on A_g^{δ} described above. For the definition of Hodge loci we refer to §2.3 in [27]. Special subvarieties contain a dense set of CM points and they are totally geodesic [27, §3.4(b)]. Conversely an algebraic totally geodesic subvariety that contains a CM point is a special subvariety ([28], [26, Thm. 4.3]).

Let us recall the definition of special subvarieties of PEL type (see [27, §3.9]). Given $J \in \mathfrak{S}(U, Q)$, set $\operatorname{End}_{\mathbb{Q}}(A_J) := \{f \in \operatorname{End}(\mathbb{Q}^{2g}) : Jf = fJ\}.$ Fix a point $J_0 \in \mathfrak{S}(U,Q)$ and set $D := \operatorname{End}_{\mathbb{Q}}(A_{J_0})$. The *PEL type* special subvariety $\mathsf{Z}(D)$ is defined as the image in A_g^{δ} of the connected component of the set $\{J \in \mathfrak{S}(U,Q) : D \subseteq \operatorname{End}_{\mathbb{Q}}(A_J)\}$ that contains J_0 . By definition $\mathsf{Z}(D)$ is irreducible.

If $G \subseteq \text{Sp}(\Lambda, Q)$ is a finite subgroup, denote by $\mathfrak{S}(U, Q)^G$ the subset of $\mathfrak{S}(U, Q)$ of fixed points by the action of G. Set

(2.2)
$$D_G := \{ f \in \operatorname{End}_{\mathbb{Q}}(\Lambda \otimes \mathbb{Q}) : Jf = fJ, \ \forall J \in \mathfrak{S}(U,Q)^G \}.$$

Let us now state a result whose proof can be found in [12, §3].

Theorem 2.3. The subset $\mathfrak{S}(U,Q)^G$ is a connected complex submanifold of $\mathfrak{S}(U,Q)$. The image of $\mathfrak{S}(U,Q)^G$ in A^{δ}_g coincides with the PEL subvariety $\mathsf{Z}(D_G)$. If $J \in \mathfrak{S}(U,Q)^G$, then $\dim \mathsf{Z}(D_G) = \dim(S^2\mathbb{R}^{2g})^G$ where \mathbb{R}^{2g} is endowed with the complex structure J.

3. Special subvarieties in the Prym Loci

Let Σ_g be a compact connected oriented surface of genus g. We denote by $T_g := \mathsf{T}(\Sigma_g)$ the Teichmüller space of Σ_g . Denote by $\mathsf{T}_{0,r}$ the Teichmüller space in genus 0 with $r \ge 4$ marked points. Let us recall the definition of $\mathsf{T}_{0,r}$ that can be found in [1, Chap. 15]. Let us fix $p_0, \ldots, p_r \in S^2$ distinct points, denote by $P := (p_1, \ldots, p_r)$. Points of $\mathsf{T}_{0,r}$ are equivalence classes of triples $(\mathbb{P}^1, x, [h])$ where $x = (x_1, \ldots, x_r)$ is an *r*-tuple of distinct points in \mathbb{P}^1 and [h] is an isotopy class of orientation preserving homeomorphisms $h : (\mathbb{P}^1, x) \to (S^2, P)$. Two such triples $(\mathbb{P}^1, x, [h]), (\mathbb{P}^1, x', [h'])$ are equivalent if there is a biholomorphism $\varphi : \mathbb{P}^1 \to \mathbb{P}^1$ such that $\varphi(x_i) = x'_i$ for any i and $[h] = [h' \circ \varphi]$.

Using the point $p_0 \in S^2 - P$ as base point we can fix an isomorphism

(3.1)
$$\Gamma_r := \langle \gamma_1, \dots, \gamma_r | \gamma_1 \cdots \gamma_r = 1 \rangle \cong \pi_1(S^2 - P, p_0).$$

Take a point $t = [(\mathbb{P}^1, x, [h])] \in \mathsf{T}_{0,r}$, and fix an epimorphism $\theta : \Gamma_r \to G$. By Riemann's existence theorem, this gives a Galois cover $C_t \to \mathbb{P}^1 = C_t/G$ ramified over x whose monodromy is given by θ , which is endowed with an isotopy class of homeomorphisms $[h_t]$ on a surface Σ_q covering S^2 .

So we get a map $\mathsf{T}_{0,r} \to \mathsf{T}_g : t \mapsto [(C_t, [h_t])]$ and the group G acts on C_t and embeds in the mapping class group Mod_g of Σ_g . We denote by G_θ the image of G in Mod_g . The image of the map $\mathsf{T}_{0,r} \to \mathsf{T}_g$ is the subset $\mathsf{T}_g^{G_\theta}$ of T_g consisting of the fixed points by the action of G_θ ([18]). By Nielsen realization theorem, we know that $\mathsf{T}_g^{G_\theta}$ is a complex submanifold of T_g of dimension r-3.

The image of $\mathsf{T}^{G_{\theta}}$ in the moduli space M_g is a (r-3)-dimensional algebraic subvariety (see e.g. [18, 5, 6] and [4, Thm. 2.1]).

We denote this image by $M_g(G,\theta)$. To determine $M_g(G,\theta)$ we have chosen the isomorphism (3.1) $\Gamma_r \cong \pi_1(S^2 - P, p_0)$. To get rid of this choice, we introduce the braid group

$$\mathbf{B}_{r} := \langle \tau_{1}, \dots, \tau_{r-1} | \tau_{i} \tau_{j} = \tau_{j} \tau_{i} \text{ for } |i-j| \ge 2, \ \tau_{i+1} \tau_{i} \tau_{i+1} = \tau_{i} \tau_{i+1} \tau_{i} \rangle.$$

There is a morphism $\varphi : \mathbf{B}_r \to \operatorname{Aut}(\Gamma_r)$ defined as follows:

$$\varphi(\tau_i)(\gamma_i) = \gamma_{i+1}, \quad \varphi(\tau_i)(\gamma_{i+1}) = \gamma_{i+1}^{-1} \gamma_i \gamma_{i+1},$$

$$\varphi(\tau_i)(\gamma_j) = \gamma_j \quad \text{for } j \neq i, i+1.$$

So we get an action of \mathbf{B}_r on the set of pairs (G, θ) , given by $\tau \cdot (G, \theta) := (G, \theta \circ \varphi(\tau^{-1}))$. Also the group $\operatorname{Aut}(G)$ acts on the set of pairs (G, θ) by $\alpha \cdot (G, \theta) := (G, \alpha \circ \theta)$. The orbits of the $\mathbf{B}_r \times \operatorname{Aut}(G)$ -action are called *Hurwitz equivalence classes* and elements in the same orbit are said to be related by a *Hurwitz move*. Data in the same orbit give rise to the same subvariety of M_g , hence the subvariety $\mathsf{M}_g(G, \theta)$ is well-defined. For more details see [32, 5, 3].

Definition 3.1. A Prym datum of type (r, b) is a triple $\Xi = (\tilde{G}, \tilde{\theta}, \sigma)$, where \tilde{G} is a finite group, $\tilde{\theta} : \Gamma_r \to \tilde{G}$ is an epimorphism and $\sigma \in \tilde{G}$ is a central involution and

$$b = \sum_{i: \sigma \in \langle \tilde{\theta}(\gamma_i) \rangle} \frac{|\tilde{G}|}{\operatorname{ord}(\tilde{\theta}(\gamma_i))}.$$

Let us fix a Prym datum $\Xi = (\tilde{G}, \tilde{\theta}, \sigma)$, and set $G = \tilde{G}/\langle \sigma \rangle$. The composition of $\tilde{\theta}$ with the projection $\tilde{G} \to G$ is an epimorphism $\theta : \Gamma_r \to G$. To a point $t \in \mathsf{T}_{0,r}$ we can associate two Galois covers

 $\tilde{C}_t \to \mathbb{P}^1 \cong \tilde{C}_t/\tilde{G}$ and $C_t \to \mathbb{P}^1 \cong C_t/G$. Denote by \tilde{g} the genus of \tilde{C}_t and by g the genus of C_t . We have a diagram

(3.2)
$$\tilde{C}_t \xrightarrow{f_t} C_t = \tilde{C}_t / \langle \sigma \rangle$$

$$\mathbb{P}^1 \checkmark$$

and by the definition of b, we immediately see that the double covering f_t has b branch points. The covering f_t is determined by its branch locus B_t and by an element $\eta_t \in \operatorname{Pic}^{\frac{b}{2}}(C_t)$ such that $\eta_t^2 = \mathcal{O}_{C_t}(B_t)$. Hence we have a map $\mathsf{T}_{0,r} \to \mathsf{R}_{g,b}$ which associates to t the class $[(C_t, \eta_t, B_t)]$. This map depends on the datum $\Xi = (\tilde{G}, \tilde{\theta}, \sigma)$. Denote by $R(\Xi)$ its image. The map $\mathsf{T}_{r,0} \to R(\Xi)$ has discrete fibres, hence dim $R(\Xi) = r - 3$.

As explained in [18, p. 79] there is an intermediate variety $\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G}, \tilde{\theta})$ such that the projection $\mathsf{T}_{\tilde{g}}^{G_{\tilde{\theta}}} \to \mathsf{M}_{\tilde{g}}(\tilde{G}, \tilde{\theta})$ factors through $\mathsf{T}_{\tilde{g}}^{\tilde{G}_{\tilde{\theta}}} \to \widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G}, \tilde{\theta}) \to \mathsf{M}_{\tilde{g}}(\tilde{G}, \tilde{\theta})$.

The variety $\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta})$ is the normalisation of $\mathsf{M}_{\tilde{g}}(\tilde{G},\tilde{\theta})$ and it parametrises equivalence classes of curves with an action of \tilde{G} of topological type determined by the datum Ξ . We also have a map $\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta}) \to R(\Xi)$, which associates to the class $[\tilde{C}]$ of a curve with a fixed \tilde{G} -action the class of the cover $\tilde{C} \to C = \tilde{C}/\langle \sigma \rangle$, where $\sigma \in \tilde{G}$ is the central involution fixed by the datum. Clearly the map $\mathsf{T}_{\tilde{g}}^{\tilde{G}\tilde{\theta}} \cong \mathsf{T}_{0,r} \to R(\Xi)$ is the composition $\mathsf{T}_{\tilde{g}}^{\tilde{G}\tilde{\theta}} \to \widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta}) \to R(\Xi)$. Moreover the Prym map $\mathsf{P}_{g,b}$ lifts to a map $\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta}) \to \mathsf{A}_{\tilde{g}-g}^{\delta}$, which sends the class of a curve $[\tilde{C}]$ with an action of \tilde{G} to $P(\tilde{C}, C = \tilde{C}/\langle \sigma \rangle)$. We still denote this map by $\mathsf{P}_{g,b} : \widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta}) \to \mathsf{A}_{\tilde{g}-g}^{\delta}$. We have the following diagram



Given a Prym datum $\Xi = (\tilde{G}, \tilde{\theta}, \sigma)$, let us fix an element \tilde{C} of the family $\mathsf{T}_{\tilde{g}}^{\tilde{G}_{\tilde{\theta}}}$ with the double covering $f : \tilde{C} \longrightarrow C$.

 Set

$$V := H^0(\tilde{C}, K_{\tilde{C}}),$$

We have an action of $\langle \sigma \rangle$ on V and an eigenspace decomposition for this action:

$$V := H^0(\tilde{C}, K_{\tilde{C}}) = V_+ \oplus V_-,$$

where $V_+ \cong H^0(C, K_C)$ and $V_- \cong H^0(C, K_C \otimes \eta)$ as *G*-modules. Similarly set $W := H^0(\tilde{C}, 2K_{\tilde{C}}) = W_+ \oplus W_-, W_+ \cong H^0(2K_C \otimes \eta^2) = H^0(2K_C(B))$ and $W_- \cong H^0(C, 2K_C \otimes \eta)$.

The multiplication map $m: S^2V \to W$ is the dual of the differential of the Torelli map $\tilde{j}: \mathsf{M}_{\tilde{g}} \to \mathsf{A}_{\tilde{g}}$ at the point $[\tilde{C}] \in \mathsf{M}_{\tilde{g}}$. This map is \tilde{G} -equivariant. We have the following isomorphisms

$$(S^2 V)^{\tilde{G}} = (S^2 V_+)^{\tilde{G}} \oplus (S^2 V_-)^{\tilde{G}}, \quad W^{\tilde{G}} = W_+^{\tilde{G}}$$

Hence the multiplication map m maps $(S^2 V)^{\tilde{G}}$ to $W^{\tilde{G}}_+$.

Notice that the codifferential of the Prym map at the point $[(C, \eta, B)]$ is the multiplication map in (2.1), which coincides with the restriction of the multiplication map m to S^2V_- . Here we are interested in the restriction of m to $(S^2V_-)^{\tilde{G}}$ that for simplicity we still denote by m:

$$(3.3) m: (S^2 V_-)^{\tilde{G}} \longrightarrow W_+^{\tilde{G}}.$$

By what we have said, this is the multiplication map

$$(S^2 H^0(C, K_C \otimes \eta))^G \longrightarrow H^0(C, 2K_C \otimes \eta^2)^G \cong H^0(2K_C)^G \cong H^0(2K_{\tilde{C}})^{\tilde{G}}.$$

Now we prove a result which is a generalisation of Theorems 3.2 and 4.2 in [10].

Theorem 3.2. Let $\Xi = (\tilde{G}, \tilde{\theta}, \sigma)$ be a Prym datum. If for some $t \in \mathsf{T}_{0,r}$ the map m in (3.3) is an isomorphism, then the closure of $\mathsf{P}_{g,b}(\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G}, \tilde{\theta}))$ in $\mathsf{A}_{q-1+\frac{k}{2}}^{\delta}$ is a special subvariety.

Proof. Over the Teichmüller space $\mathsf{T}_{0,r}$ we have the families \tilde{C}_t , C_t , η_t , B_t . We also have the lattice $H_1(\tilde{C}_t,\mathbb{Z})$, the intersection form E on $H_1(\tilde{C}_t,\mathbb{Z})$ and the sublattice $\Lambda := H_1(\tilde{C}_t,\mathbb{Z})_-$, which are independent of t.

Moreover $Q := E|_{\Lambda}$ is an integer-valued form on Λ of type $\delta = (1, \dots, 1, 2, \dots, 2)$.

Set $p := g - 1 + \frac{b}{2}$. Let $\mathfrak{S}(U, Q)$ be the Siegel space parametrizing complex structures on $U := \Lambda \otimes \mathbb{R} = H_1(\tilde{C}_t, \mathbb{R})_-$ that are compatible with Q. For any $t \in \mathsf{T}_{0,r}$ we have a decomposition $H^1(\tilde{C}_t, \mathbb{C})_- = V_{t,-} \oplus \overline{V_{t,-}}$. Dualizing we get a decomposition $H_1(\tilde{C}_t, \mathbb{C})_- = V_{t,-}^* \oplus \overline{V_{t,-}^*}$ corresponding to a complex structure J_t on $H_1(\tilde{C}_t, \mathbb{R})_-$. The complex structure J_t is compatible with Q, hence it is a point of $\mathfrak{S}(U,Q)$, that we denote by $\varphi(t)$. So we have a map $\varphi: \mathsf{T}_{0,r} \to \mathfrak{S}(U,Q)$ and a commutative diagram:

(3.4)
$$\begin{array}{ccc} \mathsf{T}_{0,r} & \stackrel{\varphi}{\longrightarrow} \mathfrak{S}(U,Q) \\ \downarrow & & \downarrow \\ \widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta}) & \stackrel{\mathsf{P}_{g,b}}{\longrightarrow} \mathsf{A}_{p}^{\delta}. \end{array}$$

The diagram commutes since

$$P(\tilde{C}_t, C_t) = V_{t,-}^* / \Lambda,$$

(see e.g. [21]).

Since the group \tilde{G} preserves E, \tilde{G} preserves Q, so \tilde{G} maps into $\operatorname{Sp}(\Lambda, Q)$. Denote by G' the image of \tilde{G} in $\operatorname{Sp}(\Lambda, Q)$. The complex structure J_t is \tilde{G} -invariant, which is equivalent to say that $\varphi(t) = J_t \in \mathfrak{S}(U,Q)^{G'}$. Hence by Theorem 2.3, $P(\tilde{C}_t, C_t)$ lies in the PEL special subvariety $\mathsf{Z}(D_{G'})$.

So we have shown that $\mathsf{P}_{g,b}(\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta})) \subset \mathsf{Z}(D_{G'})$. Since $\varphi(\mathsf{T}_{0,r}) \subset \mathfrak{S}(U,Q)^{G'}$ we can view φ as a map $\varphi : \mathsf{T}_{0,r} \to \mathfrak{S}(U,Q)^{G'}$. Now, recall that

$$\Omega^{1}_{\varphi(t)}(\mathfrak{S}(U,Q)^{G'}) \cong (S^{2}H^{0}(C_{t},K_{C_{t}}\otimes\eta_{t}))^{G} = (S^{2}V_{t,-})^{G},$$
$$\Omega^{1}_{t}\mathsf{T}_{0,r} = \Omega^{1}_{[C_{t}]}\mathsf{T}_{g}^{G_{\theta}} = H^{0}(C_{t},2K_{C_{t}})^{G} \cong H^{0}(C_{t},2K_{C_{t}}\otimes\eta_{t}^{2})^{G} = W_{t,+}^{\tilde{G}}.$$

The codifferential is simply the multiplication map

$$m = (d\varphi_t)^* : (S^2 V_{t,-})^{\tilde{G}} \longrightarrow W_{t,+}^{\tilde{G}}$$

(see [29] Prop. 3.1, or [21]). By assumption there is some $t \in \mathsf{T}_{0,r}$ such that the map m is an isomorphism at t. So $\dim(S^2V_{t,-})^{\tilde{G}} = \dim W^{\tilde{G}}_{t,+} = r-3$ and φ is an immersion at t, hence its image has dimension r-3. The vertical arrows in (3.4) are discrete maps, hence both $\mathsf{P}_{g,b}(\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta}))$ and $\mathsf{Z}(D_{G'})$ have dimension r-3. Since $\mathsf{P}_{g,b}(\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta})) \subset \mathsf{Z}(D_{G'})$ and $\mathsf{Z}(D_{G'})$ is irreducible, we conclude that $\overline{\mathsf{P}_{g,b}(\widetilde{\mathsf{M}}_{\tilde{g}}(\tilde{G},\tilde{\theta}))} = \mathsf{Z}(D_{G'})$, so it is a special subvariety of A_p^{δ} .

We would like to use Theorem 3.2 to construct Shimura curves contained in the closure of $\mathsf{P}_{g,b}(\mathsf{R}_{g,b})$ in $\mathsf{A}_{g-1+\frac{b}{2}}^{\delta}$ and intersecting $\mathsf{P}_{g,b}(\mathsf{R}_{g,b})$. So from now on we will assume that r = 4, that means that the coverings $\tilde{C} \to \tilde{C}/\tilde{G} \cong \mathbb{P}^1$ and $C \to C/G \cong \mathbb{P}^1$ in diagram (3.2) are ramified over 4 points in \mathbb{P}^1 .

Hence, as in [10], the hypothesis in Theorem 3.2 can be divided in two conditions:

$$\dim(S^2 V_-)^{\bar{G}} = 1.$$

(B)
$$m: (S^2V_-)^{\tilde{G}} \longrightarrow W^G_+ \text{ is not identically } 0.$$

Once condition (A) is true, a sufficient condition ensuring (B) is the following

(B1)
$$(S^2V_-)^G$$
 is generated by a decomposable tensor.

In fact, assume that $(S^2V_-)^{\tilde{G}}$ is generated by $s_1 \otimes s_2$ with $s_i \in V_-$, then $m(s_1 \otimes s_2) = s_1 \cdot s_2$, which is different from zero.

Remark 3.3. In [10], Remark 3.3, it is shown that if (A) holds, then (B1) is equivalent to the fact that $(S^2V_{-})^{\tilde{G}} = W_1 \otimes W_2$ where the W_i 's are 1-dimensional representations.

Hence, if \tilde{G} is abelian and condition (A) holds, then condition (B1) is automatically satisfied, since all the irreducible representations of an abelian group are 1-dimensional (see [10] Remark 3.4).

If (A) holds, (B) is equivalent to say that the Prym map is not constant. In particular this happens if the prym decomposes up to isogeny as

(B2)
$$P(\tilde{C}, C) \sim A \times JC'$$

where A is a fixed abelian variety and JC' is the jacobian of a curve $C' = \tilde{C}/N$ defined as a quotient of \tilde{C} by a normal subgroup $N \triangleleft \tilde{G}$, and the family of Galois covers $C' \rightarrow \mathbb{P}^1 = C'/(G/N)$ satisfies condition (*) of [12], hence it describes a Shimura curve.

In fact, since A is a fixed abelian variety and JC' moves in a Shimura curve, the Prym variety P(C, C) necessarily moves.

Notice that [13, Thm. 3.8]) ensures that condition (B2) implies that the family of Pryms $P(\tilde{C}, C)$ yields a totally geodesic subvariety of A_n^{δ} .

Finally in [31] it is proven that $\mathsf{P}_{g,b}$ is an embedding for all $b \ge 6$ and for all g > 0, hence if $b \ge 6$, and g > 0 condition (A) implies condition (B).

4. Examples in the Prym locus

In this section we describe some examples of families of covers satisfying conditions (A) and (B). They do not satisfy (B1), and we explicitly show that (B) holds. However, notice that the first 5 examples are in $\mathsf{R}_{q,b}$ with $b \ge 6$, hence condition (B) is automatically satisfied thanks to [31].

In the first two examples we show that condition (B2) is satisfied.

The other examples do not satisfy (B1), nor (B2).

In particular, example 6 does not satisfy (B1), nor (B2) and it has b = 4, hence we need to show that (B) holds with an ad-hoc argument.

To describe the examples we first give the values of \tilde{g} , g and b. We also indicate a number # that refers to the numbering of the examples that we found with the MAGMA script with the given values of \tilde{g}, g and b. For the group \tilde{G} we use the presentation and the name of the group given by MAGMA. We then give the decomposition of V_+ and V_- as a direct sum of irreducible representations of G and also for the irreducible representations we use the notation of MAGMA.

Example 1.

 $\tilde{g} = 10, g = 4, b = 6, \#5.$

 $\tilde{G} = G(24, 10) = \mathbb{Z}/3 \times D_4 = \langle g_3 | g_3^3 = 1 \rangle \times \langle g_1, g_2, g_4 | g_1^2 = g_2^2 = g_4^2 = 1, g_1^{-1}g_2g_1 = g_2g_4 \rangle,$

 $\tilde{\theta}(\gamma_1) = g_1, \ \tilde{\theta}(\gamma_2) = g_2 g_4, \ \tilde{\theta}(\gamma_3) = g_3^2 g_4, \ \tilde{\theta}(\gamma_4) = g_1 g_2 g_3 g_4, \ \sigma = g_4.$ $V_+ = V_5 \oplus V_6 \oplus V_7 \oplus V_{10} \text{ and } V_- = V_{13} \oplus 2V_{15}, \text{ where } \dim V_i = 1, \ \forall i < 13, \ \dim V_{13} = \dim V_{15} = 2.$ $(S^2 V_+)^{\tilde{G}} = V_5 \otimes V_7 \text{ and } (S^2 V_-)^{\tilde{G}} = (S^2 V_{13})^{\tilde{G}} \text{ have dimension } 1. \text{ Thus condition } (A) \text{ is satisfied, but }$ (B1) is not. We want to show that condition (B2) holds and how it works. If we find a curve C' such that $H^{1,0}(JC') = V_{13}$, then we obtain $P(\tilde{C}, C) \sim JC' \times A$ for some fixed abelian variety A such that $H^{1,0}(A) = 2V_{15}.$

Notice that $V_{13} = \text{Fix}\langle g_3 \rangle$ and $\langle g_3 \rangle \triangleleft \tilde{G}$, so we consider $G' = \tilde{G}/\langle g_3 \rangle$ and $C' = \tilde{C}/\langle g_3 \rangle$. Thus we have a commutative diagram



The map β has 6 branch points, and C' has genus 2, and the map $C' \to C'/G'$ has 4 critical values. Since $\dim(S^2H^{1,0}(C'))^{G'} = \dim(S^2V_{13})^{\tilde{G}} = 1$ then condition (*) of [12] is satisfied, thus the family of jacobians JC' yields a Shimura curve. In fact it is family (29) = (4) of [12].

Finally, the group algebra decomposition for $J\bar{C}$ (see [33]), allows to give a decomposition up to isogeny:

$$J\tilde{C} \sim B_5 \times B_6 \times B_{10} \times B_{13}^2 \times B_{15}^2$$

where dim $B_5 = \dim B_{15} = 2$ and the others have dimension 1. In particular $P(\tilde{C}, C) \sim B_{13}^2 \times B_{15}^2$ and $JC' \sim B_{13}^2$. In fact $JC' = E^2$ where E is the elliptic curve defined by $E := \tilde{C}/\langle g_1, g_3 \rangle = C'/\langle g_1 \rangle$, so

 $B_{13} \sim E$. Thus $P(C, C) \sim A \times JC'$, where $A := B_{15}^2$ is a fixed abelian variety and $JC' \sim E^2$ varies in the Shimura curve (29)=(4) of [12], hence condition (B2) holds.

Example 2.

$$\begin{split} \tilde{g} &= 10, \ g = 4, \ b = 6, \ \#8. \\ \tilde{G} &= G(36, 12) = \mathbb{Z}/3\mathbb{Z} \times D_6 = \langle g_3 | g_3^3 = 1 \rangle \times \langle g_1, g_2, g_4 | g_1^2 = g_2^2 = g_4^3 = 1, \ g_1^{-1}g_4g_1 = g_4^2 \rangle. \\ \tilde{\theta}(\gamma_1) &= g_1g_2, \ \tilde{\theta}(\gamma_2) = g_1g_4, \ \tilde{\theta}(\gamma_3) = g_3^2g_4^2, \ \tilde{\theta}(\gamma_4) = g_2g_3, \quad \sigma = g_2. \\ V_+ &= V_6 \oplus V_9 \oplus V_{16} \ \text{and} \ V_- = V_5 \oplus V_8 \oplus V_{14} \oplus V_{15}, \ \text{where } \dim V_i = 1, \ \text{for} \ i = 5, 6, 8, 9, \ \dim V_{14} = \dim V_{15} = \dim V_{16} = 2. \end{split}$$

 $(S^2V_+)^{\tilde{G}} = V_6 \otimes V_9$ and $(S^2V_-)^{\tilde{G}} = (S^2V_{14})^{\tilde{G}}$ have dimension 1. Thus condition (A) is satisfied, but (B1) is not. We want to show that condition (B2) holds. If we find a curve C' such that $H^{1,0}(JC') = V_{14}$, then we obtain $P(\tilde{C}, C) \sim JC' \times A$ for some fixed abelian variety A such that $H^{1,0}(A) = V_5 \oplus V_8 \oplus V_{15}$.

We have: $V_{14} = \text{Fix}\langle g_3 \rangle$ and $\langle g_3 \rangle \triangleleft \tilde{G}$, so consider $G' = \tilde{G}/\langle g_3 \rangle$ and $C' = \tilde{C}/\langle g_3 \rangle$. We have a commutative diagram



The map β has 6 branch points, C' has genus 2 and the cover $C' \to C'/G'$ has 4 critical values. Since $\dim(S^2H^{1,0}(C'))^{G'} = \dim(S^2V_{14})^{\tilde{G}} = 1$, condition (*) of [12] is satisfied, thus the family of jacobians JC' yields a Shimura curve, which corresponds to family (30) of [12].

The group algebra decomposition for $J\tilde{C}$ gives

$$J\tilde{C} \sim B_5 \times B_6 \times B_8 \times B_{14}^2 \times B_{15}^2 \times B_{16}^2$$

where dim $B_6 = 2$ and the others have dimension 1. So $P(\tilde{C}, C) \sim A \times JC' \sim B_5 \times B_8 \times B_{14}^2 \times B_{15}^2$, $JC' \sim B_{14}^2$, $A \sim B_5 \times B_8 \times B_{15}^2$.

Example 3.

$$\begin{split} \tilde{g} &= 10, \ g = 4, \ b = 6, \ \#7.\\ \tilde{G} &= G(36, 12) = \mathbb{Z}/6 \times S_3 = \langle g_2, g_3 | g_2^2 = g_3^3 = 1 \rangle \times \langle g_1, g_4 | g_1^2 = g_4^3 = 1, g_1^{-1}g_4g_1 = g_4^2 \rangle,\\ \tilde{\theta}(\gamma_1) &= g_1g_4, \ \tilde{\theta}(\gamma_2) = g_1g_2g_4^2, \ \tilde{\theta}(\gamma_3) = g_3^2, \ \tilde{\theta}(\gamma_4) = g_2g_3g_4^2, \quad \sigma = g_2.\\ V_+ &= V_6 \oplus V_9 \oplus V_{18} \text{ and } V_- = V_5 \oplus V_8 \oplus V_{15} \oplus V_{17}, \text{ where } \dim V_5 = \dim V_6 = \dim V_8 = \dim V_9 = 1 \text{ and } \\ \dim V_{15} &= \dim V_{17} = \dim V_{18} = 2.\\ (S^2 V_-)^{\tilde{G}} &= (V_{15} \otimes V_{17})^{\tilde{G}} \text{ is one-dimensional, so condition (A) is satisfied, but neither (B1) nor (B2) \end{split}$$

 $(S^2V_{-})^{\circ} = (V_{15} \otimes V_{17})^{\circ}$ is one-dimensional, so condition (A) is satisfied, but neither (B1) nor (B2) hold.

 $G = \tilde{G}/\langle g_2 \rangle \cong \mathbb{Z}/3 \times S_3 = G(18,3), (S^2V_+)^{\tilde{G}} = (V_6 \otimes V_9)$, which has dimension 1, hence the family $C \to C/G$ satisfies condition (*) of [12] and it yields a Shimura curve, corresponding to family (38) of [12].

The group algebra decomposition of $J\tilde{C}$ is the following: $J\tilde{C} \sim B_5 \times B_6 \times B_8 \times B_{15}^2 \times B_{18}^2$ where dim $B_6 = \dim B_{15} = 2$ and dim $B_5 = \dim B_8 = \dim B_{18} = 1$. Hence

$$P(\tilde{C}, C) \sim B_5 \times B_8 \times B_{15}^2.$$

The trace of $g_{1|V_{15}}$ is zero, hence V_{15} decomposes as a direct sum of the two one dimensional (± 1) eigenspaces of the action of g_1 . The same happens for the action of g_1 on V_{17} . Denote by $\langle e_1 \rangle$ (resp. $\langle f_1 \rangle$) the (+1)-eigenspace of $g_{1|V_{15}}$ (resp. of $g_{1|V_{17}}$) and by $\langle e_2 \rangle$ (resp. $\langle f_2 \rangle$) the (-1)-eigenspace of $g_{1|V_{15}}$ (resp. of $g_{1|V_{17}}$), so $V_{15} = \langle e_1, e_2 \rangle$, $V_{17} = \langle f_1, f_2 \rangle$. One easily checks that the elliptic curve B_{18} is isogenous to $\tilde{C}/\langle g_1, g_2 \rangle$ and we denote by $\langle v_1 \rangle = H^{1,0}(B_{18}) \subset V_{18}$. We define $C' := \tilde{C}/\langle g_1 \rangle$ and we calculate $H^{1,0}(C') = V_8 \oplus \langle e_1 \rangle \oplus \langle f_1 \rangle \oplus \langle v_1 \rangle$ and the decomposition $JC' \sim B_8 \times B_{15} \times B_{18}$. Similarly if we define $D' := \tilde{C}/\langle g_1 g_2 \rangle$ we have $H^{1,0}(D') = V_5 \oplus \langle e_2 \rangle \oplus \langle f_2 \rangle \oplus \langle v_1 \rangle$ and $JD' \sim B_5 \times B_{15} \times B_{18}$. Thus we obtain

$$JC' \times JD' \sim P(\tilde{C}, C) \times B_{18}^2$$

and we obtain that $P(\tilde{C}, C)$ moves if and only if the product of jacobians on the left move since B_{18} is a fixed abelian variety. Moreover it moves if and only if B_{15} does, since both B_5 and B_8 are fixed. We consider the following covers



Assume that the critical values of the cover $\tilde{C} \to \tilde{C}/\tilde{G} \cong \mathbb{P}^1$ are $\{P_1 = \lambda, P_2 = 0, P_3 = 1, P_4 = \infty\}$. The maps ε and ε' are 2:1, ramified respectively over $\{0, \infty\}$, and $\{\lambda, \infty\}$. So we can assume $\varepsilon(z) = z^2$, and $\varepsilon'(z) = (1 - \lambda)z^2 + \lambda$. Thus the covers δ, δ' are ramified over $1, -1, \mu, -\mu, \infty$ and $1, -1, \alpha, -\alpha, \infty$ respectly, where $\mu^2 = \lambda$ and $\alpha^2 = \frac{\lambda}{\lambda - 1}$. The 3:1 covers t, t' are ramified over 3 points, which we can assume to be $1, -1, \infty$ for both maps. So B_5 and B_8 are isomorphic and an equation for them as $\mathbb{Z}/3$ -covers of \mathbb{P}^1 is given by $y^3 = (x^2 - 1)$. So the critical values of the maps β (resp. β') are the 6 preimages of $\pm \mu$, (resp. $\pm \alpha$) via t (resp. t'). Since they move, varying λ , the jacobians $JC' \sim JD'$ move.

Example 4.

$$\begin{split} \tilde{g} &= 10, \ g = 4, \ b = 6, \ \#9. \\ \tilde{G} &= G(36, 12) = \mathbb{Z}/6 \times S_3 = \langle g_2, g_3 | g_2^2 = g_3^3 = 1 \rangle \times \langle g_1, g_4 | g_1^2 = g_4^3 = 1, g_1^{-1}g_4g_1 = g_4^2 \rangle, \\ \tilde{\theta}(\gamma_1) &= g_1g_4^2, \ \tilde{\theta}(\gamma_2) = g_1g_2g_4^2, \ \tilde{\theta}(\gamma_3) = g_3^2g_4, \ \tilde{\theta}(\gamma_4) = g_2g_3g_4^2, \quad \sigma = g_2. \\ V_+ &= V_6 \oplus V_9 \oplus V_{13} \text{ and } V_- = V_5 \oplus V_8 \oplus V_{14} \oplus V_{17}, \text{ where dim } V_i = 1 \text{ for } i = 5, 6, 8, 9, \text{ dim } V_{13} = \text{dim } V_{14} = \text{dim } V_{17} = 2. \\ (S^2V_+)^{\tilde{G}} &= (V_6 \otimes V_9) \oplus (S^2V_{13})^{\tilde{G}} \text{ and } (S^2V_-)^{\tilde{G}} = (S^2V_{14})^{\tilde{G}} \text{ where all summands have dimension } 1. \\ \text{Condition (A) is satisfied, but (B1) and (B2) are not. We notice that } V_{14} \text{ is fixed by } g_3 \text{ and } \text{Fix}\langle g_3 \rangle = V_{13} \oplus V_{14}. \text{ If we define } C' := \tilde{C}/\langle g_3 \rangle \text{ and } C'' := \tilde{C}/\langle g_2, g_3 \rangle = C'/\langle g_2 \rangle \text{ then we have } P(\tilde{C}, C) \sim A \times P(C', C'') \text{ where } A \text{ is a fixed abelian variety such that } H^{1,0}(A) = V_5 \oplus V_8 \oplus V_{17}. \text{ Therefore the family } P(\tilde{C}, C) \text{ describes a Shimura variety if the family of } P(C', C'') \text{ moves. And this is true as noted in Example 2 of [10].} \end{split}$$

Example 5. $\tilde{g} = 11, g = 3, b = 12, \#2.$ $\tilde{G} = G(32, 42) = (\mathbb{Z}/2 \times \mathbb{Z}/8) \rtimes \mathbb{Z}/2 = \langle g_1, g_2, g_3, g_4, g_5 | g_1^2 = g_2^2 = g_5^2 = 1, g_3^2 = g_4^2 = g_5, g_1^{-1}g_2g_1 = g_2g_4, g_1^{-1}g_4g_1 = g_4g_5, g_2^{-1}g_4g_2 = g_4g_5 \rangle,$ $\tilde{\theta}(\gamma_1) = g_2g_4g_5, \ \tilde{\theta}(\gamma_2) = g_1, \ \tilde{\theta}(\gamma_3) = g_3, \ \tilde{\theta}(\gamma_4) = g_1g_2g_3g_4, \quad \sigma = g_5.$ $V_+ = V_6 \oplus V_{10} \text{ and } V_- = V_{12} \oplus 2V_{13} \oplus V_{14} \text{ where dim } V_6 = 1 \text{ and the others have dimension } 2.$ $(S^2V_+)^{\tilde{G}} = (S^2V_6) \oplus (S^2V_{10})^{\tilde{G}} \text{ and } (S^2V_-)^{\tilde{G}} = (V_{12} \otimes V_{14})^{\tilde{G}} \text{ where all summands have dimension } 1.$ Condition (A) is satisfied, but (B1) and (B2) are not. We consider its centraliser $K = \langle g_1g_2, g_3g_4 \rangle \cong$ $\mathbb{Z}/8 \times \mathbb{Z}/2$. We have $\langle g_3 g_4 \rangle \lhd \langle g_3 g_4, g_5 \rangle \lhd K \lhd G$ and the following diagram.



where $K' = K/\langle g_3 g_4 \rangle \cong \mathbb{Z}/8 = \langle \overline{g_1 g_2} \rangle$. Set $h := \overline{g_1 g_2}$.

Assume that the critical values of the cover $\tilde{C} \to \tilde{C}/\tilde{G} \cong \mathbb{P}^1$ are $\{P_1 = 1, P_2 = \infty, P_3 = \lambda, P_4 = 0\}$. The map ε is a double cover ramified at 2 points, so we can assume $\varepsilon(z) = -z^2 + 1$, hence it is ramified at $0, \infty$ and $\varepsilon^{-1}(\lambda) = \{\pm \alpha\}, \varepsilon^{-1}(0) = \{\pm 1\}$ ($\alpha^2 = 1 - \lambda$).

The critical values of φ are $[\alpha, -\alpha, 1, -1]$ and the monodromy is given by $[h^2, h^2, h, h^3]$, so the equation of the genus 6 curves C' as $\mathbb{Z}/8$ -cyclic covers of \mathbb{P}^1 is: $y^8 = (x - \alpha)^2 (x + \alpha)^2 (x - 1)(x + 1)^3$. Hence an equation for the $\mathbb{Z}/4$ coverings $F \to C'/K'$ is $y^4 = (x - \alpha)^2 (x + \alpha)^2 (x - 1)(x + 1)^3$, F has genus 2 and this is the Shimura family (4) = (29) of [12].

Denote by $W_i := \{x \in H^0(K_{C'}) \mid h(x) = \zeta^i x\}$, where ζ is a primitive 8-th root of unity. Then one easily computes that $\dim W_1 = \dim W_4 = 0$, $\dim W_2 = \dim W_3 = \dim W_5 = \dim W_6 = 1$, $\dim W_7 = 2$. Therefore we have $H^0(K_F) \cong W_2 \oplus W_6$, $H^{1,0}(P(C', F) \cong W_3 \oplus W_5 \oplus W_7$, hence $(S^2 H^{1,0}(P(C', F))^{\langle h \rangle} \cong W_3 \otimes W_5$ is one dimensional. Thus the codifferential of the Prym map of the double coverings $C' \to F$ is the multiplication map $(S^2 H^{1,0}(P(C', F))^{\langle h \rangle} \cong W_3 \otimes W_5 \to H^0(2K_{C'})^{\langle h \rangle}$ and it is an isomorphism.

This shows that the Pryms P(C', F) vary and yield a Shimura curve in A_4^{δ} . In fact this is the first example with $\tilde{g} = 6$, g = 2, b = 6 in Table 1 and it satisfies both (A) and (B1).

Finally the group algebra decomposition gives $J\tilde{C} \sim B_6 \times B_{10}^2 \times B_{12}^2$, where dim B_6 = dim $B_{10} = 1$, dim $B_{12} = 4$. Moreover $JC \sim B_6 \times B_{10}^2$, hence $P(\tilde{C}, C) \sim B_{12}^2$, $JC' \sim B_{10}^2 \times B_{12}$, $JF \sim B_{10}^2$, hence $(P(C', F))^2 \sim B_{12}$. Therefore $P(\tilde{C}, C) \sim (P(C', F))^2$, hence it moves, and condition (B) is satisfied.

Example 6.

 $\tilde{g} = 7, g = 3, b = 4, \#4.$

 $\tilde{G} = \tilde{G}(16,6) = \mathbb{Z}/8 \rtimes \mathbb{Z}/2 = \langle g_1 \rangle \rtimes \langle g_2 \rangle = \langle g_1, g_2 \mid g_1^8 = 1, \ g_2^2 = 1, \ g_2^{-1}g_1g_2 = g_1^5 \rangle. \text{ Set } g_3 := g_1^2, \ g_4 = g_1^4.$ $\tilde{\theta}(\gamma_1) = g_2, \ \tilde{\theta}(\gamma_2) = g_2g_4, \ \tilde{\theta}(\gamma_3) = g_1g_4, \ \tilde{\theta}(\gamma_4) = g_1g_3g_4, \quad \sigma = g_4.$

 $V_+ = V_3 \oplus V_5 \oplus V_7$ and $V_- = V_9 \oplus V_{10}$ where dim $V_9 = \dim V_{10} = 2$ and the others have dimension 1. $(S^2V_+)^{\tilde{G}} = (S^2V_3) \oplus (V_5 \otimes V_7)$ and $(S^2V_-)^{\tilde{G}} = (V_9 \otimes V_{10})^{\tilde{G}}$ where all summands have dimension 1. Condition (A) is satisfied, but (B1) and (B2) are not. The trace of $g_{2|V_9}$ and of $g_{2|V_{10}}$ is zero, hence we can write $V_9 = \langle e_1, e_2 \rangle$, $V_{10} = \langle f_1, f_2 \rangle$, where $g_2(e_1) = e_1, g_2(e_2) = -e_2, g_2(f_1) = f_1, g_2(f_2) = -f_2$. Since g_4 acts as -Id on V_9 and on V_{10} , we have: $g_2g_4(e_1) = -e_1, g_2g_4(e_2) = e_2, g_2g_4(f_1) = -f_1, g_2g_4(f_2) = f_2$. The curves $C' = \tilde{C}/\langle g_2 \rangle$ and $D' = \tilde{C}/\langle g_2 g_4 \rangle$, are isomorphic since g_2 and g_2g_4 are conjugated in \tilde{G} . From the group algebra decomposition we get: $J\tilde{C} \sim B_3 \times B_5 \times B_9^2$, where dim $B_3 = 1$ and dim $B_5 = \dim B_9 = 2$. Furthemore $JC' \cong JD' \sim B_9$ and $P(\tilde{C}, C) \sim JC' \times JD' \sim (JC')^2 \sim B_9^2$. In order to study C', we consider $H := \langle g_3, g_2 \rangle \cong \mathbb{Z}/4 \times \mathbb{Z}/2$ so that $\langle g_2 \rangle \triangleleft H \triangleleft \tilde{G}$ and we get $\tilde{C}/H = \mathbb{P}^1$.



The map l is a double cover ramified in 2 points, so we may assume that $l(z) = z^2$. Hence the critical values for k are $\{\pm 1, \pm \mu, 0, \infty\}$ where $\mu^2 = \lambda$. Thus the branch locus of the map $t : C' \to C'/\mathbb{Z}/4$ is $\{1, \mu, 0, \infty\}$ with monodromy $[\bar{2}, \bar{2}, \bar{1}, \bar{1}]$, (here by \bar{n} we mean the class of n in $\mathbb{Z}/4 \cong H/\langle g_2 \rangle$). So the equation for C' is $y^4 = (x-1)^2(x-\mu)^2 x$ and we conclude that JC' moves. Hence $P(\tilde{C}, C) \sim (JC')^2$ moves and condition (B) is satisfied.

Appendix

In this appendix we briefly explain how the MAGMA script works.

The starting data is composed of the number r of critical values on \mathbb{P}^1 and the genus \tilde{g} of the curve \tilde{C} . Next we proceed by generating Prym datas with the following steps:

- (1) We are intrested in the case r > 3. We consider mainly r = 4, in this case the maximum order for the authomorphism group of a curve of genus \tilde{g} is $12(\tilde{g}-1)$. For $r \ge 5$ this limit decreases to $4(\tilde{g}-1)$.
- (2) An ammissible signature (m_1, \ldots, m_r) is formed by divisors of $|\tilde{G}|$ grater than 1 and must satisfy the Riemann-Hurwitz formula

$$2\tilde{g} - 2 = |\tilde{G}| \left(-2 + \sum_{i=1}^{r} \left(1 - \frac{1}{m_i} \right) \right).$$

We also restrict to ordered sequences.

- (3) Then we generate the data. For each even order we consider all groups of that cardinality. For each group \tilde{G} and signature we focus on elements of order m_1, \ldots, m_r . We consider all possible r-uple (x_1, \ldots, x_r) of elements of \tilde{G} such that $\operatorname{ord}(x_i) = m_i$ and $x_1 \cdots x_r = 1$. Moreover all central elements of order 2 are saved.
- (4) Finally we exclude equivalent data. Two monodromy data are equivalent in two cases: either they are isomorphic under an automorphism of \tilde{G} or under the action of the Braid group \mathbf{B}_r . One datum per orbit is saved.

From a Prym datum $(\tilde{G}, \tilde{\theta}, \sigma)$ we obtain the datum (G, θ) of the quotient curve from the quotient by σ . Now we can easly compute the genus g of C and the number b of ramification points via the Riemann-Hurwitz formula or the formula given in Definition 3.1. After that we test the hypothesis of Theorem 3.2.

- (5) We test condition (A) by computing the dimension of $S^2 H^{1,0}$ for both curves \tilde{C} , C. To do this we use the Chevalley-Weil formula to obtain the \tilde{G} -isotopic decomposition of $H^{1,0}$ for both curves. If condition (A) is verified the datum is recorded.
- (6) For these data conditions (B1) and (B2) are tested through the study of the G-isotopic decomposition of S^2V_- . If one of them is verified it means that the hypothesis of Theorem 3.2 are satisfied and therefore the datum gives rise to a special variety. Otherwise further ad-hoc analysis is needed to determine if condition (B) is satisfied.

The following tables list all the calculated data. Table 1 shows the examples that satisfy both conditions (A) and (B). For each pair \tilde{g} and g (and thus b and $p = \tilde{g} - g$) the examples are listed with a progressive index (under column #). For each example it is reported the group \tilde{G} and the id of \tilde{G} and G as small group in the MAGMA Database. A check mark or a cross tells if the conditions are met or not. The script tests condition (B2) only if (B1) doesn't hold. Different examples with same data are grouped in the same row.

In Table 2 for each \tilde{g} and g it is reported on the last column the number of non-equivalent data that satisfies condition (A). This includes all examples listed in Table 1 but also all the cases where neither (B1), nor (B2) hold and we did not check if (B) holds. There aren't others cases satisfying (A) with $\tilde{g} \leq 70$.

\tilde{g}	g	b	p	#	$ ilde{G}$	$\tilde{G}\; \operatorname{Id}$	$G\;{\rm Id}$	(B1)	(B2)	$b \ge 6$	(B)
2	0	6	2	1	$\mathbb{Z}/4$	G(4, 1)	G(2, 1)	\checkmark		\checkmark	\checkmark
2	0	6	2	2	$\mathbb{Z}/6$	G(6,2)	G(3,1)	\checkmark		\checkmark	\checkmark
2	0	6	2	3	D_4	G(8,3)	G(4, 2)	X	X	\checkmark	\checkmark
2	0	6	2	4	D_6	G(12, 4)	G(6,1)	X	X	\checkmark	\checkmark
3	2	0	1	1	$\mathbb{Z}/2 \times \mathbb{Z}/4$	G(8, 2)	G(4, 1)	\checkmark		X	\checkmark
3	2	0	1	2	$\mathbb{Z}/2 \times D_4$	G(16, 11)	G(8,3)	\checkmark		X	\checkmark
3	1	4	2	1	$\mathbb{Z}/4$	G(4, 1)	G(2, 1)	\checkmark		X	\checkmark
3	1	4	2	2	$\mathbb{Z}/6$	G(6,2)	G(3,1)	\checkmark		X	\checkmark
3	1	4	2	$3,\!4,\!6$	$\mathbb{Z}/2 \times \mathbb{Z}/4$	G(8,2)	G(4, 1)	\checkmark		X	\checkmark
3	1	4	2	5	$\mathbb{Z}/2 \times \mathbb{Z}/4$	G(8,2)	G(4,2)	\checkmark		X	\checkmark
3	1	4	2	8	$\mathbb{Z}/2 \times D_4$	G(16, 11)	G(8,5)	X	\checkmark	X	\checkmark
3	0	8	3	1	$\mathbb{Z}/2 \times \mathbb{Z}/4$	G(8,2)	G(4,2)	\checkmark		\checkmark	\checkmark
4	2	2	2	1	$\mathbb{Z}/6$	G(6,2)	G(3,1)	\checkmark		X	\checkmark
4	1	6	3	1	$\mathbb{Z}/6$	G(6, 2)	G(3, 1)	\checkmark		\checkmark	\checkmark
4	1	6	3	2,3	$\mathbb{Z}/2 \times \mathbb{Z}/6$	G(12, 5)	G(6,2)	\checkmark		\checkmark	\checkmark
4	0	10	4	1	Q_8	G(8, 4)	G(4, 2)	X	X	\checkmark	\checkmark
5	3	0	2	1 - 3	$\mathbb{Z}/2 \times \mathbb{Z}/4$	G(8, 2)	G(4, 1)	\checkmark		X	\checkmark
5	3	0	2	4	$\mathbb{Z}/2 \times \mathbb{Z}/6$	G(12, 5)	G(6,2)	\checkmark		X	\checkmark
5	3	0	2	5	$(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/4$	G(16, 3)	G(8,3)	\checkmark		X	\checkmark
5	3	0	2	8	$(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/4$	G(16, 3)	G(8,2)	X	\checkmark	X	\checkmark
5	3	0	2	9 - 11	$\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/4$	G(16, 10)	G(8,2)	\checkmark		X	\checkmark
5	3	0	2	14	$\mathbb{Z}/2 \times \mathbb{Z}/2 \times S_3$	G(24, 14)	G(12, 4)	X	\checkmark	X	\checkmark
5	3	0	2	15	$\mathbb{Z}/2 \times A_4$	G(24, 13)	G(12, 3)	\checkmark		X	\checkmark
5	3	0	2	$16,\!17$	$(\mathbb{Z}/2)^2 \wr \mathbb{Z}/2$	G(32, 27)	G(16, 11)	X	\checkmark	X	\checkmark
5	3	0	2	18	$\mathbb{Z}/4 \times D_4$	G(32, 28)	G(16, 13)	X	\checkmark	X	\checkmark
5	3	0	2	19	$\mathbb{Z}/2 \times S_4$	G(48, 48)	G(24, 12)	X	\checkmark	X	\checkmark
5	2	4	3	1,2	$(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/4$	G(16,3)	G(8,3)	X	\checkmark	X	\checkmark
5	2	4	3	3	$(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/4$	G(16,3)	G(8,3)	\checkmark		X	\checkmark
5	1	8	4	1	$\mathbb{Z}/8$	G(8,1)	G(4,1)	\checkmark		\checkmark	\checkmark
5	1	8	4	2	$\mathbb{Z}/2 \times \mathbb{Z}/4$	G(8,2)	G(4,2)	\checkmark		\checkmark	\checkmark
5	1	8	4	3	Q_8	G(8,4)	G(4,2)	X	X	\checkmark	\checkmark
5	1	8	4	4	$(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/4$	G(16,3)	G(8,3)	X	X	\checkmark	\checkmark
5	1	8	4	5	$Q_8 \rtimes \mathbb{Z}/2$	G(16,8)	G(8,3)	X	X	\checkmark	\checkmark
5	1	8	4	6	$\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/4$	G(16, 10)	G(8,5)	\checkmark		\checkmark	\checkmark
5	1	8	4	7	$\mathbb{Z}/2 \circ D_4$	G(16, 13)	G(8,5)	X	X_{-}	\checkmark	\checkmark
5	1	8	4	8	$\mathbb{Z}/4 \times D_4$	G(32, 28)	G(16, 11)	X	\checkmark	\checkmark	\checkmark
5	1	8	4	9	$\mathbb{Z}/8 \rtimes (\mathbb{Z}/2 \times \mathbb{Z}/2)$	G(32, 43)	G(16, 11)	X	X	\checkmark	\checkmark
5	0	12	5	1	$\mathbb{Z}/3 \rtimes \mathbb{Z}/4$	G(12, 1)	G(6,1)	X	X	\checkmark	\checkmark
6	3	2	3	$1,\!2$	$\mathbb{Z}/2 imes \mathbb{Z}/6$	G(12, 5)	G(6,2)	\checkmark		X	\checkmark
6	2	6	4	1	$\mathbb{Z}/8$	G(8,1)	G(4,1)	\checkmark		\checkmark	\checkmark
6	2	6	4	2	$\mathbb{Z}/10$	G(10,2)	G(5,1)	\checkmark		\checkmark	\checkmark
6	2	6	4	3	$\mathbb{Z}/3 \rtimes \mathbb{Z}/4$	G(12, 1)	G(6,1)	X	X	\checkmark	\checkmark
6	2	6	4	4	$\mathbb{Z}/3 \rtimes D_4$	G(24, 8)	G(12, 4)	X	X	\checkmark	\checkmark

Table 1: Data satisfying condition (A) and (B).

\tilde{g}	g	b	p	#	$ ilde{G}$	$ ilde{G}\; \operatorname{Id}$	$G\;{\rm Id}$	(B1)	(B2)	$b \ge 6$	(B)
7	4	0	3	1	$\mathbb{Z}/2 imes \mathbb{Z}/6$	G(12, 5)	G(6,2)	\checkmark		X	\checkmark
7	4	0	3	2	$\mathbb{Z}/4 \rtimes \mathbb{Z}/4$	G(16, 4)	G(8,4)	\checkmark		X	\checkmark
7	4	0	3	3	$\mathbb{Z}/4 \rtimes \mathbb{Z}/4$	G(16, 4)	G(8, 4)	X	\checkmark	X	\checkmark
7	4	0	3	4	$\mathbb{Z}/2 imes Q_8$	G(16, 12)	G(8, 4)	\checkmark		X	\checkmark
7	3	4	4	1	$\mathbb{Z}/8$	G(8,1)	G(4, 1)	\checkmark		X	\checkmark
7	3	4	4	2	$\mathbb{Z}/2 \times \mathbb{Z}/6$	G(12, 5)	G(6,2)	\checkmark		X	\checkmark
7	3	4	4	3	$\mathbb{Z}/12$	G(12, 2)	G(6,2)	\checkmark		X	\checkmark
7	3	4	4	4	$M_{4}(2)$	G(16,6)	G(8,2)	X	X	X	\checkmark
7	3	4	4	6,7	$\mathbb{Z}/4 imes \mathbb{Z}/4$	G(16, 2)	G(8,2)	\checkmark		X	\checkmark
7	3	4	4	8,9	$\mathbb{Z}/4 \rtimes \mathbb{Z}/4$	G(16,4)	G(8,2)	X	\checkmark	X	\checkmark
7	2	8	5	1,3	$\mathbb{Z}/12$	G(12, 2)	G(6,2)	\checkmark		\checkmark	\checkmark
7	2	8	5	2	$\mathbb{Z}/3 \rtimes \mathbb{Z}/4$	G(12, 1)	G(6,1)	X	X	\checkmark	\checkmark
7	2	8	5	4	$\mathbb{Z}/4 \rtimes \mathbb{Z}/4$	G(16, 4)	G(8,3)	X	\checkmark	\checkmark	\checkmark
7	2	8	5	5	$\mathbb{Z}/4 \rtimes \mathbb{Z}/4$	G(16, 4)	G(8,3)	\checkmark		\checkmark	\checkmark
7	1	12	6	1	$\mathbb{Z}/3 \rtimes \mathbb{Z}/4$	G(12, 1)	G(6,1)	X	X	\checkmark	\checkmark
7	1	12	6	2	$\mathbb{Z}/2 imes Q_8$	G(16, 12)	G(8,5)	X	\checkmark	\checkmark	\checkmark
7	1	12	6	3	$\mathbb{Z}/4 \times S_3$	G(24, 5)	G(12, 4)	X	X	\checkmark	\checkmark
7	1	12	6	4	$Q_8 \rtimes \mathbb{Z}/3$	G(24, 3)	G(12,3)	X	X	\checkmark	\checkmark
7	1	12	6	5	$Q_8 \rtimes_3 S_3$	G(48, 41)	G(24, 14)	X	X	\checkmark	\checkmark
8	4	2	4	1,2	$\mathbb{Z}/10$	G(10, 2)	G(5, 1)	\checkmark		X	\checkmark
8	4	2	4	3,4	$\mathbb{Z}/2 imes \mathbb{Z}/6$	G(12, 5)	G(6,2)	\checkmark		X	\checkmark
9	5	0	4	1 - 5	$\mathbb{Z}/2 \times \mathbb{Z}/8$	G(16, 5)	G(8, 1)	\checkmark		X	\checkmark
9	5	0	4	6	$\mathbb{Z}/4 \times \mathbb{Z}/4$	G(16, 2)	G(8,2)	\checkmark		X	\checkmark
9	5	0	4	8	$\mathbb{Z}/4 \rtimes \mathbb{Z}/4$	G(16, 4)	G(8, 4)	\checkmark		X	\checkmark
9	5	0	4	11 - 13	$\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/4$	G(16, 10)	G(8,2)	\checkmark		X	\checkmark
9	5	0	4	19,20	$\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/6$	G(24, 15)	G(12, 5)	\checkmark		X	\checkmark
9	5	0	4	21	$\mathbb{Z}/6 \rtimes \mathbb{Z}/4$	G(24,7)	G(12, 1)	X	\checkmark	X	\checkmark
9	5	0	4	30	$\mathbb{Z}/2 \times (\mathbb{Z}/2)^2 \rtimes \mathbb{Z}/4$	G(32, 22)	G(16, 11)	\checkmark		X	\checkmark
9	5	0	4	34	$\mathbb{Z}/4 \times D_4$	G(32, 25)	G(16, 13)	\checkmark		X	\checkmark
9	5	0	4	35	$\mathbb{Z}/4 \times D_4$	G(32, 25)	G(16, 10)	X	\checkmark	X	\checkmark
9	5	0	4	37	$(\mathbb{Z}/2)^2$. D_4	G(32, 30)	G(16, 13)	X	\checkmark	X	\checkmark
9	5	0	4	$44,\!45$	$\mathbb{Z}/2 \times \mathbb{Z}/3 \rtimes D_4$	G(48, 43)	G(24, 8)	X	\checkmark	X	\checkmark
9	5	0	4	48 - 50	$(\mathbb{Z}/2)^3 \rtimes_2 D_4$	G(64, 73)	G(32, 27)	X	\checkmark	X	\checkmark
9	5	0	4	55	$\mathbb{Z}/4 \rtimes D_8$	G(64, 140)	G(32, 43)	X	\checkmark	X	\checkmark
9	4	4	5	1	$\mathbb{Z}/10$	G(10, 2)	G(5,1)	\checkmark		X	\checkmark
9	4	4	5	2,3	$\mathbb{Z}/12$	G(12, 2)	G(6,2)	\checkmark		X	\checkmark
9	3	8	6	1	$\mathbb{Z}/12$	G(12, 2)	G(6,2)	\checkmark		\checkmark	\checkmark
9	3	8	6	2,3	$\mathbb{Z}/2 \times \mathbb{Z}/8$	G(16, 5)	G(8,2)	\checkmark		\checkmark	\checkmark
9	3	8	6	4	$\mathbb{Z}/4 \times \mathbb{Z}/4$	G(16, 2)	G(8,2)	\checkmark		\checkmark	\checkmark
9	3	8	6	5	$\mathbb{Z}/4 \rtimes \mathbb{Z}/4$	G(16, 4)	G(8,3)	X	X	\checkmark	\checkmark
9	3	8	6	6	$\mathbb{Z}/4 \times S_3$	G(24, 5)	G(12, 4)	X	X	\checkmark	\checkmark
9	3	8	6	7	$\mathbb{Z}/4 \times S_3$	G(24, 5)	G(12, 4)	\checkmark		\checkmark	\checkmark
9	3	8	6	8	$\mathbb{Z}/4 \times D_4$	G(32, 25)	G(16, 11)	\checkmark		\checkmark	\checkmark
9	3	8	6	9	$(\mathbb{Z}/2)^2$. D_4	G(32, 30)	G(16, 11)	X	X	\checkmark	\checkmark
9	2	12	7	1	$\mathbb{Z}/6 \rtimes \mathbb{Z}/4$	G(24,7)	G(12, 4)	X	\checkmark	\checkmark	\checkmark

\tilde{g}	g	b	p	#	$ ilde{G}$	$ ilde{G}\; \operatorname{Id}$	$G \; \operatorname{Id}$	(B1)	(B2)	$b \ge 6$	(B)
10	4	6	6	1,2	$\mathbb{Z}/12$	G(12, 2)	G(6, 2)	\checkmark		\checkmark	\checkmark
10	4	6	6	$3,\!4$	$\mathbb{Z}/3 \times \mathbb{Z}/6$	G(18, 5)	G(9, 2)	\checkmark		\checkmark	\checkmark
10	4	6	6	5	$\mathbb{Z}/3 imes D_4$	G(24, 10)	G(12,5)	X	\checkmark	\checkmark	\checkmark
10	4	6	6	6	$Q_8 \rtimes \mathbb{Z}/3$	G(24, 3)	G(12, 3)	X	X	\checkmark	\checkmark
10	4	6	6	7,9	$\mathbb{Z}/6 \times S_3$	G(36, 12)	G(18, 3)	X	X	\checkmark	\checkmark
10	4	6	6	8	$\mathbb{Z}/6 \times S_3$	G(36, 12)	G(18, 3)	X	\checkmark	\checkmark	\checkmark
10	4	6	6	10	$Q_8 \rtimes S_3$	G(48, 29)	G(24, 12)	X	X	\checkmark	\checkmark
11	6	0	5	1	$\mathbb{Z}/2 \times \mathbb{Z}/8$	G(16, 5)	G(8, 1)	\checkmark		X	\checkmark
11	6	0	5	2	$\mathbb{Z}/2 \times \mathbb{Z}/12$	G(24, 9)	G(12, 2)	\checkmark		X	\checkmark
11	6	0	5	3,4	$\mathbb{Z}/6 \rtimes \mathbb{Z}/4$	G(24,7)	G(12, 1)	X	\checkmark	X	\checkmark
11	3	12	8	1	$\mathbb{Z}/6 \rtimes \mathbb{Z}/4$	G(24,7)	G(12, 4)	X	\checkmark	\checkmark	\checkmark
11	3	12	8	2	$\mathbb{Z}/4 \circ D_8$	G(32, 42)	G(16, 11)	X	X	\checkmark	\checkmark
11	3	12	8	3	$\mathbb{Z}/8$. $(\mathbb{Z}/2)^2$	G(32, 44)	G(16, 11)	X	X	\checkmark	\checkmark
12	6	2	6	1.2	ℤ/14	G(14, 2)	$\overline{G(7,1)}$	\checkmark		X	\checkmark
12	6	2	6	3	$\mathbb{Z}/2 \times \mathbb{Z}/10$	G(20,5)	G(10,2)	√		X	√
13	7	0	6	1	$\mathbb{Z}/2 \times \mathbb{Z}/8$	G(16, 5)	$\frac{G(10, 2)}{G(8, 1)}$			X	· √
13	7	0	6	2	$\mathbb{Z}/2 \times \mathbb{Z}/0$ $\mathbb{Z}/2 \times \mathbb{Z}/10$	G(10, 5) G(20, 5)	G(10, 2)	•		X	•
13	7	0	6	3	$\mathbb{Z}/2 \times \mathbb{Z}/10$ $\mathbb{Z}/2 \times \mathbb{Z}/12$	G(20, 9) G(24, 9)	G(12, 2) G(12, 2)	•		X	•
13	7	0	6	6	$D_4 \rtimes \mathbb{Z}/4$	G(21,9) G(32,9)	G(12,2) G(16,7)	• •		X	•
13	7	0	6	11	$(\mathbb{Z}/4 \times \mathbb{Z}/4) \rtimes \mathbb{Z}/2$	G(32, 24)	G(16, 13)	• √		X	• √
13	7	0	6	12	$(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes Q_{\circ}$	G(32, 29)	G(16, 13) G(16, 13)	X	1	X	• √
13	7	0	6	17	$(\underline{\mathbb{Z}}/\underline{\mathbb{Z}}/\underline{\mathbb{Z}}/\underline{\mathbb{Z}})/(\underline{\mathbb{Q}})$ $D_4 \rtimes D_4$	G(64, 130)	G(32, 43)	X	· √	X	• √
13	6	4	7	1	$\mathbb{Z}/18$	$\frac{G(01, 100)}{G(18, 2)}$	$\frac{G(92, 10)}{G(9, 1)}$		•	X	· ·
13	6	4	7	2	$\mathbb{Z}/2 \times \mathbb{Z}/12$	G(10, 2) G(24, 9)	G(12, 5)	•		X	•
13	6	4	7	3	$\mathbb{Z}/2 \times \mathbb{Z}/12$ $\mathbb{Z}/2 \times \mathbb{Z}/12$	G(24, 9) G(24, 9)	G(12, 3) G(12, 2)	•		X	•
12	5	8	8	1	$\frac{\mathbb{Z}/2 \times \mathbb{Z}/12}{O_2 \times \mathbb{Z}/3}$	$\frac{G(24,3)}{G(24,3)}$	$\frac{G(12,2)}{G(12,3)}$	•	Y	1	•
13	5	8	8	2	$(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/8$	G(24, 5) G(32, 5)	G(12,3) G(16,3)	X	X	•	v .(
13	5	8	8	2	$\mathbb{Z}/8 \circ D_4$	C(32, 38)	C(16, 10)	X	X	•	•
13	5	8	8	4	$\mathbb{Z}/2 \times SD_{1c}$	G(32, 30) G(32, 40)	G(10, 10) G(16, 11)	X	X	•	•
13	5	8	8	5	$(\mathbb{Z}/2 \times \mathbb{SD}_{16})$ $(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes O_{2}$	C(32, 40) C(32, 20)	C(16, 11) C(16, 11)	X	<u></u>	•	•
13	5	8	8	6	$(\square/2 \land \square/2) \land Q_8$ $\overline{Z}/A A$	G(32, 23) G(48, 33)	G(10, 11) G(24, 13)	X	v X	•	•
13	5	8	8	7	$D_4 \rtimes D_4$	G(40, 33) G(64, 130)	G(24, 13) G(32, 27)	X	X	•	•
14	7	2	7	1	$\frac{D_4 \land D_4}{\mathbb{Z}/18}$	C(18, 2)	$\frac{C(02,21)}{C(0,1)}$	1	21	V	•
14	1	2	1	1	<u>Z/10</u>	G(16, 2)	$\frac{G(9,1)}{G(9,1)}$	v		<u>Λ</u>	v
14	0	0	8	1	<u>Z/10</u>	$\frac{G(10,1)}{G(21,0)}$	$\frac{G(8,1)}{G(12,2)}$	√		√ 	
15	8	0	7	1,2	$\mathbb{Z}/2 \times \mathbb{Z}/12$	G(24,9)	G(12,2)	✓		<i>X</i>	√
15	7	4	8	1	$\mathbb{Z}/16$	G(16, 1)	G(8,1)	\checkmark		X	√
15	7	4	8	2	$\frac{\mathbb{Z}/2 \times \mathbb{Z}/12}{\mathbb{Z}/12}$	G(24,9)	G(12,5)	√ 		X	✓
15	5	12	10	1	$\mathbb{Z}/4 \times D_5$	G(40,5)	G(20, 4)	X	X	\checkmark	√
15	5	12	10	2	$\mathbb{Z}/4 \circ D_{12}$	G(48, 37)	G(24, 14)	X	X	√	\checkmark
15	5	12	10	3	$D_4 \rtimes_2 S_3$	G(48, 39)	G(24, 14)	X	X	\checkmark	\checkmark
17	9	0	8	1	$\mathbb{Z}/2 \times \mathbb{Z}/12$	G(24,9)	G(12, 2)	\checkmark		X	\checkmark
17	9	0	8	3	$\mathbb{Z}/2 \times \mathbb{Z}/4 \times \mathbb{Z}/4$	G(32, 21)	G(16, 2)	\checkmark		X	\checkmark
17	9	0	8	4	$\mathbb{Z}/4 \times Q_8$	G(32, 26)	G(16, 10)	X	\checkmark	X	\checkmark
17	9	0	8	6	$D_6 \rtimes \mathbb{Z}/4$	G(48, 14)	G(24, 6)	✓ 		X	√
17	9	0	8	12	$A_4 \rtimes \mathbb{Z}/4$	G(48, 30)	G(24, 12)	X	\checkmark	X	\checkmark
17	9	0	8	17	$(\mathbb{Z}/2)^43 (\mathbb{Z}/2)^2$	G(64, 71)	G(32, 34)	✓ 		X	√
17	9	0	8	31	$M_4(2) \rtimes D_4$	G(128,738)	G(64, 134)	X	\checkmark	X	\checkmark
17	5	16	12	1	$D_{4 \cdot 10} D_{4}$	G(64, 137)	G(32, 27)	X	X	\checkmark	\checkmark
19	9	4	10	1	$\mathbb{Z}/20$	G(20, 2)	G(10, 2)	\checkmark		X	\checkmark
19	9	4	10	2	$\mathbb{Z}/4 \times D_5$	G(40, 5)	G(20, 4)	\checkmark		X	\checkmark

\tilde{g}	g	b	p	#	$ ilde{G}$	$ ilde{G}\; \operatorname{Id}$	$G\; {\rm Id}$	(B1)	(B2)	$b\geq 6$	(B)
21	11	0	10	1	$\mathbb{Z}/4 \times \mathbb{Z}/8$	G(32, 3)	G(16, 5)	\checkmark		X	\checkmark
21	11	0	10	5	$\mathbb{Z}/6$. D_4	G(48, 19)	G(24, 8)	X	\checkmark	X	\checkmark
21	11	0	10	6	$\mathbb{Z}/4 imes D_8$	G(64, 118)	G(32, 39)	\checkmark		X	\checkmark
21	9	8	12	1	$S_3 imes \mathbb{Z}/8$	G(48, 4)	G(24, 5)	X	X	\checkmark	\checkmark
21	9	8	12	2	$\mathbb{Z}/24 \rtimes \mathbb{Z}/2$	G(48, 6)	G(24, 6)	X	X	\checkmark	\checkmark
21	9	8	12	3	$\mathbb{Z}/8 \circ D_8$	G(64, 124)	G(32, 25)	X	X	\checkmark	\checkmark
21	9	8	12	4	$D_47 D_4$	G(64, 133)	G(32, 27)	X	X	\checkmark	\checkmark
21	9	8	12	5	$\mathbb{Z}/8$.12 D_4	G(64, 176)	G(32, 34)	X	X	\checkmark	\checkmark
21	9	8	12	6	$\mathbb{Z}/4$.6 S_4	G(96, 192)	G(48, 48)	X	X	\checkmark	\checkmark
25	13	0	12	6	$(\mathbb{Z}/2)^3 \rtimes Q_8$	G(64, 74)	G(32, 27)	X	\checkmark	X	\checkmark
25	9	16	16	1	$(\mathbb{Z}/2 \times \mathbb{Z}/8) \cdot D_4$	G(128, 749)	G(64, 73)	X	X	\checkmark	\checkmark
29	13	8	16	1	Q_{16} . D_4	G(128, 925)	G(64, 128)	X	X	\checkmark	\checkmark
31	13	12	18	1	D_6 . D_6	G(144, 141)	G(72, 46)	X	X	\checkmark	\checkmark

Table 2: Number of data satisfying condition (A).

$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	\tilde{g}	g	b	p	#	•	\tilde{g}	g	b	p	#	•	\tilde{g}	g	b	p	#
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	2	0	6	2	4	-	9	5	0	4	58	•	17	9	0	8	35
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	3	2	0	1	2	-	9	4	4	5	3		17	5	16	12	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	3	1	4	2	8		9	3	8	6	9		19	10	0	9	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	3	0	8	3	1		9	2	12	7	1		19	9	4	10	4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	4	2	2	2	2	-	10	4	6	6	10	-	21	11	0	10	10
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	4	1	6	3	3		11	6	0	5	6	-	21	9	8	12	6
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	4	0	10	4	1		11	5	4	6	2		23	11	4	12	4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	3	0	2	19	-	11	3	12	8	3		25	13	0	12	18
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	2	4	3	3		12	6	2	6	3	-	25	9	16	16	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	1	8	4	9		13	7	0	6	22	-	29	15	0	14	4
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	0	12	5	1		13	6	4	7	3		29	13	8	16	1
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	3	2	3	2		13	5	8	8	7		31	15	4	16	1
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	2	6	4	4		14	7	2	7	1	-	31	13	12	18	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	7	4	0	3	5	-	14	6	6	8	1		33	17	0	16	5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	7	3	4	4	11		15	8	0	7	2	-	39	19	4	20	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	7	2	8	5	5		15	7	4	8	3		41	21	0	20	2
8 4 2 4 5 16 8 2 8 2 49 25 0 24	7	1	12	6	5	_	15	5	12	10	3	_	47	23	4	24	1
	8	4	2	4	5	-	16	8	2	8	2	-	49	25	0	24	2

References

- E. Arbarello, M. Cornalba, and P. A. Griffiths. Geometry of algebraic curves. Vol. II, volume 268 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, New York, 2011.
- [2] Bardelli, F., Ciliberto, C., Verra, A., Curves of minimal genus on a general abelian variety, Compos. Math. 96 (1995), no. 2, 115–147.
- [3] J. S. Birman. Braids, links, and mapping class groups. Princeton University Press, Princeton, N.J., 1974. Annals of Mathematics Studies, No. 82.
- [4] S. A. Broughton. The equisymmetric stratification of the moduli space and the Krull dimension of mapping class groups. *Topology Appl.*, 37(2):101–113, 1990.
- [5] F. Catanese, M. Lönne, and F. Perroni. Irreducibility of the space of dihedral covers of the projective line of a given numerical type. Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl., 22(3):291–309, 2011.
- [6] F. Catanese, M. Lönne, and F. Perroni. Genus stabilization for the components of moduli spaces of curves with symmetries. Algebr. Geom., 3(1):23–49, 2016.
- [7] Colombo, E., Frediani P., A bound on the dimension of a totally geodesic submanifold in the Prym locus. Collectanea Mathematica, 70, n.1, (2019), 51-57. DOI: 10.1007/s13348-018-0215-0.
- [8] Colombo, E., Frediani P., Second fundamental form of the Prym map in the ramified case. To appear in "Galois Covers, Grothendieck-Teichmueller Theory and Dessins d'Enfants - Interactions between Geometry, Topology, Number Theory and Algebra", Springer Proceedings in Mathematics & Statistics. arXiv:1812.07402.
- [9] E. Colombo, P. Frediani, and A. Ghigi. On totally geodesic submanifolds in the Jacobian locus. International Journal of Mathematics, 26 (2015), no. 1, 1550005 (21 pages).
- [10] Colombo, E., Frediani, P., Ghigi, A., Penegini M., Shimura curves in the Prym locus. Communications in Contemporary Mathematics, Vol. 21, No. 2 (2019) 1850009 (34 pages). DOI: 10.1142/S0219199718500098.
- [11] E. Colombo, G. P. Pirola, and A. Tortora. Hodge-Gaussian maps. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 30(1):125–146, 2001.
- [12] P. Frediani, A. Ghigi and M. Penegini. Shimura varieties in the Torelli locus via Galois coverings. Int. Math. Res. Not. 2015, no. 20, 10595-10623.
- [13] P. Frediani, A. Ghigi and I Spelta. Infinitely many Shimura varieties in the Jacobian locus for $g \le 4$. arXiv:1910.13245. To appear in Ann. Scuola Norm. Sup. Pisa Cl. Sci.
- [14] Frediani, Paola; Pirola Gian Pietro. On the geometry of the second fundamental form of the Torelli map. arXiv:1907.11407.
- [15] Friedman, Robert; Smith, Roy, The generic Torelli theorem for the Prym map. Invent. Math. 67 (1982), no. 3, 473-490.
- [16] Ghigi, Alessandro; On some differential-geometric aspects of the Torelli map. Boll. Unione Mat. Ital. 12 (2019), no. 1-2, 133?144.
- [17] Ghigi, Alessandro; Pirola, Gian Pietro; Torelli, Sara. Roy, Totally geodesic subvarieties in the moduli space of curves. To appear on Communications in Contemporary Mathematics. https://doi.org/10.1142/S0219199720500200.
- [18] G. González Díez and W. J. Harvey. Moduli of Riemann surfaces with symmetry. In Discrete groups and geometry (Birmingham, 1991), volume 173 of London Math. Soc. Lecture Note Ser., pages 75–93. Cambridge Univ. Press, Cambridge, 1992.
- [19] A. Ikeda, Global Prym-Torelli Theorem for double coverings of elliptic curves, preprint, arXiv:1812.07837.
- [20] V. I. Kanev. A global Torelli theorem for Prym varieties at a general point. Izv. Akad. Nauk SSSR Ser. Mat., 46(2):244–268, 431, 1982.
- [21] H. Lange and A. Ortega. Prym varieties of cyclic coverings. Geom. Dedicata, 150:391–403, 2011.
- [22] MAGMA Database of Small Groups; http://magma.maths.usyd.edu.au/magma/htmlhelp/ text404.htm.script.
- [23] V.O. Marcucci, G.P. Pirola. Generic Torelli theorem for Prym varieties of ramified coverings. Compos. Math. 148 (2012), no. 4, 1147–1170.
- [24] Marcucci, V, Naranjo, J.C., Prym varieties of double coverings of elliptic curves, Int. Math. Res. Notices 6 (2014), 1689-1698.
- [25] Mohajer, A, On Shimura subvarieties of the Prym locus. arXiv:1804.10131v2.
- [26] B. Moonen. Linearity properties of Shimura varieties. I. J. Algebraic Geom., 7(3):539-567, 1998.
- [27] B. Moonen and F. Oort. The Torelli locus and special subvarieties. In Handbook of Moduli: Volume II, pages 549–94. International Press, Boston, MA, 2013.
- [28] D. Mumford. A note of Shimura's paper "Discontinuous groups and abelian varieties". Math. Ann., 181:345–351, 1969.
 [29] D. S., Nagaraj; S. Ramanan, S. Polarisations of type (1,2,...,2) on abelian varieties. Duke Math. J. 80 (1995), no. 1,
- 157-194.[30] Naranjo, J.C., Ortega, A., Verra, A., Generic injectivity of the Prym map for double ramified coverings. Trans of AMS. DOI: 10.1090/tran/7459.
- [31] J.C. Naranjo, A. Ortega, Global Prym-Torelli for double coverings ramified in at least 6 points, arXiv:2005.11108.
- [32] M. Penegini. Surfaces isogenous to a product of curves, braid groups and mapping class groups. In Beauville surfaces and groups, volume 123 of Springer Proc. Math. Stat., pages 129–148. Springer, Cham, 2015.
- [33] A. M. Rojas. Group actions on Jacobian varieties, Rev. Mat. Iber. 23 (2007), 397-420.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PAVIA, VIA FERRATA 5, I-27100 PAVIA, ITALY *E-mail address*: paola.frediani@unipv.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PAVIA, VIA FERRATA 5, I-27100, PAVIA, ITALY *E-mail address*: g.grosselli@campus.unimib.it