MINIMALITY AND UNIQUENESS FOR DECOMPOSITIONS OF SPECIFIC TERNARY FORMS

ELENA ANGELINI AND LUCA CHIANTINI

ABSTRACT. The paper deals with the computation of the rank and the identifiability of a specific ternary form. Often, one knows some short Waring decomposition of a given form, and the problem is to determine whether the decomposition is minimal and unique. We show how the analysis of the Hilbert-Burch matrix of the set of points representing the decomposition can solve this problem in the case of ternary forms. Moreover, when the decomposition is not unique, we show how the procedure of liaison can provide alternative, maybe shorter, decompositions. We give an explicit algorithm that tests our criterion of minimality for the case of ternary forms of degree 9. This is the first numerical case in which a new phaenomenon appears: the span of 18 general powers of linear forms contains points of (subgeneric) rank 18, but it also contains points whose rank is 17, due to the existence of a second shorter decomposition which is completely different from the given one.

1. INTRODUCTION

The paper deals with homogeneous polynomials (forms) F, which will be also seen as symmetric tensors, over the complex field \mathbb{C} . Our analysis starts with a given *Waring* expression of F in terms of powers of linear forms

$$F = \lambda_1 L_1^d + \dots + \lambda_r L_r^d$$

(d is the degree of F). Our target is to determine the minimality and the uniqueness of the expression, up to a scalar multiplication of the L_i 's. Indeed, in our setting minimality is a consequence of uniqueness. So, when uniqueness holds, r is the (Waring) rank of F.

The determination of the rank of a given form, as well as the the uniqueness of a Waring expression, is relevant in several aspects of tensor theory related to physics, signal processing, statistics, chemistry, artificial intelligence, etc. For some examples of these relations, taken from the huge literature on the subject, let us mention the papers [2], [3], [6], [33].

Our analysis, besides its theoretical interest, will produce the following concrete application. There are several situations in which one knows some Waring expression of a specific form F. For instance, this happens when F splits in a sum of blocks whose rank is well known, or when one applies to F some heuristic algorithm, e.g. based on local regression. In these cases, however, the question about the minimality of the expression remains open. We will see in examples (see Section 5.2) that a local analysis of the Waring expression cannot guarantee its minimality,

²⁰⁰⁰ Mathematics Subject Classification. 14N07, 14J70, 14C20, 14N05, 15A69, 15A72.

This work was supported by the National Group for Algebraic and Geometric Structures, and their Applications (GNSAGA – INdAM) and by the Italian PRIN 2015 - Geometry of Algebraic Varieties (B16J15002000005).

except when the length r is rather small (as, e.g., in the Kruskal's criterion). We will produce a method which, in principle, can solve the minimality problem, and then guarantee that r is the rank of F, for all Waring expressions of ternary forms. The consequent algorithm that one can construct depends on the degree d. In the last section, we produce a concrete example of the algorithm, for ternary forms of degree 9.

We attack the problem with tools of projective algebraic geometry. Thus, we start with the polynomial ring $R = \mathbb{C}[x_0, \ldots, x_n]$ and we indicate with R_d the homogeneous piece of degree d. Any linear form L determines a point (which, by abuse of notation, we still indicate with L) in the projective space \mathbb{P}^n over the linear space R_1 of forms of degree 1. In this notation, F corresponds to a point in the projective space over R_d . We identify $\mathbb{P}(R_d)$, of (projective) dimension $N = \binom{n+d}{d} - 1$, with $\mathbb{P}(Sym^d(R_1))$. The map $\nu_d : \mathbb{P}^n \to \mathbb{P}^N$ which sends a linear form L to its d-th power is universally known as the d-th Veronese map.

A Waring expression $F = \lambda_1 L_1^d + \cdots + \lambda_r L_r^d$ of F determines a subset $A = \{L_1, \ldots, L_r\} \subset \mathbb{P}^n$ such that F sits in the span of $\nu_d(A)$. In our notation, we will say that A is a *decomposition* of length r of F. The decomposition A is *non-redundant* if F does not lie in the span of $\nu_d(A')$, for any proper subset $A' \subset A$. The decomposition A is *minimal* (resp. *unique*) if there are no other subsets $B \subset \mathbb{P}^n$, of length r' < r (resp. $r' \leq r$) such that F belongs to the span of $\nu_d(B)$. So, when A is minimal, its length is the (Waring) rank of F.

There are methods that, in some setting, can prove that a decomposition $A = \{L_1, \ldots, L_r\}$ of F is unique, and also minimal. For instance, one can take several flattenings of the tensors associated to F, L_1^d, \ldots, L_r^d and use them to prove the uniqueness, as in [20], [21]. Alternatively, one can study the catalecticant map α defined by some partial derivatives of F, and the intersection of the image of α with a suitable Veronese variety, as in [28], [29]. The most famous method to prove uniqueness goes back to Kruskal, [27]. Kruskal used the Kruskal's rank of the matrix whose entries are the coefficients of the L_i 's (i.e. the coordinates of the L_i 's, as points in \mathbb{P}^n) to produce an inequality which, if satisfied, guarantees that the decomposition is unique, and minimal. Kruskal's criterion has been refined in [14], by taking into account the matrices defined by powers of the L_i 's.

All the aforementioned methods have a common problem: they will not provide an answer when the degree d and the length r grow.

From the point of view of projective geometry, it is clear that the Waring rank is constant in a Zariski open subset of \mathbb{P}^N , i.e. it is constant outside a subset defined by algebraic equations: a small subset, in any reasonable metric. Denote with r_g the generic value of the rank, which holds in a dense subset of \mathbb{P}^N . Then, it is almost straightforward that a decomposition cannot be unique when $r > r_g$ (too many parameters). When $r = r_g$ generic uniqueness can hold, but it is a quite rare phenomenon, completely classified (see [23]). On the contrary, for subgeneric values $r < r_g$, uniqueness holds for a minimal Waring expression of a sufficiently general form F, except for a short list of cases (see [15]). On the other hand, for a specific decomposition of a specific form F, whose rank is unknown, all the known methods can guarantee its uniqueness or minimality only when the length r is much smaller than r_g (see Section 3 of [4], for a concrete bound). The analysis that we propose overruns this difficulty, and will guarantee the minimality, and also the uniqueness, of a given decomposition, in principle for all $r < r_g$. We will give a concrete example for forms of degree 9 in three variables.

The theoretical situation can be summarized as follows. Geometrically, the Waring expression $F = \lambda_1 L_1^d + \cdots + \lambda_r L_r^d$ presents F as a point of the (linear) span of the set $\nu_d(A)$. When r is big, even if the L_i 's are general, it turns out that the span of $\nu_d(A)$ contains both points F for which the decomposition A is unique, and points $F' = \mu_1 L_1^d + \cdots + \mu_r L_r^d$ for which uniqueness, and also minimality, do not hold (though A is still a non-redundant decomposition of G). Examples of this situation are described in Section 3 below. It may happen indeed (Case 2 of Proposition 3.5) that even if F' is non-redundantly spanned by $\nu_d(A)$, yet there exists another decomposition B of F', completely different from A and with less summands. Thus, methods based on the geometrical analysis of A alone (as the Kruskal's or the catalecticant approaches) cannot distinguish between forms F, F' in the span of $\nu_d(A)$, so they will fail to guarantee e.g. minimality, when the span contains points with different behavior. The answer can be obtained only by analyzing A and the coefficients λ_i 's of the Waring expression.

Our analysis follows the guidelines introduced in [4]. We attack the problem with a set of tools typical for the study of the geometry of finite sets in projective spaces: Hilbert-Burch matrices, the Cayley-Bacharach property, and liaison. An extensive illustration of the interplay between decompositions and the geometry of finite sets can be found in the book [26]. Papers [10], [8], [9] are based on the study of Hilbert functions of points. What is really new in the present paper, as well as in [4], is the observation that given a finite set $A \subset \mathbb{P}^2$, the possible alternative (maybe even shorter) decompositions of any ternary form F in the span of $\nu_d(A)$ can be recovered geometrically, by playing with the Hilbert-Burch matrix of A and liaison. Thus, we can characterize the (algebraic) subset Θ of the span of $\nu_d(A)$ consisting of forms for which A is non-redundant, but yet they have an alternative decomposition of length r' < r. Moreover, we produce concrete algorithms which guarantee that a form F does not lie in the 'bad' set Θ .

In the paper, we illustrate in details the procedure in the case of ternary forms of degree 9. For such forms, the generic rank is $r_g = 19$, see [1]. We consider the highest sub-generic value r = 18. We show that for any generic choice of a subset Aof 18 points in \mathbb{P}^2 (the projective space of linear forms in three variables) the span Λ of $\nu_9(A)$ contains: (i) points F for which the decomposition A is minimal and unique, (ii) points F' for which A in non-redundant, but there exists an alternative decomposition $B \neq A$, of length 18, (iii) points F'' for which A is non-redundant, but there exists an alternative decomposition of length 17 (so their rank is smaller than 18). Moreover, we prove that the (closure of) the set of points satisfying (ii) is a hypersurface of Λ , and it is a birational image of a Grassmannian of lines (see Theorem 4.5 below). We also determine properties of the closure Θ of the set of points satisfying (iii), which is a subvariety of Λ .

Finally, we produce an algorithm to test if a given form $F = \lambda_1 L_1^9 + \cdots + \lambda_{18} L_{18}^9$ in the span Λ of $\nu_9(\{L_1, \ldots, L_{18}\})$ lies in the 'bad' locus Θ defined above, i.e. its rank is smaller than 18. The algorithm analyses both A and the coefficients λ_i 's of the Waring expression. When the answer provided by the algorithm is negative, we can conclude that A is a minimal decomposition of F, so that F has rank 18. Similar algorithms for detecting the uniquess of the decomposition can be easily produced, though they need more parameters. Finally, when a second decomposition of length $r' \leq r$ exists, we show how we can recover the new decomposition from the known one A (see the end of the examples in Section 5.2).

The paper is structured as follows. In Sect. 2 we introduce our setting and recall the main tools, coming from classical algebraic geometry, that characterize our analysis. In particular, we mention the Hilbert function and the Cayley-Bacharach property for finite sets in the projective space, the notion of Hilbert-Burch matrix and liason theory for sets of points in the projective plane. Sect. 3 is the core of the paper: we develop our analysis for ternary forms of degree 9, dealing with ranks that are outside the range of applicability of the celebrated Kruskal's criterion. Sect. 4 disposes on the case in which two decompositions of a ternary nonic intersect. Finally, Sect. 5 is devoted to the effective algorithm we developed according to the criterion of minimality (and uniqueness) obtained in Sect. 3. Several numerical examples are presented.

Acknowledgements. This work was supported by the National Group for Algebraic and Geometric Structures, and their Applications (GNSAGA – INdAM) and by the Italian PRIN 2015 - Geometry of Algebraic Varieties (B16J15002000005).

2. Preliminaries

2.1. Notation. For any finite set A we denote with $\ell(A)$ the cardinality. For $d, n \in \mathbb{N}$, let \mathbb{C}^{n+1} be the space of linear forms in x_0, \ldots, x_n , so that $S^d \mathbb{C}^{n+1}$ is the space of forms of degree d in n+1 variables over \mathbb{C} .

Every form $T \in S^d \mathbb{C}^{n+1}$ defines an element of $\mathbb{P}(S^d \mathbb{C}^{n+1}) \cong \mathbb{P}^N$ $(N = \binom{n+d}{d} - 1)$, which, by abuse, we still denote by T.

We denote with $\nu_d : \mathbb{P}^n \to \mathbb{P}^N$ the *Veronese embedding* of \mathbb{P}^n of degree d, which is given by

$$\nu_d([a_0x_0 + \ldots + a_nx_n]) = [(a_0x_0 + \ldots + a_nx_n)^d].$$

For any subset $Z \subset \mathbb{P}^N$ we denote with $\langle Z \rangle$ the linear span of Z. For instance, if $A = \{P_1, \ldots, P_r\} \subset \mathbb{P}^n$ is a finite set, $\langle \nu_d(A) \rangle$ is the linear space in \mathbb{P}^N spanned by the points $\nu_d(P_1), \ldots, \nu_d(P_r)$.

With the above notations we give the following definitions.

Definition 2.1. Let $A \subset \mathbb{P}^n$ be a finite set, and $T \in S^d \mathbb{C}^{n+1}$ a form of degree d. We say that A computes T if $T \in \langle \nu_d(A) \rangle$.

We say that a set A which computes T is *non-redundant* if there are no proper subsets A' of A such that A' computes T.

We say that a set A which computes T is *minimal* if there are no sets B, with $\ell(B) < \ell(A)$, such that B computes T.

If A computes T and it is *minimal*, the cardinality $\ell(A)$ is called the (Waring) rank of T. In this case we say that A computes the rank of T.

Remark 2.2. If $A \subset \mathbb{P}^n$ is a finite set that computes the rank of T then A is non-redundant, and the points of $\nu_d(A)$ are linearly independent, i.e.

$$\dim(\langle \nu_d(A) \rangle) = \ell(A) - 1.$$

Definition 2.3. A form T is *identifiable* if there exists a unique set A that computes the rank of T.

2.2. Kruskal's criterion for symmetric tensors. One of the most used criteria for detecting the identifiability of a form T is here described, with some extensions, in geometric terms and adapted to the case of forms (= symmetric tensors).

We start with a definition which is the geometric analogue of the Kruskal's rank of a matrix.

Definition 2.4. The *d*-th Kruskal's rank of a finite set $A \subset \mathbb{P}^n$ is

 $k_d(A) = \max\{k \mid \text{for all } A' \subset A \text{ with } \ell(A') \le k, \text{ then } \dim\langle \nu_d(A') \rangle = \ell(A') - 1\}.$

In other words, $k_d(A)$ is the maximal k such that the image under ν_d of any subset of A of cardinality at most k is linearly independent.

Remark 2.5. For any d, it holds that $k_d(A) \leq \min\{N+1, \ell(A)\}$.

Moreover, if $k_d(A) = \min\{N + 1, \ell(A)\}$ (i.e. $k_d(A)$ is maximal), then for all $A' \subset A$ the *d*-th Kruskal's rank $k_d(A')$ is also maximal.

If A is sufficiently general, then $k_d(A) = \min\{N+1, \ell(A)\}$ i.e. the d-th Kruskal's rank is maximal (see e.g. Lemma 4.4 of [14]).

The Kruskal's rank is fundamental in the statement of the reshaped Kruskal's criterion.

Theorem 2.6 (Reshaped Kruskal's Criterion, see [14]). Assume $d \geq 3$ and let $A \subset \mathbb{P}^n$ be a non-redundant set computing $T \in \mathbb{P}(S^d \mathbb{C}^{n+1})$. Fix a partition $d = d_1 + d_2 + d_3$ with $d_1 \geq d_2 \geq d_3 \geq 1$. If

(1)
$$\ell(A) \le \frac{k_{d_1}(A) + k_{d_2}(A) + k_{d_3}(A) - 2}{2}$$

then T has rank $\ell(A)$ and it is identifiable.

In the case of ternary forms, the reshaped Kruskal's criterion has been recently extended in [4], [7], [31].

Theorem 2.7. Let $T \in \mathbb{P}(S^d \mathbb{C}^3)$ and let $A \subset \mathbb{P}^2$ be a non-redundant set computing T. Then T is identifiable of rank r if one of the following holds:

• d = 2m is even, $k_{m-1}(A) = \min\{\binom{m+1}{2}, r\}, h_A(m) = r \le \binom{m+2}{2} - 2;$

•
$$d = 2m + 1$$
 is odd, $k_m(A) = \min\{\binom{m+2}{2}, r\}, h_A(m+1) = r \leq \binom{m+2}{2} + \lfloor \frac{m}{2} \rfloor$

Theorem 2.8. Let $T \in \mathbb{P}(S^d\mathbb{C}^3)$ and let $A \subset \mathbb{P}^2$ be a non-redundant set computing T. Then A computes the rank of T if one of the following holds:

• d = 2m is even, and $h_A(m) = r(\leq \binom{m+2}{2});$

•
$$d = 2m + 1$$
 is odd, $k_m(A) = \min\{\binom{m+2}{2}, r\}, h_A(m+1) = r \le \binom{m+2}{2} + \lceil \frac{m}{2} \rceil$

2.3. The Hilbert function of finite sets and its difference.

Definition 2.9. The *evaluation map* of degree d on an ordered finite set of vectors $Y = \{Y_1, \ldots, Y_\ell\} \subset \mathbb{C}^{n+1}$ is the linear map given by

$$ev_Y(d): S^d \mathbb{C}^{n+1} \longrightarrow \mathbb{C}^\ell$$

 $ev_Y(d)(F) = (F(Y_1), \dots, F(Y_\ell))$

Definition 2.10. Let Y be a set of homogeneous coordinates for a finite set Z of \mathbb{P}^n . The *Hilbert function* of Z is the map

$$h_Z:\mathbb{Z}\longrightarrow\mathbb{N}$$

such that $h_Z(j) = 0$, for j < 0, $h_Z(j) = rank(ev_Y(j))$, for $j \ge 0$. It is elementary indeed that the map depends only on Z, and not on the choice of a set of homogeneous coordinates for the points of Z.

We will often use the first difference of the Hilbert function Dh_Z , given by

 $Dh_Z(j) = h_Z(j) - h_Z(j-1), \ j \in \mathbb{Z}.$

We collect some useful elementary properties of h_Z and Dh_Z . They are well know and contained (sparsely) in the literature, see e.g. [26], [30]. A summary can be found in [13].

Remark 2.11. (i) $Dh_Z(j) \ge 0$ for all j; $h_Z(j) = Dh_Z(j) = 0$ for j < 0; $h_Z(0) = Dh_Z(0) = 1$.

(ii) $h_Z(j) = \ell(Z)$ for all $j \gg 0$, so that $Dh_Z(j) = 0$ for $j \gg 0$ and $\sum Dh_Z(j) = \ell(Z)$.

(iii) If $Z' \subset Z$, then for all $j \in \mathbb{Z}$ we have $h_{Z'}(j) \leq h_Z(j)$ and $Dh_{Z'}(j) \leq Dh_Z(j)$.

Proposition 2.12. For any i > 0 such that $Dh_Z(i) \leq i$, it holds

$$Dh_Z(i) \ge Dh_Z(i+1),$$

i.e. the function Dh_Z becomes non-increasing from *i* on. Therefore, if $Dh_Z(i) = 0$, then $Dh_Z(j) = 0$ for any $j \ge i$.

The following proposition is a straightforward application of the Grassmann formula in projective spaces.

Proposition 2.13. Let $A, B \subset \mathbb{P}^n$ be disjoint finite sets such that both $\nu_d(A)$ and $\nu_d(B)$ are linearly independent. Set $Z = A \cup B$. For any $d \in \mathbb{N}$,

$$\dim(\langle v_d(A)\rangle \cap \langle v_d(B)\rangle) = \ell(Z) - h_Z(d) - 1.$$

As pointed out e.g. in [4], it follows that when A, B are two disjoint sets that compute a form T, then $\langle v_d(A) \rangle \cap \langle v_d(B) \rangle$ is non empty, thus the union $Z = A \cup B$ satisfies $h_Z(d) < \ell(Z)$. It follows:

Proposition 2.14. Let $T \in S^d \mathbb{C}^{n+1}$ and let $A, B \subset \mathbb{P}^n$ be disjoint non-redundant finite sets computing T. Pose $Z = A \cup B \subset \mathbb{P}^n$. Then $Dh_Z(d+1) > 0$.

The following Theorem has been generalized to sets of points in any projective space \mathbb{P}^n (see [11]), but we will need only the case n = 2.

Theorem 2.15 (Davis, [18]). Let $Z \subset \mathbb{P}^2$ be a finite set. Assume that for some j > 0 one has $0 < Dh_Z(j) = Dh_Z(j+1) \le j$. Then Z splits in a union $Z = Z_1 \cup Z_2$ where Z_1 lies on a curve of degree $e = Dh_Z(j)$

$$Dh_{Z_1}(i) = Dh_Z(i)$$
 for $i \ge j$.

The consequences of Davis' Theorem in our analysis are resumed in the following Proposition (see [4] Proposition 2.20).

Proposition 2.16. Let $T \in S^d \mathbb{C}^3$ and let $A, B \subset \mathbb{P}^2$ be disjoint, non-redundant finite sets computing T. Pose $Z = A \cup B$. Then there are no indices $j \leq d$ such that $0 < Dh_Z(j) = Dh_Z(j+1) < j$.

and

Proof. (sketch) Assume on the contrary that j exists. Then, by Davis' Theorem and the Grassmann formula, there exists a proper subset $Z_1 \subset Z$ such that, if $A_1 = A \cap Z_1$ and $B_1 = B \cap Z_1$, then

$$\langle v_d(A) \rangle \cap \langle v_d(B) \rangle = \langle v_d(A_1) \rangle \cap \langle v_d(B_1) \rangle.$$

Thus, at least one between A, B cannot be non-redundant.

2.4. The Cayley-Bacarach property.

Definition 2.17. A finite set $Z \subset \mathbb{P}^n$ satisfies the *Cayley-Bacharach property in degree i*, abbreviated as CB(i), if for all $P \in Z$, it holds that every form of degree *i* vanishing at $Z \setminus \{P\}$ also vanishes at *P*.

The main consequence of the Cayley-Bacharach property on the Hilbert function of a set Z is contained in the following two results, proved in [5], by means of a deep result in [24].

Theorem 2.18. If a finite set $Z \subset \mathbb{P}^n$ satisfies CB(i), then for any j such that $0 \leq j \leq i+1$ we have

$$Dh_Z(0) + Dh_Z(1) + \dots + Dh_Z(j) \le Dh_Z(i+1-j) + \dots + Dh_Z(i+1).$$

Corollary 2.19. Let $T \in S^d \mathbb{C}^3$ and let $A \subset \mathbb{P}^2$ be a non-redundant finite set computing T. Let $B \subset \mathbb{P}^2$ be another non-redundant finite set computing T and assume $A \cap B = \emptyset$. Then $Z = A \cup B$ satisfies the Cayley-Bacharach property CB(d).

2.5. The Hilbert-Burch matrix of a set of points in \mathbb{P}^2 . Let $A \subset \mathbb{P}^2$ be a finite set and call I_A the homogeneous ideal of A, in the polynomial ring $R = \mathbb{C}[x_0, x_1, x_2]$. Then there exists an exact sequence of graded R-modules (called a *minimal resolution* for I_A), as follows:

(2)
$$0 \longrightarrow F_1 \xrightarrow{M} F_0 \longrightarrow I_A \longrightarrow 0$$

where F_1, F_0 are free graded *R*-modules, i.e. we have

$$F_0 = \bigoplus_{i=1}^{s} R(-d_i)$$
 $F_1 = \bigoplus_{j=1}^{s-1} R(-e_j)$

(R(i) is the graded module R with the degrees shifted by i. The shift is needed in order to get that the maps in the sequence are graded homomorphisms).

The map $F_0 \longrightarrow I_A$ sends the standard basis of F_0 to a set of minimal generators for I_A , and the numbers d_i 's correspond to the degrees of the chosen minimal generators.

The map $F_1 \longrightarrow F_0$ is given by a matrix of forms M, which describes the *first* syzygies of A, i.e. the relations between the minimal generators.

Definition 2.20. The matrix of forms M which determines the map $F_1 \longrightarrow F_0$ is called a *Hilbert-Burch matrix* for A.

The degree matrix of M is called the *degree Hilbert-Burch matrix* of A.

Theorem 2.21. (Hilbert-Burch Theorem, see [16]) The Hilbert-Burch matrix M depends on a choice of a set of minimal generators for the ideal I_A , but the degree Hilbert-Burch matrix depends only on A.

The minimal generators which determine the surjection $F_0 \longrightarrow I_A$ are precisely the $(s-1) \times (s-1)$ minors of the matrix M (taken with the corresponding sign).

Example 2.22. A set of points $Z \subset \mathbb{P}^2$ is *complete intersection* if there are two plane curves F, G such that $Z = F \cap G$. In this case, the homogeneous ideal I_Z is generated by the forms F, G which define the two curves. If d_1, d_2 denote the degrees of F, G respectively, then a minimal resolution of I_Z looks like

$$0 \longrightarrow R(-d_1 - d_2) \xrightarrow{M} R(-d_1) \oplus R(-d_2) \longrightarrow I_Z \longrightarrow 0$$

and the Hilbert-Burch matrix M is given by $M = \begin{pmatrix} G \\ -F \end{pmatrix}$.

2.6. Linked sets and mapping cone. Let $A, B \subset \mathbb{P}^2$ be two sets of points such that $Z = A \cup B$ is complete intersection of two curves F and G, of degrees respectively d_1 and d_2 . In this case, we say that A and B are *linked* by (or also that B is the *residue* of A in) a complete intersection of type (d_1, d_2) .

When the two sets are linked, a Hilbert-Burch matrix of B can be found from a Hilbert-Burch matrix of A via the *mapping cone* procedure (for more details, we refer to [22], [32], and Proposition 5.2.10 of [30]).

The homogeneous ideal I_B is given by

$$I_B = I_Z : I_A = \{ f \in R : fI_A \subset I_Z \}.$$

The inclusion $I_Z \subset I_A$ determines on the resolutions of I_A and I_Z a commutative diagram:

where the rightmost vertical map is the inclusion.

Then I_B is the image of the dual map $R(d_1) \oplus R(d_2) \oplus F_1^{\vee} \xrightarrow{(\phi', M')^{\vee}} R(d_1 + d_2)$, twisted by $-d_1 - d_2$. A resolution of I_B is given by:

$$0 \to F_0^{\vee}(-d_1 - d_2) \xrightarrow{(\phi, M)^{\vee}} R(-d_2) \oplus R(-d_1) \oplus F_1^{\vee}(-d_1 - d_2) \xrightarrow{(\phi', M')^{\vee}} I_B \to 0.$$

Notice that the resolution of I_B obtained by the mapping cone procedure needs not to be minimal, in the sense that some summands of the map $F_0^{\vee}(-d_1 - d_2) \rightarrow R(-d_2) \oplus R(-d_1) \oplus F_1^{\vee}(-d_1 - d_2)$ could be factored out.

Example 2.23. Let A be a general set of three points in \mathbb{P}^2 . The ideal I_A is generated by three quadrics, and a minimal resolution is given by

$$0 \longrightarrow R(-3)^2 \xrightarrow{M} R(-2)^3 \longrightarrow I_A \longrightarrow 0,$$

where the Hilbert-Burch matrix M is a 3×2 matrix of linear forms ℓ_{ij} . If we take a general quadric F and a general cubic G containing A, we get a linkage between A and another set B of three points in the plane. Diagram (3) looks like

and the mapping cone gives the resolution

$$0 \to R(-3)^3 \xrightarrow{(\phi,M)^{\vee}} R(-3) \oplus R(-2) \oplus R(-2)^2 \longrightarrow I_B \to 0.$$

The matrix of $(\phi, M)^{\vee}$ is obtained as follows. Since the minimal generators of I_A are the minors of M, F, G have a representation as determinants of matrices as follows

$$F = \det \begin{pmatrix} \ell_{11} & \ell_{12} & c_{13} \\ \ell_{21} & \ell_{22} & c_{23} \\ \ell_{31} & \ell_{32} & c_{33} \end{pmatrix} \quad G = \det \begin{pmatrix} \ell_{11} & \ell_{12} & h_{13} \\ \ell_{21} & \ell_{22} & h_{23} \\ \ell_{31} & \ell_{32} & h_{33} \end{pmatrix}$$

for some choice of constants c_{i3} and linear forms h_{i3} . Thus the matrix of $(\phi, M)^{\vee}$ is given by

$$\begin{pmatrix} c_{13} & c_{23} & c_{33} \\ h_{13} & h_{23} & h_{33} \\ \ell_{11} & \ell_{21} & \ell_{31} \\ \ell_{12} & \ell_{22} & \ell_{32} \end{pmatrix}$$

This resolution is not minimal, because we can choose the resolution of I_A so that F is the first generator. In this case the row (c_{13}, c_{23}, c_{33}) is equal to (1, 0, 0) and can be factored out. So, we can drop the cubic generator of I_B . The resolution becomes

$$0 \to R(-3)^2 \xrightarrow{N} R(-2) \oplus R(-2)^2 \longrightarrow I_B \to 0$$

and the Hilbert-Burch matrix \boldsymbol{N} is:

$$N = \begin{pmatrix} h_{23} & h_{33} \\ \ell_{21} & \ell_{31} \\ \ell_{22} & \ell_{32} \end{pmatrix}.$$

2.7. Grassmannian Varieties. Let $r, s \in \mathbb{N}$ such that $r \leq s$ and let $\mathbb{G}(r, s)$ be the Grassmannian variety of r-dimensional linear spaces in \mathbb{C}^s . $\mathbb{G}(r, s)$ is an algebraic subvariety of a projective space by means of the Plücker embedding

(4) $i: \mathbb{G}(r,s) \hookrightarrow \mathbb{P}(\Lambda^r \mathbb{C}^s)$

defined by

$$i(\langle v_1, \ldots, v_r \rangle) = [v_1 \land \ldots \land v_r]$$

where v_1, \ldots, v_r are r linearly independent vectors of \mathbb{C}^s . The homogeneous coordinates of $\mathbb{G}(r,s)$ on $\mathbb{P}^{\binom{s}{r}-1} \cong \mathbb{P}(\Lambda^r \mathbb{C}^s)$ are called Plücker coordinates. We recall that

$$\dim \mathbb{G}(r,s) = r(s-r).$$

Let $W \in \mathbb{G}(r, s)$. We can associate to W the $r \times s$ matrix with complex entries M_W whose rows contain the coordinates of a basis $\{v_1, \ldots, v_r\}$ of W. We notice that this representation is not unique: if we multiply for an element $U \in GL(r, \mathbb{C})$, the new matrix represents the same point of $\mathbb{G}(r, s)$. The Plücker coordinates of $\mathbb{G}(r, s)$ are the minors of size r of M_W , which are simply multiplied by a number when we substitute M_W with UM_W .

The embedding $i : \mathbb{G}(r,s) \hookrightarrow \mathbb{P}(\Lambda^r \mathbb{C}^s)$ is defined by a divisor $h_{r,s}$ of $\mathbb{G}(r,s)$. According to Lemma 11.1 of [19], the divisor $h_{r,s} \subset \mathbb{G}(r,s)$ associated to i is

$$h_{r,s} = \{ \Lambda \subset \mathbb{C}^s | \Lambda \cap \tilde{L} \neq \emptyset \}$$

where $\tilde{L} \cong \mathbb{C}^{s-r-1}$ is a fixed subspace of \mathbb{C}^s .

3. Forms of degree 9 in three variables

In this section, let us assume that n = 2, d = 9 and let $T \in S^9 \mathbb{C}^3$. Thus T is a form of degree 9 in three variables, which is associated to a curve of degree 9 in \mathbb{P}^2 .

Let $A = \{P_1, \ldots, P_r\} \subset \mathbb{P}^2$ be a finite set that computes T.

Our target is to determine conditions on T such that A is the unique set that computes the rank of T.

We will assume in this section that A is sufficiently general, in a very precise sense. We assume indeed that:

- (i) A is non-redundant;
- (ii) $k_4(A) = \min\{15, r\};$
- (iii) $h_A(5) = r$.

It is a standard fact indeed that when $r \leq 21$, all A in a Zariski open subset of $(\mathbb{P}^2)^r$ satisfy the previous conditions.

Remark 3.1. For a general $T \in S^9 \mathbb{C}^3$, according to the Alexander-Hirschowitz Theorem [1], the rank is 19 and identifiability does not hold.

On the other hand, if $r \leq 17$ and A satisfies the previous conditions, then, by Theorem 2.7, T is identifiable of rank r.

Therefore, we will assume in this section that r = 18 (so that $h_A(5) = 18$ and $k_4(A) = 15$).

Since A verifies properties (ii) and (iii), then the Hilbert function of A and its first difference verify, respectively,

Remark 3.2. The previous values of the Hilbert function imply that A is not contained in quartic curves (it is clear, since we are assuming $k_4(A) = 15$), moreover A is contained in 3 independent quintics $Q_1, Q_2, Q_3 \in S^5 \mathbb{C}^3$. The quintics Q_i 's are elements of minimal degree in the homogeneous ideal I_A of A. Thus, they are among the minimal generators. If we multiply the Q_i 's by 3 independent linear forms, which generate $S^1 \mathbb{C}^3$, we obtain a set of nine forms of degree 6 contained in the ideal I_A . These sextics span a subspace Λ_6 of dimension exactly nine in the homogeneous piece of degree 6 of I_A . Indeed, the condition $h_A(5) = r = 18$ implies $h_A(6) = 18$, so that $(I_A)_6$, which is the kernel of the evaluation map in degree 6, has affine dimension 28 - 18 = 10, hence projective dimension 9.

In section 5) we produce an algorithm which, starting with the coordinates of the points of A, texts whether or not A satisfies the generality conditions (i) - (iii). Conditions (i) - (iii) determine the shape of the degree Hilbert-Burch matrix of A.

Proposition 3.3. Assume A satisfies conditions (i) - (iii). Then there exist three forms $Q_1, Q_2, Q_3 \in S^5 \mathbb{C}^3$ of degree 5 and one form $S \in S^6 \mathbb{C}^3$ of degree 6 such that the ideal I_A is minimally generated by Q_1, Q_2, Q_3, S . The resolution of I_A , determined by this choice of generators, has the form

$$0 \longrightarrow R(-7)^{\oplus 3} \xrightarrow{M} R(-5)^3 \oplus R(-6) \longrightarrow I_A \longrightarrow 0.$$

Proof. Since the space $(I_A)_5$ of quintics through A is three-dimensional, a basis Q_1, Q_2, Q_3 for $(I_A)_5$ determines three minimal generators for I_A . Since $h_A(5) = 18$, by the Castelnuovo-Mumford regularity Theorem (see [12]) I_A is generated in degree 6. By Remark 3.2, the three quintics Q_1, Q_2, Q_3 determine a subspace Λ_6 of dimension 9 in $(I_A)_6$. $(I_A)_6$ has dimension 10, since $h_A(6) = 18$. It follows that by taking $S \in (I_A)_6 \setminus \Lambda_6$, we get a minimal set of 4 generators for I_A , which thus determines a surjection $\alpha : R(-5)^3 \oplus R(-6) \longrightarrow I_A$. The graded piece $(I_A)_7$ has dimension $\dim(R_7) - h_A(7) = 36 - 18 = 18$. By multiplying each Q_i by a basis for $S^2 \mathbb{C}^3$ and S by a basis for $S^1 \mathbb{C}^3$, we get 21 elements in $(I_A)_7$. Thus we have at least 3 independent relations of degree 7 among Q_1, Q_2, Q_3, S . Since 3 = 4 - 1, by the Hilbert-Burch Theorem 2.21, the independent relations of degree 7 are exactly 3 and the kernel of the map α is $R^3(-7)$.

By the Hilbert-Burch Theorem and by Proposition 3.3, there exist $c_{uv} \in S^2 \mathbb{C}^3$ and $\ell_j \in S^1 \mathbb{C}^3$, for $u, v, j \in \{1, 2, 3\}$, such that A has a Hilbert-Burch matrix M:

(6)
$$M = \begin{pmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \\ \ell_1 & \ell_2 & \ell_3 \end{pmatrix}$$

 Q_1, Q_2, Q_3, S coincide, respectively, with $(-1)^i$ times the minor obtained by leaving out the *i*-th row of $M, i \in \{1, 2, 3, 4\}$. In other words, we have:

(7)
$$Q_1 = -\begin{vmatrix} c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \\ \ell_1 & \ell_2 & \ell_3 \end{vmatrix}$$
, $Q_2 = \begin{vmatrix} c_{11} & c_{12} & c_{13} \\ c_{31} & c_{32} & c_{33} \\ \ell_1 & \ell_2 & \ell_3 \end{vmatrix}$,
 $Q_3 = -\begin{vmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ \ell_1 & \ell_2 & \ell_3 \end{vmatrix}$, $S = \begin{vmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{vmatrix}$.

Now, assume that $B = \{P'_1, \dots, P'_{\ell(B)}\} \subset \mathbb{P}^2$ is another finite set computing T such that

(i) $\ell(B) \le 18;$ (ii) *B* is non-redundant

and set $Z = A \cup B \subset \mathbb{P}^2$.

We assume, for the rest of the section, that the intersection $A \cap B$ is empty.

We will analyze the case in which the intersection is non-empty in the next section. Observe that, by Corollary 2.19, the last assumption: implies that Z satisfies the Cayley-Bacharach property CB(9).

Remark 3.4. Since Z satisfies CB(9), then, by Theorem 2.18, Remark 2.11 (iii) and (5), we get that

$$Dh_Z(6) + Dh_Z(7) + Dh_Z(8) + Dh_Z(9) + Dh_Z(10) \ge \ge Dh_A(0) + Dh_A(1) + Dh_A(2) + Dh_A(3) + Dh_A(4) = 15.$$

Moreover, being $Dh_A(5) = 3$, then by Remark 2.11 (iii) we obtain that

(8)
$$36 \ge \ell(Z) \ge 15 + Dh_Z(5) + Dh_Z(6) + Dh_Z(7) + Dh_Z(8) + Dh_Z(9) + Dh_Z(10)$$

 $\ge 18 + Dh_Z(6) + Dh_Z(7) + Dh_Z(8) + Dh_Z(9) + Dh_Z(10).$

Therefore

(9)
$$Dh_Z(6) + Dh_Z(7) + Dh_Z(8) + Dh_Z(9) + Dh_Z(10) \le 18$$

We immediately get the following chain of inequalities

(10) $15 \le Dh_Z(6) + Dh_Z(7) + Dh_Z(8) + Dh_Z(9) + Dh_Z(10) \le 18.$

Proposition 3.5. Assume that $A \cap B = \emptyset$. Then one the following cases occurs for the first difference of the Hilbert function of Z: Case 1:

	j	0	1	2	3	4	5	6	7	8	9	10	11	
	$Dh_Z(j)$	1	2	3	4	5	6	5	4	3	2	1	0	•••
and $\ell(B) = 18, \ \ell(Z) = 36;$														
Case 2:														
	j	0	1	2	3	4	5	6	7	8	9	10	11	
	$Dh_Z(j)$	1	2	3	4	5	5	5	4	3	2	1	0	
and $\ell(B) = 17, \ \ell(Z) = 35;$														
Case 3:														
	j	0	1	2	3	4	5	6	7	8	9	10	11	
	$Dh_Z(j)$	1	2	3	4	5	5	5	5	3	2	1	0	
and $\ell(B) = 18, \ \ell(Z) = 36.$														

Proof. By Proposition 2.14, we know that $Dh_Z(10) \ge 1$. Assume that $Dh_Z(10) \ge 2$. Then, by Proposition 2.16, necessarily $Dh_Z(9) \ge Dh_Z(10) + 1 \ge 3$, $Dh_Z(8) \ge Dh_Z(9) + 1 \ge 4$, $Dh_Z(7) \ge Dh_Z(8) + 1 \ge 5$, which implies, by (9), that $Dh_Z(6) \le 4$, a contradiction. Therefore

$$Dh_Z(10) = 1$$

Moreover, for any $j \ge 11$ we get $Dh_Z(j) = 0$, and also $Dh_Z(9) \ge 2$, otherwise we violate Proposition 2.16.

Assume that $Dh_Z(9) \geq 3$. Then, by arguing as before, $Dh_Z(8) \geq Dh_Z(9) + 1 \geq 4$, $Dh_Z(7) \geq Dh_Z(8) + 1 \geq 5$ and $Dh_Z(6) \leq 5$. If $Dh_Z(6) = 5$, then $Dh_Z(7) = 5$ and, by the first line of (8), $Dh_Z(5) + Dh_Z(8) + Dh_Z(9) \leq 10$, so that $Dh_Z(5) \leq 3$, a contradiction. If $Dh_Z(6) \leq 4$, then, by Proposition 2.12, $Dh_Z(7) \leq 4$, a contradiction. Therefore

$$Dh_Z(9) = 2$$

and $Dh_Z(8) \ge 3$, by Proposition 2.16.

Assume that $Dh_Z(8) \ge 4$. Then, with the same arguments as above, $Dh_Z(7) \ge 5$. If $Dh_Z(7) = 5$, then $Dh_Z(6) \ge 5$ and, being $\sum_{i=0}^{10} Dh_Z(i) \le 36$, then $Dh_Z(5) \le 4$, a contradiction; if $Dh_Z(7) > 5$, then, by(9), $Dh_Z(6) < 6$, which provides a contradiction to Proposition 2.12. As a consequence, we get that

$$Dh_Z(8) = 3$$

so that, by Proposition 2.16 again, $Dh_Z(7) \ge 4$. **Case 3.** Assume that $Dh_Z(7) \ge 5$. If $Dh_Z(7) > 5$, then, by Proposition 2.12, $Dh_Z(6) > 5$ and so, by (9), $Dh_Z(6) + Dh_Z(7) + Dh_Z(8) + Dh_Z(9) + Dh_Z(10) = 18$. In particular, it turns to be $Dh_Z(5) = 3 < Dh_Z(6)$, a contradiction. Hence we have

$$Dh_Z(7) = 5.$$

By Proposition 2.12 we know that $Dh_Z(6) \ge 5$. In particular, if $Dh_Z(6) > 5$, then direct computations show that $Dh_Z(5) < 5$, which violates Proposition 2.12. Thus, necessarily,

$$Dh_Z(6) = 5$$

which implies that

$$Dh_Z(5) = 5.$$

We get Case 3 of the statement. In particular, we get $\ell(Z) = 36$ and $\ell(B) = 18$. Cases 1, 2. Now assume that

$$Dh_Z(7) = 4$$

Therefore, by Proposition 2.16, we get that $Dh_Z(6) \ge 5$. If $Dh_Z(6) > 5$, then, by (8), $Dh_Z(5) \le 5$, a contradiction. Thus

$$Dh_Z(6) = 5$$

As a consequence, by (8) and Proposition 2.12 we get that either

$$Dh_{Z}(5) = 6$$

and we are in Case 1, with $\ell(Z) = 36$, $\ell(B) = 18$, or

$$Dh_Z(5) = 5,$$

and we are in Case 2, with $\ell(Z) = 35$, $\ell(B) = 17$.

We analyze in details the three cases that appear in Proposition 3.5.

3.1. Case 1.

Let us assume to be in the first case of Proposition 3.5. In fact, we will prove below that this is the general case of the three, in the sense that Case 2 and Case 3 are limits of Case 1.

Recall that the ideal I_A is generated by three quintics Q_1, Q_2, Q_3 and one sextic S, and its minimal resolution, with Hilbert-Burch matrix M, is described in Proposition 3.3.

The Hilbert function tells us that Z is contained in no quintics and it lies in two irreducible sextics. Moreover, we know that Z satisfies CB(9), and its Hilbert function is the same as the one of a complete intersection of type (6, 6). By the Main Theorem of [17], it follows that Z is a complete intersection of type (6, 6). So I_Z is generated by F, F', where $F, F' \in (I_A)_6$. We get a commutative diagram

where M is described in (6), and the matrices M_1, M_2 are defined by writing

$$F = L_1Q_1 + L_2Q_2 + L_3Q_3 + aS \quad F' = L'_1Q_1 + L'_2Q_2 + L'_3Q_3 + a'S$$

so that

$$M_2 = \begin{pmatrix} L_1 & L_1' \\ L_2 & L_2' \\ L_3 & L_3' \\ a & a' \end{pmatrix}, \quad M_1 = \begin{pmatrix} q_1 \\ q_2 \\ q_3 \end{pmatrix},$$

with $q_i \in S^5 \mathbb{C}^3$ for $i \in \{1, 2, 3\}$.

By the mapping cone, a resolution of I_B is given by

(11)
$$0 \to R(-7)^3 \oplus R(-6) \xrightarrow{N} R(-6)^2 \oplus R(-5)^3 \to I_B \to 0$$

where

$$N = \begin{pmatrix} L_1 & L_2 & L_3 & a \\ L'_1 & L'_2 & L'_3 & a' \\ c_{11} & c_{21} & c_{31} & \ell_1 \\ c_{12} & c_{22} & c_{32} & \ell_2 \\ c_{13} & c_{23} & c_{33} & \ell_3 \end{pmatrix}.$$

and notice that the last three lines represent the transpose of M.

Remark 3.6. Let S' be any form of degree 6 in I_A . Then S' corresponds to an element of the homogeneous piece $(I_A)_6$, which is a 10-dimensional linear space. By the description of the generators of I_A , there are linear forms u_1, u_2, u_3 and a constant $c \in \mathbb{C}$ such that S' is the determinant of the matrix:

$$\begin{pmatrix} u_1 & u_2 & u_3 & c \\ c_{11} & c_{21} & c_{31} & \ell_1 \\ c_{12} & c_{22} & c_{32} & \ell_2 \\ c_{13} & c_{23} & c_{33} & \ell_3 \end{pmatrix}$$

If we pose

 $u_1 = a_0 x_0 + a_1 x_1 + a_2 x_2, u_2 = a_3 x_0 + a_4 x_1 + a_5 x_2, u_3 = a_6 x_0 + a_7 x_1 + a_8 x_2$ then we can associate to S' the 10 coordinates (a_0, \ldots, a_8, c) . This provides a set

of coordinates of S' in the linear space $(I_A)_6$.

In particular, we have

(12)
$$F = - \begin{vmatrix} L_1 & L_2 & L_3 & a \\ c_{11} & c_{21} & c_{31} & \ell_1 \\ c_{12} & c_{22} & c_{32} & \ell_2 \\ c_{13} & c_{23} & c_{33} & \ell_3 \end{vmatrix}$$
(13)
$$F' = \begin{vmatrix} L'_1 & L'_2 & L'_3 & a' \\ c_{11} & c_{21} & c_{31} & \ell_1 \\ c_{12} & c_{22} & c_{32} & \ell_2 \\ c_{13} & c_{23} & c_{33} & \ell_3 \end{vmatrix}$$

so that, if we pose

(14) $L_1 = a_0x_0 + a_1x_1 + a_2x_2, L_2 = a_3x_0 + a_4x_1 + a_5x_2, L_3 = a_6x_0 + a_7x_1 + a_8x_2$ $L'_1 = a_{10}x_0 + a_{11}x_1 + a_{12}x_2, L'_2 = a_{13}x_0 + a_{14}x_1 + a_{15}x_2, L'_3 = a_{16}x_0 + a_{17}x_1 + a_{18}x_2,$ then we can associate to the pair (F, F') the matrix

$$W = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & a_4 & a_5 & a_6 & a_7 & a_8 & a_9 \\ a_{10} & a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} & a_{17} & a_{18} & a_{19} \end{pmatrix}$$

where $a_9 = a, a_{19} = a'$. The pair (F, F') determines a two dimensional linear subspace Λ of $(I_A)_6$, hence an element of the Grassmannian of 2-dimensional linear subspaces in a (10)-dimensional space. The Plücker coordinates of Λ are precisely the 2×2 minors of W, in the sense of (4).

Since the residue scheme B depends only on the linear space Λ spanned by F, F', we can summarize the analysis in the following remark.

Remark 3.7. Let $T \in S^9 \mathbb{C}^3$ be a form with a decomposition $A \subset \mathbb{P}^2$ which satisfies conditions (i)-(iii) above. Assume that there exists a second decomposition B for T, such that $Z = A \cup B$ has the difference Hilbert function of Case 1. Then there exists a 2-dimensional subspace $\Lambda \subset (I_A)_6$ such that B is linked to A by two sextics which give a basis of Λ .

We can prove that the converse holds.

Remark 3.8. Choose a general 2-dimensional subspace $\Lambda \subset (I_A)_6$ and a basis F, F' of Λ . Since I_A is generated in degree 6 and Λ is general, the residue B of A in the complete intersection Z of F, F' is a set of 18 distinct points which does not intersect A (i.e. Z is smooth, as a consequence of the classical Bertini's Theorem). Moreover, since $h_Z(9) = 35 < \ell(Z) = 36$, then by the Grassmann formula the linear spaces $\langle v_9(A) \rangle$ and $\langle v_9(B) \rangle$ meet in exactly one point T. Thus, T is a form of degree 9 which has two different (disjoint) decompositions of length 18.

Remark 3.8 tells us that we have a (rational) map

(15)
$$f: \mathbb{G}(2, 10) \dashrightarrow \langle v_9(A) \rangle$$

whose image contains tensors T with a second decomposition B of length 18, such that $Z = A \cup B$ has a Hilbert function as in Case 1 above. Since the Grassmannian of 2-dimensional subspaces of $(I_A)_6$ has dimension 2(10-2) = 16, while the span of $v_9(A)$ is 17-dimensional, it turns out that a general element $T \in \langle v_9(A) \rangle$ cannot belong to the (closure of) the image of f.

Next target is to analyze the map f and determine when a given T belongs to the closure of im(f).

From the Hilbert functions of A and B, we know that both $(I_A)_9$ and $(I_B)_9$ determine 37-dimensional subspaces of the linear space R_9 , which has dimension 55. By standard facts on the intersection of ideals, $I_A \cap I_B$ is the ideal of the union $Z = A \cup B$. Thus, from the Hilbert function of Z one knows that

$$\dim((I_A)_9 \cap (I_B)_9) = \dim((I_Z)_9) = 55 - 35 = 20.$$

From the Grassmann formula one computes:

$$\dim((I_A)_9 + (I_B)_9) = 37 + 37 - 20 = 54.$$

It follows that $(I_A)_9 + (I_B)_9$ is a hyperplane in R_9 , thus it determines a point in the dual projective space $\mathbb{P}^{\vee} = \mathbb{P}(R_9)^{\vee}$, of dimension 54.

Before going on, we need to make a remark on the relation between points and forms in projective spaces, clarifying the roles of elements of \mathbb{P}^N and its dual. The following remark collects standard facts for the relations between projective geometry and linear algebra.

Remark 3.9. Let $\mathbb{P}(V)$ be a projective space and let $\mathbb{P}(V)^{\vee}$ be its dual, defined as the set of hyperplanes in $\mathbb{P}(V)$. If we fix coordinates in $\mathbb{P}(V)$, then there are dual coordinates in $\mathbb{P}(V)^{\vee}$, defined as follows: for any hyperplane H in $\mathbb{P}(V)$, the coefficients of an equation H in the fixed coordinates of $\mathbb{P}(V)$ are dual coordinates for the point [H] representing H in $\mathbb{P}(V)^{\vee}$. Thus, for a point $T \in \mathbb{P}(V)$, the coordinates of T are coefficients for an equation of the hyperplane dual to T, in the dual coordinates of $\mathbb{P}(V)^{\vee}$.

If Λ is a linear subspace of $\mathbb{P}(V)$, the dual subspace $\Lambda^{\vee} \subset \mathbb{P}(V)^{\vee}$ is the set of

hyperplanes containing Λ . Thus for $\Lambda_1, \Lambda_2 \subset \mathbb{P}(V)$ subspaces, the intersection $\Lambda_1^{\vee} \cap \Lambda_2^{\vee}$ corresponds to the dual of the linear subspace $\Lambda_1 + \Lambda_2$.

Coming to our situation, up to now we indicated with \mathbb{P}^2 the projective space over the space of linear forms in the ring $R = \mathbb{C}[x_0, x_1, x_2]$, i.e. $\mathbb{P}^2 = \mathbb{P}(R_1)$. There is a natural interpretation of $\mathbb{P}(R_9)^{\vee} = (\mathbb{P}^{54})^{\vee}$ as $\mathbb{P}(Sym_9(R_1^{\vee}))$. If $T \in R_9$ is a form, the coefficients of T are coordinates of T in the natural frame defined by monomials. Thus T represents an equation for the hyperplane of $\mathbb{P}(R_9)^{\vee}$ in the dual set of coordinates.

In this interpretation, forms in the span Λ of powers L_1^9, \ldots, L_r^9 of linear forms are equations for hyperplanes in $\mathbb{P}(R_9)^{\vee}$ associated to points of Λ .

It follows that, if $A = \{L_1, \ldots, L_r\} \subset \mathbb{P}^2$, then the space $\mathbb{P}((I_A)_9)$ has a natural interpretation as the set of hyperplanes in \mathbb{P}^{54} which are dual to points of Λ .

A consequence of the previous remark is the interpretation of $(I_A)_9 + (I_B)_9$ expressed in the following:

Proposition 3.10. There is a natural interpretation of $\mathbb{P}(R_9)^{\vee}$ so that, in the notation above, the hyperplane $(I_A)_9 + (I_B)_9$ is the dual of the point T.

Proof. T belongs to the span of the L_i^{9} 's exactly when the hyperplane dual of T contains the intersection of the hyperplanes dual to the L_i^{9} . In the dual space, the intersection of the hyperplanes dual to the L_i^{9} is defined by forms in $(I_A)_{9}$. The claim follows.

Next result describes birationally the geometric locus (hypersurface) of forms of degree 9 in the span of $\nu_9(A)$ for which a second decomposition exists.

Theorem 3.11. Let $f : \mathbb{G}(2, 10) \dashrightarrow \langle v_9(A) \rangle$ be the rational map defined in (15) and let $D \subset \mathbb{G}(2, 10)$ be the divisor of the Grassmannian which defines f. Then $D = h_{2,10}$. Therefore f is a linear projection of the Plücker embedding of $\mathbb{G}(2, 10)$.

Proof. Our aim is to show that, if H is a hyperplane of \mathbb{P}^{54} not containing $\langle v_9(A) \rangle$, then $f^{-1}(H)$ is linearly equivalent to the divisor $h_{2,10}$. We will do that for a special hyperplane H.

In the dual space $(\mathbb{P}^{54})^{\vee}$, H corresponds to a point [H]. Therefore, we want to describe for which elements $W \in \mathbb{G}(2, 10)$ the hyperplane $H_{f(W)}$ associated to the point f(W) contains the point [H].

As in Remark 3.9, an equation for H, whose coefficients are a set of homogeneous coordinates for the dual point [H], determines a form of degree 9 in R, which we call G_H . By construction,

$$H_{f(W)} = \mathbb{P}((I_A)_9 + (I_{B(W)})_9)$$

where $B(W) = \{P'_1(W), \ldots, P'_{18}(W)\} \subset \mathbb{P}^2$ is such that $Z(W) = A \cup B(W)$ is the complete intersection of type (6, 6) associated to W. Thus

(16) $[H] \in H_{f(W)}$ if and only if there exist $G_A \in (I_A)_9$, $G_{B(W)} \in (I_{B(W)})_9$ such that $G_H = G_A + G_{B(W)}$.

The condition now becomes purely algebraic.

Take a set Y of homogeneous coordinates of the points of A, and call $\rho : R \to \mathbb{C}^{18}$ the evaluation map of forms on the set Y. The natural inclusion of $I_{Z(W)}$ in $I_{B(W)}$ gives rise to an exact sequence

(17)
$$0 \to (I_{Z(W)})_9 \longrightarrow (I_{B(W)})_9 \xrightarrow{\rho} \mathbb{C}^{18}$$

where the righmost map is not surjective.

Claim 3.12. The following are equivalent:

(i) there exist $G_A \in (I_A)_9$, $G_{B(W)} \in (I_{B(W)})_9$ such that $G_H = G_A + G_{B(W)}$, (ii) $\rho(G_H)$ belongs to $\rho((I_{B(W)})_9)$.

Proof. If (i) holds, then

$$\rho(G_H) = \rho(G_A + G_{B(W)}) = \rho(G_A) + \rho(G_{B(W)}) = \rho(G_{B(W)})$$

which implies (ii).

On the other side, assume (*ii*) so that there exists $G_{B(W)} \in (I_{B(W)})_9$ such that $\rho(G_{B(W)}) = \rho(G_H)$. Therefore $\rho(G_H - G_{B(W)}) = 0$, so that $G_H - G_{B(W)}$ belongs to $(I_A)_9$ and (*i*) holds.

By combining (16) with Claim 3.12, our aim turns out to be the following: given a form G_H of R_9 , we want to describe the elements $W \in \mathbb{G}(2, 10)$ such that $\rho(G_H)$ belongs to $\rho((I_{B(W)})_9)$.

Claim 3.13. For any $W \in \mathbb{G}(2, 10)$ set $Z'(W) = Z(W) \setminus \{P_{17}, P_{18}\}$. Then for any form G of degree 9 in $I_{Z'(W)}$, the residue $\rho(G)$ is fixed, up to scalar multiplication.

Proof. We apply Proposition 5.2.10 of [30] (Mapping cone) to the commutative diagram (see Section 2.6)

$$0 \longrightarrow R(-3) \xrightarrow{\begin{pmatrix} -c \\ \ell \end{pmatrix}} R(-1) \oplus R(-2) \longrightarrow I_{\{P_{17}, P_{18}\}} \longrightarrow 0$$

$$\uparrow \qquad \uparrow \qquad 0$$

$$0 \longrightarrow R(-12) \xrightarrow{\begin{pmatrix} -F' \\ F \end{pmatrix}} R(-6)^{\oplus 2} \longrightarrow I_{Z(W)} \longrightarrow 0$$

where c, ℓ are generators for the ideal of $\{P_{17}, P_{18}\}$. We get that $I_{Z'(W)}$ admits a resolution of the form

(18)
$$0 \to R(-11) \oplus R(-10) \xrightarrow{\begin{pmatrix} \phi_1 & \psi_1 \\ \phi_2 & \psi_2 \\ -c & \ell \end{pmatrix}} R(-6)^{\oplus 2} \oplus R(-9) \to I_{Z'(W)} \to 0$$

where $\phi_1, \phi_2 \in S^5 \mathbb{C}^3, \psi_1, \psi_2 \in S^4 \mathbb{C}^3$ are defined by the central vertical map of the diagram. The ideal $I_{Z'(W)}$ is thus generated by F, F', G_0 with

(19)
$$F = \begin{vmatrix} \phi_1 & \psi_1 \\ -c & \ell \end{vmatrix} = \ell \phi_1 + c \psi_1$$

(20)
$$F' = \begin{vmatrix} \phi_2 & \psi_2 \\ -c & \ell \end{vmatrix} = \ell \phi_2 + c \psi_2$$

(21)
$$G_0 = \begin{vmatrix} \phi_1 & \psi_1 \\ \phi_2 & \psi_2 \end{vmatrix} = \phi_1 \psi_2 - \phi_2 \psi_1.$$

Then for any $G \in (I_{Z'(W)})_9$ we get $G = U_1F_1 + U_2F_2 + qG_0, q \in \mathbb{C}$, where U_1, U_2 are suitable cubics. Since F_1, F_2 vanish at Z(W), then the residue $\rho(G)$ is a multiple of $\rho(G_0)$.

In particular, for $W \in \mathbb{G}(2, 10)$, we can find scalars α_W, β_W such that the residue $\rho(G)$ is a scalar multiple of $(0, \ldots, 0, \alpha_W, \beta_W)$ for all $G \in (I_{Z'(W)})_9$.

Fix two non-zero scalars $\alpha, \beta \in \mathbb{C} - \{0\}$. Let $v_{\alpha,\beta} = (v_1, \ldots, v_{18}) \in \mathbb{C}^{18}$ be defined by

$$v_j = \begin{cases} 0 & j \in \{1, \dots, 16\} \\ \alpha & j = 17 \\ \beta & j = 18 \end{cases}$$

Since the evaluation map ρ surjects, we can find H such that the form G_H satisfies $\rho(G_H) = v_{\alpha,\beta}$. We compute $f^{-1}(H)$ for this hyperplane H. Notice that H cannot contain $\langle v_9(A) \rangle$, for G_H does not vanish at P_{17}, P_{18} .

By Claim 3.12, W belongs to $f^{-1}(H)$ if and only if there exists $G \in (I_{B(W)})_9$ with $\rho(G) = \rho(G_H)$. Such a form G vanishes at the points P_1, \ldots, P_{16} , thus it belongs to $(I_{Z'(W)})_9$. We obtain by Claim 3.13 that $W \in f^{-1}(H)$ if and only if

(22)
$$\det \begin{pmatrix} \alpha & \beta \\ \alpha_W & \beta_W \end{pmatrix} = 0.$$

Now we compare the two expressions that we have for F, from (12) and (19). We get that

$$\ell\phi_1 + c\psi_1 = L_1Q_1 + L_2Q_2 + L_3Q_3 + aS,$$

but Q_1, Q_2, Q_3, S vanish at P_{17}, P_{18} , thus they belong to the ideal spanned by ℓ, c . In particular, we can write $Q_i = M_i c + N_i \ell$ and $S = \overline{M}c + \overline{N}\ell$. Since ℓ, c give a complete intersection, so they have only trivial syzygies, we conclude that there exists a form U of degree 3 such that

$$\phi_1 = Uc + L_1N_1 + L_2N_2 + L_3N_3 + aN$$

$$\psi_1 = -U\ell + L_1M_1 + L_2M_2 + L_3M_3 + a\bar{M}.$$

Similarly, if we compare the two expressions that we have for F', from (13) and (20), we get that

$$\phi_2 = U'c + L'_1N_1 + L'_2N_2 + L'_3N_3 + a'\bar{N}$$

$$\psi_2 = -U'\ell + L'_1M_1 + L'_2M_2 + L'_3M_3 + a'\bar{M}.$$

for some form U' of degree 3.

Now put the previous expressions in the formula (21) for G_0 . When we compute the residue of G_0 at Z(W), the contribution of U, U' disappears, since c, ℓ vanish at the points P_{17}, P_{18} . Moreover the $M_i, N_i, \overline{M}, \overline{N}$ are fixed and do not depend on the choice of W. It follows that (α_W, β_W) is an expression in terms of the 2×2 determinants of the coefficients of the L_i 's and a and the coefficients of the L'_i 's and a'. That is: (α_W, β_W) is a linear expression in the Plücker coordinates of W. This concludes the proof of the theorem. \Box

Of course, we do not know which projection of the Grassmannian determines the rational map f. It is likely that the projection is highly special.

Claim 3.14. The map $f : \mathbb{G}(2, 10) \dashrightarrow \langle v_9(A) \rangle$ defined in (15) is birational onto the image.

Proof. We aim to show that, for some $P \in im(f) \subset \langle v_9(A) \rangle$, the set $f^{-1}(P)$ is finite and has degree 1.

We proceed via a computational approach in Macaulay2 [25] (over a finite field, but

then the proof holds also over \mathbb{C}). For a detailed description of our procedure, we refer to the ancillary file nonics3.txt.

We start by selecting a finite set $A = \{P_1, \ldots, P_{18}\} \subset \mathbb{P}^2$, whose elements have random coefficients. Then, we construct the Hilbert-Burch matrix of A and we fix 2 forms F, F' of degree 6 in I_A , so that one is not the multiple of the other. This is equivalent to a choice of 6 linear forms $L_1, L_2, L_3, L'_1, L'_2, L'_3$ and 2 scalars a, a' (since the choice is general, we may assume that $\partial_{x_0}L_1 = 1, \partial_{x_0}L'_1 = 0, a = 0, a' = 1$). We get a residual set $B_{F,F'}$, whose ideal admits a resolution as in (11). By means of (15), we compute f(F, F') and we pose P = f(F, F'). Let (p_0, \ldots, p_{54}) be a representative vector for P.

In order to obtain $f^{-1}(P)$, in the first and second row of the Hilbert-Burch matrix N of $B_{F,F'}$ we change L_j (resp. L'_j) with $L_j = a_{3j-3}x_0 + a_{3j-2}x_1 + a_{3j-1}x_2$ (resp. with $L'_j = a_{3j+7}x_0 + a_{3j+8}x_1 + a_{3j+9}x_2$), for $j \in \{1, 2, 3\}$ (with $a_0 = 1$ and $a_{10} = 0$) and we consider the 55 × 54 matrix MFix" whose columns provide a set of generators for $(I_A)_9 + (I_{B_{F,F'}})_9$. Notice that MFix" is divided in 2 blocks: the first 37 columns have integer entries, while in the last 17 the entries depend linearly on the 2x2 minors of the matrix

$$W = \begin{pmatrix} 1 & a_1 & a_2 & a_3 & a_4 & a_5 & a_6 & a_7 & a_8 & 0\\ 0 & a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} & a_{17} & a_{18} & 1 \end{pmatrix},$$

i.e. on the Plücker coordinates of W in the sense of (4). According to the previous remark, write $MFix'' = (A_1|A_2)$. Therefore

$$f^{-1}(P) = \{(a_1, \dots, a_8, a_{11}, \dots, a_{18}) \in \mathbb{C}^{16} \mid (p_0, \dots, p_{54}) \cdot MFix'' = 0_{1 \times 55} \}.$$

Since $(p_0, \ldots, p_{54}) \cdot A_1 = 0_{1 \times 37}$ provide trivial conditions, then

(23)
$$f^{-1}(P) = \{(a_1, \dots, a_8, a_{11}, \dots, a_{18}) \in \mathbb{C}^{16} \mid (p_0, \dots, p_{54}) \cdot A_2 = 0_{1 \times 17} \}.$$

We notice that the 17 equations appearing in (23) provide a linear system in the Plücker coordinates of W. Our computations in Macaulay2 show that $f^{-1}(P)$ has dimension 0 and degree 1, which concludes the proof.

Claim 3.14 implies the following:

Remark 3.15. If $T \in S^9 \mathbb{C}^3$ of rank 18 is a general point in $\operatorname{im}(f)$, i.e. a general unidentifiable nonic of rank 18, then there exist *exactly two* finite sets computing the rank of T.

In other words, the space $\langle v_9(A) \rangle$ contains a variety Θ , which is the closure of a linear projection of $\mathbb{G}(2, 10)$, whose general points consist of forms in $S^9\mathbb{C}^3$ of rank 18, that admit two finite sets computing the rank.

Remark 3.16. Given $T \in S^9 \mathbb{C}^3$ of rank 18 with a non-redundant finite set $A = \{P_1, \ldots, P_{18}\} \subset \mathbb{P}^2$ computing it, such that $k_4(A) = 15$ and $h_A(5) = 18$, by following the proof of 3.14, in principle one can develop a criterion that establishes the uniqueness of such an A.

The corresponding algorithm is explained in more details in Section 5.

3.2. Case 2.

Let us assume to be in the second case of Proposition 3.5. This is the unique case in which the new decomposition has less than 18 summands, so that the rank of T is 17, not 18. In Section 5 we will provide a software which can exclude that a given form T falls in this case. Notice that, by Theorem 2.15, Z is contained in a plane quintic. Moreover, passing to cohomology in the exact sequence

$$0 \to (I_Z)_s \to R(s) \to \mathbb{C}^{35} \to 0$$

for $s \in \{5, 7, 12\}$, we get that Z is contained in a unique quintic Q, and there exists a septic G containing Z and not containing Q. Since, Z satisfies CB(9) and the Hilbert function of Z is the same as the Hilbert function of a complete intersection of type (5,7), then, by the Main Theorem of [17], Z is a complete intersection of type (5,7). In particular, $I_Z = (Q, G)$, with $Q \in S^5 \mathbb{C}^3$ and $G \in S^7 \mathbb{C}^3$ and a minimal resolution of I_Z is given by

$$0 \to R(-12) \xrightarrow{\begin{pmatrix} -G\\Q \end{pmatrix}} R(-5) \oplus R(-7) \to I_Z \to 0$$

Again, fix three quintics Q_1, Q_2, Q_3 and one sextic S that generate the ideal of A, so that a minimal resolution of I_A , with Hilbert-Burch matrix M, is as described in Proposition 3.3.

Remark 3.17. For any choice of the quintic generators Q_1, Q_2, Q_3 , the ideal I_A coincides with the ideal generated by the Q_i 's in degree 7.

Indeed, it follows from the description of the Hilbert-Burch matrix M that, for a general choice of the quintic forms, there are no relations of degree 7 involving only Q_1, Q_2, Q_3 . In other words, there are no non-trivial quadrics q_1, q_2, q_3 such that $\sum q_i Q_i$ is the zero polynomial. It follows that the ideal J generated by Q_1, Q_2, Q_3 satisfies dim $J_7 = 18$, which is exactly the dimension of $(I_A)_7$, as computed from the Hilbert function.

Remark 3.18. In the space $\mathbb{P}^{20} = \mathbb{P}(R_5)$ which parameterizes quintic forms (up to scalar multiplication) the generators Q_1, Q_2, Q_3 of I_A determine a plane $\mathbb{P}^2 = \Pi$. Fix a general quintic Q that belongs to the plane, so that in particular Q belongs to I_A . Call J the ideal generated by Q. The quotient $(I_A)_7/J_7$ is a vector space of dimension 12.

The variety E of sets Z containing A and complete intersection of type (5,7) is thus a $\mathbb{P}((I_A)_7/J_7)$ -bundle, i.e. a \mathbb{P}^{11} -bundle, over an open subset of Π . It has dimension 13.

Sets B of 17 points of \mathbb{P}^2 linked to A by a complete intersection of type (5,7) are parameterized by E. As in Remark 3.8, we get that a general such set B determines one point of the span of $v_9(A)$ with a second decomposition of length 17. Hence there exists a rational map

(24)
$$f': E \dashrightarrow \langle v_9(A) \rangle$$

such that the closure Θ' of the image of f' (of dimension at most 13) is the closure of the locus of forms $T \in \langle v_9(A) \rangle$ for which A is non-redundant but the rank is 17.

We will provide in Section 5 an algorithm which guarantees that a given form T does not lie in the subvariety Θ' , so it has rank 18. In order to produce the algorithm, we need a description of the form T which is the intersection of $\langle v_9(A) \rangle$ and $\langle v_9(B) \rangle$, where B is linked to A by a complete intersection $Q \cap G$ of type (5, 7).

Remark 3.19. Fix a quintic $Q \in I_A$, which can be written as

$$Q = a_1 Q_1 + a_2 Q_2 + a_3 Q_3,$$

for a choice of the scalars a_i 's. Notice that if $a_3 \neq 0$ (resp. $a_2 \neq 0$. $a_1 \neq 0$), then the ideal generated by Q_1, Q_2, Q_3, Q coincides with the ideal generated by Q_1, Q_2, Q (resp. Q_1, Q, Q_3, Q, Q_2, Q_3). By Remark 3.17, a general septic $G \in I_A$ can be written as

$$G = q_1 Q_1 + q_2 Q_2 + q_3 Q_3,$$

for a choice of quadrics q_1, q_2, q_3 .

Let B = B(Q, G) be the residue of A in the complete intersection $Q \cap G$. As in case 1, from the Hilbert functions of A, B and $Z = A \cup B$, we know that $(I_A)_9$, resp. $(I_B)_9$, determines a 37-dimensional, resp. a 38-dimensional, subspace of the linear space R_9 , which has dimension 55. Moreover dim $((I_A)_9 + (I_B)_9) = 54$, so that $(I_A)_9 + (I_B)_9$ is a hyperplane in R_9 .

As in proposition 3.10, the hyperplane $(I_A)_9 + (I_B)_9$ is dual to the point T of intersection between $\langle v_9(A) \rangle$ and $\langle v_9(B) \rangle$.

As in case 1, the mapping cone procedure provides an effective way of computing $(I_B)_9$, thus also the sum $(I_A)_9 + (I_B)_9$. Namely, the ideal I_B is defined by the minors of the Hilbert-Burch matrix M, to which we add the columns that define Q, G. We get the matrix:

(25)
$$M' = \begin{pmatrix} a_1 & a_2 & a_3 & 0\\ q_1 & q_2 & q_3 & 0\\ c_{11} & c_{21} & c_{31} & \ell_1\\ c_{12} & c_{22} & c_{32} & \ell_2\\ c_{13} & c_{23} & c_{33} & \ell_3 \end{pmatrix}.$$

In particular, B depends on the choice of forms in the first two rows

$$\begin{pmatrix} a_1 & a_2 & a_3 & 0 \ q_1 & q_2 & q_3 & 0 \end{pmatrix}.$$

Theorem 3.20. The map $f': E \dashrightarrow \langle v_9(A) \rangle$ defined in (24) is birational onto the image.

Proof. As in Theorem 3.14, we proceed via a computational approach in Macaulay2 [25]. For a detailed description of our procedure, we refer to the ancillary file nonics1.txt.

Select a finite set $A = \{P_1, \ldots, P_{18}\} \subset \mathbb{P}^2$, and construct the Hilbert-Burch matrix M of A. The choice of a quintic form Q and a septic form G in I_A , not multiple of Q, is equivalent to a choice of scalars a_1, a_2, a_3 and quadrics q_1, q_2, q_3 .

Assume first that $a_1 \neq 0$, so that, after rescaling, we may assume $a_1 = 1$. In this case notice that Q_1 is generated by Q_2, Q_3, Q . Thus the residue B = B(Q, G) of A in the complete intersection $Q \cap G$ is also the residual in the complete intersection of $Q \cap G'$ where $G' = (q_2 - a_2q_1)Q_2 + (q_3 - a_3q_1)Q_3 = q'_2Q_2 + q'_3Q_3$. I.e., we may assume $q_1 = 0$. In particular B = B(Q, G) depends on the choice of a_2, a_3 and the 12 coefficients a_4, \ldots, a_{15} of the quadrics q_2, q_3 .

Now we can compute how many choices of parameters $a_2, \ldots a_{15}$ determine a given hyperplane $P = (I_A)_9 + (I_{B(Q_0,G_0)})_9$. This can be done by Macaulay2.

We choose a nonic P corresponding to a hyperplane $(I_A)_9 + (I_{B(Q_0,G_0)})_9$ where Q_0, G_0 are determined randomly, with no coefficients a_1, \ldots, a_{15} equal to 0. The

coefficients (p_0, \ldots, p_{54}) of P in $(\mathbb{P}^{54})^{\vee}$ can be easily computed via the mapping cone. Take the 55 × 54 matrix N whose columns provide a set of generators for $(I_A)_9 + (I_{B(Q,G)})_9$, for Q, G general. Then, in the subset of Π in which $a_1 \neq 0$

(26)
$$f'^{-1}(P) = \{(a_2, \dots, a_{15}) \in \mathbb{C}^{14} | (p_0, \dots, p_{54}) \cdot N = 0_{1 \times 54} \}$$

In practice, in order to simplify the computation, observe that in an open set of E, we may also assume that one among a_4, \ldots, a_{15} is equal to 1. Thus, in the open set, we get a parametrization of E with 13 parameters, corresponding to the dimension of E. By varying the coefficients among a_1, a_2, a_3 and among a_4, \ldots, a_{15} which are set equal to 1, we obtain a complete scan of $f'^{-1}(P)$.

Our computations in Macaulay2 show that, in any case, $f'^{-1}(P)$ has dimension 0 and degree 1, i.e. it is a point. This concludes the proof.

In the rest of the section, we will prove that Case 2 is a degeneration of Case 1. We keep all the previous notation.

Proposition 3.21. Let $P_0 \in \{G = 0\}$ be a general point. Then $Z \cup \{P_0\}$ is a limit for t = 0 of a family $\{\tilde{Z}_t\}$ (over a small disc), with \tilde{Z}_t complete intersection of type (6,6) containing A for all $t \neq 0$.

Proof. The proof is direct. Let ℓ_1, ℓ_2 be general generators of I_{P_0} . Since $P_0 \in \{G = 0\}$ then there are forms E_1, E_2 of degree 6 such that $S = E_1\ell_1 + E_2\ell_2$. Since G is irreducible, for $t \in \mathbb{C}$ general the forms $\ell_1Q + tE_1, \ell_2Q + tE_2$ determine a complete intersection \tilde{Z}_t of type (6,6). By taking the flat limit for t = 0, the ideal of \tilde{Z}_t degenerates to the ideal generated by ℓ_1Q, ℓ_2Q, G , which is the ideal of $Z \cup \{P_0\}$.

It remains to prove that we can assume $A \subset \tilde{Z}_t$. Let A_t be the subset of \tilde{Z}_t which degenerates to A (note that A_t is well defined since t moves in a small disc). Since the ideal of A is generated in degree 6, the same holds for the ideal of A_t . Moreover $(I_A)_6$ and $(I_{A_t})_6$ have the same dimension. Thus the space of sextic curves containing A_t degenerates to the space of sextic curves containing A. It follows that the set of complete intersections of type (6, 6) containing A_t degenerates to the set of complete intersections of type (6, 6) containing A. In particular, Z belongs to the closure of the set of complete intersections of type (6, 6) containing A.

Proposition 3.22. Let T be a form of degree 9 with decompositions A, B such that $Z = A \cup B$ has Hilbert function as in Case 2. Then there exists a family of tensors T_t , over a small disc Δ , such that $T_0 = T$, T_t belongs to span of $v_9(A)$ for all $t \in \Delta$ and for $t \neq 0$ there exists a second decomposition B_t of T_t , which degenerates to B, with $A \cup B_t$ complete intersection of type (6,6).

Proof. The claim is an immediate consequence of Proposition 3.21, because for Z_t belonging to both Case 1 and Case 2 we have $h_9^1(Z_t) = 1$, thus the spans of $v_9(A)$ and $v_9(B_t)$ intersect in one point, for all $t \in \Delta$.

It follows that nonic forms with two decompositions A, B such that $A \cup B$ is as in Case 2 are degeneration of forms with two decompositions A, B_t with $A \cup B_t$ as in Case 1.

3.3. Case 3.

Let us consider the third case of Proposition 3.5. We will prove again that the tensor T is the limit of a family tensors $T_t \in v_9(A)$ with two decompositions whose union is complete intersection of two sextic curves.

As in the previous situation, Theorem 2.15 implies that Z is contained in a plane quintic. Moreover, the evaluation map determines an exact sequence

$$0 \to (I_Z)_s \to R(s) \to \mathbb{C}^{36} \to 0$$

for $s \in \{5, 8, 9, 12\}$. We get that Z is contained in a unique quintic Q, and there exists a pencil of plane curves of degree 8 containing Z and not containing Q. Fix two curves $O_i \in S^8 \mathbb{C}^3$, i = 1, 2 such that $Z \subset Q \cap O_1 \cap O_2$.

Claim 3.23. The set Z as above is limit for t = 0 of a family $\{Z_t\}$, containing A, such that for $t \neq 0$ the set Z_t belongs to an irreducible quintic Q containing A.

Proof. Assume that the quintic Q which contains Z is reducible. In any case, the Hilbert function of Z coincides with the Hilbert polynomial in degree 10, thus, by the Castelnuovo-Mumford regularity Theorem (see [12]), the ideal of Z is generated in degree 11. Link Z first with a complete intersection 5, 11, then link the residue with a complete intersection 5, 7. By using twice the mapping cone (see Section 2.6, one realizes that the final residue W is complete intersection of 2 quartics.

There exists a family $\{W_t\}$ of sets of 16 points, with $W_0 = W$, whose generic element is complete intersection of two quartics and it is contained in an irreducible quintic curve which contains A.

Indeed, if X, X' are the equations of two quartics whose intersection is W, then Q = zX + z'X', where z, z' are two linear forms which intersect in some point $P \in Q$. Now, move Q in a family of quintics $\{Q_t\}$, containing A, whose general element is irreducible. Move P in a family of points P_t such that $P_t \in Q_t$. The linear forms z, z' generalize to two families of linear forms $\{z_t\}, \{z'_t\}$ such that z_t, z'_t define P_t . Then, there are two families of quartics $\{X_t\}, \{X'_t\}$ such that $Q_t = z_t X_t + z'_t X'_t$ and the families specialize to X, X' for t = 0. So, just take W_t to be the intersection of X_t, X'_t .

Once the existence of the family $\{W_t\}$ is established, the existence of $\{Z_t\}$ follows immediately by linking back the general element of $\{W_t\}$, first with a complete intersection of type 5, 7, and then with a complete intersection of type 5, 11.

Thus, since we want to find Z as a limit, we may assume that Q is irreducible. The residue X of Z in the complete intersection $Q \cap O_1$ is a scheme of length 4, whose Hilbert function can be computed from the Hilbert functions of Z and the complete intersection (see Section 2.6). It follows that X is contained in a line. Thus X is complete intersection of a line ℓ and a quartic q. The mapping cone implies thus that the homogeneous ideal of Z is generated by Q, O_1, O_2 , and its minimal resolution is given by

(27)
$$0 \to R(-12) \oplus R(-9) \xrightarrow{M} R(-5) \oplus R(-8)^{\oplus 2} \to I_Z \to 0$$

where

(28)
$$M = \begin{pmatrix} S & q_1 \\ q_2 & \ell_1 \\ q & \ell \end{pmatrix}$$

denotes the Hilbert-Burch matrix of I_Z .

Note that $S \in S^7 \mathbb{C}^3$, $q_i \in S^4 \mathbb{C}^3$ for $i \in \{1, 2\}$, $\ell_1 \in S^1 C^3$, and Q, O_1, O_2 coincide, respectively, with $(-1)^i$ times the minor obtained by leaving out the *i*-th row of M, $i \in \{1, 2, 3\}$.

Let W be the residue of Z in the intersection O_1, O_2 . The fact that O_1, O_2 intersect properly, for a general choice of the two forms of degree 8 in the ideal of Z, follows from the resolution above. By the mapping cone, W is a set of length 28, complete intersection of S and q_1 . Thus, the residue of W in the intersection of q_1 and O_1 is the set X.

Move the set X in O_1 , to a general set X_t of 4 points, q_1 moves to a quartic containing X_t . Taking residues, we obtain a family of sets $\{W_t\}$ of length 28 in O_1 such that $W_0 = W$. For $t \neq 0$ the set W_t is the residue of a general set of 4 points, which is complete intersection of 2 quadrics. Thus the resolution of the ideal of W_t is:

$$0 \to R(-10)^{\oplus 2} \to R(-4) \oplus R(-8)^{\oplus 2} \to I_{W_t} \to 0.$$

The ideal of W_t has two minimal generators in degree 8, one of which is O_1 (fixed).

Claim 3.24. We can move X to X_t so that the second generator O'_t of degree 8 moves to O_2 as t goes to 0.

Proof. By the mapping cone, for t general the residue of X_t in a complete intersection (4, 8) is contained in a pencil of curves of degree 8. The limit of this pencil determines a pencil in the 2-dimensional space of curves of degree 8 through W. We need to prove that we can choose the family $\{X_t\}$ so that the limit contains O_2 . But this is clear for all the limits contain O_1 and the limit changes if we vary the family.

Notice that, for t general, the residue of W_t with respect to $O_1 \cap O'_t$ is complete intersection of two sextics. It follows that Z is limit of a family of 36 points, whose general element is a complete intersection of two sextics.

Collecting all the previous claims, we get:

Proposition 3.25. In Case 3, the form T is the limit of a family tensors $T_t \in v_9(A)$ with two decompositions whose union is complete intersection of two sextic curves.

Proof. We know that Z is limit of a family $\{Z_t\}$ whose general element is complete intersection of 2 sextics. We can conclude that Z is limit of a family $\{Z_t\}$ as above, whose general element contains A, because the family of complete intersections of type (6,6) containing Z is irreducible, and it is the limit of the set of complete intersections of type (6,6) containing a general set of 18 points.

4. The case of non-empty intersection

In this section we assume that A satisfies the genericity conditions (i)-(iii), but we drop the assumption that $A \cap B$ is empty. We will see that the case can be characterized in terms of the Case 2 of the previous section.

By arguing as in the proof of Claim 4.2 of [4], we have the following:

Proposition 4.1. If $A \cap B \neq \emptyset$, then the cardinality of B is 18 and the intersection contains only one point.

Proof. Assume $A \cap B = \{P_1, \ldots, P_s\}$, with $1 \leq s < 18$. Fix coordinates T_1, \ldots, T_{18} for the points of $\nu_9(A)$ and coordinates $T'_{s+1}, \ldots, T'_{\ell(B)}$ for the points of $\nu_9(B \setminus A)$.

Then there is a choice of scalars such that

$$T = a_1 T_1 + \ldots + a_{18} T_{18} =$$

$$= b_1 T_1 + \ldots + b_s T_s + b_{s+1} T'_{s+1} + \ldots + b_{\ell(B)} T'_{\ell(B)}.$$

Define

$$T_0 = (a_1 - b_1)T_1 + \ldots + (a_s - b_s)T_s + a_{s+1}T_{s+1} + \ldots + a_{18}T_{18} = b_{s+1}T'_{s+1} + \ldots + b_{\ell(B)}T'_{\ell(B)}.$$

 T_0 is an element of $S^9 \mathbb{C}^3$ admitting two disjoint decompositions: A and $B_0 = B \setminus A$. Notice that, since

(29)
$$T = T_0 + b_1 T'_1 + \ldots + b_s T'_s,$$

necessarily B_0 is non-redundant for T_0 , otherwise B is redundant for T. Denote by $A' \subset A$ a non-redundant decomposition of T_0 . Notice that $P_i \in A'$ for all $i \in \{s+1,\ldots,18\}$, otherwise, by (29), A is a redundant decomposition for T. It turns out that T_0 has two non-redundant decompositions, A' and B_0 , with $\ell(A') \leq 18$ and $\ell(B_0) = \ell(B) - s \leq 18 - s \leq \ell(A')$. Since A satisfies properties (ii) and (iii), then, by Remark 2.5, $k_4(A') = \min\{15, \ell(A')\}$ and $h_{A'}(5) = \ell(A') \leq 18$, and so, by Theorem 2.7, B_0 cannot exist, unless $\ell(A') = 18$, i.e. A = A'. Thus also A is non-redundant for T_0 .

The tensor T_0 has two non-redundant decompositions A, B_0 , with $\ell(B_0) \leq \ell(B) - s$, and $A \cap B_0 = \emptyset$. Now assume that either $s \geq 2$ or $\ell(B) < 18$, hence $\ell(B_0) \leq 16$. Thus Z, which is also equal to $A \cup B_0$, is a set of at most 34 points which, from Corollary 2.19, must satisfy the property CB(9). It follows that:

$$\ell(Z) \ge \sum_{i=0}^{4} Dh_A(i) + Dh_Z(5) + \sum_{i=6}^{10} Dh_Z(i) \ge 2\sum_{i=0}^{4} Dh_A(i) + Dh_Z(5) = 30 + Dh_Z(5).$$

It follows $Dh_Z(5) \leq 4$. Thus, by Proposition 2.12, $Dh_Z(i) \leq 4$ for all i > 4. Since $Dh_Z(10) > 0$, the difference Hilbert function Dh_Z cannot be strictly decreasing from 5 to 10. This contradicts Proposition 2.16.

Remark 4.2. Thus, if $A \cap B$ is non-empty, there exists a unique point, say $P_1 \in A$, which also belongs to B. Put $B_0 = B \setminus \{P_1\}$. With the notation of the proof of Proposition 4.1, we obtain that T_0 has two non-redundant decompositions: A and B_0 , with $\ell(B_0) = 17$. It follows from Proposition 3.5 applied to T_0 that $Z = A \cup B_0$ has Hilbert function as in Case 2. Thus, the previous analysis of Case 2 shows that A and B_0 are linked in a complete intersection of type 5, 7, and T_0 is the unique point of intersection of the spans of $\nu_9(A)$ and $\nu_9(B_0)$.

Moreover, the tensor T belongs to the line joining $\nu_9(P_1)$ to T_0 .

Remark 4.3. Let A, B be as in Case 1 of the previous section. We know that $Z = A \cup B$ is the complete intersection of two sextics F, F', and the spans of $\nu_9(A)$, $\nu_9(B)$ meet in one point. Assume that F moves in a family, so that the limit curves F_0 is still irreducible, and it is tangent to F in one point of A, say in P_1 . Then B moves in a family of finite sets whose limit B_0 is linked to A by a complete intersection F_0, F'_0 . It is clear from the construction that A, B_0 share the point P_1 , so that $\nu_9(P_1)$ is the unique point of intersection of the spans of $\nu_9(A), \nu_9(B_0)$.

Notice that since the ideal of A is generated in degree 6, for all sextics F containing A one can find infinitely many sextics F'_0 , containing A and tangent to F at P_1 .

Now, we are ready to prove that this case can also be seen as a degeneration of the previous Case 1.

Proposition 4.4. When $A \cap B \neq \emptyset$, the form T is the limit of a family of tensors $T_t \in v_9(A)$ with two decompositions whose union is complete intersection of two sextic curves.

Proof. We know, by Proposition 4.1, that $A \cap B$ is one point, say P_1 , moreover $B_0 = B \setminus \{P_1\}$ is a set of 17 points, linked to A by a complete intersection of type 5,7. Moreover the span of $\nu_9(A)$ and $\nu_9(B_0)$ meet in a form T_0 of rank 17 and T sits in the line joining $\nu_9(P_1)$ and T_0 .

By Case 2 of the previous section and by Proposition 3.22, we know that T_0 is a limit of a family $\{T_t\}$, whose general element T_t is the intersection of the spans of $\nu_9(A)$ and $\nu_9(B_t)$, where $\{B_t\}$ is a family of finite sets of cardinality 18, which are linked to A by a complete intersection of two families of sextics $\{F_t\}, \{F'_t\}$. The limits of the two families F_0, F'_0 intersect in a common quintic Q, containing $A \cup B_0 = Z$. In particular, $F_0 = Q \cup L$, where L is a general line.

Now, take a family $\{G_t\}$ of sextics through A, such that G_t is tangent to F_t for t general (hence for all t). Such a family exists because the ideal of A is generated in degree 6. For each $a \in \mathbb{C}$ consider the family $\{H(a)_t\}$, where $H(a)_t = \{G_t + aF'_t\}$. For t, a general the curves $H(a)_t$ and F'_t link A to a finite set $B(a)_t$ of cardinality 18. By the construction of Case 1 of the previous section, the generators of the ideal $I_{B(a)_t}$ depend linearly on a. Thus the hyperplane defined by $I_A + I_{B(a)_t}$ determines a pencil of hyperplanes, when a moves. For a = 0 the hyperplane $I_A + I_{B(a)_t}$ corresponds to the intersection T(a, t) of the spans of $\nu_9(A)$ and $\nu_9(B(0)_t)$. Since, for a = 0, $H_t = G_t$ is tangent to F_t at P_1 , the intersection is P_1^9 . When a goes to infinity and t goes to 0, the limit of T(a, t) corresponds to the limit of the points defined by the families $\{F_t\}, \{F'_t\}$, hence to T_0 . For any intermediate a, we get as a limit for t = 0 a point corresponding to hyperplanes of the pencil, thus points of the line joining P^9 and T_0 .

At the end of the analysis of Case 1,2,3 and the case of non empty intersection, we see that all the forms for which there exists a second decomposition are limits of forms described in Case 1 of the previous section. Since the image of a Grassmannian in a projection is Zariski closed, we can summarize the result of the section in the following

Theorem 4.5. Let F be a ternary form of degree 9, with a Waring expression

$$F = \lambda_1 L_1^9 + \dots + \lambda_{18} L_{18}^9$$

and call A the set of points induced by $\{L_1, \ldots, L_{18}\}$, so that A is a decomposition of F. Assume that A satisfies the following genericity properties of the beginning of Section 3:

- (i) A is non-redundant;
- (ii) $k_4(A) = 15;$
- (iii) $h_A(5) = 18$.

Then there exists a second decomposition B of length ≤ 18 for F only if F belongs to a fixed hypersurface Θ in the span of $\nu_9(A)$, which is the closed image of a birational projection of the Grassmannian of lines in \mathbb{P}^9 in its Plücker embedding.

Therefore, if F does not belong to the hypersurface Θ , then the rank of F is 18 and A is the unique decomposition of F.

Remark 4.6. (*The Panforte Challange*) The bad locus Θ is thus a hypersurface in a projective space \mathbb{P}^{17} .

As in [4], Remark 4.10, geometrically Θ is composed of points in which two folds of the secant variety Sec_{18} to the Veronese variety $\nu_9(\mathbb{P}^2)$ cross each other. Thus, the points of Θ are singular points of the secant variety, which is a hypersurface of \mathbb{P}^{54} . It turns out that the secant variety has a singular locus of codimension 1.

We do not know an equation, or even the degree of Θ in \mathbb{P}^{17} , not even when the points of A are general. A computer based calculation did not provide an answer, in a reasonable time.

As a challenge, the second author offers a Panforte (traditional cake of Siena) to the first who will determine an equation for Θ , for a general choice of the decomposition A.

5. The Algorithm

This section is devoted to the algorithm we developed based on the criterion of minimality explained in Remark 3.18, Remark 3.19, and Theorem 3.20. Our criterion is effective in the sense of [14].

5.1. The algorithm. Fix a finite set $A = \{P_1, \ldots, P_{18}\} \subset \mathbb{P}^2$ and a ternary nonic T in the linear span of $\nu_9(A)$, i.e.

$$T = \sum_{i=1}^{18} \lambda_i L_i^9$$

where, according to Remark 3.9, L_i is the linear form associated with P_i in $\mathbb{P}^2 = \mathbb{P}(R_1)$ and $\lambda_i \in \mathbb{C}$, for $i \in \{1, \ldots, 18\}$. For any *i*, denote by \mathbf{v}_i a representative vector for P_i and by (t_0, \ldots, t_{54}) the coefficients of *T* in the standard monomial basis of degree 9 in 3 variables.

In order to establish that the given T has exactly rank 18, proceed as follows. First, check that the tests

- 1) non-redundancy test: dim $\langle \nu_9(\mathbf{v}_1), \ldots, \nu_9(\mathbf{v}_{18}) \rangle = 18$
- 2) fourth Kruskal's rank test: $k_4(A) = 15$
- 3) fifth Hilbert function test: $h_5(A) = 18$

provide positive answers.

If so, construct the ideal I_A and its Hilbert-Burch matrix M as in (6). Add to the transpose of M the rows

$$\begin{pmatrix} a_1 & a_2 & a_3 & 0 \\ q_1 & q_2 & q_3 & 0 \end{pmatrix}$$

so that we get the Hilbert-Burch matrix M' of a hypothetical non-redundant decomposition B of T with $\ell(B) = 17$, as in (25).

By arguing as in the proof of Theorem 3.20 and by using the same notation, assume that $a_1 = 1$ and $q_1 = 0$ (the cases $a_2 = 1$, $q_2 = 0$ and $a_3 = 1$, $q_3 = 0$ are similar),

take the matrix N_1 whose columns yield a set of generators for $(I_A)_9 + (I_{B(Q,G)})_9$, for Q, G general and compute

$$d_1 = \max_{4 \le i \le 15} \dim\{(a_2, \dots, a_{15}) \in \mathbb{C}^{14} \mid (t_0, \dots, t_{54}) \cdot N_1 = \underline{0} \cap a_i = 0\}.$$

Equivalently, in the cases $a_2 = 1$, $q_2 = 0$ and $a_3 = 1$, $q_3 = 0$ compute, respectively,

$$d_2 = \max_{4 \le i \le 15} \dim\{(a_1, a_3 \dots, a_{15}) \in \mathbb{C}^{14} \mid (t_0, \dots, t_{54}) \cdot N_2 = \underline{0} \cap a_i = 0\}$$

and

$$d_3 = \max_{4 \le i \le 15} \dim\{(a_1, a_2, \dots, a_{15}) \in \mathbb{C}^{14} \mid (t_0, \dots, t_{54}) \cdot N_3 = \underline{0} \cap a_i = 0\}.$$

Thus, if the also the next test is successful:

4) $d_1 = d_2 = d_3 = -1$, i.e. $\dim f'^{-1}(T) = -1$

1

then $f'^{-1}(T)$ is empty and so T has rank 18.

The algorithm has been implemented in Macaulay2, over the finite field \mathbb{Z}_{31991} . The detailed procedure is contained in the ancillary file nonics2.txt.

Some examples of ternary nonics of rank 18 and 17, with a non-redundant decomposition of length 18, are presented in the following subsection.

5.2. Examples. In Macaulay2, we generated a random collection of 18 points:

$$\mathbf{A} = \begin{bmatrix} \mathbf{v}_i \end{bmatrix}_{i=1}^{18} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ -1 & 2 & 1 \\ 1 & 2 & 3 \\ 1 & -2 & 0 \\ 2 & 1 & 4 \\ 4 & 2 & -3 \\ 1 & 5 & 1 \\ 5 & 2 & 3 \\ 1 & 5 & 1 \\ 5 & 2 & 3 \\ 6 & 2 & 3 \\ 1 & 7 & 7 \\ 1 & 7 & 3 \\ 6 & 5 & 4 \\ -7 & 2 & 3 \\ 3 & 7 & 4 \\ 2 & -5 & 1 \\ 6 & 3 & -4 \\ -7 & 6 & 6 \end{bmatrix}$$

By abuse of notation, define $A = \{P_1, \ldots, P_{18}\} \subset \mathbb{P}^2$, where P_i is the projective class of \mathbf{v}_i and denote by L_i the linear form whose coefficients are given by \mathbf{v}_i .

Test 1) and test 3) show, respectively, that $\dim \langle \nu_9(A) \rangle = \operatorname{rank}([\nu_9(\mathbf{v}_i)]_{i=1}^{18}) = 18$ and $h_A(5) = \operatorname{rank}([\nu_5(\mathbf{v}_i)]_{i=1}^{18}) = 18$, as required. Moreover, all the 816 subsets of 15 columns of $[\nu_4(\mathbf{v}_i)]_{i=1}^{18}$ are of rank 15, so that $k_4(A) = 15$.

A case of rank 18. Let

$$T_2 = \sum_{i=1}^{18} L_i^9 =$$

$$\begin{split} &= [4283x_0^9 - 14212x_0^8x_1 + 2365x_0^7x_1^2 - 11335x_0^6x_1^3 + 10354x_0^5x_1^4 - 7342x_0^4x_1^5 + 11432x_0^3x_1^6 + \\ &- 15881x_0^2x_1^7 - 10204x_0x_1^8 - 663x_1^9 - 10837x_0^8x_2 - 6573x_0^7x_1x_2 + 6070x_0^6x_1^2x_2 - 12124x_0^5x_1^3x_2 + \\ &+ 8455x_0^4x_1^4x_2 - 9097x_0^3x_1^5x_2 + 200x_0^2x_1^6x_2 + 11563x_0x_1^7x_2 + 11173x_1^8x_2 + 2810x_0^7x_2^2 + \\ &+ 5187x_0^6x_1x_2^2 - 1688x_0^5x_1^2x_2^2 - 3089x_0^4x_1^3x_2^2 + 8745x_0^3x_1^4x_2^2 + 12508x_0^2x_1^5x_2^2 + 151x_0x_1^6x_2^2 + \\ &+ 11119x_1^7x_2^2 + 11414x_0^6x_2^3 + 2714x_0^5x_1x_2^3 + 11939x_0^4x_1^2x_2^3 + 5024x_0^3x_1^3x_2^3 + 10884x_0^2x_1^4x_2^3 + \\ &+ 8404x_0x_1^5x_2^3 + 755x_1^6x_2^3 + 15891x_0^5x_2^4 - 1013x_0^4x_1x_2^4 - 11790x_0^3x_1^2x_2^4 + 14982x_0^2x_1^3x_2^4 + \\ &- 8411x_0x_1^4x_2^4 - 5236x_1^5x_2^4 + 4416x_0^4x_2^5 - 11481x_0^3x_1x_2^5 + 14698x_0^2x_1^2x_2^5 + 5309x_0x_1^3x_2^5 + \\ &+ 11614x_1^4x_2^5 - 9777x_0^3x_2^6 - 2702x_0^2x_1x_2^6 - 5846x_0x_1^2x_2^6 - 10960x_1^3x_2^6 - 8430x_0^2x_2^7 + 7085x_0x_1x_2^7 + \\ &+ 12763x_1^2x_2^7 - 14136x_0x_2^8 - 9808x_1x_2^8 + 9194x_2^9]. \end{split}$$

Since tests 1), 2) and 3) are successful, then T_2 is general enough, so that our criterion applies. Moreover, test 4) provides positive answer too. Therefore we conclude that A is minimal for T_2 , i.e. T_2 has rank 18 (ancillary nonics2.txt).

A case of lower rank. In the same span of $v_9(A)$ one can find forms for which the decomposition A is non-redundant, yet there is another decomposition of length 17. For instance, one can take

$$(\lambda_1, \dots, \lambda_{18}) = (10308, -9437, -13956, -12270, 2135, -4854, -2213, 1755, -13629, 7308, -8496, 2940, 11348, -12437, -6712, 4086, -823, -2818)$$

so that

$$T_1 = \sum_{i=1}^{18} \lambda_i L_i^9 =$$

$$= [9666x_{0}^{9} + 13004x_{0}^{8}x_{1} + 12463x_{0}^{7}x_{1}^{2} - 13235x_{0}^{6}x_{1}^{3} - 15442x_{0}^{5}x_{1}^{4} + 15509x_{0}^{4}x_{1}^{5} + -6311x_{0}^{3}x_{1}^{6} + 2390x_{0}^{2}x_{1}^{7} + 547x_{0}x_{1}^{8} - 119x_{1}^{9} - 14916x_{0}^{8}x_{2} + 1822x_{0}^{7}x_{1}x_{2} - 8022x_{0}^{6}x_{1}^{2}x_{2} - 9386x_{0}^{5}x_{1}^{3}x_{2} + 2742x_{0}^{4}x_{1}^{4}x_{2} + 10541x_{0}^{3}x_{1}^{5}x_{2} + 1156x_{0}^{2}x_{1}^{6}x_{2} - 12023x_{0}x_{1}^{7}x_{2} + 4417x_{1}^{8}x_{2} - 11823x_{0}^{7}x_{3}^{2} - 737x_{0}^{6}x_{1}x_{1}^{2} + -7616x_{0}^{5}x_{1}^{2}x_{2}^{2} + 11293x_{0}^{4}x_{1}^{3}x_{2}^{2} - 8260x_{0}^{3}x_{1}^{4}x_{2}^{2} - 9332x_{0}^{2}x_{1}^{5}x_{2}^{2} + 7078x_{0}x_{1}^{6}x_{2}^{2} - 4553x_{1}^{7}x_{2}^{2} - 15941x_{0}^{6}x_{2}^{3} + 44339x_{0}^{5}x_{1}x_{2}^{3} - 4251x_{0}^{4}x_{1}^{2}x_{2}^{3} + 9854x_{0}^{3}x_{1}^{3}x_{2}^{3} - 22x_{0}^{2}x_{1}^{4}x_{2}^{3} + 8408x_{0}x_{1}^{5}x_{2}^{3} + 11858x_{1}^{6}x_{2}^{3} + 9161x_{0}^{5}x_{4}^{4} - 9854x_{0}^{4}x_{1}x_{2}^{4} - 13165x_{0}^{3}x_{1}^{2}x_{2}^{2} - 2105x_{0}^{2}x_{1}^{3}x_{2}^{4} - 8715x_{0}x_{1}^{4}x_{4}^{4} + 390x_{1}^{5}x_{4}^{4} - 9955x_{0}^{4}x_{2}^{5} + 11013x_{0}^{3}x_{1}x_{2}^{5} - 10651x_{0}^{2}x_{1}^{2}x_{2}^{5} - 3850x_{0}x_{1}^{3}x_{2}^{5} + 4029x_{1}^{4}x_{2}^{5} - 11735x_{0}^{3}x_{0}^{2}x_{1} - 12427x_{0}^{2}x_{1}x_{2}^{6} + 12255x_{0}x_{1}^{2}x_{2}^{6} + 3608x_{1}^{3}x_{2}^{6} - 2271x_{0}^{2}x_{2}^{7} + 5939x_{0}x_{1}x_{2}^{7} - 3402x_{1}^{2}x_{2}^{7} + 13298x_{0}x_{2}^{8} + 6455x_{1}x_{2}^{8} + x_{2}^{9}].$$

 T_1 has been obtained as the point of intersection of the span of $\nu_9(A)$ with $\nu_9(B)$, where B is a set of 17 points, linked to A with two curves Q_0 and G_0 of degree 5 and 7 in I_A . In particular, Q (resp. G_0) is defined by the determinant of the 4×4 matrix obtained by adding the row $R_1 = (1, 10399, 13534, 0)$ (resp. the row $R_2 = (0, -633x_0^2 - 11455x_0x_1 + 2134x_0x_2 + 11038x_1^2 - 8888x_1x_2 - 588x_2^2, 1927x_0^2 + 4114x_0x_1 + 11328x_0x_2 + 13814x_1^2 - 10664x_1x_2 - 1749x_2^2, 0))$ to the transpose of the Hilbert-Burch matrix M of A, i.e.

 $(a_1, a_2, \dots, a_{15}) = (1, 10399, 13534, -633, -11455, 2134, 11038, -8888, -588, 1927, 4114, 11328, 13814, -10664, -1749).$

Notice that, according to the notation of test 4), in this case $d_1 = d_2 = d_3 = 0$ and

$$\max_{4 \le i \le 15} \deg\{(a_2, \dots, a_{15}) \in \mathbb{C}^{14} \mid (t_0, \dots, t_{54}) \cdot N_1 = \underline{0} \cap a_i = 0\} = 0,$$
$$\max_{4 \le i \le 15} \deg\{(a_1, a_3, \dots, a_{15}) \in \mathbb{C}^{14} \mid (t_0, \dots, t_{54}) \cdot N_2 = \underline{0} \cap a_i = 0\} = 0,$$
$$\max_{4 \le i \le 15} \deg\{(a_1, a_2, \dots, a_{15}) \in \mathbb{C}^{14} \mid (t_0, \dots, t_{54}) \cdot N_3 = \underline{0} \cap a_i = 0\} = 0.$$

Therefore, the rank of T_1 is at most 17. Indeed, the rank is exactly 17, by Proposition 3.20 and its proof (ancillary file nonics1.txt).

In order to get coordinates for the points in B, one needs to solve the polynomial system given by the maximal minors of the 5×4 matrix $\begin{pmatrix} R_1 \\ R_2 \\ M^t \end{pmatrix}$. This can be achieved with Macaulay2 software system: indeed, by computing the eigenvalues

and eigenvectors of certain *companion* matrices (ancillary file nonics4.txt) one can find the following representative vectors in \mathbb{C}^2 for the points in B

(1, 62.6659, 29.7378)

```
(1, 13.368 + 38.1825 i, -19.099 + 7.53788 i)
    (1, 13.368 - 38.1825 i, -19.099 - 7.53788 i)
                (1, 35.333, 40.797)
               (1, 14.7061, 27.8538)
               (1, 10.7119, 4.95399)
              (1, -0.796312, 2.23381)
   (1, 1.06064 + 0.13583 i, 1.62951 - 0.563286 i)
   (1, 1.06064 - 0.13583 i, 1.62951 + 0.563286 i)
             (1, 0.737271, -0.0631582)
             (1, -0.245331, -0.76262)
             (1, -0.187307, 0.100519)
            (1, -0.0870499, -0.126324)
            (1, 0.00104432, 0.00164595)
(1, 0.306581 + 0.0182712 i, -0.877193 - 0.031211 i)
(1, 0.306581 - 0.0182712 i, -0.877193 + 0.031211 i)
              (1, 0.390447, 0.585521).
```

Remark 5.1. In principle one can try to use our analysis also to determine the identifiability of a form T. In this case one starts by adding, in the previous algorithm, the following test:

5) check that the solution set of the polynomial system $(t_0, \ldots, t_{54}) \cdot A_2 = 0_{1 \times 17}$ introduced in (23) has dimension -1. If the answer is negative, we can conclude that there are other solutions of the system (23), so one can guess that T has other decompositions.

Unfortunately, if the answer is positive, one cannot immediately conclude the identifiability of T. Namely, test 5) checks that there exists no other set B of length 18, linked to A by a complete intersection of two sextics F, F', such that T also sits in the span of $\nu_9(B)$, but only when $A \cap B = \emptyset$ and F, F' have no common components.

However, we know that there are limit cases (the case $A \cap B$ non-empty, or the case 3 of Proposition 3.5) in which B exists but the intersection $F \cap F'$ is not proper. To exclude these cases one needs new ad hoc tests, which can be constructed in principle, but then the procedure becomes quite laborious.

We believe that the natural way to prove identifiability is to produce equations for the locus Θ described in remark 4.6, and test the vanishing of the equation for T.

An unidentifiable case. Of course, one can use the construction of Case 1 of Proposition 3.5 to produce many examples of unidentifiable forms. Let

$$(\lambda_1, \dots, \lambda_{18}) = (5864, 9496, 11539, 1233, -13315, -14222, 10709, -5067, 13797, 13169, -10531, 1592, 12589, 1728, -4725, -4784, -8696, 7515)$$

and let

$$T_3 = \sum_{i=1}^{18} a_i L_i^9 =$$

 $= [11096x_{9}^{0} + 14876x_{0}^{8}x_{1} + 14398x_{0}^{7}x_{1}^{2} - 11088x_{0}^{6}x_{1}^{3} - 3441x_{0}^{5}x_{1}^{4} + 11138x_{0}^{4}x_{1}^{5} - 3819x_{0}^{3}x_{1}^{6} + \\ + 6626x_{0}^{2}x_{1}^{7} - 9525x_{0}x_{1}^{8} - 9028x_{1}^{9} + 11951x_{0}^{8}x_{2} - 14433x_{0}^{7}x_{1}x_{2} + 15878x_{0}^{6}x_{1}^{2}x_{2} + 3683x_{0}^{5}x_{1}^{3}x_{2} + \\ + 12902x_{0}^{4}x_{1}^{4}x_{2} + 9968x_{0}^{3}x_{1}^{5}x_{2} + 1167x_{0}^{2}x_{1}^{6}x_{2} - 1011x_{0}x_{1}^{7}x_{2} + 11114x_{1}^{8}x_{2} - 1174x_{0}^{7}x_{2}^{2} + \\ - 10039x_{0}^{6}x_{1}x_{2}^{2} + 15571x_{0}^{5}x_{1}^{2}x_{2}^{2} - 1797x_{0}^{4}x_{1}^{3}x_{2}^{2} + 7799x_{0}^{3}x_{1}^{4}x_{2}^{2} + 3353x_{0}^{2}x_{1}^{5}x_{2}^{2} - 9008x_{0}x_{1}^{6}x_{2}^{2} + \\ + 7892x_{1}^{7}x_{2}^{2} + 8863x_{0}^{6}x_{3}^{3} + 12538x_{0}^{5}x_{1}x_{2}3 + 4156x_{0}^{4}x_{1}^{2}x_{2}^{3} + 5014x_{0}^{3}x_{1}^{3}x_{2}^{3} + 15217x_{0}^{2}x_{1}^{4}x_{2}^{3} + \\ + 10693x_{0}x_{1}^{5}x_{2}^{3} - 14254x_{1}^{6}x_{2}^{3} - 12480x_{0}^{5}x_{2}^{4} + 15094x_{0}^{4}x_{1}x_{2}^{4} + 11796x_{0}^{3}x_{1}^{2}x_{2}^{4} - 11496x_{0}^{2}x_{1}^{3}x_{2}^{4} + \\ - 3087x_{0}x_{1}^{4}x_{2}^{4} - 7767x_{1}^{5}x_{2}^{4} + 1751x_{0}^{4}x_{2}^{5} + 9059x_{0}^{3}x_{1}x_{2}^{5} + 14238x_{0}^{2}x_{1}^{2}x_{2}^{5} - 640x_{0}x_{1}^{3}x_{2}^{5} + \\ - 14792x_{1}^{4}x_{2}^{5} - 14262x_{0}^{3}x_{2}^{6} - 6895x_{0}^{2}x_{1}x_{2}^{6} + 13550x_{0}x_{1}^{2}x_{2}^{6} + 7631x_{1}^{3}x_{2}^{6} + 9523x_{0}^{2}x_{2}^{7} + \\ - 2161x_{0}x_{1}x_{2}^{7} - 3449x_{1}^{2}x_{2}^{7} - 7220x_{0}x_{2}^{8} + 395x_{1}x_{2}^{8} + x_{2}^{9}].$

As in the previous cases, tests 1), 2), and 3) are successful for T_3 . On the other hand, in this case our computations show that the polynomial system $(t_0, \ldots, t_{54}) \cdot A_2 =$ $0_{1 \times 17}$ defined in (23) has a solution set of dimension 0 and degree 1. In particular, T_3 admits at least two decompositions of length 18, A and another set B. Indeed, $T_3 = f(F, F')$, where F (resp. F') is the ternary form of degree 6 defined by the determinant of the 4×4 matrix obtained by adding the row $(x_0 + 14307x_1 +$ $13416x_2, 11657x_0 + 9248x_1 + 8324x_2, -13193x_0 - 1403x_1 + 12171x_2, 0)$ (resp. the row $(7694x_1 + 12549x_2, -12983x_0 + 538x_1 + 11728x_2, 743x_0 - 12966x_1 + 12870x_2, 1)$) to the transpose of the Hilbert-Burch matrix M of A. Therefore, T_3 is computed by two non-redundant finite sets of length 18: A and its residual set, $B = B_{F,F'}$, in the complete intersection (6, 6) given by F and F'. According to our theory, test 5) fails for T_2 . Notice that test 4) is successful for T_3 , which means that T_3 has rank 18. CITARE ancillary nonice3.txt con referenza arXiv

A non trivial solution for the previous system determines the Hilbert-Burch matrix for B. Then, coordinates for the points of B can be found by solving (e.g. with

the software Macaulay2) the polynomial system given by the maximal minors of its Hilbert-Burch matrix. Indeed, by computing the eigenvalues and eigenvectors of certain *companion* matrices (for more details on the procedure see the ancillary file nonics5.txt), we get that the following representative vectors in \mathbb{C}^3 for the points of B:

$$\begin{array}{l} (1,-7.96881+4.74847\,i,29.737-8.31447+5.39065\,i)\\ (1,-7.96881-4.74847\,i,-8.31447-5.39065\,i)\\ (1,-8.88473,-21.3598)\\ (1,-8.88473,-21.3598)\\ (1,-3.19251,-1.98613)\\ (1,2.29572+0.339361\,i,2.08576+2.10835\,i)\\ (1,2.29572-0.339361\,i,2.08576-2.10835\,i)\\ (1,1.1725,0.789914)\\ (1,-0.662147+0.268568\,i,1.41128+0.060661\,i)\\ (1,-0.662147-0.268568\,i,1.41128+0.060661\,i)\\ (1,-0.676455+0.162048\,i,0.269336-0.242414\,i)\\ (1,-0.676455-0.162048\,i,0.269336+0.242414\,i)\\ (1,0.299266+0.586034\,i,0.441543+0.153418\,i)\\ (1,0.299266-0.586034\,i,0.441543-0.153418\,i)\\ (1,0.299266-0.586034\,i,0.441543-0.153418\,i)\\ (1,0.209511,0.479921)\\ (1,0.483517+0.0585949\,i,-0.520821-0.078578\,i)\\ (1,0.483517-0.0585949\,i,-0.520821+0.078578\,i)\\ (1,0.503511,0.553533)\end{array}$$

References

- J. Alexander and A. Hirschowitz, *Polynomial interpolation in several variables*, J. Algebraic Geom. 4 (1995), 201–222.
- E.S. Allman, C. Matias, and J.A. Rhodes, *Identifiability of parameters in latent structure models with many observed variables*, Ann. Statistics **37** (2009), 3099–3132.
- A. Anandkumar, R. Ge, D. Hsu, S.M. Kakade, and M. Telgarsky, Tensor decompositions for learning latent variable models, J. Machine Learn. Res. 15 (2014), 2773–2832.
- 4. E. Angelini and L. Chiantini, On the identifiability of ternary forms, to appear on Linear Algebra Appl.
- 5. E. Angelini, L. Chiantini, and N. Vannieuwenhoven, *Identifiability beyond Kruskal's bound for symmetric tensors of degree 4*, Rend. Lincei Mat. Applic. **29** (2018), 465–485.
- C.J. Appellof and E.R. Davidson, Strategies for analyzing data from video fluorometric monitoring of liquid chromatographic effluents, Anal. Chem. 53 (1981), 2053–2056.
- E. Ballico, An effective criterion for the additive decompositions of forms, Rend. Ist. Matem. Trieste 51 (2019), 1–12.
- E. Ballico and A. Bernardi, Decomposition of homogeneous polynomials with low rank, Math. Zeit. 271 (2012), 1141–1149.
- Unique decomposition for a polynomial of low rank, Ann. Polonici Math. 108 (2013), 219–224.
- E. Ballico and L. Chiantini, Sets computing the symmetric tensor rank, Mediterranean J. Math. 10 (2013), 643—654.
- A.M. Bigatti, A.V. Geramita, and J. Migliore, Geometric consequences of extremal behavior in a theorem of Macaulay, Trans. Amer. Math. Soc. 346 (1994), 203–235.
- G. Castelnuovo, Sui multipli di una serie lineare di gruppi di punti appartenente ad una curva algebrica, Rend. Circolo Mat. Palermo 7 (1893), 89–110.

- L. Chiantini, *Hilbert functions and tensor analysis*, Quantum Physics and Geometry, Lecture Notes of the Unione Matematica Italiana, vol. 25, Springer, Berlin, New York NY, 2019, pp. 125–151.
- L. Chiantini, G. Ottaviani, and N. Vannieuwenhoven, Effective criteria for specific identifiability of tensors and forms, SIAM J. Matrix Anal. Appl. 38 (2017), 656–681.
- On generic identifiability of symmetric tensors of subgeneric rank, Trans. Amer. Math. Soc. 369 (2017), 4021–4042.
- C. Ciliberto, A.V. Geramita, and F. Orecchia, *Remarks on a theorem of Hilbert-Burch*, Boll. UMI 7 (1988), 463–483.
- E. Davis, Hilbert functions and complete intersections, Rend. Seminario Mat. Univ. Politecnico Torino 42 (1984), 333–353.
- 18. _____, Complete intersections of codimension 2 in \mathbb{P}^r : the Bezout-Jacobi-Segre theorem revisited, Rend. Seminario Mat. Univ. Politecnico Torino **43** (1985), 333–353.
- I. Dolgachev, Lectures on invariant theory, London Mathematical Society Lecture Notes Series, vol. 296, Cambridge University press, Cambridge, UK, 2010.
- I. Domanov and L. De Lathauwer, On the uniqueness of the canonical polyadic decomposition of third-order tensors tensors- part i: Basic results and uniqueness of one factor matrix, SIAM J. Matrix Anal. Appl. 34 (2013), 855–875.
- Canonical polyadic decomposition of third-order tensors: relaxed uniqueness conditions and algebraic algorithm, Linear Algebra Appl. 513 (2017), 342–375.
- 22. D. Ferrand, Courbes gauches et fibres de rang 2, C. R. Acad. Sci. Paris 281 (1975), 345-347.
- F. Galuppi and M. Mella, Identifiability of homogeneous polynomials and Cremona transformations, Available online arXiv:1606.06895, 2016.
- A.V. Geramita, M. Kreuzer, and L. Robbiano, Cayley-Bacharach schemes and their canonical modules, Trans. Amer. Math. Soc. 339 (1993), 443–452.
- 25. D. Grayson and M. Stillman, Macaulay 2, a software system for research in algebraic geometry, Available at http://www.math.uiuc.edu/Macaulay2/.
- A. Iarrobino and V. Kanev, Power sums, Gorenstein algebras, and determinantal loci, Lecture Notes in Mathematics, vol. 1721, Springer, Berlin, New York NY, 1999.
- J.B. Kruskal, Three-way arrays: rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics, Linear Algebra Appl. 18 (1977), 95–138.
- J.M. Landsberg and G. Ottaviani, Equations for secant varieties of Veronese and other varieties, Ann. Mat. Pura Appl. 192 (2013), 569–606.
- A. Massarenti, M. Mella, and G. Staglianó, Effective identifiability criteria for tensors and polynomials, J. Symbolic Comput. 87 (2018), 227–237.
- J. Migliore, Introduction to liaison theory and deficiency modules, Progress in Mathematics, vol. 165, Birkäuser, Basel, Boston MA, 1998.
- B. Mourrain and A. Oneto, On minimal decompositions of low rank symmetric tensors, Available online arXiv:1805.11940, 2018.
- 32. C. Peskine and L. Szpiro, Liaison des variétés algébriques, Invent. Math. 26 (1974), 271–302.
- W. Rao, D. Li, and J.Q. Zhang, A tensor-based approach to L-shaped arrays processing with enhanced degrees of freedom, IEEE Signal Proc. Lett. 25 (2018), 1–5.

DIPARTIMENTO DI INGEGNERIA DELL'INFORMAZIONE E SCIENZE MATEMATICHE, UNIVERSITÀ DI SIENA, ITALY

E-mail address: elena.angelini@unisi.it

DIPARTIMENTO DI INGEGNERIA DELL'INFORMAZIONE E SCIENZE MATEMATICHE, UNIVERSITÀ DI SIENA, ITALY

E-mail address: luca.chiantini@unisi.it