

Null Geodesics and Thermodynamic Phase Transition of Four-Dimensional Gauss-Bonnet AdS Black Hole

Kartheek Hegde,^{*} Naveena Kumara A.,[†] Ahmed Rizwan C.L.,[‡] and Ajith K.M.[§]

*Department of Physics, National Institute of
Technology Karnataka, Surathkal 575 025, India*

Md Sabir Ali[¶]

*Department of Physics, Indian Institute of
Technology, Ropar, Rupnagar, Punjab 140 001, India*

Abstract

Modified gravity theories are of great interest in both observational and theoretical studies. In this article we study the correlation between the null geodesics in the background of a four-dimensional Einstein-Maxwell-Gauss-Bonnet AdS black hole, a modification of Einstein gravity, and its thermodynamic phase transition. We study the phase structure of the black hole, using the coexistence and spinodal curves, to understand the phase transition in an extended phase space. The imprints of this phase transition features are observed in the behaviour of photon orbit radius and minimum impact parameter with respect to the Hawking temperature and pressure. The change in these two quantities during the phase transition serve as order parameter which characterises the critical behaviour. The correlation shows that thermodynamic phase transition can be studied by observing the effects of strong gravitational field and vice versa.

PACS numbers:

Keywords: 4D Gauss-Bonnet gravity, Null geodesics, Black hole thermodynamics, Phase transition

^{*}Electronic address: hegde.kartheek@gmail.com

[†]Electronic address: naviphsics@gmail.com

[‡]Electronic address: ahmedrizwancl@gmail.com

[§]Electronic address: ajith@nitk.ac.in

[¶]Electronic address: alimd.sabir3@gmail.com

I. INTRODUCTION

The existence of black hole singularity is one of the most fundamental question in physics. The Penrose cosmic censorship hypothesis asserts that the spacetime singularities need to be hidden from an observer at infinity by an event horizon, which blocks all of the information within it [1, 2]. Generally all the electrovacuum solutions of classical general relativity are consistent with this conjecture. However, the conjecture does not restrain us from considering black hole spacetimes which are free from singularity, within classical general relativity. In this context, recently proposed theory of four dimensional Gauss-Bonnet gravity is quite interesting one [3]. It was demonstrated that for a positive Gauss-Bonnet coupling parameter α , the static and spherically symmetric solution of the theory is free from the much debated singularity problem. The theory is captivating for other reasons too, for example, the obtained black hole solution appears in the setting of the gravity with a conformal anomaly [4, 5], and also in the context of quantum corrections [6, 7]. However, the black hole solution in four dimensional Gauss-Bonnet theory is attractive, as it is a modified theory of classical gravity, and hence is on an equal footing with general relativity. These captivating features of this novel theory resulted in a surge of various investigations around this theory, including the theoretical aspects, viability of the solution and physical properties [8–36].

It is well known that black holes are not merely strong gravity systems, but also a thermal systems. Particularly, the establishment of laws of black hole thermodynamics has made the phase transition of these compact objects appealing in every sense [37, 38]. In recent times, anti-de Sitter black hole thermodynamics have gained more interest, as the identification of cosmological constant as the thermodynamic variable pressure, leads to the modification in first law, which has a conventional PdV term [39, 40]. In this extended phase space, AdS black holes exhibit variety of phase transition features, of which, the van der Waals like transition is of great interest [41–43]. As in the case of a conventional van der Waals fluid, the black hole shows a first order phase transition between two phases, namely, the large black hole phase and the small black hole phase. The authors have studied the thermodynamics of the four dimensional Gauss-Bonnet AdS black hole for both the charged and uncharged cases [44], and it was observed that a vdW like phase transition exists. Having said so, as the black hole is a gravitational and also a thermal system, it is quite natural to seek a connection between the effects of strong gravity and phase transition.

It is customary to seek the details of a gravitating object, especially a compact object with strong gravity, by observing the characteristic features of a test particle moving along the geodesics around it. For a particle moving in the vicinity of a black hole, the black hole features are expected to be encoded in the behaviour of the particle motion. These notions are exploited in connecting the unseen attributes of black hole to observational aspects, for example, black hole shadow and quasinormal modes [45, 46]. Along with this, the phase transition signature of a charged AdS black hole can be obtained using the quasinormal mode (QNM) studies [47]. It was reported that during the van der Waals like phase transition of the black hole, the slope of the quasinormal mode changes drastically, which is an observable phenomenon.

These initial findings motivated to investigate a more concrete relationship between the gravitational and phase transition features of the AdS black holes using the null geodesics [48, 49]. By studying the photon orbits in the background of a charged AdS black hole, the phase transition properties are observed from the behaviour of radius r_{ps} and minimum impact parameter u_{ps} of the circular orbit. The behaviour of r_{ps} and u_{ps} with the Hawking temperature T and pressure P , mimics the isobar and isotherms found in thermodynamics counterpart. Below the critical values, the first order phase transition is reflected by these orbital parameters. During the phase transition, these two parameters change by a finite amount, which serves as order parameters to characterise the black hole phase transition, with a critical exponent $1/2$. Originally, this was observed in a charged AdS black holes [48], this correlation between gravity and thermodynamics, via photon orbits, can be seen in different black hole spacetimes, namely, Kerr-AdS [49], Born-Infeld AdS background [50], regular AdS black holes [51], massive gravity [52], Born-Infeld-dilaton black hole [53], five-dimensional Gauss-Bonnet black holes [54] etc. Related studies in other contexts have also appeared in subsequent works [55–57]. In this article we seek a similar correlation for the novel four-dimensional Gauss-Bonnet AdS black hole.

The article is organised as follows. In the next section (II) we briefly present the 4D Gauss-Bonnet AdS black hole solution and its thermodynamics. In section III we investigate the phase transition features of the black hole, wherein, the phase structure is probed using the coexistence and metastable curves. This is followed by section IV, where we consider the null geodesics on the equatorial plane, and hence obtain the photon orbit radius r_{ps} and minimum impact parameter u_{ps} . In section V, we study the critical behaviour of r_{ps} and

u_{ps} , where the order parameters are presented. Finally, we conclude the paper in section VI.

II. 4D GAUSS-BONNET ADS BLACK HOLE: METRIC AND THERMODYNAMICS

In this section we briefly present the black hole metric and its thermodynamics. The D -dimensional Einstein-Maxwell-Gauss-Bonnet theory with a negative cosmological constant Λ is described by the action [12],

$$\mathcal{I} = \frac{1}{16\pi} \int d^D x \sqrt{-g} [R + 2\Lambda + \alpha \mathcal{G} - F^{ab} F_{ab}], \quad (1)$$

where g is the determinant of the metric g_{ab} , $F_{ab} = \partial_a A_b - \partial_b A_a$, is the Maxwell field tensor and α is the Gauss-Bonnet coupling coefficient. The Gauss Bonnet term is given by,

$$\mathcal{G} = R^2 - 4R_{ab}R^{ab} + R_{abcd}R^{abcd}, \quad (2)$$

where R is the Ricci scalar, R_{ab} is the Ricci tensor, R_{abcd} is the Riemann tensor. The cosmological constant is related to the AdS radius l as,

$$\Lambda = -\frac{(D-1)(D-2)}{2l^2}. \quad (3)$$

In four dimensions the Gauss-Bonnet term does not contribute to the dynamics of the system, as the integral over that term is a topological invariant. However, recently a genuine four dimensional Einstein-Gauss-Bonnet gravity was obtained by scaling α as [3],

$$\alpha \rightarrow \frac{\alpha}{D-4}, \quad (4)$$

and then taking the limit $D \rightarrow 4$. The spherically symmetric solution for the action (1) is,

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2 d\Omega_{D-2}^2. \quad (5)$$

In the limit $D \rightarrow 4$ the metric function has the form,

$$f(r) = 1 + \frac{r^2}{2\alpha} \left(1 - \sqrt{1 + 4\alpha \left(-\frac{1}{l^2} + \frac{2M}{r^3} - \frac{Q^2}{r^4} \right)} \right), \quad (6)$$

where M is the ADM mass and Q is the total charge of the black hole. The validity of the theory from which we obtained the above static spherically symmetric solution has been

scrutinised in detail in several propositions [27, 33–35, 58–60]. However, these does not rule out the possibility of having a spherically symmetric solution as it can be obtained from consistent formulations [13, 17, 32, 61, 62]. Therefore we approach the solution (5) as a self-reliant one. Interestingly, this solution also appears in the context of a conformal anomaly gravity [4, 5].

The horizon of the black hole (r_+) is defined by the condition $f(r_+) = 0$. Using this condition we obtain the mass of the black hole to be,

$$M = \frac{r_+^3}{2l^2} + \frac{Q^2}{2r_+} + \frac{\alpha}{2r_+} + \frac{r_+}{2}. \quad (7)$$

We present the thermodynamics of the black hole in an extended phase space, where the cosmological constant (Λ) is treated as the thermodynamic pressure (P), and they are related as $P = -\frac{\Lambda}{8\pi}$. The Hawking temperature of the black hole is associated with the surface gravity κ , which is,

$$T = \frac{\kappa}{2\pi} = \frac{f'(r)}{4\pi} \Big|_{r=r_+} = -\frac{\alpha - 8\pi P r_+^4 + Q^2 - r_+^2}{4\pi r_+^3 + 8\pi \alpha r_+}. \quad (8)$$

The first law of black hole can be written, considering the GB coupling parameter α to be a thermodynamic variable [63, 64], as,

$$dM = TdS + VdP + \Phi dQ + \mathcal{A} d\alpha \quad (9)$$

where the potentials Φ and \mathcal{A} are conjugate to Q and α , respectively. Likewise, the thermodynamic volume V is a conjugate to pressure P ,

$$V = \left(\frac{\partial M}{\partial P} \right)_{S, Q, \alpha} = \frac{4}{3} \pi r_+^3. \quad (10)$$

The entropy of the black hole can be obtained as follows,

$$S = \int_0^{r_+} \frac{1}{T} dM = \frac{A}{4} + 2\pi\alpha \ln \left(\frac{A}{A_0} \right), \quad (11)$$

where $A = 4\pi r_+^2$ is the horizon area and A_0 is the integration constant, which has the dimension of $[length]^2$. It is clear that the Gauss Bonnet coupling parameter α modifies the Bekenstein-Hawking entropy-area law. In general, the black hole entropy is independent of the charge Q and cosmological constant Λ , therefore, the integration constant can be set as $A_0 = 4\pi|\alpha|$ [21]. With this identification, the entropy reads,

$$S = \pi r_+^2 + 4\pi\alpha \ln \left(\frac{r_+}{\sqrt{|\alpha|}} \right). \quad (12)$$

We emphasise that the black hole entropy has a logarithmic correction, whereas, the thermodynamic volume remains same as the geometric volume. Before concluding the thermodynamics of the black hole, we also mention that, the variables presented above satisfy the Smarr relation in addition to the first law,

$$M = 2TS + \Phi Q - 2PV + 2\alpha\mathcal{A}. \quad (13)$$

III. PHASE TRANSITION OF 4D GAUSS BONNET ADS BLACK HOLE

The phase transition of the 4D Gauss-Bonnet black hole has been well studied by the authors [44]. Here we recall them to analyse the phase structure using the coexistence and spinodal curves. The state equation of the system is obtained by inverting the expression for Hawking temperature,

$$P = \frac{Q^2}{8\pi r_+^4} + \frac{\alpha}{8\pi r_+^4} + \frac{\alpha T}{r_+^3} - \frac{1}{8\pi r_+^2} + \frac{T}{2r_+}. \quad (14)$$

In terms of volume we have,

$$P = \frac{(6\pi)^{2/3}(\alpha + Q^2)}{18\pi^{1/3}V^{4/3}} + \frac{4\pi\alpha T}{3V} + \frac{\pi^{1/3}T}{6V^{1/3}} - \frac{1}{2 \times 6^{2/3}\pi^{1/3}V^{2/3}}. \quad (15)$$

The critical behaviour of the black hole can be easily seen in the $P - V$ isotherms, where a first order phase transition exists between a small black hole phase (SBH) and a large black hole phase (LBH). This phase transition property is exhibited by both the charged and neutral black holes. The critical point of the phase transitions is determined by using the condition,

$$\left(\frac{\partial P}{\partial V}\right)_{T,Q,\alpha} = \left(\frac{\partial^2 P}{\partial V^2}\right)_{T,Q,\alpha} = 0. \quad (16)$$

The critical values of the thermodynamic variables are [44],

$$T_c = \frac{(8\alpha + 3Q^2 - \rho) \sqrt{6\alpha + 3Q^2 + \rho}}{48\pi\alpha^2}; \quad (17)$$

$$P_c = \frac{9\alpha + 6Q^2 + \rho}{24\pi(6\alpha + 3Q^2 + \rho)^2}; \quad (18)$$

$$V_c = \frac{4}{3}\pi(6\alpha + 3Q^2 + \rho)^{3/2}; \quad (19)$$

where $\rho = \sqrt{48\alpha^2 + 9Q^4 + 48\alpha Q^2}$. Making use of these quantities we define the reduced thermodynamic variables,

$$\tilde{T} = \frac{T}{T_c} \quad \tilde{P} = \frac{P}{P_c} \quad \tilde{V} = \frac{V}{V_c}. \quad (20)$$




Figure 1: The coexistence curve (red solid line) and Spinodal curve (blue dashed line) in $\tilde{P} - \tilde{T}$ and $\tilde{T} - \tilde{V}$ plane. The black dot at (1,1) denotes the critical point.

By observing the phase structure, we can have a better understanding of the phase transition. In the extended phase space, the black hole mass plays the role of enthalpy, which is evident from the first law (9). With this understanding, the Gibbs free energy of the black hole is calculated to be $G = M - TS$, which reads,

$$G = \frac{4}{3}\pi P r_+^3 + \frac{Q^2}{2r_+} - T \left[\pi r_+^2 + 4\pi\alpha \log \left(\frac{r_+}{\sqrt{\alpha}} \right) \right] + \frac{\alpha}{2r_+} + \frac{r_+}{2}. \quad (21)$$

Here r_+ is regarded as a function of (P, T) from equation of state. We obtain the coexistence curve in the $\tilde{P} - \tilde{T}$ plane, by using the swallow tail behaviour of the Gibbs free energy. The coexistence expression is also translated into $\tilde{T} - \tilde{V}$ plane. The results are shown in fig. 1. In the $\tilde{P} - \tilde{T}$ plane, the coexistence line (red solid line) partitions the SBH and LBH phases below the critical point. It terminates at the second order phase transition point, above which the phase is supercritical. The figures also display the metastable curves (blue dashed lines) which satisfy,

$$(\partial_V P)_T = 0, \quad (\partial_V T)_P = 0. \quad (22)$$

The region between the coexistence curve and metastable curve are the metastable phases, namely, superheated SBH and supercooled LBH phases. In the $\tilde{T} - \tilde{V}$ plane the region under the metastable curve corresponds to the coexistence phase of SBH and LBH.

IV. GEODESIC EQUATIONS OF MOTION

In this section we establish the relationship between the thermodynamics and the null geodesics. Consider a photon which is orbiting the black hole freely on the equatorial plane described by $\theta = \pi/2$. The Lagrangian which characterises this motion can be directly written from the metric (5),

$$2\mathcal{L} = -f(r)\dot{t}^2 + \frac{\dot{r}^2}{f(r)} + r^2\dot{\phi}^2. \quad (23)$$

Here, the dots represent the differentiation with respect to an affine parameter. The 4D Gauss-Bonnet AdS black hole spacetime has two Killing fields, ∂_t and ∂_ϕ , which leads to two constants of motion, E and L , which are the conserved quantities, the energy and orbital angular momentum of the photon, respectively. The generalised momenta corresponding to the Lagrangian (23) can be obtained by using $p_a = g_{ab}\dot{x}^b$ as,

$$p_t = -f(r)\dot{t} \equiv E \quad (24)$$

$$p_\phi = r^2\dot{\phi} \equiv L \quad (25)$$

$$p_r = \dot{r}/f(r). \quad (26)$$

The t and r motion of the photon can now be described as,

$$\dot{t} = \frac{E}{f(r)} \quad (27)$$

$$\dot{\phi} = \frac{L}{r^2 \sin^2 \theta}. \quad (28)$$

The Hamiltonian for the system is obtained from the standard definition, and it vanishes,

$$2\mathcal{H} = -E\dot{t} + L\dot{\phi} + \dot{r}^2/f(r) = 0. \quad (29)$$

Employing r and ϕ motion, we can rewrite the expression for the radial r motion as,

$$\dot{r}^2 + V_{eff} = 0, \quad (30)$$

with the effective potential given by,

$$V_{eff} = \frac{L^2}{r^2}f(r) - E^2. \quad (31)$$

The photon can only move in the region where $V_{eff} < 0$, since $\dot{r}^2 > 0$. A photon approaching the black hole will be absorbed if it has a smaller angular momentum L and get scattered

if the angular momentum is large enough. The absorption and scattering are separated by a critical angular momentum, which defines a unstable circular photon orbit. Expressions governing this orbit are,

$$V_{eff} = 0 \quad , \quad V'_{eff} = 0 \quad , \quad V''_{eff} < 0, \quad (32)$$

where prime denotes a differentiation with respect to r . The radial velocity \dot{r} of the photon is zero in this unstable circular orbit. The corresponding value of r is the radius of photon orbit. Expanding the second equation in (32) we have,

$$2f(r_{ps}) - r_{ps}\partial_r f(r_{ps}) = 0. \quad (33)$$

Substituting the metric function (6) into this equation and solving, we obtain the expression for the radius of photon sphere r_{ps} , which is a complicated expression and a function of black hole parameters (M, Q, P, α) . Solving the first equation in (32), ($V_{eff} = 0$), we obtain the minimum impact parameter of the photon as,

$$u_{ps} = \frac{L_c}{E} = \frac{r}{\sqrt{f(r)}} \Big|_{r_{ps}}. \quad (34)$$

To investigate the correlation between the photon sphere and the black hole phase transition, we observe the behaviour of the radius r_{ps} and minimum impact parameter u_{ps} with respect to the Hawking temperature and pressure, in the reduced parameter space. Apparently this investigation is motivated by the observations in the phenomena of black hole lensing, where the impact parameter u has a close connection with the deflection angle. The deflection angle is small for a large impact parameter. Yet, in the limit $u \rightarrow u_{ps}$, the deflection angle is unbounded [65].

In fig. 2, the Hawking temperature T is shown as a function of the photon orbit radius r_{ps} and minimum impact parameter u_{ps} , separately, with fixed pressures. The isobars in this figure imply the typical van der Waals like phase transition. For pressures below the critical value, the isobars first increase, then decrease, and finally increase with respect to the photon sphere radius r_{ps} and minimum impact parameter u_{ps} . In fig. 3 the pressure P is seen as function of r_{ps} , first, and then of u_{ps} , keeping the temperature as a constant. The behaviour of pressure here (fig. 3) is opposite to that of temperature (fig. 2). For example, when temperature \tilde{T} increases with r_{ps} or u_{ps} , the pressure \tilde{P} decreases. In summary, from the behaviour of the photon orbit radius and minimum impact parameter along the




Figure 2: The behaviour of photon sphere radius and minimum impact parameter of unstable null geodesic with Hawking temperature in reduced parameter space. We take $Q = 1$ and $\alpha = 0.5$




Figure 3: The behaviour of photon sphere radius and minimum impact parameter of unstable null geodesic with pressure in reduced parameter space. We take $Q = 1$ and $\alpha = 0.5$

isothermal and isobaric curves of 4D Gauss-Bonnet AdS black hole, the van der Waals like phase transition can be clearly identified. This affirms that there exists a correlation between null geodesics and phase transition of the black hole.

V. CRITICAL BEHAVIOUR OF THE PHOTON SPHERE

The black hole exhibits a first order vdW like phase transition which terminates at the critical point, which corresponds to second order phase transition. As we have seen, there is a connection between the photon sphere and phase transition, it is worth examining the behaviour of change in photon orbit and minimum impact parameter during the phase




Figure 4: The variation of the radius of photon sphere and the minimum impact parameter, for unstable null geodesic, with respect to the Hawking temperature (in reduced parameter space). The SBH (blue dashed line) and LBH (red solid line) meet at the critical point ($\tilde{T} = 1$).




Figure 5: The change in photon sphere radius and minimum impact parameter of unstable null geodesic during the phase transition of the black hole. The concavity of the curve changes near the critical point, which is shown in an enlarged form in inlets.

transition. We construct the equal area law for the $\tilde{T} - \tilde{r}_{ps}$ and $\tilde{T} - \tilde{u}_{ps}$ isobars, similar to the isobars in the $\tilde{T} - \tilde{S}$ plane of the black hole. From the result, we study the behaviour of the photon orbit radius r_{ps} along the coexistence curve (Fig. 4). As the temperature increases, the radius r_{ps} for the coexistence LBH phase decreases, whereas for the coexistence SBH phase it increases. The r_{ps} of both coexistence phases attain same value at the critical point $\tilde{T} = 1$. Same behaviour is observed for the minimum impact parameter u_{ps} . In fig 5 we display the differences of the quantities r_{ps} and u_{ps} with the phase transition temperature. Both Δr_{ps} and Δu_{ps} behaves acutely like the order parameter. They have non-zero value

corresponding to first-order phase transition and vanish at the second-order phase transition. The behaviour in the neighbourhood of critical point is shown in the insets. Near the critical point a change in concavity is observed. We numerically obtain the critical exponent of these differences near the critical point to be,

$$\Delta\tilde{r}_{ps} = 3.57249(1 - \tilde{T})^{0.510839} \quad (35)$$

and

$$\Delta\tilde{u}_{ps} = 2.54786(1 - \tilde{T})^{0.506096}. \quad (36)$$

This behaviour, i.e. $\Delta\tilde{r}_{ps} \sim (1 - \tilde{T})^{1/2}$ and $\Delta\tilde{u}_{ps} \sim (1 - \tilde{T})^{1/2}$, show that $\Delta\tilde{r}_{ps}$ and $\Delta\tilde{u}_{ps}$ can serve as the order parameters to characterise the black hole phase transition. These results strongly confirm our previous assertion that photon orbits and thermodynamic phase transitions are related to each other.

VI. CONCLUDING REMARKS

In this article we show that the unstable circular photon orbit around the four dimensional Gauss-Bonnet AdS black hole reflects the phase transition information of the black hole. The radius of the photon orbit r_{ps} and the minimum impact parameter u_{ps} are studied in detail. The study establishes a link between the gravity and thermodynamics in the background of Gauss-Bonnet AdS strong gravity.

In the first part of the article we presented the thermodynamics and phase transition of the black hole. The phase structure of the black hole is analysed using the coexistence curve and the metastable curves. These curves are the boundaries that separates different stable and metastable phases of the black holes, using which a clear understanding of phase transition features are obtained. The first-order and second order phase transition details are sought in this study, which are influenced by the Gauss-Bonnet coupling parameter α . Throughout our study we keep in mind that, the extended phase space thermodynamics features are same for both the charged and neutral Gauss-Bonnet AdS black holes, as it was reported in our previous work [44].

In the second part of the article, using the Lagrangian of a photon moving freely in the equatorial plane of the black hole we investigated the null geodesics. Using the effective potential, we solve the photon orbit radius r_{ps} and the minimum impact parameter u_{ps} for

the 4D Gauss-Bonnet AdS black hole. These two key quantities depend on the black hole parameters, especially the charge Q and Gauss-Bonnet coupling parameter α . To establish the relationship between the photon sphere and black hole phase transition we study the behaviour of r_{ps} and u_{ps} along the isobar and the isotherms of the system. The first order phase transition is revealed from these plots. When the pressure or temperature is below the critical value there exists two extreme values for r_{ps} and u_{ps} , which coincide to form one extreme point for the critical values of pressure or temperature. Above the critical value of pressure or temperature the r_{ps} and u_{ps} do not exhibit any extremum. Thus they increase monotonically. This behaviour of the photon orbit isobar and isotherm are consistent with that of the black hole thermodynamics. Finally we probe the behaviour of r_{ps} and u_{ps} along the coexistence curve. The two coexistence branches, namely, small black hole and large black hole, have different r_{ps} and u_{ps} values. Their differences Δr_{ps} and Δu_{ps} serve as order parameters for the black hole phase transition. They vanish near the critical point, which corresponds to the second order phase transition. In the neighbourhood of this critical point, Δr_{ps} and Δu_{ps} have a critical exponent of $1/2$, which is obtained numerically. Our results show that in the background of Einstein-Maxwell-Gauss-Bonnet AdS spacetime, the black hole thermodynamics can be probed via the strong gravity effects and vice versa.

Acknowledgments

K.H. , N.K.A. and A.R.C.L. would like to thank U.G.C. Govt. of India for financial assistance under UGC-NET-SRF scheme.

-
- [1] S. W. Hawking and R. Penrose, *The Singularities of gravitational collapse and cosmology*, *Proc. Roy. Soc. Lond.* **A314** (1970) 529.
 - [2] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*, Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2011, [10.1017/CBO9780511524646](https://doi.org/10.1017/CBO9780511524646).
 - [3] D. Glavan and C. Lin, *Einstein-Gauss-Bonnet gravity in 4-dimensional space-time*, *Phys. Rev. Lett.* **124** (2020) 081301 [[1905.03601](https://arxiv.org/abs/1905.03601)].

- [4] R.-G. Cai, L.-M. Cao and N. Ohta, *Black Holes in Gravity with Conformal Anomaly and Logarithmic Term in Black Hole Entropy*, *JHEP* **04** (2010) 082 [[0911.4379](#)].
- [5] R.-G. Cai, *Thermodynamics of Conformal Anomaly Corrected Black Holes in AdS Space*, *Phys. Lett.* **B733** (2014) 183 [[1405.1246](#)].
- [6] Y. Tomozawa, *Quantum corrections to gravity*, [1107.1424](#).
- [7] G. Cognola, R. Myrzakulov, L. Sebastiani and S. Zerbini, *Einstein gravity with Gauss-Bonnet entropic corrections*, *Phys. Rev. D* **88** (2013) 024006 [[1304.1878](#)].
- [8] R. A. Konoplya and A. F. Zinhailo, *Quasinormal modes, stability and shadows of a black hole in the novel 4D Einstein-Gauss-Bonnet gravity*, [2003.01188](#).
- [9] M. Guo and P.-C. Li, *The innermost stable circular orbit and shadow in the novel 4D Einstein-Gauss-Bonnet gravity*, [2003.02523](#).
- [10] A. Casalino, A. Colleaux, M. Rinaldi and S. Vicentini, *Regularized Lovelock gravity*, [2003.07068](#).
- [11] R. Konoplya and A. Zhidenko, *Black holes in the four-dimensional Einstein-Lovelock gravity*, [2003.07788](#).
- [12] P. G. Fernandes, *Charged Black Holes in AdS Spaces in 4D Einstein Gauss-Bonnet Gravity*, *Phys. Lett. B* **805** (2020) 135468 [[2003.05491](#)].
- [13] H. Lu and Y. Pang, *Horndeski Gravity as $D \rightarrow 4$ Limit of Gauss-Bonnet*, [2003.11552](#).
- [14] R. A. Konoplya and A. Zhidenko, *BTZ black holes with higher curvature corrections in the 3D Einstein-Lovelock theory*, [2003.12171](#).
- [15] S. G. Ghosh and R. Kumar, *Generating black holes in the novel 4D Einstein-Gauss-Bonnet gravity*, [2003.12291](#).
- [16] R. A. Konoplya and A. Zhidenko, *(In)stability of black holes in the 4D Einstein-Gauss-Bonnet and Einstein-Lovelock gravities*, [2003.12492](#).
- [17] T. Kobayashi, *Effective scalar-tensor description of regularized Lovelock gravity in four dimensions*, *JCAP* **07** (2020) 013 [[2003.12771](#)].
- [18] C.-Y. Zhang, P.-C. Li and M. Guo, *Greybody factor and power spectra of the Hawking radiation in the novel 4D Einstein-Gauss-Bonnet de-Sitter gravity*, [2003.13068](#).
- [19] S. A. Hosseini Mansoori, *Thermodynamic geometry of novel 4-D Gauss Bonnet AdS Black Hole*, [2003.13382](#).
- [20] A. Kumar and R. Kumar, *Bardeen black holes in the novel 4D Einstein-Gauss-Bonnet*

- gravity, [2003.13104](#).
- [21] S.-W. Wei and Y.-X. Liu, *Extended thermodynamics and microstructures of four-dimensional charged Gauss-Bonnet black hole in AdS space*, *Phys. Rev. D* **101** (2020) 104018 [[2003.14275](#)].
 - [22] M. S. Churilova, *Quasinormal modes of the Dirac field in the novel 4D Einstein-Gauss-Bonnet gravity*, [2004.00513](#).
 - [23] S. U. Islam, R. Kumar and S. G. Ghosh, *Gravitational lensing by black holes in 4D Einstein-Gauss-Bonnet gravity*, [2004.01038](#).
 - [24] C. Liu, T. Zhu and Q. Wu, *Thin Accretion Disk around a four-dimensional Einstein-Gauss-Bonnet Black Hole*, [2004.01662](#).
 - [25] R. Konoplya and A. F. Zinhailo, *Grey-body factors and Hawking radiation of black holes in 4D Einstein-Gauss-Bonnet gravity*, [2004.02248](#).
 - [26] X.-h. Jin, Y.-x. Gao and D.-j. Liu, *Strong gravitational lensing of a 4D Einstein-Gauss-Bonnet black hole in homogeneous plasma*, [2004.02261](#).
 - [27] W.-Y. Ai, *A note on the novel 4D Einstein-Gauss-Bonnet gravity*, [2004.02858](#).
 - [28] M. Heydari-Fard, M. Heydari-Fard and H. R. Sepangi, *Bending of light in novel 4D Gauss-Bonnet-de Sitter black holes by Rindler-Ishak method*, [2004.02140](#).
 - [29] S.-L. Li, P. Wu and H. Yu, *Stability of the Einstein Static Universe in 4D Gauss-Bonnet Gravity*, [2004.02080](#).
 - [30] S.-W. Wei and Y.-X. Liu, *Testing the nature of Gauss-Bonnet gravity by four-dimensional rotating black hole shadow*, [2003.07769](#).
 - [31] R. Kumar and S. G. Ghosh, *Rotating black holes in the novel 4D Einstein-Gauss-Bonnet gravity*, [2003.08927](#).
 - [32] R. A. Hennigar, D. Kubiznak, R. B. Mann and C. Pollack, *On Taking the $D \rightarrow 4$ limit of Gauss-Bonnet Gravity: Theory and Solutions*, [2004.09472](#).
 - [33] S. Mahapatra, *A note on the total action of 4D Gauss-Bonnet theory*, [2004.09214](#).
 - [34] F.-W. Shu, *Vacua in novel 4D Einstein-Gauss-Bonnet Gravity: pathology and instability?*, [2004.09339](#).
 - [35] M. Gurses, T. C. Sisman and B. Tekin, *Is there a novel Einstein-Gauss-Bonnet theory in four dimensions?*, [2004.03390](#).
 - [36] A. Naveena Kumara, C. A. Rizwan, K. Hegde, M. S. Ali and A. K. M., *Rotating 4D*

- Gauss-Bonnet black hole as particle accelerator*, [2004.04521](#).
- [37] J. D. Bekenstein, *Black holes and entropy*, *Phys. Rev.* **D7** (1973) 2333.
 - [38] J. M. Bardeen, B. Carter and S. Hawking, *The Four laws of black hole mechanics*, *Commun. Math. Phys.* **31** (1973) 161.
 - [39] D. Kastor, S. Ray and J. Traschen, *Enthalpy and the Mechanics of AdS Black Holes*, *Class. Quant. Grav.* **26** (2009) 195011 [[0904.2765](#)].
 - [40] B. P. Dolan, *Pressure and volume in the first law of black hole thermodynamics*, *Class. Quant. Grav.* **28** (2011) 235017 [[1106.6260](#)].
 - [41] D. Kubiznak and R. B. Mann, *P-V criticality of charged AdS black holes*, *JHEP* **07** (2012) 033 [[1205.0559](#)].
 - [42] S. Gunasekaran, R. B. Mann and D. Kubiznak, *Extended phase space thermodynamics for charged and rotating black holes and Born-Infeld vacuum polarization*, *JHEP* **11** (2012) 110 [[1208.6251](#)].
 - [43] D. Kubiznak, R. B. Mann and M. Teo, *Black hole chemistry: thermodynamics with Lambda*, *Class. Quant. Grav.* **34** (2017) 063001 [[1608.06147](#)].
 - [44] K. Hegde, A. Naveena Kumara, C. A. Rizwan, A. K. M. and M. S. Ali, *Thermodynamics, Phase Transition and Joule Thomson Expansion of novel 4-D Gauss Bonnet AdS Black Hole*, [2003.08778](#).
 - [45] V. Cardoso, A. S. Miranda, E. Berti, H. Witek and V. T. Zanchin, *Geodesic stability, Lyapunov exponents and quasinormal modes*, *Phys. Rev.* **D79** (2009) 064016 [[0812.1806](#)].
 - [46] I. Z. Stefanov, S. S. Yazadjiev and G. G. Gyulchev, *Connection between Black-Hole Quasinormal Modes and Lensing in the Strong Deflection Limit*, *Phys. Rev. Lett.* **104** (2010) 251103 [[1003.1609](#)].
 - [47] Y. Liu, D.-C. Zou and B. Wang, *Signature of the Van der Waals like small-large charged AdS black hole phase transition in quasinormal modes*, *JHEP* **09** (2014) 179 [[1405.2644](#)].
 - [48] S.-W. Wei and Y.-X. Liu, *Photon orbits and thermodynamic phase transition of d-dimensional charged AdS black holes*, *Phys. Rev.* **D97** (2018) 104027 [[1711.01522](#)].
 - [49] S.-W. Wei, Y.-X. Liu and Y.-Q. Wang, *Probing the relationship between the null geodesics and thermodynamic phase transition for rotating Kerr-AdS black holes*, *Phys. Rev.* **D99** (2019) 044013 [[1807.03455](#)].
 - [50] Y.-M. Xu, H.-M. Wang, Y.-X. Liu and S.-W. Wei, *Photon sphere and reentrant phase*

- transition of charged Born-Infeld-AdS black holes, *Phys. Rev. D* **100** (2019) 104044 [[1906.03334](#)].
- [51] N. K. A., C. A. Rizwan, A. K. M. and M. S. Ali, *Photon Orbits and Thermodynamic Phase Transition of Regular AdS Black Holes*, [1912.11909](#).
- [52] M. Chabab, H. El Moumni, S. Iraoui and K. Masmar, *Probing correlation between photon orbits and phase structure of charged AdS black hole in massive gravity background*, [1902.00557](#).
- [53] H. Li, Y. Chen and S.-J. Zhang, *Photon orbits and phase transitions in Born-Infeld-dilaton black holes*, [1908.09570](#).
- [54] S.-Z. Han, J. Jiang, M. Zhang and W.-B. Liu, *Photon orbits and thermodynamic phase transition in Gauss-Bonnet AdS black holes*, [1812.11862](#).
- [55] M. Zhang, S.-Z. Han, J. Jiang and W.-B. Liu, *Circular orbit of a test particle and phase transition of a black hole*, *Phys. Rev. D* **99** (2019) 065016 [[1903.08293](#)].
- [56] B. Chandrasekhar and S. Mohapatra, *A Note on Circular Geodesics and Phase Transitions of Black Holes*, *Phys. Lett. B* **791** (2019) 367 [[1805.05088](#)].
- [57] S.-W. Wei and Y.-X. Liu, *Null Geodesics, Quasinormal Modes, and Thermodynamic Phase Transition for Charged Black Holes in Asymptotically Flat and dS Spacetimes*, [1909.11911](#).
- [58] S. Tian and Z.-H. Zhu, *Comment on "Einstein-Gauss-Bonnet Gravity in Four-Dimensional Spacetime"*, [2004.09954](#).
- [59] J. Bonifacio, K. Hinterbichler and L. A. Johnson, *Amplitudes and 4D Gauss-Bonnet Theory*, *Phys. Rev. D* **102** (2020) 024029 [[2004.10716](#)].
- [60] J. Arrechea, A. Delhom and A. Jimnez-Cano, *Yet another comment on four-dimensional Einstein-Gauss-Bonnet gravity*, [2004.12998](#).
- [61] P. G. Fernandes, P. Carrilho, T. Clifton and D. J. Mulryne, *Derivation of Regularized Field Equations for the Einstein-Gauss-Bonnet Theory in Four Dimensions*, *Phys. Rev. D* **102** (2020) 024025 [[2004.08362](#)].
- [62] K. Aoki, M. A. Gorji and S. Mukohyama, *A consistent theory of $D \rightarrow 4$ Einstein-Gauss-Bonnet gravity*, [2005.03859](#).
- [63] R.-G. Cai, L.-M. Cao, L. Li and R.-Q. Yang, *P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space*, *JHEP* **09** (2013) 005 [[1306.6233](#)].
- [64] S.-W. Wei and Y.-X. Liu, *Triple points and phase diagrams in the extended phase space of*

- charged Gauss-Bonnet black holes in AdS space*, *Phys. Rev. D* **90** (2014) 044057 [[1402.2837](#)].
- [65] V. Bozza, *Gravitational lensing in the strong field limit*, *Phys. Rev.* **D66** (2002) 103001 [[gr-qc/0208075](#)].