

NEW BOUNDS ON MAXIMAL LINKLESS GRAPHS

RAMIN NAIMI, ANDREI PAVELESCU, AND ELENA PAVELESCU

ABSTRACT. We construct a family of maximal linklessly embeddable graphs on n vertices and $3n - 5$ edges for all $n \geq 10$, and another family on n vertices and $m < \frac{25n}{12} - \frac{1}{4}$ edges for all $n \geq 13$. The latter significantly improves the lowest edge-to-vertex ratio for any previously known infinite family. We construct a family of graphs showing that the class of maximal linklessly embeddable graphs differs from the class of graphs that are maximal without a K_6 minor studied by L. Jørgensen. We give necessary and sufficient conditions for when the clique sum of two maximal linklessly embeddable graphs over K_2 , K_3 , or K_4 is a maximal linklessly embeddable graph, and use these results to prove our constructions yield maximal linklessly embeddable graphs.

1. INTRODUCTION

All graphs in this paper are finite and simple. A graph is *intrinsically linked* (IL) if every embedding of it in \mathbb{R}^3 (or, equivalently, S^3) contains a nontrivial 2-component link. A graph is *linklessly embeddable* if it is not intrinsically linked (nIL). A nIL graph G is *maxnil* if it is not a proper subgraph of a nIL graph of the same order. The combined work of Conway and Gordon [2], Sachs [11] and Robertson, Seymour and Thomas [9] fully characterized IL graphs: a graph is IL if and only if it contains a graph in the Petersen family as a minor. The Petersen family consists of seven graphs obtained from K_6 by ∇Y -moves and $Y\nabla$ -moves, as described in Figure 1. The ∇Y -move and the $Y\nabla$ -move preserve the IL property.




FIGURE 1. ∇Y - and $Y\nabla$ -moves

The property of being maxnil is, in a way, analogous to the property of being maximal planar. While it is well known that every maximal planar graph with n vertices has $3n - 6$ edges, an analogous statement for maxnil graphs does not exist. For example, start with a maximal planar graph G and add one vertex v

together with all the edges from v to the vertices of G . Such a graph is maxnil by [11], and if it has n vertices, then it has $4n - 10$ edges. In fact, $4n - 10$ is an upper bound on the number of edges of a maxnil graph on n vertices. This follows from work of Mader [7] who proved that having more than $4n - 10$ edges implies the existence of a K_6 minor, which implies the graph is IL.

On the other hand, Jørgensen [5] and Dehkordi and Farr [3] constructed maxnil graphs with n vertices and $3n - 3$ edges. Jørgensen's maxnil graphs are obtained from the Jørgensen graph in Figure 2(a) by subdividing the highlighted edge incident to the vertex y and then adding edges that connect every new vertex to u and v . We denote the graph obtained this way through i subdivisions by J_i , $i \geq 1$. See Figure 2(b).




FIGURE 2. (a) The Jørgensen graph; (b) The graph J_i in Jørgensen's $3n - 3$ family.

Recently, Aires [1] found a family of graphs with fewer than $3n - 3$ edges. For each value $n \geq 13$ with $n \equiv 3 \pmod{10}$, he constructed a maxnil graph with $\frac{14n}{5} - \frac{27}{5}$ edges. He also proved that if G is a maxnil graph with $n \geq 5$ vertices and m edges, then $m \geq 2n$. This bound is sharp; the maxnil graph $Q(13, 3)$ described by Maharry [8] has 26 edges and 13 vertices.

In Section 2, we present two constructions of maxnil graphs. The first one is a family of maxnil graphs with $n \geq 10$ vertices and $3n - 5$ edges. This construction builds upon a maxnil graph on 10 vertices and 25 edges and uses edge subdivisions. The second construction significantly improves on Aires' result on the number of edges. Using clique sums of copies of $Q(13, 3)$, we construct examples with a smaller “edge-to-vertex ratio,” as in the following theorem.

Theorem. For each $n \geq 13$, there exists a maxnil graph G with n vertices and $m < \frac{25n}{12} - \frac{1}{4}$ edges.

In Section 3, we study the properties of maxnil graphs under clique sums. Some of these results are used in the constructions of Section 2. We give sufficient and necessary conditions for when the clique sum of two maxnil graphs over K_2 , K_3 or K_4 is maxnil. Jørgensen [5] studied clique sums of graphs that are maximal without a K_6 minor. We give examples showing that the class of maxnil graphs and the class of graphs that are maximal without a K_6 minor are distinct.

2. TWO FAMILIES OF MAXNIL GRAPHS

We note that the Jørgensen graph is 2-apex, i.e., removing the vertices u and v leaves a planar graph P . Furthermore, the embedding of P in \mathbb{R}^2 shown in Figure 2(a) has no separating cycles, i.e., for every cycle C in P , one of the components of $\mathbb{R}^2 \setminus C$ contains no vertices of P . These properties are generalized in the next Lemma, which we use to prove the graphs in the $3n - 5$ family are nIL.

Lemma 1. *Let G be a graph with two nonadjacent vertices u, v such that there exists an embedding Σ of $G - \{u, v\}$ in \mathbb{R}^2 where for every cycle C in Σ , $\mathbb{R}^2 \setminus C$ has a component X such that $X \cup C$ separates u and v (i.e., every path in G from u to v contains a vertex in $X \cup C$). Then embedding u as $(0, 0, 1)$ and v as $(0, 0, -1)$ and connecting each of them to its neighbors in Σ with straight edges yields a linkless embedding of G in \mathbb{R}^3 .*

Proof. Let Γ denote the embedding of G as described in the lemma, and let $K \cup K'$ be a 2-component link in Γ . We consider two cases.

Case 1. Neither K nor K' contains both u and v . Then we have three subcases: zero, one, or both of K and K' are in Σ . In each of these three subcases it is easy to see that $K \cup K'$ is a trivial link. We prove this for one of the three subcases here; the other two are similar and easier. Suppose K contains u but not v , and $K' \subset \Sigma$. Then K consists of two edges incident to u and a path $P \subset \Sigma$. Connecting u with straight line segments to every point in P gives us a Γ -panel for K . On the other hand, K' bounds a disk D in \mathbb{R}^2 . We isotop D , while keeping its boundary fixed, by pushing its interior slightly below \mathbb{R}^2 , to make it disjoint from K (since K contains no points below \mathbb{R}^2). It follows that $K \cup K'$ is a trivial link.

Case 2. One of the link's components, say K , contains both u and v . Then $K' \subset \Sigma$. So $\mathbb{R}^2 \setminus K'$ has two components such that one of them, X , separates u and v . Therefore all vertices of K except u and v lie in X . Now, K has exactly two vertices, call them a, b , that are adjacent to u , and two vertices, c, d , adjacent to v . Note that $\{a, b\}$ is not necessarily disjoint, or even distinct, from $\{c, d\}$. Furthermore, $K \cap X$ consists of two components, P_1 and P_2 , each of which is a path of length zero or greater. We can assume $a, c \in P_1$ and $b, d \in P_2$. We consider three subcases.




FIGURE 3. (a) Configuration for Case 2.1 ; (b) Configuration for Case 2.2

Case 2.1. $a = c, b = d$. Join a to b by an arc $\beta \subset X$ (not necessarily in Σ), and then connect each of u and v by straight line segments to every point in β . See Figure 3(a). This gives us a disk bounded by K and disjoint from K' . Similarly to Case 1 above, K' also bounds a disk disjoint from K . Hence $K \cup K'$ is a trivial link.

Case 2.2. $a = c, b \neq d$. Join a to each of b and d by disjoint arcs β and δ respectively, both in X , such that $\beta \cup \delta \cup P_2$ is a simple closed curve. See Figure 4(b). Connect each of u and v by straight line segments to every point in β and δ respectively. This gives us two disks whose union with the disk bounded by $\beta \cup \delta \cup P_2$ in X is a disk bounded by K and disjoint from K' . And, as before, K' bounds a disk disjoint from K . Hence $K \cup K'$ is a trivial link.

Case 2.3. $a \neq c, b \neq d$. This case is similar to Case 2.2, except that we join a to b and c to d by disjoint arcs β and δ in X such that $\beta \cup \delta \cup P_1 \cup P_2$ is a simple closed curve.

□

2.1. The $3n - 5$ family. We construct a family of graphs with n vertices and $3n - 5$ edges, for $n \geq 10$. This family is obtained from the graph G pictured in Figure 4(a) through a sequence of subdivisions and edge additions. The graph G is obtained from the Jørgensen graph by splitting (the opposite of contracting edges) the vertices a and b into the edges ad and bc . See Figures 2(a) and 4(a). With the notation in Figure 4(a), construct the graph G_1 by subdividing the edge xy with a new vertex z_1 , then adding edges z_1u and z_1v . Construct graphs G_i , for $i \geq 2$, as follows: subdivide the edge $z_{i-1}y$ of G_{i-1} with a new vertex z_i , then add edges z_iu and z_iv to G_{i-1} . Notice that G_i has one more vertex and three more

edges than G_{i-1} . The graph G_i has $10 + i$ vertices and $25 + 3i = 3(10 + i) - 5$ edges. We note that the graphs G_i can also be obtained by successive splittings of the vertex y into the edge yz_i .




FIGURE 4. (a) The graph G is maxnil with 10 vertices and 25 edges; (b) The graph G_i is obtained through i edge subdivisions and edge additions.

Proposition 2. *The graphs G and G_i in Figures 4(a) and 4(b) are linklessly embeddable.*

Proof. It is straightforward to check that these graphs satisfy the hypotheses of Lemma 1 and hence are nIL. \square

Proposition 3. *The graph G in Figure 4(a) is maxnil.*

Proof. Since G is linklessly embeddable, it remains to show that adding any edge to G gives an IL graph. We note that by contracting the edges ad and bc , we obtain the Jørgensen graph as a minor of G . If an edge e other than bd is added to $G - \{u, v\}$, such edge is preserved by these two edge contractions. Thus $G + e$ contains a minor that itself contains the Jørgensen graph added an edge. Since the Jørgensen graph is maxnil, $G + e$ is IL. Same holds if $e = uv$ is added to G . If the edge bd is added, then contracting the edges dt, cz, ux and vy creates a K_6 minor of $G + bd$. If an edge from u to $G - \{u, v\}$ is added, say ua , then contracting the edges cd, dt, by and uz creates a K_6 minor of $G + ua$. If an edge from v to $G - \{u, v\}$ is added, say vb , then contracting the edges ax, cz, du and dt creates a K_6 minor of $G + vb$. \square

Proposition 4. *All graphs G_i , $i \geq 1$ are maxnil.*

Proof. Since G_i is linklessly embeddable, it remains to show that adding any edge to G_i gives an IL graph. Adding any edge e different from xy and disjoint from $\{z_1, z_2, \dots, z_i\}$ to G_i gives a graph $G_i + e$ that contains $G + e$ as a minor (obtained by contracting the path $xz_1z_2\dots z_i$). Since G is maxnil, $G + e$ is IL and so is $G_i + e$. Adding an edge e that is either xy or has at least one endpoint in $\{z_1, z_2, \dots, z_i\}$ to G_i , gives a graph $G_i + e$ that contains $J_i + e$ as a minor (obtained by contracting the edges ad and bc). Since J_i is maxnil, $J_i + e$ is IL and so is $G_i + e$. \square

2.2. The $Q(13, 3)$ family. A graph G is called *triangular* if each edge of G belongs to at least a triangle. In a non-triangular graph, an edge that is not part of a triangle is a *non-triangular edge*. In Section 3, we study the properties of maxnil graphs under the operation of clique sum (defined in Section 3). For the construction presented in the next theorem we use the result of Lemma 10 about clique sums of maxnil graphs over K_2 .

Theorem 5. *For each $n \geq 13$, there exists a maxnil graph G with n vertices and $m < \frac{25n}{12} - \frac{1}{4}$ edges.*

Proof. The construction is based on the maxnil graph $Q_{13,3}$ described by Maharry [8]. See Figure 5(a). This graph has 13 vertices and 26 edges, and it is triangle free.

For each n with $13 \leq n \leq 39$, we construct a set of maxnil graphs with n vertices and $2n$ edges by adding $n - 13$ new vertices, and then choosing $n - 13$ edges in $Q_{13,3}$ and connecting the two endpoints of each of them to one of the new vertices. Equivalently, we are taking the clique sum of $Q_{13,3}$ with $n - 13$ disjoint triangles, over $n - 13$ copies of K_2 . See Figure 5(b). By Lemma 10, the resulting graph is maxnil.

The graph on 39 vertices obtained this way is triangular, so the construction cannot proceed further. To build graphs with a larger number of vertices we use multiple copies of $Q_{13,3}$ joined along an edge (clique sum over K_2). Consider $k \geq 1$ copies of $Q_{13,3}$ and choose one edge in each copy. Then join the k graphs together by identifying the k chosen edges into one edge. This graph, which we denote by H_k , is maxnil (by repeated application of Lemma 10) and has $11k + 2$ vertices and $25k + 1$ edges. All edges of H_k are non-triangular and adding vertices of degree 2 (as above) along any subset of the edges of H_k gives a maxnil graph.

For $n \geq 13$, let $k = \lceil \frac{n-3}{36} \rceil$ and add $n - (11k + 2)$ vertices of degree 2 along any $n - (11k + 2)$ edges of H_k . With every added vertex of degree 2, the number of edges is increased by 2. This gives a maxnil graph with n vertices and $m = (25k + 1) + 2[n - (11k + 2)] = 2n + 3k - 3$ edges. Moreover,

$$m = 2n + 3\lceil \frac{n-3}{36} \rceil - 3 < 2n + 3(\frac{n-3}{36} + 1) - 3 = \frac{25n}{12} - \frac{1}{4}.$$

\square




FIGURE 5. (a) $Q_{13,3}$ is a maxnil graph with 13 vertices and 26 edges; (b) A maxnil graph with 17 vertices and 34 edges obtained from $Q_{13,3}$ by adding four vertices of degree 2 and eight edges.

Remark 6. The above shows there exist maxnil graphs of arbitrarily large order n with an edge-to-vertex ratio of less than $\frac{25}{12} - \frac{1}{4n}$. Whether this edge-to-vertex ratio can be lowered further is an open question.

3. CLIQUE SUMS OF MAXNIL GRAPHS

In this section we study the properties of maxnil graphs under taking clique sums. A set $S \subset V(G)$ is a *vertex cut set* of a connected graph G if $G - S$ is disconnected. We say a vertex cut set $S \subset V(G)$ is *minimal* if no proper subset of S is a vertex cut set of G . A graph G is the *clique sum* of G_1 and G_2 over K_t if $V(G) = V(G_1) \cup V(G_2)$, $E(G) = E(G_1) \cup E(G_2)$, and the subgraphs induced by $V(G_1) \cap V(G_2)$ in both G_1 and G_2 are complete of order t . Since the vertices of the clique over which a clique sum is taken is a vertex cut set in the resulting graph, the vertex connectivity of a clique sum over K_t is at most t . For a set of vertices $\{v_1, v_2, \dots, v_k\} \subset V(G)$, $\langle v_1, v_2, \dots, v_k \rangle_G$ denotes the subgraph of G induced by this set of vertices. By abuse of notation, the subgraph induced in G by the union of the vertices of subgraphs H_1, H_2, \dots, H_k is denoted by $\langle H_1, H_2, \dots, H_k \rangle_G$.

Holst, Lovász, and Schrijver [4] studied the behavior of the Colin de Verdière μ -invariant for graphs under clique sums (Theorem 2.10). Since a graph G is nIL if and only if $\mu(G) \leq 4$ ([6], [10]), their theorem implies the following.

Theorem 7 (Holst, Lovász, and Schrijver [4]). *If G is the clique sum over S of two nIL graphs, then G is IL if and only if one can contract two or three components of $G - S$ so that the contracted nodes together with S form a K_7 minus a triangle.*

Theorem 7 implies that for $t \leq 3$, the clique sum over K_t of nIL graphs is nIL. While Theorem 7 shows when a clique sum is nIL, it does not establish when a clique sum of maxnil graphs is maxnil.

Lemma 8. *Any maxnil graph is 2-connected.*

Proof. Let G be a maxnil graph. If G is disconnected, let A and B denote two of its connected components. Let $a \in V(A)$ and $b \in V(B)$. Then $G + ab$ is a nIL graph, as it can be obtained by performing two consecutive clique sums over K_1 of nIL summands, namely

$$G + ab = A \cup_{\{a\}} ab \cup_{\{b\}} (G - A).$$

But this contradicts the maximality of G .

If the vertex connectivity of G is one, assume $x \in V(G)$ is a cut vertex, that is $G - x = A \sqcup B$, with A and B nonempty, and no edges between vertices of A and vertices of B . Let $a \in V(A)$ and $b \in V(B)$ be neighbors of x in G . Then $G + ab$ is nIL, as it can be obtained by performing two consecutive clique sums over K_2 of nIL summands. If Δ denotes the triangle axb ,

$$G + ab = \langle A, x \rangle_G \cup_{ax} \Delta \cup_{xb} \langle B, x \rangle_G.$$

But this contradicts the maximality of G . □

Lemma 9. *Let G be a maxnil graph with a vertex cut set $S = \{x, y\}$, and let G_1, G_2, \dots, G_r denote the connected components of $G - S$. Then $xy \in E(G)$ and $\langle G_i, S \rangle_G$ is maxnil for all $1 \leq i \leq r$.*

Proof. By Lemma 8, x and y are distinct and each of them has at least one neighbor in each G_i . Suppose $xy \notin E(G)$. Let $G' = G + xy$ and $G'_i = \langle G_i, S \rangle_{G'}$. Then, for every i , G'_i is a minor of G since if we pick a $j \neq i$ and in $\langle G_i, G_j, S \rangle_G$ contract G_j to x , we get a graph isomorphic to G'_i . So G'_i is nIL. Then, by Theorem 7, $G' = G'_1 \cup_{xy} \dots \cup_{xy} G'_r$ is nIL, contradicting the assumption that G is maxnil. So $xy \in E(G)$.

For each i , we repeatedly add new edges to $\langle G_i, S \rangle_G$, if necessary, to get a maxnil graph H_i . Then $H := H_1 \cup_{xy} \dots \cup_{xy} H_r$ is nIL and contains G as a subgraph, so $H = G$ and every $\langle G_i, S \rangle_G$ is maxnil. □

Lemma 10. *Let G_1 and G_2 be maxnil graphs. Pick an edge in each G_i and label it e . Then $G = G_1 \cup_e G_2$ is maxnil if and only if e is non-triangular in at least one G_i .*

Proof. The graph G is nIL by Theorem 7. Suppose e is non-triangular in at least one G_i , say G_2 . To prove G is maxnil, it is enough to show that for all $b_i \in V(G_i)$, $G + b_1 b_2$ is IL. Denote the endpoints of e in G by x, y . By Lemma 8, G_1 is 2-connected, so each of x, y has at least one neighbor in G_1 . So if we contract G_1 to b_1 and then contract $b_1 b_2$ to b_2 , we obtain a graph G'_2 that contains G_2 as a

proper subgraph since $b_2x \in E(G'_2)$. So G'_2 is IL since G_2 is maxnil. But G'_2 is a minor of G , which is nIL, so we have a contradiction.

To prove the converse, suppose e is triangular in G_1 and G_2 . Let $t_i \in V(G_i)$ be adjacent to both endpoints of e . Let K be a complete graph on four vertices, with vertices labeled x, y, t_1, t_2 . Denote the triangles induced by x, y, t_i in K and in G_i by Δ_i . Then by Theorem 7, $G' := G_1 \cup_{\Delta_1} K \cup_{\Delta_2} G_2$ is nIL. But G' is isomorphic to $G + t_1t_2$, so G is not maxnil. \square

Lemma 11. *Let G be a maxnil graph with vertex connectivity 3 and a vertex cut set $S = \{x, y, z\}$. Let G_1, G_2, \dots, G_r denote the connected components of $G - S$. Then $\langle S \rangle_G \simeq K_3$ and $\langle G_i, S \rangle_G$ is maxnil for all $1 \leq i \leq r$.*

Proof. Suppose $\langle S \rangle_G \not\simeq K_3$. Let G' be the graph obtained from G by adding one or more edges to $\langle S \rangle_G$ so that S induces a triangle T in G' . For $1 \leq i \leq r$, let $G'_i = \langle G_i, T \rangle_{G'}$. We see as follows that G'_i is nIL. Pick any $j \neq i$, and in the graph $\langle G_i, G_j, S \rangle_G$, contract G_j to an arbitrary vertex v in G_j . Then v is connected to each of x, y, z since G is 3-connected and hence each of x, y, z has at least one neighbor in G_j . The graph M_i obtained this way is minor of G , and hence is nIL. Performing a ∇Y -move on $T \subset G'_i$ we obtain a subgraph of M_i . Since M_i is nIL, so is G'_i . By Theorem 7, $G' = G'_1 \cup_T \dots \cup_T G'_r$ is nIL, which contradicts the maximality of G . So $T = \langle S \rangle_G \simeq K_3$.

To show $\langle G_i, S \rangle_G$ is maxnil, repeatedly add new edges to it, if necessary, to get a maxnil graph H_i . Then $H := H_1 \cup_T \dots \cup_T H_r$ is nIL by Theorem 7 and contains G as a subgraph, so $H = G$ and every $\langle G_i, S \rangle_G$ is maxnil. \square

Let G be a graph and let $T = \langle x, y, z, t \rangle_G$ be an induced K_4 subgraph (*tetrahedral graph*). We say T is *strongly separating* if $G - T$ has at least two connected components C_1, C_2 such that every vertex of T has a neighbor in each C_i .

Lemma 12. *Let G_1, G_2 be maxnil graphs and let $G = G_1 \cup_{\Delta} G_2$ be the clique sum of G_1 and G_2 over a K_3 subgraph $\Delta = \langle x, y, z \rangle_G$. Assume Δ is a minimal vertex cut set in G . Then G is maxnil if and only if for some $i \in \{1, 2\}$, every induced K_4 subgraph of the form $\langle x, y, z, t \rangle_{G_i}$ is strongly separating.*

Proof. By Theorem 7, $G := G_1 \cup_{\Delta} G_2$ is nIL. Then G is maxnil if and only if for every $t_1 \in V(G_1) - V(\Delta)$ and $t_2 \in V(G_2) - V(\Delta)$, $G' := G + t_1t_2$ is IL.

First, suppose for some i at least one of x, y, z is not connected to t_i , say $xt_2 \notin E(G_2)$. Contracting $G_1 - \{y, z\}$ to x produces $G_2 + t_2x$ as a minor of G' . Since G_2 is maxnil, this minor is IL, and hence G' is IL, as desired. So we can assume $\langle x, y, z, t_i \rangle_{G_i}$ is a tetrahedral graph for both $i = 1, 2$.

Assume every tetrahedral graph in G_2 that contains Δ is strongly separating. So $G_2 - \langle x, y, z, t_2 \rangle_{G_2}$ has at least two connected components each of which, when contracted to a single vertex, is adjacent to all four vertices x, y, z, t_2 . In Figure 6

these vertices are denoted by c_1 and c_2 . Now, if the component of $G_1 - \Delta$ that contains t_1 is contracted to t_1 , this vertex too will be adjacent to x, y, z, t_2 . So we get a minor of G isomorphic to K_7 minus a triangle, which is IL since it contains a Petersen family graph (the one obtained by one ∇Y -move on K_6) as a minor. It follows that G' is IL, and therefore G is maxnil.




FIGURE 6. A K_7 minus a triangle minor of the graph G .

To prove the converse, for $i = 1, 2$ let t_i be a vertex in G_i such that $T_i := \langle x, y, z, t_i \rangle_{G_i}$ is a tetrahedral graph that is not strongly separating. Let $G' = G + t_1 t_2$. Then $G' = G_1 \cup_{T_1} \langle x, y, z, t_1, t_2 \rangle_{G'} \cup_{T_2} G_2$. Each of these click sums is over a K_4 , each summand is nIL, and each of T_1, T_2 is non-strongly separating; so, by Theorem 7, G' is nIL, and hence G is not maxnil. \square

Unlike the vertex connectivity 2 and 3 cases, it is not true that a minimal vertex cut set in a 4-connected maxnil graph must be a clique. The four neighbors of b in the graph depicted in Figure 4(a) form a vertex cut set, but the graph induced by its vertices has exactly 2 edges. The four neighbors of any vertex in the graph $Q_{13,3}$ in Figure 5(a) form a discrete vertex cut set. However, if a maxnil graph G has vertex connectivity 4, the following lemma provides some restrictions on the shape of the subgraph induced by the vertices of any minimal vertex cut set.

Lemma 13. *Let G be a maxnil graph and assume $\{x, y, z, t\}$ is a minimal vertex cut. Let $S = \langle x, y, z, t \rangle_G$. Then S is either a clique or a subgraph of C_4 (a 4-cycle).*

Proof. Assume that S is neither a clique nor a subgraph of a 4-cycle. This implies that if S has a vertex of degree at least 3, then it contains $K_{1,3}$ as a subgraph; and if every vertex of S has degree less than 3, then S contains K_3 as a subgraph. Below, we consider these two cases separately.

Case 1. S has a K_3 subgraph. We can assume that x, y, z induce a triangle in G . If $G - S$ has at least three connected components, contracting each of them

to a single node would produce a minor of G isomorphic to K_7 minus a triangle, contradicting that G is nIL. So $G - S = G_1 \sqcup G_2$, with G_1 and G_2 each connected.

Since $\{x, y, z, t\}$ is a minimal vertex cut set in G , each of x, y, z, t has at least one neighbor in each G_i , $i = 1, 2$. Contracting $\langle G_i, t \rangle_G$ to t produces a minor of G , denoted by G'_i , which must be nIL. Since xyz is a triangle and each of x, y, z has at least one neighbor in each G_i , $\{x, y, z, t\}$ induces a clique T in both G'_1 and G'_2 . By [4], the clique sum $G' = G'_1 \cup_T G'_2$ is nIL since $G' - T = G_1 \sqcup G_2$; but G' strictly contains G as a subgraph, a contradiction.

Case 2. S has a $K_{1,3}$ subgraph. We can assume that t is adjacent to x, y and z in G . If $G - S$ has at least three connected components, contracting each of them to a single node would produce a minor of G containing a subgraph isomorphic to $K_{3,3,1}$, thus G is IL. So $G - S = G_1 \sqcup G_2$, with G_1 and G_2 connected. For $i = 1, 2$, contracting each of G_i to a single node t_i , deleting the edge $t_i t$, deleting any existing edges of $\langle x, y, z \rangle_G$, and then performing a $Y\nabla$ -move at t_i , produces a nIL graph, denoted by G'_i . Let $G' = G'_1 \cup_{K_4} G'_2$ be the clique sum over the complete graph with vertices x, y, z, t . By Theorem 7, G' is nIL since $G' - S = G_1 \sqcup G_2$; but G' strictly contains G as a subgraph, a contradiction. \square


Lemma 14. *Let $G = G_1 \cup_S G_2$ be the clique sum of maxnil graphs G_1 and G_2 over $S = \langle x, y, z, t \rangle_G \simeq K_4$. Assume S is a minimal vertex cut set in G . Then G is maxnil if and only if, in both G_1 and G_2 , S is not strongly separating.*

Proof. If S is strongly separating in G_1 or G_2 , then $G - S$ has at least three connected components and contracting each of them to a single node produces a minor isomorphic to K_7 minus a triangle.


If, in both G_1 and G_2 , S is not strongly separating, then $G - S$ has only two connected components. Contracting each of the two components to a single node produces K_6 minus an edge as a minor (not K_7 minus a triangle); hence G is nIL by Theorem 7. Adding an edge between a vertex in $G_1 - S$ and a vertex in $G_2 - S$ and contracting $G_1 - S$ and $G_2 - S$ to single nodes produces a K_6 minor. It follows that G is maxnil in this case. \square

The graph G of Figure 7 is maxnil since $G - u$ is a maximal planar graph. If $S = \langle x, y, z, t, u \rangle$, $G_1 = \langle a, x, y, z, t, u \rangle$, and $G_2 = \langle b, x, y, z, t, u \rangle$, then $S \simeq K_5$, $G_1 \simeq G_2 \simeq K_6^-$ (K_6 minus one edge), and $G = G_1 \cup_S G_2$. This shows it is possible for the clique sum of two maxnil graphs over $S \simeq K_5$ to be nIL (and maxnil). However, no clique S of order 5 can be a minimal vertex cut set in a nIL graph G , since then any connected component of $G - S$ would form a K_6 -minor together with S , which would imply G is IL. For $t \geq 6$, any clique sum over K_t is IL since K_6 is IL.

Jørgensen studied clique sums of graphs that are maximal without a K_6 minor [5]. These are graphs that do not contain a K_6 minor and a K_6 minor is created

FIGURE 7. A maxnil graph that is a clique sum over K_5 .

by the addition of any edge. The class of maxnil graphs and the class of graphs that are maximal without a K_6 minor are not the same, as shown in the following proposition.

FIGURE 8. A maxnil graph G (left) that is not maximal without a K_6 minor is obtained by adding two vertices to a plane triangulation with nine vertices (right)

Proposition 15. *The graph in Figure 8 is maxnil, and it is not maximal without a K_6 minor.*

Proof. The graph G in Figure 8 is obtained by adding vertices v and w to the plane triangulation H : vertex v connects to all nine vertices of H and vertex w connects to vertices a , b and c of H . The graph $H + v$ is maxnil, since it is a cone over a maximal planar graph [11]. The graph G is the clique sum over $K_3 = \langle a, b, c \rangle_G$ of maxnil graphs $H + v$ and $K_4 = \langle a, b, c, w \rangle_G$. The graph

$\langle a, b, c, v \rangle_{H+v}$ is the only induced K_4 subgraph in $H + v$ containing a, b and c and it is strongly separating in $H + v$. So, by Lemma 12, G is maxnil; in particular it has no K_6 minor. The graph $G + vw$ is a clique sum over $K_4 = \langle a, b, c, v \rangle_G$ of graphs $H + v$ and $K_5 = \langle a, b, c, v, w \rangle$, both of which are K_6 minor free. Hence, by [5], $G + vw$ is K_6 minor free, so G is not maximal without a K_6 minor. \square

Remark 16. Starting with the graph G in Proposition 15, one can construct graphs G_n with $n \geq 11$ vertices that are maxnil but not maximal without a K_6 minor. Take $G_{11} = G$ and construct G_{11+k} from G by triangulating the disk bounded by the triangle efg with k new vertices and adding edges between v and these new vertices. The argument used in the proof of Proposition 15 shows that G_n , $n \geq 11$, is maxnil but not maximal without a K_6 minor. We conjecture $n = 11$ is the minimal order of a graph with this property, i.e., every maxnil graph with $n \leq 10$ vertices is maximal without a K_6 minor.

REFERENCES

1. M. Aires. *On the number of edges in maximally linkless graphs*, preprint, arXiv:1911.08552v1
2. J. Conway and C. Gordon. *Knots and links in spatial graphs*. J. Graph Theory **7** (1983), No. 4, 445–453
3. H. R. Dehkordi and G. Farr. *Non-separating planar graphs*, preprint, arXiv:1907.09817
4. H. van der Holst, L. Lovász and A. Schrijver. *The Colin de Verdière graph parameter*, Graph Theory and Combinatorial Biology (Balatonlelle, 1996), Bolyai Soc. Math. Stud. **7**, Budapest: János Bolyai Math. Soc., pp. 29–85
5. L. K. Jørgensen. *Some maximal graphs that are not contractible to K_6* , Aalborg Universitetscenter. Institut for Elektroniske Systemer - Rapport, Nr. 1989 : R 89-28 (1989)
6. L. Lovász and A. Schrijver. *A Borsuk theorem for antipodal links and a spectral characterization of linklessly embeddable graphs*, Proc. Amer. Math. Soc. **126**, Number 2 (1998), 1275–1285
7. W. Mader, *Homomorphiesätze für Graphen*, Mathematische Annalen, Vol. 178, Iss. 2, (1968), 154–168
8. J. Maharry. *A splitter for graphs with no Petersen family minor*, Journal of Combinatorial Theory, Series **B** 72. 136–139 (1998)
9. N. Robertson, P. Seymour and R. Thomas. *Linkless embeddings of graphs in 3-space*. Bulletin of the Amer. Math. Soc. **28** (1993), No. 1, 84–89
10. N. Robertson, P. Seymour and R. Thomas. *A survey of linkless embeddings*. Graph Structure Theory (N. Robertson, P. Seymour, eds.), Contemporary Mathematics, American Mathematical Society, Providence, Rhode Island, 1993, pp. 125–136
11. H. Sachs. *On spatial representations of finite graphs* A. Hajnal, L. Lovasz, V.T. Sós (Eds.), Colloq. Math. Soc. János Bolyai, Vol. **37**, North-Holland, Amsterdam, 1984, 649–662