

Secondary component of gravitational-wave signal GW190814 as an anisotropic neutron star

Zacharias Roupas¹

¹*Centre for Theoretical Physics, The British University in Egypt, Sherouk City
11837, Cairo, Egypt*

Abstract

The gravitational-wave signal GW190814 involves a compact object with mass $(2.50 - 2.67)M_\odot$ within the so-called low mass gap. As yet, a general consensus on its nature, being a black hole, a neutron star or an exotic star, has not been achieved. We investigate the possibility this compact object to be an anisotropic neutron star. Anisotropies in a neutron star core arise naturally by effects such as superfluidity, hyperons, strong magnetic fields and allow the maximum mass to exceed that of the ideally isotropic stars. We consider the Krori-Barua ansatz to model an anisotropic core and constrain the equation of state with LIGO/Virgo observations GW170817 and GW190814. We find that the GW190814 secondary component can be an anisotropic neutron star compatible with LIGO/Virgo constraints if the radius attains a value in the range $(13.2 - 14.0)$ km with the anisotropic core's boundary density in the range $(3.5 - 4.0) \cdot 10^{14} \text{g/cm}^3$.

1 Introduction

The LIGO/Virgo Collaboration announced recently the observation of a merger of a black hole with mass $23.2^{+1.1}_{-1.0}M_\odot$ with a compact object with mass $2.59^{+0.08}_{-0.09}M_\odot$ [1]. The mass of the secondary component lies within the so-called low mass gap [2–4]. Theoretical and observational evidence suggest that black holes of mass less than $\sim 5M_\odot$ may not be produced by stellar evolution [2, 3, 5]. On the other hand while some candidate equations of state of neutron stars allow for a neutron star maximum mass $M_{\text{max}} \sim 3M_\odot$ [6–10], the relatively small tidal deformabilities measured in gravitational-wave signal GW170817 do not favor such large values of M_{max} but rather suggest it is of the order of $2.5M_\odot$ [1, 11, 12]. The heaviest neutron star observed to date has a mass of $2.01 \pm 0.04M_\odot$ [13], while there was a recent claim that PSR J0740+6620 may host a $2.14^{+0.10}_{-0.09}M_\odot$ neutron star [14]. Therefore the existence and nature of compact objects in the mass regime $\sim [2.5, 5]M_\odot$ are highly uncertain.

These theoretical and observational uncertainties regarding the maximum mass of neutron stars and lower mass of black holes render it challenging to conclude with certainty on the nature of the secondary GW190814 component. Several analyses seem to favor a stellar black hole [1, 15–17]. Other proposals include a primordial black hole [18–21], a fast pulsar [22–24], a heavy neutron star with stiff equation of state [25], a neutron star with accretion disk [26].

Independent of the LIGO/Virgo results, neutron star properties have been reported recently by the Neutron Star Interior Composition Explorer (NICER) team. In particular, observations of the isolated millisecond pulsar PSR J0030+0451 indicate a stiffer equation of state [27, 28] than those mostly favored by the LIGO/Virgo collaboration [1, 11].

Besides a stiff equation of state, anisotropies in the core of a neutron star [29] may also allow higher maximum neutron star masses [30]. The anisotropies inside the star can grow due to superfluidity [31–33], solidification [34–37], hyperons [38], quarks [39] as well as pion and kaon condensates [40, 41]. In addition, nuclear matter in a magnetic field becomes anisotropic, with different pressures in directions along and transverse to the field [42, 43]. The electromagnetic energy-momentum tensor is naturally anisotropic. Thus, an anisotropic core is more realistic than an ideally isotropic one.

An anisotropic compact object is subject to a tangential pressure $p_t = p_\theta = p_\phi$ in the angular directions that is different than the radial pressure $p_r \neq p_t$ [44, 45]. If the anisotropy parameter is positive $\Delta \equiv p_t - p_r > 0$ an additional repulsive anisotropic force enables more compact stable configurations to appear in the anisotropic than in the isotropic case [46]. The maximum mass of

anisotropic compact neutron stars has been estimated to be $M_{\text{max}} \sim 4M_{\odot}$ [30]. Several anisotropic solutions include Refs [47–58].

Here, we will investigate the possibility that the GW190814 secondary component is an anisotropic neutron star. We will work in a metric ansatz introduced by Krori & Barua (KB) [59]. Anisotropic compact star models in the KB-spacetime have also been studied in Refs. [60–77]. With input the total mass we will calculate the boundary density, radius, and equation of state compatible with LIGO's constraints imposed by gravitational-wave signals GW170817 and GW190814 [1].

In the next section 2 we impose the LIGO constraints of the neutron star equation of state to an anisotropic neutron star with mass $2.6M_{\odot}$, that is equal to that of the secondary GW190814 component. In the final section we discuss our conclusions.

2 Anisotropic star subject to LIGO constraints

Let us first review briefly the KB-ansatz. We write the spherically symmetric metric in General Relativity in spherical coordinates (t, r, θ, ϕ) as

$$ds^2 = -e^{\alpha(r)} c^2 dt^2 + e^{\beta(r)} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (1)$$

for some functions of r , $\alpha(r)$ and $\beta(r)$. If we denote ρ , p_r , p_t , the mass density, the radial pressure and the tangential pressure, respectively, the spherically symmetric, anisotropic energy momentum tensor is

$$T_{\nu}^{\mu} = \left(\frac{p_t}{c^2} + \rho\right) u^{\mu} u_{\nu} + p_t \delta_a^{\mu} + (p_r - p_t) \xi^{\mu} \xi_{\nu}, \quad (2)$$

where u^{μ} is the four-velocity and ξ^{μ} the unit space-like vector in the radial direction.

Krori & Barua [59] introduced the following ansatz for the metric potentials

$$\alpha(x) = a_0 x^2 + a_1, \quad \beta(x) = a_2 x^2, \quad (3)$$

where however we use the dimensionless variable

$$x \equiv \frac{r}{R} \in [0, 1]. \quad (4)$$

We will use the KB-ansatz to model the core of an anisotropic neutron star. We denote the radius of the core as $r = R$. Using the characteristic density

$$\rho_{\star} \equiv \frac{c^2}{8\pi G R^2} \quad (5)$$

we get the dimensionless variables

$$\tilde{\rho} = \frac{\rho}{\rho_{\star}}, \quad \tilde{p}_r = \frac{p_r}{\rho_{\star} c^2}, \quad \tilde{p}_t = \frac{p_t}{\rho_{\star} c^2}. \quad (6)$$

Einstein equations give for the dimensionless variables

$$\tilde{\rho} = \frac{e^{-a_2 x^2}}{x^2} \left(e^{a_2 x^2} - 1 + 2a_2 x^2 \right), \quad (7)$$

$$\tilde{p}_r = \frac{e^{-a_2 x^2}}{x^2} \left(1 - e^{a_2 x^2} + 2a_0 x^2 \right), \quad (8)$$

$$\tilde{p}_t = e^{-a_2 x^2} \left(2a_0 - a_2 + a_0(a_0 - a_2)x^2 \right). \quad (9)$$

Integrating $\mathcal{M}' = 4\pi\rho r^2$ we get for the mass $\mathcal{M}(r)$ contained within r that

$$\mathcal{M}(x) = M C^{-1} x \left(1 - e^{-a_2 x^2} \right). \quad (10)$$

where M denotes the total mass of the core and C is the compactness

$$C = \frac{2GM}{Rc^2}. \quad (11)$$




Figure 1: The radial pressure with respect to the density for the two marginal anisotropic cores with $M = 2.6M_\odot$ and $R = 13.2$ (dash-dotted line), $R = 14.0$ km (continuous line) compatible with LIGO's constraints (green shaded region) from GW170817, GW190814.

We match the KB-solution with the Tolman-Oppenheimer-Volkoff metric [78] at the boundary of the anisotropic core

$$ds_{\text{TOV}}^2 = -e^{\alpha_{\text{TOV}}(r)} c^2 dt^2 + \left(1 - \frac{2GM(r)}{rc^2}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (12)$$

$$\alpha_{\text{TOV}}(r) = \frac{2}{c^2} \int_r^\infty d\xi \left(\frac{GM(\xi)}{\xi^2} + \frac{4\pi G}{c^2} P(\xi) \xi \right) \left(1 - \frac{2GM(\xi)}{\xi c^2}\right)^{-1}, \quad r \geq R, \quad (13)$$

that is assumed to describe the outer layers with a different equation of state $P = P(\rho)$. The boundary pressure $p_R \equiv p_r(R) = P(R)$ is a free parameter. We have

$$\beta(r=R) = \ln(1-C)^{-1}, \quad \tilde{p}_r(r=R) = \tilde{p}_R. \quad (14)$$

For our purposes we only need a_0 , a_2 as is evidenced from Eqs. (7)-(9). By use of Eqs. (3), (8), (14) we get

$$a_0(C) = \frac{1}{2} \frac{C + \tilde{p}_R}{1-C}, \quad a_2(C) = \ln(1-C)^{-1}. \quad (15)$$

These equations along with (7)-(9) allow us parametrize density and pressure solely with respect to compactness, $\tilde{\rho} = \tilde{\rho}(x; C)$, $\tilde{p} = \tilde{p}(x; C)$, as was firstly shown in [77]. Stars with the same compactness acquire the same core profile and properties. The parameter a_1 , not required in our analysis, can be determined from the boundary condition $\alpha_{\text{TOV}}(r=R) = a_0 + a_1$.

The boundary density ρ_R and boundary radial pressure p_R are free parameters which we constrain by LIGO's observations. These are summarized in Figure 8 of [1] and reproduced here as the green shaded region of Figure 1. We find that an anisotropic neutron core with mass equal to the secondary GW190814 component $M = 2.6M_\odot$ satisfies the constraints on the equation of state compatible with signals GW170817 and GW190814 given by LIGO only for radius with values in the range (13.2 – 14.0) km, given in detail in Table 1. The boundary density is $(3.5 - 4.0)10^{14} \text{ g/cm}^3$ with the lower value corresponding to the bigger radius. In Figure 1 we depict that the two marginal solutions $R = 13.2$ km and $R = 14.0$ km satisfy LIGO's constraints of the equation of state. It seems plausible that the bigger star $R = 14.0$ km is a more probable candidate, because of the more realistic core's boundary density $\rho_R = 3.5 \cdot 10^{14} \text{ g/cm}^3$.

The acceptable radii correspond to compactness values $C = (0.55 - 0.58)$. These are not only realistic for a neutron star, but also are lower than $C < 0.71$. This is the stability and physical requirement condition for an anisotropic star in KB-spacetime calculated by Roupas & Nashed [77]. Note that this limit was calculated assuming zero boundary pressure. Nevertheless, it is straightforward to verify that the solutions of Table 1 satisfy the stricter condition that gave the limit 0.71, that is the strong energy condition [79, 80] $\rho c^2 - p_r - 2p_t > 0$. It is also straightforward to verify that there are satisfied all other conditions of Roupas & Nashed [77] such as the stability condition for the adiabatic index [81, 82] $\Gamma \equiv \frac{\rho + p_r}{p_r} \frac{dp_r}{d\rho} > \frac{4}{3}$, the causality conditions $v_r < c$, $v_t < c$ and also the condition of stability against cracking [83, 84] $0 < v_r^2 - v_t^2 < c^2$.

$R[\text{km}]$	$\rho_R[10^{14}\text{g/cm}^3]$	$p_R/c^2[10^{14}\text{g/cm}^3]$	$\frac{dp_r}{d\rho}[c^2]$	$\frac{dp_t}{d\rho}[c^2]$
13.2	4.03	(0.24 – 0.26)	0.49	0.34
13.3	3.95	(0.18 – 0.23)	(0.46 – 0.48)	(0.32 – 0.34)
13.4	3.88	(0.14 – 0.21)	(0.45 – 0.47)	(0.31 – 0.33)
13.5	3.81	(0.13 – 0.18)	(0.44 – 0.46)	(0.30 – 0.32)
13.6	3.73	(0.12 – 0.16)	(0.44 – 0.45)	(0.30 – 0.31)
13.7	3.66	(0.11 – 0.14)	(0.43 – 0.44)	0.30
13.8	3.59	(0.10 – 0.12)	(0.42 – 0.43)	(0.29 – 0.30)
13.9	3.53	(0.09 – 0.10)	0.42	0.29
14.0	3.46	0.08	0.41	0.28

Table 1: The radius R , boundary density ρ_R , boundary radial pressure p_R and slopes of $p_r(\rho)$, $p_t(\rho)$ linear fits, compatible with LIGO’s constraints for an anisotropic core with mass $2.6M_\odot$.

3 Conclusions

We have used the constraints on a neutron star’s core equation of state given by LIGO as derived from the gravitational-wave signals GW170817, GW190814 [1] in order to investigate if an anisotropic neutron core is compatible with the secondary GW190817 component. We found intriguingly that not only it is compatible, but also that the parameters imposed by these constraints give physical, stable solutions. There is no a-priori reason why this would be the case, especially since LIGO’s analysis [1] favours a black hole as a candidate for the secondary GW190814 component.

In particular, we conclude that an anisotropic neutron star core with $M = 2.6M_\odot$ in the KB-ansatz is compatible with LIGO constraints for a radius $R = (13.2 - 14.0)$ km. The corresponding boundary density for the $R = 14.0$ km solution is $\rho_R = 3.5 \cdot 10^{14}\text{g/cm}^3$ that is very close to the nuclear saturation density. For this solution the linear fit of the equation of state gives $p_r \propto 0.41\rho c^2$ and $p_t \propto 0.28\rho c^2$.

References

- [1] R. Abbott, et al. GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object. *ApJL*, 896(2):L44, June 2020.
- [2] Charles D. Bailyn, et al. The mass distribution of stellar black holes. *The Astrophysical Journal*, 499(1):367374, May 1998.
- [3] Feryal zel, et al. The black hole mass distribution in the galaxy. *The Astrophysical Journal*, 725(2):19181927, Dec 2010.
- [4] Krzysztof Belczynski, et al. Missing black holes unveil the supernova explosion mechanism. *The Astrophysical Journal*, 757(1):91, Sep 2012.
- [5] Will M. Farr, et al. The mass distribution of stellar-mass black holes. *The Astrophysical Journal*, 741(2):103, Oct 2011.
- [6] Horst Müller and Brian D. Serot. Relativistic mean-field theory and the high-density nuclear equation of state. *Nuc. Phys. A*, 606:508–537, February 1996.
- [7] Clifford E. Rhoades and Remo Ruffini. Maximum mass of a neutron star. *Phys. Rev. Lett.*, 32:324–327, Feb 1974.
- [8] Feryal zel, et al. On the mass distribution and birth masses of neutron stars. *The Astrophysical Journal*, 757(1):55, Sep 2012.
- [9] Blent Kiziltan, et al. The neutron star mass distribution. *The Astrophysical Journal*, 778(1):66, Nov 2013.

- [10] Justin Alsing, Hector O Silva, and Emanuele Berti. Evidence for a maximum mass cut-off in the neutron star mass distribution and constraints on the equation of state. *Monthly Notices of the Royal Astronomical Society*, 478(1):13771391, Apr 2018.
- [11] B. P. Abbott, et al. Gw170817: Measurements of neutron star radii and equation of state. *Phys. Rev. Lett.*, 121:161101, Oct 2018.
- [12] B. P. Abbott, et al. Properties of the Binary Neutron Star Merger GW170817. *Physical Review X*, 9(1):011001, January 2019.
- [13] J. Antoniadis, et al. A massive pulsar in a compact relativistic binary. *Science*, 340(6131):12332321233232, Apr 2013.
- [14] H. T. Cromartie, et al. Relativistic shapiro delay measurements of an extremely massive millisecond pulsar. *Nature Astronomy*, 4(1):7276, Sep 2019.
- [15] F. J. Fattoyev, et al. GW190814: Impact of a 2.6 solar mass neutron star on nucleonic equations of state. *arXiv e-prints*, arXiv:2007.03799, July 2020.
- [16] Ingo Tews, et al. On the nature of GW190814 and its impact on the understanding of supranuclear matter. *arXiv e-prints*, arXiv:2007.06057, July 2020.
- [17] Armen Sedrakian, Fridolin Weber, and Jia-Jie LI. Confronting GW190814 with hyperonization in dense matter and hypernuclear compact stars. *arXiv e-prints*, arXiv:2007.09683, July 2020.
- [18] Kyriakos Vattis, Isabelle S. Goldstein, and Savvas M. Koushiappas. Could the 2.6 M_{\odot} object in GW190814 be a primordial black hole? *arXiv e-prints*, arXiv:2006.15675, June 2020.
- [19] Benjamin V. Lehmann, Stefano Profumo, and Jackson Yant. Model-independent discovery prospects for primordial black holes at ligo, 2020.
- [20] Sebastien Clesse and Juan Garcia-Bellido. GW190425 and GW190814: Two candidate mergers of primordial black holes from the QCD epoch. *arXiv e-prints*, arXiv:2007.06481, July 2020.
- [21] Karsten Jedamzik. Evidence for primordial black hole dark matter from LIGO/Virgo merger rates. *arXiv e-prints*, arXiv:2007.03565, July 2020.
- [22] Nai-Bo Zhang and Bao-An Li. GW190814’s secondary component with mass $(2.50 - 2.67) M_{\odot}$ as a super-fast pulsar. *arXiv e-prints*, arXiv:2007.02513, July 2020.
- [23] Elias R. Most, et al. A lower bound on the maximum mass if the secondary in GW190814 was once a rapidly spinning neutron star. *arXiv e-prints*, arXiv:2006.14601, June 2020.
- [24] Antonios Tsokaros, Milton Ruiz, and Stuart L. Shapiro. GW190814: Spin and equation of state of a neutron star companion. *arXiv e-prints*, arXiv:2007.05526, July 2020.
- [25] Hung Tan, Jacquelyn Noronha-Hostler, and Nico Yunes. Kinky neutron stars in light of GW190814. *arXiv e-prints*, arXiv:2006.16296, June 2020.
- [26] Mohammadtaher Safarzadeh and Abraham Loeb. Formation of mass-gap objects in highly asymmetric mergers. *arXiv e-prints*, arXiv:2007.00847, July 2020.
- [27] M. C. Miller, et al. PSR J0030+0451 Mass and Radius from NICER Data and Implications for the Properties of Neutron Star Matter. *ApJL*, 887(1):L24, December 2019.
- [28] T. E. Riley, et al. A NICER View of PSR J0030+0451: Millisecond Pulsar Parameter Estimation. *ApJL*, 887(1):L21, December 2019.
- [29] P. Haensel, A. Y. Potekhin, and D. G. Yakovlev. *Neutron Stars 1 : Equation of State and Structure*, volume 326. 2007.
- [30] H. Heintzmann and W. Hillebrandt. Neutron stars with an anisotropic equation of state: mass, redshift and stability. *Astron. Astrophys.*, 38(1):51–55, January 1975.
- [31] Gordon Baym, Christopher Pethick, and David Pines. Superfluidity in Neutron Stars. *Nature*, 224(5220):673–674, November 1969.

- [32] M. Hoffberg, et al. Anisotropic Superfluidity in Neutron Star Matter. *Phys. Rev. Lett.*, 24(14):775–777, April 1970.
- [33] Armen Sedrakian and John W. Clark. Superfluidity in nuclear systems and neutron stars. *European Physical Journal A*, 55(9):167, September 2019.
- [34] P. W. Anderson and R. G. Palmer. Solidification Pressure of Nuclear and Neutron Star Matter. *Nature Physical Science*, 231(24):145–146, June 1971.
- [35] J. W. Clark and N. C. Chao. The Crystallization of Neutronic Matter. *Nature Physical Science*, 236(64):37–39, March 1972.
- [36] V. Canuto and S. M. Chitre. Solidification of Neutron Matter. *Phys. Rev. Lett.*, 30(20):999–1002, May 1973.
- [37] Daniel Schiff. Crystallization Density of Cold Dense Neutron Matter. *Nature Physical Science*, 243(130):130–133, June 1973.
- [38] Shmuel Balberg and Nir Barnea. S-wave pairing of Λ hyperons in dense matter. *Phys. Rev. C*, 57(1):409–416, January 1998.
- [39] D. Bailin and A. Love. Superfluidity and superconductivity in relativistic fermion systems. *Phys. Rep.*, 107(6):325–385, May 1984.
- [40] R. F. Sawyer. Condensed π^- Phase in Neutron-Star Matter. *Phys. Rev. Lett.*, 29(6):382–385, August 1972.
- [41] T. Takatsuka and R. Tamagaki. Nucleon Superfluidity in Kaon-Condensed Neutron Stars. *Progress of Theoretical Physics*, 94(3):457–461, September 1995.
- [42] Efrain J. Ferrer, et al. Equation of state of a dense and magnetized fermion system. *Phys. Rev. C*, 82(6):065802, December 2010.
- [43] Efrain J. Ferrer and Aric Hackebill. Equation of State of a Magnetized Dense Neutron System. *Universe*, 5(5):104, May 2019.
- [44] H. Bondi. Massive Spheres in General Relativity. *Proceedings of the Royal Society of London Series A*, 282(1390):303–317, November 1964.
- [45] Richard L. Bowers and E. P. T. Liang. Anisotropic Spheres in General Relativity. *Astrophys. J.*, 188:657–665, 1974.
- [46] B. V. Ivanov. Maximum bounds on the surface redshift of anisotropic stars. *Phys. Rev. D*, 65(10):104011, May 2002.
- [47] Selçuk Ş. Bayin. Anisotropic fluid spheres in general relativity. *Phys. Rev. D*, 26:1262–1274, Sep 1982.
- [48] L. Herrera, J. Ospino, and A. Di Prisco. All static spherically symmetric anisotropic solutions of einstein’s equations. *Phys. Rev. D*, 77:027502, Jan 2008.
- [49] S Thirukkanesh and S D Maharaj. Charged anisotropic matter with a linear equation of state. *Classical and Quantum Gravity*, 25(23):235001, Nov 2008.
- [50] Ksh. Newton Singh, Piyali Bhar, and Neeraj Pant. Solutions of the Einstein’s field equations with anisotropic pressure compatible with cold star model. *Astrophys. Space Sci.*, 361(10):339, October 2016.
- [51] S. K. Maurya, et al. Generalised model for anisotropic compact stars. *The European Physical Journal C*, 76(12), Dec 2016.
- [52] S. K. Maurya and M. Govender. Generating physically realizable stellar structures via embedding. *The European Physical Journal C*, 77(5), May 2017.
- [53] Gabino Estevez-Delgado and Joaquin Estevez-Delgado. On the effect of anisotropy on stellar models. *European Physical Journal C*, 78(8):673, August 2018.

- [54] Francisco Tello-Ortiz, et al. Anisotropic relativistic fluid spheres: an embedding class I approach. *European Physical Journal C*, 79(11):885, October 2019.
- [55] Shyam Das, Farook Rahaman, and Lipi Baskey. A new class of compact stellar model compatible with observational data. *European Physical Journal C*, 79(10):853, October 2019.
- [56] Joaquin Estevez-Delgado, et al. A possible representation for the neutron star PSR J0437-4715. *European Physical Journal Plus*, 134(12):600, December 2019.
- [57] Piyali Bhar. Anisotropic compact star model: a brief study via embedding. *European Physical Journal C*, 79(2):138, February 2019.
- [58] B. V. Ivanov. Maximum bounds on the surface redshift of anisotropic stars. *Physical Review D*, 65(10), Apr 2002.
- [59] K. D. Krori and J. Barua. A singularity-free solution for a charged fluid sphere in general relativity. *Journal of Physics A Mathematical General*, 8(4):508–511, April 1975.
- [60] Victor Varela, et al. Charged anisotropic matter with linear or nonlinear equation of state. *Phys. Rev. D*, 82:044052, Aug 2010.
- [61] Farook Rahaman, et al. Strange stars in Krori-Barua space-time. *European Physical Journal C*, 72:2071, July 2012.
- [62] Mehedi Kalam, et al. Central density dependent anisotropic compact stars. *The European Physical Journal C*, 73(4), Apr 2013.
- [63] Piyali Bhar. Strange quintessence star in Krori–Barua spacetime. *Astrophys. & Space Sci.*, 356(2):365–373, 2015.
- [64] Piyali Bhar. Singularity-free anisotropic strange quintessence star. *Astrophys. Space Sci.*, 356(2):309–318, April 2015.
- [65] Piyali Bhar, et al. Anisotropic charged fluids with chaplygin equation of state in $(2 + 1)$ dimension. *Astrophysics and Space Science*, 360, 11 2015.
- [66] G. Abbas, Shahid Qaisar, and Abdul Jawad. Strange stars in $f(T)$ gravity with MIT bag model. *Astrophys. Space Sci.*, 359:17, October 2015.
- [67] M. Zubair, G. Abbas, and I. Noureen. Possible formation of compact stars in $f(R, T)$ gravity. *Astrophys. Space Sci.*, 361:8, January 2016.
- [68] M. Zubair and G. Abbas. Some interior models of compact stars in $f(R)$ gravity. *Astrophys. Space Sci.*, 361(10):342, October 2016.
- [69] M. Sharif and Arfa Waseem. Physical behavior of anisotropic compact stars in $f(R, T, R_{\mu\nu}T^{\mu\nu})$ gravity. *Canadian Journal of Physics*, 94(10):1024–1039, October 2016.
- [70] M. Ilyas, et al. Existence of relativistic structures in $f(R, T)$ gravity. *Astrophys. Space Sci.*, 362(12):237, December 2017.
- [71] Z. Yousaf, M. Zaeem-ul-Haq Bhatti, and M. Ilyas. Existence of compact structures in $f(R, T)$ gravity. *European Physical Journal C*, 78(4):307, April 2018.
- [72] Ines G. Salako, Abdul Jawad, and Hooman Moradpour. Anisotropic compact stars in non-conservative theory of gravity. *International Journal of Geometric Methods in Modern Physics*, 15(6):1850093-595, January 2018.
- [73] M. Farasat Shamir and Adnan Malik. Behavior of Anisotropic Compact Stars in $f(R, \phi)$ Gravity. *Communications in Theoretical Physics*, 71(5):599, May 2019.
- [74] Parneli Saha and Ujjal Debnath. Study of anisotropic compact stars with quintessence field and modified chaplygin gas in $f(T)$ gravity. *European Physical Journal C*, 79(11):919, November 2019.
- [75] M. Farasat Shamir and I. Fayyaz. Charged anisotropic compact stars in Logarithmic-Corrected R^2 gravity. *International Journal of Modern Physics A*, 35(4):2050013, February 2020.

- [76] M. Farasat Shamir and Iffat Fayyaz. Analysis of compact stars in logarithmic-corrected R^2 gravity. *Modern Physics Letters A*, 35(2):1950354, January 2020.
- [77] Zacharias Roupas and Gamal G. L. Nashed. Anisotropic Neutron Stars Modelling: Constraints in Krori-Barua Spacetime. *arXiv e-prints*, arXiv:2007.09797, July 2020.
- [78] Zacharias Roupas. Gravitational potential from maximum entropy principle. *Classical and Quantum Gravity*, 37(9):097001, May 2020.
- [79] Charalampos A. Kolassis, Nilton O. Santos, and Dimitri Tsoubelis. Energy conditions for an imperfect fluid. *Classical and Quantum Gravity*, 5(10):1329–1338, October 1988.
- [80] B. V. Ivanov. Analytical study of anisotropic compact star models. *European Physical Journal C*, 77(11):738, November 2017.
- [81] R. Chan, L. Herrera, and N. O. Santos. Dynamical Instability for Shearing Viscous Collapse. *MNRAS*, 267:637, April 1994.
- [82] L. Herrera and N. O. Santos. Local anisotropy in self-gravitating systems. *Phys. Rep.* , 286:53–130, July 1997.
- [83] L. Herrera. Cracking of self-gravitating compact objects. *Physics Letters A*, 165(3):206 – 210, 1992.
- [84] H. Abreu, H. Hernández, and L. A. Núñez. Sound speeds, cracking and the stability of self-gravitating anisotropic compact objects. *Classical and Quantum Gravity*, 24(18):4631–4645, September 2007.