

Generalized Scalar Field Cosmologies: A Global Dynamical Systems Formulation

Genly Leon

Departamento de Matemáticas, Universidad Católica del Norte, Avda. Angamos 0610,
Casilla 1280 Antofagasta, Chile.

E-mail: genly.leon@ucn.cl

Felipe Orlando Franz Silva

Departamento de Matemáticas, Universidad Católica del Norte, Avda. Angamos 0610,
Casilla 1280 Antofagasta, Chile.

E-mail: felipe.franz@alumnos.ucn.cl

Abstract. Local and global phase-space descriptions and averaging methods are used to find qualitative features of solutions for the FLRW and the Bianchi I metrics in the context of scalar field cosmologies with arbitrary potentials and arbitrary couplings to matter. The stability of the equilibrium points in a phase-space as well as the dynamics in the regime where the scalar field diverges are studied. Equilibrium points that represent some solutions of cosmological interest such as: several types of scaling solutions, a kinetic dominated solution representing a stiff fluid, a solution dominated by an effective energy density of geometric origin, a quintessence scalar field dominated solution, the vacuum de Sitter solution associated to the minimum of the potential, and a non-interacting matter dominated solution are obtained. All reveal a very rich cosmological phenomenology.

PACS numbers: 98.80.-k, 98.80.Jk, 95.36.+x

1. Introduction

Gravitational scalar field theories of special interest are Jordan-Brans-Dicke Theories [1, 2], Horndeski Theories [4], Teleparallel Analogue of Horndeski Theories [5, 6, 7], Inflationary Models [3], Extended Quintessence, Modified Gravity, Hořava-Lifschitz and the Galileons, etc., [8, 9, 10, 11, 12, 13, 14, 15, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69]. There are several studies in literature that provide both global and local dynamical systems analysis for scalar field cosmologies with arbitrary potentials and arbitrary couplings, say [70, 71, 72, 73, 74, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92] (see paper [93] and discussions therein).

In [93], the action for a general class of Scalar Tensor Theory of Gravity written in the so-called Einstein's frame was studied. This action is given by [94, 95]:

$$\mathcal{L} = \int d^4x \sqrt{|g|} \left\{ \frac{1}{2}R - \frac{1}{2}g^{\mu\nu}\nabla_\mu\phi\nabla_\nu\phi - V(\phi) - \Lambda + \chi(\phi)^{-2}\mathcal{L}(\mu, \nabla\mu, \chi(\phi)^{-1}g_{\alpha\beta}) \right\}, \quad (1)$$

where a system of units in which $8\pi G = c = \hbar = 1$ is used. In this equation R is the curvature scalar, ϕ is the scalar field, ∇_α is the covariant derivative, $V(\phi)$ is the self-interaction potential, $\Lambda \geq 0$ is the Cosmological Constant, $\chi(\phi)^{-2}$ is the coupling function. \mathcal{L} is the matter Lagrangian, and μ is the collective name for the matter degrees of freedom.

For the action (1), the matter energy-momentum tensor is defined by:

$$T_{\alpha\beta} = -\frac{2}{\sqrt{|g|}}\frac{\delta}{\delta g^{\alpha\beta}} \left\{ \sqrt{|g|}\chi^{-2}\mathcal{L}(\mu, \nabla\mu, \chi^{-1}g_{\alpha\beta}) \right\}. \quad (2)$$

The “energy exchange” vector is defined by:

$$Q_\beta \equiv \nabla^\alpha T_{\alpha\beta} = -\frac{1}{2}T\frac{1}{\chi(\phi)}\frac{d\chi(\phi)}{d\phi}\nabla_\beta\phi, \quad T = T^\alpha_\alpha, \quad (3)$$

where T is the trace of the energy-momentum tensor. The matter is assumed in the form of a perfect fluid with energy density $\rho_m \geq 0$ and pressure $p_m = (\gamma - 1)\rho_m$, $\gamma \in [1, 2]$. The expression (3) implies that the scalar field ϕ is non-minimally coupled to matter with coupling function $\chi(\phi)$. Additionally, the geometric properties of the metric are incorporated in the form of the function:

$$G_0(a) = \begin{cases} -3\frac{k}{a^2}, k = 0, \pm 1, & \text{FLRW} \\ \frac{\sigma_0^2}{a^6}, & \text{Bianchi I} \end{cases}. \quad (4)$$

The following field equations holds:

$$\dot{H} = -\frac{1}{2}(\gamma\rho_m + y^2) + \frac{1}{6}aG'_0(a), \quad (5a)$$

$$\dot{\rho}_m = -3\gamma H\rho_m - \frac{1}{2}(4 - 3\gamma)\rho_my\frac{d\ln\chi(\phi)}{d\phi}, \quad (5b)$$

$$\dot{a} = aH, \quad (5c)$$

$$\dot{y} = -3Hy - \frac{dV(\phi)}{d\phi} + \frac{1}{2}(4 - 3\gamma)\rho_m\frac{d\ln\chi(\phi)}{d\phi}, \quad (5d)$$

$$\dot{\phi} = y, \quad (5e)$$

$$3H^2 = \rho_m + \frac{1}{2}\dot{\phi}^2 + V(\phi) + \Lambda + G_0(a), \quad (5f)$$

The phase-space can be defined using equation (5f) as follows:

$$\left\{ (H, \rho_m, a, y, \phi) \in \mathbb{R}^5 : 3H^2 = \rho_m + \frac{1}{2}y^2 + V(\phi) + \Lambda + G_0(a) \right\}. \quad (6)$$

In paper [93] some theorems related to the asymptotic behavior of a very general cosmological model given by system (5) were presented. In particular, the conditions of a scalar field potential $V \in C^2(\mathbb{R})$ under which $\lim_{t \rightarrow \infty} \dot{\phi} = 0$ were discussed; they are: non-negativity of the potential which is zero only at the origin and the boundedness of both $V'(\phi)$ and $V(\phi)$ (Theorem 2.1.3 in [93]). Furthermore, some extra conditions for having $\lim_{t \rightarrow \infty} \phi(t) \in \{-\infty, 0, +\infty\}$ were presented. They are the previous conditions with the addition of $V'(\phi) > 0$ for $\phi > 0$ and $V'(\phi) < 0$ for $\phi < 0$ (Theorem 2.2.3 in [93]). Mild conditions under the potential for having $\lim_{t \rightarrow +\infty} \dot{\phi} = 0$ and $\lim_{t \rightarrow +\infty} \phi(t) = +\infty$ (Theorem 2.3.3 in [93]) were considered. It was examined to which extent the hypotheses of the theorems proved in [93] can be relaxed in order to obtain the same conclusions or provide a counterexample, by means of generalized harmonic self-interacting potentials: $V_1(\phi) = \mu^3 \left[\frac{\phi^2}{\mu} + bf \cos \left(\delta + \frac{\phi}{f} \right) \right]$, $b \neq 0$ and $V_2(\phi) = \mu^3 \left[bf \left(\cos(\delta) - \cos \left(\delta + \frac{\phi}{f} \right) \right) + \frac{\phi^2}{\mu} \right]$, $b \neq 0$. Harmonic potentials plus cosine corrections were introduced in the context of inflation in loop-quantum cosmology in [96]. Finally, the Hubble-normalized formulation for a scalar field cosmology with generalized harmonic potential $V(\phi) = \frac{\phi^2}{2} + f \left[1 - \cos \left(\frac{\phi}{f} \right) \right]$, $f > 0$, nonminimally coupled to matter with coupling function $\chi = \chi_0 e^{\frac{\lambda \phi}{4-3\gamma}}$, where λ is a constant and $0 \leq \gamma \leq 2$, $\gamma \neq \frac{4}{3}$ was performed for FLRW metrics and for Bianchi I metric. Using Theorem 2.4 in [93], the late time attractors correspond to the non zero local minimums of the potential, i.e., $\phi = \phi_*$, satisfying $\sin(\phi^*/f) + \phi^* = 0$, $0 < |\phi^*| \leq 1$, $\cos \left(\frac{\phi^*}{f} \right) + f > 0$. Whenever ϕ^* is a local non zero minimum of $V(\phi)$:

$$\lim_{t \rightarrow \infty} \phi(t) = \phi^*, \lim_{t \rightarrow \infty} \dot{\phi}(t) = 0, \lim_{t \rightarrow \infty} H(t) = \frac{\sqrt{V(\phi^*)}}{\sqrt{3}}, \text{ and } \lim_{t \rightarrow \infty} G_0(a) = 0.$$

The global minimum of $V(\phi)$ at $\phi = 0$ is unstable to curvature perturbations for $\gamma > \frac{2}{3}$ in case of negatively curved FLRW model. This confirms the result in [97] that for $\gamma > 2/3$, in a non-degenerated minima with zero critical value the curvature has a dominant effect on the late evolution of the universe and will eventually dominate both the perfect fluid and the scalar field. For Bianchi I model the global minimum $V(0) = 0$ is unstable to shear perturbations.

In this companion paper complementary formulations based on [98] will be presented, including the analysis of the oscillations entering the system via Klein–Gordon equation [99].

It is useful to define the quantities $\lambda = -\frac{V'(\phi)}{V(\phi)}$, $f \equiv \frac{V''(\phi)}{V(\phi)} - \frac{V'(\phi)^2}{V(\phi)^2}$ to obtain qualitative information about the past and future asymptotic structure of the equations's solutions in scalar field theories with potential $V(\phi)$. This is the basis of the so-called method of f -devisers. For the common scalar field potentials the function $f(\lambda)$ is found as follows. The monomial potential $V(\phi) = \frac{1}{2n}(\mu\phi)^{2n}$, $\mu > 0$, $n = 1, 2, \dots$ [100] has $f(\lambda) = -\frac{\lambda^2}{2n}$. The so-called E -model studied from the dynamical systems point of view in [101] has potential $V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3\alpha}}\phi} \right)^{2n}$. The corresponding f -deviser is $f(\lambda) = -\frac{\lambda(\lambda-\sqrt{6}\mu)}{2n}$, where $\mu = \frac{n}{3\sqrt{\alpha}}$. The exponential potential plus a cosmological constant $V(\phi) = V_0 e^{-l\phi} + V_1$ [102, 103, 104] has $f(\lambda) = -\lambda(\lambda-l)$. The hyperbolic potentials: $V(\phi) = V_0 (\cosh(\xi\phi) - 1)$ [103, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114] with $f(\lambda) = -\frac{1}{2}(\lambda^2 - \xi^2)$ and $V(\phi) = V_0 \sinh^{-\alpha}(\beta\phi)$ [103, 105, 106, 108, 109, 112, 115] with $f(\lambda) = \frac{\lambda^2}{\alpha} - \alpha\beta^2$. The double exponential potential $V(\phi) = V_0 (e^{\alpha\phi} + e^{\beta\phi})$ [25, 116, 117] has $f(\lambda) = -(\lambda + \alpha)(\lambda + \beta)$.

This paper's proposal, for a scalar field non-minimally coupled to matter, relies on defining $\lambda = -\frac{W'(\phi)}{W(\phi)}$, $f = \frac{W''(\phi)}{W(\phi)} - \frac{W'(\phi)^2}{W(\phi)^2}$, $g = -\frac{\chi'(\phi)}{\chi(\phi)}$, where $W(\phi) = \Lambda + V(\phi)$, $W(\phi) \geq 0$. Then, it is assumed that f and g can be explicitly written as functions of λ , as in the original f -deviser method. This is a significant advantage due to the analysis for arbitrary potential can be performed as a first step; then, just substitute the desired forms instead of repeating the whole procedure for every distinct potential. Similar approaches were used in [58, 68] for two-field Jordan-Brans-Dicke cosmologies.

This paper is organized as follows. A local dynamical systems analysis using Hubble normalized equations is performed in Section 2. Then, a dynamical systems formulation using global dynamical systems variables based on Alho & Uggla's approach [98] is given in Section 3. Section 3.1 is devoted to the asymptotic analysis as $\phi \rightarrow \infty$ for arbitrary $V(\phi)$ and $\chi(\phi)$. In Section 3.2, the physical interpretation of the solutions that were found is discussed. Section 4 is devoted to equations analysis of a scalar field model with potential $V(\phi) = V_0 e^{-\lambda\phi}$ in a vacuum. In Section 5, the potential of the so-called E -model: $V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3\alpha}}\phi}\right)^{2n}$ is investigated. This potential was discussed in [101] for a conventional scalar field cosmology and in [59] for HoÅŽavaâŞLifshitz cosmology. In Sections 6 and 7, scalar field cosmologies under the potentials $V_1(\phi) = \mu^3 \left[\frac{\phi^2}{\mu} + bf \cos \left(\delta + \frac{\phi}{f} \right) \right]$, $b \neq 0$ and $V_2(\phi) = \mu^3 \left[bf \left(\cos(\delta) - \cos \left(\delta + \frac{\phi}{f} \right) \right) + \frac{\phi^2}{\mu} \right]$, $b \neq 0$ are respectively studied. Sections 6.1 and 7.1 are devoted to the analysis of the dynamics as $\phi \rightarrow \infty$. In Sections 6.2 and 7.2, the study of the oscillatory behavior is presented. In Section 8, the main results are summarized. Finally, Section 9 is devoted to conclusions.

2. Local dynamical systems analysis for arbitrary $V(\phi)$ and $\chi(\phi)$

For the analysis of the system (5) for an arbitrary $V(\phi)$ and arbitrary $\chi(\phi)$ are defined:

$$\lambda(\phi) = -\frac{W'(\phi)}{W(\phi)}, \quad f(\phi) = \frac{W''(\phi)}{W(\phi)} - \frac{W'(\phi)^2}{W(\phi)^2}, \quad g(\phi) = -\frac{\chi'(\phi)}{\chi(\phi)}, \quad (7)$$

where $W(\phi) = \Lambda + V(\phi)$, $W(\phi) \geq 0$. The conditions (7) can be alternatively written as:

$$\frac{d\lambda}{d\phi} = -f(\phi), \quad \frac{dW}{d\phi} = -\lambda(\phi)W(\phi), \quad \frac{d\chi}{d\phi} = -g(\phi)\chi(\phi). \quad (8)$$

The idea is to assume that f and g can be explicitly written as functions of λ . Hence, (8) is written as:

$$\frac{d\phi}{d\lambda} = -\frac{1}{f(\lambda)}, \quad \frac{dW}{d\lambda} = \frac{\lambda W(\lambda)}{f(\lambda)}, \quad \frac{d\chi}{d\lambda} = \frac{g(\lambda)\chi(\lambda)}{f(\lambda)}, \quad (9)$$

which can be integrated in quadrature as:

$$\phi(\lambda) = \phi(1) - \int_1^\lambda \frac{1}{f(s)} ds, \quad W(\lambda) = W(1)e^{\int_1^\lambda \frac{s}{f(s)} ds}, \quad \chi(\lambda) = \chi(1)e^{\int_1^\lambda \frac{g(s)}{f(s)} ds}. \quad (10)$$

The last equations can be used to generate the potentials and couplings by giving the functions $f(\lambda)$ and $g(\lambda)$ as input.

Defining the new variables:

$$x = \frac{\dot{\phi}}{\sqrt{6}H}, \quad \Omega_m = \frac{\rho_m}{3H^2}, \quad \Omega_0 = \frac{G_0(a)}{3H^2}, \quad \lambda = -\frac{V'(\phi)}{\Lambda + V(\phi)}, \quad (11)$$

where $G_0(a) = qa^{-p}$, $p \geq 0$, and the time variable $\tau = \ln a$, the following dynamical system is obtained:

$$\frac{dx}{d\tau} = \frac{1}{2}x(3\gamma\Omega_m + p\Omega_0 + 6x^2 - 6) - \sqrt{\frac{3}{2}}\lambda(x^2 + \Omega_0 + \Omega_m - 1) + \frac{\sqrt{6}}{4}(3\gamma - 4)\Omega_m g(\lambda), \quad (12a)$$

$$\frac{d\Omega_m}{d\tau} = \frac{1}{2}\Omega_m \left(\sqrt{6}(4 - 3\gamma)xg(\lambda) + 2(3\gamma(\Omega_m - 1) + p\Omega_0 + 6x^2) \right), \quad (12b)$$

$$\frac{d\Omega_0}{d\tau} = \Omega_0(3\gamma\Omega_m + p(\Omega_0 - 1) + 6x^2), \quad (12c)$$

$$\frac{d\lambda}{d\tau} = -\sqrt{6}xf(\lambda), \quad (12d)$$

with the restriction

$$x^2 + \Omega_m + \Omega_0 = 1 - \frac{W(\phi)}{3H^2} \leq 1. \quad (13)$$

The equilibrium points of the system (12) are the following:

$A_1(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{(4-3\gamma)g(\hat{\lambda})}{\sqrt{6}(\gamma-2)}, 1 - \frac{(4-3\gamma)^2g(\hat{\lambda})^2}{6(\gamma-2)^2}, 0, \hat{\lambda} \right)$, where $\hat{\lambda}$ denotes any value of λ satisfying $f(\hat{\lambda}) = 0$, represents a matter - kinetic scaling solution. For $1 \leq \gamma < 2$, it follows that $A_1(\hat{\lambda})$ is a sink under one of the conditions (i)- (x) in [Appendix A](#). If it exists, it will never be a source.

$A_2(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{\sqrt{\frac{2}{3}}(p-3\gamma)}{(3\gamma-4)g(\hat{\lambda})}, \frac{2(6-p)(p-3\gamma)}{3(4-3\gamma)^2g(\hat{\lambda})^2}, \frac{2(p-3\gamma)(\gamma-2)}{(4-3\gamma)^2g(\hat{\lambda})^2} + 1, \hat{\lambda} \right)$ represents a matter- scalar field - geometric “fluid” scaling solution. For $1 \leq \gamma < 2$, it follows that $A_2(\hat{\lambda})$ is a sink under one of the conditions (i) - (viii) in [Appendix A](#). It is non-hyperbolic for $p = 6$, and a saddle for $p = 2$. If exists, it will never be a source.

$A_3(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{-\sqrt{6}\gamma}{(4-3\gamma)g(\hat{\lambda})+2\hat{\lambda}}, \frac{2(4-3\gamma)g(\hat{\lambda})\hat{\lambda}+4(\hat{\lambda}^2-3\gamma)}{((3\gamma-4)g(\hat{\lambda})-2\hat{\lambda})^2}, 0, \hat{\lambda} \right)$ represents a matter - scalar field scaling solution. In this case, it can be proceeded semi-analytically, that is, for non-minimal coupling $g \equiv 0$ and assuming $1 \leq \gamma < 2$, $A_3(\hat{\lambda})$ is a sink for:

i) $1 \leq \gamma < 2, p > 3\gamma, -\frac{2\sqrt{6}\gamma}{\sqrt{9\gamma-2}} \leq \hat{\lambda} < -\sqrt{3}\sqrt{\gamma}, f'(\hat{\lambda}) < 0$, or

ii) $1 \leq \gamma < 2, p > 3\gamma, \sqrt{3}\sqrt{\gamma} < \hat{\lambda} \leq \frac{2\sqrt{6}\gamma}{\sqrt{9\gamma-2}}, f'(\hat{\lambda}) > 0$.

Otherwise, it is a saddle. For non-minimal coupling the analysis has to be done numerically.

$A_4(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (-1, 0, 0, \hat{\lambda})$ is a kinetic dominated solution representing a stiff fluid.

$A_4(\hat{\lambda})$ is a source for:

i) $1 \leq \gamma < \frac{4}{3}, 0 \leq p < 6, g(\hat{\lambda}) < \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > -\sqrt{6}$, or

ii) $\gamma = \frac{4}{3}, 0 \leq p < 6, f'(\hat{\lambda}) > 0, \hat{\lambda} > -\sqrt{6}$, or

iii) $\frac{4}{3} < \gamma < 2, 0 \leq p < 6, g(\hat{\lambda}) > \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > -\sqrt{6}$.

$A_4(\hat{\lambda})$ is a sink for:

i) $1 \leq \gamma < \frac{4}{3}, p > 6, g(\hat{\lambda}) > \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < -\sqrt{6}$, or

ii) $\frac{4}{3} < \gamma < 2, p > 6, g(\hat{\lambda}) < \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < -\sqrt{6}$.

$A_5(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (0, 0, 1, \hat{\lambda})$ is a solution dominated by the effective energy density of $G_0(a)$. It is non-hyperbolic with a 3D unstable manifold for $p > 6$.

$A_6(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (1, 0, 0, \hat{\lambda})$ is a kinetic dominated solution representing a stiff fluid.

$A_6(\hat{\lambda})$ is a source for:

- i) $1 \leq \gamma < \frac{4}{3}, p < 6, g(\hat{\lambda}) > -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < \sqrt{6}$, or
- ii) $\gamma = \frac{4}{3}, p < 6, f'(\hat{\lambda}) < 0, \hat{\lambda} < \sqrt{6}$, or
- iii) $\frac{4}{3} < \gamma < 2, p < 6, g(\hat{\lambda}) < -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < \sqrt{6}$.

$A_6(\hat{\lambda})$ is a sink for:

- i) $1 \leq \gamma < \frac{4}{3}, p > 6, g(\hat{\lambda}) < -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > \sqrt{6}$, or
- ii) $1 \leq \gamma > \frac{4}{3}, p > 6, g(\hat{\lambda}) > -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > \sqrt{6}$.

$A_7(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{p}{\sqrt{6}\hat{\lambda}}, 0, 1 - \frac{p}{\hat{\lambda}^2}, \hat{\lambda}\right)$ is a scaling solution where the energy density of the scalar field and the effective energy density from $G_0(a)$ scale with the same order of magnitude. For $1 \leq \gamma \leq 2$, $A_7(\hat{\lambda})$, it is a sink for one of the conditions (i) - (xiv) in [Appendix A](#). It will never be a source.

$A_8(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{\hat{\lambda}}{\sqrt{6}}, 0, 0, \hat{\lambda}\right)$ represents the typical quintessence scalar field dominated solution. Assuming $1 \leq \gamma \leq 2$, $A_8(\hat{\lambda})$ is a sink for:

- i) $1 \leq \gamma < \frac{4}{3}, p > \hat{\lambda}^2, f'(\hat{\lambda}) > 0, 0 < \hat{\lambda} < \sqrt{6}, g(\hat{\lambda}) < \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}$, or
- ii) $1 \leq \gamma < \frac{4}{3}, p > \hat{\lambda}^2, g(\hat{\lambda}) > \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}, -\sqrt{6} < \hat{\lambda} < 0, f'(\hat{\lambda}) < 0$, or
- iii) $\gamma = \frac{4}{3}, p > \hat{\lambda}^2, -2 < \hat{\lambda} < 0, f'(\hat{\lambda}) < 0$, or
- iv) $\gamma = \frac{4}{3}, p > \hat{\lambda}^2, f'(\hat{\lambda}) > 0, 0 < \hat{\lambda} < 2$, or
- v) $\frac{4}{3} < \gamma < 2, p > \hat{\lambda}^2, g(\hat{\lambda}) > \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}, f'(\hat{\lambda}) > 0, 0 < \hat{\lambda} < \sqrt{6}$, or
- vi) $\frac{4}{3} < \gamma < 2, p > \hat{\lambda}^2, -\sqrt{6} < \hat{\lambda} < 0, g(\hat{\lambda}) < \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}, f'(\hat{\lambda}) < 0$.

It will never be a source.

A_9 : $(x, \Omega_m, \Omega_0, \lambda) = (0, 0, 0, 0)$ represents the vacuum de Sitter solution associated to the minimum of the potential. It is a sink for $p > 0, \gamma > 0, f(0) > 0$. Otherwise, it is a saddle.

$A_{10}(\tilde{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (0, 1, 0, \tilde{\lambda})$, where we denote by $\tilde{\lambda}$, the values of λ for which $g(\lambda) = 0$. It represents a non-interacting matter dominated solution. It is a saddle.

3. Global dynamical analysis for arbitrary $V(\phi)$ and $\chi(\phi)$

In Section 2, the stability of the equilibrium points using Hubble-normalized equations, which is essentially based on the Copeland, Liddle & Wands's approach [69] was investigated. It is well-known that this procedure is well-suited to investigate local stability features of the equilibrium points. However, it does not provide a global description of the phase space when generically ϕ diverges or when $H \rightarrow 0$; in these cases the method fails. For this reason, a new global systems analysis for arbitrary $V(\phi)$ and $\chi(\phi)$ as $|\phi| \rightarrow \infty$ is presented based on Alho & Uggla's approach [98]. Doing so, setting $\Lambda = 0$ for simplicity.

Now, it is assumed that the following limits exist:

$$N = \lim_{\phi \rightarrow +\infty} \frac{V'(\phi)}{V(\phi)}, \text{ and } M = \lim_{\phi \rightarrow +\infty} \frac{\chi'(\phi)}{\chi(\phi)}. \quad (14)$$

This permits the definition of the new functions $W_V(\phi), W_\chi(\phi)$:

$$W_V(\phi) = \frac{V'(\phi)}{V(\phi)} - N, \quad W_\chi(\phi) = \frac{\chi'(\phi)}{\chi(\phi)} - M, \quad (15)$$

where the constants N and M are such that

$$\lim_{\phi \rightarrow +\infty} W_V(\phi) = 0, \quad \lim_{\phi \rightarrow +\infty} W_\chi(\phi) = 0. \quad (16)$$

After, the following variables are defined:

$$\begin{aligned} T &= \frac{m}{m+H}, \quad \theta = \tan^{-1} \left(\frac{\dot{\phi}}{\sqrt{2V(\phi) + 2\rho_m + 2G_0(a)}} \right), \\ \Omega_m &= \frac{\rho_m}{3H^2}, \quad \Omega_0 = \frac{G_0(a)}{3H^2}, \quad m > 0, \end{aligned} \quad (17)$$

where $G_0(a) = qa^{-p}$, $p \geq 0$, such that:

$$\begin{aligned} V(\phi) &= \frac{3m^2(1-T)^2(\cos(2\theta) - 2\Omega_0 - 2\Omega_m + 1)}{2T^2}, \\ H &= m \left(\frac{1-T}{T} \right), \quad \dot{\phi} = \frac{\sqrt{6}m(1-T)\sin(\theta)}{T}, \quad \rho_m = \frac{3m^2(1-T)^2\Omega_m}{T^2}. \end{aligned} \quad (18)$$

Finally, the time variable $\tau = \ln a$ is defined; resulting the following unconstrained dynamical system:

$$\frac{dT}{d\tau} = \frac{1}{2}(1-T)T(3\gamma\Omega_m + 6\sin^2(\theta) + p\Omega_0), \quad (19a)$$

$$\begin{aligned} \frac{d\theta}{d\tau} &= -\frac{\sec(\theta)}{4\sqrt{6}} \left\{ 3\sqrt{6}\sin(3\theta) + 6(\cos(2\theta) - 2\Omega_0 + 1)(N + W_V(\phi)) \right. \\ &\quad \left. + \Omega_m(-6\sqrt{6}\sin(\theta) - 12(2M + N + W_V(\phi) + 2W_\chi(\phi)) + 18\gamma(M + W_\chi(\phi))) \right. \\ &\quad \left. + \sqrt{6}\sin(\theta)(3 - 2p\Omega_0) \right\}, \end{aligned} \quad (19b)$$

$$\begin{aligned} \frac{d\Omega_m}{d\tau} &= -\frac{1}{2}\Omega_m \left\{ 6\cos(2\theta) - \sqrt{6}(3\gamma - 4)\sin(\theta)(M + W_\chi(\phi)) \right. \\ &\quad \left. - 2(3\gamma(\Omega_m - 1) + p\Omega_0 + 3) \right\}, \end{aligned} \quad (19c)$$

$$\frac{d\Omega_0}{d\tau} = -\Omega_0(-3\gamma\Omega_m + 3\cos(2\theta) - p\Omega_0 + p - 3), \quad (19d)$$

$$\frac{d\phi}{d\tau} = \sqrt{6}\sin(\theta). \quad (19e)$$

Rather to discuss all the equilibrium points of (19) (which is a complementary formulation of the system (12)), the focus will be to study what happens in the region of $\phi \rightarrow +\infty$. Due to the symmetries of the model, it is not necessary to study the limit $\phi \rightarrow -\infty$, which can be accomplished by taking the reversal $\phi \rightarrow -\phi$.

3.1. Asymptotic analysis as $\phi \rightarrow \infty$

In this section the stability of the equilibrium points of (19) as $\phi \rightarrow \infty$ (for functions V and χ well-behaved at infinity of exponential orders of N and M , respectively) is discussed. This analysis at infinity complements the analysis of Section 2.

Definition 1 (Definition 1 [70]) Let $V : \mathbb{R} \rightarrow \mathbb{R}$ be a $C^2(\mathbb{R})$ nonnegative function. Let there exist some $\phi_0 > 0$ for which $V(\phi) > 0$ for all $\phi > \phi_0$ and some number $N < \infty$, such that the function:

$$W_V : [\phi_0, \infty) \rightarrow \mathbb{R}, \quad \phi \mapsto \frac{V'(\phi)}{V(\phi)} - N \quad (20)$$

is well-defined and satisfies

$$\lim_{\phi \rightarrow \infty} W_V(\phi) = 0, \quad (21)$$

then, V is said to be well-behaved at infinity (WBI) of exponential order N .

Definition 2 (Definition 2 [70]) A $C^k(\mathbb{R})$ function $V(\phi)$ is a class k WBI function, if it is WBI of exponential order N , and there are $\phi_0 > 0$, and a coordinate transformation $\varphi = h(\phi)$ which maps the interval $[\phi_0, \infty)$ onto $(0, \epsilon]$, where $\epsilon = h(\phi_0)$, satisfying $\lim_{\phi \rightarrow +\infty} h(\phi) = 0$, and with the following additional properties:

(i) h is C^{k+1} and strictly decreasing.

(ii) The functions:

$$\bar{W}_V = \begin{cases} \frac{V'(h^{-1}(\varphi))}{V(h^{-1}(\varphi))} - N, & \varphi > 0, \\ 0, & \varphi = 0 \end{cases} \quad (22)$$

and

$$\bar{h}'(\varphi) = \begin{cases} h'(h^{-1}(\varphi)), & \varphi > 0, \\ \lim_{\phi \rightarrow \infty} h'(\phi), & \varphi = 0 \end{cases} \quad (23)$$

are C^k on the closed interval $[0, \epsilon]$ and

(iii)

$$\frac{d\bar{W}_V}{d\varphi}(0) = \frac{d\bar{h}'}{d\varphi}(0) = 0. \quad (24)$$

The last hypotheses are equivalent to:

$$\lim_{\varphi \rightarrow 0} \frac{W'_\chi(h^{-1}(\varphi))}{h'(h^{-1}(\varphi))} \text{ and } \lim_{\varphi \rightarrow 0} \frac{h''(h^{-1}(\varphi))}{h'(h^{-1}(\varphi))} = 0.$$

Assuming the C^2 functions $V(\phi)$ and $\chi(\phi)$ are class 2 WBI, the unconstrained dynamical system is obtained:

$$\frac{dT}{d\tau} = \frac{1}{2}(1-T)T \left(3\gamma\Omega_m + 6\sin^2(\theta) + p\Omega_0 \right), \quad (25a)$$

$$\begin{aligned} \frac{d\theta}{d\tau} = & -\frac{\sec(\theta)}{4\sqrt{6}} \left\{ 3\sqrt{6}\sin(3\theta) + 6(\cos(2\theta) - 2\Omega_0 + 1)(N + \bar{W}_V(\varphi)) \right. \\ & + \Omega_m \left(-6\sqrt{6}\gamma\sin(\theta) - 12(2M + N + \bar{W}_V(\varphi) + 2\bar{W}_\chi(\varphi)) + 18\gamma(M + \bar{W}_\chi(\varphi)) \right) \\ & \left. + \sqrt{6}\sin(\theta)(3 - 2p\Omega_0) \right\}, \end{aligned} \quad (25b)$$

$$\frac{d\Omega_m}{d\tau} = -\frac{1}{2}\Omega_m \left\{ 6\cos(2\theta) - \sqrt{6}(3\gamma - 4)\sin(\theta)(M + \bar{W}_\chi(\varphi)) - 2(3\gamma(\Omega_m - 1) + p\Omega_0 + 3) \right\}, \quad (25c)$$

$$\frac{d\Omega_0}{d\tau} = -\Omega_0(-3\gamma\Omega_m + 3\cos(2\theta) - p\Omega_0 + p - 3), \quad (25d)$$

$$\frac{d\varphi}{d\tau} = \sqrt{6}\bar{h}'(\varphi)\sin(\theta), \quad (25e)$$

defined on the phase space:

$$\left\{ (T, \theta, \Omega_m, \Omega_0, \varphi) \in \mathbb{R}^5 : 0 \leq T \leq 1, -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}, 0 \leq \varphi \leq h(\phi_0), 2T^2 V(h^{-1}(\varphi)) = 3m^2(1-T)^2(\cos(2\theta) - 2\Omega_0 - 2\Omega_m + 1) \right\}. \quad (26)$$

θ is unique modulo 2π . It has been chosen such that $\cos \theta \geq 0$. In the following list $\tan^{-1}[x, y]$ gives the arc tangent of y/x , taking into account on which quadrant the point (x, y) is in. When $x^2 + y^2 = 1$, $\tan^{-1}[x, y]$ gives the number θ such as $x = \cos \theta$ and $y = \sin \theta$.

The equilibrium points of system (25) with $\varphi = 0$ (i.e., corresponding to $\phi \rightarrow \infty$) are the following:

B_1 : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{N^2}{6}}, -\frac{N}{\sqrt{6}} \right] + 2\pi c_1, 0, 0, 0 \right)$, $c_1 \in \mathbb{Z}$, represents a scalar field dominated solution satisfying $H \rightarrow \infty$. It is always a non-hyperbolic saddle.

B_2 : $\left(1, \tan^{-1} \left[\sqrt{1 - \frac{N^2}{6}}, -\frac{N}{\sqrt{6}} \right] + 2\pi c_1, 0, 0, 0 \right)$, $c_1 \in \mathbb{Z}$, represents a scalar field dominated solution, satisfying $H \rightarrow 0$. The case of physical interest is when B_2 is non-hyperbolic with a 4D stable manifold under one of the conditions (i)- (x) of [Appendix B](#).

B_3 : $(0, 2\pi c_1, 0, 1, 0)$, $c_1 \in \mathbb{Z}$, with eigenvalues $\{0, \frac{p-6}{2}, \frac{p}{2}, p, p - 3\gamma\}$. It represents a geometric “fluid” dominated solution with $H \rightarrow \infty$. The physical interesting situation is when B_3 is non-hyperbolic with a 4D unstable manifold for $p > 6, 1 \leq \gamma \leq 2$. It is a non-hyperbolic saddle otherwise.

B_4 : $(1, 2\pi c_1, 0, 1, 0)$, $c_1 \in \mathbb{Z}$, represents a geometric “fluid” dominated solution with $H \rightarrow 0$. It is a non-hyperbolic saddle.

B_5 : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{p^2}{6N^2}}, -\frac{p}{\sqrt{6}N} \right] + 2\pi c_1, 0, 1 - \frac{p}{N^2}, 0 \right)$, $c_1 \in \mathbb{Z}$, represents a scaling solution where neither the energy density of the geometric “fluid”, nor the energy density of the scalar field completely dominates, that satisfies $H \rightarrow \infty$. It is a non-hyperbolic saddle.

B_6 : $\left(1, \tan^{-1} \left[\sqrt{1 - \frac{p^2}{6N^2}}, -\frac{p}{\sqrt{6}N} \right] + 2\pi c_1, 0, 1 - \frac{p}{N^2}, 0 \right)$, $c_1 \in \mathbb{Z}$, represents a scaling solution where neither the energy density of geometric “fluid” nor the energy density of the scalar field completely dominates, that satisfies $H \rightarrow 0$. The situation of physical interest is when B_6 is non-hyperbolic with a 4D stable manifold under one of the conditions (i)- (xl) in [Appendix B](#).

B_7 : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{2(p-3\gamma)^2}{3M^2(4-3\gamma)^2}}, \frac{2(p-3\gamma)}{\sqrt{6}M(4-3\gamma)} \right] + 2\pi c_1, \frac{2(6-p)(p-3\gamma)}{3(4-3\gamma)^2 M^2}, \frac{(4-3\gamma)^2 M^2 + 2(\gamma-2)(p-3\gamma)}{(4-3\gamma)^2 M^2}, 0, 0\right), c_1 \in \mathbb{Z}$, represents a matter- scalar field - geometric “fluid” scaling solution with $H \rightarrow \infty$. It is a non-hyperbolic saddle.

B_8 : $\left(1, \tan^{-1} \left[\sqrt{1 - \frac{2(p-3\gamma)^2}{3M^2(4-3\gamma)^2}}, \frac{2(p-3\gamma)}{\sqrt{6}M(4-3\gamma)} \right] + 2\pi c_1, \frac{2(6-p)(p-3\gamma)}{3(4-3\gamma)^2 M^2}, \frac{(4-3\gamma)^2 M^2 + 2(\gamma-2)(p-3\gamma)}{(4-3\gamma)^2 M^2}, 0, 0\right), c_1 \in \mathbb{Z}$, represents a matter- scalar field - geometric “fluid” scaling solution with $H \rightarrow 0$. The situation of physical interest is when B_8 is non-hyperbolic with a 4D stable manifold under the conditions (i) - (xi) of [Appendix B](#).

B_9 : $\left(0, \tan^{-1} \left[\frac{\sqrt{6(2-\gamma)^2 - (4-3\gamma)^2 M^2}}{\sqrt{6}(2-\gamma)}, \frac{(4-3\gamma)M}{\sqrt{6}(2-\gamma)} \right] + 2\pi c_1, \frac{6(2-\gamma)^2 - (4-3\gamma)^2 M^2}{6(2-\gamma)^2}, 0, 0, 0\right), c_1 \in \mathbb{Z}$, represents a matter- scalar field scaling solution with $H \rightarrow \infty$. It is a non-hyperbolic saddle.

B_{10} : $\left(1, \tan^{-1} \left[\frac{\sqrt{6(2-\gamma)^2 - (4-3\gamma)^2 M^2}}{\sqrt{6}(2-\gamma)}, \frac{(4-3\gamma)M}{\sqrt{6}(2-\gamma)} \right] + 2\pi c_1, \frac{6(2-\gamma)^2 - (4-3\gamma)^2 M^2}{6(2-\gamma)^2}, 0, 0, 0\right), c_1 \in \mathbb{Z}$, represents a matter- scalar field scaling solution with $H \rightarrow 0$. The situation of physical interest is when B_{10} is non-hyperbolic with a 4D stable manifold for:

$$\text{i) } 1 \leq \gamma < \frac{4}{3}, \quad N < -\sqrt{6}, \quad \frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} < M < \frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4}, \quad p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}, \text{ or}$$

$$\text{ii) } \frac{4}{3} < \gamma < 2, \quad N < -\sqrt{6}, \quad \frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4} < M < \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4}, \quad p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}, \text{ or}$$

$$\text{iii) } 1 \leq \gamma < \frac{4}{3}, \quad N > \sqrt{6}, \quad \frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4} < M < \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4}, \quad p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}, \text{ or}$$

$$\text{iv) } \frac{4}{3} < \gamma < 2, \quad N > \sqrt{6}, \quad \frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} < M < \frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4}, \quad p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}.$$

B_{11} : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{6\gamma^2}{(2N+M(4-3\gamma))^2}}, -\frac{\sqrt{6}\gamma}{2N+M(4-3\gamma)} \right] + 2\pi c_1, \frac{4N(2M+N)-6(2+MN)\gamma}{(2N+M(4-3\gamma))^2}, 0, 0, 0\right), c_1 \in \mathbb{Z}$, represents a matter- scalar field scaling solution with $H \rightarrow \infty$. It is a hyperbolic saddle.

B_{12} : $\left(1, \tan^{-1} \left[\sqrt{1 - \frac{6\gamma^2}{(2N+M(4-3\gamma))^2}}, -\frac{\sqrt{6}\gamma}{2N+M(4-3\gamma)} \right] + 2\pi c_1, \frac{4N(2M+N)-6(2+MN)\gamma}{(2N+M(4-3\gamma))^2}, 0, 0, 0\right), c_1 \in \mathbb{Z}$ represents a matter- scalar field scaling solution with $H \rightarrow 0$. The situation of physical interest is when B_{12} is non-hyperbolic with a 4D stable manifold under one of the conditions (i) - (lviii) in [Appendix B](#).

3.2. Discussion

Firstly, a dynamical system analysis of the system (5) for arbitrary $V(\phi)$ and arbitrary $\chi(\phi)$ formulated as system (12), was provided. Secondly, to complement the analysis of system (12), the dynamics of the system (5) at the limit $\phi \rightarrow +\infty$ formulated as system (25), was analyzed. In both cases, the equilibrium points in the finite region of the phase space (local analysis) as well as in the limit $\phi \rightarrow +\infty$ (global analysis at infinity) were exhaustively examined; obtaining equilibrium points that represent some solutions of cosmological interest, such as several types of scaling solutions. I.e., a kinetic dominated solution representing a stiff fluid, a solution dominated by an effective energy density of geometric origin, a quintessence scalar field dominated solution, the vacuum de Sitter solution associated to the minimum of the potential, and a non-interacting matter dominated solution.

4. Example: a scalar field model with potential $V(\phi) = V_0 e^{-\lambda\phi}$ in vacuum

In this section, a scalar field model with potential $V(\phi) = V_0 e^{-\lambda\phi}$ in vacuum for the flat FLRW metric is considered. That is, $N = -\lambda$, $M = 0$, $W_V \equiv 0$, $W_\chi \equiv 0$, $\Omega_m \equiv 0$, $\Omega_0 \equiv 0$, $\rho_m \equiv 0$, $G_0 \equiv 0$ and $\chi \equiv 1$.

The traditional formulation of the stability analysis of the equilibrium points in a cosmological setup uses Hubble-normalized equations, which is essentially based on the Copeland, Liddle & Wands's approach [69]. It is well-known that this procedure is well-suited to investigate local stability features of the equilibrium points.

Defining the variables:

$$x = \frac{\dot{\phi}}{\sqrt{6}H}, \quad y = \frac{\sqrt{V(\phi)}}{\sqrt{3}H}, \quad (27)$$

and $\tau = \ln a$, through $d\tau = Hdt$, $H > 0$, the following dynamical system is obtained:

$$\frac{dx}{d\tau} = 3x^3 - 3x + \sqrt{\frac{3}{2}}\lambda y^2, \quad \frac{dy}{d\tau} = 3xy \left(x - \frac{\sqrt{6}}{6}\lambda \right). \quad (28)$$

This system can be reduced in one dimension using the relation $x^2 + y^2 = 1$:

$$\frac{dx}{d\tau} = f(x) := -3(1-x^2) \left(x - \frac{\sqrt{6}}{6}\lambda \right). \quad (29)$$

Equation (29) is integrable and leads to

$$\tau := \ln a = c_1 + \ln \left[(1-x)^{\frac{1}{6-\sqrt{6}\lambda}} (x+1)^{\frac{1}{\sqrt{6}\lambda+6}} \left| 6x - \sqrt{6}\lambda \right|^{\frac{2}{\lambda^2-6}} \right]. \quad (30)$$

That is, the scaling factor a can be expressed as a function of x by

$$a(x) = e^{c_1} (1-x)^{\frac{1}{6-\sqrt{6}\lambda}} (x+1)^{\frac{1}{\sqrt{6}\lambda+6}} \left| 6x - \sqrt{6}\lambda \right|^{\frac{2}{\lambda^2-6}}, \quad x \in [-1, 1]. \quad (31)$$

Additionally,

$$\frac{d\phi}{dx} = \frac{\frac{d\phi}{d\tau}}{\frac{dx}{d\tau}} = -\frac{2\sqrt{6}x}{(1-x^2)(6x-\sqrt{6}\lambda)}, \quad (32)$$

from which it follows by integration that:

$$\phi(x) = c_2 + \ln \left[(1-x)^{\frac{1}{\sqrt{6}-\lambda}} (x+1)^{-\frac{1}{\lambda+\sqrt{6}}} \left| 6x - \sqrt{6}\lambda \right|^{\frac{2\lambda}{\lambda^2-6}} \right]. \quad (33)$$

On the other hand, from (27) with $V(\phi) = V_0 e^{-\lambda\phi}$ holds:

$$H(x) = e^{-\frac{c_2\lambda}{2}} \sqrt{\frac{V_0}{3-3x^2}} (1-x)^{\frac{1}{\sqrt{6}-\lambda}} (x+1)^{-\frac{1}{\lambda+\sqrt{6}}} \left| 6x - \sqrt{6}\lambda \right|^{\frac{2\lambda}{\lambda^2-6}}. \quad (34)$$

Finally, it follows that:

$$\frac{dt}{dx} = \frac{2\sqrt{3}e^{\frac{c_2\lambda}{2}} (1-x)^{\frac{1}{\lambda-\sqrt{6}}} (x+1)^{\frac{1}{\lambda+\sqrt{6}}} \left| 6x - \sqrt{6}\lambda \right|^{-\frac{2\lambda}{\lambda^2-6}}}{\sqrt{V_0(1-x^2)} (6x-\sqrt{6}\lambda)}, \quad (35)$$




Figure 1. $a(x)$ given by (31) with $c_1 = 0$, $\phi(x)$ given by (33) with $c_2 = 0$ and $H(x)$ given by (34) with $c_2 = 0$, $V_0 = 1$, for several choices of λ .

$$t = 2\sqrt{3}e^{\frac{c_2\lambda}{2}} \int \frac{(1-x)^{\frac{1}{\lambda-\sqrt{6}}}(x+1)^{\frac{1}{\lambda+\sqrt{6}}} |6x - \sqrt{6}\lambda|^{-\frac{2\lambda}{\lambda^2-6}}}{\sqrt{V_0(1-x^2)} (6x - \sqrt{6}\lambda)} dx. \quad (36)$$

Figure 1 show $a(x)$, given by (31) for $c_1 = 0$, represented by the solid thick red line; $\phi(x)$ given by (33) for $c_2 = 0$, represented by solid thinner blue line; and $H(x)$, given by (34) for $c_2 = 0, V_0 = 1$, represented by the dashed black line for several choices of λ . The cosmological time t is related with x by the quadrature (36).

Figure 1(a) shows:

$$\lim_{x \rightarrow -1} (a(x), \phi(x), H(x)) = (0, +\infty, +\infty), \quad (37)$$

$$\lim_{x \rightarrow 1} (a(x), \phi(x), H(x)) = (0, -\infty, +\infty), \quad (38)$$

$$\lim_{x \rightarrow \frac{\sqrt{6}\lambda}{6}} (a(x), \phi(x), H(x)) = (+\infty, -\infty, 0). \quad (39)$$

Figure 1(b) shows:

$$\lim_{x \rightarrow -1} (a(x), \phi(x), H(x)) = (0, +\infty, +\infty), \quad (40)$$

$$\lim_{x \rightarrow 1} (a(x), \phi(x), H(x)) = (0, -\infty, +\infty), \quad (41)$$

$$\lim_{x \rightarrow 0} (a(x), \phi(x), H(x)) = \left(\infty, 0, \sqrt{\frac{V_0}{3}} \right). \quad (42)$$

Figure 1(c) shows:

$$\lim_{x \rightarrow -1} (a(x), \phi(x), H(x)) = (0, +\infty, +\infty), \quad (43)$$

$$\lim_{x \rightarrow 1} (a(x), \phi(x), H(x)) = (0, -\infty, 0), \quad (44)$$

$$\lim_{x \rightarrow \frac{\sqrt{6}\lambda}{6}} (a(x), \phi(x), H(x)) = (+\infty, +\infty, +\infty). \quad (45)$$

Figure 1(d) shows:

$$\lim_{x \rightarrow -1} (a(x), \phi(x), H(x)) = (0, 0, +\infty), \quad (46)$$

$$\lim_{x \rightarrow 1} (a(x), \phi(x), H(x)) = (0, -\infty, 0), \quad (47)$$

$$\lim_{x \rightarrow \frac{\sqrt{6}\lambda}{6}} (a(x), \phi(x), H(x)) = (+\infty, +\infty, +\infty). \quad (48)$$

The above limits show that at the equilibrium points the scalar field eventually diverges and the Hubble parameter can be zero, infinity or it tends to a finite non-zero number. But, when the scalar field is divergent or H goes to zero, the Hubble-normalization fails. Therefore, although this procedure is well-suited to investigate local stability features of the equilibrium points, it does not provide a global description of the phase space when generically ϕ diverges or when $H \rightarrow 0$. In such case the method fails. For this reason, a new global systems analysis motivated by Alho & Uggla's approach [98] is presented.

Using

$$T = \frac{m}{m + H}, \quad \theta = \tan^{-1} \left(\frac{\dot{\phi}}{\sqrt{2V(\phi)}} \right), \quad (49)$$

Label	(T, θ)	Existence	Stability
P_1	$(0, -\frac{\pi}{2} + 2c_1\pi)$	$\forall \lambda$	Saddle for $\lambda < -\sqrt{6}$. Non-hyperbolic for $\lambda = -\sqrt{6}$. Source for $\lambda > -\sqrt{6}$.
P_2	$(0, \frac{\pi}{2} + 2c_1\pi)$	$\forall \lambda$	Saddle for $\lambda > \sqrt{6}$. Non-hyperbolic for $\lambda = \sqrt{6}$. Source for $\lambda < \sqrt{6}$.
P_3	$(0, \arcsin(\frac{\lambda}{\sqrt{6}}))$	$-\sqrt{6} \leq \lambda \leq \sqrt{6}$	Non-hyperbolic for $\lambda \in \{-\sqrt{6}, 0, \sqrt{6}\}$. Saddle for $-\sqrt{6} < \lambda < 0$ or $0 < \lambda < \sqrt{6}$.
P_4	$(1, -\frac{\pi}{2} + 2c_1\pi)$	$\forall \lambda$	Sink for $\lambda < -\sqrt{6}$. Non-hyperbolic for $\lambda = -\sqrt{6}$. Saddle for $\lambda > -\sqrt{6}$.
P_5	$(1, \frac{\pi}{2} + 2c_1\pi)$	$\forall \lambda$	Sink for $\lambda > \sqrt{6}$. Non-hyperbolic for $\lambda = \sqrt{6}$. Saddle for $\lambda < \sqrt{6}$.
P_6	$(1, \arcsin(\frac{\lambda}{\sqrt{6}}))$	$-\sqrt{6} \leq \lambda \leq \sqrt{6}$	Non-hyperbolic for $\lambda \in \{-\sqrt{6}, 0, \sqrt{6}\}$. Sink for $-\sqrt{6} < \lambda < 0$ or $0 < \lambda < \sqrt{6}$.

Table 1. Existence conditions and stability conditions of the equilibrium points of equations (50), $c_1 \in \mathbb{Z}$.

the following unconstrained dynamical system is deduced:

$$\frac{dT}{d\tau} = 3(1-T)T \sin^2(\theta), \quad \frac{d\theta}{d\tau} = \frac{1}{2} \cos(\theta) (\sqrt{6}\lambda - 6 \sin(\theta)), \quad (50)$$

defined in the finite cylinder \mathbf{S} with boundaries $T = 0$ and $T = 1$.

The variable T is suitable for global analysis [98], due to

$$\left. \frac{dT}{d\tau} \right|_{\sin \theta=0} = 0, \quad \left. \frac{d^2 T}{d\tau^2} \right|_{\sin \theta=0} = 0, \quad \left. \frac{d^3 T}{d\tau^3} \right|_{\sin \theta=0} = 9\lambda^2(1-T)T. \quad (51)$$

From the first equation of (50) and equation (51), T is a monotonically increasing function on \mathbf{S} . As a consequence, all orbits originate from the invariant subset $T = 0$ (which contains the α -limit), which is classically related to the initial singularity with $H \rightarrow \infty$, and ends on the invariant boundary subset $T = 1$, which corresponds asymptotically to $H = 0$.

The equilibrium points of equations (50) are the following:

- (i) $P_1 : (T, \theta) = (0, -\frac{\pi}{2} + 2c_1\pi)$, $c_1 \in \mathbb{Z}$, with eigenvalues $\{3, \sqrt{\frac{3}{2}}\lambda + 3\}$. It represents a kinetic dominated solution with $H \rightarrow \infty$. The stability conditions of P_1 are the following:
 - i) Saddle for $\lambda < -\sqrt{6}$.
 - ii) Non-hyperbolic for $\lambda = -\sqrt{6}$.
 - iii) Source for $\lambda > -\sqrt{6}$.




Figure 2. Unwrapped solution space (left panel) - Projection over the cylinder S (right panel) solution space of system (50) for different values of λ .

- (ii) $P_2 : (T, \theta) = (0, \frac{\pi}{2} + 2c_1\pi)$, $c_1 \in \mathbb{Z}$ with eigenvalues $\left\{3, 3 - \sqrt{\frac{3}{2}}\lambda\right\}$. It represents a kinetic dominated solution with $H \rightarrow \infty$. The stability conditions of P_2 are the following:
- i) Source for $\lambda < \sqrt{6}$.
 - ii) Non-hyperbolic for $\lambda = \sqrt{6}$.
 - iii) Saddle for $\lambda > \sqrt{6}$.
- (iii) $P_3 : (T, \theta) = \left(0, \arcsin\left(\frac{\lambda}{\sqrt{6}}\right)\right)$ with eigenvalues $\left\{\frac{\lambda^2}{2}, \frac{1}{2}(\lambda^2 - 6)\right\}$. It represents a scalar field dominated solution with $H \rightarrow \infty$. The stability conditions of P_3 are the following:
- i) Non-hyperbolic for $\lambda \in \{-\sqrt{6}, 0, \sqrt{6}\}$.
 - ii) Saddle for $-\sqrt{6} < \lambda < 0$ or $0 < \lambda < \sqrt{6}$.
- (iv) $P_4 : (T, \theta) = \left(1, -\frac{\pi}{2} + 2c_1\pi\right)$, $c_1 \in \mathbb{Z}$ with eigenvalues $\left\{-3, \sqrt{\frac{3}{2}}\lambda + 3\right\}$. It represents a kinetic dominated solution with $H \rightarrow 0$. The stability conditions of P_4 are the following:
- i) Sink for $\lambda < -\sqrt{6}$.
 - ii) Non-hyperbolic for $\lambda = -\sqrt{6}$.
 - iii) Saddle for $\lambda > -\sqrt{6}$.
- (v) $P_5 : (T, \theta) = \left(1, \frac{\pi}{2} + 2c_1\pi\right)$, $c_1 \in \mathbb{Z}$ with eigenvalues $\left\{-3, 3 - \sqrt{\frac{3}{2}}\lambda\right\}$. It represents a kinetic dominated solution with $H \rightarrow 0$. The stability conditions of P_5 are the following:
- i) It is a sink for $\lambda > \sqrt{6}$.
 - ii) Non-hyperbolic for $\lambda = \sqrt{6}$.
 - iii) Saddle for $\lambda < \sqrt{6}$.
- (vi) $P_6 : (T, \theta) = \left(1, \arcsin\left(\frac{\lambda}{\sqrt{6}}\right)\right)$ with eigenvalues $\left\{-\frac{\lambda^2}{2}, \frac{1}{2}(\lambda^2 - 6)\right\}$. It represents a scalar field dominated solution with $H \rightarrow 0$. The stability conditions of P_6 are the following:
- i) Non-hyperbolic for $\lambda \in \{-\sqrt{6}, 0, \sqrt{6}\}$.
 - ii) Sink for $-\sqrt{6} < \lambda < 0$ or $0 < \lambda < \sqrt{6}$.

In table 1, the existence conditions and stability conditions of the equilibrium points of equations (50) are summarized.

In the Figure 2, some orbits of the flow of equations (50) (left panel) and a projection over the cylinder S (right panel) for different values of λ are presented.

5. Example: a scalar field model with E -potential $V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3\alpha}}\phi}\right)^{2n}$ in vacuum

Considering the E -model with potential $V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3\alpha}}\phi}\right)^{2n}$. This is a nonnegative potential with a single minimum located at $(\phi, V(\phi)) = (0, 0)$. Therefore, the model admits a Minkowski solution represented by the equilibrium point $(H, \dot{\phi}, \phi) = (0, 0, 0)$. The potential has a plateau $V = V_0$, when $\phi \rightarrow +\infty$, while $V \sim V_0 e^{-2n\sqrt{\frac{2}{3\alpha}}\phi}$ as $\phi \rightarrow -\infty$ [101]. At small ϕ the E -potential behaves as ϕ^{2n} .

Defining the Hubble normalized variables:

$$x = \frac{\dot{\phi}}{\sqrt{6}H}, \quad Y = \left(\frac{V(\phi)}{3H^2}\right)^{\frac{1}{2n}} = \tilde{T} \left(1 - e^{-\sqrt{\frac{2}{3\alpha}}\phi}\right), \quad \tilde{T} = \left[\frac{V_0}{3H^2}\right]^{\frac{1}{2n}}, \quad (52)$$

the dynamical system:

$$\frac{dx}{dN} = 6\mu Y^{2n-1}(Y - \tilde{T}) + (q - 2)x, \quad (53a)$$

$$\frac{dY}{dN} = \frac{Y(-6\mu x + q + 1) + 6\mu x \tilde{T}}{n}, \quad (53b)$$

$$\frac{d\tilde{T}}{dN} = \frac{(q + 1)\tilde{T}}{n}, \quad (53c)$$

where $\mu = \frac{n}{3\sqrt{\alpha}}$, and $q = 2 - 3Y^{2n}$, is studied. This system has been extensively studied in [101] in the context of a canonical scalar field cosmology, and it was extended in [59] to Hořava-Lifshitz cosmology. Now, the most relevant features of the solution space according to [59] will be discussed. It can be easily proven that \tilde{T} is monotonically increasing towards the future and decreasing towards the past. The phase space is limited to the past by the invariant subset $\tilde{T} = 0$ for $Y \leq 0$, and by $\tilde{T} - Y = 0$ for $Y \geq 0$. The state space is bounded when $\tilde{T} > 0, \tilde{T} - Y > 0$. The two past boundaries are intersected at the two massless scalar field points $M_{\pm} = (\Sigma, Y) = (\pm 1, 0)$. The subset $\tilde{T} - Y = 0$, on the other hand, is divided in two disconnected regions separated by the de Sitter equilibrium point $dS = (\tilde{T}, Y) = (1, 1)$.

Introducing the complementary global transformation (which for $n = 1$ is exactly the definition of θ in Section 4):

$$x = F(\theta) \sin(\theta), \quad Y = \cos(\theta), \quad F(\theta) = \sqrt{\frac{1 - \cos^{2n}(\theta)}{1 - \cos^2(\theta)}} = \sqrt{\sum_{k=0}^{n-1} \cos^{2k}(\theta)}, \quad (54)$$

the following regular unconstrained 2D dynamical system:

$$\frac{d\theta}{dN} = -\frac{6\mu F(\theta)(\tilde{T} - \cos(\theta))}{n} - \frac{3F(\theta)^2 \sin(2\theta)}{2n}, \quad (55a)$$

$$\frac{d\tilde{T}}{dN} = \frac{3\tilde{T}(1 - \cos^{2n}(\theta))}{n}, \quad (55b)$$

is obtained, and the deceleration parameter becomes

$$q = 2 - 3\cos^{2n}(\theta). \quad (56)$$

In Figure 3, the unwrapped solution space (left panel) and the projection over the cylinder S (right panel) of the system (57a), (57b) for some values of n, μ, α are represented. This plot clearly shows that the future boundary is $T = 1$ which corresponds to $H = 0$, and the final state is the Minkowski point given by a limit cycle. Introducing the new compact variable $T = \frac{\tilde{T}}{1+\tilde{T}}$ and the new time derivative $\frac{d\bar{\tau}}{d\ln a} = 1 + \tilde{T} = (1 - T)^{-1}$; the regular system:

$$\frac{d\theta}{d\bar{\tau}} = \frac{3(T - 1)F(\theta)^2 \sin(\theta) \cos(\theta)}{n} - \frac{6\mu F(\theta)((T - 1) \cos(\theta) + T)}{n}, \quad (57a)$$

$$\frac{dT}{d\bar{\tau}} = -\frac{3(T - 1)^2 T (\cos^{2n}(\theta) - 1)}{n}, \quad (57b)$$

is obtained. The past boundary is attached to the phase-space, and in the new variables (θ, T) is defined by $\{T = 0, \cos(\theta) \leq 0\} \cup \{T - (1 - T) \cos \theta = 0, \cos(\theta) > 0\}$. It is also included the future boundary $T = 1$ which corresponds to $H = 0$, and the final state is the Minkowski point. The region $\{T - (1 - T) \cos \theta < 0, \cos(\theta) > 0\}$ is forbidden.




Figure 3. Unwrapped solution space (left panel). Projection over the cylinder S (right panel) of system (57a), (57b) for some values of n, μ, α .

The equilibrium points of (57a), (57b) are given by:

M_{\pm} : $\tilde{T} = T = 0; x = \pm 1, Y = 0; \theta = \pm \frac{\pi}{2} + 2k\pi, k = 0, 1, 2, \dots$. They are massless scalar field solutions. They are saddle and source, respectively, as it is confirmed in Figure 3.

dS : $\tilde{T} = 1, T = \frac{1}{2}; x = 0, Y = 1; \theta = 2k\pi, k = 0, 1, 2, \dots$. They are de Sitter solutions.

PL : $\tilde{T} = T = 0; x = 2\mu; Y = -(1 - 4\mu^2)^{\frac{1}{2n}}; \theta = \pm \arccos Y$. It exists for $\mu < 1/2$, and corresponds to a powerlaw selfsimilar solution for the exponential potential.

6. Example: a scalar-field cosmology with generalized harmonic potential

$$V(\phi) = \mu^3 \left[\frac{\phi^2}{\mu} + bf \cos \left(\delta + \frac{\phi}{f} \right) \right], b \neq 0 \text{ in vacuum}$$

In this section, the asymptotic analysis as $\phi \rightarrow \infty$ of a scalar-field cosmology with generalized harmonic potential

$$V(\phi) = \mu^3 \left[\frac{\phi^2}{\mu} + bf \cos \left(\delta + \frac{\phi}{f} \right) \right], \quad b \neq 0, \quad (58)$$

in a vacuum is presented for a flat FLRW model. Let be $N = 0, M = 0, W_\chi \equiv 0, \Omega_m \equiv 0, \Omega_0 \equiv 0, \rho_m \equiv 0, G_0 \equiv 0$ and $\chi \equiv 1$.

Furthermore,

$$W_V(\phi) = \frac{2\phi - b\mu \sin \left(\delta + \frac{\phi}{f} \right)}{bf\mu \cos \left(\delta + \frac{\phi}{f} \right) + \phi^2}. \quad (59)$$

6.1. Analysis as $\phi \rightarrow \infty$

In this section the dynamics as $\phi \rightarrow \infty$ of a scalar-field cosmology with generalized harmonic potential (58) in a vacuum is analyzed.

Defining the transformation:

$$\varphi = h(\phi) = \varphi = \left(\delta + \frac{\phi}{f} \right)^{-\frac{1}{4}}, \quad (60)$$

it follows that $V(\phi)$ is 2 WBI with exponential order $N = 0$.

$$\bar{W}_V(\varphi) = \begin{cases} \frac{-b\mu\varphi^8 \sin \left(\frac{1}{\varphi^4} \right) - 2\delta f\varphi^8 + 2f\varphi^4}{f \left(b\mu\varphi^8 \cos \left(\frac{1}{\varphi^4} \right) + f(\delta\varphi^4 - 1)^2 \right)}, & \varphi > 0 \\ 0, & \varphi = 0 \end{cases}, \quad (61)$$

$$\bar{h}'(\varphi) = \begin{cases} -\frac{\varphi^5}{4f}, & \varphi > 0, \\ 0, & \varphi = 0 \end{cases}, \quad (62)$$

that satisfy the conditions (ii) and (iii) of Definition 2. Hence, the dynamical system:

$$\frac{dT}{d\tau} = 3(1 - T)T \sin^2(\theta), \quad (63a)$$

$$\frac{d\theta}{d\tau} = -\frac{1}{2} \cos(\theta) \left(6 \sin(\theta) + \sqrt{6} \bar{W}_V(\varphi) \right), \quad (63b)$$

$$\frac{d\varphi}{d\tau} = \sqrt{6} \bar{h}'(\varphi) \sin(\theta), \quad (63c)$$

is obtained, where we have used the new time variable $\tau = \ln a$, defined on a phase space which consists of the vector product $\mathbf{S} \times J$ of the finite cylinder \mathbf{S} with boundaries $T = 0$ and $T = 1$ with the interval $J = \left[0, \left(\delta + \frac{\phi_0}{f}\right)^{-\frac{1}{4}}\right]$. The variable T is suitable for global analysis [98], due to:

$$\begin{aligned} \frac{dT}{d\tau} \Big|_{\sin \theta=0} &= 0, \quad \frac{d^2T}{d\tau^2} \Big|_{\sin \theta=0} = 0, \\ \frac{d^3T}{d\tau^3} \Big|_{\sin \theta=0} &= \frac{9(1-T)Tv^8(b\mu v^4 \sin(\frac{1}{v^4}) + 2f(\delta v^4 - 1))^2}{f^2(b\mu v^8 \cos(\frac{1}{v^4}) + f(\delta v^4 - 1)^2)}. \end{aligned} \quad (64)$$

From equation (63a) and equation (64), T is a monotonically increasing function on $\mathbf{S} \times J$. As a consequence, all orbits are originated from the invariant subset $T = 0$ (which contains the α -limit), which is classically related to the initial singularity with $H \rightarrow \infty$ and ends at the invariant boundary subset $T = 1$, which corresponds to the asymptotically $H = 0$.

The curves of the equilibrium points of (63) are the following:




Figure 4. Evaluation of $(\cos \theta, \sin \theta, T)$ at some orbits of the system (63) for $(b, f, \delta, \mu) = (0.1, 0.33, 0, 1)$. In the plot, the points (iii) and (iv) are sources; the points (v) and (vi) are saddles. The vertical plane represents the invariant set $\{(T, \theta, \varphi) : \varphi = 0, \sin(\theta) = 0\}$. The vertical sides (i) and (ii) of this rectangle are the local sinks.

- (i) $(T, \theta, \varphi) = (T_c, 2n\pi, 0)$ with eigenvalues $\{-3, 0, 0\}$. It is non-hyperbolic.
- (ii) $(T, \theta, \varphi) = (T_c, \pi + 2n\pi, 0)$ with eigenvalues $\{-3, 0, 0\}$. It is non-hyperbolic.
- (iii) $(T, \theta, \varphi) = (0, -\frac{\pi}{2} + 2n\pi, 0)$ with eigenvalues $\{3, 3, 0\}$. It is non-hyperbolic.
- (iv) $(T, \theta, \varphi) = (0, \frac{\pi}{2} + 2n\pi, 0)$ with eigenvalues $\{3, 3, 0\}$. It is non-hyperbolic.
- (v) $(T, \theta, \varphi) = (1, -\frac{\pi}{2} + 2n\pi, 0)$ with eigenvalues $\{-3, 3, 0\}$. It behaves as saddle.
- (vi) $(T, \theta, \varphi) = (1, \frac{\pi}{2} + 2n\pi, 0)$ with eigenvalues $\{-3, 3, 0\}$. It behaves as saddle.

In Figure 4, the functions $(\cos \theta, \sin \theta, T)$ are evaluated at some orbits of the system (63) for $(b, f, \delta, \mu) = (0.1, 0.33, 0, 1)$. In the plot the points (iii) and (iv), which are sources, and the points (v) and (vi), which are saddles, are represented. The vertical plane represents the invariant set $\{(T, \theta, \varphi) : \varphi = 0, \sin(\theta) = 0\}$. The vertical sides (i) and (ii) of this rectangle are the local sinks.

6.2. Oscillating regime.

In the reference [118], oscillating scalar field models with potential $\frac{1}{2}\phi^2$ and potentials $\frac{1}{2}\phi^2 + W(\phi)$ with W smooth and $W(\phi) = o(\phi^3)$ were studied. Improved asymptotic expansions for the solution in homogeneous and isotropic spaces were derived. Various generalizations for non-linear massive scalar fields, k -essence models and $f(R)$ -gravity were obtained. In this section the potential $V(\phi) = \mu^3 \left[\frac{\phi^2}{\mu} + bf \cos \left(\delta + \frac{\phi}{f} \right) \right]$, $b \neq 0$ is investigated looking for oscillatory behavior, as expected in numerical investigations. Asymptotic expansions are derived as well. Noticing that the cosine corrections are $O(bf\mu^3)$; therefore, they do not fall in the potential class studied by [118].

The pair

$$\left(\frac{\sqrt{2}\mu\phi}{\sqrt{\dot{\phi}^2 + 2\mu\phi^2}}, \quad \frac{\dot{\phi}}{\sqrt{\dot{\phi}^2 + 2\mu\phi^2}} \right) \quad (65)$$

defines a function that depends on the time variable t , with values in the unit circle. Hence, an angular function $\vartheta(t)$, that is unique under identification module 2π , can be defined by

$$\vartheta = \tan^{-1} \left(\frac{\dot{\phi}}{\sqrt{2}\mu\phi} \right), \quad (66)$$

together with the radial variable

$$r = \sqrt{\dot{\phi}^2 + 2\mu^2\phi^2}, \quad (67)$$

which satisfy

$$-bf\mu^3 \cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) + 3H^2 - \frac{r^2}{2} = 0. \quad (68)$$

The inverse transformation $(r, \vartheta) \mapsto (\phi, \dot{\phi})$ is

$$\phi = \frac{r \cos(\vartheta)}{\sqrt{2}\mu}, \quad \dot{\phi} = r \sin(\vartheta). \quad (69)$$

For expanding universes ($H > 0$) the following holds:

$$\dot{r} = -\sqrt{\frac{3}{2}}r \sin^2(\vartheta) \sqrt{2bf\mu^3 \cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) + r^2} + b\mu^3 \sin(\vartheta) \sin \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right), \quad (70a)$$

$$\begin{aligned} \dot{\vartheta} &= -\sqrt{2}\mu - \sqrt{\frac{3}{2}} \sin(\vartheta) \cos(\vartheta) \sqrt{2bf\mu^3 \cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) + r^2} \\ &\quad + \frac{b\mu^3 \cos(\vartheta) \sin \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right)}{r}. \end{aligned} \quad (70b)$$

As $b \rightarrow 0$ the last equations reduce to:

$$\dot{r} = -\sqrt{\frac{3}{2}}r^2 \sin^2(\vartheta), \quad \dot{\vartheta} = -\sqrt{2}\mu - \sqrt{\frac{3}{2}}r \sin(\vartheta) \cos(\vartheta). \quad (71)$$

The solutions of the limiting equations admit the asymptotic expansions [118]

$$r(t) = \frac{4}{\sqrt{6t}} + O(t^{-2} \ln t), \quad \vartheta(t) = -\sqrt{2}\mu t + O(\ln t). \quad (72)$$

Hence,

$$\phi(t) = \frac{4 \cos t}{\sqrt{6t}} + O(t^{-2} \ln t), \quad \dot{\phi}(t) = \frac{4 \sin t}{\sqrt{6t}} + O(t^{-2} \ln t). \quad (73)$$

These expansions can be improved to the order $O(t^{-3} \ln t)$ as in [118]. Here, asymptotic expansions of the full problem ($b \neq 0$) are derived using averaging techniques.

Note that:

$$\dot{r} = b\mu^3 \sin(\delta) \sin(\vartheta) + \left(\frac{b\mu^2 \cos(\delta) \cos(\vartheta) \sin(\vartheta)}{\sqrt{2}f} - \sqrt{3} \sqrt{bf\mu^3 \cos(\delta)} \sin^2(\vartheta) \right) r + O(r^2), \quad (74a)$$

$$\begin{aligned} \dot{\vartheta} + \sqrt{2}\mu &= \frac{b\mu^3 \sin(\delta) \cos(\vartheta)}{r} + \left(\frac{b\mu^2 \cos(\delta) \cos^2(\vartheta)}{\sqrt{2}f} - \sqrt{3} \sin(\vartheta) \cos(\vartheta) \sqrt{bf\mu^3 \cos(\delta)} \right) \\ &\quad \frac{r \left(\sqrt{6}f \tan(\delta) \sin(\vartheta) \cos^2(\vartheta) \sqrt{bf\mu^3 \cos(\delta)} - b\mu^2 \sin(\delta) \cos^3(\vartheta) \right)}{4f^2\mu} + O(r^2). \end{aligned} \quad (74b)$$

In order to obtain an approximated solution near the oscillatory regime, the average with respect to ϑ over any orbit of period 2π given by

$$\langle f \rangle = \frac{1}{2\pi} \int_c^{c+2\pi} f(\vartheta) d\vartheta, \quad \cos(c) \geq 0, \quad (75)$$

is applied to $f = (\dot{r}, \dot{\vartheta})$, leading to

$$\dot{r} = -\frac{1}{2} \sqrt{3}r \sqrt{bf\mu^3 \cos(\delta)}, \quad \dot{\vartheta} = \frac{b\mu^2 \cos(\delta)}{2\sqrt{2}f} - \sqrt{2}\mu. \quad (76)$$

The averaged equations have solutions

$$r(t) = r_0 e^{-\frac{1}{2}\sqrt{3}t\sqrt{bf\mu^3 \cos(\delta)}}, \quad \vartheta(t) = \left(\frac{b\mu^2 \cos(\delta)}{2\sqrt{2}f} - \sqrt{2}\mu \right) t + \vartheta_0. \quad (77)$$

Introducing along with ϑ and r , the new variable

$$\varepsilon = \frac{H}{\mu + H}, \quad (78)$$

with inverse

$$H = \frac{\mu\varepsilon}{1 - \varepsilon}, \quad (79)$$

satisfying:

$$-bf\mu^3(\varepsilon - 1)^2 \cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) - \frac{1}{2}r^2(\varepsilon - 1)^2 + 3\mu^2\varepsilon^2 = 0, \quad (80)$$




Figure 5. Phase portrait of equations (74) (left panel). Projection over the cylinder S (right panel) for $(b, f, \delta) = (0.1, 0.33, 0)$ and different values of μ .

along with the time derivative $\hat{\tau}$ given by

$$\frac{d\hat{\tau}}{dt} := \mu + H, \quad (81)$$

the following equations hold:

$$\varepsilon' = -\frac{r^2(1-\varepsilon)^3 \sin^2(\vartheta)}{2\mu^2}, \quad (82a)$$

$$r' = -3r\varepsilon \sin^2(\vartheta) + b\mu^2(1-\varepsilon) \sin(\vartheta) \sin\left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu}\right) \quad (82b)$$

$$\vartheta' = -\sqrt{2}(1-\varepsilon) - 3\varepsilon \sin(\vartheta) \cos(\vartheta) + \frac{b\mu^2(1-\varepsilon) \cos(\vartheta) \sin\left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu}\right)}{r}. \quad (82c)$$

To obtain an approximated solution near the oscillatory regime the average with respect to ϑ




Figure 6. Phase portrait of equations (74) (left panel). Projection over the cylinder S (right panel) for $(b, f, \delta) = (0.99, 0.09, 0)$ and different values of μ .

over any orbit of period 2π , $\langle (\varepsilon', r', \vartheta') \rangle$ is taken, leading to:

$$\varepsilon' = \frac{r^2(\varepsilon - 1)^3}{4\mu^2}, \quad (83a)$$

$$r' = -\frac{3r\varepsilon}{2}, \quad (83b)$$

$$\vartheta' = \sqrt{2}(\varepsilon - 1) - \left\langle \frac{b\mu^2(\varepsilon - 1) \cos(\vartheta) \sin\left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu}\right)}{r} \right\rangle. \quad (83c)$$

But, as $r \rightarrow 0$,

$$\begin{aligned} & \frac{1}{2\pi} \int_c^{c+2\pi} \frac{b\mu^2(\varepsilon - 1) \cos(\vartheta) \sin\left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu}\right)}{r} d\vartheta \\ & \sim \frac{b\mu(\varepsilon - 1) \cos(\delta)}{2\sqrt{2}f} - \frac{r^2(b(\varepsilon - 1) \cos(\delta))}{32(\sqrt{2}f^3\mu)} + O(r^3). \end{aligned} \quad (84)$$

Finally, it follows that:

$$\varepsilon' = -\frac{r^2(1-\varepsilon)^3}{4\mu^2}, \quad (85a)$$

$$r' = -\frac{3r\varepsilon}{2}, \quad (85b)$$

$$\vartheta' = \left[-\sqrt{2} + \frac{b\mu \cos(\delta)}{2\sqrt{2}f} - \frac{r^2(b \cos(\delta))}{32(\sqrt{2}f^3\mu)} \right] (1-\varepsilon), \quad (85c)$$

with the averaged constraint:

$$\mu^2 (3\varepsilon^2 - bf\mu(1-\varepsilon)^2 \cos(\delta)) + \frac{r^2(1-\varepsilon)^2(b\mu \cos(\delta) - 4f)}{8f} = 0, \quad (86)$$

as $r \rightarrow 0$.

The above system is integrable yielding:

$$r(\varepsilon) = \frac{\sqrt{2}\sqrt{c_1(\varepsilon-1)^2 + \mu^2(6\varepsilon-3)}}{1-\varepsilon}, \quad (87a)$$

$$\vartheta(\varepsilon) = \frac{\mu \tanh^{-1} \left(\frac{c_1(\varepsilon-1)+3\mu^2}{\mu\sqrt{9\mu^2-3c_1}} \right) (b\mu \cos(\delta) - 4f)}{f\sqrt{18\mu^2-6c_1}} + \frac{b\mu \cos(\delta)}{8\sqrt{2}f^3(1-\varepsilon)} + c_2, \quad (87b)$$

and

$$3(t-t_0) = \ln \left(\frac{(1-\varepsilon)^2 \left(\frac{3\mu\sqrt{3\mu^2-c_1}+\sqrt{3}c_1\varepsilon-\sqrt{3}c_1+3\sqrt{3}\mu^2}{3\mu\sqrt{3\mu^2-c_1}-\sqrt{3}c_1\varepsilon+\sqrt{3}c_1-3\sqrt{3}\mu^2} \right) \frac{\mu}{\sqrt{\mu^2-\frac{c_1}{3}}} }{c_1(\varepsilon-1)^2 + \mu^2(6\varepsilon-3)} \right) \\ \sim \ln \left(\frac{\left(\frac{2\mu(\sqrt{9\mu^2-3c_1}+3\mu)-c_1}{c_1} \right) \frac{\mu}{\sqrt{\mu^2-\frac{c_1}{3}}}}{c_1-3\mu^2} \right) - \frac{6\mu^2\varepsilon}{c_1-3\mu^2} + O(\varepsilon^2), \quad (87c)$$

as $\varepsilon \rightarrow 0$.

The phase portrait of equations (74) (left panel) and the projection over the cylinder **S** (right panel) for $(b, f, \delta) = (0.1, 0.33, 0)$ for different values of μ are presented in figure 5. Figure 6 shows the phase portrait of equations (74) (left panel) and the projection over the cylinder **S** (right panel) for $(b, f, \delta) = (0.99, 0.09, 0)$ and different values of μ . The plots show the oscillatory behavior of the solutions.

7. Example: a scalar-field cosmology with generalized harmonic potential

$$V(\phi) = \mu^3 \left[bf \left(\cos(\delta) - \cos \left(\delta + \frac{\phi}{f} \right) \right) + \frac{\phi^2}{\mu} \right], b \neq 0, \text{in vacuum}$$

In this section the asymptotic analysis as $\phi \rightarrow \infty$ of a scalar-field cosmology with generalized harmonic potential:

$$V(\phi) = \mu^3 \left[bf \left(\cos(\delta) - \cos \left(\delta + \frac{\phi}{f} \right) \right) + \frac{\phi^2}{\mu} \right], \quad b \neq 0, \quad (88)$$

in a vacuum for a flat FLRW model is performed. Let be $N = 0, M = 0, W_\chi \equiv 0, \Omega_m \equiv 0, \Omega_0 \equiv 0, \rho_m \equiv 0, G_0 \equiv 0$ and $\chi \equiv 1$.

Furthermore,

$$W_V(\phi) = \frac{b\mu \sin\left(\delta + \frac{\phi}{f}\right) + 2\phi}{bf\mu \left(\cos(\delta) - \cos\left(\delta + \frac{\phi}{f}\right)\right) + \phi^2}. \quad (89)$$

7.1. Analysis as $\phi \rightarrow \infty$

In this section, the dynamics as $\phi \rightarrow \infty$ of a scalar-field cosmology with generalized harmonic potential (88) in a vacuum is analyzed.

Using again the transformation (60) and the new time variable $\tau = \ln a$, a system of the form (63) holds with the definitions:

$$\bar{W}_V(\varphi) = \begin{cases} \frac{b\mu\varphi^8 \sin\left(\frac{1}{\varphi^4}\right) - 2\delta f\varphi^8 + 2f\varphi^4}{f\left(b\mu\varphi^8 \left(\cos(\delta) - \cos\left(\frac{1}{\varphi^4}\right)\right) + f(\delta\varphi^4 - 1)^2\right)}, & \varphi > 0, \\ 0, & \varphi = 0 \end{cases}, \quad (90)$$

$$\bar{h}'(\varphi) = \begin{cases} -\frac{\varphi^5}{4f}, & \varphi > 0, \\ 0, & \varphi = 0 \end{cases}, \quad (91)$$

that satisfy the conditions (ii) and (iii) of Definition 2. The phase space is the vector product $\mathbf{S} \times J$ of the finite cylinder \mathbf{S} with boundaries $T = 0$ and $T = 1$ with the interval $J = \left[0, \left(\delta + \frac{\phi_0}{f}\right)^{-\frac{1}{4}}\right]$. The variable T is suitable for global analysis [98], due to

$$\begin{aligned} \frac{dT}{d\tau} \Big|_{\sin \theta=0} &= 0, \quad \frac{d^2T}{d\tau^2} \Big|_{\sin \theta=0} = 0, \\ \frac{d^3T}{d\tau^3} \Big|_{\sin \theta=0} &= \frac{9(1-T)Tv^8 \left(b\mu v^4 \sin\left(\frac{1}{v^4}\right) + 2f(\delta v^4 - 1)\right)^2}{f^2 \left(b\mu v^8 \cos\left(\frac{1}{v^4}\right) + f(\delta v^4 - 1)^2\right)^2}. \end{aligned} \quad (92)$$

As in section 6.1, T is a monotonically increasing function on $\mathbf{S} \times J$. As a consequence, all orbits are originated from the invariant subset $T = 0$ (which contains the α -limit), which is classically related to the initial singularity with $H \rightarrow \infty$, and ends on the invariant boundary subset $T = 1$, which corresponds to asymptotically $H = 0$.

The curves of the equilibrium points are the same as in (63) with essentially the same dynamics as in Figure 4. That is, for $(b, f, \delta, \mu) = (0.1, 0.33, 0, 1)$ the points (iii) and (iv) are sources, and the points (v) and (vi) are saddles. The vertical plane represents the invariant set $\{(T, \theta, \varphi) : \varphi = 0, \sin(\theta) = 0\}$. The vertical sides (i) and (ii) of this rectangle are the local sinks.

7.2. Oscillating regime.

In this section, the potential $V(\phi) = \mu^3 \left[bf \left(\cos(\delta) - \cos\left(\delta + \frac{\phi}{f}\right)\right) + \frac{\phi^2}{\mu}\right]$, $b \neq 0$ is analyzed; looking for oscillatory behavior as expected from the numerical investigations. Asymptotic expansions are derived as well. As before, the pair

$$\left(\frac{\sqrt{2\mu\phi}}{\sqrt{\dot{\phi}^2 + 2\mu\phi^2}}, \quad \frac{\dot{\phi}}{\sqrt{\dot{\phi}^2 + 2\mu\phi^2}} \right), \quad (93)$$

defines a function of t with values in the unit circle. Therefore, the angular function $\vartheta(t)$, which is unique under identification modulo 2π , can be defined by

$$\vartheta = \tan^{-1} \left(\frac{\dot{\phi}}{\sqrt{2}\mu\phi} \right), \quad (94)$$

together with the radial function

$$r = \sqrt{\dot{\phi}^2 + 2\mu^2\phi^2}, \quad (95)$$

which satisfy:

$$bf\mu^3 \left(\cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) - \cos(\delta) \right) + 3H^2 - \frac{r^2}{2} = 0. \quad (96)$$

The inverse transformation $(r, \vartheta) \mapsto (\phi, \dot{\phi})$ is

$$\phi = \frac{r \cos(\vartheta)}{\sqrt{2}\mu}, \quad \dot{\phi} = r \sin(\vartheta). \quad (97)$$

For expanding universes ($H > 0$) the equations hold:

$$\begin{aligned} \dot{r} &= -\sqrt{\frac{3}{2}}r \sin^2(\vartheta) \sqrt{2bf\mu^3 \left(\cos(\delta) - \cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) \right) + r^2} \\ &\quad - b\mu^3 \sin(\vartheta) \sin \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right), \end{aligned} \quad (98a)$$

$$\begin{aligned} \dot{\vartheta} &= -\sqrt{2}\mu - \sqrt{\frac{3}{2}} \sin(\vartheta) \cos(\vartheta) \sqrt{2bf\mu^3 \left(\cos(\delta) - \cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) \right) + r^2} \\ &\quad - \frac{b\mu^3 \cos(\vartheta) \sin \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right)}{r}. \end{aligned} \quad (98b)$$

In the limit $b \rightarrow 0$ the solutions of the limiting equation admit the asymptotic expansions [118]:

$$\vartheta(t) = -\sqrt{2}\mu t + O(\ln t), \quad r(t) = \frac{4}{\sqrt{6}t} + O(t^{-2} \ln t). \quad (99)$$

Hence, when $b = 0$:

$$\phi(t) = \frac{4 \cos t}{\sqrt{6}t} + O(t^{-2} \ln t), \quad \dot{\phi}(t) = \frac{4 \sin t}{\sqrt{6}t} + O(t^{-2} \ln t). \quad (100)$$

The asymptotic expansions of the full problem ($b \neq 0$) are derived as follows.
Note that:

$$\begin{aligned} \dot{r} &= -b\mu^3 \sin(\delta) \sin(\vartheta) - \frac{b\mu^2 \cos(\delta) \cos(\vartheta) \sin(\vartheta) r}{\sqrt{2}f} \\ &\quad - \frac{\sqrt{3} \sqrt{b\mu^2 \cos(\vartheta) \sin(\delta)} \sin^2(\vartheta) r^{3/2}}{\sqrt[4]{2}} + \frac{b\mu \cos^2(\vartheta) \sin(\delta) \sin(\vartheta) r^2}{4f^2} + O(r^{5/2}), \end{aligned} \quad (101a)$$




Figure 7. Phase portrait of equations (98) (left panel). Projection over the cylinder S (right panel) for $(b, f, \delta) = (0.1, 0.33, 0)$ and different values of μ .

$$\begin{aligned} \dot{\vartheta} + \sqrt{2}\mu &= -\frac{\sqrt{3}\cos(\vartheta)\sqrt{b\mu^2\cos(\vartheta)\sin(\delta)}\sin(\vartheta)\sqrt{r}}{\sqrt[4]{2}} + \frac{b\mu\cos^3(\vartheta)\sin(\delta)r}{4f^2} \\ &\quad - \frac{(\sqrt{3}\cos(\vartheta)(b\mu\cos(\delta)\cos^2(\vartheta) + 2f)\sin(\vartheta))r^{3/2}}{4(2^{3/4}f\sqrt{b\mu^2\cos(\vartheta)\sin(\delta)})} + \frac{b\cos(\delta)\cos^4(\vartheta)r^2}{12\sqrt{2}f^3} + O(r^{5/2}). \end{aligned} \quad (101b)$$

To obtain an approximated solution near the oscillatory regime the average with respect to ϑ over any orbit of period 2π , $\langle (\dot{r}, \dot{\vartheta}) \rangle$ is taken, leading to

$$\dot{r} = kr^{3/2}, \quad k = \frac{2^{3/4}\sqrt{3}\sqrt{b}\mu(E(\frac{c}{2}|2) - E(\frac{c}{2} + \pi|2))\sqrt{\sin(\delta)}}{5\pi}, \quad (102a)$$

$$\dot{\vartheta} + \sqrt{2}\mu = \frac{b\cos(\delta)(r - 4f\mu)(r + 4f\mu)}{32\sqrt{2}f^3}. \quad (102b)$$




Figure 8. Phase portrait of equations (98) (left panel). Projection over the cylinder \mathbf{S} (right panel) for $(b, f, \delta) = (0.99, 0.09, 0)$ and different values of μ .

where $E(\phi|m)$ gives the elliptic integral of the second kind:

$$E(\phi|m) = \int_0^\phi (1 - m \sin^2(\theta))^{\frac{1}{2}} d\theta, \quad -\frac{\pi}{2} < \phi < \frac{\pi}{2}. \quad (103)$$

The averaged equations have solutions

$$r(t) = \frac{4}{(c_1 + kt)^2}, \vartheta(t) = -\frac{b \cos(\delta) \left(3f^2 \mu^2 (c_1 + kt) + \frac{1}{(c_1 + kt)^3} \right) + 12f^3 \mu (c_1 + kt)}{6\sqrt{2}f^3k} + \vartheta_0. \quad (104)$$

Introducing along with ϑ and r , the new variable

$$\varepsilon = \frac{H}{\mu + H}, \quad (105)$$

with inverse

$$H = \frac{\mu\varepsilon}{1-\varepsilon}, \quad (106)$$

satisfying:

$$bf\mu^3(\varepsilon - 1)^2 \left(\cos \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right) - \cos(\delta) \right) - \frac{1}{2}r^2(\varepsilon - 1)^2 + 3\mu^2\varepsilon^2 = 0, \quad (107)$$

along with the time derivative $\hat{\tau}$ given by

$$\frac{d\hat{\tau}}{dt} = \mu + H, \quad (108)$$

the following equations are obtained:

$$\varepsilon' = \frac{r^2(\varepsilon - 1)^3 \sin^2(\vartheta)}{2\mu^2}, \quad (109a)$$

$$r' = -3r\varepsilon \sin^2(\vartheta) + b\mu^2(\varepsilon - 1) \sin(\vartheta) \sin \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right), \quad (109b)$$

$$\vartheta' = -\sqrt{2}(1 - \varepsilon) - 3\varepsilon \sin(\vartheta) \cos(\vartheta) + \frac{b\mu^2(\varepsilon - 1) \cos(\vartheta) \sin \left(\delta + \frac{r \cos(\vartheta)}{\sqrt{2}f\mu} \right)}{r}. \quad (109c)$$

In order to obtain an approximated solution near the oscillatory regime the average with respect to ϑ over any orbit of period 2π , $\langle (\varepsilon', r', \vartheta') \rangle$, is taken, leading to:

$$\varepsilon' = \frac{(\varepsilon - 1)^3 r^2}{4\mu^2}, \quad (110a)$$

$$r' = -\frac{3\varepsilon r}{2}, \quad (110b)$$

$$\vartheta' = \frac{(\varepsilon - 1)(4f + b\mu \cos(\delta))}{2\sqrt{2}f} - \frac{(b(\varepsilon - 1) \cos(\delta))r^2}{32(\sqrt{2}f^3\mu)}, \quad (110c)$$

with the averaged constraint:

$$3\mu^2\varepsilon^2 - \frac{r^2((\varepsilon - 1)^2(b\mu \cos(\delta) + 4f))}{8f} = 0, \quad (111)$$

as $r \rightarrow 0$.

The above system is integrable yielding:

$$r(\varepsilon) = \frac{\sqrt{2}\sqrt{c_1(\varepsilon - 1)^2 + \mu^2(6\varepsilon - 3)}}{1 - \varepsilon}, \quad (112a)$$

$$\vartheta(\varepsilon) = -\frac{\mu \left(\frac{b \cos(\delta)}{\varepsilon - 1} - \frac{8f^2 \tanh^{-1} \left(\frac{c_1(\varepsilon - 1) + 3\mu^2}{\mu\sqrt{9\mu^2 - 3c_1}} \right) (b\mu \cos(\delta) + 4f)}{\sqrt{9\mu^2 - 3c_1}} \right)}{8\sqrt{2}f^3} + c_2, \quad (112b)$$

and

$$3(t - t_0) = \ln \left(\frac{(1 - \varepsilon)^2 \left(\frac{3\mu\sqrt{3\mu^2 - c_1} + \sqrt{3}c_1\varepsilon - \sqrt{3}c_1 + 3\sqrt{3}\mu^2}{3\mu\sqrt{3\mu^2 - c_1} - \sqrt{3}c_1\varepsilon + \sqrt{3}c_1 - 3\sqrt{3}\mu^2} \right) \frac{\mu}{\sqrt{\mu^2 - \frac{c_1}{3}}} }{c_1(\varepsilon - 1)^2 + \mu^2(6\varepsilon - 3)} \right)$$

$$\sim \ln \left(\frac{\left(\frac{2\mu(\sqrt{9\mu^2 - 3c_1} + 3\mu) - c_1}{c_1} \right) \frac{\mu}{\sqrt{\mu^2 - \frac{c_1}{3}}}}{c_1 - 3\mu^2} \right) - \frac{6\mu^2\varepsilon}{c_1 - 3\mu^2} + O(\varepsilon^2), \quad (112c)$$

as $\varepsilon \rightarrow 0$.

In Figure 7, the phase portrait of equations (98) (left panel) and the projection over the cylinder \mathbf{S} (right panel) for $(b, f, \delta) = (0.1, 0.33, 0)$ and different values of μ are presented. In Figure 8, the phase portrait of equations (98) (left panel) and the projection over the cylinder \mathbf{S} (right panel) for $(b, f, \delta) = (0.99, 0.09, 0)$ and different values of μ are shown. The plots show the periodic nature of the solutions.

8. Discussion

In this research, a local dynamical systems analysis for arbitrary $V(\phi)$ and $\chi(\phi)$ using Hubble normalized equations was provided. The analysis relies on two arbitrary functions $f(\lambda)$ and $g(\lambda)$ which encode the potential and the coupling function through the quadrature

$$\phi(\lambda) = \phi(1) - \int_1^\lambda \frac{1}{f(s)} ds, \quad V(\lambda) + \Lambda = W(1)e^{\int_1^\lambda \frac{s}{f(s)} ds}, \quad \chi(\lambda) = \chi(1)e^{\int_1^\lambda \frac{g(s)}{f(s)} ds}.$$

After that, a global dynamical systems formulation using the Alho & Uggla's approach [98] was implemented in Section 3. The equilibrium points that represent some solutions of cosmological interest were obtained: a matter–kinetic scaling solution, a matter–scalar field scaling solution, a kinetic dominated solution representing a stiff fluid, a solution dominated by the effective energy density of the geometric term $G_0(a)$, a scaling solution where the kinetic term and the effective energy density from $G_0(a)$ scales with the same order of magnitude, a quintessence scalar field dominated solution, the vacuum de Sitter solution associated to the minimum of the potential and a non-interacting matter dominated solution. All of which reveal a very rich cosmological behavior.

For the exponential potential $V(\phi) = V_0 e^{-\lambda\phi}$ in a vacuum analyzed in Section 4, the asymptotic states using either the Hubble–normalized equations or the Alho & Uggla's approach [98] are the same, but in some examples the asymptotic behavior can be better explained using Alho & Uggla's approach [98]. E.g., for the E -model: $V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3\alpha}}\phi}\right)^{2n}$ [101, 59] or the (generalized) harmonic potential. Recalling in Hubble–normalized equations the evolution equation for H , which is given by the Raychaudhuri equation, decouples. In particular, due it has the symmetry such that its derivative is also an exponential function, the Raychaudhuri equation always decouples for a scalar field with exponential potential. The asymptotics of the remaining reduced system is then typically given by the equilibrium points and often can be determined by a dynamical system analysis [15, 16, 17]. For other potentials that do not satisfy the above symmetry, like the harmonic potential $V(\phi) = \mu^2\phi^2$, the Raychaudhuri equation fails to decouple [98]. The resulting

Hubble normalized equations prove to be difficult to analyze in the usual dynamical systems approach due to the oscillations entering the system via the Klein–Gordon equation [99]. In reference [99], the oscillations and future asymptotics of locally rotationally symmetric Bianchi type III cosmologies with a massive scalar field with potential $V(\phi) = \frac{1}{2}\phi^2$ were studied.

According to the previous discussions the advantage of the Alho & Uggla's approach [98] is taken to study scalar-field cosmologies with generalized harmonic potentials of the type $V(\phi) = \mu^2\phi^2 + \text{cosine corrections}$.

In Section 5, the potential of the so-called E –model: $V(\phi) = V_0 \left(1 - e^{-\sqrt{\frac{2}{3\alpha}}\phi}\right)^{2n}$ with $f(s) = -\frac{s(s-\sqrt{6}\mu)}{2n}$, $\mu = \frac{2n}{3\sqrt{\alpha}}$, discussed in [101] for a conventional scalar field cosmology and in HoÅžavaâšLifshitz cosmology in [59], was studied. Observing that the dynamics of the latter potential is equivalent to that of the exponential potential plus a cosmological constant, $V = V_0 e^{-\sqrt{6}\mu\phi} + \Lambda$ having the f -deviser: $f(s) = -s(s - \sqrt{6}\mu)$, up to a rescaling in the independent variable.

In Sections 6 and 7, scalar field cosmologies under the potentials $V_1(\phi) = \mu^3 \left[\frac{\phi^2}{\mu} + b f \cos \left(\delta + \frac{\phi}{f} \right) \right]$, $b \neq 0$ and $V_2(\phi) = \mu^3 \left[b f \left(\cos(\delta) - \cos \left(\delta + \frac{\phi}{f} \right) \right) + \frac{\phi^2}{\mu} \right]$, $b \neq 0$ were respectively studied. The Alho & Uggla's approach [98] was used later in Sections 6.1 and 7.1 to find qualitative features and also to present the asymptotic analysis as $\phi \rightarrow \infty$ for the harmonic potentials $V_1(\phi)$ and $V_2(\phi)$ in a vacuum, respectively. In Section 6.2, the oscillatory regime for the scalar field under the potential $V_1(\phi)$ was investigated. Meanwhile, in Section 7.2, the oscillations of the scalar field under the potential $V_2(\phi)$ were studied.

9. Conclusions

In this paper, both local and global phase-space descriptions and averaging methods were used to find qualitative features of solutions for the FLRW and the Bianchi I metrics in the context of scalar field cosmologies with arbitrary potentials and arbitrary couplings to matter. The stability of the equilibrium points in a phase-space, as well as the dynamics in the regime where the scalar field diverges was studied. Equilibrium points that represent some solutions of cosmological interest such as: several types of scaling solutions, a kinetic dominated solution representing a stiff fluid, a solution dominated by an effective energy density of geometric origin, a quintessence scalar field dominated solution, the vacuum de Sitter solution associated to the minimum of the potential and a non-interacting matter dominated solution were found. All these reveal a very rich cosmological phenomenology.

The preliminary analysis of oscillations in scalar-field cosmologies with generalized harmonic potentials of type $V(\phi) = \mu^2\phi^2 + \text{cosine corrections}$ (implemented in Sections 6.2 and 7.2) will be improved in a forthcoming paper using averaging techniques similar to those used in [99] for a family of generalized harmonic potentials when H monotonically tends to zero. In this approach, the Hubble scalar plays the role of a time dependent perturbation parameter which controls the magnitude of error between full and the time-averaged solutions. The oscillations can be viewed as perturbations that can be smoothed out; and the Raychaudhuri equation of the averaged equations decouples. At the end, the analysis of the system is reduced to study the corresponding averaged equations.

Acknowledgments

This research was funded by Agencia Nacional de Investigación y Desarrollo- ANID through the program FONDECYT Iniciación grant no. 11180126 and by Vicerrectoría de Investigación y Desarrollo Tecnológico at Universidad Católica del Norte. Ellen de los M. Fernández Flores, Alfredo Millano, and Joey Latta are acknowledged for proofreading this manuscript and improving the English. Thanks to Alan Coley for his encouraging suggestions.

Appendix A. Existence and stability conditions of the equilibrium points of the system (12) as $\phi \rightarrow \infty$

The equilibrium points of the system (12) are the following:

$A_1(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{(4-3\gamma)g(\hat{\lambda})}{\sqrt{6(\gamma-2)}}, 1 - \frac{(4-3\gamma)^2 g(\hat{\lambda})^2}{6(\gamma-2)^2}, 0, \hat{\lambda} \right)$, where $\hat{\lambda}$ denotes the values of λ for which $f(\lambda) = 0$. It exists for $1 - \frac{(4-3\gamma)^2 g(\hat{\lambda})^2}{6(\gamma-2)^2} \geq 0$. The eigenvalues are $\left\{ \frac{3(\gamma-2)}{2} - \frac{(4-3\gamma)^2 g(\hat{\lambda})^2}{4(\gamma-2)}, 3\gamma - \frac{(4-3\gamma)^2 g(\hat{\lambda})^2}{2(\gamma-2)} - p, \frac{6(\gamma-2)\gamma + (3\gamma-4)g(\hat{\lambda})((4-3\gamma)g(\hat{\lambda})+2\hat{\lambda})}{2(\gamma-2)}, \frac{(3\gamma-4)g(\hat{\lambda})f'(\hat{\lambda})}{\gamma-2} \right\}$.

For $1 \leq \gamma < 2$, it follows that $A_1(\hat{\lambda})$ is a sink for:

- i) $3 < p \leq 4$, $1 \leq \gamma < \frac{p}{3}$, $-\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < g(\hat{\lambda}) < 0$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- ii) $3 < p \leq 4$, $1 \leq \gamma < \frac{p}{3}$, $0 < g(\hat{\lambda}) < \sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- iii) $4 < p \leq 6$, $1 \leq \gamma < \frac{4}{3}$, $-\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < g(\hat{\lambda}) < 0$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- iv) $4 < p \leq 6$, $1 \leq \gamma < \frac{4}{3}$, $0 < g(\hat{\lambda}) < \sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- v) $4 < p \leq 6$, $\frac{4}{3} < \gamma < \frac{p}{3}$, $-\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < g(\hat{\lambda}) < 0$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- vi) $4 < p \leq 6$, $\frac{4}{3} < \gamma < \frac{p}{3}$, $0 < g(\hat{\lambda}) < \sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- vii) $p > 6$, $1 \leq \gamma < \frac{4}{3}$, $-\frac{\sqrt{6}(\gamma-2)}{3\gamma-4} < g(\hat{\lambda}) < 0$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- viii) $p > 6$, $1 \leq \gamma < \frac{4}{3}$, $0 < g(\hat{\lambda}) < \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- ix) $p > 6$, $\frac{4}{3} < \gamma < 2$, $\frac{\sqrt{6}(\gamma-2)}{3\gamma-4} < g(\hat{\lambda}) < 0$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or
- x) $p > 6$, $\frac{4}{3} < \gamma < 2$, $0 < g(\hat{\lambda}) < -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > (\frac{3\gamma}{2} - 2)g(\hat{\lambda}) - \frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}$, or

$$\frac{3(\gamma-2)\gamma}{(3\gamma-4)g(\hat{\lambda})}.$$

If it exists, it will never be a source.

$$A_2(\hat{\lambda}): (x, \Omega_m, \Omega_0, \lambda) = \left(\frac{\sqrt{\frac{2}{3}(p-3\gamma)}}{(3\gamma-4)g(\hat{\lambda})}, \frac{2(6-p)(p-3\gamma)}{3(4-3\gamma)^2g(\hat{\lambda})^2}, \frac{2(p-3\gamma)(\gamma-2)}{(4-3\gamma)^2g(\hat{\lambda})^2} + 1, \hat{\lambda} \right). \text{ It exists for } \frac{2(6-p)(p-3\gamma)}{3(4-3\gamma)^2g(\hat{\lambda})^2} \geq 0. \text{ The eigenvalues are} \\ \left\{ -\frac{1}{4} \left(6 - p - \frac{\sqrt{p-6}\sqrt{3(4-3\gamma)^2g(\hat{\lambda})^2(-8\gamma+3p-2)+16(\gamma-2)(p-3\gamma)^2}}{(3\gamma-4)g(\hat{\lambda})} \right), \right. \\ \left. -\frac{1}{4} \left(6 - p + \frac{\sqrt{p-6}\sqrt{3(4-3\gamma)^2g(\hat{\lambda})^2(-8\gamma+3p-2)+16(\gamma-2)(p-3\gamma)^2}}{(3\gamma-4)g(\hat{\lambda})} \right), p - \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)g(\hat{\lambda})}, -\frac{2(p-3\gamma)f'(\hat{\lambda})}{(3\gamma-4)g(\hat{\lambda})} \right\}.$$

For $1 \leq \gamma < 2$, it follows that $A_2(\hat{\lambda})$ is a sink for:

- i) $1 < \gamma < \frac{4}{3}$, $3\gamma < p \leq \frac{2}{3}(4\gamma+1)$, $g(\hat{\lambda}) < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$, or
- ii) $1 < \gamma < \frac{4}{3}$, $3\gamma < p \leq \frac{2}{3}(4\gamma+1)$, $g(\hat{\lambda}) > \sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$, or
- iii) $1 < \gamma < \frac{4}{3}$, $\frac{2}{3}(4\gamma+1) < p < 6$, $-\frac{4\sqrt{-\frac{(\gamma-2)(p-3\gamma)^2}{(3\gamma-4)^2(-8\gamma+3p-2)}}}{\sqrt{3}} \leq g(\hat{\lambda}) < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$, or
- iv) $1 < \gamma < \frac{4}{3}$, $\frac{2}{3}(4\gamma+1) < p < 6$, $\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < g(\hat{\lambda}) \leq \frac{4\sqrt{-\frac{(\gamma-2)(p-3\gamma)^2}{(3\gamma-4)^2(-8\gamma+3p-2)}}}{\sqrt{3}}$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$, or
- v) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq \frac{2}{3}(4\gamma+1)$, $g(\hat{\lambda}) < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$, or
- vi) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq \frac{2}{3}(4\gamma+1)$, $g(\hat{\lambda}) > \sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$, or
- vii) $\frac{4}{3} < \gamma < 2$, $\frac{2}{3}(4\gamma+1) < p < 6$, $-\frac{4\sqrt{-\frac{(\gamma-2)(p-3\gamma)^2}{(3\gamma-4)^2(-8\gamma+3p-2)}}}{\sqrt{3}} \leq g(\hat{\lambda}) < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, $f'(\hat{\lambda}) < 0$, $\hat{\lambda} < \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$, or
- viii) $\frac{4}{3} < \gamma < 2$, $\frac{2}{3}(4\gamma+1) < p < 6$, $\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < g(\hat{\lambda}) \leq \frac{4\sqrt{-\frac{(\gamma-2)(p-3\gamma)^2}{(3\gamma-4)^2(-8\gamma+3p-2)}}}{\sqrt{3}}$, $f'(\hat{\lambda}) > 0$, $\hat{\lambda} > \frac{(3\gamma-4)pg(\hat{\lambda})}{2(p-3\gamma)}$.

It is non-hyperbolic for $p = 6$, saddle for $p = 2$. If it exists, it will never be a source.

$$A_3(\hat{\lambda}): (x, \Omega_m, \Omega_0, \lambda) = \left(\frac{\sqrt{6}\gamma}{(4-3\gamma)g(\hat{\lambda})+2\hat{\lambda}}, \frac{2(4-3\gamma)g(\hat{\lambda})\hat{\lambda}+4(\hat{\lambda}^2-3\gamma)}{((3\gamma-4)g(\hat{\lambda})-2\hat{\lambda})^2}, 0, \hat{\lambda} \right). \text{ Assuming } 1 \leq \gamma \leq 2, \text{ it exists for:}$$

- i) $1 \leq \gamma \leq 2$, $p > 0$, $\hat{\lambda} \leq \frac{1}{4}(-4g(\hat{\lambda}) - \sqrt{48\gamma+9\gamma^2g(\hat{\lambda})^2-24\gamma g(\hat{\lambda})^2+16g(\hat{\lambda})^2+3\gamma g(\hat{\lambda})})$, or
- ii) $1 \leq \gamma \leq 2$, $p > 0$, $\hat{\lambda} \geq \frac{1}{4}(-4g(\hat{\lambda}) + \sqrt{48\gamma+9\gamma^2g(\hat{\lambda})^2-24\gamma g(\hat{\lambda})^2+16g(\hat{\lambda})^2+3\gamma g(\hat{\lambda})})$.

The eigenvalues are $\left\{ \frac{p(4-3\gamma)g(\hat{\lambda})+2(p-3\gamma)\hat{\lambda}}{(-4+3\gamma)g(\hat{\lambda})-2\hat{\lambda}}, \frac{1}{(-8+6\gamma)g(\hat{\lambda})-4\hat{\lambda}} \left((12-9\gamma)g(\hat{\lambda}) - \right. \right.$

$$\begin{aligned}
& 3(-2 + \gamma)\hat{\lambda} - \sqrt{3}\sqrt{\left(2(-4 + 3\gamma)^3g(\hat{\lambda})^3\hat{\lambda} + (4 - 3\gamma)^2g(\hat{\lambda})^2\left(3 + 12\gamma - 8\hat{\lambda}^2\right)\right)} - \\
& 2(-4 + 3\gamma)g(\hat{\lambda})\hat{\lambda}\left(6 - 3\gamma + 6\gamma^2 - 4\hat{\lambda}^2\right) - 3(-2 + \gamma)\left(24\gamma^2 + (2 - 9\gamma)\hat{\lambda}^2\right)\Big), \\
& \frac{1}{(-8 + 6\gamma)g(\hat{\lambda}) - 4\hat{\lambda}}\left((12 - 9\gamma)g(\hat{\lambda}) - 3(-2 + \gamma)\hat{\lambda} + \right. \\
& \left.\sqrt{3}\sqrt{\left(2(-4 + 3\gamma)^3g(\hat{\lambda})^3\hat{\lambda} + (4 - 3\gamma)^2g(\hat{\lambda})^2\left(3 + 12\gamma - 8\hat{\lambda}^2\right)\right)} - 2(-4 + 3\gamma)g(\hat{\lambda})\hat{\lambda}\right. \\
& \left.\left(6 - 3\gamma + 6\gamma^2 - 4\hat{\lambda}^2\right) - 3(-2 + \gamma)\left(24\gamma^2 + (2 - 9\gamma)\hat{\lambda}^2\right)\right), \frac{6\gamma f'(\hat{\lambda})}{(-4 + 3\gamma)g(\hat{\lambda}) - 2\hat{\lambda}}\}.
\end{aligned}$$

In this case using a semi-analytically procedure for non-minimal coupling $g \equiv 0$ the eigenvalues are reduced to

$$\begin{cases} -\frac{-3\sqrt{(2-\gamma)(24\gamma^2+(2-9\gamma)\hat{\lambda}^2)}-3(\gamma-2)\hat{\lambda}}{4\hat{\lambda}}, -\frac{3\sqrt{(2-\gamma)(24\gamma^2+(2-9\gamma)\hat{\lambda}^2)}-3(\gamma-2)\hat{\lambda}}{4\hat{\lambda}}, \\ 3\gamma - p, -\frac{3\gamma f'(\hat{\lambda})}{\hat{\lambda}} \end{cases}.$$

Hence, assuming that $1 \leq \gamma < 2$, $A_3(\hat{\lambda})$ is a sink for:

- i) $1 \leq \gamma < 2, p > 3\gamma, -\frac{2\sqrt{6}\gamma}{\sqrt{9\gamma-2}} \leq \hat{\lambda} < -\sqrt{3}\sqrt{\gamma}, f'(\hat{\lambda}) < 0$, or
- ii) $1 \leq \gamma < 2, p > 3\gamma, \sqrt{3}\sqrt{\gamma} < \hat{\lambda} \leq \frac{2\sqrt{6}\gamma}{\sqrt{9\gamma-2}}, f'(\hat{\lambda}) > 0$.

Otherwise, it is a saddle. For non-minimal coupling the analysis is more complicated, so a numerical study is more reliable.

$A_4(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (-1, 0, 0, \hat{\lambda})$. The eigenvalues are

$$\left\{6 - p, -3\gamma + \sqrt{\frac{3}{2}}(3\gamma - 4)g(\hat{\lambda}) + 6, \sqrt{6}\hat{\lambda} + 6, \sqrt{6}f'(\hat{\lambda})\right\}.$$

$A_4(\hat{\lambda})$ is a source for:

- i) $1 \leq \gamma < \frac{4}{3}, 0 \leq p < 6, g(\hat{\lambda}) < \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > -\sqrt{6}$, or
- ii) $\gamma = \frac{4}{3}, 0 \leq p < 6, f'(\hat{\lambda}) > 0, \hat{\lambda} > -\sqrt{6}$, or
- iii) $\frac{4}{3} < \gamma < 2, 0 \leq p < 6, g(\hat{\lambda}) > \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > -\sqrt{6}$.

$A_4(\hat{\lambda})$ is a sink for

- i) $1 \leq \gamma < \frac{4}{3}, p > 6, g(\hat{\lambda}) > \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < -\sqrt{6}$, or
- ii) $\frac{4}{3} < \gamma < 2, p > 6, g(\hat{\lambda}) < \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < -\sqrt{6}$.

$A_5(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (0, 0, 1, \hat{\lambda})$. The eigenvalues are $\{0, \frac{p-6}{2}, p, p - 3\gamma\}$. It has a non-hyperbolic 3D unstable manifold for $p > 6$.

$A_6(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (1, 0, 0, \hat{\lambda})$. The eigenvalues are

$$\left\{6 - p, -3\gamma + \sqrt{\frac{3}{2}}(4 - 3\gamma)g(\hat{\lambda}) + 6, 6 - \sqrt{6}\hat{\lambda}, -\sqrt{6}f'(\hat{\lambda})\right\}.$$

$A_6(\hat{\lambda})$ is a source for:

- i) $1 \leq \gamma < \frac{4}{3}, p < 6, g(\hat{\lambda}) > -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < \sqrt{6}$, or
- ii) $\gamma = \frac{4}{3}, p < 6, f'(\hat{\lambda}) < 0, \hat{\lambda} < \sqrt{6}$, or
- iii) $\frac{4}{3} < \gamma < 2, p < 6, g(\hat{\lambda}) < -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) < 0, \hat{\lambda} < \sqrt{6}$.

$A_6(\hat{\lambda})$ is a sink for:

- i) $1 \leq \gamma < \frac{4}{3}, p > 6, g(\hat{\lambda}) < -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > \sqrt{6}$, or
- ii) $1 \leq \gamma > \frac{4}{3}, p > 6, g(\hat{\lambda}) > -\frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, f'(\hat{\lambda}) > 0, \hat{\lambda} > \sqrt{6}$.

$A_7(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{p}{\sqrt{6}\hat{\lambda}}, 0, 1 - \frac{p}{\hat{\lambda}^2}, \hat{\lambda} \right)$. It exists for $\hat{\lambda} < 0, 0 \leq p \leq 6$, or $\hat{\lambda} > 0, 0 \leq p \leq 6$. The eigenvalues are $\left\{ -\frac{1}{4} \left(6 - p - \frac{\sqrt{(p-6)(\hat{\lambda}^2(9p-6)-8p^2)}}{\hat{\lambda}} \right), -\frac{1}{4} \left(6 - p + \frac{\sqrt{(p-6)(\hat{\lambda}^2(9p-6)-8p^2)}}{\hat{\lambda}} \right), -3\gamma + \frac{(4-3\gamma)p\hat{\lambda}}{2\hat{\lambda}} + p, -\frac{pf'(\hat{\lambda})}{\hat{\lambda}} \right\}$.

For $1 \leq \gamma \leq 2$, $A_7(\hat{\lambda})$, it is a sink for:

- i) $1 \leq \gamma < \frac{4}{3}$, $\hat{\lambda} \leq -\sqrt{6}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) < 0$, or
- ii) $1 \leq \gamma < \frac{4}{3}$, $\hat{\lambda} \geq \sqrt{6}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- iii) $1 \leq \gamma < \frac{4}{3}$, $\hat{\lambda} > \frac{8}{3\sqrt{3}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- iv) $1 \leq \gamma < \frac{4}{3}$, $-\sqrt{6} < \hat{\lambda} < -\frac{8}{3\sqrt{3}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) < 0$, or
- v) $1 \leq \gamma < \frac{4}{3}$, $-\frac{8}{3\sqrt{3}} \leq \hat{\lambda} < 0$, $0 < p < \hat{\lambda}^2$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) < 0$, or
- vi) $1 \leq \gamma < \frac{4}{3}$, $-\sqrt{6} < \hat{\lambda} < -\frac{8}{3\sqrt{3}}$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) < 0$, or
- vii) $1 \leq \gamma < \frac{4}{3}$, $0 < \hat{\lambda} \leq \frac{8}{3\sqrt{3}}$, $0 < p < \hat{\lambda}^2$, $g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- viii) $1 \leq \gamma < \frac{4}{3}$, $\frac{8}{3\sqrt{3}} < \hat{\lambda} < \sqrt{6}$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2$, $g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- ix) $\gamma = \frac{4}{3}$, $\hat{\lambda} \geq \frac{8}{\sqrt{15}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $f'(\hat{\lambda}) > 0$, or
- x) $\gamma = \frac{4}{3}$, $\hat{\lambda} > 2$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $f'(\hat{\lambda}) > 0$, or
- xi) $\gamma = \frac{4}{3}$, $\hat{\lambda} \leq -\frac{8}{\sqrt{15}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $f'(\hat{\lambda}) < 0$, or
- xii) $\gamma = \frac{4}{3}$, $\frac{8}{3\sqrt{3}} < \hat{\lambda} \leq 2$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right)$, $f'(\hat{\lambda}) > 0$, or
- xiii) $\gamma = \frac{4}{3}$, $0 < \hat{\lambda} \leq \frac{8}{3\sqrt{3}}$, $0 < p < \hat{\lambda}^2$, $f'(\hat{\lambda}) > 0$, or
- xiv) $\gamma = \frac{4}{3}$, $2 < \hat{\lambda} < \frac{8}{\sqrt{15}}$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right) \leq p < 4$, $f'(\hat{\lambda}) > 0$, or
- xv) $\gamma = \frac{4}{3}$, $\frac{8}{3\sqrt{3}} < \hat{\lambda} \leq 2$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3} \sqrt{\hat{\lambda}^2 (27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2$, $f'(\hat{\lambda}) > 0$, or

- xvi) $\gamma = \frac{4}{3}$, $-\frac{8}{\sqrt{15}} < \hat{\lambda} < -2$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $f'(\hat{\lambda}) < 0$, or
- xvii) $\gamma = \frac{4}{3}$, $-2 \leq \hat{\lambda} < -\frac{8}{3\sqrt{3}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $f'(\hat{\lambda}) < 0$, or
- xviii) $\gamma = \frac{4}{3}$, $-\frac{8}{3\sqrt{3}} \leq \hat{\lambda} < 0$, $0 < p < \hat{\lambda}^2$, $f'(\hat{\lambda}) < 0$, or
- xix) $\gamma = \frac{4}{3}$, $-\frac{8}{\sqrt{15}} < \hat{\lambda} < -2$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < 4$, $f'(\hat{\lambda}) < 0$, or
- xx) $\gamma = \frac{4}{3}$, $-2 \leq \hat{\lambda} < -\frac{8}{3\sqrt{3}}$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2$, $f'(\hat{\lambda}) < 0$, or
- xxi) $\frac{4}{3} < \gamma < 2$, $\hat{\lambda} \geq \sqrt{6}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- xxii) $\frac{4}{3} < \gamma < 2$, $\hat{\lambda} = 2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- xxiii) $\frac{4}{3} < \gamma < 2$, $\hat{\lambda} > 2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- xxiv) $\frac{4}{3} < \gamma < 2$, $\hat{\lambda} = 2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}$, $3\gamma \leq p < \hat{\lambda}^2$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- xxv) $\frac{4}{3} < \gamma < 2$, $\hat{\lambda} \leq -\sqrt{6}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) < 0$, or
- xxvi) $\frac{4}{3} < \gamma < 2$, $\hat{\lambda} = -2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) < 0$, or
- xxvii) $\frac{4}{3} < \gamma < 2$, $\hat{\lambda} = -2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}$, $3\gamma \leq p < \hat{\lambda}^2$, $g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) < 0$, or
- xxviii) $\frac{4}{3} < \gamma < 2$, $\frac{8}{3\sqrt{3}} < \hat{\lambda} < 2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}$, $0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right)$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- xxix) $\frac{4}{3} < \gamma < 2$, $0 < \hat{\lambda} \leq \frac{8}{3\sqrt{3}}$, $0 < p < \hat{\lambda}^2$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- xxx) $\frac{4}{3} < \gamma < 2$, $\frac{8}{3\sqrt{3}} < \hat{\lambda} < 2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or
- xxxi) $\frac{4}{3} < \gamma < 2$, $2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}} < \hat{\lambda} < \sqrt{6}$, $\frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2$, $g(\hat{\lambda}) > \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}$, $f'(\hat{\lambda}) > 0$, or

- xxxii) $\frac{4}{3} < \gamma < 2, -\sqrt{6} < \hat{\lambda} < -2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}, 0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right), g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}, f'(\hat{\lambda}) < 0, \text{ or}$
- xxxiii) $\frac{4}{3} < \gamma < 2, -2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}} < \hat{\lambda} < -\frac{8}{3\sqrt{3}}, 0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right), g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}, f'(\hat{\lambda}) < 0, \text{ or}$
- xxxiv) $\frac{4}{3} < \gamma < 2, -\frac{8}{3\sqrt{3}} \leq \hat{\lambda} < 0, 0 < p < \hat{\lambda}^2, g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}, f'(\hat{\lambda}) < 0, \text{ or}$
- xxxv) $\frac{4}{3} < \gamma < 2, -\sqrt{6} < \hat{\lambda} < -2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}}, \frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2, g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}, f'(\hat{\lambda}) < 0, \text{ or}$
- xxxvi) $\frac{4}{3} < \gamma < 2, -2\sqrt{6}\sqrt{\frac{\gamma^2}{9\gamma-2}} < \hat{\lambda} < -\frac{8}{3\sqrt{3}}, \frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2, g(\hat{\lambda}) < \frac{2\hat{\lambda}(p-3\gamma)}{(3\gamma-4)p}, f'(\hat{\lambda}) < 0, \text{ or}$
- xxxvii) $\gamma = 2, \hat{\lambda} \geq \sqrt{6}, 0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right), g(\hat{\lambda}) > \frac{\hat{\lambda}(p-6)}{p}, f'(\hat{\lambda}) > 0, \text{ or}$
- xxxviii) $\gamma = 2, \hat{\lambda} > \frac{8}{3\sqrt{3}}, 0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right), g(\hat{\lambda}) > \frac{\hat{\lambda}(p-6)}{p}, f'(\hat{\lambda}) > 0, \text{ or}$
- xxxix) $\gamma = 2, \hat{\lambda} \leq -\sqrt{6}, 0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right), g(\hat{\lambda}) < \frac{\hat{\lambda}(p-6)}{p}, f'(\hat{\lambda}) < 0, \text{ or}$
- xli) $\gamma = 2, 0 < \hat{\lambda} \leq \frac{8}{3\sqrt{3}}, 0 < p < \hat{\lambda}^2, g(\hat{\lambda}) > \frac{\hat{\lambda}(p-6)}{p}, f'(\hat{\lambda}) > 0, \text{ or}$
- xlii) $\gamma = 2, \frac{8}{3\sqrt{3}} < \hat{\lambda} < \sqrt{6}, \frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2, g(\hat{\lambda}) > \frac{\hat{\lambda}(p-6)}{p}, f'(\hat{\lambda}) > 0, \text{ or}$
- xliii) $\gamma = 2, -\sqrt{6} < \hat{\lambda} < -\frac{8}{3\sqrt{3}}, 0 < p \leq \frac{1}{16} \left(9\hat{\lambda}^2 - \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right), g(\hat{\lambda}) < \frac{\hat{\lambda}(p-6)}{p}, f'(\hat{\lambda}) < 0, \text{ or}$
- xliv) $\gamma = 2, -\sqrt{6} < \hat{\lambda} < -\frac{8}{3\sqrt{3}}, \frac{1}{16} \left(9\hat{\lambda}^2 + \sqrt{3}\sqrt{\hat{\lambda}^2(27\hat{\lambda}^2 - 64)} \right) \leq p < \hat{\lambda}^2, g(\hat{\lambda}) < \frac{\hat{\lambda}(p-6)}{p}, f'(\hat{\lambda}) < 0.$

It will never be a source.

$A_8(\hat{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = \left(\frac{\hat{\lambda}}{\sqrt{6}}, 0, 0, \hat{\lambda} \right)$. It exists for $\hat{\lambda}^2 \leq 6$. The eigenvalues are

$$\left\{ \frac{1}{2} \left(\hat{\lambda}^2 - 6 \right), \hat{\lambda}^2 - p, -3\gamma + \frac{1}{2}(4 - 3\gamma)\hat{\lambda}g(\hat{\lambda}) + \hat{\lambda}^2, -\hat{\lambda}f'(\hat{\lambda}) \right\}.$$

$A_8(\hat{\lambda})$ is a sink for:

i) $1 \leq \gamma < \frac{4}{3}, p > \hat{\lambda}^2, f'(\hat{\lambda}) > 0, 0 < \hat{\lambda} < \sqrt{6}, g(\hat{\lambda}) < \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}$, or

ii) $1 \leq \gamma < \frac{4}{3}, p > \hat{\lambda}^2, g(\hat{\lambda}) > \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}, -\sqrt{6} < \hat{\lambda} < 0, f'(\hat{\lambda}) < 0$, or

- iii) $\gamma = \frac{4}{3}, p > \hat{\lambda}^2, -2 < \hat{\lambda} < 0, f'(\hat{\lambda}) < 0$, or
- iv) $\gamma = \frac{4}{3}, p > \hat{\lambda}^2, f'(\hat{\lambda}) > 0, 0 < \hat{\lambda} < 2$, or
- v) $\frac{4}{3} < \gamma < 2, p > \hat{\lambda}^2, g(\hat{\lambda}) > \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}, f'(\hat{\lambda}) > 0, 0 < \hat{\lambda} < \sqrt{6}$, or
- vi) $\frac{4}{3} < \gamma < 2, p > \hat{\lambda}^2, -\sqrt{6} < \hat{\lambda} < 0, g(\hat{\lambda}) < \frac{6\gamma-2\hat{\lambda}^2}{4\hat{\lambda}-3\gamma\hat{\lambda}}, f'(\hat{\lambda}) < 0$.

It will never be a source.

A_9 : $(x, \Omega_m, \Omega_0, \lambda) = (0, 0, 0, 0)$. The eigenvalues are

$$\left\{ -p, -3\gamma, -\frac{1}{2} \left(3 + \sqrt{9 - 12f(0)} \right), -\frac{1}{2} \left(3 - \sqrt{9 - 12f(0)} \right) \right\}. \text{ Sink for } p > 0, \gamma > 0, f(0) > 0. \text{ Otherwise, it is a saddle.}$$

$A_{10}(\tilde{\lambda})$: $(x, \Omega_m, \Omega_0, \lambda) = (0, 1, 0, \tilde{\lambda})$, where we denote by $\tilde{\lambda}$, the values of λ for which $g(\lambda) = 0$. The eigenvalues are $\left\{ \frac{1}{4} \left(3\gamma - 6 - \sqrt{24(4-3\gamma)f(\tilde{\lambda})g'(\tilde{\lambda}) + 9(\gamma-2)^2} \right), \frac{1}{4} \left(3\gamma - 6 + \sqrt{24(4-3\gamma)f(\tilde{\lambda})g'(\tilde{\lambda}) + 9(\gamma-2)^2} \right), 3\gamma, 3\gamma - p \right\}$. It is a saddle.

Appendix B. Existence and stability conditions of the equilibrium points of system (25) as $\phi \rightarrow \infty$

The equilibrium points of system (25) with $\varphi = 0$ (i.e., corresponding to $\phi \rightarrow \infty$) are the following:

B_1 : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{N^2}{6}}, -\frac{N}{\sqrt{6}} \right] + 2\pi c_1, 0, 0, 0 \right)$, $c_1 \in \mathbb{Z}$ with eigenvalues $\left\{ 0, \frac{N^2}{2}, \frac{1}{2} (N^2 - 6), N^2 - p, N(2M + N) - \frac{3}{2}\gamma(MN + 2) \right\}$. It exists for $N^2 \leq 6$. It is always a non-hyperbolic saddle for $N^2 \leq 6$.

B_2 : $\left(1, \tan^{-1} \left[\sqrt{1 - \frac{N^2}{6}}, -\frac{N}{\sqrt{6}} \right] + 2\pi c_1, 0, 0, 0 \right)$, $c_1 \in \mathbb{Z}$ with eigenvalues $\left\{ 0, -\frac{N^2}{2}, \frac{1}{2} (N^2 - 6), N^2 - p, N(2M + N) - \frac{3}{2}\gamma(MN + 2) \right\}$. It exists for $N^2 \leq 6$.

The case of physical interest is when it is non-hyperbolic with a 4D stable manifold for:

- i) $M \in \mathbb{R}, -2 < N < 0, p > N^2, \gamma = \frac{4}{3}$, or
- ii) $M \in \mathbb{R}, 0 < N < 2, p > N^2, \gamma = \frac{4}{3}$, or
- iii) $-\sqrt{6} < N \leq -2, p > N^2, 1 \leq \gamma < \frac{4}{3}, M > \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$, or
- iv) $-2 < N < 0, p > N^2, 1 \leq \gamma < \frac{4}{3}, M > \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$, or
- v) $0 < N < 2, p > N^2, \frac{4}{3} < \gamma \leq 2, M > \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$, or
- vi) $2 \leq N < \sqrt{6}, p > N^2, \frac{4}{3} < \gamma \leq 2, M > \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$, or
- vii) $-\sqrt{6} < N \leq -2, p > N^2, \frac{4}{3} < \gamma \leq 2, M < \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$, or
- viii) $0 < N < 2, p > N^2, 1 \leq \gamma < \frac{4}{3}, M < \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$, or
- ix) $2 \leq N < \sqrt{6}, p > N^2, 1 \leq \gamma < \frac{4}{3}, M < \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$, or
- x) $-2 < N < 0, p > N^2, \frac{4}{3} < \gamma \leq 2, M < \frac{2(N^2-3\gamma)}{N(3\gamma-4)}$.

B_3 : $(0, 2\pi c_1, 0, 1, 0)$, $c_1 \in \mathbb{Z}$ with eigenvalues $\{0, \frac{p-6}{2}, \frac{p}{2}, p, p - 3\gamma\}$. The physical interesting situation is when it is non-hyperbolic with a 4D unstable manifold for $p > 6, 1 \leq \gamma \leq 2$. It is a non-hyperbolic saddle otherwise.

B_4 : $(1, 2\pi c_1, 0, 1, 0), c_1 \in \mathbb{Z}$ with eigenvalues $\{0, \frac{p-6}{2}, -\frac{p}{2}, p, p-3\gamma\}$. It is a non-hyperbolic saddle.

B_5 : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{p^2}{6N^2}}, -\frac{p}{\sqrt{6}N} \right] + 2\pi c_1, 0, 1 - \frac{p}{N^2}, 0\right), c_1 \in \mathbb{Z}$ with eigenvalues $\left\{0, \frac{p}{2}, \frac{p((4-3\gamma)M+2N)}{2N} - 3\gamma, \frac{1}{4} \left(p - 6 + \frac{\sqrt{(p-6)(N^2(9p-6)-8p^2)}}{N}\right), \frac{1}{4} \left(p - 6 - \frac{\sqrt{(p-6)(N^2(9p-6)-8p^2)}}{N}\right)\right\}$. It exists for $p > 0, N^2 > \frac{p^2}{6}$. Furthermore, it satisfies $\Omega_0 \geq 0$ for $0 < p < 6, N^2 \geq p$, or $p \geq 6, N^2 > \frac{p^2}{6}$. It is a non-hyperbolic saddle.

B_6 : $\left(1, \tan^{-1} \left[\sqrt{1 - \frac{p^2}{6N^2}}, -\frac{p}{\sqrt{6}N} \right] + 2\pi c_1, 0, 1 - \frac{p}{N^2}, 0\right), c_1 \in \mathbb{Z}$ with eigenvalues $\left\{0, -\frac{p}{2}, \frac{p((4-3\gamma)M+2N)}{2N} - 3\gamma, \frac{1}{4} \left(p - 6 + \frac{\sqrt{(p-6)(N^2(9p-6)-8p^2)}}{N}\right), \frac{1}{4} \left(p - 6 - \frac{\sqrt{(p-6)(N^2(9p-6)-8p^2)}}{N}\right)\right\}$. It exists for $p > 0, N^2 > \frac{p^2}{6}$. Furthermore, it satisfies $\Omega_0 \geq 0$ for $0 < p < 6, N^2 \geq p$, or $p \geq 6, N^2 > \frac{p^2}{6}$. The situation of physical interest is when it is non-hyperbolic with a 4D stable manifold for:

- i) $N > \frac{8}{\sqrt{15}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), \gamma = \frac{4}{3}$, or
- ii) $N < -\frac{8}{3\sqrt{3}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), \gamma = \frac{4}{3}$, or
- iii) $N = \frac{8}{\sqrt{15}}, 0 < p \leq \frac{4}{5}, \gamma = \frac{4}{3}$, or
- iv) $0 < N \leq \frac{8}{3\sqrt{3}}, 0 < p < N^2, \gamma = \frac{4}{3}$, or
- v) $-\frac{8}{3\sqrt{3}} \leq N < 0, 0 < p < N^2, \gamma = \frac{4}{3}$, or
- vi) $\frac{8}{3\sqrt{3}} < N < \frac{8}{\sqrt{15}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), \gamma = \frac{4}{3}$, or
- vii) $\frac{8}{3\sqrt{3}} < N \leq 2, \frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2-64)}\right) \leq p < N^2, \gamma = \frac{4}{3}$, or
- viii) $-2 < N < -\frac{8}{3\sqrt{3}}, \frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2-64)}\right) \leq p < N^2, \gamma = \frac{4}{3}$, or
- ix) $2 < N < \frac{8}{\sqrt{15}}, \frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2-64)}\right) \leq p < 4, \gamma = \frac{4}{3}$, or
- x) $-\frac{8}{\sqrt{15}} < N \leq -2, \frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2-64)}\right) \leq p < 4, \gamma = \frac{4}{3}$, or
- xi) $\frac{8}{3\sqrt{3}} < N < \frac{8}{\sqrt{15}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), 1 \leq \gamma < \frac{4}{3}, M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xii) $\frac{8}{3\sqrt{3}} < N < \frac{8}{\sqrt{15}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), \frac{4}{3} < \gamma \leq 2, M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xiii) $N < -\frac{8}{3\sqrt{3}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), 1 \leq \gamma < \frac{4}{3}, M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xiv) $N < -\frac{8}{3\sqrt{3}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), \frac{4}{3} < \gamma \leq 2, M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xv) $N > \frac{8}{\sqrt{15}}, 0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2-64)}\right), 1 \leq \gamma < \frac{4}{3}, M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or

- xvi) $N > \frac{8}{\sqrt{15}}$, $0 < p \leq \frac{1}{16} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right)$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xvii) $N = \frac{8}{\sqrt{15}}$, $0 < p \leq \frac{4}{5}$, $1 \leq \gamma < \frac{4}{3}$, $M < \frac{16(p-3\gamma)}{\sqrt{15}(3\gamma-4)p}$, or
- xviii) $N = \frac{8}{\sqrt{15}}$, $0 < p \leq \frac{4}{5}$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{16(p-3\gamma)}{\sqrt{15}(3\gamma-4)p}$, or
- xix) $N = \frac{8}{\sqrt{15}}$, $4 \leq p < \frac{64}{15}$, $1 \leq \gamma < \frac{4}{3}$, $M < \frac{16(p-3\gamma)}{\sqrt{15}(3\gamma-4)p}$, or
- xx) $N = \frac{8}{\sqrt{15}}$, $4 \leq p < \frac{64}{15}$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{16(p-3\gamma)}{\sqrt{15}(3\gamma-4)p}$, or
- xxi) $-2 < N < -\frac{8}{3\sqrt{3}}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxii) $-\sqrt{6} < N \leq -\frac{8}{\sqrt{15}}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxiii) $\frac{8}{3\sqrt{3}} < N \leq 2$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxiv) $\frac{8}{\sqrt{15}} < N < \sqrt{6}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxv) $\frac{8}{3\sqrt{3}} < N \leq 2$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxvi) $\frac{8}{\sqrt{15}} < N < \sqrt{6}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxvii) $-2 < N < -\frac{8}{3\sqrt{3}}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxviii) $-\sqrt{6} < N \leq -\frac{8}{\sqrt{15}}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxix) $2 < N < \frac{8}{\sqrt{15}}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < 4$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxx) $-\frac{8}{\sqrt{15}} < N \leq -2$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < 4$, $1 \leq \gamma < \frac{4}{3}$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxxi) $2 < N < \frac{8}{\sqrt{15}}$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < 4$, $1 \leq \gamma < \frac{4}{3}$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxxii) $-\frac{8}{\sqrt{15}} < N \leq -2$, $\frac{1}{16} \left(9N^2 + \sqrt{3}\sqrt{N^2(27N^2 - 64)} \right) \leq p < 4$, $\frac{4}{3} < \gamma \leq 2$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxxiii) $-\frac{8}{3\sqrt{3}} \leq N < 0$, $0 < p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxxiv) $0 < N \leq \frac{8}{3\sqrt{3}}$, $0 < p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- xxxv) $2 < N < \frac{8}{\sqrt{15}}$, $4 \leq p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or

xxxvi) $-\frac{8}{\sqrt{15}} < N < -2$, $4 \leq p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M > \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or

xxxvii) $-\frac{8}{3\sqrt{3}} \leq N < 0$, $0 < p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or

xxxviii) $0 < N \leq \frac{8}{3\sqrt{3}}$, $0 < p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or

xxxix) $2 < N < \frac{8}{\sqrt{15}}$, $4 \leq p < N^2$, $1 \leq \gamma < \frac{4}{3}$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or

xl) $-\frac{8}{\sqrt{15}} < N < -2$, $4 \leq p < N^2$, $\frac{4}{3} < \gamma \leq 2$, $M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$.

B_7 : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{2(p-3\gamma)^2}{3M^2(4-3\gamma)^2}}, \frac{2(p-3\gamma)}{\sqrt{6}M(4-3\gamma)} \right] + 2\pi c_1, \frac{2(6-p)(p-3\gamma)}{3(4-3\gamma)^2 M^2}, \frac{(4-3\gamma)^2 M^2 + 2(\gamma-2)(p-3\gamma)}{(4-3\gamma)^2 M^2}, 0\right)$, $c_1 \in \mathbb{Z}$, with eigenvalues

$$\left\{ 0, \frac{p}{2}, \frac{-3\gamma Mp + 4Mp - 6\gamma N + 2Np}{M(4-3\gamma)}, \frac{1}{4} \left(p - 6 + \frac{\sqrt{(p-6)(3(4-3\gamma)^2 M^2(-8\gamma+3p-2)+16(\gamma-2)(p-3\gamma)^2)}}{(3\gamma-4)M} \right), \frac{1}{4} \left(p - 6 - \frac{\sqrt{(p-6)(3(4-3\gamma)^2 M^2(-8\gamma+3p-2)+16(\gamma-2)(p-3\gamma)^2)}}{(3\gamma-4)M} \right) \right\}. \text{ It exists for:}$$

i) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M > 0$, or

ii) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M > 0$, or

iii) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M > 0$, or

iv) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M > 0$, or

v) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M < 0$, or

vi) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M < 0$, or

vii) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M < 0$, or

viii) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M < 0$, or

ix) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \geq \frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$, or

x) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \geq -\frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$, or

xi) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \geq \frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$, or

xii) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \geq -\frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$, or

xiii) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \leq -\frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$, or

xiv) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \leq \frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$, or

xv) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \leq -\frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$, or

xvi) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \leq \frac{\sqrt{\frac{2}{3}(3\gamma-p)}}{3\gamma-4}$.

It is a non-hyperbolic saddle.

B_8 : $\left(1, \tan^{-1} \left[\sqrt{1 - \frac{2(p-3\gamma)^2}{3M^2(4-3\gamma)^2}}, \frac{2(p-3\gamma)}{\sqrt{6}M(4-3\gamma)} \right] + 2\pi c_1, \frac{2(6-p)(p-3\gamma)}{3(4-3\gamma)^2 M^2}, \frac{(4-3\gamma)^2 M^2 + 2(\gamma-2)(p-3\gamma)}{(4-3\gamma)^2 M^2}, 0\right)$, $c_1 \in \mathbb{Z}$, with eigenvalues

$$\left\{ 0, -\frac{p}{2}, \frac{-3\gamma Mp + 4Mp - 6\gamma N + 2Np}{M(4-3\gamma)}, \frac{1}{4} \left(p - 6 + \frac{\sqrt{(p-6)(3(4-3\gamma)^2 M^2(-8\gamma+3p-2)+16(\gamma-2)(p-3\gamma)^2)}}{(3\gamma-4)M} \right), \frac{1}{4} \left(p - 6 - \frac{\sqrt{(p-6)(3(4-3\gamma)^2 M^2(-8\gamma+3p-2)+16(\gamma-2)(p-3\gamma)^2)}}{(3\gamma-4)M} \right) \right\}. \text{ It exists for:}$$

i) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M > 0$, or

ii) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M > 0$, or

iii) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M > 0$, or

iv) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M > 0$, or

v) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M < 0$, or

vi) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M < 0$, or

- vii) $1 \leq \gamma < \frac{4}{3}$, $p = 3\gamma$, $M < 0$, or
- viii) $\frac{4}{3} < \gamma \leq 2$, $p = 3\gamma$, $M < 0$, or
- ix) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \geq \frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$, or
- x) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \geq -\frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$, or
- xi) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \geq \frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$, or
- xii) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \geq -\frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$, or
- xiii) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \leq -\frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$, or
- xiv) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \leq \frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$, or
- xv) $1 \leq \gamma < \frac{4}{3}$, $3\gamma < p \leq 6$, $M \leq -\frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$, or
- xvi) $\frac{4}{3} < \gamma < 2$, $3\gamma < p \leq 6$, $M \leq \frac{\sqrt{\frac{2}{3}}(3\gamma-p)}{3\gamma-4}$.

The situation of physical interest is when it is non-hyperbolic with a 4D stable manifold for:

- i) $N < -3\sqrt{6}$, $p = \frac{1}{16} \left(27N^2 - \sqrt{3}\sqrt{N^2(243N^2 - 1216)} \right)$, $\frac{4}{3} < \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $\frac{2N(p-3\gamma)}{(3\gamma-4)p} < M < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, or
- ii) $N < -3\sqrt{6}$, $N^2 - \sqrt{N^2(N^2-6)} < p \leq \frac{1}{8} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2 - 160)} \right)$, $\gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $-\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- iii) $N < -3\sqrt{6}$, $\frac{1}{2} \left(3N^2 - \sqrt{3}\sqrt{N^2(3N^2 - 16)} \right) < p < \frac{1}{16} \left(27N^2 - \sqrt{3}\sqrt{N^2(243N^2 - 1216)} \right)$, $\frac{4}{3} < \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $\frac{2N(p-3\gamma)}{(3\gamma-4)p} < M < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, or
- iv) $N < -3\sqrt{6}$, $\frac{1}{8} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2 - 160)} \right) < p \leq \frac{1}{2} \left(3N^2 - \sqrt{3}\sqrt{N^2(3N^2 - 16)} \right)$, $\frac{9N^2p-6N^2-8p^2}{4(6N^2-p^2)} \leq \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $-\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- v) $-3\sqrt{6} \leq N < -\sqrt{6}$, $p = \frac{1}{16} \left(27N^2 - \sqrt{3}\sqrt{N^2(243N^2 - 1216)} \right)$, $\frac{4}{3} < \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $\frac{2N(p-3\gamma)}{(3\gamma-4)p} < M < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, or
- vi) $-3\sqrt{6} \leq N < -\sqrt{6}$, $N^2 - \sqrt{N^2(N^2-6)} < p \leq \frac{1}{8} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2 - 160)} \right)$, $\gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $-\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- vii) $-3\sqrt{6} \leq N < -\sqrt{6}$, $\frac{1}{2} \left(3N^2 - \sqrt{3}\sqrt{N^2(3N^2 - 16)} \right) < p < \frac{1}{16} \left(27N^2 - \sqrt{3}\sqrt{N^2(243N^2 - 1216)} \right)$, $\frac{4}{3} < \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $\frac{2N(p-3\gamma)}{(3\gamma-4)p} < M < -\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, or
- viii) $-3\sqrt{6} \leq N < -\sqrt{6}$, $\frac{1}{8} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2 - 160)} \right) < p < \frac{1}{2} \left(3N^2 - \sqrt{3}\sqrt{N^2(3N^2 - 16)} \right)$, $\frac{9N^2p-6N^2-8p^2}{4(6N^2-p^2)} \leq \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $-\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or
- ix) $N > \sqrt{6}$, $p = \frac{1}{16} \left(27N^2 - \sqrt{3}\sqrt{N^2(243N^2 - 1216)} \right)$, $\frac{4}{3} < \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$, or

- x) $N > \sqrt{6}$, $N^2 - \sqrt{N^2(N^2 - 6)} < p \leq \frac{1}{8} \left(9N^2 - \sqrt{3}\sqrt{N^2(27N^2 - 160)} \right)$, $\gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $\frac{2N(p-3\gamma)}{(3\gamma-4)p} < M < \sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}}$, or
xi) $N > \sqrt{6}$, $\frac{1}{2} \left(3N^2 - \sqrt{3}\sqrt{N^2(3N^2 - 16)} \right) < p < \frac{1}{16} \left(27N^2 - \sqrt{3}\sqrt{N^2(243N^2 - 1216)} \right)$, $\frac{4}{3} < \gamma < \frac{2p(N^2-p)}{6N^2-p^2}$, $\sqrt{2}\sqrt{\frac{(\gamma-2)(3\gamma-p)}{(3\gamma-4)^2}} < M < \frac{2N(p-3\gamma)}{(3\gamma-4)p}$.

B_9 : $\left(0, \tan^{-1} \left[\frac{\sqrt{6(2-\gamma)^2-(4-3\gamma)^2M^2}}{\sqrt{6}(2-\gamma)}, \frac{(4-3\gamma)M}{\sqrt{6}(2-\gamma)} \right] + 2\pi c_1, \frac{6(2-\gamma)^2-(4-3\gamma)^2M^2}{6(2-\gamma)^2}, 0, 0 \right)$, $c_1 \in \mathbb{Z}$, with eigenvalues

$$\left\{ 0, \frac{6(\gamma-2)^2-(4-3\gamma)^2M^2}{4(\gamma-2)}, -\frac{(4-3\gamma)^2M^2-6(\gamma-2)\gamma}{4(\gamma-2)}, \gamma \left(3 - \frac{9M^2}{2} \right) + \frac{2M(N-M)}{\gamma-2} + 3M(M+N), -\frac{(4-3\gamma)^2M^2+2(\gamma-2)(p-3\gamma)}{2(\gamma-2)} \right\}. \text{ It exists for:}$$

- i) $\gamma = \frac{4}{3}$, $p > 0$, or
- ii) $1 \leq \gamma < \frac{4}{3}$, $\frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4} \leq M \leq \frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4}$, $p > 0$, or
- iii) $\frac{4}{3} < \gamma < 2$, $\frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4} \leq M \leq \frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4}$, $p > 0$.

It is a non-hyperbolic saddle.

B_{10} : $\left(1, \tan^{-1} \left[\frac{\sqrt{6(2-\gamma)^2-(4-3\gamma)^2M^2}}{\sqrt{6}(2-\gamma)}, \frac{(4-3\gamma)M}{\sqrt{6}(2-\gamma)} \right] + 2\pi c_1, \frac{6(2-\gamma)^2-(4-3\gamma)^2M^2}{6(2-\gamma)^2}, 0, 0 \right)$, $c_1 \in \mathbb{Z}$ with eigenvalues $\left\{ 0, \frac{6(\gamma-2)^2-(4-3\gamma)^2M^2}{4(\gamma-2)}, \frac{(4-3\gamma)^2M^2-6(\gamma-2)\gamma}{4(\gamma-2)}, \gamma \left(3 - \frac{9M^2}{2} \right) + \frac{2M(N-M)}{\gamma-2} + 3M(M+N), -\frac{(4-3\gamma)^2M^2+2(\gamma-2)(p-3\gamma)}{2(\gamma-2)} \right\}$. It exists for:

- i) $\gamma = \frac{4}{3}$, $p > 0$, or
- ii) $1 \leq \gamma < \frac{4}{3}$, $\frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4} \leq M \leq \frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4}$, $p > 0$, or
- iii) $\frac{4}{3} < \gamma < 2$, $\frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4} \leq M \leq \frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4}$, $p > 0$.

The situation of physical interest is when it is non-hyperbolic with a 4D stable manifold for:

- i) $1 \leq \gamma < \frac{4}{3}$, $N < -\sqrt{6}$, $\frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} < M < \frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4}$, $p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}$, or
- ii) $\frac{4}{3} < \gamma < 2$, $N < -\sqrt{6}$, $\frac{\sqrt{6}\gamma-2\sqrt{6}}{3\gamma-4} < M < \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4}$, $p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}$, or
- iii) $1 \leq \gamma < \frac{4}{3}$, $N > \sqrt{6}$, $\frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4} < M < \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4}$, $p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}$, or
- iv) $\frac{4}{3} < \gamma < 2$, $N > \sqrt{6}$, $\frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} < M < \frac{2\sqrt{6}-\sqrt{6}\gamma}{3\gamma-4}$, $p > \frac{6\gamma^2-12\gamma-9\gamma^2M^2+24\gamma M^2-16M^2}{2\gamma-4}$.

B_{11} : $\left(0, \tan^{-1} \left[\sqrt{1 - \frac{6\gamma^2}{(2N+M(4-3\gamma))^2}}, -\frac{\sqrt{6}\gamma}{2N+M(4-3\gamma)} \right] + 2\pi c_1, \frac{4N(2M+N)-6(2+MN)\gamma}{(2N+M(4-3\gamma))^2}, 0, 0 \right)$, $c_1 \in \mathbb{Z}$, with eigenvalues $\left\{ 0, \frac{3N\gamma}{4M+2N-3M\gamma}, \frac{Mp(4-3\gamma)+2N(p-3\gamma)}{-2N+M(-4+3\gamma)}, \frac{1}{2(2N+M(4-3\gamma))^2}(3(2N+M(4-3\gamma))(N(-2+\gamma)+M(-4+3\gamma))+\sqrt{3}\sqrt{((2N+M(4-3\gamma))^2(-M^2(-3+8N^2-12\gamma)(4-3\gamma)^2+2M^3N(-4+3\gamma)^3+2MN(-4+3\gamma)(-6+4N^2+3\gamma-6\gamma^2)-3(-2+\gamma)(N^2(2-9\gamma)+24\gamma^2)))}, \frac{1}{2(2N+M(4-3\gamma))^2}(3(2N+M(4-3\gamma))(N(-2+\gamma)+M(-4+3\gamma))- \right.$

$\sqrt{3}\sqrt{\left((2N+M(4-3\gamma))^2 \left(-M^2 (-3+8N^2-12\gamma) (4-3\gamma)^2 + 2M^3 N(-4+3\gamma)^3 + 2MN(-4+3\gamma) (-6+4N^2+3\gamma-6\gamma^2) - 3(-2+\gamma) (N^2(2-9\gamma)+24\gamma^2)\right)\right)}\}. It exists for:$

- i) $M \in \mathbb{R}, \gamma = \frac{4}{3}, N < -2, \text{ or}$
- ii) $M \in \mathbb{R}, \gamma = \frac{4}{3}, N > 2, \text{ or}$
- iii) $\frac{4}{3} < \gamma \leq 2, N = -\sqrt{6}, M > \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, \text{ or}$
- iv) $\frac{4}{3} < \gamma \leq 2, 0 < N \leq \sqrt{6}, M < \frac{2(N^2-3\gamma)}{(3\gamma-4)N}, \text{ or}$
- v) $\frac{4}{3} < \gamma \leq 2, N < -\sqrt{6}, M \geq \frac{\sqrt{6}\gamma+2N}{3\gamma-4}, \text{ or}$
- vi) $\frac{4}{3} < \gamma \leq 2, -\sqrt{6} < N < 0, M > \frac{2(N^2-3\gamma)}{(3\gamma-4)N}, \text{ or}$
- vii) $\frac{4}{3} < \gamma \leq 2, N > \sqrt{6}, M \leq -\frac{\sqrt{6}\gamma-2N}{3\gamma-4}, \text{ or}$
- viii) $1 \leq \gamma < \frac{4}{3}, N = -\sqrt{6}, M < \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, \text{ or}$
- ix) $1 \leq \gamma < \frac{4}{3}, N > \sqrt{6}, M \geq -\frac{\sqrt{6}\gamma-2N}{3\gamma-4}, \text{ or}$
- x) $1 \leq \gamma < \frac{4}{3}, N < -\sqrt{6}, M \leq \frac{\sqrt{6}\gamma+2N}{3\gamma-4}, \text{ or}$
- xi) $1 \leq \gamma < \frac{4}{3}, 0 < N \leq \sqrt{6}, M > \frac{2(N^2-3\gamma)}{(3\gamma-4)N}, \text{ or}$
- xii) $1 \leq \gamma < \frac{4}{3}, -\sqrt{6} < N < 0, M < \frac{2(N^2-3\gamma)}{(3\gamma-4)N}$

It is an hyperbolic saddle.

$B_{12}: \left(1, \tan^{-1} \left[\sqrt{1 - \frac{6\gamma^2}{(2N+M(4-3\gamma))^2}}, -\frac{\sqrt{6}\gamma}{2N+M(4-3\gamma)} \right] + 2\pi c_1, \frac{4N(2M+N)-6(2+MN)\gamma}{(2N+M(4-3\gamma))^2}, 0, 0\right), c_1 \in \mathbb{Z} \text{ with eigenvalues } \left\{0, -\frac{3N\gamma}{4M+2N-3M\gamma}, \frac{Mp(4-3\gamma)+2N(p-3\gamma)}{-2N+M(-4+3\gamma)}, \right.$
 $\left.\frac{1}{4N+M(8-6\gamma)}(3N(-2+\gamma)+3M(-4+3\gamma)+\sqrt{(-3M^2(-3+8N^2-12\gamma)(4-3\gamma)^2+6M^3N(-4+3\gamma)^3+6MN(-4+3\gamma)(-6+4N^2+3\gamma-6\gamma^2)-9(-2+\gamma)(N^2(2-9\gamma)+24\gamma^2))}), \frac{1}{4N+M(8-6\gamma)}(3N(-2+\gamma)+3M(-4+3\gamma)-\sqrt{(-3M^2(-3+8N^2-12\gamma)(4-3\gamma)^2+6M^3N(-4+3\gamma)^3+6MN(-4+3\gamma)(-6+4N^2+3\gamma-6\gamma^2)-9(-2+\gamma)(N^2(2-9\gamma)+24\gamma^2))})\right\}. It exists for:$

- i) $M \in \mathbb{R}, \gamma = \frac{4}{3}, N < -2, \text{ or}$
- ii) $M \in \mathbb{R}, \gamma = \frac{4}{3}, N > 2, \text{ or}$
- iii) $\frac{4}{3} < \gamma \leq 2, N = -\sqrt{6}, M > \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, \text{ or}$
- iv) $\frac{4}{3} < \gamma \leq 2, 0 < N \leq \sqrt{6}, M < \frac{2(N^2-3\gamma)}{(3\gamma-4)N}, \text{ or}$
- v) $\frac{4}{3} < \gamma \leq 2, N < -\sqrt{6}, M \geq \frac{\sqrt{6}\gamma+2N}{3\gamma-4}, \text{ or}$
- vi) $\frac{4}{3} < \gamma \leq 2, -\sqrt{6} < N < 0, M > \frac{2(N^2-3\gamma)}{(3\gamma-4)N}, \text{ or}$
- vii) $\frac{4}{3} < \gamma \leq 2, N > \sqrt{6}, M \leq -\frac{\sqrt{6}\gamma-2N}{3\gamma-4}, \text{ or}$
- viii) $1 \leq \gamma < \frac{4}{3}, N = -\sqrt{6}, M < \frac{\sqrt{6}(\gamma-2)}{3\gamma-4}, \text{ or}$
- ix) $1 \leq \gamma < \frac{4}{3}, N > \sqrt{6}, M \geq -\frac{\sqrt{6}\gamma-2N}{3\gamma-4}, \text{ or}$
- x) $1 \leq \gamma < \frac{4}{3}, N < -\sqrt{6}, M \leq \frac{\sqrt{6}\gamma+2N}{3\gamma-4}, \text{ or}$
- xi) $1 \leq \gamma < \frac{4}{3}, 0 < N \leq \sqrt{6}, M > \frac{2(N^2-3\gamma)}{(3\gamma-4)N}, \text{ or}$
- xii) $1 \leq \gamma < \frac{4}{3}, -\sqrt{6} < N < 0, M < \frac{2(N^2-3\gamma)}{(3\gamma-4)N}.$

The situation of physical interest is when it is non-hyperbolic with a 4D stable manifold for:

- i) $\gamma = \frac{4}{3}$, $2 < N \leq 2.06559$, $p > 4$, or
- ii) $\gamma = \frac{4}{3}$, $-2.06559 \leq N < -2$, $p > 4$, or
- iii) $\gamma = 1$, $N = -2.42061$, $M = M_{1,1}$, $p > \frac{6N}{M+2N}$, or
- iv) $\gamma = 1$, $N = 2.42061$, $M = -2.13014$, $p > \frac{6N}{M+2N}$, or
- v) $\gamma = 1$, $N = 2.44949$, $\frac{6-2N^2}{N} < M \leq M_{1,3}$, $p > \frac{6N}{M+2N}$, or
- vi) $\gamma = 1$, $N \leq -2.44949$, $M_{1,1} \leq M < -\sqrt{N^2 - 6} - N$, $p > \frac{6N}{M+2N}$, or
- vii) $\gamma = 1$, $N > 2.44949$, $\sqrt{N^2 - 6} - N < M \leq M_{1,1}$, $p > \frac{6N}{M+2N}$, or
- viii) $\gamma = 1$, $2.42061 < N < 2.44949$, $M_{1,2} \leq M \leq M_{1,3}$, $p > \frac{6N}{M+2N}$, or
- ix) $\gamma = 1$, $-2.44949 < N < -2.42061$, $M_{1,1} \leq M \leq M_{1,2}$, $p > \frac{6N}{M+2N}$, or
- x) $\gamma = 1$, $-2.42061 < N < 0$, $M_{1,1} \leq M < \frac{6-2N^2}{N}$, $p > \frac{6N}{M+2N}$, or
- xi) $\gamma = 1$, $-2.44949 < N \leq -2.42061$, $M_{1,3} \leq M < \frac{6-2N^2}{N}$, $p > \frac{6N}{M+2N}$, or
- xii) $\gamma = 1$, $0 < N < 2.44949$, $\frac{6-2N^2}{N} < M \leq M_{1,1}$, $p > \frac{6N}{M+2N}$, or
- xiii) $\gamma = 1.62562$, $N > 2.44949$, $M_{2,1} \leq M < \frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xiv) $1 < \gamma < \frac{9}{8}$, $N = N_{3,2}$, $M = M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xv) $1 < \gamma < \frac{9}{8}$, $N = N_{3,3}$, $M = M_{2,2}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xvi) $\gamma = 1.62562$, $N < N_{3,1}$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{-3\gamma M+4M+2N}$, or
- xvii) $\gamma = 1.62562$, $-2 < N < 0$, $\frac{2N^2-6\gamma}{3\gamma N-4N} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xviii) $\gamma = 1.62562$, $N_{3,1} \leq N < -2.44949$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xix) $\gamma = 1.62562$, $0 < N < 2.44949$, $M_{2,1} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{-3\gamma M+4M+2N}$, or
- xx) $1 < \gamma < \frac{9}{8}$, $N = -2.44949$, $M_{2,1} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxi) $\frac{4}{3} < \gamma < 1.62562$, $N > 2.44949$, $M_{2,1} \leq M < \frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxii) $1.62562 < \gamma < 2$, $N > 2.44949$, $M_{2,1} \leq M < \frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxiii) $1 < \gamma < \frac{9}{8}$, $N < -2.44949$, $M_{2,1} \leq M < \frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxiv) $\frac{9}{8} < \gamma < \frac{4}{3}$, $N < -2.44949$, $M_{2,1} \leq M < \frac{N}{3\gamma-4} - \sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxv) $1 < \gamma < \frac{9}{8}$, $N = N_{3,2}$, $M_{2,3} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxvi) $\gamma = 1.62562$, $-2.44949 < N \leq -2$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxvii) $1 < \gamma < \frac{9}{8}$, $N > N_{3,4}$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$,

- or
- xxviii) $1 < \gamma < \frac{9}{8}$, $N = N_{3,3}$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxix) $1 < \gamma < \frac{9}{8}$, $N = 2.44949$, $\frac{2N^2-6\gamma}{3\gamma N-4N} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxx) $\frac{9}{8} < \gamma < \frac{4}{3}$, $N > N_{3,4}$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$,
or
- xxxi) $\frac{4}{3} < \gamma < 1.62562$, $N < N_{3,1}$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{-3\gamma M+4M+2N}$, or
- xxxii) $1.62562 < \gamma < 2$, $N < N_{3,1}$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxxiii) $\frac{4}{3} < \gamma < 1.62562$, $N_{3,1} \leq N < -2.44949$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxxiv) $1.62562 < \gamma < 2$, $N_{3,1} \leq N < -2.44949$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{-3\gamma M+4M+2N}$, or
- xxxv) $\frac{4}{3} < \gamma < 1.62562$, $0 < N < 2.44949$, $M_{2,1} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxxvi) $1.62562 < \gamma < 2$, $0 < N < 2.44949$, $M_{2,1} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$,
or
- xxxvii) $\frac{9}{8} < \gamma < \frac{4}{3}$, $-2.44949 < N < 0$, $M_{2,1} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxxviii) $1 < \gamma < \frac{9}{8}$, $N_{3,2} < N < 0$, $M_{2,1} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xxxix) $1 < \gamma < \frac{9}{8}$, $-2.44949 < N < N_{3,2}$, $M_{2,1} \leq M \leq M_{2,2}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xli) $1 < \gamma < \frac{9}{8}$, $0 < N < N_{3,3}$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{-3\gamma M+4M+2N}$, or
- xlii) $\frac{9}{8} < \gamma < \frac{4}{3}$, $0 < N < N_{3,3}$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xlii) $1 < \gamma < \frac{9}{8}$, $-2.44949 < N < N_{3,2}$, $M_{2,3} \leq M < \frac{2N^2-6\gamma}{N(3\gamma-4)}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$,
or
- xliii) $1 < \gamma < \frac{9}{8}$, $2.44949 < N \leq N_{3,4}$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xliv) $\frac{9}{8} < \gamma < \frac{4}{3}$, $2.44949 < N \leq N_{3,4}$, $\sqrt{\frac{6\gamma^2-12\gamma+N^2}{(3\gamma-4)^2}} + \frac{N}{3\gamma-4} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xlv) $\frac{4}{3} < \gamma < 1.62562$, $-2.44949 < N \leq N_{3,2}$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xlvi) $1.62562 < \gamma < 2$, $-2.44949 < N \leq N_{3,2}$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xlvii) $1 < \gamma < \frac{9}{8}$, $N_{3,3} < N < 2.44949$, $M_{2,2} \leq M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xlviii) $1 < \gamma < \frac{9}{8}$, $N_{3,3} < N < 2.44949$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- xlix) $\frac{4}{3} < \gamma < 1.62562$, $N_{3,2} < N < 0$, $\frac{2N^2-6\gamma}{3\gamma N-4N} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- l) $1.62562 < \gamma < 2$, $N_{3,2} < N < 0$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,1}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or
- li) $\frac{9}{8} < \gamma < \frac{4}{3}$, $N_{3,3} \leq N < 2.44949$, $\frac{2N^2-6\gamma}{N(3\gamma-4)} < M \leq M_{2,3}$, $p > \frac{6\gamma N}{M(4-3\gamma)+2N}$, or

- lii) $\gamma = \frac{9}{8}$, $N > 2.61227$, $\frac{1}{5}\sqrt{64N^2 - 378} - \frac{8}{5}N < M \leq M_{4,1}$, $p > \frac{54N}{5M+16N}$, or
- liii) $\gamma = \frac{9}{8}$, $N < -2.44949$, $M_{4,1} \leq M < -\frac{1}{5}\sqrt{64N^2 - 378} - \frac{8}{5}N$, $p > \frac{54N}{5M+16N}$, or
- liv) $\gamma = \frac{9}{8}$, $2.44949 < N \leq 2.61227$, $\frac{1}{5}\sqrt{64N^2 - 378} - \frac{8}{5}N < M \leq M_{4,3}$, $p > \frac{54N}{5M+16N}$, or
- lv) $\gamma = \frac{9}{8}$, $-2.44949 < N < 0$, $M_{4,1} \leq M < \frac{(54-16N^2)}{5N}$, $p > \frac{54N}{5M+16N}$, or
- lvi) $\gamma = \frac{9}{8}$, $0 < N < 2.44949$, $\frac{(54-16N^2)}{5N} < M \leq M_{4,1}$, $p > \frac{54N}{5M+16N}$, or
- lvii) $\gamma = 2$, $0 < N < 2.44949$, $\frac{(8N^2-27)}{8N} - \frac{1}{8}\sqrt{\frac{729-48N^2}{N^2}} \leq M < \frac{N^2-6}{N}$, $p > \frac{6N}{N-M}$, or
- lviii) $\gamma = 2$, $-2.44949 < N < 0$, $\frac{N^2-6}{N} < M \leq \frac{1}{8}\sqrt{\frac{729-48N^2}{N^2}} + \frac{(8N^2-27)}{8N}$, $p > \frac{6N}{N-M}$.

Where $M_{1,1}$, $M_{1,2}$, $M_{1,3}$ are the first, the second and the third root of the polynomial $P_1(M) = 2M^3N + M^2(8N^2 - 15) + M(8N^3 - 18N) + 21N^2 - 72$, respectively. $M_{2,1}$, $M_{2,2}$ and $M_{2,3}$ are the first, the second and the third root of the polynomial $P_2(M) = -72\gamma^3 + 144\gamma^2 + M^3(54\gamma^3N - 216\gamma^2N + 288\gamma N - 128N) + M^2(108\gamma^3 - 261\gamma^2 + 120\gamma - 72\gamma^2N^2 + 192\gamma N^2 - 128N^2 + 48) + M(24\gamma N^3 - 32N^3 - 36\gamma^3N + 66\gamma^2N - 60\gamma N + 48N) + 27\gamma^2N^2 - 60\gamma N^2 + 12N^2$, respectively. $N_{3,1}$, $N_{3,2}$, $N_{3,3}$ and $N_{3,4}$ are the first, the second, the third and the fourth root of the polynomial: $P_3(N) = -1152\gamma^4 + 1440\gamma^3 + 1512\gamma^2 + 414\gamma + 64N^4 + (321\gamma^2 - 1176\gamma + 96)N^2 + 36$, respectively. Finally, $M_{4,1}$ is first root and $M_{4,3}$ is the third root of the polynomial: $P_4(M) = 125M^3N + M^2(800N^2 - 1650) + M(1280N^3 - 3270N) + 5460N^2 - 20412$.

- [1] P. Jordan, "Research on the Theory of General Relativity". 1958.
- [2] C. Brans and R. H. Dicke, Phys. Rev. **124**, 925 (1961).
- [3] A. H. Guth, Phys. Rev. D **23**, 347 (1981) [Adv. Ser. Astrophys. Cosmol. **3**, 139 (1987)].
- [4] G. W. Horndeski, Int. J. Theor. Phys. **10**, 363 (1974).
- [5] S. Bahamonde, K. F. Dialektopoulos and J. Levi Said, Phys. Rev. D **100**, no.6, 064018 (2019).
- [6] S. Bahamonde, K. F. Dialektopoulos, V. Gakis and J. Levi Said, Phys. Rev. D **101**, no.8, 084060 (2020).
- [7] S. Bahamonde, K. F. Dialektopoulos, M. Hohmann and J. Levi Said, [arXiv:2003.11554 [gr-qc]].
- [8] E. J. Copeland, E. W. Kolb, A. R. Liddle and J. E. Lidsey, Phys. Rev. D **48**, 2529 (1993).
- [9] J. Ibanez, R. J. van den Hoogen and A. A. Coley, Phys. Rev. D **51**, 928 (1995).
- [10] L. P. Chimento and A. S. Jakubi, Int. J. Mod. Phys. D **5**, 71 (1996).
- [11] J. E. Lidsey, A. R. Liddle, E. W. Kolb, E. J. Copeland, T. Barreiro and M. Abney, Rev. Mod. Phys. **69** (1997) 373.
- [12] A. A. Coley, J. Ibanez and R. J. van den Hoogen, J. Math. Phys. **38**, 5256 (1997).
- [13] E. J. Copeland, I. J. Grivell, E. W. Kolb and A. R. Liddle, Phys. Rev. D **58**, 043002 (1998).
- [14] A. A. Coley and R. J. van den Hoogen, Phys. Rev. D **62**, 023517 (2000).
- [15] A. A. Coley, "Dynamical systems in cosmology," gr-qc/9910074.
- [16] A. Coley, "Dynamical systems and cosmology," Astrophys. Space Sci. Libr. **291** (2003).
- [17] Wainwright, J., and Ellis, G. F. R., Eds. Dynamical Systems in Cosmology. Cambridge Univ. Press, Cambridge, 1997.
- [18] A. Coley and M. Goliath, Class. Quant. Grav. **17**, 2557 (2000).
- [19] A. Coley and M. Goliath, Phys. Rev. D **62**, 043526 (2000).
- [20] A. Coley and Y. J. He, Gen. Rel. Grav. **35**, 707 (2003).
- [21] E. Elizalde, S. Nojiri and S. D. Odintsov, Phys. Rev. D **70** (2004) 043539.
- [22] S. Capozziello, S. Nojiri and S. D. Odintsov, Phys. Lett. B **632** (2006) 597.
- [23] R. Curbelo, T. Gonzalez, G. Leon and I. Quiros, Class. Quant. Grav. **23**, 1585 (2006).
- [24] T. Gonzalez, G. Leon and I. Quiros, astro-ph/0502383.
- [25] T. Gonzalez, G. Leon and I. Quiros, Class. Quant. Grav. **23**, 3165 (2006).
- [26] R. Lazkoz and G. Leon, Phys. Lett. B **638**, 303-309 (2006).
- [27] R. Lazkoz, G. Leon and I. Quiros, Phys. Lett. B **649**, 103 (2007).

- [28] E. Elizalde, S. Nojiri, S. D. Odintsov, D. Saez-Gomez and V. Faraoni, Phys. Rev. D **77** (2008) 106005.
- [29] G. Leon and E. N. Saridakis, Phys. Lett. B **693**, 1 (2010).
- [30] G. Leon and E. N. Saridakis, JCAP **0911**, 006 (2009).
- [31] G. Leon, Y. Leyva, E. N. Saridakis, O. Martin and R. Cardenas, Falsifying Field-based Dark Energy Models, in Dark Energy: Theories, Developments, and Implications, Nova Science Publishers, (2010). arXiv:0912.0542 [gr-qc].
- [32] G. Leon and E. N. Saridakis, Class. Quant. Grav. **28**, 065008 (2011).
- [33] S. Basilakos, M. Tsamparlis and A. Paliathanasis, Phys. Rev. D **83**, 103512 (2011).
- [34] C. Xu, E. N. Saridakis and G. Leon, JCAP **1207**, 005 (2012).
- [35] G. Leon and E. N. Saridakis, JCAP **1303**, 025 (2013).
- [36] G. Leon, J. Saavedra and E. N. Saridakis, Class. Quant. Grav. **30**, 135001 (2013).
- [37] C. R. Fadragas, G. Leon and E. N. Saridakis, Class. Quant. Grav. **31**, 075018 (2014).
- [38] G. Kofinas, G. Leon and E. N. Saridakis, Class. Quant. Grav. **31**, 175011 (2014).
- [39] G. Leon and E. N. Saridakis, JCAP **1504**, 031 (2015).
- [40] A. Paliathanasis and M. Tsamparlis, Phys. Rev. D **90**, no. 4, 043529 (2014).
- [41] R. De Arcia, T. Gonzalez, G. Leon, U. Nucamendi and I. Quiros, Class. Quant. Grav. **33**, no. 12, 125036 (2016).
- [42] A. Paliathanasis, M. Tsamparlis, S. Basilakos and J. D. Barrow, Phys. Rev. D **91**, no. 12, 123535 (2015).
- [43] G. Leon and E. N. Saridakis, JCAP **1511**, 009 (2015).
- [44] J. D. Barrow and A. Paliathanasis, Phys. Rev. D **94**, no. 8, 083518 (2016).
- [45] J. D. Barrow and A. Paliathanasis, Gen. Rel. Grav. **50**, no. 7, 82 (2018).
- [46] M. Cruz, A. Ganguly, R. Gannouji, G. Leon and E. N. Saridakis, Class. Quant. Grav. **34**, no. 12, 125014 (2017).
- [47] A. Paliathanasis, Mod. Phys. Lett. A **32**, no. 37, 1750206 (2017).
- [48] B. Alhulaimi, R. J. Van Den Hoogen and A. A. Coley, JCAP **1712**, 045 (2017).
- [49] N. Dimakis, A. Giacomini, S. Jamal, G. Leon and A. Paliathanasis, Phys. Rev. D **95**, no. 6, 064031 (2017).
- [50] A. Giacomini, S. Jamal, G. Leon, A. Paliathanasis and J. Saavedra, Phys. Rev. D **95**, no. 12, 124060 (2017).
- [51] L. Karpathopoulos, S. Basilakos, G. Leon, A. Paliathanasis and M. Tsamparlis, Gen. Rel. Grav. **50**, no. 7, 79 (2018).
- [52] R. De Arcia, T. Gonzalez, F. A. Horta-Rangel, G. Leon, U. Nucamendi and I. Quiros, Class. Quant. Grav. **35**, no. 14, 145001 (2018).
- [53] M. Tsamparlis and A. Paliathanasis, Symmetry **10**, no. 7, 233 (2018).
- [54] A. Paliathanasis, G. Leon and S. Pan, Gen. Rel. Grav. **51**, no. 9, 106 (2019).
- [55] S. Basilakos, G. Leon, G. Papagiannopoulos and E. N. Saridakis, Phys. Rev. D **100**, no. 4, 043524 (2019).
- [56] R. J. Van Den Hoogen, A. A. Coley, B. Alhulaimi, S. Mohandas, E. Knighton and S. O’Neil, JCAP **1811**, 017 (2018).
- [57] G. Leon, A. Paliathanasis and J. L. Morales-Martinez, Eur. Phys. J. C **78**, no. 9, 753 (2018).
- [58] G. Leon, A. Paliathanasis and L. Velázquez Abab, Gen Relativ Gravit **52**, 71 (2020).
- [59] G. Leon and A. Paliathanasis, Eur. Phys. J. C **79**, no. 9, 746 (2019).
- [60] A. Paliathanasis and G. Leon, Zeitschrift fÄijr Naturforschung A, Volume 75, Issue 6, Pages 523–532 (2020).
- [61] G. Leon, A. Coley and A. Paliathanasis, Annals Phys. **412**, 168002 (2020).
- [62] A. Paliathanasis, G. Papagiannopoulos, S. Basilakos and J. D. Barrow, Eur. Phys. J. C **79**, no. 8, 723 (2019).
- [63] J. D. Barrow and A. Paliathanasis, Eur. Phys. J. C **78**, no. 9, 767 (2018).
- [64] I. Quiros, Int. J. Mod. Phys. D **28**, no. 07, 1930012 (2019).
- [65] G. Leon and F. O. F. Silva, [arXiv:1912.09856 [gr-qc]].
- [66] A. Giacomini, G. Leon, A. Paliathanasis and S. Pan, [arXiv:2008.01395 [gr-qc]].
- [67] A. Paliathanasis and G. Leon, [arXiv:2007.13223 [gr-qc]].
- [68] A. Giacomini, G. Leon, A. Paliathanasis and S. Pan, Eur. Phys. J. C **80**, no. 3, 184 (2020).
- [69] E. J. Copeland, A. R. Liddle and D. Wands, Phys. Rev. D **57**, 4686–4690 (1998).
- [70] S. Foster, Class. Quant. Grav. **15**, 3485 (1998).
- [71] J. Miritzis, Class. Quant. Grav. **20**, 2981 (2003).
- [72] D. Gonzalez Morales, Y. Napoles Alvarez, Quintaesencia con acoplamiento no minimo a la materia oscura desde la perspectiva de los sistemas dinamicos, Bachelor Thesis, Universidad Central Marta Abreu de Las Villas, 2008.
- [73] G. Leon, Class. Quant. Grav. **26**, 035008 (2009).
- [74] R. Giambò and J. Miritzis, Class. Quant. Grav. **27**, 095003 (2010).
- [75] M. Shahalam, R. Myrzakulov and M. Y. Khlopov, Gen. Rel. Grav. **51**, no. 9, 125 (2019).
- [76] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, Annals Phys. **418**, 168186 (2020).
- [77] F. Humieja and M. SzydÅCwski, Eur. Phys. J. C **79**, no. 9, 794 (2019).
- [78] J. Matsumoto and S. V. Sushkov, JCAP **1801**, 040 (2018).
- [79] J. Matsumoto and S. V. Sushkov, JCAP **1511**, 047 (2015).
- [80] A. R. Solomon, “Cosmology Beyond Einstein,” doi:10.1007/978-3-319-46621-7 arXiv:1508.06859 [gr-qc].

- [81] T. Harko, F. S. N. Lobo, J. P. Mimoso and D. Pavon, Eur. Phys. J. C **75**, 386 (2015).
- [82] O. Minazzoli and A. Hees, Phys. Rev. D **90**, 023017 (2014).
- [83] M. A. Skugoreva, S. V. Sushkov and A. V. Toporensky, Phys. Rev. D **88**, 083539 (2013) Erratum: [Phys. Rev. D **88**, no. 10, 109906 (2013)].
- [84] M. Jamil, D. Momeni and R. Myrzakulov, Eur. Phys. J. C **72**, 2075 (2012).
- [85] J. Miritzis, J. Phys. Conf. Ser. **283**, 012024 (2011).
- [86] O. Hrycyna and M. Szydlowski, Phys. Rev. D **76**, 123510 (2007).
- [87] R. Giambò, F. Giannoni and G. Magli, Gen. Rel. Grav. **41**, 21 (2009).
- [88] G. Leon, P. Silveira and C. R. Fadragas, arXiv:1009.0689 [gr-qc]. In book: Classical and Quantum Gravity: Theory, Analysis and Applications. Chapter: 10 Publisher: Nova Science Publishing Inc. Editors: Vincent R. Frignanni, New York, 2010.
- [89] G. Leon Torres, “Qualitative analysis and characterization of two cosmologies including scalar fields,” arXiv:1412.5665 [gr-qc]. Phd Thesis, Universidad Central Marta Abreu de Las Villas, 2010.
- [90] G. Leon and C. R. Fadragas, Dynamical Systems: And Their Applications. LAP Lambert Academic Publishing, 2012, arXiv:1412.5701 [gr-qc].
- [91] C. R. Fadragas and G. Leon, Class. Quant. Grav. **31**, no. 19, 195011 (2014).
- [92] K. Tzanni and J. Miritzis, Phys. Rev. D **89**, no. 10, 103540 (2014) Addendum: [Phys. Rev. D **89**, no. 12, 129902 (2014)].
- [93] G. Leon, F. O. F. Silva, Generalized scalar field cosmologies: Theorems on asymptotic behavior.
- [94] N. Kaloper and K. A. Olive, Phys. Rev. D **57**, 811 (1998).
- [95] T. Gonzalez and I. Quiros, Class. Quant. Grav. **25**, 175019 (2008).
- [96] M. Sharma, M. Shahalam, Q. Wu and A. Wang, JCAP **1811**, 003 (2018).
- [97] R. Giambà, J. Miritzis and A. Pezzola, Eur. Phys. J. Plus **135**, no.4, 367 (2020)
- [98] A. Alho and C. Uggla, J. Math. Phys. **56**, no. 1, 012502 (2015).
- [99] D. Fajman, G. Heijzel and M. Maliborski, Class. Quant. Grav. **37**, no.13, 135009 (2020).
- [100] A. Alho, J. Hell and C. Uggla, Class. Quant. Grav. **32**, no.14, 145005 (2015).
- [101] A. Alho and C. Uggla, Phys. Rev. D **95**, no.8, 083517 (2017).
- [102] J. Yearsley and J. D. Barrow, Class. Quant. Grav. **13**, 2693-2706 (1996).
- [103] S. A. Pavluchenko, Phys. Rev. D **67**, 103518 (2003).
- [104] R. Cardenas, T. Gonzalez, Y. Leiva, O. Martin and I. Quiros, Phys. Rev. D **67**, 083501 (2003).
- [105] B. Ratra and P. Peebles, Phys. Rev. D **37**, 3406 (1988).
- [106] C. Wetterich, Nucl. Phys. B **302**, 668-696 (1988).
- [107] T. Matos, J. R. Luevano, I. Quiros, L. Urena-Lopez and J. A. Vazquez, Phys. Rev. D **80**, 123521 (2009).
- [108] E. J. Copeland, S. Mizuno and M. Shaeri, Phys. Rev. D **79**, 103515 (2009).
- [109] Y. Leyva, D. Gonzalez, T. Gonzalez, T. Matos and I. Quiros, Phys. Rev. D **80**, 044026 (2009).
- [110] S. del Campo, C. R. Fadragas, R. Herrera, C. Leiva, G. Leon and J. Saavedra, Phys. Rev. D **88**, 023532 (2013).
- [111] V. Sahni and L. M. Wang, Phys. Rev. D **62**, 103517 (2000).
- [112] V. Sahni and A. A. Starobinsky, Int. J. Mod. Phys. D **9**, 373-444 (2000).
- [113] J. E. Lidsey, T. Matos and L. A. Urena-Lopez, Phys. Rev. D **66**, 023514 (2002).
- [114] T. Matos and L. Urena-Lopez, Class. Quant. Grav. **17**, L75-L81 (2000).
- [115] L. Urena-Lopez and T. Matos, Phys. Rev. D **62**, 081302 (2000).
- [116] T. Barreiro, E. J. Copeland and N. Nunes, Phys. Rev. D **61**, 127301 (2000).
- [117] T. Gonzalez, R. Cardenas, I. Quiros and Y. Leyva, Astrophys. Space Sci. **310**, 13-18 (2007).
- [118] A. D. Rendall, Class. Quant. Grav. **24**, 667-678 (2007).