

Optimal sample size for the Birnbaum-Saunders distribution under a decision-theoretic approach

Eliardo G. Costa^{1*} and Manoel Santos-Neto²

¹Departamento de Estatística, Universidade Federal do Rio Grande do Norte, Brazil

²Departamento de Estatística, Universidade Federal de Campina Grande, Brazil

April 9, 2024

Abstract

The Birnbaum-Saunders distribution has been widely applied in several areas of science and although several methodologies related to this distribution have been proposed, the problem of determining the optimal sample size for estimating its mean has not yet been studied. For this purpose, we propose a methodology to determine the optimal sample size under a decision-theoretic approach. In this approach, we consider loss functions for point and interval inference. Finally, computational tools in the R language were developed to use in practice.

Keywords: inverse gamma distribution; loss function; Bayes risk; sampling cost.

1 Introduction

Birnbaum & Saunders (1969b) introduced a family of distributions to model failure times for metals subject to periodic stress and provided a natural physical justification for this family. This family is the so-called Birnbaum-Saunders (BS) distribution. In the last decades, this distribution has received considerable attention in the literature and many methodologies have been proposed for parameter inference. Such attention is justified by its wide applicability

*Corresponding Author: Eliardo G. Costa. Departamento de Estatística, Universidade Federal do Rio Grande do Norte, Brazil. Email: eliardocosta@ccet.ufrn.br

and its variations have been applied in several areas including finance, business, engineering, environmental, medicine, quality control and many others. A detailed review of the BS distribution including methodologies under the frequentist and Bayesian approaches is presented in Balakrishnan & Kundu (2019).

Although several methodologies related to this distribution have been proposed, the problem of determining an optimal size for estimating the mean of the BS distribution has not yet been studied. Recently, Bourguignon et al. (2020) presented guidelines about the minimum sample size for monitoring the median parameter of the BS distribution in the context of quality control under a frequentist approach. In this way, we develop a methodology via a Bayesian decision-theoretic approach based on a criterion that minimizes the sum of the Bayes risk and the sampling cost. The proposed approach depends on an *ad hoc* loss function defined to accommodate the implications of a decision. We consider four different loss functions for point and interval inference, two for each type of inference. Using the same approach but for other models there is a considerable literature, see for example Etzioni & Kadane (1993), Sahu & Smith (2006), Parmigiani & Inoue (2009), Islam & Pettit (2012, 2014), De Santis & Gubbiotti (2016), Costa (2017) and references therein.

The paper unfolds as follows. In Section 2 we discuss the Bayesian model and the inference of the parameters of the BS distribution. In Section 3 we present the methodology to obtain the optimal sample size under a decision-theoretic approach. Finally, we conclude with a discussion of the results in Section 4.

2 Bayesian model

Much of the information about the BS distribution presented in this section has been gathered from other works, for example, Birnbaum & Saunders (1969b,a), Leiva (2016) and Balakrishnan & Kundu (2019). Let X be a BS distribution with a scale parameter β and a shape parameter α , we denote by $X \sim \text{BS}(\alpha, \beta)$. Then, the respective probability density function is given by

$$f_X(x|\alpha, \beta) = \frac{1}{\sqrt{2\pi}} \exp \left[-\frac{1}{2\alpha^2} \left(\frac{x}{\beta} + \frac{\beta}{x} - 2 \right) \right] \frac{(x + \beta)}{2\alpha\sqrt{\beta x^3}}, \quad x \in \mathbb{R}_{>0}, \quad \alpha, \beta \in \mathbb{R}_{>0}.$$

Besides being the scale parameter, the parameter β is also the median of this distribution. Furthermore, the mean and the variance of the BS distribution are given by

$$\theta := \mathbb{E}[X] = \beta \left(1 + \frac{\alpha^2}{2} \right) \quad \text{and} \quad \text{Var}[X] = (\alpha\beta)^2 \left(1 + \frac{5\alpha^2}{4} \right). \quad (1)$$

Also, if X is Birnbaum-Saunders distributed then

$$X = \frac{\beta}{4} \left(\alpha Z + \sqrt{(\alpha Z)^2 + 4} \right)^2, \quad (2)$$

where Z follows a standard normal distribution, which is useful to draw values from the $\text{BS}(\alpha, \beta)$ distribution. Given a sample $\mathbf{x}_n = (x_1, \dots, x_n)$, the likelihood function from the $\text{BS}(\alpha, \beta)$ satisfies

$$\mathcal{L}(\alpha, \beta; \mathbf{x}_n) \propto \frac{1}{(\alpha\beta)^n} \prod_{i=1}^n \left[\left(\frac{\beta}{x_i} \right)^{1/2} + \left(\frac{\beta}{x_i} \right)^{3/2} \right] \exp \left[-\frac{1}{2\alpha^2} \sum_{i=1}^n \left(\frac{x_i}{\beta} + \frac{\beta}{x_i} - 2 \right) \right].$$

For the parameters α and β of the model, we consider proper prior distributions because the use of noninformative prior distributions yields an improper posterior distribution and continuous conjugate priors do not exist (Wang et al., 2016). A possible choice for a prior distribution for β is the inverse gamma distribution in which the density satisfies

$$\pi(\beta) \propto \beta^{-(a_1+1)} \exp \left(-\frac{b_1}{\beta} \right), \quad \beta \in \mathfrak{R}_{>0},$$

where a_1 and b_1 are positive and known constants (hyperparameters), we denote by $\beta \sim \text{IG}(a_1, b_1)$. We also assume a inverse gamma prior distribution for α^2 with hyperparameters a_2 and b_2 . Thus, we may write the model hierarchically as follows

$$\begin{aligned} X_i | \alpha, \beta &\stackrel{\text{iid}}{\sim} \text{BS}(\alpha, \beta), \quad i = 1, 2, \dots, n; \\ \beta &\sim \text{IG}(a_1, b_1) \quad \text{and} \quad \alpha^2 \sim \text{IG}(a_2, b_2). \end{aligned}$$

In this context, the conditional posterior distribution of α^2 given β and \mathbf{x}_n is

$$\alpha^2 | \beta, \mathbf{x}_n \sim \text{IG} \left(\frac{n+1}{2} + a_2, \frac{1}{2} \sum_{i=1}^n \left(\frac{x_i}{\beta} + \frac{\beta}{x_i} - 2 \right) + b_2 \right), \quad (3)$$

and the marginal posterior distribution of β given \mathbf{x}_n satisfies

$$\pi(\beta | \mathbf{x}_n) \propto \beta^{-(n+a_1+1)} \exp \left(\frac{b_1}{\beta} \right) \prod_{i=1}^n \left[\left(\frac{\beta}{x_i} \right)^{1/2} + \left(\frac{\beta}{x_i} \right)^{3/2} \right] \left[\frac{1}{2} \sum_{i=1}^n \left(\frac{x_i}{\beta} + \frac{\beta}{x_i} - 2 \right) + b_2 \right]^{-\frac{n+1}{2}-a_2}, \quad (4)$$

which is not a known distribution (Wang et al., 2016). In this way, we use the random walk Metropolis-Hastings algorithm (Metropolis et al., 1953, Hastings, 1970) to draw samples from the marginal posterior distribution of β given \mathbf{x}_n . Using this sampling algorithm and the

posterior distribution in (3) we may draw values from the joint posterior distribution of α^2 and β . For a given \mathbf{x}_n , first we draw values of β from (4) and given these values we draw values of α^2 using (3). Note that the parameter of interest θ is the mean of the BS distribution and is a function of α^2 and β . In order to obtain a random sample of the posterior distribution of θ given \mathbf{x}_n , we may draw values from the joint posterior of α^2 and β , then apply (1) in each sampled pair of values.

3 Optimal sample size

We may approach the problem of determining the optimal sample size as a decision problem (see Raiffa & Schlaifer, 1961, Parmigiani & Inoue, 2009, for example). Given that θ is the parameter of interest, we specify a loss function $L(\theta, d_n)$ based on a sample $\mathbf{X}_n = (X_1, \dots, X_n)$ and a decision function $d_n \equiv d_n(\mathbf{X}_n)$. For a given n and depending on the adopted loss function, the action $d_n(\mathbf{x}_n)$ consists of the specification of one quantity (point inference case) representing an estimate for θ , or two quantities (interval inference case) representing the lower and upper limits of a credible interval for θ . Let π be a prior distribution for the unknown parameter θ and d_n a decision function; the Bayes risk is (Parmigiani & Inoue, 2009)

$$r(\pi, d_n) := \int_{\Theta} \int_{\mathcal{X}^n} L(\theta, d_n) g(\mathbf{x}_n | \theta) \pi(\theta) d\mathbf{x}_n d\theta, \quad (5)$$

where $g(\cdot)$ is the sampling distribution for \mathbf{X}_n given θ , Θ is the parameter space, and \mathcal{X}_n is the sample space. The decision d_n^* that minimizes $r(\pi, d_n)$ among all the possible decisions d_n is called the Bayes rule. In this context, we define the optimal sample size as the one that minimizes the total cost

$$TC(n) := r(\pi, d_n^*) + C(n),$$

where $C(n)$ is the sampling cost function. Here, we take $C(n) = cn$, where c is the per-unit cost for observing a unit in the population. Since it is not possible to compute $r(\pi, d_n^*)$ analytically, we use Monte Carlo simulations as an alternative to estimate $TC(n)$ for each n . Suppose that the order of the integration may be reverted in (5), then we have

$$r(\pi, d_n^*) = \int_{\mathcal{X}^n} \mathbb{E}[L(\theta, d_n^*) | \mathbf{x}_n] g(\mathbf{x}_n) d\mathbf{x}_n,$$

so that we may estimate the minimized Bayes risk through the posterior expected value of loss function applied to the Bayes rule d_n^* . This may be done as follows in the Algorithm 1.

After obtaining an estimate of $r(\pi, d_n^*)$ we add the respective cost sampling cn , which finally gives us an estimate of the total cost $TC(n)$ for a given n . We applied this procedure for

a grid of plausible values of n . For example, if we set this grid of values as $n = 2, 12, \dots, 82, 92$, then we obtain an estimate for $TC(2), TC(12), \dots, TC(82), TC(92)$. The choice of the grid of values is arbitrary and the smaller the span between its consecutive elements, the better to visualize the behavior of the total cost, but as we decrease this span the required computer processing power also increases, as well as the time to compute all these estimates. Thus, the choice of this grid must take into account all these settings.

In Step 4 of the Algorithm 1, when sampling from the marginal posterior distribution (4), we consider a burn-in of 500 iterations and a thinning of 20 with a final number of iterations of 500. We use these 500 iterations to compute an estimate of the Bayes risk. We inspect a trace and autocorrelation plot for a lower value of the grid used for n , we expect the same or better behavior as the n increases in the grid. All the trace plots showed a random behavior around a value and in all the autocorrelation plots the autocorrelations for almost every lag were zero. In each value of n in the grid, we estimate the Bayes risk ten times.

Costa (2017) propose to fit the following curve to the grid of values of n and the respective estimates of $TC(n)$, denoted by $tc(n)$

$$tc(n) = \frac{E}{(1+n)^G} + cn,$$

where E and G are parameters to be estimated. This curve may be linearized as a linear regression as follows

$$\log[tc(n) - cn] = \log E - G \log(1+n),$$

and the estimates of E and G may be computed by least squares. In this setting, the optimal

Algorithm 1:

- 1 Set values for the hyperparameters;
 - 2 Draw one value of α^2 and one value of β from the respective prior distributions, compute the square root of α^2 ;
 - 3 Given α and β , draw a value of X_i from the $BS(\alpha, \beta)$ using (2), for $i = 1, \dots, n$. This generates a sample $\mathbf{x}_n = (x_1, \dots, x_n)$;
 - 4 Given \mathbf{x}_n , draw a sample of size N (as large as possible) from the joint posterior distribution of α^2 and β as explained in Section 2. This generates values (α_j^2, β_j) , $j = 1, \dots, N$;
 - 5 For $j = 1, \dots, N$, compute the posterior values θ_j using the generated values in Step 4 and (1);
 - 6 Obtain the respective Bayes rule d_n^* using the sample of the posterior distribution of θ obtained in Step 5;
 - 7 Use the values generated in Step 5 to compute an estimate of $E[L(\theta, d_n^*)|\mathbf{x}_n]$;
 - 8 Repeat the Steps 1-7 K times (as large as possible), this generates K estimates of $E[L(\theta, d_n^*)|\mathbf{x}_n]$;
 - 9 Take the average of the K estimates obtained in Step 8, this is an estimate of $r(\pi, d_n^*)$.
-

sample size (n_o) is the nearest integer closest to

$$\left(\frac{\hat{E} \hat{G}}{c}\right)^{1/(\hat{G}+1)} - 1,$$

where \hat{E} and \hat{G} are, respectively, the least square estimates of E and G .

3.1 Loss functions

We adopted four loss functions, the loss functions 1 and 2 may be used for point inference, *i.e.*, the decision d_n provides a quantity representing an estimate for the parameter of interest θ . The loss functions 3 and 4 may be used for interval inference, in this case, a decision provides two quantities, the lower (say, a) and the upper (say, b) limits of a credible interval for θ .

3.1.1 Loss function 1 (L1)

The first loss function is

$$L(\theta, d_n) = |\theta - d_n|,$$

which is known as the absolute loss function. For this loss function the Bayes rule d_n^* is the median of the posterior distribution of θ . Given a sample θ_j , $j = 1, \dots, N$, of the posterior distribution of θ , an estimate of $\mathbb{E}[L(\theta, d_n^*)|\mathbf{x}_n]$ may be obtained from $N^{-1} \sum_{j=1}^N |\theta_j - d_n^*|$.

3.1.2 Loss function 2 (L2)

Second, we consider the well-known quadratic loss function

$$L(\theta, d_n) = (\theta - d_n)^2,$$

for this loss function the Bayes rule d_n^* corresponds to the posterior expected value of θ and in this case $\mathbb{E}[L(\theta, d_n^*)|\mathbf{x}_n] = \text{Var}(\theta|\mathbf{x}_n)$. Given a sample θ_j , $j = 1, \dots, N$, of the posterior distribution of θ , an estimate of $\mathbb{E}[L(\theta, d_n^*)|\mathbf{x}_n]$ may be obtained from the respective sample variance.

3.1.3 Loss function 3 (L3)

The third loss function is

$$L(\theta, d_n) = \rho\tau + (a - \theta)^+ + (\theta - b)^+, \tag{6}$$

where $0 < \rho < 1$ is a weight, $\tau = (b - a)/2$ is the half-length of the desired interval, the function x^+ is equal to x if $x > 0$ and equal to zero, otherwise. The smaller is τ the narrower the interval. The terms $(a - \theta)^+$ and $(\theta - b)^+$ are included to penalize intervals that do not contain the parameter of interest θ . These terms are equal to zero if $\theta \in [a, b]$ and increase as θ moves away from the interval. Note that the loss function (6) is a weighted sum of two terms, τ and $(a - \theta)^+ + (\theta - b)^+$, where the weights are ρ and 1, respectively. The Bayes rule d_n^* corresponds to taking a and b as the quantiles of probabilities $\rho/2$ and $1 - \rho/2$ of the posterior distribution of θ . For more details see Rice et al. (2008) or Costa (2017). If we consider this loss function applied to the Bayes rule, we have

$$\mathbb{E}[L(\theta, d_n^*)|\mathbf{x}_n] = \mathbb{E}[\theta\delta_\theta(A_{b^*})|\mathbf{x}_n] - \mathbb{E}[\theta\delta_\theta(A_{a^*})|\mathbf{x}_n],$$

where $A_{b^*} = [b^*, \infty)$, $A_{a^*} = (0, a^*]$, a^* and b^* are the corresponding bounds of the Bayes rule d_n^* and $\delta_\theta(\cdot)$ is the indicator function. Given a sample θ_j , $j = 1, \dots, N$, of the posterior distribution of θ , an estimate of $\mathbb{E}[L(\theta, d_n^*)|\mathbf{x}_n]$ may be obtained from $N^{-1} \sum_{j=1}^N [\theta_j \delta_{\theta_j}(A_{b^*}) - \theta_j \delta_{\theta_j}(A_{a^*})]$.

3.1.4 Loss function 4 (L4)

The last loss function is

$$L(\theta, d_n) = \gamma\tau + (\theta - m)^2/\tau,$$

where $\gamma > 0$ is a fixed constant and $m = (a + b)/2$ is the center of the credible interval. The first term involves the half-width of the interval and the second, the square of the distance between the parameter of interest θ and the center of the interval, which is divided by the half-width to maintain the same measurement unit of the first term.

The weights attributed to each term are γ and 1, respectively. If $\gamma < 1$, we attribute the largest weight to the second term; if $\gamma > 1$, the situation is reversed and if $\gamma = 1$ the two terms have the same weight. For this loss function, the Bayes rule d_n^* corresponds to the quantities which define the interval $[a^*, b^*] = [m^* - \text{SD}_\gamma, m^* + \text{SD}_\gamma]$, where $m^* = \mathbb{E}[\theta|\mathbf{x}_n]$ and $\text{SD}_\gamma = \gamma^{-1/2}[\text{Var}(\theta|\mathbf{x}_n)]^{1/2}$. For more details see Rice et al. (2008), Parmigiani & Inoue (2009) or Costa (2017). For this loss function, we have

$$\mathbb{E}[L(\theta, d_n^*)|\mathbf{x}_n] = 2\gamma^{1/2}\sqrt{\text{Var}(\theta|\mathbf{x}_n)}.$$

Given a sample θ_j , $j = 1, \dots, N$, of the posterior distribution of θ , an estimate of $\mathbb{E}[L(\theta, d_n^*)|\mathbf{x}_n]$ may be obtained from the respective sample variance and the previous equation.

For the hyperparameters of the prior distribution of β , we consider $b_1 = 50$ and $a_1 = 8, 10, 13$ and 15 , with these values we have different degrees of prior information, see Figure 1. For the prior distribution of α^2 , we set $a_2 = a_1$ and $b_2 = b_1$. We consider $c = 0.001, 0.01$ and 0.1 for the per-unit cost. For the loss function L3 we consider $\rho = 0.01, 0.05$ and 0.10 , while for L4 we consider $\gamma = 0.25, 0.50$ and 1.00 . For each combination of these values we compute the optimal sample size n_o for estimating θ . The average acceptance rate for the Metropolis-Hastings algorithm in all these combinations was $\approx 70\%$. Since the proposed methodology is based on simulation methods, we obtain n_o as triplicate and observe the difference between the three values. In Table 1 we present the optimal sample sizes computed with these settings.

An implementation of the proposed methodology is provided in the R language. The n_o may be obtained using the R package `samplesizeBS` (Costa & Santos-Neto, 2020). Also, the n_o may be obtained via the following link <https://santosneto.shinyapps.io/samplesizeBSapp/>, which also presents a graph with the fitted curve.




Figure 1: Density for different values of the hyperparameter a_1 ($b_1 = 50$) of the inverse gamma prior distribution for β .

4 Discussion

We propose a methodology to compute the optimal sample size for estimating the mean of the Birnbaum-Saunders distribution, a widely applied and studied distribution in several areas of science. We consider four different loss functions which allow to make both point and interval inference for the parameter of interest.

An advantage of the proposed methodology is that the per-unit cost, represented by c , is explicitly taken into account. When the cost c is fixed and $b_1 = 50$, the optimal sample size n_o decreases as the a_1 increases (or prior variance decreases) as expected, since in this case the prior knowledge increases as the a_1 increases. This occurs with all loss functions. For $b_1 = 50$ and a_1 fixed, the n_o also decreases as the c increases; however, the total sampling cost decreases. For example, if we take the loss function L1, $a_1 = 8$ and $c = 0.001$, the corresponding n_o is 651 (Table 1), which generates a total cost of $C(651) = 0.001 \times 651 = 0.651$, whereas if we take $c = 0.1$, the corresponding n_o is 23 (Table 1), which generates a total cost of $C(23) = 0.1 \times 23 = 2.3$. For the loss function L3, when ρ increases the n_o also increases, if we consider a_1 e c fixed. This makes sense because ρ is the weight attributed to the term τ in L3, this term is related to the length of the credible interval and when we increases ρ we expect longer credible intervals, consequently the probability of the respective interval decreases. The same is valid for γ in the loss function L4, but in this case the decreasing of the respective credible interval is easily noted by the presence of the term $\gamma^{-\frac{1}{2}}$ in the expression of the respective Bayes rule. When γ increases this term shrinks the length of the interval.

Since the proposed methodology is based on simulations, we obtain the n_o in triplicate for each scenario of values of a_1 , c , ρ and γ . We observe that the largest discrepancies in the scenarios occur for $a_1 = 8$, these discrepancies decrease as the a_1 increases, or when the prior variance decreases. This also occurs when $c = 0.001$ and/or when we consider the loss function L2. In general the discrepancy is close to zero, but if a large discrepancy occurs we suggest to inspect visually the graph of the fitted curves and take the value of n_o which corresponds to the best fit. However, if all the curves fit visually well, we suggest to use the median of the values obtained for n_o . In our case we obtained the values of n_o in triplicate. For example, in Figure 2 under the loss function L4 with $a_1 = 8$, $c = 0.001$ and $\gamma = 0.50$ the values of n_o were 1208, 1179 and 1183. Since there is a discrepancy between these values and the fitting of the curves were visually well, in this case we suggest to use $n_o = 1183$.

Finally, note that we have no entry in Table 1 in some scenarios, which means that it is not worth sampling in these cases because the sampling cost outweighs the decreasing of the minimized Bayes risk. This was also observed by Etzioni & Kadane (1993) and Islam & Pettit (2014).




Figure 2: Fitted curves with the respective optimal sample sizes obtained via the loss function L4 with $a_1 = 8$, $c = 0.001$ and $\gamma = 0.50$.

Acknowledgements

Research carried out using the computational resources of the Center for Mathematical Sciences Applied to Industry (CeMEAI) funded by Fundação de Amparo à Pesquisa do Estado de São Paulo (grant 2013/07375-0).

Table 1: Optimal sample sizes n_o (in triplicate) for estimating the mean of the BS distribution via four different loss functions.

ρ/γ	$a_1 = 8$			$a_1 = 10$			$a_1 = 13$			$a_1 = 15$		
	$c = 0.001$	$c = 0.01$	$c = 0.1$	$c = 0.001$	$c = 0.01$	$c = 0.1$	$c = 0.001$	$c = 0.01$	$c = 0.1$	$c = 0.001$	$c = 0.01$	$c = 0.1$
L1 loss												
	651	117	23	436	80	13	267	47	9	210	36	6
	641	121	22	429	77	14	267	47	8	209	36	6
	627	140	21	429	77	13	268	47	8	209	37	6
L2 loss												
	2096	641	176	1130	317	108	542	144	33	381	88	21
	2129	697	200	1198	326	97	558	138	42	380	89	23
	2075	622	218	1182	292	81	530	139	32	360	89	21
L3 loss												
$\rho = 0.10$	279	53	9	175	31	7	103	18	3	79	14	2
	271	54	10	171	31	6	106	18	3	79	14	2
	284	53	10	168	33	5	103	18	3	80	14	2
$\rho = 0.05$	187	37	8	118	22	4	70	13	2	54	9	
	197	37	8	121	22	4	71	12	2	55	9	
	184	40	7	121	22	4	71	13	2	55	9	
$\rho = 0.01$	82	18	3	54	9		30	5		22	4	
	85	19	3	52	9		29	5		22	4	
	83	18	3	49	9		30	5		22	4	
L4 loss												
$\gamma = 1.00$	1461	271	51	899	171	30	556	103	18	441	78	13
	1472	292	55	942	162	31	561	99	18	438	78	13
	1460	282	56	883	162	30	554	101	18	433	78	13
$\gamma = 0.50$	1208	203	39	684	132	23	427	78	14	337	59	10
	1179	201	38	690	130	23	434	78	14	335	59	10
	1183	213	42	693	134	24	436	80	14	338	60	10
$\gamma = 0.25$	796	166	32	538	106	18	333	59	10	259	46	8
	859	171	30	540	99	19	331	62	11	260	47	8
	894	167	32	531	101	18	333	60	10	260	46	8

References

- Balakrishnan, N., & Kundu, D. 2019. Birnbaum-Saunders distribution: A review of models, analysis, and applications. Applied Stochastic Models in Business and Industry, **35**(1), 4–49. 2
- Birnbaum, Z. W., & Saunders, S. C. 1969a. Estimation for a family of life distributions with applications to fatigue. Journal of Applied Probability, **6**(2), 328–347. 2
- Birnbaum, Z. W., & Saunders, S. C. 1969b. A new family of life distributions. Journal of Applied Probability, **6**(2), 319–327. 1, 2
- Bourguignon, M., Lee Ho, L., & Fernandes, F. H. 2020. Control charts for monitoring the median parameter of Birnbaum-Saunders distribution. Quality and Reliability Engineering International, **36**(4), 1333–1363. 2
- Costa, E. G. 2017. Tamanho amostral para estimar a concentração de organismos em água de lastro: uma abordagem bayesiana. Ph.D. thesis, Departamento de Estatística, Universidade de São Paulo, São Paulo. In Portuguese. DOI: 10.11606/T.45.2018.tde-05072018-164225. 2, 5, 7
- Costa, Eliado G., & Santos-Neto, Manoel. 2020. samplesizeBS: Bayesian sample size in a decision-theoretic approach for the Birnbaum-Saunders. www.github.com/santosneto/samplesizeBS. R package version 0.1.5. 8
- De Santis, F., & Gubbiotti, S. 2016. A decision-theoretic approach to sample size determination under several priors. Applied Stochastic Models in Business and Industry,. doi: 10.1002/asmb.2211. 2
- Etzioni, R., & Kadane, J. B. 1993. Optimal experimental design for another’s analysis. Journal of the American Statistical Association, **88**(424), 1404–1411. 2, 9
- Hastings, W. K. 1970. Monte Carlo sampling methods using Markov chains and their applications. Biometrika, **57**(1), 97–109. 3
- Islam, A. F. M. S., & Pettit, L. I. 2012. Bayesian Sample Size Determination Using Linex Loss and Linear Cost. Communications in Statistics - Theory and Methods, **41**(2), 223–240. 2
- Islam, A. F. M. S., & Pettit, L. I. 2014. Bayesian sample size determination for the bounded linex loss function. Journal of Statistical Computation and Simulation, **84**(8), 1644–1653. 2, 9

- Leiva, V. 2016. The Birnbaum-Saunders distribution. New York: Academic Press. 2
- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., & Teller, E. 1953. Equation of state calculations by fast computing machines. Journal of Chemical Physics, **21**(6), 1087–1092. 3
- Parmigiani, G., & Inoue, L. 2009. Decision theory: principles and approaches. New York: John Wiley & Sons. 2, 4, 7
- Raiffa, H., & Schlaifer, R. 1961. Applied statistical decision theory. Boston: Harvard University Press. 4
- Rice, K. M., Lumley, T., & Szpiro, A. A. 2008. Trading bias for precision: decision theory for intervals and sets. <http://www.bepress.com/uwbiostat/paper336>. Working Paper 336, UW Biostatistics. 7
- Sahu, S. K., & Smith, T. M. F. 2006. A Bayesian method of sample size determination with practical applications. Journal of the Royal Statistical Society: Series A (Statistics in Society), **169**(2), 235–253. 2
- Wang, M., Sun, X., & Park, C. 2016. Bayesian analysis of Birnbaum-Saunders distribution via the generalized ratio-of-uniforms method. Computational Statistics, **31**(1), 207–225. 3