

ON THE TRANSITION OF THE RAYLEIGH-TAYLOR INSTABILITY IN 2D WATER WAVES

QINGTANG SU

ABSTRACT. In this paper we prove the existence of water waves with sign-changing Taylor sign coefficients, that is, the strong Taylor sign holds initially, while breaks down at a later time, and vice versa. Such a phenomenon can be regarded as the transition between the stable and unstable regime in the sense of Rayleigh-Taylor of water waves. As a byproduct, we prove the sharp wellposedness of 2d water waves in Gevrey-2 spaces.

CONTENTS

1. Introduction	2
1.1. The background and the main result	2
1.2. The difficulty and the strategy	4
1.3. Outline of the paper	8
2. Preliminaries	8
2.1. Governing equation for the free boundary	8
2.2. Function spaces	14
2.3. Commutator estimates in Gevrey spaces	17
3. The main results: quantitative statements	20
3.1. The initial data	20
3.2. The main theorems	21
4. A quasilinear system	22
5. Estimates	27
5.1. A priori assumptions	27
5.2. Estimates	28
6. Proof of Theorem 3	38
6.1. The wellposedness of the quasilinear system (2.29)	38
7. Applications: sign changing of the Taylor sign coefficient	43
7.1. The Taylor sign coefficient	44
7.2. The case $\lambda_1 = -\lambda_2 > 0$ (travel upward)	45
7.3. The case $\lambda_1 = -\lambda_2 < 0$ (Travel downward).	50
7.4. Conclude the proof of Theorem 4	51
Acknowledgement	51
Appendix A. The Taylor sign	51
A.1. The proof of Corollary 7.1	53
References	55

1. INTRODUCTION

1.1. The background and the main result. We consider the motion of an inviscid and incompressible ideal fluid with a free surface in two space dimensions (that is, the interface separating the fluid and the vacuum is one dimensional), such as the surface waves in the ocean. We refer such fluid as *water waves*. We denote the fluid region by $\Omega(t)$, with a free interface $\Sigma(t)$. The equations of motion are Euler's equations, coupled to the motion of the boundary, and with vanishing boundary condition for the pressure. It is assumed that the fluid region is below the air region. Assume that the density of the fluid is 1, the gravitational field is $-\vec{k}$, where $\vec{k}$ is the unit vector pointing in the upward vertical direction. In two dimensions, if the surface tension is zero, then the motion of the fluid is described by

$$\begin{cases} v_t + v \cdot \nabla v = -\nabla P - (0, 1) \\ \operatorname{div} v = 0, \\ P \equiv 0 \\ (1, v) \text{ is tangent to the free surface } (t, \Sigma(t)). \end{cases} \quad \begin{matrix} \Omega(t) \\ \Sigma(t) \end{matrix} \quad (1.1)$$

where v is the fluid velocity, P is the fluid pressure.

This system, and many variants and generalizations, has been extensively studied in the literature. The so-called *Taylor-sign condition* (also referred as *Rayleigh-Taylor sign condition* in many literature) $-\frac{\partial P}{\partial n} > 0$ on the pressure is an important stability condition for the water waves problem. We shall call $-\frac{\partial P}{\partial n}$ the *Taylor-sign coefficient*. If the *Taylor-sign condition* fails, the system is, in general, unstable, see, for example, [7], [9], [45], [21]. We refer such instability as the *Rayleigh-Taylor instability*. In the irrotational case and without a bottom the validity of the *Taylor-sign condition* was shown by Wu [52, 53], and was the key to obtaining the first local-in-time existence results for large data in Sobolev spaces. In the case of non-trivial vorticity or with a bottom it is widely believed that the *Taylor-sign condition* can fail and the sign condition has to be part of the assumptions for the initial data. In the irrotational case, Nalimov [38], Yosihara [57] and Craig [18] proved local well-posedness for 2d water waves equation for small initial data. In S. Wu's breakthrough works [52, 53] she proved local-in-time well-posedness without smallness assumption. Since then, a lot of interesting local well-posedness results were obtained, see for example [5], [6], [14], [15], [30], [33], [34], [39], [42], [58], [1], [2], [37], and the references therein. See also [41, 50, 49, 51] for water waves with non-smooth interface. For the formation of splash singularities, see for example [11], [10], [16], [17]. Regarding the local-in-time wellposedness with regular vorticity, see [29], [39], [40], [14], [14], [34], [58], and [44] for water waves with point vortices. In the irrotational case, almost global and global well-posedness for water waves were proved in [54, 55], [23], [31], [4], and see also [26, 27], [47], [59]. In the rotational case, see [28], [8], [24], and [44].

All the aforementioned results require the *Taylor-sign condition* to hold. Shinbrot [43], Kano, and Nishida [32] justify the Friedrichs expansion for the water-waves equations in terms of the shallowness parameter using the Cauchy-Kowalevski theorem. They take analytic initial data, and the *Taylor-sign condition* is not assumed. Very recently, Alazard, Burq, and Zuily [3] revisit the analysis of the water-problem with analytic data, using tools and methods that they developed previously to study the Cauchy problem with rough initial data. Their result allows a bottom with Sobolev regularity. Yet none of these results proved the existence of solutions with the *Taylor sign*

hold initially while failing at a later time. As we know, the *Taylor sign condition* is in some sense a criterion for the stability of the water waves. It is natural to ask the following question:

Question 1: Is there a solution to the system (1.1) such that the strong *Taylor-sign condition* holds at $t = 0$, while fails at some $t_0 > 0$?

The rigorous mathematical analysis of the transition of the *Rayleigh-Taylor instability* is an interesting yet less-understood subject. From the technical point of view, the break down of the *Taylor sign condition* corresponds to the loss of derivatives of the solution. In [12], Castro, Cordoba, Fefferman, and Gancedo showed that the Rayleigh-Taylor condition for the Muskat problem may hold initially but break down in finite time. In their case, the solution has instant analyticity, that is, even the initial data has finite smoothness, the solution will simultaneously become real analytic after the initial time. Such instant analyticity can compensate for the loss of derivative caused by the break down of the Rayleigh-Taylor condition. For the water waves, there is no such instant analyticity, which makes the problem more difficult. To the best of our knowledge, such a transition of the Rayleigh-Taylor instability/stability was not known for the water waves. Of course, for infinite depth and irrotational water waves, the *Taylor sign condition* always holds, as long as the free interface is smooth and non-self-intersecting. In order that (1.1) admits a solution with the *Taylor sign* breaks down, we must assume that the initial data has nontrivial vorticity.

The interaction of the free surface and the vorticity is in general very complicated. In order to control the motion of the vorticity and therefore make the interaction between the free surface and the vorticity predictable, we assume that the initial vorticity is highly concentrated in a small region. In idealized cases, we can assume that the vorticity is given by the *point vortices*, that is, the vorticity distribution $\omega := \text{curl } v$ is a linear combination of dirac masses, i.e.,

$$\omega(\cdot, t) = \sum_{j=1}^N \lambda_j \delta_{z_j(t)}(\cdot), \quad (1.2)$$

where $\{z_j(t)\} \subset \Omega(t)$, and $\lambda_j \in \mathbb{R}$. Since the vorticity of the 2d Euler is transported following the fluid, if the initial vorticity of (1.1) is a Dirac delta mass, then ω might remain as such a point vortex for all t . The question is, as the singular vorticity $\lambda_j \delta_{z_j(t)}$ generates a singularly rotational part $\frac{\lambda_j}{\pi} \nabla^\perp \log |z - z_j(t)|$ of v (Here, $z = (x, y)$ and $\nabla^\perp = (-\partial_y, \partial_x)$), following what vector field should $z_j(t)$ move? Since the above singular vector field is purely rotational about $z_j(t)$ and does not move that particle $z_j(t)$, it is reasonable to expect that the dynamics of $z_j(t)$ is governed only by the remaining smooth part of v , that is,

$$\dot{z}_j(t) = \left(v(z, t) - \sum_{j=1}^N \frac{\lambda_j}{\pi} \nabla^\perp \log |z - z_j(t)| \right) \Big|_{z=z_j(t)}. \quad (1.3)$$

This well-known result is rigorously established by considering a family of vortex patch solutions whose initial vorticity limiting (weakly) to a Dirac delta mass (see [35] Theorem 4.1, 4.2 for more details).

The system (1.1)-(1.2)-(1.3) is a model for the motion of submerged bodies (see e.g. [13],[20]) and it is believed to give some insight into the problem of turbulence ([35], chap 4, §4.6). There have been many numerical studies on this system (see for example, [19], [25], [22], [36], [46], [48]) and studies on the local in time well-posedness of the Cauchy problem in the irrotational case (that is, no vorticity), and for regular vortex distributions. In [44], the author proved local wellposedness of (1.1) with point vortices under the strong Taylor sign assumption, and obtain extended lifespan with small initial data when there is a pair of counter-rotating point vortices travelling downward.

In the case of point vortices analyzed in this paper, we give an affirmative answer to *Question 1*. The main result of this paper is

Theorem 1. *For any given constants $0 < \eta_0 < 1$ and $\eta_1 > 0$, there exist $T_0 > 0$ and a nonempty open set $\mathcal{O}$ in a category of infinite smoothness such that for each $\mathbf{u}_0 \in \mathcal{O}$, the system (1.1)-(1.2)-(1.3) with initial value $\mathbf{u}_0$ admits a unique smooth solution on $[0, T_0]$. Moreover, the Taylor sign coefficient $-\frac{\partial P}{\partial n}$ of the corresponding solution satisfies:*

- (1) At $t = 0$, $\inf_{\alpha \in \mathbb{R}}(-\frac{\partial P}{\partial n}(\alpha, 0)) \geq 1 - \eta_0$;
- (2) At $t = T_0$, $\inf_{\alpha \in \mathbb{R}}(-\frac{\partial P}{\partial n}(\alpha, T_0)) \leq -\eta_1$.

In another words, the infimum of the Taylor sign coefficient can transit from almost the constant one to an arbitrary large yet negative number.

Remark 1.1. *The category of infinite smoothness in Theorem 1 is indeed the Gevrey-2 space, see Definition 2.4 for the definition of such spaces, and Theorem 4 for the precise quantitative version of Theorem 1.*

Such a result is our first step to investigate systematically the transition of the Rayleigh-Taylor instability (from stable regime to unstable regime, and vice versa) in water waves. Since (1.1)-(1.2)-(1.3) is reversible in time, *Theorem 1* implies that there exists a nonempty open set of initial data in a category of infinite smoothness with a pair of point vortices such that $\inf_{\alpha \in \mathbb{R}}(-\frac{\partial P}{\partial n}(\alpha, 0)) < -\eta_1$, while $\inf_{\alpha \in \mathbb{R}}(-\frac{\partial P}{\partial n}(\alpha, T_0)) > 1 - \eta_0$ at the finite time $0 < T_0 < \infty$.

Remark 1.2. *Our method actually works if the pair of point vortices are replaced by a pair of concentrated vortex patches.*

1.2. The difficulty and the strategy. In Riemann mapping variables (See §2.1.2 for the precise derivations), the water waves system can be written in the form

$$\begin{cases} D_t U = -\Re\{D_t Q\} + A\Lambda W \\ D_t W = -U + \Re\{Q\} - \Re\{[D_t Z, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\} - 2\Re\{(I - \mathbb{H})Q\}, \\ Z - \alpha = (I + \mathbb{H})W, \\ F = (I + \mathbb{H})U. \end{cases} \quad (1.4)$$

where U, W are real-valued functions, $D_t = \partial_t + b\partial_\alpha$, A and b are real-valued functions depending on U, W and Q , and $Q = -\sum_{j=1}^N \frac{\lambda_j i}{2\pi} \frac{1}{Z(\alpha, t) - z_j(t)}$. $\mathbb{H}$ is the standard Hilbert transform. A plays the role of the *Taylor-sign condition*. If $\inf_{\alpha \in \mathbb{R}} A \geq 0$, then the *Taylor-sign condition* holds; otherwise, the *Taylor-sign condition* fails.

1.2.1. A mechanism for solution with A changes sign: the interaction of vorticity and the free interface. Let $A_1 := A|Z_\alpha|^2$. It suffices to construct solutions to the water waves with a sign-changing A_1 . In [44], the author derived a formula for A_1 , which we record as follows.

$$A_1 = 1 + \frac{1}{2\pi} \int \frac{|D_t Z(\alpha, t) - D_t Z(\beta, t)|^2}{(\alpha - \beta)^2} d\beta - \sum_{j=1}^N \frac{\lambda_j}{2\pi} \operatorname{Re} \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) (D_t Z - \dot{z}_j(t)) \right\}. \quad (1.5)$$

Here, $D_t Z$ is the trace of the velocity field along the free boundary. We split A_1 as

$$A_1 = A_1^{pv} + A_1^{int},$$

where A_1^{pv} represents the unperturbed case, i.e., corresponding to $Z(\alpha, t) = \alpha$, $F = 0$; A_1^{int} represents the contribution due to the interaction of the wave F and the point vortices. Our idea is to find a solution to the water wave system such that, for given $\eta_0 \in (0, 1)$ and $\eta_1 > 0$,

- (R1) $\inf_{\alpha \in \mathbb{R}} A_1^{pv}(\alpha, 0) > 1 - \eta_0$,
(R2) $\inf_{\alpha \in \mathbb{R}} A_1^{pv}(\alpha, T_0) < -\eta_1$, for some $T_0 > 0$,
(R3) $\sup_{\alpha \in \mathbb{R}} |A_1^{int}(\alpha, t)| \ll \left| \inf_{\alpha \in \mathbb{R}} A_1^{pv}(\alpha, t) \right|$, for $t = 0, T_0$.

(R1) and (R2) can be guaranteed by analyzing the motion of a pair of counter-rotating point vortices. Indeed, we have the following observation.

The Taylor sign of the 2d water waves with a vortex pair

Idealized case: the motion of the water waves is completely given by the motion of the point vortices. Suppose there exists a smooth solution to the water waves such that at time t , the free interface $\Sigma_t = \mathbb{R}$, and the velocity field is generated by a pair of counter-rotating point vortices only, that is, the initial vorticity $\omega := \lambda \delta_{z_1(t)} - \lambda \delta_{z_2(t)}$, with $z_1(t) = -x(t) + iy(t)$ and $z_2(t) = x(t) + iy(t)$ symmetric about the vertical axis. In this case, $A^{pv} = A_1$. Then we have

- (C1) Fix λ and $x(t)$. If $|y(t)|$ is sufficiently large, then $A_1 = 1 + O(\frac{1}{|y(t)|})$.
(C2) Fix $x(t)$. Let $\frac{\lambda^2}{|y(t)|^3} \rightarrow \infty$ and require $|y(t)| \gg 1$, then $A_1 \rightarrow -\infty$.

(C1) and (C2) will be justified from the calculation in §7 and the appendix.

Another observation: If $\lambda > 0$, then the vortex pair $z_1(t), z_2(t)$ travel toward the free interface (so $y(t)$ becomes smaller).

A natural idea: Construct water waves with a symmetric pair of point vortices far away from the free interface initially and traveling toward the free interface rapidly. Then we expect A_1 to satisfy (R1)-(R2)-(R3).

To guarantee (R3), the main difficulty is to justify that the water waves really remain a small perturbation of the motion of the vortex pair (for a period of time that allows the vortex pair to travel sufficiently close to the free interface such that the Taylor-sign fails), which can be guaranteed if we can control the growth of U and W . This forces us to prove the local existence in some reasonable spaces of such water waves. Nevertheless, even in the irrotational case, all the existing local wellposedness results for the water wave system with finite smoothness require the strong *Taylor-sign condition* to hold. As we mentioned earlier, if the *Taylor-sign condition* fails, then the water waves can be subject to the *Rayleigh-Taylor instability* and a local wellposedness is not expected, at least in spaces with finite smoothness.

Roughly speaking, if the *Taylor-sign condition* fails, the system (1.4) losses $\frac{1}{2}$ derivatives. In Sobolev spaces H^s , such a loss of derivative can be overcome if the strong *Taylor-sign condition* holds. If the *Taylor-sign condition* fails, then the Fourier mode can grow exponentially, which prevents a local wellposedness in Sobolev spaces.

Instead of seeking for water waves with finite regularity (e.g., Sobolev spaces), we perturb the vortex pair in a category with infinite smoothness. Such an idea is certainly not new. Even in the context of water waves, by expanding the solution as a power series in time, Shinbrot [43], Kano and Nishida's [32] obtained local existence in time for the general initial value problem with real analytic data. Their tool is the Cauchy-Kowalevski theorem. In [3], Alazard, Burq, and Zuily proved the local wellposedness of water waves in real analytic spaces via the energy method. Their main analytical tool is the paradifferential calculus. One should be able to adapt their method to prove the local in time wellposedness of water waves with point vortices. In this paper, working in Riemann mapping variables, we prove the local wellposedness in *Gevrey-2 space* (The *Gevrey-2 space* is a space of infinite smoothness that extends the real-analytic space. See *Definition 2.4* for the precise definition) by using the energy method directly. The key to prove *Theorem 1* is the following.

Theorem 2. *The water wave system (1.4) is locally wellposed in Gevrey-2 spaces.*

Such a local wellposedness is expected to be sharp, in the sense that if the initial data is in a space rougher than the Gevrey-2 space, then the system (1.4) is illposed. See *Remark 3.2* for more quantitative discussions.

To illustrate our strategy, we consider the following toy model:

$$\begin{cases} \partial_t U = A\Lambda W, \\ \partial_t W = -U, \\ (U, W) = (U_0, W_0), \end{cases} \quad (1.6)$$

for some real-valued function A that might depend on u and W .

1.2.2. The Gevrey framework. Assume that $A(\alpha_0, 0) < 0$ for some given $\alpha_0 \in \mathbb{R}$. Heuristically, by considering solutions U and W concentrating near α_0 , we can then without loss of generality (although it is not trivial to rigorously justify this) assume that $A(\alpha, t) \equiv A(\alpha_0, 0) < 0$. In Fourier variables, we have

$$\partial_t \begin{bmatrix} \hat{U} \\ \hat{W} \end{bmatrix} = \begin{bmatrix} 0 & A|\xi| \\ -1 & 0 \end{bmatrix} \begin{bmatrix} \hat{U} \\ \hat{W} \end{bmatrix}. \quad (1.7)$$

The matrix $\begin{bmatrix} 0 & A|\xi| \\ -1 & 0 \end{bmatrix}$ has real eigenvalues $\lambda_1 = \sqrt{-A|\xi|}$ and $\lambda_2 = -\sqrt{-A|\xi|}$, with eigenvectors $V_1 = \begin{bmatrix} -\sqrt{-A|\xi|} \\ 1 \end{bmatrix}$ and $V_2 = \begin{bmatrix} \sqrt{-A|\xi|} \\ 1 \end{bmatrix}$, respectively. Let $S = [V_1, V_2]$, and let $\mathcal{Y} = S^{-1} \begin{bmatrix} \hat{U} \\ \hat{W} \end{bmatrix}$, then

$$\partial_t \mathcal{Y} = \begin{bmatrix} \sqrt{-A|\xi|} & 0 \\ 0 & -\sqrt{-A|\xi|} \end{bmatrix} \mathcal{Y}. \quad (1.8)$$

So

$$\mathcal{Y}(\xi, t) = \begin{bmatrix} e^{t\sqrt{-A|\xi|}} & 0 \\ 0 & e^{-t\sqrt{-A|\xi|}} \end{bmatrix} \mathcal{Y}(\xi, 0),$$

which implies that for any $t > 0$, we have ¹

$$\int_{\mathbb{R}} e^{\delta|\xi|} |\mathcal{Y}(\xi, t)|^2 d\xi = \infty \quad \Leftrightarrow \quad \sum_{n=0}^{\infty} \frac{\delta^{2n}}{(n!)^4} \int_{\mathbb{R}} |\xi|^{2n} |\mathcal{Y}(\xi, t)|^2 d\xi = \infty, \quad (1.9)$$

provided that $\mathcal{Y}(\xi, 0) \geq C e^{\delta|\xi|^{1/2}}$ for some $\delta > -\sqrt{-A}$. Therefore, it is natural to impose the assumption $|\mathcal{Y}(\xi, 0)| \leq C e^{-\sqrt{-A|\xi|}}$. Such $\mathcal{Y}(\cdot, 0)$ lies in the Gevrey-2 spaces. See *Definition 2.4* for the more precise definition. We denote

$$\|f\|_{X_{L_0}}^2 := \sum_{n=0}^{\infty} \frac{L_0^{2n}}{(n!)^4} \int_{\mathbb{R}} |\xi|^{2n} |\hat{f}(\xi)|^2 d\xi \quad (= \sum_{n=0}^{\infty} \frac{L_0^{2n}}{(n!)^4} \|\partial_{\alpha}^n f\|_{L^2}^2).$$

L_0 is called the *radius of analyticity*. We seek for solution (U, W) such that the initial data (U_0, W_0) satisfies

$$\|\Lambda^{1/2} U_0\|_{X_{L_0}} < \infty, \quad \|\partial_{\alpha} W_0\|_{X_{L_0}} < \infty.$$

Of course, in general, A is not a constant. In particular, in the context of water waves, A depends on U and W . Moreover, in the first equation of (1.4), W losses one half derivative, due to the term

¹The reason that we use the infinite series to define the Gevrey norm is as follows: we need to estimate the Gevrey norm of functions of the form $Q_j := \frac{1}{Z(\alpha, t) - z_j(t)}$. It is easier to estimate $\sum \frac{\sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n Q_j\|_{L^2}^2$ than $\int_{\mathbb{R}} e^{2\sigma|\partial_{\alpha}|^{1/2}} |Q(\alpha, t)|^2 d\alpha$.

$A\Lambda W$. To compensate for this loss, we allow the *radius of analyticity* of the solution to decay linearly in time, that is, we seek for solution (U, W) to (1.6) with initial data (U_0, W_0) such that

$$\|\Lambda^{1/2}U(\cdot, t)\|_{X_{L_0-\delta t}} < \infty, \quad \|\partial_\alpha W\|_{X_{L_0-\delta t}} < \infty. \quad (1.10)$$

Such a linear decay in the radius of analyticity has the same effect of introducing a viscosity term $\approx \Lambda^{1/2} \begin{bmatrix} U \\ W \end{bmatrix}$, that is, in the $X_{L_0-\delta t}$ norm, the toy model behaves like

$$\begin{cases} \partial_t U + c\Lambda^{1/2}U = A\Lambda W, \\ \partial_t W + c\Lambda^{1/2}W = -U, \end{cases} \quad (1.11)$$

for some positive constant $c > 0$. Clearly, (1.11) does not loss any derivative. A more apparent way to explain the compensation of the loss of derivative is by energy estimates. We obtain

$$\begin{aligned} & \frac{d}{dt} (\|\Lambda^{1/2}U\|_{X_{L_0-\delta t}}^2 + \|\partial_\alpha W\|_{X_{L_0-\delta t}}^2) \\ & \leq C (\|\Lambda^{1/2}U\|_{X_{L_0-\delta t}}^2 + \|\partial_\alpha W\|_{X_{L_0-\delta t}}^2) \\ & \quad - \left(\frac{2\delta}{L_0 - \delta t} \sum_{n=1}^{\infty} \frac{n(L_0 - \delta t)^{2n}}{(n!)^4} \|\partial_\alpha^n \Lambda^{1/2}U(\cdot, t)\|_{L^2}^2 + \frac{2\delta}{L_0 - \delta t} \sum_{n=1}^{\infty} \frac{n(L_0 - \delta t)^{2n}}{(n!)^4} \|\partial_\alpha^n \Lambda^{1/2}U(\cdot, t)\|_{L^2}^2 \right) \\ & \quad + \sum_{n=0}^{\infty} \frac{(L_0 - \delta t)^{2n}}{(n!)^4} \left| \int_{\mathbb{R}} \partial_\alpha^n \partial_\alpha U(\alpha, t) \partial_\alpha^n W_\alpha(\alpha, t) d\alpha \right| \end{aligned}$$

By choosing δ sufficiently large, one can bound the last summation by

$$\left(\frac{2\delta}{L_0 - \delta t} \sum_{n=1}^{\infty} \frac{n(L_0 - \delta t)^{2n}}{(n!)^4} \|\partial_\alpha^n \Lambda^{1/2}U(\cdot, t)\|_{L^2}^2 + \frac{2\delta}{L_0 - \delta t} \sum_{n=1}^{\infty} \frac{n(L_0 - \delta t)^{2n}}{(n!)^4} \|\partial_\alpha^n \Lambda^{1/2}U(\cdot, t)\|_{L^2}^2 \right).$$

So we can obtain closed energy estimates.

Using this idea, we are able to prove the local wellposedness of (1.4) in Gevrey-2 spaces, regardless of the sign of A .

The lifespan of the perturbation. As we mentioned earlier, it is critical that the water waves remain a small perturbation of the vortex pair for sufficiently long time².

- (I1) By (C1), in order that A_1 is sufficiently close to 1 at $t = 0$, we need $|\lambda|^2/|y(0)|^3 \ll 1$. So $|y(0)| \gg |\lambda|^{2/3}$.
- (I2) By (C2), in order that $A_1(T_0)$ takes a large negative number, we need $\frac{\lambda^2}{|y(t)|^3}$ to be sufficiently large, which suggests that $|y(T_0)| \ll |y(0)|$. For simplicity, we take $y(T_0) \approx -|y(0)|^{9/10}$.
- (I3) In a perturbative regime, the velocity of the point vortex pair is $\frac{\lambda}{4\pi x(0)}$, plus a negligible error term (the details will be provided in §7). So we have

$$T_0 = \frac{|y(0) - y(T_0)|}{\frac{\lambda}{4\pi x(0)}}. \quad (1.12)$$

In particular, $T_0 \gg \frac{4\pi x(0)}{|y_0|^{1/2}}$.

It is not difficult to prove the local well-posedness of (1.4), yet to obtain a lifespan with lower bound at least $\frac{4\pi x(0)}{|y_0|^{1/2}}$ is another story, because if $\frac{\lambda^2}{|y(t)|^3}$ and $y(0)$ are large, then the quantity $-\Re\{D_t Q\}$ on the right hand side of the first equation of (1.4) is also large, of size roughly speaking

²By *sufficiently long time*, we do not mean a long time existence for water waves. It merely means that the lifespan is sufficient for the *Taylor sign coefficient* to transit from a positive number to an arbitrary large negative number.

$\frac{\lambda^2}{|y(t)|^2}$. Thus we need to analyze the interaction of the free interface and the point vortices to exclude the possibility of rapid growth of the water waves, from which we can conclude the proof of our main theorem.

1.3. Outline of the paper. In §2, we formulate the water wave system in Riemann variables, define the function spaces, and derive some estimates in Gevrey spaces. In §3, we state a quantitative version of the main results. In §4, using the energy method, we prove the local wellposedness in Gevrey spaces of a specific quasilinear system. In §5, we derive a priori estimates. In §6, we use the picard iteration method to prove the existence and uniqueness of solutions of the system (2.29) and therefore conclude the proof of Theorem 3. In §7, using Theorem 3, we prove Theorem 4. In the appendix, we provide the details of some calculations for the Taylor sign coefficient.

2. PRELIMINARIES

Throughout this paper, we denote the fluid region at time t by $\Omega(t)$, with a nonself-intersect free surface $\Sigma(t)$. We identify a point $(x, y) \in \mathbb{R}$ as a point $x+iy \in \mathbb{C}$. For a point $x+iy \in \Sigma(t)$, we assume $|y| \rightarrow 0$ as $|x| \rightarrow \infty$. That is, $\Sigma(t)$ approaches the real axis at $\pm\infty$. Define $\langle u, v \rangle := \int_{\mathbb{R}} u(\alpha) \bar{v}(\alpha)$. Let $z = x + iy \in \mathbb{C}$, $\Re\{z\} := x$, $\Im\{z\} := y$.

2.1. Governing equation for the free boundary. The system (1.1)-(1.2)-(1.3) is completely determined by the free surface $\Sigma(t)$, the trace of the velocity v along the free surface, and the position of the point vortices. We shall first formulate the system (1.1)-(1.2)-(1.3) in Lagrangian coordinates. Because of the moving boundary, it is convenient to use the Riemann mapping variables to study the Taylor sign and construct special solutions.

2.1.1. Lagrangian formulation. We parametrize the free surface by Lagrangian coordinates, i.e., let α be such that

$$z_t(\alpha, t) = v(z(\alpha, t), t). \quad (2.1)$$

$P|_{\Sigma(t)} \equiv 0$ implies that $\nabla P|_{\Sigma(t)}$ is along the normal direction, so we can write ∇P as $-iaz_\alpha$, where $a = -\frac{\partial P}{\partial \bar{n}} \frac{1}{|z_\alpha|}$ is real valued. Here, $\bar{n} = i \frac{z_\alpha}{|z_\alpha|}$ is the unit outward normal to $\Omega(t)$. So the trace of the momentum equation $v_t + v \cdot \nabla v = -\nabla P - (0, 1)$ can be written as

$$z_{tt} - iaz_\alpha = -i. \quad (2.2)$$

Definition 2.1 (Hilbert transform). Assume that $z(\alpha)$ satisfies

$$\beta_0 |\alpha - \beta| \leq |z(\alpha) - z(\beta)| \leq \beta_1 |\alpha - \beta|, \quad \forall \alpha, \beta \in \mathbb{R}, \quad (2.3)$$

where $0 < \beta_0 < \beta_1 < \infty$ are constants. The Hilbert transform associates to the curve $z(\alpha)$ is defined as

$$\mathfrak{H}f(\alpha) := \frac{1}{\pi i} p.v. \int_{-\infty}^{\infty} \frac{z_\beta(\beta)}{z(\alpha) - z(\beta)} f(\beta) d\beta. \quad (2.4)$$

The standard Hilbert transform is the Hilbert transform associated to $z(\alpha) = \alpha$, that is,

$$\mathbb{H}f(\alpha) := \frac{1}{\pi i} p.v. \int_{-\infty}^{\infty} \frac{1}{\alpha - \beta} f(\beta) d\beta. \quad (2.5)$$

We can use the Hilbert transform to characterize the boundary value of holomorphic functions. Such characterization is classical, the reader can see for example Proposition 2.1 in [56].

Lemma 2.1. Let $\Omega \subset \mathbb{C}$ be a domain with C^1 boundary $\Sigma : z = z(\alpha), \alpha \in \mathbb{R}$, oriented clockwise. Let $\mathfrak{H}$ be the Hilbert transform associated to Ω .

(a.) Let $g \in L^p$ for some $1 < p < \infty$. Then g is the boundary value of a holomorphic function G on Ω with $G(z) \rightarrow 0$ at infinity if and only if

$$(I - \mathfrak{H})g = 0. \quad (2.6)$$

(b.) Let $f \in L^p$ for some $1 < p < \infty$. Then $\frac{1}{2}(I + \mathfrak{H})f$ is the boundary value of a holomorphic function $\mathfrak{G}$ on Ω , with $\mathfrak{G}(z) \rightarrow 0$ as $|z| \rightarrow \infty$.

(c.) $\mathfrak{H}1 = 0$.

We decompose $\bar{z}_t$ as $\bar{z}_t = f + p$, where $p = -\sum_{j=1}^n \frac{\lambda_j i}{2\pi} \frac{1}{z(\alpha, t) - z_j(t)}$. Note that f is holomorphic in $\Omega(t)$ with the value at the boundary $\Sigma(t)$ given by $\bar{z}_t + \sum_{j=1}^N \frac{\lambda_j i}{2\pi(z(\alpha, t) - z_j(t))}$. Let $f \in L^2(\mathbb{R})$, then f is the boundary value of a holomorphic function in $\Omega(t)$ if and only if

$$(I - \mathfrak{H})f = 0, \quad (2.7)$$

where $\mathfrak{H}$ is the Hilbert transform associated with the curve $z(\alpha, t)$, i.e.,

$$\mathfrak{H}f(\alpha) := \frac{1}{\pi i} p.v. \int_{-\infty}^{\infty} \frac{z\beta}{z(\alpha, t) - z(\beta, t)} f(\beta) d\beta. \quad (2.8)$$

Because of the singularity of the velocity at the point vortices, we don't have $(I - \mathfrak{H})\bar{z}_t = 0$. However, the following lemma asserts that $\bar{z}_t$ is almost holomorphic, in the sense that $(I - \mathfrak{H})\bar{z}_t$ consists of lower order terms.

Lemma 2.2 (Almost holomorphicity). *Assume that $z(\cdot, t) \in L^2(\mathbb{R})$ satisfies (2.3) and $\bar{z}_t$ is the boundary value of a velocity field $\bar{v}$ in $\Omega(t)$ such that $\bar{v} + \sum_{j=1}^N \frac{\lambda_j i}{2\pi(z - z_j(t))}$ is holomorphic in $\Omega(t)$. Then we have*

$$(I - \mathfrak{H})\bar{z}_t = -\frac{i}{\pi} \sum_{j=1}^N \frac{\lambda_j}{z(\alpha, t) - z_j(t)}. \quad (2.9)$$

Proof. Since $\bar{z}_t + \sum_{j=1}^N \frac{\lambda_j i}{2\pi(z(\alpha, t) - z_j(t))}$ is the boundary value of a holomorphic function in $\Omega(t)$, by lemma 2.1,

$$(I - \mathfrak{H})(\bar{z}_t + \sum_{j=1}^N \frac{\lambda_j i}{2\pi(z(\alpha, t) - z_j(t))}) = 0,$$

we have

$$(I - \mathfrak{H})\bar{z}_t = -\sum_{j=1}^N (I - \mathfrak{H}) \frac{\lambda_j i}{2\pi(z(\alpha, t) - z_j(t))}. \quad (2.10)$$

Since $\frac{1}{z(\alpha, t) - z_j(t)}$ is boundary value of the holomorphic function $\frac{1}{z - z_j(t)}$ in $\Omega(t)^c$, by lemma 2.1 again, we have

$$(I - \mathfrak{H}) \frac{1}{z(\alpha, t) - z_j(t)} = \frac{2}{z(\alpha, t) - z_j(t)}. \quad (2.11)$$

(2.10) together with (2.11) complete the proof of the lemma. $\square$

So the system (1.1) is reduced to a system of equations for the free boundary coupled with the dynamic equation for the motion of the point vortices:

$$\begin{cases} z_{tt} - iaz_\alpha = -i \\ \frac{d}{dt}z_j(t) = (v - \frac{\lambda_j i}{2\pi(z - z_j)}) \Big|_{z=z_j} \\ (I - \mathfrak{H})f = 0. \end{cases} \quad (2.12)$$

Note that v can be recovered from (2.12). Indeed, we have

$$\bar{v}(z, t) + \sum_{j=1}^N \frac{\lambda_j i}{2\pi(z - z_j(t))} = \frac{1}{2\pi i} \int \frac{z_\beta}{z - z(\beta)} \left(\bar{z}_t(\beta, t) + \sum_{j=1}^N \frac{\lambda_j i}{2\pi(z(\beta, t) - z_j(t))} \right) d\beta. \quad (2.13)$$

So the system (1.1) and the system (2.12) are equivalent.

The quantity $a|z_\alpha|$ plays an important role in the study of water waves³.

Definition 2.2. (*The Taylor-sign condition and the strong Taylor sign condition*)

- (1) *If $a|z_\alpha| \geq 0$ pointwisely, then we say the Taylor-sign condition holds.*
- (2) *If there is some positive constant c_0 such that $a|z_\alpha| \geq c_0 > 0$ pointwisely, then we say the strong Taylor sign condition holds.*

In order to derive a useful formula for the *Taylor sign coefficient* and use the iteration method to construct solutions to (2.12), we use the Riemann mapping formulation.

2.1.2. The Riemann mapping formulation. Let $\mathbb{P}_- = \{z \in \mathbb{C} : \Im\{z\} < 0\}$. Let $\Phi(\cdot, t) : \Omega(t) \rightarrow \mathbb{P}_-$ be the Riemann mapping such that $\Phi_z \rightarrow 1$ as $z \rightarrow \infty$. Denote

$$\begin{cases} h(\alpha, t) := \Phi(z(\alpha, t), t), \\ Z(\alpha, t) := z \circ h^{-1}(\alpha, t), \\ b = h_t \circ h^{-1}, \\ D_t := \partial_t + b\partial_\alpha, \\ A := (ah_\alpha) \circ h^{-1}, \\ F := f \circ h^{-1}. \end{cases} \quad (2.14)$$

In Riemann mapping variables, the system (2.12) becomes

$$\begin{cases} (D_t^2 - iA\partial_\alpha)Z = -i \\ \frac{d}{dt}z_j(t) = \left(v - \frac{\lambda_j i}{2\pi(\bar{z} - \bar{z}_j)}\right)\Big|_{z=z_j} \\ (I - \mathbb{H})F = 0. \end{cases} \quad (2.15)$$

Denote

$$A_1 := A|Z_\alpha|^2. \quad (2.16)$$

Since $(a|z_\alpha|) \circ h^{-1} = A|Z_\alpha| = \frac{A_1}{|Z_\alpha|}$, it's clear that the *Taylor-sign condition* holds if and only if

$$\inf_{\alpha \in \mathbb{R}} \frac{A_1}{|Z_\alpha|} \geq 0, \quad (2.17)$$

and the *strong Taylor sign condition* holds if and only if

$$\inf_{\alpha \in \mathbb{R}} \frac{A_1}{|Z_\alpha|} > 0. \quad (2.18)$$

Definition 2.3. We call A_1 the Taylor sign coefficient corresponding to the solution $(Z, F, \{z_j\})$ of the water waves.

Remark 2.1. Note that $A_1 = -|Z_\alpha| \frac{\partial P}{\partial \bar{n}}$.

³Indeed, $a|z_\alpha| = -\frac{\partial P}{\partial \bar{n}}\Big|_{\Sigma(t)}$.

Formula for b : Recall that $h(\alpha, t) = \Phi(z(\alpha, t), t)$, where Φ is the Riemann mapping. So we have

$$h_t = \Phi_t + \Phi_z z_t, \quad \Phi_z = \frac{h_\alpha}{z_\alpha}. \quad (2.19)$$

Precomposite with h^{-1} on both sides of the above,

$$\begin{aligned} b &= h_t \circ h^{-1} = \Phi_t \circ Z + \frac{D_t Z}{Z_\alpha} \\ &= \Phi_t \circ Z + D_t Z \left(\frac{1}{Z_\alpha} - 1 \right) + D_t Z \\ &= \Phi_t \circ Z + D_t Z \left(\frac{1}{Z_\alpha} - 1 \right) + \bar{Q} + \bar{F}. \end{aligned}$$

Applying $I - \mathbb{H}$, using $(I - \mathbb{H})\Phi_t \circ Z = 0$ and $(I - \mathbb{H})\left(\frac{1}{Z_\alpha} - 1\right) = 0$, $(I - \mathbb{H})\bar{F} = 2F$, then taking real part, we obtain

$$b = \Re\{[D_t Z, \mathbb{H}]\left(\frac{1}{Z_\alpha} - 1\right)\} + 2\Re\{D_t Z\} = \Re\{[D_t Z, \mathbb{H}]\left(\frac{1}{Z_\alpha} - 1\right)\} + 2\Re\{(I - \mathbb{H})\bar{Q}\} + 2\Re\{F\}. \quad (2.20)$$

Decomposing b as $b = b_0 + b_1$, where

$$b_0 = 2\Re\{F\} + \Re\{[\bar{F}, \mathbb{H}]\left(\frac{1}{Z_\alpha} - 1\right)\}, \quad (2.21)$$

and

$$b_1 = \Re\{[\bar{Q}, \mathbb{H}]\left(\frac{1}{Z_\alpha} - 1\right)\} + 2\Re\{(I - \mathbb{H})\bar{Q}\}. \quad (2.22)$$

Note that b_1 is more regular than b_0 .

Formula for A_1 : In §4 of [44], the author derived a formula for A_1 , which we record as follows:

$$A_1 = 1 + \frac{1}{2\pi} \int \frac{|D_t Z(\alpha, t) - D_t Z(\beta, t)|^2}{(\alpha - \beta)^2} d\beta - \sum_{j=1}^N \frac{\lambda_j}{2\pi} \operatorname{Re} \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) (D_t Z - \dot{z}_j(t)) \right\}. \quad (2.23)$$

Let $u = D_t \bar{Z} = F + Q$, recall that $F(\alpha, t) = \mathcal{U}(Z(\alpha, t), t)$ for some holomorphic function $\mathcal{U}$ in $\mathbb{P}_-$, and Q is given by

$$Q = - \sum_{j=1}^N \frac{\lambda_j i}{2\pi} \frac{1}{Z(\alpha, t) - z_j(t)}, \quad (2.24)$$

We have

$$\mathcal{U}(z, t) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{Z_\beta}{z - Z(\beta, t)} F(\beta, t) d\beta. \quad (2.25)$$

So we obtain a system in Riemann variables:

$$\begin{cases} D_t F = -D_t Q - iA\bar{Z}_\alpha + i, \\ D_t(\bar{Z} - \alpha) = F + Q - b, \\ F, Z - \alpha \text{ holomorphic.} \end{cases} \quad (2.26)$$

(2.26) is equivalent to (2.12). Let $U = \Re\{F\}$, $W = \Re\{Z - \alpha\}$. Since F and $Z - \alpha$ are holomorphic, U and W determine F and $Z - \alpha$, respectively. Moreover, we have

$$-iA\bar{Z}_\alpha + i = A\Lambda(\bar{Z} - \alpha) - i(A - 1), \quad (2.27)$$

where $\Lambda = |\partial_\alpha|$. Taking real parts on both sides of the first and the second equation of (2.26), we obtain

$$\begin{cases} D_t U = A\Lambda W - \Re\{D_t Q\} \\ D_t W = U + \Re\{Q\} - b, \\ Z - \alpha = (I - \mathbb{H})W, \\ F = (I - \mathbb{H})U. \end{cases} \quad (2.28)$$

Using (2.20), and decompose $b = b_0 + b_1$ for b_0, b_1 given by (2.21) and (2.22), respectively, we obtain⁴

$$\begin{cases} D_t U = -\Re\{D_t Q\} + A\Lambda W \\ D_t W = -U + \Re\{Q\} - \Re\{[D_t \bar{F}, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\} - b_1, \\ Z - \alpha = (I + \mathbb{H})W, \\ F = (I + \mathbb{H})U. \end{cases} \quad (2.29)$$

The motion of $z_j(t)$ is given by

$$\frac{d}{dt} z_j(t) = \overline{\mathcal{U}(z_j(t), t)} + \sum_{\substack{1 \leq k \leq N \\ k \neq j}} \frac{\lambda_k i}{2\pi} \frac{1}{z_j(t) - z_k(t)}, \quad (2.30)$$

where $\mathcal{U}$ is given by (2.25). Now we obtain a system (2.29)-(2.24)-(2.30), which is equivalent to (1.1). Let's denote

$$G := -\Re\{D_t Q\}, \quad (2.31)$$

$$R := \Re\{Q\} - b_1 = \Re\{Q\} - 2\Re\{(I - \mathbb{H})\bar{Q}\} - \Re\{[\bar{Q}, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\}. \quad (2.32)$$

(2.29) then becomes

$$\begin{cases} D_t U = A\Lambda W + G \\ D_t W = -U - \Re\{[\bar{F}, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\} + R, \\ Z - \alpha = (I - \mathbb{H})W, \\ F = (I - \mathbb{H})U. \end{cases} \quad (2.33)$$

Note that G is determined by Z, F and $\{z_j\}$ (and therefore determined by W, U , and $\{z_j\}$), so we can write G as $G(W, U, \{z_j(t)\})$. Here, $\{z_j\}$ means $\{z_1(t), \dots, z_N(t)\}$. For the same reason, $D_t F, b_1, A, R$ are determined by W, U and $\{z_j\}$. Therefore, we write

$$\begin{aligned} G &= G(W, U, \{z_j(t)\}) \\ R &= R(W, U, \{z_j(t)\}) \\ b_1 &= b_1(W, U, \{z_j(t)\}) \\ A &= A(W, U, \{z_j(t)\}) \end{aligned}$$

⁴We need to estimate $\|W_\alpha\|_{X_{\phi(t)}}$ (see *Definition 2.4* for the definition of the Gevrey-2 norm $\|\cdot\|_{X_\sigma}$), so we need to estimate $\|b_\alpha\|_{X_{\phi(t)}}$. We can prove that $\|\partial_\alpha b_1\|_{X_{\phi(t)}} < \infty$. However, $\|\partial_\alpha b_0\|_{X_{\phi(t)}}$ is not necessarily finite. So we need to treat $\partial_\alpha b_0$ carefully.

2.1.3. *Some notations.* For the reader's convenience, we list some notations as follows

$\Sigma(t)$	The free surface at time t ,
$\Omega(t)$	The fluid region at time t , $\Omega(t)$ is bounded above by $\Sigma(t)$,
$\Phi(\cdot, t)$	The Riemann mapping from $\Omega(t)$ to $\mathbb{P}_-$,
$Z(\alpha, t)$	The free interface in Riemann variables,
$h(\alpha, t)$	The trace of $\Phi(\cdot, t)$, that is, $h(\alpha, t) = \Phi(Z(\alpha, t), t)$,
$b(\alpha, t)$	$b(\cdot, t) = h_t \circ h^{-1}$,
D_t	$D_t = \partial_t + b\partial_\alpha$,
$D_t Z$	The trace of the velocity field,
$A Z_\alpha $	The Taylor sign coefficient in Riemann mapping variable,
$z_j(t)$	The coordinates of the j -th point vortex, $z_j(t) = x_j(t) + iy_j(t)$,
$\dot{z}_j$	The time derivative of the point vortex,
$d_I(t)$	The distance between the point vortices and the free interface at time t , that is, $d_I(t) := \inf_{\alpha \in \mathbb{R}} \min_{1 \leq j \leq N} Z(\alpha, t) - z_j(t) $,
$Q(\alpha, t)$	The conjugate of the velocity field generated by the point vortices, $Q(\alpha, t) = - \sum_{j=1}^N \frac{\lambda_j i}{2\pi} \frac{1}{Z(\alpha, t) - z_j(t)}$
$F(\alpha, t)$	The wave part of the conjugate of the velocity field, that is, $F = D_t \bar{Z} - Q$,
$\mathcal{U}(\cdot, t)$	The holomorphic extension of $F(\cdot, t)$, that is, $F(\alpha, t) = \mathcal{U}(Z(\alpha, t), t)$,
$A_1(\alpha, t)$	$A_1(\alpha, t) = A Z_\alpha ^2$,
U	The real part of the velocity field, that is, $U = \Re\{D_t Z\}$,
$W(\alpha, t)$	The real part of $Z(\alpha, t) - \alpha : W = \Re\{Z - \alpha\}$.

We parametrize Σ_0 by Riemann mapping, that is, $h_0(\alpha) = \alpha$, so $z_0(\alpha) = Z_0(\alpha)$. Let $U_0, W_0, z_{j,0}$, Q_0, Q_1 denote the initial value of $U, W, z_j, Q, D_t Q$, respectively.

Using (2.24), we obtain

$$D_t Q = \sum_{j=1}^N \frac{\lambda_j i}{2\pi} \frac{D_t Z - \dot{z}_j(t)}{(Z(\alpha, t) - z_j(t))^2}.$$

Using (2.30), we have

$$\dot{z}_{j,0} = \sum_{\substack{1 \leq k \leq N \\ k \neq j}} \frac{\lambda_j i}{2\pi} \frac{1}{z_{k,0} - z_{j,0}} + \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\partial_\beta Z_0(\beta)}{z_{j,0} - Z_0(\beta)} F_0(\beta) d\beta. \quad (2.34)$$

$$Q_1 = \sum_{j=1}^N \frac{\lambda_j i}{2\pi} \frac{\bar{u}_0 - \dot{z}_{j,0}}{(Z_0(\alpha) - z_{j,0})^2}. \quad (2.35)$$

$$A_{1,0} = 1 + \frac{1}{2\pi} \int \frac{|\bar{u}_0(\alpha) - \bar{u}_0(\beta)|^2}{(\alpha - \beta)^2} d\beta - \sum_{j=1}^N \frac{\lambda_j}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{\partial_\alpha Z_0(\alpha)}{(Z_0(\alpha) - z_j(t))^2} \right) (\bar{u}_0(\alpha) - \dot{z}_{j,0}) \right\}. \quad (2.36)$$

The initial interface Z_0 satisfies

$$|Z_0(\alpha) - Z_0(\beta)| \geq C_0|\alpha - \beta|, \quad \forall \alpha, \beta \in \mathbb{R}. \quad (2.37)$$

2.2. Function spaces. We define the Gevrey-2 spaces as follows.

Definition 2.4. Let $\sigma > 0$, we define

$$\dot{X}_\sigma = \{f \in C^\infty(\mathbb{R}) \mid \|f\|_{\dot{X}_\sigma}^2 := \sum_{j=1}^{\infty} \frac{\sigma^{2j}}{(j!)^4} \|\partial_\alpha^j f\|_{L^2}^2 < \infty\}. \quad (2.38)$$

$$X_\sigma = \{f \in C^\infty(\mathbb{R}) \mid \|f\|_{X_\sigma}^2 := \sum_{j=0}^{\infty} \frac{\sigma^{2j}}{(j!)^4} \|\partial_\alpha^j f\|_{L^2}^2 < \infty\}. \quad (2.39)$$

$$\dot{Y}_\sigma = \{f \in C^\infty(\mathbb{R}) \mid \|f\|_{\dot{Y}_\sigma}^2 := \sum_{j=1}^{\infty} \frac{j^2 \sigma^{2j}}{(j!)^4} \|\partial_\alpha^j f\|_{L^2}^2 < \infty\}. \quad (2.40)$$

$$Y_\sigma = \{f \in C^\infty(\mathbb{R}) \mid \|f\|_{Y_\sigma}^2 := \|f\|_{L^2}^2 + \sum_{j=0}^{\infty} \frac{j^2 \sigma^{2j}}{(j!)^4} \|\partial_\alpha^j f\|_{L^2}^2 < \infty\}. \quad (2.41)$$

It is not difficult to verify that $\dot{X}_\sigma, X_\sigma, \dot{Y}_\sigma$ and Y_σ are Banach spaces (actually, Hilbert spaces.), we shall call these spaces the *Gevrey spaces*, or *Gevrey-2 spaces*, and call σ the *radius of convergence*. Moreover, we have the Sobolev type embedding.

Lemma 2.3. 1. Let $f \in X_\sigma$ and $n \geq 0$ be an integer, then there is an absolute constant $C > 0$ such that

$$\|\partial_\alpha^n f\|_{L^\infty} \leq C \left(\frac{((n+1)!)^2}{\sigma^{n+1}} + \frac{(n!)^2}{\sigma^n} \right) \|f\|_{X_\sigma}. \quad (2.42)$$

2. Let $f \in Y_\sigma$ and $n \geq 1$ be an integer, then

$$\|\partial_\alpha^n f\|_{L^\infty} \leq C \left(\frac{((n+1)!)^2}{(n+1)\sigma^{n+1}} + \frac{(n!)^2}{n\sigma^n} \right) \|f\|_{Y_\sigma}. \quad (2.43)$$

Proof. The proof follows from the standard Sobolev embedding $\|f\|_{L^\infty} \leq C\|f\|_{H^1}$. $\square$

The Hilbert transform is a unitary operator on these spaces.

$$\mathbb{H}f(\alpha) := \frac{1}{\pi i} \int_{\mathbb{R}} \frac{1}{\alpha - \beta} f(\beta) d\beta. \quad (2.44)$$

Lemma 2.4. Let $\sigma > 0$ and $f \in X_\sigma$, we have

$$\|\mathbb{H}f\|_{X_\sigma} = \|f\|_{X_\sigma}, \quad (2.45)$$

$$\|\mathbb{H}f\|_{\dot{X}_\sigma} = \|f\|_{\dot{X}_\sigma}, \quad (2.46)$$

$$\|\mathbb{H}f\|_{Y_\sigma} = \|f\|_{Y_\sigma}. \quad (2.47)$$

and

$$\|\mathbb{H}f\|_{\dot{Y}_\sigma} = \|f\|_{\dot{Y}_\sigma}. \quad (2.48)$$

Proof. We prove (2.45) only. (2.47) and (2.48) are proved similarly. Using the fact that $\|\partial_\alpha^n \mathbb{H}f\|_{L^2} = \|\mathbb{H}\partial_\alpha^n f\|_{L^2} = \|\partial_\alpha^n f\|_{L^2}$, we have

$$\|\mathbb{H}f\|_{X_\sigma}^2 = \sum_{n \geq 0} \frac{\sigma^{2n}}{(n!)^4} \|\partial_\alpha^n \mathbb{H}f\|_{L^2}^2 = \sum_{n \geq 0} \frac{\sigma^{2n}}{(n!)^4} \|\partial_\alpha^n f\|_{L^2}^2 = \|f\|_{X_\sigma}^2.$$

$\square$

Lemma 2.5 (Product estimates). *Let $\sigma \geq \sigma_0 > 0$. We have*

$$\|fg\|_{X_\sigma} \leq C\|f\|_{X_\sigma}\|g\|_{X_\sigma}, \quad (2.49)$$

$$\|fg\|_{Y_\sigma} \leq C\|f\|_{Y_\sigma}\|g\|_{Y_\sigma}, \quad (2.50)$$

$$\|fg\|_{\dot{X}_\sigma} \leq C\|f\|_{\dot{X}_\sigma}\|g\|_{X_\sigma} + C\|f\|_{X_\sigma}\|g\|_{\dot{X}_\sigma}, \quad (2.51)$$

$$\|fg\|_{\dot{Y}_\sigma} \leq C\|f\|_{\dot{Y}_\sigma}\|g\|_{Y_\sigma} + C\|f\|_{Y_\sigma}\|g\|_{\dot{Y}_\sigma}, \quad (2.52)$$

for some constant $C > 0$ depends linearly on $1 + \frac{1}{\sigma_0}$.

Proof. We prove (2.49) only.

First, note that

$$\left(\sum_{n=0}^{10} \frac{\sigma^{2n}}{(n!)^4} \|\partial_\alpha^n(fg)\|_{L^2}^2 \right)^{1/2} \leq C\|f\|_{X_\sigma}\|g\|_{X_\sigma}. \quad (2.53)$$

Second, for $n > 10$, we estimate $\|\partial_\alpha^n(fg)\|_{L^2}$ by

$$\|\partial_\alpha^n(fg)\|_{L^2} \leq \sum_{k=0}^n \frac{n!}{k!(n-k)!} \|f^{(k)}g^{(n-k)}\|_{L^2} \quad (2.54)$$

$$\leq \sum_{k=0}^{[n/2]} \frac{n!}{k!(n-k)!} \|f^{(k)}g^{(n-k)}\|_{L^2} + \sum_{k=[n/2]+1}^n \frac{n!}{k!(n-k)!} \|f^{(k)}g^{(n-k)}\|_{L^2} \quad (2.55)$$

$$:= I + II. \quad (2.56)$$

Here, $f^{(k)}, g^{(n-k)}$ represent the k -th and $(n-k)$ -th derivative of f and g , respectively. For I , we estimate $f^{(k)}$ in L^∞ and estimate $g^{(n-k)}$ in L^2 , and we have

$$\begin{aligned} \|f^{(k)}g^{(n-k)}\|_{L^2} &\leq \|f^{(k)}\|_{L^\infty}\|g^{(n-k)}\|_{L^2} \leq C\left(\frac{(k!)^2}{\sigma^k} + \frac{((k+1)!)^2}{\sigma^{k+1}}\right)\|f\|_{X_\sigma} \frac{((n-k)!)^2}{\sigma^{n-k}}\|g\|_{X_\sigma} \\ &= C\left(1 + \frac{(k+1)^2}{\sigma}\right) \frac{(k!)^2((n-k)!)^2}{\sigma^n} \|f\|_{X_\sigma}\|g\|_{X_\sigma}. \end{aligned} \quad (2.57)$$

Moreover, we further decompose I as

$$I = \sum_{k=0}^4 \frac{n!}{k!(n-k)!} \|f^{(k)}g^{(n-k)}\|_{L^2} + \sum_{k=5}^{[n/2]} \frac{n!}{k!(n-k)!} \|f^{(k)}g^{(n-k)}\|_{L^2} := I_1 + I_2. \quad (2.58)$$

For $n > 10$ and $0 \leq k \leq 4$, we have

$$\frac{\sigma^n}{(n!)^2} I_1 \leq Cn^{-2} \|f\|_{X_\sigma}\|g\|_{X_\sigma}. \quad (2.59)$$

For $k \geq 5$ and $n > 10$, we have

$$\frac{n!}{k!(n-k)!} = \frac{(n-k)!}{(n-k)!} \frac{\prod_{j=0}^{k-1} (n-j)}{k!} \geq Cn^4. \quad (2.60)$$

Using (2.57) and (2.60), we obtain

$$\begin{aligned} \frac{\sigma^n}{(n!)^2} I_2 &\leq C \sum_{k=5}^{[n/2]} \frac{k!(n-k)!}{n!} \frac{1}{(k!)^2((n-k)!)^2} \left(1 + \frac{k^2}{\sigma}\right) \frac{(k!)^2((n-k)!)^2}{\sigma^n} \|f\|_{X_\sigma}\|g\|_{X_\sigma} \\ &\leq C \sum_{k=5}^{[n/2]} \frac{1}{n^4} \left(1 + \frac{k^2}{\sigma}\right) \|f\|_{X_\sigma}\|g\|_{X_\sigma} \\ &\leq Cn^{-1} \|f\|_{X_\sigma}\|g\|_{X_\sigma}. \end{aligned}$$

So we obtain

$$\frac{\sigma^n}{(n!)^2} I \leq C n^{-1} \|f\|_{X_\sigma} \|g\|_{X_\sigma}.$$

Similarly,

$$\frac{\sigma^n}{(n!)^2} II \leq C n^{-1} \|f\|_{X_\sigma} \|g\|_{X_\sigma}.$$

Then we obtain

$$\begin{aligned} \|fg\|_{X_\sigma}^2 &= \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \|\partial_\alpha^n(fg)\|_{L^2}^2 \\ &\leq \sum_{n=0}^{\infty} (1+n^2)^{-1} \|f\|_{X_\sigma}^2 \|g\|_{X_\sigma}^2 \\ &\leq C \|f\|_{X_\sigma}^2 \|g\|_{X_\sigma}^2. \end{aligned}$$

□

We have also the following estimates.

Lemma 2.6. *Let $\sigma \geq \sigma_0 > 0$ and $f, g \in X_\sigma$. If in addition*

$$\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n f\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n g\|_{L^2}^2 < \infty,$$

then

$$\left| \sum_{n=1}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \langle \partial_\alpha^{n+1}(fg), \partial_\alpha^n g \rangle \right| \leq d_1 \|f\|_{\dot{X}_\sigma} \|g\|_{\dot{X}_\sigma}^2 + d_1 \|g\|_{X_\sigma} \left(\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n f\|_{L^2}^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n g\|_{L^2}^2 \right)^{1/2}, \quad (2.61)$$

where $d_1 = d(\sigma_0^{-1} + 1)$ for some absolute constant $d > 0$. In particular, if $\sigma_0 \geq 1$, then d_1 is an absolute constant.

Proof. We have

$$\partial_\alpha^{n+1}(fg) = g \partial_\alpha^{n+1} f + f \partial_\alpha^{n+1} g + \sum_{k=1}^{n+1} \frac{(n+1)!}{k!(n+1-k)!} \partial_\alpha^k f \partial_\alpha^{n+1-k} g.$$

Using the same proof as in Lemma 2.5, we obtain

$$\left| \sum_{n=1}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \left\langle \sum_{k=1}^{n+1} \frac{(n+1)!}{k!(n+1-k)!} \partial_\alpha^k f \partial_\alpha^{n+1-k} g, \partial_\alpha^n g \right\rangle \right| \leq d_1 \|f\|_{\dot{X}_\sigma} \|g\|_{\dot{X}_\sigma}^2.$$

By Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\left| \sum_{n=1}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \langle g \partial_{\alpha}^{n+1} f, \partial_{\alpha}^n g \rangle \right| &\leq \left(\sum_{n=1}^{\infty} \frac{\sigma^{2n}}{n(n!)^4} \|g \partial_{\alpha}^{n+1} f\|_{L^2}^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n g\|_{L^2}^2 \right)^{1/2} \\
&\leq \|g\|_{L^{\infty}} \left(\sum_{n=1}^{\infty} \frac{\sigma^{2n}}{n(n!)^4} \|\partial_{\alpha}^{n+1} f\|_{L^2}^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n g\|_{L^2}^2 \right)^{1/2} \\
&= \|g\|_{L^{\infty}} \left(\sum_{n=1}^{\infty} \frac{\sigma^{2(n+1)}}{((n+1)!)^4} \frac{((n+1)!)^4}{\sigma^2 n(n!)^4} \|\partial_{\alpha}^{n+1} f\|_{L^2}^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n g\|_{L^2}^2 \right)^{1/2} \\
&\leq d_1 \|g\|_{L^{\infty}} \left(\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n f\|_{L^2}^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n g\|_{L^2}^2 \right)^{1/2},
\end{aligned}$$

where we can take $d_1 = 2\sigma^{-1} + K$, for some absolute constant $K > 0$. Integration by parts, we obtain

$$\left| \sum_{n=1}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \langle f \partial_{\alpha}^{n+1} g, \partial_{\alpha}^n g \rangle \right| = \left| \sum_{n=1}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \langle f_{\alpha} \partial_{\alpha}^n g, \partial_{\alpha}^n g \rangle \right| \leq \|f_{\alpha}\|_{L^{\infty}} \|g\|_{\dot{X}_{\sigma}}^2 \leq d_1 \|f\|_{\dot{X}_{\sigma}} \|g\|_{\dot{X}_{\sigma}}^2.$$

Using $\|g\|_{L^{\infty}} \leq 2(1 + \sigma^{-1})\|g\|_{X_{\sigma}}$, we conclude the proof of the lemma. $\square$

2.3. Commutator estimates in Gevrey spaces. Let $f, g \in X_{\sigma}$. Define

$$S_1(g, f)(\alpha, t) := \frac{1}{\pi} p.v. \int_{\mathbb{R}} \frac{g(\alpha, t) - g(\beta, t)}{(\alpha - \beta)^2} f(\beta, t) d\beta. \quad (2.62)$$

$$S_2(g, f)h(\alpha, t) := \frac{1}{\pi} p.v. \int_{\mathbb{R}} \frac{(g(\alpha, t) - g(\beta, t))(f(\alpha, t) - f(\beta, t))}{(\alpha - \beta)^2} \frac{h_{\beta}}{Z_{\beta}} d\beta. \quad (2.63)$$

The following commutator estimates are the Gevrey version of the Sobolev counterpart.

Lemma 2.7. 1. Let $f \in Y_{\sigma}$, $g \in X_{\sigma}$, then

$$\|[f, \mathbb{H}]g_{\alpha}\|_{X_{\sigma}} \leq C \|f\|_{\dot{X}_{\sigma}} \|g\|_{X_{\sigma}}. \quad (2.64)$$

$$\|[f, \mathbb{H}]g_{\alpha}\|_{Y_{\sigma}} \leq C \|f\|_{\dot{Y}_{\sigma}} \|g\|_{X_{\sigma}}. \quad (2.65)$$

2. Let $f, g, h \in X_{\sigma}$, $Z - \alpha \in \dot{Y}_{\sigma}$. There holds

$$\|S_2(g, f)h\|_{X_{\sigma}} \leq C \|f\|_{X_{\sigma}} \|g\|_{X_{\sigma}} \|h\|_{X_{\sigma}} (1 + \|Z - \alpha\|_{\dot{Y}_{\sigma}}). \quad (2.66)$$

(3.) If in addition $\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n f\|_{L^2}^2 < \infty$, and $\sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n g\|_{L^2}^2 < \infty$, then

$$\|[f_{\alpha}, \mathbb{H}]g\|_{X_{\sigma}} \leq C \left(\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n f\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n g\|_{L^2}^2 \right) + C \|g\|_{L^2}^2. \quad (2.67)$$

Proof. We prove (2.64) only. It suffices to notice that

$$\|[f, \mathbb{H}]g_{\alpha}\|_{L^2} \leq C \min\{\|f\|_{L^2} \|g\|_{H^1}, \|f'\|_{L^{\infty}} \|g\|_{L^2}\}. \quad (2.68)$$

(2.68) follows easily from Fourier analysis, we omit the proof. Using (2.68), we obtain

(1) For $n = 0$:

$$\|[f, \mathbb{H}]g_{\alpha}\|_{L^2} \leq C \|f'\|_{L^{\infty}} \|g\|_{L^2} \leq C \|f\|_{\dot{X}_{\sigma}} \|g\|_{X_{\sigma}}. \quad (2.69)$$

(2) For $n \geq 1$, we have

$$\|\partial_{\alpha}^n [f, \mathbb{H}]g_{\alpha}\|_{L^2} \leq C (\|\partial_{\alpha}^n f\|_{L^2} \|g\|_{X_{\sigma}}). \quad (2.70)$$

Indeed, let $n \geq 1$, using $|\xi|^n |sgn(\xi) - sgn(\eta)| \leq |\xi - \eta|^n$, we have

$$\begin{aligned} \left| \widehat{\partial_\alpha^n [f, \mathbb{H}] g_\alpha}(\xi) \right| &\leq |\xi|^n \int |sgn(\eta) - sgn(\xi)| |\eta| \hat{f}(\xi - \eta) \hat{g}(\eta) d\eta \\ &\leq \int |\widehat{\partial_\alpha^n f}(\xi - \eta) \widehat{\partial_\alpha g}(\eta)| d\eta \\ &= |\widehat{\partial_\alpha^n f}| * |\widehat{\partial_\alpha g}|(\xi) \end{aligned}$$

So

$$\|\partial_\alpha^n [f, \mathbb{H}] g_\alpha\|_{L^2} \leq \| |\widehat{\partial_\alpha^n f}| * |\widehat{\partial_\alpha g}| \|_{L^2} \leq \|\widehat{\partial_\alpha^n f}\|_{L^2} \|\widehat{\partial_\alpha g}\|_{L^1}.$$

By Plancherel Theorem, $\| |\widehat{\partial_\alpha^n f}| * |\widehat{\partial_\alpha g}| \|_{L^2} = \|\partial_\alpha^n f\|_{L^2}$. Using

$$\widehat{g_\alpha}(\xi) = \frac{1}{|\xi|} |\xi| \widehat{g_\alpha}(\xi) = \frac{1}{|\xi|} \widehat{\partial_\alpha^2 g},$$

we have

$$\|\widehat{\partial_\alpha g}\|_{L^1} \leq \int_{|\xi| \leq 1} |\widehat{\partial_\alpha g}(\xi)| d\xi + \int_{|\xi| \geq 1} \frac{1}{|\xi|} |\widehat{\partial_\alpha^2 g}(\xi)| d\xi \leq \|g_\alpha\|_{H^1} \leq C \|g\|_{X_\sigma}.$$

So we obtain (2.70), and therefore conclude the proof of the lemma. $\square$

Lemma 2.8. *Let $H \in X_\sigma$ be such that $1 + H \geq c_0$ for some constant $c_0 > 0$. Then*

(1) $\frac{1}{H+1} - 1 \in X_\sigma$, and

$$\left\| \frac{1}{H+1} - 1 \right\|_{X_\sigma} \leq C(c_0) \|H\|_{X_\sigma}. \quad (2.71)$$

(2) If in addition $\sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_\alpha^n H\|_{L^2}^2 < \infty$, then

$$\sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \left\| \partial_\alpha^n \left(\frac{1}{H+1} - 1 \right) \right\|_{L^2}^2 \leq C(c_0) \sum_{n=1}^{\infty} \frac{n\sigma^{2n}}{(n!)^4} \|\partial_\alpha^n H\|_{L^2}^2. \quad (2.72)$$

Here $C(c_0)$ is a constant depending on c_0 .

The proof of Lemma 2.8 is similar to that of Lemma 2.5, so we omit the proof.

Lemma 2.9. *Let $w \in \mathbb{P}_-$ and define $h(\alpha) = \frac{1}{(\alpha-w)^2}$. Then $h \in X_\sigma$ for any $\sigma \in \mathbb{R}$, and*

$$\|h\|_{X_\sigma} \leq C \frac{1}{\sqrt{|Im\{w\}|}} e^{\frac{|\sigma|}{4\pi|Im\{w\}|}} \quad (2.73)$$

Proof. The Fourier transform of h is

$$\hat{h}(\xi) = -2\pi i (i\xi) e^{-w|\xi|}. \quad (2.74)$$

For $\sigma \in \mathbb{R}$,

$$\begin{aligned}
\|h\|_{\dot{X}_\sigma}^2 &= \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \|\partial_\alpha^n h\|_{L^2}^2 \\
&= \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \|\xi|^n \hat{h}(\xi)\|_{L^2}^2 \\
&= \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \int |\xi|^{2n+2} e^{2\pi \operatorname{Im}\{w\}|\xi|} d\xi \\
&= 2 \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \frac{1}{|2\pi \operatorname{Im}\{w\}|^{2n+1}} \int_0^\infty |\xi|^{2n} e^{-|\xi|} d\xi \\
&= 2 \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \frac{1}{|2\pi \operatorname{Im}\{w\}|^{2n+1}} (2n+1)! \\
&\leq \frac{1}{2\pi |\operatorname{Im}\{w\}|} \sum_{n=0}^{\infty} \left(\frac{\sigma}{2\pi |\operatorname{Im}\{w\}|} \right)^{2n} \frac{2^n}{(n!)^2} \\
&\leq C \frac{1}{|\operatorname{Im}\{w\}|} e^{\frac{|\sigma|}{2\pi |\operatorname{Im}\{w\}|}},
\end{aligned}$$

for some absolute constant $C > 0$. □

Lemma 2.10. *Let $\sigma > 0$ and let $f \in X_\sigma$, $g, h \in Y_\sigma$. Then*

(1)

$$\sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \left| \langle \partial_\alpha^n (f \Lambda g), \partial_\alpha^n h \rangle \right| \leq d_0 \|f\|_{X_\sigma} \|g\|_{Y_\sigma} \|h\|_{Y_\sigma}, \quad (2.75)$$

$$\sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \left| \langle \partial_\alpha^n (f \partial_\alpha g), \partial_\alpha^n h \rangle \right| \leq d_0 \|f\|_{X_\sigma} \|g\|_{Y_\sigma} \|h\|_{Y_\sigma}, \quad (2.76)$$

$$\sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \left| \langle \partial_\alpha^n (f \partial_\alpha g), \partial_\alpha^n g \rangle \right| \leq d_0 \|f\|_{\dot{X}_\sigma} \|g\|_{\dot{X}_\sigma}^2, \quad (2.77)$$

where $d_0 = K(1 + \sigma^{-1})$ for some absolute constant $K > 0$. In particular, if $\sigma \geq 1$, then d_0 is an absolute constant.

(2) If in addition $\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n g\|_{L^2}^2 < \infty$, and $\sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n h\|_{L^2}^2 < \infty$, then

$$\sum_{n=1}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \left| \langle \partial_\alpha^n (f \Lambda g), \partial_\alpha^n h \rangle \right| \leq d_0 \|f\|_{X_\sigma} \left(\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n g\|_{L^2}^2 \right)^{1/2} \left(\|h\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n h\|_{L^2}^2 \right)^{1/2} \quad (2.78)$$

Proof. Let $n \geq 1$, we have

$$\begin{aligned}
\left| \langle \partial_\alpha^n (f \Lambda g), \partial_\alpha^n h \rangle \right| &\leq \left| \langle f \partial_\alpha^n \Lambda g, \partial_\alpha^n h \rangle \right| + \left| \langle \partial_\alpha^n (f \Lambda g) - f \partial_\alpha^n \Lambda g, \partial_\alpha^n h \rangle \right| \\
&:= I_1 + I_2.
\end{aligned}$$

For I_1 , we have

$$\begin{aligned}
|I_1| &= \left| \int f(\partial_\alpha^n \Lambda g) \overline{\partial_\alpha^n h} \right| \\
&\leq \int \left| f(\partial_\alpha^n \Lambda^{1/2} g) \overline{\Lambda^{1/2} \partial_\alpha^n h} \right| + \int \left| g[f, \Lambda^{1/2}] \overline{\partial_\alpha^n h} \right| \\
&\leq d_0 \|f\|_{X_\sigma} \|\Lambda^{1/2} \partial_\alpha^n g\|_{L^2} \|\Lambda^{1/2} \partial_\alpha^n h\|_{L^2}.
\end{aligned}$$

So we obtain

$$\sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} I_1 \leq C \|f\|_{X_\sigma} \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} \|\Lambda^{1/2} \partial_\alpha^n g\|_{L^2} \|\Lambda^{1/2} \partial_\alpha^n h\|_{L^2} \leq d_0 \|f\|_{X_\sigma} \|g\|_{Y_\sigma} \|h\|_{Y_\sigma}.$$

The term I_2 can be handled in the same way as in the proof of Lemma 2.5, we omit the details. We have

$$\sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} I_1 \leq d_0 \|f\|_{X_\sigma} \|g\|_{Y_\sigma} \|h\|_{Y_\sigma}.$$

So we complete the proof of (2.75). The proof of (2.76) is the same as that of (2.75).

The proof of (2.77) is similar:

$$\begin{aligned} \left| \langle \partial_\alpha^n (f \Lambda g), \partial_\alpha^n g \rangle \right| &\leq \left| \langle f \partial_\alpha^n \Lambda g, \partial_\alpha^n g \rangle \right| + \left| \langle \partial_\alpha^n (f \Lambda g) - f \partial_\alpha^n \Lambda g, \partial_\alpha^n g \rangle \right| \\ &:= II_1 + II_2. \end{aligned}$$

The treatment of II_2 is the same as that for I_2 . For II_1 , we have

$$\begin{aligned} II_1 &= \left| \operatorname{Re} \int f \partial_\alpha^n \Lambda g \overline{\partial_\alpha^n g} \right| = \left| \operatorname{Re} \int |\partial_\alpha^n g|^2 \Lambda f \right| \\ &\leq d_0 \|\Lambda f\|_{L^\infty} \|\partial_\alpha^n g\|_{L^2}^2 \\ &\leq d_0 \|f\|_{\dot{X}_\sigma} \|\partial_\alpha^n g\|_{L^2}^2. \end{aligned}$$

So $\sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^4} I_1 \leq C \|f\|_{\dot{X}_\sigma} \|g\|_{\dot{X}_\sigma}^2$. So we obtain (2.77).

For the proof of (2.78), it suffices to notice that

$$\begin{aligned} \left| \int f(\partial_\alpha^n \Lambda g) \overline{\partial_\alpha^n h} \right| &\leq \int \left| f(\partial_\alpha^n \Lambda^{3/4} g) \overline{\Lambda^{1/4} \partial_\alpha^n h} \right| + \int \left| g[f, \Lambda^{1/4}] \overline{\partial_\alpha^n h} \right| \\ &\leq d_0 \|f\|_{X_\sigma} \|\Lambda \partial_\alpha^n g\|_{L^2} \|(1 + \Lambda^{1/4}) \partial_\alpha^n h\|_{L^2}. \end{aligned}$$

□

3. THE MAIN RESULTS: QUANTITATIVE STATEMENTS

Throughout the rest of this paper, for brevity, we consider a pair of symmetric and counter-rotating point vortices embedded in the water waves, the general situation can actually be treated similarly. We assume

$$\omega(\cdot, t) = \lambda \delta_{z_1(t)} - \lambda \delta_{z_2(t)}, \quad (3.1)$$

where $\lambda \in \mathbb{R}$, and $z_1(t) = -x(t) + iy(t)$, $z_2(t) = x(t) + iy(t)$, with $x(t) > 0$ and $z_1(t), z_2(t) \in \Omega(t)$. We assume also that $\Omega(t)$ is symmetric about the vertical axis⁵. Without loss of generality, we assume $x(0) = 1$. Moreover, we assume $|y(0)| \gg 1$. It should be clear from the proof that regarding the local wellposedness in Gevrey spaces, these assumptions are unnecessary.

3.1. The initial data. Let the initial fluid region be given by Ω_0 , with a nonself-intersect smooth free surface Σ_0 . We parametrize Σ_0 by Riemann variable, that is, we choose α such that $h_0(\alpha) = \alpha$.

Let $Z_0(\alpha) := Z(\alpha, 0)$. $U_0 = \Re\{F_0\}$, and $W_0 = \Re\{Z_0 - \alpha\}$. Recall that

$$d_I(t) = \inf_{\alpha \in \mathbb{R}} \min_{1 \leq j \leq 2} |Z(\alpha, t) - z_j(t)|,$$

and $d_{I,0} := d_I(0)$, $x_0 := x(0) = 1$. Without loss of generality, we assume

$$d_{I,0} = \inf_{\alpha \in \mathbb{R}} \min_{j=1,2} |\Im\{Z(\alpha, 0) - z_j(0)\}|.$$

⁵Such symmetry assumption can be removed, we assume it for convenience.

Let x_0, y_0, λ , and $L_0 \geq 4$ be given constants. Denote $\phi(t) = L_0 - \delta_0 t$. We assume the following:

(H1) (W_0, U_0) is given such that $(\partial_\alpha W_0, U_0) \in X_{L_0} \times Y_{L_0}$. Without loss of generality, we assume for simplicity that⁶

$$\|(U_0, \partial_\alpha W_0)\|_{Y_{L_0} \times X_{L_0}} \leq 1/2. \quad (3.2)$$

(H2) δ_0 is chosen such that

$$\frac{\delta_0}{L_0} - 4 - 2d_0(\|A(\cdot, 0)\|_{L^\infty} + \|A(\cdot, 0)\|_{\dot{Y}_{\phi(t)}}) - 4d_1\|(U_0, \partial_\alpha W_0)\|_{Y_{\phi(0)} \times X_{\phi(0)}} \geq 0. \quad (3.3)$$

Here, $A(\cdot, 0) = \frac{A_{1,0}}{|\partial_\alpha Z_0|^2}$, with $A_{1,0}$ given by (2.36), and d_0 and d_1 are the constants given in Lemma 2.10 and Lemma 2.6, respectively.

(H3) $z_{1,0} = -x_0 + iy_0$, $z_{2,0} = x_0 + iy_0$, where x_0 and y_0 are constants, satisfying $x_0 = 1$, $y_0 < 0$, and $|y_0| \gg 1$.

(H4) $|Z_0(\alpha) - Z_0(\beta)| \geq C_0|\alpha - \beta|$, for some absolute constant $C_0 > 0$.

(H5) $Z_0(\alpha) := \alpha + (I + \mathbb{H})W_0$, $F_0 = (I + \mathbb{H})U_0$.

(H6) Ω_0 is symmetric about the y -axis, W_0 is an odd function. U_0 is also an odd function.

For $0 \leq t \leq \frac{L_0}{2\delta_0}$, we have $\phi(t) \geq \frac{1}{2}L_0$. Denote

$$\tau_0 := \frac{1}{4 + 2d_0(\|A(\cdot, 0)\|_{L^\infty} + \|A(\cdot, 0)\|_{\dot{Y}_{\phi(t)}}) + 4d_1\|(U_0, \partial_\alpha W_0)\|_{Y_{\phi(0)} \times X_{\phi(0)}}}. \quad (3.4)$$

In particular, $\delta_0 \geq 4L_0 \geq 16$.

Remark 3.1. The assumption that $|y_0| \gg 1$ is not essential in terms of the local wellposedness in Gevrey-2 spaces for the system (2.33). We can certainly remove it. Nevertheless, for the purpose of proving Theorem 1, we do need to assume $|y_0| \gg 1$. Therefore, for convenience, we make such an assumption.

3.2. The main theorems.

Theorem 3. Let $W_0, U_0, \{z_{j,0}\}$, L_0, δ_0, Z_0 , and F_0 be given such that (H1)-(H6) hold. Then there exists a constant $T > 0$, such that the system (2.33) with initial data $(W_0, U_0, \{z_{j,0}\})$ admits a unique solution $(W, U, \{z_j\})$ satisfying

(a) $(W_\alpha, U) \in C([0, T]; X_{\phi(t)} \times Y_{\phi(t)})$, and $z_j \in C^1([0, T]; \Omega(t))$, $j = 1, 2$. Moreover,

$$\sup_{0 \leq t \leq T} \left(\|U(\cdot, t)\|_{\dot{Y}_{\phi(t)}}^2 + \|W_\alpha\|_{X_{\phi(t)}}^2 \right) \leq \begin{cases} \|(U_0, \partial_\alpha W_0)\|_{\dot{Y}_{\phi(t)} \times X_{\phi(t)}}^2 + C|\lambda|(d_{I,0} - \frac{|\lambda|}{4\pi x(0)}T)^{-5/2}, & \lambda > 0, \\ \|(U_0, \partial_\alpha W_0)\|_{\dot{Y}_{\phi(t)} \times X_{\phi(t)}}^2 + C|\lambda|d_{I,0}^{-5/2}, & \lambda < 0 \end{cases}$$

and

$$\sup_{0 \leq t \leq T} \|U(\cdot, t)\|_{L^2}^2 \leq 1, \quad \sup_{0 \leq t \leq T} \|U(\cdot, t)\|_{L^\infty}^2 \leq Cd_{I,0}^{-1/3}.$$

Here, the constant C depends on C_0 and $\frac{1}{L_0}$.

(b) $|Z(\alpha, t) - Z(\beta, t)| \geq \frac{1}{2}C_0|\alpha - \beta|$, $\alpha, \beta \in \mathbb{R}$, $t \in [0, T]$.

(c) $d_I(t) \geq \frac{1}{2}|y_0|^{9/10}$, $\frac{1}{2}x(0) \leq x(t) \leq 2x(0)$, $t \in [0, T]$.

(d) $\Omega(t)$ is symmetric about the y -axis. For each fixed $t \in [0, T]$, $W(\cdot, t)$ and $U(\cdot, t)$ are odd functions. $\Re\{z_1(t)\} = -\Re\{z_2(t)\}$, $\Im\{z_1(t)\} = \Im\{z_2(t)\}$.

⁶In application, we will take $\|(U_0, \partial_\alpha W_0)\|_{Y_{L_0} \times X_{L_0}}$ to be small.

Here ⁷,

$$T = \begin{cases} \min\{T_1(C_0, \|(U_0, \partial_\alpha W_0)\|_{\dot{Y}_{L_0} \times X_{L_0}}), \frac{L_0}{2\delta_0}, \frac{4\pi x(0)(|y_0| - |y_0|^{9/10})}{|\lambda|}\}, & \lambda > 0, \\ \min\{T_2(C_0, \|(U_0, \partial_\alpha W_0)\|_{\dot{Y}_{L_0} \times X_{L_0}}), \frac{L_0}{2\delta_0}\} & \lambda < 0, \end{cases}$$

with T_1, T_2 depend continuously on its parameters. In particular, if $|y_0|$ is sufficiently large, then if $\lambda > 0$, we can take $T = \frac{4\pi x(0)(|y_0| - |y_0|^{9/10})}{|\lambda|}$; if $\lambda < 0$, we can take $T = O(1)$.

Remark 3.2. (1) Using the same proof as in [44], we can prove that the solutions to the water waves preserve the symmetries in (H6). We shall omit the proof of (d) and refer the readers to Theorem 5 in [44].

(2) From the discussion in §1.2.2, in particular, the equation (1.9), our wellposedness result is sharp in the Gevrey spaces. That is, if we define

$$X_{\sigma,k} := \{f \in C^\infty : \|f\|_{X_{\sigma,k}}^2 = \sum_{n=0}^{\infty} \frac{\sigma^{2n}}{(n!)^{2k}} \|\partial_\alpha^n f\|_{L^2}^2\},$$

and $\dot{X}_{\sigma,k}, Y_{\sigma,k}, \dot{Y}_{\sigma,k}$ similarly. Then (2.33) is illposedness in space $(U, W_\alpha) \in Y_{\phi(t),k} \times X_{\phi(t),k}$ for $k > 2$. We shall prove this in a separate paper.

Using Theorem 3, we are able to prove the following.

Theorem 4. Let $(W, U, \{z_1(t), z_2(t)\})$ be the unique solution to (2.33) on $[0, T]$ constructed in Theorem 3 with initial data $(W_0, U_0, \{z_1(0), z_2(0)\})$. For any given constants $0 < \eta_0 < 1$, $\eta_1 > 0$, $\gamma > 0$ and $0 < \epsilon_0 \ll 1$, there exist constants $N_0 \gg 1$, $\delta_0, \delta_1, \delta_2 \ll 1$ such that for all

$$y(0) \leq -N_0 \quad \lambda = \gamma|y_0|^{3/2-\epsilon_0}, \quad |x(0) - 1| \leq \epsilon_1, \quad \|(U_0, \partial_\alpha W_0)\|_{Y_{L_0} \times X_{L_0}} \leq \epsilon_2,$$

there holds,

$$\inf_{\alpha \in \mathbb{R}} A_1(\alpha, 0) \geq 1 - \eta_0, \quad \inf_{\alpha \in \mathbb{R}} A_1(\alpha, T) < -\eta_1,$$

For example, we can take $\epsilon_1 = \frac{1}{N_0}$, $\epsilon_2 = \frac{1}{N_0}$, $\epsilon_0 = \frac{1}{10}$, $L_0 = 10$, $\delta_0 = 1000$.

Remark 3.3. The choice of the parameters $\epsilon_0, \epsilon_1, \epsilon_2$, etc are certainly not optimal. It is not our primary goal in this paper to obtain those sharp bounds.

4. A QUASILINEAR SYSTEM

Given b_1, G, R, A, U_0 and W_0 , we consider the following quasilinear system (with G and R as external forces):

$$\begin{cases} D_t U = A \Lambda W + G, \\ D_t W = -U - \Re\{[\bar{F}, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\} + R, \\ (U, W)(\cdot, 0) = (U_0, W_0). \end{cases} \quad (4.1)$$

Here, $Z_\alpha = (I + \mathbb{H})W_\alpha + 1$. $D_t = \partial_t + b\partial_\alpha$, with

$$b = b_0 + b_1, \quad b_0 = 2U + \Re\{[(I - \mathbb{H})U, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\},$$

⁷The power $\frac{9}{10}$ for $|y_0|^{9/10}$ is not optimal. We can use any $|y_0|^{1-\epsilon}$, where $\epsilon \in (0, 1)$.

where b_1 is an a priori given real valued function. Using $\partial_\alpha[f, \mathbb{H}]g = [f_\alpha, \mathbb{H}]g + [f, \mathbb{H}]g_\alpha$, and by (2.64) and (2.67), we have

$$\begin{aligned} & \|\partial_\alpha \Re\{(I - \mathbb{H})U, \mathbb{H}\}(\frac{1}{Z_\alpha} - 1)\|_{X_{\phi(t)}} \\ & \leq C(\|U\|_{\dot{X}_{\phi(t)}}^2 + \|\frac{1}{Z_\alpha} - 1\|_{X_{\phi(t)}}^2) + C\left(\sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n}}{(n!)^4} \|\partial_\alpha^n U\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n \phi(t)^{2n}}{(n!)^4} \|\partial_\alpha^n (\frac{1}{Z_\alpha} - 1)\|_{L^2}^2\right), \end{aligned} \quad (4.2)$$

for some constant $C > 0$ depending on $\frac{1}{\phi(t)}$.

Denote

$$H := (I + \mathbb{H})W_\alpha.$$

Let $\phi(t) = \phi_{\delta_0, L_0}(t) := L_0 - \delta_0 t$. Let

$$E_{L_0, \delta_0}(t) := \frac{1}{2} \|(U(\cdot, t), \partial_\alpha W(\cdot, t))\|_{\dot{Y}_{\phi(t)} \times X_{\phi(t)}}^2. \quad (4.3)$$

By Lemma 2.8, if $\inf_{\alpha \in \mathbb{R}}(H + 1) \geq c_0/2$, then

$$\left\| \frac{1}{H + 1} - 1 \right\|_{X_{\phi(t)}} \leq C(c_0) \|H\|_{X_{\phi(t)}} \leq C(c_0) \|W_\alpha\|_{X_{\phi(t)}}, \quad (4.4)$$

and

$$\sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \left\| \partial_\alpha^n \left(\frac{1}{H + 1} - 1 \right) \right\|_{L^2}^2 \leq C(c_0) \sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n H\|_{L^2}^2 \leq C(c_0) \sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_\alpha^n W_\alpha\|_{L^2}^2. \quad (4.5)$$

Theorem 5. *Let $c_0 > 0$ and $L_0 \geq 4$ be given constants. Assume that $b_1 \in C([0, T]; Y_{\phi(t)})$, $A - 1, R_\alpha \in C([0, T]; X_{\phi(t)})$, $G \in C([0, T]; Y_{\phi(t)})$ and $(U_0, \partial_\alpha W_0) \in Y_{\phi(0)} \times X_{\phi(0)}$. Assume further that $\inf_{\alpha \in \mathbb{R}}(1 + H(\alpha, 0)) \geq c_0$. Let d_0 and d_1 be the constant given in Lemma 2.10 and Lemma 2.6, respectively. Assume that*

$$\sup_{0 \leq t \leq T} \left(\frac{\delta_0}{L_0} - 4 - 2d_0(\|A\|_{L^\infty} + \|A\|_{\dot{Y}_{\phi(t)}}) - 4d_1\|(U_0, \partial_\alpha W_0)\|_{Y_{\phi(0)} \times X_{\phi(0)}} \right) \geq 0. \quad (4.6)$$

Then there exists $0 < T_0 \leq T$ such that (4.1) admits a unique solution $(U, \partial_\alpha W) \in C^0([0, T]; Y_{\phi(t)} \times X_{\phi(t)})$. Moreover,

$$E_{L_0, \delta_0}(t) \leq E_{L_0, \delta_0}(0) e^{B_{T_0} t} + \int_0^t e^{B_{T_0} \tau} \mathcal{N}(\tau) d\tau, \quad (4.7)$$

$$\|U(\cdot, t)\|_{L^2}^2 \leq \|U(\cdot, 0)\|_{L^2}^2 e^{\gamma(t)} + \int_0^t e^{\gamma(t) - \gamma(s)} (\|G(s)\|_{L^2}^2 + \|A(s)\|_{L^\infty} E_{L_0, \delta_0}(s)) ds, \quad (4.8)$$

where

$$B_{T_0} := C(1 + \|b_1\|_{C([0, T_0]; Y_{\phi(t)})}), \quad (4.9)$$

and

$$\mathcal{N}(t) := \|R_\alpha\|_{X_{\phi(t)}}^2 + \|G\|_{\dot{Y}_{\phi(t)}}^2 + C\|b_1\|_{\dot{Y}_{\phi(t)}}^2, \quad (4.10)$$

$$\gamma(t) := \int_0^t C(E_{L_0, \delta_0}^{1/2}(\tau) + \|b_1(\cdot, \tau)\|_{\dot{Y}_{\phi(\tau)}} + \|A(\cdot, t)\|_{L^\infty}) d\tau.$$

Here, $T_0 = \min\{T, \frac{L_0}{2\delta_0}\}$, and C is a constant depending on c_0 and $\frac{1}{L_0}$.

We prove *Theorem 5* by the energy method. We provide closed a priori energy estimates only. Note that $T_0 \leq \frac{L_0}{2\delta_0}$ implies that $\phi(t) \geq \frac{L_0}{2} \geq 2$ for $t \in [0, T_0]$, so d_0 and d_1 are absolute constants. By Lemma 2.7 and (4.5), under the assumption of *Theorem 5*, we have the a priori estimate

$$\begin{aligned} \|b_0\|_{\dot{Y}_{\phi(t)}}^2 &= \sum_{n=1}^{\infty} \frac{n^2 \phi(t)^{2n}}{(n!)^4} \left\| \partial_{\alpha}^n \left(2U + \Re\{[(I - \mathbb{H})U, \mathbb{H}](\frac{1}{Z_{\alpha}} - 1)\} \right) \right\|_{L^2}^2 \\ &\leq C \|U\|_{\dot{Y}_{\phi(t)}}^2 + C \|U\|_{\dot{Y}_{\phi(t)}} \|\partial_{\alpha} W\|_{X_{\phi(t)}} \\ &\leq C E_{L_0, \delta_0}, \end{aligned} \quad (4.11)$$

for some constant $C > 0$ depending on c_0 and $\frac{1}{L_0}$.

Proof. Applying ∂_{α} on both sides of $D_t W = -U - \Re\{[\bar{F}, \mathbb{H}](\frac{1}{Z_{\alpha}} - 1)\} + R$, we obtain

$$D_t W_{\alpha} = -U_{\alpha} + R_{\alpha} - b_{\alpha} W_{\alpha} - \partial_{\alpha} \Re\{\bar{F}, \mathbb{H}](\frac{1}{Z_{\alpha}} - 1)\}. \quad (4.12)$$

Using (4.12) and $U_t = A\Lambda W - bU_{\alpha} + G$, we have

$$\begin{aligned} \frac{d}{dt} E_{L_0, \delta_0}(t) &= \frac{1}{2} \frac{d}{dt} \left(\sum_{n=1}^{\infty} \frac{n^2 \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n U(\cdot, t)\|_{L^2}^2 \right) + \frac{1}{2} \frac{d}{dt} \sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}(\cdot, t)\|_{L^2}^2 \\ &= -\delta_0 \sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n-1}}{(n!)^4} \|\partial_{\alpha}^n U(\cdot, t)\|_{L^2}^2 + \Re \sum_{n=1}^{\infty} \frac{n^2 \phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n U_t, \partial_{\alpha}^n U \rangle \\ &\quad - \delta_0 \sum_{n=1}^{\infty} \frac{n \phi(t)^{2n-1}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}(\cdot, t)\|_{L^2}^2 + \Re \sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n \partial_{\alpha} W_t, \partial_{\alpha}^n W_{\alpha} \rangle \\ &= -\delta_0 \sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n-1}}{(n!)^4} \|\partial_{\alpha}^n U(\cdot, t)\|_{L^2}^2 \\ &\quad + \Re \sum_{n=1}^{\infty} \frac{n^2 \phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n (A\Lambda W - bU_{\alpha} + G), \partial_{\alpha}^n U \rangle \\ &\quad - \delta_0 \sum_{n=1}^{\infty} \frac{n \phi(t)^{2n-1}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}(\cdot, t)\|_{L^2}^2 \\ &\quad + \Re \sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n (-U_{\alpha} + R_{\alpha} - b_{\alpha} W_{\alpha} - \partial_{\alpha} \Re\{[\bar{F}, \mathbb{H}](\frac{1}{Z_{\alpha}} - 1)\} - b \partial_{\alpha} W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \end{aligned}$$

Using Lemma 2.10, we have ⁸

$$\begin{aligned} &\left| \sum_{n=1}^{\infty} \frac{n^2 \phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n (A\Lambda W), \partial_{\alpha}^n U \rangle \right| \\ &\leq d_0 (\|A\|_{L^{\infty}} + \|A\|_{\dot{X}_{\phi(t)}}) \left(\sum_{n=1}^{\infty} \frac{n \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n \Lambda W\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 \right), \end{aligned}$$

⁸It turns out that if we choose $|y_0|$ large and $|\lambda| \sim |y_0|^{3/2}$, then $\|A - 1\|_{L^{\infty}} \sim 1$, while $\|A - 1\|_{L^2} \sim |y_0|^{1/2}$, which is large. So we avoid to bound $A - 1$ in L^2 .

for some absolute constant $d_0 > 0$. Using Lemma 2.10 again, we obtain

$$\begin{aligned} \left| \Re \sum_{n=1}^{\infty} \frac{n^2 \phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n (A \Lambda W - b U_{\alpha} + G), \partial_{\alpha}^n U \rangle \right| &\leq C((\|b\|_{L^{\infty}} + \|b\|_{\dot{Y}_{\phi(t)}}) \|U\|_{\dot{Y}_{\phi(t)}}^2 + \|U\|_{\dot{Y}_{\phi(t)}}^2 + \|G\|_{\dot{Y}_{\phi(t)}}^2) \\ &\quad + d_0(\|A\|_{L^{\infty}} + \|A\|_{\dot{Y}_{\phi(t)}}) \left(\sum_{n=1}^{\infty} \frac{n \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n \Lambda W\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 \right) \\ &\leq C(1 + \|b\|_{L^{\infty}} + \|b\|_{\dot{Y}_{\phi(t)}}) E_{L_0, \delta_0}(t) + \|G\|_{\dot{Y}_{\phi(t)}}^2 \\ &\quad + d_0(\|A\|_{L^{\infty}} + \|A\|_{\dot{Y}_{\phi(t)}}) \left(\sum_{n=1}^{\infty} \frac{n \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n \Lambda W\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 \right). \end{aligned}$$

Cauchy-Schwarz implies

$$\left| \Re \sum_{n=1}^{\infty} \frac{n^2 \phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n U_{\alpha}, \partial_{\alpha}^n W_{\alpha} \rangle \right| \leq \sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}\|_{L^2}^2.$$

Decomposing $b = b_0 + b_1$, we obtain

$$\begin{aligned} \left| \Re \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n (b \partial_{\alpha} W_{\alpha} + b_{\alpha} W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \right| &= \left| \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^{n+1} (b W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \right| \\ &\leq \left| \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^{n+1} (b_0 W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \right| + \left| \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^{n+1} (b_1 W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \right|. \end{aligned}$$

Since we can estimate $\partial_{\alpha} b_1$ in $X_{\phi(t)}$, expressing $\partial_{\alpha}^{n+1} (b_1 W_{\alpha}) = \partial_{\alpha}^n (\partial_{\alpha} b_1 W_{\alpha} + b_1 \partial_{\alpha} W_{\alpha})$ and using (2.77), we obtain

$$\left| \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^{n+1} (b_1 W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \right| \leq \|\partial_{\alpha} b_1\|_{X_{\phi(t)}} \|\partial_{\alpha} W\|_{X_{\phi(t)}}^2.$$

The formula for b_0 implies

$$\begin{aligned} &\left| \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^{n+1} (b_0 W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \right| \\ &\leq \left| \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^{n+1} (2U W_{\alpha}), \partial_{\alpha}^n W_{\alpha} \rangle \right| + \left| \sum_{n=1}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^{n+1} (W_{\alpha} \Re[(I - \mathbb{H})U, \mathbb{H}](\frac{1}{Z_{\alpha}} - 1)), \partial_{\alpha}^n W_{\alpha} \rangle \right| \\ &:= I + II. \end{aligned}$$

Using Lemma 2.6, we have

$$\begin{aligned} I &\leq d_1 \|U\|_{\dot{X}_{\sigma}} \|W_{\alpha}\|_{\dot{X}_{\sigma}}^2 + d_1 \|W_{\alpha}\|_{X_{\sigma}} \left(\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 \right)^{1/2} \left(\sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}\|_{L^2}^2 \right)^{1/2} \\ &\leq d_1 E_{L_0, \delta_0}^{3/2} + d_1 E_{L_0, \delta_0}^{1/2} \left(\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}\|_{L^2}^2 \right), \end{aligned}$$

where d_1 is the constant in Lemma 2.6. Similarly, using Lemma 2.6, (4.4) and (4.5), and $\partial_{\alpha}[f, \mathbb{H}]g = [f_{\alpha}, \mathbb{H}]g + [f, \mathbb{H}]g_{\alpha}$, we obtain

$$II \leq d_1 E_{L_0, \delta_0}^{3/2} + d_1 E_{L_0, \delta_0}^{1/2} \left(\sum_{n=1}^{\infty} \frac{n^3 \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n \sigma^{2n}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}\|_{L^2}^2 \right).$$

By Cauchy-Schwarz,

$$\left| \Re \sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \langle \partial_{\alpha}^n R_{\alpha}, \partial_{\alpha}^n W_{\alpha} \rangle \right| \leq \|R_{\alpha}\|_{\dot{X}_{\phi(t)}}^2 + \|W_{\alpha}\|_{\dot{X}_{\phi(t)}}^2.$$

Combining the above estimates together with (4.11), bounding $\|b_1\|_{L^{\infty}}$ by $C\|b_1\|_{Y_{\phi(t)}}$, we obtain

$$\begin{aligned} & \frac{d}{dt} E_{L_0, \delta_0}(t) \\ & + \left(\frac{\delta_0}{\phi(t)} - 2 - d_0(\|A\|_{L^{\infty}} + \|A\|_{\dot{X}_{\phi(t)}}) - d_1 E_{L_0, \delta_0}^{1/2} \right) \left(\sum_{n=1}^{\infty} \frac{n^3 \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n U\|_{L^2}^2 + \sum_{n=1}^{\infty} \frac{n \phi(t)^{2n}}{(n!)^4} \|\partial_{\alpha}^n W_{\alpha}\|_{L^2}^2 \right) \\ & \leq C(1 + \|b\|_{L^{\infty}} + \|b\|_{\dot{Y}_{\phi(t)}}) E_{L_0, \delta_0}(t) + \|R_{\alpha}\|_{\dot{X}_{\phi(t)}}^2 + d_1 E_{L_0, \delta_0}^{3/2} + \|G\|_{\dot{Y}_{\phi(t)}}^2 + C\|b\|_{\dot{Y}_{\phi(t)}}^2 \\ & \leq C(1 + \|b_1\|_{Y_{\phi(t)}} + E_{L_0, \delta_0}(t)^{1/2}) E_{L_0, \delta_0}(t) + \|R_{\alpha}\|_{\dot{X}_{\phi(t)}}^2 + d_1 E_{L_0, \delta_0}^{3/2} + \|G\|_{\dot{Y}_{\phi(t)}}^2 + C\|b_1\|_{\dot{Y}_{\phi(t)}}^2 + C E_{L_0, \delta_0}(t). \end{aligned}$$

We choose $T_0 = \min\{T, \frac{L_0}{2\delta_0}, \tau_0\}$. With this choice, $\phi(t) \geq \frac{L_0}{2} \geq 2$. Using (4.6), by a bootstrap argument,

$$\frac{\delta_0}{\phi(t)} - 2 - d_0(\|A\|_{L^{\infty}} + \|A\|_{\dot{X}_{\phi(t)}}) - d_1 E_{L_0, \delta_0}^{1/2} \geq 0, \quad \forall t \in [0, T_0].$$

By choosing d_1 larger if necessary, we obtain

$$\frac{d}{dt} E_{L_0, \delta_0}(t) \leq C(1 + \|b_1\|_{Y_{\phi(t)}}) E_{L_0, \delta_0}(t) + d_1 E_{L_0, \delta_0}^{3/2} + \|R_{\alpha}\|_{\dot{X}_{\phi(t)}}^2 + \|G\|_{\dot{Y}_{\phi(t)}}^2 + C\|b_1\|_{\dot{Y}_{\phi(t)}}^2. \quad (4.13)$$

By the method of continuity, we obtain

$$E_{L_0, \delta_0}(t) \leq E_{L_0, \delta_0}(0) e^{B_{T_0} t} + \int_0^t e^{B_{T_0} \tau} \mathcal{N}(\tau) d\tau, \quad (4.14)$$

where $\mathcal{N}$ is defined in (4.10).

To estimate $\|U\|_{L^2}$, using energy estimates, we have

$$\begin{aligned} \frac{d}{dt} \|U(\cdot, t)\|_{L^2}^2 &= 2\langle U_t, U \rangle \\ &= 2\langle -b\partial_{\alpha} U + G + A\Lambda W, U \rangle \\ &\leq 2\|b_{\alpha}\|_{L^{\infty}} \|U\|_{L^2}^2 + \|G\|_{L^2}^2 + \|U\|_{L^2}^2 + \|A\|_{L^{\infty}} \|\Lambda W\|_{L^2}^2 + \|A\|_{L^{\infty}} \|U\|_{L^2}^2 \\ &\leq C(E_{L_0, \delta_0}^{1/2} + \|b_1\|_{\dot{Y}_{\phi(t)}}) \|U\|_{L^2}^2 + \|G\|_{L^2}^2 + \|U\|_{L^2}^2 + \|A\|_{L^{\infty}}^2 E_{L_0, \delta_0}(t) + \|A\|_{L^{\infty}} \|U\|_{L^2}^2. \end{aligned}$$

So we obtain

$$\|U(\cdot, t)\|_{L^2}^2 \leq \|U(\cdot, 0)\|_{L^2}^2 e^{\gamma(t)} + \int_0^t e^{\gamma(t)-\gamma(s)} (\|G(s)\|_{L^2}^2 + \|A(s)\|_{L^{\infty}} E_{L_0, \delta_0}(s)) ds, \quad (4.15)$$

where

$$\gamma(t) := \int_0^t C(E_{L_0, \delta_0}^{1/2}(\tau) + \|b_1(\cdot, \tau)\|_{\dot{Y}_{\phi(\tau)}} + \|A(\cdot, \tau)\|_{L^{\infty}}) d\tau.$$

□

5. ESTIMATES

In §6, we use the Picard iteration to prove *Theorem 3*. For each iteration, we need to solve the quasilinear system (4.1) with G, R, b_1, A constructed in the previous iteration. Hence we need to derive estimates for these quantities. Most of the estimates in this section hold for the general case: without symmetry, and applies to an arbitrary number of point vortices. The only place that the symmetry plays a role is the estimates for the velocity of the point vortices, which is in Lemma 5.6.

5.1. A priori assumptions. We'll derive estimates under the following a priori assumptions. These assumptions are verified during the process of the Picard iteration in the next section.

Without loss of generality, we assume $0 < C_0 < C_1$. Denote

$$\frac{1}{4}m_0^2 := \|U_0\|_{Y_{\phi(0)}}^2 + \|\partial_\alpha W_0\|_{X_{\phi(0)}}^2. \quad (5.1)$$

Let's denote ⁹

$$M_{\lambda,h}^2 := \begin{cases} m_0^2 + C|\lambda|(|y_0| - \frac{|\lambda|}{4\pi x(0)}T)^{-5/2}, & \lambda > 0; \\ m_0^2 + |\lambda|d_{I,0}^{-5/2}, & \lambda < 0. \end{cases} \quad (5.2)$$

$$M_\infty = (d_{I,0} + C|\lambda|(|y_0| - \frac{|\lambda|}{4\pi x(0)}T)^{-5/2})^{-1/2}. \quad (5.3)$$

Here, $C > 0$ is a constant depending on C_0 and $\frac{1}{L_0}$ only¹⁰. We will choose $T = \frac{4\pi x(0)(|y_0| - |y_0|^{9/10})}{|\lambda|}$ if $\lambda > 0$ and $T = O(1)$ if $\lambda < 0$. So we can take

$$M_{\lambda,h}^2 = \begin{cases} m_0^2 + C|y_0|^{-3/4}, & \lambda > 0 \\ m_0^2 + C|y_0|^{-1} & \lambda < 0. \end{cases} \quad (5.4)$$

Without loss of generality, we assume

$$\begin{cases} m_0^2 \leq |y_0|^{-3/4}, & \lambda > 0; \\ m_0^2 \leq |y_0|^{-1}, & \lambda < 0. \end{cases} \quad (5.5)$$

Definition 5.1. Given $W, U, z_j(t)$, we say that $(U, W, \{z_j(t)\})$ satisfies AS, if the following hold:

(AS1) $W \in C([0, T]; X_{\phi(t)})$, $U \in C([0, T]; Y_{\phi(t)})$, $z_j(t) \in C^1([0, T]; \Omega(t))$.

(AS2) $\sup_{0 \leq t \leq T} \left(\|U(\cdot, t)\|_{Y_{\phi(t)}}^2 + \|W_\alpha\|_{X_{\phi(t)}}^2 \right) \leq M_{\lambda,h}^2 \leq 1$, and

$$\sup_{0 \leq t \leq T} \|U(\cdot, t)\|_{L^2}^2 \leq 1, \quad \sup_{0 \leq t \leq T} \|U(\cdot, t)\|_{L^\infty} \leq M_\infty.$$

(AS3) $\frac{1}{2}C_0|\alpha - \beta| \leq |Z(\alpha, t) - Z(\beta, t)|$, $\forall t \in [0, T]$.

(AS4) $d_I(t) \geq \frac{1}{2}d_{I,0}^{9/10} \geq 1$, $x(t) \geq \frac{1}{2}x(0)$, $t \in [0, T]$.

(AS5) $\sup_{t \in [0, T]} \phi(t) \geq \frac{1}{2}L_0$.

We derive a priori estimates under the a priori assumptions (AS1)-(AS5). Note that (AS2) implies

$$\sup_{0 \leq t \leq T} (\|F\|_{Y_{\phi(t)}}^2 + \|Z_\alpha - 1\|_{X_{\phi(t)}}^2) \leq 4M_{\lambda,h}^2. \quad (5.6)$$

⁹In the definition of $M_{\lambda,h}$, we will choose $d_{I,0}$ large. We do not claim that $|d_{I,0}|^{3/2}$ is the optimal choice. The letter h in $M_{\lambda,h}$ refers to *homogeneous*.

¹⁰Since we choose $L_0 \geq 4$, the constant can be chosen such that it depends on C_0 only. At the moment let's keep track of its dependence on $\frac{1}{L_0}$.

Convention: If not specified, in this section, a constant C will depend on $C_0, C_1, M_{\lambda,h}, d_I(t)^{-1}$, and $\frac{1}{L_0}$.

5.2. Estimates.

Lemma 5.1. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS. Then any $k \geq 2$, there holds*

$$\int_{-\infty}^{\infty} \frac{1}{|Z(\beta, t) - z_j(t)|^k} d\beta \leq \frac{8}{C_0} d_I(t)^{-k+1} + \frac{8}{(k-1)C_0} \left(\frac{4C_1}{d_I(t)C_0} \right)^{k-1}. \quad (5.7)$$

In particular, we have

$$\int_{-\infty}^{\infty} \frac{1}{|Z(\beta, t) - z_j(t)|^k} d\beta \leq \frac{16}{C_0} \left(\frac{4C_1}{d_I(t)C_0} \right)^{k-1}. \quad (5.8)$$

Proof. We may assume that $d_I(t) = d(z_j(t), z(0, t))$.

$$\begin{aligned} & \int_{-\infty}^{\infty} \frac{1}{|z_j(t) - Z(\beta, t)|^k} d\beta \\ &= \int_{|Z(0,t) - Z(\beta,t)| \leq 2d_I(t)} \frac{1}{|z_j(t) - Z(\beta, t)|^k} d\beta + \int_{|z(0,t) - Z(\beta,t)| \geq 2d_I(t)} \frac{1}{|z_j(t) - Z(\beta, t)|^k} d\beta \\ &:= I + II. \end{aligned}$$

Denote

$$E := \{\beta : |Z(0, t) - Z(\beta, t)| \leq 2d_I(t)\}.$$

Since

$$\frac{1}{2}C_0|\beta - 0| \leq |Z(\beta, t) - Z(0, t)|,$$

we have for $\beta \in E$,

$$|\beta - 0| \leq \frac{4}{C_0} d_I(t).$$

Therefore

$$I \leq 8C_0^{-1} d_I(t)^{-k+1}.$$

For $\beta \in E^c$, we have

$$|Z(\beta, t) - Z(0, t) - d_I(t)| \geq |Z(\beta, t) - Z(0, t)| - d_I(t) \geq \frac{1}{2}|Z(\beta, t) - Z(0, t)| \geq \frac{C_0}{4}|\beta - 0|. \quad (5.9)$$

Also, we have

$$2C_1|\beta - 0| \geq |Z(\beta, t) - Z(0, t)| \geq 2d_I(t). \quad (5.10)$$

So

$$|\beta| \geq \frac{1}{C_1} d_I(t) \quad (5.11)$$

Therefore, for II , we have

$$II \leq \frac{4^k}{C_0^k} \int_{|\beta| \geq \frac{1}{C_1} d_I(t)} |\beta|^{-k} d\beta = 2 \frac{4^k}{(k-1)C_0^k} C_1^{k-1} d_I(t)^{-k+1} = \frac{8}{(k-1)C_0} \left(\frac{4C_1}{d_I(t)C_0} \right)^{k-1}.$$

□

Lemma 5.2. *Denote $\tilde{Q}_j := \frac{1}{Z(\alpha, t) - z_j(t)}$. Assume that $(W, U, \{z_j(t)\})$ satisfies AS. Then*

$$\|\partial_\alpha \tilde{Q}_j\|_{X_{\phi(t)}} \leq C d_I(t)^{-3/2}, \quad (5.12)$$

for some constant $C > 0$ depending on $M_{\lambda,h}, C_0, C_1$, and $\frac{1}{L_0}$.

Proof. We have $\partial_\alpha \tilde{Q}_j = -\frac{Z_\alpha - 1}{(Z(\alpha, t) - z_j(t))^2} - \frac{1}{(Z(\alpha, t) - z_j(t))^2}$. Since

$$\left\| \frac{Z_\alpha - 1}{(Z(\alpha, t) - z_j(t))^2} \right\|_{X_{\phi(t)}} \leq C \|Z_\alpha - 1\|_{X_{\phi(t)}} \left\| \frac{1}{(Z(\alpha, t) - z_j(t))^2} \right\|_{X_{\phi(t)}},$$

it suffices to prove $\left\| \frac{1}{(Z(\alpha, t) - z_j(t))^2} \right\|_{X_{\phi(t)}} \leq C d_I(t)^{-3/2}$.

Denote $Q_j := \frac{1}{(Z(\alpha, t) - z_j(t))^2}$, and $g(\alpha) := \frac{1}{(\alpha - z_j(t))^2}$. The Faa di Bruno formula implies

$$\partial_\alpha^n Q_j(\alpha, t) = \sum_{k=1}^n \sum_{\Lambda_{n,k}} \frac{n!}{k_1! k_2! \cdots k_n!} g^{(k)}(Z(\alpha, t), t) \prod_{j=1}^n \left(\frac{\partial_\alpha^j Z(\alpha, t)}{j!} \right)^{k_j},$$

where

$$\Lambda_{n,k} := \{(k_1, \dots, k_n) \in (\mathbb{N} \cup \{0\})^n : \sum_{j=1}^n k_j = k, \sum_{j=1}^n j k_j = n\}.$$

Using (AS2) and (AS5), by Sobolev embedding, we estimate Z_α by

$$\begin{aligned} |Z_\alpha(\alpha, t)| &\leq 1 + |Z_\alpha - 1| \leq 1 + \|Z_\alpha - 1\|_{H^1} \\ &\leq 1 + \left(1 + \frac{2}{\phi(t)}\right) 2M_{\lambda,h} \\ &\leq 1 + 6M_{\lambda,h} \\ &:= K_0. \end{aligned}$$

For $2 \leq j \leq n-3$, we estimate $\partial_\alpha^j Z(\alpha, t)$ by

$$\begin{aligned} |\partial_\alpha^j Z(\alpha, t)| &\leq \|\partial_\alpha^j (Z - \alpha)\|_{H^1} \\ &\leq \frac{(j!)^2}{\phi^j(t)} \left(1 + \frac{(j+1)^2}{\phi(t)}\right) \|Z - \alpha\|_{\dot{Y}_{\phi(t)}} \\ &\leq 4 \frac{((j+1)!)^2}{\phi(t)^j} M_{\lambda,h} \leq \frac{K_0((j+1)!)^2}{\phi(t)^j}. \end{aligned} \tag{5.13}$$

So we obtain

$$\begin{aligned} \prod_{j=1}^{n-1} \left(\frac{\partial_\alpha^j Z(\alpha, t)}{j!} \right)^{k_j} &\leq \prod_{j=1}^n \left(\frac{K_0 j!}{\phi(t)^{j-1}} \right)^{k_j} \\ &= K_0^{\sum_{j=1}^n k_j} \phi(t)^{-(\sum_{j=1}^n j k_j - \sum_{j=1}^n k_j)} \prod_{j=1}^n (j!)^{k_j} \\ &= K_0^k \phi(t)^{-n+k} \prod_{j=1}^n (j!)^{k_j}. \end{aligned}$$

For $n-2 \leq j \leq n$, we use

$$\|\partial_\alpha^j Z(\alpha, t)\|_{L^2} = \|\partial_\alpha^{j-1} (Z_\alpha - 1)\|_{L^2} \leq 2 \frac{((j-1)!)^2}{\phi(t)^{j-1}} M_{\lambda,h}. \tag{5.14}$$

For $n \leq 5$, there holds

$$\sum_{n=0}^5 \frac{\phi(t)^{2n}}{(n!)^4} \|\partial_\alpha^n Q_j\|_{L^2}^2 \leq C d_I(t)^{-3/2}, \tag{5.15}$$

for some constant C depending on $C_0, C_1, M_{\lambda,h}$, and $\frac{1}{L_0}$.

Note that for $n \geq 6$ and $k \geq [n/2]$, we must have $j \leq n - 3$. So for $n \geq 6$,

$$\begin{aligned} \|\partial_\alpha^n Q_j(\alpha, t)\|_{L^2} &\leq \sum_{k=3}^{[n/2]} \sum_{\Lambda_{n,k}} \frac{n!k!}{k_1! \cdots k_n!} K_0^k \phi(t)^{-n+k} \prod_{j=1}^n (j!)^{k_j} \left\| \frac{1}{|Z(\alpha, t) - z_j(t)|^{k+1}} \right\|_{L^2} \\ &\quad + \sum_{k=[n/2]+1}^n \sum_{\Lambda_{n,k}} \frac{n!k!}{k_1! \cdots k_n!} K_0^k \phi(t)^{-n+k} \prod_{j=1}^n (j!)^{k_j} \left\| \frac{1}{|Z(\alpha, t) - z_j(t)|^{k+1}} \right\|_{L^2} \\ &\quad + \left\| \sum_{k=1}^2 \sum_{\Lambda_{n,k}} \frac{n!}{k_1! k_2! \cdots k_n!} g^{(k)}(Z(\alpha, t), t) \prod_{j=1}^n \left(\frac{\partial_\alpha^j Z(\alpha, t)}{j!} \right)^{k_j} \right\|_{L^2} \\ &:= II_1 + II_2 + II_3. \end{aligned}$$

Assume that $n \geq 6$ and $k \geq 3$. Let $(k_1, \dots, k_n) \in \Lambda_{n,k}$. Note that

$$\frac{\prod_{j=2}^n (j!)^{k_j}}{n!} = (n!)^{-1} \prod_{\substack{2 \leq j \leq n \\ k_j \neq 0}} (j!)^{k_j} \leq \frac{C}{n^2 \prod_{j=0}^{k-1} (n-j)}, \quad (5.16)$$

for some absolute constant $C > 0$. For example, we can take $C = 100$. Note that $n \geq 6$ and $1 \leq j \leq [n/2]$ imply $n - j \geq \frac{1}{2}n$. Therefore, for $3 \leq k \leq [n/2]$,

$$\prod_{j=0}^{k-1} (n-j) \geq 2^{-k} n^k. \quad (5.17)$$

For $k \geq [n/2] + 1$, we have

$$\prod_{j=0}^{k-1} (n-j) \geq 2^{-[n/2]} n^{[n/2]}. \quad (5.18)$$

Lemma 5.1 and (5.17) imply

$$\begin{aligned} II_1 &\leq C \frac{(n!)^2}{n^2 \phi(t)^n} \sum_{k=3}^{[n/2]} \sum_{\Lambda_{n,k}} \frac{k!}{k_1! \cdots k_n!} \left(\frac{4K_0 C_1 \phi(t)}{nC_0 d_I(t)} \right)^k \\ &\leq C \frac{(n!)^2}{n^2 \phi(t)^n} \sum_{k=1}^n \sum_{\Lambda_{n,k}} \frac{k!}{k_1! \cdots k_n!} \left(\frac{4K_0 C_1 \phi(t)}{nC_0 d_I(t)} \right)^k \\ &= C \frac{(n!)^2}{n^2 \phi(t)^n} \frac{4K_0 C_1 \phi(t)}{nC_0 d_I(t)} \left(1 + \frac{4K_0 C_1 \phi(t)}{nC_0 d_I(t)} \right)^n \\ &\leq C d_I(t)^{-1} \frac{(n!)^2}{n^2 \phi(t)^n}, \end{aligned}$$

for some constant $C > 0$ depending on $C_0, C_1, M_{\lambda, h}, \frac{1}{L_0}$. Here we've used the identity

$$\sum_{k=1}^n \sum_{\Lambda_{n,k}} \frac{k!}{(k_1)! \cdots (k_n)!} R^k = R(1+R)^{n-1}, \quad (5.19)$$

and the estimate

$$\sup_{n \geq 1} \left(1 + \frac{4K_0 C_1 \phi(t)}{nC_0 d_I(t)} \right)^n \leq e^{\frac{4K_0 C_1 \phi(t)}{C_0 d_I(t)}} \leq e^{\frac{4K_0 C_1 L_0}{C_0 d_I(t)}}. \quad (5.20)$$

So we have

$$\sum_{n=0}^{\infty} \frac{(1+n^2)(\phi(t))^{2n}}{(n!)^4} II_1^2 \leq C d_I(t)^{-1}, \quad (5.21)$$

for some constant $C > 0$ depending on $K_0, C_0, C_1, M_{\lambda, h}, \frac{1}{L_0}$.

For II_2 , using Lemma 5.1 and (5.18), we obtain

$$\begin{aligned} II_2 &\leq C \frac{(n!)^2}{\phi(t)^n} \sum_{k=[n/2]+1}^n \sum_{\Lambda_{n,k}} \frac{k!}{k_1! \cdots k_n!} \frac{(\frac{2K_0 C_1 \phi(t)}{C_0 d_I(t)})^k}{\prod_{j=0}^{k-1} (n-j)} \\ &\leq C \frac{(n!)^2}{\phi(t)^n} \sum_{k=[n/2]+1}^n \sum_{\Lambda_{n,k}} \frac{k!}{k_1! \cdots k_n!} \frac{(\frac{2K_0 C_1 \phi(t)}{C_0 d_I(t)})^k}{(n/2)^{n/2}} \\ &\leq C d_I(t)^{-1} \frac{(n!)^2}{\phi(t)^n} \frac{(1 + \frac{2K_0 C_1 \phi(t)}{C_0 d_I(t)})^n}{(n/2)^{n/2}} \\ &\leq C d_I(t)^{-1} \frac{(n!)^2}{n^2 \phi(t)^n}, \end{aligned}$$

for some constant $C > 0$ depending on $K_0, C_0, C_1, M_{\lambda, h}, \frac{1}{L_0}$. Here, we've used the identity

$$\sum_{k=1}^n \sum_{\Lambda_{n,k}} \frac{k!}{(k_1)! \cdots (k_n)!} R^k = R(1+R)^{n-1}, \quad (5.22)$$

and the estimate

$$\sup_{n \geq 1} \frac{1}{n^{[n/2]-2}} \left(1 + \frac{2K_0 C_1 \phi(t)}{C_0 d_I(t)} \right)^n \leq C, \quad (5.23)$$

for some constant $C > 0$ depending on $K_0, C_0, C_1, M_{\lambda, h}$ and $\frac{1}{L_0}$.

So we have

$$\sum_{n=1}^{\infty} \frac{(\phi(t))^{2n}}{(n!)^4} II_2^2 \leq C d_I(t)^{-1} \sum_{n=0}^{\infty} \frac{(\phi(t))^{2n}}{(n!)^4} \frac{(n!)^4}{\phi(t)^{2n}} \frac{1}{n^4} \leq C d_I(t)^{-3/2}. \quad (5.24)$$

For II_3 , when $k = 1$, we must have $\Lambda_{n,1} = (0, \dots, 0, 1)$. In this case,

$$\begin{aligned} &\sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \left\| \sum_{\Lambda_{n,1}} \frac{n!}{k_1! k_2! \cdots k_n!} g^{(k)}(Z(\alpha, t), t) \prod_{j=1}^n \left(\frac{\partial_{\alpha}^j Z(\alpha, t)}{j!} \right)^{k_j} \right\|_{L^2}^2 \\ &= \sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \left\| \frac{\partial_{\alpha}^{n-1}(Z_{\alpha} - 1)}{|Z(\alpha, t) - z_j(t)|^2} \right\|_{L^2}^2 \\ &\leq \sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} \frac{1}{d_I(t)^2} \frac{((n-1)!)^4}{\phi(t)^{2(n-1)}} \left(\frac{\phi(t)^{2(n-1)}}{((n-1)!)^4} \|\partial_{\alpha}^{n-1}(Z_{\alpha} - 1)\|_{L^2}^2 \right) \\ &\leq C d_I(t)^{-2}. \end{aligned}$$

for some constant $C > 0$ depending on $C_0, C_1, M_{\lambda, h}$, and $\frac{1}{L_0}$.

The same argument applies to $k = 2$. We obtain

$$\sum_{n=0}^{\infty} \frac{\phi(t)^{2n}}{(n!)^4} II_3^2 \leq C d_I(t)^{-2}. \quad (5.25)$$

This concludes the proof of the lemma. $\square$

Similarly, we can prove

Lemma 5.3. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

$$\left\| \frac{1}{(Z(\alpha, t) - z_j(t))^m} \right\|_{Y_{\phi(t)}} \leq C d_I(t)^{-m+1/2}, \quad (5.26)$$

$$\left\| \frac{1}{Z(\alpha, t) - z_1(t)} - \frac{1}{Z(\alpha, t) - z_2(t)} \right\|_{Y_{\phi(t)}} \leq C \frac{|z_1(t) - z_2(t)|}{d_I(t)^{3/2}}, \quad (5.27)$$

$$\left\| \frac{1}{Z(\alpha, t) - z_1(t)} - \frac{1}{Z(\alpha, t) - z_2(t)} \right\|_{\dot{Y}_{\phi(t)}} \leq C \frac{|z_1(t) - z_2(t)|}{d_I(t)^{5/2}}, \quad (5.28)$$

for some constant $C > 0$ depending on $C_0, C_1, M_{\lambda, h}$, and $\frac{1}{L_0}$.

As a consequence,

Corollary 5.1. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS. Let $g \in X_{\phi(t)}$, then ¹¹*

1. *For arbitrary N , let $\lambda_0 := N \max_{1 \leq j \leq N} |\lambda_j|$, we have*

$$\left\| \sum_{j=1}^N \frac{\lambda_j g}{(Z(\alpha, t) - z_j(t))^m} \right\|_{X_{\phi(t)}} \leq C \lambda_0 d_I(t)^{-m} \|g\|_{X_{\phi(t)}}, \quad (5.29)$$

2. *Assume $N = 2$ and $\lambda_1 = -\lambda_2 = \lambda$, then $Q := \frac{\lambda i}{2\pi} \frac{1}{Z(\alpha, t) - z_1(t)} - \frac{\lambda i}{2\pi} \frac{1}{Z(\alpha, t) - z_2(t)}$ satisfies*

$$\|Q\|_{Y_{\phi(t)}} \leq C |\lambda| |z_1(t) - z_2(t)| d_I(t)^{-3/2}, \quad (5.30)$$

$$\|Q\|_{\dot{Y}_{\phi(t)}} \leq C |\lambda| |z_1(t) - z_2(t)| d_I(t)^{-5/2}, \quad (5.31)$$

$$\|gQ\|_{X_{\phi(t)}} \leq C |\lambda| |z_1(t) - z_2(t)| d_I(t)^{-2} \|g\|_{X_{\phi(t)}}, \quad (5.32)$$

$$\|(I + \mathbb{H})Q\|_{Y_{\phi(t)}} \leq C |\lambda| |z_1(t) - z_2(t)| d_I(t)^{-5/2}, \quad (5.33)$$

for some constant $C > 0$ depending on $C_0, C_1, M_{\lambda, h}$, and $\frac{1}{L_0}$.

Proof. For (5.29), we estimate $\left\| \sum_{j=1}^N \frac{\lambda_j g}{(Z(\alpha, t) - z_j(t))^m} \right\|_{L^2}$ by

$$\left\| \sum_{j=1}^N \frac{\lambda_j g}{(Z(\alpha, t) - z_j(t))^m} \right\|_{L^2} \leq C \|g\|_{L^2} \left\| \sum_{j=1}^N \frac{\lambda_j g}{(Z(\alpha, t) - z_j(t))^m} \right\|_{L^\infty} \leq C d_I(t)^{-m} \|g\|_{X_{\phi(t)}}.$$

By Lemma 2.5 and Lemma 5.3,

$$\left\| \sum_{j=1}^N \frac{\lambda_j g}{(Z(\alpha, t) - z_j(t))^m} \right\|_{\dot{X}_{\phi(t)}} \leq C \lambda_0 d_I(t)^{-m-1/2} \|g\|_{X_{\phi(t)}}.$$

So

$$\left\| \sum_{j=1}^N \frac{\lambda_j g}{(Z(\alpha, t) - z_j(t))^m} \right\|_{X_{\phi(t)}} \leq C \lambda_0 d_I(t)^{-m-1/2} \|g\|_{X_{\phi(t)}} + C d_I(t)^{-m} \|g\|_{X_{\phi(t)}} \leq C d_I(t)^{-m} \|g\|_{X_{\phi(t)}}.$$

Here, we use the assumption that $d_I(t) \geq 1$. So we obtain (5.29). (5.30)-(5.32) follow from (5.29) and Lemma 5.3. $\square$

Using the same proof as for Lemma 5.2, we obtain the following.

¹¹We will construct solutions with $d_I(t)$ large. Hence larger k gives smaller quantity $d_I(t)^{-k}$.

Lemma 5.4. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS. Then*

$$\left\| \frac{1}{Z_\alpha} - 1 \right\|_{X_{\phi(t)}} \leq CM_{\lambda,h}, \quad (5.34)$$

for some constant $C > 0$ depending on C_0 and $M_{\lambda,h}$.

Lemma 5.5. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

$$|\mathcal{U}(z, t)| \leq C \min\{M_\infty, (1 + d(z, \Sigma(t)))^{-1/2}\}, \quad (5.35)$$

$$|\mathcal{U}_z(z, t)| \leq C \min\{M_{\lambda,h}, (1 + d(z, \Sigma(t)))^{-3/2}\}, \quad (5.36)$$

for some constant $C > 0$ depending on C_0, C_1 . Here, $d(z, \Sigma(t)) := \inf_{\beta \in \mathbb{R}} |z - Z(\beta, t)|$.

Proof. Recall that $\mathcal{U}$ is holomorphic in $\Omega(t)$ with boundary value F . So we have for $z \in \Omega(t)$,

$$\mathcal{U}(z, t) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{Z_\beta(\beta, t)}{z - Z(\beta, t)} F(\beta, t) d\beta.$$

If $d(z, \Sigma(t)) \leq 1$, then by (AS2)

$$|\mathcal{U}(z, t)| \leq \|\mathcal{U}(\cdot, t)\|_{L^\infty(\Omega(t))} \leq M_\infty.$$

If $d(z, \Sigma(t)) \geq 1$, we have

$$|\mathcal{U}(z, t)| \leq \frac{1}{2\pi} \left(\int_{-\infty}^{\infty} \frac{1}{|z - Z(\beta, t)|^2} d\beta \right)^{1/2} \|Z_\alpha\|_{L^\infty} \|F\|_{L^2} \leq CM_{\lambda,h} z^{-1/2}.$$

So we obtain (5.35). Notice that $\mathcal{U}_z$ has boundary values $\frac{F_\alpha}{Z_\alpha}$. Using the same proof as for (5.35), we obtain (5.36). $\square$

Lemma 5.6. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

$$|\dot{z}_j(t)| \leq C(d_I(t)^{-1/2} + \frac{|\lambda|}{x(t)}), \quad j = 1, 2. \quad (5.37)$$

$$|\Re\{\dot{z}_j(t)\}| \leq Cd_I(t)^{-3/2}. \quad (5.38)$$

$$|\dot{z}_1(t) - \dot{z}_2(t)| \leq Cd_I(t)^{-3/2} x(t), \quad (5.39)$$

Proof. The main tool is the maximum principle for holomorphic functions. Recall that for $j = 1, 2$,

$$\dot{z}_j(t) = \sum_{1 \leq k \leq 2, k \neq j} \frac{\lambda_k i}{2\pi(\overline{z_j(t)} - \overline{z_k(t)})} + \bar{\mathcal{U}}(z_j(t), t),$$

where $\mathcal{U}(Z, t) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{Z_\beta(\beta, t)}{Z - Z(\beta, t)} F(\beta, t) d\beta$ is holomorphic in $\Omega(t)$ with boundary value F on $\Sigma(t)$.

Estimate $\dot{z}_j$: Clearly,

$$\frac{\lambda_2}{2\pi(\overline{z_1(t)} - \overline{z_2(t)})} \leq \frac{|\lambda|}{4\pi x(t)}.$$

By (5.35) and the assumption that $d_I(t) \geq 1$, we have

$$|\bar{\mathcal{U}}(z_j(t), t)| \leq C(1 + d(z_j(t), \Sigma(t)))^{-1/2} = Cd_I(t)^{-1/2}.$$

So we obtain

$$|\dot{z}_j(t)| \leq C(d_I(t)^{-1/2} + \frac{|\lambda|}{x(t)}). \quad (5.40)$$

Estimate $|\dot{z}_1(t) - \dot{z}_2(t)|$: By (5.36) of Lemma 5.5 we have

$$|\dot{z}_1(t) - \dot{z}_2(t)| = |\mathcal{U}(z_1(t), t) - \mathcal{U}(z_2(t), t)| \leq \|\mathcal{U}_z\|_{L^\infty(\Omega(t))} |z_1(t) - z_2(t)| \leq C d_I(t)^{-3/2} x(t).$$

(5.38) follows immediately from

$$\Re\{\dot{z}_j(t)\} = \Re\{\bar{\mathcal{U}}(z_j(t), t)\} \leq |\mathcal{U}_z(\tilde{x}(t) + iy(t), t)| x(t) \leq C d_I(t)^{-3/2}.$$

□

The following lemma follows by direct calculation.

Lemma 5.7. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then for $t \in [0, T]$,*

$$|\frac{d}{dt}x(t)| \leq C d_I(t)^{-3/2} x(t), \quad (5.41)$$

1. *If $\lambda < 0$, then*

$$\frac{d}{dt}y(t) \leq -\frac{|\lambda|}{8\pi x(0)}, \quad (5.42)$$

2. *If $\lambda > 0$, then*

$$\frac{d}{dt}y(t) \geq \frac{|\lambda|}{8\pi x(0)}, \quad (5.43)$$

Lemma 5.8. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

$$\|A_1 - 1\|_{X_{\phi(t)}} \leq C(1 + |\lambda|^2 d_I(t)^{-5/2}). \quad (5.44)$$

$$\|A_1 - 1\|_{\dot{X}_{\phi(t)}} \leq C(M_{\lambda,h}^2 + |\lambda|^2 d_I(t)^{-7/2}), \quad (5.45)$$

$$\|A_1 - 1\|_{L^\infty} \leq C(1 + \lambda^2 d_I(t)^{-3}). \quad (5.46)$$

Proof. We prove (5.44) only. We have

$$A_1 = 1 + \frac{1}{2\pi} \int \frac{|D_t Z(\alpha, t) - D_t Z(\beta, t)|^2}{(\alpha - \beta)^2} d\beta - \sum_{j=1}^2 \frac{\lambda_j}{2\pi} \operatorname{Re} \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) (D_t Z - \dot{z}_j(t)) \right\}.$$

Splitting $D_t Z = \bar{F} + \bar{Q}$. Using (AS2), (5.35), Lemma 2.5, Lemma 2.7 and (5.30), we have

$$\begin{aligned} \left\| \int \frac{|D_t Z(\alpha, t) - D_t Z(\beta, t)|^2}{(\alpha - \beta)^2} d\beta \right\|_{X_{\phi(t)}} &\leq \left\| \int \frac{|F(\alpha, t) - F(\beta, t)|^2}{(\alpha - \beta)^2} d\beta \right\|_{X_{\phi(t)}} + \left\| \int \frac{|Q(\alpha, t) - Q(\beta, t)|^2}{(\alpha - \beta)^2} d\beta \right\|_{X_{\phi(t)}} \\ &\leq C(\|F\|_{X_{\phi(t)}}^2 + \|Q\|_{X_{\phi(t)}}^2) \\ &\leq C(1 + M_{\lambda,h}^2 + |\lambda|^2 x(t)^2 d_I(t)^{-3}). \end{aligned}$$

Similarly, we obtain

$$\left\| \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) D_t Z \right\} \right\|_{X_{\phi(t)}} \leq C(1 + M_{\lambda,h}^2 + |\lambda|^2 x(t)^2 d_I(t)^{-3}).$$

For $\left\| \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) \dot{z}_j(t) \right\} \right\|_{X_{\phi(t)}}$, we use $\dot{z}_j(t) = \frac{\lambda_j i}{4\pi x(t)} + \bar{\mathcal{U}}(z_j(t), t)$. We have

$$\begin{aligned} &\sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) \frac{\lambda_j i}{4\pi x(t)} \right\} \\ &= -\frac{\lambda^2}{4\pi^2} \Re \left\{ (I - \mathbb{H}) \left(\frac{Z_\alpha}{(Z(\alpha, t) - z_1(t))^2 (Z(\alpha, t) - z_2(t))} + \frac{Z_\alpha}{(Z(\alpha, t) - z_1(t))^2 (Z(\alpha, t) - z_2(t))} \right) \right\} \end{aligned}$$

Using Lemma 5.3 (and similar proofs, if necessary), we obtain

$$\left\| \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) \frac{\lambda i}{4\pi x(t)} \right\} \right\|_{X_{\phi(t)}} \leq C |\lambda|^2 d_I(t)^{-5/2},$$

$$\left\| \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) \frac{\lambda i}{4\pi x(t)} \right\} \right\|_{L^\infty} \leq C |\lambda|^2 d_I(t)^{-3},$$

and

$$\left\| \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) \frac{\lambda i}{4\pi x(t)} \right\} \right\|_{\dot{X}_{\phi(t)}} \leq C |\lambda|^2 d_I(t)^{-7/2},$$

Using (5.35), we obtain

$$\left\| \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \Re \left\{ \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) \bar{U}(z_j(t), t) \right\} \right\|_{X_{\phi(t)}} \leq C |\lambda| d_I(t)^{-2}.$$

So we conclude the proof of the lemma. $\square$

Since $A - 1 = \frac{A_1}{|Z_\alpha|^2} - 1 = \frac{A_1 - 1}{|Z_\alpha|^2} - \frac{|Z_\alpha|^2 - 1}{|Z_\alpha|^2}$, a direct consequence of Lemma 5.8 and Lemma 2.8 yields the following estimates for $A - 1$.

Corollary 5.2. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

$$\|A - 1\|_{L^\infty} \leq C(1 + M_{\lambda, h}^2 + |\lambda|^2 d_I(t)^{-3}). \quad (5.47)$$

$$\|A - 1\|_{X_{\phi(t)}} \leq C(1 + M_{\lambda, h}^2 + |\lambda|^2 d_I(t)^{-5/2}). \quad (5.48)$$

$$\|A - 1\|_{\dot{X}_{\phi(t)}} \leq C(M_{\lambda, h}^2 + |\lambda|^2 d_I(t)^{-4}). \quad (5.49)$$

In particular, if $m_0 \leq C|y_0|^{-1/2}$, then

$$\sup_{0 \leq t \leq T} \|A - 1\|_{L^\infty} \leq \begin{cases} C(1 + |y_0|^{\frac{3}{10}}), & \lambda > 0, \\ C, & \lambda < 0. \end{cases} \quad (5.50)$$

and

$$\sup_{0 \leq t \leq T} \|A - 1\|_{\dot{X}_{\phi(t)}} \leq \begin{cases} C|y_0|^{-3/8}, & \lambda > 0, \\ C|y_0|^{-1/2}, & \lambda < 0. \end{cases} \quad (5.51)$$

Note that $\|A - 1\|_{L^2} \leq C|d_I(t)^{3/4}$, which is significantly larger than $\|A - 1\|_{L^\infty}$.

Lemma 5.9. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

$$\|D_t Q\|_{Y_{\phi(t)}} \leq C \frac{|\lambda|}{d_I(t)^2} + C \frac{\lambda^2}{d_I(t)^{5/2}}, \quad (5.52)$$

$$\|D_t Q\|_{\dot{Y}_{\phi(t)}} \leq C M_{\lambda, h} \frac{|\lambda|}{d_I(t)^2} + C \frac{\lambda^2}{d_I(t)^{7/2}}, \quad (5.53)$$

$$\|D_t Q\|_{L^\infty} \leq C M_{\lambda, \infty} \frac{|\lambda|}{d_I(t)^2} + C \frac{\lambda^2}{d_I(t)^3}, \quad (5.54)$$

for some constant $C > 0$ depending on $C_0, C_1, M_{\lambda, h}, \frac{1}{L_0}$.

Proof. To prove (5.52), note that

$$D_t Q = \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \frac{D_t Z - \dot{z}_j(t)}{(Z(\alpha, t) - z_j(t))^2}. \quad (5.55)$$

Decomposing $D_t Z = \bar{F} + \bar{Q}$, we estimate $D_t Q$ by

$$\begin{aligned} \|D_t Q\|_{Y_{\phi(t)}} &\leq \sum_{j=1}^2 \frac{|\lambda_j|}{2\pi} \left\{ \left\| \frac{\bar{F}}{(Z(\alpha, t) - z_j(t))^2} \right\|_{Y_{\phi(t)}} + \left\| \frac{\bar{Q}}{(Z(\alpha, t) - z_j(t))^2} \right\|_{Y_{\phi(t)}} \right\} + \left\| \sum_{j=1}^2 \frac{\lambda_j}{2\pi} \frac{\dot{z}_j(t)}{(Z(\alpha, t) - z_j(t))^2} \right\|_{Y_{\phi(t)}} \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

Note that Lemma 5.3, Lemma 5.6 and (AS2) imply that

$$I_1 + I_2 \leq C \frac{|\lambda|}{d_I(t)^2} + C \frac{\lambda^2}{d_I(t)^{7/2}}.$$

For I_3 , using $\dot{z}_j(t) = \frac{\lambda i}{4\pi x(t)} + \bar{U}(z_j(t), t)$, we obtain

$$\begin{aligned} I_3 &\leq \left\| \sum_{j=1}^2 \frac{\lambda_j}{2\pi} \frac{\frac{\lambda i}{4\pi x(t)}}{(Z(\alpha, t) - z_j(t))^2} \right\|_{Y_{\phi(t)}} + \left\| \sum_{j=1}^2 \frac{\lambda_j}{2\pi} \frac{\bar{U}(z_j(t), t)}{(Z(\alpha, t) - z_j(t))^2} \right\|_{Y_{\phi(t)}} \\ &:= I_{31} + I_{32}. \end{aligned}$$

By exploring the cancellations, we have

$$\begin{aligned} I_{31} &= \frac{\lambda^2}{4\pi^2} \left\| \frac{1}{(Z(\alpha, t) - z_1(t))^2 (Z(\alpha, t) - z_2(t))} + \frac{1}{(Z(\alpha, t) - z_1(t))^2 (Z(\alpha, t) - z_2(t))} \right\|_{Y_{\phi(t)}} \\ &\leq C \lambda^2 d_I(t)^{-5/2}, \end{aligned}$$

and

$$I_{32} \leq C |\lambda|^2 d_I(t)^{-3}. \quad (5.56)$$

We obtain (5.52) by combining the estimates for I_1, I_2, I_{31}, I_{32} . Similarly,

$$\begin{aligned} \|D_t Q\|_{L^\infty} &\leq \sum_{j=1}^2 \frac{|\lambda_j|}{2\pi} \left\{ \left\| \frac{\bar{F}}{(Z(\alpha, t) - z_j(t))^2} \right\|_{L^\infty} + \left\| \frac{\bar{Q}}{(Z(\alpha, t) - z_j(t))^2} \right\|_{L^\infty} \right\} + \left\| \sum_{j=1}^2 \frac{\lambda_j}{2\pi} \frac{\dot{z}_j(t)}{(Z(\alpha, t) - z_j(t))^2} \right\|_{L^\infty} \\ &\leq C M_{\lambda, \infty} \frac{|\lambda|}{d_I(t)^2} + C \frac{\lambda^2}{d_I(t)^3}. \end{aligned}$$

To prove (5.53), it suffices to notice that

$$\left\| \sum_{j=1}^2 \frac{\lambda_j}{2\pi} \frac{\dot{z}_j(t)}{(Z(\alpha, t) - z_j(t))^2} \right\|_{\dot{Y}_{\phi(t)}} \leq C \frac{\lambda^2}{d_I(t)^{7/2}}.$$

So we obtain (5.53). $\square$

Lemma 5.10. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

$$\|b_1\|_{Y_{\phi(t)}} \leq C(1 + M_{\lambda, h}) |\lambda| d_I(t)^{-5/2}, \quad (5.57)$$

$$\|b_1\|_{\dot{Y}_{\phi(t)}} \leq C M_{\lambda, h} |\lambda| d_I(t)^{-5/2}, \quad (5.58)$$

$$\|b_1\|_{L^\infty} \leq C(1 + M_{\lambda, \infty}) |\lambda| d_I(t)^{-3}, \quad (5.59)$$

for some constant $C > 0$ depending on $C_0, C_1, M_{\lambda, h}$, and $\frac{1}{L_0}$.

Proof. Recall that

$$b_1 = \Re\{[\bar{Q}, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\} + 2\Re\{(I - \mathbb{H})\bar{Q}\}. \quad (5.60)$$

We prove (5.57) only. The proof of (5.58) is similar. By Lemma 2.7, Corollary 5.1, Lemma 5.4, we have

$$\begin{aligned} \|b_1\|_{Y_{\phi(t)}} &= \left\| \Re\{[\bar{Q}, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\} + 2\Re\{(I - \mathbb{H})\bar{Q}\} \right\|_{Y_{\phi(t)}} \\ &\leq \left\| [\bar{Q}, \mathbb{H}](\frac{1}{Z_\alpha} - 1) \right\|_{Y_{\phi(t)}} + 2\|(I + \mathbb{H})Q\|_{Y_{\phi(t)}} \\ &\leq C\|Q\|_{\dot{Y}_{\phi(t)}} \left\| \frac{1}{Z_\alpha} - 1 \right\|_{X_{\phi(t)}} + 2\|(I + \mathbb{H})Q\|_{Y_{\phi(t)}} \\ &\leq C(1 + M_{\lambda,h})|\lambda|d_I(t)^{-5/2}, \end{aligned}$$

for some constant $C > 0$ depending on $C_0, C_1, M_{\lambda,h}, \frac{1}{L_0}$. $\square$

The following two lemmas are consequence of the previous estimates.

Lemma 5.11. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS, then*

(a) $G = G(W, U, \{z_j\}) \in C([0, T]; Y_{\phi(t)})$, and

$$\|G(W, U, \{z_j\})\|_{\dot{Y}_{\phi(t)}} \leq CM_{\lambda,h} \frac{|\lambda|}{d_I(t)^2} + C \frac{\lambda^2}{d_I(t)^{7/2}}, \quad (5.61)$$

$$\|G(W, U, \{z_j\})\|_{Y_{\phi(t)}} \leq C \frac{|\lambda|}{d_{I,t}^2} + C \frac{\lambda^2}{d_I(t)^{5/2}}, \quad (5.62)$$

(b) $R = R(W, U, \{z_j\}) \in C([0, T]; X_{\phi(t)})$, and

$$\|\partial_\alpha R(W, U, \{z_j\})\|_{X_{\phi(t)}} \leq C(1 + M_{\lambda,h})|\lambda|d_I(t)^{-5/2}. \quad (5.63)$$

Proof. Recall that $G = -\Re\{D_t Q\}$. Invoking Lemma 5.9, we obtain

$$\|G\|_{Y_{\phi(t)}} \leq C \frac{|\lambda|}{d_I(t)^2} + C \frac{\lambda^2}{d_I(t)^{7/2}}.$$

Recall that $R = \Re\{Q\} - 2\Re\{(I - \mathbb{H})\bar{Q}\} - \Re\{[\bar{Q}, \mathbb{H}](\frac{1}{Z_\alpha} - 1)\}$. Then (5.63) follows immediately from Lemma 2.7, Lemma 5.10 and Corollary 5.3. $\square$

Lemma 5.12. *Assume that $(W, U, \{z_j(t)\})$ satisfies AS.*

1. *If $\lambda < 0$, then*

$$d_I(t) \geq d_{I,0} + \frac{|\lambda|}{8\pi}t. \quad (5.64)$$

2. *If $\lambda > 0$, then*

$$d_I(t) \geq d_{I,0} - |\lambda|t. \quad (5.65)$$

Proof. If $\lambda < 0$, then we have

$$\begin{aligned} \Im\{Z(\alpha, t) - z_j(t)\} &= -|\Im\{Z(\alpha, 0) - z_j(0)\}| - \int_0^t |\Im\{\frac{d}{d\tau}(Z(\alpha, \tau) - z_j(\tau))\}|d\tau \\ &\leq -|\Im\{Z(\alpha, 0) - z_j(0)\}| - \frac{|\lambda|}{8\pi x(0)}t. \end{aligned}$$

Therefore ¹²

$$d_I(t) \geq |\inf_{\alpha \in \mathbb{R}} \Im\{Z(\alpha, t) - z_1(t)\}| \geq d_{I,0} + \frac{|\lambda|}{2\pi x(0)}$$

If $\lambda > 0$, then we have

$$\begin{aligned} \Im\{Z(\alpha, t) - z_j(t)\} &= |\Im\{Z(\alpha, 0) - z_j(0)\}| - \int_0^t |\Im\{\frac{d}{d\tau}(Z(\alpha, \tau) - z_j(\tau))\}| d\tau \\ &\geq d_{I,0} - \frac{|\lambda|}{x(0)}t. \end{aligned}$$

Then we have

$$d_I(t) \geq d_{I,0} - \frac{|\lambda|}{x(0)}t.$$

Recall that we assume $x(0) = 1$. So we conclude the proof of the lemma. $\square$

6. PROOF OF THEOREM 3

6.1. The wellposedness of the quasilinear system (2.29). In this subsection we prove Theorem 3. Without loss of generality, we assume $d_{I,0} \geq 4$.

The proof of Theorem 3. Let $T > 0$ to be determined, define

$$S_T = \left\{ (W_\alpha, U, \{z_j\}) \left| \begin{array}{l} \text{(AS1)-(AS5) hold for } (W, U, \{z_j\}) \end{array} \right. \right\}.$$

So

$$S_T \subset C([0, T]; X_{\phi(t)}) \times C([0, T]; Y_{\phi(t)}) \times \{C^1([0, T]; \mathbb{C})\}.$$

We denote $D_t^{(n)}$ by $\partial_t + b^n \partial_\alpha$, where b^n is the n -th approximation of b , which will be constructed shortly.

The zero-th approximation. We take $U^0 = \Re\{F_0\}$, $\{z_j^0\} = \{z_{j,0}\}$, $Z^0 = Z_0$, $W^0 := \Re\{Z_0 - \alpha\}$, $G^0 := G(W^0, U^0, \{z_j^0\})$, $R^0 := R(W^0, U^0, \{z_j^0\})$, $b_1^0 := b_1(W^0, U^0, \{z_j^0\})$, $A^0 := A(W^0, U^0, \{z_j^0\})$. For arbitrary $T > 0$,

$$(W^0, U^0, \{z_j^0\}) \in S_T.$$

The n -th approximation. Assume we have constructed $(W^n, U^n, \{z_j^n\})$ such that

$$(W^n, U^n, \{z_j^n\}) \in S_T.$$

Define Z^n and F^n by

$$Z^n(\alpha, t) = \alpha + (I + \mathbb{H})W^n, \quad F^n = (I + \mathbb{H})U^n. \quad (6.1)$$

¹²Recall that we assume $d_{I,0} = \inf_{\alpha \in \mathbb{R}} \min_{j=1,2} |\Im\{Z(\alpha, 0) - z_j(0)\}|$.

The $(n+1)$ -th approximation. Let's construct $(W^{n+1}, U^{n+1}, \{z_j^{n+1}\})$ as follows.

Step 1. Define

$$\begin{cases} G^n := G(W^n, U^n, \{z_j^n(t)\}), \\ R^n := R(W^n, U^n, \{z_j^n(t)\}), \\ b_1^n := b_1(W^n, U^n, \{z_j^n(t)\}), \\ b_0^n := \Re\{\bar{F}^{n+1}, \mathbb{H}\}(\frac{1}{Z_\alpha^{n+1}} - 1)\} + 2F^{n+1}, \\ b^n = b_1^n + b_0^n, \\ A^n = A(W^n, U^n, \{z_j^n(t)\}), \\ D_t^{(n)} := \partial_t + b^n \partial_\alpha, \\ Q^n := -\sum_{j=1}^2 \frac{\lambda_{ji}}{2\pi} \frac{1}{Z^n(\alpha, t) - z_j^n(t)}, \\ d_{I,n}(t) := \inf_{\alpha \in \mathbb{R}} \min_{j=1,2} |Z^n(\alpha, t) - z_j(t)|, \end{cases} \quad (6.2)$$

and define $\mathcal{U}^n$ by

$$\mathcal{U}^n(Z, t) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\partial_\beta Z^n(\beta, t)}{Z - Z^n(\beta, t)} F^n(\beta, t) d\beta.$$

Let $\Sigma^n(t)$ be the curve parametrized by $Z^n(\alpha, t)$, and $\Omega(t)^n$ the region bounded above by Z^n . Note that b_0^n depends on the unknowns F^{n+1} and Z^{n+1} .

Step 2. (U^{n+1}, W^{n+1}) is defined as the solution of

$$\begin{cases} D_t^{(n)} U^{n+1} = A \Lambda W^{n+1} + G^n, \\ D_t^{(n)} W^{n+1} = -U^{n+1} - \Re[\bar{F}^{n+1}, \mathbb{H}](\frac{1}{\partial_\alpha Z^n} - 1)) + R^n, \\ F^{n+1} = (I + \mathbb{H})U^{n+1}, \\ W^{n+1}(\cdot, 0) = W_0, \quad U^{n+1}(\cdot, 0) = U_0. \end{cases} \quad (6.3)$$

Define Z^{n+1} and $\{z_j^{n+1}\}$ by

$$Z^{n+1}(\alpha, t) := \alpha + (I + \mathbb{H})W^{n+1}, \quad (6.4)$$

$$\begin{cases} \frac{d}{dt} z_j^{n+1}(t) = \overline{\mathcal{U}^n(z_j^n(t), t)} + \sum_{\substack{1 \leq k \leq 2 \\ k \neq j}} \frac{\lambda_{ki}}{2\pi} \frac{1}{\overline{z_k^n(t) - z_j^n(t)}}, \\ z_j^{n+1}(0) = z_j(0). \end{cases} \quad (6.5)$$

We show that $(W^{n+1}, U^{n+1}, \{z_j^{n+1}\})$ satisfies (AS1)-(AS5).

We choose T such that

$$T = \begin{cases} \min\{T_1(C_0, \|(U_0, \partial_\alpha W_0)\|_{\dot{Y}_{L_0} \times X_{L_0}}), \frac{L_0}{2\delta_0}, \tau_0, \frac{4\pi x(0)(|y_0| - |y_0|^{9/10})}{|\lambda|}\}, & \lambda > 0 \\ \min\{T_2(C_0, \|(U_0, \partial_\alpha W_0)\|_{\dot{Y}_{L_0} \times X_{L_0}}), \tau_0, \frac{L_0}{2\delta_0}\} & \lambda < 0 \end{cases}.$$

Here, T_1, T_2 depend continuously on its parameters.

By Lemma 5.2, Lemma 5.10, Lemma 5.11, we have $A^n - 1 \in C([0, T]; X_{\phi(t)})$, $b^n \in C([0, T]; Y_{\phi(t)})$, $G^n \in C([0, T]; Y_{\phi(t)})$ and $\partial_\alpha R^n \in C([0, T]; X_{\phi(t)})$. Moreover, for $t \in [0, T]$,

$$\begin{aligned} \|b_1^n\|_{Y_{\phi(t)}} &\leq C(1 + M_{\lambda,h})|\lambda|d_{I,n}(t)^{-5/2} \leq C|\lambda|d_{I,n}^{-5/2} \leq \begin{cases} C|y_0|^{-3/4} & \lambda > 0, \\ C|y_0|^{-1}, & \lambda < 0. \end{cases} \\ \|A^n - 1\|_{X_{\phi(t)}} &\leq C(1 + |\lambda|^2 d_{I,n}(t)^{-5/2}) \leq \begin{cases} C|y_0|^{3/4}, & \lambda > 0 \\ C|y_0|^{1/2}, & \lambda < 0. \end{cases} \\ \|A^n - 1\|_{L^\infty} &\leq C(1 + |\lambda|^2 d_{I,n}(t)^{-3}) \leq \begin{cases} C(1 + |y_0|^{\frac{3}{10}}), & \lambda > 0, \\ C, & \lambda < 0. \end{cases} \\ \|A^n - 1\|_{\dot{X}_{\phi(t)}} &\leq C(M_{\lambda,h}^2 + |\lambda|^2 d_I(t)^{-4}) \leq \begin{cases} C|y_0|^{-3/4}, & \lambda > 0 \\ C|y_0|^{-1}, & \lambda < 0. \end{cases} \end{aligned} \quad (6.6)$$

for some constant C depending only on C_0, C_1, L_0^{-1} and $M_{\lambda,h}$. Since $M_{\lambda,h} \leq 1$, and $C_1 \leq \|Z_\alpha^n\|_{L^\infty} \leq 1 + \|Z_\alpha^n\|_{X_{\phi(t)}} \leq 1$, $L_0^{-1} \leq 1$, we can chose C depending only on C_0 . Since we require

$$\sup_{0 \leq t \leq T} \left(\frac{\delta_0}{L_0} - 4 - 2d_0(\|A\|_{L^\infty} + \|A\|_{\dot{Y}_{\phi(t)}}) - 4d_1\|(U_0, \partial_\alpha W_0)\|_{Y_{\phi(0)} \times X_{\phi(0)}} \right) \geq 0,$$

we choose $\delta_0 = \begin{cases} |y_0|^{7/16}, & \lambda > 0 \\ M_2, & \lambda < 0 \end{cases}$. Here, M_2 is a large but absolute constant. Basing on these arguments, we can take

$$T = \begin{cases} \tau, & \lambda < 0 \\ T = \frac{4\pi x(0)(|y_0| - |y_0|^{9/10})}{|\lambda|}, & \lambda > 0, \end{cases}$$

where $\tau > 0$ is a small but absolute constant.

By analyzing (6.5), using Lemma 5.6, we obtain

$$|x^{n+1}(t) - x(0)| \leq C \int_0^t |\Re\{\dot{x}^{n+1}(\tau)\}| d\tau \leq CT d_I(t)^{-3/2} \leq C|y_0|^{-1}, \quad (6.7)$$

and

$$|y^{n+1}(t) - y(0) - \frac{\lambda}{4\pi x(0)}| \leq CT d_I(t)^{-3/2} \leq C|y_0|^{-1}. \quad (6.8)$$

By Theorem 5, there is a unique solution $(U^{n+1}, \partial_\alpha W^{n+1}) \in C([0, T]; Y_{\phi(t)}) \times C([0, T]; X_{\phi(t)})$, to the system (6.3), such that

$$\sup_{t \in [0, T]} (\|(U^{n+1}, \partial_\alpha W^{n+1})\|_{\dot{Y}_{\phi(t)} \times X_{\phi(t)}}^2) \leq \frac{1}{4} M_{\lambda,h}^2 e^{B^n T} + \int_0^T e^{B^n(T-\tau)} \mathcal{N}^n(\tau) d\tau, \quad (6.9)$$

where by (6.6),

$$B^n := C(1 + \|b_1^n\|_{C([0, T]; Y_{\phi(t)})}) \leq C + C \sup_{0 \leq t \leq T} |\lambda| d_I^{(n)}(t)^{-5/2} := \gamma^n(T),$$

and

$$\begin{aligned} \mathcal{N}^n(t) &:= \|R_\alpha^n\|_{X_{\phi(t)}}^2 + \|G^n\|_{\dot{Y}_{\phi(t)}}^2 + C\|b_1^n\|_{\dot{Y}_{\phi(t)}}^2 \\ &\leq CM_{\lambda,h}^2 + C\lambda^2 d_{I,n}(t)^{-3} \leq Cm_0^2 + C\lambda^2 d_{I,n}(t)^{-7/2} := \beta^n(t). \end{aligned}$$

Case 1: $\lambda < 0$. By Lemma 5.12, we have $d_{I,n}(t) \geq \frac{1}{2}(d_{I,0} + \frac{|\lambda|}{4\pi x(0)}t)$. So

$$\gamma^n(t) \leq Cm_0 + C\lambda^2(d_{I,0} + |\lambda|t)^{-3} \leq Cm_0 + C\lambda^2 d_{I,0}^{-3}, \quad \beta^n(t) \leq Cm_0^2 + C\lambda^2(d_{I,0} + |\lambda|t)^{-7/2}.$$

Denote

$$\gamma_0 := Cm_0 + C\lambda^2 d_{I,0}^{-3}.$$

So we have

$$\begin{aligned} \sup_{t \in [0, T]} (\| (U^{(n+1)}, \partial_\alpha W^{n+1}) \|_{Y_{\phi(t)} \times X_{\phi(t)}}^2 &\leq \frac{1}{4} m_0^2 e^{\gamma_0 T} + C \int_0^T e^{\gamma_0(T-\tau)} \frac{\lambda^2}{(d_{I,0} + |\lambda|t)^{7/2}} d\tau \\ &\leq \frac{1}{4} M_{\lambda,h}^2 e^{\gamma_0 T} + C e^{\gamma_0 T} \frac{|\lambda|}{d_{I,0}^{5/2}}. \end{aligned}$$

Choosing $T \leq \frac{1}{\gamma_0}$ and $|\lambda|^2 \approx |d_{I,0}|^{3/2} \gg 1$. Then $C e^{\gamma_0 T} \frac{|\lambda|}{d_{I,0}^{5/2}} \leq \frac{C}{d_{I,0}}$. So we have

$$\sup_{t \in [0, T]} (\| (U^{(n+1)}, \partial_\alpha W^{n+1}) \|_{Y_{\phi(t)} \times X_{\phi(t)}}^2 \leq m_0^2 + C \frac{1}{d_{I,0}}. \quad (6.10)$$

Similarly, we obtain

$$\sup_{0 \leq t \leq T} \|U^{n+1}(\cdot, t)\|_{L^2}^2 \leq 1, \quad \sup_{0 \leq t \leq T} \|U^{n+1}(\cdot, t)\|_{L^\infty} \leq M_\infty.$$

Therefore, $(W^{n+1}, U^{n+1}, \{z_j^{n+1}\})$ satisfies (AS2) and (AS5).

Case 2: $\lambda > 0$. Let's again consider the situation when $|y_0| \gg 1$. So we take $T = \frac{4\pi x(0)(|y_0| - |y_0|^{9/10})}{|\lambda|}$.

Using the similar argument, we have

$$\sup_{t \in [0, T]} d_I(t) \geq \frac{1}{2} d_{I,0}^{9/10}, \quad (6.11)$$

and

$$\sup_{t \in [0, T]} (\| (U^{(n+1)}, \partial_\alpha W^{n+1}) \|_{Y_{\phi(t)} \times X_{\phi(t)}}^2 \leq m_0^2 + C \frac{1}{d_{I,0}^{3/4}}. \quad (6.12)$$

So $(W^{n+1}, U^{n+1}, \{z_j^{n+1}\})$ satisfies (AS2) and (AS5).

Let $\Sigma^{n+1}(t)$ be the curve parametrized by $Z^{n+1}(\alpha, t)$, and let $\Omega^{n+1}(t)$ be the region bounded above by $\Sigma^{n+1}(t)$. Define $\mathcal{U}^{n+1}$ by

$$\mathcal{U}^{n+1}(Z, t) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{Z_\beta^{n+1}(\beta, t)}{Z - Z^{n+1}(\beta, t)} F^{n+1}(\beta, t) d\beta.$$

Estimate $|Z^{n+1}(\alpha, t) - Z^{n+1}(\beta, t)|$: Since $Z^{n+1}(\alpha, t) = \alpha + (I + \mathbb{H})U^{n+1}$, we have

$$\begin{aligned} \partial_t(Z^{n+1}(\alpha, t) - Z^{n+1}(\beta, t)) &= [(\bar{F}^n(\alpha, t) - \bar{F}^n(\beta, t)) + (\bar{Q}^n(\alpha, t) - \bar{Q}^n(\beta, t))] \\ &\quad - (b^n(\alpha, t)Z_\alpha^{n+1}(\alpha, t) - b^n(\beta, t)Z_\beta^{n+1}(\beta, t)). \end{aligned}$$

Note that

$$\begin{aligned} &\left| [(\bar{F}^n(\alpha, t) - \bar{F}^n(\beta, t)) + (\bar{Q}^n(\alpha, t) - \bar{Q}^n(\beta, t))] - (b^n(\alpha, t)Z_\alpha(\alpha, t) - b^n(\beta, t)Z_\beta(\beta, t)) \right| \\ &\leq CM_{\lambda,h} |\alpha - \beta|. \end{aligned} \quad (6.13)$$

So we obtain

$$|Z_0(\alpha) - Z_0(\beta)| - CM_{\lambda,h}|\alpha - \beta|t \leq |Z^{n+1}(\alpha, t) - Z^{n+1}(\beta, t)| \leq |Z_0(\alpha) - Z_0(\beta)|t + CM_{\lambda,h}|\alpha - \beta|. \quad (6.14)$$

So we obtain

$$\frac{1}{2}C_0|\alpha - \beta| \leq |Z^{n+1}(\alpha, t) - Z^{n+1}(\beta, t)| \quad \forall \alpha, \beta \in \mathbb{R}, \quad t \in [0, T]. \quad (6.15)$$

So $(Z^{n+1}, F^{n+1}, \{z_j^{n+1}\})$ satisfies (AS3).

By (6.7), (6.8), and (6.15), $(W^{n+1}, U^{n+1}, \{z_j^{n+1}\})$ satisfies (AS4). So $(W^{n+1}, U^{n+1}, \{z_j^{n+1}\}) \in S_T$.

Error estimates and convergence of the approximate solutions. We show that $\{(W^k, F^k, \{z_j^k\})\}$ is a Cauchy sequence in some Banach space. Let

$$V^{k+1} := -\Re\{[\bar{F}^{k+1}, \mathbb{H}](\frac{1}{\partial_\alpha Z^k} - 1)\} + R^k. \quad (6.16)$$

Denote

$$\begin{cases} \hat{U} = & U^{k+1} - U^k, \\ \hat{G} = & -(b^k - b^{k-1})\partial_\alpha U^k - (A^k - A^{k-1})\Lambda W^k + G^k - G^{k-1}, \\ \hat{V} = & -(b^k - b^{k-1})\partial_\alpha W^k + V^{k+1} - V^k, \\ \hat{Z} = & Z^{k+1} - Z^k, \\ \hat{W} = & W^{k+1} - W^k, \\ \hat{z}_j = & z_j^{k+1} - z_j^k. \end{cases} \quad (6.17)$$

Then $\hat{U}$ and $\hat{W}$ satisfy

$$\begin{cases} D_t^{(k)}\hat{U} = -A^k\Lambda\hat{W} + \hat{G}, \\ D_t^{(k)}\hat{W} = -\hat{U} + \hat{V}, \\ \hat{U}(\cdot, 0) = 0, \quad \hat{W}(\cdot, 0) = 0. \end{cases} \quad (6.18)$$

And $\hat{z}_j$ satisfies

$$\begin{cases} \frac{d}{dt}\hat{z}_j = \mathcal{U}_{error}^k + \mathcal{SP}_{error}^k, \\ \hat{z}_j(0) = 0. \end{cases} \quad (6.19)$$

Here,

$$\mathcal{U}_{error}^k = \overline{\mathcal{U}^k(z_j^k(t), t)} - \overline{\mathcal{U}^{k-1}(z_j^{k-1}(t), t)},$$

and

$$\mathcal{SP}_{error}^k = \sum_{\substack{1 \leq l \leq 2 \\ l \neq j}} \frac{\lambda_l i}{2\pi} \left(\frac{1}{z_l^k(t) - z_j^k(t)} - \frac{1}{z_l^{k-1}(t) - z_j^{k-1}(t)} \right).$$

Denote

$$\mathfrak{E}^k(t) := \|(\hat{F}, \hat{W})\|_{H^4 \times H^4}^2 + |\{\hat{z}_j\}|^2. \quad (6.20)$$

It's elementary to check that

$$|\hat{z}_j(t)|^2 \leq CM_{\lambda,h}^2 t \mathfrak{E}^{k-1}(t). \quad (6.21)$$

Using Lemma 5.11, we have

$$\|\hat{V}\|_{H^4} + \|\hat{G}\|_{H^4} \leq CM_{\lambda,h} \mathfrak{E}^{k-1}. \quad (6.22)$$

Using energy estimates, we obtain

$$\|(\hat{F}, \hat{H})\|_{H^4 \times H^4}^2 \leq CM_{\lambda,h}^2 t \mathfrak{E}^{k-1}. \quad (6.23)$$

To estimate $\hat{Z}$, using (6.4), we obtain

$$\|\hat{Z}\|_{H^4} \leq CM_{\lambda,h}^2 t \mathfrak{E}^{k-1}. \quad (6.24)$$

So we obtain

$$\mathfrak{E}^k(t) \leq CM_{\lambda,h}^2 t \mathfrak{E}^{k-1}. \quad (6.25)$$

By (5.4) and (5.5) and the assumption that $|y_0|$ large, we have

$$\sup_{0 \leq t \leq T} \mathfrak{E}^k(t) \leq c \sup_{0 \leq t \leq T} \mathfrak{E}^{k-1}(t), \quad (6.26)$$

for some constant $0 < c < 1$. So $(W^n, U^n, \{z_j^n\})$ and therefore $(Z^n - \alpha, F^n, \{z_j^n\})$ is a Cauchy sequence in $C([0, T]; H^4 \times H^4 \times \mathbb{C}^2)$. So

$$(W^n, U^n, \{z_j^n\}) \rightarrow (W, U, \{z_j\}) \quad (6.27)$$

in $C([0, T]; H^4 \times H^4 \times \mathbb{C}^2)$. Since $(\partial_\alpha W^n, U^n)$ is bounded in $C([0, T]; X_{\phi(t)} \times Y_{\phi(t)})$, we have

$$(W_\alpha, U) \in C([0, T]; X_{\phi(t)} \times Y_{\phi(t)}).$$

Also,

$$Q^n \rightarrow - \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \frac{1}{Z(\alpha, t) - z_j(t)} \quad \text{in } C([0, T]; Y_{\phi(t)}). \quad (6.28)$$

$$b^n \rightarrow b = \Re\{[D_t Z, \mathbb{H}]\left(\frac{1}{Z_\alpha} - 1\right)\} + 2\Re\{Q\} + 2\Re\{F\} \quad \text{in } C([0, T]; Y_{\phi(t)}), \quad (6.29)$$

Let $\Sigma(t)$ be the curve parametrized by $Z(\alpha, t)$ and $\Omega(t)$ the region bounded above by $\Sigma(t)$. Then we have

$$\mathcal{U}^n(z, t) \rightarrow \frac{1}{2\pi i} \int \frac{Z_\beta(\beta, t)}{z - Z(\beta, t)} F(\beta, t) d\beta \quad \text{uniformly in } \Omega(t). \quad (6.30)$$

So we can verify that

$$\dot{z}_j(t) = \overline{\mathcal{U}(z_j(t), t)} + \sum_{\substack{1 \leq k \leq 2 \\ k \neq j}} \frac{\lambda_k i}{2\pi} \frac{1}{\overline{z_k(t) - z_j(t)}}, \quad (6.31)$$

and $\{z_j(t)\} \in C^2([0, T]; \Omega(t))$. So $(W, U, \{z_j\})$ is the unique solution to (2.29) on $[0, T]$. So we complete the proof of the theorem. $\square$

7. APPLICATIONS: SIGN CHANGING OF THE TAYLOR SIGN COEFFICIENT

In this section, we apply *Theorem 3* to prove *Theorem 4*.

Given appropriate initial data $(W_0, U_0, \{z_{1,0}, z_{2,0}\})$ and $\lambda > 0$, we solve the water waves backward and forward in time. Note that the backward evolution is equivalent to solving the forward in time water waves with the same initial data $(W_0, U, \{z_{1,0}, \{z_{2,0}\})$ but with λ replaced by $-\lambda$. We choose $|\lambda| = O(|y_0|^{3/2})$.

- (1) If $\lambda > 0$, the point vortices travel toward the free interface. At time $T_0 = \frac{4\pi x(0)(|y_0| - |y_0|^{9/10})}{|\lambda|}$, the distance between the point vortices and the free interface is $\approx |y_0|^{9/10}$. This distance is significantly smaller than the initial one, so we can show that $\inf_{\alpha \in \mathbb{R}} A_1(\alpha, T_0) \leq -\eta_1$, for arbitrary large $\eta_1 > 0$, provided that we choose $|y_0|$ sufficiently large.
- (2) If $\lambda < 0$, the point vortices travel away from the free interface. At time $T_0 = O(1)$, the distance between the point vortices and the free interface is $\approx |y_0|^{3/2}$. This distance is significantly larger than the initial one, so we can show that $\inf_{\alpha \in \mathbb{R}} A_1(\alpha, T_0) \geq 1 - \eta_0$, for arbitrary small $\eta_0 > 0$, provided that we choose $|y_0|$ sufficiently large.

7.1. The Taylor sign coefficient. Let $A_1 := A|Z_\alpha|^2$. we have

Proposition 1. (Corollary 4.2 in [44]) Let $(Z, F, \{z_j\})$ be a solution to the water waves system such that $(Z_\alpha - 1, D_t Z) \in C([0, T_0]; H^2 \times H^2)$, then we have

$$A_1 = 1 + \frac{1}{2\pi} \int \frac{|D_t Z(\alpha, t) - D_t Z(\beta, t)|^2}{(\alpha - \beta)^2} d\beta - \text{Im} \left\{ \sum_{j=1}^N \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} \right) (D_t Z - \dot{z}_j(t)) \right\}. \quad (7.1)$$

Corollary 7.1. Let $(Z, F, \{z_j\})$ be a solution to the water waves system such that $(Z_\alpha - 1, D_t Z) \in C([0, T_0]; H^2 \times H^2)$. Fix a time $t \in [0, T_0]$. Assume that $Z(\alpha, t) = \alpha$, $D_t Z = \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \frac{1}{\alpha - z_j(t)}$. Assume $z_1(t) = -x(t) + iy(t)$, $z_2(t) = x(t) + iy(t)$, with $x(t) > 0$, $y(t) < 0$, and $\lambda_1 = -\lambda_2 := \lambda$. We simply write $x(t)$, $y(t)$ as x , y . Then

$$\begin{aligned} A_1(\alpha, t) = & 1 + \frac{\lambda^2}{\pi^2} \frac{3y\alpha^4 + (x^2 + y^2)y(3x^2 - y^2 + 2\alpha^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2} \\ & + \frac{\lambda^2}{4\pi^2} \frac{\alpha^2 x^2 + x^4 + 5x^2 y^2}{((\alpha + x)^2 + y^2)((\alpha - x)^2 + y^2)(x^2 + y^2)|y|}. \end{aligned} \quad (7.2)$$

The detail calculation for (7.2) is in §A.1.

To simplify (7.2), we take $x \approx 1$ and $|y| \gg 1$. We will take $|\lambda| = O(|y|^{3/2})$. Direct calculation yields the following lemma.

Lemma 7.1. Let $x \leq 1$ and $|y| \gg 1$ and $\lambda^2 \leq C|y|^3$, where $C > 0$ is a given constant. Then

$$\sup_{\alpha \in \mathbb{R}} \frac{\lambda^2}{4\pi^2} \frac{\alpha^2 x^2 + x^4 + 5x^2 y^2}{((\alpha + x)^2 + y^2)((\alpha - x)^2 + y^2)(x^2 + y^2)|y|} \leq \frac{C'}{y^2}, \quad (7.3)$$

$$\sup_{\alpha \in \mathbb{R}} \frac{\lambda^2}{\pi^2} \left| \frac{3y\alpha^4 + (x^2 + y^2)y(3x^2 - y^2 + 2\alpha^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2} - \frac{3y\alpha^4 + y^3(-y^2 + 2\alpha^2)}{(\alpha^4 + y^4 + 2\alpha^2 y^2)^2} \right| \leq \frac{C'}{y^2}, \quad (7.4)$$

for some constant C' depending on C only. In particular, we have

$$|A_1(\alpha, 0) - G(\alpha; y, \lambda)| \leq \frac{C'}{y^2}, \quad (7.5)$$

where

$$G(\alpha; y, \lambda) := 1 + \frac{\lambda^2}{\pi^2} \frac{3y\alpha^4 + y^3(-y^2 + 2\alpha^2)}{(\alpha^4 + y^4 + 2\alpha^2 y^2)^2}. \quad (7.6)$$

Denote $\gamma := \frac{\lambda^2}{\pi^2 |y|^3}$. Assume $\alpha = k|y|$, we have ¹³

$$f(\gamma, k) := G(\alpha; y, \lambda) = 1 - \gamma g(k), \quad (7.7)$$

where

$$g(k) = \frac{3k^4 + 2k^2 - 1}{(k^2 + 1)^4}.$$

By routine calculus, we have

1. $g(k)$ obtains its absolute maximum $\frac{1}{4}$ at $k = \pm 1$.
2. $g(k)$ obtains its absolute minimum -1 at $k = 0$.

So we conclude that

- i. For $\gamma < 4$, we have $f(\gamma, k) > 0$ for all $k \in \mathbb{R}$.

¹³Recall that $y < 0$.

- ii. For $\gamma > 4$, we have $f(\gamma, \pm 1) < 0$.
- iii. $f(4, \pm 1) = 0$, which is equivalent to $G(\pm|y|; -|y|, 2\pi|y|^{3/2}) = 0$.

7.2. The case $\lambda_1 = -\lambda_2 > 0$ (travel upward).

7.2.1. *Initial data.* Fix $|y_0| \gg 1$, $x_0 = 1$, and $\gamma > 0$ (independent of $|y_0|$). Assume that $\lambda > 0$ and let $\lambda = \gamma^{1/2}\pi|y_0|^{3/2}$. Let

$$z_1(0) = -x_0 + i(y_0 - 1), \quad z_2(0) = x_0 + i(y_0 - 1). \quad (7.8)$$

Choosing the initial W_0 and U_0 such that

$$\|(\partial_\alpha W_0, U_0)\|_{X_{10} \times Y_{10}} + \|W_0\|_{L^2} \leq \frac{1}{|y_0|}. \quad (7.9)$$

We assume that U_0 and W_0 are odd functions. Denote $Q(\alpha, t) = \frac{\lambda}{2\pi i} \frac{1}{Z(\alpha, t) - z_1(t)} - \frac{\lambda}{2\pi i} \frac{1}{Z(\alpha, t) - z_2(t)}$. We use $L_0 = 10$ and $\delta_0 = 1000$. In this case, $d_{I,0} > |y_0|$.

7.2.2. *Some basic estimates.* By Theorem 3, there exists $T_0 > 0$ such that (2.15) admits a unique solution $(W, U, \{z_1(t), z_2(t)\})$ on $[0, T_0]$ such that

- (1) $T_0 = \frac{|y_0| - |y_0|^{9/10}}{\frac{|\lambda|}{4\pi x_0}}$.
- (2) Define $Z = \alpha + (I + \mathbb{H})W$ and $F = (I + \mathbb{H})U$. For all $t \in [0, T_0]$,

$$\|(Z_\alpha(\cdot, t) - 1, F(\cdot, t))\|_{C([0, T_0]; H^4 \times \dot{H}^4)} \leq \frac{C}{|y_0|^{3/8}}, \quad \|F\|_{C([0, T_0]; L^2)} \leq 1. \quad (7.10)$$

Here, $\dot{H}^4$ represents the homogeneous Sobolev space $\dot{H}^4$.

- (3) For each fixed $t \in [0, T_0]$, $W(\cdot, t)$ and $U(\cdot, t)$ are odd functions, and $\Re\{z_1(t)\} = -\Re\{z_2(t)\}$, $\Im\{z_1(t)\} = \Im\{z_2(t)\}$.
- (4) $\sup_{0 \leq t \leq T_0} d_I(t) \geq \frac{1}{2}|y_0|^{9/10}$.
- (5) For $0 \leq t \leq T_0$,

$$\|Q\|_{L^\infty} = \left\| \frac{\lambda}{2\pi} \left(\frac{1}{Z(\alpha, t) - z_1(t)} - \frac{1}{Z(\alpha, t) - z_2(t)} \right) \right\|_{L^\infty} \leq C|\lambda|d_I(t)^{-2} \leq C|y_0|^{-3/10}.$$

We simply bound $\|D_t Z(\cdot, t)\|_{L^\infty}$ by

$$\sup_{t \in [0, T_0]} \|D_t Z(\cdot, t)\|_{L^\infty} \leq \sup_{0 \leq t \leq T_0} (\|F(\cdot, t)\|_{L^\infty} + \|Q(\cdot, t)\|_{L^\infty}) \leq 1, \quad (7.11)$$

and simply bound b by

$$\sup_{0 \leq t \leq T_0} \|b(\cdot, t)\|_{L^\infty} \leq 1. \quad (7.12)$$

Using

$$\begin{aligned} Z(\alpha, t) - \alpha &= \int_0^t \partial_\tau (Z(\alpha, \tau) - \alpha) d\tau \\ &= \int_0^t D_\tau Z(\alpha, \tau) - b(\alpha, \tau) Z_\alpha(\alpha, \tau) d\tau, \end{aligned}$$

we obtain

$$\|Z(\alpha, t) - \alpha\|_{H^4} \leq C \sup_{t \in [0, T_0]} (\|D_t Z(\cdot, t)\|_{L^\infty} + \|b(\cdot, t)\|_{L^\infty}) t \leq C|y_0|^{-1/2} \leq C|y_0|^{-1/2}. \quad (7.13)$$

7.2.3. *The velocity of the point vortices.* We have

$$\begin{aligned}\frac{d}{dt}z_1(t) &= -\frac{\lambda i}{2\pi} \frac{1}{z_1(t) - z_2(t)} + \mathcal{U}(z_1(t), t) = \frac{\lambda i}{4\pi x(t)} + \mathcal{U}(z_1(t), t), \\ \frac{d}{dt}z_2(t) &= \frac{\lambda i}{2\pi} \frac{1}{z_2(t) - z_1(t)} + \mathcal{U}(z_2(t), t) = \frac{\lambda i}{4\pi x(t)} + \mathcal{U}(z_2(t), t).\end{aligned}\tag{7.14}$$

(5.35) implies

$$|\mathcal{U}(z_j(t), t)| \leq C d_I(t)^{-1/2} \leq C |y_0|^{-9/20}.\tag{7.15}$$

An application of (5.38) yields

$$\left| \frac{d}{dt}x(t) \right| \leq C |d_I(t)|^{-3/2} \leq C |y_0|^{-27/20}, \quad \text{for } 0 \leq t \leq T_0.\tag{7.16}$$

So

$$1 - C |y_0|^{-27/20} t \leq x(t) \leq 1 + C |y_0|^{-27/20} t, \quad 0 \leq t \leq T_0.\tag{7.17}$$

Integrating (7.14) in time, using (7.15) and (7.17), we obtain

$$\begin{cases} z_1(t) = -x_0 + i(y_0 + \frac{\lambda}{4\pi x_0} t) + O(t|y_0|^{-9/20}), \\ z_2(t) = x_0 + i(y_0 + \frac{\lambda}{4\pi x_0} t) + O(t|y_0|^{-9/20}) \end{cases}\tag{7.18}$$

In particular, at $T_0 = \frac{4\pi x_0(|y_0| - |y_0|^{9/10})}{|y_0|^{3/2}}$,

$$z_1(T_0) = -|x_0| - i|y_0|^{9/10} + O\left(\frac{1}{|y_0|^{19/20}}\right), \quad z_2(T_0) = |x_0| - i|y_0|^{9/10} + O\left(\frac{1}{|y_0|^{19/20}}\right)\tag{7.19}$$

7.2.4. *The Taylor sign at T_0 .* By (7.1), at T_0 ,

$$\begin{aligned}A_1(\alpha, T_0) &= 1 + \frac{1}{2\pi} \int \frac{|D_t Z(\alpha, T_0) - D_t Z(\beta, T_0)|^2}{(\alpha - \beta)^2} d\beta \\ &\quad - \Im \left\{ \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, T_0) - z_j(t_0))^2} \right) (D_t Z - \dot{z}_j(T_0)) \right\}.\end{aligned}$$

Denote

$$G_1(\alpha; x, y, \lambda) := \frac{\lambda^2}{\pi^2} \frac{3y\alpha^4 + (x^2 + y^2)y(3x^2 - y^2 + 2\alpha^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2}.\tag{7.20}$$

$$G_2(\alpha; x, y, \lambda) := \frac{\lambda^2}{4\pi^2} \frac{\alpha^2 x^2 + x^4 + 5x^2 y^2}{((\alpha + x)^2 + y^2)((\alpha - x)^2 + y^2)(x^2 + y^2)|y|}.\tag{7.21}$$

Proposition 2. *Choosing the initial data as in §7.2.1, then*

$$A_1(\alpha, T_0) = 1 + G_1(\alpha; 0, -|y_0|^{9/10}, \lambda) + O\left(\frac{1}{|y_0|^{1/2}}\right).\tag{7.22}$$

Decomposing the integral term in (7.1) as follows:

$$\begin{aligned}\frac{1}{2\pi} \int \frac{|D_t Z(\alpha, t) - D_t Z(\beta, t)|^2}{(\alpha - \beta)^2} d\beta &= \frac{1}{2\pi} \int \frac{|Q(\alpha, t) - Q(\beta, t)|^2}{(\alpha - \beta)^2} d\beta + \frac{1}{2\pi} \int \frac{|F(\alpha, t) - F(\beta, t)|^2}{|\alpha - \beta|^2} d\beta \\ &\quad + \frac{1}{\pi} \int \Re \frac{(F(\alpha, t) - F(\beta, t)) \overline{Q(\alpha, t) - Q(\beta, t)}}{(\alpha - \beta)^2} d\beta \\ &:= I_1(\alpha, t) + I_2(\alpha, t) + I_3(\alpha, t).\end{aligned}$$

First, direct calculation yields the following estimates for I_1, I_2 , and I_3 .

Lemma 7.2. *For $0 \leq t \leq T_0$, we have*

$$|I_1(\alpha, t)| + |I_2(\alpha, t)| + |I_3(\alpha, t)| = O\left(\frac{1}{|y_0|^{1/2}}\right). \quad (7.23)$$

Proof. For I_1 , we have

$$I_1 = \int_{|\alpha| \leq |y_0|^{1/2}} \frac{|Q(\alpha, t) - Q(\beta, t)|^2}{(\alpha - \beta)^2} d\beta + \int_{|\alpha| \geq |y_0|^{1/2}} \frac{|Q(\alpha, t) - Q(\beta, t)|^2}{(\alpha - \beta)^2} d\beta := I_{11} + I_{12}.$$

We use (5.30) to bound $\frac{Q(\alpha, t) - Q(\beta, t)}{\alpha - \beta}$ by

$$\left| \frac{Q(\alpha, t) - Q(\beta, t)}{\alpha - \beta} \right| \leq \|\partial_\alpha Q(\cdot, t)\|_{L^\infty} \leq C|\lambda|d_I(t)^{-5/2} \leq C|y_0|^{-3/4}.$$

Therefore,

$$\begin{aligned} I_{11} &\leq \int_{|\alpha| \leq |y_0|^{1/2}} \|\partial_\alpha Q(\cdot, t)\|_{L^\infty}^2 d\alpha \\ &\leq C|y_0|^{1/2}|y_0|^{-3/2} = C|y_0|^{-1}. \end{aligned}$$

For I_{12} ,

$$I_{12} \leq 2\|Q\|_{L^\infty}^2 \int_{|\alpha - \beta| \geq |y_0|^{1/2}} \frac{1}{(\alpha - \beta)^2} d\beta \leq C|y_0|^{-1}.$$

So we obtain $|I_1(\alpha, t)| \leq C|y_0|^{-1}$. Using the same argument, we obtain

$$I_2 \leq C|y_0|^{-1/2}.$$

For I_3 , using Cauchy-Schwarz inequality, we have $I_3 \leq 2(I_1 + I_2)$. So we conclude the proof of the lemma. $\square$

Second, we manipulate $-Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2}\right) (D_t Z - \dot{z}_j(t))\right\}$.

Lemma 7.3. *For $0 \leq t \leq T_0$, we have*

$$-Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2}\right) (D_t Z - \dot{z}_j(t))\right\} = G_1(\alpha; 0, y(t), \lambda) + O\left(\frac{1}{|y_0|^{1/2}}\right).$$

Proof. Indeed, by Corollary A.2,

$$-Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2}\right) (D_t Z - \dot{z}_j(t))\right\} = -\sum_{j=1}^2 \frac{\lambda_j}{\pi} Re\left\{\frac{D_t Z - \dot{z}_j}{c_0^j(\alpha - w_0^j)^2}\right\}, \quad (7.24)$$

where

$$c_0^j = (\Phi^{-1})_z(\omega_0^j), \quad \omega_0^j = \Phi(z_j). \quad (7.25)$$

We claim that ¹⁴

$$c_0^j = 1 + O\left(\frac{1}{|y_0|}\right), \quad \omega_0^j = z_j(t) + O\left(\frac{1}{|y_0|^{19/20}}\right), \quad t \in [0, T_0]. \quad (7.26)$$

Indeed, since Φ^{-1} has boundary value $Z(\alpha, t)$, we have

$$\Phi^{-1}(z, t) - z = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1}{z - \beta} (Z(\beta, t) - \beta) d\beta \quad (7.27)$$

¹⁴(7.26) is certainly not optimal. Indeed, most of the estimates in this section are quite rough. It is not our goal to obtain optimal estimates.

Expanding $\Phi(z, t)$ about $z_j(t)$, we obtain

$$\begin{aligned}\omega_0^j - z_j(t) &= \Phi(z_j(t), t) - z_j(t) = \Phi(z_j(t), t) - \Phi(\Phi^{-1}(z_j(t), t), t) \\ &\leq \|\Phi_z\|_{L^\infty} |\Phi^{-1}(z_j(t), t) - z_j(t)| \\ &\leq \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{1}{|z_j(t) - \beta|} |Z(\beta, t) - \beta| d\beta \\ &\leq C d_I(t)^{-1/2} \left(\int_{\mathbb{R}} |Z(\beta, t) - \beta|^2 d\beta \right)^{1/2}.\end{aligned}$$

Using (7.13) and (7.18), we obtain

$$|\omega_0^j - z_j(t)| \leq C(|y_0|^{9/10})^{-1/2} |y_0|^{-1/2} \leq C|y_0|^{-19/20}. \quad (7.28)$$

Taking ∂_z on both sides of (7.27) yields

$$(\Phi^{-1})_z(z, t) = 1 - \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1}{(z - \beta)^2} (Z(\beta, t) - \beta) d\beta.$$

By (7.13), Cauchy-Schwarz and $\omega_0^j = z_j(t) + O(\frac{1}{|y_0|^{19/20}})$, we obtain

$$|\Phi_z^{-1}(\omega_0^j, t) - 1| \leq \left| \int_{-\infty}^{\infty} \frac{1}{(\omega_0^j(t) - \beta)^2} (Z(\beta, t) - \beta) d\beta \right| \leq C|y_0|^{-1}.$$

So we obtain $|c_0^j - 1| \leq C|y_0|^{-1}$ and therefore verify (7.26).

Decomposing $D_t Z = \bar{F} + \bar{Q}$. We use the following rough estimate

$$\left| \sum_{j=1}^2 \frac{\lambda_j}{\pi} \frac{\bar{Q}}{c_0^j(\alpha - \omega_0^j)^2} \right| \leq C|y|^{-1}.$$

Using the symmetry $\lambda_1 = -\lambda_2$, the estimate $\|F\|_{L^\infty} \leq C|y_0|^{-1/3}$ (from Theorem 3), (7.25), we obtain

$$\left| \sum_{j=1}^2 \frac{\lambda_j}{\pi} \frac{\bar{F}}{c_0^j(\alpha - \omega_0^j)^2} \right| \leq C|\lambda| d_I(t)^{-3} \|F\|_{L^\infty} \leq C|y_0|^{-1}.$$

The same argument gives

$$\left| \sum_{j=1}^2 \frac{\lambda_j}{\pi} \frac{\mathcal{U}(z_j(t), t)}{c_0^j(\alpha - \omega_0^j)^2} \right| \leq C|y_0|^{-1}.$$

Using (7.26) and the symmetry $\lambda_1 = -\lambda_2$, it is straightforward to verify that

$$\frac{1}{c_0^j(\alpha - \omega_0^j)^2} = \frac{1}{(\alpha - z_j(t))^2} + O\left(\frac{1}{d_I(t)^3} |y_0|^{-19/20}\right).$$

using $\lambda = O(|y_0|^{3/2})$ and $d_I(t) \geq C|y_0|^{9/10}$, we have

$$\sum_{j=1}^2 \frac{\lambda_j}{\pi} \operatorname{Re} \left\{ \frac{\frac{\lambda_j}{4\pi x(t)}}{c_0^j(\alpha - \omega_0^j)^2} \right\} = \sum_{j=1}^2 \frac{\lambda_j}{\pi} \operatorname{Re} \left\{ \frac{\frac{\lambda_j}{4\pi x(t)}}{(\alpha - z_j(t))^2} \right\} + O(|y_0|^{-1/2}).$$

Since $\dot{z}_j(t) = \frac{\lambda_i}{4\pi x(t)} + \bar{\mathcal{U}}(z_j(t), t)$, we obtain

$$\begin{aligned}
& -Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2}\right) (D_t Z - \dot{z}_j(t))\right\} \\
&= -\sum_{j=1}^2 \frac{\lambda_j}{\pi} \Re\left\{\frac{-\dot{z}_j}{c_0^j(\alpha - w_0^j)^2}\right\} + O(|y_0|^{-1}) \\
&= -\sum_{j=1}^2 \frac{\lambda_j}{\pi} \Re\left\{\frac{-\frac{\lambda_i}{4\pi x(t)}}{(\alpha - z_j(t))^2}\right\} + O(|y_0|^{-1}) \\
&= \frac{\lambda^2}{\pi^2} \frac{3y(t)\alpha^4 + (x(t)^2 + y(t)^2)y(t)(3x(t)^2 - y(t)^2 + 2\alpha^2)}{(\alpha^4 + (x(t)^2 + y(t)^2)^2 + 2\alpha^2(y(t)^2 - x(t)^2))^2} + O(|y_0|^{-1}),
\end{aligned}$$

where the last equality follows from

$$-\sum_{j=1}^2 \frac{\lambda_j}{\pi} \Re\left\{\frac{-\frac{\lambda_i}{4\pi x(t)}}{(\alpha - z_j(t))^2}\right\} = \frac{\lambda^2}{\pi^2} \frac{3y(t)\alpha^4 + (x(t)^2 + y(t)^2)y(t)(3x(t)^2 - y(t)^2 + 2\alpha^2)}{(\alpha^4 + (x(t)^2 + y(t)^2)^2 + 2\alpha^2(y(t)^2 - x(t)^2))^2}. \quad (7.29)$$

For the calculation of (7.29), see §A.1. So we obtain

$$\begin{aligned}
& -Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2}\right) (D_t Z - \dot{z}_j(t))\right\} \\
&= G_1(\alpha; x(t), y(t), \lambda) + O(|y_0|^{-1}).
\end{aligned}$$

Finally, using (7.4) and $|y(t)| \geq C|y_0|^{9/10}$, we obtain

$$G_1(\alpha; x(t), y(t), \lambda) = G_1(\alpha; 0, y(t), \lambda) + O\left(\frac{1}{|y_0|^{1/2}}\right). \quad (7.30)$$

So we conclude the proof of the lemma. $\square$

Proof of Proposition 2. Proposition 2 follows from Lemma 7.2, $y(T_0) = -|y_0|^{9/10}$, and Lemma 7.3. $\square$

Now we are able to calculate $A_1(\cdot, T_0)$. Note that

$$1 + G_1(|y_0|^{9/10}; 0, -|y_0|^{9/10}, \lambda) = G(|y_0|^{9/10}; -|y_0|^{9/10}, \lambda) = 1 - \frac{\lambda^2}{\pi^2(|y_0|^{9/10})^3} \times \frac{1}{4}.$$

Therefore, (7.30) and Proposition 2, we have

$$\begin{aligned}
A_1(|y_0|^{9/10}, T_0) &= 1 + G_1(|y_0|^{9/10}; 0, -|y_0|^{9/10}, \lambda) + O\left(\frac{1}{|y_0|^{1/2}}\right) \\
&= 1 - \frac{\lambda^2}{(\pi^2|y_0|^{9/10})^3} \times \frac{1}{4} + O\left(\frac{1}{|y_0|^{1/2}}\right) \\
&= 1 - \frac{\gamma|y_0|^3}{4\pi^2|y_0|^{27/10}} + O\left(\frac{1}{|y_0|^{1/2}}\right) \\
&= 1 - \frac{\gamma|y_0|^{3/10}}{4\pi^2} + O\left(\frac{1}{|y_0|^{1/2}}\right).
\end{aligned}$$

So for any given η_1 , choosing $|y_0| \geq \left(\frac{4\pi^2(\eta_1+1)}{\gamma}\right)^{10/3}$, we have

$$A_1 \leq -\eta_1.$$

7.3. The case $\lambda_1 = -\lambda_2 < 0$ (Travel downward). To distinguish from the case that $\lambda > 0$, we write the solution as $(W_-, U_-, \{z_{j,-}\})$, and denote the corresponding Taylor sign coefficient by $A_{1,-}$. Denote the strength of the j -th point vortex by $\lambda_{j,-}$, $j = 1, 2$. We take

$$\lambda_{1,-} = -|\lambda|, \quad \lambda_{2,-} = |\lambda|,$$

so the point vortices travel downward. Z_- and F_- are defined as

$$Z_-(\alpha, t) = \alpha + (I + \mathbb{H})W_-, \quad F_- = (I + \mathbb{H})U_-.$$

7.3.1. The initial data. Choosing the same initial data as in §7.2.1: Let x_0, y_0, U_0, W_0 and γ be the same as in §7.2.1. Let

$$z_{1,-}(0) = -x_0 + i(y_0 - 1), \quad z_{2,-}(0) = x_0 + i(y_0 - 1). \quad (7.31)$$

We use $L_0 = 10$ and $\delta_0 = 1000$.

By Theorem 3, there exists $T_{0,-} > 0$ such that (2.15) admits a unique solution

$$(\partial_\alpha W_-, U_-, \{z_{1,-}(0), z_{2,-}(0)\})$$

satisfying

- (1) $T_{0,-} = O(1)$. In particular, $T_{0,-}$ does not depend on $|y_0|$.
- (2) At $T_{0,-}$,

$$d_I(T_{0,-}) = O(|\lambda|) = O(|y_0|^{3/2}).$$

- (3) W and U are odd functions, and $\Re\{z_1(t)\} = -\Re\{z_2(t)\}$, $\Im\{z_1(t)\} = \Im\{z_2(t)\}$.
- (4) For $0 \leq t \leq T_{0,-}$, we have $\frac{1}{2} \leq x(t) \leq 2$, and $y(T_{0,-}) \geq |y_0| + \frac{|\lambda|}{16\pi}T_{0,-}$. Then we have

$$Q(T_{0,-}) \leq C \frac{|\lambda|}{d_I(T_{0,-})^2} \leq C \frac{|\lambda|}{|\lambda|^2} \leq C|y_0|^{-3/2}.$$

We simply bound $D_t Z_-$ by the rough estimate

$$\|D_t Z_-(\cdot, T_{0,-})\|_{L^\infty} \leq \|F_-(\cdot, T_{0,-})\|_{L^\infty} + \|Q(\cdot, T_{0,-})\|_{L^\infty} \leq 1.$$

Using

$$|\dot{z}_{j,-}(t)| = \frac{|\lambda|}{4\pi x_0} + O(1),$$

we obtain

$$\begin{aligned} & \left| \operatorname{Im} \left\{ \sum_{j=1}^2 \frac{\lambda_{j,-} i}{2\pi} \left((I - \mathbb{H}) \frac{\partial_\alpha Z_-}{(Z_-(\alpha, t) - z_{j,-}(t))^2} \right) (D_t Z_- - \dot{z}_{j,-}(t)) \right\} \right| \\ & \leq \left| \operatorname{Im} \left\{ \sum_{j=1}^2 \frac{\lambda_{j,-} i}{2\pi} \left((I - \mathbb{H}) \frac{\partial_\alpha Z_-}{(Z_-(\alpha, t) - z_{j,-}(t))^2} \right) \right\} \right| \sum_{j=1}^2 |D_t Z_- - \dot{z}_{j,-}(t)| \\ & \leq C \frac{|\lambda|}{d_I(T_{0,-})^3} (\|F_-\|_{L^\infty} + \|Q\|_{L^\infty} + |\lambda|) \\ & \leq C \frac{|\lambda|^2}{(|\lambda|T_{0,-})^3} \\ & \leq C \frac{1}{|\lambda|T_{0,-}^3}. \end{aligned}$$

Since $T_{0,-}$ does not depend on $|\lambda|$, for given $\eta_0 \in (0, 1)$, we can take $|\lambda|$ sufficiently large such that $C \frac{1}{|\lambda|T_{0,-}^3} \leq \eta_0$. So we have

$$A_{1,-}(\alpha, T_{0,-}) \geq 1 - \left| \operatorname{Im} \left\{ \sum_{j=1}^2 \frac{\lambda_j i}{2\pi} \left((I - \mathbb{H}) \frac{\partial_\alpha Z_-}{(Z_-(\alpha, t) - z_{j,-}(t))^2} \right) (D_t Z_- - \dot{z}_{j,-}(t)) \right\} \right| \geq 1 - \eta_0. \quad (7.32)$$

7.4. Conclude the proof of Theorem 4. Let $\lambda > 0$. Let $(W, U, \{z_j\})$ and $(W_-, U_-, \{z_{j,-}\})$ be the solution constructed in §7.2 and §7.3, respectively. Let $(\tilde{W}, \tilde{U}, \{\tilde{z}_j\})$ be the solution to (2.33) with initial data

$$\begin{cases} \tilde{W}(\cdot, -T_{0,-}) := W_-(\cdot, T_{0,-}), \\ \tilde{U}(\cdot, -T_{0,-}) := U_-(\cdot, T_{0,-}), \\ \tilde{z}_j(-T_{0,-}) := z_{j,-}(T_{0,-}). \end{cases} \quad (7.33)$$

Since (2.33) is time reversible and translation invariant in time, by the uniqueness of solutions, we have

$$\tilde{W}(\cdot, t) = W_-(\cdot, T_{0,-} - t), \quad \tilde{U}(\cdot, t) = U_-(\cdot, T_{0,-} - t), \quad \tilde{z}_j(t) = z_{j,-}(T_{0,-} - t). \quad (7.34)$$

In particular,

$$\tilde{W}(\cdot, T_{0,-}) = W_-(\cdot, 0), \quad \tilde{U}(\cdot, T_{0,-}) = U_-(\cdot, 0), \quad \tilde{z}_j(T_{0,-}) = z_{j,-}(0).$$

By the uniqueness of solution again, we have

$$\tilde{W}(\cdot, t) = W(\cdot, t), \quad \tilde{U}(\cdot, t) = U(\cdot, t), \quad \tilde{z}_j(t) = z_j(t), \quad t \in [0, T_0]. \quad (7.35)$$

Denote $\tilde{A}_1$ the Taylor sign coefficient corresponding to $(\tilde{W}, \tilde{U}, \{\tilde{z}_j\})$, then

- (1) $\inf_{\alpha \in \mathbb{R}} \tilde{A}_1(\alpha, -T_{0,-}) \geq 1 - \eta_0$.
- (2) $\inf_{\alpha \in \mathbb{R}} \tilde{A}_1(\alpha, T_0) \leq -\eta_1$.

Up to a time translation $t \mapsto t + T_{0,-}$, we conclude the proof of Theorem 4.

ACKNOWLEDGEMENT

The author would like to thank Prof. Sijue Wu for suggesting this problem and thank Gong Chen for invaluable comments.

APPENDIX A. THE TAYLOR SIGN

Lemma A.1. *Let $w_1, w_2 \in \mathbb{P}_-$. Then*

$$\int_{-\infty}^{\infty} \frac{1}{(\beta - w_1)(\beta - \overline{w_2})} d\beta = \frac{2\pi i}{\overline{w_2} - w_1} \quad (A.1)$$

Proof. $\overline{w_2}$ is the only residue of $\frac{1}{(\beta - w_1)(\beta - \overline{w_2})}$ in $\mathbb{P}_+$. By residue Theorem,

$$\int_{-\infty}^{\infty} \frac{1}{(\beta - w_1)(\beta - \overline{w_2})} d\beta = \frac{2\pi i}{\overline{w_2} - w_1}.$$

□

Corollary A.1. *Assume further that $Z(\alpha, t) = \alpha$, $\bar{Q} = \sum_{j=1}^N \frac{\lambda_j i}{2\pi} \frac{1}{\alpha - z_j(t)}$. Then we have*

$$\frac{1}{2\pi} \int \frac{|\bar{Q}(\alpha, t) - \bar{Q}(\beta, t)|^2}{(\alpha - \beta)^2} d\beta = \sum_{1 \leq j, k \leq N} \frac{\lambda_j \lambda_k}{(2\pi)^2} \frac{1}{(\alpha - z_j)(\alpha - z_k)} \frac{i}{\bar{z}_k - z_j}. \quad (A.2)$$

Proof. We have

$$\bar{Q}(\alpha, t) - \bar{Q}(\beta, t) = \sum_{j=1}^N \frac{\lambda_j i}{2\pi} \frac{\beta - \alpha}{(\alpha - z_j)(\beta - z_j)}. \quad (\text{A.3})$$

So we have

$$\begin{aligned} \left| \frac{\bar{Q}(\alpha, t) - \bar{Q}(\beta, t)}{\alpha - \beta} \right|^2 &= \left| \sum_{j=1}^N \frac{\lambda_j}{2\pi} \frac{1}{(\alpha - z_j)(\beta - z_j)} \right|^2 \\ &= \sum_{j=1}^N \sum_{k=1}^N \frac{\lambda_j \lambda_k}{(2\pi)^2} \frac{1}{(\alpha - z_j)(\beta - z_j)(\alpha - z_k)(\beta - z_k)} \end{aligned} \quad (\text{A.4})$$

Apply lemma A.1, we have

$$\int_{-\infty}^{\infty} \frac{1}{(\alpha - z_j)(\beta - z_j)(\alpha - z_k)(\beta - z_k)} d\beta = \frac{1}{(\alpha - z_j)(\alpha - z_k)} \frac{2\pi i}{\bar{z}_k - z_j}.$$

So we have

$$\begin{aligned} \frac{1}{2\pi} \int \frac{|\bar{Q}(\alpha, t) - \bar{Q}(\beta, t)|^2}{(\alpha - \beta)^2} d\beta &= \sum_{1 \leq j, k \leq N} \frac{\lambda_j \lambda_k}{(2\pi)^3} \frac{1}{(\alpha - z_j)(\alpha - z_k)} \frac{2\pi i}{\bar{z}_k - z_j} \\ &= \sum_{1 \leq j, k \leq N} \frac{\lambda_j \lambda_k}{(2\pi)^2} \frac{1}{(\alpha - z_j)(\alpha - z_k)} \frac{i}{\bar{z}_k - z_j}. \end{aligned}$$

□

Lemma A.2. Let $z_0 \in \Omega(t)$. Assume that $Z(\alpha, t) = \Phi^{-1}(\alpha, t)$, where $\Phi : \Omega(t) \rightarrow \mathbb{P}_-$ is the Riemann mapping, we have

$$(I - \mathbb{H}) \frac{1}{Z(\alpha, t) - z_0} = \frac{2}{c_1(\alpha - w_0)}, \quad c_1 = (\Phi^{-1})_z(w_0), \quad w_0 = \Phi(z_0, t). \quad (\text{A.5})$$

Proof. Note that $Z(\alpha, t) = \Phi^{-1}(\alpha, t)$. So $Z(\alpha, t) - z_0$ is the boundary value of $\Phi^{-1}(z, t) - z_0$ in the lower half plane. Since Φ^{-1} is 1-1 and onto, $\Phi^{-1}(z, t) - z_0$ has a unique zero $w_0 := \Phi(z_0)$, so $\frac{1}{Z(\alpha, t) - z_0}$ has a exactly one pole of multiplicity one. For z near w_0 , we have

$$\Phi^{-1}(z, t) - z_0 = c_1(z - w_0) + \sum_{n=2}^{\infty} c_n(z - w_0)^n, \quad \text{where } c_1 = (\Phi^{-1})_z(w_0) \neq 0. \quad (\text{A.6})$$

Therefore, we have $\frac{1}{Z(\alpha, t) - z_0} - \frac{1}{c_1(\alpha - w_0)}$ is holomorphic in $\mathbb{P}_-$, and hence

$$(I - \mathbb{H}) \left(\frac{1}{Z(\alpha, t) - z_0} - \frac{1}{c_1(\alpha - w_0)} \right) = 0. \quad (\text{A.7})$$

Since $\frac{1}{c_1(\alpha - w_0)}$ is holomorphic in $\mathbb{P}_+$, we have

$$(I - \mathbb{H}) \frac{1}{Z(\alpha, t) - z_0} = (I - \mathbb{H}) \frac{1}{c_1(\alpha - w_0)} = \frac{2}{c_1(\alpha - w_0)}. \quad (\text{A.8})$$

□

Corollary A.2. Let $z_j(t) \in \Omega(t)$. Assume that $Z(\alpha, t) = \Phi^{-1}(\alpha, t)$, where $\Phi : \Omega(t) \rightarrow \mathbb{P}_-$ is the Riemann mapping, then

$$(I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} = \frac{2}{(\Phi^{-1})_z(\Phi(z_j(t)))(\alpha - \Phi(z_j(t)))^2} \quad (\text{A.9})$$

Proof. We have

$$\begin{aligned} (I - \mathbb{H}) \frac{Z_\alpha}{(Z(\alpha, t) - z_j(t))^2} &= -\partial_\alpha (I - \mathbb{H}) \frac{1}{Z(\alpha, t) - z_j(t)} \\ &= -\partial_\alpha \frac{2}{(\Phi^{-1})_z(\Phi(z_j(t)))(\alpha - \Phi(z_j(t)))} \\ &= \frac{2}{(\Phi^{-1})_z(\Phi(z_j(t)))(\alpha - \Phi(z_j(t)))^2}. \end{aligned}$$

□

Corollary A.3. Assume that $Z(\alpha, t) = \Phi^{-1}(\alpha, t)$, where $\Phi : \Omega(t) \rightarrow \mathbb{P}_-$ is the Riemann mapping

$$A_1 = 1 + \frac{1}{2\pi} \int \frac{|\mathcal{D}_t Z(\alpha, t) - \mathcal{D}_t Z(\beta, t)|^2}{(\alpha - \beta)^2} d\beta - \sum_{j=1}^N \frac{\lambda_j}{\pi} \operatorname{Re} \left\{ \frac{\mathcal{D}_t Z - \dot{z}_j}{c_0^j(\alpha - w_0^j)^2} \right\}, \quad (\text{A.10})$$

where

$$c_0^j = (\Phi^{-1})_z(\omega_0^j), \quad \omega_0^j = \Phi(z_j). \quad (\text{A.11})$$

A.1. The proof of Corollary 7.1.

Proof. we calculate $-Im\{\sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} (D_t Z(\alpha) - \dot{z}_j)\}$. We have

$$\dot{z}_j = \frac{\lambda i}{4\pi x}.$$

We have

$$\begin{aligned} \sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} &= \frac{\lambda i}{\pi} \left(\frac{1}{(\alpha - z_1)^2} - \frac{1}{(\alpha - z_2)^2} \right) = \frac{\lambda i}{\pi} \frac{(\alpha - z_2)^2 - (\alpha - z_1)^2}{(\alpha - z_1)^2 (\alpha - z_2)^2} \\ &= \frac{\lambda i}{\pi} \frac{(z_1 - z_2)(2\alpha - z_1 - z_2)}{(\alpha - z_1)^2 (\alpha - z_2)^2} \end{aligned}$$

We have

$$z_t(\alpha) = \frac{\lambda i}{2\pi} \left(\frac{1}{\alpha - z_1} - \frac{1}{\alpha - z_2} \right) = \frac{\lambda i}{2\pi} \frac{\overline{z_1 - z_2}}{(\alpha - z_1)(\alpha - z_2)}$$

So we have

$$\begin{aligned} \sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} z_t(\alpha) &= -\frac{\lambda^2}{2\pi^2} \frac{|z_1 - z_2|^2 (2\alpha - 2yi)}{|\alpha - z_1|^2 |\alpha - z_2|^2 (\alpha - z_1)(\alpha - z_2)} \\ &= -\frac{\lambda^2}{2\pi^2} \frac{8x^2(\alpha - yi)(\alpha - z_1)(\alpha - z_2)}{|\alpha - z_1|^4 |\alpha - z_2|^4} \\ &= -\frac{4\lambda^2 x^2}{\pi^2} \frac{(\alpha - yi)(\alpha^2 - x^2 - y^2 + 2y\alpha i)}{((\alpha + x)^2 + y^2)^2 ((\alpha - x)^2 + y^2)^2}, \end{aligned}$$

here, we've used

$$(\alpha - z_1)(\alpha - z_2) = \alpha^2 - \alpha(z_1 + z_2) + z_1 z_2 = \alpha^2 - x^2 - y^2 - 2y\alpha i.$$

Use also that

$$((\alpha + x)^2 + y^2)^2 ((\alpha - x)^2 + y^2)^2 = (\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2,$$

we have

$$\begin{aligned}
-Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} z_t(\alpha)\right\} &= Im\left\{\frac{4\lambda^2 x^2}{\pi^2} \frac{(\alpha - yi)(\alpha^2 - x^2 - y^2 + 2y\alpha i)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2}\right\} \\
&= \frac{4\lambda^2 x^2}{\pi^2} \frac{2y\alpha^2 - y(\alpha^2 - x^2 - y^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2} \\
&= \frac{4\lambda^2 x^2}{\pi^2} \frac{y\alpha^2 + y(x^2 + y^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2}
\end{aligned}$$

We have

$$\begin{aligned}
\sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} (-\dot{z}_j) &= -\frac{\lambda i}{4\pi x} \sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} \\
&= -\frac{\lambda i}{\pi} \frac{(z_1 - z_2)(2\alpha - z_1 - z_2)}{(\alpha - z_1)^2(\alpha - z_2)^2} \frac{\lambda i}{4\pi x} \\
&= -\frac{\lambda^2}{\pi^2} \frac{\alpha - yi}{(\alpha - z_1)^2(\alpha - z_2)^2} \\
&= -\frac{\lambda^2}{\pi^2} \frac{(\alpha - yi)(\alpha - z_1)^2(\alpha - z_2)^2}{|\alpha - z_1|^4 |\alpha - z_2|^4} \\
&= -\frac{\lambda^2}{\pi^2} \frac{(\alpha - yi)(\alpha^2 - x^2 - y^2 - 2y\alpha i)^2}{|\alpha - z_1|^4 |\alpha - z_2|^4} \\
&= -\frac{\lambda^2}{\pi^2} \frac{(\alpha - yi)((\alpha^2 - x^2 - y^2)^2 - 4y^2\alpha^2 + 4y\alpha(\alpha^2 - x^2 - y^2)i)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2}
\end{aligned}$$

So

$$-Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} (-\dot{z}_j)\right\} = \frac{\lambda^2}{\pi^2} \frac{4y\alpha^2(\alpha^2 - x^2 - y^2) - y((\alpha^2 - x^2 - y^2)^2 - 4y^2\alpha^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2}.$$

So we obtain

$$\begin{aligned}
&-Im\left\{\sum_{j=1}^2 \frac{\lambda_j i}{\pi} \frac{1}{(\alpha - z_j)^2} (D_t Z(\alpha) - \dot{z}_j)\right\} \\
&= \frac{4\lambda^2 x^2}{\pi^2} \frac{y\alpha^2 + y(x^2 + y^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2} \\
&\quad + \frac{\lambda^2}{\pi^2} \frac{4y\alpha^2(\alpha^2 - x^2 - y^2) - y((\alpha^2 - x^2 - y^2)^2 - 4y^2\alpha^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2} \\
&= \frac{\lambda^2}{\pi^2} \frac{3y\alpha^4 + (x^2 + y^2)y(3x^2 - y^2 + 2\alpha^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2}
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
& \sum_{1 \leq j, k \leq 2} \frac{\lambda_j \lambda_k}{(2\pi)^2} \frac{1}{(\alpha - z_j)(\alpha - \bar{z}_k)} \frac{i}{\bar{z}_k - z_j} \\
&= \frac{\lambda^2}{4\pi^2} \frac{1}{|\alpha - z_1|^2} \frac{i}{\bar{z}_1 - z_1} + \frac{\lambda^2}{4\pi^2} \frac{1}{|\alpha - z_2|^2} \frac{i}{\bar{z}_2 - z_2} - \frac{\lambda^2}{4\pi^2} \frac{1}{(\alpha - z_1)(\alpha - \bar{z}_2)} \frac{i}{\bar{z}_2 - z_1} \\
&\quad - \frac{\lambda^2}{4\pi^2} \frac{1}{(\alpha - z_2)(\alpha - \bar{z}_1)} \frac{i}{\bar{z}_1 - z_2} \\
&= \frac{\lambda^2}{4\pi^2} \frac{1}{(\alpha + x)^2 + y^2} \frac{1}{-2y} + \frac{\lambda^2}{4\pi^2} \frac{1}{(\alpha - x)^2 + y^2} \frac{1}{-2y} \\
&\quad + \frac{\lambda^2}{4\pi^2} \frac{(\alpha^2 - x^2 + y^2)y - 2x^2y}{((\alpha + x)^2 + y^2)((\alpha - x)^2 + y^2)(x^2 + y^2)} \\
&= \frac{\lambda^2}{4\pi^2} \frac{\alpha^2 x^2 + x^4 + 5x^2 y^2}{((\alpha + x)^2 + y^2)((\alpha - x)^2 + y^2)(x^2 + y^2)|y|}
\end{aligned}$$

So we obtain

$$\begin{aligned}
A_1(\alpha) &= 1 + \frac{\lambda^2}{\pi^2} \frac{3y\alpha^4 + (x^2 + y^2)y(3x^2 - y^2 + 2\alpha^2)}{(\alpha^4 + (x^2 + y^2)^2 + 2\alpha^2(y^2 - x^2))^2} \\
&\quad + \frac{\lambda^2}{4\pi^2} \frac{\alpha^2 x^2 + x^4 + 5x^2 y^2}{((\alpha + x)^2 + y^2)((\alpha - x)^2 + y^2)(x^2 + y^2)|y|}.
\end{aligned}$$

□

REFERENCES

- [1] Albert Ai. Low regularity solutions for gravity water waves. *Water Waves*, 1(1):145–215, 2019.
- [2] Albert Ai. Low regularity solutions for gravity water waves ii: The 2d case. *Annals of PDE*, 6:4, 2020.
- [3] Thomas Alazard, Nicolas Burq, and Claude Zuily. Cauchy theory for the water waves system in an analytic framework. *arXiv preprint arXiv:2007.08329*, 2018.
- [4] Jean-Marc Alazard, Thomas Delort. Global solutions and asymptotic behavior for two dimensional gravity water waves. *Ann. Sci. c. Norm. Supr. (4)*, 48(5):1149–1238, 2015.
- [5] Thomas Alazard, Nicolas Burq, and Claude Zuily. On the cauchy problem for gravity water waves. *Inventiones mathematicae*, 198(1):71–163, 2014.
- [6] D Ambrose and Nader Masmoudi. The zero surface tension limit two-dimensional water waves. *Communications on pure and applied mathematics*, 58(10):1287–1315, 2005.
- [7] J Thomas Beale, Thomas Y Hou, and John S Lowengrub. Growth rates for the linearized motion of fluid interfaces away from equilibrium. *Communications on Pure and Applied Mathematics*, 46(9):1269–1301, 1993.
- [8] Lydia Bieri, Shuang Miao, Sohrab Shahshahani, and Sijue Wu. On the motion of a self-gravitating incompressible fluid with free boundary. *Communications in Mathematical Physics*, 355(1):161–243, 2017.
- [9] Garrett Birkhoff. Helmholtz and taylor instability. In *Proc. Symp. Appl. Math.*, volume 13, pages 55–76, 1962.
- [10] Angel Castro, Diego Córdoba, Charles Fefferman, Francisco Gancedo, and Javier Gómez-Serrano. Finite time singularities for the free boundary incompressible euler equations. *Annals of Mathematics*, pages 1061–1134, 2013.
- [11] Angel Castro, Diego Córdoba, Charles Fefferman, Francisco Gancedo, and Javier Gómez-Serrano. Finite time singularities for water waves with surface tension. *Journal of Mathematical Physics*, 53(11):115622, 2012.
- [12] Ángel Castro, Diego Córdoba, Charles Fefferman, Francisco Gancedo, and María López-Fernández. Rayleigh-taylor breakdown for the muskat problem with applications to water waves. *Annals of Mathematics*, pages 909–948, 2012.
- [13] Kuang-An Chang, Tian-Jian Hsu, and Philip L-F Liu. Vortex generation and evolution in water waves propagating over a submerged rectangular obstacle: Part i. solitary waves. *Coastal Engineering*, 44(1):13–36, 2001.

- [14] Demetrios Christodoulou and Hans Lindblad. On the motion of the free surface of a liquid. *Communications on Pure and Applied Mathematics*, 53(12):1536–1602, 2000.
- [15] Daniel Coutand and Steve Shkoller. Well-posedness of the free-surface incompressible euler equations with or without surface tension. *Journal of the American Mathematical Society*, 20(3):829–930, 2007.
- [16] Daniel Coutand and Steve Shkoller. On the finite-time splash and splat singularities for the 3-d free-surface euler equations. *Communications in Mathematical Physics*, 325(1):143–183, 2014.
- [17] Daniel Coutand and Steve Shkoller. On the impossibility of finite-time splash singularities for vortex sheets. *Archive for Rational Mechanics and Analysis*, 221(2):987–1033, 2016.
- [18] W. Craig. An existence theory for water waves and the boussinesq and korteweg-devries scaling limits. *Comm. in P.D.E.*, 10(8):787–1003, 1985.
- [19] Christopher W Curtis and Henrik Kalisch. Vortex dynamics in nonlinear free surface flows. *Physics of Fluids*, 29(3):032101, 2017.
- [20] RA Dalrymple and BD Rogers. Numerical modeling of water waves with the sph method. *Coastal engineering*, 53(2-3):141–147, 2006.
- [21] David G Ebin. The equations of motion of a perfect fluid with free boundary are not well posed. *Communications in Partial Differential Equations*, 12(10):1175–1201, 1987.
- [22] S Fish. Vortex dynamics in the presence of free surface waves. *Physics of Fluids A: Fluid Dynamics*, 3(4):504–506, 1991.
- [23] Pierre Germain, Nader Masmoudi, and Jalal Shatah. Global solutions for the gravity water waves equation in dimension 3. *Annals of Mathematics*, pages 691–754, 2012.
- [24] Daniel Ginsberg. On the lifespan of three-dimensional gravity water waves with vorticity. *arXiv preprint arXiv:1812.01583*, 2018.
- [25] FM Hill. A numerical study of the descent of a vortex pair in a stably stratified atmosphere. *Journal of Fluid Mechanics*, 71(1):1–13, 1975.
- [26] John K Hunter, Mihaela Ifrim, and Daniel Tataru. Two dimensional water waves in holomorphic coordinates. *Communications in Mathematical Physics*, 346(2):483–552, 2016.
- [27] Mihaela Ifrim and Daniel Tataru. Two dimensional water waves in holomorphic coordinates ii: global solutions. *arXiv preprint arXiv:1404.7583*, 2014.
- [28] Mihaela Ifrim and Daniel Tataru. Two dimensional gravity water waves with constant vorticity: I. cubic lifespan. *arXiv preprint arXiv:1510.07732*, 2015.
- [29] T Iguchi, N Tanaka, and A Tani. On a free boundary problem for an incompressible ideal fluid in two space dimensions. *ADVANCES IN MATHEMATICAL SCIENCES AND APPLICATIONS*, 9:415–472, 1999.
- [30] Tatsuo Iguchi. Well-posedness of the initial value problem for capillary-gravity waves. *FUNKCIALAJ EKVACIOJ SERIO INTERNACIA*, 44(2):219–242, 2001.
- [31] Alexandru D. Ionescu and Fabio Pusateri. Global solutions for the gravity water waves system in 2d. *Inventiones mathematicae*, 199(3):653–804, 2015.
- [32] Tadayoshi Kano, Takaaki Nishida, et al. Sur les ondes de surface de leau avec une justification mathématique des équations des ondes en eau peu profonde. *Journal of Mathematics of Kyoto University*, 19(2):335–370, 1979.
- [33] David Lannes. Well-posedness of the water-waves equations. *Journal of the American Mathematical Society*, 18(3):605–654, 2005.
- [34] Hans Lindblad. Well-posedness for the motion of an incompressible liquid with free surface boundary. *Annals of mathematics*, pages 109–194, 2005.
- [35] Carlo Marchioro and Mario Pulvirenti. *Mathematical theory of incompressible nonviscous fluids*, volume 96. Springer Science & Business Media, 2012.
- [36] Daniel L Marcus and Stanley A Berger. The interaction between a counter-rotating vortex pair in vertical ascent and a free surface. *Physics of Fluids A: Fluid Dynamics*, 1(12):1988–2000, 1989.
- [37] Shuang Miao, Sohrab Shahshahani, and Sijue Wu. Well-posedness for free boundary hard phase fluids with minkowski background. *arXiv preprint arXiv:2003.02987*, 2020.
- [38] V. I. Nalimov. The cauchy-poisson problem (in russian). *Dynamika Splosh*, Sredy(18):104–210, 1974.
- [39] Masao Ogawa and Atusi Tani. Free boundary problem for an incompressible ideal fluid with surface tension. *Mathematical Models and Methods in Applied Sciences*, 12(12):1725–1740, 2002.
- [40] Masao Ogawa and Atusi Tani. Incompressible perfect fluid motion with free boundary of finite depth. *Advances in Mathematical Sciences and Applications*, 13(1):201–223, 2003.
- [41] Sijue Wu Rafe Kinsey. A priori estimates for two-dimensional water waves with angled crests. *preprint 2014*, *arXiv1406:7573*.

- [42] Jalal Shatah and Chongchun Zeng. Geometry and a priori estimates for free boundary problems of the euler's equation. *arXiv preprint math/0608428*, 2006.
- [43] Marvin Shinbrot. The initial value problem for surface waves under gravity, i: the simplest case. *Indiana University Mathematics Journal*, 25(3):281–300, 1976.
- [44] Qingtang Su. Long time behavior of 2d water waves with point vortices. *arXiv preprint arXiv:1812.00540*, 2018.
- [45] Geoffrey Ingram Taylor. The instability of liquid surfaces when accelerated in a direction perpendicular to their planes. i. *Proc. R. Soc. Lond. A*, 201(1065):192–196, 1950.
- [46] John G Telste. Potential flow about two counter-rotating vortices approaching a free surface. *Journal of Fluid Mechanics*, 201:259–278, 1989.
- [47] Xuecheng Wang. Global infinite energy solutions for the 2d gravity water waves system. *Communications on Pure and Applied Mathematics*, 71(1):90–162, 2018.
- [48] William W Willmarth, Grétar Tryggvason, Amir Hirs, and D Yu. Vortex pair generation and interaction with a free surface. *Physics of Fluids A: Fluid Dynamics*, 1(2):170–172, 1989.
- [49] Sijue Wu. A blow-up criteria and the existence of 2d gravity water waves with angled crests. *preprint 2015*, *arXiv:1502.05342*.
- [50] Sijue Wu. On a class of self-similar 2d surface water waves. *preprint 2012*, *arXiv:1206.2208*.
- [51] Sijue Wu. Wellposedness of the 2d full water wave equation in a regime that allows for non- c^1 interfaces. *arXiv:1803.08560*.
- [52] Sijue Wu. Well-posedness in sobolev spaces of the full water wave problem in 2-d. *Inventiones mathematicae*, 130(1):39–72, 1997.
- [53] Sijue Wu. Well-posedness in sobolev spaces of the full water wave problem in 3-d. *J. Amer. Math. Soc.*, 12:445–495, 1999.
- [54] Sijue Wu. Almost global wellposedness of the 2-d full water wave problem. *Inventiones mathematicae*, 177(1):45, 2009.
- [55] Sijue Wu. Global wellposedness of the 3-d full water wave problem. *Inventiones mathematicae*, 184(1):125–220, 2011.
- [56] Sijue Wu. Wellposedness and singularities of the water wave equations. *Lectures on the theory of water waves*, 426:171–202, 2016.
- [57] H. Yosihara. Gravity waves on the free surface of an incompressible perfect fluid of finite depth. *RIMS Kyoto*, 18:49–96, 1982.
- [58] Ping Zhang and Zhifei Zhang. On the free boundary problem of three-dimensional incompressible euler equations. *Communications on Pure and Applied Mathematics*, 61(7):877–940, 2008.
- [59] Fan Zheng. Long-term regularity of 3d gravity water waves. *arXiv preprint arXiv:1910.01912*, 2019.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA, 90089, USA
 E-mail address: qingtang@usc.edu