

CYLINDERS IN FANO VARIETIES

IVAN CHELTISOV, JIHUN PARK, YURI PROKHOROV, AND MIKHAIL ZAIDENBERG

ABSTRACT. This paper is a survey about cylinders in Fano varieties and related problems.

CONTENTS

1. Introduction	1
Notations.	7
Acknowledgments.	7
2. Cylinders in del Pezzo surfaces	7
2.1. Cylinders in rational surfaces	8
2.2. Absence of polar cylinders	11
2.3. Construction of polar cylinders	15
3. Cylinders in higher-dimensional varieties	19
3.1. Cylindrical Fano threefolds	20
3.2. Cylindrical Fano fourfolds	25
3.3. Cylinders in Mori fibrations	28
4. Beyond cylindricity	30
4.1. Flexible affine varieties	30
4.2. Cylinders in complements to hypersurfaces	34
4.3. Compactifications of \mathbb{C}^n	35
Appendix A. Singularities of pairs	38
References	43

Throughout this paper except for Section 4.3, we always assume that all varieties are defined over an algebraically closed field \mathbb{k} of characteristic 0.

1. INTRODUCTION

A *cylinder* in a projective variety X is a Zariski open subset $U \subset X$ such that

$$U \cong \mathbb{A}^1 \times Z$$

for an affine variety Z . If X contains a cylinder, we say that X is *cylindrical*. Since cylindrical varieties have negative Kodaira dimension, we will focus our attention on cylindrical Fano varieties, because they are building blocks of projective varieties with negative Kodaira dimension.

Example 1.1. For positive integers m, n with $m < n$, let X be the Grassmannian $\mathrm{Gr}(m, n)$ of m -dimensional subspaces of an n -dimensional vector space over \mathbb{k} . Then X is a smooth projective variety of dimension $m(n - m)$, and $-K_X \sim nH$, where H is an ample generator of the group $\mathrm{Pic}(X)$. Since X contains an open Schubert cell isomorphic to $\mathbb{A}^{m(n-m)}$, it is a cylindrical Fano variety.

2010 *Mathematics Subject Classification.* 14J50, 14J45, 14R20, 14R25, 14E05, 14E08, 14E30.

Key words and phrases. cylinder, Fano variety, unipotent group action.

However, not all Fano varieties are cylindrical, e.g. smooth cubic threefolds and smooth quartic threefolds do not contain cylinders, because they are irrational [50, 113]. On the other hand, every smooth rational projective surface contains a cylinder (see, for example, [123, Proposition 3.13]). In particular, all smooth del Pezzo surfaces (two-dimensional Fano varieties) are also cylindrical. Therefore, one can expect that all rational Fano varieties are cylindrical. However, the following example shows that this is not the case:

Example 1.2. Let X be a hypersurface of degree 6 in $\mathbb{P}(1, 1, 2, 3)$ that is given by

$$x_3^2 = x_2(x_2 + x_0x_1)(x_2 + \lambda x_0x_1),$$

for some $\lambda \in \mathbb{k} \setminus \{0, 1\}$, where x_0, x_1, x_2 and x_3 are coordinates of weights 1, 1, 2 and 3 respectively. Then X is a del Pezzo surface that has exactly two Du Val singular points of type D_4 , it is rational, has Picard number 1, and does not contain cylinders by [39, Theorem 1.5], see also Theorem 2.10 of the present survey and its proof.

The surface in Example 1.2 is singular. There are other examples of singular non-cylindrical rational surfaces (see Examples 1.27, 2.5, 2.6 below). What about *smooth* rational varieties?

Question 1.3. *Does every smooth rational Fano variety contain a cylinder?*

We do not know the answer to this question even in dimension three despite the fact that smooth three-dimensional Fano varieties (Fano threefolds) are completely classified and well studied [114]. Nevertheless, we believe that the answer to Question 1.3 is negative (see Conjectures 3.9 and 3.13). In fact, we do not know the answer to the following generalization of Question 1.3:

Question 1.4 ([33]). *Is it true that any smooth rational variety is cylindrical?*

A cylindrical variety X is birationally equivalent to a product $\mathbb{A}^1 \times Z$. Thus, if X is rationally connected, then Z is also rationally connected. In particular, if X is a cylindrical Fano threefold with Kawamata log terminal singularities, then X must be rational [218]. Moreover, we have

Proposition 1.5. *Let X be a cylindrical smooth Fano variety with $\rho(X) = 1$. Then X is birational to the product $Y \times \mathbb{A}^2$ for some rationally connected variety Y .*

Proof. Let U be a cylinder in the Fano variety X . Then $U \cong Z \times \mathbb{A}^1$ for some affine variety Z . Let \overline{Z} be a projective completion of the variety Z . Consider the natural completion

$$\overline{Z} \times \mathbb{A}^1 \subset \overline{Z} \times \mathbb{P}^1,$$

let $D = (\overline{Z} \times \mathbb{P}^1) \setminus (\overline{Z} \times \mathbb{A}^1)$, and let $\psi: \overline{Z} \times \mathbb{P}^1 \dashrightarrow X$ be the birational map induced by the open embedding $Z \times \mathbb{A}^1 \subset X$. Since $\rho(X) = 1$ by assumption, the divisor D must be ψ -exceptional, which implies that D is birational to $Y \times \mathbb{A}^1$ for some variety Y . Then X is birational to $Y \times \mathbb{A}^2$. Since X is rationally connected (see [26, 133]), the variety Y is rationally connected as well. \square

Corollary 1.6. *Let X be a cylindrical smooth Fano fourfold with $\rho(X) = 1$. Then X is rational.*

However, we do not know cylindricity of many rational smooth Fano fourfolds of Picard rank 1. For instance, we do not know whether any smooth rational cubic fourfold in \mathbb{P}^5 is cylindrical or not (see Question 3.18 and Remark 3.19). Keeping in mind Corollary 1.6, we ask

Question 1.7. *Is it true that all cylindrical smooth Fano varieties of Picard rank one are rational?*

In the paper [90], Gromov asked whether every smooth rational variety is uniformly rational? Recall from [20, 149, 178] that a smooth rational variety is said to be *uniformly rational* if its every point has a Zariski open neighborhood isomorphic to an open subset of the space \mathbb{A}^n (cf. [19]). Similarly, a smooth cylindrical projective variety is said to be *uniformly cylindrical* if its every

point is contained in a (Zariski open) cylinder (see Section 4.1 for the motivation and examples). It is easy to see that all smooth rational surfaces are uniformly rational and uniformly cylindrical. On the other hand, we do not know the answer to Gromov's question for varieties of higher dimensions, and we do not know the answer to

Question 1.8. *Is it true that any cylindrical smooth projective variety is uniformly cylindrical?*

In Section 3, we will present several cylindrical smooth Fano threefolds whose Picard groups are generated by their anticanonical divisors. We do not know such examples in any other dimension. The counter-examples to [196, Conjecture 5.1] found in [27] made us believe that such examples should exist in any dimension ≥ 4 . Therefore, we pose

Problem 1.9. *Find a cylindrical smooth Fano variety of dimension ≥ 4 whose Picard group is generated by its anticanonical divisor.*

One can also define cylindricity and uniform cylindricity for affine varieties in the same way we did this for projective varieties. Note that [123, Definition 3.4] asks that the cylinder should be principal, that is, its complement should be a principal divisor, which is not automatic.

Remark 1.10 (cf. Question 1.8). There are cylindrical smooth affine varieties that are not uniformly cylindrical. Indeed, let V be the Koras-Russell cubic threefold in \mathbb{A}^4 that is given by

$$x_1 + x_1^2 x_2 + x_3^2 + x_4^3 = 0,$$

where x_1, x_2, x_3 and x_4 are coordinates on \mathbb{A}^4 . Then V is a cylindrical smooth affine variety [135]. Since $\text{Pic}(V) = 0$, every cylinder in V is principal, that is, the complement of a principal divisor. It follows from [64, Corollary 4.5] that $(0, 0, 0, 0)$ is fixed by any element of $\text{Aut}(V)$, which implies that this point is not contained in any cylinder in V . Indeed, otherwise the origin would be moved by a suitable \mathbb{G}_a -action on V , cf. Theorem 1.13 below.

Like in the projective case, every cylindrical affine variety X has negative log Kodaira dimension. Moreover, a smooth affine surface contains a cylinder if and only if its log Kodaira dimension is negative [154, Ch. 2, Theorem 2.1.1], cf. [155]. However, this is no longer true in higher dimensions:

Example 1.11. Let X be a smooth hypersurface in \mathbb{P}^n of degree $n \geq 3$. Then $\mathbb{P}^n \setminus X$ is a smooth affine n -fold of negative Kodaira dimension that does not contain cylinders [33, 58].

The problem of existence of cylinders in projective varieties is closely related to unipotent actions on the affine cones over them. To illustrate this link, consider the following

Question 1.12 ([66, Question 2.22]). *Let V be the affine cone in \mathbb{A}^4 over the Fermat cubic surface, which is given by*

$$x_1^3 + x_2^3 + x_3^3 + x_4^3 = 0,$$

where x_1, x_2, x_3 and x_4 are coordinates on \mathbb{A}^4 . Does V admit an effective \mathbb{G}_a -action?

The answer to this question is negative [38], see also [53, Theorem 7.1] for a purely algebraic proof. The geometric proof of this fact is based on the following result:

Theorem 1.13 ([123, Proposition 3.1.5]). *An affine variety V admits an effective \mathbb{G}_a -action if and only if V contains a principal effective divisor D such that $V \setminus \text{Supp}(D)$ is a cylinder.*

Using this criterion, we can formulate the corresponding criterion for projective varieties, which requires the following refined notion of cylindricity:

Definition 1.14. Let X be a projective normal variety that contains a Zariski open cylinder U , and let H be an ample \mathbb{Q} -Cartier \mathbb{Q} -divisor on X . The cylinder U is said to be H -polar if

$$U = X \setminus \text{Supp}(D)$$

for some effective \mathbb{Q} -divisor D on the variety X such that $D \sim_{\mathbb{Q}} H$.

Now, we are in position to state the following criterion discovered in [124].

Theorem 1.15. *Let X be a projective normal variety, let H be an ample Cartier divisor on it, let*

$$V = \text{Spec} \left(\bigoplus_{n \geq 0} H^0(\mathcal{O}_X(nH)) \right).$$

Then V admits an effective \mathbb{G}_a -action $\iff X$ contains an H -polar cylinder.

Corollary 1.16. *Let X be a smooth rational projective surface. Then there is an embedding $X \hookrightarrow \mathbb{P}^n$ such that the affine cone in \mathbb{A}^{n+1} over X admits an effective \mathbb{G}_a -action.*

Corollary 1.17. *Let X be a projective normal variety in \mathbb{P}^n whose divisor class group is of rank 1. Then the affine cone in \mathbb{A}^{n+1} over X admits an effective \mathbb{G}_a -action $\iff X$ is cylindrical.*

Remark 1.18. Let X , H and V be as in Theorem 1.15. If V is \mathbb{Q} -Gorenstein and admits an effective action of the additive group \mathbb{G}_a , then X is a Fano variety and $H \sim_{\mathbb{Q}} -\lambda K_X$ for some $\lambda \in \mathbb{Q}_{>0}$ [123, (3.18)]. This explains our primary interest in the affine cones over Fano varieties.

The problem of existence of an effective \mathbb{G}_a -action on affine varieties is interesting on its own. If an affine variety V admits a non-trivial \mathbb{G}_a -action and $\dim(V) \geq 2$, then $\text{Aut}(V)$ is infinite dimensional and non-algebraic [67]. On the other hand, if it does not admit non-trivial \mathbb{G}_a -actions, then $\text{Aut}(V)$ contains a unique maximal torus \mathbb{T} , and $\text{Aut}(V)$ is an extension of its centralizer by a discrete subgroup in $\text{GL}_r(\mathbb{Z})$ (see [9] for details).

Example 1.19. Let V be the Pham–Brieskorn surface in \mathbb{A}^3 , which is given by

$$x_1^{a_1} + x_2^{a_2} + x_3^{a_3} = 0,$$

where a_1, a_2, a_3 are integers such that $2 \leq a_1 \leq a_2 \leq a_3$, and x_1, x_2, x_3 are coordinates on \mathbb{A}^3 . By [121, Lemma 4], the affine variety V admits an effective \mathbb{G}_a -action $\iff a_1 = a_2 = 2$.

Affine varieties that do not admit effective \mathbb{G}_a -actions are often called *rigid* [6, 7, 24, 67, 84, 121]. Applying [123, Corollary 2.1.4] and [9, Proposition 4.1] to affine cones over projective varieties, we obtain the following result:

Theorem 1.20. *Let V be the affine cone in \mathbb{A}^{n+1} over a projectively normal subvariety $X \subset \mathbb{P}^n$. Suppose that V is rigid and $\text{Aut}(X)$ is finite. Then there exists an exact sequence of groups*

$$1 \longrightarrow \mathbb{G}_m \longrightarrow \text{Aut}(V) \longrightarrow \text{Aut}(X),$$

so that $\text{Aut}(V)$ is a finite extension of the torus \mathbb{G}_m by a finite subgroup in $\text{Aut}(X)$.

In particular, combining this result with the negative answer to Question 1.12, we obtain

Corollary 1.21. *If V is the affine hypersurface from Question 1.12, then $\text{Aut}(V) = \mathbb{G}_m \times (\mu_3^3 \rtimes \mathbb{S}_4)$.*

Both Question 1.12 and Example 1.19 are very special cases of the following old conjecture, which has been confirmed in many cases (see [49] and Remark 3.19).

Conjecture 1.22 ([66, 121]). *Let V the Pham–Brieskorn hypersurface in \mathbb{A}^n with $n \geq 3$ given by*

$$x_1^{a_1} + x_2^{a_2} + \cdots + x_n^{a_n} = 0,$$

where a_1, \dots, a_n are integers such that $2 \leq a_1 \leq \cdots \leq a_n$, and x_0, x_1, \dots, x_n are coordinates on \mathbb{A}^n . Suppose that $a_2 \geq 3$. Then the affine hypersurface V is rigid.

In fact, using Theorem 1.15, we can restate Question 1.12 as follows:

Question 1.23. *Let X be the Fermat cubic surface. Does X contain $(-K_X)$ -polar cylinder?*

As we already mentioned, this question has a negative answer. Moreover, we will see later that the answer is also negative for any smooth cubic surface (cf. Theorem 2.8). This brings us to

Problem 1.24. *Describe Fano varieties that do not contain anticanonical polar cylinders.*

This problem has been solved for del Pezzo surfaces with Du Val singularities in [38, 39, 123]. However, it is still open for smooth Fano threefolds and singular del Pezzo surfaces with quotient singularities. For Fano varieties whose divisor class groups is of rank 1, Problem 1.24 is equivalent to the cylindricity problem (the problem of existence of cylinders).

Remark 1.25. One can consider Problem 1.24 for Fano varieties defined over an arbitrary possibly algebraically non-closed field. In Section 3.3, we will give a motivation for doing this.

Let us present one obstruction for the existence of anticanonical polar cylinders in Fano varieties. Recall from [44, 208] that the α -invariant of Tian of the Fano variety X is the number

$$\alpha(X) = \sup \left\{ \lambda \in \mathbb{Q} \mid \begin{array}{l} \text{the log pair } (X, \lambda D) \text{ is log canonical} \\ \text{for any effective } \mathbb{Q}\text{-divisor } D \sim_{\mathbb{Q}} -K_X \end{array} \right\}.$$

This number plays an important role in K-stability of Fano varieties, since X is K-stable if

$$\alpha(X) > \frac{\dim(X)}{\dim(X) + 1}.$$

On the other hand, we have the following result.

Theorem 1.26. *Let X be a Fano variety that has at most Kawamata log terminal singularities. If $\alpha(X) \geq 1$, then X does not contain $(-K_X)$ -polar cylinders.*

Proof. Suppose X contains a $(-K_X)$ -polar cylinder. Then $U \cong Z \times \mathbb{A}^1$ for an affine variety Z , and

$$U = X \setminus \text{Supp}(D)$$

for some effective \mathbb{Q} -divisor D on X such that $D \sim_{\mathbb{Q}} -K_X$. Arguing as in the proof Corollary 2.7, we see that the log pair (X, D) is not log canonical in this case, so that $\alpha(X) < 1$. \square

Let us show how to use this obstruction.

Example 1.27. Let X be a del Pezzo surface with Du Val singularities of degree $K_X^2 = 1$ such that one of the following two conditions holds:

- (1) either X has 2 singular points of type A_3 and 2 singular points of type A_1 ;
- (2) or the surface X has 4 singular points of type A_2 .

By [217, Theorem 1.2], the surface X exists, and it is uniquely determined by its singularities. Moreover, it follows from [217, Table 4.1] that the pencil $|-K_X|$ contains exactly 4 singular fibers. They are singular fibers of types I_4 and I_2 (in the first case) or of types I_2 (in the second case). This gives $\alpha(X) = 1$ by [34, Theorem 1.25], so that X contains no anticanonical polar cylinders. Since the group $\text{Cl}(X)$ is of rank 1, the surface X contains no cylinders at all.

Remark 1.28. Implicitly, Theorem 1.26 has been already used by many people for quite some time. For instance, Miyanishi conjectured in [94] that the smooth locus of a del Pezzo surface with quotient singularities and Picard rank 1 admits a finite unramified covering that contains a cylinder. It turned out to be wrong. Namely, in [122, Example 21.3.3], Keel and McKernan have constructed a singular del Pezzo surface X with quotient singularities such that $\rho(X) = 1$ and $\alpha(X) \geq 1$, but its smooth locus has trivial algebraic fundamental group. Thus, its smooth locus does not admit non-trivial unramified coverings, and X does not contain cylinders by Theorem 1.26.

Using Theorem 1.26, we can create many rational Fano varieties without anticanonical polar cylinders. Indeed, if X and Y are Fano varieties that have Kawamata log terminal singularities, then it follows from [44, Lemma 2.29] and [136, Proposition 8.11] that

$$\alpha(X \times Y) = \min\{\alpha(X), \alpha(Y)\}.$$

Thus, if S is a general smooth del Pezzo surface with $K_S^2 = 1$, then $\alpha(S) = 1$ by [30, Theorem 1.7], which implies that we also have $\alpha(X) = 1$ for the $2n$ -dimensional smooth Fano variety

$$X = \underbrace{S \times S \times \cdots \times S}_{n \text{ times}},$$

so that X does not contain $(-K_X)$ -polar cylinders, but X is cylindrical, because S is cylindrical. We can construct many similar examples using [34, 36, 37, 46, 197].

Example 1.29. Let S be a general smooth del Pezzo surface with $K_S^2 = 1$, and let Y be a general smooth hypersurface in $\mathbb{P}(1^{n+1}, n)$ of degree $2n$ for $n \geq 3$. Then $\alpha(S) = 1$ by [30, Theorem 1.7], and $\alpha(Y) = 1$ by [197, Theorem 2] (see also [37]). Let $X = S \times Y$. Then $\dim(X) = 2 + n \geq 5$ and

$$\alpha(X) = \min\{\alpha(S), \alpha(Y)\} = 1,$$

so that X contains no $(-K_X)$ -polar cylinder by Theorem 1.26. But X is cylindrical.

Surprisingly, we do not know a single example of a cylindrical smooth Fano threefold that contains no anticanonical polar cylinder (cf. Examples 3.14, 3.15, 3.16 and 3.17).

Problem 1.30. *Find a cylindrical smooth Fano threefold without anticanonical polar cylinder.*

Note that there are Fano varieties without cylinders whose α -invariant of Tian is smaller than 1. For instance, if X is the del Pezzo surface from Example 1.2, then $\alpha(X) = \frac{1}{2}$ by [34, Theorem 1.25]. On the other hand, this surface does not contain cylinders [39]. Note that it is K-polystable [165]. Surprisingly, all known K-unstable Fano varieties are also cylindrical.

Example 1.31 ([70–72, 114]). Let X be a smooth Fano variety of dimension $n \geq 2$ such that

$$-K_X \sim (n-1)H,$$

where H is an ample divisor such that $H^n = 5$. Then $n \in \{2, 3, 4, 5, 6\}$, and X is unique for each n . The divisor H is very ample, and the linear system $|H|$ gives an embedding $X \hookrightarrow \mathbb{P}^{n+3}$ such that the image is a section of the Grassmannian $\text{Gr}(2, 5) \subset \mathbb{P}^9$ by a linear subspace of dimension $3 + n$. Moreover, if $n \neq 2$, then $\text{Pic}(X) = \mathbb{Z}[H]$. Furthermore, the following assertions hold.

- The variety X contains a Zariski open subset isomorphic to \mathbb{A}^n , so that it is cylindrical. If $n \neq 5$, this follows from Example 1.1 and Theorems 3.6 and 3.20 (see also [71, 189]). If $n = 5$, then X contains a plane Π such that there exists the following Sarkisov link:

$$\begin{array}{ccc} & \tilde{X} & \\ \alpha \swarrow & & \searrow \beta \\ X & & \mathbb{P}^5 \end{array}$$

where α is the blowup of the plane Π , and β is the blowup of a smooth cubic scroll in \mathbb{P}^5 . This easily implies that X contains a Zariski open subset isomorphic to \mathbb{A}^5 .

- If $n \in \{2, 3, 6\}$, then X is known to be K-polystable (see, for example, [30, 45, 173, 209, 219]). On the other hand, if $n \in \{4, 5\}$, then X is K-unstable by [69].

Keeping in mind Theorem 1.26 and examples of K-stable Fano varieties without anticanonical polar cylinders (for example, smooth del Pezzo surfaces of degree 1, 2 and 3), we pose

Conjecture 1.32. *Let X be a Fano variety that has at most Kawamata log terminal singularities. If X does not contain $(-K_X)$ -polar cylinders, then X is K -polystable.*

For a projective variety X , consider the following subset of the cone of ample \mathbb{Q} -divisors on X :

$$\text{Amp}^{cyl}(X) = \left\{ H \in \text{Amp}(X) \mid \text{there is an } H\text{-polar cylinder on } X \right\}.$$

Let us call it the *cone of cylindrical ample divisors* of the variety X . We have seen in Examples 1.2 that $\text{Amp}^{cyl}(X)$ can be empty even if X is a Fano variety. Thus, we can enhance Problem 1.24 by

Problem 1.33. *For a given Fano variety X , describe the cone $\text{Amp}^{cyl}(X)$.*

This problem is not yet solved even for smooth del Pezzo surfaces. However, we know the answer for many of them (see [40]). Namely, if X is a smooth del Pezzo surface such that $K_X^2 \geq 4$, then

$$\text{Amp}^{cyl}(X) = \text{Amp}(X).$$

On the other hand, if $K_X^2 \leq 3$, then $-K_X \notin \text{Amp}^{cyl}(X)$. This gives an evidence for

Conjecture 1.34. *If X is a Fano variety, then $-K_X \in \text{Amp}^{cyl}(X) \iff \text{Amp}^{cyl}(X) = \text{Amp}(X)$.*

Let us describe the structure of this survey. In Section 2 we review results about polar cylinders in rational surfaces. In Section 3, we describe results about cylinders in smooth Fano threefolds, smooth Fano fourfolds, and del Pezzo fibrations. In Section 4, we survey results on three topics that are closely related to the main topic of this survey: flexibility of affine varieties with a special accent on the flexibility of affine cones over Fano varieties, cylinders in the complements to hypersurfaces in weighted projective spaces, and compactifications of \mathbb{C}^n . Finally, in Appendix A, we present some results about singularities of two-dimensional log pairs, which are used in Section 2 to prove the absence of polar cylinders in some del Pezzo surfaces.

Notations. Throughout this paper, we will use the following notation:

- μ_n is a cyclic subgroup of order n ;
- \mathbb{G}_a is a one-dimensional unipotent additive group;
- \mathbb{G}_m is a one-dimensional algebraic torus;
- \mathbb{F}_n is the Hirzebruch surface;
- \mathbb{P}^n is the n -dimensional projective space over \mathbb{k} ;
- \mathbb{A}^n is the n -dimensional affine space over \mathbb{k} ;
- $\mathbb{P}(a_1, \dots, a_n)$ is the weighted projective space;
- for a variety X , we denote by $\rho(X)$ the rank of its Picard group.

Acknowledgments. This work was supported by the Royal Society grant No. IES\R1\180205 and by the Russian Academic Excellence Project 5-100. The second author has been supported by IBS-R003-D1, Institute for Basic Science in Korea.

The authors would like to thank Adrien Dubouloz and Sasha Perepechko for useful comments.

2. CYLINDERS IN DEL PEZZO SURFACES

In this section, we review results about cylinders in del Pezzo surfaces. A *del Pezzo surface* means here a two-dimensional Fano variety with at most quotient singularities. Recall that a smooth del Pezzo surface is either $\mathbb{P}^1 \times \mathbb{P}^1$, or a blowup of \mathbb{P}^2 in at most 8 points such that

- at most 2 points are contained in a line;
- at most 5 points are contained in a conic;
- there is no singular cubic in \mathbb{P}^2 that contains 8 points and is singular in one of them.

A Gorenstein del Pezzo surface is a del Pezzo surface whose anticanonical divisor is Cartier, equivalently a del Pezzo surface with only Du Val singularities. Such surface is either a quadric, or its minimal resolution of singularities can be obtained by blowing up \mathbb{P}^2 in at most 8 points such that at most 3 of them are contained in a line, and at most 6 of them are contained in a conic.

First, let us go over basic facts about cylinders in rational surfaces.

2.1. Cylinders in rational surfaces. Observe that every smooth rational surface is cylindrical. This immediately follows from the fact that \mathbb{P}^2 contains a cylinder and the following

Lemma 2.1. *Let C be an irreducible curve in \mathbb{F}_n that is a section of the natural projection $\mathbb{F}_n \rightarrow \mathbb{P}^1$, and let F_1, \dots, F_r be fibers of this projection, where $r \geq 1$. Then $\mathbb{F}_n \setminus (C \cup F_1 \cup \dots \cup F_r)$ is a cylinder.*

Proof. Performing appropriate elementary birational transformations, we may assume that $C^2 = 0$, so that $n = 0$. In this case, the required assertion is obvious. \square

However, as we have seen already in Example 1.2, there are singular rational surfaces that contain no cylinders. Let us explain how to find many such rational surfaces and provide an obstruction for the existence of cylinders (see Remark 2.3 below), which will be used in Section 2.2 to show the absence of anticanonical polar cylinders in smooth del Pezzo surfaces of degree 1, 2 and 3.

Let S be a rational surface with quotient singularities and suppose that S contains a cylinder U . Then U is a Zariski open subset in S such that $U \cong \mathbb{A}^1 \times Z$ for some affine curve Z . We then have the following commutative diagram

$$\begin{array}{ccccccc}
 \mathbb{P}^1 \times \mathbb{P}^1 & \hookleftarrow & \mathbb{A}^1 \times \mathbb{P}^1 & \hookleftarrow & \mathbb{A}^1 \times Z \cong U & \hookrightarrow & S \xleftarrow{\pi} \tilde{S} \\
 & \searrow & \downarrow p_2 & & \downarrow p_Z & \nearrow \psi & \nearrow \varphi \\
 & & \mathbb{P}^1 & & Z & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathbb{P}^1 & & \mathbb{P}^1 & &
 \end{array}$$

(Note: The diagram shows a commutative diagram with nodes $\mathbb{P}^1 \times \mathbb{P}^1$, $\mathbb{A}^1 \times \mathbb{P}^1$, $\mathbb{A}^1 \times Z \cong U$, S , \tilde{S} , \mathbb{P}^1 , and Z . Arrows include \bar{p}_2 , p_2 , p_Z , ψ , and φ . Dashed arrows indicate birational maps.)

where p_Z , p_2 and \bar{p}_2 are the natural projections to the second factors, ψ is the rational map induced by p_Z , π is a birational morphism resolving the indeterminacy of ψ and φ is a morphism. By construction, a general fiber of φ is \mathbb{P}^1 . Let C_1, \dots, C_n be the irreducible curves in S such that

$$S \setminus U = \bigcup_{i=1}^n C_i.$$

The curves C_1, \dots, C_n generate the divisor class group $\text{Cl}(S)$ of the surface S , because $\text{Cl}(U) = 0$. In particular, one has

$$(2.2) \quad \text{rank Cl}(S) \leq n.$$

Let E_1, \dots, E_r be all exceptional curves of the morphism π (if any), and let $\Gamma = \mathbb{P}^1 \times \mathbb{P}^1 \setminus \mathbb{A}^1 \times \mathbb{P}^1$. Denote by $\tilde{C}_1, \dots, \tilde{C}_n$ and $\tilde{\Gamma}$ the proper transforms \tilde{S} of the curves C_1, \dots, C_n and Γ , respectively. Then $\tilde{\Gamma}$ is a section of the conic bundle φ , and $\tilde{\Gamma}$ is one of the curves $\tilde{C}_1, \dots, \tilde{C}_n$ and E_1, \dots, E_r . Moreover, all other curves among $\tilde{C}_1, \dots, \tilde{C}_n$ and E_1, \dots, E_r are components of some fibers of φ . Thus, we may assume that either $\tilde{\Gamma} = \tilde{C}_1$ or $\tilde{\Gamma} = E_r$. Then ψ is a morphism $\iff \tilde{\Gamma} = \tilde{C}_1$.

Let $\lambda_1, \dots, \lambda_n$ be arbitrary rational numbers, and let $D = \lambda_1 C_1 + \dots + \lambda_n C_n$. Then

$$K_{\tilde{S}} + \sum_{i=1}^n \lambda_i \tilde{C}_i + \sum_{i=1}^r \mu_i E_i \sim_{\mathbb{Q}} \pi^* (K_S + D)$$

8

for some real numbers μ_1, \dots, μ_r . Let \tilde{F} be a general fiber of φ . Then $K_{\tilde{S}} \cdot \tilde{F} = -2$ by the adjunction formula. Put $F = \pi(\tilde{F})$. If $\tilde{\Gamma} = E_r$, then

$$-2 + \mu_r = \left(K_{\tilde{S}} + \sum_{i=1}^n \lambda_i \tilde{C}_i + \sum_{i=1}^r \mu_i E_i \right) \cdot \tilde{F} = \pi^* (K_S + D) \cdot \tilde{F} = (K_S + D) \cdot F$$

Similarly, if $\tilde{\Gamma} = C_1$, then

$$-2 + \lambda_1 = \left(K_{\tilde{S}} + \sum_{i=1}^n \lambda_i \tilde{C}_i + \sum_{i=1}^r \mu_i E_i \right) \cdot \tilde{F} = \pi^* (K_S + D) \cdot \tilde{F} = (K_S + D) \cdot F.$$

On the other hand, if $K_S + D$ is pseudo-effective, then $(K_S + D) \cdot F \geq 0$.

Remark 2.3. We are therefore able to draw the following conclusions:

- if $K_S + D$ is pseudo-effective, then (S, D) is not log canonical;
- if $K_S + D$ is pseudo-effective and $\lambda_i < 2$ for each $i \in \{1, \dots, n\}$, then ψ is not a morphism.

Corollary 2.4. *A rational surface with quotient singularities and pseudo-effective canonical divisor cannot contain any cylinder.*

Now we present two examples of rational singular surfaces with nef canonical divisors, which do not contain cylinders by Corollary 2.4. For more examples, see [104, 145, 146, 167–170, 214].

Example 2.5 (cf. [166]). Let E be the Fermat cubic curve in \mathbb{P}^2 . Take $\sigma \in \text{Aut}(E)$ of order 6 that fixes a point in E . Let $S = E \times E / \langle \sigma \rangle$, where σ acts on $E \times E$ diagonally. Then S is rational. Moreover, it has quotient singularities and $6K_S \sim 0$. Then S contains no cylinder by Corollary 2.4.

Example 2.6 ([132]). Let $a_0, a_1, a_2, a_3, w_0, w_1, w_2, w_3$ be positive integers such that

- $a_0 \geq 4, a_1 \geq 4, a_2 \geq 4, a_3 \geq 4$;
- $a_0 w_0 + w_1 = a_1 w_1 + w_2 = a_2 w_2 + w_3 = a_3 w_3 + w_0$;
- $\gcd(w_0, w_2) = 1, \gcd(w_1, w_3) = 1$.

From the first condition above we obtain

$$\begin{cases} w_0 = a_1 a_2 a_3 - a_2 a_3 + a_3 - 1, \\ w_1 = a_0 a_2 a_3 - a_0 a_3 + a_0 - 1, \\ w_2 = a_0 a_1 a_3 - a_0 a_1 + a_1 - 1, \\ w_3 = a_0 a_1 a_2 - a_1 a_2 + a_2 - 1. \end{cases}$$

Let S be the hypersurface in $\mathbb{P}(w_0, w_1, w_2, w_3)$ defined by the following equation:

$$x_0^{a_0} x_1 + x_1^{a_1} x_2 + x_2^{a_2} x_3 + x_3^{a_3} x_0 = 0,$$

where x_0, x_1, x_2 and x_3 are coordinates of weights w_0, w_1, w_2, w_3 , respectively. Then

$$K_S = \mathcal{O}_S(a_0 a_1 a_2 a_3 - w_0 - w_1 - w_2 - w_3 - 1)$$

and $a_0 a_1 a_2 a_3 - w_0 - w_1 - w_2 - w_3 - 1 > 0$, so that K_S is ample. But S is rational by [132, Theorem 39]. By Corollary 2.4, the surface S cannot contain any cylinder.

We are mostly interested in cylinders in del Pezzo surfaces. Applying our Remark 2.3 to them, we obtain the following special case of Theorem 1.26, which we already applied in Example 1.27.

Corollary 2.7. *Suppose that $-K_S$ is ample, and U is a $(-K_S)$ -polar cylinder. Then $\alpha(S) < 1$.*

Proof. There exists an effective \mathbb{Q} -divisor D' on the surface S such that $D' \sim_{\mathbb{Q}} -K_S$ and

$$D' = \sum_{i=1}^n a_i C_i,$$

for some positive rational numbers a_1, \dots, a_n . Let $D = D'$. Then $K_S + D \sim_{\mathbb{Q}} 0$ is pseudo-effective, so that (S, D) is not log canonical by Remark 2.3, which implies that $\alpha(S) < 1$. \square

Now, we state main result of this section, which implies negative answer to Question 1.12.

Theorem 2.8 ([38,39,123,126]). *Let S be a del Pezzo surface that has at most Du Val singularities. Then S does not contain $(-K_S)$ -polar cylinders exactly when*

- $K_S^2 = 1$ and S allows at most singular points of types A_1, A_2, A_3, D_4 if any;
- $K_S^2 = 2$ and S allows at most singular points of type A_1 if any;
- $K_S^2 = 3$ and S is smooth.

Corollary 2.9. *A smooth del Pezzo surface S contains a $(-K_S)$ -polar cylinder $\iff K_S^2 \geq 4$.*

In the next two subsections, we will explain how to prove Theorem 2.8. Now let us use this result to find all del Pezzo surfaces with Du Val singularities that contain no cylinder.

Theorem 2.10 ([16, Theorem 1.6]). *Let S be a del Pezzo surface that has Du Val singularities. Then S contains no cylinder \iff it is one of the surfaces described in Examples 1.2 and 1.27.*

Proof. If S is one of the surfaces from Examples 1.2 and 1.27, then $\rho(S) = 1$, so that it does not contain cylinders by Theorem 2.8. To prove the converse assume that S contains no cylinder. Let us show that S is one of the singular del Pezzo surfaces described in Examples 1.2 and 1.27. If $\rho(S) = 1$, this follows from Theorem 2.8 and [217, Theorem 1.2].

We may assume that $\rho(S) \geq 2$. Let us seek for a contradiction. Since every smooth rational surface contains a cylinder, we see that S is singular. Then $K_S^2 \leq 2$ by Theorem 2.8.

Let $\pi: S \rightarrow Y$ be the contraction of an extremal ray of the Mori cone $\overline{\text{NE}}(S)$ of the surface S . Then it follows from [161] that one of the following cases hold:

- either π is a conic bundle, $Y = \mathbb{P}^1$ and $\rho(S) = 2$;
- or π is birational, Y is a del Pezzo surface with Du Val singularities, $\rho(Y) = \rho(S) + 1$; the morphism π is a weighted blowup of a smooth point in Y with weights $(1, k)$ for $k \geq 1$, and $K_Y^2 = K_S^2 + k$.

Suppose that π is a conic bundle. Then we have the following commutative diagram:

$$\begin{array}{ccc} & \tilde{S} & \\ \alpha \swarrow & & \searrow \beta \\ S & & \mathbb{F}_n \\ \pi \downarrow & & \downarrow \\ \mathbb{P}^1 & \xlongequal{\quad} & \mathbb{P}^1 \end{array}$$

where α is a minimal resolution of singularities, β is a birational map, and $\mathbb{F}_n \rightarrow \mathbb{P}^1$ is a natural projection. On the other hand, it follows from Tsen's theorem that S contains a smooth irreducible curve Z that is a section of the conic bundle π . Let C be its proper transform on \mathbb{F}_n . Then

$$S \setminus (Z \cup T_1 \cup \dots \cup T_r) \cong S \setminus (C \cup F_1 \cup \dots \cup F_r)$$

where T_1, \dots, T_r are fibers of π that contain singular points of the surface S , and F_1, \dots, F_r are fibers of the projection $\mathbb{F}_n \rightarrow \mathbb{P}^1$ over the points $\pi(T_1), \dots, \pi(T_r)$, respectively. Then S contains a cylinder by Lemma 2.1, which is a contradiction.

We see that π is birational. Let E be the π -exceptional curve. If Y contains a cylinder U , then it also contains a cylinder $U' \subset U$ such that $\pi(E) \not\subset U'$, so that its preimage in S is a cylinder as well. Thus, the surface Y does not contain cylinders. Then Y is singular and $K_Y^2 \leq 2$ by Theorem 2.8.

We see that $K_Y^2 = 2$ and π is a blowup of a smooth point in Y . If $\rho(Y) \geq 2$, then we can apply the same arguments to Y to show that it contains a cylinder. Hence, we conclude that $\rho(Y) = 1$. On the other hand, all singularities of the surface Y are ordinary double points by Theorem 2.8. We see that $K_Y^2 = 2$ and Y has 7 singular points of type A_1 . But such a surface does not exist. \square

Let us conclude this subsection by presenting few results about polar cylinders in arbitrary rational surfaces. To do this, fix an ample \mathbb{Q} -divisor H on the surface S . If S contains an H -polar cylinder, we say that H is *cylindrical*. The cylindrical ample \mathbb{Q} -divisors on S form a cone, which we denoted earlier by $\text{Amp}^{cyl}(S)$. To investigate this cone, consider the following number:

$$\mu_H = \inf \left\{ \lambda \in \mathbb{R}_{>0} \mid \text{the } \mathbb{R}\text{-divisor } K_S + \lambda H \text{ is pseudo-effective} \right\}.$$

Remark 2.11. The number μ_H is known as the Fujita invariant of the divisor H , because it was implicitly used by Fujita in [73–76]. It plays an essential role in Manin’s conjecture (see [14, 98]).

Let Δ_H be the smallest extremal face of the Mori cone $\overline{\text{NE}}(S)$ that contains the divisor $K_S + \mu_H H$. Put $r_H = \dim(\Delta_H)$. Observe that $r_H = 0$ if and only if S is a del Pezzo surface and $\mu_H H \sim_{\mathbb{Q}} -K_S$.

Theorem 2.12 ([28]). *Suppose that S is smooth, $r_H + K_S^2 \leq 3$, and the self-intersection of every smooth rational curve in S is at least -1 . Then S does not contain H -polar cylinders.*

Note that if S is smooth del Pezzo surface, then the self-intersection of every smooth rational curve in S is at least -1 . Moreover, it follows from [54, Proposition 2.4] that this condition also holds if S is obtained by blowing up \mathbb{P}^2 at any number of points in general position.

Corollary 2.13 ([40]). *If S is a smooth del Pezzo surface and $r_H + K_S^2 \leq 3$, then $H \notin \text{Amp}^{cyl}(S)$.*

On the other hand, we have the following complimentary result:

Theorem 2.14 ([40, 150]). *Suppose that S is a smooth rational surface. If $K_S^2 \geq 4$, then*

$$\text{Amp}^{cyl}(S) = \text{Amp}(S).$$

If $K_S^2 = 3$ and $-K_S$ is not ample, then $\text{Amp}^{cyl}(S) = \text{Amp}(S)$. If $K_S^2 = 3$ and $-K_S$ is ample, then

$$\text{Amp}^{cyl}(S) = \text{Amp}(S) \setminus \mathbb{Q}_{>0}[-K_S].$$

If S is a smooth rational surface and $K_S^2 \leq 2$, then $\text{Amp}^{cyl}(S)$ is poorly understood (see [40]).

2.2. Absence of polar cylinders. Now, we show that smooth del Pezzo surfaces of degree ≤ 3 does not contain any anticanonical polar cylinders, which is one way implication of Corollary 2.9. For singular del Pezzo surfaces of degree ≤ 2 with types of singular points listed in Theorem 2.8, the same implication can be verified in a similar way (see [39] for the details).

Let S be a smooth del Pezzo surface of degree $K_S^2 = d \leq 3$, and let D be an effective \mathbb{Q} -divisor on the surface S , i.e., we have

$$D = \sum_{i=1}^r a_i C_i,$$

where every C_i is an irreducible curve on S , and every a_i is a non-negative rational number. Suppose that $D \sim_{\mathbb{Q}} -K_S$. If $d \in \{2, 3\}$, then each a_i does not exceed 1 by Lemmas A.9 and A.10. Similarly, if $d = 1$, we have

$$1 = d = K_S^2 = D \cdot (-K_S) = \sum_{i=1}^r a_i C_i \cdot (-K_S) \geq a_i C_i \cdot (-K_S),$$

which immediately implies that $a_i \leq 1$ for each i .

Theorem 2.15. *Let P be a point in S . Suppose that the log pair (S, D) is not log canonical at P . Then there exists a curve $T \in |-K_S|$ such that*

- the curve T is singular at P ;
- the log pair (S, T) is not log canonical at P ;
- $\text{Supp}(T) \subseteq \text{Supp}(D)$.

Proof. We consider the cases $d = 1$, $d = 2$ and $d = 3$ separately. See the proof of [38, Theorem 1.12] for an alternative proof in the case $d = 3$.

Suppose that $K_S^2 = 1$. Let C be a curve in $|-K_S|$ that passes through P . Then C is irreducible. If C is not contained in the support of D , then it follows from Lemma A.3 that

$$1 = d \geq K_S^2 = D \cdot C \geq \text{mult}_P(D) > 1.$$

This shows that $C \subset \text{Supp}(D)$. If (S, C) is not log canonical at P , then we can put $T = C$ and we are done. Thus, we may assume that (S, C) is log canonical at P . Then Remark A.2 implies the existence of an effective \mathbb{Q} -divisor D' such that $D' \sim_{\mathbb{Q}} -K_S$, the curve C is not contained in the support of D' , and (S, D') is not log canonical at P . Now Lemma A.3 implies that

$$1 = d \geq K_S^2 = D' \cdot C \geq \text{mult}_P(D') > 1,$$

which is absurd.

Now, we suppose that $K_S^2 = 2$. In this case there exists a double cover $\tau: S \rightarrow \mathbb{P}^2$ branched over a smooth quartic curve C . Moreover, we have

$$D \sim_{\mathbb{Q}} -K_S \sim \tau^*(L),$$

where L is a line in \mathbb{P}^2 . By Lemma A.10, we have $\tau(P) \in C$. Now let us choose L to be the tangent line to C at the point $\tau(P)$, and let R be the curve in $|-K_S|$ such that $\tau(R) = L$. Then $\text{mult}_P(R) = 2$. If R is irreducible and is not contained in the support of D , then Lemma A.3 gives

$$2 = d \geq K_S^2 = D \cdot R \geq \text{mult}_P(D) \text{mult}_P(R) \geq 2 \text{mult}_P(D) > 2.$$

Note that either R is irreducible or R consists of two (-1) -curves that both pass through P . Therefore, if one component of the curve R is not contained in the support of the divisor D , then we obtain a contradiction in a similar way by intersecting D with this irreducible component. Thus, we may assume that all irreducible component of the curve R are contained in $\text{Supp}(D)$. Now we can use Remark A.2 as in the case $d = 1$ to conclude that (S, R) is not log canonical at P . Hence, we can let $T = R$.

Finally, we suppose that $K_S^2 = 3$. Then S is a smooth cubic surface in \mathbb{P}^3 , and $-K_S$ is rationally equivalent to its hyperplane section. Let T_P be the intersection of the surface S with the hyperplane that is tangent to S at the point P . Then T_P is a reduced cubic curve that is singular at P . If (S, T_P) is not log canonical at P and $\text{Supp}(T_P) \subseteq \text{Supp}(D)$, we can let $T = T_P$ and we are done. Therefore, we may assume that at least one of the following two conditions hold:

- (1) the log pair (S, T_P) is log canonical at P ;
- (2) $\text{Supp}(D)$ does not contain at least one irreducible components of the curve T_P .

To obtain a contradiction, we may assume by Remark A.2 that at least one irreducible component of the curve T_P is not contained in $\text{Supp}(D)$.

If L_P is a line that passes through P , then $L_P \subseteq \text{Supp}(D)$, since otherwise we would get

$$1 \geq D \cdot L_P \geq \text{mult}_P(D) \text{mult}_P(L_P) \geq \text{mult}_P(D) > 1$$

by Lemma A.3. Thus, we see that $\text{mult}_P(T_P) = 2$.

Let $f: \tilde{S} \rightarrow S$ be the blowup of the point P , let E be the exceptional curve of the blowup f , and let \tilde{D} be the proper transform on \tilde{S} of the \mathbb{Q} -divisor D . Then $\text{mult}_P(D) > 1$ by Lemma A.3. Moreover, it follows from Lemma A.5 that the log pair

$$(\tilde{S}, \tilde{D} + (\text{mult}_P(D) - 1)E)$$

is not log canonical at some point $Q \in E$. Moreover, there is a commutative diagram

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{g} & \bar{S} \\ f \downarrow & & \downarrow h \\ S & \xrightarrow{\psi} & \mathbb{P}^2, \end{array}$$

where ψ is a projection from P , the morphism g is a contraction of the proper transforms of all lines in S that pass through P , and h is a double cover branched over a quartic curve. This quartic curve has at most two ordinary double points, because $\text{mult}_P(T_P) \neq 3$.

Let \tilde{T}_P be the proper transform on \tilde{S} of the curve T_P . Then $Q \in E \cap \tilde{T}_P$ by Lemma A.10.

Note that T_P is one of the following curves: an irreducible cubic curve, a union of a conic and a line, a union of three lines. Let us consider these cases separately.

Suppose that T_P is a union of a conic and a line, so that $T_P = L_P + C_P$, where L_P is a line, and C_P is an irreducible conic. Then $L_P \subset \text{Supp}(D)$, so that C_P is not contained in $\text{Supp}(D)$. Thus, we write $D = aL_P + \Omega$, where $a \in \mathbb{Q}_{>0}$, and Ω is an effective \mathbb{Q} -divisor on S whose support contains none of the curves L_P and C_P . Put $m = \text{mult}_P(\Omega)$. Then $\text{mult}_P(D) = m + a$ and

$$2 - 2a = \Omega \cdot C_P \geq m,$$

which gives $m + 2a \leq 2$. Similarly, we obtain $1 + a \geq m$ by using

$$1 + a = L_P \cdot D = \Omega \cdot L_P \geq m.$$

Denote by \tilde{C}_P the proper transform of the conic C_P on the surface \tilde{S} , denote by \tilde{L}_P the proper transform of the line L_P on the surface \tilde{S} , and denote by $\tilde{\Omega}$ the proper transform of the divisor Ω on the surface \tilde{S} . Put $\tilde{m} = \text{mult}_Q(\tilde{\Omega})$. Then the log pair

$$(2.16) \quad (\tilde{S}, a\tilde{L}_P + \tilde{\Omega} + (m + a - 1)E)$$

is not log canonical at P . Now, applying Lemma A.3 to this log pair, we obtain $2a + m + \tilde{m} > 2$. On the other hand, if $Q \in \tilde{C}_P$, then

$$2 - 2a - m = \tilde{\Omega} \cdot \tilde{C}_P \geq \tilde{m},$$

so that $Q \notin \tilde{C}_P$. Since $Q \in \tilde{T}_P$, we see that $Q \in \tilde{L}_P$. Then we have

$$1 + a - m = \tilde{\Omega} \cdot \tilde{L}_P \geq \tilde{m},$$

so that $2 \geq 1 + a \geq m + \tilde{m} \geq 2\tilde{m}$, which gives $\tilde{m} \leq 1$. Thus, we can apply Theorem A.8 to the log pair (2.16) at the point Q . This gives

$$m = \tilde{\Omega} \cdot E \geq (\tilde{\Omega} \cdot E)_Q > 2(2 - a - m)$$

or

$$1 + a - m = \tilde{\Omega} \cdot \tilde{L} \geq (\tilde{\Omega} \cdot \tilde{L})_Q > 2(1 - a),$$

so that we get $3a + m > 3$ or $2a + m > 2$, which is impossible since $a \leq 1$ and $m + 2a \leq 2$.

Therefore, we conclude that the curve T_P is a union of three lines. Hence, we have $T_P = L_1 + L_2 + L_3$, where L_1, L_2, L_3 are lines in S such that $P = L_1 \cap L_2$ and $P \notin L_3$. Then $L_1 \subset \text{Supp}(D) \supset L_2$. Therefore, we can write $D = a_1 L_1 + a_2 L_2 + \Delta$, where a_1 and a_2 are some positive rational numbers, and Δ is an effective \mathbb{Q} -divisor whose support does not contain L_1 and L_2 . Put $\mathbf{m} = \text{mult}_P(\Delta)$. Then

$$\mathbf{m} \leq \Delta \cdot L_1 = (H - a_1 L_1 - a_2 L_2) \cdot L_1 = 1 + a_1 - a_2,$$

because $L_1 \cdot L_2 = 1$ and $L_1^2 = -1$ on the surface S . Similarly, we see that

$$\mathbf{m} \leq \Delta \cdot L_2 = (H - a_1 L_1 - a_2 L_2) \cdot L_2 = 1 - a_1 + a_2.$$

This gives $\mathbf{m} \leq 1$. Thus, we can apply Theorem A.8 to the log pair (S, D) at the point P . Then

$$1 + a_1 - a_2 = \Delta \cdot L_1 \geq (\Delta \cdot L_1)_P > 2(1 - a_2)$$

or

$$1 - a_1 + a_2 = \Delta \cdot L_2 \geq (\Delta \cdot L_2)_P > 2(1 - a_1),$$

which implies that $a_1 + a_2 > 1$. On the other hand, we have

$$0 \leq \Delta \cdot L_3 = (H - a_1 L_1 - a_2 L_2) \cdot L_3 = 1 - a_1 - a_2,$$

which implies that $a_1 + a_2 \leq 1$. The obtained contradiction completes the solution. \square

We now claim that a smooth del Pezzo surface of degree $d \leq 3$ cannot contain a $(-K_S)$ -cylinder. If $d \leq 2$, the claim is [126, Proposition 5.1]. Similarly, if $d = 3$, then the claim is [38, Theorem 1.7]. Let us show how to derive the claim from Theorem 2.15 and Remark 2.3.

Suppose that S contains a $(-K_S)$ -polar cylinder U . Then

$$S \setminus U = C_1 \cup \cdots \cup C_n$$

for some irreducible curves C_1, \dots, C_n in S , and there are positive rational numbers $\lambda_1, \dots, \lambda_n$ such that

$$\sum_{i=1}^n \lambda_i C_i \sim_{\mathbb{Q}} -K_S.$$

Put $D = \lambda_1 C_1 + \cdots + \lambda_n C_n$. Then (S, D) is not log canonical at some point $P \in S$ by Remark 2.3. Hence, by Theorem 2.15, there exists a curve $T \in |-K_S|$ such that

- the log pair (S, T) is not log canonical at P ;
- and $\text{Supp}(T) \subseteq \text{Supp}(D)$.

Then $D \neq T$, because $n > 3$ by (2.2), and T does not have more than $d \leq 3$ irreducible components. Thus, there exists a rational number $\mu > 0$ such that $(1 + \mu)D - \mu T$ is effective, and its support does not contain at least one irreducible component of the curve T . Then $(S, (1 + \mu)D - \mu T)$ is not log canonical at P by Remark 2.3, which contradicts to Theorem 2.15, since

$$(1 + \mu)D - \mu T \sim_{\mathbb{Q}} -K_S.$$

2.3. Construction of polar cylinders. Now, we show how to construct anticanonical polar cylinders in singular del Pezzo surfaces with Du Val singularities. We start with

Lemma 2.17 ([123, Theorem 3.19]). *Let S is a smooth del Pezzo surface. Suppose that $K_S^2 \geq 4$. Then the surface S contains a $(-K_S)$ -polar cylinder.*

Proof. We may assume that $S \neq \mathbb{P}^1 \times \mathbb{P}^1$. Then there exists a birational map $\sigma: S \rightarrow \mathbb{P}^2$ that blows up $k \leq 5$ distinct points. Let E_1, \dots, E_k be the σ -exceptional curves, let C be an irreducible conic in \mathbb{P}^2 that contains all points $\sigma(E_1), \dots, \sigma(E_k)$, and let L be a line in \mathbb{P}^2 that is tangent to the conic C at some point that is different from $\sigma(E_1), \dots, \sigma(E_k)$. Denote by \tilde{C} and \tilde{L} the proper transforms on S of the curves C and L , respectively. Then

$$-K_S \sim \sigma^*(-K_{\mathbb{P}^2}) - \sum_{i=1}^k E_i \sim_{\mathbb{Q}} (1 + \varepsilon)\tilde{C} + (1 - 2\varepsilon)\tilde{L} + \varepsilon \sum_{i=1}^k E_i$$


for every positive $\varepsilon < \frac{1}{2}$. On the other hand, we have

$$S \setminus (\tilde{C} \cup \tilde{L} \cup E_1 \cup \dots \cup E_k) \cong \mathbb{P}^2 \setminus (C \cup L) \cong (\mathbb{A}^1 \setminus \{0\}) \times \mathbb{A}^1,$$

so that the surface S contains a $(-K_S)$ -polar cylinder. \square

Now let us present an example of a singular del Pezzo surface of degree 2 that has one singular point of type A_2 and contains an anticanonical polar cylinder.

Example 2.18. Let $h: \hat{S} \rightarrow \mathbb{P}^2$ be a composition of 10 blowups, let E_1, \dots, E_{10} be the exceptional curves of the birational morphism h , let L_1 and L_2 be two distinct lines in \mathbb{P}^2 , and let \hat{L}_1 and \hat{L}_2 be their proper transforms on \hat{S} , respectively. Now, let us choose h such that the intersections of these twelve curves are depicted as follows:



Note that the 10 blowups are arranged in the order indicated by the indices of their exceptional curves E_i . To describe the intersection form of the curves $\hat{L}_1, \hat{L}_2, E_1, \dots, E_{10}$, observe that

$$\hat{L}_1^2 = -1, \quad \hat{L}_2^2 = -5, \quad E_1^2 = -3, \quad E_2^2 = -2, \quad E_3^2 = -2, \quad E_4^2 = \dots = E_{10}^2 = -1.$$

Let $g: \hat{S} \rightarrow \tilde{S}$ be the contraction of the curves \hat{L}_1, E_2, E_3 , and let $\tilde{L}_2, \tilde{E}_1, \tilde{E}_4, \dots, \tilde{E}_{10}$ be the proper transforms on \tilde{S} of the curves $\hat{L}_2, \hat{E}_1, \hat{E}_4, \dots, \hat{E}_{10}$, respectively. Then \tilde{S} is smooth and $K_{\tilde{S}}^2 = 2$.

Moreover, the divisor $-K_{\tilde{S}}$ is nef. To show this, fix an arbitrary positive rational number $\epsilon < \frac{1}{3}$, let $D_{\tilde{S}}$ be the following \mathbb{Q} -divisor:

$$(2-\epsilon)\hat{L}_1 + (1+\epsilon)\hat{L}_2 + (1-\epsilon)E_1 + (2-2\epsilon)E_2 + (2-3\epsilon)E_3 + (1-3\epsilon)E_4 + \epsilon(E_5 + E_6 + E_7 + E_8 + E_9 + E_{10}),$$

and denote by $D_{\tilde{S}}$ its proper transform on \tilde{S} . Then $D_{\tilde{S}}$ is effective, $D_{\tilde{S}} \sim_{\mathbb{Q}} -K_{\tilde{S}}$ and $D_{\tilde{S}} \sim_{\mathbb{Q}} -K_{\tilde{S}}$. Moreover, we have $\tilde{L}_2^2 = \tilde{E}_1 = -2$, $\tilde{E}_4 = 0$ and $\tilde{E}_5^2 = \dots = \tilde{E}_{10}^2 = -1$, so that

$$-K_{\tilde{S}} \cdot \tilde{L}_2 = -K_{\tilde{S}} \cdot \tilde{E}_1 = 0, -K_{\tilde{S}} \cdot \tilde{E}_4 = 2, -K_{\tilde{S}} \cdot \tilde{E}_5 = \dots = -K_{\tilde{S}} \cdot \tilde{E}_{10} = 1.$$

This shows that $-K_{\tilde{S}}$ is nef. Moreover, we also see that \tilde{L}_2 and \tilde{E}_1 are the only (-2) -curves in \tilde{S} . Let $f: \tilde{S} \rightarrow S$ be the birational contraction of these two (-2) -curves. Then S is a del Pezzo surface with one singular point of type A_2 such that $K_S^2 = 2$. Let $D_S = f \circ g(D_{\tilde{S}})$. Then $D_S \sim_{\mathbb{Q}} -K_S$ and

$$S \setminus \text{Supp}(D_S) \cong \mathbb{P}^2 \setminus \text{Supp}(D_{\mathbb{P}^2}) \cong \mathbb{A}^1 \times (\mathbb{A}^1 \setminus \{0\}),$$

so that S contains $(-K_S)$ -polar cylinder.

One can use the construction in Example 2.18 to construct an anticanonical polar cylinder in *every* del Pezzo surface of degree 2 that has a single singular point of type A_2 (see [39, §4.3]). Similarly, we can prove the existence part of Theorem 2.8. However, there is an alternative proof, which is more algebraic. Let us describe it following [33].

Let S be a singular del Pezzo surface of degree $K_S^2 \leq 3$ that has at most Du Val singularities, and let P be its singular point. Suppose, in addition, that the following conditions hold:

- the singular point P is not of type A_1 if $K_S^2 = 2$;
- the singular point is not of types A_1, A_2, A_3, D_4 if $K_S^2 = 1$.

Now, let us prove that S contains a $(-K_S)$ -polar cylinder (cf. Theorem 2.8).

Denote by \mathbb{P} the three-dimensional weighted projective space in which S sits as a hypersurface. Note that $\mathbb{P} = \mathbb{P}^3$ (respectively, $\mathbb{P}(1, 1, 1, 2)$, $\mathbb{P}(1, 1, 2, 3)$) if $K_S^2 = 3$ (respectively, $K_S^2 = 2$, $K_S^2 = 1$). For the quasi-homogenous coordinate system for \mathbb{P} , we use $[x : y : z : w]$. By a coordinate change, we may assume that $P = [1 : 0 : 0 : 0]$. Then the equation of S can be described as follows:

- if $K_S^2 = 3$, then S is given by

$$(2.19) \quad x f_2(y, z, w) + f_3(y, z, w) = 0,$$

where f_2 and f_3 are polynomials of degrees 2 and 3, respectively;

- if $K_S^2 = 2$, then S is given by

$$(2.20) \quad w^2 + x(ayw + f_3(y, z)) + f_4(y, z) = 0,$$

where f_3 and f_4 are polynomials of degrees 3 and 4, respectively, and $a \in \mathbb{k}$;

- if $K_S^2 = 1$, then S is given by

$$(2.21) \quad w^2 + x(ay^2w + f_5(y, z)) + f_6(y, z) = 0$$

or

$$(2.22) \quad w^2 + x(zw + f_5(y, z)) + f_6(y, z) = 0,$$

where f_5 and f_6 are polynomials of degrees 5 and 6, respectively, and $a \in \mathbb{k}$.

Let Π be the hyperplane in \mathbb{P} defined by $x = 0$, and let $\pi: S \dashrightarrow \Pi$ be the map given by

$$[x : y : z : w] = [0 : y : z : w].$$

The hyperplane Π is isomorphic to \mathbb{P}^2 , $\mathbb{P}(1, 1, 2)$, $\mathbb{P}(1, 2, 3)$ according to $K_S^2 = 3, 2, 1$, respectively. We denote by $g(y, z, w)$ the coefficient of x in each of equations (2.19), (2.20), (2.21) and (2.22). Namely, if $K_S^2 = 3$, then $g(y, z, w) = f_2(y, z, w)$. Similarly, if $K_S^2 = 2$, then

$$g(y, z, w) = ayw + f_3(y, z).$$

Finally, if $K_S^2 = 1$, then $g(y, z, w) = zw + f_5(y, z)$ or

$$g(y, z, w) = ay^2w + f_5(y, z).$$

Let D be the divisor on S that is cut out by $g(y, z, w) = 0$. If $K_S^2 = 3$, then D consists of the lines that contains P . There are at most six such lines and they are defined in \mathbb{P}^3 by

$$\begin{cases} g(y, z, w) = 0, \\ f_3(y, z, w) = 0. \end{cases}$$

Similarly, if $K_S^2 = 2$, then the divisor D consists of at most six curves passing through the point P . They are defined in $\mathbb{P}(1, 1, 1, 2)$ by

$$\begin{cases} g(y, z, w) = 0, \\ w^2 + f_4(y, z) = 0. \end{cases}$$

Finally, if $K_S^2 = 1$, then the divisor D consists of at most five curves passing through the point P , which are defined in $\mathbb{P}(1, 1, 2, 3)$ by

$$\begin{cases} g(y, z, w) = 0, \\ w^2 + f_6(y, z) = 0. \end{cases}$$

In each case, the number of curves in D is the same as the number of points determined by the corresponding system of equations in Π . We denote these curves by L_1, \dots, L_r in each case. The map π contracts each curve L_i to a point on Π .

The equations (2.19), (2.20), (2.21) and (2.22) immediately imply that π is a birational map. Moreover, it induces an isomorphism

$$\tilde{\pi}: S \setminus (L_1 \cup \dots \cup L_r) \cong \text{Im}(\tilde{\pi}) \subset \Pi.$$

Let \mathcal{C} be the curve on Π defined by $g(y, z, w) = 0$. Then \mathcal{C} can be reducible or non-reduced.

Lemma 2.23. *Suppose that $K_S^2 = 3$. Then there is a hyperplane section H of the surface S such that the complement $S \setminus (H \cup L_1 \cup \dots \cup L_r)$ is a $(-K_S)$ -polar cylinder.*

Proof. Observe that $\text{Im}(\tilde{\pi}) = \Pi \setminus \mathcal{C}$. Let $\varphi: \bar{S} \rightarrow S$ be the blowup of the point P . Then there exists a commutative diagram

$$\begin{array}{ccc} & \bar{S} & \\ \varphi \swarrow & & \searrow \psi \\ S & \xrightarrow{\pi} & \Pi \end{array}$$

where ψ is the birational morphism that contracts the proper transforms of the lines L_1, \dots, L_r . Let E be the exceptional curve of the blowup φ . Then $\psi(E) = \mathcal{C}$, and \mathcal{C} contains each point $\pi(L_i)$.

If P is an ordinary double point of the cubic surface S , then the curve \mathcal{C} is a smooth conic. Similarly, if P is a singular point of type A_n for $n \geq 2$, then \mathcal{C} splits as a union of two distinct lines. Finally, if P is either of type D_n or of type E_6 , then \mathcal{C} is a double line.

If \mathcal{C} is smooth, let ℓ be a general line in Π that is tangent to \mathcal{C} . If \mathcal{C} is singular, let ℓ be a general line in Π that passes through a singular point of the conic \mathcal{C} . By a suitable coordinate change, we may assume that ℓ is defined by $x = y = 0$. Let H be the curve in S cut out by $y = 0$. Then

$$S \setminus (H \cup L_1 \cup \cdots \cup L_r) \cong \Pi \setminus (\mathcal{C} \cup \ell) \cong \begin{cases} (\mathbb{A}^1 \setminus \{0, 1\}) \times \mathbb{A}^1 & \text{if } \mathcal{C} \text{ is a union of two distinct lines,} \\ (\mathbb{A}^1 \setminus \{0\}) \times \mathbb{A}^1 & \text{otherwise.} \end{cases}$$

Therefore, $S \setminus (H \cup L_1 \cup \cdots \cup L_r)$ is a cylinder. But $H + D \sim -3K_S$ and $L_1 \cup \cdots \cup L_r = \text{Supp}(D)$. Thus, the complement $S \setminus (H \cup L_1 \cup \cdots \cup L_r)$ is a $(-K_S)$ -polar cylinder. \square

To deal with the cases $K_S^2 = 1$ and $K_S^2 = 2$, let ℓ_y be the curve in \mathbb{P} that is given by $x = y = 0$, and let H_y be the curve in the surface S that is cut out by $y = 0$.

Lemma 2.24. *If $K_S^2 = 2$ or if $K_S^2 = 1$ and the surface S is defined by equation (2.21), then the complement $S \setminus (H_y \cup L_1 \cup \cdots \cup L_r)$ is a $(-K_S)$ -polar cylinder.*

Proof. Observe that the morphism $\tilde{\pi}$ gives an isomorphism $S \setminus (H_y \cup L_1 \cup \cdots \cup L_r) \cong \Pi \setminus (\mathcal{C} \cup \ell_y)$. But π maps $S \setminus H_y$ onto $\Pi \setminus \ell_y \cong \mathbb{A}^2$. Thus, if $K_S^2 = 2$, then $S \setminus (H_y \cup L_1 \cup \cdots \cup L_r)$ is isomorphic to the complement in \mathbb{A}^2 of the curve defined by

$$aw + f_3(1, z) = 0.$$

Similarly, if $K_S^2 = 1$ and S is defined by (2.21), then $S \setminus (H_y \cup L_1 \cup \cdots \cup L_r)$ is isomorphic to the complement in \mathbb{A}^2 of the curve defined by

$$aw + f_5(1, z) = 0.$$

Therefore, in both cases, the complement $S \setminus (H_y \cup L_1 \cup \cdots \cup L_r)$ is a cylinder. Now, arguing as in the proof of Lemma 2.23, we see that $S \setminus (H_y \cup L_1 \cup \cdots \cup L_r)$ is a $(-K_S)$ -polar cylinder. \square

Finally, to deal with the remaining case, let ℓ_z be the curve in \mathbb{P} that is given by $x = z = 0$, and let H_z be the hyperplane section of S that is cut by $z = 0$.

Lemma 2.25. *Suppose that $K_S^2 = 1$ and the del Pezzo surface S is given by equation (2.22). Then the complement $S \setminus (H_z \cup L_1 \cup \cdots \cup L_r)$ is a $(-K_S)$ -polar cylinder.*

Proof. Observe that the morphism $\tilde{\pi}$ gives an isomorphism $S \setminus (H_z \cup L_1 \cup \cdots \cup L_r) \cong \Pi \setminus (\mathcal{C} \cup \ell_z)$. But π maps $S \setminus H_z$ onto $\Pi \setminus \ell_z$. Then $\Pi \setminus (\mathcal{C} \cup \ell_z)$ is the complement of the curve defined by

$$w + f_5(y, 1) = 0$$

in $\Pi \setminus \ell_z \cong \mathbb{A}^2/\mu_2$, where the μ_2 -action is given by $(y, w) \mapsto (-y, -w)$.

Since $f_5(y, 1)$ is an odd polynomial in y , the isomorphism $\mathbb{A}^2 \rightarrow \mathbb{A}^2$ defined by

$$(y, w) \mapsto (y, w + f_5(y, 1))$$

is μ_2 -equivariant and gives an isomorphism between the complement $\Pi \setminus (\mathcal{C} \cup \ell_z)$ and the complement in \mathbb{A}^2/μ_2 of the image of the curve defined by $w = 0$, which is isomorphic to $\mathbb{A}^1 \setminus \{0\} \times \mathbb{A}^1$.

We see that $S \setminus (H_z \cup L_1 \cup \cdots \cup L_r)$ is a cylinder. Now, arguing as in the proof of Lemma 2.23, we conclude that $S \setminus (H_z \cup L_1 \cup \cdots \cup L_r)$ is a $(-K_S)$ -polar cylinder. \square

3. CYLINDERS IN HIGHER-DIMENSIONAL VARIETIES

In this section, we describe known results about cylinders in smooth Fano threefolds and fourfolds, and varieties fibred into del Pezzo surfaces. Let us say few words about Fano varieties [108, 114, 163].

Let V be a smooth Fano variety of dimension $n \geq 3$. The number $(-K_V)^n$ is known as the degree of the Fano variety V . Put

$$\iota(V) = \max \left\{ t \in \mathbb{N} \mid -K_V \sim tH \text{ for } H \in \text{Pic}(V) \right\}.$$

Then $\iota(V)$ is known as the (Fano) index of the variety V . It is well known that $1 \leq \iota(V) \leq n + 1$. Moreover, one has

$$\iota(V) = n + 1 \iff V \cong \mathbb{P}^n.$$

Similarly, we have $\iota(V) = n$ if and only if V is a quadric (see [114, 128]).

Remark 3.1 ([70–72, 114]). Suppose that $\iota(V) = n - 1$. Then

$$-K_V \sim (n - 1)H$$

for some ample divisor $H \in \text{Pic}(V)$. In this case, the variety V is usually called a *del Pezzo variety*. If $\rho(V) = 1$, then there are just the following possibilities:

- $H^n = 1$ and $V = V_6$ is a weighted hypersurface in $\mathbb{P}(1^n, 2, 3)$ of degree 6;
- $H^n = 2$ and $V = V_4$ is a weighted hypersurface in $\mathbb{P}(1^{n+1}, 2)$ of degree 4;
- $H^n = 3$ and $V = V_3$ is a cubic hypersurface in \mathbb{P}^{n+1} ;
- $H^n = 4$ and $V = V_{2,2}$ is a complete intersection of two quadrics in \mathbb{P}^{n+2} ;
- $H^n = 5$, $n \in \{3, 4, 5, 6\}$ and V is described in Example 1.31.

If $\dim(V) = 3$ and $\rho(V) = 1$, then the values of the Hodge number $h^{1,2}(V)$ are given in

H^3	1	2	3	4	5
$h^{1,2}(V)$	21	10	5	2	0

Let us prove cylindricity of any higher-dimensional smooth intersection of two quadrics.

Lemma 3.2 ([123]). *Let V be a smooth complete intersection of two quadric hypersurfaces in \mathbb{P}^{n+2} . Then V is cylindrical.*

Proof. Let ℓ be a line in V , let D be an irreducible divisor in X swept out by lines meeting ℓ , let $\sigma: \tilde{V} \rightarrow V$ be the blowup of the line ℓ , let E be its exceptional divisor, and let \tilde{D} be the proper transform on \tilde{V} of the divisor D . There exists the following commutative diagram:

$$\begin{array}{ccc} & \tilde{V} & \\ \sigma \swarrow & & \searrow \varphi \\ V & \overset{\psi}{\dashrightarrow} & \mathbb{P}^n \end{array}$$

where φ is a birational morphism that contracts \tilde{D} , and ψ is the projection from ℓ . Thus, we have

$$V \setminus D \cong \mathbb{P}^n \setminus \varphi(E).$$

But $\varphi(E)$ is a quadric that contains a one-parameter family of linear subspaces of dimension $n - 2$. Hence, this quadric is singular, so that $\mathbb{P}^n \setminus \varphi(E)$ contains a cylinder. \square

Smooth Fano varieties of dimension $n \geq 3$ and index $n - 2$ are known as *Fano–Mukai varieties*. If V is a Fano–Mukai variety and $H \in \text{Pic}(V)$ such that $-K_V \sim (n - 2)H$, then the number

$$g(V) = \frac{1}{2}H^n + 1$$

is integral and is called the *genus* of the Fano–Mukai variety V . The possible values of the genus are given in the following table:

$g(V)$	$2 \leq g(V) \leq 5$	6	7	8	9	10	12
$\dim(V)$	any	≤ 6	≤ 10	≤ 8	≤ 6	≤ 5	3

Moreover, the following result has been recently proved in [142].

Theorem 3.3. *Let V be a smooth Fano–Mukai variety such that $\rho(V) = 1$ and $g(V) \in \{7, 8, 9, 10\}$. Suppose that $\dim(V) \geq 5$. Then V is cylindrical.*

In Subsection 3.1, we will outline several known results about cylindrical smooth Fano threefolds. Then, in Subsection 3.2, we will present constructions of cylinders in some smooth Fano fourfolds. In particular, we will explain how to prove the following result:

Theorem 3.4. *For every $g \in \{7, 8, 9, 10\}$, there is a cylindrical Fano–Mukai fourfold of genus g .*

Finally, in Subsection 3.3, we will present results about cylinders in Mori fibrations.

3.1. Cylindrical Fano threefolds. Let X be a smooth Fano variety that has dimension three. Then X belongs to one of 105 families, which have been explicitly described in [106–108, 110, 157–160]. Their automorphism groups have been studied in [42, 140, 143, 164, 182]. In particular, we have

Theorem 3.5. *Let X be a smooth Fano threefold such that $\rho(X) = 1$ and $\text{Aut}(X)$ is infinite. Then X and $\text{Aut}(X)$ can be described as follows:*

- (1) $X = \mathbb{P}^3$ and $\text{Aut}(X) \cong \text{PGL}_4(\mathbb{k})$;
- (2) X is a smooth quadric in \mathbb{P}^4 and $\text{Aut}(X) \cong \text{PSO}_5(\mathbb{k})$;
- (3) X is the quintic del Pezzo threefold described in Example 1.31 and $\text{Aut}(X) \cong \text{PGL}_2(\mathbb{k})$;
- (4) X is one of the following Fano threefolds in \mathbb{P}^{13} of degree 22 and genus 12:
 - (a) the Mukai–Umemura threefold X_{22}^{mu} with $\text{Aut}(X_{22}^{\text{mu}}) \cong \text{PGL}_2(\mathbb{k})$;
 - (b) the unique special threefold X_{22}^{a} with $\text{Aut}(X_{22}^{\text{a}}) \cong \mathbb{G}_a \rtimes \mu_4$;
 - (c) a threefold X_{22}^{m} in one-parameter family with $\text{Aut}(X_{22}^{\text{m}}) \cong \mathbb{G}_m \rtimes \mu_2$.

Before we describe some cylindrical smooth Fano threefolds, observe that we have the following implications:

$$X \text{ contains } (-K_X)\text{-polar cylinder} \implies X \text{ is cylindrical} \implies X \text{ is rational.}$$

Moreover, the rationality problem for smooth Fano threefolds is *almost* completely solved (see [114]). In particular, for *general* member of every family, we know whether it is rational or irrational. It is expected that the same answer holds for *every* smooth member in each family.

If $\iota(X) \geq 3$, then either $X \cong \mathbb{P}^3$ or X is a smooth quadric in \mathbb{P}^4 , so that X is cylindrical.

If $\iota(X) = 2$, then $-K_X \sim 2H$ for $H \in \text{Pic}(X)$, and we have the following possibilities:

- $H^3 = 1$ and $X = V_1$ is a sextic hypersurface in $\mathbb{P}(1, 1, 1, 2, 3)$;
- $H^3 = 2$ and $X = V_2$ is quartic hypersurface in $\mathbb{P}(1, 1, 1, 1, 2)$;
- $H^3 = 3$ and $X = V_3$ is a cubic hypersurface in \mathbb{P}^4 ;
- $H^3 = 4$ and $X = V_4$ is an intersection of two quadrics in \mathbb{P}^5 ;
- $H^3 = 5$ and $X = V_5$ is the quintic del Pezzo threefold described in Example 1.31;
- $H^3 = 6$ and X is a divisor in $\mathbb{P}^2 \times \mathbb{P}^2$ of degree $(1, 1)$;

- $H^3 = 6$ and $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$;
- $H^3 = 7$ and $X = V_7$ is a blowup of \mathbb{P}^3 at a point.

In this case, if $H^3 \leq 3$, then X is irrational (see [5, 43, 50, 88, 89, 213]), so that it is not cylindrical. On the other hand, if $H^3 \geq 4$, then X contains a $(-K_X)$ -polar cylinder. Indeed, if $H^3 = 4$, this follows from Lemma 3.2. If $H^3 \geq 6$, this is obvious. Finally, if $H^3 = 5$, this follows from

Theorem 3.6. *Let V_5 be the quintic del Pezzo threefold in \mathbb{P}^6 that is described in Example 1.31. Then V_5 contains a hyperplane section H such that $V_5 \setminus H \cong \mathbb{A}^3$.*

Proof. Let us give two constructions of the required hyperplane section. First, let L be a line in X . Let $\alpha: \tilde{V}_5 \rightarrow V_5$ be the blowup of the line L . Then we have the following commutative diagram:

$$\begin{array}{ccc} & \tilde{V}_5 & \\ \alpha \swarrow & & \searrow \beta \\ V_5 & & Q \end{array}$$

where Q is a smooth quadric in \mathbb{P}^4 , and β is the blowup of a twisted cubic curve C contained in Q . Let H_C be the unique hyperplane section of Q that contains C , and let H_L be the unique hyperplane section of V_5 that is singular along L . Then H_L is the proper transform of the β -exceptional surface, and H_C is the proper transform of the α -exceptional surface. Note that H_L is swept out by the lines that intersect the line L . Moreover, it follows from [114, 143] that

$$\mathcal{N}_{L/V_5} \cong \begin{cases} \mathcal{O}_L \oplus \mathcal{O}_L & L \text{ is a line of type } (0, 0), \\ \mathcal{O}_L(1) \oplus \mathcal{O}_L(-1) & L \text{ is a line of type } (1, -1). \end{cases}$$

The lines in V_5 are parameterized by \mathbb{P}^2 , and the lines of the type $(1, -1)$ are parameterized by a smooth conic in this plane (see [83, 108, 143]). Furthermore, the surface H_C is smooth if and only if L is a line of type $(1, -1)$. Thus, if we choose L to be a line of type $(1, -1)$ and put $H = H_L$, then $V_5 \setminus H \cong Q \setminus H_C \cong \mathbb{A}^3$ as required.

To present the second construction, let P be a point in V_5 . Recall that $\text{Aut}(V_5) \cong \text{PGL}_2(\mathbb{k})$. Moreover, it follows from [47, 83, 108, 143, 164] that $\text{Aut}(V_5)$ has exactly three orbits on V_5 :

- (1) a closed one-dimensional orbit \mathcal{C} , which is a twisted rational sextic curve in \mathbb{P}^6 ;
- (2) a two-dimensional orbit $\mathring{\mathcal{S}}$ whose closure is a surface $\mathcal{S} \sim -K_V$ which is singular along \mathcal{C} ;
- (3) an open orbit $V_5 \setminus \mathcal{S}$.

Furthermore, let k_P be the number of lines in V_5 passing through P . Then

$$k_P = \begin{cases} 1 & \text{if } P \in \mathcal{C}, \\ 2 & \text{if } \mathcal{S} \setminus \mathcal{C}, \\ 3 & \text{if } V_5 \setminus \mathcal{S}. \end{cases}$$

Observe also that \mathcal{S} is swept out by the lines of type $(1, -1)$.

Let $\sigma: \widehat{V}_5 \rightarrow V_5$ be the blowup of the point P . Then it follows from [82] that there exists the following Sarkisov link:

$$\begin{array}{ccc} & \widehat{V}_5 & \overset{\chi}{\dashrightarrow} \overline{V}_5 \\ \sigma \swarrow & & \searrow \varphi \\ V_5 & \overset{\psi}{\dashrightarrow} & \mathbb{P}^2 \end{array}$$

where χ is a composition of flops of the proper transforms of lines in V_5 that pass through P , the morphism φ is a \mathbb{P}^1 -bundle, and ψ is given by the linear system of hyperplane sections that are singular at the point P . Now we suppose that $P \in \mathcal{C}$.

Let E be the σ -exceptional surface, and let \overline{E} be its proper transform on the threefold \overline{V}_5 . Then \overline{E} is a del Pezzo surface of degree 6 with at most Du Val singularities, and its singular locus consists of one singular point of type A_2 . Moreover, the \mathbb{P}^1 -bundle $\varphi: \overline{V}_5 \rightarrow \mathbb{P}^2$ induces a birational map $\overline{E} \rightarrow \mathbb{P}^2$ that contracts a single curve $\Gamma \subset \overline{E}$ to a point in \mathbb{P}^2 .

Let \mathcal{L} be a line in \mathbb{P}^2 that passes through the point $\varphi(\Gamma)$, let \overline{H} be its preimage in \overline{V}_5 via φ , let \widehat{H} be its proper transform on \widehat{V}_5 , and let $H = \sigma(\widehat{H})$. Then

$$V_5 \setminus H \cong \overline{V}_5 \setminus (\overline{E} \cup \overline{H}),$$

and H is a hyperplane section of the threefold V_5 that is singular at P . Furthermore, one can show that the surface H is smooth away from P , and H has Du Val singularity of type A_4 at this point. Then the \mathbb{P}^1 -bundle φ induces a morphism $\overline{V}_5 \setminus (\overline{E} \cup \overline{H}) \rightarrow \mathbb{P}^2 \setminus \mathcal{L}$ that is an \mathbb{A}^1 -bundle over \mathbb{A}^2 . This implies that $V_5 \setminus H \cong \overline{V}_5 \setminus (\overline{E} \cup \overline{H}) \cong \mathbb{A}^3$ as required. \square

Now, we assume that $\iota(X) = 1$. This leaves us 95 families of smooth Fano threefolds [114, 157]. If $\rho(X) = 1$, $\iota(X) = 1$ and $g(X) \leq 6$, then we have the following possibilities:

- (1) $g(X) = 2$ and X is a sextic hypersurface in $\mathbb{P}(1^4, 3)$;
- (2) $g(X) = 3$ and X is an intersection of a quadric and a quartic in $\mathbb{P}(1^5, 2)$;
- (3) $g(X) = 4$ and X is a complete intersection of a quadric and a cubic in \mathbb{P}^5 ;
- (4) $g(X) = 5$ and X is a complete intersection of three quadrics in \mathbb{P}^6 ;
- (5) $g(X) = 6$ and X is a section of the cone in \mathbb{P}^8 over the smooth quintic del Pezzo fourfold described in Example 1.31 by a quadric and a hyperplane.

All these deformation families are irreducible. General members of the family (2) are smooth quartic hypersurfaces in \mathbb{P}^4 , and special members are double covers of the quadric threefold branched over octic surfaces. Similarly, general members of the family (5) are sections of the smooth quintic del Pezzo fourfold in \mathbb{P}^7 by quadrics, and special members are double covers of the smooth quintic del Pezzo threefold branched over anticanonical surfaces.

In the first two cases, the Fano threefold X is known to be irrational even if we allow mild isolated singularities [35, 105, 109, 113, 144, 152, 193, 206]. In the case (4), the threefold X is also irrational [15]. General threefolds of the families (3) and (5) are irrational [15, 99, 112, 115, 198], and every smooth member is also expected to be irrational. Therefore, in all these cases, the threefold X is either non-cylindrical or it is expected to be irrational and, thus, non-cylindrical.

Remark 3.7. Let V_5 be the smooth quintic del Pezzo threefold, see Example 1.31, and let $\pi: X \rightarrow V_5$ be a double cover branched over a surface $S \in |-K_{V_5}|$. If S has an isolated ordinary double point, then X is rationally connected [218], it is \mathbb{Q} -factorial [52], and it follows from [187] that there exists the following Sarkisov link:

$$\begin{array}{ccc} & \widetilde{X} & \\ \alpha \swarrow & & \searrow \beta \\ X & & \mathbb{P}^2, \end{array}$$

where α is the blow up of the singular point of X , and β is a standard conic bundle, whose discriminant curve has degree 6. Hence, in this case, the threefold X is irrational by [205, Theorem 10.2]. Now, using [131, Theorem IV.1.8.3], we conclude that X is also irrational if S is a very general surface in the linear system $|-K_{V_5}|$.

If $\rho(X) = 1$, $\iota(X) = 1$ and $g(X) \geq 7$, then $g(X) \in \{7, 8, 9, 10, 12\}$. Moreover, if $g(X) = 8$, then the threefold X is birational to a smooth cubic hypersurface in \mathbb{P}^4 (see, for example, [109, 114, 207]), so that it is irrational [50]. On the other hand, we know that X is rational if

$$g(X) \in \{7, 9, 10, 12\}.$$

In these cases, the divisor $-K_X$ is very ample, and $|-K_X|$ gives an embedding $X \hookrightarrow \mathbb{P}^{g(X)+1}$. Moreover, all the known constructions of cylinders in X use the *double projection* from a line in X (see [110]). Recall from [114, 125, 183] that X can contain two types of lines depending on their normal bundles. Namely, for a line $\ell \subset X$, we have the following two possibilities:

$$\mathcal{N}_{\ell/X} \cong \begin{cases} \mathcal{O}_{\ell} \oplus \mathcal{O}_{\ell}(-1) & \ell \text{ is of type } (0, -1), \\ \mathcal{O}_{\ell}(1) \oplus \mathcal{O}_{\ell}(-2) & \ell \text{ is of type } (1, -2). \end{cases}$$

If X is a sufficiently general member of one of these three families of smooth Fano threefolds, then X does not contain lines of type $(1, -2)$. Moreover, one can show that the threefolds containing lines of type $(1, -2)$ form a codimension one subset in the corresponding moduli spaces. On the other hand, we have the following result:

Theorem 3.8 ([125, Theorem 0.1]). *Suppose that $\rho(X) = 1$, $\iota(X) = 1$, and $g(X) = 9$ or $g(X) = 10$. If X contains a line of type $(1, -2)$, then X is cylindrical.*

Proof. Let ℓ be a line in the Fano threefold X , and let $\sigma: \tilde{X} \rightarrow X$ be the blowup of the line ℓ . Then it follows from [108, 110, 114, 184] that there is the Sarkisov link:

$$\begin{array}{ccc} & \tilde{X} & \xrightarrow{\chi} \hat{X} \\ \sigma \swarrow & & \searrow \varphi \\ X & & Y, \end{array}$$

where Y is a smooth Fano threefold described below, the morphism φ is the blowup of a smooth rational curve Γ , and χ is a composition of flops of the proper transforms of the lines that meet ℓ . Moreover, we have the following options:

- if $g(X) = 9$, then $Y = \mathbb{P}^3$, and Γ is a curve of degree 7 and genus 3;
- if $g(X) = 10$, then Y is a smooth quadric in \mathbb{P}^4 , and Γ is a curve of degree 7 and genus 2.

Let E be the σ -exceptional surface, let \hat{E} be its proper transform on \hat{X} , and let $\mathcal{S} = \varphi(\hat{E})$. Then \mathcal{S} is a (maybe singular or non-normal) del Pezzo surface of degree $g(X) - 3$ that contains Γ . Similarly, let S be the proper transform of the φ -exceptional surface on the Fano threefold X . Then S is a hyperplane section of X such that $\text{mult}_{\ell}(S) = 3$. Using this, we conclude that

$$X \setminus S \cong Y \setminus \mathcal{S}.$$

Moreover, if ℓ is a line of type $(1, -2)$, then the surface \mathcal{S} is not normal. This implies that the complement $Y \setminus \mathcal{S}$ contains a cylinder, so that X is cylindrical. \square

In fact, we believe that the following is true:

Conjecture 3.9. *Let X be a very general smooth Fano threefold such that $\rho(X) = 1$, $\iota(X) = 1$, and $g(X) = 9$ or $g(X) = 10$. Then X is not cylindrical.*

Using a similar Sarkisov link as in the proof of Theorem 3.8, we obtain

Theorem 3.10 ([123]). *Suppose that $\rho(X) = 1$, $\iota(X) = 1$ and $g(X) = 12$. Then X is cylindrical.*

Proof. Let ℓ be a line in X . Then there exists a unique surface $S \in |-K_X|$ such that $\text{mult}_\ell(S) = 3$. Moreover, it follows from [108, 110, 114, 184] that there exists the following Sarkisov link:

$$(3.11) \quad \begin{array}{ccc} & \tilde{X} & \xrightarrow{\chi} \hat{X} \\ \sigma \swarrow & & \searrow \varphi \\ X & & V_5, \end{array}$$

where σ is the blowup of the line ℓ , the variety V_5 is a smooth quintic del Pezzo threefold in \mathbb{P}^6 , the morphism φ is the blowup of a rational quintic curve Γ , and χ is a composition of flops.

Let E be the σ -exceptional surface, let \hat{E} be its proper transform on \hat{X} , and let $\mathcal{S} = \varphi(\hat{E})$. Then \mathcal{S} is a hyperplane section of the threefold V_5 that contains the curve Γ , and S is the proper transform of the φ -exceptional surface. Moreover, we have

$$X \setminus S \cong V_5 \setminus \mathcal{S}.$$

Let us show that $V_5 \setminus \mathcal{S}$ contains a cylinder. In fact, this follows from the proof of Theorem 3.6. We will use the notation and assumptions introduced in this proof.

Let L be a line in V_5 that is contained in \mathcal{S} (it does exist). If $\mathcal{S} \neq H_L$, let S be the proper transform on Q of the surface \mathcal{S} . Otherwise, we let $S = H_C$. Then the surface S is a hyperplane section of the quadric Q . Thus, we see that

$$V_5 \setminus (\mathcal{S} \cup H_L) \cong Q \setminus (S \cup H_C).$$

Now taking the linear projection $Q \dashrightarrow \mathbb{P}^3$ from a sufficiently general point in $S \cap H_C$, one can easily show that the complement $Q \setminus (S \cup H_C)$ contains a cylinder, so that X is cylindrical. \square

Remark 3.12 ([184]). In the notation and assumptions of the proof of Theorem 3.10, let ℓ be a line of type $(-1, 2)$. Then \mathcal{S} is a non-normal surface whose singular locus is a line in V_5 . Letting L to be this line gives $\mathcal{S} = H_L$, so that

$$X \setminus S \cong V_5 \setminus \mathcal{S} \cong Q \setminus H_C.$$

Thus, if we also have $\mathcal{N}_{L/V_5} \cong \mathcal{O}_L(1) \oplus \mathcal{O}_L(-1)$, then H_C is singular (see the proof of Theorem 3.6), so that $X \setminus S \cong \mathbb{A}^3$. We can always find such ℓ and L if $\text{Aut}(X)$ is infinite (see Theorem 3.5).

We do not know examples of cylindrical smooth Fano threefolds of Picard rank 1 and genus 7. In fact, we believe that any such threefold is not cylindrical.

Conjecture 3.13. *Let X be a smooth Fano threefold such that $\rho(X) = 1$, $\iota(X) = 1$, and $g(X) = 7$. Then X is not cylindrical.*

Before we close this section, let us mention that most of smooth Fano threefolds with $\rho(X) \geq 2$ are rational [112, 114, 188], and many of them are known to be cylindrical. However, we do not know the existence of anticanonical polar cylinders in majority of cylindrical smooth Fano threefolds. Let us list few examples.

Example 3.14. Let Y be a smooth Fano threefold such that Y is a del Pezzo threefold or $Y = \mathbb{P}^3$. Take $H \in \text{Pic}(Y)$ on Y such that $-K_Y \sim 2H$. Choose a smooth curve $\mathcal{C} \subset Y$ that is a complete intersection of two surfaces from $|H|$. Suppose that X is a blowup of the threefold Y along \mathcal{C} . Then X is a smooth Fano threefold. Moreover, if $H^3 \geq 4$, then X is cylindrical.

Example 3.15. Suppose that X is a blowup of \mathbb{P}^3 along a smooth curve that is a complete intersection of two cubic surfaces. Then X is a cylindrical smooth Fano threefold.

Example 3.16. Suppose that X is a blowup of \mathbb{P}^3 along a smooth curve of degree 6 and genus 3, which is an intersection of cubic hypersurfaces. Then X is a cylindrical smooth Fano threefold.

Example 3.17. Let Q be a smooth quadric threefold in \mathbb{P}^4 , and let H be its hyperplane section. Suppose that X is a blowup of Q along a smooth curve that is a complete intersection of two surfaces from $|2H|$. Then X is a cylindrical smooth Fano threefold.

Each smooth Fano threefold described in Examples 3.14, 3.15, 3.16 and 3.17 is cylindrical, but we do not know whether any of these threefolds contains anticanonical polar cylinders or not.

3.2. Cylindrical Fano fourfolds. Now, let X be a smooth Fano fourfold such that $\rho(X) = 1$. By Corollary 1.6 we have the following implications:

$$X \text{ is cylindrical} \implies X \text{ is rational.}$$

If $\iota(X) = 5$ or $\iota(X) = 4$, then $X = \mathbb{P}^4$ or X is a smooth quadric fourfold, so that X is cylindrical. Similarly, if $\iota(X) = 3$, then it follows from Remark 3.1 that X is one of the following fourfolds:

- (1) a smooth sextic hypersurface in $\mathbb{P}(1, 1, 1, 1, 2, 3)$;
- (2) a smooth quartic hypersurface in $\mathbb{P}(1, 1, 1, 1, 1, 2)$;
- (3) a smooth cubic fourfold in \mathbb{P}^5 ;
- (4) a smooth complete intersection of two quadrics in \mathbb{P}^6 ;
- (5) the quintic del Pezzo fourfold described in Example 1.31.

In the first two cases, we expect that X is always irrational. In fact, we know that a very general quartic hypersurface in $\mathbb{P}(1, 1, 1, 1, 1, 2)$ is irrational [97], so that it is definitely not cylindrical. Similarly, general cubic fourfold in \mathbb{P}^5 is expected to be irrational. But there are rational smooth cubic fourfolds (see [95, 96, 199, 210]), so that it is very natural to ask the following question:

Question 3.18. *Are there smooth rational cylindrical cubic fourfolds?*

Remark 3.19. Every smooth cubic fourfold in \mathbb{P}^5 containing two skew planes is rational (see [96]). In particular, the Fermat cubic fourfold is rational. If it is cylindrical, then the affine cone over it admits an effective action of the group \mathbb{G}_a by Theorem 1.15, which contradicts Conjecture 1.22.

By Lemma 3.2, we know that a smooth complete intersection of two quadrics in \mathbb{P}^6 is cylindrical. Let us prove that the quintic del Pezzo fourfold described in Example 1.31 is cylindrical as well. To do this, let us present a detailed description of this fourfold given in [186].

Let V_5 be the quintic del Pezzo fourfold in \mathbb{P}^7 . By [179, Theorem 6.6], we have the following exact sequence of groups:

$$1 \longrightarrow (\mathbb{G}_a)^4 \rtimes \mathbb{G}_m \longrightarrow \text{Aut}(V_5) \longrightarrow \text{PGL}_2(\mathbb{C}) \longrightarrow 1,$$

so that the group $\text{Aut}(V_5)$ is not reductive. In particular, the fourfold V_5 is not K-polystable [1]. The planes on V_5 belong to one of the following two classes:

- (i) a unique plane Ξ which is a Schubert variety of type $\sigma_{2,2}$;
- (ii) a one-parameter family of planes Π_t that are Schubert varieties of type $\sigma_{3,1}$.

We say that Ξ is the plane of type $\sigma_{2,2}$, and Π_t are planes of type $\sigma_{3,1}$. They are distinguished by the types of the normal bundles: $c_2(\mathcal{N}_{\Xi/X}) = 2$ and $c_2(\mathcal{N}_{\Pi_t/X}) = 1$. Moreover, there is a hyperplane section \mathcal{H} of the fourfold V_5 that contains all planes in V_5 . Furthermore, one has $\text{Sing}(\mathcal{H}) = \Xi$, the threefold \mathcal{H} is the union of all the $\sigma_{3,1}$ -planes in V_5 , and Ξ contains a special conic \mathcal{C} such that

- the intersection $\Pi_t \cap \Xi$ is a tangent line to the conic \mathcal{C} ;
- two distinct $\sigma_{3,1}$ -planes Π_{t_1} and Π_{t_2} meet in a point in $\Xi \setminus \mathcal{C}$.

The automorphism group $\text{Aut}(V_5)$ has the following orbits in V_5 :

- (1) the open orbit $X \setminus \mathcal{H}$;
- (2) the three-dimensional orbit $\mathcal{H} \setminus \Xi$;
- (3) the two-dimensional orbit $\Xi \setminus \mathcal{C}$;
- (4) the one-dimensional closed orbit \mathcal{C} .

The Hilbert scheme of lines on the del Pezzo fourfold V_5 is smooth, irreducible, and four-dimensional. Moreover, if ℓ is a line in V_5 , then ℓ belongs to one of the following five classes:

- (a) $\ell \not\subset \mathcal{H}$, $\ell \cap \Xi = \emptyset$, and $\ell \cap \mathcal{H}$ is a point;
- (b) $\ell \subset \mathcal{H}$, $\ell \cap \Xi$ is a point, and $\ell \cap \mathcal{C} = \emptyset$;
- (c) $\ell \subset \mathcal{H}$, and $\ell \cap \Xi = \ell \cap \mathcal{C}$ is a point;
- (d) $\ell \subset \Xi$, and the intersection $\ell \cap \mathcal{C}$ consists of two points;
- (e) $\ell \subset \Xi$ and ℓ is tangent to \mathcal{C} .

The group $\text{Aut}(V_5)$ acts transitively on the lines in each of these classes. For a line $\ell \subset V_5$, the lines meeting ℓ sweep out a hyperplane section H_ℓ of the fourfold V_5 that is singular along the line ℓ . Vice versa, if H is a hyperplane section of the quintic del Pezzo fourfold V_5 that has non-isolated singularities, then $H = H_\ell$ for some line $\ell \subset V_5$.

Theorem 3.20 ([186]). *Let ℓ be a line in V_5 that is not a line of type (b). Then $V_5 \setminus H_\ell \cong \mathbb{A}^4$.*

Proof. If ℓ is a line of type (d) or (e), then $H_\ell = \mathcal{H}$. On the other hand, there exists the following $\text{Aut}(V_5)$ -equivariant Sarkisov link:

$$\begin{array}{ccc} & \tilde{V}_5 & \\ \sigma \swarrow & & \searrow \varphi \\ V_5 & \text{---} \text{---} \text{---} \psi \text{---} \text{---} \text{---} & \mathbb{P}^4 \end{array}$$

where σ is the blowup of the plane Ξ , φ is the blowup of a twisted cubic curve C , and ψ is the linear projection from Ξ . Then the φ -exceptional divisor is the proper transform of the threefold \mathcal{H} . Moreover, if E is the σ -exceptional divisor, then $\varphi(E)$ is the hyperplane in \mathbb{P}^4 that contains C . Thus, if ℓ is a line of type (d) or (e), then $V_5 \setminus H_\ell = V_5 \setminus \mathcal{H} \cong \mathbb{P}^4 \setminus \varphi(E) \cong \mathbb{A}^5$.

Let $\pi: \widehat{V}_5$ be the blowup of the line ℓ . Then there exists the following Sarkisov link:

$$(3.21) \quad \begin{array}{ccc} & \widehat{V}_5 & \\ \pi \swarrow & & \searrow \eta \\ V_5 & \text{---} \text{---} \text{---} \zeta \text{---} \text{---} \text{---} & Q \end{array}$$

where Q is an irreducible quadric in \mathbb{P}^5 , the map ζ is the projection from ℓ , and η is a birational morphism that contracts the proper transform of the hyperplane section H_ℓ to a surface of degree 3. Let \widehat{H}_ℓ be the proper transform on \widehat{V}_5 of the threefold H_ℓ , and let F be the π -exceptional divisor. Then $V_5 \setminus H_\ell \cong Q \setminus \eta(F)$, and $\eta(F)$ is a singular hyperplane section of the quadric Q .

If ℓ is a line of type (a), then all fibers of η are one-dimensional, so that Q is smooth (see [2]). Thus, in this case, we have $V_5 \setminus H_\ell \cong Q \setminus \eta(F) \cong \mathbb{A}^4$.

To complete the proof, we may assume that ℓ is of type (c). Then ℓ is contained in a plane in V_5 , so that η has a two-dimensional fiber. Hence, in this case, the quadric Q can be singular (cf. [3]). Analyzing the situation more carefully, we see that $V_5 \setminus H_\ell \cong Q \setminus \eta(F) \cong \mathbb{A}^4$. \square

Corollary 3.22. *The quintic del Pezzo fourfold is cylindrical.*

In the remaining part of this subsection, we present known constructions of cylinders in some smooth Fano–Mukai fourfolds. Basically, our main goal is to explain how to prove Theorem 3.4. Thus, we suppose that X is a smooth Fano–Mukai fourfold, $\rho(X) = 1$ and $g(X) \in \{7, 8, 9, 10\}$.

Let H be an ample Cartier divisor on X such that

$$-K_X \sim 2H.$$

Then $H^4 = 2g(X) - 2 \in \{12, 14, 16, 18\}$. Moreover, the divisor H is very ample, and the linear system $|H|$ gives an embedding $X \hookrightarrow \mathbb{P}^{g(X)+2}$. Let us deal with four cases separately.

If $g(X) = 10$, then $X = X_{18}$ is a hyperplane section of the homogeneous fivefold $G_2/P \subset \mathbb{P}^{13}$, where G_2 is the simple algebraic group of exceptional type G_2 , and P is its parabolic subgroup that corresponds to a short root (see [162, 163]). The family \mathfrak{X} of all such fourfolds is one-dimensional. Moreover, if $X = X_{18}$ is a general member of \mathfrak{X} , then $\text{Aut}(X) \cong \mathbb{G}_m^2 \rtimes \mu_2$. Besides, there are three distinguished fourfolds in this family:

- (0) X_{18}^r such that $\text{Aut}(X_{18}^r) \cong \mathbb{G}_m^2 \rtimes \mu_6$;
- (1) X_{18}^s such that $\text{Aut}(X_{18}^s) \cong \text{GL}_2(\mathbb{C}) \rtimes \mu_2$;
- (2) X_{18}^a such that $\text{Aut}(X_{18}^a) \cong (\mathbb{G}_a \times \mathbb{G}_m) \rtimes \mu_2$.

See [191] for details, where the following result has been proved:

Theorem 3.23 ([191]). *Let X be a smooth Fano–Mukai fourfold in \mathbb{P}^{12} of genus 10 with $\rho(X) = 1$. Then there exists an $\text{Aut}^0(X)$ -invariant hyperplane section H of X such that the complement $X \setminus H$ is $\text{Aut}^0(X)$ -equivariantly isomorphic to \mathbb{A}^4 .*

This theorem implies, in particular, that any smooth Fano–Mukai fourfolds of genus 10 is cylindrical. See also Example 4.16 for another application of Theorem 3.23.

If $g(X) = 8$, then $X = X_{14}$ is a section of the Grassmannian $\text{Gr}(2, 6) \subset \mathbb{P}^{14}$ by a linear subspace of dimension 10 (see [162, 163]). Some of these fourfolds are cylindrical.

Example 3.24 ([189]). Suppose that $g(X) = 8$ and X contains a plane Π which is a Schubert variety of type $\sigma_{4,2}$, and X does not contain planes meeting Π along a line. Such fourfolds do exist and form a subspace of codimension one in the moduli space of all Fano–Mukai fourfolds of genus 8. Then it follows from [185] that there exists the following Sarkisov link:

$$\begin{array}{ccc} & \tilde{X} & \\ \sigma \swarrow & & \searrow \varphi \\ X & & V_5 \end{array}$$

where V_5 is the del Pezzo quintic fourfold in \mathbb{P}^7 (see Theorem 3.20), σ is the blowup of the plane Π , and φ is the blowup of a smooth rational surface S of degree 7 such that $K_S^2 = 3$. Then

$$X \setminus H_X \cong V_5 \setminus H_{V_5},$$

where H_{V_5} is the proper transform on V_5 of the σ -exceptional divisor, and H_X is the proper transform on X of the φ -exceptional divisor. On the other hand, the divisor H_{V_5} is a hyperplane section of the fourfold V_5 that contains S , and H_X is a hyperplane section of X containing Π . Thus, the set $V_5 \setminus H_{V_5}$ contains a cylinder by [189, Theorem 4.1], so that X is cylindrical.

If $g(X) = 7$, then $X = X_{12}$ is a section of the orthogonal Grassmannian $\text{OGr}(4, 9) \subset \mathbb{P}^{15}$ by a linear subspace of dimension 9 (see [162, 163]). In this case, we also have cylindrical fourfolds.

Example 3.25 ([189]). Suppose that $g(X) = 7$ and X contains a plane Π . Such fourfolds do exist. Suppose that X is a sufficiently general Fano–Mukai fourfold of genus 7 that contains the plane Π . Then by [185] there exists the Sarkisov link

$$\begin{array}{ccc} & \tilde{X} & \\ \sigma \swarrow & & \searrow \varphi \\ X & & V_4 \end{array}$$

where V_4 is a smooth complete intersection of two quadrics in \mathbb{P}^6 , σ is the blowup of the plane Π , and φ is the blowup of a smooth del Pezzo surface S such that $K_S^2 = 5$. Arguing as in Example 3.24, we conclude that X is cylindrical.

If $g(X) = 9$, then $X = X_{16}$ is a section of the Lagrangian Grassmannian $\mathrm{LGr}(3, 6) \subset \mathbb{P}^{13}$ by a linear subspace of dimension 11 (see [162, 163]). There are cylindrical fourfolds in this family.

Example 3.26 ([190]). Suppose that $g(X) = 9$. Then X_{16} contains an irreducible two-dimensional quadric surface S . Suppose, for simplicity, that X_{16} is a general Fano–Mukai fourfold of genus 9 that contains S . Then there exists the following Sarkisov link:

$$\begin{array}{ccc} & \tilde{X} & \\ \sigma \swarrow & & \searrow \varphi \\ X & & V_5 \end{array}$$

where V_5 is the del Pezzo quintic fourfold, σ is the blowup of the surface S , and φ is the blowup along a smooth del Pezzo surface of degree 6. Arguing as in Example 3.24, we see that X is cylindrical.

The interested reader can consult also the recent preprint [103] for further examples of cylindrical Fano fourfolds.

3.3. Cylinders in Mori fibrations. This subsection is inspired by the following

Question 3.27. *Given a family of cylindrical varieties, when its total space is cylindrical?*

For example, irrational three-dimensional conic bundles are not cylindrical, though their general fibers are. In general, this question is very subtle and has birational nature, so that it is natural to consider it for Mori fibred spaces first.

Let V be a projective variety with terminal \mathbb{Q} -factorial singularities, let $\pi: V \rightarrow B$ be a dominant projective non-birational morphism such that $-K_V$ is π -ample, $\pi_*\mathcal{O}_V = \mathcal{O}_B$ and $\rho(V) = \rho(B) + 1$. Let X_η be the fiber of the morphism π over the (scheme-theoretic) generic point η of the base B . Then X_η is a Fano variety that has at most terminal singularities, which is defined over $\mathbb{K} = \mathbb{k}(B)$, i.e. the field of rational functions on B . Over the (algebraically non-closed) field \mathbb{K} , the divisor class group of the Fano variety X_η is of rank 1, because we assume that $\rho(V) = \rho(B) + 1$.

Definition 3.28 ([61]). If the variety V contains a (Zariski open) cylinder $U = \mathbb{A}^1 \times Z$, we say that the cylinder U is *vertical* (with respect to π) if there is a morphism $h: Z \rightarrow B$ such that the restriction $\pi|_U: U \rightarrow B$ is a composition $h \circ \mathrm{pr}_Z$, where $\mathrm{pr}_Z: U \rightarrow Z$ is the natural projection. In this case, we have commutative diagram:

$$(3.29) \quad \begin{array}{ccc} \mathbb{A}^1 \times Z = U & \xhookrightarrow{\quad} & V \\ \downarrow \mathrm{pr}_Z & & \downarrow \pi \\ Z & \xrightarrow{\quad h \quad} & B \end{array}$$

A cylinder in V which is not vertical is called *twisted*.

If V contains a vertical cylinder $U = \mathbb{A}^1 \times Z$, then the Fano variety X_η contains a cylinder

$$U_\eta = \mathbb{A}^1 \times Z_\eta,$$

where U_η and Z_η are generic (scheme) fibers of the morphisms $h \circ \mathrm{pr}_Z$ and h in (3.29), respectively. Vice versa, if the Fano variety X_η contains a cylinder defined over the field \mathbb{K} , then V does contain a vertical cylinder by [61, Lemma 3]. This gives a motivation to study cylinders in Fano varieties defined over arbitrary fields (cf. [17, 100, 141, 142]) The first step in this direction is

Theorem 3.30 ([61]). *Let S be a geometrically irreducible smooth del Pezzo surface defined over a field \mathbb{F} of characteristic 0. Suppose that $\rho(S) = 1$. Then the following conditions are equivalent:*

- (i) *the surface S contains a cylinder defined over \mathbb{F} ;*
- (ii) *the surface S is rational over \mathbb{F} ;*
- (iii) *$K_S^2 \geq 5$ and S has an \mathbb{F} -point.*

Proof. It is commonly known that the conditions (ii) and (iii) are equivalent (see, for example, [111]). Moreover, the implication (iii) \Rightarrow (i) can be shown using well-known Sarkisov links that start at S , which are described in [111]. For details, see the proof of [61, Proposition 12]. Thus, we just have to show that (i) implies (iii). This can also be shown using Sarkisov links, but we present another proof.

Suppose that S contains a cylinder U which is defined over \mathbb{F} . Then $U \cong \mathbb{A}^1 \times Z$ for some affine curve Z defined over \mathbb{F} . Let \overline{Z} be the completion of the curve Z . Then \overline{Z} is a geometrically irreducible curve. Moreover, we have the following commutative diagram

$$\begin{array}{ccccccc}
 \mathbb{P}^1 \times \overline{Z} & \xleftarrow{\quad} & \mathbb{A}^1 \times \overline{Z} & \xleftarrow{\quad} & \mathbb{A}^1 \times Z \cong U & \xrightarrow{\quad} & S \xleftarrow{\pi} \tilde{S} \\
 & \searrow \bar{p}_2 & \searrow p_2 & \downarrow p_Z & \searrow \psi & \searrow \varphi & \\
 & & & Z & & & \\
 & & & \downarrow & & & \\
 & & & \overline{Z} & & &
 \end{array}$$

where p_Z , p_2 and \bar{p}_2 are the natural projections to the second factors, ψ is the rational map induced by p_Z , π is a birational morphism resolving the indeterminacy of ψ and φ is a morphism. By construction, a general fiber of φ is isomorphic to \mathbb{P}^1 .

Let Γ be the section of \bar{p}_2 that is the complement of $\mathbb{A}^1 \times \overline{Z}$ in $\mathbb{P}^1 \times \overline{Z}$, and let $\tilde{\Gamma}$ be the proper transform on \tilde{S} of the curve Γ . Then $\tilde{\Gamma} \cong \Gamma \cong \overline{Z}$, the curve $\tilde{\Gamma}$ is a section of φ , and the curve $\tilde{\Gamma}$ is π -exceptional, because $\rho(S) = 1$. Let $P = \pi(\tilde{\Gamma})$. Then P is an \mathbb{F} -point.

Now, we can proceed in two (slightly different) ways. First, as in the proof of [61, Theorem 1], we can let \mathcal{M} to be the linear system on S that gives the map ψ . Then, arguing as in Section 2.2, we conclude that $(S, \lambda\mathcal{M})$ is not log canonical at P for a some $\lambda \in \mathbb{Q}_{>0}$ such that $\lambda\mathcal{M} \sim_{\mathbb{Q}} -K_S$. Such number exists, since $\rho(S) = 1$. Let M_1 and M_2 be two general curves in \mathcal{M} . Then

$$\frac{K_S^2}{\lambda^2} = M_1 \cdot M_2 \geq (M_1 \cdot M_2)_P > \frac{4}{\lambda^2}$$

by [51, Theorem 3.1]. This gives $K_S^2 \geq 5$, so that (i) implies (iii).

Alternatively, we can use Corollary 2.9. Let C_1, \dots, C_n be the irreducible curves in S that lie in the complement $S \setminus U$. Then we put $D = \lambda(C_1 + \dots + C_n)$ for $\lambda \in \mathbb{Q}_{>0}$ such that $D \sim_{\mathbb{Q}} -K_S$. Therefore, we conclude that S contains a $(-K_S)$ -polar cylinder, so that $K_S^2 \geq 4$ by Corollary 2.9. Thus, we may assume that $K_S^2 = 4$. Then our point P is not contained in any (-1) -curve in $S \otimes_{\mathbb{F}} \overline{\mathbb{F}}$, where $\overline{\mathbb{F}}$ is an algebraic closure of the field \mathbb{F} . Indeed, otherwise the $\text{Gal}(\overline{\mathbb{F}}/\mathbb{F})$ -orbit of this curve would consist of at least four (-1) -curves that all pass through the point P , which is impossible. Let $\xi: \hat{S} \rightarrow S$ be the blowup of the point P , and let E be the exceptional curve of the blowup ξ . Then \tilde{S} is a smooth del Pezzo surface of degree $K_{\tilde{S}}^2 = 3$ and

$$\tilde{D} + (\text{mult}_P(D) - 1)E \sim_{\mathbb{Q}} -K_{\tilde{S}},$$

where $\text{mult}_P(D) > 1$ by Remark 2.3 and Lemma A.3. Then \tilde{S} contains a $(-K_{\tilde{S}})$ -polar cylinder, which is impossible by Corollary 2.9. This again shows that (i) implies (iii). \square

Corollary 3.31 ([61, Theorem 1]). *Suppose that X_η is a del Pezzo surface. Then V contains a vertical cylinder $\iff K_{X_\eta}^2 \geq 5$ and π has a rational section.*

Note that if \mathbb{k} is uncountable and the general fiber of π contains a cylinder, then it follows from [63, 116] that the total space of the family $V \times_B B' \rightarrow B$ contains a vertical cylinder for an appropriate finite base change $B' \rightarrow B$. This basically means that $X_\eta \otimes_{\mathbb{K}} \mathbb{K}'$ contains a cylinder defined over \mathbb{K}' for an appropriate finite extension of fields $\mathbb{K} \subset \mathbb{K}'$.

Remark 3.32. If X_η is a del Pezzo surface and $K_{X_\eta}^2 \leq 4$, then V can contain twisted cylinders. In fact, there are three-dimensional examples constructed in [60, 61] such that $K_{X_\eta}^2 \leq 3$, $B = \mathbb{P}^1$, and V contains a Zariski open subset isomorphic to \mathbb{A}^3 . See also colored[59, 91, 201, 202].

Now let us mention one relevant result about forms of the quintic del Pezzo threefold defined over a non-algebraically closed field (cf. [141, Theorem 3.3]).

Theorem 3.33 ([62]). *Let X be a smooth Fano threefold defined over a field \mathbb{F} of characteristic 0. Suppose that $X \otimes_{\mathbb{F}} \overline{\mathbb{F}} \cong V_5$, where V_5 is the quintic del Pezzo threefold described in Example 1.31, where $\overline{\mathbb{F}}$ is the algebraic closure of the field \mathbb{F} . Then the following assertions hold:*

- X contains a Zariski open subset $U \cong \mathbb{A}^2 \times Z$ for some affine curve Z ;
- X contains a Zariski open subset isomorphic to \mathbb{A}^3 if and only if X contains a smooth rational curve ℓ defined over \mathbb{F} such that $-K_{V_5} \cdot \ell = 2$ and $\mathcal{N}_{\ell/X} \cong \mathcal{O}_{\ell}(-1) \oplus \mathcal{O}_{\ell}(1)$.

Let us conclude this section with the following generalization of Theorem 3.3.

Theorem 3.34 ([142]). *Let X be a smooth Fano threefold defined over a field \mathbb{F} of characteristic 0. Suppose that $X \otimes_{\mathbb{F}} \overline{\mathbb{F}} \cong X_{2g-2}$, where X_{2g-2} is a Fano–Mukai variety of genus g with $\rho(X_{2g-2}) = 1$, where $\overline{\mathbb{F}}$ be the algebraic closure of \mathbb{F} . Suppose that the following conditions hold:*

- (1) $\dim(X) \geq 5$;
- (2) $g \in \{7, 8, 9, 10\}$;
- (3) X has an \mathbb{F} -point.

Then X is cylindrical over \mathbb{F} .

4. BEYOND CYLINDRICITY

4.1. Flexible affine varieties. Let X be an affine variety. Given a \mathbb{G}_a -action on X , it induces a representation of the group \mathbb{G}_a on the structure \mathbb{k} -algebra $\mathcal{O}(X)$ of the form

$$(t, f) \longmapsto \exp(t\partial)(f)$$

for $t \in \mathbb{G}_a$ and $f \in \mathcal{O}(X)$, where the infinitesimal generator ∂ of the \mathbb{G}_a -subgroup is a *locally nilpotent derivation* of $\mathcal{O}(X)$, which means that every element $f \in \mathcal{O}(X)$ is annihilated by $\partial^{(m)}$ for some sufficiently large m that depends on the element f . Conversely, any locally nilpotent derivation of the \mathbb{k} -algebra $\mathcal{O}(X)$ generates a \mathbb{G}_a -action on X (see [67]).

Recall that the derivations of $\mathcal{O}(X)$ correspond to the regular vector fields on X . We say that a vector field on X is *locally nilpotent* if the corresponding derivation is.

If an open variety X admits a \mathbb{G}_a -action, then the log-Kodaira dimension of X is negative. However, the converse does not hold, in general. Indeed, there are smooth affine surfaces of negative log-Kodaira dimension which admit no effective \mathbb{G}_a -action [92]. Let us stay on this in more detail.

As we mentioned already, any smooth affine surface X of negative log-Kodaira dimension contains a cylinder [154, Ch. 2, Theorem 2.1.1]. Moreover, X is affine-ruled, that is, there is a morphism $X \rightarrow C$ onto a smooth curve C with general fiber \mathbb{A}^1 . The base curve C could be affine or projective. However, a smooth affine surface X admits an effective \mathbb{G}_a -action if and only if it

admits an \mathbb{A}^1 -ruling $X \rightarrow C$ over an affine curve C , or, which is equivalent, a *principal* cylinder, see Theorem 1.13. In [92] there are examples of smooth rational affine surfaces \mathbb{A}^1 -ruled over \mathbb{P}^1 and with no \mathbb{A}^1 -ruling over an affine curve. Hence, such a surface admits no effective \mathbb{G}_a -action.

To construct such a surface X we start with the quadric $\mathbb{P}^1 \times \mathbb{P}^1$ endowed with the first projection to \mathbb{P}^1 . We blow up three distinct points on the section $S = \mathbb{P}^1 \times \{\infty\}$ and infinitesimally near points in such a way that each of the 3 resulting reducible fibers has a unique (-1) -component of multiplicity 2 and the union of the section S and the remaining components of the reducible fibers forms a connected divisor D . The complement of D in the resulting projective surface is a smooth affine surface X . It comes equipped with an \mathbb{A}^1 -fibration $X \rightarrow \mathbb{P}^1$. Each fiber of this fibration is irreducible, and three of them are multiple of multiplicity 2. According to [93, Theorem 4.1], such a surface X does not carry any \mathbb{A}^1 -fibration over an affine curve. Hence, X admits no \mathbb{G}_a -action.

Definition 4.1. A point $P \in X$ is said to be *flexible* if locally nilpotent vector fields on X span the tangent space $T_P X$. The variety X is said to be *flexible* if every smooth point of X is flexible. We also say that X is *generically flexible* if every point in a non-empty Zariski open subset of X is flexible.

Let $\text{SAut}(X)$ be the subgroup of $\text{Aut}(X)$ generated by all the \mathbb{G}_a -subgroups. The flexibility of X is ultimately related to the transitivity of the action of the group $\text{SAut}(X)$. Indeed, we have the following criteria of flexibility.

Theorem 4.2 ([8]). *Suppose that $\dim(X) \geq 2$. Then the following conditions are equivalent:*

- (1) *the variety X is flexible;*
- (2) *the group $\text{SAut}(X)$ acts transitively on the smooth locus of X ;*
- (3) *the group $\text{SAut}(X)$ acts highly transitively on the smooth locus of X .*

One says that a group acts *highly transitively*¹ on an infinite set if it acts m -transitively for any natural number m .

Remark 4.3. A dimension count shows that an algebraic group cannot act highly transitively on an affine variety. Moreover, it cannot act even 3-transitively on an affine variety [22, 127].

Let us present examples of flexible affine varieties; see e.g. [11–13] for further examples.

Example 4.4. Let $X = \mathbb{A}^n$, where $n \geq 2$. Then the subgroup of translations in $\text{SAut}(\mathbb{A}^n)$ acts transitively on the variety X , so that X is flexible by Theorem 4.2 (cf. [120]).

Example 4.5. Let X be the n th Calogero–Moser space defined as follows:

$$\left\{ (A, B) \in \text{Mat}_n(\mathbb{k}) \times \text{Mat}_n(\mathbb{k}) \mid \text{rk}([A, B] + I_n) = 1 \right\} // \text{PGL}_n(\mathbb{k}),$$

where $\text{PGL}_n(\mathbb{k})$ acts via $g.(A, B) = (gAg^{-1}, gBg^{-1})$. Then X is a smooth rational irreducible affine algebraic variety of dimension $2n$ [181, 215], and it follows from [18, 138] that $\text{Aut}(X)$ acts highly transitively on X for every $n \geq 1$. Moreover, the variety X is flexible by [4, Proposition 2.9].

There are several constructions producing new flexible varieties from given ones (see [8, 12, 65, 120]). For instance, the product of flexible varieties is flexible. Some further examples of flexible varieties are as follows.

Example 4.6. Suppose that X is an affine G -variety of dimension ≥ 2 , where G is a connected linear algebraic group that acts on X with an open orbit. Then X is flexible in the following cases:

- X is a normal toric variety with no torus factor [12, Theorem 0.2.2];
- $X = G/H$ is a homogeneous space and G has no nontrivial character [8, Theorem 5.4];

¹Or *infinitely transitively* in another terminology

- X is smooth and G is semisimple [8, Theorem 5.6];
- X is smooth with only constant invertible functions and G is reductive [85, Theorem 2];
- X is normal and $G = \mathrm{SL}_2(\mathbb{k})$ [8, Theorem 5.7];
- X is normal horospherical and G is semisimple [204, Theorem 2];
- X is normal horospherical with no non-constant invertible regular function [85, Theorem 3].

See also [21, 56, 57, 87, 117, 139].

If we replace the smooth locus of X in Theorem 4.2 by the open orbit of the group $\mathrm{SAut}(X)$, we obtain a criterion for the generic flexibility [8]. If X contains \mathbb{A}^n as a principal Zariski open set, then X is generically flexible. Generically flexible varieties are unirational, but they are not always stably rational (see [148, Proposition 4.9] and [180, Example 1.22]).

Example 4.7. Suppose that X is a normal affine surface such that X can be completed by a simple normal crossing chain of rational curves. Then X is often called a *Gizatullin surface*. If $X \not\cong \mathbb{A}^1 \times (\mathbb{A}^1 \setminus \{0\})$, then it is generically flexible [86], but it is not necessarily flexible [137].

Affine cones over cylindrical Fano varieties often provide examples of flexible affine varieties.

Example 4.8. Let $V = G/P$, where G is a semisimple algebraic group, and P is its parabolic subgroup. Then V is a smooth Fano variety. Let $V \hookrightarrow \mathbb{P}^n$ be any projectively normal embedding, and let \widehat{V} be the affine cone in \mathbb{A}^{n+1} over V . If $\dim(V) \geq 2$, then \widehat{V} is flexible by [12, Theorem 1.1].

To explain why this is the case, let us present two explicit criteria of flexibility of affine cones. To do this, fix a smooth projective variety V . Let H be a very ample divisor on the variety X . Then the linear system $|H|$ gives an embedding $V \hookrightarrow \mathbb{P}^n$. Let \widehat{V} be the affine cone in \mathbb{A}^{n+1} over V . We are interested in the case when V is a smooth cylindrical Fano variety.

If the variety V is uniformly cylindrical, then each point of V is contained in a cylinder, so that the variety V admits a covering

$$(4.9) \quad V = \bigcup_{i \in I} U_i,$$

where each U_i is a Zariski open subset in V such that $U_i \cong \mathbb{A}^1 \times Z_i$ for some affine variety Z_i . In this case, a subset $Y \subset V$ is said to be *invariant* with respect to a cylinder U_i if

$$Y \cap U_i = \pi_i^{-1}(\pi_i(Y \cap U_i)),$$

where $\pi_i: U_i \rightarrow Z_i$ is the natural projection.

Definition 4.10. If V is uniformly cylindrical, then we say that the covering (4.9) is *transversal* if no proper subset $Y \subset X$ is invariant with respect to every cylinder U_i in the covering (4.9).

Now, we are ready to state the first flexibility criterion for affine cones.

Theorem 4.11 ([174]). *Suppose that V is uniformly cylindrical and has a covering (4.9) such that*

- (i) *the covering (4.9) is transversal;*
- (ii) *each cylinder in the covering (4.9) is H -polar.*

Then the affine cone \widehat{V} is flexible.

The second useful criterion is given by the following

Theorem 4.12 ([153]). *The affine cone \widehat{V} is flexible if the variety V is uniformly cylindrical and admits a covering*

$$V = \bigcup_{j \in J} W_j$$

where each W_j is a flexible affine Zariski open subset in V such that $W_j = V \setminus \mathrm{Supp} D_j$ for some effective \mathbb{Q} -divisor D_j on the variety V that satisfies $D_j \sim_{\mathbb{Q}} H$.

Using these criteria and the proof of Lemma 2.17, one can prove the following result:

Theorem 4.13 ([172, 174]). *Suppose that V is a smooth del Pezzo surface such that $K_V^2 \geq 4$. Then the affine cone \hat{V} is flexible for every very ample divisor H on the surface V .*

Unfortunately, we cannot apply Theorems 4.11 and 4.12 to the affine cone in \mathbb{A}^4 over a smooth cubic surface in \mathbb{P}^3 , simply because its anticanonical divisor is not cylindrical by Corollary 2.9. On the other hand, in this case, we know from Theorem 2.14 that every ample \mathbb{Q} -divisor that is not a multiple of the anticanonical divisor is cylindrical. Using this and the construction of cylinders given in the proof of Theorem 2.14, Perepechko very recently proved the following result:

Theorem 4.14 ([175]). *If V is a smooth cubic surface, then the affine cone \hat{V} is generically flexible for every very ample divisor H on the surface V such that $H \notin \mathbb{Z}_{>0}[-K_V]$.*

Now, let us consider the flexibility of affine cones over some cylindrical smooth Fano threefolds. Many of them are flexible by Theorem 4.12, because the underlying Fano threefolds admit covering like in Theorem 4.12 with each Zariski open subset W_j isomorphic to \mathbb{A}^3 . A possibly non-complete list of such smooth Fano threefolds is given in [10, Proposition 4]. This gives

Corollary 4.15. *Suppose that V is a smooth Fano threefold admitting an effective $\mathrm{PSL}_2(\mathbb{k})$ -action. If $\rho(V) = 1$, then the affine cone \hat{V} is flexible.*

Proof. If $\rho(V) = 1$, then it follows from Theorem 3.5 that one of the following four cases occurs:

- (i) $V = \mathbb{P}^3$;
- (ii) V is the smooth quadric threefold in \mathbb{P}^4 ;
- (iii) V is the smooth quintic del Pezzo threefold $V_5 \subset \mathbb{P}^6$ described in Example 1.31;
- (iv) V is the Mukai–Umemura threefold $X = X_{22}^{\mathrm{mu}} \subset \mathbb{P}^{13}$.

We may assume that we are in the case (iii) or (iv), because the required assertion is clear in the remaining cases. Then it follows from the proofs of Theorems 3.6 and 3.10 that V contains a one-parameter family of hyperplane sections H_ℓ such that each H_ℓ is singular along a line ℓ and

$$V \setminus H_\ell \cong \mathbb{A}^3.$$

The group $\mathrm{PSL}_2(\mathbb{k})$ acts transitively on this family. So, to apply Theorem 4.12, we need to check that the intersection of all these hyperplane sections is empty. Suppose that this is not the case. Then this intersection is $\mathrm{PSL}_2(\mathbb{k})$ -invariant, so that it contains a closed $\mathrm{PSL}_2(\mathbb{k})$ -orbit of minimal dimension. But the variety V does not contain $\mathrm{PSL}_2(\mathbb{k})$ -fixed points, and the only one-dimensional closed $\mathrm{PSL}_2(\mathbb{k})$ -orbit in V is not contained in any hyperplane section singular along a line. \square

For more examples of smooth Fano threefolds with flexible affine cones, see [153, Theorem 4.5]. Now, let us present examples of smooth Fano fourfolds with flexible affine cones.

Example 4.16 ([191]). It follows from Theorems 3.20 and 3.23 that the following smooth cylindrical Fano fourfolds admit coverings by affine charts isomorphic to \mathbb{A}^4 :

- (1) the quintic del Pezzo fourfold V_5 described in Example 1.31 (see Theorem 3.20);
- (2) the Fano–Mukai fourfold X_{18}^s of genus 10 with $\mathrm{Aut}(X_{18}^s) \cong \mathrm{GL}_2(\mathbb{C}) \rtimes \mu_2$;
- (3) the Fano–Mukai fourfolds X_{18} of genus 10 with $\mathrm{Aut}^0(X_{18}) \cong \mathbb{G}_m^2$ (there is a one-parameter family of these, up to isomorphism).

Hence, all of them have flexible affine cones.

By Theorem 3.23, every smooth Fano–Mukai fourfold in \mathbb{P}^{12} of genus 10 contains a Zariski open subset isomorphic to \mathbb{A}^4 . Moreover, the following result has been recently proved in [192].

Theorem 4.17. *The affine cones over any smooth Fano–Mukai fourfold of genus 10 are flexible.*

For more higher-dimensional examples of flexible affine cones, see [153].

4.2. Cylinders in complements to hypersurfaces. This section is motivated by the following folklore conjecture that first appeared in 2005 [68].

Conjecture 4.18. *Let S be a smooth cubic surface in \mathbb{P}^3 . Then any automorphism of the affine variety $\mathbb{P}^3 \setminus S$ is induced by an automorphism of \mathbb{P}^3 , i.e., we have*

$$\mathrm{Aut}(\mathbb{P}^3 \setminus S) = \mathrm{Aut}(\mathbb{P}^3, S).$$

If S is smooth surface in \mathbb{P}^3 of degree ≥ 4 , then it is easy to see that $\mathrm{Aut}(\mathbb{P}^3 \setminus S) = \mathrm{Aut}(\mathbb{P}^3, S)$. Vice versa, if S is either a smooth quadric surface or a plane in \mathbb{P}^3 , then $\mathrm{Aut}(\mathbb{P}^3 \setminus S) \neq \mathrm{Aut}(\mathbb{P}^3, S)$. Moreover, it is not hard to see that Conjecture 4.18 fails for **some** singular cubic surfaces.

Example 4.19. Let S be one of the three cubic surfaces with Du Val singularities in \mathbb{P}^3 that admits an effective \mathbb{G}_a -action (see [41, 151, 200]). Then $\mathrm{Aut}(\mathbb{P}^3, S)$ contains a subgroup isomorphic to \mathbb{G}_a , so that $\mathrm{Aut}(\mathbb{P}^3 \setminus S)$ also contains a subgroup isomorphic to \mathbb{G}_a . Then $\mathrm{Aut}(\mathbb{P}^3 \setminus S)$ must be infinite dimensional (see [67]), so that $\mathrm{Aut}(\mathbb{P}^3 \setminus S) \neq \mathrm{Aut}(\mathbb{P}^3, S)$, because $\mathrm{Aut}(\mathbb{P}^3, S)$ is algebraic.

Based on the results in [33, 38, 39], we may generalize the problem to del Pezzo surfaces that are hypersurfaces in weighted projective spaces. To be precise, let S be a del Pezzo surface that has at most Du Val singularities such that $K_S^2 \leq 3$. Then we have one of the following three cases:

- (1) $K_S^2 = 1$, and S is a hypersurface of degree 6 in $\mathbb{P}(1, 1, 2, 3)$;
- (2) $K_S^2 = 2$, and S is a hypersurface of degree 4 in $\mathbb{P}(1, 1, 1, 2)$;
- (3) $K_S^2 = 3$, and S is a hypersurface of degree 3 in \mathbb{P}^3 .

Denote by \mathbb{P} the weighted projective space in these three cases: $\mathbb{P}(1, 1, 2, 3)$, $\mathbb{P}(1, 1, 1, 2)$ or \mathbb{P}^3 . Then, very surprisingly, we have the following result:

Theorem 4.20 ([33, 171]). *The following three conditions are equivalent:*

- *the surface S contains a $(-K_S)$ -polar cylinder;*
- *the complement $\mathbb{P} \setminus S$ is cylindrical;*
- *the group $\mathrm{Aut}(\mathbb{P} \setminus S)$ contains a unipotent subgroup.*

Combining this result with Theorem 2.8, we obtain

Corollary 4.21 ([171, Corollary 1.6]). *The group $\mathrm{Aut}(\mathbb{P} \setminus S)$ contains no unipotent subgroup exactly when S is one of the surfaces listed in Theorem 2.8.*

Corollary 4.22 ([33, Corollary 4.10]). *Suppose that the surface S contains a $(-K_S)$ -polar cylinder. Then $\mathrm{Aut}(\mathbb{P} \setminus S) \neq \mathrm{Aut}(\mathbb{P}, S)$.*

Proof. By Theorem 4.20, the group $\mathrm{Aut}(\mathbb{P} \setminus S)$ contains a unipotent subgroup, so that it is infinite dimensional, which implies that $\mathrm{Aut}(\mathbb{P} \setminus S) \neq \mathrm{Aut}(\mathbb{P}, S)$, because $\mathrm{Aut}(\mathbb{P}, S)$ is algebraic. \square

This corollary together with Theorem 2.8 show that Conjecture 4.18 fails for **all** singular cubic surfaces that have Du Val singularities. On the other hand, we have

Theorem 4.23 ([33, Theorem 4.1]). *Suppose that S is smooth. If $K_S^2 = 1$, then*

$$\mathrm{Aut}(\mathbb{P} \setminus S) = \mathrm{Aut}(\mathbb{P}, S).$$

If $K_S^2 = 2$ or $K_S^2 = 3$, then $\mathrm{Aut}(\mathbb{P} \setminus S)$ does not contain nontrivial connected algebraic groups.

The proof of this result depends on irrationality of some del Pezzo threefolds (see [50, 88, 89, 213]). Taking into account Theorem 4.23, Corollary 2.9 and Corollary 4.22, we propose the following

Conjecture 4.24. *The surface S contains no $(-K_S)$ -polar cylinder $\iff \mathrm{Aut}(\mathbb{P} \setminus S) = \mathrm{Aut}(\mathbb{P}, S)$.*

If S is a smooth cubic surface, then it does not contain any $(-K_S)$ -polar cylinder by Theorem 2.8. In this case, Conjecture 4.24 claims that $\text{Aut}(\mathbb{P} \setminus S) = \text{Aut}(\mathbb{P}, S)$, which is Conjecture 4.18.

In [171], Theorem 4.20 has been generalized as follows. Let X be a normal projective variety, and let D be an ample Cartier divisor on X . Suppose that the following conditions are satisfied:

- (1) the section ring of (X, D) is a hypersurface, i.e., one has

$$\bigoplus_{m=0}^{\infty} H^0(X, \mathcal{O}_X(mD)) \cong \mathbb{k}[x_0, x_1, \dots, x_n]/(F),$$

where $\mathbb{k}[x_0, \dots, x_n]$ is a polynomial ring in variables x_0, \dots, x_n with weights

$$a_0 = \text{wt}(x_0) \leq a_1 = \text{wt}(x_1) \leq \dots \leq a_n = \text{wt}(x_n),$$

and F is a quasi-homogeneous polynomial of degree d , so that X is a hypersurface in the weighted projective space $\mathbb{P}(a_0, a_1, \dots, a_n) = \text{Proj}(\mathbb{k}[x_0, x_1, \dots, x_n])$;

- (2) the Veronese map $v_d : \mathbb{P}(a_0, a_1, \dots, a_n) \dashrightarrow \mathbb{P}^N$ given by $|\mathcal{O}_{\mathbb{P}(a_0, a_1, \dots, a_n)}(d)|$ is an embedding.

Recall from [123, Proposition 3.5] that the complement $\mathbb{P}(a_0, a_1, \dots, a_n) \setminus X$ admits a nontrivial \mathbb{G}_a -action if and only if it is cylindrical. On the other hand, we have the following result:

Theorem 4.25 ([171, Theorem 3.1]). *Suppose that $\mathbb{P}(a_0, a_1, \dots, a_n) \setminus X$ has a nontrivial \mathbb{G}_a -action. Then X contains a D -polar cylinder.*

Based on the results on non-ruledness of smooth hypersurfaces of low degrees in the projective spaces such as [29, 50, 55, 113, 130, 194, 195, 203] one can extend Conjecture 4.18 as follows:

Conjecture 4.26. *Let X be a smooth hypersurface in \mathbb{P}^n of degree $d \geq 3$. Then*

$$\text{Aut}(\mathbb{P}^n \setminus X) = \text{Aut}(\mathbb{P}^n, X).$$

The conjecture holds when $d > n$ since the hypersurface X has non-negative Kodaira dimension. It remains true if $d = n \geq 4$ and $(n, d) = (4, 3)$ due to the results by [29, 50, 55, 113, 194, 195].

4.3. Compactifications of \mathbb{C}^n . In this subsection, we assume that varieties are defined over \mathbb{C} . In this case, the problem of existence of (Zariski open) cylinders in smooth Fano varieties is closely related to the following famous problem posed by Hirzebruch 65 years ago in [102].

Problem 4.27. *Find all complex analytic compactifications of \mathbb{C}^n with second Betti number 1.*

This problem asks to describe all compact complex manifolds X with $b_2(X) = 1$ that contain an open subset U which is biholomorphic to \mathbb{C}^n and whose complement $A = X \setminus U$ is a closed complex analytic subspace. Thus, we call a *compactification* of \mathbb{C}^n a pair (X, A) consisting of

- a compact complex manifold X with $b_2(X) = 1$;
- and a closed complex analytic subset $A \subset X$ such that $X \setminus A \xrightarrow{\text{bihol}} \mathbb{C}^n$.

A compactification (X, A) of \mathbb{C}^n is said to be *algebraic* if X is a smooth projective variety, and the biholomorphism $X \setminus A \xrightarrow{\text{bihol}} \mathbb{C}^n$ is an algebraic isomorphism. Thus, we see that

(X, A) is an algebraic compactification of $\mathbb{C}^n \implies X$ is a cylindrical Fano variety.

Proposition 4.28 ([25, 211]). *Let (X, A) be a compactification of \mathbb{C}^n . Then the following holds.*

- (1) A is purely 1-codimensional and irreducible;
- (2) $H^i(X, \mathbb{Z}) \cong H^i(A, \mathbb{Z})$, $H_i(X, \mathbb{Z}) \cong H_i(A, \mathbb{Z})$ for every $i \leq 2n - 2$;
- (3) $H^1(X, \mathbb{Z}) = 0$ and $H_1(X, \mathbb{Z}) = 0$;
- (4) the class of A generates the groups $H^2(X, \mathbb{Z}) \cong \mathbb{Z}$ and $H^2(A, \mathbb{Z}) \cong \mathbb{Z}$;
- (5) if X is Moishezon, then $H^1(X, \mathcal{O}_X) = 0$ and $H^2(X, \mathcal{O}_X) = 0$, so that $\text{Pic}(X) \cong H^2(X, \mathbb{Z})$.

The following deep result is due to Kodaira [129, Theorem 3]:

Theorem 4.29. *If (X, A) is a compactification of \mathbb{C}^n , then*

$$h^0(X, \omega_X^{\otimes m}) = 0$$

for every $m > 0$, where ω_X is the sheaf of holomorphic n -forms on X .

Thus, if (X, A) is a compactification of \mathbb{C}^n and X is projective, then X is a smooth Fano variety, and A is an ample divisor on X that generates $\text{Pic}(X)$.

Example 4.30. Let (X, A) be one of the following polarized smooth Fano varieties:

- (1) $X = \mathbb{P}^n$ and A is a hyperplane;
- (2) X is a smooth quadric in \mathbb{P}^{n+1} and A is its singular hyperplane section;
- (3) $X = \text{Gr}(m, k)$ and A is its Schubert subvariety of codimension 1, where $n = m(k - m)$;
- (4) $X = G/P$ and A is its open cell isomorphic to \mathbb{C}^n (such a cell does exist by [23, 131]), where G is a semisimple connected complex linear algebraic group, and P is its maximal parabolic subgroup.

Then (X, A) is a compactification of \mathbb{C}^n .

In two-dimensional case, Problem 4.27 has an easy solution: if (X, A) is a compactification of \mathbb{C}^2 , then $X = \mathbb{P}^2$ and A is a line in X . In the three-dimensional case, Problem 4.27 has been solved in the series of papers [77–82, 176, 177, 184]. In particular, we have the following result:

Theorem 4.31. *Let (X, A) be a compactification of \mathbb{C}^3 . Suppose that X is a projective threefold. Then this compactification is algebraic and (X, A) can be described as follows:*

- (1) $X = \mathbb{P}^3$ and A is a plane;
- (2) X is a smooth quadric in \mathbb{P}^4 and A is its singular hyperplane section;
- (3) X is the quintic del Pezzo threefold in \mathbb{P}^5 described in Example 1.31 and A is its singular hyperplane section that can be described as follows:
 - (a) a surface whose singular locus is a line L with normal bundle $\mathcal{N}_{L/X} = \mathcal{O}_L(1) \oplus \mathcal{O}_L(-1)$;
 - (b) a normal del Pezzo surface that has a unique singular point of type A_4 ;
- (4) X is a smooth Fano threefold of index 1 and genus 12 in \mathbb{P}^{13} and A is its certain hyperplane section whose singular locus is a line ℓ with normal bundle $\mathcal{N}_{\ell/X} = \mathcal{O}_\ell(1) \oplus \mathcal{O}_\ell(-2)$.

Proof. We know that X is a smooth Fano threefold, and the surface A generates $\text{Pic}(X)$, so that

$$-K_X \sim \iota(X)A,$$

where $\iota(X)$ is the Fano index of the threefold X . If $\iota(X) = 4$, then $X = \mathbb{P}^3$ and A is a plane. Similarly, if $\iota(X) = 3$, then X is a smooth quadric threefold in \mathbb{P}^4 , and A is its hyperplane section. In this case, the surface A must be singular, since $H^2(A, \mathbb{Z}) = \mathbb{Z}$ by Proposition 4.28.

If $\iota(X) = 1$, then the surface A must be a non-normal K3 surface, and the proof uses a delicate analysis of its singularities. As a result, one can show that X is a Fano threefold of genus 12 in \mathbb{P}^{13} , and A is its hyperplane section that is singular along a line of type $(1, -2)$. One construction of such compactification is described in Remark 3.12. We will not dwell into further details in this case.

Suppose that $\iota(X) = 2$. Let us show that X is the quintic del Pezzo threefold in \mathbb{P}^5 , and A is its singular hyperplane section described above. Note that in this case (X, A) is indeed a compactification of \mathbb{C}^3 , which follows from the proof of Theorem 3.6.

By Proposition 4.28, we have $H^2(A, \mathbb{Z}) = \mathbb{Z}$ and

$$(4.32) \quad 4 + 2h^{1,2}(X) = \chi_{\text{top}}(X) = \chi_{\text{top}}(A) + 1.$$

First, we suppose that the surface A is normal. Then $-K_A$ is ample by the adjunction formula, so that A is a del Pezzo surface with isolated Gorenstein singularities. If its singularities are worse

than Du Val, then A must be a (generalized) cone over an elliptic curve [101], so that $\chi_{\text{top}}(A) = 1$. The latter contradicts (4.32). Thus, we see that A is a del Pezzo surface with Du Val singularities. Then $\rho(A) = 1$, because $H^2(A, \mathbb{Z}) = \mathbb{Z}$. Then $\chi_{\text{top}}(A) = 3$, so that we have $h^{1,2}(X) = 0$ by (4.32). Now, using Remark 3.1, we conclude that X is the quintic del Pezzo threefold in \mathbb{P}^5 as required. Moreover, we have $K_A^2 = 5$, so that A is a quintic del Pezzo surface that has Du Val singularities. Since $\rho(A) = 1$, it follows from [77, 156] that A has a unique singular point of type A_4 .

Now, we suppose that A is non-normal, so that it has a singular locus of positive dimension. It is easy to show that any hyperplane section of a smooth complete intersection has only isolated singularities, and the same result holds for hyperplane sections of weighed smooth hypersurfaces. Therefore, using Remark 3.1, we conclude again that X is the quintic del Pezzo threefold in \mathbb{P}^5 , and A is its hyperplane section. Using the adjunction formula, we see that a general hyperplane section of the surface A is an irreducible singular curve of arithmetic genus 1, so that it has one singular point. Thus, the non-normal locus of the surface A is some line L . Hence, it follows from the proof of Theorem 3.6 that $\text{Sing}(A) = L$ and

$$X \setminus A \cong Q \setminus H$$

where Q is a smooth quadric threefold in \mathbb{P}^4 , and H is its hyperplane section. Since $X \setminus A \cong \mathbb{C}^3$, we conclude that the surface H is singular. As we already mentioned in the proof of Theorem 3.6, this implies that $\mathcal{N}_{L/X} = \mathcal{O}_L(1) \oplus \mathcal{O}_L(-1)$ as required. \square

Corollary 4.33. *Let (X, A) be a compactification of \mathbb{C}^3 . Suppose that X is a projective threefold. Then $H^k(X, \mathbb{Z}) \cong H^k(\mathbb{P}^3, \mathbb{Z})$ for all k .*

It would be interesting to find an alternative proof of Theorem 4.31 that does not heavily rely on the classification of smooth Fano threefolds.

Remark 4.34. Let X be a smooth Fano threefold such that $\rho(X) = 1$, $\iota(X) = 1$ and $g(X) = 12$. If X is a compactification of \mathbb{C}^4 , then X contains a line ℓ such that $\mathcal{N}_{\ell/X} = \mathcal{O}_\ell(1) \oplus \mathcal{O}_\ell(-2)$. However, this condition does not always guarantee that X is a compactification of \mathbb{C}^4 (see [184]).

Remark 4.35. The list in Theorem 4.31 is similar to the list in Theorem 3.5.

In higher dimensions, we know very few results on Problem 4.27. Let us present one of them, which follows from Theorem 3.20 and its proof. We use here the notation introduced in Section 3.2.

Theorem 4.36 ([186]). *Let (X, A) be a compactification of \mathbb{C}^4 , where X is a smooth Fano fourfold. Suppose that $\iota(X) = 3$. Then X is the quintic del Pezzo fourfold in \mathbb{P}^7 and*

- (1) *either $A = H_\ell$, where ℓ is a line in X that is not a line of type (b);*
- (2) *or A is a singular hyperplane section of the del Pezzo fourfold X such that its singular locus consists of a single ordinary double point that is not contained in the divisor \mathcal{H} .*

Each of these compactifications is algebraic and unique up to isomorphism.

Proof. We prove the existence part only. In the first case, the existence follows from Theorem 3.20. To deal with the second case, let us use the notation introduced in the proof of Theorem 3.20. Consider the Sarkisov link (3.21) with ℓ being a line of type (a). We already know that Q is smooth, and so we may assume that it is given in \mathbb{P}^4 by

$$x_2x_3 + x_1x_4 + x_0x_5 = 0.$$

Similarly, we may assume that $\eta(F)$ is cut out by $x_0 = 0$. Moreover, the surface $\eta(\widehat{H}_\ell)$ is a smooth cubic scroll in this case. Hence, we may assume that it is cut out on Q by

$$\begin{cases} x_0 = 0, \\ x_2x_4 + x_1x_5 = 0, \\ x_4^2 - x_3x_5 = 0. \end{cases}$$

Let D be the hyperplane section of the quadric Q that is cut out by $x_3 = 0$, and let \widehat{D} be its proper transform on \widehat{V}_5 . Then D is singular. We claim that $\widehat{V}_5 \setminus (\widehat{D} \cup \widehat{H}_\ell) \cong \mathbb{A}^4$. Indeed, let $U = Q \setminus D$. Then $U \cong \mathbb{A}^4$ with coordinates $y_0 = \frac{x_0}{x_3}$, $y_1 = \frac{x_1}{x_3}$, $y_4 = \frac{x_4}{x_3}$, $y_5 = \frac{x_5}{x_4}$, so that $\widehat{V}_5 \setminus \widehat{D}$ is given by

$$y_0 z_0 = (y_5 - y_4^2) z_1$$

in $\mathbb{A}^4 \times \mathbb{P}^1$, where z_0 and z_1 are coordinates on \mathbb{P}^1 . Then $\widehat{V}_5 \setminus (\widehat{D} \cup \widehat{H}_\ell)$ is given in $\mathbb{A}^4 \times \mathbb{A}^1$ by

$$y_0 z = y_5 - y_4^2,$$

where $z = \frac{z_0}{z_1}$. This implies that $\widehat{V}_5 \setminus (\widehat{D} \cup \widehat{H}_\ell) \cong \mathbb{A}^4$. Now, observe that $\pi(\widehat{D})$ is a hyperplane section of V_5 whose singular locus consists of a single ordinary double point not contained in \mathcal{H} . \square

In dimension 4, we know very few compactifications (X, A) of \mathbb{C}^4 . They can be listed as follows:

- $X = \mathbb{P}^4$ and A is a hyperplane;
- X is a smooth quadric and A is its singular hyperplane section;
- X is the del Pezzo quintic fourfold and A is described in Theorem 4.36;
- X is a smooth Fano–Mukai fourfold of genus 10 and A is described in Theorem 3.23.

In particular, in every known example of a compactification (X, A) of \mathbb{C}^4 with $X \not\cong \mathbb{P}^4$, one has

$$H^k(X, \mathbb{Z}) \cong H^k(Q, \mathbb{Z})$$

for all k , where Q is a smooth quadric in \mathbb{P}^5 . We wonder whether this is just a coincidence.

Question 4.37. *Does there exist a smooth Fano fourfold of index 1 that is a compactification of \mathbb{C}^4 ?*

Before we conclude this survey, let us set the following question:

Question 4.38. *Is it true that any compactification of \mathbb{C}^n is rational?*

Note that the answer to this question is not obvious, since the isomorphism $X \setminus A \cong \mathbb{C}^n$ in the definition of a compactification of \mathbb{C}^n is a biholomorphism, which is not necessarily algebraic.

APPENDIX A. SINGULARITIES OF PAIRS

Let S be a surface with at most quotient singularities, let D be an effective non-zero \mathbb{Q} -divisor on S , let P be a point of S , and let

$$D = \sum_{i=1}^r a_i C_i,$$

where C_1, \dots, C_r are distinct irreducible curves on S , and each a_i is a non-negative rational number. We call (S, D) a log pair.

Let $\pi: \widetilde{S} \rightarrow S$ be a birational morphism such that \widetilde{S} is smooth. For each C_i , denote by \widetilde{C}_i its proper transform on the surface \widetilde{S} . Let F_1, \dots, F_n be π -exceptional curves. Then

$$K_{\widetilde{S}} + \sum_{i=1}^r a_i \widetilde{C}_i + \sum_{j=1}^n b_j F_j \sim_{\mathbb{Q}} \pi^*(K_S + D)$$

for some rational numbers b_1, \dots, b_n . Suppose that $\widetilde{C}_1 + \dots + \widetilde{C}_2 + F_1 + \dots + F_n$ is a divisor with simple normal crossings. Then we say that $\pi: \widetilde{S} \rightarrow S$ is a log resolution of the log pair (S, D) .

Definition A.1. The log pair (S, D) is said to be log canonical at the point P if the following two conditions are satisfied:

- $a_i \leq 1$ for every C_i such that $P \in C_i$;
- $b_j \leq 1$ for every F_j such that $\pi(F_j) = P$.

The log pair (S, D) is called log canonical if it is log canonical at every point of S .

This definition does not depend on the choice of the log resolution $\pi: \tilde{S} \rightarrow S$.

Remark A.2. Let R be an effective \mathbb{Q} -divisor on S such that $R \sim_{\mathbb{Q}} D$. For a rational number ϵ , let

$$D_{\epsilon} = (1 + \epsilon)D - \epsilon R.$$

Then $D_{\epsilon} \sim_{\mathbb{Q}} D$. Suppose that $R \neq D$. Then there exists the greatest rational number $\epsilon_0 \geq 0$ such that the divisor D_{ϵ_0} is effective. By construction, the support of the divisor D_{ϵ_0} does not contain at least one curve contained in the support of the divisor R . Moreover, if (S, D) is not log canonical at P , but (S, R) is log canonical at P , then (S, D_{ϵ_0}) is not log canonical at P .

Now, we suppose that the surface S is smooth at P .

Lemma A.3. *Suppose that (S, D) is not log canonical at P . Then $\text{mult}_P(D) > 1$.*

Proof. Left to the reader. \square

Let $f: \bar{S} \rightarrow S$ be a blowup of the point P , and let E be the f -exceptional curve. Denote by \bar{D} the proper transform of the \mathbb{Q} -divisor D on the surface \bar{S} via f . Then the log pair

$$(A.4) \quad (\bar{S}, \bar{D} + (\text{mult}_P(D) - 1)E)$$

is called the log pull back of the log pair (S, D) on the surface \bar{S} .

Lemma A.5. *Suppose that the log pair (S, D) is not log canonical at P . Then*

- (i) *the \mathbb{Q} -divisor $\bar{D} + (\text{mult}_P(D) - 1)E$ is effective;*
- (ii) *the log pair (A.4) is not log canonical at some point $Q \in E$.*

Proof. The required assertion follows from Definition A.1 and Lemma A.3. \square

The following handy statement is a very special case of a much more general result, which is known as *Inversion of Adjunction* (see, for example, [134, Theorem 6.29]).

Lemma A.6 ([134, Exercise 6.31]). *Suppose that C_1 is smooth at P , the log pair (S, D) is not log canonical at P , and $a_1 \leq 1$. Let $\Delta = a_2 C_2 + \cdots + a_r C_r$. Then $(C_1 \cdot \Delta)_P > 1$.*

Proof. Let $m = \text{mult}_P(\Delta)$. If $m > 1$, then we are done, since

$$(C_1 \cdot \Delta)_P \geq m.$$

Therefore, we may assume that $m \leq 1$. This implies that the log pair (S, D) is log canonical in a punctured neighborhood of the point $P \in S$. Since the log pair (S, D) is not log canonical at P , there exists a birational morphism $h: \hat{S} \rightarrow S$ that is a composition of $s \geq 1$ blowups of points dominating P such that $e_s > 1$, where e_s is a rational number determined by

$$K_{\hat{S}} + a_1 \hat{C}_1 + \hat{\Delta} + \sum_{i=1}^s e_i E_i \sim_{\mathbb{Q}} h^*(K_S + D),$$

where each e_i is a rational number, each E_i is an h -exceptional divisor, $\hat{\Delta}$ is a proper transform on the surface \hat{S} of the divisor Δ , and \hat{C}_1 is a proper transform on \hat{S} of the curve C_1 .

Let $\bar{\Delta}$ and \bar{C}_1 be the proper transforms on \bar{S} of the divisor Δ and the curve C_1 , respectively. Then $(\bar{S}, a_1 \bar{C}_1 + (a_1 + m - 1)E + \bar{\Delta})$ is not log canonical at some point $Q \in E$ by Lemma A.5.

Let us prove the inequality $(C_1 \cdot \Delta)_P > 1$ by induction on s . If $s = 1$, then

$$a_1 + m - 1 > 1,$$

which implies that $m > 2 - a_1 \geq 1$, so that $(C_1 \cdot \Delta)_P \geq m > 1$ as required. Thus, we may assume that $s \geq 2$ and $a_1 + m - 1 \leq 2$. Since

$$(C_1 \cdot \Delta)_P \geq m + (\bar{C}_1 \cdot \bar{\Delta})_Q,$$

it is enough to show that $m + (\overline{C}_1 \cdot \overline{\Delta})_Q > 1$. We may also assume that $m \leq 1$, since $(C_1 \cdot \Delta)_P \geq m$.

If $Q \notin \overline{C}_1$, then $(\overline{S}, (a_1 + m - 1)E + \overline{\Delta})$ is not log canonical at the point Q , which gives

$$m = \overline{\Delta} \cdot E \geq (\overline{\Delta} \cdot E)_Q > 1$$

by induction. The latter implies that $Q = \overline{C}_1 \cap E$, since $m \leq 1$. Then

$$a_1 + m - 1 + (\overline{C}_1 \cdot \overline{\Delta})_Q = \left(((a_1 + m - 1)E + \overline{\Delta}) \cdot \overline{C}_1 \right)_Q > 1$$

by induction. This gives $(\overline{C} \cdot \overline{\Delta})_Q > 2 - a_1 - m$. Then

$$m + (\overline{C}_1 \cdot \overline{\Delta})_Q > 2 - a_1 \geq 1$$

as required. \square

Corollary A.7. *In the notation and assumptions of Lemma A.5, suppose that $\text{mult}_P(D) \leq 2$. Then there exists a unique point $Q \in E$ such that (A.4) is not log canonical at Q .*

Proof. If (A.4) is not log canonical at two distinct points P_1 and P_2 , then

$$2 \geq \text{mult}_P(D) = \overline{D} \cdot E \geq (\overline{D} \cdot E)_{P_1} + (\overline{D} \cdot E)_{P_2} > 2$$

by Lemma A.6. Now use Lemma A.5. \square

The following result plays an essential role in the proof of Theorem 2.15 given in Section 2.2. In fact, this theorem has been discovered [31] in an attempt to give a simple proof of Theorem 2.15, since its original proof in [38] is very technical. For other applications of Theorem 2.15, see [32, 212].

Theorem A.8 ([31]). *Suppose that $(C_1 \cdot C_2)_P = 1$, and the log pair (S, D) is not log canonical at P . Let $\Delta = a_3 C_3 + \cdots + a_r C_r$ and $m = \text{mult}_P(\Delta)$. Suppose also that $m \leq 1$. Then*

$$(C_1 \cdot \Delta)_P > 2(1 - a_2)$$

or

$$(C_2 \cdot \Delta)_P > 2(1 - a_1).$$

Proof. We may assume that $a_1 \leq 1$ and $a_2 \leq 1$. There is a morphism $h: \widehat{S} \rightarrow S$ that is a composition of $s \geq 1$ blowups of points dominating P such that $e_s > 1$ for $e_s \in \mathbb{Q}$ that is determined by

$$K_{\widehat{S}} + a_1 \widehat{C}_1 + a_2 \widehat{C}_2 + \widehat{\Delta} + \sum_{i=1}^r e_i E_i = h^*(K_S + a_1 C_1 + a_2 C_2 + \Delta),$$

where each e_i is a rational number, each E_i is an h -exceptional divisor, \widehat{C}_1 and \widehat{C}_2 , are proper transforms on \widehat{S} of the curves C_1 and C_2 , respectively, and $\widehat{\Delta}$ is a proper transform of the divisor Δ .

Let $\overline{\Delta}$, \overline{C}_1 , \overline{C}_2 be the proper transforms on \overline{S} of the divisors Δ , C_1 and C_2 , respectively. Then

$$(\overline{S}, a_1 \overline{C}_1 + a_2 \overline{C}_2 + (a_1 + a_2 + m - 1)E + \overline{\Delta})$$

is not log canonical at some point $Q \in E$ by Lemma A.5.

If $s = 1$, then $a_1 + a_2 + m - 1 > 1$. If $m > 2 - a_1 - a_2$, then $m > 2(1 - a_1)$ or $m > 2(1 - a_2)$, because otherwise we would have

$$2m \leq 4 - 2(a_1 + a_2),$$

which contradicts to $m > 2 - a_1 - a_2$. Then $(\Delta \cdot C_1)_P > 2(1 - a_2)$ or $(\Delta \cdot C_2)_P > 2(1 - a_1)$ if $s = 1$.

Let us prove the required assertion by induction on s . The case $s = 1$ is already done, so that we may assume that $s \geq 2$ and $a_1 + a_2 + m \leq 2$. If $Q \neq E \cap \overline{C}_1$ and $Q \neq E \cap \overline{C}_2$, then

$$m = \overline{\Delta} \cdot E > 1$$

by Lemma A.6, which is impossible by assumption. Thus, either $Q = E \cap \overline{C}_1$ or $Q = E \cap \overline{C}_2$. Without loss of generality, we may assume that $Q = E \cap \overline{C}_1$.

By induction, we can apply the lemma to $(\overline{S}, a_1 \overline{C}_1 + (a_1 + a_2 + m - 1)E + \overline{\Delta})$ at the point Q . This implies that either

$$(\overline{\Delta} \cdot \overline{C}_1)_Q > 2(1 - (a_1 + a_2 + m - 1)) = 4 - 2a_1 - 2a_2 - 2m$$

or $(\overline{\Delta} \cdot E)_Q > 2(1 - a_1)$. In the latter case, we have

$$(\Delta \cdot C_2)_P \geq m = \overline{\Delta} \cdot E \geq (\overline{\Delta} \cdot E)_Q > 2(1 - a_1),$$

which is exactly what we want. Therefore, we may assume that $(\overline{\Delta} \cdot \overline{C}_1)_Q > 4 - 2a_1 - 2a_2 - 2m$. If $(\Delta \cdot C_2)_P > 2(1 - a_1)$, then we are done. Hence, we may assume $(\Delta \cdot C_2)_P \leq 2(1 - a_1)$. Then

$$m \leq (\Delta \cdot C_2)_P \leq 2(1 - a_1).$$

This gives

$$(\Delta \cdot C_1)_P \geq m + (\overline{\Delta} \cdot \overline{C}_1)_Q > m + 4 - 2a_1 - 2a_2 - 2m > 2(1 - a_2),$$

because $m \leq 2(1 - a_1)$. □

Almost all results we have considered so far in this subsection are local (except for Remark A.2). Let us conclude this subsection by two global statements. The first of them is due to Puhlikov:

Lemma A.9 ([134, Lemma 5.36]). *Suppose that S is a smooth surface in \mathbb{P}^3 , and D is \mathbb{Q} -linearly equivalent to its hyperplane section. Then each a_i does not exceed 1.*

Proof. Let X be a cone over the curve C_i whose vertex is a general enough point in \mathbb{P}^3 . Then

$$X \cap S = C_i + \widehat{C}_i,$$

where \widehat{C}_i is an irreducible curve of degree $(\deg(S) - 1)\deg(C_i)$. Moreover, \widehat{C}_i is not contained in the support of the divisor D , and the intersection $C_i \cap \widehat{C}_i$ consists of exactly $\deg(\widehat{C}_i)$ points. Then

$$\deg(\widehat{C}_i) = D \cdot \widehat{C}_i \geq a_i C_i \cdot \widehat{C}_i \geq a_i \deg(\widehat{C}_i),$$

which implies that $a_i \leq 1$. □

The second global result we want to mention is the following lemma about del Pezzo surfaces of degree 2 that have at most two ordinary double points.

Lemma A.10. *Suppose that there is a double cover $\tau: S \rightarrow \mathbb{P}^2$ branched over an irreducible quartic curve B that has at most two ordinary double points, and*

$$D \sim_{\mathbb{Q}} -K_S.$$

Then each a_i does not exceed 1. Moreover, if (S, D) is not log canonical at P , then $\tau(P) \in B$.

Proof. Write $D = a_1 C_1 + \Delta$, where $\Delta = a_2 C_2 + \cdots + a_r C_r$. Suppose that $a_1 > 1$. Let us seek for a contradiction. Since

$$2 = -K_S \cdot D = -K_S \cdot (a_1 C_1 + \Delta) = -a_1 K_S \cdot C_1 - K_S \cdot \Delta \geq -a_1 K_S \cdot C_1 > -K_S \cdot C_1,$$

we have $-K_S \cdot C_1 = 1$. Then $\tau(C_1)$ is a line. Hence, the surface S contains an irreducible curve Z_1 such that $C_1 + Z_1 \sim -K_S$ and $\tau(C_1) = \tau(Z_1)$. Note that the curves C_1 and Z_1 are interchanged by the biregular involution of the surface S induced by the double cover τ . Then

$$2 = (-K_S)^2 = (C_1 + Z_1)^2 = 2C_1^2 + 2C_1 \cdot Z_1,$$

which implies that $C_1 \cdot Z_1 = 1 - C_1^2$. Since C_1 and Z_1 are smooth rational curves, we have

$$C_1^2 = Z_1^2 = -1 + \frac{k}{2},$$

where k is the number of singular points of S that lie on C_1 . Now we write $D = a_1 C_1 + b_1 Z_1 + \Theta$, where b_1 is a non-negative rational number, and Θ is an effective \mathbb{Q} -divisor whose support does not contain the curves C_1 and Z_1 . Then

$$1 = C_1 \cdot (a_1 C_1 + b_1 Z_1 + \Theta) \geq a_1 C_1^2 + b_1 C_1 \cdot Z_1 = a_1 C_1^2 + b_1 (1 - C_1^2),$$

and hence $1 \geq a_1 C_1^2 + b_1 (1 - C_1^2)$. Similarly, from $Z_1 \cdot D = 1$, we obtain

$$1 \geq b_1 C_1^2 + a_1 (1 - C_1^2).$$

The obtained two inequalities imply that $a_1 \leq 1$ and $b_1 \leq 1$, because $C_1^2 = -1 + \frac{k}{2}$ and $k \leq 2$. Since $a_1 > 1$ by our assumption, this is a contradiction.

We see that $a_1 \leq 1$. Similarly, we see that $a_i \leq 1$ for every i .

Now we suppose that the log pair (S, D) is not log canonical at P . Let us show that $\tau(P) \in B$. Suppose that $\tau(P) \notin B$. Then S is smooth at P . Let us seek for a contradiction.

Let H be a general curve in $| -K_S |$ that passes through the point P . Then

$$2 = H \cdot D \geq \text{mult}_P(H) \text{mult}_P(D) \geq \text{mult}_P(D),$$

so that $\text{mult}_P(D) \leq 2$. But the pair (A.4) is not log canonical at some point $Q \in E$ by Lemma A.5. Applying Lemma A.3 to (A.4), we get $\text{mult}_P(D) + \text{mult}_Q(\overline{D}) > 2$.

Since $\tau(P) \notin B$, there exists a unique (possibly reducible) curve $R \in | -K_S |$ such that its proper transform on \overline{S} passes through the point Q . Note that R is smooth at P . This enables us to assume that the support of D does not contain at least one irreducible component of R by Remark A.2. Denote by \overline{R} the proper transform of R on the surface \overline{R} . If the curve R is irreducible, then

$$2 - \text{mult}_P(D) = 2 - \text{mult}_P(C) \text{mult}_P(D) = \overline{R} \cdot \overline{D} \geq \text{mult}_Q(\overline{R}) \text{mult}_Q(\overline{D}) = \text{mult}_Q(\overline{D}),$$

which is impossible, since $\text{mult}_P(D) + \text{mult}_Q(\overline{D}) > 2$. Thus, the curve R must be reducible.

Write $R = R_1 + R_2$, where R_1 and R_2 are irreducible smooth curves. Without loss of generality we may assume that the curve R_1 is not contained in $\text{Supp}(D)$. Then $P \in R_2$, because otherwise we would have

$$1 = D \cdot R_1 \geq \text{mult}_P(D) > 1,$$

since $\text{mult}_P(D) > 1$ by Lemma A.3. Thus, we put $D = aR_2 + \Omega$, where a is a non-negative rational number and Ω is an effective \mathbb{Q} -divisor whose support does not contain the curve R_2 . Then

$$1 = R_1 \cdot D = \left(2 - \frac{1}{2}l\right) a + R_1 \cdot \Omega \geq \left(2 - \frac{1}{2}l\right) a,$$

where l is the number of singular points of the surface S contained in R_1 . Denote by \overline{R}_2 the proper transform on \overline{S} of the curve R_2 , and denote by $\overline{\Omega}$ the proper transform on \overline{S} of the divisor Ω . Then the log pair

$$\left(\overline{S}, a\overline{R}_2 + \overline{\Omega} + (\text{mult}_P(D) - 1)E\right)$$

is not log canonical at Q . Note that we already proved that $a \leq 1$. Thus, using Lemma A.6, we get

$$\left(2 - \frac{1}{2}l\right) a = \overline{R}_2 \cdot \left(\overline{\Omega} + (\text{mult}_P(D) - 1)E\right) > 1.$$

This is a contradiction. □

REFERENCES

- [1] J. ALPER, H. BLUM, D. HALPERN-LEISTNER, AND C. XU, *Reductivity of the automorphism group of K -polystable Fano varieties*, Arxiv e-print, arXiv:1906.03122 (2019).
- [2] T. ANDO, *On extremal rays of the higher-dimensional varieties*, Invent. Math., 81 (1985), pp. 347–357.
- [3] M. ANDREATTA AND J. WIŚNIEWSKI, *On contractions of smooth varieties*, J. Algebraic Geom., 7 (1998), pp. 253–312.
- [4] R. ANDRIST, *The density property for Calogero–Moser spaces*, Arxiv e-print, arXiv:2006.11936 (2020).
- [5] M. ARTIN AND D. MUMFORD, *Some elementary examples of unirational varieties which are not rational*, Proc. London Math. Soc., 25 (1972), pp. 75–95.
- [6] I. ARZHANTSEV, *On rigidity of factorial trinomial hypersurfaces*, Internat. J. Algebra Comput., 26 (2016), pp. 1061–1070.
- [7] ———, *Polynomial curves on trinomial hypersurfaces*, Acta Arith., 186 (2018), pp. 87–99.
- [8] I. ARZHANTSEV, H. FLENNER, S. KALIMAN, F. KUTZSCHEBAUCH, AND M. ZAIDENBERG, *Flexible varieties and automorphism groups*, Duke Math. J., 162 (2013), pp. 767–823.
- [9] I. ARZHANTSEV AND S. GAIFULLIN, *The automorphism group of a rigid affine variety*, Math. Nachr., 290 (2017), pp. 662–671.
- [10] I. ARZHANTSEV, A. PEREPECHKO, AND H. SÜSS, *Infinite transitivity on universal torsors*, J. Lond. Math. Soc., 89 (2014), pp. 762–778.
- [11] I. ARZHANTSEV AND M. ZAIDENBERG, *Tits-type alternative for groups acting on toric affine varieties*, Int. Math. Res. Not. (to appear), doi:10.1093/imrn/rnaa342; Arxiv e-print, arXiv:2003.00037 (2020).
- [12] I. ARZHANTSEV, M. ZAIDENBERG, AND K. KUYUMZHIYAN, *Flag varieties, toric varieties, and suspensions: three examples of infinite transitivity*, Mat. Sb., 203 (2012), pp. 3–30.
- [13] ———, *Infinite transitivity, finite generation, and Demazure roots*, Adv. Math. 351 (2019), pp. 1–32.
- [14] V. BATYREV AND Y. MANIN, *Sur le nombre des points rationnels de hauteur borné des variétés algébriques*, Math. Ann., 286 (1990), pp. 27–43.
- [15] A. BEAUVILLE, *Varieties de Prym et jacobienne intermédiaires*, Ann. Sci. Ecole Norm. Sup., 10 (1977), pp. 309–391.
- [16] G. BELOUSOV, *Cylinders in del Pezzo surfaces with du Val singularities*, Bull. Korean Math. Soc., 54 (2017), pp. 1655–1667.
- [17] O. BENOIST AND O. WITTENBERG, *Intermediate jacobians and rationality over arbitrary fields*, Arxiv e-print, arXiv:1909.12668 (2019).
- [18] Y. BEREST, A. ESHMATOV, AND F. ESHMATOV, *Multitransitivity of Calogero–Moser spaces*, Transform. Groups, 21 (2016), pp. 35–50.
- [19] G. BODNAR, H. HAUSER, J. SCHICHO, AND O. VILLAMAYOR, *Plain varieties*, Bull. Lond. Math. Soc., 40 (2008), pp. 965–971.
- [20] F. BOGOMOLOV AND C. BÖHNING, *On uniformly rational varieties*, Providence, RI: American Mathematical Society, 2014, pp. 33–48.
- [21] I. BOLDYREV AND S. GAIFULLIN, *Automorphisms of nonnormal toric varieties*, Arxiv e-print, arXiv:2012.03346 (2020).
- [22] A. BOREL, *Les bouts des espaces homogènes de groupes de Lie*, Ann. Math. (2), 58 (1953), pp. 443–457.
- [23] ———, *Kählerian coset spaces of semisimple Lie groups*, Proc. Natl. Acad. Sci. USA, 40 (1954), pp. 1147–1151.
- [24] V. BOROVIK AND S. GAIFULLIN, *m-suspensions over rigid varieties*, Arxiv e-print, arXiv:2007.07882 (2020).
- [25] L. BRENTON AND J. MORROW, *Compactifications of \mathbf{C}^n* , Trans. Amer. Math. Soc., 246 (1978), pp. 139–153.
- [26] F. CAMPANA, *Connexité rationnelle des variétés de Fano*, Ann. Sci. École Norm. Sup. (4), 25 (1992), pp. 539–545.
- [27] A.-M. CASTRAVET, *Examples of Fano varieties of index one that are not birationally rigid*, Proc. Amer. Math. Soc., 135 (2007), pp. 3783–3788.
- [28] I. CHELTSOV, *Cylinders in rational surfaces*, Mat. Sb., 212:3 (2021), pp. 139–156.
- [29] ———, *On smooth quintic 4-fold*, Sb. Math., 191 (2000), pp. 1399–1419.
- [30] ———, *Log canonical thresholds of del Pezzo surfaces*, Geom. Funct. Anal., 11 (2008), pp. 1118–1144.
- [31] ———, *Del Pezzo surfaces and local inequalities*, vol. 79 of Springer Proc. Math. Stat., Springer, 2014, pp. 83–101.

- [32] I. CHELTSOV, Y. AHMADINEZHAD, AND J. SCHICHO, *On a conjecture of Tian*, Math. Z., 288 (2018), pp. 217–241.
- [33] I. CHELTSOV, A. DUBOULOZ, AND J. PARK, *Super-rigid affine Fano varieties*, Compos. Math., 154 (2018), pp. 2462–2484.
- [34] I. CHELTSOV AND D. KOSTA, *Computing α -invariants of singular del Pezzo surfaces*, J. Geom. Anal., 24 (2014), pp. 798–842.
- [35] I. CHELTSOV AND J. PARK, *Sextic double solids*, vol. 282 of Progress in Math., Birkhauser Boston, 2010, pp. 75–132.
- [36] I. CHELTSOV, J. PARK, AND C. SHRAMOV, *Exceptional del Pezzo hypersurfaces*, J. Geom. Anal., 20 (2010), pp. 787–816.
- [37] I. CHELTSOV, J. PARK, AND J. WON, *Log canonical thresholds of certain Fano hypersurfaces*, Math. Z., 276 (2014), pp. 51–79.
- [38] ———, *Affine cones over smooth cubic surfaces*, J. Eur. Math. Soc. (JEMS), 18 (2016), pp. 1537–1564.
- [39] ———, *Cylinders in singular del Pezzo surfaces*, Compos. Math., 152 (2016), pp. 1198–1224.
- [40] ———, *Cylinders in del Pezzo surfaces*, Int. Math. Res. Not., (2017), pp. 1179–1230.
- [41] I. CHELTSOV AND Y. PROKHOROV, *Del Pezzo surfaces with infinite automorphism groups*, Algebr. Geom., 8 (2021), pp. 319–357.
- [42] I. CHELTSOV, Y. PRZYJALKOWSKI, AND C. SHRAMOV, *Fano threefolds with infinite automorphism groups*, Izv. Math., 83 (2019), pp. 860–907.
- [43] ———, *Which quartic double solids are rational?*, J. Alg. Geom., 28 (2019), pp. 201–243.
- [44] I. CHELTSOV AND C. SHRAMOV, *Log canonical thresholds of smooth Fano threefolds*, Russian Math. Surveys, 63 (2008), pp. 71–178.
- [45] ———, *Extremal metrics on del Pezzo threefolds*, Proc. of the Steklov Inst. of Math., 264 (2009), pp. 30–44.
- [46] ———, *Del Pezzo zoo*, Experimental Math., 22 (2013), pp. 313–326.
- [47] ———, *Cremona groups and the icosahedron*, Boca Raton, FL: CRC Press, 2016.
- [48] X.-X. CHEN, S. DONALDSON, AND S. SUN, *Kähler–Einstein metrics on Fano manifolds. I, II, III*, J. Amer. Math. Soc., 28 (2015), pp. 183–197, 199–234, 235–278.
- [49] M. CHITAYAT AND D. DAIGLE, *On the rigidity of certain Pham–Brieskorn rings*, Arxiv e-print, arXiv:1907.13259 (2019).
- [50] C. CLEMENS AND P. GRIFFITHS, *The intermediate Jacobian of the cubic threefold*, Ann. of Math., 95 (1972), pp. 281–356.
- [51] A. CORTI, *Singularities of linear systems and 3-fold birational geometry*, L.M.S. Lecture Note Series, 281 (2000), pp. 259–312.
- [52] S. CYNK, *Defect of a nodal hypersurface*, Manuscripta Math., 104 (2001), pp. 325–331.
- [53] D. DAIGLE, G. FREUDENBURG, AND L. MOSER-JAUSLIN, *Locally nilpotent derivations of rings graded by an abelian group*, vol. 75, Math. Soc. Japan, Tokyo, 2017, pp. 29–48.
- [54] T. DE FERNEX, *Negative curves on very general blow-ups of \mathbb{P}^2* , de Gruyter, Berlin, 2005, pp. 199–207.
- [55] ———, *Birationally rigid hypersurfaces*, Invent. Math., 192 (2013), pp. 533–566.
- [56] A. DUBOULOZ, *Flexible bundles over rigid affine surfaces*, Comment. Math. Helv. 90 (2015), pp. 121–137.
- [57] ———, *Exotic \mathbb{G}_a -quotients of $\mathrm{SL}_2 \times \mathbb{A}^1$* , Eur. J. Math. 5 (2019), pp. 828–844.
- [58] A. DUBOULOZ AND T. KISHIMOTO, *Log-uniruled affine varieties without cylinder-like open subsets*, Bull. Soc. Math. France, 143 (2015), pp. 383–401.
- [59] ———, *Families of affine ruled surfaces: existence of cylinders*, Nagoya Math. J., 223 (2016), pp. 1–20.
- [60] ———, *Explicit biregular/birational geometry of affine threefolds: completions of \mathbb{A}^3 into del Pezzo fibrations and Mori conic bundles*, vol. 75 of Adv. Stud. Pure Math., Math. Soc. Japan, Tokyo, 2017, pp. 49–71.
- [61] ———, *Cylinders in del Pezzo fibrations*, Israel J. Math., 225 (2018), pp. 797–815.
- [62] ———, *Cylindres dans les fibrations de Mori: formes du volume quintique de del Pezzo*, Ann. Inst. Fourier, 69 (2019), pp. 2377–2393.
- [63] ———, *Deformations of \mathbb{A}^1 -cylindrical varieties*, Math. Ann., 373 (2019), pp. 1135–1149.
- [64] A. DUBOULOZ, L. MOSER-JAUSLIN, AND P.-M. POLONI, *Inequivalent embeddings of the Koras–Russell cubic 3-fold*, Michigan Math. J., 59 (2010), pp. 679–694.
- [65] H. FLENNER, S. KALIMAN, AND M. ZAIDENBERG, *A Gromov–Winkelmann type theorem for flexible varieties*, J. Eur. Math. Soc., 18 (2016), pp. 2483–2510.

- [66] H. FLENNER AND M. ZAIDENBERG, *Rational curves and rational singularities*, Math. Z., 244 (2003), pp. 549–575.
- [67] G. FREUDENBURG, *Algebraic theory of locally nilpotent derivations*, vol. 136 of Encyclopaedia of Mathematical Sciences, Springer-Verlag, Berlin, second ed., 2017.
- [68] G. FREUDENBURG AND P. RUSSELL, *Open problems in affine algebraic geometry*, vol. 369 of Contemp. Math., Amer. Math. Soc., Providence, RI, 2005, pp. 1–30.
- [69] K. FUJITA, *Examples of K -unstable Fano manifolds with the Picard number 1*, Proc. the Edinburgh Math. Soc., 60 (2017), pp. 881–891.
- [70] T. FUJITA, *On the structure of polarized manifolds with total deficiency one. I*, J. Math. Soc. Japan, 32 (1980), pp. 709–725.
- [71] ———, *On the structure of polarized manifolds with total deficiency one. II*, J. Math. Soc. Japan, 33 (1981), pp. 415–434.
- [72] ———, *On the structure of polarized manifolds with total deficiency one. III*, J. Math. Soc. Japan, 36 (1984), pp. 75–89.
- [73] ———, *On polarized manifolds whose adjoint bundles are not semipositive*, vol. 10 of Adv. Stud. Pure Math., North-Holland, Amsterdam, 1987, pp. 167–178.
- [74] ———, *On Kodaira energy and adjoint reduction of polarized manifolds*, Manuscr. Math., 76 (1992), pp. 59–84.
- [75] ———, *On Kodaira energy of polarized log varieties*, J. Math. Soc. Japan, 48 (1996), pp. 1–12.
- [76] ———, *On Kodaira energy and adjoint reduction of polarized threefolds*, Manuscr. Math., 94 (1997), pp. 211–229.
- [77] M. FURUSHIMA, *Singular del Pezzo surfaces and analytic compactifications of 3-dimensional complex affine space \mathbb{C}^3* , Nagoya Math. J., 104 (1986), pp. 1–28.
- [78] ———, *Complex analytic compactifications of \mathbb{C}^3* , Compos. Math., 76 (1990), pp. 163–196.
- [79] ———, *Mukai-Umemura’s example of the Fano threefold with genus 12 as a compactification of \mathbb{C}^3* , Nagoya Math. J., 127 (1992), pp. 145–165.
- [80] ———, *The complete classification of compactifications of \mathbb{C}^3 which are projective manifolds with the second Betti number one*, Math. Ann., 297 (1993), pp. 627–662.
- [81] ———, *A new example of a compactification of \mathbb{C}^3* , Math. Z., 212 (1993), pp. 395–399.
- [82] M. FURUSHIMA AND N. NAKAYAMA, *A new construction of a compactification of \mathbb{C}^3* , Tohoku Math. J. (2), 41 (1989), pp. 543–560.
- [83] ———, *The family of lines on the Fano threefold V_5* , Nagoya Math. J., 116 (1989), pp. 111–122.
- [84] S. GAIFULLIN, *On rigidity of trinomial hypersurfaces and factorial trinomial varieties*, ArXiv e-print, arXiv:1902.06136 (2019).
- [85] S. GAIFULLIN AND A. SHAFAREVICH, *Flexibility of normal affine horospherical varieties*, Proc. Amer. Math. Soc., 147 (2019), pp. 3317–3330.
- [86] M. GIZATULLIN, *Quasihomogeneous affine surfaces*, Math. USSR Izv., 5 (1971), pp. 1057–1081.
- [87] ———, *Two examples of affine homogeneous varieties*, Eur. J. Math. 4 (2018), pp. 1035–1064.
- [88] M. GRINENKO, *On the double cone over the Veronese surface*, Izv. Ross. Akad. Nauk Ser. Mat., 67 (2003), pp. 5–22.
- [89] ———, *Mori structures on a Fano threefold of index 2 and degree 1*, Tr. Mat. Inst. Steklova, 246 (2004), pp. 116–141.
- [90] M. GROMOV, *Oka’s principle for holomorphic sections of elliptic bundles*, J. Amer. Math. Soc., 2 (1989), pp. 851–897.
- [91] R. GURJAR, K. MASUDA, AND M. MIYANISHI, *Deformations of \mathbb{A}^1 -fibrations*, vol. 79 of Springer Proc. Math. Stat., Springer, Cham, 2014, pp. 327–361.
- [92] R. GURJAR, K. MASUDA, M. MIYANISHI, AND P. RUSSELL, *Affine lines on affine surfaces and the Makar-Limanov invariant*, Canadian J. Math., 60 (2018), pp. 109–139.
- [93] R. GURJAR AND M. MIYANISHI, *Automorphisms of affine surfaces with \mathbb{A}^1 -fibrations*, Michigan Math. J., 53 (2005), 33–55.
- [94] R. GURJAR AND D.-Q. ZHANG, *π_1 of smooth points of a log del Pezzo surface is finite. I*, J. Math. Sci. Univ. Tokyo, 1 (1994), pp. 137–180.
- [95] B. HASSETT, *Some rational cubic fourfolds*, J. Algebraic Geom., 8 (1999), pp. 103–114.
- [96] ———, *Cubic fourfolds, $K3$ surfaces, and rationality questions*, Lecture Notes in Math., 2172 (2016), pp. 29–66.

- [97] B. HASSETT, A. PIRUTKA, AND Y. TSCHINKEL, *A very general quartic double fourfold is not stably rational*, Arxiv e-print, arXiv:1605.03220 (2016).
- [98] B. HASSETT, S. TANIMOTO, AND Y. TSCHINKEL, *Balanced line bundles and equivariant compactifications of homogeneous spaces*, Int. Math. Res. Not., (2015), pp. 6375–6410.
- [99] B. HASSETT AND Y. TSCHINKEL, *On stable rationality of Fano threefolds and del Pezzo fibrations*, J. Reine Angew. Math., 751 (2018), pp. 275–287.
- [100] ———, *Rationality of Fano threefolds of degree 18 over nonclosed fields*, Arxiv e-print, arXiv:1910.13816 (2019).
- [101] F. HIDAKA AND K. WATANABE, *Normal Gorenstein surfaces with ample anti-canonical divisor*, Tokyo J. Math., 4 (1981), pp. 319–330.
- [102] F. HIRZEBRUCH, *Some problems on differentiable and complex manifolds*, Ann. Math. (2), 60 (1954), pp. 213–236.
- [103] N. T. A. HANG, M. HOFF, AND H. L. TRUONG, *On cylindrical smooth rational Fano fourfolds*, Arxiv e-print, arXiv:2101.04441 (2021).
- [104] D. HWANG AND J. KEUM, *Construction of singular rational surfaces of Picard number one with ample canonical divisor*, Proc. Amer. Math. Soc., 140 (2012), pp. 1865–1879.
- [105] A. ILIEV, L. KATZARKOV, AND V. PRZYJALKOWSKI, *Double solids, categories and non-rationality*, Proc. Edinb. Math. Soc., 57 (2014), pp. 145–173.
- [106] V. ISKOVSKIKH, *Fano 3-folds I*, Math. of the USSR, Izv., 11 (1977).
- [107] ———, *Fano 3-folds II*, Math. of the USSR, Izv., 12 (1978).
- [108] ———, *Anticanonical models of three-dimensional algebraic varieties*, J. Soviet Math., 13 (1980), pp. 745–814.
- [109] ———, *Birational automorphisms of three-dimensional algebraic varieties*, J. Soviet Math., 13 (1980), pp. 815–868.
- [110] ———, *A double projection from a line on Fano threefolds of the first kind*, Math. USSR, Sb., 66 (1990), pp. 265–284.
- [111] ———, *Factorization of birational maps of rational surfaces from the viewpoint of Mori theory*, Russian Math. Surveys, 51 (1996), pp. 585–652.
- [112] ———, *On the rationality problem for three-dimensional algebraic varieties*, Proc. of the Steklov Inst. of Math., 2018 (1997), pp. 186–227.
- [113] V. ISKOVSKIKH AND Y. MANIN, *Three-dimensional quartics and counterexamples to the Lüroth problem*, Mat. Sb., 86(128) (1971), pp. 140–166.
- [114] V. ISKOVSKIKH AND Y. PROKHOROV, *Fano varieties*, Springer, Berlin, 1999.
- [115] V. ISKOVSKIKH AND A. PUKHLIKOV, *Birational automorphisms of multidimensional algebraic varieties*, Contemp. Math. Appl., 19 (1994), pp. 3–76.
- [116] S. KALIMAN, *The Kraft-Russell generic equivalence theorem and its application*, Michigan Math. J., 69 (2020), pp. 751–764.
- [117] ———, *Surfaces with big automorphism groups*, in: Kuchment, Peter (ed.) et al., Functional analysis and geometry. Selim Grigorievich Krein centennial. Providence, RI: American Mathematical Society (AMS). Contemp. Math. 733 (2019), pp. 185–196.
- [118] ———, *Extensions of isomorphisms of subvarieties in flexible varieties*, Transform. Groups 25 (2020), pp. 517–575.
- [119] S. KALIMAN, F. KUTZSCHEBAUCH, AND M. LEUENBERGER, *Complete algebraic vector fields on affine surfaces*, Internat. J. Math. 31:2050018 (2020), 50 pp.
- [120] S. KALIMAN AND M. ZAIDENBERG, *Affine modifications and affine hypersurfaces with a very transitive automorphism group*, Transform. Groups, 4 (1999), pp. 53–95.
- [121] ———, *Miyanishi’s characterization of the affine 3-space does not hold in higher dimensions*, Ann. Inst. Fourier, 50 (2000), pp. 1649–1669.
- [122] S. KEEL AND J. MCKERNAN, *Rational curves on quasi-projective surfaces*, Mem. Amer. Math. Soc., 140 (1999), pp. viii+153.
- [123] T. KISHIMOTO, Y. PROKHOROV, AND M. ZAIDENBERG, *Group actions on affine cones*, vol. 54 of Centre de Recherches Mathématiques CRM Proc. and Lect. Notes, 2011, pp. 123–163.
- [124] ———, *G_a -actions on affine cones*, Transform. Groups, 18 (2013), pp. 1137–1153.
- [125] ———, *Affine cones over Fano threefolds and additive group actions*, Osaka J. Math., 51 (2014), pp. 1093–1113.
- [126] ———, *Unipotent group actions on del Pezzo cones*, Algebr. Geom., 1 (2014), pp. 46–56.

- [127] F. KNOP, *Mehrfach transitive Operationen algebraischer Gruppen*, Arch. Math. (Basel), 41 (1983), pp. 438–446.
- [128] S. KOBAYASHI AND T. OCHIAI, *Characterizations of complex projective spaces and hyperquadrics*, J. Math. Kyoto Univ., 13 (1973), pp. 31–47.
- [129] K. KODAIRA, *Holomorphic mappings of polydiscs into compact complex manifolds*, J. Diff. Geom., 6 (1971/72), pp. 33–46.
- [130] J. KOLLÁR, *Nonrational hypersurfaces*, J. Amer. Math. Soc., 8 (1995), pp. 241–249.
- [131] ———, *Rational curves on algebraic varieties*, vol. 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, Springer-Verlag, Berlin, 1996.
- [132] ———, *Is there a topological Bogomolov-Miyaoka-Yau inequality?*, Pure Appl. Math. Q., 4 (2008), pp. 203–236.
- [133] J. KOLLÁR, M. MIYAOKA, AND S. MORI, *Rational curves on Fano varieties*, vol. 1515 of *Lecture Notes in Math.*, Springer, 1992, pp. 100–105.
- [134] J. KOLLÁR, K. SMITH, AND A. CORTI, *Rational and nearly rational varieties*, vol. 92 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, 2004.
- [135] M. KORAS AND P. RUSSELL, *Contractible threefolds and \mathbb{C}^* -actions on \mathbb{C}^3* , J. Algebraic Geom., 6 (1997), pp. 671–695.
- [136] S. KOVACS AND Z. PATAKFALVI, *Projectivity of the moduli space of stable log-varieties and subadditivity of log-Kodaira dimension*, J. Amer. Math. Soc., 30 (2017), pp. 959–1021.
- [137] S. KOVALENKO, *Transitivity of automorphism groups of Gizatullin surfaces*, Int. Math. Res. Not., 21 (2015), pp. 11433–11484.
- [138] K. KUYUMZHIYAN, *Infinite transitivity for Calogero–Moser spaces*, Proc. Amer. Math. Soc., 148 (2020), pp. 3723–3731.
- [139] K. KUYUMZHIYAN AND F. MANGOLTE, *Infinitely transitive actions on real affine suspensions*, J. Pure Appl. Algebra 216 (2012), pp. 2106–2112.
- [140] A. KUZNETSOV AND Y. PROKHOROV, *Prime Fano threefolds of genus 12 with a \mathbf{G}_m -action*, Épijournal Géom. Algébrique, 2 (2018).
- [141] ———, *Rationality of Fano threefolds over non-closed fields*, Arxiv e-print, arXiv:1911.08949 (2019).
- [142] ———, *Rationality of Mukai varieties over non-closed fields*, Arxiv e-print, arXiv:2003.10761 (2020).
- [143] A. KUZNETSOV, Y. PROKHOROV, AND C. SHRAMOV, *Hilbert schemes of lines and conics and automorphism groups of Fano threefolds*, Japanese J. Math., 13 (2018), pp. 109–185.
- [144] A. KUZNETSOVA, *Sextic double solids with Artin–Mumford obstructions to rationality*, Arxiv e-print, arXiv:1901.05055 (2019).
- [145] Y. LEE AND N. NAKAYAMA, *Simply connected surfaces of general type in positive characteristic via deformation theory*, Proc. Lond. Math. Soc., 106 (2013), pp. 225–286.
- [146] Y. LEE AND J. PARK, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 2$* , Invent. Math., 170 (2007), pp. 483–505.
- [147] C. LI, G. TIAN, AND F. WANG, *On Yau–Tian–Donaldson conjecture for singular Fano varieties*, Comm. Pure Appl. Math. (to appear), doi:10.1002/cpa.21936.
- [148] A. LIENDO, *Affine T -varieties of complexity one and locally nilpotent derivations*, Transform. Groups, 15 (2010), pp. 389–425.
- [149] A. LIENDO AND C. PETITJEAN, *Uniformly rational varieties with torus action*, Transform. Groups, 24 (2019), pp. 149–153.
- [150] L. MARQUAND AND J. WON, *Cylinders in rational surfaces*, Eur. J. Math., 4 (2018), pp. 1161–1196.
- [151] G. MARTIN AND C. STADLMAYR, *Weak del Pezzo surfaces with global vector fields*, Arxiv e-print, arXiv:2007.03665 (2020).
- [152] M. MELLA, *Birational geometry of quartic 3-folds II: the importance of being \mathbb{Q} -factorial*, Math. Ann., 330 (2004), pp. 107–126.
- [153] M. MICHAŁEK, A. PEREPECHKO, AND H. SÜSS, *Flexible affine cones and flexible coverings*, Math. Z., 290 (2018), pp. 1457–1478.
- [154] M. MIYANISHI, *Open algebraic surfaces*, CRM Monograph Series 12. Providence, RI: American Mathematical Society (AMS), 2000.
- [155] M. MIYANISHI AND T. SUGIE, *Affine surfaces containing cylinderlike open sets*, J. Math. Kyoto Univ., 20 (1980), pp. 11–42.
- [156] M. MIYANISHI AND D.-Q. ZHANG, *Gorenstein log del Pezzo surfaces of rank one*, J. Algebra, 118 (1988), pp. 63–84.

- [157] S. MORI AND S. MUKAI, *Classification of Fano 3-folds with $B_2 \geq 2$* , Manuscr. Math., 36 (1981/82), pp. 147–162.
- [158] ———, *On Fano 3-folds with $B_2 \geq 2$* , vol. 1 of Adv. Stud. Pure Math., North-Holland, Amsterdam, 1983, pp. 101–129.
- [159] ———, *Classifications of Fano 3-folds with $B_2 \geq 2$* , I. Algebraic and topological theories. Papers from the symposium dedicated to the memory of Dr. Takehiko Miyata held in Kinosaki, October 30 – November 9, 1984, Kinokuniya, 1986, pp. 496–545.
- [160] ———, *Erratum: “Classification of Fano 3-folds with $B_2 \geq 2$ ”* [Manuscr. Math. **36** (1981/82), no. 2, 147–162], Manuscr. Math., 110 (2003), p. 407.
- [161] D. MORRISON, *The birational geometry of surfaces with rational double points*, Math. Ann., 271 (1985), pp. 415–438.
- [162] S. MUKAI, *Curves, K3 surfaces and Fano 3-folds of genus ≤ 10* , Kinokuniya, Tokyo, 1988, pp. 357–377.
- [163] ———, *Biregular classification of Fano 3-folds and Fano manifolds of coindex 3*, Proc. Nat. Acad. Sci. U.S.A., 86 (1989), pp. 3000–3002.
- [164] S. MUKAI AND H. UMEMURA, *Minimal rational threefolds*, vol. 1016 of Lecture Notes in Math., Springer, Berlin, 1983, pp. 490–518.
- [165] Y. ODAKA, C. SPOTTI, AND S. SUN, *Compact moduli spaces of del Pezzo surfaces and Kähler–Einstein metrics*, J. Differential Geom., 102 (2016), pp. 127–172.
- [166] K. OGUSO AND T. TRUONG, *Explicit examples of rational and Calabi–Yau threefolds with primitive automorphisms of positive entropy*, J. Math. Sci. Univ. Tokyo, 22 (2015), pp. 361–385.
- [167] K. OGUSO AND D.-Q. ZHANG, *On the most algebraic K3 surfaces and the most extremal log Enriques surfaces*, Amer. J. Math., 118 (1996), pp. 1277–1297.
- [168] ———, *On the complete classification of extremal log Enriques surfaces*, Math. Z., 231 (1999), pp. 23–50.
- [169] H. PARK, J. PARK, AND D. SHIN, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 3$* , Geom. Topol., 13 (2009), pp. 743–767.
- [170] ———, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 4$* , Geom. Topol., 13 (2009), pp. 1483–1494.
- [171] J. PARK, *\mathbf{G}_a -actions on the complements of hypersurfaces*, Transform. Groups (to appear), doi:10.1007/s00031-020-09589-x.
- [172] J. PARK AND J. WON, *Flexible affine cones over del Pezzo surfaces of degree 4*, Eur. J. Math., 2 (2016), pp. 304–318.
- [173] ———, *K-stability of smooth del Pezzo surfaces*, Math. Ann., 372 (2018), pp. 1239–1276.
- [174] A. PEREPECHKO, *Flexibility of affine cones over del Pezzo surfaces of degree 4 and 5*, Funct. Anal. Appl., 47 (2013), pp. 284–289.
- [175] ———, *Affine cones over cubic surfaces are flexible in codimension one*, Forum Math., 33 (2021), pp. 339–348.
- [176] P. PETERNELL, *Compactifications of \mathbb{C}^3 . II*, Math. Ann., 283 (1989), pp. 121–137.
- [177] P. PETERNELL AND M. SCHNEIDER, *Compactifications of \mathbb{C}^3 . I*, Math. Ann., 280 (1988), pp. 129–146.
- [178] C. PETITJEAN, *Equivariantly uniformly rational varieties*, Michigan Math. J., 66 (2017), pp. 245–268.
- [179] J. PIONTKOWSKI AND A. VAN DE VEN, *The automorphism group of linear sections of the Grassmannians $\mathbf{G}(1, N)$* , Doc. Math., 4 (1999), pp. 623–664.
- [180] V. L. POPOV, *On the Makar–Limanov, Derksen invariants, and finite automorphism groups of algebraic varieties*, vol. 54 of Centre de Recherches Mathématiques CRM Proc. and Lect. Notes, 2011, pp. 289–311.
- [181] ———, *On infinite dimensional algebraic transformation groups*, Transform. Groups, 19 (2014), pp. 549–568.
- [182] Y. PROKHOROV, *Automorphism groups of Fano 3-folds*, Russian Math. Surveys, 45 (1990), pp. 222–223.
- [183] ———, *Geometrical properties of Fano threefolds*, PhD thesis, Moscow State Univ., 1990. in Russian.
- [184] ———, *Fano threefolds of genus 12 and compactifications of \mathbb{C}^3* , St. Petersburg. Math. J., 3 (1991), pp. 855–864.
- [185] ———, *Rationality constructions of some Fano fourfolds of index 2*, Moscow Univ. Math. Bull., 48 (1993), pp. 32–35.
- [186] ———, *Compactifications of \mathbf{C}^4 of index 3*, Aspects Math., E25, Vieweg, 1994, pp. 159–169.
- [187] ———, *On the number of singular points of terminal factorial Fano threefolds*, Math. Notes, 101 (2017), pp. 1068–1073.
- [188] ———, *The rationality problem for conic bundles*, Russ. Math. Surveys, 73 (2018), pp. 375–456.

- [189] Y. PROKHOROV AND M. ZAIDENBERG, *Examples of cylindrical Fano fourfolds*, Eur. Jour. Math., 2 (2016), pp. 262–282.
- [190] ———, *New examples of cylindrical Fano fourfolds*, vol. 75 of Adv. Stud. Pure Math., Mathematical Society of Japan, Kinokuniya, Tokyo, 2017, pp. 443–463.
- [191] ———, *Fano-Mukai fourfolds of genus 10 as compactifications of \mathbf{C}^4* , Eur. J. Math., 4 (2018), pp. 1197–1263.
- [192] ———, *Affine cones over Fano-Mukai fourfolds of genus 10 are flexible*, Arxiv e-print, arXiv:2005.12092 (2020).
- [193] V. PRZYJALKOWSKI AND C. SHRAMOV, *Double quadrics with large automorphism groups*, Proc. Steklov Inst. Math., 294 (2016), pp. 154–175.
- [194] A. PUKHLIKOV, *Birational isomorphisms of four-dimensional quintics*, Invent. Math., 87 (1987), pp. 303–329.
- [195] ———, *Birational automorphisms of Fano hypersurfaces*, Invent. Math., 134 (1998), pp. 401–426.
- [196] ———, *Birationally rigid Fano varieties*, Torino: Università di Torino, Dipartimento di Matematica, 2004, pp. 659–681.
- [197] ———, *Birational geometry of Fano direct products*, Izv. Math., 69 (2005), pp. 1225–1255.
- [198] ———, *Birationally rigid varieties*, vol. 190 of Mathematical Surveys and Monographs, American Mathematical Society, 2013.
- [199] F. RUSSO AND G. STAGLIANO, *Explicit rationality of some cubic fourfolds*, Arxiv e-print, arXiv:1811.03502 (2018).
- [200] Y. SAKAMAKI, *Automorphism groups on normal singular cubic surfaces with no parameters*, Trans. Amer. Math. Soc., 362 (2010), pp. 2641–2666.
- [201] M. SAWAHARA, *Cylinders in weak del Pezzo fibrations*, Arxiv e-print, arXiv:1912.09016 (2019).
- [202] ———, *Cylinders in canonical del Pezzo fibrations*, Arxiv e-print, arXiv:2012.10062 (2020).
- [203] S. SCHREIEDER, *Stably irrational hypersurfaces of small slopes*, J. Amer. Math. Soc., 32 (2019), pp. 1171–1199.
- [204] A. SHAFAREVICH, *Flexibility of s -varieties of semisimple groups*, Mat. Sb., 208 (2017), pp. 121–148.
- [205] V. SHOKUROV, *Prym varieties: theory and applications*, Izv. Akad. Nauk SSSR Ser. Mat., 47 (1983), pp. 785–855.
- [206] C. SHRAMOV, *Birational rigidity and \mathbb{Q} -factoriality of a singular double quadric*, Arxiv e-print, arXiv:math/0701522 (2007).
- [207] K. TAKEUCHI, *Some birational maps of Fano 3-folds*, Compos. Math., 71 (1989), pp. 265–283.
- [208] G. TIAN, *On Kähler–Einstein metrics on certain Kähler manifolds with $c_1(m) > 0$* , Invent. Math., 89 (1987), pp. 225–246.
- [209] ———, *On Calabi’s conjecture for complex surfaces with positive first Chern class*, Invent. Math., 101 (1990), pp. 101–172.
- [210] S. TREGUB, *Three constructions of rationality of a cubic fourfold*, Vestnik Moskov. Univ. Ser. I Mat. Mekh., 3 (1984), pp. 8–14.
- [211] A. VAN DE VEN, *Analytic compactifications of complex homology cells*, Math. Ann., 147 (1962), pp. 189–204.
- [212] N. VISWANATHAN, *Lowest log canonical thresholds of a reduced plane curve of degree d* , Eur. Jour. of Math., 5 (2019), pp. 2199–6768.
- [213] C. VOISIN, *Sur la jacobienne intermédiaire du double solide d’indice deux*, Duke Math. J., 57 (1988), pp. 629–646.
- [214] F. WANG, *The classification of log Enriques surfaces of rank 18*, J. Korean Math. Soc., 48 (2011), pp. 797–822.
- [215] G. WILSON, *Collisions of Calogero–Moser particles and an adelic Grassmannian (with an Appendix by I. G. Macdonald)*, Invent. Math., 133 (1998), pp. 1–41.
- [216] C. XU, *K -stability of Fano varieties: an algebro-geometric approach*, Arxiv e-print, arXiv:2011.10477.
- [217] Q. YE, *On Gorenstein log del Pezzo surfaces*, Japanese J. of Math., 28 (2002), pp. 87–136.
- [218] Q. ZHANG, *Rational connectedness of log \mathbb{Q} -Fano varieties*, J. Reine Angew. Math., 590 (2006), pp. 131–142.
- [219] Z. ZHUANG, *Optimal destabilizing centers and equivariant K -stability*, Arxiv e-print, arXiv:2004.09413 (2020).

Ivan Cheltsov

School of Mathematics, The University of Edinburgh, Edinburgh, Scotland
Laboratory of Mirror Symmetry, NRU HSE, 6 Usacheva street, Moscow, Russia
Email address: I.Cheltsov@ed.ac.uk

Jihun Park

Center for Geometry and Physics, Institute for Basic Science
77 Cheongam-ro, Nam-gu, Pohang, Gyeongbuk, 37673, Korea
Department of Mathematics, POSTECH
77 Cheongam-ro, Nam-gu, Pohang, Gyeongbuk, 37673, Korea
Email address: wlog@postech.ac.kr

Yuri Prokhorov

Steklov Mathematical Institute, 8 Gubkina street, Moscow, Russia
Laboratory of Algebraic Geometry, NRU HSE, 6 Usacheva street, Moscow, Russia
Email address: prokhorov@mi-ras.ru

Mikhail Zaidenberg

Université Grenoble Alpes, CNRS, Institut Fourier, F-38000 Grenoble, France
Email address: Mikhail.Zaidenberg@univ-grenoble-alpes.fr