

# Mean-square convergence rates of implicit Milstein type methods for SDEs with non-Lipschitz coefficients \*

Xiaojie Wang<sup>†</sup>

School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha, China

March 21, 2023

## Abstract

A class of implicit Milstein type methods is introduced and analyzed in the present article for stochastic differential equations (SDEs) with non-globally Lipschitz drift and diffusion coefficients. By incorporating a pair of method parameters  $\theta, \eta \in [0, 1]$  into both the drift and diffusion parts, the new schemes are indeed a kind of drift-diffusion double implicit methods. Within a general framework, we offer upper mean-square error bounds for the proposed schemes, based on certain error terms only getting involved with the exact solution processes. Such error bounds help us to easily analyze mean-square convergence rates of the schemes, without relying on a priori high-order moment estimates of numerical approximations. Putting further globally polynomial growth condition, we successfully recover the expected mean-square convergence rate of order one for the considered schemes with  $\theta \in [\frac{1}{2}, 1], \eta \in [0, 1]$ . Also, some of the proposed schemes are applied to solve three SDE models evolving in the positive domain  $(0, \infty)$ . More specifically, the particular drift-diffusion implicit Milstein method ( $\theta = \eta = 1$ ) is utilized to approximate the Heston  $\frac{3}{2}$ -volatility model and the stochastic Lotka-Volterra competition model. The semi-implicit Milstein method ( $\theta = 1, \eta = 0$ ) is used to solve the Ait-Sahalia interest rate model. Thanks to the previously obtained error bounds, we reveal the optimal mean-square convergence rate of the positivity preserving schemes under more relaxed conditions, compared with existing relevant results in the literature. Numerical examples are also reported to confirm the previous findings.

**AMS subject classification:** 60H35, 60H15, 65C30.

**Key Words:** stochastic differential equations, implicit Milstein type methods, mean-square convergence rates, Heston  $\frac{3}{2}$ -volatility model, Ait-Sahalia interest rate model, stochastic Lotka-Volterra competition model, positivity preserving schemes

## 1 Introduction

Stochastic differential equations (SDEs) find applications in a wide range of scientific areas such as finance, chemistry, biology, engineering and many other branches of science. In general, analytical solutions to nonlinear SDEs are usually not available and development and analysis of numerical

---

<sup>†</sup> x.j.wang7@csu.edu.cn, x.j.wang7@gmail.com

methods for simulation of SDEs are of significant interest in practice. To analyze the numerical approximations, a global Lipschitz condition is often imposed on the coefficient functions of SDEs [30, 43]. Nevertheless, SDEs arising from applications rarely obey such a traditional but restrictive condition. Notable examples of SDEs with non-globally Lipschitz continuous coefficients include numerous models such as the  $\frac{3}{2}$ -volatility model [19, 33],

$$dX_t = X_t(\mu - \alpha X_t)dt + \beta X_t^{3/2} dW_t, \quad X_0 = x_0 > 0, \quad \mu, \alpha, \beta > 0, \quad (1.1)$$

and the Ait-Sahalia interest rate model [1],

$$dX_t = (\alpha_{-1}X_t^{-1} - \alpha_0 + \alpha_1 X_t - \alpha_2 X_t^\kappa) dt + \sigma X_t^\rho dW_t, \quad X_0 = x_0 > 0, \quad (1.2)$$

from mathematical finance, where  $\alpha_{-1}, \alpha_0, \alpha_1, \alpha_2, \sigma > 0$  are positive constants and  $\kappa > 1, \rho > 1$ . Evidently, coefficients of these two models violate the global Lipschitz condition. As already shown in [25], the popularly used Euler-Maruyama method produces divergent numerical approximations when used to solve a large class of SDEs with super-linearly growing coefficients, such as (1.1) and (1.2). Therefore special care must be taken to design and analyze convergent numerical schemes in the absence of the Lipschitz regularity of coefficients. Recent years have witnessed a prosper growth of relevant works devoted to the numerical analysis of SDEs under non-globally Lipschitz conditions, with an emphasis on analyzing implicit schemes [2–5, 10, 20–22, 39, 40, 44, 52, 55, 57], and devising explicit methods based on modifications of traditionally explicit schemes [9, 12, 13, 18, 23, 24, 26–29, 32, 37, 38, 46, 47, 49–51, 56], to just mention a few. Although explicit methods such as tamed methods [26, 47] and truncated schemes [18, 37], computationally more efficient than implicit ones for one time step, are able to well tackle non-stiff SDEs with super-linearly growing coefficients, they usually face a severe stepsize restriction due to stability issues when used to solve stiff SDE systems [43]. Moreover, explicit time stepping schemes like tamed methods, similarly to the classical explicit Euler/Milstein schemes, are usually not positivity preserving when applied to approximate financial models whose solutions naturally remain positive (see, e.g., [8, 21, 48]).

In this article we are concerned with implicit Milstein schemes for mean-square approximations of Itô SDEs with non-globally Lipschitz continuous coefficients, in the form of

$$dX_t = f(X_t) dt + g(X_t) dW_t, \quad t \in (0, T], \quad X_0 = x_0, \quad (1.3)$$

where  $W: [0, T] \times \Omega \rightarrow \mathbb{R}^m$  stands for the  $\mathbb{R}^m$ -valued standard Brownian motion,  $f: \mathbb{R}^d \rightarrow \mathbb{R}^d$  the drift coefficient function, and  $g: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$  the diffusion coefficient function. Mean-square approximations are of particular importance for the computation of statistical quantities of the solution process of (1.3) through computationally efficient multilevel Monte Carlo (MLMC) methods [15]. Recall that Milstein-type schemes achieve a higher mean-square convergence rate than the Euler-type schemes and can be combined with the MLMC approach to reduce computational costs further [14–16]. In the literature, various Milstein type methods [5–7, 18, 21, 28, 31, 32, 34, 51, 53, 56] have been studied and the present work proposes a class of implicit Milstein-type schemes and establish a mean-square convergence theory for the new schemes. On a uniform mesh constructed over  $[0, T]$  with a uniform time stepsize  $h = \frac{T}{N}, N \in \mathbb{N}$ , we develop a family of double implicit Milstein-type methods with a pair of method parameters  $(\theta, \eta)$  for (1.3) as follows:

$$\begin{aligned} Y_{n+1} = & Y_n + \theta f(Y_{n+1})h + (1 - \theta)f(Y_n)h + g(Y_n)\Delta W_n + \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(Y_n) I_{j_1, j_2}^{t_n, t_{n+1}} \\ & + \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(Y_n)h - \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(Y_{n+1})h, \quad Y_0 = X_0, \end{aligned} \quad (1.4)$$

where  $\theta, \eta \in [0, 1]$ ,  $\Delta W_n := W_{t_{n+1}} - W_{t_n}$ ,  $n \in \{0, 1, 2, \dots, N-1\}$  and  $\mathcal{L}^{j_1} g_{j_2}, I_{j_1, j_2}^{t_n, t_{n+1}}$  are precisely defined by (2.3). When  $d = m = 1$ , the schemes (1.4) coincide with the proposed ones in [21], where the authors used the positivity preserving schemes to solve the  $\frac{3}{2}$ -volatility model (1.1) and proved its strong convergence with no convergence rate revealed. After assigning  $\eta = 0$ , the proposed scheme reduces to the classical  $\theta$ -Milstein method, which has been studied in [7, 30, 57]. But in the regime of possibly super-linearly growing diffusion coefficients  $g$ , the strong convergence rate of the  $\theta$ -Milstein method is, up to the best of our knowledge, still an open problem. This paper shall fill the gap.

Also, we mention that an order reduction would be caused due to additional costs of approximating multiple stochastic integrals  $I_{j_1, j_2}^{t_n, t_{n+1}}$  when the multi-dimensional SDEs are driven by non-commutative noise. As clarified in [45, section 7], the effective order of Milstein methods should be  $\frac{2}{3}$  in the case of non-commutative noise when the multiple stochastic integrals are efficiently approximated. We refer to the simulation method proposed by Wiktorsson [54] and see also [17] for implementation issues. Compared to the order 0.5 strong Euler-type schemes, which attains the effective order 0.5, there is still a significantly improved convergence for the Milstein methods in the non-commutative noise setting. Furthermore, we mention that the application of fully implicit (stochastically implicit) methods are unavoidable for stiff systems where the stochastic part plays an essential role (see [43, pp.33] and [42] for detailed comments and an illustrative example). By incorporating a pair of method parameters  $\theta, \eta \in [0, 1]$  into the drift and diffusion parts, here we construct a kind of fully implicit methods for general multi-dimension SDE systems with non-Lipschitz coefficients. Finally, we point out that proving the expected convergence rate of the proposed schemes for SDEs in non-Lipschitz settings, especially for the above two financial models, is highly non-trivial and remains an unsolved problem. The present work aims to fill these gaps by successfully establishing a first order of mean-square convergence for the scheme (1.4) in different settings, covering the two aforementioned financial models.

By formulating certain generalized monotonicity conditions in a domain  $D \subset \mathbb{R}^d$  (Assumption 3.1), we develop an easy and novel approach to derive upper mean-square error bounds for the proposed schemes, which only get involved with the exact solution processes (Theorem 3.3). The framework is broad and covers the two aforementioned SDE financial models. Such error bounds are powerful as they help us to easily analyze mean-square convergence rates of the schemes, without relying on a priori high-order moment estimates of numerical approximations. Putting further globally polynomial growth and coercivity conditions in  $\mathbb{R}^d$  (Assumption 4.1), we utilize the derived upper error bound to successfully identify a mean-square convergence rate of order one for the schemes (1.4) solving general SDEs (1.3) (see Theorem 4.2, Corollaries 4.3, 4.4).

Later in section 5, we turn our attention to two scalar SDE models (1.1) and (1.2) arising in mathematical finance and a stochastic Lotka-Volterra (LV) competitive model (5.21) from ecology. Since the considered models evolve in the positive domain  $D = (0, \infty)$ , instead of the whole space  $\mathbb{R}$ , the convergence theory developed in section 4 cannot be applied in this situation. In order to address such issues, we apply two particular schemes covered by (1.4) to approximate these specific models, which are capable of preserving positivity of the continuous models. More precisely, the drift-diffusion double implicit Milstein method with parameters  $\theta = \eta = 1$  is utilized to approximate the Heston  $\frac{3}{2}$ -volatility model (1.1) and the stochastic LV competitive model (5.21), resulting in a recurrence of a quadratic equation with an explicit solution. And the semi-implicit Milstein method with a pair of parameters  $\theta = 1, \eta = 0$  is used to solve the Ait-Sahalia interest rate model (1.2) in both a standard and a critical regime. Both schemes are able to preserve positivity of the underlying models and their mean-square convergence rates are carefully analyzed. With the aid of the previously obtained error bounds, we prove a first order of mean-square convergence

for both schemes under mild assumptions for the first time, which fills the gap left by [21, 48]. Compared with existing relevant results for first order schemes, more relaxed conditions are put here. Specifically, the drift-diffusion double implicit Milstein scheme is shown to achieve a mean-square convergence rate of order one when used to solve the Heston  $\frac{3}{2}$ -volatility model (1.1) with model parameters obeying  $\frac{\alpha}{\beta^2} \geq \frac{5}{2}$  (Theorem 5.2). Also, the semi-implicit Milstein method is proved to retain a mean-square convergence rate of order one, when solving the Ait-Sahalia interest rate model (1.2), for full model parameters in the standard regime  $\kappa + 1 > 2\rho$  (Theorem 5.10) and for model parameters obeying  $\frac{\alpha_2}{\sigma^2} \geq 2\kappa - \frac{3}{2}$  and  $\frac{\alpha_2}{\sigma^2} > \frac{\kappa+1}{2\sqrt{2}}$  in the general critical case  $\kappa + 1 = 2\rho$  (Theorem 5.13).

Recall that a kind of Lamperti-backward Euler method was proposed and analyzed in [44] for a class of scalar SDEs defined in a domain, covering the above two financial models. There a mean-square convergence rate of order one was proved for the scheme applied to the  $\frac{3}{2}$ -volatility model with parameters satisfying  $\frac{\alpha}{\beta^2} \geq 5$  (see [44, Proposition 3.2]). Also, the scheme used to approximate the Ait-Sahalia interest rate model owns a first mean-square convergence order for full model parameters in the case  $\kappa + 1 > 2\rho$  and for parameters obeying  $\frac{\alpha_2}{\sigma^2} > 5$  in a special critical case  $\kappa = 2, \rho = 1.5$  (see Propositions 3.5, 3.6 from [44]). Unlike the Lamperti transformed scheme introduced in [44], we propose and analyze the implicit Milstein-type schemes applied to SDEs directly. From the above discussions, one can easily detect that our convergence results improve relevant ones in [44]. On the one hand, we prove the expected convergence rate for the  $\frac{3}{2}$ -volatility model on the condition  $\frac{\alpha}{\beta^2} \geq \frac{5}{2}$ , also improving the restriction  $\frac{\alpha}{\beta^2} \geq 5$  required in [44]. On the other hand, our approach is able to treat the Ait-Sahalia model in the general critical case  $\kappa + 1 = 2\rho$ , with a first mean-square convergence order identified under conditions  $\frac{\alpha_2}{\sigma^2} > 2\kappa - \frac{3}{2}$  and  $\frac{\alpha_2}{\sigma^2} > \frac{\kappa+1}{2\sqrt{2}}$ , which is, as far as we know, missing in the literature. For the special critical case  $\kappa = 2, \rho = 1.5$  studied in [44], the restriction  $\frac{\alpha_2}{\sigma^2} > 5$  there is moderately relaxed to  $\frac{\alpha_2}{\sigma^2} > \frac{5}{2}$  here.

To conclude, the main contributions of the article are summarized as follows: (i) a family of double implicit Milstein-type schemes is introduced for multi-dimension SDE systems with non-Lipschitz coefficients; (ii) a novel approach of the error analysis is developed to recover the mean-square convergence rate of order one for the schemes, which fills several gaps in the literature; (iii) the optimal mean-square convergence rate of the positivity preserving schemes applied to two financial models is obtained for the first time and more relaxed conditions are required, compared with existing relevant results for first order schemes in the literature. Therefore, this work can justify an efficient Multilevel Monte Carlo method [15] for SDEs with non-globally Lipschitz coefficients including the above financial models.

The remainder of this article is structured as follows. In the forthcoming section, a setting is formulated and a family of new Milstein-type schemes are introduced. Upper mean-square error bounds of the proposed schemes are then elaborated in section 3. Equipped with the obtained error bounds, mean-square convergence rates of the schemes are analyzed in section 4 for a general class of SDEs, under further globally polynomial growth conditions. Additionally, applications of the error bounds to two schemes for several SDE models in practice are examined in section 5, with an optimal convergence rate revealed. Further, some numerical tests are provided to confirm the theoretical findings and a brief conclusion is made at the end of the article.

## 2 SDEs and the proposed schemes

Throughout this paper, we use  $\mathbb{N}$  to denote the set of all positive integers and let  $d, m \in \mathbb{N}$ ,  $T \in (0, \infty)$  be given. Let  $\|\cdot\|$  and  $\langle \cdot, \cdot \rangle$  denote the Euclidean norm and the inner product of vectors in

$\mathbb{R}^d$ , respectively. Adopting the same notation as the vector norm, we denote  $\|A\| := \sqrt{\text{trace}(A^T A)}$  as the trace norm of a matrix  $A \in \mathbb{R}^{d \times m}$ . Given a filtered probability space  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$ , we use  $\mathbb{E}$  to mean the expectation and  $L^r(\Omega; \mathbb{R}^d)$ ,  $r \geq 1$ , to denote the family of  $\mathbb{R}^d$ -valued random variables  $\xi$  satisfying  $\mathbb{E}[\|\xi\|^r] < \infty$ . Let us consider the following SDEs of Itô type:

$$\begin{cases} dX_t = f(X_t) dt + g(X_t) dW_t, & t \in (0, T], \\ X_0 = x_0, \end{cases} \quad (2.1)$$

where  $f: \mathbb{R}^d \rightarrow \mathbb{R}^d$  is the drift coefficient function, and  $g: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$  is the diffusion coefficient function, frequently written as  $g = (g_{i,j})_{d \times m} = (g_1, g_2, \dots, g_m)$  for  $g_{i,j}: \mathbb{R}^d \rightarrow \mathbb{R}$  and  $g_j: \mathbb{R}^d \rightarrow \mathbb{R}^d$ ,  $i \in \{1, 2, \dots, d\}$ ,  $j \in \{1, 2, \dots, m\}$ . Moreover,  $W: [0, T] \times \Omega \rightarrow \mathbb{R}^m$  stands for the  $\mathbb{R}^m$ -valued standard Brownian motions with respect to  $\{\mathcal{F}_t\}_{t \in [0, T]}$  and the initial data  $X_0: \Omega \rightarrow \mathbb{R}^d$  is assumed to be  $\mathcal{F}_0$ -measurable.

In general, the system of SDEs (2.1) does not have a closed-form solution. In order to approximate (2.1), we construct a uniform mesh on  $[0, T]$  with  $h = \frac{T}{N}$  being the stepsize, for any  $N \in \mathbb{N}$ . On the uniform mesh, we propose a family of double implicit Milstein methods with a pair of method parameters  $(\theta, \eta)$ , given by

$$\begin{aligned} Y_{n+1} = & Y_n + \theta f(Y_{n+1})h + (1 - \theta)f(Y_n)h + g(Y_n)\Delta W_n + \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(Y_n) I_{j_1, j_2}^{t_n, t_{n+1}} \\ & + \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(Y_n)h - \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(Y_{n+1})h, \quad Y_0 = X_0, \end{aligned} \quad (2.2)$$

where  $\theta, \eta \in [0, 1]$ ,  $\Delta W_n := W_{t_{n+1}} - W_{t_n}$ ,  $n \in \{0, 1, 2, \dots, N-1\}$ , and

$$\mathcal{L}^{j_1} := \sum_{k=1}^d g_{k, j_1} \frac{\partial}{\partial x^k}, \quad I_{j_1, j_2}^{t_n, t_{n+1}} := \int_{t_n}^{t_{n+1}} \int_{t_n}^{s_2} dW_{s_1}^{j_1} dW_{s_2}^{j_2}, \quad j_1, j_2 \in \{1, 2, \dots, m\}. \quad (2.3)$$

In the following we use  $\frac{\partial \phi}{\partial x}$  to denote the Jacobian matrix of the vector function  $\phi: \mathbb{R}^d \rightarrow \mathbb{R}^d$  and one can observe that, for  $g_{j_2}: \mathbb{R}^d \rightarrow \mathbb{R}^d$ ,  $j_2 \in \{1, 2, \dots, m\}$ ,

$$\mathcal{L}^{j_1} g_{j_2}(x) = \sum_{k=1}^d g_{k, j_1} \frac{\partial g_{j_2}(x)}{\partial x^k} = \frac{\partial g_{j_2}(x)}{\partial x} g_{j_1}(x), \quad x \in \mathbb{R}^d. \quad (2.4)$$

By incorporating a pair of method parameters  $\theta, \eta \in [0, 1]$  into the drift and diffusion coefficients, the newly proposed schemes are implicitly defined when  $\theta + \eta \neq 0$  and their well-posedness will be discussed later. Taking  $\eta = 0$  in (2.2), the above double implicit Milstein methods (2.2) reduce to the classic  $\theta$  Milstein methods [30], which are drift implicit and given by

$$Y_{n+1} = Y_n + \theta f(Y_{n+1})h + (1 - \theta)f(Y_n)h + g(Y_n)\Delta W_n + \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(Y_n) I_{j_1, j_2}^{t_n, t_{n+1}}, \quad Y_0 = X_0. \quad (2.5)$$

In general, a straightforward introduction of implicitness into approximations of the diffusion term containing random variables suffers from unbounded numerical approximations with positive probability, see [43, Chapter 1.3.4] for clarifications. When the diffusion coefficient  $g$  fulfills the so-called commutativity condition, namely,

$$\mathcal{L}^{j_1} g_{j_2} = \mathcal{L}^{j_2} g_{j_1}, \quad j_1, j_2 \in \{1, \dots, m\}, \quad (2.6)$$



by recalling

$$I_{j_1, j_2}^{t_n, t_{n+1}} + I_{j_2, j_1}^{t_n, t_{n+1}} = \Delta W_n^{j_1} \Delta W_n^{j_2}, \quad j_1, j_2 \in \{1, \dots, m\}, j_1 \neq j_2 \quad (2.7)$$

and

$$I_{j, j}^{t_n, t_{n+1}} = \frac{1}{2}(|\Delta W_n^j|^2 - h), \quad j \in \{1, \dots, m\}, \quad (2.8)$$

one can recast the proposed double implicit Milstein method (2.2) as

$$\begin{aligned} Y_{n+1} = & Y_n + \theta f(Y_{n+1})h + (1 - \theta)f(Y_n)h + g(Y_n)\Delta W_n + \frac{1}{2} \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(Y_n) \Delta W_n^{j_1} \Delta W_n^{j_2} \\ & - \frac{(1-\eta)}{2} \sum_{j=1}^m \mathcal{L}^j g_j(Y_n)h - \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(Y_{n+1})h, \quad Y_0 = X_0. \end{aligned} \quad (2.9)$$

Here an implicit approximation is introduced with an additional method parameter  $\eta \in [0, 1]$  only in the last term that does not contain any random variable. In [7], such schemes were applied to scalar linear SDEs with several multiplicative noise terms ( $m > 1$ ) and their mean-square stability properties were studied. In particular, the commutativity condition (2.6) is fulfilled when  $m = 1$  and the newly proposed schemes (2.2) (or (2.9) equivalently) applied to the scalar SDEs ( $d = m = 1$ ) reduce to

$$\begin{aligned} Y_{n+1} = & Y_n + \theta f(Y_{n+1})h + (1 - \theta)f(Y_n)h + g(Y_n)\Delta W_n + \frac{1}{2}g'(Y_n)\Delta W_n^2 \\ & - \frac{(1-\eta)}{2}g'(Y_n)h - \frac{\eta}{2}g'(Y_{n+1})h, \quad Y_0 = X_0. \end{aligned} \quad (2.10)$$

Such schemes have been examined in [21], where the authors recovered the strong convergence rate only under globally Lipschitz conditions. Moreover, the authors used (2.10) to solve the  $\frac{3}{2}$ -volatility model (1.1) and proved its strong convergence with no convergence rate revealed. Roughly speaking, the main difficulty of recovering the convergence rate is caused by the super-linearly growing diffusion coefficients of SDEs. In the literature, a lot of researchers [3–5, 13, 18, 23, 24, 32, 34, 38–40, 47, 49, 50, 52, 56] attempt to analyze strong approximations of SDEs with super-linearly growing diffusion coefficients. However, the strong convergence rate of the classical  $\theta$ -Milstein method in the regime of possibly super-linearly growing diffusion coefficients is, up to the best of our knowledge, still an open problem. The present article aims to establish a mean-square convergence theory for the generalized  $\theta$ -Milstein schemes (1.4) within a general framework, which fills several gaps in the literature and provides improved convergence results for computational finance. Finally, it is worthwhile to emphasize that the newly proposed double implicit Milstein methods (2.2) do not require the commutativity condition (2.6) and thus work for non-commutative noise driven SDEs.

### 3 Upper mean-square error bounds for the schemes

The aim of the present section is to derive upper mean-square error bounds of the implicit Milstein type methods for SDEs taking values in a domain  $D \subset \mathbb{R}^d$ , which will help us to easily analyze the mean-square convergence rate of the schemes later. To this end, we set up a general framework by making two key assumptions as follows.

**Assumption 3.1** (Generalized monotonicity conditions in a domain). *Assume that the diffusion coefficients  $g_j: D \rightarrow \mathbb{R}^d, j \in \{1, 2, \dots, m\}$  are differentiable in a domain  $D \subset \mathbb{R}^d$  and that the*

drift coefficient  $f: D \rightarrow \mathbb{R}^d$  and the diffusion coefficient  $g = (g_1, g_2, \dots, g_m): D \rightarrow \mathbb{R}^{d \times m}$  of SDEs (2.1) satisfy certain monotonicity conditions in  $D \subset \mathbb{R}^d$ . More accurately, for method parameters  $\theta, \eta \in [0, 1]$  there exist constants  $q \in (2, \infty)$ ,  $\varrho \in (1, \infty)$ ,  $L_1, L_2 \in [0, \infty)$  and  $h_0 \in (0, T]$  such that,  $\forall x, y \in D$ ,  $h = \frac{T}{N} \in (0, h_0)$ ,

$$2\langle x - y, f(x) - f(y) \rangle + (q - 1)\|g(x) - g(y)\|^2 + \frac{\varrho}{2}h \sum_{j_1, j_2=1}^m \|\mathcal{L}^{j_1} g_{j_2}(x) - \mathcal{L}^{j_1} g_{j_2}(y)\|^2 \quad (3.1)$$

$$+ \eta h \left\langle \sum_{j=1}^m [\mathcal{L}^j g_j(x) - \mathcal{L}^j g_j(y)], f(x) - f(y) \right\rangle + (1 - 2\theta)h\|f(x) - f(y)\|^2 \leq L_1\|x - y\|^2,$$

$$\left\langle x - y, \theta[f(x) - f(y)] - \frac{\eta}{2} \sum_{j=1}^m [\mathcal{L}^j g_j(x) - \mathcal{L}^j g_j(y)] \right\rangle \leq L_2\|x - y\|^2. \quad (3.2)$$

Conditions in Assumption 3.1 are crucial to the error analysis for the proposed schemes and are called generalized monotonicity conditions in a domain  $D$ . When  $\eta = 0$ , the implicit methods (2.2) reduce to the classic  $\theta$  Milstein methods (2.5) and the above two conditions are satisfied as  $\theta \in [\frac{1}{2}, 1]$ ,  $g, \mathcal{L}^{j_1} g_{j_2}, j_1, j_2 \in \{1, 2, \dots, m\}$  satisfy the globally Lipschitz condition

$$\|g(x) - g(y)\|^2 + \sum_{j_1, j_2=1}^m \|\mathcal{L}^{j_1} g_{j_2}(x) - \mathcal{L}^{j_1} g_{j_2}(y)\|^2 \leq L\|x - y\|^2, \quad \forall x, y \in D, \quad (3.3)$$

and  $f$  obeys the monotonicity condition

$$\langle x - y, f(x) - f(y) \rangle \leq L\|x - y\|^2, \quad \forall x, y \in D. \quad (3.4)$$

Such a global monotonicity condition (3.4) is frequently used in the literature, to ensure the well-posedness of drift-implicit methods and to derive their strong convergence rates. When the diffusion  $g$  is not globally Lipschitz, which is the case for the aforementioned models (1.1), (1.2), things become much more involved. As one can see later, Assumption 3.6 and Assumption 3.8 below provide sufficient conditions that imply Assumption 3.1 and allow for non-globally Lipschitz diffusion coefficient. Since Assumption 3.1 alone does not suffice to guarantee the well-posedness of SDEs and the considered schemes in the domain  $D$ , we additionally require the following assumptions.

**Assumption 3.2** (Well-posedness of SDEs and schemes). *Assume  $\sum_{j_1, j_2=1}^m \mathbb{E}[\|\mathcal{L}^{j_1} g_{j_2}(X_0)\|^2] < \infty$  and SDE (2.1) possesses a unique  $\{\mathcal{F}_t\}_{t \in [0, T]}$ -adapted  $D$ -valued global solution with continuous sample paths,  $X: [0, T] \times \Omega \rightarrow D \subset \mathbb{R}^d$ , satisfying  $\sup_{s \in [0, T]} \mathbb{E}[\|X_s\|^2] + \sup_{s \in [0, T]} \mathbb{E}[\|f(X_s)\|^2] < \infty$ . Moreover, for  $\theta, \eta \in [0, 1]$  specified in Assumption 3.1 suppose the proposed scheme (2.2) admits a unique  $\{\mathcal{F}_{t_n}\}_{n=0}^N$ -adapted solution  $\{Y_n\}_{n=0}^N$ ,  $N \in \mathbb{N}$ , taking values in the domain  $D$ .*

We mention that Assumption 3.2 is necessary but not strict. For example, by taking  $D$  to be the whole space  $\mathbb{R}^d$ , i.e.,  $D = \mathbb{R}^d$ , Assumption 3.1 and Assumption 4.1 below together suffice to imply Assumption 3.2. In addition, some models in practice taking values in  $D = (0, \infty)$  are also given in Section 5 to satisfy the above assumptions. Under the above two assumptions, we are able to formulate the following main result of this section that offers upper mean-square error bounds for the underlying schemes.

**Theorem 3.3** (Upper mean-square error bounds). *Let Assumptions 3.1, 3.2 hold with  $\theta \in [\frac{1}{2}, 1]$  and  $2L_2h \leq \nu$  for some  $\nu \in (0, 1)$ . Let  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$  be solutions to (2.1) and (2.2), respectively. Then there exists a uniform constant  $C$  such that, for any  $n \in \{1, 2, \dots, N\}$ ,  $N \in \mathbb{N}$ ,*

$$\mathbb{E}[\|X_{t_n} - Y_n\|^2] \leq C \left( \sum_{i=1}^n \mathbb{E}[\|R_i\|^2] + \frac{1}{h} \sum_{i=1}^n \mathbb{E}[\|\mathbb{E}(R_i | \mathcal{F}_{i-1})\|^2] \right), \quad (3.5)$$

where we denote

$$\begin{aligned} R_i := & \theta \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_i}) ds + (1 - \theta) \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_{i-1}}) ds + \int_{t_{i-1}}^{t_i} g(X_s) - g(X_{t_{i-1}}) dW_s \\ & - \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}}) I_{j_1, j_2}^{t_n, t_{n+1}} + \frac{\eta}{2} h \sum_{j=1}^m [\mathcal{L}^j g_j(X_{t_i}) - \mathcal{L}^j g_j(X_{t_{i-1}})], \quad i \in \{1, 2, \dots, n\}. \end{aligned} \quad (3.6)$$

Throughout this paper, by  $C$  we denote a generic deterministic positive constant, which might vary for each appearance but is independent of the time stepsize  $h = \frac{T}{N} > 0$ ,  $N \in \mathbb{N}$ . It is interesting to observe that the term  $R_i, i \in \{1, 2, \dots, N\}$ ,  $N \in \mathbb{N}$  defined by (3.6) only gets involved with the exact solutions to SDEs. Such error bounds can be used to analyze mean-square convergence rates of the schemes without relying on a priori high-order moment estimates of numerical approximations. The proof of Theorem 3.3 is postponed, which requires the following two lemmas.

**Lemma 3.4.** *Let Assumptions 3.1, 3.2 hold and let  $R_i, i \in \{1, 2, \dots, N\}$ ,  $N \in \mathbb{N}$  be defined by (3.6). Then it holds*

$$\mathbb{E}[\|R_i\|^2] < \infty, \quad \forall i \in \{1, 2, \dots, N\}, N \in \mathbb{N}. \quad (3.7)$$

*Proof of Lemma 3.4.* In light of (3.1), one can show,  $\forall x, y \in D$ ,

$$\begin{aligned} & (q-1)\|g(x) - g(y)\|^2 + \frac{\varrho}{2} h \sum_{j_1, j_2=1}^m \|\mathcal{L}^{j_1} g_{j_2}(x) - \mathcal{L}^{j_1} g_{j_2}(y)\|^2 \\ & \leq L_1 \|x - y\|^2 - 2\langle x - y, f(x) - f(y) \rangle - (1 - 2\theta)h \|f(x) - f(y)\|^2 \\ & \quad - \eta h \left\langle \sum_{j=1}^m [\mathcal{L}^j g_j(x) - \mathcal{L}^j g_j(y)], f(x) - f(y) \right\rangle \\ & \leq (L_1 + 1) \|x - y\|^2 + \frac{\varrho h}{4} \sum_{j=1}^m \|\mathcal{L}^j g_j(x) - \mathcal{L}^j g_j(y)\|^2 + \frac{\varrho + (m + \varrho)h}{\varrho} \|f(x) - f(y)\|^2, \end{aligned} \quad (3.8)$$

and thus,  $\forall x, y \in D$ ,

$$(q-1)\|g(x) - g(y)\|^2 + \frac{\varrho}{4} h \sum_{j_1, j_2=1}^m \|\mathcal{L}^{j_1} g_{j_2}(x) - \mathcal{L}^{j_1} g_{j_2}(y)\|^2 \leq (L_1 + 1) \|x - y\|^2 + \frac{\varrho + (m + \varrho)h}{\varrho} \|f(x) - f(y)\|^2. \quad (3.9)$$

Combining this with Assumption 3.2 guarantees, for any  $i \in \{1, 2, \dots, N\}$ ,  $N \in \mathbb{N}$  and  $s \in [t_{i-1}, t_i]$ ,

$$\mathbb{E}[\|g(X_s) - g(X_{t_{i-1}})\|^2] < \infty, \quad (3.10)$$

$$\begin{aligned} h \sum_{j_1, j_2=1}^m \mathbb{E}[\|\mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}})\|^2] & \leq 2h \sum_{j_1, j_2=1}^m \mathbb{E}[\|\mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}}) - \mathcal{L}^{j_1} g_{j_2}(X_0)\|^2] \\ & \quad + 2h \sum_{j_1, j_2=1}^m \mathbb{E}[\|\mathcal{L}^{j_1} g_{j_2}(X_0)\|^2] < \infty. \end{aligned} \quad (3.11)$$



This in turn implies, for any  $i \in \{1, 2, \dots, N\}$ ,  $N \in \mathbb{N}$  and  $s \in [t_{i-1}, t_i]$ ,

$$\begin{aligned} \mathbb{E} \left[ \left\| \int_{t_{i-1}}^{t_i} g(X_s) - g(X_{t_{i-1}}) dW_s \right\|^2 \right] &= \int_{t_{i-1}}^{t_i} \mathbb{E} [\|g(X_s) - g(X_{t_{i-1}})\|^2] ds < \infty, \\ \mathbb{E} \left[ \left\| \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}}) I_{j_1, j_2}^{t_k, t_{k+1}} \right\|^2 \right] &= \frac{h^2}{2} \sum_{j_1, j_2=1}^m \mathbb{E} [\|\mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}})\|^2] < \infty, \\ \mathbb{E} \left[ \left\| \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(X_{t_i}) h - \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(X_{t_{i-1}}) h \right\|^2 \right] &< \infty. \end{aligned} \quad (3.12)$$

The desired assertion follows, by taking (3.12) and the assumption  $\sup_{s \in [0, T]} \mathbb{E} [\|f(X_s)\|^2] < \infty$  into account.  $\square$

Based on the boundedness of  $\mathbb{E} [\|R_i\|^2]$ ,  $i \in \{1, 2, \dots, N\}$ , one can arrive at the subsequent moment bounds.

**Lemma 3.5.** *Let  $\theta \in (0, 1]$ ,  $\eta \in [0, 1]$ ,  $2L_2 h \leq \nu$  for some  $\nu \in (0, 1)$  and let Assumptions 3.1, 3.2 hold. Then it holds for all  $k \in \{0, 1, 2, \dots, N\}$ ,  $N \in \mathbb{N}$  that*

$$\begin{aligned} \mathbb{E} \left[ \left\| e_k - \theta \Delta f_k^{X, Y} h + \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_{k+1}^{X, Y} h \right\|^2 \right] &< \infty, \quad \mathbb{E} [\|e_k\|^2] < \infty, \\ \mathbb{E} [\|\Delta f_k^{X, Y}\|^2] &< \infty, \quad \mathbb{E} [\|\Delta g_k^{X, Y}\|^2] < \infty, \quad \sum_{j_1, j_2=1}^m \mathbb{E} [\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X, Y}\|^2] < \infty, \end{aligned} \quad (3.13)$$

where for any  $k \in \{0, 1, \dots, N\}$ ,  $j_1, j_2 \in \{1, 2, \dots, m\}$  we denote

$$\begin{aligned} e_k &:= X_{t_k} - Y_k, \quad \Delta f_k^{X, Y} := f(X_{t_k}) - f(Y_k), \quad \Delta g_k^{X, Y} := g(X_{t_k}) - g(Y_k), \\ \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X, Y} &:= \mathcal{L}^{j_1} g_{j_2}(X_{t_k}) - \mathcal{L}^{j_1} g_{j_2}(Y_k). \end{aligned} \quad (3.14)$$

*Proof of Lemma 3.5.* We first note that, for any  $k \in \{0, 1, \dots, N-1\}$ ,  $j_1, j_2 \in \{1, 2, \dots, m\}$ ,

$$\begin{aligned} X_{t_{k+1}} &= X_{t_k} + \int_{t_k}^{t_{k+1}} f(X_s) ds + \int_{t_k}^{t_{k+1}} g(X_s) dW_s \\ &= X_{t_k} + \theta f(X_{t_{k+1}}) h + (1 - \theta) f(X_{t_k}) h + g(X_{t_k}) \Delta W_k + \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(X_{t_k}) I_{j_1, j_2}^{t_n, t_{n+1}} \\ &\quad + \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(X_{t_k}) h - \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(X_{t_{k+1}}) h + R_{k+1}, \end{aligned} \quad (3.15)$$

where  $R_{k+1}$  is defined by (3.6). Using the short-hand notation (3.14), we subtract (2.2) from (3.15) to get

$$\begin{aligned} e_{k+1} &= e_k + \theta \Delta f_{k+1}^{X, Y} h + (1 - \theta) \Delta f_k^{X, Y} h + \Delta g_k^{X, Y} \Delta W_k + \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X, Y} I_{j_1, j_2}^{t_n, t_{n+1}} \\ &\quad + \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X, Y} h - \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_{k+1}^{X, Y} h + R_{k+1}, \quad k \in \{0, 1, \dots, N-1\}. \end{aligned} \quad (3.16)$$

Denoting further

$$\mathcal{J}_k^{X,Y} := e_k - \theta \Delta f_k^{X,Y} h + \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} h, \quad k \in \{0, 1, \dots, N\}, \quad (3.17)$$

one can recast (3.16) as

$$\mathcal{J}_{k+1}^{X,Y} = \mathcal{J}_k^{X,Y} + \Delta f_k^{X,Y} h + \Delta g_k^{X,Y} \Delta W_k + \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} + R_{k+1}. \quad (3.18)$$

Squaring both sides of the above equality yields

$$\begin{aligned} \|\mathcal{J}_{k+1}^{X,Y}\|^2 &= \left\| \mathcal{J}_k^{X,Y} + \Delta f_k^{X,Y} h + \Delta g_k^{X,Y} \Delta W_k + \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} + R_{k+1} \right\|^2 \\ &= \|\mathcal{J}_k^{X,Y}\|^2 + h^2 \|\Delta f_k^{X,Y}\|^2 + \|\Delta g_k^{X,Y} \Delta W_k\|^2 + \left\| \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} \right\|^2 + \|R_{k+1}\|^2 \\ &\quad + 2h \langle \mathcal{J}_k^{X,Y}, \Delta f_k^{X,Y} \rangle + 2 \langle \mathcal{J}_k^{X,Y}, \Delta g_k^{X,Y} \Delta W_k \rangle + 2 \left\langle \mathcal{J}_k^{X,Y}, \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} \right\rangle \\ &\quad + 2 \langle \mathcal{J}_k^{X,Y}, R_{k+1} \rangle + 2h \langle \Delta f_k^{X,Y}, \Delta g_k^{X,Y} \Delta W_k \rangle + 2h \left\langle \Delta f_k^{X,Y}, \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} \right\rangle \\ &\quad + 2h \langle \Delta f_k^{X,Y}, R_{k+1} \rangle + 2 \left\langle \Delta g_k^{X,Y} \Delta W_k, \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} \right\rangle \\ &\quad + 2 \langle \Delta g_k^{X,Y} \Delta W_k, R_{k+1} \rangle + 2 \left\langle \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}}, R_{k+1} \right\rangle. \end{aligned} \quad (3.19)$$

With this at hand, we first prove  $\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] < \infty$  for all  $k \in \{0, 1, \dots, N\}$  based on an induction argument. Noting that  $X_0 = Y_0$  we thus have  $\mathbb{E}[\|\mathcal{J}_0^{X,Y}\|^2] = 0$ . We assume  $\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] < \infty$  for some  $k \in \{0, 1, \dots, N-1\}$ , which together with (3.2) implies

$$\begin{aligned} \infty &> \mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] = \mathbb{E} \left[ \left\| e_k - \theta \Delta f_k^{X,Y} h + \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} h \right\|^2 \right] \\ &= \mathbb{E}[\|e_k\|^2] + h^2 \mathbb{E} \left[ \left\| \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} - \theta \Delta f_k^{X,Y} \right\|^2 \right] \\ &\quad - 2\theta h \mathbb{E}[\langle e_k, \Delta f_k^{X,Y} \rangle] + \eta h \mathbb{E} \left[ \left\langle e_k, \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} \right\rangle \right] \\ &\geq (1 - 2hL_2) \mathbb{E}[\|e_k\|^2] + h^2 \mathbb{E} \left[ \left\| \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} - \theta \Delta f_k^{X,Y} \right\|^2 \right]. \end{aligned} \quad (3.20)$$

Therefore, for  $2hL_2 \leq \nu < 1$ ,  $\theta \in (0, 1]$  and for some  $k \in \{0, 1, 2, \dots, N-1\}$  it holds

$$\mathbb{E}[\|e_k\|^2] < \infty, \quad \mathbb{E} \left[ \left\| \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} - \theta \Delta f_k^{X,Y} \right\|^2 \right] < \infty. \quad (3.21)$$

This along with the generalized monotonicity condition (3.1) shows, for some  $k \in \{0, 1, 2, \dots, N-1\}$ ,

$$\begin{aligned}
& (q-1)\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] + \frac{\rho}{2}h \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] \\
& \leq L_1 \mathbb{E}[\|e_k\|^2] - 2\mathbb{E}[\langle \mathcal{J}_k^{X,Y}, \Delta f_k^{X,Y} \rangle] - h\mathbb{E}[\|\Delta f_k^{X,Y}\|^2] \\
& \leq L_1 \mathbb{E}[\|e_k\|^2] + \frac{1}{h}\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] < \infty.
\end{aligned} \tag{3.22}$$

In view of (3.17), (3.21), (3.22) and the assumption  $\theta > 0$ , one can easily see

$$\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] < \infty, \quad \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] < \infty, \quad \mathbb{E}[\|\Delta f_k^{X,Y}\|^2] < \infty \tag{3.23}$$

for some  $k \in \{0, 1, 2, \dots, N-1\}$ . These bounded moments suffice to ensure, for some  $k \in \{0, 1, 2, \dots, N-1\}$ ,

$$\begin{aligned}
& \mathbb{E}\left[\left\|\sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}}\right\|^2\right] = \frac{h^2}{2} \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] < \infty, \\
& \mathbb{E}[\|\Delta g_k^{X,Y} \Delta W_k\|^2] = h\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] < \infty,
\end{aligned} \tag{3.24}$$

and

$$\begin{aligned}
& \mathbb{E}\left[\left\langle \mathcal{J}_k^{X,Y}, \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} \right\rangle\right] = 0, \\
& \mathbb{E}\left[\left\langle \Delta f_k^{X,Y}, \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} \right\rangle\right] = 0, \quad \mathbb{E}[\langle \Delta f_k^{X,Y}, \Delta g_k^{X,Y} \Delta W_k \rangle] = 0 \\
& \mathbb{E}\left[\left\langle \Delta g_k^{X,Y} \Delta W_k, \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}} \right\rangle\right] = 0, \quad \mathbb{E}[\langle \mathcal{J}_k^{X,Y}, \Delta g_k^{X,Y} \Delta W_k \rangle] = 0.
\end{aligned} \tag{3.25}$$

Equipped with these estimates and taking expectations on both sides of (3.19), one can derive

$$\begin{aligned}
\mathbb{E}[\|\mathcal{J}_{k+1}^{X,Y}\|^2] &= \mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] + h^2\mathbb{E}[\|\Delta f_k^{X,Y}\|^2] + h\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] \\
&+ \frac{h^2}{2} \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] + \mathbb{E}[\|R_{k+1}\|^2] + 2h\mathbb{E}[\langle \mathcal{J}_k^{X,Y}, \Delta f_k^{X,Y} \rangle] \\
&+ 2\mathbb{E}[\langle \mathcal{J}_k^{X,Y}, R_{k+1} \rangle] + 2h\mathbb{E}[\langle \Delta f_k^{X,Y}, R_{k+1} \rangle] + 2\mathbb{E}[\langle \Delta g_k^{X,Y} \Delta W_k, R_{k+1} \rangle] \\
&+ 2\mathbb{E}\left[\left\langle \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}}, R_{k+1} \right\rangle\right].
\end{aligned} \tag{3.26}$$

Owing to the assumption that  $\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] < \infty$  for some  $k \in \{0, 1, 2, \dots, N-1\}$  and its consequence (3.23) as well as (3.7), one can use the Cauchy-Schwarz inequality to infer

$$\begin{aligned}
\mathbb{E}[\|\mathcal{J}_{k+1}^{X,Y}\|^2] &\leq 3\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] + 3h^2\mathbb{E}[\|\Delta f_k^{X,Y}\|^2] + 2h\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] \\
&+ h^2 \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] + 5\mathbb{E}[\|R_{k+1}\|^2] < \infty.
\end{aligned} \tag{3.27}$$

Based on the induction argument, the assertion  $\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] < \infty$  holds for all  $k \in \{0, 1, 2, \dots, N\}$ . Following the same lines as used in (3.20)-(3.23), the boundedness of  $\mathbb{E}[\|\mathcal{J}_n^{X,Y}\|^2]$  for all  $n \in \{0, 1, 2, \dots, N\}$  ensures that

$$\mathbb{E}[\|e_k\|^2] < \infty, \quad \mathbb{E}[\|\Delta g_k^{X,Y}\|^2] < \infty, \quad \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] < \infty, \quad \mathbb{E}[\|\Delta f_k^{X,Y}\|^2] < \infty \quad (3.28)$$

hold for all  $k \in \{0, 1, 2, \dots, N\}$ . The desired assertion are thus justified.  $\square$

Before proceeding further, we point out that the moment bounds in (3.13), depending on  $N$ , are not proved to be uniformly bounded with respect to  $N$ . This means that the moment bounds might depend on the step number  $N$ . However, such moment bounds are enough for the subsequent error analysis, which does not rely on the precise uniform moment bounds of the numerical approximations. Now we are well prepared to prove Theorem 3.3.

*Proof of Theorem 3.3.* Recalling  $\mathcal{J}_k^{X,Y} := e_k - \theta \Delta f_k^{X,Y} h + \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} h$  and using its consequence  $\Delta f_k^{X,Y} h = \frac{1}{\theta}(e_k - \mathcal{J}_k^{X,Y} + \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y} h)$  and (3.13), we derive from (3.26) that, for any  $k \in \{0, 1, 2, \dots, N-1\}$ ,

$$\begin{aligned} \mathbb{E}[\|\mathcal{J}_{k+1}^{X,Y}\|^2] &= \mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] + (1-2\theta)h^2\mathbb{E}[\|\Delta f_k^{X,Y}\|^2] + h\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] \\ &\quad + \frac{h^2}{2} \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] + \mathbb{E}[\|R_{k+1}\|^2] + 2h\mathbb{E}[\langle e_k, \Delta f_k^{X,Y} \rangle] \\ &\quad + \eta h^2 \mathbb{E}\left[\left\langle \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y}, \Delta f_k^{X,Y} \right\rangle\right] + 2\mathbb{E}[\langle \mathcal{J}_k^{X,Y}, R_{k+1} \rangle] + \frac{2}{\theta} \mathbb{E}[\langle e_k, R_{k+1} \rangle] \\ &\quad - \frac{2}{\theta} \mathbb{E}[\langle \mathcal{J}_k^{X,Y}, R_{k+1} \rangle] + \frac{\eta h}{\theta} \mathbb{E}\left[\left\langle \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y}, R_{k+1} \right\rangle\right] \\ &\quad + 2\mathbb{E}[\langle \Delta g_k^{X,Y} \Delta W_k, R_{k+1} \rangle] + 2\mathbb{E}\left[\left\langle \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}}, R_{k+1} \right\rangle\right] \\ &= \mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] + (1-2\theta)h^2\mathbb{E}[\|\Delta f_k^{X,Y}\|^2] + h\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] \\ &\quad + \frac{h^2}{2} \sum_{j_1, j_2=1}^m \mathbb{E}\left[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2\right] + \mathbb{E}[\|R_{k+1}\|^2] + 2h\mathbb{E}[\langle e_k, \Delta f_k^{X,Y} \rangle] \\ &\quad + \eta h^2 \mathbb{E}\left[\left\langle \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y}, \Delta f_k^{X,Y} \right\rangle\right] + \frac{2\theta-2}{\theta} \mathbb{E}[\langle \mathcal{J}_k^{X,Y}, \mathbb{E}(R_{k+1}|\mathcal{F}_{t_k}) \rangle] \\ &\quad + \frac{2}{\theta} \mathbb{E}[\langle e_k, \mathbb{E}(R_{k+1}|\mathcal{F}_{t_k}) \rangle] + \frac{\eta h}{\theta} \mathbb{E}\left[\left\langle \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y}, R_{k+1} \right\rangle\right] \\ &\quad + 2\mathbb{E}[\langle \Delta g_k^{X,Y} \Delta W_k, R_{k+1} \rangle] + 2\mathbb{E}\left[\left\langle \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}}, R_{k+1} \right\rangle\right]. \end{aligned} \quad (3.29)$$

Using the Cauchy-Schwarz inequality and the Young inequality gives

$$\begin{aligned} \frac{2\theta-2}{\theta} \mathbb{E}[\langle \mathcal{J}_k^{X,Y}, \mathbb{E}(R_{k+1}|\mathcal{F}_{t_k}) \rangle] &\leq h\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] + \frac{(\theta-1)^2}{\theta^2 h} \mathbb{E}[\|\mathbb{E}(R_{k+1}|\mathcal{F}_{t_k})\|^2], \\ \frac{2}{\theta} \mathbb{E}[\langle e_k, \mathbb{E}(R_{k+1}|\mathcal{F}_{t_k}) \rangle] &\leq h\mathbb{E}[\|e_k\|^2] + \frac{1}{\theta^2 h} \mathbb{E}[\|\mathbb{E}(R_{k+1}|\mathcal{F}_{t_k})\|^2], \\ \frac{\eta h}{\theta} \mathbb{E}\left[\left\langle \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y}, R_{k+1} \right\rangle\right] &\leq \frac{\eta-1}{4} h^2 \sum_{j=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^j g_j)_k^{X,Y}\|^2] + \frac{\eta^2 m}{(\eta-1)\theta^2} \mathbb{E}[\|R_{k+1}\|^2], \end{aligned}$$

$$2\mathbb{E}[\langle \Delta g_k^{X,Y} \Delta W_k, R_{k+1} \rangle] \leq (q-2)h\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] + \frac{1}{q-2}\mathbb{E}[\|R_{k+1}\|^2], \quad (3.30)$$

$$2\mathbb{E}\left[\left\langle \sum_{j_1, j_2=1}^m \Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y} I_{j_1, j_2}^{t_k, t_{k+1}}, R_{k+1} \right\rangle\right] \leq \frac{q-1}{4}h^2 \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] + \frac{2}{q-1}\mathbb{E}[\|R_{k+1}\|^2].$$

Taking these estimates into consideration and recalling (3.1) yield

$$\begin{aligned} \mathbb{E}[\|\mathcal{J}_{k+1}^{X,Y}\|^2] &\leq (1+h)\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] + (1-2\theta)h^2\mathbb{E}[\|\Delta f_k^{X,Y}\|^2] + h(q-1)\mathbb{E}[\|\Delta g_k^{X,Y}\|^2] \\ &\quad + \frac{\varrho h^2}{2} \sum_{j_1, j_2=1}^m \mathbb{E}[\|\Delta(\mathcal{L}^{j_1} g_{j_2})_k^{X,Y}\|^2] + \eta h^2 \mathbb{E}\left[\left\langle \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_k^{X,Y}, \Delta f_k^{X,Y} \right\rangle\right] \\ &\quad + \left(\frac{q-1}{q-2} + \frac{\eta^2 m}{(\varrho-1)\theta^2} + \frac{2}{\varrho-1}\right)\mathbb{E}[\|R_{k+1}\|^2] + 2h\mathbb{E}[\langle e_k, \Delta f_k^{X,Y} \rangle] \\ &\quad + h\mathbb{E}[\|e_k\|^2] + \frac{\theta^2-2\theta+2}{\theta^2 h} \mathbb{E}[\|\mathbb{E}(R_{k+1}|\mathcal{F}_{t_k})\|^2] \\ &\leq (1+h)\mathbb{E}[\|\mathcal{J}_k^{X,Y}\|^2] + (1+L_1)h\mathbb{E}[\|e_k\|^2] \\ &\quad + \left(\frac{q-1}{q-2} + \frac{\eta^2 m}{(\varrho-1)\theta^2} + \frac{2}{\varrho-1}\right)\mathbb{E}[\|R_{k+1}\|^2] + \frac{\theta^2-2\theta+2}{\theta^2 h} \mathbb{E}[\|\mathbb{E}(R_{k+1}|\mathcal{F}_{t_k})\|^2]. \end{aligned} \quad (3.31)$$

By iteration and observing  $\mathcal{J}_0^{X,Y} = e_0 - \theta \Delta f_0^{X,Y} h + \frac{\eta}{2} h \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_0^{X,Y} = 0$  we deduce

$$\begin{aligned} \mathbb{E}[\|\mathcal{J}_{k+1}^{X,Y}\|^2] &\leq (1+L_1)h \sum_{i=0}^k (1+h)^{(k-i)} \mathbb{E}[\|e_i\|^2] \\ &\quad + \left(\frac{q-1}{q-2} + \frac{\eta^2 m}{(\varrho-1)\theta^2} + \frac{2}{\varrho-1}\right) \sum_{i=0}^k (1+h)^{(k-i)} \mathbb{E}[\|R_{i+1}\|^2] \\ &\quad + \frac{\theta^2-2\theta+2}{\theta^2 h} \sum_{i=0}^k (1+h)^{(k-i)} \mathbb{E}[\|\mathbb{E}(R_{i+1}|\mathcal{F}_{t_i})\|^2] \\ &\leq (1+L_1)h e^T \sum_{i=0}^k \mathbb{E}[\|e_i\|^2] + \left(\frac{q-1}{q-2} + \frac{\eta^2 m}{(\varrho-1)\theta^2} + \frac{2}{\varrho-1}\right) e^T \sum_{i=0}^k \mathbb{E}[\|R_{i+1}\|^2] \\ &\quad + \frac{\theta^2-2\theta+2}{\theta^2 h} e^T \sum_{i=0}^k \mathbb{E}[\|\mathbb{E}(R_{i+1}|\mathcal{F}_{t_i})\|^2]. \end{aligned} \quad (3.32)$$

Additionally, the assumption (3.2) ensures

$$\begin{aligned} \mathbb{E}[\|\mathcal{J}_{k+1}^{X,Y}\|^2] &= \mathbb{E}\left[\left\|e_{k+1} - \theta \Delta f_{k+1}^{X,Y} h + \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_{k+1}^{X,Y} h\right\|^2\right] \\ &= \mathbb{E}[\|e_{k+1}\|^2] + \mathbb{E}\left[\left\|\theta h \Delta f_{k+1}^{X,Y} - \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_{k+1}^{X,Y} h\right\|^2\right] \\ &\quad - 2h\mathbb{E}\left[\left\langle e_{k+1}, \theta \Delta f_{k+1}^{X,Y} - \frac{\eta}{2} \sum_{j=1}^m \Delta(\mathcal{L}^j g_j)_{k+1}^{X,Y} h \right\rangle\right] \\ &\geq (1-2L_2 h)\mathbb{E}[\|e_{k+1}\|^2]. \end{aligned} \quad (3.33)$$

Inserting this into (3.32) yields

$$\begin{aligned}
(1 - 2L_2h)\mathbb{E}[\|e_{k+1}\|^2] \leq & (1 + L_1)he^T \sum_{i=0}^k \mathbb{E}[\|e_i\|^2] + \left(\frac{q-1}{q-2} + \frac{\eta^2 m}{(\varrho-1)\theta^2} + \frac{2}{\varrho-1}\right)e^T \sum_{i=0}^k \mathbb{E}[\|R_{i+1}\|^2] \\
& + \frac{\theta^2 - 2\theta + 2}{\theta^2 h} e^T \sum_{i=0}^k \mathbb{E}[\|\mathbb{E}(R_{i+1}|\mathcal{F}_{t_i})\|^2].
\end{aligned} \tag{3.34}$$

Owing to  $2L_2h \leq \nu < 1$  by assumption and bearing the moment bounds (3.13) in mind, one can apply Gronwall's inequality to acquire the desired assertion.  $\square$

It is worthwhile to point out that, conditions in Assumption 3.1 are not difficult to be fulfilled. For instance, the following assumption suffices to imply Assumption 3.1.

**Assumption 3.6.** *Assume that the diffusion coefficients  $g_j: \mathbb{R}^d \rightarrow \mathbb{R}^d, j \in \{1, 2, \dots, m\}$  are differentiable in a domain  $D \subset \mathbb{R}^d$ . There exist constants  $q \in (2, \infty)$ ,  $\zeta \in (0, \infty)$  and  $L_3 \in [0, \infty)$  such that,  $\forall x, y \in D$ ,  $h \in (0, 2\zeta)$ , the drift and diffusion coefficients of SDEs (2.1) obey*

$$2\langle x - y, f(x) - f(y) \rangle + (q-1)\|g(x) - g(y)\|^2 + \zeta \sum_{j_1, j_2=1}^m \|\mathcal{L}^{j_1} g_{j_2}(x) - \mathcal{L}^{j_1} g_{j_2}(y)\|^2 \leq L_3 \|x - y\|^2. \tag{3.35}$$

We mention that such a condition was also used in [5, Theorem 2.3] for the backward Milstein method ( $\theta = 1, \eta = 0$ ) and  $D = \mathbb{R}^d$ . It is not difficult to check that, when the above condition (3.35) holds, all conditions in Assumption 3.1 are satisfied with  $\theta \in [\frac{1}{2}, 1], \eta = 0$ ,  $L_1 = L_3$  and  $L_2 = \frac{\theta L_3}{2}$ . As a direct consequence of Theorem 3.3, we get the following corollary.

**Corollary 3.7.** *Let Assumptions 3.2, 3.6 be fulfilled with  $\theta L_3 h \leq \nu$  for some  $\nu \in (0, 1)$  and  $\theta \in [\frac{1}{2}, 1], \eta = 0$ . Let  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$  be solutions to SDEs (2.1) and the semi-implicit Milstein method (2.5), respectively. Then the mean-square error upper bound (3.5) holds.*

Observe that the condition (3.35) would impose a strict restriction on the polynomial growth of the diffusion coefficient, which excludes practical models such as the  $\frac{3}{2}$ -volatility model (1.1) and the Ait Sahalia model (1.2). This can be remedied by utilizing the following assumption.

**Assumption 3.8.** *Assume that the diffusion coefficients  $g_j: D \rightarrow \mathbb{R}^d, j \in \{1, 2, \dots, m\}$  are differentiable in the domain  $D \subset \mathbb{R}^d$ . For method parameters  $\theta \in (\frac{1}{2}, 1], \eta \in [0, 1]$ , there exist constants  $q \in (2, \infty)$ ,  $\varrho \in (1, \infty)$  and  $L_4, L_5, L_6 \in [0, \infty)$  such that,  $\forall x, y \in D$ , the drift and diffusion coefficients of SDEs (2.1) obey*

$$\begin{aligned}
2\langle x - y, f(x) - f(y) \rangle + (q-1)\|g(x) - g(y)\|^2 & \leq L_4 \|x - y\|^2, \\
\frac{\varrho}{2} \sum_{j_1, j_2=1}^m \|\mathcal{L}^{j_1} g_{j_2}(x) - \mathcal{L}^{j_1} g_{j_2}(y)\|^2 + \eta \left\langle \sum_{j=1}^m [\mathcal{L}^j g_j(x) - \mathcal{L}^j g_j(y)], f(x) - f(y) \right\rangle \\
& + (1 - 2\theta)\|f(x) - f(y)\|^2 \leq L_5 \|x - y\|^2, \\
\left\langle x - y, \theta[f(x) - f(y)] - \frac{\eta}{2} \sum_{j=1}^m [\mathcal{L}^j g_j(x) - \mathcal{L}^j g_j(y)] \right\rangle & \leq L_6 \|x - y\|^2.
\end{aligned} \tag{3.36}$$

One can straightforwardly verify that Assumption 3.8 implies Assumption 3.1 and one gets the following corollary, as a direct consequence of Theorem 3.3.



**Corollary 3.9.** *Let Assumptions 3.2, 3.8 hold with  $\theta \in (\frac{1}{2}, 1]$ ,  $\eta \in [0, 1]$  and  $2L_6h \leq \nu$  for some  $\nu \in (0, 1)$ . Let  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$  be solutions to SDEs (2.1) and the double implicit Milstein method (2.2), respectively. Then the mean-square error upper bound (3.5) holds.*

In section 5, we will show that the above two financial models and their numerical schemes fulfill Assumption 3.8 and one can thus rely on Corollary 3.9 to obtain the desired convergence rate. Before closing this section, we would like to mention that, the previously obtained mean-square error bound (3.5) is powerful as it helps us to easily analyze mean-square convergence rates of the schemes, without relying on a priori high-order moment estimates of numerical approximations. This will be seen in the forthcoming two sections, where we shall use the error bounds to recover the expected mean-square convergence rates of the proposed schemes in various circumstances.

## 4 Mean-square convergence rates under globally polynomial growth conditions

Equipped with the previously derived upper mean-square error bounds, the present section aims to identify the expected mean-square convergence rate of the underlying schemes (2.2) for SDEs in the whole space  $\mathbb{R}^d$  under further globally polynomial assumptions. To this end, we make the following globally polynomial growth and coercivity conditions on the drift and diffusion coefficients.

**Assumption 4.1** (Globally polynomial growth and coercivity conditions in  $\mathbb{R}^d$ ). *Assume both the drift coefficient  $f: \mathbb{R}^d \rightarrow \mathbb{R}^d$  and the diffusion coefficients  $g_j: \mathbb{R}^d \rightarrow \mathbb{R}^d, j \in \{1, 2, \dots, m\}$  of SDEs (2.1) are continuously differentiable in  $\mathbb{R}^d$ , and there exist some positive constants  $\gamma \in [1, \infty)$  and  $p^* \in [6\gamma - 4, \infty)$  such that,*

$$\langle x, f(x) \rangle + \frac{p^*-1}{2} \|g(x)\|^2 \leq C(1 + \|x\|^2), \quad \forall x \in \mathbb{R}^d, \quad (4.1)$$

$$\left\| \left( \frac{\partial f}{\partial x}(x) - \frac{\partial f}{\partial x}(\tilde{x}) \right) y \right\| \leq C(1 + \|x\| + \|\tilde{x}\|)^{\gamma-2} \|x - \tilde{x}\| \cdot \|y\|, \quad \forall x, \tilde{x}, y \in \mathbb{R}^d, \quad (4.2)$$

$$\left\| \left( \frac{\partial g_j}{\partial x}(x) - \frac{\partial g_j}{\partial x}(\tilde{x}) \right) y \right\|^2 \leq C(1 + \|x\| + \|\tilde{x}\|)^{\gamma-3} \|x - \tilde{x}\|^2 \cdot \|y\|^2, \quad \forall x, \tilde{x}, y \in \mathbb{R}^d, j \in \{1, 2, \dots, m\}. \quad (4.3)$$

Additionally we assume that the vector functions  $\eta \mathcal{L}^j g_j: \mathbb{R}^d \rightarrow \mathbb{R}^d, \eta \in [0, 1], j \in \{1, 2, \dots, m\}$  are continuously differentiable and

$$\eta \left\| \left( \frac{\partial(\mathcal{L}^j g_j)}{\partial x}(x) - \frac{\partial(\mathcal{L}^j g_j)}{\partial x}(\tilde{x}) \right) y \right\| \leq C(1 + \|x\| + \|\tilde{x}\|)^{\gamma-2} \|x - \tilde{x}\| \cdot \|y\|, \quad \forall x, \tilde{x}, y \in \mathbb{R}^d. \quad (4.4)$$

Moreover, the initial data  $X_0$  is supposed to be  $\mathcal{F}_0$ -adapted, satisfying

$$\|X_0\|_{L^{p^*}(\Omega; \mathbb{R}^d)} < \infty. \quad (4.5)$$

Recall that we use  $\frac{\partial \phi}{\partial x}$  to denote the Jacobian matrix of a vector function  $\phi: \mathbb{R}^d \rightarrow \mathbb{R}^d$ . We mention that the condition (4.1) is usually called a coercivity condition, which is a classical one in the literature to guarantee that the exact solution has finite  $p^*$ -th moments, i.e.,  $\sup_{t \in [0, T]} \|X_t\|_{L^{p^*}(\Omega; \mathbb{R}^d)} < \infty$ . The remaining conditions (4.2)-(4.4) are a kind of polynomial growth conditions, which have been also used in [5, 32] to carry out the error analysis of Milstein type methods. In subsection 4.2, we present a system of SDEs that fulfill the above conditions. Note that the condition (4.2) immediately implies

$$\left\| \frac{\partial f}{\partial x}(x) y \right\| \leq C(1 + \|x\|)^{\gamma-1} \|y\|, \quad \forall x, y \in \mathbb{R}^d, \quad (4.6)$$

which in turn implies

$$\|f(x) - f(\tilde{x})\| \leq C(1 + \|x\| + \|\tilde{x}\|)^{\gamma-1} \|x - \tilde{x}\|, \quad \forall x, \tilde{x} \in \mathbb{R}^d, \quad (4.7)$$

$$\|f(x)\| \leq C(1 + \|x\|)^\gamma, \quad \forall x \in \mathbb{R}^d. \quad (4.8)$$

Likewise, the assumption (4.3) ensures

$$\left\| \frac{\partial g_j}{\partial x}(x)y \right\|^2 \leq C(1 + \|x\|)^{\gamma-1} \|y\|^2, \quad \forall x, y \in \mathbb{R}^d, j \in \{1, 2, \dots, m\}, \quad (4.9)$$

and therefore

$$\|g_j(x) - g_j(\tilde{x})\|^2 \leq C(1 + \|x\| + \|\tilde{x}\|)^{\gamma-1} \|x - \tilde{x}\|^2, \quad \forall x, \tilde{x} \in \mathbb{R}^d, j \in \{1, 2, \dots, m\}. \quad (4.10)$$

This in turn gives

$$\|g(x)\|^2 \leq C(1 + \|x\|)^{\gamma+1}, \quad \forall x \in \mathbb{R}^d. \quad (4.11)$$

Similarly as above, the assumption (4.4) promises

$$\eta \left\| \frac{\partial \mathcal{L}^j g_j}{\partial x}(x)y \right\| \leq C(1 + \|x\|)^{\gamma-1} \|y\|, \quad \forall x, y \in \mathbb{R}^d, \quad (4.12)$$

and hence

$$\begin{aligned} \eta \left\| \mathcal{L}^j g_j(x) - \mathcal{L}^j g_j(\tilde{x}) \right\| &\leq C(1 + \|x\| + \|\tilde{x}\|)^{\gamma-1} \|x - \tilde{x}\|, \quad \forall x, \tilde{x} \in \mathbb{R}^d, \\ \eta \left\| \mathcal{L}^j g_j(x) \right\| &\leq C(1 + \|x\|)^\gamma, \quad \forall x \in \mathbb{R}^d. \end{aligned} \quad (4.13)$$

Further, Assumption 4.1 together with Assumption 3.1 in  $D = \mathbb{R}^d$  suffices to guarantee Assumption 3.2 holds in  $D = \mathbb{R}^d$ . More formally, under these assumptions, the SDE (2.1) possesses a unique adapted solution with continuous sample paths,  $X: [0, T] \times \Omega \rightarrow \mathbb{R}^d$ , satisfying

$$\sup_{t \in [0, T]} \|X_t\|_{L^p(\Omega; \mathbb{R}^d)} \leq C(1 + \|X_0\|_{L^p(\Omega; \mathbb{R}^d)}) < \infty, \quad p \in [2, p^*], \quad (4.14)$$

and thus  $\sup_{s \in [0, T]} \mathbb{E}[\|X_s\|^2] + \sup_{s \in [0, T]} \mathbb{E}[\|f(X_s)\|^2] + \sum_{j_1, j_2=1}^m \mathbb{E}[\|\mathcal{L}^{j_1} g_{j_2}(X_0)\|^2] < \infty$ , where  $p^* \in [6\gamma - 4, \infty)$  comes from Assumption 4.1. Further, the condition (3.2) in  $\mathbb{R}^d$  from Assumption 3.1 ensures that the implicit Milstein type methods (2.2) are well-defined in  $\mathbb{R}^d$ . Thanks to Assumption 4.1 as well as the above implications, one can straightforwardly show

$$\|X_{t_1} - X_{t_2}\|_{L^\delta(\Omega; \mathbb{R}^d)} \leq C \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\gamma\delta}(\Omega; \mathbb{R}^d)}^\gamma \right) |t_1 - t_2|^{\frac{1}{2}}, \quad \forall \delta \in [1, \frac{p^*}{\gamma}], t_1, t_2 \in [0, T]. \quad (4.15)$$

## 4.1 Analysis of the mean-square convergence rate

We are now ready to give the main result of this section that reveals the optimal mean-square convergence rate of the considered schemes under Assumptions 3.1, 4.1.

**Theorem 4.2** (Mean-square convergence rates of the schemes). *Let coefficients of SDEs (2.1) and method parameters of the schemes (2.2) obey Assumption 3.1 in the whole space  $D = \mathbb{R}^d$ . Let Assumption 4.1 be fulfilled and let the step-size  $h = \frac{T}{N} \in (0, \frac{1}{2\theta L_2})$  with  $\theta \in [\frac{1}{2}, 1], N \in \mathbb{N}$ . Then SDEs (2.1) and the schemes (2.2) admit unique adapted solutions in  $\mathbb{R}^d$ , denoted by  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$ , respectively. Furthermore, there exists a constant  $C > 0$ , independent of  $N \in \mathbb{N}$ , such that, for any  $N \in \mathbb{N}$ ,*

$$\sup_{0 \leq n \leq N} \|X_{t_n} - Y_n\|_{L^2(\Omega; \mathbb{R}^d)} \leq C \left( 1 + \|X_0\|_{L^{\max\{2\gamma, 3\gamma-2\}}_{\max\{4\gamma, 6\gamma-4\}}(\Omega; \mathbb{R}^d)} \right) h. \quad (4.16)$$

*Proof of Theorem 4.2.* The above discussion reminds us that all conditions in Assumptions 3.1, 3.2 hold in  $D = \mathbb{R}^d$ . Therefore, Theorem 3.3 is applicable here and we only need to properly estimate two error terms  $\mathbb{E}[\|R_i\|^2]$  and  $\mathbb{E}[\|\mathbb{E}(R_i|\mathcal{F}_{t_{i-1}})\|^2]$  before arriving at the expected mean-square convergence rate. Recalling the definition of  $\{R_i\}_{0 \leq i \leq N}$  given by (3.6) and using a triangle inequality yield

$$\begin{aligned}
\|R_i\|_{L^2(\Omega; \mathbb{R}^d)} &\leq \theta \left\| \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_i}) \, ds \right\|_{L^2(\Omega; \mathbb{R}^d)} \\
&\quad + (1 - \theta) \left\| \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_{i-1}}) \, ds \right\|_{L^2(\Omega; \mathbb{R}^d)} \\
&\quad + \left\| \int_{t_{i-1}}^{t_i} g(X_s) - g(X_{t_{i-1}}) \, dW_s - \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}}) I_{j_1, j_2}^{t_{i-1}, t_i} \right\|_{L^2(\Omega; \mathbb{R}^d)} \\
&\quad + \left\| \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(X_{t_i}) h - \frac{\eta}{2} \sum_{j=1}^m \mathcal{L}^j g_j(X_{t_{i-1}}) h \right\|_{L^2(\Omega; \mathbb{R}^d)} \\
&=: \mathbb{I}_1 + \mathbb{I}_2 + \mathbb{I}_3 + \mathbb{I}_4.
\end{aligned} \tag{4.17}$$

Next we handle the first term in (4.17) and the second term can be treated similarly. Using the Hölder inequality, (4.7), (4.14) and (4.15) shows

$$\begin{aligned}
\mathbb{I}_1 &\leq \theta \int_{t_{i-1}}^{t_i} \|f(X_s) - f(X_{t_i})\|_{L^2(\Omega; \mathbb{R}^d)} \, ds \\
&\leq C \int_{t_{i-1}}^{t_i} \left( 1 + \|X_s\|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{\gamma-1} + \|X_{t_i}\|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{\gamma-1} \right) \|X_s - X_{t_i}\|_{L^{(4\gamma-2)/\gamma}(\Omega; \mathbb{R}^d)} \, ds \\
&\leq C \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{2\gamma-1} \right) h^{\frac{3}{2}}.
\end{aligned} \tag{4.18}$$

In the same way, one can also obtain

$$\mathbb{I}_2 \leq C \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{2\gamma-1} \right) h^{\frac{3}{2}}. \tag{4.19}$$

Before coming to the estimate of  $\mathbb{I}_3$ , we note that, for any differentiable functions  $\phi: \mathbb{R}^d \rightarrow \mathbb{R}^d$ ,

$$\begin{aligned}
\phi(X_t) - \phi(X_s) &= \frac{\partial \phi}{\partial x}(X_s)(X_t - X_s) + \mathcal{R}_\phi(X_s, X_t) \\
&= \frac{\partial \phi}{\partial x}(X_s) \left( \int_s^t f(X_\xi) \, d\xi + \int_s^t g(X_\xi) \, dW_\xi \right) + \mathcal{R}_\phi(X_s, X_t), \quad s < t,
\end{aligned} \tag{4.20}$$

where for short we denote

$$\mathcal{R}_\phi(X_s, X_t) := \int_0^1 \left[ \frac{\partial \phi}{\partial x}(X_s + r(X_t - X_s)) - \frac{\partial \phi}{\partial x}(X_s) \right] (X_t - X_s) \, dr. \tag{4.21}$$

As a direct consequence of (2.4) and (4.20), one can show

$$\begin{aligned}
& \int_{t_{i-1}}^{t_i} g(X_s) - g(X_{t_{i-1}}) dW_s - \sum_{j_1, j_2=1}^m \mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}}) I_{j_1, j_2}^{t_{i-1}, t_i} \\
&= \sum_{j_2=1}^m \int_{t_{i-1}}^{t_i} \left[ g_{j_2}(X_s) - g_{j_2}(X_{t_{i-1}}) - \sum_{j_1=1}^m \mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}}) (W_s^{j_1} - W_{t_{i-1}}^{j_1}) \right] dW_s^{j_2} \\
&= \sum_{j_2=1}^m \int_{t_{i-1}}^{t_i} \left[ g_{j_2}(X_s) - g_{j_2}(X_{t_{i-1}}) - \sum_{j_1=1}^m \frac{\partial g_{j_2}}{\partial x}(X_{t_{i-1}}) g_{j_1}(X_{t_{i-1}}) (W_s^{j_1} - W_{t_{i-1}}^{j_1}) \right] dW_s^{j_2} \quad (4.22) \\
&= \sum_{j=1}^m \int_{t_{i-1}}^{t_i} \left[ \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) \left( \int_{t_{i-1}}^s f(X_\xi) d\xi + \int_{t_{i-1}}^s [g(X_\xi) - g(X_{t_{i-1}})] dW_\xi \right) \right. \\
&\quad \left. + \mathcal{R}_{g_j}(X_{t_{i-1}}, X_s) \right] dW_s^j.
\end{aligned}$$

Bearing this in mind, one can utilize the Itô isometry to obtain

$$\begin{aligned}
|\mathbb{I}_3|^2 &= \sum_{j_2=1}^m \int_{t_{i-1}}^{t_i} \mathbb{E} \left[ \left\| g_{j_2}(X_s) - g_{j_2}(X_{t_{i-1}}) - \sum_{j_1=1}^m \mathcal{L}^{j_1} g_{j_2}(X_{t_{i-1}}) (W_s^{j_1} - W_{t_{i-1}}^{j_1}) \right\|^2 \right] ds \\
&= \sum_{j=1}^m \int_{t_{i-1}}^{t_i} \mathbb{E} \left[ \left\| \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) \left( \int_{t_{i-1}}^s f(X_\xi) d\xi + \int_{t_{i-1}}^s [g(X_\xi) - g(X_{t_{i-1}})] dW_\xi \right) + \mathcal{R}_{g_j}(X_{t_{i-1}}, X_s) \right\|^2 \right] ds \\
&\leq 3 \sum_{j=1}^m \int_{t_{i-1}}^{t_i} \mathbb{E} \left[ \left\| \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) \int_{t_{i-1}}^s f(X_\xi) d\xi \right\|^2 \right] ds + 3 \sum_{j=1}^m \int_{t_{i-1}}^{t_i} \mathbb{E} [\| \mathcal{R}_{g_j}(X_{t_{i-1}}, X_s) \|^2] ds \\
&\quad + 3 \sum_{j=1}^m \int_{t_{i-1}}^{t_i} \mathbb{E} \left[ \left\| \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) \int_{t_{i-1}}^s g(X_\xi) - g(X_{t_{i-1}}) dW_\xi \right\|^2 \right] ds. \quad (4.23)
\end{aligned}$$

In the following we cope with the above three items separately. Thanks to (4.8), (4.9) and the Hölder inequality, we first get

$$\begin{aligned}
\left\| \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) \int_{t_{i-1}}^s f(X_\xi) d\xi \right\|_{L^2(\Omega; \mathbb{R}^d)} &\leq C \int_{t_{i-1}}^s \left\| (1 + \|X_{t_{i-1}}\|)^{\frac{\gamma-1}{2}} \|f(X_\xi)\| \right\|_{L^2(\Omega; \mathbb{R})} d\xi \\
&\leq C \int_{t_{i-1}}^s \left\| (1 + \|X_{t_{i-1}}\|)^{\frac{\gamma-1}{2}} (1 + \|X_\xi\|)^\gamma \right\|_{L^2(\Omega; \mathbb{R})} d\xi \quad (4.24) \\
&\leq Ch \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\frac{3\gamma-1}{2}}(\Omega; \mathbb{R}^d)} \right).
\end{aligned}$$

Again, using the Itô isometry, the Hölder inequality, (4.9), (4.10) and (4.15) yields

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) \int_{t_{i-1}}^s g(X_\xi) - g(X_{t_{i-1}}) dW_\xi \right\|^2 \right] \\
&= \sum_{l=1}^m \int_{t_{i-1}}^s \left\| \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) [g_l(X_\xi) - g_l(X_{t_{i-1}})] \right\|_{L^2(\Omega; \mathbb{R}^d)}^2 d\xi \\
&\leq C \sum_{l=1}^m \int_{t_{i-1}}^s \left\| (1 + \|X_{t_{i-1}}\|)^{\frac{\gamma-1}{2}} \|g_l(X_\xi) - g_l(X_{t_{i-1}})\| \right\|_{L^2(\Omega; \mathbb{R})}^2 d\xi \\
&\leq C \int_{t_{i-1}}^s \left\| (1 + \|X_{t_{i-1}}\|)^{\frac{\gamma-1}{2}} (1 + \|X_\xi\| + \|X_{t_{i-1}}\|)^{\frac{\gamma-1}{2}} \|X_\xi - X_{t_{i-1}}\| \right\|_{L^2(\Omega; \mathbb{R})}^2 d\xi \\
&\leq C \int_{t_{i-1}}^s \left\| (1 + \|X_\xi\| + \|X_{t_{i-1}}\|)^{\gamma-1} \|X_\xi - X_{t_{i-1}}\| \right\|_{L^2(\Omega; \mathbb{R})}^2 d\xi \\
&\leq Ch^2 \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{4\gamma-2} \right).
\end{aligned} \tag{4.25}$$

In light of (4.3), (4.14) and (4.15), one can further use the Hölder inequality to acquire

$$\begin{aligned}
& \|\mathcal{R}_{g_j}(X_{t_{i-1}}, X_s)\|_{L^2(\Omega; \mathbb{R}^d)} \\
&\leq \int_0^1 \left\| \left[ \frac{\partial g_j}{\partial x}(X_{t_{i-1}} + r(X_s - X_{t_{i-1}})) - \frac{\partial g_j}{\partial x}(X_{t_{i-1}}) \right] (X_s - X_{t_{i-1}}) \right\|_{L^2(\Omega; \mathbb{R}^d)} dr \\
&\leq C \int_0^1 \left\| (1 + \|rX_s + (1-r)X_{t_{i-1}}\| + \|X_{t_{i-1}}\|)^{\frac{\gamma-3}{2}} \|X_s - X_{t_{i-1}}\|^2 \right\|_{L^2(\Omega; \mathbb{R})} dr \\
&\leq Ch \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\max\{4\gamma, 5\gamma-3\}}(\Omega; \mathbb{R}^d)}^{\frac{\max\{4\gamma, 5\gamma-3\}}{2}} \right).
\end{aligned} \tag{4.26}$$

Plugging the above three estimates (4.24)-(4.26) into (4.23) gives

$$\mathbb{I}_3 \leq Ch^{\frac{3}{2}} \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\max\{4\gamma, 5\gamma-3\}}(\Omega; \mathbb{R}^d)}^{\frac{\max\{4\gamma, 5\gamma-3\}}{2}} \right). \tag{4.27}$$

With regard to  $\mathbb{I}_4$ , we utilize (4.13)-(4.15) and the Hölder inequality to obtain

$$\begin{aligned}
\mathbb{I}_4 &\leq \frac{\eta h}{2} \sum_{j=1}^m \left\| \mathcal{L}^j g_j(X_{t_i}) - \mathcal{L}^j g_j(X_{t_{i-1}}) \right\|_{L^2(\Omega; \mathbb{R}^d)} \\
&\leq Ch \left\| (1 + \|X_{t_i}\| + \|X_{t_{i-1}}\|)^{\gamma-1} \|X_{t_i} - X_{t_{i-1}}\| \right\|_{L^2(\Omega; \mathbb{R})} \\
&\leq Ch^{\frac{3}{2}} \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{2\gamma-1} \right).
\end{aligned} \tag{4.28}$$

Putting all the above estimates together we derive from (4.17) that

$$\|R_i\|_{L^2(\Omega; \mathbb{R}^d)} \leq Ch^{\frac{3}{2}} \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\max\{4\gamma, 5\gamma-3\}}(\Omega; \mathbb{R}^d)}^{\frac{\max\{4\gamma, 5\gamma-3\}}{2}} \right). \tag{4.29}$$

Noting that the stochastic integral vanishes under the conditional expectation, one can, similarly as in (4.17), infer that

$$\begin{aligned}
\|\mathbb{E}(R_i|\mathcal{F}_{t_{i-1}})\|_{L^2(\Omega;\mathbb{R}^d)} &\leq \theta \left\| \mathbb{E} \left( \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_i}) ds \middle| \mathcal{F}_{t_{i-1}} \right) \right\|_{L^2(\Omega;\mathbb{R}^d)} \\
&\quad + (1-\theta) \left\| \mathbb{E} \left( \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_{i-1}}) ds \middle| \mathcal{F}_{t_{i-1}} \right) \right\|_{L^2(\Omega;\mathbb{R}^d)} \\
&\quad + \frac{\eta h}{2} \sum_{j=1}^m \left\| \mathbb{E} \left( [\mathcal{L}^j g_j(X_{t_i}) - \mathcal{L}^j g_j(X_{t_{i-1}})] \middle| \mathcal{F}_{t_{i-1}} \right) \right\|_{L^2(\Omega;\mathbb{R}^d)} \\
&=: \mathbb{I}_5 + \mathbb{I}_6 + \mathbb{I}_7.
\end{aligned} \tag{4.30}$$

In order to estimate  $\mathbb{I}_5$ , we first note that

$$\begin{aligned}
\mathbb{E} \left( \int_{t_{i-1}}^{t_i} \frac{\partial f}{\partial x}(X_s) \int_s^{t_i} g(X_\xi) dW_\xi ds \middle| \mathcal{F}_{t_{i-1}} \right) &= \int_{t_{i-1}}^{t_i} \mathbb{E} \left( \int_s^{t_i} \frac{\partial f}{\partial x}(X_s) g(X_\xi) dW_\xi \middle| \mathcal{F}_{t_{i-1}} \right) ds \\
&= \int_{t_{i-1}}^{t_i} \mathbb{E} \left( \mathbb{E} \left( \int_s^{t_i} \frac{\partial f}{\partial x}(X_s) g(X_\xi) dW_\xi \middle| \mathcal{F}_s \right) \middle| \mathcal{F}_{t_{i-1}} \right) ds \\
&= 0.
\end{aligned} \tag{4.31}$$

Using this and (4.20) with  $\phi = f$  ensures

$$\mathbb{E} \left( \int_{t_{i-1}}^{t_i} f(X_{t_i}) - f(X_s) ds \middle| \mathcal{F}_{t_{i-1}} \right) = \mathbb{E} \left( \int_{t_{i-1}}^{t_i} \left[ \frac{\partial f}{\partial x}(X_s) \int_s^{t_i} f(X_\xi) d\xi + \mathcal{R}_f(X_s, X_{t_i}) \right] ds \middle| \mathcal{F}_{t_{i-1}} \right), \tag{4.32}$$

and thus

$$\begin{aligned}
\mathbb{I}_5 &= \theta \left\| \mathbb{E} \left( \int_{t_{i-1}}^{t_i} \left[ \frac{\partial f}{\partial x}(X_s) \int_s^{t_i} f(X_\xi) d\xi + \mathcal{R}_f(X_s, X_{t_i}) \right] ds \middle| \mathcal{F}_{t_{i-1}} \right) \right\|_{L^2(\Omega;\mathbb{R}^d)} \\
&\leq \theta \left\| \int_{t_{i-1}}^{t_i} \left[ \frac{\partial f}{\partial x}(X_s) \int_s^{t_i} f(X_\xi) d\xi + \mathcal{R}_f(X_s, X_{t_i}) \right] ds \right\|_{L^2(\Omega;\mathbb{R}^d)} \\
&\leq \theta \int_{t_{i-1}}^{t_i} \int_s^{t_i} \left\| \frac{\partial f}{\partial x}(X_s) f(X_\xi) \right\|_{L^2(\Omega;\mathbb{R}^d)} d\xi ds + \theta \int_{t_{i-1}}^{t_i} \left\| \mathcal{R}_f(X_s, X_{t_i}) \right\|_{L^2(\Omega;\mathbb{R}^d)} ds,
\end{aligned} \tag{4.33}$$

where the Jensen inequality was used for the second step. Here we employ (4.6), (4.8) and the Hölder inequality to show

$$\begin{aligned}
\left\| \frac{\partial f}{\partial x}(X_s) f(X_\xi) \right\|_{L^2(\Omega;\mathbb{R}^d)} &\leq C \left\| (1 + \|X_s\|)^{\gamma-1} (1 + \|X_\xi\|)^\gamma \right\|_{L^2(\Omega;\mathbb{R})} \\
&\leq C \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{4\gamma-2}(\Omega;\mathbb{R}^d)}^{2\gamma-1} \right),
\end{aligned} \tag{4.34}$$



and employ (4.2), (4.15) and the Hölder inequality to arrive at

$$\begin{aligned}
\|\mathcal{R}_f(X_s, X_{t_i})\|_{L^2(\Omega; \mathbb{R}^d)} &\leq \int_0^1 \left\| \left[ \frac{\partial f}{\partial x}(X_s + r(X_{t_i} - X_s)) - \frac{\partial f}{\partial x}(X_s) \right] (X_{t_i} - X_s) \right\|_{L^2(\Omega; \mathbb{R}^d)} dr \\
&\leq C \int_0^1 \left\| (1 + \|rX_{t_i} + (1-r)X_s\| + \|X_s\|)^{\gamma-2} \|X_{t_i} - X_s\|^2 \right\|_{L^2(\Omega; \mathbb{R})} dr \\
&\leq Ch \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\max\{2\gamma, 3\gamma-2\}}(\Omega; \mathbb{R}^d)}^{\max\{2\gamma, 3\gamma-2\}} \right).
\end{aligned} \tag{4.35}$$

Inserting (4.34) and (4.35) into (4.33) implies

$$\mathbb{I}_5 \leq Ch^2 \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\max\{4\gamma, 6\gamma-4\}}(\Omega; \mathbb{R}^d)}^{\max\{2\gamma, 3\gamma-2\}} \right). \tag{4.36}$$

The estimates of  $\mathbb{I}_6$  and  $\mathbb{I}_7$  are similar and one can also get

$$\mathbb{I}_6 + \mathbb{I}_7 \leq Ch^2 \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\max\{4\gamma, 6\gamma-4\}}(\Omega; \mathbb{R}^d)}^{\max\{2\gamma, 3\gamma-2\}} \right). \tag{4.37}$$

Therefore, from (4.30) it immediately follows that

$$\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R}^d)} \leq Ch^2 \left( 1 + \sup_{t \in [0, T]} \|X_t\|_{L^{\max\{4\gamma, 6\gamma-4\}}(\Omega; \mathbb{R}^d)}^{\max\{2\gamma, 3\gamma-2\}} \right). \tag{4.38}$$

In view of Theorem 3.3 and (4.14), we validate the desired assertion (4.16).  $\square$

As already mentioned at the end of Section 3, Theorem 3.3 still holds when Assumption 3.1 is replaced by Assumption 3.6 or Assumption 3.8. Therefore, the following two corollaries follow directly from Corollaries 3.7, 3.9.

**Corollary 4.3.** *Let Assumption 3.6 be fulfilled with  $D = \mathbb{R}^d$  and let  $\theta L_3 h \leq \nu$  for some  $\theta \in [\frac{1}{2}, 1]$ ,  $\nu \in (0, 1)$ . Let conditions in Assumption 4.1 be all satisfied. Then SDEs (2.1) and the semi-implicit Milstein methods (2.5) admit unique adapted solutions in  $\mathbb{R}^d$  and (4.16) holds, namely, the schemes (2.5) retain a mean-square convergence rate of order one.*

**Corollary 4.4.** *Let Assumption 3.8 be fulfilled with  $D = \mathbb{R}^d$  and let conditions in Assumption 4.1 be all satisfied. Let  $\theta L_6 h \leq \nu$  for some  $\theta \in (\frac{1}{2}, 1]$ ,  $\eta \in [0, 1]$  and for some  $\nu \in (0, 1)$ . Then SDEs (2.1) and the proposed schemes (2.2) admit unique adapted solutions in  $\mathbb{R}^d$  and (4.16) holds, namely, the schemes (2.2) retain a mean-square convergence rate of order one.*

## 4.2 An example with numerical simulations

In this subsection, we aim to give an example SDE that satisfies Assumptions 3.1, 4.1. To this end, let us first consider the following semi-linear stochastic partial differential equation (SPDE) [35, 36]:

$$\begin{cases} du(t, x) = [\frac{\partial^2}{\partial x^2} u(t, x) + u(t, x) - u^3(t, x)] dt + g(u(t, x)) dW_t, & t \in (0, T], x \in (0, 1), \\ u(t, 0) = u(t, 1) = 0, \\ u(0, x) = u_0(x), \end{cases} \tag{4.39}$$

where  $g: \mathbb{R} \rightarrow \mathbb{R}$  and  $W: [0, T] \times \Omega \rightarrow \mathbb{R}$  is the real-valued standard Brownian motions. Such an SPDE is usually termed as the stochastic Allen-Cahn equation. Next we want to spatially discretize the above SPDE to obtain an SDE system. On the interval  $[0, 1]$  we construct a uniform mesh with stepsize  $\Delta x := \frac{1}{K}$  and denote  $x_i = i\Delta x$ ,  $i = 1, 2, \dots, K-1$ . Discretizing the SPDE (4.39) spatially by a finite difference method yields a system of SDEs:

$$dX_t = [\mathbb{A}X_t + \mathbb{F}(X_t)] dt + \mathbb{G}(X_t) dW_t, \quad t \in (0, T], \quad X_0 = x_0, \quad (4.40)$$

where  $X_t = (X_{1,t}, X_{2,t}, \dots, X_{K-1,t})^T := (u(t, x_1), u(t, x_2), \dots, u(t, x_{K-1}))^T$ ,  $\mathbb{A} \in R^{(K-1) \times (K-1)}$ ,  $x_0 = (u_0(x_1), u_0(x_2), \dots, u_0(x_{K-1}))^T$  and

$$\mathbb{A} = K^2 \begin{bmatrix} -2 & 1 & 0 & \cdots & 0 & 0 \\ 1 & -2 & 1 & \cdots & 0 & 0 \\ 0 & 1 & -2 & \cdots & 0 & 0 \\ & \cdots & & \cdots & & \\ 0 & 0 & 0 & \cdots & -2 & 1 \\ 0 & 0 & 0 & \cdots & 1 & -2 \end{bmatrix}, \mathbb{F}(X) = \begin{bmatrix} X_1 - (X_1)^3 \\ X_2 - (X_2)^3 \\ \vdots \\ X_{K-1} - (X_{K-1})^3 \end{bmatrix}, \mathbb{G}(X) = \begin{bmatrix} g(X_1) \\ g(X_2) \\ \vdots \\ g(X_{K-1}) \end{bmatrix}.$$

We do not consider the error caused by the spatial discretization but focus on the temporal discretization of the SDE system (4.40), done by the semi-implicit Milstein method ( $\theta = 1, \eta = 0$ ). Moreover, we assume  $g \in C_b^3(\mathbb{R}, \mathbb{R})$ , i.e.,  $g$  is three times differentiable with derivatives bounded. It is easy to check all conditions in Assumption 4.1 are fulfilled in  $D = \mathbb{R}^{K-1}$  with  $\gamma = 3$  and for any  $p^* \geq 14$ . By setting  $\theta = 1, \eta = 0$ , conditions (3.1), (3.2) in Assumption 3.1 are also both satisfied in  $D = \mathbb{R}^{K-1}$ . Therefore, Theorem 4.2 is applicable, with the first convergence rate obtained for the semi-implicit Milstein method. Since the SDE system has commutative noise, the Milstein type methods do not involve the Levy area [30, 43] and can be implemented as easily as the Euler type methods.

In what follows we set  $g(u) = \sin(u) + 1$  and  $u_0(x) \equiv 1$  and do some numerical experiments. In Figure 1, we plot mean-square errors of the semi-implicit Milstein method ( $\theta = 1, \eta = 0$ ) for the SDE system (4.40) with  $K = 4$ . There one can observe a convergence rate of order one, as the step-sizes shrink. Here and below numerical approximations are performed using six different stepsizes  $h = 2^{-i}, i = 2, 3, \dots, 7$ . The “exact” solution is identified as the numerical one using a fine stepsize  $h_{\text{exact}} = 2^{-12}$  and the expectations are approximated by computing averages over  $10^4$  samples. For comparison, we also discretize (4.40) by the tamed Milstein method for non-Lipschitz SDEs [32, 51]. Tables 1-3 provide mean-square approximation errors of these two methods for three cases  $K = 4, 8, 16$ . Clearly, the tamed Milstein method gives satisfactory results in the low dimension case  $K = 4$  when the time stepsize is small, i.e.,  $h = 2^{-5}, 2^{-6}, 2^{-7}$ . As the dimension  $K$  increases ( $K = 8, 16$ ), the tamed Milstein method gives large errors and the approximations become unreliable for even small stepsizes. However, the semi-implicit Milstein method performs much better, even in high dimension case  $K = 16$ . This happens because the eigenvalues  $\{\lambda_i\}_{i=1}^{K-1}$  of  $A$  are  $\lambda_i = -4K^2 \sin^2(\frac{i\pi}{2K}) < 0$  and the problem (4.40) turns to be a very stiff system [43] as  $K$  increases. As a kind of explicit method, the tamed Milstein method applied to solve stiff system, faces severe time step-size reduction due to the stability issue. On the contrary, the semi-implicit Milstein method has excellent stability property and is well suited for such stiff system.




Figure 1: Mean-square convergence rate of the semi-implicit Milstein method for (4.40) ( $K = 4$ ).

Table 1: Mean-square approximation errors for two schemes ( $K = 4$ ).

Stepsizes $h$	Semi-implicit Milstein	Tamed Milstein
$h = 2^{-2}$	0.228228472003678	1.334521881473836
$h = 2^{-3}$	0.142671496841737	0.669681337348534
$h = 2^{-4}$	0.092138829109993	0.304845858687944
$h = 2^{-5}$	0.050402455908956	0.104293855220492
$h = 2^{-6}$	0.026477850950294	0.044846913728710
$h = 2^{-7}$	0.014040231850694	0.023917375308279

Table 2: Mean-square approximation errors for two schemes ( $K = 8$ ).

Stepsizes $h$	Semi-implicit Milstein	Tamed Milstein
$h = 2^{-2}$	0.337954132405219	3.127906338055271
$h = 2^{-3}$	0.215927776446030	2.264234907688349
$h = 2^{-4}$	0.143858604122065	1.393606951102787
$h = 2^{-5}$	0.082829804151649	0.792573908322782
$h = 2^{-6}$	0.045812280417151	0.375592176659368
$h = 2^{-7}$	0.025766349691283	0.060932654697185

Table 3: Mean-square approximation errors for two schemes ( $K = 16$ ).

Stepsizes $h$	Semi-implicit Milstein	Tamed Milstein
$h = 2^{-2}$	0.483493085665317	5.551900376818316
$h = 2^{-3}$	0.310800712759207	4.923675828972162
$h = 2^{-4}$	0.209040389203629	4.088007760382119
$h = 2^{-5}$	0.122832739545349	3.113087309657318
$h = 2^{-6}$	0.070290414827683	2.028019061710102
$h = 2^{-7}$	0.041888961361398	1.128188804503420

## 5 Convergence rates of positivity preserving schemes for SDEs with non-globally Lipschitz coefficients

In the present section, we turn our attention to the aforementioned scalar SDE models (1.1) and (1.2) arising from mathematical finance. Unlike general SDEs studied in the previous section, the considered financial models do not evolve in the whole space  $\mathbb{R}$ , but only in the positive domain  $D = (0, \infty)$ . This thus makes the convergence theory developed in the previous section not applicable in this situation. Moreover, preservation of positivity is usually a desirable modeling property and positivity of the approximation is, in many cases, necessary in order for the numerical scheme to be well defined (see, e.g., (5.2) and (5.30) below). However, numerical schemes are, in general, not able to preserve positivity. For example, the classical Euler-Maruyama method fails to preserve positivity for any scalar SDE [8]. In this section we choose two particular schemes from (2.10) to approximate these two models, which are capable of preserving positivity of the continuous models. By means of the previously obtained error bound, we carefully analyze the expected mean-square convergence rate of the resulting numerical approximations.

### 5.1 The double implicit Milstein scheme for the Heston $\frac{3}{2}$ -volatility model

As the first considered financial model, let us look at the Heston  $\frac{3}{2}$ -volatility model [19, 33]:

$$dX_t = X_t(\mu - \alpha X_t)dt + \beta X_t^{3/2} dW(t), \quad X_0 = x_0 > 0, \quad \mu, \alpha, \beta > 0, \quad t > 0, \quad (5.1)$$

which can be viewed as an inverse of a Cox-Ingersoll-Ross (CIR) process [44]. Such an equation is also used for modelling term structure dynamics [11]. Recently, some researchers [9, 21, 44] proposed and analyzed different positivity-preserving numerical schemes for strong approximations of the  $\frac{3}{2}$ -process. Similarly to [21], we choose a particular double implicit Milstein scheme (2.10) with  $\theta = \eta = 1$  to approximate the above  $\frac{3}{2}$ -process. Furthermore, we attempt to prove the expected convergence rate for the scheme, which is missing in [21].

Given  $T \in (0, \infty)$  and  $N \in \mathbb{N}$ , one can construct a uniform mesh on the interval  $[0, T]$  with the uniform stepsize  $h = \frac{T}{N}$ . Based on the uniform mesh, we apply the drift-diffusion double implicit Milstein scheme (2.10) with  $\theta = \eta = 1$  to the model (5.1), resulting in, for  $n \in \{0, 1, 2, \dots, N-1\}$ ,

$$Y_{n+1} = Y_n + Y_{n+1}(\mu - \alpha Y_{n+1})h + \beta Y_n^{3/2} \Delta W_n + \frac{3\beta^2}{4} Y_n^2 |\Delta W_n|^2 - \frac{3\beta^2}{4} Y_{n+1}^2 h, \quad Y_0 = X_0, \quad (5.2)$$

which is a quadratic equation and has a unique positive solution explicitly given by  $Y_0 = X_0$  and

$$Y_{n+1} = \left( \sqrt{(1 - \mu h)^2 + 4h(\alpha + \frac{3}{4}\beta^2)(Y_n + \beta|Y_n|^{\frac{3}{2}}\Delta W_n + \frac{3}{4}\beta^2|Y_n|^2|\Delta W_n|^2)} - (1 - \mu h) \right) / (2\alpha h + \frac{3}{2}\beta^2 h) > 0, \quad (5.3)$$

given that  $Y_n > 0$ ,  $n \in \{0, 1, 2, \dots, N-1\}$ . We mention that no additional restriction is put on the stepsize  $h > 0$  to ensure the positivity of the above approximations. In order to carry out the error analysis for the scheme using Theorem 3.3, we should first justify all conditions required in Assumptions 3.1, 3.2, which are clarified in the forthcoming lemma.

**Lemma 5.1.** *Let  $\mu, \alpha, \beta > 0$ ,  $X_0 > 0$ . Then the Heston  $-\frac{3}{2}$  volatility model (5.1) has a unique global solution in  $(0, \infty)$  and the scheme (5.2) produces unique positivity preserving approximations given by (5.3). When  $\alpha > \frac{3}{2}\beta^2$ , the SDE model (5.1) and the scheme (5.2) obey Assumptions 3.1, 3.2 in the domain  $D = (0, \infty)$  for some  $2 < q < 1 + \frac{8\alpha}{9\beta^2}$ .*

*Proof of Lemma 5.1.* The well-posedness of the considered model (5.1) and the scheme (5.2) in the positive domain  $(0, \infty)$  can be found in [21, 44]. It remains to validate the other conditions in Assumptions 3.1, 3.2. For brevity, we denote the drift and diffusion coefficients of SDE (5.1) by

$$f(x) := x(\mu - \alpha x), \quad g(x) := \beta x^{\frac{3}{2}}, \quad x \in \mathbb{R}_+. \quad (5.4)$$

As a result,  $g'(x)g(x) = \frac{3}{2}\beta^2 x^2$ ,  $x \in \mathbb{R}_+$  and one can find a positive constant  $\tilde{c} > 0$  such that

$$\begin{aligned} \Xi(x, y, h) &:= \varrho^{\frac{h}{2}} \|g'(x) - g'(y)\|^2 + \eta h \langle g'(x) - g'(y), f(x) - f(y) \rangle - h \|f(x) - f(y)\|^2 \\ &= \frac{9}{8} \varrho \beta^4 h (x^2 - y^2)^2 + \frac{3}{2} \beta^2 \eta \mu h (x^2 - y^2)(x - y) - \frac{3}{2} \beta^2 \eta \alpha h (x^2 - y^2)^2 \\ &\quad - h [\mu^2 (x - y)^2 - 2\mu\alpha(x - y)(x^2 - y^2) + \alpha^2 (x^2 - y^2)^2] \\ &= \left\{ \left[ \frac{9}{8} \varrho \beta^4 - \frac{3}{2} \beta^2 \alpha - \alpha^2 \right] (x + y)^2 + \left[ \frac{3}{2} \beta^2 \mu + 2\mu\alpha \right] (x + y) - \mu^2 \right\} (x - y)^2 h \\ &\leq \tilde{c} (x - y)^2 h, \quad \forall x, y \in \mathbb{R}_+, \end{aligned} \quad (5.5)$$

where we used the facts that  $\eta = 1$  and that  $\frac{9}{8} \varrho \beta^4 - \frac{3}{2} \beta^2 \alpha - \alpha^2 < 0$  for some  $\varrho > 1$  since  $\alpha > \frac{3}{2} \beta^2$  by assumption. Further, we take some  $2 < q < 1 + \frac{8\alpha}{9\beta^2}$  to promise  $\frac{9}{4}(q-1)\beta^2 - 2\alpha \leq 0$  and hence

$$\begin{aligned} &2 \langle x - y, f(x) - f(y) \rangle + (q-1) \|g(x) - g(y)\|^2 + \Xi(x, y, h) \\ &= 2\mu|x - y|^2 - 2\alpha(x^2 - y^2)(x - y) + (q-1)\beta^2(x^{\frac{3}{2}} - y^{\frac{3}{2}})^2 + \Xi(x, y, h) \\ &\leq 2\mu|x - y|^2 + \left[ \frac{9}{4}(q-1)\beta^2 - 2\alpha \right] (x + y)(x - y)^2 + \tilde{c}(x - y)^2 h \\ &\leq (2\mu + \tilde{c}T)|x - y|^2, \quad \forall x, y \in \mathbb{R}_+, \end{aligned} \quad (5.6)$$

which means the condition (3.1) in Assumption 3.1 is fulfilled. Now we validate (3.2) as follows:

$$\begin{aligned} &\langle x - y, f(x) - f(y) - \frac{1}{2}[g'(x)g(x) - g'(y)g(y)] \rangle \\ &= \mu|x - y|^2 - \alpha(x^2 - y^2)(x - y) - \frac{3}{4}\beta^2(x^2 - y^2)(x - y) \\ &\leq \mu|x - y|^2, \quad \forall x, y \in \mathbb{R}_+. \end{aligned} \quad (5.7)$$

Next we note that for any  $p^* \leq 4$

$$\langle x, f(x) \rangle + \frac{p^*-1}{2} \|g(x)\|^2 = \mu x^2 - (\alpha - \frac{p^*-1}{2}\beta^2)x^3 \leq \mu x^2, \quad \forall x \in \mathbb{R}_+, \quad (5.8)$$

where the assumption  $\alpha > \frac{3}{2}\beta^2$  was again used. This assures  $\sup_{t \in [0, T]} \|X_t\|_{L^4(\Omega; \mathbb{R}^d)} < \infty$  and thus  $\sup_{s \in [0, T]} \mathbb{E}[\|X_s\|^2] + \sup_{s \in [0, T]} \mathbb{E}[\|f(X_s)\|^2] < \infty$ , as required in Assumption 3.2.  $\square$

Now we are able to apply Theorem 3.3 to deduce the convergence rate of the numerical scheme.

**Theorem 5.2.** Let  $X_0 > 0$  and let  $\mu, \alpha, \beta > 0$  satisfy  $\alpha \geq \frac{5}{2}\beta^2$ . Let  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$  be uniquely given by (5.1) and (5.3), respectively. Let  $h \in (0, \frac{1}{2\mu})$ . Then there exists a constant  $C > 0$ , independent of  $N \in \mathbb{N}$ , such that

$$\sup_{0 \leq n \leq N} \|Y_n - X_{t_n}\|_{L^2(\Omega; \mathbb{R})} \leq Ch. \quad (5.9)$$

*Proof of Theorem 5.2.* As already clarified in the proof of Lemma 5.1, the considered model and the scheme obey Assumptions 3.1, 3.2 in the domain  $D = (0, \infty)$ . Therefore, Theorem 3.3 is applicable here and it remains to estimate two error terms  $\mathbb{E}[\|R_i\|^2]$  and  $\mathbb{E}[\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|^2]$ ,  $i \in \{1, 2, \dots, N\}$ . before attaining the convergence rate. First of all, we recall  $f(x) := x(\mu - \alpha x)$ ,  $g(x) := \beta x^{\frac{3}{2}}$ ,  $x \in \mathbb{R}_+$ . Following the notation used in Theorem 3.3, one can easily see

$$\begin{aligned} \|R_i\|_{L^2(\Omega; \mathbb{R})} &\leq \left\| \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_i}) \, ds \right\|_{L^2(\Omega; \mathbb{R})} + \frac{h}{2} \|g'(X_{t_i}) - g'(X_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R})} \\ &\quad + \left\| \int_{t_{i-1}}^{t_i} [g(X_s) - g(X_{t_{i-1}}) - g'(X_{t_{i-1}})(W_s - W_{t_{i-1}})] \, dW_s \right\|_{L^2(\Omega; \mathbb{R})} \\ &=: I_1 + I_2 + I_3. \end{aligned} \quad (5.10)$$

Applying the Itô formula to the quadratic polynomial  $f(x) = x(\mu - \alpha x)$ ,  $x \in \mathbb{R}_+$  yields

$$f(X_{t_i}) - f(X_s) = \int_s^{t_i} [f'(X_r)f(X_r) + \frac{1}{2}f''(X_r)g^2(X_r)] \, dr + \int_s^{t_i} f'(X_r)g(X_r) \, dW_r \quad (5.11)$$

and thus

$$\begin{aligned} I_1 &\leq \int_{t_{i-1}}^{t_i} \int_s^{t_i} \|f'(X_r)f(X_r) + \frac{1}{2}f''(X_r)g^2(X_r)\|_{L^2(\Omega; \mathbb{R})} \, dr \, ds \\ &\quad + \int_{t_{i-1}}^{t_i} \left( \int_s^{t_i} \mathbb{E}[\|f'(X_r)g(X_r)\|^2] \, dr \right)^{\frac{1}{2}} \, ds \\ &\leq Ch^{\frac{3}{2}} \left( 1 + \sup_{s \in [0, T]} \|X_s\|_{L^6(\Omega; \mathbb{R})}^3 \right), \end{aligned} \quad (5.12)$$

where one used the Itô isometry and computed that  $f'(x)f(x) = x(\mu - \alpha x)(\mu - 2\alpha x)$ ,  $f''(x)g^2(x) = -2\alpha\beta^2x^3$  and  $f'(x)g(x) = \beta(\mu - 2\alpha x)x^{\frac{3}{2}}$ ,  $x \in \mathbb{R}_+$ . Since  $g'(x)g(x) = \frac{3}{2}\beta^2x^2$ ,  $x \in \mathbb{R}_+$  is also a quadratic polynomial, one can repeat the same lines as above to arrive at

$$I_2 \leq Ch^{\frac{3}{2}} \left( 1 + \sup_{s \in [0, T]} \|X_s\|_{L^6(\Omega; \mathbb{R})}^3 \right). \quad (5.13)$$

Also, applying the Itô formula applied to  $g(x) = \beta x^{\frac{3}{2}}$  and  $g'(x)g(x) = \frac{3}{2}\beta^2x^2$ ,  $x \in \mathbb{R}_+$ , using the



Itô isometry and considering (5.13), one can show

$$\begin{aligned}
|I_3|^2 &= \int_{t_{i-1}}^{t_i} \|g(X_s) - g(X_{t_{i-1}}) - g'(X_{t_{i-1}})(W_s - W_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R})}^2 ds \\
&= \int_{t_{i-1}}^{t_i} \left\| \int_{t_{i-1}}^s [g'(X_r)f(X_r) + \frac{1}{2}g''(X_r)g^2(X_r)] dr \right. \\
&\quad \left. + \int_{t_{i-1}}^s [g'g(X_r) - g'g(X_{t_{i-1}})] dW_r \right\|_{L^2(\Omega; \mathbb{R})}^2 ds \\
&\leq 2h \int_{t_{i-1}}^{t_i} \int_{t_{i-1}}^s \mathbb{E}[\|g'(X_r)f(X_r) + \frac{1}{2}g''(X_r)g^2(X_r)\|^2] dr \\
&\quad + 2 \int_{t_{i-1}}^{t_i} \int_{t_{i-1}}^s \mathbb{E}[\|g'g(X_r) - g'g(X_{t_{i-1}})\|^2] dr \\
&\leq Ch^3 \left(1 + \sup_{s \in [0, T]} \|X_s\|_{L^6(\Omega; \mathbb{R})}^6\right),
\end{aligned} \tag{5.14}$$

where we computed that  $g'(x)f(x) = \frac{3}{2}\beta x^{\frac{3}{2}}(\mu - \alpha x)$ ,  $g''(x)g^2(x) = \frac{3}{4}\beta^3 x^{\frac{5}{2}}$ ,  $x \in \mathbb{R}_+$ . Gathering the above three estimates together, we derive from (5.10) that

$$\|R_i\|_{L^2(\Omega; \mathbb{R})} \leq Ch^{\frac{3}{2}} \left(1 + \sup_{s \in [0, T]} \|X_s\|_{L^6(\Omega; \mathbb{R})}^3\right). \tag{5.15}$$

At the moment it remains to bound  $\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R}^d)}$ , which, similarly to (4.30), can be decomposed into two terms by a triangle inequality:

$$\begin{aligned}
\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R})} &\leq \left\| \int_{t_{i-1}}^{t_i} \mathbb{E}([f(X_s) - f(X_{t_i})] | \mathcal{F}_{t_{i-1}}) ds \right\|_{L^2(\Omega; \mathbb{R})} \\
&\quad + \frac{h}{2} \|\mathbb{E}([g'g(X_{t_i}) - g'g(X_{t_{i-1}})] | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R})}.
\end{aligned} \tag{5.16}$$

Keeping (5.11) in mind, recalling that the Itô integral vanishes under the conditional expectation (see (4.31) for clarification) and utilizing the Jensen inequality, we derive

$$\begin{aligned}
&\left\| \int_{t_{i-1}}^{t_i} \mathbb{E}([f(X_s) - f(X_{t_i})] | \mathcal{F}_{t_{i-1}}) ds \right\|_{L^2(\Omega; \mathbb{R})} \\
&= \left\| \int_{t_{i-1}}^{t_i} \int_s^{t_i} \mathbb{E}[(f'(X_r)f(X_r) + \frac{1}{2}f''(X_r)g^2(X_r)) | \mathcal{F}_{t_{i-1}}] dr ds \right\|_{L^2(\Omega; \mathbb{R})} \\
&\leq \int_{t_{i-1}}^{t_i} \int_s^{t_i} \|f'(X_r)f(X_r) + \frac{1}{2}f''(X_r)g^2(X_r)\|_{L^2(\Omega; \mathbb{R})} dr ds \\
&\leq Ch^2 \left(1 + \sup_{s \in [0, T]} \|X_s\|_{L^6(\Omega; \mathbb{R})}^3\right).
\end{aligned} \tag{5.17}$$

Following the same arguments as before, one can derive

$$\frac{h}{2} \|\mathbb{E}([g'g(X_{t_i}) - g'g(X_{t_{i-1}})] | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R})} \leq Ch^2 \left(1 + \sup_{s \in [0, T]} \|X_s\|_{L^6(\Omega; \mathbb{R})}^3\right). \tag{5.18}$$

Plugging these two estimates into (5.16) results in

$$\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R})} \leq Ch^2 \left(1 + \sup_{s \in [0, T]} \|X_s\|_{L^6(\Omega; \mathbb{R})}^3\right). \tag{5.19}$$

Analogously to (5.8), the assumption  $\alpha \geq \frac{5}{2}\beta^2$  ensures

$$\sup_{t \in [0, T]} \|X_t\|_{L^6(\Omega; \mathbb{R})} < \infty. \quad (5.20)$$

Thanks to (5.20) and Theorem 3.3, the assertion (5.9) follows based on (5.15) and (5.19).  $\square$

**Remark 5.3.** Recall that strong convergence of the implicit Milstein scheme (5.2) for the  $\frac{3}{2}$  process was analyzed by Higham et al. [21], with no convergence rates recovered. Later in [44], with the aid of the Lamperti transformation, Neuenkirch and Szpruch [44] proposed a Lamperti transformed backward Euler method for a class of scalar SDEs in a domain including the  $\frac{3}{2}$  process as a special case. There a mean-square convergence rate of order 1 was proved for the Lamperti-backward Euler method solving the  $\frac{3}{2}$  process when the model parameters obey  $\frac{\alpha}{\beta^2} > 5$  (see Propositions 3.2 from [44]). In this work we turn to the implicit Milstein scheme (5.2), covered by (2.2) and also studied in [21], and successfully prove a mean-square convergence rate of order 1 for the scheme on the condition  $\frac{\alpha}{\beta^2} > \frac{5}{2}$ . This not only fills the gap left by [21], but also significantly relaxes the restriction put on the model parameters as required in [44].

## 5.2 The double implicit Milstein scheme for the stochastic Lotka-Volterra competition model

In this subsection, we consider the scalar stochastic Lotka-Volterra (LV) competitive model [41]

$$dX_t = [bX_t - aX_t^2] dt + \sigma X_t dW_t, \quad X_0 = x_0 > 0 \quad (5.21)$$

for a single species, where individuals within the species are competitive and  $b, a, \sigma$  are all positive numbers. The well-posedness of the model in the positive domain  $(0, \infty)$  is known in the paper [41], where a positivity-preserving scheme is proposed, but with no convergence rate revealed. On the uniform mesh, we apply the double implicit Milstein scheme (2.10) with  $\theta = \eta = 1$  to numerically solve the model (5.21) as follows:

$$Y_{n+1} = Y_n + (bY_{n+1} - aY_{n+1}^2)h + \sigma Y_n \Delta W_n + \frac{1}{2}\sigma^2 Y_n |\Delta W_n|^2 - \frac{1}{2}\sigma^2 Y_{n+1} h, \quad Y_0 = X_0. \quad (5.22)$$

Obviously, it is a quadratic equation and has a unique positive solution:

$$Y_{n+1} = \frac{-(1 - bh + \frac{\sigma^2}{2}h) + \sqrt{(1 - bh + \frac{\sigma^2}{2}h)^2 + 4ahY_n(1 + \sigma\Delta W_n + \frac{\sigma^2}{2}|\Delta W_n|^2)}}{2ah} > 0, \quad (5.23)$$

given that  $Y_n > 0$ ,  $n \in \{0, 1, 2, \dots, N-1\}$ . We highlight that no additional restriction is put on the stepsize  $h > 0$  to ensure the positivity of the above approximations. Also, one can easily verify that Assumptions 3.1, 3.2 are both fulfilled in the domain  $D = (0, \infty)$ .

**Lemma 5.4.** Let  $b, a, \sigma > 0$ ,  $X_0 > 0$ . Then the stochastic LV competitive model (5.21) has a unique global solution in  $(0, \infty)$  satisfying

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |X_t|^p \right] < \infty, \quad \forall p \geq 2. \quad (5.24)$$

Moreover, the scheme (5.22) produces unique positivity preserving approximations given by (5.23). The SDE model (5.21) and the scheme (5.22) obey Assumptions 3.1, 3.2 in the domain  $D = (0, \infty)$ .

*Proof of Lemma 5.4.* The well-posedness of the model in the positive domain  $(0, \infty)$  is known in [41] and the moment bound (5.24) comes from [41, Lemma 2.2]. As discussed above, the scheme (5.22) has a unique positive solution given by (5.23). Consequently, all conditions in Assumption 3.2 are satisfied with  $D = (0, \infty)$ . Now it remains to validate conditions in Assumption 3.1. For brevity, we denote the drift and diffusion coefficients of SDE (5.1) by

$$f(x) := bx - ax^2, \quad g(x) := \sigma x, \quad x > 0. \quad (5.25)$$

By setting  $\theta = \eta = 1$ , the conditions (3.1), (3.2) in Assumption 3.1 reduce to

$$2\langle x - y, f(x) - f(y) \rangle + (q - 1)\|g(x) - g(y)\|^2 + \frac{q}{2}h\|g'(x)g(x) - g'(y)g(y)\|^2 \quad (5.26)$$

$$- h\|f(x) - f(y)\|^2 \leq L_1\|x - y\|^2,$$

$$\langle x - y, [f(x) - f(y)] - \frac{1}{2}[g'(x)g(x) - g'(y)g(y)] \rangle \leq L_2\|x - y\|^2, \quad \forall x, y \in (0, \infty). \quad (5.27)$$

Note that the diffusion  $g$  is a linear function and  $g'(x)g(x) = \sigma^2 x$ ,  $x \in \mathbb{R}_+$  is also linear. Further, note that  $f'(x) = b - 2ax \leq b$ ,  $\forall x \in (0, \infty)$ . These facts ensure that conditions (5.26)-(5.27) are both satisfied, which validates Assumption 3.1.  $\square$

Thanks to Lemma 5.4 and similarly to the proof of Theorem 5.2, we are now able to apply Theorem 3.3 to deduce the convergence rate of the numerical scheme (5.22).

**Theorem 5.5.** *Let  $b, a, \sigma > 0$ ,  $X_0 > 0$ . Let  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$  be uniquely given by (5.21) and (5.22), respectively. For  $h > 0$  satisfying  $(2b - \sigma^2)h < 1$ , there exists a constant  $C > 0$ , independent of  $N \in \mathbb{N}$ , such that*

$$\sup_{0 \leq n \leq N} \|Y_n - X_{t_n}\|_{L^2(\Omega; \mathbb{R})} \leq Ch. \quad (5.28)$$

By estimating  $\mathbb{E}[\|R_i\|^2]$  and  $\mathbb{E}[\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|^2]$ ,  $i \in \{1, 2, \dots, N\}$ , the proof of Theorem 5.5 is similar to that of Theorem 5.2 and omitted here. Different from Theorem 5.2 for the Heston- $\frac{3}{2}$  volatility model, no further restriction is put on the parameters of the model (5.21) because the diffusion coefficient is linear and all required conditions are satisfied for full parameters  $b, a, \sigma > 0$ .

### 5.3 The semi-implicit Milstein scheme for the Ait-Sahalia-type interest rate model

The next SDE financial model that we aim to numerically investigate is the generalized Ait-Sahalia-type interest rate model [1], described by

$$dX_t = (\alpha_{-1}X_t^{-1} - \alpha_0 + \alpha_1 X_t - \alpha_2 X_t^\kappa) dt + \sigma X_t^\rho dW_t, \quad t > 0, \quad X_0 = x_0 > 0, \quad (5.29)$$

where  $\alpha_{-1}, \alpha_0, \alpha_1, \alpha_2, \sigma > 0$  are positive constants and  $\kappa > 1, \rho > 1$ . Compared with the previous financial model (5.1), a complication in (5.29) is due to the drift containing a term  $\alpha_{-1}X_t^{-1}$  that does not behave well near the origin. The well-posedness of the model (5.29) has been already shown in [48, Theorem 2.1] and we repeat it as follows.

**Proposition 5.6.** *Let  $\alpha_{-1}, \alpha_0, \alpha_1, \alpha_2, \sigma > 0$  be positive constants and  $\kappa > 1, \rho > 1$ . Given any initial data  $X_0 = x_0 > 0$ , there exists a unique, positive global solution  $\{X_t\}_{t \geq 0}$  to (5.29).*

Recently, such a model has been numerically studied by many authors [9, 44, 48, 52], with an emphasis on introducing and analyzing various positivity preserving strong approximation schemes (see Remark 5.14 for more details). Different from numerical schemes introduced in [9, 44, 48, 52], we apply the newly proposed Milstein scheme to the model (5.29) with  $\kappa + 1 \geq 2\rho$ , covering both the standard regime  $\kappa + 1 > 2\rho$  and the critical regime  $\kappa + 1 = 2\rho$ , and successfully recover the expected mean-square convergence rate, by use of the previously obtained error bounds. Given a uniform mesh on the interval  $[0, T]$  with the uniform stepsize  $h = \frac{T}{N}$ ,  $N \in \mathbb{N}$ ,  $T \in (0, \infty)$ , we apply the proposed Milstein type scheme (2.2) with  $\theta = 1$ ,  $\eta = 0$  (called the semi-implicit Milstein method) to the above model (5.29) and obtain numerical approximations, given by  $Y_0 = X_0$  and

$$Y_{n+1} = Y_n + h[\alpha_{-1}Y_{n+1}^{-1} - \alpha_0 + \alpha_1Y_{n+1} - \alpha_2Y_{n+1}^\kappa] + \sigma Y_n^\rho \Delta W_n + \frac{1}{2}\rho\sigma^2 Y_n^{2\rho-1}(|\Delta W_n|^2 - h), \quad n \in \{0, 1, 2, \dots, N-1\}. \quad (5.30)$$

The next lemma concerns the well-posedness of the scheme (5.30), which can be easily checked based on the observation that the drift coefficient function satisfies a monotonicity condition (consult [48, Lemma 3.1] and (5.38)).

**Lemma 5.7.** *Let conditions in Proposition 5.6 are all satisfied. For  $h \in (0, \frac{1}{\alpha_1}]$ , the semi-implicit Milstein scheme (5.30) is well-defined in the sense that it admits a unique solution, preserving positivity of the underlying model (5.29).*

For simplicity of notation in the following analysis, we update the definitions of functions  $f, g$  in subsection 5.1 and denote the coefficients of SDE (5.29) by

$$f(x) := \alpha_{-1}x^{-1} - \alpha_0 + \alpha_1x - \alpha_2x^\kappa, \quad g(x) := \sigma x^\rho, \quad x \in \mathbb{R}_+. \quad (5.31)$$

It is easy to check that

$$g'(x)g(x) = \rho\sigma^2x^{2\rho-1}, \quad f(x) - f(y) = \left(-\frac{\alpha_{-1}}{xy} + \alpha_1 - \alpha_2\frac{x^\kappa - y^\kappa}{x-y}\right)(x-y). \quad (5.32)$$

In addition, for  $\iota \in [1, \infty)$  we introduce a function  $z_\iota: \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$  defined by

$$z_\iota(x, y) := \frac{x^\iota - y^\iota}{x-y}, \quad x, y \in \mathbb{R}_+. \quad (5.33)$$

In the following error analysis for the numerical approximations, we cope with the standard case  $\kappa + 1 > 2\rho$  and the critical case  $\kappa + 1 = 2\rho$  separately, since different cases own different model properties.

### 5.2.1 The standard case $\kappa + 1 > 2\rho$

At first, we focus on the standard case  $\kappa + 1 > 2\rho$  and recall a lemma concerning (inverse) moment bounds of the solution to (5.29), quoted from [48, Lemma 2.1].

**Lemma 5.8.** *Let conditions in Proposition 5.6 be fulfilled with  $\kappa + 1 > 2\rho$  and let  $\{X_t\}_{t \geq 0}$  be the unique solution to (5.29). Then for any  $p \geq 2$  it holds that*

$$\sup_{t \in [0, \infty)} \mathbb{E}[|X_t|^p] < \infty, \quad \sup_{t \in [0, \infty)} \mathbb{E}[|X_t|^{-p}] < \infty. \quad (5.34)$$

In order to achieve the mean-square convergence rate of the scheme by means of Theorem 3.3, we need to check all conditions required in Assumptions 3.1, 3.2, which are clarified in the forthcoming lemma.

**Lemma 5.9.** *Let conditions in Proposition 5.6 be all fulfilled with  $\kappa + 1 > 2\rho$  and let  $h \in (0, \frac{1}{\alpha_1}]$ . Then the Ait-Sahalia model (5.29) and the scheme (5.30) obey Assumptions 3.1, 3.2 in the domain  $D = (0, \infty)$ .*

*Proof of Lemma 5.9.* Note first that the well-posedness of the model and the scheme in  $D = (0, \infty)$  has been proven in Proposition 5.6 and Lemma 5.7. It remains to confirm the other conditions. We first claim that, for any  $c > 0$  there exists  $a_0 \in [0, \infty)$  such that  $z_{2\rho-1} \leq cz_\kappa + a_0$ , where we recall that  $z_i$  is defined by (5.33). Clearly,  $z_{2\rho-1}(x, y) > 0$  and  $z_\kappa(x, y) > 0$  for all  $x, y > 0$ . Without loss of generality, we assume  $x > y > 0$ . Since  $\kappa - 1 > 2\rho - 2$ , for any  $c > 0$  one can find  $a_0 \in [0, \infty)$  such that  $\sup_{v>0} ((2\rho - 1)v^{2\rho-2} - c\kappa v^{\kappa-1}) \leq a_0$ . As a consequence,

$$\begin{aligned} (x - y)(z_{2\rho-1} - cz_\kappa) &= x^{2\rho-1} - y^{2\rho-1} - c(x^\kappa - y^\kappa) \\ &= (x - y) \int_0^1 [(2\rho - 1)(y + \xi(x - y))^{2\rho-2} - c\kappa(y + \xi(x - y))^{\kappa-1}] d\xi \quad (5.35) \\ &\leq a_0(x - y), \quad \forall x > y > 0. \end{aligned}$$

The claim is thus validated. So one can choose  $c < \frac{\sqrt{2}\alpha_2}{\sqrt{\varrho}\rho\sigma^2}$  for some  $\varrho > 1$  such that  $\frac{\varrho}{2}\rho^2\sigma^4c^2 - \alpha_2^2 < 0$  and thus

$$\begin{aligned} &\varrho \frac{h}{2} \|g'(x)g(x) - g'(y)g(y)\|^2 - h\|f(x) - f(y)\|^2 \\ &= \left( h \frac{\varrho}{2} \rho^2 \sigma^4 \left( \frac{x^{2\rho-1} - y^{2\rho-1}}{x - y} \right)^2 - h \left( \frac{\alpha_1}{xy} - \alpha_1 + \alpha_2 \frac{x^\kappa - y^\kappa}{x - y} \right)^2 \right) (x - y)^2 \\ &= h \left[ \frac{\varrho}{2} \rho^2 \sigma^4 z_{2\rho-1}^2 - \left( \frac{\alpha_1}{xy} - \alpha_1 + \alpha_2 z_\kappa \right)^2 \right] (x - y)^2 \\ &= h \left[ \frac{\varrho}{2} \rho^2 \sigma^4 z_{2\rho-1}^2 - \left( \frac{\alpha_1}{xy} - \alpha_1 \right)^2 - \alpha_2^2 z_\kappa^2 - 2 \left( \frac{\alpha_1}{xy} - \alpha_1 \right) \alpha_2 z_\kappa \right] (x - y)^2 \quad (5.36) \\ &\leq h \left[ \frac{\varrho}{2} \rho^2 \sigma^4 z_{2\rho-1}^2 - \alpha_2^2 z_\kappa^2 + 2\alpha_1 \alpha_2 z_\kappa \right] (x - y)^2 \\ &\leq h \left[ \frac{\varrho}{2} \rho^2 \sigma^4 (cz_\kappa + a_0)^2 - \alpha_2^2 z_\kappa^2 + 2\alpha_1 \alpha_2 z_\kappa \right] (x - y)^2 \\ &= h \left[ \left( \frac{\varrho}{2} \rho^2 \sigma^4 c^2 - \alpha_2^2 \right) z_\kappa^2 + (\varrho \rho^2 \sigma^4 ca_0 + 2\alpha_1 \alpha_2) z_\kappa + \frac{\varrho}{2} \rho^2 \sigma^4 a_0^2 \right] (x - y)^2 \\ &\leq Ch(x - y)^2, \quad \forall x, y \in \mathbb{R}_+. \end{aligned}$$

Furthermore, one can readily compute that, for any  $\kappa + 1 > 2\rho$  and for some  $q > 2$ ,

$$\sup_{x>0} (f'(x) + \frac{q-1}{2}|g'(x)|^2) = \sup_{x>0} \left( -\alpha_{-1}x^{-2} + \alpha_1 - \alpha_2\kappa x^{\kappa-1} + \frac{(q-1)\sigma^2\rho^2}{2}x^{2\rho-2} \right) =: L < \infty. \quad (5.37)$$

This implies that

$$\begin{aligned} &\langle x - y, f(x) - f(y) \rangle + \frac{q-1}{2} \|g(x) - g(y)\|^2 \\ &= \int_0^1 f'(y + \xi(x - y)) d\xi \cdot (x - y)^2 + \frac{q-1}{2} \left| \int_0^1 g'(y + \xi(x - y)) d\xi \right|^2 \cdot (x - y)^2 \quad (5.38) \\ &\leq \int_0^1 [f'(y + \xi(x - y)) + \frac{q-1}{2}|g'(y + \xi(x - y))|^2] d\xi \cdot (x - y)^2 \\ &\leq L(x - y)^2, \quad \forall x, y \in \mathbb{R}_+. \end{aligned}$$

Gathering (5.36) and (5.38) together, the condition (3.1) is hence justified in the domain  $D = (0, \infty)$  with  $\theta = 1, \eta = 0$ . From (5.38), one can assert that (3.2) is satisfied in  $D = (0, \infty)$  with  $\theta = 1, \eta = 0, L_2 = \alpha_1$ . Thus all conditions in Assumption 3.1 are fulfilled in the domain  $D = (0, \infty)$ . Assumption 3.2 follows by taking Proposition 5.6, Lemmas 5.7, 5.8 into consideration.  $\square$

At the moment, we are well prepared to carry out the error analysis for the numerical approximations with the help of Theorem 3.3.

**Theorem 5.10.** Let  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$  be solutions to (5.29) and (5.30), respectively. Let  $q \in (2, \infty)$ ,  $\varrho \in (1, \infty)$ , let  $\alpha_{-1}, \alpha_0, \alpha_1, \alpha_2, \sigma > 0$ , let  $\kappa > 1, \rho > 1$  obey  $\kappa + 1 > 2\rho$ , and let  $h \in (0, \frac{1}{2\alpha_1})$ . Then there exists a constant  $C > 0$ , independent of  $N \in \mathbb{N}$ , such that

$$\sup_{1 \leq n \leq N} \|X_{t_n} - Y_n\|_{L^2(\Omega; \mathbb{R})} \leq Ch. \quad (5.39)$$

*Proof of Theorem 5.10.* As implied by Lemma 5.9, all conditions in Assumptions 3.1, 3.2 are fulfilled in  $D = (0, \infty)$ . Based on Theorem 3.3, one just needs to properly estimate  $\|R_i\|_{L^2(\Omega; \mathbb{R})}$  and  $\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R}^d)}$ ,  $i \in \{1, 2, \dots, N\}$ . Following the notation used in (3.6) and (5.31), we first split the estimate of  $\|R_i\|_{L^2(\Omega; \mathbb{R})}$  as follows:

$$\begin{aligned} \|R_i\|_{L^2(\Omega; \mathbb{R})} &\leq \left\| \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_i}) \, ds \right\|_{L^2(\Omega; \mathbb{R})} \\ &\quad + \left\| \int_{t_{i-1}}^{t_i} g(X_s) - g(X_{t_{i-1}}) - g'(X_{t_{i-1}})(W_s - W_{t_{i-1}}) \, dW_s \right\|_{L^2(\Omega; \mathbb{R})} \\ &=: I_4 + I_5. \end{aligned} \quad (5.40)$$

Repeating the same arguments as used in (5.12), we apply the Itô formula to  $f(x) = \alpha_{-1}x^{-1} - \alpha_0 + \alpha_1x - \alpha_2x^\kappa$ ,  $x \in \mathbb{R}_+$  and use Lemma 5.8 to derive

$$\begin{aligned} I_4 &\leq \int_{t_{i-1}}^{t_i} \|f(X_s) - f(X_{t_i})\|_{L^2(\Omega; \mathbb{R})} \, ds \\ &\leq Ch^{\frac{3}{2}} \left( 1 + \sup_{s \in [0, T]} \|X_s\|_{L^{4\kappa-2}(\Omega; \mathbb{R})}^{2\kappa-1} + \sup_{s \in [0, T]} \|X_s^{-1}\|_{L^6(\Omega; \mathbb{R})}^3 \right) \\ &\leq Ch^{\frac{3}{2}}. \end{aligned} \quad (5.41)$$

Similarly to (5.14), by means of the Itô isometry and the Itô formula applied to  $g(x) = \sigma x^\rho$  and  $g'(x)g(x) = \rho\sigma^2x^{2\rho-1}$ ,  $x \in \mathbb{R}_+$  one can show

$$\begin{aligned} |I_5|^2 &\leq 2h \int_{t_{i-1}}^{t_i} \int_{t_{i-1}}^s \mathbb{E}[\|g'(X_r)f(X_r) + \frac{1}{2}g''(X_r)g^2(X_r)\|^2] \, dr \, ds \\ &\quad + 2 \int_{t_{i-1}}^{t_i} \int_{t_{i-1}}^s \mathbb{E}[\|g'g(X_r) - g'g(X_{t_{i-1}})\|^2] \, dr \, ds \\ &\leq Ch^3, \end{aligned} \quad (5.42)$$

where the (inverse) moment bounds in Lemma 5.8 were also used for the last step. Inserting (5.41) and (5.42) into (5.40) implies

$$\|R_i\|_{L^2(\Omega; \mathbb{R})} \leq Ch^{\frac{3}{2}}. \quad (5.43)$$

In the same spirit of (5.17), we rely on the use of Itô formula applied to  $f(x) = \alpha_{-1}x^{-1} - \alpha_0 + \alpha_1x - \alpha_2x^\kappa$  to show

$$\begin{aligned} \|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R}^d)} &\leq \left\| \int_{t_{i-1}}^{t_i} \mathbb{E}([f(X_s) - f(X_{t_i})] | \mathcal{F}_{t_{i-1}}) \, ds \right\|_{L^2(\Omega; \mathbb{R}^d)} \\ &\leq \int_{t_{i-1}}^{t_i} \int_s^{t_i} \|f'(X_r)f(X_r) + \frac{1}{2}f''(X_r)g^2(X_r)\|_{L^2(\Omega; \mathbb{R})} \, dr \, ds \\ &\leq Ch^2, \end{aligned} \quad (5.44)$$



where we recalled that the Itô integral vanishes under the conditional expectation and also used the Jensen inequality and Lemma 5.8. Armed with these two estimates, one can apply Theorem 3.3 to arrive at the desired assertion.  $\square$

### 5.2.2 The critical case $\kappa + 1 = 2\rho$

In what follows we turn to the general critical case  $\kappa + 1 = 2\rho$  and present first a lemma concerning (inverse) moment bounds of the solution process, which can be proved by following the same lines in the proof of Lemma 5.8 (cf. [48, Lemma 2.1]).

**Lemma 5.11.** *Let conditions in Proposition 5.6 be all fulfilled with  $\kappa + 1 = 2\rho$  and let  $\{X_t\}_{t \geq 0}$  be the unique solution to (5.29). Then we have, for any  $2 \leq p_1 \leq \frac{\sigma^2 + 2\alpha_2}{\sigma^2}$  and for any  $p_2 \geq 2$ ,*

$$\sup_{t \in [0, \infty)} \mathbb{E}[|X_t|^{p_1}] < \infty, \quad \sup_{t \in [0, \infty)} \mathbb{E}[|X_t|^{-p_2}] < \infty. \quad (5.45)$$

For the purpose of analyzing the convergence rate of the numerical approximations, we validate all conditions of Assumptions 3.1, 3.2 in the next lemma, which is required by Theorem 3.3.

**Lemma 5.12.** *Let conditions in Proposition 5.6 be all fulfilled with  $\kappa + 1 = 2\rho$  and let  $h \in (0, \frac{1}{\alpha_1}]$ . Let the model parameters obey  $\frac{\alpha_2}{\sigma^2} \geq 2\kappa - \frac{3}{2}$  and  $\frac{\alpha_2}{\sigma^2} > \frac{\kappa+1}{2\sqrt{2}}$ . Then the SDE model (5.29) and the scheme (5.30) satisfy Assumptions 3.1, 3.2 in the domain  $D = (0, \infty)$ .*

*Proof of Lemma 5.12.* Recall that the well-posedness of the model and the scheme in  $D = (0, \infty)$  has been proven in Proposition 5.6 and Lemma 5.7. It remains to verify the other conditions. Thanks to the assumptions  $\kappa + 1 = 2\rho$  and  $\frac{\alpha_2}{\sigma^2} > \frac{\kappa+1}{2\sqrt{2}}$ , one can find  $\varrho > 1$  such that  $\alpha_2^2 > \frac{\varrho}{2}\rho^2\sigma^4 = \frac{\varrho}{8}(\kappa + 1)^2\sigma^4$  and thus

$$\begin{aligned} & h \frac{\varrho}{2} \|g'(x) - g'(y)\|^2 - h \|f(x) - f(y)\|^2 \\ &= h \left( \frac{\varrho}{2} \rho^2 \sigma^4 z_\kappa^2 - \left( \frac{\alpha_{-1}}{xy} - \alpha_1 + \alpha_2 z_\kappa \right)^2 \right) (x - y)^2 \\ &= h \left[ \frac{\varrho}{2} \rho^2 \sigma^4 z_\kappa^2 - \left( \frac{\alpha_{-1}}{xy} - \alpha_1 \right)^2 - \alpha_2^2 z_\kappa^2 - 2 \left( \frac{\alpha_{-1}}{xy} - \alpha_1 \right) \alpha_2 z_\kappa \right] (x - y)^2 \\ &\leq h \left[ \left( \frac{\varrho}{2} \rho^2 \sigma^4 - \alpha_2^2 \right) z_\kappa^2 + 2\alpha_1 \alpha_2 z_\kappa \right] (x - y)^2 \\ &\leq Ch(x - y)^2, \quad \forall x, y \in \mathbb{R}_+. \end{aligned} \quad (5.46)$$

Noting  $\kappa + 1 = 2\rho$  again, one can deduce from (5.37) that

$$\sup_{x > 0} \left( f'(x) + \frac{q-1}{2} |g'(x)|^2 \right) \leq \sup_{x > 0} \left( \alpha_1 - \left( \alpha_2 \kappa - \frac{(q-1)\sigma^2 \rho^2}{2} \right) x^{\kappa-1} \right), \quad \forall x \in \mathbb{R}_+. \quad (5.47)$$

Since  $\frac{\alpha_2}{\sigma^2} \geq 2\kappa - \frac{3}{2} > \frac{\kappa+3}{8} > \frac{1}{8}(\kappa + 2 + \frac{1}{\kappa})$  for  $\kappa > 1$ , one can find  $q > 2$  such that  $\frac{\alpha_2}{\sigma^2} \geq \frac{q-1}{8}(\kappa + 2 + \frac{1}{\kappa})$ , i.e.,  $\alpha_2 \kappa - \frac{(q-1)\sigma^2 \rho^2}{2} = \alpha_2 \kappa - \frac{(q-1)}{8} \sigma^2 (\kappa + 1)^2 \geq 0$  in (5.47), and thus, similarly to (5.38),

$$\langle x - y, f(x) - f(y) \rangle + \frac{q-1}{2} \|g(x) - g(y)\|^2 \leq \alpha_1 (x - y)^2, \quad \forall x, y \in \mathbb{R}_+. \quad (5.48)$$

Combining this with (5.46) ensures that the condition (3.1) is fulfilled in  $D = (0, \infty)$  with  $\theta = 1, \eta = 0$ . The condition (3.2) follows from (5.48) directly. Finally, since  $\frac{\sigma^2 + 2\alpha_2}{\sigma^2} \geq 4\kappa - 2 > 2\kappa$  by assumption  $\frac{\alpha_2}{\sigma^2} \geq 2\kappa - \frac{3}{2}$ ,  $\kappa > 1$ , in view of Lemma 5.11 one can infer  $\sup_{t \in [0, T]} \|X_t\|_{L^2(\Omega; \mathbb{R})} < \infty$  and

$$\sup_{t \in [0, T]} \|f(X_t)\|_{L^2(\Omega; \mathbb{R})} \leq \sup_{t \in [0, T]} \left( \alpha_{-1} \|X_t^{-1}\|_{L^2(\Omega; \mathbb{R})} + \alpha_0 + \alpha_1 \|X_t\|_{L^2(\Omega; \mathbb{R})} + \alpha_2 \|X_t\|_{L^{2\kappa}(\Omega; \mathbb{R})}^\kappa \right) < \infty. \quad (5.49)$$

Therefore, all conditions in Assumptions 3.1, 3.2 are confirmed in the domain  $D = (0, \infty)$ .  $\square$

Now we are in a position to derive the convergence order with the aid of Theorem 3.3.

**Theorem 5.13.** *Let  $\{X_t\}_{t \in [0, T]}$  and  $\{Y_n\}_{0 \leq n \leq N}$  be solutions to (5.29) and (5.30), respectively. Let conditions in Proposition 5.6 be all fulfilled with  $\kappa + 1 = 2\rho$  and let  $h \in (0, \frac{1}{2\alpha_1})$ . Let the model parameters  $\alpha_{-1}, \alpha_0, \alpha_1, \alpha_2, \sigma > 0$ ,  $\kappa > 1, \rho > 1$  obey  $\frac{\alpha_2}{\sigma^2} \geq 2\kappa - \frac{3}{2}$  and  $\frac{\alpha_2}{\sigma^2} > \frac{\kappa+1}{2\sqrt{2}}$ . Then there exists a constant  $C > 0$ , independent of  $N \in \mathbb{N}$ , such that*

$$\sup_{1 \leq n \leq N} \|X_{t_n} - Y_n\|_{L^2(\Omega; \mathbb{R})} \leq Ch. \quad (5.50)$$

*Proof of Theorem 5.13.* As already verified in Lemma 5.12, all conditions in Assumptions 3.1, 3.2 are fulfilled in  $D = (0, \infty)$ . Based on Theorem 3.3, one only needs to properly estimate  $\|R_i\|_{L^2(\Omega; \mathbb{R})}$  and  $\|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R}^d)}$ . Similarly as above, we split the error term  $\|R_i\|_{L^2(\Omega; \mathbb{R})}$  into two parts:

$$\begin{aligned} \|R_i\|_{L^2(\Omega; \mathbb{R})} &\leq \left\| \int_{t_{i-1}}^{t_i} f(X_s) - f(X_{t_i}) \, ds \right\|_{L^2(\Omega; \mathbb{R})} \\ &\quad + \left\| \int_{t_{i-1}}^{t_i} g(X_s) - g(X_{t_{i-1}}) - g'(X_{t_{i-1}})(W_s - W_{t_{i-1}}) \, dW_s \right\|_{L^2(\Omega; \mathbb{R})} \\ &=: I_6 + I_7, \end{aligned} \quad (5.51)$$

where the coefficients  $f, g$  are defined by (5.31). The Itô formula applied to  $f(x) = \alpha_{-1}x^{-1} - \alpha_0 + \alpha_1x - \alpha_2x^\kappa$ ,  $x \in \mathbb{R}_+$  gives

$$\begin{aligned} I_6 &\leq \int_{t_{i-1}}^{t_i} \|f(X_s) - f(X_{t_i})\|_{L^2(\Omega; \mathbb{R})} \, ds \\ &\leq Ch^{\frac{3}{2}} \left( 1 + \sup_{s \in [0, T]} \|X_s\|_{L^{4\kappa-2}(\Omega; \mathbb{R})}^{2\kappa-1} + \sup_{s \in [0, T]} \|X_s^{-1}\|_{L^6(\Omega; \mathbb{R})}^3 \right). \end{aligned} \quad (5.52)$$

Following the same lines as in (5.42), one can similarly show

$$|I_7|^2 \leq Ch^3 \left( 1 + \sup_{s \in [0, T]} \mathbb{E}[|X_s|^{4\kappa-2}] + 1_{\{\rho < 2\}} \sup_{s \in [0, T]} \mathbb{E}[|X_s|^{-(4-2\rho)}] \right), \quad (5.53)$$

where we set  $1_{\{\rho < 2\}} = 1$  for  $\rho < 2$  and  $1_{\{\rho < 2\}} = 0$  for  $\rho \geq 2$ . Since  $\frac{\sigma^2 + 2\alpha_2}{\sigma^2} \geq 4\kappa - 2$  by the assumption  $\frac{\alpha_2}{\sigma^2} \geq 2\kappa - \frac{3}{2}$ , we can plug these two estimates into (5.51) and use Lemma 5.11 to get

$$\|R_i\|_{L^2(\Omega; \mathbb{R})} \leq Ch^{\frac{3}{2}}. \quad (5.54)$$

Moreover, similarly to (5.44), applying the Itô formula to  $f(x) = \alpha_{-1}x^{-1} - \alpha_0 + \alpha_1x - \alpha_2x^\kappa$  and noting the Itô integral vanishes under the conditional expectation we deduce

$$\begin{aligned} \|\mathbb{E}(R_i | \mathcal{F}_{t_{i-1}})\|_{L^2(\Omega; \mathbb{R}^d)} &\leq \left\| \int_{t_{i-1}}^{t_i} \mathbb{E}([f(X_s) - f(X_{t_i})] | \mathcal{F}_{t_{i-1}}) \, ds \right\|_{L^2(\Omega; \mathbb{R}^d)} \\ &\leq \int_{t_{i-1}}^{t_i} \int_s^{t_i} \|f'(X_r)f(X_r) + \frac{1}{2}f''(X_r)g^2(X_r)\|_{L^2(\Omega; \mathbb{R})} \, dr \, ds \\ &\leq Ch^3 \left( 1 + \sup_{s \in [0, T]} \|X_s\|_{L^{4\kappa-2}(\Omega; \mathbb{R})}^{2\kappa-1} + \sup_{s \in [0, T]} \|X_s^{-1}\|_{L^6(\Omega; \mathbb{R})}^3 \right). \end{aligned} \quad (5.55)$$

In light of Lemma 5.11 and with the help of Theorem 3.3, one can obtain the assertion (5.50).  $\square$

**Remark 5.14.** Recall that Szpruch et al. [48] examined the backward Euler method for the Ait-Sahalia model (5.29) and proved its strong convergence only when  $\kappa+1 > 2\rho$ , but without revealing a rate of convergence. Very recently, the authors of [52] fill the gap by identifying the expected mean-square convergence rate of order  $\frac{1}{2}$  for stochastic theta methods applied to the Ait-Sahalia model under conditions  $\kappa+1 \geq 2\rho$ . In 2014, a kind of Lamperti-backward Euler method was introduced in [44] for the Ait-Sahalia model, with a mean-square convergence rate of order 1 identified for the full parameter range in the general standard case  $\kappa+1 > 2\rho$  and for a particular critical case  $\kappa=2, \rho=1.5$  when  $\frac{\alpha_2}{\sigma^2} > 5$  (see Propositions 3.5, 3.6 from [44]). As shown above, we apply the semi-implicit Milstein method (5.30) to the Ait-Sahalia model, which is able to treat both the general standard case and a more general critical case  $\kappa+1 = 2\rho$  for any  $\kappa, \rho > 1$ . Moreover, we prove a mean-square convergence rate of order 1 for the full parameter range in the general standard case and for parameters satisfying  $\frac{\alpha_2}{\sigma^2} > 2\kappa - \frac{3}{2}$  and  $\frac{\alpha_2}{\sigma^2} > \frac{\kappa+1}{2\sqrt{2}}$  in the general critical case. For the special critical case  $\kappa=2, \rho=1.5$ , the restriction on parameters reduces into  $\frac{\alpha_2}{\sigma^2} > \frac{5}{2}$ , which is moderately more relaxed than  $\frac{\alpha_2}{\sigma^2} > 5$  as required in [44].

## 5.4 Numerical tests

The aim of this subsection is to illustrate the above theoretical findings by providing several numerical examples. Two different schemes covered by (2.2) are utilized to simulate the two previously studied financial models. The resulting mean-square approximation errors are computed at the endpoint  $T=1$  and the desired expectations are approximated by averages over 10000 samples. Moreover, the “exact” solutions are identified as numerical ones using a fine stepsize  $h_{\text{exact}} = 2^{-12}$ .

As the first example, let us first look at the following SDE,

$$dX_t = X_t(\mu - \alpha X_t) dt + (\beta X_t^{3/2} + \sigma X_t) dW(t), \quad X_0 = 1, \quad t \in (0, 1]. \quad (5.56)$$

When  $\sigma = 0$  and  $\beta = 0$ , the considered SDE (5.56) reduces to the Heston  $\frac{3}{2}$ -volatility model (5.1) and the stochastic LV competitive model (5.21), respectively. We choose the parameters  $(\mu, \alpha, \beta, \sigma) = (2, \frac{5}{2}, 1, 0)$  such that  $\alpha \geq \frac{5}{2}\beta^2$  for the  $\frac{3}{2}$ -model (5.1) and  $(\mu, \alpha, \beta, \sigma) = (2, 1, 0, 1)$  for the stochastic LV competitive model (5.21). By taking  $\theta = \eta = 1$ , we discrete these two models by the drift-diffusion double implicit Milstein method (2.2), which is explicitly solvable here (see (5.3) and (5.23)). In the following simulations, the expectations are approximated by computing averages over  $10^4$  samples and the “exact” solutions are identified as approximations using a fine stepsize  $h_{\text{exact}} = 2^{-12}$ . It turns out that the resulting numerical approximations always remain positive for all  $10^4$  paths. In Figure 2, we present one-path simulations of the drift-diffusion double implicit Milstein method for the Heston  $\frac{3}{2}$ -volatility model (Left) and the stochastic LV model (Right), which are shown to be positive. To test the mean-square convergence rates, we depict in Figure 3 mean-square approximation errors  $e_h$  against six different stepsizes  $h = 2^{-i}, i = 4, 5, \dots, 9$  on a log-log scale. Also, two reference lines of slope 1 and  $\frac{1}{2}$  are given there. From Figure 3 one can easily detect that the approximation errors decrease at a slope close to 1 when stepsizes shrink, coinciding with the predicted convergence order obtained in Theorem 5.2 and Theorem 5.5. Suppose that the approximation errors  $e_h$  obey a power law relation  $e_h = Ch^\delta$  for  $C, \delta > 0$ , so that  $\log e_h = \log C + \delta \log h$ . Then we do a least squares power law fit for  $\delta$  and get the value 0.9923 for the rate  $\delta$  with residual of 0.0719. Again, this confirms the expected convergence rate in Theorem 5.2 and Theorem 5.5.

As the second example model, we look at the Ait-Sahalia interest rate model, given by

$$dX_t = (\alpha_{-1}X_t^{-1} - \alpha_0 + \alpha_1X_t - \alpha_2X_t^\kappa) dt + \sigma X_t^\rho dW_t, \quad X_0 = 1, \quad t \in (0, 1]. \quad (5.57)$$




Figure 2: One-path simulations of the drift-diffusion double implicit Milstein method for the Heston  $\frac{3}{2}$ -volatility model (Left) and the stochastic LV model (Right).




Figure 3: Mean-square convergence rates of the drift-diffusion double implicit Milstein method for the Heston  $\frac{3}{2}$ -volatility model (Left) and the stochastic LV model (Right).




Figure 4: Mean-square convergence rates of the semi-implicit Milstein method (5.30) for the Ait-Sahalia interest rate model (Left for Case I and right for Case II).

Let us consider both the standard case  $\kappa + 1 > 2\rho$  and the critical case  $\kappa + 1 = 2\rho$ , by taking two sets of model parameters:

- Case I:  $\kappa = 4$ ,  $\rho = 2$ ,  $\alpha_{-1} = \frac{3}{2}$ ,  $\alpha_0 = 2$ ,  $\alpha_1 = 1$ ,  $\alpha_2 = 1$ ,  $\sigma = 1$ ;
- Case II:  $\kappa = 3$ ,  $\rho = 2$ ,  $\alpha_{-1} = \frac{3}{2}$ ,  $\alpha_0 = 2$ ,  $\alpha_1 = 1$ ,  $\alpha_2 = \frac{9}{2}$ ,  $\sigma = 1$ .

It is easy to check that Case I corresponds to the standard case and Case II corresponds to the critical case  $\kappa + 1 = 2\rho$  satisfying  $\frac{\alpha_2}{\sigma^2} \geq 2\kappa - \frac{3}{2}$  and  $\frac{\alpha_2}{\sigma^2} > \frac{\kappa+1}{2\sqrt{2}}$ . The semi-implicit Milstein scheme (5.30) is used to simulate the model (5.57) for these two cases. As shown in Figure 4, the mean-square approximation error lines have slopes close to 1 for both cases. A least squares fit produces a rate 0.9798 with residual of 0.0929 for Case I and a rate 1.0129 with residual of 0.0968 for Case II. Hence, numerical results are consistent with strong order of convergence equal to one, as already revealed in Theorem 5.10 and Theorem 5.13.

## 6 Conclusion

The present work introduces a family of implicit Milstein type methods for strong approximations of stochastic differential equations (SDEs) with non-globally Lipschitz drift and diffusion coefficients. An easy and direct approach of the error analysis is developed to recover the expected mean-square convergence rate of order one for the proposed schemes. In particular, the optimal convergence rate of the positivity preserving schemes applied to three models in practice is obtained for the first time and more relaxed conditions are required, compared with existing results for first order schemes in the literature. In the future, we attempt to identify the general  $L^p$  rate

of convergence with  $p \geq 2$  for the schemes, which is highly non-trivial.

**Funding** This work was supported by Natural Science Foundation of China (12071488, 11971488) and Natural Science Foundation of Hunan Province for Distinguished Young Scholars (2020JJ2040).

## Declarations

**Conflict of interest** The author declares no competing interests.

## References

- [1] Y. Ait-Sahalia. Testing continuous-time models of the spot interest rate. *The review of financial studies*, 9(2):385–426, 1996.
- [2] A. Alfonsi. Strong order one convergence of a drift implicit Euler scheme: Application to the CIR process. *Statistics & Probability Letters*, 83(2):602–607, 2013.
- [3] A. Andersson and R. Kruse. Mean-square convergence of the BDF2-Maruyama and backward Euler schemes for SDE satisfying a global monotonicity condition. *BIT Numerical Mathematics*, 57(1):21–53, 2017.
- [4] W.-J. Beyn, E. Isaak, and R. Kruse. Stochastic C-stability and B-consistency of explicit and implicit Euler-type schemes. *Journal of Scientific Computing*, 67(3):955–987, 2016.
- [5] W.-J. Beyn, E. Isaak, and R. Kruse. Stochastic C-stability and B-consistency of explicit and implicit Milstein-type schemes. *Journal of Scientific Computing*, 70(3):1042–1077, 2017.
- [6] M. Bossy and H. O. Quinteros. Strong convergence of the symmetrized Milstein scheme for some CEV-like SDEs. *Bernoulli*, 24(3):1995–2042, 2015.
- [7] E. Buckwar and T. Sickenberger. A comparative linear mean-square stability analysis of Maruyama- and Milstein-type methods. *Mathematics & Computers in Simulation*, 81(6):1110–1127, 2011.
- [8] C. Kahl, M. Gunther and T. Rosberg. Structure preserving stochastic integration schemes in interest rate derivative modeling. *Applied Numerical Mathematics*, 58:284–295, 2008.
- [9] J. Chassagneux, A. Jacquier, and I. Mihaylov. An explicit Euler scheme with strong rate of convergence for financial SDEs with non-lipschitz coefficients. *SIAM Journal on Financial Mathematics*, 7(1):993–1021, 2016.
- [10] J. Cui, J. Hong, and D. Sheng. Convergence in density of splitting AVF scheme for stochastic Langevin equation. *arXiv:1906.03439*, 2019.
- [11] D. H. Ahn and B. Gao. A parametric nonlinear model of term structure dynamics. *Review of Financial Studies*, 12:721, 1999.
- [12] W. Fang and M. B. Giles. Adaptive Euler–Maruyama method for SDEs with nonglobally Lipschitz drift. *Ann. Appl. Probab.*, 30(2):526–560, 2020.

- [13] S. Gan, Y. He, and X. Wang. Tamed Runge-Kutta methods for SDEs with super-linearly growing drift and diffusion coefficients. *Applied Numerical Mathematics*, 152:379–402, 2020.
- [14] M. B. Giles. Improved multilevel Monte Carlo convergence using the Milstein scheme. In: *Keller A., Heinrich S., Niederreiter H. (eds) Monte Carlo and Quasi-Monte Carlo Methods 2006. Springer, Berlin, Heidelberg*, 2008.
- [15] Giles M. Multilevel monte carlo path simulation. *Oper. Res.*, 56:607–617, 2008.
- [16] Giles M., Debrabant K., Rößler A. Analysis of multilevel monte carlo path simulation using the milstein discretisation. *Discrete Cont Dyn-B*, 24(8), 2019.
- [17] H. Gilling and T. Shardlow. Sdelab: A package for solving stochastic differential equations in matlab. *Journal of Computational and Applied Mathematics*, 205(2):1002–1018, 2007.
- [18] Q. Guo, W. Liu, X. Mao, and R. Yue. The truncated Milstein method for stochastic differential equations with commutative noise. *Journal of Computational and Applied Mathematics*, 338:298–310, 2018.
- [19] S. Heston. A simple new formula for options with stochastic volatility. *Course Notes of Washington University in St. Louis, Missouri*, 1997.
- [20] D. J. Higham, X. Mao, and A. M. Stuart. Strong convergence of Euler-type methods for nonlinear stochastic differential equations. *SIAM Journal on Numerical Analysis*, 40(3):1041–1063, 2002.
- [21] D. J. Higham, X. Mao, and L. Szpruch. Convergence, non-negativity and stability of a new Milstein scheme with applications to finance. *Discrete & Continuous Dynamical Systems-Series B*, 18(8):2083–2100, 2013.
- [22] Y. Hu. Semi-implicit euler-maruyama scheme for stiff stochastic equations. in *Stochastic Analysis and Related Topics V: The Silvri Workshop, Progr. Probab.*, 38:183–202, 1996.
- [23] M. Hutzenthaler and A. Jentzen. Numerical approximation of stochastic differential equations with non-globally Lipschitz continuous coefficients. *Mem. Amer. Math. Soc.*, 236(1112), 2015.
- [24] M. Hutzenthaler and A. Jentzen. On a perturbation theory and on strong convergence rates for stochastic ordinary and partial differential equations with non-globally monotone coefficients. *Ann. Probab.*, 48(1):53–93, 2020.
- [25] M. Hutzenthaler, A. Jentzen, and P. E. Kloeden. Strong and weak divergence in finite time of Euler’s method for stochastic differential equations with non-globally Lipschitz continuous coefficients. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 467(2130):1563–1576, 2011.
- [26] M. Hutzenthaler, A. Jentzen, and P. E. Kloeden. Strong convergence of an explicit numerical method for SDEs with nonglobally Lipschitz coefficients. *Ann. Appl. Probab.*, 22(4):1611–1641, 2012.
- [27] M. Hutzenthaler, A. Jentzen, and X. Wang. Exponential integrability properties of numerical approximation processes for nonlinear stochastic differential equations. *Math. Comput.*, 87(311):1353–1413, 2018.



- [28] C. Kelly, G. Lord, and F. Sun. Strong convergence of an adaptive time-stepping Milstein method for SDEs with one-sided Lipschitz drift. *arXiv preprint arXiv:1909.00099*, 2019.
- [29] C. Kelly and G. J. Lord. Adaptive time-stepping strategies for nonlinear stochastic systems. *IMA Journal of Numerical Analysis*, 38(3):1523–1549, 2017.
- [30] P. E. Kloeden and E. Platen. *Numerical solution of stochastic differential equations*, volume 23. Springer Science & Business Media, 1992.
- [31] R. Kruse and Y. Wu. A randomized Milstein method for stochastic differential equations with non-differentiable drift coefficients. *Discrete & Continuous Dynamical Systems-B*, 24(8):3475–3502, 2019.
- [32] C. Kumar and S. Sabanis. On Milstein approximations with varying coefficients: the case of super-linear diffusion coefficients. *BIT Numerical Mathematics*, 59(4):929–968, 2019.
- [33] A. L. Lewis. *Option Valuation Under Stochastic Volatility*. Finance Press, 2000.
- [34] X. Li and G. Yin. Explicit Milstein schemes with truncation for nonlinear stochastic differential equations: convergence and its rate. *Journal of Computational and Applied Mathematics*, 374:112771, 2020.
- [35] Z. Liu and Z. Qiao. Strong approximation of monotone stochastic partial differential equations driven by multiplicative noise. *Stochastics and Partial Differential Equations: Analysis and Computations*, 9(3):559–602, 2021.
- [36] A. K. Majee and A. Prohl. Optimal strong rates of convergence for a space-time discretization of the stochastic allen–cahn equation with multiplicative noise. *Computational Methods in Applied Mathematics*, 18(2):297–311, 2018.
- [37] X. Mao. The truncated Euler–Maruyama method for stochastic differential equations. *J. Comput. Appl. Math.*, 290:370–384, 2015.
- [38] X. Mao. Convergence rates of the truncated Euler–Maruyama method for stochastic differential equations. *J. Comput. Appl. Math.*, 296:362–375, 2016.
- [39] X. Mao and L. Szpruch. Strong convergence and stability of implicit numerical methods for stochastic differential equations with non-globally Lipschitz continuous coefficients. *Journal of Computational and Applied Mathematics*, 238:14–28, 2013.
- [40] X. Mao and L. Szpruch. Strong convergence rates for backward Euler–Maruyama method for non-linear dissipative-type stochastic differential equations with super-linear diffusion coefficients. *Stochastics An International Journal of Probability and Stochastic Processes*, 85(1):144–171, 2013.
- [41] X. Mao, F. Wei, and T. Wiriyakraikul. Positivity preserving truncated euler–maruyama method for stochastic lotka–volterra competition model. *Journal of Computational and Applied Mathematics*, 394:113566, 2021.
- [42] G. N. Milstein, E. Platen, and H. Schurz. Balanced implicit methods for stiff stochastic systems. *SIAM Journal on Numerical Analysis*, 35(3):1010–1019, 1998.

- [43] G. N. Milstein and M. V. Tretyakov. *Stochastic numerics for mathematical physics*. Springer Science & Business Media, 2013.
- [44] A. Neuenkirch and L. Szpruch. First order strong approximations of scalar sdes defined in a domain. *Numerische Mathematik*, 128(1):103–136, 2014.
- [45] A. Rößler. Runge–kutta methods for the strong approximation of solutions of stochastic differential equations. *SIAM Journal on Numerical Analysis*, 48(3):922–952, 2010.
- [46] S. Sabanis. A note on tamed Euler approximations. *Electron. Commun. Probab*, 18(47):1–10, 2013.
- [47] S. Sabanis. Euler approximations with varying coefficients: the case of super-linearly growing diffusion coefficients. *Ann. Appl. Probab.*, 26(4):2083–2105, 2016.
- [48] L. Szpruch, X. Mao, D. J. Higham, and J. Pan. Numerical simulation of a strongly nonlinear Ait-Sahalia-type interest rate model. *BIT Numerical Mathematics*, 51(2):405–425, 2011.
- [49] L. Szpruch and X. Zhang. V-integrability, asymptotic stability and comparison property of explicit numerical schemes for non-linear SDEs. *Math. Comput.*, 87(310):755–783.
- [50] M. V. Tretyakov and Z. Zhang. A fundamental mean-square convergence theorem for SDEs with locally Lipschitz coefficients and its applications. *SIAM J. Numer. Anal.*, 51(6):3135–3162, 2013.
- [51] X. Wang and S. Gan. The tamed Milstein method for commutative stochastic differential equations with non-globally Lipschitz continuous coefficients. *Journal of Difference Equations and Applications*, 19(3):466–490, 2013.
- [52] X. Wang, J. Wu, and B. Dong. Mean-square convergence rates of stochastic theta methods for SDEs under a coupled monotonicity condition. *BIT Numer. Math.*, 60(3):759–790, 2020.
- [53] Wang X., Gan S., Wang D. A family of fully implicit milstein methods for stiff stochastic differential equations with multiplicative noise. *BIT Numer. Math.*, 52(3):741–772, 2012.
- [54] M. Wiktorsson. Joint characteristic function and simultaneous simulation of iterated itô integrals for multiple independent brownian motions. *The Annals of Applied Probability*, 11(2):470–487, 2001.
- [55] J. Yao and S. Gan. Stability of the drift-implicit and double-implicit milstein schemes for nonlinear sdes. *Applied Mathematics and Computation*, 339:294–301, 2018.
- [56] Z. Zhang and H. Ma. Order-preserving strong schemes for SDEs with locally Lipschitz coefficients. *Appl. Numer. Math.*, 112:1–16, 2017.
- [57] X. Zong, F. Wu, and G. Xu. Convergence and stability of two classes of theta-Milstein schemes for stochastic differential equations. *Journal of Computational and Applied Mathematics*, 336:8–29, 2018.