

Causal inference methods for small non-randomized studies: Methods and an application in COVID-19

Sarah Friedrich and Tim Friede

Department of Medical Statistics, University Medical Center Göttingen, Germany*
sarah.friedrich@med.uni-goettingen.de, tim.friede@med.uni-goettingen.de

August 3, 2020

Abstract

The usual development cycles are too slow for the development of vaccines, diagnostics and treatments in pandemics such as the ongoing SARS-CoV-2 pandemic. Given the pressure in such a situation, there is a risk that findings of early clinical trials are overinterpreted despite their limitations in terms of size and design. Motivated by a non-randomized open-label study investigating the efficacy of hydroxychloroquine in patients with COVID-19, we describe in a unified fashion various alternative approaches to the analysis of non-randomized studies and apply them to the example study exploring the question whether different methods might have led to different conclusions. A widely used tool to reduce the impact of treatment-selection bias are so-called propensity score (PS) methods. Conditioning on the propensity score allows one to replicate the design of a randomized controlled trial, conditional on observed covariates. Extensions include the doubly robust g-computation, which is less frequently applied, in particular in clinical studies. Here, we investigate the properties of propensity score based methods including g-computation in small sample settings, typical for early trials, in a simulation study. We conclude that the doubly robust g-computation has some desirable properties and should be more frequently applied in clinical research. In the hydroxychloroquine study, g-computation resulted in a very wide confidence interval indicating much uncertainty. We speculate that application of the method might have prevented some of the hype surrounding hydroxychloroquine in the early stages of the SARS-CoV-2 pandemic. R code for the g-computation is provided.

Keywords: COVID-19; Causal Inference; Propensity Score; Small samples

*Humboldtallee 32, 37073 Göttingen, Germany

1 Introduction

Pandemic situations such as the currently ongoing SARS-CoV-2 pandemic require the fast development of diagnostics, vaccines and treatments. As the usual development programs are too long in these situations, more efficient development pathways are sought. These include more innovative approaches such as platform trials and adaptive designs [34]. Furthermore, in situations of desperate medical need such as with COVID-19, early clinical trials might receive more attention than they would normally do. In March 2020, for instance, [17] published a report of a small open-label non-randomized controlled study suggesting that “hydroxychloroquine treatment is significantly associated with viral load reduction/disappearance in COVID-19 patients”. Although typically not much notice would have been taken of such a small-scale study with its methodological limitations, the treatment was hailed “a game changer” by the US president putting pressure on the regulatory authorities to license the drug for COVID-19 [27].

In particular when a lot of importance is placed on non-randomized studies, their analyses and interpretation must be robust. Non-randomized studies might be prone to bias due to confounding. Besides covariate adjustments in regression models a wide range of methods were proposed to deal with this. A widely used tool to reduce the impact of treatment-selection bias in observational data are so-called propensity score (PS) methods. The propensity score is defined as a participant’s probability of receiving treatment given the observed covariates [28, 29]. Conditioning on the propensity score allows one to replicate the design of a randomized controlled trial, conditional on *observed* covariates. Extensions include the doubly robust g-computation [18, 25], which is less frequently applied, in particular in clinical studies. Doubly robust refers here to the property that it is sufficient that either the outcome or the propensity score model is correctly specified. Hence, g-computation does not rely on correct specification of both models.

[17] did not apply any of these methods for non-randomized studies, but analyzed the trial as if it was randomized. Here, we describe in a unified fashion various alternative approaches and apply them to the data by [17]. We explore the question whether different methods might have led to different conclusions. New evidence has emerged in the meanwhile and we now know that hydroxychloroquine is not an appropriate therapy in COVID-19 [12, 30]. Here, we wonder whether a more appropriate analysis of the study by [17] could have prevented much of the hype and as a result might have saved some resources.

In the context of the analysis of clinical registries and routine data including electronic health records some of the methods described above have widely been applied and their characteristics explored in simulation studies. Given the applications, simulation experiments naturally considered large-scale data sets [7]. Here, we investigate the properties of propensity score based methods including g-computation in small sample settings, typical for early trials, in a simulation study. The doubly robust g-computation has some desirable properties as the simulations will demonstrate, but has so far not gained the attention deserved in clinical applications.

The manuscript is organized as follows. In Section 2 we introduce the study by [17], which motivated our investigations, in more detail. In Section 3 several approaches to the analysis of non-randomized trials are described and applied to the motivating example. Their properties are assessed in a simulation study, in particular in the setting of small sample sizes, in Section 4. We close with a brief discussion of the findings and the limitations of our study (Section 5).

2 Motivating example in COVID-19

[17] conducted an open-label non-randomized study investigating the efficacy and safety of hydroxychloroquine in addition to standard of care in comparison to standard of care alone. The patients in the hydroxychloroquine group were all from the coordinating centre whereas the controls were recruited from several centres including the coordinating centre. In the coordinating centre, however, only those patients refusing therapy with hydroxychloroquine were included as controls. A total of 36 patients were included in the analyses, 20 patients receiving hydroxychloroquine and 16 control patients. Out of the 20 patients on hydroxychloroquine, 6 patients received in addition also azithromycin. For the purpose of illustration, we only consider two treatment groups, i.e. with and without hydroxychloroquine. The primary outcome was virological clearance at Day 6 (with Day 0 being baseline). The individual participant data of the study are reported in Supplementary Table 1 of [17]. The variables included in the table include the patient's age, sex, clinical status (asymptomatic, upper respiratory infection or lower respiratory infection), duration of symptoms, and results of daily PCR testing for Days 0 to 6. [17] report virological cure at Day 6 for 14 out of 20 patients treated with hydroxychloroquine and for 2 out of 16 in the control group, resulting in a p-value of 0.001 in an analysis not adjusted for any covariates.

The study by [17] has not been without criticism, mainly due to its limitations in design including the small sample size, choice of control patients, open label treatment and study discontinuations [1]. Although some preclinical data suggested potentially beneficial effects [13], there were also some early warnings regarding some potentially harmful effects [15]. In the meanwhile, data from large-scale randomized controlled trials are available suggesting that hydroxychloroquine is not suitable for post-exposure prophylaxis for or the treatment of COVID-19 [11, 12, 19]. The timeline of events is nicely depicted in Figure 1 of a review by [30].

3 Alternative analysis methods and their application to the data example

3.1 Notation and some causal background

We consider a binary outcome Y as well as a binary treatment A (1: experimental treatment, 0: control) and a vector of observed covariates L . In a randomized controlled trial, one would assume that due to randomization, the influence of the covariates L is the same for treated and control patients. Thus, the effect of interest in this situation would be the odds ratio

$$OR = \frac{P(Y = 1|A = 1)/P(Y = 0|A = 1)}{P(Y = 1|A = 0)/P(Y = 0|A = 0)}, \quad (3.1)$$

i. e. the ratio of the odds of having the outcome under treatment and the odds of experiencing the outcome in the control group. One way to model this is using a logistic regression model, i. e. we model the probabilities in (3.1) as

$$P(Y = 1|a) = \frac{\exp(\beta_0 + \beta_1 a)}{1 + \exp(\beta_0 + \beta_1 a)}, \quad (3.2)$$

yielding an estimator of the odds ratio $\widehat{OR} = \exp(\widehat{\beta}_1)$.

In observational studies, where allocation of the treatment is not in the hand of the investigator, this direct comparison of the treatments may no longer be fair due to the influence of other confounding factors, i.e., the distribution of the other risk factors L may differ between treated and controls. In order to imitate an RCT and to get valid estimates in this situation, one has to consider the so-called potential or counterfactual outcomes [18]: Let $Y^{a=1}$ denote the outcome that would have been observed under treatment value $a = 1$, and $Y^{a=0}$ the outcome that would have been observed under control ($a = 0$). A causal effect is now defined as follows: we say that A has a causal effect on Y if $Y^{a=1} \neq Y^{a=0}$ for an individual. In practice, however, only one of these outcomes is observed for an individual under study. Therefore, we can only ever estimate an *average causal effect*, which is present if $P(Y^{a=1} = 1) \neq P(Y^{a=0} = 1)$, i. e. the probability of the outcome under treatment is different from that under control, in the population of interest [18]. In our situation with a binary outcome and a binary treatment, we thus consider the causal odds ratio

$$OR_c := \frac{P(Y^{a=1} = 1)/(1 - P(Y^{a=1} = 1))}{P(Y^{a=0} = 1)/(1 - P(Y^{a=0} = 1))}, \quad (3.3)$$

as our primary outcome measure. Different approaches have been proposed to estimate OR_c and we will discuss the most commonly used ones in the following.

3.2 Covariate adjustment of outcome model

The conventional method to correct for baseline differences between groups is adjusting for all relevant patient characteristics in the outcome regression model. Thus, (3.1) becomes

$$OR = \frac{P(Y = 1|A = 1, L_1, \dots, L_p)/P(Y = 0|A = 1, L_1, \dots, L_p)}{P(Y = 1|A = 0, L_1, \dots, L_p)/P(Y = 0|A = 0, L_1, \dots, L_p)}$$

for a p -dimensional vector of baseline covariates $L = (L_1, \dots, L_p)$. The model for these probabilities then becomes

$$P(Y = 1|a, L) = \frac{\exp(\beta_0 + \beta_1 a + \beta_2 \ell_1 + \dots + \beta_{p+1} \ell_p)}{1 + \exp(\beta_0 + \beta_1 a + \beta_2 \ell_1 + \dots + \beta_{p+1} \ell_p)}, \quad (3.4)$$

where ℓ_1, \dots, ℓ_p denote the observed values of the covariates L_1, \dots, L_p .

In our data example, adjusting for all baseline covariates in the outcome model was not favorable due to the small number of subjects in the two groups. For comparison, we therefore included the special case of a simple logistic regression model with treatment assignment as the only covariate, i. e. model (3.2).

3.3 Propensity score based methods

Several different methods have been proposed to estimate OR_c in the literature, see e. g. [18] for a concise introduction. Many of these methods are based on the propensity score. The propensity score of individual i is defined as $\hat{p}_i := \hat{P}(A_i = 1|L_i)$, i. e., the estimated probability of receiving treatment given the covariates. For all methods considered in this paper, we estimate the propensity score using a logistic regression model for treatment allocation based on all observed covariates, i. e.

$$P(A = 1|L) = \frac{\exp(\beta_0 + \beta_1 \ell_1 + \dots + \beta_p \ell_p)}{1 + \exp(\beta_0 + \beta_1 \ell_1 + \dots + \beta_p \ell_p)}.$$

In a practical data analysis, there are several possibilities for taking the propensity score into account. We will describe the most common methods in the following and apply them to the data example.

PS covariate adjustment According to [36] and [32], covariate adjustment using the propensity score was the most commonly used PS method in clinical literature. In this approach, the outcome Y is regressed on the estimated propensity score \hat{p} and the treatment exposure A . The estimated treatment effect \widehat{OR}_c is the odds ratio for treatment exposure obtained from this logistic regression, that is we model

$$P(Y = 1|A, \hat{p}) = \frac{\exp(\beta_0 + \beta_1 a + \beta_2 \hat{p})}{1 + \exp(\beta_0 + \beta_1 a + \beta_2 \hat{p})}$$

and an estimator of the causal odds ratio is given by $\exp(\hat{\beta}_1)$.

Matching on the propensity score Another possibility to balance treatment allocation is to match subjects on the propensity score. The idea is to find individuals with a similar propensity score in the treatment and the control group. There are various methods to match individuals. Particularly in small sample studies, it is impossible in practice to find exact matches. Thus, one needs to define an acceptable difference between the propensity scores of treated individuals and controls that will be used for matching. These differences are called *calipers* and should be small enough to allow for “a practical but meaningful equation of pairs” [2]. Following the recently published recommendations by [3], who investigated propensity matching in small sample sizes, we performed a 1:1 nearest neighbor matching without replacement on the logit of the propensity score using calipers with a maximum width of 0.2 standard deviations. In this modification of classical nearest neighbor matching, subjects are only matched if the absolute difference of their propensity scores is within the pre-specified caliper distance [7]. This distance is usually defined as a proportion of the standard deviation of the logit of the propensity score. In R, this can e. g. be performed using the *MatchIt*-package, where the PS-model, the method used for matching and the caliper can be specified. A caliper of 0.2 avoids matching dissimilar individuals. Note, however, that this setting differs from the default setting in R, where the caliper is set to 0.

When it comes to analyzing the matched data set, recommendations as to whether a matched-pair analysis is required or not are not entirely clear, see e. g. [6, 9, 14, 35] for discussions of this point. Thus, we compared three different approaches to analyze the matched data set:

1. **Match unadjusted:** A logistic regression model for the outcome conditional on treatment exposure as in (3.2) was implemented. This method does not account for the matched pairs.
2. **Match conditional:** A conditional logistic regression model accounting for the matched pairs was implemented. This is achieved by the *clogit* function in R.
3. **Match GEE:** The logistic regression model was estimated using generalized estimating equations (GEE), which allows for specification of the matched pairs and estimation of robust standard errors. This approach was implemented using the *geepack*-package in R. There are different ways to specify the correlation structure, e. g. using an exchangeable correlation matrix.

Note that since we match individuals without replacement, the matched data set will usually be smaller than the original study, sometimes even discarding treated individuals.

Inverse probability of treatment weighting (IPTW) Inverse probability weighting uses the whole data set, but weighs each individual with his or her (inverse) probability of receiving the actually given treatment. This way, it generates a pseudo-population with (almost) perfect covariate balance between treatment groups. More specifically, IPTW assigns weight $w_i = 1/\hat{p}_i$ to treated subjects and weight $w_i = 1/(1 - \hat{p}_i)$ to controls. The resulting pseudo-population is analyzed using weighted logistic regression with robust standard errors obtained from the *sandwich*-package in R. Note that these standard errors could also be obtained by fitting a GEE model instead. In contrast to the GEE fitted for the matched sample above, however, we do not have clustered data here.

3.4 Doubly robust g-computation

The fourth possibility to account for covariate unbalance that we consider is *g-computation* [18, 25], also known as *the parametric g-formula* or *direct standardization*. The idea is that the marginal counterfactual risk

$$P(Y^a = 1) = \sum_{\ell} P(Y^a = 1|L = \ell)P(L = \ell) = \sum_{\ell} P(Y = 1|L = \ell, A = a)P(L = \ell).$$

Here, the sum is over all values ℓ of the confounder(s) L that occur in the population. The right-hand side of this equation can now be estimated using the available data on Y , A and L . More precisely, we have to predict the outcome for every person i in the population assuming

1. i was treated
2. i was a control

irrespective of the treatment actually received. Thus, we first fit a so-called Q-model to the data relating the outcome Y to the exposure A and to confounders L . For a binary outcome as in our situation, this is usually a logistic regression model as in (3.4). Based on the Q-model, we then predict $\hat{P}(Y = 1|L = \ell, A = 1)$ and $\hat{P}(Y = 1|L = \ell, A = 0)$ for all individuals by artificially creating two new data sets: One where $A = 1$ for all individuals and one where $A = 0$ for all individuals, respectively. The causal OR \widehat{OR}_c can then be estimated using Equation (3.3), i. e. we calculate the marginal OR as

$$\frac{\bar{p}_1/(1 - \bar{p}_1)}{\bar{p}_0/(1 - \bar{p}_0)}.$$

Here $\bar{p}_1 = \hat{P}(Y^1 = 1)$ is the mean probability of the outcome if everyone were treated and $\bar{p}_0 = \hat{P}(Y^0 = 1)$ is the mean probability of the outcome if no-one received the treatment.

While IP weighting requires the propensity model to be correct, i. e. a correct model for the treatment A conditional on confounders L , the g-formula requires a correct model for the outcome Y conditional on treatment A and the confounders L , the Q-model. A doubly robust estimator, in contrast, is consistent if at least one of the two models is correctly specified. There are many types of doubly robust estimators, but we will focus on a very simple one here [18]. First, we estimate the weights w_i as described above. We then fit our Q-model to the data including an additional covariate z , where $z_i = w_i$ if $A_i = 1$ and $z_i = -w_i$ if $A_i = 0$. Finally, we again obtain a causal OR from Equation (3.3). Confidence intervals for g-computation are usually obtained by a nonparametric bootstrap approach, i. e. by drawing with

replacement from the data and analyzing each bootstrap data set like we analyzed the original data. Here, the number of bootstrap repetitions should be chosen reasonably large. We used 10,000 repetitions in the analysis of the data example and 1,000 bootstrap repetitions in the simulation study. Upper and lower 95% confidence intervals are obtained using the 2.5 and 97.5 percentiles of the bootstrap distribution. Note that a statistical test can be obtained similarly by calculating the test statistic in each bootstrap sample and then comparing the original test statistic to the empirical $(1 - \alpha)$ -quantile of the bootstrap distribution. A p -value is obtained by counting how often the original test statistic is smaller than the bootstrap statistic and dividing this number by the conducted bootstrap replications.

3.5 Re-analysis of the COVID-19 study

We will now re-analyze the data example described in Section 2 using the methods detailed above. Following the authors' example, we imputed missing outcomes on Day 6 by a last-observation-carried-forward-approach, i. e. patients with missing PCR were considered positive on Day 6, if they were actually positive the day(s) before [17]. Moreover, after setting time since onset of symptoms to zero for asymptomatic patients, two patients with missing time since onset remained. Both patients were classified as URTI and we used the median time since onset (3 days) for URTI-patients to impute these missing values. Our PS-model included sex, age, clinical status and time since onset of symptoms as explanatory variables. The Q-model for the g-computation additionally included treatment status and the covariate z described above. We used 10,000 bootstrap iterations for the calculation of the confidence interval. The matching procedure resulted in a data set with 11 controls and 11 treated patients, thus discarding 5 controls and 9 treated patients from the analysis. Due to the small remaining sample size, the conditional logistic regression model did not converge at all, while the GEE procedure (using an exchangeable correlation structure) did not produce estimates of the standard errors and the simple model (match unadjusted) returned confidence intervals ranging from 0 to ∞ . Thus, the matching procedure did not yield useful estimates and is discarded from the results displayed below. The code for this analysis is freely available on Github (https://github.com/smn74/CIM_COVID-19).

The distribution of the propensity scores in the two groups –estimated by a kernel density estimator– is displayed in Figure 1. As we can see, the propensity scores show a good overlap between the treatment and control group. Moreover, propensity scores are reasonably far from 0 and 1 to ensure stable estimation of the IPT weights [37].

As we can see in Figure 2, all four methods (the crude unadjusted as well as the three causal inference methods) yield similar point estimates, which are, however, extremely large. Moreover, the methods differ with respect to the width of the confidence intervals and statistical significance: The very wide confidence interval obtained by the doubly robust g-computation approach includes 1 whereas the other methods yielded statistically significant treatment effects. So the question remains: Which method can we trust the most in this situation? In order to answer this, we have conducted a large simulation study which will be presented in the next section.

4 Simulation study

The set-up of our simulation study closely followed [5]. The data-generating process is as follows: First, we generate n covariates x_1, \dots, x_n . We then generate the treatment status for each subject $i = 1, \dots, N$




Figure 1: Propensity score distribution in the treatment (red curve) and control group (blue curve), respectively.

according to the model

$$\text{logit}(p_{i,\text{treatment}}) = \beta_0 + \beta_1 x_1 + \dots + \beta_n x_n. \quad (4.1)$$

Treatment is then randomly assigned to each subject following a Bernoulli distribution with subject-specific probability of treatment assignment $A_i \sim \text{Bernoulli}(p_{i,\text{treatment}})$. Next, the outcome Y_i of each subject is simulated conditional on treatment assignment A_i and the covariates associated with the outcome according to

$$\text{logit}(p_{i,\text{outcome}}) = \alpha_0 + \beta_{\text{trt}} A_i + \alpha_1 x_1 + \dots + \alpha_n x_n \quad (4.2)$$

and $Y_i \sim \text{Bernoulli}(p_{i,\text{outcome}})$. Here, β_{trt} denotes the conditional treatment effect on the log-odds scale.

Note that this data-generating process allows for introduction of a *conditional* odds ratio given the covariates. With the propensity score methods, however, we estimate a marginal odds ratio, which is in




Figure 2: Estimated Odds Ratios with 95% confidence intervals obtained by the different methods. Here, "Crude OR" refers to the simple logistic regression of treatment on the outcome, while "PS covariate" additionally includes the estimated propensity score as covariate in the logistic regression. The ORs are displayed on a logarithmic scale.

general different from the conditional OR due to non-collapsibility of the odds ratio, see e. g. [16, 26, 33]. The marginal and conditional OR coincide if there's no treatment effect, i. e., if $\beta_{trt} = 0$. For situations where the desired marginal OR is not 1, we have determined the corresponding conditional OR for a given marginal OR as described by [5]. In particular, we use Monte Carlo simulation to estimate the conditional OR. Therefore, we randomly generate 1,000 data sets of size $N = 10,000$. For each individual, we generate the counterfactual outcomes under treatment ($A_i = 1$) and control ($A_i = 0$) and calculate the marginal OR as

$$\frac{\bar{p}_1/(1 - \bar{p}_1)}{\bar{p}_0/(1 - \bar{p}_0)},$$

where \bar{p}_1 denotes the mean probability of the outcome if everyone were treated and \bar{p}_0 denotes the mean probability of the outcome if no-one received the experimental treatment, see also Equation (3.3). Using an iterative process, we modified β_{trt} until we got close enough to the desired marginal OR.

Concerning the covariates, we considered three different scenarios:

4.1 Scenario 1: The COVID-19 example

The first scenario aimed at mimicking the data example. Thus, we generated four covariates:

1. x_1 (representing sex) followed a Bernoulli distribution with parameter 0.5

2. x_2 (representing age) was drawn from a $N(45, 15)$ distribution and rounded to integers
3. x_3 (clinical status) was simulated as a categorical covariate with three categories, i. e. a $\text{Bin}(2, 0.5)$ distribution
4. x_4 (time since onset of disease) was generated from a uniform distribution on $[0, 10]$ and rounded to integers.

Treatment status was then generated according to Equation (4.1) with

$$(\beta_0, \beta_1, \beta_2, \beta_{3,1}, \beta_{3,2}, \beta_4) = (-2.3, 0.31, 0.03, 1.099, -0.1054, 0.1031).$$

Here, $\beta_{3,1}$ and $\beta_{3,2}$ correspond to the dummy-coded categories $x_{3,1}$ and $x_{3,2}$ for $x_3 = 1$ and $x_3 = 2$, respectively. The parameters were obtained from the data by univariate logistic regression. Note that this implies a moderate association of treatment with $x_1, x_{3,2}$ and x_4 , a weak association with x_2 and a strong association with $x_{3,1}$.

Similarly, the outcome was generated following Equation (4.2) with

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_{3,1}, \alpha_{3,2}, \alpha_4) = (-1.06, 0.619, 0.0077, 0.9461, -1.3499, 0.0896),$$

implying a moderate association with x_1 and x_4 , a strong association with x_3 and a weak association with x_2 . The parameter β_{trt} was varied to generate different marginal ORs in the following way: For $\beta_{trt} = 0$, marginal and conditional OR coincide and are equal to 1. For $\beta_{trt} = 0.8678$ we get a marginal OR of 2 and for $\beta_{trt} = 2.7565$ the true marginal OR equals 10. Finally, $\beta_0 = -2.3$ resulted in a similar distribution of treated individuals and controls as in the original data, yielding an average of 55.21% of individuals in the treatment group. To study the influence of more or less unbalanced treatment groups, we also varied this parameter in the simulations. In particular, we additionally considered a treatment allocation of approx. 2:1 and 4:1.

4.2 Scenario 2: Unmeasured confounder

The parameters in this setting are identical to Scenario 1, but we additionally added an unmeasured confounder. Thus, we simulated a covariate x_5 following an $N(0, 1)$ distribution with a strong effect on both treatment assignment and outcome. Therefore, β_5 and α_5 were set to $\log(5)$. However, x_5 entered neither the propensity score model nor the Q-model for the g-computation. For a marginal OR of 2 and 10, β_{trt} was set to 1.1111 and 3.4793, respectively.

4.3 Scenario 3: Following Austin's design

This scenario aims at reproducing some of the findings of [5]. Therefore, we used the same set-up as he did, namely simulating 9 binary covariates with different association to treatment assignment and outcome as described in Table 1.

Here, a strong association is represented by a coefficient of $\log(5)$, i. e. $\beta_1 = \beta_4 = \beta_7 = \alpha_1 = \alpha_2 = \alpha_3 = \log(5)$, while a moderate association has a coefficient of $\log(2)$, i.e. $\beta_2 = \beta_5 = \beta_8 = \alpha_4 = \alpha_5 = \alpha_6 = \log(2)$. We chose $\beta_0 = -3.5$ to obtain a balanced design with respect to treatment and α_0 was set

Table 1: Association to treatment assignment and outcome used in the simulation scenario motivated by [5] (Scenario 3).

	Strongly associated with treatment	Moderately associated with treatment	Not associated with treatment
Strongly associated with outcome	x_1	x_2	x_3
Moderately associated with outcome	x_4	x_5	x_6
Not associated with outcome	x_7	x_8	x_9

to -5 . For more details on the simulation set-up, see [5]. The propensity score model and the Q-model included all 9 covariates. For a marginal OR of 2 and 10, β_{trt} was set to 0.9707 and 3.2625, respectively. In addition to Austin’s setting with an equal treatment allocation of 1:1, we also considered a situation with approx. 4:1 treated patients.

An overview of all simulated scenarios is given in Table 2.

Table 2: Overview of the simulated scenarios and where to find the results.

	β_{trt}	true marginal OR	β_0	Percent treated on average	simulated for true OR
Scenario 1	0	1	-2.3	55.2%	1, 2, 10
	0.8678	2	-1.8	65.6%	1
	2.7565	10	-1	79.7%	1
Scenario 2	0	1	-2.3	53.8%	1, 2, 10
	1.1111	2			
	3.4793	10			
Scenario 3	0	1	-3.5	49.4%	1, 2, 10
	0.9707	2	-1.5	80.1%	1
	3.2625	10			
Results:	Figures 3–5, Tables 3–5		Figures 6 and 7		

4.4 Simulation results

To study the influence of small sample sizes on the methods, we simulated $N = 40, 100, 1000$ individuals for each scenario. Simulations were performed in R Version 3.6.3 with 2,000 simulation runs and the bootstrap confidence intervals for the g-computation are based on 1,000 bootstrap replications. Note that while 1,000 bootstrap replications suffice in simulations, we would recommend a higher number, say 10,000, in real-life applications.

We used different measures to compare the results. With respect to the point estimators, we considered the bias on the log-scale, i. e. the difference between the true marginal log OR and the estimated log OR. Since the methods often resulted in extreme estimators of the treatment effect, we considered both the mean bias (difference between true OR and mean estimated treatment effect) and the median bias

(difference between true OR and the median of the estimated treatment effect). The results are displayed in Figures 3 and 4, respectively. Moreover, the mean square error of each estimated marginal OR (MSE) is displayed in Tables 3–5.

Concerning the confidence intervals, we considered the percentage of 95% confidence intervals that contained the true odds ratio (coverage probability) as well as the median length of the 95% confidence interval. These measures are displayed in Figure 5 and Tables 3–5, respectively. Finally, we also reported how often the chosen models did not converge or yielded an estimated OR ≥ 3000 . These were excluded from the calculations and reported as failures in the tables.

For comparison, we again included the crude OR estimated by simple logistic regression including only treatment status as a covariate.




Figure 3: Displayed is the mean bias on the log-scale for the three scenarios (rows) and the three simulated marginal odds ratios (columns).

Across all scenarios considered here, we note that the matching procedure is the most prone to failure. Especially for the small sample sizes, this approach often fails as we have also seen in the data example. These results are in line with the findings of [3], who stress the need for development of appropriate matching methods in small sample studies. Overall, there’s not much difference between the different matching approaches. Furthermore, we found that using the default caliper, which is 0 in R, leads to




Figure 4: Displayed is the median bias on the log-scale for the three scenarios (rows) and the three simulated marginal odds ratios (columns).

extremely biased results with coverage probabilities dropping below 1% in some situations (results not shown).

As expected, the number of failures decreases with increasing sample size across scenarios. Moreover, we note that the mean bias of all methods decreases with growing sample sizes. This is, however, not the case for the median bias, see Figure 3 and 4, respectively. This finding implies that with growing sample sizes, the methods lead to less extreme results in the estimation. The largest mean bias is observed for g-computation and the unadjusted crude OR. Considering the median bias, however, the unadjusted crude OR remains positively biased through all scenarios, while g-computation does not differ much from the other methods anymore. In particular, it leads to the best results for Scenario 2.

In all of the considered scenarios, the CIs for the unadjusted crude OR greatly undercover the nominal 95% level, see Figure 5. This is especially apparent for growing sample size, which reduces the length of the CIs, thus centering them more around the wrong point estimate.

In Scenario 2, where we have an additional unobserved confounder, all methods lead to biased results even for large sample sizes and coverage probabilities are far from the 95% level, see Figure 5. In this situation, the doubly robust g-computation yields by far the best results with respect to coverage.




Figure 5: Displayed is the coverage probability (in %) for the three scenarios (rows) and the three simulated marginal odds ratios (columns).

Moreover, it also has the least median bias (Figure 4). These results are especially important, since you can never be sure to have included all relevant confounders in a practical data analysis.

For Scenario 3, we again see that the matching methods fail for $N = 40$ (Table 5). The results obtained are similar to [5], but one has to keep in mind that he simulated data sets of size $N = 10,000$. Moreover, he did not include the g-computation approach in his comparisons.

Figures 6 and 7 show the median bias and coverage probabilities of the methods in Scenario 1 and 3, respectively, where we varied the proportion of individuals who receive treatment. In Scenario 1 (Figure 6), we don't see much difference with respect to the median bias. The coverage, on the other hand, decreases with increasing imbalance for almost all methods. In Scenario 3 (Figure 7), the difference between the methods becomes more apparent with increasing imbalance. In particular, the performance of IPTW is decreasing with increasing imbalance as demonstrated by the higher median bias and the lower coverage probability (especially for smaller sample sizes), while g-computation seems to improve slightly with growing imbalance.




Figure 6: Median bias and coverage probabilities for Scenario 1 with a true OR of 1 and different proportions of treated individuals. Note that the coverage is truncated to $\geq 75\%$ implying that the unadjusted method is not displayed for $N = 1000$.

5 Discussion

In ongoing pandemics there is an urgent unmet medical need to develop vaccines, diagnostics and treatments in a very timely fashion. Despite the time pressure, however, the standards of evidence should not unduly be lowered [10, 27]. Using a small-scale non-randomized study in COVID-19 [17] as a motivating example, we demonstrate how robust analyses can be conducted by use of appropriate causal inference methods. We speculate that application of doubly-robust g-computation in the motivating study might have cast doubt on the efficacy of hydroxychloroquine and consequently, in the context of other criticism voiced, might have dampened the early enthusiasm regarding the use of this drug in COVID-19. Ultimately this might have saved some resources that now were wasted on clinical trials investigating hydroxychloroquine in patients suffering from COVID-19. More importantly, this might have prevented a shortage of hydroxychloroquine in the treatment of rheumatological disorders, which some tried to counter by revised treatment schedules to maintain the standard of care [31].

The conventional method to correct for baseline differences between groups is adjusting for all relevant patient characteristics in the outcome regression model. This is, however, not favorable for different




Figure 7: Median bias and coverage probabilities for Scenario 3 with a true OR of 1 and different proportions of treated individuals. Note that the coverage is truncated to $\geq 75\%$ implying that the unadjusted method is not displayed for some settings.

reasons. As [28] point out, covariate adjustment works poorly in cases where e. g. the variance of a covariate is unequal in the treatment and the control group. A commonly applied alternative in observational studies are propensity score methods. Since these methods were derived from a formal model for causal inference, their use allows for well-defined causal questions [23]. Moreover, propensity score methods also work as a dimension reduction tool by combining multiple covariates into a single score [23, 29]. This is especially important in situations with a large number of covariates compared to the number of subjects under study. In our data example, adjusting for all baseline covariates in the outcome model was not favorable due to the small number of subjects in the two groups. For comparison, we therefore included the special case of a simple logistic regression model with treatment assignment as the only covariate.

Some comments on the estimands obtained by the different methods are in place: First, as already explained in the set-up of the simulation study, we aimed at estimating *marginal* treatment effects here. Due to non-collapsibility of the odds ratio, these are different from *conditional* treatment effects, i. e. effects at subject level [8, 22]. Second, our aim was to estimate the average causal effect in our study population. Propensity score matching, however, creates a population where treated individuals, who cannot

be matched to any control patients, are excluded. Thus, the effect estimate obtained here corresponds to a subset of the population, which is hard to describe. Since the matched population is not very well characterized, it is difficult to generalize results obtained there to the general population [18]. Finally, it is worth noting that among the methods we discussed here, only IPT weighting and g-computation can be generalized to more complex situations involving time-varying treatments [18].

Motivated by the study conducted by [17] we investigated the properties of a range of causal inference methods in small samples. As expected this posed additional challenges to the various approaches. Interestingly, it turned out that the default settings in software implementations are often more suitable for large sample sizes and need to be adjusted for applications in small-scale studies. For example, we found that the matching procedure in R using the default calipers of 0 resulted in extremely biased results in our small sample simulations. SAS software, in contrast, uses a default caliper width of 0.25.

In our investigations, we focused here on the non-randomized nature of the study by [17]. However, the study suffers also from other weaknesses including the small sample size, open label treatment and study discontinuations [1]. For instance, we did not address the problems in the interpretation caused by study discontinuations here, but used the last-observation-carried-forward approach as [17] although this approach has gone out of fashion due to its limitations, in particular in underestimating the standard errors, see e. g. [24] and the references cited therein.

Besides the design of efficient trials to develop treatments for COVID-19 [34], one concern to trialists these days is the threat posed by the SARS-CoV-2 pandemic to clinical trials in non-COVID-19 indications [4, 21]. SARS-CoV-2 infections of patients in these trials, or merely the increased risk thereof, might lead to post-randomization events (or intercurrent events in the language of the ICH E9 addendum [20]) such as treatment or study discontinuations as well as adverse events that ultimately invalidate an analysis relying on randomization. In such situations, the causal inference approach discussed here might provide a suitable alternative analysis strategy either as primary or sensitivity analysis.

Acknowledgement

Support by the DFG (grant FR 4121/2-1) is gratefully acknowledged.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1] Paul Elias Alexander, Victoria Borg Debono, Manoj J. Mammen, Alfonso Iorio, Komal Aryala, Dianna Deng, Eva Brocard, and Waleed Alhazzani. COVID-19 coronavirus research has overall low methodological quality thus far: case in point for chloroquine/hydroxychloroquine. *Journal of Clinical Epidemiology*, 123:120–126, 2020.
- [2] Robert P Althausser and Donald Rubin. The computerized construction of a matched sample. *American Journal of Sociology*, 76(2):325–346, 1970.

- [3] Anais Andrillon, Romain Pirracchio, and Sylvie Chevret. Performance of propensity score matching to estimate causal effects in small samples. *Statistical Methods in Medical Research*, 29(3):644–658, 2020.
- [4] Stefan D Anker, Javed Butler, Muhammad Shahzeb Khan, William T Abraham, Johann Bauersachs, Edimar Bocchi, Biykem Bozkurt, Eugene Braunwald, Vijay K Chopra, John G Cleland, Justin Ezekowitz, Gerasimos Filippatos, Tim Friede, Adrian F Hernandez, Carolyn S P Lam, JoAnn Lindenfeld, John J V McMurray, Mandeep Mehra, Marco Metra, Milton Packer, Burkert Pieske, Stuart J Pocock, Piotr Ponikowski, Giuseppe M C Rosano, John R Teerlink, Hiroyuki Tsutsui, Dirk J Van Veldhuisen, Subodh Verma, Adriaan A Voors, Janet Wittes, Faiez Zannad, Jian Zhang, Petar Seferovic, and Andrew J S Coats. Conducting clinical trials in heart failure during (and after) the COVID-19 pandemic: an Expert Consensus Position Paper from the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *European Heart Journal*, 41(22):2109–2117, 2020. URL: <https://doi.org/10.1093/eurheartj/ehaa461>, arXiv:<https://academic.oup.com/eurheartj/article-pdf/41/22/2109/33368354/ehaa461.pdf>, doi:10.1093/eurheartj/ehaa461.
- [5] Peter C. Austin. The performance of different propensity score methods for estimating marginal odds ratios. *Statistics in Medicine*, 26(16):3078–3094, 2007. URL: <https://doi.org/10.1002%2Fsim.2781>, doi:10.1002/sim.2781.
- [6] Peter C Austin. A critical appraisal of propensity-score matching in the medical literature between 1996 and 2003. *Statistics in Medicine*, 27(12):2037–2049, 2008.
- [7] Peter C Austin. A comparison of 12 algorithms for matching on the propensity score. *Statistics in Medicine*, 33(6):1057–1069, 2014.
- [8] Peter C. Austin, Paul Grootendorst, Sharon-Lise T. Normand, and Geoffrey M. Anderson. Conditioning on the propensity score can result in biased estimation of common measures of treatment effect: a monte carlo study. *Statistics in Medicine*, 26(4):754–768, 2007. URL: <https://doi.org/10.1002%2Fsim.2618>, doi:10.1002/sim.2618.
- [9] Seunghee Baek, Seong Ho Park, Eugene Won, Yu Rang Park, and Hwa Jung Kim. Propensity score matching: a conceptual review for radiology researchers. *Korean Journal of Radiology*, 16(2):286–296, 2015.
- [10] Howard Bauchner and Phil B. Fontanarosa. Randomized Clinical Trials and COVID-19: Managing Expectations. *JAMA*, 323(22):2262–2263, 06 2020. URL: <https://doi.org/10.1001/jama.2020.8115>, arXiv:https://jamanetwork.com/journals/jama/articlepdf/2765696/jama_bauchner_2020_ed_200043.pdf, doi:10.1001/jama.2020.8115.
- [11] David R. Boulware, Matthew F. Pullen, Ananta S. Bangdiwala, Katelyn A. Pastick, Sarah M. Lofgren, Elizabeth C. Okafor, Caleb P. Skipper, Alanna A. Nascene, Melanie R. Nicol, Mahsa Abassi, Nicole W. Engen, Matthew P. Cheng, Derek LaBar, Sylvain A. Lothar, Lauren J. MacKenzie, Glen Drobot, Nicole Marten, Ryan Zarychanski, Lauren E. Kelly, Ilan S. Schwartz, Emily G. McDonald, Radha Rajasingham, Todd C. Lee, and Kathy H. Hullsiek. A randomized trial of hydroxychloroquine as postexposure prophylaxis for covid-19. *New England Journal of Medicine*,

2020. URL: <https://doi.org/10.1056/NEJMoa2016638>, arXiv:<https://doi.org/10.1056/NEJMoa2016638>, doi:10.1056/NEJMoa2016638.
- [12] Alexandre B. Cavalcanti, Fernando G. Zampieri, Regis G. Rosa, Luciano C.P. Azevedo, Viviane C. Veiga, Alvaro Avezum, Lucas P. Damiani, Aline Marcadenti, Leticia Kawano-Dourado, Thiago Lisboa, Debora L. M. Junqueira, Pedro G.M. de Barros e Silva, Lucas Tramujas, Erlon O. Abreu-Silva, Ligia N. Laranjeira, Aline T. Soares, Leandro S. Echenique, Adriano J. Pereira, Flavio G.R. Freitas, Otavio C.E. Gebara, Vicente C.S. Dantas, Remo H.M. Furtado, Eveline P. Milan, Nicole A. Golin, Fabio F. Cardoso, Israel S. Maia, Conrado R. Hoffmann Filho, Adrian P.M. Kormann, Roberto B. Amazonas, Monalisa F. Bocchi de Oliveira, Ary Serpa-Neto, Maicon Falavigna, Renato D. Lopes, Flavia R. Machado, and Otavio Berwanger. Hydroxychloroquine with or without azithromycin in mild-to-moderate covid-19. *New England Journal of Medicine*, 2020. URL: <https://doi.org/10.1056/NEJMoa2019014>, arXiv:<https://doi.org/10.1056/NEJMoa2019014>, doi:10.1056/NEJMoa2019014.
- [13] Andrea Cortegiani, Giulia Ingoglia, Mariachiara Ippolito, Antonino Giarratano, and Sharon Einav. A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19. *Journal of Critical Care*, 57:279–283, 2020.
- [14] Markus C Elze, John Gregson, Usman Baber, Elizabeth Williamson, Samantha Sartori, Roxana Mehran, Melissa Nichols, Gregg W Stone, and Stuart J Pocock. Comparison of propensity score methods and covariate adjustment: evaluation in 4 cardiovascular studies. *Journal of the American College of Cardiology*, 69(3):345–357, 2017.
- [15] Christian Funck-Brentano, Lee S Nguyen, and Joe-Elie Salem. Retraction and republication: cardiac toxicity of hydroxychloroquine in COVID-19. *The Lancet*, 2020. doi:10.1016/S0140-6736(20)31528-2.
- [16] Mitchell H Gail, S Wieand, and Steven Piantadosi. Biased estimates of treatment effect in randomized experiments with nonlinear regressions and omitted covariates. *Biometrika*, 71(3):431–444, 1984.
- [17] Philippe Gautret, Jean-Christophe Lagier, Philippe Parola, Line Meddeb, Morgane Mailhe, Barbara Doudier, Johan Courjon, Valérie Giordanengo, Vera Esteves Vieira, Hervé Tissot Dupont, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International Journal of Antimicrobial Agents*, 56(1), 2020. doi:<https://doi.org/10.1016/j.ijantimicag.2020.105949>.
- [18] Miguel Hernan and James Robins. *Causal Inference: What If*. Chapman & Hall/CRC, Boca Raton, 2020.
- [19] Peter Horby, Marion Mafham, Louise Linsell, Jennifer L Bell, Natalie Staplin, Jonathan R Emberson, Martin Wiselka, Andrew Ustianowski, Einas Elmahi, Benjamin Prudon, Anthony Whitehouse, Timothy Felton, John Williams, Jakki Faccenda, Jonathan Underwood, J Kenneth Baillie, Lucy Chappell, Saul N Faust, Thomas Jaki, Katie Jeffery, Wei Shen Lim, Alan Montgomery, Kathryn Rowan, Joel Tarning, James A Watson, Nicholas J White, Edmund Juszczak, Richard Haynes, and Martin J Landray. Effect of hydroxychloroquine in hospitalized patients with covid-19: Preliminary results from a multi-centre, randomized, controlled trial. *medRxiv*, 2020. URL: <https://doi.org/10.1101/2020.05.06.20091729>.

//www.medrxiv.org/content/early/2020/07/15/2020.07.15.20151852,
arXiv:https://www.medrxiv.org/content/early/2020/07/15/2020.07.
15.20151852.full.pdf, doi:10.1101/2020.07.15.20151852.

- [20] ICH. ICH E9 (R1) addendum on estimands and sensitivity analysis in clinical trials to the guideline on statistical principles for clinical trials. 2019. URL: <https://www.ema.europa.eu/en/ich-e9-statistical-principles-clinical-trials>.
- [21] Cornelia Ursula Kunz, Silke Jörgens, Frank Bretz, Nigel Stallard, Kelly Van Lancker, Dong Xi, Sarah Zohar, Christoph Gerlinger, and Tim Friede. Clinical trials impacted by the covid-19 pandemic: Adaptive designs to the rescue? *Statistics in Biopharmaceutical Research*, 2020. URL: <https://doi.org/10.1080/19466315.2020.1799857>, arXiv:<https://doi.org/10.1080/19466315.2020.1799857>, doi: 10.1080/19466315.2020.1799857.
- [22] Huzhang Mao and Liang Li. Flexible regression approach to propensity score analysis and its relationship with matching and weighting. *Statistics in Medicine*, 2020. doi:<https://doi.org/10.1002/sim.8526>.
- [23] Daniel F McCaffrey, Beth Ann Griffin, Daniel Almirall, Mary Ellen Slaughter, Rajeev Ramchand, and Lane F Burgette. A tutorial on propensity score estimation for multiple treatments using generalized boosted models. *Statistics in Medicine*, 32(19):3388–3414, 2013.
- [24] Geert Molenberghs, Herbert Thijs, Ivy Jansen, Caroline Beunckens, Michael G Kenward, Craig Mallinckrodt, and Raymond J Carroll. Analyzing incomplete longitudinal clinical trial data. *Biostatistics*, 5(3):445–464, 2004.
- [25] James Robins. A new approach to causal inference in mortality studies with a sustained exposure period – application to control of the healthy worker survivor effect. *Mathematical Modelling*, 7(9-12):1393–1512, 1986.
- [26] Laurence D Robinson and Nicholas P Jewell. Some surprising results about covariate adjustment in logistic regression models. *International Statistical Review/Revue Internationale de Statistique*, pages 227–240, 1991.
- [27] Bejamnin N. Rome and Jerry Avorn. Drug evaluation during the covid-19 pandemic. *New England Journal of Medicine*, 382(24):2282–2284, 2020.
- [28] Paul R Rosenbaum and Donald B Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55, 1983.
- [29] Paul R Rosenbaum and Donald B Rubin. Reducing bias in observational studies using subclassification on the propensity score. *Journal of the American Statistical Association*, 79(387):516–524, 1984.
- [30] Sebastian E. Sattui, Jean W. Liew, Elizabeth R. Graef, Ariella Coler-Reilly, Francis Berenbaum, Ali Duarte-Garcia, Carly Harrison, Maximilian F. Konig, Peter Korsten, Michael S. Putman, Philip C. Robinson, Emily Sirotych, Manuel F. Ugarte-Gil, Kate Webb, Kristen J. Young, Alfred H.J. Kim, and Jeffrey A. Sparks. Swinging the pendulum: lessons learned from public discourse concerning

- hydroxychloroquine and COVID-19. *Expert Review of Clinical Immunology*, 2020. doi:<https://doi.org/10.1080/1744666X.2020.1792778>.
- [31] Marc H. Scheetz, Maximilian F. Konig, Philip C. Robinson, Jeffrey A. Sparks, and Alfred H.J. Kim. A Pharmacokinetics-Informed Approach to Navigating Hydroxychloroquine Shortages in Patients With Rheumatic Disease During the COVID-19 Pandemic. *ACR Open Rheumatology*, 2020. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/acr2.11164>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1002/acr2.11164>, doi:10.1002/acr2.11164.
- [32] Bajju R. Shah, Andreas Laupacis, Janet E. Hux, and Peter C. Austin. Propensity score methods gave similar results to traditional regression modeling in observational studies: a systematic review. *Journal of Clinical Epidemiology*, 58(6):550–559, jun 2005. URL: <https://doi.org/10.1016%2Fj.jclinepi.2004.10.016>, doi:10.1016/j.jclinepi.2004.10.016.
- [33] Arvid Sjölander, Elisabeth Dahlqwist, and Johan Zetterqvist. A note on the noncollapsibility of rate differences and rate ratios. *Epidemiology*, 27(3):356–359, 2016.
- [34] Nigel Stallard, Lisa Hampson, Norbert Benda, Werner Brannath, Thomas Burnett, Tim Friede, Peter K. Kimani, Franz Koenig, Johannes Krisam, Pavel Mozgunov, Martin Posch, James Wason, Gernot Wassmer, John Whitehead, S. Faye Williamson, Sarah Zohar, and Thomas Jaki. Efficient adaptive designs for clinical trials of interventions for COVID-19. *Statistics in Biopharmaceutical Research*, 2020. arXiv:arXiv:2005.13309v1.
- [35] Elizabeth A. Stuart. Developing practical recommendations for the use of propensity scores: Discussion of 'A critical appraisal of propensity score matching in the medical literature between 1996 and 2003' by Peter Austin, *Statistics in Medicine*. *Statistics in Medicine*, 27(12):2062–2065, 2008. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/sim.3207>, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1002/sim.3207>, doi:10.1002/sim.3207.
- [36] Sherry Weitzen, Kate L Lapane, Alicia Y Toledano, Anne L Hume, and Vincent Mor. Principles for modeling propensity scores in medical research: a systematic literature review. *Pharmacoepidemiology and Drug Safety*, 13(12):841–853, 2004.
- [37] Yunji Zhou, Roland A Matsouaka, and Laine Thomas. Propensity score weighting under limited overlap and model misspecification. *Statistical Methods in Medical Research*, 2020. PMID: 32693715. URL: <https://doi.org/10.1177/0962280220940334>, arXiv:<https://doi.org/10.1177/0962280220940334>, doi:10.1177/0962280220940334.

Table 3: Median length of the 95% confidence intervals (Length CI), mean square error of the estimated treatment effect (MSE) and number of models (out of 2000 simulation runs) that did not converge or resulted in an OR > 3000 for Scenario 1: COVID-19. Crude OR refers to the simple logistic regression adjusting only for treatment assignment, PS covariate denotes the method including the PS in the outcome regression model and g-comp refers to the doubly robust g-computation method.

True OR	N	Crude OR	PS Covariate	IPTW	g-comp	Match unadjusted	Match conditional	Match GEE	
1	N = 40	CI Length	5.912	4.519	4.195	28.36	6.958	6.169	
		MSE	6.982	2.333	4.124	82.07	4.003	1.404	3.571
		failures	2	2	2	1	23	358	107
	N = 100	CI Length	3.05	2.116	2.044	15.54	2.542	2.737	2.498
		MSE	1.387	0.276	0.2825	26.6	0.4405	0.9441	17.35
		failures	0	0	0	0	0	1	51
	N = 1000	CI Length	0.8548	0.5757	0.5546	4.016	0.6576	0.7004	0.6597
		MSE	0.5304	0.01933	0.01553	1.421	0.03618	0.04705	0.04198
		failures	0	0	0	0	0	0	8
2	N = 40	CI Length	14.14	11.1	10.33	75.63	15.2	12.4	13.77
		MSE	37.82	45.31	30.76	3719	21.03	2.049	45.81
		failures	4	4	4	3	87	542	192
	N = 100	CI Length	6.713	4.863	4.55	31.41	5.572	6.613	5.589
		MSE	7.117	1.706	1.352	104.6	2.642	6.326	2.637
		failures	0	0	0	0	0	25	58
	N = 1000	CI Length	1.863	1.299	1.184	8.301	1.391	1.681	1.396
		MSE	2.344	0.1502	0.07487	5.786	0.1741	0.4285	0.1762
		failures	0	0	0	0	0	0	1
10	N = 40	CI Length	113	91.59	77.36	8.8 · 10 ¹¹	62.14	35.34	51.42
		MSE	459.1	3052	705.6	22242	55.8	42.9	53.79
		failures	521	522	521	373	946	1441	1021
	N = 100	CI Length	62.98	46.12	40.95	23153036	48.55	76.84	47.83
		MSE	602.8	306.7	303	6854	127.1	19.97	127.4
		failures	74	74	74	69	191	720	247
	N = 1000	CI Length	14.42	10.8	8.993	54.91	10.52	17.99	10.58
		MSE	81.08	17.56	5.278	253.8	10.76	65.29	10.86
		failures	0	0	0	0	0	0	1

Table 4: Median length of the 95% confidence intervals (Length CI), mean square error of the estimated treatment effect (MSE) and number of models (out of 2000 simulation runs) that did not converge or resulted in an OR > 3000 for Scenario 2: Unmeasured Confounder. Crude OR refers to the simple logistic regression adjusting only for treatment assignment, PS covariate denotes the method including the PS in the outcome regression model and g-comp refers to the doubly robust g-computation method.

True OR	N	Crude OR	PS Covariate	IPTW	g-comp	Match unadjusted	Match conditional	Match GEE	
1	N = 40	CI Length	15.93	16.42	15.98	80.72	23.17	21.88	22.11
		MSE	101.9	307.6	127.4	653.2	88.55	8.701	828.8
		failures	6	6	6	6	72	558	201
	N = 100	CI Length	8.405	7.646	7.422	39.2	8.648	10.49	8.88
		MSE	19.54	13.68	12.29	205.8	16.83	26.12	4730
		failures	0	0	0	0	0	46	103
	N = 1000	CI Length	2.285	1.991	1.924	18.84	2.125	2.553	2.162
		MSE	11.21	7.146	6.238	43.79	7.34	8.707	289.5
		failures	0	0	0	0	0	0	48
2	N = 40	CI Length	45.63	46.45	43.65	350.8	56.03	35.34	54.46
		MSE	586.3	1581	805.5	4492	244.4	10.28	1158
		failures	39	40	39	27	285	975	425
	N = 100	CI Length	21.89	20.42	19.32	101.2	24.4	32.95	24.42
		MSE	159.5	124.6	104.4	1283	170.7	94.92	177.4
		failures	0	0	0	0	2	265	56
	N = 1000	CI Length	5.714	5.066	4.754	42.14	5.294	7.534	5.295
		MSE	60.46	41.71	33.66	224.3	39.54	54.12	39.67
		failures	0	0	0	0	0	0	17
10	N = 40	CI Length	282.1	290.8	262.3	2.2 · 10 ¹²	159.5	49.11	136.1
		MSE	2646	11888	4497	42624	333.5	22.25	5111
		failures	1075	1077	1075	709	1439	1784	1482
	N = 100	CI Length	297.4	283.5	267.4	7.2 · 10 ⁹	328.1	146.3	309.2
		MSE	5672	5054	4800	52680	2192	101.4	2150
		failures	404	404	404	370	658	1385	690
	N = 1000	CI Length	75.46	67.68	61.92	352.3	69.42	146.3	69.26
		MSE	4282	3285	2450	16564	2960	7944	2964
		failures	0	0	0	0	0	24	1

Table 5: Median length of the 95% confidence intervals (Length CI), mean square error of the estimated treatment effect (MSE) and number of models (out of 2000 simulation runs) that did not converge or resulted in an OR > 3000 for Scenario 3: Austin (2007). Crude OR refers to the simple logistic regression adjusting only for treatment assignment, PS covariate denotes the method including the PS in the outcome regression model and g-comp refers to the doubly robust g-computation method.

True OR	N	Crude OR	PS Covariate	IPTW	g-comp	Match unadjusted	Match conditional	Match GEE	
1	N = 40	CI Length	9.028	9.211	7.77	52.76	.	.	
		MSE	13.62	207.3	17.14	216.9	.	.	
		failures	56	66	56	29	2000	2000	2000
	N = 100	CI Length	4.504	3.022	3.23	16.76	4.046	3.967	4.053
		MSE	3.756	0.6415	1.654	19.76	1.955	1.621	1.976
		failures	0	0	0	0	10	76	92
N = 1000	CI Length	1.218	0.7266	0.8238	1.462	0.8583	0.876	0.861	
	MSE	1.259	0.02995	0.03788	0.1572	0.05978	0.06768	0.0809	
	failures	0	0	0	0	0	0	48	
2	N = 40	CI Length	18.26	17.15	16.41	89.43	.	.	
		MSE	64.2	4782	102.1	631.8	.	.	
		failures	56	72	56	24	2000	2000	2000
	N = 100	CI Length	8.968	6.018	6.701	27.4	7.804	8.42	7.82
		MSE	15.24	2.453	6.258	68.3	7.68	5.197	7.665
		failures	0	0	0	0	6	93	84
N = 1000	CI Length	2.367	1.446	1.625	2.925	1.656	1.817	1.657	
	MSE	5.065	0.137	0.1431	0.5701	0.2294	0.378	0.2411	
	failures	0	0	0	0	0	0	7	
10	N = 40	CI Length	150.2	133.7	133.8	9.3·10 ¹¹	.	.	
		MSE	2845	15204	6961	17302	.	.	
		failures	125	177	125	57	2000	2000	2000
	N = 100	CI Length	59.41	40.1	44.18	178.1	50.5	69.9	50.29
		MSE	1249	389.1	724.6	2050	425.9	20.47	560.7
		failures	1	1	1	1	14	700	155
N = 1000	CI Length	14.77	9.183	9.804	16.95	9.786	16.39	9.934	
	MSE	173.2	13.63	5.445	19.68	9.661	49.29	16.21	
	failures	0	0	0	0	0	0	66	