

# Refining constraints from Borexino measurements on a light $Z'$ -boson coupled to $L_\mu$ - $L_\tau$ current

Sergei Gninenko<sup>a</sup>, Dmitry Gorbunov<sup>a,b</sup>

<sup>a</sup>*Institute for Nuclear Research of Russian Academy of Sciences, 117312 Moscow, Russia*

<sup>b</sup>*Moscow Institute of Physics and Technology, 141700 Dolgoprudny, Russia*

---

## Abstract

The recent confirmation by FNAL of the  $(g-2)_\mu$  muon anomaly gives strong evidence for the possible existence of new physics beyond the Standard Model in the muon sector. Thus it is worthy to revisit the existing experimental constraints on models suggesting theoretically consistent explanations of the anomaly. In this work, we point out that accounting for the loss of coherence between the wave packets (mass states) of solar neutrinos is important for setting limits on any model with new flavor-sensitive couplings in the neutrino sector. By taking into account this effect and considering more accurately the experimental constraints from the BOREXINO measurement of the  $^7\text{Be}$  solar neutrino interaction rate we corrected the limits previously placed on the coupling of the light  $Z'$  to  $L_\mu$ - $L_\tau$  current in the parameter space relevant to the muon  $(g-2)_\mu$ .

---

1. A longstanding problem in particle physics naturally stimulates theorists to invent solutions beyond the Standard Model of particle physics (SM). Discrepancies between measurements and SM predictions can indeed be because of the incompleteness of the SM (though, overlooked experimental issues and problems with theoretical calculations are not excluded generically) and indicate explicit effects of new physics. A well known example is a tension in the muon anomalous magnetic moment,  $a_\mu = (g-2)_\mu/2$ , see, e.g. [1], inspiring this activity for two decades. The recent measurement of the  $a_\mu$  value by the E989 experiment at Fermilab shows a discrepancy with respect to the theoretical prediction of the SM [2] which when combined with the previous Brookhaven results [3] leads to a  $4.2\sigma$  tension of  $\Delta a_\mu = 251(59) \times 10^{-11}$ .

One of the possible solutions of  $(g-2)_\mu$  puzzle is a new massive vector boson  $Z'$  coupled to non-anomalous difference of muon and tau lepton currents,  $L_\mu$ - $L_\tau$  [4]-[9],

$$\mathcal{L} = g_{Z'} Z'_\rho (L_\mu^\rho - L_\tau^\rho) \equiv g_{Z'} Z'_\rho (\bar{\tau} \gamma^\rho \tau + \bar{\nu}_\tau \gamma^\rho \nu_\tau - \bar{\mu} \gamma^\rho \mu - \bar{\nu}_\mu \gamma^\rho \nu_\mu) . \quad (1)$$

---

*Email addresses:* Sergei.Gninenko@cern.ch (Sergei Gninenko), gorbys@ms2.inr.ac.ru (Dmitry Gorbunov)

The electrophobic nature of  $Z'$  prevents it from contributing to the similar  $(g-2)_e$  observable of electron, which otherwise would clash with measurements. While limited by physical constraints from many experiments in particle physics and astrophysics revealing no anomalies,  $Z'$  boson can play a role of messenger of the dark matter sector, which makes the model interesting in cosmology and deserving to be tested in a wider range of model parameters [10]–[16]. In particular, the promising for  $(g-2)_\mu$  part of model parameter space can be fully explored by the NA64 experiment operating at the CERN SPS [10–12].

**2.** One important limit [15] comes from measurements of  $^7\text{Be}$  solar neutrino interaction rate inside the BOREXINO detector [17]: virtual intermediate  $Z'$  boson may contribute to  $\nu_\mu - e$  and  $\nu_\tau - e$  scatterings off electrons in the detector volume. The  $Z'$  coupling to fermions of the first generation emerge due to the  $\gamma - Z'$  kinetic mixing, which is generated through the loop involving the muon and tau leptons [12, 15, 16]. Consistency of the measured rate with SM predictions has allowed for imposing limits from above on the magnitude of  $Z'$  boson coupling  $g_{Z'}$  to  $L_\mu - L_\tau$ .

However, in doing it, the authors, e.g., of Ref. [15] have implicitly assumed that  $^7\text{Be}$  solar neutrinos, being originally pure electron neutrinos  $\nu_e$ , oscillate on their way to Earth and arrive there as equipartition combination of pure flavor states  $\nu_e$ ,  $\nu_\mu$  and  $\nu_\tau$ , so that each of the neutrino flavor *interacts coherently* with matter of BOREXINO detector. The interaction rate then is given by

$$\Gamma \propto F_{\nu_\mu} + F_{\nu_\tau} = \frac{2}{3} F_{^7\text{Be}}, \quad (2)$$

where  $F_{\nu_\alpha}$  are flavor fluxes and  $F_{^7\text{Be}}$  is electron neutrino flux from  $^7\text{Be}$  to be measured in absence of neutrino oscillations.

In fact, that is not the case, since the neutrino wave packets lose the coherence on their way to Earth [18] and arrive as pure mass states  $\nu_i$ ,  $i = 1, 2, 3$  with fluxes

$$F_i = |U_{ei}|^2 F_{^7\text{Be}}, \quad (3)$$

where  $U_{\alpha i}$  are elements of the PMNS matrix [18]. Each of the mass state is a known combination of flavor states,  $\nu_i = \sum_\alpha U_{\alpha i} \nu_\alpha$ , which hence couple coherently when exchanging  $Z'$  with matter. Since  $\nu_e$  does not couple to  $Z'$  boson (except through the mentioned mixing with  $Z$  boson which is small) and  $\nu_\mu$  and  $\nu_\tau$  coupling constants are equal in magnitude but of opposite signs, the interaction of the neutrino mass states with  $Z'$  following from (1) upon rotation of the neutrino states reads

$$\mathcal{L} = g_{Z'} Z_\rho \bar{\nu}_i \gamma^\rho \nu_j (U_{\tau i}^* U_{\tau j} - U_{\mu i}^* U_{\mu j}). \quad (4)$$

Hence, the expected impact of  $Z'$  on the  $^7\text{Be}$  solar neutrino interaction rate in BOREXINO should be summed over all the three initial mass states and all the three final neutrino mass

states, which reveals

$$\begin{aligned}
\Gamma &\propto \sum_i F_i \times \sum_j |U_{\mu i}^* U_{\mu j} - U_{\tau i}^* U_{\tau j}|^2 = F_{7\text{Be}} \times \sum_i |U_{ei}|^2 \times \sum_j |U_{\mu i}^* U_{\mu j} - U_{\tau i}^* U_{\tau j}|^2 \\
&= F_{7\text{Be}} \times \sum_i |U_{ei}|^2 \times (|U_{\mu i}|^2 + |U_{\tau i}|^2) = F_{7\text{Be}} \times \sum_i |U_{ei}|^2 \times (1 - |U_{ei}|^2) \\
&= F_{7\text{Be}} \times \left(1 - \sum_i |U_{ei}|^4\right),
\end{aligned} \tag{5}$$

where we simplify the formula by making use of the unitarity of the PMNS matrix. Instead of (2), it gives numerically

$$\approx 0.45 \times F_{7\text{Be}}$$

for the central values of the neutrino mixing parameters taken from [18].

One observes, that the combination of the PMNS matrix elements emerging in the results (5) is equal to the electron survival probability  $P_{ee} = \sum_i |U_{ei}|^4$  in case of pure vacuum oscillations. Recall, that  ${}^7\text{Be}$  solar neutrino flux is expected to be a bit suppressed by scattering off electrons inside the Sun: neutrino mixing angles in matter is somewhat different for  ${}^7\text{Be}$  neutrino energy as compared to pure vacuum case used for the numerical estimate above. However the theoretical expectation and observations by BOREXINO [17] are both consistent within the errors with pure vacuum case, which we adopt here to be conservative in placing limits on the new physics.

Note in passing that the same formulas (5) can be obtained by considering the neutrino interaction in the flavor basis. There the scattering of muon and tau neutrinos from a particular wave packet (pure mass state) must be sum up incoherently, which gives

$$\Gamma \propto \sum_i F_i \times (|U_{\mu i}|^2 + |U_{\tau i}|^2)$$

coinciding with (5) for the fluxes (3). Consequently, the limits of  $Z'$  boson couplings  $g_{Z'}$  from Ref.[15] must be weakened by a factor  $\approx (0.66/0.45)^{1/4} \approx 1.11$ .

**3.** The BOREXINO limit [17] has been applied in Ref. [14] in a different way to infer constraints on the  $L_\mu - L_\tau$  model parameters. The authors took the central value of the measured by BOREXINO survival probability for  ${}^7\text{Be}$  solar neutrinos,  $P_{ee} = 0.51$  (so the  $\nu_\mu$  and  $\nu_\tau$  flux is  $0.49 \times F_{7\text{Be}}$ ) and stated that the new physics contribution should not exceed 8% [14]. This estimate should be also corrected. First, for the best fit values of the neutrino sector parameters and Standard Solar Model (SSM) the theoretical evaluation of the electron survival probability is somewhat higher than the measured number, resulting in the the  $\nu_\mu$  and  $\nu_\tau$  flux of  $0.47 \times F_{7\text{Be}}$ , see Fig. 2 in Ref. [17], which is still consistent with pure vacuum oscillations as mentioned above. Second, the BOREXINO estimate for the survival  $\nu_e$  probability including both theoretical SSM and experimental uncertainties,  $P_{ee} = 0.51 \pm 0.07$ , has the total error of 14% [17]. This bigger error accounts also for the uncertainties in the flux prediction associated with the Standard Solar Model, which apparently have been ignored in Ref. [14]. Hence, to place a limit on the possible  $L_\mu - L_\tau$   $Z'$

contribution, one should adopt that is should not exceed 23% at 90% C.L.. Consequently, the estimate in Ref. [14] on  $g'$  coupling has to be corrected by a factor

$$\left(\frac{0.23}{0.08} \times \frac{0.49}{0.47}\right)^{1/4} = 1.32. \quad (6)$$

This results in a new 90% C.L. exclusion region shown in the  $(m_{Z'}, g_{Z'})$  plane in Fig. 1,




Figure 1: Shown are the 90% C.L. exclusion regions (shaded areas above the curves) from the analysis of BOREXINO data performed in [14] and in this work (light grey area) in the  $(m_{Z'}, g_{Z'})$  plane together with the favourable  $(g - 2)_\mu$  parameter space in the  $L_\mu - L_\tau$  model. The 90% C.L. expected exclusion regions (dashed curves) from the measurements with the electron beam with  $\simeq 10^{13}$  EOT [12, 19] and muons with  $\simeq 10^{12}$  MOT [10, 11] in the NA64 experiment at CERN, as well as constraints from the CCFR [20], and BABAR [21] experiments, and the region of parameters explaining the XENON1T excess [22] are also shown. Two symbols + and x indicate parameters  $(m_{Z'}, g_{Z'}) = (11 \text{ MeV}, 5 \times 10^{-4})$  and  $(9 \text{ MeV}, 4 \times 10^{-4})$ , respectively, which are used to explain the characteristic features of cosmic neutrino spectrum reported by the IceCube Collaboration, see Ref. [16] for details.

together with the BOREXINO limit from Ref. [14] and the favourable  $(g - 2)_\mu$  parameter space of the  $L_\mu - L_\tau$  model. The existing constraints from the CCFR [20] and BABAR [21] experiments, as well as expected limits from the measurements with the electron beam for  $\simeq 4 \cdot 10^{13}$  electrons on target (EOT) [19] and the muon beam for  $\simeq 10^{12}$  muons on target (MOT) [10] with the NA64 experiment at CERN are also shown. Interestingly, in a specifically extended model there is a region in the  $(m_{Z'}, g_{Z'})$  plane which could simultaneously explain the  $(g - 2)_\mu$  and an excess of low energy electrons recently observed by the XENON1T experiment [22]. It is worth mentioning that particular variants of the model with light  $Z'$ -bosons are generically constrained from cosmology, e.g. due to possible impact of light  $Z'$ -bosons on the Big Bang Nucleosynthesis and Cosmic Microwave Background

anisotropies [14, 15, 23]. Nevertheless, this observation makes searches for  $Z'$  with NA64 quite interesting, as the experiment can probe the parameter space shown in Fig.1 and provide important complementary results in the near future [11]. As shown in Fig.1 with about  $\sim 10^{13}$  EOT NA64e will probe the area of parameters explaining both, the  $(g-2)_\mu$  anomaly and the XENON1t excess. Interestingly, it will also probe both reference points of the model suggested in Ref. [16] for explaining characteristic features of cosmic neutrino spectrum observed by IceCube, which are also shown in Fig. 1.

To conclude, from the theoretical point of view we expect the effect of losing the coherence is particularly important for the phenomenology of any model of new physics with non-universal interactions with neutrino flavors: e.g. in models with neutrino dipole moments different in values for electron, muon and tau neutrinos. While from the experiment view point, given the significance of the  $(g-2)_\mu$  anomaly for a possible discovery of new physics, correction (6) enlarges the  $(m_{Z'}, g_{Z'})$  parameter space for the  $L_\mu - L_\tau$   $Z'$  explanation of the muon  $(g-2)_\mu$  anomaly, and at the same time increases motivation for searching the  $Z'$  in NA64e and NA64 $_\mu$  experiments at CERN [10–12] and M<sup>3</sup> [13] at Fermilab, in particular in the parameter space, which could explain the  $(g-2)_\mu$  anomaly and the XENON1T excess simultaneously. This makes these searches quite exciting. Despite of the fact that BOREXINO bounds have already probed indirectly the scenario under consideration, it would be of great importance to perform in these experiments independent direct searches for the  $Z'$ -boson in the mass range  $m_{Z'} \lesssim 10$  MeV. An estimate shows that, e.g., the NA64e constraints on the couplings  $g_{Z'}$  in this mass range, which can be obtained from the analysis of data collected for the invisible decays of dark photons [12, 19], are comparable with those presented above.

We thank Mikhail Shaposhnikov for useful discussions. The estimate of the interaction rate at BOREXINO was supported by the RSF Grant No. 17-12-01547. The work on NA64 was supported in part by the Ministry of Science and Higher Education of Russia, the Agreement No 05.613.21.0098 ID No RFMEFI61320X0098 on July 23, 2020.

## References

- [1] T. Aoyama et al., "The anomalous magnetic moment of the muon in the Standard Model", arXiv:2006.04822 [hep-ph].
- [2] B. Abi et al. (Muon g-2), "Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm", Phys. Rev. Lett. **126**, 141801 (2021), 2104.03281, arXiv: 2104.03281 [hep-ex].
- [3] G.W. Bennett et al. (Muon g-2 Collaboration), "Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL", Phys. Rev. D **73**, 072003 (2006), hep-ex/0602035 [hep-ex].
- [4] X. He, G. C. Joshi, H. Lew, and R. Volkas, "New Z-prime phenomenology", Phys. Rev. D **43**, (1991) 22.
- [5] R. Foot, "New Physics From Electric Charge Quantization?", Mod. Phys. Lett. A **6** (1991) 527.
- [6] X.-G. He, G. C. Joshi, H. Lew, and R. Volkas, "Simplest Z-prime model", Phys. Rev. D **44** (1991) 2118.
- [7] S.N. Gninenko and N.V. Krasnikov, "The Muon anomalous magnetic moment and a new light gauge boson", Phys. Lett. B **513** (2001) 119, hep-ph/0102222 [hep-ph].
- [8] S. Baek, N.G. Deshpande, X.G. He, "Muon anomalous g-2 and gauged  $L_\mu - L_\tau$  models" Phys. Rev. D **64** (2001) 055006, hep-ph/0104141 [hep-ph]

- [9] E. Ma, D. Roy and S. Roy, "Gauged  $L_\mu - L_\tau$  with large muon anomalous magnetic moment and the bimaximal mixing of neutrinos", Phys. Lett. **B 525** (2002) 101, hep-ph/0110146 [hep-ph].
- [10] S.N. Gninenko, N.V. Krasnikov, V.A. Matveev, "Muon  $g-2$  and searches for a new leptophobic sub-GeV dark boson in a missing-energy experiment at CERN", Phys. Rev. D **91** (2015) 095015, arXiv: 1412.1400 [hep-ph].
- [11] D. Banerjee et al. (NA64 Collaboration), "Proposal for an experiment to search for dark sector particles weakly coupled to muon at the CERN SPS", CERN-SPSC-2019-002/SPSC-P-359 (CERN, Geneva, 2019).
- [12] S.N. Gninenko and N.V. Krasnikov, "Probing the muon  $g_\mu-2$  anomaly,  $L_\mu - L_\tau$  gauge boson and Dark Matter in dark photon experiments", Phys. Lett. **B 783** (2018) 24, arXiv:1801.10448 [hep-ph].
- [13] Y. Kahn, G. Krnjaic, N. Tran, A. Whitbeck, " $M^3$ : a new muon missing momentum experiment to probe  $(g-2)_\mu$  and dark matter at Fermilab", JHEP **09** (2018) 153; arXiv: 1804.03144 [hep-ph].
- [14] A. Kamada, K. Kaneta, K. Yanagi, H-B Yu, "Self-interacting dark matter and muon  $g-2$  in a gauged  $U(1)_{L_\mu-L_\tau}$  model", JHEP **06** (2018) 117, arXiv:1805.00651 [hep-ph].
- [15] A. Kamada and H-B Yu, "Coherent propagation of PeV neutrinos and the dip in the neutrino spectrum at IceCube" Phys. Rev. **D 92** (2015) 113004, arXiv:1504.00711 [hep-ph].
- [16] T. Araki, F. Kaneko, T. Ota, J. Sato, and T. Shimomura, "MeV scale leptonic force for cosmic neutrino spectrum and muon anomalous magnetic moment," Phys. Rev. **D 93** (2016) 013014, arXiv:1508.07471 [hep-ph].
- [17] G. Bellini et al., "Precision measurement of the  $^7\text{Be}$  solar neutrino interaction rate in Borexino," Phys. Rev. Lett. **107** (2011) 141302, arXiv:1104.1816 [hep-ex].
- [18] Particle Data Group Collaboration, M. Tanabashi et al., "Review of Particle Physics", Phys. Rev. **D 98** no. 3, (2018) 030001.
- [19] NA64 Collaboration, D. Banerjee et al., "Dark Matter Search in Missing Energy Events with NA64", Phys. Rev. Lett. **123** (2019) 121801, arXiv: 1906.00176 [hep-ex].
- [20] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, "Neutrino Trident Production: A Powerful Probe of New Physics with Neutrino Beams", Phys. Rev. Lett. **113**, (2014) 091801, arXiv:1406.2332 [hep-ph].
- [21] J.P. Lees et al. (BaBar Collaboration), "Search for a muonic dark force at BABAR", Phys. Rev. D **94**, (2016) 011102, arXiv:1606.03501 [hep-ex].
- [22] D. Borah, M. Dutta, S. Mahapatra, and N. Sahu, "Muon ( $g-2$ ) and XENON1T excess with boosted dark matter in  $L_\mu - L_\tau$  model", arXiv: 2104.05656v1 [hep-ph].
- [23] M. Escudero, D. Hooper, G. Krnjaic, and M. Pierre, "Cosmology With a Very Light  $L_\mu - L_\tau$  Gauge Boson", JHEP **03** (2019) 071, arXiv:1901.02010 [hep-ph].