

# Emergent Universe Scenario in Modified Gauss-Bonnet Gravity

B. C. Paul · S. D. Maharaj · A. Beesham

Received: date / Accepted: date

**Abstract** We present modified Gauss-Bonnet gravity without matter in four dimensions which accommodates flat emergent universe (EU) obtained in Einstein's general theory of gravity with a non-linear equation of state. The EU model is interesting which is free from big-bang singularity with other observed features of the universe. It is assumed that the present universe emerged out from a static Einstein universe phase exists in the infinite past. To obtain a flat EU model we reconstructed mimetic modified  $f(G)$ -gravity ( $G$  representing Gauss-Bonnet terms) without matter. The functional form of  $f(G)$ -gravity is determined which accommodates the early inflation and late accelerating phases without matter.

## 1. Introduction

In modern cosmology it is believed that the present universe emerged from an inflationary phase in the early universe [1,2,3]. The observational cosmology in recent decades predicted that our universe emerged out from an early inflationary epoch. In spite of success of hot Big

Bang model of the universe it fails to find no solution in Einstein's general theory of relativity with perfect fluid. The concept of early inflation introduced in cosmology resolve some of these issues in cosmology. It is known that inflation can be obtained in a semiclassical theory of gravity or in a higher order theory of gravity. Recent cosmological observations [4,5,6] predicted another interesting feature of the universe that the present universe is passing through an accelerating phase. Theoretically it is challenging to describe the origin of the late accelerating phase as the physics of the inflation [7,8,9,10,11] and the introduction of a small cosmological constant for late acceleration are among the features that are not yet understood [12,13]. A number of proposal put forwarded in understanding the late accelerating universe similar in this direction to realize the early inflation either by a modification of the gravitational sector [14,15,16] or by a modification of the matter sector of the Einstein's general theory of relativity. Thus there is enough motivation to explore for an alternative cosmological model. Astashenok *et al.* [17] in a mimetic modified gravity models investigated various scenarios of cosmological evolution, with or without extra matter fluids to obtain different cosmological evolutions. Considering a simplest formulation based on the use of the Lagrange multiplier constraint it is shown that in some theories, it is possible to realise accelerated expansion of the Universe or even unified evolution, which includes inflation with dark energy, and dark matter described by the theory. Adopting general reconstruction schemes, it is found that mimetic gravity also generates a specific cosmological evolution namely, bouncing cosmologies. Recently, in the literature [17,18,19] cosmological models with a number of known evolutionary features are examined. The motivation of the paper is to reconstruct modified Gauss-Bonnet gravity which per-

---

B. C. Paul  
E-mail: bcpaul@associates.iucaa.in  
Department of Physics, University of North Bengal, Siliguri,  
Dist. : Darjeeling 734 014, West Bengal, India  
S. D. Maharaj  
E-mail: maharaj@ukzn.ac.za  
Astrophysics and Cosmology Research Unit, School of Mathematics, Statistics and Computer Science, University of KwaZulu-Natal, Private Bag X54001, Durban 4000, South Africa  
A. Beesham  
E-mail: abeesham@yahoo.com  
Department of Mathematical Sciences, University of Zululand  
Private Bag X1001, Kwa-Dlangezwa 3886, South Africa

mits a singularity free cosmology namely, flat emergent universe model which encompass the unified evolutions of the universe.

The possibility of a cosmological model [20] in which there is no Big Bang singularity, no beginning of time, and the universe effectively avoids a quantum regime for space-time by staying large at all times was considered first by Ellis and Maartens [21] satisfactorily. In the model the universe started out in the infinite past in an almost static Einstein universe and subsequently it entered in an expanding phase slowly, eventually evolving into a hot Big Bang era. Later Ellis *et al.* [22] constructed an emergent universe scenario for a spatially closed universe with a minimally coupled scalar field  $\phi$ , using a special form for the interaction potential  $V(\phi)$ . However, it was shown [23] immediately that such a potential can be recovered starting from a modified gravitational action with a polynomial Lagrangian  $L = R + \alpha R^2$  making use of a suitable conformal transformation and identification of the field as  $\phi = -\sqrt{3} \ln(1 + 2\alpha R)$ . In the above however one requires to have a negative coupling term  $\alpha$  for a consistent emergent universe (EU) scenario. Later in a flat space-time Mukherjee *et al.* [24] obtained a new cosmological solutions with a static Einstein universe in the framework of a semiclassical gravity [25].

It is known that EU scenario promises to solve several conceptual and technical issues of the Big Bang model. Recent cosmological observations predict that the universe is most likely to be spatially flat. Therefore, it is important to explore a flat EU model. A decade ago Mukherjee *et al.* [26] in the framework of Einstein gravity obtained an EU scenario in a flat universe making use of cosmic fluid that describe by a non-linear equation of state  $p = A\rho - B\sqrt{\rho}$  where  $A$  and  $B$  are arbitrary constants.

The possibility of emergent cosmology using the effective potential formalism has been further explored recently showing that new models of emergent cosmology can be realized within a limit set by the constraints imposed by the cosmic microwave background (CMB) radiation [27]. It has also been demonstrated that within the framework of modified gravity, the emergent scenario may be obtained in a spatially open or closed universe. It is shown that a static Einstein universe solution permitted in a modified theories of gravity with perfect fluid [22] may be useful for constructing an alternative cosmological model without Big Bang singularity. It is also shown later that such an EU scenario can be realized even in a flat universe in a modified theory of gravity by including higher order Gauss-Bonnet terms in the presence of a dilaton coupling [28]. Subsequently the EU model in a flat universe is explored

in various theoretical framework, namely Brane world [29,30,31], Brans-Dicke theory [32], as well as in the context of the non-linear sigma model [33] and found that it can be obtained with all the features of the observed universe. The EU model accommodates late time de-Sitter expansion and thus, naturally permits late time acceleration of the universe. Such a scenario is promising from the perspective of offering unified early as well as late time dynamics of the universe. Note however, that the focal point of unification in such emergent universe models lies in the choice of equation of state for the polytropic fluid, while several other models of unification rely more on the scalar field dynamics in the presence of a suitable field potential [34,35,36]. A number of issues pertaining to different models of EU have been discussed in the literature [37,27]. It is known that Gauss-Bonnet term without dilaton field does not contribute in the dynamics [28] of the universe in four dimensions as it is a Euler number. The motivation of the paper is to obtain EU scenario in a modified gravity with a modified Gauss-Bonnet gravity term without dilaton field which can accommodate a flat EU model of the universe *i.e.*, we reconstruct theoretically an acceptable modified gravity with GB terms in the gravitational action for accommodating EU scenario. The Emergent Universe model is interesting because of its various observed features. It permits a non-singular solution of cosmology, an important and new property in understanding the early evolution of the universe, which is different from that obtained recently in the literature [17,38] where a singular bounce solution is obtained in a  $F(G)$  theory of gravity where  $G = R^2 - 4R_{\mu\delta}R^{\mu\delta} + R_{\mu\alpha\nu\beta}R^{\mu\alpha\nu\beta}$  is the Gauss-Bonnet combination.

The paper is organized as follows: In Sec. 2, we set up the relevant field equation starting from a gravitational action for modified Gauss-Bonnet gravity. In Sec. 3, we present the Gauss-Bonnet modified theory for emergent universe, In Sec. 4, non-Singular bounce from mimetic  $F(G)$  gravity is presented and in Sec. 5, we present a brief discussion.

## 2 . Gravitational Action and the Field Equations

The gravitational action for modified Gauss-Bonnet gravity theory [39,40,41,42,43,44,45,46,57] is given by

$$I = -\frac{1}{2\kappa^2} \int (R + F(G)) \sqrt{-g} \, d^4x + I_m \quad (1)$$

where  $\kappa^2 = \frac{1}{M_P^2}$  with the Planck mass represented by  $M_P$ ,  $R$  represents the Ricci scalar and  $I_m$  represents the

action for matter fields. The field equation is obtained by varying the metric which is given by

$$\begin{aligned}
R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R - \frac{1}{2}g_{\mu\nu}F(G) - (-2RR_{\mu\nu} + 4R_{\mu\delta}R_{\nu}^{\delta} - 2R_{\mu}^{\delta\sigma\tau}R_{\nu\delta\sigma\tau} \\
+ 4g^{\alpha\delta}g^{\beta\sigma}R_{\mu\alpha\nu\beta}R_{\delta\sigma})F'(G) - 2(\nabla_{\mu}\nabla_{\nu}F'(G))R \\
+ 2g_{\mu\nu}(\square F'(G))R - 4(\square F'(G))R_{\mu\nu} + 4(\nabla_{\mu}\nabla_{\nu}F'(G))R_{\nu}^{\delta} \\
+ 4(\nabla_{\delta}\nabla_{\nu}F'(G))R_{\mu}^{\delta} - 4g_{\mu\nu}(\nabla_{\delta}\nabla_{\sigma}F'(G))R^{\delta\sigma} \\
+ 4(\nabla_{\delta}\nabla_{\sigma}F'(G))g^{\alpha\delta}g^{\beta\sigma}R_{\mu\alpha\nu\beta} = \kappa^2 T_{\mu\nu}
\end{aligned} \quad (2)$$

where  $T_{\mu\nu}$  is the energy momentum tensor.

We consider a Robertson-Walker metric for a spatially homogeneous and isotropic space-time which is given by

$$ds^2 = -dt^2 + a^2(t) \left[ \frac{dr^2}{1 - kr^2} + r^2 (d\theta^2 + \sin^2\theta d\phi^2) \right] \quad (3)$$

in the unit  $c = 1$  and  $\hbar = 1$ , where  $a(t)$  represents the scale factor of the universe and  $k = 0, +1, -1$  represent flat, close and open universe respectively. As we explore emergent universe in a flat space-time [26] we consider here  $k = 0$ . The Ricci scalar and the Gauss-Bonnet invariant in a flat space-time are given by

$$R = 6\dot{H} + 12H^2, \quad G = 24H^2(\dot{H} + H^2) \quad (4)$$

where  $H = \frac{\dot{a}}{a}$  is the Hubble parameter and  $\dot{}$  is the time derivative. The gravitational field equations are given by

$$6H^2 + F(G) - GF'(G) + 24H^3\dot{G}F''(G) = 2\kappa^2\rho \quad (5)$$

$$4\dot{H} + 6H^2 + F(G) - GF'(G) + 16H\dot{G}(\dot{H} + H^2)f''(G)$$

$$+ 8H^2\ddot{G}F''(G) + 8H^2\dot{G}^2F'''(G) = -2\kappa^2 p \quad (6)$$

In four dimensions emergent universe (EU) can be obtained with a non-linear equation of state in Einstein gravity. The EU scenario in a flat universe was obtained by Mukherjee *et al.* [26] which evolves out of a static phase of the universe in the infinitely past time. It is free from big bang singularity and exhibits the features of an observable universe [47, 48, 49, 50, 51, 52].

It has been shown that the EU model can be implemented in a modified gravity with Gauss-Bonnet terms coupled with dilaton field [28]. In this paper we explore EU model in vacuum in a modified Gauss-Bonnet gravity and determine the functional form of  $f(G)$  in the action. It is known that in 4 dimensions only GB terms in the action is a Euler number which is not contributing in the dynamics. However, it is shown that modified Gauss-Bonnet gravity plays an important role in understanding the evolution of the universe ([53]-[18]). The form of  $F(G)$ -gravity will be reconstructed by using the method given in Ref. [58, 59, 19] in the next section.

## 2.1 Reconstruction Method of $F(G)$ gravity

We reconstruct  $F(G)$  gravity models by using the method of Ref. ([42, 58, 59]). Consider the gravitational action [18] without matter

$$I = -\frac{1}{2\kappa^2} \int (R + P(t)G + Q(t)) \sqrt{-g} d^4x \quad (7)$$

where  $P(t)$  and  $Q(t)$  are two proper functions of a scalar field  $t$ , which is interpreted as the cosmic time. Now varying the action with respect to  $t$ , we obtain [42, 58, 59, 19, 60, 61]

$$\frac{dP(t)}{dt}G + \frac{dQ(t)}{dt} = 0. \quad (8)$$

The solution of the differential equation can be expressed as  $t = t(G)$ . Consequently substitution of the above solution in eq. (7), yields the modified Gauss-Bonnet gravity functional in the Einstein-Hilbert action

$$F(G) = P(t)G + Q(t). \quad (9)$$

It is interesting to determine  $P(t)$  and  $Q(t)$  in terms of Hubble parameter to study observed cosmological consequences. The modified action may work well to get early inflation as well as late accelerating phase.

Using eqs. (5) and (9) one obtains

$$Q(t) = -6H^2(t) - 24H^3(t)\frac{dP(t)}{dt}. \quad (10)$$

Finally using eqs. (6), (9) and (10), one obtains a second order differential equation given by

$$2H^2(t)\frac{d^2P(t)}{dt^2} + 2H(t)\left(2\dot{H}(t) - H^2(t)\right)\frac{dP(t)}{dt} + \dot{H} = 0. \quad (11)$$

## 3 . Gauss-Bonnet Modified Gravity and The Emergent Universe

The energy-density and pressure obtained from the Einstein field equations in 4-dimensions are given by

$$\rho = 3H^2, \quad p = -(2\dot{H} + 2H^2) \quad (12)$$

where  $H$  represents Hubble parameter where  $\kappa^2 = 1$ . The field equations with a non-linear equation of state (EoS) :  $p = A\rho - B\sqrt{\rho}$  was considered by Mukherjee *et al.* [26] to obtain an emergent universe (EU) model. The important feature of the EU model is that there is no-singularity and it describes the observed features of the universe. The conservation equation is given by

$$\frac{d\rho}{dt} + 3H(\rho + p) = 0. \quad (13)$$

The conservation equation can be integrated using the non-linear EoS which determines the energy density in terms of the scale factors. The energy density is

$$\rho(a) = \frac{1}{(A+1)^2} \left( B + \frac{K}{a^{\frac{3(A+1)}{2}}} \right)^2 \quad (14)$$

which is new and interesting it depends on the EoS parameters  $A$  and  $B$  and  $K$  is an integration constant. It may be mentioned here that Chaplygin gas [62] and its modified forms [63,64,65] given by  $p = A\rho - \frac{B}{\rho^\alpha}$  with  $\alpha > 0$  which is also a promising candidate in cosmology does not permit such an expression for energy density which can be identified with different types of cosmic fluids. In the non-linear EoS considered by us it permits three different types of fluids as the composition of cosmic fluids determined by the parameter  $A$ . Expanding the above equations, the total energy density can be identified with three different fluids corresponding to three different types of EoS. Therefore, one can express

$$\rho(a) = \Sigma_{i=1}^3 \rho_i \text{ and } p(a) = \Sigma_{i=1}^3 p_i. \quad (15)$$

It is noted that the parameter  $A$  plays an important role in understanding different types of cosmic fluids present in the universe [26]. Later considering interactions among the fluids it is shown that a physically viable emergent universe can be obtained for different strengths of interactions among them accommodating observed universe [51]. The scale factor of the universe is obtained from eq. (12) which is

$$a(t) = \left[ \frac{3K(A+1)}{2} \left( \sigma + \frac{2}{\sqrt{3}B} e^{\frac{\sqrt{3}}{2} Bt} \right) \right]^{\frac{2}{3(A+1)}} \quad (16)$$

where  $\sigma$  and  $K$  are integration constants. It is evident that at infinitely past *i.e.*  $t \rightarrow -\infty$ , the scale factor becomes  $a = a_o$  a static Einstein universe which evolves later to a dynamical universe scenario [26,51] encompassing the present observed universe. The interesting aspect of the emergent universe is that there is no singularity, early inflationary universe can be realized in addition to present accelerating phase. The universe however emerged out from an Einstein static phase. It is also shown that the static Einstein phase can be realized in the context of wormhole. The throat of the wormhole expands to a big universe after a pretty long time determined by a cosmological term  $\Lambda$  [66].

For an emergent universe with scale factor given by eq. (16), the differential equation for the Hubble parameter is expressed as

$$\dot{H} = \alpha H - \beta H^2 \quad (17)$$

with  $\alpha = \frac{\sqrt{3}B}{2}$  and  $\beta = \frac{3(A+1)}{2}$ . Now the scale factor given by eq. (16) can be rewritten in a simple form as [28]

$$a(t) = a_o [\eta + e^{\alpha t}]^{\frac{1}{\beta}} \quad (18)$$

denoting  $a_o = \left( \frac{3K(A+1)}{2} \right)^{\frac{2}{3(A+1)}} \frac{2}{\sqrt{3}B}$ ,  $\eta = \frac{\sqrt{3}B\sigma}{2}$ . In this case at infinitely past *i.e.*  $t \rightarrow -\infty$ , the scale factor becomes  $a \rightarrow a_o \eta^{\frac{1}{\beta}}$ , thus a static Einstein universe emerged which later evolves to a dynamical universe [26,51] encompassing the present observed universe. The Hubble parameter is given by  $H = \frac{\alpha e^{\alpha t}}{\beta(\eta + e^{\alpha t})}$ .

The general solution of eq. (11) using eq. (12) can be obtained which accommodates emergent universe solution in a modified gravity in vacuum which is given by

$$P(t) = -\frac{\eta\beta^2(\eta + 2e^{\alpha t})}{\alpha^2(1+\beta)e^{2\alpha t}} + C_1 a_o^{1+2\beta} \int \frac{(\eta + e^{\alpha t})^{2+\frac{1}{\beta}}}{e^{2\alpha t}} dt + C_2 \quad (19)$$

and consequently from eq. (10) one obtains

$$Q(t) = -\frac{6\alpha^2 e^{\alpha t}}{\beta^2(\eta + e^{\alpha t})^2} [8\eta\beta + e^{\alpha t}] + \frac{24C_1 a_o^{1+2\beta} \alpha^3 e^{\alpha t}}{\beta^3(\eta + e^{\alpha t})^{1-\frac{1}{\beta}}} \quad (20)$$

where  $C_1$  and  $C_2$  are integration constants. The Gauss-Bonnet combination can be expressed as a function of cosmic time  $t$  which is now given by

$$G(t) = \frac{24\alpha^4}{\beta^4} \frac{e^{3\alpha t}(\eta\beta + e^{\alpha t})}{(\eta + e^{\alpha t})^4}. \quad (21)$$

The Gauss-Bonnet terms  $G$  is a function of  $t$  which is highly non-linear. An inverse function of  $t$  in terms of  $G$  namely,  $t = t(G)$  can be obtained which is given by

$$t = \pm \left[ \frac{\ln \eta}{\alpha} - \frac{1}{\alpha} \ln \left( \left( \frac{24\alpha^4}{\beta^4 G} \right)^{\frac{1}{4}} \mp 1 \right) \right]. \quad (22)$$

Finally, the modified Gauss-Bonnet function is obtained which is given by

$$F(G) = -\sqrt{\frac{3G}{2}} - \frac{2\beta^2}{\alpha^2(1+\beta)} \left( \frac{\alpha}{\beta} \left( \frac{24}{G} \right)^{\frac{1}{4}} - 1 \right) G - \frac{4\beta\eta^2\sqrt{6G}}{\left( \frac{\alpha}{\beta} \left( \frac{24}{G} \right)^{\frac{1}{4}} - 1 \right)} - \frac{\beta^2 G}{\alpha^2(1+\beta) \left( \frac{\alpha}{\beta} \left( \frac{24}{G} \right)^{\frac{1}{4}} - 1 \right)} \quad (23)$$

for  $C_1 = 0$  and  $C_2 = 0$ . The interesting feature of the emergent universe solution is that the initial singularity does not arise here. The universe in course of its evolution admits observed features of the universe scenario as obtained in Ref. [26,51].

#### 4 . Emergent Universe from Mimetic $F(G)$ Gravity

To obtain the mimetic  $F(G)$ -gravity the following parametrization of metric is assumed in the gravitational action given by eq. (1) : [60]-[73]

$$g_{\mu\nu} = -\bar{g}^{\rho\tau} \partial_\rho \phi \partial_\tau \phi \bar{g}_{\mu\nu} \quad (24)$$

Varying the metric one gets

$$\delta g_{\mu\nu} = \bar{g}^{\rho\tau} \delta g_{\tau\omega} \bar{g}^{\omega\sigma} \partial_\rho \phi \partial_\sigma \phi \bar{g}_{\mu\nu} - \bar{g}^{\rho\sigma} \partial_\rho \phi \partial_\sigma \phi \delta \bar{g}_{\mu\nu}$$

$$-2\bar{g}^{\rho\sigma} \partial_\rho \phi \partial_\sigma (\delta \phi) \bar{g}_{\mu\nu}. \quad (25)$$

Consequently one obtains following field equations [60,61,67,68,69,70,71,72,73] varying the action with respect to the redefined metric  $\bar{g}_{\rho\sigma}$  instead of the standard Jordan frame metric  $g_{\mu\nu}$  one obtains

$$\begin{aligned} R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \\ 8[R_{\mu\delta\nu\sigma} + R_{\delta\nu}g_{\sigma\mu} - R_{\delta\sigma}g_{\nu\mu} - R_{\mu\nu}g_{\sigma\delta} + R_{\mu\sigma}g_{\nu\delta}] \nabla^\delta \nabla^\sigma F_G \\ + 4(g_{\mu\sigma} - g_{\nu\delta})R \nabla^\delta \nabla^\sigma F_G + (F_G G - F(G))g_{\mu\nu} + \partial_\mu \phi \partial_\nu \phi \\ \left[ -R + 8 \left( -R_{\delta\sigma} + \frac{1}{2}g_{\delta\sigma}R \right) \nabla^\delta \nabla^\sigma F_G + 4(F_G G - F(G)) \right] \\ = T_{\mu\nu} + \nabla_\mu \phi \nabla_\nu \phi T \end{aligned} \quad (26)$$

where  $F_G = \frac{dF}{dG}$  and  $\nabla_\mu f_\nu = \partial_\mu f_\nu - \Gamma_{\mu\nu}^\lambda f_\lambda$  with the energy momentum tensor  $T_{\mu\nu} = \text{diagonal}(\rho, -p, -p, -p)$ ,  $\rho$  the energy density,  $p$  the cosmic pressure and  $\phi$  mimetic scalar field. Now varying the action w.r.t.  $\phi$  one obtains

$$\begin{aligned} \nabla^\mu \left[ \partial_\mu \phi \left( -R - 8 \left( R_{\delta\sigma} - \frac{1}{2}g_{\delta\sigma}R \right) \nabla^\delta \nabla^\sigma F_G \right) \right] \\ + 4(F_G G - F(G)) - T = 0. \end{aligned} \quad (27)$$

From eq. (24) one obtains

$$g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi = -1,$$

where homogeneous scalar field  $\phi$  depends on cosmic time with a constraint  $\phi = t$ . For a flat universe the time-time component of the field equation in eq. (26) is expressed as

$$\begin{aligned} 2\dot{H} + 3H^2 + 16H(\dot{H} + H^2) \frac{dF_G}{dt} \\ + 8H^2 \frac{d^2 F_G}{dt^2} - (F_G G - F(G)) = -p. \end{aligned} \quad (28)$$

On integration eq. (27), one obtains

$$-R - 8 \left( R_{\delta\sigma} - \frac{R}{2}g_{\delta\sigma} \right) \partial^\delta \partial^\sigma F_G +$$

$$4(F_G G - F(G)) - \rho - 3p = -\frac{C}{a^3} \quad (29)$$

$$2\dot{H} + 2H^2 + \frac{2}{3}(F_G G - F(G)) +$$

$$4H(2\dot{H} + 3H^2) \frac{dF_G}{dt} + 4H^2 \frac{d^2 F_G}{dt^2} = -\frac{\rho}{6} + \frac{p}{2} - \frac{C}{a^3} \quad (30)$$

here  $C$  is an integration constant. Using eqs. (28) and (30) a final equation is obtained which is

$$\begin{aligned} \dot{H} + 2H^2 + \frac{2}{3}(F_G G - F(G)) + \\ 4H(2\dot{H} - H^2) \frac{dF_G}{dt} + 4H^2 \frac{d^2 F_G}{dt^2} = -\frac{1}{2}(\rho + p) - \frac{C}{a^3}. \end{aligned} \quad (31)$$

Here we denote  $g(t) = \frac{dF_G}{dt}$  and the above equation can be written as

$$4H^2 \frac{dg(t)}{dt} + 4H(2\dot{H} - H^2)g(t) = -\dot{H} - \frac{(\rho + p)}{2} - \frac{C}{a^3}. \quad (32)$$

To obtain an emergent universe model discussed in sec.(II), we use the eq. (17), it will generate the modified gravity. The general solution now can be obtained from eq.(32) corresponding to a functional form given by

$$g(t) = -\frac{C_1}{H^2}a(t) \quad (33)$$

where  $C_1$  is an integration constant. In this section, we reconstruct the modified Gauss-Bonnet gravity to realize the emergent universe scenario obtained by Mukherjee *et. al* [26] in Einstein's General theory of relativity with non-linear equation of state describing the matter. Now emergent universe model is obtained when the right hand side of eq. (32) in the absence of dark radiation ( $C = 0$ ) is replaced by

$$\dot{H} + \frac{1}{2}(\rho + p) = \frac{1}{2}H^2 \quad (34)$$

which determines the modified Gauss-Bonnet terms. The modified GB terms will be determined in the next section.

The Hubble parameter for emergent universe model is

$$H = \frac{H_0 e^{\alpha t}}{\eta + e^{\alpha t}}, \quad H_0 = \frac{\alpha}{\beta}.$$

Thus the general solution in terms of  $t$  is given by

$$g(t) = \frac{C_1 a_0 \beta^2}{\alpha^2 e^{2\alpha t}} (\eta + e^{\alpha t})^{2+\frac{2}{\beta}}. \quad (35)$$

The modified Gauss-Bonnet gravity is then determined in terms of  $t$  as

$$F_G(t) = \int g(t) dt. \quad (36)$$

It can be integrated for different values of  $\beta$ .

- For  $\beta = 1$  one obtains

$$F_G(t) = g_o \left( \frac{e^{2\alpha t}}{2} + 4\eta e^{\alpha t} + 6\eta^2 \alpha t - 4\eta^3 e^{-\alpha t} - \frac{\eta^4 e^{-2\alpha t}}{2} \right) + F_o \quad (37)$$

where  $g_o = \frac{C_1 a_o \beta^2}{\alpha^3}$  and  $F_o$  is a constant.

- For  $\beta = 2$

$$F_G(t) = g_o \left( e^{\alpha t} + 3\eta \alpha t - 3\eta^2 e^{-\alpha t} - \frac{\eta^3}{2} e^{-2\alpha t} \right) + F_1 \quad (38)$$

where  $F_1$  is an integration constant. It is noted that  $\beta$  is playing here an important role in determining the modified Gauss-Bonnet gravity to accommodate emergent universe solution. Using the GB terms with the scale factor for emergent universe,  $t$  can be replaced by  $G$  as

$$t = \frac{1}{\alpha} \frac{\eta}{\left(\frac{G_o}{G}\right)^{1/4} - 1} \quad (39)$$

where  $G_o = 24H_0^4(1 + \beta\eta)$ . The functional form of modified Gauss-Bonnet gravity can be determined from

- (i) when  $\beta = 1$

$$F_G(G) = g_o \left( \frac{\eta^2}{2 \left[ \left(\frac{G_o}{G}\right)^{1/4} - 1 \right]^2} + \frac{4\eta^2}{\left(\frac{G_o}{G}\right)^{1/4} - 1} \right) - 6\eta^2 g_o \ln \left[ \left(\frac{G_o}{G}\right)^{1/4} - 1 \right] - 4g_o \eta^2 \left[ \left(\frac{G_o}{G}\right)^{1/4} - 1 \right] - \frac{g_o \eta^4}{2} \left[ \left(\frac{G_o}{G}\right)^{1/4} - 1 \right]^2 + f_o \quad (40)$$

- (ii) when  $\beta = 2$

$$F_G(G) = g_o \left( \frac{\eta}{\left(\frac{G_o}{G}\right)^{1/4} - 1} - 3\eta \ln \left[ \left(\frac{G_o}{G}\right)^{1/4} - 1 \right] \right) - g_o \eta \left[ \left(\frac{G_o}{G}\right)^{1/4} - 1 \right] - \frac{g_o \eta}{2} \left[ \left(\frac{G_o}{G}\right)^{1/4} - 1 \right]^2 + f_1 \quad (41)$$

where  $f_o$  and  $f_1$  are constants. The above modified gravitational actions accommodates EU scenario fairly well. For  $G \gg G_o$  one gets

$$\begin{aligned} \bullet F(G) &= \lambda_1 + \left( f_o - 10g_o \eta^2 - \frac{1}{2} g_o \eta^4 - \frac{3\eta^2 g_o \ln G_o}{2} \right) G + \\ &\frac{3\eta^2 g_o}{2} G \ln G - \frac{13g_o \eta^2 G_o}{2} \ln G + 8\eta^2 g_o G_o^{\frac{3}{4}} G^{\frac{1}{4}} - (2\eta^2 + \\ &11) g_o \eta^2 \sqrt{G_o G} + 4g_o \eta^2 (\eta^2 - 1) G_o^{\frac{1}{4}} G^{\frac{3}{4}}; \\ \bullet F(G) &= \left( f_1 - \frac{3g_o \eta}{2} - \frac{3g_o \eta}{4} \ln G_o \right) G + \frac{3g_o \eta}{4} G \ln G - \\ &g_o \eta G_o \ln G + 4g_o \eta G_o^{\frac{3}{4}} G^{\frac{1}{4}} - 3g_o \eta \sqrt{G_o} \sqrt{G} + \\ &\frac{4g_o \eta G_o^{\frac{1}{4}}}{3} G^{\frac{3}{4}} + \lambda_2 \end{aligned}$$

where  $\lambda_1$  and  $\lambda_2$  are integration constants for  $\beta = 1$  and 2 respectively.

## 5. Discussion

In this paper modified gravity with Gauss-Bonnet terms is reconstructed which accommodates flat emergent universe model. Earlier a number of cosmological models with various evolution having a bouncing solution are explored with a reconstructed mimetic gravity [17, 18, 19], but the present work is different. It is known that the Gauss-Bonnet terms in four dimensions is equivalent to a Euler number which does not contribute in the dynamics of a matter dominated universe. However, in the presence of a dilaton field coupled to Gauss-Bonnet combination is capable of producing the evolutionary features of the observed universe. It is also shown that an Emergent universe scenario can be realized in this framework which describes early universe and late universe fairly well [28]. The non-linear functional form of the Gauss-Bonnet gravity is obtained which accommodates the emergent universe scenario with all the features of the observed universe. We note that in the absence of a dilaton field a non-linear combination of GB terms accommodates EU solution. We reconstructed the modified gravity action with  $F(G)$  which is relevant in the early and also in the late universe. The non-linear EoS parameters  $A$  and  $B$  of GR is replaced here by  $\alpha$  and  $\beta$ . It is noted that a number of modified gravity can be predicted with different values of  $\beta$ -parameter which admits EU model here. It is found that for a given  $\alpha$ , the Hubble parameter decreases with an increase in  $\beta$  at the present epoch. Thus Gauss-Bonnet terms play an important role in understanding the observed universe which is described by dark matter and dark energy in Einstein General theory of relativity. The parameter  $\eta$  is also playing a significant role which analogously compared with the non-linear EoS parameter  $A$  in GR for obtaining EU scenario. Thus a rich structure of GR modified with non-linear combination of Gauss-Bonnet terms in the Einstein-Hilbert action is explored here which plays an important role in cosmology.

## 6 Acknowledgements:

BCP would like to acknowledge support for a visit to the University of Zululand, South Africa and University of Kwa-Zulu Natal, South Africa where the work has been initiated. BCP would like to thank IUCAA Pune, India for warm hospitality and DST-SERB for supporting project (EMR/2016/005734).

## References

1. E. W. Kolb and M. S. Turner, *The Early Universe* (Addison-Wesley, Mass. , 1990).
2. S. Dodelson, *Modern Cosmology* (Academic press, 2003).
3. S. Weinberg *Cosmology*, (Oxford University Press, 2008).
4. A. G. Riess *et al*, *Astron. J.* **116** 1009 (1998).
5. S. Perlmutter *et al*, *Nature* **51** 391 (1998).
6. S. Perlmutter *et al*, *Astrophys. J.* **517** 565 (1999).
7. A. H. Guth, *Phys. Rev. D* **23** 347 (1981).
8. K. Sato, *Mon. Not. Roy. Astron. Soc.* **195** 467 (1981).
9. A. D. Linde, *Phys. Lett. B* **108** 389 (1982).
10. A. Albrecht and P. J. Steinhardt, *Phys. Rev. Lett.* **48** 1220 (1982).
11. A. D. Linde, *Phys. Lett. B* **129** 177 (1983).
12. R. H. Dicke, P. J. E. Peebles, P. J. Roll and D. T. Wilkinson, *Astrophys. J. Lett.* **142** 414 (1965).
13. S. M Carroll, *Living Rev. Rel.* **4** 1 (2001).
14. S. M. Carroll, V. Duvvuri, M. Trodden and M. S. Turner, *Phys. Rev. D* **70** 043528 (2004).
15. S. Nojiri and S. D. Odintsov, *J. Phys. Conf. Ser.* **66** 012005 (2007).
16. S. Nojiri and S. D. Odintsov, *J. Phys. A* **40** 6725 (2007).
17. A. V. Astashenok, S. D. Odintsov and V. K. Oikonomou, *Class. Quantum Grav.* **32** 185007 (2015).
18. K. Bamba, A. N. Makarenko, A. N. Myagky and S. D. Odintsov, *Phys. Lett. B* **732** 349 (2014).
19. K. Bamba, S. Nojiri and S. D. Odintsov, *JCAP* **0810** 045 (2008).
20. E. R. Harrison, *Mon. Not. Roy. Astron. Soc.* **69** 137 (1967).
21. G. F. R. Ellis and R. Maartens, *Class. Quant. Grav.* **21** 223 (2004).
22. G. F. R. Ellis, J. Murugan and C. G. Tsagas, *Class. Quant. Grav.* **21** 233 (2004).
23. D. J. Mulryne, R. Tavakol, J. E. Lidsey and G. F. R. Ellis, *Phys. Rev. D* **71** 123512 (2005).
24. S. Mukherjee, B. C. Paul, S. D. Maharaj and A. Beesham, arXiv:gr-qc/0505103 (2005).
25. A. A. Starobinsky, *Phys. Letts.* **91 B** 99 (1980).
26. S. Mukherjee, B. C. Paul, N. Dadhich, S. D. Maharaj and A. Beesham, *Class. Quant. Grav.* **23** 6927 (2006).
27. S. Bag, V. Sahni, Y. Shtanov and S. Unnikrishnan, *JCAP* **07** 034 (2014).
28. B. C. Paul and S. Ghose, *Gen. Rel. Grav.* **42** 795 (2010).
29. A. Banerjee, T. Bandyopadhyay and S. Chakraborty, *Gen. Rel. Grav.* **40** 1603 (2008).
30. A. Banerjee, T. Bandyopadhyay and S. Chakraborty, *Grav. & Cosmology* **13** 290 (2007).
31. U. Debnath, *Class. Quant. Grav.* **25** 205019 (2008).
32. S. del Campo, R. Herrera and P. Labrana, *J. Cosmo. Astropart. Physics* **30** 0711 (2007).
33. A. Beesham, S. V. Chervon and S. D. Maharaj, *Class. Quantum Grav.* **26** 075017 (2009).
34. A. S. Majumdar, *Phys. Rev. D* **64** 083503 (2001).
35. N. Bose and A. S. Majumdar, *Phys. Rev. D* **79** 103517 (2009).
36. N. Bose and A. S. Majumdar, *Phys. Rev. D* **80** 103508 (2009).
37. L. P. Chimento, *Phys. Rev. D* **81** 043525 (2010).
38. V. K. Oikonomou, *Phys. Rev. D* **92**, 124027 (2015)'
39. K. Bamba, Zong-Kuan Guo and N. Ohta, *Prog. Theor. Phys.* **118** 879 (2007).
40. G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, *Phys. Rev. D* **75** 086002 (2007).
41. G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, *Euro. Phys. J C* **64** 483 (2009).
42. K. Bamba, S. D. Odintsov, L. Sebastiani and S. Zerbini, *Euro. Phys. J C* **67** 295 (2010).
43. S. Nojiri and S. D. Odintsov, *Phys. Rep.* **505** 59 (2011).
44. S. Nojiri and S. D. Odintsov, *Int. J. Geom. Methods Mod. Phys.* **11** 1460006 (2014).
45. V. Faraoni and S. Capozziello, *Beyond Einstein Gravity: A Survey of Gravitational Theories for Cosmology and Astrophysics, Fundamental Theories of Physics* Vol. 170 (Springer, New York, 2010) .
46. S. Capozziello and M. De Laurentis, *Phys. Rep.* **509** 167 (2011).
47. B. C. Paul, P. Thakur and S. Ghose *Mon. Not. Roy. Astron. Soc.* **407** 415 (2010).
48. B. C. Paul, S. Ghose and P. Thakur, *Mon. Not. Roy. Astron. Soc.* **413** 686 (2011).
49. B. C. Paul and P. Thakur, *Astrophysics and Space Sci.* **362** 73 (2017).
50. S. Ghose, P. Thakur and B. C. Paul, *Mon. Not. Roy. Astron. Soc.* **421** 20 (2012).
51. B. C. Paul and A. S. Majumdar, *Class. Quantum Grav.* **32** 115001 (2015).
52. B. C. Paul, P. Thakur and A. Beesham, *Astrophys. Space. Sci.* **361** 336 (2016).
53. S. Nojiri , S. D. Odintsov and M. Sasaki, *Phys. Rev.* **71** 123509 (2005).
54. B. Li, J. D. Barrow and D.F. Mota, *Phys. Rev.* **76** 044027 (2007).
55. S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **631** 1 (2005).
56. S. Nojiri, S.D. Odintsov and O. G. Gorbunova, *J. Phys. A* **39** 6627 (2006).
57. G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov and S. Zerbini, *Phys. Rev. D* **73** 084007 (2006).
58. S. Capozziello, S. Nojiri, S. D. Odintsov and A. Troisi, *Phys. Lett. B* **639** 135 (2006).
59. S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **74** 086005 (2006).
60. A. H. Chamseddine and V. Mukhanov, *J. High Energy Phys. JHEP* **11**, 135 (2013).
61. A. H. Chamseddine, V. Mukhanov and A. Vikman, *J. Cosmol. Astropart. Phys. JCAP* **06**, 017 (2014).
62. S. Chaplygin, *Sci. Mem. Moscow Univ. Math. Phys.* **21** 1 (1904).
63. N. Blic, G. B. Tupper and R. D. Viollier, *Phys. Lett. B* **535** 17 (2002).
64. M. C. Bento, O. Bertolamu and A. Sen, *Phys. Rev. D* **66** 043507 (2002).
65. B. C. Paul and P. Thakur, *J. Cosmo. Astroparticle Phys.* **11** 052 (2013).
66. B. C. Paul and A. S. Majumdar, *Class. & Quantum Gravity* **35** 065001 (2018).
67. A. Golovnev, *Phys. Lett. B* **728** 39 (2014).
68. N. Deruelle and J. Rua, *J. Cosmol. Astropart. Phys. JCAP* **04** 002 (2014).
69. D. Momeni, A. Altaibayeva and R. Myrzakulov, *Int. J. Geom. Methods Mod. Phys.* **11** 1450091 (2014).

- 
70. J. Matsumoto, S. D. Odintsov and S. V. Sushkov, *Phys. Rev. D* **91** 064062 (2015).
  71. G. Leon and E. N. Saridakis, *J. Cosmol. Astropart. Phys. JCAP* **04** 031 (2015).
  72. R. Myrzakulov and L. Sebastiani, *Gen. Rel. Grav.* **47** 89 (2015).
  73. D. Momeni, R. Myrzakulov and E. Gdekli, *Int. J. Geom. Methods Mod. Phys.* **12** 1550101 (2015).