

THE NONLINEAR SCHRÖDINGER EQUATION IN THE HALF-SPACE

ANTONIO J. FERNÁNDEZ AND TOBIAS WETH

ABSTRACT. The present paper is concerned with the half-space Dirichlet problem

$$(P_c) \quad -\Delta v + v = |v|^{p-1}v, \quad \text{in } \mathbb{R}_+^N, \quad v = c, \quad \text{on } \partial\mathbb{R}_+^N, \quad \lim_{x_N \rightarrow \infty} v(x', x_N) = 0 \text{ uniformly in } x' \in \mathbb{R}^{N-1},$$

where $\mathbb{R}_+^N := \{x \in \mathbb{R}^N : x_N > 0\}$ for some $N \geq 1$ and $p > 1$, $c > 0$ are constants. We analyse the existence, non-existence and multiplicity of bounded positive solutions to (P_c) . We prove that the existence of bounded positive solutions to (P_c) depend on a striking way on the value of $c > 0$ and also on the dimension N . We find an explicit number $c_p \in (1, \sqrt{e})$, depending only on p , which determines the threshold between existence and non-existence. In particular, in dimensions $N \geq 2$, we prove that, for $0 < c < c_p$, problem (P_c) admits infinitely many bounded positive solutions, whereas, for $c > c_p$, there are no bounded positive solutions to (P_c) .

1. INTRODUCTION

Due to its relevance within several models arising in physics and biology, the nonlinear stationary Schrödinger equation

$$(1.1) \quad -\Delta v + v = |v|^{p-1}v, \quad \text{in } \mathbb{R}^N,$$

received extensive attention in the last four decades. In particular, let us mention that the study of solitary wave solutions for the (focusing) NLS

$$i\partial_t \varphi + \Delta \varphi + |\varphi|^{p-1} \varphi = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^N,$$

is reduced to problem (1.1) via a time-harmonic ansatz. For classic existence and multiplicity results, we refer the reader e.g. to the seminal papers [2, 3, 6, 8, 9, 27, 28] and the monographs [29, 31]. We also recall the fundamental works [16, 21] where the radial symmetry and uniqueness, up to translations, of positive solutions to (1.1) satisfying the decay condition

$$(1.2) \quad v(x) \rightarrow 0, \quad \text{as } |x| \rightarrow \infty,$$

are proved in the case $1 < p < 2^* - 1$. In particular, these results imply the uniqueness, up to translations, of positive finite energy solutions $u \in H^1(\mathbb{R}^N)$. In contrast, (1.1) admits an abundance of sign-changing finite energy solutions satisfying (1.2), see e.g. [2, 3, 24, 25] and the references therein. Moreover, more recent geometric constructions of different solution shapes highlight the rich structure of the set of positive solutions which do not satisfy the decay assumption (1.2), see e.g. [1, 11, 23, 25] and the references therein.

Whereas it seems impossible to provide an exhaustive list of references for the full space problem (1.1), much less is known regarding the half-space Dirichlet problem

$$(1.3) \quad \begin{cases} -\Delta v + v = |v|^{p-1}v, & \text{in } \mathbb{R}_+^N, \\ v = c, & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

where $\mathbb{R}_+^N := \{x \in \mathbb{R}^N : x_N > 0\}$ for some $N \geq 1$ and $c \geq 0$ is a constant. In the case $c = 0$, general nonexistence results are available for (1.3). More precisely, the non-existence of finite energy solutions $u \in H_0^1(\mathbb{R}_+^N)$ to (1.3) in the case $c = 0$ follows from [14, Theorem I.1], while [4, Corollary 1.3] yields, in particular, the non-existence of positive solutions to (1.3) with $c = 0$ and the the decay property

$$(1.4) \quad \lim_{x_N \rightarrow \infty} v(x', x_N) = 0, \quad \text{uniformly in } x' \in \mathbb{R}^{N-1}.$$

The aim of the present paper is to analyse the existence, non-existence and multiplicity of bounded positive solutions v to the problem (1.3)-(1.4) in the case $c > 0$, for which we are not aware of any previous result in general dimensions N . As we shall see below, the multiplicity of positive solutions depends on a striking way

on the value $c > 0$ and, somewhat surprisingly, also on the dimension N . Let us stress that we cannot expect the existence of finite energy solutions $u \in H^1(\mathbb{R}_+^N)$ to (1.3) in the case $N \geq 2$. Actually, we cannot expect solutions to (1.3) belonging to $L^p(\mathbb{R}_+^N)$ for any $1 \leq p < \infty$. The one-dimensional decay condition (1.4) therefore seems natural.

Not surprisingly, problem (1.3) is completely understood in the case $N = 1$. This is due to the fact that the one-dimensional equation

$$(1.5) \quad -w'' + w = w^p$$

admits a first integral, and from this one can easily deduce that, for $1 < p < \infty$, (1.5) admits, up to sign and translation, a unique global non trivial solution satisfying $w(t) \rightarrow 0$ as $t \rightarrow \pm\infty$. See e.g. [8, Theorem 5] and [31, Theorem 3.16]. By direct computations, one can verify that this solution is precisely given by

$$(1.6) \quad t \mapsto w_0(t) = c_p \left[\cosh\left(\frac{p-1}{2}t\right) \right]^{-\frac{2}{p-1}}, \quad \text{with} \quad c_p := \left(\frac{p+1}{2}\right)^{\frac{1}{p-1}} = w_0(0) = \sup_{t \in \mathbb{R}} w_0(t).$$

As we shall see below, the value c_p will be of key importance also for the higher dimensional version of (1.3). The following complete characterization of the one dimensional case is an immediate consequence of these facts.

Proposition 1.1. *Let $N = 1$, $p > 1$ and $c > 0$. Then:*

- (i) *If $0 < c < c_p$, problem (1.3)-(1.4) admits exactly two positive solutions given by $t \mapsto w_0(t + t_{c,p})$ and $t \mapsto w_0(t - t_{c,p})$, where*

$$(1.7) \quad t_{c,p} := \frac{2}{p-1} \ln \left(\sqrt{\frac{p+1}{2c^{p-1}}} + \sqrt{\frac{p+1}{2c^{p-1}} - 1} \right).$$

- (ii) *If $c = c_p$, the function w_0 is the unique positive solution to (1.3)-(1.4).*
- (iii) *If $c > c_p$, problem (1.3)-(1.4) does not admit solutions.*

Our main result concerns dimensions $N \geq 2$ and reads as follows.

Theorem 1.2. *Let $N \geq 2$, $p > 1$ and $c > 0$. Then:*

- (i) *If $0 < c < c_p$, there exist infinitely many bounded positive solutions to (1.3)-(1.4).*
- (ii) *If $c > c_p$, there are no bounded positive solutions to (1.3)-(1.4).*

Remark 1.1.

- (a) By a bounded positive solution to (1.3)-(1.4), we mean a positive function $v \in C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N}) \cap L^\infty(\mathbb{R}_+^N)$ satisfying (1.3) in the pointwise sense and such that (1.4) holds.
- (b) The nonexistence part (ii) of Theorem 1.2 is proved with a variant of the sliding method based on a comparison with x_N -translates of the function $x \mapsto w_0(x_N)$. We shall comment on this in more detail further below.
- (c) As we have indicated above already, Theorem 1.2 (i) highlights the rich structure of solutions to (1.3)-(1.4) in the case $N \geq 2$, $0 < c < c_p$, which is in striking contrast to the case $N = 1$.
- (d) It suffices to prove Theorem 1.2 (i) in the case $N = 2$ since every positive solution v to (1.3)-(1.4) in the case $N = 2$ gives rise to a corresponding solution \tilde{v} to (1.3)-(1.4) in general dimension $N \geq 3$ by simply setting $\tilde{v}(x) = v(x_1, x_N)$.
- (e) Problem (1.3)-(1.4) is invariant under translations and rotations parallel to the boundary $\partial\mathbb{R}_+^N = \mathbb{R}^{N-1}$. In particular, if $N \geq 2$, any positive solution v to (1.3)-(1.4) which is not merely a function of the x_N -variable already gives rise to infinitely many solutions of the type $x \mapsto v(x' + \tau, x_N)$, $\tau \in \mathbb{R}^{N-1}$, where, here and in the following, we write $x = (x', x_N)$ for $x \in \mathbb{R}_+^N$ with $x' \in \mathbb{R}^{N-1}$.
- (f) It remains as an interesting open question whether the function $x \mapsto w_0(x_N)$ is the unique bounded positive solution to (1.3)-(1.4) in the case $c = c_p$. At first glance, it seems natural to establish such a uniqueness result also with the help of a sliding argument as mentioned in (b) above, but additional difficulties appear in the case $c = c_p$, and non-uniqueness remains a possibility for now.

The following result provides some information on the shape of the solutions we construct.

Theorem 1.3. *Let $N \geq 2$, $1 < p < 2^* - 1$ and $0 < c < c_p$. Then there exists a positive solution to (1.3)-(1.4) of the form*

$$(1.8) \quad x \mapsto w_0(x_N + t_{c,p}) + u(x),$$

with a nonnegative function $u \in H_0^1(\mathbb{R}_+^N) \setminus \{0\}$.

Here and in the following, 2^* denotes the critical Sobolev exponent, i.e. $2^* = \frac{2N}{N-2}$ for $N \geq 3$ and $2^* = 2^* - 1 = \infty$ for $N = 1, 2$. By the remarks above and since all exponents $p < \infty$ are subcritical in the case $N = 2$, Theorem 1.3 implies Theorem 1.2 (i) in the case $N = 2$ and therefore for all $N \geq 2$. It also allows us to distinguish different solution orbits under translations and rotations in \mathbb{R}^{N-1} . In the following, we call two solutions *geometrically distinct* if they do not belong to the same orbit of solutions under translations and rotations in \mathbb{R}^{N-1} .

Corollary 1.4. *Let $N \geq 3$, $1 < p < \frac{M+2}{M-2}$ for some $M \in \{3, \dots, N\}$ and $0 < c < c_p$. Then problem (1.3)-(1.4) admits at least $M + 1$ geometrically distinct positive solutions.*

This result is a rather immediate corollary of Theorem 1.3. Indeed, under the given assumptions, for every dimension $\tilde{N} \in \{2, 3, \dots, M\}$, Theorem 1.3 yields the existence of a solution to (1.3)-(1.4) of the form

$$x \mapsto w_0(x_N + t_{c,p}) + u(x_1, \dots, x_{\tilde{N}-1}, x_{\tilde{N}})$$

with a nonnegative $u \in H_0^1(\mathbb{R}_+^{\tilde{N}}) \setminus \{0\}$. Clearly, these ansatzs give rise to solutions that are geometrically different.

It is natural to guess that the change of the solution set when passing from $c > c_p$ to $c < c_p$ is a bifurcation phenomenon. More precisely, one may guess that the solutions constructed in Theorem 1.3 have the property that $u = u_c \rightarrow 0 \in H_0^1(\mathbb{R}_+^N)$ as $c \nearrow c_p$ for the functions u in the ansatz (1.8). This remains an open question, and the answer could even depend on the value of p . We note that standard results from bifurcation theory do not apply here since the linearized problem

$$\begin{cases} -\Delta v + v - p|w_0(x_N)|^{p-1}v = 0 & \text{in } \mathbb{R}_+^N, \\ v = 0 & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

at the parameter value $c = c_p$ has purely essential spectrum due to its invariance with respect to translations in directions parallel to the boundary $\partial\mathbb{R}_+^N = \mathbb{R}^{N-1}$. Bifurcation from the essential spectrum has been observed successfully in other contexts (see e.g. the survey paper [30] and the references therein), but there is still no general functional analytic framework which provides sufficient abstract conditions.

We now give some ideas of the proof of Theorem 1.3. For this we fix $c \in (0, c_p)$ and define the functions

$$t \mapsto z_c(t) := w_0(t + t_{c,p}) \quad \text{and} \quad t \mapsto \tilde{z}_c(t) := w_0(t - t_{c,p}),$$

where $t_{c,p}$ is given in (1.7). We recall that z_c and \tilde{z}_c are the unique positive solutions to (1.5) such that $z_c(0) = \tilde{z}_c(0) = c$. Moreover, we define $u_c : \overline{\mathbb{R}_+^N} \rightarrow \mathbb{R}$ and $\tilde{u}_c : \overline{\mathbb{R}_+^N} \rightarrow \mathbb{R}$ as

$$(1.9) \quad u_c(x) := z_c(x_N) \quad \text{and} \quad \tilde{u}_c(x) := \tilde{z}_c(x_N),$$

and we directly notice that u_c and \tilde{u}_c are both solutions to (1.3)-(1.4). Furthermore, it follows that

$$(1.10) \quad u_c(x) \sim e^{-x_N} \quad \text{and} \quad \tilde{u}_c(x) \sim e^{-x_N}, \quad \text{as } x_N \rightarrow \infty.$$

Proving Theorem 1.3 now amounts to find a nonnegative solution $u \in H_0^1(\mathbb{R}_+^N) \setminus \{0\}$ to the non-autonomous Schrödinger type equation

$$(1.11) \quad -\Delta u + u = f(x, u), \quad u \in H_0^1(\mathbb{R}_+^N),$$

with

$$(1.12) \quad f(x, s) := |u_c(x) + s|^{p-1}(u_c(x) + s) - (u_c(x))^p,$$

because in this case $v = u_c + u$ is of the form (1.8), solves (1.3) and it is easy to see that also satisfies (1.4). Since we are interested in finding non-negative solutions to (1.11), we truncate the nonlinearity and define

$$(1.13) \quad g(x, s) := (u_c(x) + s^+)^p - (u_c(x))^p = \begin{cases} f(x, s), & \text{if } s \geq 0, \\ 0, & \text{if } s \leq 0, \end{cases}$$

with f given in (1.12). We then consider the auxiliary problem

$$(1.14) \quad -\Delta u + u = g(x, u), \quad u \in H_0^1(\mathbb{R}_+^N).$$

Considering $u^- \in H_0^1(\mathbb{R}_+^N)$ as test function in (1.14), one can easily check that every solution to (1.14) is non-negative and so, that every solution to (1.14) is a non-negative solution to (1.11). It might be worth pointing out that the one-dimensional function $\tilde{u} := \tilde{u}_c - u_c \in C^2(\mathbb{R}_+^N)$ is a positive solution to the equation in (1.14) and also satisfies $\tilde{u} = 0$ on $\partial\mathbb{R}_+^N$. However, $\tilde{u} \notin H_0^1(\mathbb{R}_+^N)$ since it only depends on the x_N variable. Hence, \tilde{u} is not a solution to (1.14).

We shall look for a non-trivial solution to (1.14) as a critical point of the associated functional

$$(1.15) \quad E : H_0^1(\mathbb{R}_+^N) \rightarrow \mathbb{R}, \quad E(u) = \frac{1}{2} \int_{\mathbb{R}_+^N} (|\nabla u|^2 + u^2) dx - \int_{\mathbb{R}_+^N} G(x, u) dx,$$

where

$$(1.16) \quad G(x, u) := \int_0^u g(x, s) ds = \frac{1}{p+1} \left((u_c + u^+)^{p+1} - u_c^{p+1} - (p+1)u_c^p u^+ \right).$$

More precisely, we are going to prove the existence of a non-trivial critical point of mountain pass type. This requires new and subtle estimates. The key difficulties in the variational approach are the non-standard shape of the nonlinearity g in (1.14) and the lack of compactness due to the unboundedness of \mathbb{R}_+^N . To overcome these difficulties, we need new estimates within the analysis of Cerami sequences and for comparing the mountain pass energy value for E with the corresponding one of the limit energy functional

$$(1.17) \quad E_\infty : H^1(\mathbb{R}^N) \rightarrow \mathbb{R}, \quad E_\infty(u) = \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + u^2) dx - \frac{1}{p+1} \int_{\mathbb{R}^N} |u|^{p+1} dx.$$

In particular, we shall use the asymptotic decay properties of the unique positive radial solution to (1.1) in order to build suitable test functions.

We now comment on the proofs of the non-existence part (ii) of Theorem 1.2. We argue by contradiction and use a suitable modification of the so-called sliding method introduced by H. Berestycki and L. Nirenberg and further developed by H. Berestycki, L. Caffarelli and L. Nirenberg among others. Specifically, our proofs are inspired by [15, Section 2] and [7, Section 4].

We finally comment on the boundedness of positive solutions to (1.3)-(1.4). As stated in the following proposition, all the positive solutions to (1.3)-(1.4) are bounded in the case where $1 < p < 2^* - 1$. Hence, the fact that we are considering bounded solutions is not a restriction in this case.

Proposition 1.5. *Let $N \geq 2$, $1 < p < 2^* - 1$ and $c > 0$. Any positive $v \in C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N})$ -solution to (1.3)-(1.4) belongs to $L^\infty(\mathbb{R}_+^N)$.*

The proof of Proposition 1.5 follows by a rather standard blow up argument based on the doubling lemma by P. Poláčik, P. Quittner and P. Souplet in [26]. For the convenience of the reader, we include the proof in Section 5 below.

Organization of the paper. In Section 2, we collect estimates related to the nonlinearity g in (1.13) and the functional E associated with (1.14). With the help of these estimates, we establish the mountain pass geometry of E in Section 3, and we show that Cerami sequences at nontrivial energy levels are bounded and admit nontrivial weak limits after suitable translation. In Section 4, we then prove a key energy estimate which shows that, in dimensions $N \geq 2$, the mountain pass energy of the functional E is strictly smaller than the corresponding one for the limit energy functional E_∞ given in (1.17). With the help of this energy estimate, we then complete the proof of Theorem 1.3 in Section 5. Finally, we give the proof of Theorem 1.2 (ii) in Section 6.

Notation. For $1 \leq p < \infty$, we let $\|\cdot\|_{L^p(\mathbb{R}_+^N)}$ denote the standard norm on the usual Lebesgue space $L^p(\mathbb{R}_+^N)$. The Sobolev space $H_0^1(\mathbb{R}_+^N)$ is endowed with the standard norm

$$\|u\|^2 = \int_{\mathbb{R}_+^N} (|\nabla u|^2 + |u|^2) dx.$$

Also, for a function v , we define $v^+ := \max\{v, 0\}$ and $v^- := \max\{-v, 0\}$ and we write $x = (x', x_N)$ for $x \in \mathbb{R}_+^N$ with $x' \in \mathbb{R}^{N-1}$. We denote by $' \rightarrow'$, respectively by $' \rightharpoonup'$, the strong convergence, respectively the weak convergence in corresponding space and denote by $B_R(x)$ the open ball in \mathbb{R}^N of center x and radius $R > 0$. Also, we shall denote by $C_i > 0$ different constants which may vary from line to line but are not essential to the analysis of the

problem. Finally, at various places, we have to distinguish the cases $p \leq 2$ and $p > 2$. For this it is convenient to introduce the special constant

$$1_{\{p>2\}} := \begin{cases} 0, & p \leq 2, \\ 1, & p > 2. \end{cases}$$

Acknowledgements. Part of this work was done while the first author was visiting the Goethe-Universität Frankfurt. He wishes to thank his hosts for the warm hospitality and the financial support.

2. PRELIMINARIES

In this section we collect some estimates related to the transformed nonlinearity g defined in (1.13), its primitive G and the functional E defined in (1.15). For this we fix, throughout Sections 2–Section 5, $c \in (0, c_p)$, $p \in (1, 2^*)$, and we let u_c be given in (1.9). We recall that we have the uniform estimate

$$(2.1) \quad 0 \leq u_c \leq c_p \quad \text{in } \mathbb{R}_+^N.$$

We start with an elementary inequality for nonnegative real numbers which will be used in the energy estimates in Section 4 below.

Lemma 2.1. *For every $q > 2$ there exists $\kappa_q > 0$ with*

$$(2.2) \quad (a+b)^q - a^q - b^q \geq qa^{q-1}b + \kappa_q ab^{q-1} \quad \text{for all } a, b \geq 0.$$

Remark 2.1. If $q \geq 3$, then (2.2) holds in symmetric form with $\kappa_q = q$, see e.g. [19, Theorem 1]. If $q \in (2, 3)$, it is easy to see that one has to choose $\kappa_q < q$.

Proof of Lemma 2.1. We first note that, since $q-1 > 1$, we have, by convexity of the function $\tau \mapsto (1+\tau)^{q-1}$,

$$(2.3) \quad (s+t)^{q-1} - (s^{q-1} + t^{q-1}) = s^{q-1} \left[\left(1 + \frac{t}{s}\right)^{q-1} - \left(1 + \left(\frac{t}{s}\right)^{q-1}\right) \right] \geq s^{q-1} \left((q-1) \frac{t}{s} - \left(\frac{t}{s}\right)^{q-1} \right) = (q-1)ts^{q-2} - t^{q-1}$$

for $s \geq t > 0$. Now, to prove the claim, it suffices to consider $a, b > 0$, since the inequality holds trivially if $a = 0$ or $b = 0$. Moreover, it suffices to prove that the inequality holds for $b \geq a > 0$ with some $\kappa_q \in (0, q]$, since then it also follows for arbitrary $a, b > 0$. For fixed $a > 0$, we consider the function

$$\ell : [0, \infty) \rightarrow \mathbb{R}, \quad \ell(t) = (a+t)^q - a^q - t^q - qa^{q-1}t$$

Then we have $\ell(0) = 0$ and

$$\ell'(t) = q \left[(a+t)^{q-1} - (a^{q-1} + t^{q-1}) \right]$$

Consequently, by (2.3) we have, for $b \geq a$,

$$\begin{aligned} \ell(b) &= \int_0^a \ell'(t) dt + \int_a^b \ell'(t) dt \geq q \left[\int_0^a ((q-1)ta^{q-2} - t^{q-1}) dt + \int_a^b ((q-1)at^{q-2} - a^{q-1}) dt \right] \\ &= qa(\kappa_{q,1}a^{q-1} + b^{q-1} - a^{q-2}b) \quad \text{with} \quad \kappa_{q,1} := \frac{q-1}{2} - \frac{1}{q} > 0. \end{aligned}$$

Since, by Young's inequality,

$$a^{q-2}b \leq \frac{q-2}{q-1}a^{q-1} + \frac{1}{q-1}b^{q-1},$$

we deduce that

$$\ell(b) \geq qa \left[\left(\kappa_{q,1} - \frac{q-2}{q-1} \right) a^{q-1} + \frac{q-2}{q-1} b^{q-1} \right].$$

If $\kappa_{q,1} \geq \frac{q-2}{q-1}$, we conclude that $\ell(b) \geq \frac{q(q-2)}{q-1} ab^{q-1}$. On the other hand, if $0 < \kappa_{q,1} < \frac{q-2}{q-1}$, we use again that $b \geq a$ and conclude that

$$\ell(b) \geq qa \left[\left(\kappa_{q,1} - \frac{q-2}{q-1} \right) b^{q-1} + \frac{q-2}{q-1} b^{q-1} \right] = q\kappa_{q,1} ab^{q-1}.$$

Hence, (2.2) holds for $b \geq a > 0$ with $\kappa_q = q \min\{\frac{q-2}{q-1}, \kappa_{q,1}\} \in (0, q)$. The proof is finished. \square

Next we provide basic but important estimates for the nonlinearity g defined in (1.13) and its primitive G .

Lemma 2.2.(i) For $(x, s) \in \mathbb{R}_+^N \times \mathbb{R}$ we have

$$(2.4) \quad 0 \leq g(x, s) - s^+ p u_c(x)^{p-1} \leq C_{1,p} [s^+]^p + 1_{\{p>2\}} C_{2,p} [s^+]^2,$$

and

$$(2.5) \quad 0 \leq G(x, s) - \frac{p}{2} [s^+]^2 u_c(x)^{p-1} \leq \frac{C_{1,p}}{p+1} [s^+]^{p+1} + 1_{\{p>2\}} \frac{C_{2,p}}{3} [s^+]^3,$$

with $C_{1,p} := 1 + 2^{p-3} p(p-1)$ and $C_{2,p} := p(p-1) 2^{p-3} c_p^{p-2}$.

(ii) Let

$$H(x, s) := \frac{1}{2} g(x, s) s - G(x, s) \quad \text{for } x \in \mathbb{R}_+^N, s \in \mathbb{R}.$$

Then we have

$$(2.6) \quad H(x, s) \geq \max \left\{ 0, \frac{p-1}{2(p+1)} [s^+]^{p+1} - u_c(x)^{p-1} D_{1,p} [s^+]^2 - u_c(x) 1_{\{p>2\}} D_{2,p} [s^+]^p \right\}$$

with $D_{1,p} := \frac{p(p-1)}{p+1} (1 + 2^{p-2})$ and $D_{2,p} := \frac{p 2^{p-2}}{p+1}$.Moreover, the function $H(x, \cdot)$ is non-decreasing in s for every $x \in \mathbb{R}_+^N$.

Remark 2.2. The constants $C_{i,p}$ and $D_{i,p}$, $i = 1, 2$, are not optimal. However, this choice simplifies the presentation. Moreover, they do not play an important role in our proofs below.

Proof of Lemma 2.2. (i) Since $g(\cdot, s) \equiv 0$ and $G(\cdot, s) \equiv 0$ for $s \leq 0$, it suffices to consider $s > 0$. Fix $x \in \mathbb{R}_+^N$. Since $g(x, \cdot)$ is of class C^1 on $(0, \infty)$, we have

$$(2.7) \quad g(x, s) = (u_c(x) + s)^p - u_c(x)^p = s p u_c(x)^{p-1} + p \int_0^s [(u_c(x) + \tau)^{p-1} - u_c(x)^{p-1}] d\tau, \quad \text{for } s > 0.$$

We now distinguish two cases. If $p \in (1, 2]$, we have

$$0 \leq (a + \tau)^{p-1} - a^{p-1} \leq \tau^{p-1} \quad \text{for } \tau > 0, a \geq 0,$$

and therefore, if $p \in (1, 2]$,

$$(2.8) \quad 0 \leq g(x, s) - s p u_c(x)^{p-1} \leq p \int_0^s \tau^{p-1} d\tau \leq s^p, \quad \text{for } s \geq 0.$$

If $p > 2$, we have

$$0 \leq (a + \tau)^{p-1} - a^{p-1} \leq (p-1) \tau (a + \tau)^{p-2}, \quad \text{for } \tau > 0, a \geq 0,$$

by the convexity of the function $a \mapsto (a + \tau)^{p-1}$ and therefore, using also (2.1),

$$\begin{aligned} 0 \leq g(x, s) - s p u_c(x)^{p-1} &\leq p \int_0^s [(u_c(x) + \tau)^{p-1} - u_c(x)^{p-1}] d\tau \\ &\leq p(p-1) \int_0^s \tau (u_c(x) + \tau)^{p-2} d\tau \leq \frac{p(p-1)}{2} s^2 (u_c(x) + s)^{p-2} \leq \frac{p(p-1)}{2} s^2 (c_p + s)^{p-2}. \end{aligned}$$

Note also that, since $p > 2$,

$$(2.9) \quad (c_p + s)^{p-2} \leq \left(2 \max\{c_p, s\} \right)^{p-2} \leq (2c_p)^{p-2} + (2s)^{p-2}, \quad \text{for } s \geq 0.$$

Consequently, if $p > 2$,

$$(2.10) \quad 0 \leq g(x, s) - s p u_c(x)^{p-1} \leq 2^{p-3} p(p-1) s^p + p(p-1) 2^{p-3} c_p^{p-2} s^2, \quad \text{for } s \geq 0.$$

Now (2.4) follows by combining (2.8) and (2.10). Moreover, (2.5) follows by integrating (2.4).

(ii) We first note that $H(x, s) \equiv 0$ for all $s \leq 0$. Thus, we just have to prove the result for $s > 0$. Directly observe that, for all $x \in \mathbb{R}_+^N$, we have $H(x, \cdot) \in C^1(\mathbb{R})$ and

$$\begin{aligned} \frac{\partial}{\partial s} H(x, s) &= \frac{p}{2}(u_c(x) + s)^{p-1}s + \frac{1}{2}(u_c(x) + s)^p - \frac{1}{2}u_c^p(x) - (u_c(x) + s)^p + u_c^p(x) \\ &= \frac{1}{2} \left[p(u_c(x) + s)^{p-1}s - (u_c(x) + s)^p + u_c^p(x) \right] \quad \text{for } (x, s) \in \mathbb{R}_+^N \times (0, +\infty). \end{aligned}$$

On the other hand, since $p > 1$, we have, by the mean value theorem,

$$(u_c(x) + s)^p - u_c^p(x) \leq p(u_c(x) + s)^{p-1}s \quad \text{for } (x, s) \in \mathbb{R}_+^N \times (0, +\infty).$$

Hence $\frac{\partial}{\partial s} H(x, s) \geq 0$ for all $s > 0$, so the function $H(x, \cdot)$ is non-decreasing in $s \in [0, \infty)$. This also implies that $H(x, s) \geq 0$ for $s \geq 0$. It thus remains to prove (2.6) for $s \geq 0$. For this we first note that

$$\begin{aligned} H(x, s) &= \frac{g(x, s)s}{2} - G(x, s) = \frac{1}{2} \left((u_c(x) + s)^p - u_c(x)^p \right) s - \frac{1}{p+1} \left((u_c(x) + s)^{p+1} - u_c(x)^{p+1} - (p+1)u_c(x)^p s \right) \\ &= \left(\frac{1}{2} - \frac{1}{p+1} \right) \left((u_c(x) + s)^p - u_c(x)^p \right) s - \frac{1}{p+1} \left((u_c(x) + s)^p - u_c(x)^p - pu_c(x)^{p-1}s \right) u_c(x) \\ &\geq \frac{p-1}{2(p+1)} s^{p+1} - \frac{1}{p+1} u_c(x) \left(g(x, s) - spu_c(x)^{p-1} \right). \end{aligned}$$

It therefore remains to show that

$$(2.11) \quad u_c(x) \left(g(x, s) - spu_c^{p-1}(x) \right) \leq (p+1) \left(u_c^{p-1}(x) D_{1,p} s^2 + u_c(x) 1_{\{p>2\}} D_{2,p} s^p \right), \quad \text{for } s > 0.$$

By (2.7) and integration by parts we have

$$\begin{aligned} u_c(x) \left(g(x, s) - spu_c^{p-1}(x) \right) &= pu_c(x) \int_0^s \left[(u_c(x) + \tau)^{p-1} - u_c(x)^{p-1} \right] d\tau \\ (2.12) \quad &= p(p-1)u_c(x) \int_0^s (s-\tau) \left[(u_c(x) + \tau)^{p-2} \right] d\tau. \end{aligned}$$

If $p \in (1, 2]$, we have $(u_c(x) + \tau)^{p-2} \leq u_c(x)^{p-2}$ for $\tau > 0$ and therefore

$$(2.13) \quad u_c(x) \left(g(x, s) - spu_c(x)^{p-1} \right) \leq p(p-1)u_c^{p-1}(x) \int_0^s (s-\tau) d\tau \leq p(p-1)u_c^{p-1}(x)s^2.$$

If $p > 2$, arguing as (2.9), we have

$$(u_c(x) + \tau)^{p-2} \leq 2^{p-2} \left(u_c^{p-2}(x) + \tau^{p-2} \right), \quad \text{for } \tau \geq 0,$$

and therefore (2.12) yields

$$\begin{aligned} u_c(x) \left(g(x, s) - spu_c(x)^{p-1} \right) &\leq 2^{p-2} p(p-1)u_c(x) \int_0^s (s-\tau) \left[u_c^{p-2}(x) + \tau^{p-2} \right] d\tau \\ (2.14) \quad &= 2^{p-2} p(p-1)u_c(x) \left(u_c^{p-2}(x) \int_0^s (s-\tau) d\tau + \int_0^s (s-\tau) \tau^{p-2} d\tau \right) \\ &\leq 2^{p-2} p(p-1)u_c^{p-1}(x)s^2 + 2^{p-2} pu_c(x)s^p, \quad \text{for } s > 0. \end{aligned}$$

Now (2.11) follows by combining (2.13) and (2.14). The proof is finished. \square

Remark 2.3.

- (a) From the growth estimates given in Lemma 2.2 (i) and the fact that g is continuous, it follows in a standard way that the functional E is well-defined on $H_0^1(\mathbb{R}_+^N)$ and of class C^1 .
- (b) Part (ii) of Lemma 2.2 will be useful in the analysis of Cerami sequences of the functional E , see Section 3 below.

Next, we consider then the quadratic form $q_c : H_0^1(\mathbb{R}_+^N) \rightarrow \mathbb{R}$ given by

$$(2.15) \quad q_c(u) := \int_{\mathbb{R}_+^N} \left(|\nabla u|^2 + V_c(x)u^2 \right) dx,$$

with

$$(2.16) \quad V_c \in L^\infty(\mathbb{R}_+^N), \quad V_c(x) = 1 - pu_c^{p-1}(x_N).$$

As we show in the following lemma, q_c is positive definite on $H_0^1(\mathbb{R}_+^N)$.

Proposition 2.3. *We have*

$$(2.17) \quad \tilde{q}_c := \inf_{u \in H_0^1(\mathbb{R}_+^N) \setminus \{0\}} \frac{q_c(u)}{\|u\|^2} > 0.$$

Proof. Since $V_c \in L^\infty(\mathbb{R}_+^N)$, it suffices to show there exists $C > 0$ such that

$$(2.18) \quad q_c(u) \geq C\|u\|_{L^2(\mathbb{R}_+^N)}^2 \quad \text{for all } u \in H_0^1(\mathbb{R}_+^N).$$

Indeed, if (2.18) holds, then for $\delta \in (0, 1)$ we have

$$q_c(u) \geq \delta q_c(u) + (1 - \delta)C\|u\|_{L^2(\mathbb{R}_+^N)}^2 \geq \delta\|u\|^2 + \left[(1 - \delta)C - \delta(1 + \|V_c\|_{L^\infty(\mathbb{R}_+^N)}) \right] \|u\|_{L^2(\mathbb{R}_+^N)}^2.$$

Choosing δ sufficiently small, we have $(1 - \delta)C - \delta(1 + \|V_c\|_{L^\infty(\mathbb{R}_+^N)}) \geq 0$ and therefore (2.17) holds with $\tilde{q}_c \geq \delta$. To show (2.18), we first consider the case $N = 1$. Arguing by contradiction, we assume that

$$\lambda := \inf \{ q_c(u) : u \in H_0^1(\mathbb{R}_+) \text{ and } \|u\|_{L^2(\mathbb{R}_+)} = 1 \} < 0,$$

(note that $\lambda > -\infty$ since $V_c \in L^\infty(\mathbb{R}_+)$). Then, there exists a sequence $(u_n)_n$ such that $\|u_n\|_{L^2(\mathbb{R}_+)} = 1$ for all $n \in \mathbb{N}$ and $q_c(u_n) \rightarrow \lambda$ as $n \rightarrow \infty$. Hence, $(u_n)_n$ is a bounded sequence in $H_0^1(\mathbb{R}_+)$, and thus $u_n \rightharpoonup u_*$ weakly in $H_0^1(\mathbb{R}_+)$ after passing to a subsequence. Moreover, with $v_n := u_n - u_*$, we have $v_n \rightarrow 0$ in $H_0^1(\mathbb{R}_+)$ and therefore $v_n \rightarrow 0$ in $L_{loc}^2(\mathbb{R})$. Since $V_c(t) \rightarrow 1$ as $t \rightarrow \infty$, this implies that

$$q_c(v_n) \geq \int_{\mathbb{R}_+} V_c(t)v_n^2 dt \geq \|v_n\|_{L^2(\mathbb{R}_+)}^2 + o(1), \quad \text{as } n \rightarrow \infty,$$

and therefore

$$\begin{aligned} \lambda + o(1) &= q_c(u_n) = q_c(u_*) + q_c(v_n) + o(1) \\ &\geq \lambda\|u_*\|_{L^2(\mathbb{R}_+)}^2 + \|v_n\|_{L^2(\mathbb{R}_+)}^2 + o(1) = \lambda(\|u_*\|_{L^2(\mathbb{R}_+)}^2 + \|v_n\|_{L^2(\mathbb{R}_+)}^2) + (1 - \lambda)\|v_n\|_{L^2(\mathbb{R}_+)}^2 + o(1) \\ &= \lambda + (1 - \lambda)\|v_n\|_{L^2(\mathbb{R}_+)}^2 + o(1) \end{aligned}$$

It thus follows that $v_n \rightarrow 0$ in $L^2(\mathbb{R}_+)$ and hence $u_n \rightarrow u_*$ in $L^2(\mathbb{R}_+)$, which yields that $\|u_*\|_{L^2} = 1$. Moreover, by weak lower semicontinuity of q_c and the definition of λ , it follows that $q_c(u_*) = \lambda$, so u_* is a constrained minimizer for q_c . A standard argument (based on replacing u_* by $|u_*|$) shows that $u_* \in H_0^1(\mathbb{R}_+)$ is a positive or negative solution of

$$-u_*'' + V_c(t)u_* = \lambda u_* \quad \text{in } \mathbb{R}_+, \quad u_*(0) = 0.$$

Without loss of generality, we may assume that u_* is positive, which implies that $u_*'(0) > 0$. We also recall that $w_* := -u_*'$ satisfies

$$-w_*'' + V_c(t)w_* = 0 \quad \text{in } \mathbb{R}_+, \quad w_* > 0 \quad \text{in } \overline{\mathbb{R}_+}.$$

Consequently, we have

$$0 < -\lambda \int_{\mathbb{R}_+} w_* u_* dx = \int_{\mathbb{R}_+} (w_* u_*'' - u_* w_*'') dx = -w_*(0)u_*'(0) < 0,$$

a contradiction. Hence, we conclude that (2.18) holds in the case $N = 1$. To show (2.18) for general $N \geq 2$, we remark that, by density, we only have to show it for $u \in C_c^\infty(\mathbb{R}_+^N)$. For any such function we then have, writing $x = (x', t) \in \mathbb{R}_+^N$ with $x' \in \mathbb{R}^{N-1}$, $t > 0$:

$$\begin{aligned} q_c(u) &\geq \int_{\mathbb{R}_+^N} (|\partial_t u|^2 + V_c u^2) dx \\ &= \int_{\mathbb{R}^{N-1}} \int_{\mathbb{R}_+} (|\partial_t u(x', t)|^2 + V_c u^2(x', t)) dt dx' \end{aligned}$$

$$\geq C \int_{\mathbb{R}^{N-1}} \int_{\mathbb{R}_+} u^2(x', t) dt dx' = C \|u\|_{L^2(\mathbb{R}_+^N)}^2.$$

Here we have used the result in the case $N = 1$ and the fact that $u(x', \cdot) \in C_c^\infty(\mathbb{R}_+) \subset H_0^1(\mathbb{R}_+)$ for every $x' \in \mathbb{R}^{N-1}$. We thus have proved (2.18) for general $N \geq 1$, and the proof is complete. \square

Remark 2.4. From Proposition 2.3 it follows that $(q_c(\cdot))^{1/2}$ is an equivalent norm to $\|\cdot\|$ in $H_0^1(\mathbb{R}_+^N)$.

Having at hand Proposition 2.3, we prove a lower estimate on the functional E given in (1.15) that will be useful at several points below.

Corollary 2.4. *We have*

$$E(u) \geq \frac{\tilde{q}_c}{2} \|u\|^2 - \frac{C_{1,p}}{p+1} \|u\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} \frac{C_{2,p}}{3} \|u\|_{L^3(\mathbb{R}_+^N)}^3 \quad \text{for all } u \in H_0^1(\mathbb{R}_+^N).$$

with \tilde{q}_c given in (2.17) and $C_{1,p}, C_{2,p}$ given in Lemma 2.2 (i).

Proof. For all $u \in H_0^1(\mathbb{R}_+^N)$, we have, by (2.5) and Proposition 2.3,

$$\begin{aligned} E(u) &= \frac{1}{2} \|u\|^2 - \int_{\mathbb{R}_+^N} G(x, u) dx \\ &\geq \frac{1}{2} \|u\|^2 - \frac{1}{2} \int_{\mathbb{R}_+^N} p u_c^{p-1} (u^+)^2 dx - \frac{C_{1,p}}{p+1} \|u^+\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} \frac{C_{2,p}}{3} \|u^+\|_{L^3(\mathbb{R}_+^N)}^3 \\ &\geq \frac{1}{2} q_c(u) - \frac{C_{1,p}}{p+1} \|u\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} \frac{C_{2,p}}{3} \|u\|_{L^3(\mathbb{R}_+^N)}^3 \\ &\geq \frac{\tilde{q}_c}{2} \|u\|^2 - \frac{C_{1,p}}{p+1} \|u\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} \frac{C_{2,p}}{3} \|u\|_{L^3(\mathbb{R}_+^N)}^3. \end{aligned}$$

\square

3. MOUNTAIN-PASS GEOMETRY AND BOUNDEDNESS OF THE CERAMI SEQUENCES

This section is devoted to show that the functional E has a Mountain-pass geometry and that, for any $d \in \mathbb{R}$, the Cerami sequences for E and level d are bounded. We keep using the notation of the introduction and of Section 2, which depends on the fixed quantities $c \in (0, c_p)$ and $p \in (1, 2^*)$. We begin by proving that the functional E has indeed a Mountain-pass geometry.

Lemma 3.1. *The functional E has the following properties.*

- (i) $E(0) = 0$.
- (ii) *There exist $\rho_0 > 0$ and $\delta_0 > 0$ such that $E(u) \geq \delta_0$ for all $u \in H_0^1(\mathbb{R}_+^N)$ such that $\|u\| = \rho_0$.*
- (iii) *There exists $\psi \in H_0^1(\mathbb{R}_+^N)$ such that $\|\psi\|_{H_0^1(\mathbb{R}_+^N)} > \rho_0$ and $E(\psi) < 0$.*

Proof. Since (i) is obvious, we concentrate on proving (ii) and (iii). We first prove (ii). Let $u \in H_0^1(\mathbb{R}_+^N)$ with $\|u\| = \rho_0$. By Corollary 2.4, we have

$$E(u) \geq \frac{\tilde{q}_c}{2} \rho_0^2 - \frac{C_{1,p}}{p+1} \|u\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} \frac{C_{2,p}}{3} \|u\|_{L^3(\mathbb{R}_+^N)}^3.$$

Applying then Sobolev embeddings, we deduce that

$$E(u) \geq \frac{\tilde{q}_c}{2} \rho_0^2 - C \left(\rho_0^{p+1} + 1_{\{p>2\}} \rho_0^3 \right),$$

with a constant $C > 0$. Here we point out that $3 \leq p+1 < 2^*$ if $p > 2$. Since $p > 1$, Claim (ii) follows by taking ρ_0 sufficiently small. It then remains to prove (iii). Let $\varphi \in C_c^\infty(\mathbb{R}_+^N)$ with $\varphi \not\equiv 0$ and $\psi := t\varphi$ with $t \in (0, +\infty)$. Directly observe that

$$E(\psi) = \frac{t^2}{2} \|\varphi\|^2 + t \int_{\mathbb{R}_+^N} u_c^p \varphi dx - \frac{1}{p+1} \int_{\mathbb{R}_+^N} ((u_c + t\varphi)^{p+1} - u_c^{p+1}) dx$$

Then, since

$$\int_{\mathbb{R}_+^N} \left((u_c + t\varphi)^{p+1} - u_c^{p+1} \right) dx \geq t^{p+1} \|\varphi\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1},$$

we have that

$$E(\psi) \leq \frac{t^2}{2} \|\varphi\|^2 + tc^p \|\varphi\|_{L^1(\mathbb{R}_+^N)} - \frac{t^{p+1}}{p+1} \|\varphi\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1}.$$

Claim (iii) follows taking t sufficiently large and thus the proof is complete. \square

We now prove the boundedness of Cerami sequences of the functional E .

Proposition 3.2. *Cerami sequences for E at any level $d \in \mathbb{R}$ are bounded.*

Remark 3.1.

(a) Recall that $(\varphi_n)_n \subset H_0^1(\mathbb{R}_+^N)$ is a Cerami sequence for E at level $d \in \mathbb{R}$ if

$$E(\varphi_n) \rightarrow d \quad \text{and} \quad (1 + \|\varphi_n\|) \|E'(\varphi_n)\|_{H^{-1}(\mathbb{R}_+^N)} \rightarrow 0.$$

(b) The proof of Proposition 3.2 is inspired by [20, Section 3]. However, since our problem is not invariant under translations in \mathbb{R}^N and our nonlinearity g has a non-standard shape, several difficulties appear.

Proof of Proposition 3.2. Let $d \in \mathbb{R}$ be an arbitrary but fixed constant and let $(u_n)_n \subset H_0^1(\mathbb{R}_+^N)$ be a Cerami sequence for E at level $d \in \mathbb{R}$. First of all, observe that

$$\|u_n^-\|^2 = -\langle E'(u_n), u_n^- \rangle \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

In particular, we deduce that $(u_n^-)_n$ is bounded. It then remains to prove that $(u_n^+)_n$ is bounded. We assume by contradiction that $\|u_n\| \rightarrow \infty$ and we set $v_n := u_n / \|u_n\|$ for all $n \in \mathbb{N}$. Since $(v_n)_n$ and $(u_n^-)_n$ are bounded, up to a subsequence if necessary, we have

$$(3.1) \quad v_n \rightharpoonup v \text{ in } H_0^1(\mathbb{R}_+^N), \quad v_n \rightarrow v \text{ in } L_{loc}^q(\mathbb{R}_+^N) \text{ for } 1 \leq q < 2^* \quad \text{and} \quad v_n \rightarrow v \text{ a.e. in } \mathbb{R}_+^N,$$

for some $v \in H_0^1(\mathbb{R}_+^N)$ with $v \geq 0$. We now consider separately two different cases:

Case 1 (Vanishing): For all $R > 0$, it follows that

$$(3.2) \quad \lim_{n \rightarrow \infty} \sup_{y \in \mathbb{R}_+^N} \int_{B_R(y) \cap \mathbb{R}_+^N} v_n^2 dx = 0.$$

Case 2 (Non-vanishing): There exist $R > 0$, $\delta > 0$ and a sequence of points $(y^n)_n \subset \mathbb{R}_+^N$ such that

$$(3.3) \quad \lim_{n \rightarrow \infty} \int_{B_R(y^n) \cap \mathbb{R}_+^N} v_n^2 dx \geq \delta.$$

Case 1 (Vanishing): First of all, observe that, by (3.2) and Lions' Lemma [22, Lemma I.1], $v_n \rightarrow 0$ in $L^q(\mathbb{R}_+^N)$ for all $2 < q < 2^*$, and so, by uniqueness of the limit we have $v \equiv 0$. We define then the sequence $(z_n)_n \subset H_0^1(\mathbb{R}_+^N)$ by $z_n := t_n u_n$ with $t_n \in [0, 1]$ satisfying

$$E(z_n) = \max_{t \in [0, 1]} E(tu_n),$$

(if, for $n \in \mathbb{N}$, t_n is not unique, we choose the smallest value) and we split the proof in the vanishing case (Case 1) into three steps.

Step 1.1: $\lim_{n \rightarrow \infty} E(z_n) = +\infty$.

We argue by contradiction. Suppose there exists $M < +\infty$ such that

$$\liminf_{n \rightarrow \infty} E(z_n) \leq M,$$

and define $(k_n)_n \subset H_0^1(\mathbb{R}_+^N)$ as

$$k_n := \left(\frac{4M}{q_c} \right)^{\frac{1}{2}} v_n = \left(\frac{4M}{q_c} \right)^{\frac{1}{2}} \frac{1}{\|u_n\|} u_n, \quad \text{for all } n \in \mathbb{N},$$

where $\tilde{q}_c > 0$ is the constant given by Proposition 2.3. First, observe that

$$(3.4) \quad k_n \rightarrow 0 \text{ in } H_0^1(\mathbb{R}_+^N), \quad k_n \rightarrow 0 \text{ in } L^q(\mathbb{R}_+^N), \text{ for } 2 < q < 2^*, \quad \text{and} \quad k_n \rightarrow 0 \text{ a.e. in } \mathbb{R}_+^N.$$

Then, by Corollary 2.4 and (3.4), we obtain that

$$(3.5) \quad E(k_n) \geq \frac{\tilde{q}_c}{2} \|k_n\|^2 - C \left(\|k_n\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} + 1_{\{p>2\}} \|k_n\|_{L^3(\mathbb{R}_+^N)}^3 \right) = 2M + o(1).$$

Taking M bigger if necessary, we have that, for all $n \in \mathbb{N}$ large enough,

$$E(k_n) > \frac{3}{2}M.$$

On the other hand, observe that, for $n \in \mathbb{N}$ large enough, $\left(\frac{4M}{\tilde{q}_c}\right)^{\frac{1}{2}} \frac{1}{\|u_n\|} \in [0, 1]$. Hence, we have that

$$\frac{3}{2}M \leq \liminf_{n \rightarrow \infty} E(k_n) \leq \liminf_{n \rightarrow \infty} E(z_n) \leq M,$$

which is a contradiction. Thus, the Step 1.1 follows.

Step 2.1: $\langle E'(z_n), z_n \rangle = 0$ for all $n \in \mathbb{N}$ large enough.

By Step 1.1 we know that $E(z_n) \rightarrow \infty$ as $n \rightarrow \infty$. On the other hand, $E(0) = 0$ and $E(u_n) \rightarrow d$ as $n \rightarrow \infty$. Hence, for $n \in \mathbb{N}$ large enough, $t_n \in (0, 1)$ and so, by the definition of z_n , the Step 2.1 follows.

Step 3.1: Conclusion Case 1.

Observe that, by Step 2.1, for all $n \in \mathbb{N}$ large enough,

$$E(z_n) = E(z_n) - \frac{1}{2} \langle E'(z_n), z_n \rangle = \int_{\mathbb{R}_+^N} H(x, z_n) dx,$$

where H is given in Lemma 2.2 (ii). By Step 1.1, we have that

$$(3.6) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}_+^N} H(x, z_n) dx = +\infty.$$

On the other hand, since $(u_n)_n$ is a Cerami sequence,

$$d + o(1) = E(u_n) - \frac{1}{2} \langle E'(u_n), u_n \rangle = \int_{\mathbb{R}_+^N} H(x, u_n) dx.$$

Then, using the definition of z_n and the fact that $H(x, s)$ is non-decreasing in s by Lemma 2.2 (ii), we obtain that

$$\int_{\mathbb{R}_+^N} H(x, z_n) dx \leq \int_{\mathbb{R}_+^N} H(x, u_n) dx = d + o(1),$$

which clearly contradicts (3.6). Hence, the proof of the result in the vanishing case (Case 1) is concluded.

Case 2 (Non-vanishing): We split the proof into two steps.

Step 1.2: There exists $M > 0$ such that $y_N^n := \text{dist}(y^n, \partial\mathbb{R}_+^N) \leq M$ for all $n \in \mathbb{N}$.

We assume by contradiction that $y_N^n \rightarrow +\infty$ as $n \rightarrow +\infty$. Then, for all $n \in \mathbb{N}$, we introduce $w_n := v_n(\cdot + y^n)$ and observe that

$$(3.7) \quad w_n \rightharpoonup w \text{ in } H^1(\mathbb{R}^N), \quad w_n \rightarrow w \text{ in } L_{loc}^q(\mathbb{R}^N) \text{ for } 1 \leq q < 2^*, \quad \text{and} \quad w_n \rightarrow w \text{ a.e. in } \mathbb{R}^N,$$

for some $w \in H^1(\mathbb{R}^N)$ with $w \not\equiv 0$ (by (3.3)) and $w \geq 0$. Now, observe that, since $(u_n)_n$ is a Cerami sequence, Lemma 2.2 (ii) implies that

$$\begin{aligned} o(1) &= \frac{1}{\|u_n\|^{p+1}} \left(E(u_n) - \frac{1}{2} \langle E'(u_n), u_n \rangle \right) = \frac{1}{\|u_n\|^{p+1}} \int_{\mathbb{R}_+^N} H(x, u_n(x)) dx \\ &\geq \frac{1}{\|u_n\|^{p+1}} \left[\frac{p-1}{2(p+1)} \|u_n^+\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - \int_{\mathbb{R}_+^N} \left(u_c^{p-1}(x) D_{1,p} [u_n^+]^2 + u_c(x) 1_{\{p>2\}} D_{2,p} [u_n^+]^p \right) dx \right] \\ &\geq \frac{1}{\|u_n\|^{p+1}} \left[\frac{p-1}{2(p+1)} \|u_n^+\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - \max\{c_p^{p-1}, c_p\} \left(D_{1,p} \|u_n^+\|_{L^2(\mathbb{R}_+^N)}^2 + 1_{\{p>2\}} D_{2,p} \|u_n^+\|_{L^p(\mathbb{R}_+^N)}^p \right) \right] \end{aligned}$$

$$\geq \frac{p-1}{2(p+1)} \|v_n^+\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - \frac{C}{\|u_n\|^{p+1}} (\|u_n\|^2 + 1_{\{p>2\}} \|u_n\|^p) \geq \frac{p-1}{2(p+1)} \|v_n^+\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} + o(1),$$

where $C > 0$ is a constant independent of n . Here we also used Sobolev embeddings and the fact that $\|u_n\| \rightarrow \infty$ as $n \rightarrow \infty$. Since $p > 1$, we thus conclude by Fatou's Lemma that

$$0 = \lim_{n \rightarrow \infty} \|v_n^+\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} = \liminf_{n \rightarrow \infty} \int_{\{x_N \geq -y_N^n\}} (w_n^+)^{p+1} dx \geq \int_{\mathbb{R}^N} (w^+)^p dx.$$

Hence $w = w^+ \equiv 0$, which clearly is a contradiction. Thus, Step 1.2 follows.

Step 2.2: Conclusion Case 2.

By Step 1.2 we know there exists $M > 0$ such that $y_N^n \leq M$ for all $n \in \mathbb{N}$. We then define, for all $n \in \mathbb{N}$, $\tilde{w}_n := v_n(\cdot + \xi_n)$, where $\xi_n = (y_1^n, \dots, y_{N-1}^n, 0)$. Again by (3.3), we have

$$(3.8) \quad \tilde{w}_n \rightharpoonup \tilde{w} \text{ in } H_0^1(\mathbb{R}_+^N), \quad \tilde{w}_n \rightarrow \tilde{w} \text{ in } L_{loc}^q(\mathbb{R}_+^N) \text{ for } 1 \leq q < 2^*, \quad \text{and} \quad \tilde{w}_n \rightarrow \tilde{w} \text{ a.e. in } \mathbb{R}_+^N,$$

for some $\tilde{w} \in H_0^1(\mathbb{R}_+^N)$ with $\tilde{w} \not\equiv 0$ and $\tilde{w} \geq 0$. For $n \in \mathbb{N}$, let $\varphi_n := \tilde{w}(\cdot - \xi_n) \in H_0^1(\mathbb{R}_+^N)$. Since $(u_n)_n$ is a Cerami sequence with $\|u_n\| \rightarrow \infty$ as $n \rightarrow \infty$, we have

$$\begin{aligned} o(1) &= \frac{\langle E'(u_n), \varphi_n \rangle}{\|u_n\|} = \frac{1}{\|u_n\|} \int_{\mathbb{R}_+^N} (\nabla u_n \nabla \varphi_n + u_n \varphi_n - g(x, u_n) \varphi_n) dx \\ &= \int_{\mathbb{R}_+^N} (\nabla v_n \nabla \varphi_n + v_n \varphi_n) dx - \int_{\mathbb{R}_+^N} \left(\frac{(u_c + u_n^+)^p - u_c^p}{\|u_n\|} \right) \varphi_n dx \\ &= \int_{\mathbb{R}_+^N} (\nabla \tilde{w}_n \nabla \tilde{w} + \tilde{w}_n \tilde{w}) dx - \int_{\mathbb{R}_+^N} \left(\frac{(u_c + \|u_n\| \tilde{w}_n^+)^p - u_c^p}{\|u_n\|} \right) \tilde{w} dx \\ &= \|\tilde{w}\|^2 + o(1) - \int_{\mathbb{R}_+^N} \left(\frac{(u_c + \|u_n\| \tilde{w}_n^+)^p - u_c^p}{\|u_n\|} \right) \tilde{w} dx. \end{aligned}$$

On the other hand, since $p > 1$, we have that

$$\liminf_{n \rightarrow \infty} \frac{(u_c + \|u_n\| \tilde{w}_n^+)^p - u_c^p}{\|u_n\| \tilde{w}_n^+} \tilde{w}_n^+ \tilde{w} = +\infty, \quad \text{a.e. in } \{\tilde{w} > 0\}$$

and therefore, since $\tilde{w} \geq 0$ and $\tilde{w} \not\equiv 0$,

$$\liminf_{n \rightarrow \infty} \int_{\mathbb{R}_+^N} \left(\frac{(u_c + u_n^+(\cdot + \xi_n))^p - u_c^p}{\|u_n\|} \right) \tilde{w} dx = \liminf_{n \rightarrow \infty} \int_{\{\tilde{w} > 0\}} \left(\frac{(u_c + u_n^+(\cdot + \xi_n))^p - u_c^p}{\|u_n\|} \right) \tilde{w} dx = +\infty$$

by Fatou's Lemma. This yields a contradiction. The proof of the result in the non-vanishing case (Case 2) is thus finished and the result follows. \square

Lemma 3.3. *Let $(u_n)_n$ be a Cerami sequence for E at level $d \in \mathbb{R} \setminus \{0\}$. Then, there exist $R > 0$, $\delta > 0$ and a sequence of points $(y^n)_n \subset \mathbb{R}_+^N$ such that*

$$\liminf_{n \rightarrow \infty} \int_{B_R(y^n) \cap \mathbb{R}_+^N} u_n^2 dx \geq \delta.$$

Proof. We assume by contradiction that, for all $R > 0$,

$$\liminf_{n \rightarrow \infty} \sup_{y \in \mathbb{R}_+^N} \int_{B_R(y) \cap \mathbb{R}_+^N} u_n^2 dx = 0.$$

Then, by Lions' [22, Lemma I.1], we have that $u_n \rightarrow 0$ in $L^q(\mathbb{R}_+^N)$ for all $2 < q < 2^*$. Now, since $(u_n)_n$ is a Cerami sequence, using Lemma 2.2 (i), we get

$$\begin{aligned} o(1) &= \langle E'(u_n), u_n \rangle = \|u_n\|^2 - \int_{\mathbb{R}_+^N} g(x, u_n(x)) u_n dx \\ &\geq \|u_n\|^2 - \int_{\mathbb{R}_+^N} p u_c^{p-1} (u_n^+)^2 dx - C_{1,p} \|u_n^+\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} C_{2,p} \|u_n^+\|_{L^3(\mathbb{R}_+^N)}^3 \\ &\geq q_c(u_n) - C_{1,p} \|u_n\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} C_{2,p} \|u_n\|_{L^3(\mathbb{R}_+^N)}^3 \end{aligned}$$

$$\geq \tilde{q}_c \|u_n\|^2 - C_{1,p} \|u_n\|_{L^{p+1}(\mathbb{R}_+^N)}^{p+1} - 1_{\{p>2\}} C_{2,p} \|u_n\|_{L^3(\mathbb{R}_+^N)}^3.$$

Hence, since $u_n \rightarrow 0$ in $L^q(\mathbb{R}_+^N)$ for all $2 < q < 2^*$, we deduce that $\|u_n\| \rightarrow 0$. Since E is continuous, this implies that $E(u_n) \rightarrow 0$ as $n \rightarrow \infty$, contradicting our assumption that $d \neq 0$. The proof is finished. \square

4. ENERGY ESTIMATES

We keep using the notation of the introduction and of Section 2, which depends on the fixed quantities $c \in (0, c_p)$ and $p \in (1, 2^*)$. Moreover, we will assume $N \geq 2$ throughout this section, which will be of key importance in order to derive the energy estimates we need. The mountain pass value associated to (1.14) is given by

$$(4.1) \quad b := \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} E(\gamma(t)),$$

where

$$\Gamma := \left\{ \gamma \in C([0,1], H_0^1(\mathbb{R}_+^N)) : \gamma(0) = 0, E(\gamma(1)) < 0 \right\}.$$

We note that $b > 0$ by Lemma 3.1. We also note that the functional E (given in (1.15)) can be written as

$$(4.2) \quad E(u) = E_\infty^+(u) - \frac{1}{p+1} \int_{\mathbb{R}_+^N} \left((u_c + u^+)^{p+1} - u_c^{p+1} - (u^+)^{p+1} - (p+1)u_c^p u^+ \right) dx,$$

where $E_\infty^+ : H_0^1(\mathbb{R}_+^N) \rightarrow \mathbb{R}$ is given by

$$E_\infty^+(u) = \frac{1}{2} \|u\|^2 - \frac{1}{p+1} \int_{\mathbb{R}_+^N} (u^+)^{p+1} dx.$$

Now, we introduce the auxiliary (limit) problem

$$(4.3) \quad -\Delta u + u = |u|^{p-1} u, \quad u \in H^1(\mathbb{R}^N),$$

and its associated energy $E_\infty : H^1(\mathbb{R}^N) \rightarrow \mathbb{R}$ given by

$$(4.4) \quad E_\infty(u) = \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + u^2) dx - \frac{1}{p+1} \int_{\mathbb{R}^N} |u|^{p+1} dx.$$

Also, we define

$$(4.5) \quad b_\infty := \inf_{\mathcal{K}_\infty} E_\infty, \quad \text{where} \quad \mathcal{K}_\infty := \left\{ u \in H^1(\mathbb{R}^N) \setminus \{0\} : E'_\infty(u) = 0 \right\}.$$

According to [8, Theorem 1], [6, Théorème 1] and [16, Theorem 2], there exists a ground-state solution $\psi \in C^2(\mathbb{R}^N)$ to (4.3) which is positive, radially symmetric, and such that

$$(4.6) \quad \psi(x) \leq C_{GS} |x|^{-\frac{N-1}{2}} e^{-|x|} \quad \text{and} \quad |\nabla \psi(x)| \leq C_{GS} |x|^{-\frac{N-1}{2}} e^{-|x|}, \quad \text{as } |x| \rightarrow \infty,$$

for some $C_{GS} > 0$ depending only on N and p . Moreover,

$$(4.7) \quad \psi \text{ is strictly decreasing in the radial variable.}$$

Let us also emphasize that

$$(4.8) \quad 0 < b_\infty = E_\infty(\psi) = \left(\frac{1}{2} - \frac{1}{p+1} \right) \|\psi\|_{L^{p+1}(\mathbb{R}^N)}^{p+1} = \frac{p-1}{2(p+1)} \|\psi\|_{L^{p+1}(\mathbb{R}^N)}^{p+1} = \frac{p-1}{2(p+1)} \inf_{u \in \mathcal{K}_\infty} \|u\|_{L^{p+1}(\mathbb{R}^N)}^{p+1}.$$

The aim of this section is to show, based on the assumption $N \geq 2$, that

$$(4.9) \quad b < b_\infty.$$

This strict inequality will be crucial to prove the existence result to (1.14) contained in Section 5. To this end, let us recall that $u_c(x) \sim e^{-x_N}$ as $x_N \rightarrow \infty$. More precisely, it follows from (1.6) and the definition of u_c that

$$(4.10) \quad m_{c,1} e^{-x_N} \leq u_c(x) \leq m_{c,2} e^{-x_N}, \quad \text{for } x \in \mathbb{R}_+^N, \quad \text{with} \quad m_{c,1} := c_p e^{-t_{c,p}}, \quad m_{c,2} := c_p 2^{\frac{2}{p-1}} e^{-t_{c,p}}.$$

Moreover, for $r > 0$, we introduce the function

$$x \mapsto \psi_r(x) := (\psi(x - r e_N) - \varepsilon_r)^+,$$

where $e_N = (0, \dots, 0, 1)$ is the N -th coordinate vector and $\varepsilon_r > 0$ is uniquely defined by (4.7) and the property that

$$\psi > \varepsilon_r \quad \text{in } B_r(0) \quad \text{and} \quad \psi \leq \varepsilon_r \quad \text{in } \mathbb{R}^N \setminus B_r(0).$$

We note that, as a consequence of (4.6), we have

$$(4.11) \quad \varepsilon_r \leq C_{GS} r^{-\frac{N-1}{2}} e^{-r}$$

We also note that $\psi_r \in H_0^1(\mathbb{R}_+^N)$ for every $r \geq 0$.

The rest of the section is devoted to prove the following result from which (4.9) immediately follows.

Proposition 4.1. *There exists $R > 0$ and $k > 0$ with the following properties:*

- i) $E(t\psi_r) < b_\infty$ for all $r \geq R$, $t \in [0, k]$.
- ii) $E(k\psi_r) < 0$ for all $r \geq R$.

We split the proof of this proposition into several lemmas.

Lemma 4.2. *There exists $C_1 > 0$ with*

$$(4.12) \quad E_\infty^+(t\psi_r) \leq b_\infty + C_1 e^{-r} r^{-\frac{N-1}{2}} t^{p+1} \quad \text{for all } t, r > 0.$$

Proof. Let $t, r > 0$. Directly observe that, by the definition of ψ_r ,

$$(4.13) \quad E_\infty^+(t\psi_r) = \frac{t^2}{2} \|\psi_r\|^2 - \frac{t^{p+1}}{p+1} \int_{\mathbb{R}^N} \psi_r^{p+1} dx = \frac{t^2}{2} \int_{B_r(0)} (|\nabla\psi|^2 + (\psi - \varepsilon_r)^2) dx - \frac{t^{p+1}}{p+1} \int_{B_r(0)} (\psi - \varepsilon_r)^{p+1} dx.$$

On the other hand, since ψ is a solution to (4.3) and $(\psi - \varepsilon_r)^+ \in H^1(\mathbb{R}^N)$, using $(\psi - \varepsilon_r)^+$ as test function in (4.3), we obtain that

$$\begin{aligned} \int_{B_r(0)} |\nabla\psi|^2 dx &= \int_{\mathbb{R}^N} \nabla\psi \nabla(\psi - \varepsilon_r)^+ dx \\ &= \int_{\mathbb{R}^N} (-\psi(\psi - \varepsilon_r)^+ + \psi^p(\psi - \varepsilon_r)^+) dx = \int_{B_r(0)} (-\psi(\psi - \varepsilon_r) + \psi^p(\psi - \varepsilon_r)) dx. \end{aligned}$$

Substituting the above identity into (4.13) and using the mean value theorem, we find that

$$\begin{aligned} E_\infty^+(t\psi_r) &= \frac{t^2}{2} \int_{B_r(0)} (-\psi(\psi - \varepsilon_r) + \psi^p(\psi - \varepsilon_r) + (\psi - \varepsilon_r)^2) dx - \frac{t^{p+1}}{p+1} \int_{B_r(0)} (\psi - \varepsilon_r)^{p+1} dx \\ &= -\frac{\varepsilon_r t^2}{2} \int_{B_r(0)} (\psi - \varepsilon_r) dx + \frac{1}{p+1} \int_{B_r(0)} (\psi - \varepsilon_r) \left[\frac{p+1}{2} t^2 \psi^p - t^{p+1} (\psi - \varepsilon_r)^p \right] dx \\ &\leq \frac{1}{p+1} \int_{B_r(0)} (\psi - \varepsilon_r) \left[\frac{p+1}{2} t^2 \psi^p - t^{p+1} (\psi - \varepsilon_r)^p \right] dx \\ &= \frac{1}{p+1} \left(\frac{p+1}{2} t^2 - t^{p+1} \right) \int_{B_r(0)} (\psi - \varepsilon_r) \psi^p dx + \frac{t^{p+1}}{p+1} \int_{B_r(0)} (\psi - \varepsilon_r) [\psi^p - (\psi - \varepsilon_r)^p] dx \\ &\leq \frac{p-1}{2(p+1)} \int_{B_r(0)} (\psi - \varepsilon_r) \psi^p dx + \frac{\varepsilon_r p t^{p+1}}{p+1} \int_{B_r(0)} (\psi - \varepsilon_r) \psi^{p-1} dx \\ &\leq \frac{p-1}{2(p+1)} \int_{\mathbb{R}^N} \psi^{p+1} dx + \frac{\varepsilon_r p t^{p+1}}{p+1} \int_{\mathbb{R}^N} (\psi - \varepsilon_r)^+ \psi^{p-1} dx. \end{aligned}$$

Using (4.8) and (4.11), we deduce that

$$E_\infty^+(t\psi_r) \leq b_\infty + \frac{\varepsilon_r p t^{p+1}}{p+1} \int_{\mathbb{R}^N} (\psi - \varepsilon_r)^+ \psi^{p-1} dx \leq b_\infty + \frac{p}{p+1} C_{GS} r^{-\frac{N-1}{2}} e^{-r} t^{p+1} \int_{\mathbb{R}^N} \psi^p dx.$$

Hence (4.12) holds with $C_1 = \frac{p C_{GS}}{p+1} \int_{\mathbb{R}^N} \psi^p dx$. □

Lemma 4.3. *There exists $R' \geq 1$ and $C_2 > 0$ with*

$$(4.14) \quad E_\infty^+(t\psi_r) - E(t\psi_r) \geq C_2 e^{-r} t^p \quad \text{for all } t > 0, r \geq R'.$$

Proof. Let $t > 0$ be arbitrary but fixed. Using Lemma 2.1 with $q = p + 1$, $\kappa := \kappa_q > 0$, the identity (4.2), the lower bound in (4.10) and the mean value theorem, we deduce that, for all $r \geq 1$,

$$\begin{aligned}
E_\infty^+(t\psi_r) - E(t\psi_r) &= \frac{1}{p+1} \int_{\mathbb{R}_+^N} \left((u_c + t\psi_r)^{p+1} - u_c^{p+1} - (t\psi_r)^{p+1} - (p+1)u_c^p t\psi_r \right) dx \geq \kappa t^p \int_{\mathbb{R}_+^N} u_c \psi_r^p dx \\
&= \kappa t^p \int_{\mathbb{R}_+^N} u_c \left((\psi(\cdot - re_N) - \varepsilon_r)^+ \right)^p dx = \kappa t^p \int_{B_r(0)} u_c(\cdot + re_N) (\psi - \varepsilon_r)^p dx \\
&\geq \kappa t^p \int_{B_r(0)} u_c(\cdot + re_N) (\psi^p - \varepsilon_r p \psi^{p-1}) dx \geq \kappa t^p \left(\int_{B_r^-(0)} u_c(\cdot + re_N) \psi^p dx - \varepsilon_r p c_p \int_{\mathbb{R}^N} \psi^{p-1} dx \right) \\
&\geq \kappa t^p \left(m_{c,1} e^{-r} \int_{B_r^-(0)} \psi^p dx - p c_p C_{GS} r^{-\frac{N-1}{2}} e^{-r} \int_{\mathbb{R}^N} \psi^{p-1} dx \right) \\
&\geq \kappa t^p e^{-r} \left(\frac{m_{c,1}}{2} \int_{B_r(0)} \psi^p dx - p c_p C_{GS} r^{-\frac{N-1}{2}} \int_{\mathbb{R}^N} \psi^{p-1} dx \right) \\
&\geq \kappa t^p e^{-r} \left(\frac{m_{c,1}}{2} \int_{B_1(0)} \psi^p dx - p c_p C_{GS} r^{-\frac{N-1}{2}} \int_{\mathbb{R}^N} \psi^{p-1} dx \right)
\end{aligned}$$

where we have set $B_r^-(0) := \{x \in B_r(0) : x_N \leq 0\}$. Since $N \geq 2$, we may choose $R' \geq 1$ sufficiently large to guarantee that

$$p c_p C_{GS} r^{-\frac{N-1}{2}} \int_{\mathbb{R}^N} \psi^{p-1} dx \leq \frac{m_{c,1}}{4} \int_{B_1(0)} \psi^p dx, \quad \text{for } r \geq R',$$

and therefore

$$E_\infty^+(t\psi_r) - E(t\psi_r) \geq C_2 e^{-r} t^p, \quad \text{for } r \geq R', \quad \text{with } C_2 := \frac{\kappa m_{c,1}}{4} \int_{B_1(0)} \psi^p dx.$$

Hence the claim follows. \square

Lemma 4.4. *Let $R' \geq 1$ be given as in Lemma 4.3. Then there exist $k > 0$ with $E(k\psi_r) < 0$ for all $r \geq R'$.*

Proof. Let $k > 0$. For $r \geq R' \geq 1$ we have, by Lemma 4.3 and since the map $r \mapsto \varepsilon_r$ is strictly decreasing by (4.7),

$$\begin{aligned}
E(k\psi_r) &\leq E_\infty^+(k\psi_r) = \frac{k^2}{2} \|\psi_r\|^2 - \frac{k^{p+1}}{p+1} \int_{\mathbb{R}_+^N} \psi_r^{p+1} dx \leq \frac{k^2}{2} \|\psi\|_{H^1(\mathbb{R}^N)}^2 - \frac{k^{p+1}}{p+1} \int_{B_r(0)} (\psi - \varepsilon_r)^{p+1} dx \\
&\leq \frac{k^2}{2} \|\psi\|_{H^1(\mathbb{R}^N)}^2 - \frac{k^{p+1}}{p+1} \int_{B_1(0)} (\psi - \varepsilon_1)^{p+1} dx
\end{aligned}$$

Since $\int_{B_1(0)} (\psi - \varepsilon_1)^{p+1} dx > 0$ by (4.7), we may choose

$$k > \left(\frac{(p+1) \|\psi\|_{H^1(\mathbb{R}^N)}^2}{2 \int_{B_1(0)} (\psi - \varepsilon_1)^{p+1} dx} \right)^{\frac{1}{p-1}}$$

which implies that $E(k\psi_r) < 0$ for $r \geq R'$, as claimed. \square

Proof of Proposition 4.1. Let $R' \geq 1$ be given by Lemma 4.3, and let $k > 0$ be given by Lemma 4.4. For $r \geq R'$ and $t \in [0, k]$ we then have, by Lemmas 4.2 and 4.3,

$$E(t\psi_r) = E_\infty^+(t\psi_r) - (E_\infty^+(t\psi_r) - E(t\psi_r)) \leq b_\infty + e^{-r} \left(t^{p+1} C_1 r^{-\frac{N-1}{2}} - t^p C_2 \right) \leq b_\infty + e^{-r} t^p \left(k C_1 r^{-\frac{N-1}{2}} - C_2 \right).$$

Since $N \geq 2$, we may fix $R \geq R'$ with the property that $k C_1 r^{-\frac{N-1}{2}} \leq \frac{C_2}{2}$ for $r \geq R$, which implies that

$$E(t\psi_r) \leq b_\infty - \frac{C_2}{2} e^{-r} t^p < b_\infty \quad \text{for } t \in (0, k], r \geq R.$$

Since also $E(0) = 0 < b_\infty$, we thus obtain that $E(t\psi_r) < b_\infty$ for $t \in [0, k]$, $r \geq R$. Moreover, by Lemma 4.4 we have $E(k\psi_r) < 0$ for all $r \geq R$ since $R \geq R'$. The proof is finished. \square

5. THE EXISTENCE RESULT

We keep using the notation of the introduction and of Section 2, which depends on the fixed quantities $c \in (0, c_p)$ and $p \in (1, 2^*)$. Moreover, we will assume $N \geq 2$ throughout this section, which will allow us to prove the existence of a non-trivial solution to (1.14). This will conclude the proof of Theorem 1.3.

Proposition 5.1. *Let $N \geq 2$. Then there exists a non-trivial solution $u \in H_0^1(\mathbb{R}_+^N)$ to (1.14).*

Proof. Since the functional E has a mountain pass geometry (see Lemma 3.1), there exists a Cerami sequence for E at the corresponding mountain pass level b defined in (4.1) (see e.g. [10] or [13, Theorem 6, Section 1, Chapter IV]), i.e. there exists $(u_n)_n \subset H_0^1(\mathbb{R}_+^N)$ such that

$$E(u_n) \rightarrow b \quad \text{and} \quad (1 + \|u_n\|) \|E'(u_n)\|_{H^{-1}(\mathbb{R}_+^N)} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

By Proposition 3.2 we know that $(u_n)_n$ is bounded in $H_0^1(\mathbb{R}_+^N)$. Moreover,

$$(5.1) \quad \|u_n^-\|^2 = \langle E'(u_n), u_n^- \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Let $(y^n)_n \subset \mathbb{R}_+^N$ be the sequence of points obtained in Lemma 3.3 applied to $(u_n)_n$, i.e., we have

$$(5.2) \quad \liminf_{n \rightarrow \infty} \int_{B_R(y^n) \cap \mathbb{R}_+^N} u_n^2 dx \geq \delta \quad \text{for some } \delta > 0.$$

We split the argument into two steps.

Step 1: *There exists $M > 0$ such that $y_N^n = \text{dist}(y^n, \partial \mathbb{R}_+^N) \leq M$ for all $n \in \mathbb{N}$.*

We assume by contradiction that

$$(5.3) \quad \lim_{n \rightarrow \infty} y_N^n = +\infty.$$

Then, let us define, for all $n \in \mathbb{N}$, $w_n := u_n(\cdot + y^n)$. By Lemma 3.3 and (5.1), it follows that

$$(5.4) \quad w_n \rightharpoonup w \text{ in } H^1(\mathbb{R}^N), \quad w_n \rightarrow w \text{ in } L_{loc}^q(\mathbb{R}^N) \text{ for } 1 \leq q < 2^*, \quad \text{and} \quad w_n \rightarrow w \text{ a.e. in } \mathbb{R}^N,$$

for some $w \in H^1(\mathbb{R}^N)$ with $w \geq 0$, $w \not\equiv 0$. We also observe that

$$(5.5) \quad b + o(1) = E(u_n) - \frac{1}{2} \langle E'(u_n), u_n \rangle = \int_{\mathbb{R}_+^N} H(x, u_n(x)) dx = \int_{\{x_N \geq -y_N^n\}} H(x + y_n, w_n^+(x)) dx, \quad \text{as } n \rightarrow \infty,$$

with the function H defined in Lemma 2.2 (ii). Next, we note that

$$H(x + y_n, w_n^+(x)) \geq 0, \quad \text{for } x \in \{x_N \geq -y_N^n\},$$

and

$$\begin{aligned} \liminf_{n \rightarrow \infty} H(x + y_n, w_n^+(x)) &\geq \liminf_{n \rightarrow \infty} \left(\frac{p-1}{2(p+1)} [w_n^+(x)]^{p+1} - u_c(x + y_n)^{p-1} D_{1,p}[w_n^+(x)]^2 - u_c(x + y_n) 1_{\{p>2\}} D_{2,p}[w_n^+(x)]^p \right) \\ &= \frac{p-1}{2(p+1)} [w^+(x)]^{p+1} = \frac{p-1}{2(p+1)} w^{p+1}(x), \quad \text{for } x \in \{x_N \geq -y_N^n\}, \end{aligned}$$

by Lemma 2.2 (ii) and (4.10). Thus, (5.5) and Fatou's Lemma imply that

$$(5.6) \quad \frac{p-1}{2(p+1)} \|w\|_{L^{p+1}(\mathbb{R}^N)}^{p+1} \leq b.$$

Next we claim that $w \in \mathcal{K}_\infty$, i.e., w is a nontrivial solution of (4.3). To see this, we fix an arbitrary $\varphi \in C_c^\infty(\mathbb{R}^N)$, and we show that

$$(5.7) \quad \int_{\mathbb{R}^N} (\nabla w \nabla \varphi + w \varphi) dx = \int_{\mathbb{R}^N} (w^+)^p \varphi dx.$$

Since (5.3) holds, we have that $\text{supp}(\varphi) \subset \{x_N \geq -y_N^n\}$ for $n \in \mathbb{N}$ sufficiently large. Hence, for $n \in \mathbb{N}$ large enough, we have that

$$\begin{aligned}
(5.8) \quad o(1) &= \langle E'(u_n), \varphi(\cdot - y^n) \rangle \\
&= \int_{\{x_N \geq -y_N^n\}} (\nabla w_n \nabla \varphi + w_n \varphi) dx - \int_{\{x_N \geq -y_N^n\}} (u_c(\cdot + y^n) + w_n^+)^p \varphi dx - \int_{\{x_N \geq -y_N^n\}} u_c^p(\cdot + y^n) \varphi dx \\
&= \int_{\mathbb{R}^N} (\nabla w_n \nabla \varphi + w_n \varphi) dx - \int_{\mathbb{R}^N} (u_c(\cdot + y^n) + w_n^+)^p \varphi dx + o(1) \\
&= \int_{\mathbb{R}^N} (\nabla w_n \nabla \varphi + w_n \varphi) dx - \int_{\mathbb{R}^N} (w_n^+)^p \varphi dx - \int_{\mathbb{R}^N} ((u_c(\cdot + y^n) + w_n^+)^p - (w_n^+)^p) \varphi dx + o(1) \\
&= \int_{\mathbb{R}^N} (\nabla w_n \nabla \varphi + w_n \varphi) dx - \int_{\mathbb{R}^N} (w_n^+)^p \varphi dx + o(1) \\
&= \int_{\mathbb{R}^N} (\nabla w \nabla \varphi + w \varphi) dx - \int_{\mathbb{R}^N} (w^+)^p \varphi dx + o(1), \quad \text{as } n \rightarrow \infty.
\end{aligned}$$

Hence (5.7) follows, and therefore $w \in \mathcal{K}_\infty$. Together with (4.8) and (5.6) it then follows that $b \geq b_\infty$, but this contradicts (4.9). Hence, (5.3) cannot happen and Step 1 follows.

Step 2: Conclusion.

Let us define, for all $n \in \mathbb{N}$, $v_n := u_n(\cdot + \xi_n)$ with $\xi_n := (y_1^n, \dots, y_{N-1}^n, 0)$ and observe that, after passing to a subsequence

$$(5.9) \quad v_n \rightharpoonup v \text{ in } H_0^1(\mathbb{R}_+^N), \quad v_n \rightarrow v \text{ in } L_{loc}^q(\mathbb{R}_+^N) \text{ for } 1 \leq q < 2^* \quad \text{and} \quad v_n \rightarrow v \text{ a.e. in } \mathbb{R}_+^N,$$

for some $v \in H_0^1(\mathbb{R}_+^N)$. Also, note that $(v_n)_n \subset H_0^1(\mathbb{R}_+^N)$ is a Cerami sequence for E at level b . Hence, if $v \not\equiv 0$, we will have that v is a non-trivial solution to (1.14). Since $v_n \rightarrow v$ in $L_{loc}^q(\mathbb{R}_+^N)$ and $y_N^n \leq M$ for all $n \in \mathbb{N}$, the lower integral bound (5.2) implies that $v \not\equiv 0$, and the result follows. \square

We now prove Proposition 1.5. Let us first state a technical lemma due to Poláčik, Quittner and Souplet that will be key to prove this result.

Lemma 5.2. (Particular case of [26, Lemma 5.1]) *Let (X, d) be a complete metric space and let $M : X \rightarrow (0, +\infty)$ be continuous. For any $\delta < \sup_X M$ and any $k > 0$ there exists $y \in X$ such that*

- $M(y) \geq \delta$.
- $M(z) \leq 2M(y)$ for all $z \in X$ with $d(z, y) \leq \frac{k}{M(y)}$.

The following proof is inspired by [12, Lemma 2.5].

Proof of Proposition 1.5. We assume by contradiction that there exists $v \in C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N})$ unbounded solving (1.3)-(1.4). By Lemma 5.2 applied with $X = \overline{\mathbb{R}_+^N}$ and $M = v^{\frac{p-1}{2}}$, there exists a sequence $(y^k)_k \subset \mathbb{R}_+^N$ such that

$$(5.10) \quad M(y^k) \rightarrow \infty, \quad \text{as } k \rightarrow \infty,$$

$$(5.11) \quad M(z) \leq 2M(y^k), \quad \text{for all } z \in \mathbb{R}_+^N \text{ with } d(z, y^k) \leq \frac{k}{M(y^k)} \text{ and all } k \in \mathbb{N}.$$

Note that, without loss of generality, we can suppose that $M(y^k) \geq 1$ for all $k \in \mathbb{N}$. We then define, for all $k \in \mathbb{N}$, $d_k := y_N^k M(y^k)$, the half-space $H_k := \{x \in \mathbb{R}_+^N : x_N > -d_k\}$ and

$$v^k : H_k \rightarrow \mathbb{R}_+^N \quad \text{given by} \quad v^k(z) := \frac{1}{v(y^k)} v\left(y^k + \frac{z}{M(y^k)}\right).$$

Note that, for all $k \in \mathbb{N}$, v^k is a positive solution to

$$(5.12) \quad \begin{cases} -\Delta v^k + \frac{1}{M^2(y^k)} v^k = (v^k)^p, & \text{in } H_k, \\ v^k = \frac{c}{M^{\frac{2}{p-1}}(y^k)}, & \text{on } \partial H_k, \end{cases}$$

and, by its definition and (5.11), it satisfies

$$(5.13) \quad v^k(0) = 1 \quad \text{and} \quad v^k(z) \leq 2^{\frac{2}{q-1}} \quad \text{for all } z \in H_k \cap B_k(0).$$

We now consider two cases separately.

Case 1: $d_k \rightarrow \infty$ as $k \rightarrow \infty$.

Using standard L^q estimates (see e.g. [18, Chapter 9]), (5.12) and (5.13), we get (taking a subsequence if necessary) that $(v^k)_k$ is locally $W^{2,q}$ -bounded in \mathbb{R}^N for arbitrarily large $q < +\infty$. Hence, up to a subsequence, $v^k \rightarrow \bar{v}$ in $C_{loc}^1(\mathbb{R}^N)$, where $\bar{v} \in C^1(\mathbb{R}^N)$ is a non-trivial positive solution to

$$(5.14) \quad \Delta \bar{v} + \bar{v}^p = 0, \quad \text{in } \mathbb{R}^N.$$

By (5.13) we infer that \bar{v} is bounded in \mathbb{R}^N with $\bar{v}(0) = 1$. Hence $\bar{v} \in C^2(\mathbb{R}^N)$ by standard elliptic regularity. Then, since by [17, Theorem 1.2] we know that the only $C^2(\mathbb{R}^N)$ non-negative solution to (5.14) is $\bar{v} \equiv 0$, we obtain a contradiction and deduce that *Case 1* cannot happen.

Case 2: $d_k \rightarrow d \geq 0$ as $k \rightarrow \infty$.

Let us define, for all $k \in \mathbb{N}$,

$$(5.15) \quad w^k : \mathbb{R}_+^N \rightarrow \mathbb{R} \quad \text{as} \quad w^k(z) = v^k(z - d_k e_N),$$

where $e_N := (0, \dots, 0, 1)$ is the N -th coordinate vector. Note that, for all $k \in \mathbb{N}$, w^k is a positive solution to

$$(5.16) \quad \begin{cases} -\Delta w^k + \frac{1}{M^2(y^k)} w^k = (w^k)^p, & \text{in } \mathbb{R}_+^N, \\ w^k = \frac{c}{M^{\frac{2}{p-1}}(y^k)}, & \text{on } \partial \mathbb{R}_+^N, \end{cases}$$

and satisfies

$$(5.17) \quad w^k(d_k e_N) = 1 \quad \text{and} \quad w^k(z) \leq 2^{\frac{2}{p-1}} \quad \text{for all } z \in \overline{\mathbb{R}_+^N} \cap B_k(d_k e_N).$$

Now, arguing as in the proof of [18, Theorem 9.13] (with auxiliary functions $\varphi^k = w^k - c M^{-\frac{2}{p-1}}(y^k)$) and taking into account (5.16) and (5.17), we get (taking a subsequence if necessary) that $(w^k)_k$ is locally $W^{2,q}$ -bounded in $\overline{\mathbb{R}_+^N}$ for arbitrarily large $q < +\infty$ and therefore also locally $C^{1,\beta}$ -bounded in $\overline{\mathbb{R}_+^N}$ for all $\beta \in (0, 1)$. In particular, $|\nabla w^k|$ remains bounded pointwise independently of k in a neighbourhood of the origin. Taking into account (5.10), the boundary conditions in (5.16) and (5.17), we infer that $d = \lim_{k \rightarrow \infty} d_k > 0$. Hence, up to a subsequence, $w^k \rightarrow \bar{w}$ in $C_{loc}^1(\overline{\mathbb{R}_+^N})$ with $\bar{w} \in C^1(\overline{\mathbb{R}_+^N})$ a non-trivial positive solution to

$$(5.18) \quad \begin{cases} \Delta \bar{w} + \bar{w}^p = 0, & \text{in } \mathbb{R}_+^N, \\ \bar{w} = 0, & \text{on } \partial \mathbb{R}_+^N. \end{cases}$$

By (5.17) we have that \bar{w} is bounded with $\bar{w}(d e_N) = 1$. Hence $\bar{w} \in C^2(\mathbb{R}^N) \cap C(\overline{\mathbb{R}_+^N})$ by standard elliptic regularity. Then, since by [17, Theorem 1.3] we know that the only $C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N})$ non-negative solution to (5.18) is $\bar{w} \equiv 0$, we obtain a contradiction and deduce that *Case 2* cannot happen either. Hence, the result follows. \square

Proof of Theorem 1.3. Let $u \in H_0^1(\mathbb{R}_+^N)$ be the non-negative and non-trivial solution to (1.11) obtained in Proposition 5.1. By standard elliptic regularity we have that $u \in C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N}) \cap L^\infty(\mathbb{R}_+^N)$, and $v := u_c + u$ is a bounded positive solution to (1.3)-(1.4) of the form (1.8). \square

Remark 5.1. As explained in the introduction, Theorem 1.2 (i) and Corollary 1.4 are direct consequences of Theorem 1.3.

6. THE NON-EXISTENCE RESULT

In this section we prove Part (ii) of Theorem 1.2, which is concerned with the non-existence of bounded positive solutions to (1.3)-(1.4) in the case $c > c_p$. Recall that

$$w_0(t) = c_p \left[\cosh\left(\frac{p-1}{2}t\right) \right]^{-\frac{2}{p-1}},$$

is the unique even non-trivial positive solution to (1.5). Throughout this section, we will use the following notation. We define $v_0 : \mathbb{R}^N \rightarrow \mathbb{R}$ as

$$(6.1) \quad v_0(x) = w_0(x_N), \quad \text{for } x \in \mathbb{R}^N.$$

Also, recall that for a bounded positive solution to (1.3)-(1.4), we mean a function $v \in C^2(\mathbb{R}_+^N) \cap C(\overline{\mathbb{R}_+^N}) \cap L^\infty(\mathbb{R}_+^N)$, positive, satisfying (1.3) in the pointwise sense and such that (1.4) holds.

Theorem 6.1. *For $N \geq 1$, $p > 1$ and $c > c_p$, there are no bounded positive solutions to (1.3)-(1.4).*

Proof. Let us fix an arbitrary $c > c_p$. We assume by contradiction that there exists a bounded positive solution v to (1.3)-(1.4) and we define, for all $t \in \mathbb{R}$, $v_t := v_0(\cdot + te_N)$ where v_0 is given in (6.1) and $e_N = (0, \dots, 0, 1)$ is the N -th coordinate vector. We split the proof into three steps.

Step 1: *There exists $t_0 > 0$ such that $v > v_t$ in $\overline{\mathbb{R}_+^N}$ for all $t \geq t_0$.*

First of all, fixed an arbitrary $x \in \overline{\mathbb{R}_+^N}$, observe that

$$v_t(x) \rightarrow 0, \quad \text{as } t \rightarrow \infty, \quad \text{and} \quad v_{t_1}(x) > v_{t_2}(x), \quad \text{for all } 0 < t_1 < t_2.$$

Hence, there exists $t_0 > 0$ such that, for all $t \geq t_0$,

$$(6.2) \quad v_t \leq \left(\frac{1}{2p}\right)^{\frac{1}{p-1}} \quad \text{in } \overline{\mathbb{R}_+^N}.$$

We fix $t_0 > 0$ such that (6.2) holds and we are going to prove the Step 1 for this t_0 . To that end, we fix an arbitrary $t \geq t_0 > 0$. First, we are going to prove that $v \geq v_t$ in $\overline{\mathbb{R}_+^N}$. Since $c > c_p \geq \max_{x \in \mathbb{R}^N} v_t(x)$, we have that

$$(6.3) \quad \begin{cases} -\Delta(v - v_t) + (v - v_t) = |v|^{p-1}v - |v_t|^{p-1}v_t, & \text{in } \mathbb{R}_+^N, \\ v - v_t > 0, & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

or equivalently

$$(6.4) \quad \begin{cases} -\Delta(v - v_t) + c_t(x)(v - v_t) = 0, & \text{in } \mathbb{R}_+^N, \\ v - v_t > 0, & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

where

$$(6.5) \quad c_t(x) := \begin{cases} 1 - \frac{(v(x))^p - (v_t(x))^p}{v(x) - v_t(x)}, & \text{if } v(x) - v_t(x) \neq 0, \\ 1, & \text{if } v(x) - v_t(x) = 0. \end{cases}$$

We assume by contradiction that

$$(6.6) \quad \{x \in \mathbb{R}_+^N : v(x) < v_t(x)\} \neq \emptyset.$$

Then, using the mean value theorem and (6.2), we deduce that, for all $x \in \{x \in \mathbb{R}_+^N : v(x) < v_t(x)\}$,

$$(6.7) \quad c_t(x) \geq 1 - p(v_t(x))^{p-1} \geq \frac{1}{2}.$$

Hence, in each connected component D of $\{x \in \mathbb{R}_+^N : v(x) < v_t(x)\}$ we have that

$$(6.8) \quad \begin{cases} -\Delta(v - v_t) + c_t(x)(v - v_t) = 0, & \text{in } D, \\ v - v_t = 0, & \text{on } \partial D, \end{cases}$$

with c_t satisfying (6.7). Then, applying the weak maximum principle [5, Lemma 2.1], we obtain that $v \geq v_t$ in \overline{D} which contradicts the fact that $D \subset \{x \in \mathbb{R}_+^N : v(x) < v_t(x)\}$. Hence, we conclude that $\{x \in \mathbb{R}_+^N : v(x) < v_t(x)\} = \emptyset$ and so, that $v \geq v_t$ in \mathbb{R}_+^N . Having this at hand and substituting in (6.3), we deduce that

$$(6.9) \quad \begin{cases} -\Delta(v - v_t) + (v - v_t) \geq 0, & \text{in } \mathbb{R}_+^N, \\ v - v_t > 0, & \text{on } \partial\mathbb{R}_+^N, \end{cases}$$

and so, the Step 1 follows from the strong maximum principle and the fact that $t \geq t_0$ is arbitrary.

Step 2: $v > v_t$ in $\overline{\mathbb{R}_+^N}$ for all $t \in \mathbb{R}$.

Note that, if we prove that $v \geq v_t$ in \mathbb{R}_+^N for all $t \in \mathbb{R}$, then the claim follows from the Strong Maximum principle. Also, by the Step 1, we know that

$$\{t \in \mathbb{R} : v \geq v_s \text{ in } \mathbb{R}_+^N \text{ for all } s \geq t\} \neq \emptyset.$$

Hence, we can define

$$(6.10) \quad t_\star := \inf\{t \in \mathbb{R} : v \geq v_s \text{ in } \mathbb{R}_+^N \text{ for all } s \geq t\} \in [-\infty, t_0].$$

We argue by contradiction and suppose that $t_\star > -\infty$. First note that, by continuity, $v \geq v_{t_\star}$ in \mathbb{R}_+^N . Also, $t_\star > -\infty$ implies the existence of $M > 0$ such that

$$(6.11) \quad v_t(x', x_N) \leq \left(\frac{1}{2p}\right)^{\frac{1}{p-1}}, \quad \text{for all } t \in [t_\star - 1, t_\star], \quad x' \in \mathbb{R}^{N-1} \text{ and } x_N \geq M.$$

We now consider separately two cases.

Case 1: $\inf_{x \in \mathbb{R}^{N-1} \times [0, M]} (v - v_{t_\star}) =: \delta_M > 0$.

First, taking into account that $\|w'_0\|_{L^\infty(\mathbb{R})} \leq c_p$, we infer that, for all $t \leq t_\star$ and $x \in \mathbb{R}^{N-1} \times [0, M]$,

$$\begin{aligned} v(x', x_N) - v_t(x', x_N) &= v(x', x_N) - v_{t_\star}(x', x_N) + (v_{t_\star}(x', x_N) - v_t(x', x_N)) \\ &\geq \delta_M - |w_0(x_N + t_\star) - w_0(x_N + t)| \\ &\geq \delta_M - c_p |t_\star - t|. \end{aligned}$$

Hence, there exists $\eta_0 \in (0, 1)$ such that, for all $t_\star \geq t \geq t_\star - \eta_0$,

$$(6.12) \quad v(x', x_N) - v_t(x', x_N) > 0, \quad \text{for all } x' \in \mathbb{R}^{N-1} \text{ and } x_N \in [0, M].$$

In particular, if we define $\Sigma_M := \{x \in \mathbb{R}^N : x_N > M\}$, we have

$$v - v_t > 0, \quad \text{on } \partial\Sigma_M, \quad \text{for all } t \in [t_\star - \eta_0, t_\star].$$

Next, we are going to prove that, for all $t \in [t_\star - \eta_0, t_\star]$, it follows $v \geq v_t$ in Σ_M . To that end, we fix an arbitrary $t \in [t_\star - \eta_0, t_\star]$. Arguing as in Step 1, we have that

$$(6.13) \quad \begin{cases} -\Delta(v - v_t) + c_t(x)(v - v_t) = 0, & \text{in } \Sigma_M, \\ v - v_t > 0, & \text{on } \partial\Sigma_M, \end{cases}$$

where

$$(6.14) \quad c_t(x) := \begin{cases} 1 - \frac{(v(x))^p - (v_t(x))^p}{v(x) - v_t(x)}, & \text{if } v(x) - v_t(x) \neq 0, \\ 1, & \text{if } v(x) - v_t(x) = 0. \end{cases}$$

We assume by contradiction that

$$(6.15) \quad \{x \in \Sigma_M : v(x) < v_t(x)\} \neq \emptyset.$$

Then, using the mean value theorem and (6.11), we deduce that, for all $x \in \{x \in \Sigma_M : v(x) < v_t(x)\}$,

$$(6.16) \quad c_t(x) \geq 1 - p(v_t(x))^{p-1} \geq \frac{1}{2}.$$

Hence, in each connected component D of $\{x \in \Sigma_M : v(x) < v_t(x)\}$ we have that

$$(6.17) \quad \begin{cases} -\Delta(v - v_t) + c_t(x)(v - v_t) = 0, & \text{in } D, \\ v - v_t = 0, & \text{on } \partial D, \end{cases}$$

with c_t satisfying (6.16). Then, applying the weak maximum principle [5, Lemma 2.1], we obtain that $v \geq v_t$ in \overline{D} which contradicts the fact that $D \subset \{x \in \Sigma_M : v(x) < v_t(x)\}$. Hence, we conclude that $\{x \in \Sigma_M : v(x) < v_t(x)\} = \emptyset$ and so, that $v \geq v_t$ in Σ_M . Taking into account (6.12), we infer that, for all $\eta \in [0, \eta_0]$, $v \geq v_{t_\star - \eta}$ in \mathbb{R}_+^N . This is in contradiction with the definition of t_\star . Hence, *Case 1* cannot happen.

$$\text{Case 2: } \inf_{x \in \mathbb{R}^{N-1} \times [0, M]} (v - v_{t_\star}) = 0.$$

In this case there exists a sequence of points $(x^n)_n \subset \mathbb{R}^{N-1} \times [0, M]$ such that

$$(6.18) \quad v(x^n) - v_{t_\star}(x^n) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Up to a subsequence, it follows that $x_N^n \rightarrow \bar{x}_N$ for some $\bar{x}_N \in [0, M]$. We define then

$$v^n(x) = v(x' + (x^n)', x_N), \quad \text{for all } n \in \mathbb{N},$$

and, for all $n \in \mathbb{N}$, we have $v^n \geq v_{t_\star}$ in \mathbb{R}_+^N and

$$\begin{cases} -\Delta v^n + v^n = (v^n)^p, & \text{in } \mathbb{R}_+^N, \\ v^n = c, & \text{on } \partial \mathbb{R}_+^N. \end{cases}$$

Moreover, for all $n \in \mathbb{N}$, it follows that

$$\begin{cases} -\Delta(v^n - v_{t_\star}) + (v^n - v_{t_\star}) \geq 0, & \text{in } \mathbb{R}_+^N, \\ v^n - v_{t_\star} > 0, & \text{on } \partial \mathbb{R}_+^N, \end{cases}$$

and so, by the Strong Maximum principle, we have that

$$v^n - v_{t_\star} > 0, \quad \text{in } \overline{\mathbb{R}_+^N}, \quad \text{for all } n \in \mathbb{N}.$$

Now, arguing as in [15, Proof of Theorem 2.1, Step 1], we deduce that the sequence $(v^n)_n$ admits a subsequence, still denoted by $(v^n)_n$, converging to a function \bar{v} in $C_{loc}^2(\mathbb{R}_+^N)$. This function \bar{v} still solves

$$\begin{cases} -\Delta \bar{v} + \bar{v} = \bar{v}^p, & \text{in } \mathbb{R}_+^N, \\ \bar{v} = c, & \text{on } \partial \mathbb{R}_+^N, \end{cases}$$

and satisfies $\bar{v} \geq v_{t_\star}$ in \mathbb{R}_+^N , and

$$(6.19) \quad \bar{v}(0', \bar{x}_N) = v_{t_\star}(0', \bar{x}_N).$$

Note that (6.19) and $\bar{v} = c > c_p \geq v_{t_\star}$ on $\partial \mathbb{R}_+^N$ imply $\bar{x}_N > 0$. Since $\bar{v} \geq v_{t_\star}$ in \mathbb{R}_+^N , we have

$$\begin{cases} -\Delta(\bar{v} - v_{t_\star}) + (\bar{v} - v_{t_\star}) \geq 0, & \text{in } \mathbb{R}_+^N, \\ \bar{v} - v_{t_\star} > 0, & \text{on } \partial \mathbb{R}_+^N, \end{cases}$$

Hence, by the Strong Maximum principle, it follows that $\bar{v} > v_{t_\star}$ in \mathbb{R}_+^N which gives a contradiction with (6.19). *Case 2* cannot happen either and hence the Step 2 follows.

Step 3: Conclusion.

Observe that $v > v_t$ in \mathbb{R}_+^N for all $t \in \mathbb{R}$ implies that $v \geq v_0(0) = c_p$ in \mathbb{R}_+^N . This gives a contradiction with (1.4) and so the proof is complete. \square

REFERENCES

- [1] W. Ao, M. Musso, F. Pacard, and J. Wei. Solutions without any symmetry for semilinear elliptic problems. *J. Funct. Anal.*, 270(3):884–956, 2016.
- [2] T. Bartsch and M. Willem. Infinitely many nonradial solutions of a Euclidean scalar field equation. *J. Funct. Anal.*, 117(2):447–460, 1993.
- [3] T. Bartsch and M. Willem. Infinitely many radial solutions of a semilinear elliptic problem on \mathbf{R}^N . *Arch. Rational Mech. Anal.*, 124(3):261–276, 1993.
- [4] H. Berestycki, L. Caffarelli, and L. Nirenberg. Further qualitative properties for elliptic equations in unbounded domains. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 25(1-2):69–94, 1997.
- [5] H. Berestycki, L. A. Caffarelli, and L. Nirenberg. Monotonicity for elliptic equations in unbounded Lipschitz domains. *Comm. Pure Appl. Math.*, 50(11):1089–1111, 1997.
- [6] H. Berestycki, T. Gallouët, and O. Kavian. Équations de champs scalaires euclidiens non linéaires dans le plan. *C. R. Acad. Sci. Paris Sér. I Math.*, 297(5):307–310, 1983.
- [7] H. Berestycki, F. Hamel, and R. Monneau. One-dimensional symmetry of bounded entire solutions of some elliptic equations. *Duke Math. J.*, 103(3):375–396, 2000.
- [8] H. Berestycki and P.-L. Lions. Nonlinear scalar field equations. I. Existence of a ground state. *Arch. Ration. Mech. Anal.*, 82(4):313–345, 1983.
- [9] H. Berestycki and P.-L. Lions. Nonlinear scalar field equations. II. Existence of infinitely many solutions. *Arch. Rational Mech. Anal.*, 82(4):347–375, 1983.
- [10] G. Cerami. Un criterio di esistenza per i punti critici su varietà illimitate. *Istit. Lombardo Accad. Sci. Lett. Rend. A*, 112(2):332–336, 1978.
- [11] E. N. Dancer. New solutions of equations on \mathbf{R}^n . *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 30(3-4):535–563 (2002), 2001.
- [12] E. N. Dancer and T. Weth. Liouville-type results for non-cooperative elliptic systems in a half-space. *J. Lond. Math. Soc. (2)*, 86(1):111–128, 2012.
- [13] I. Ekeland. *Convexity methods in Hamiltonian mechanics*. Springer-Verlag, Berlin, 1990.
- [14] M. J. Esteban and P.-L. Lions. Existence and nonexistence results for semilinear elliptic problems in unbounded domains. *Proc. Roy. Soc. Edinburgh Sect. A*, 93(1-2):1–14, 1982/83.
- [15] A. Farina. Rigidity and one-dimensional symmetry for semilinear elliptic equations in the whole of \mathbf{R}^N and in half spaces. *Adv. Math. Sci. Appl.*, 13(1):65–82, 2003.
- [16] B. Gidas, W. M. Ni, and L. Nirenberg. Symmetry of positive solutions of nonlinear elliptic equations in \mathbf{R}^n . *Adv. in Math. Suppl. Stud.*, 7:369–402, 1981.
- [17] B. Gidas and J. Spruck. A priori bounds for positive solutions of nonlinear elliptic equations. *Comm. Partial Differential Equations*, 6(8):883–901, 1981.
- [18] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Springer-Verlag, Berlin, 2001.
- [19] G. J. O. Jameson. Inequalities comparing $(a+b)^p - a^p - b^p$ and $a^{p-1}b + ab^{p-1}$. *Elem. Math.*, 69(3):156–161, 2014.
- [20] L. Jeanjean. On the existence of bounded Palais-Smale sequences and application to a Landesman-Lazer-type problem set on \mathbf{R}^N . *Proc. Roy. Soc. Edinburgh Sect. A*, 129(4):787–809, 1999.
- [21] M. K. Kwong. Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbf{R}^n . *Arch. Rational Mech. Anal.*, 105(3):243–266, 1989.
- [22] P.-L. Lions. The concentration-compactness principle in the calculus of variations. The locally compact case. II. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 1(4):223–283, 1984.
- [23] Y. Liu and J. Wei. On the Helmholtz equation and Dancer’s-type entire solutions for nonlinear elliptic equations. *Proc. Amer. Math. Soc.*, 147(3):1135–1148, 2019.
- [24] S. Lorca and P. Ubilla. Symmetric and nonsymmetric solutions for an elliptic equation on \mathbf{R}^N . *Nonlinear Anal.*, 58(7-8):961–968, 2004.
- [25] M. Musso, F. Pacard, and J. Wei. Finite-energy sign-changing solutions with dihedral symmetry for the stationary nonlinear Schrödinger equation. *J. Eur. Math. Soc. (JEMS)*, 14(6):1923–1953, 2012.
- [26] P. Poláčik, P. Quittner, and P. Souplet. Singularity and decay estimates in superlinear problems via Liouville-type theorems. I. Elliptic equations and systems. *Duke Math. J.*, 139(3):555–579, 2007.
- [27] P. H. Rabinowitz. On a class of nonlinear Schrödinger equations. *Z. Angew. Math. Phys.*, 43(2):270–291, 1992.
- [28] W. A. Strauss. Existence of solitary waves in higher dimensions. *Comm. Math. Phys.*, 55(2):149–162, 1977.
- [29] M. Struwe. *Variational methods, Applications to nonlinear partial differential equations and Hamiltonian systems*, volume 34. Springer-Verlag, Berlin, fourth edition, 2008.
- [30] C. A. Stuart. Bifurcation from the essential spectrum. In *Topological nonlinear analysis, II (Frascati, 1995)*, volume 27 of *Progr. Nonlinear Differential Equations Appl.*, pages 397–443. Birkhäuser Boston, Boston, MA, 1997.
- [31] M. Willem. *Minimax theorems*. Birkhäuser Boston, Inc., Boston, MA, 1996.

ANTONIO J. FERNÁNDEZ

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH BA2 7AY, UK.

E-mail address: a.jf77@bath.ac.uk

TOBIAS WETH

INSTITUT FÜR MATHEMATIK, GOETHE-UNIVERSITÄT FRANKFURT, ROBERT-MAYER-STR. 10, D-60629 FRANKFURT AM MAIN, GERMANY.

E-mail address: weth@math.uni-frankfurt.de