

HQET vertex diagram: ε expansion

Andrey G. Grozin^{1,2,*}

¹*Budker Institute of Nuclear Physics, Lavrentyev St. 11, Novosibirsk 630090, Russia*

²*Novosibirsk State University, Pirogov St. 1, Novosibirsk 630090, Russia*

(Dated:)

Differential equations for the one-loop HQET vertex diagram with arbitrary self-energy insertions and arbitrary residual energies are reduced to the ε form and used to obtain the ε expansion in terms of Goncharov polylogarithms.




FIG. 1. The one-loop HQET vertex diagram

We consider the one-loop vertex diagram (Fig. 1) with arbitrary degrees of all 3 denominators:

$$I_{n_1, n_2, n_3}(\vartheta; \omega_1, \omega_2) = \frac{1}{i\pi^{d/2}} \int \frac{d^d k}{D_1^{n_1} D_2^{n_2} D_3^{n_3}},$$

$$D_1 = -2(k + p_1) \cdot v_1, \quad D_2 = -2(k + p_2) \cdot v_2,$$

$$D_3 = -k^2, \quad (1)$$

where $\omega_{1,2} = p_{1,2} \cdot v_{1,2}$, $\cosh \vartheta = v_1 \cdot v_2$. It has obvious properties

$$I_{n_1, n_2, n_3}(\vartheta; \omega_1, \omega_2) = I_{n_1, n_2, n_3}(-\vartheta; \omega_1, \omega_2), \quad (2)$$

$$I_{n_1, n_2, n_3}(\vartheta; \omega_1, \omega_2) = I_{n_2, n_1, n_3}(\vartheta; \omega_2, \omega_1), \quad (3)$$

$$I_{n_1, 0, n_3}(\vartheta; \omega_1, \omega_2) = I_{n_1, n_3}(-2\omega_1)^{d-n_1-2n_3}, \quad (4)$$

where

$$I_{n_1, n_2} = \frac{\Gamma(n_1 + 2n_2 - d)\Gamma(d/2 - n_2)}{\Gamma(n_1)\Gamma(n_2)} \quad (5)$$

is the one-loop HQET self-energy diagram.

Results exact in ε are known for $\omega_1 = \omega_2$ [1]

$$I_{n_1, n_2, n_3}(\vartheta; \omega, \omega) = I_{n_1 + n_2, n_3}(-2\omega)^{d-n_1-n_2-2n_3}$$

$$\times {}_3F_2\left(\begin{matrix} n_1, n_2, \frac{d}{2} - n_3 \\ \frac{n_1 + n_2}{2}, \frac{n_1 + n_2 + 1}{2} \end{matrix} \middle| \frac{1 - \cosh \vartheta}{2}\right) \quad (6)$$

and $\vartheta = 0$ [2]

$$I_{n_1, n_2, n_3}(0; \omega_1, \omega_2) = I_{n_1 + n_2, n_3}(-2\omega_2)^{d-n_1-n_2-2n_3}$$

$$\times {}_2F_1\left(\begin{matrix} n_1, n_1 + n_2 + 2n_3 - d \\ n_1 + n_2 \end{matrix} \middle| 1 - y\right). \quad (7)$$

(the symmetry (3) follows from a hypergeometric identity). Here and below we use $d = 4 - 2\varepsilon$,

$$x = e^\vartheta, \quad y = \frac{\omega_1}{\omega_2}. \quad (8)$$

We consider the one-loop vertex (Fig. 1) with any numbers of self-energy insertions into each of 3 lines, provided that all lines in these insertions are massless. If the full number of loops in all self-energy insertions into the line i is l_i , then $n_{1,2} = m_{1,2} + 2l_{1,2}\varepsilon$, $n_3 = m_3 + l_3\varepsilon$, where all m_i are integer. All integrals with a given set l_i can be reduced [1], using IBP, to 3 master integrals with $m_i = (0, 1, 1)$, $(1, 0, 1)$ and $(1, 1, 1)$. We choose the column of the basis integrals $(f_1, f_2, f_3)^T$, where

$$I_{2l_1\varepsilon, 1+2l_2\varepsilon, 1+l_3\varepsilon}(\vartheta; \omega_1, \omega_2) = I_{1+2(l_1+l_2)\varepsilon, 1+l_3\varepsilon}$$

$$\times (-2\omega_1)^{-l\varepsilon} (-2\omega_2)^{1-l\varepsilon} f_1(x, y),$$

$$I_{1+2l_1\varepsilon, 2l_2\varepsilon, 1+l_3\varepsilon}(\vartheta; \omega_1, \omega_2) = I_{1+2(l_1+l_2)\varepsilon, 1+l_3\varepsilon}$$

$$\times (-2\omega_1)^{1-l\varepsilon} (-2\omega_2)^{-l\varepsilon} f_2(x, y),$$

$$I_{1+2l_1\varepsilon, 1+2l_2\varepsilon, 1+l_3\varepsilon}(\vartheta; \omega_1, \omega_2) = I_{2+2(l_1+l_2)\varepsilon, 1+l_3\varepsilon}$$

$$\times (-2\omega_1)^{-l\varepsilon} (-2\omega_2)^{-l\varepsilon} f_3(x, y), \quad (9)$$

where $l = l_1 + l_2 + l_0$ is the total number of loops, $l_0 = l_3 + 1$. They have symmetry properties

$$f(x^{-1}, y) = f(x, y), \quad (10)$$

$$f(x, y^{-1}) = S_y[f(x, y)]_{l_1 \leftrightarrow l_2}, \quad S_y = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The initial condition is $f(1, 1) = (1, 1, 1)^T$. If $l_1 = 0$, f_1 is trivial (4); if $l_2 = 0$, f_2 is trivial; if $l_1 = l_2 = 0$, there is only one non-trivial master integral f_3 . If $l_1 = l_2$, $f_2(x, y) = f_1(x, y^{-1})$ (10), and there are only 2 unknown functions f_1 and f_3 .

We shall use the method of differential equations [3]. Using

$$\sinh \vartheta \frac{\partial}{\partial \vartheta} I_{n_1, n_2, n_3}$$

$$= n_1 [1^+ 2^- - 2(\omega_1 \cosh \vartheta - \omega_2) 1^+ - \cosh \vartheta] I_{n_1, n_2, n_3}$$

$$= n_2 [2^+ 1^- - 2(\omega_2 \cosh \vartheta - \omega_1) 2^+ - \cosh \vartheta] I_{n_1, n_2, n_3},$$

$$\frac{\partial}{\partial \omega_1} I_{n_1, n_2, n_3} = 2n_1 1^+ I_{n_1, n_2, n_3},$$

$$\frac{\partial}{\partial \omega_2} I_{n_1, n_2, n_3} = 2n_2 2^+ I_{n_1, n_2, n_3} \quad (11)$$

and the IBP reduction, we can derive the differential equations

$$\partial_x f = M_x f, \quad \partial_y f = M_y f, \quad (12)$$

where the matrices $M_{x,y}$ (depending on x, y and ε) satisfy

$$\partial_x M_y - \partial_y M_x - [M_x, M_y] = 0 \quad (13)$$

* A.G.Grozin@inp.nsk.su

because $\partial_x \partial_y f = \partial_y \partial_x f$. The symmetries (10) lead to

$$\begin{aligned} M_x(x^{-1}, y) + x^2 M_x(x, y) &= 0, \quad M_y(x^{-1}, y) = M_y(x, y); \\ M_x(x, y^{-1}) &= S_y [M_x]_{l_1 \leftrightarrow l_2} S_y, \\ M_y(x, y^{-1}) + y^2 S_y [M_y]_{l_1 \leftrightarrow l_2} S_y &= 0. \end{aligned} \quad (14)$$

The differential equations (12) can be reduced to the canonical form [4] by a linear transformation $f = TF$ (the matrix T depends on x, y, ϵ),

$$dF = \epsilon dMF, \quad M(x, y) = \sum_i M_i \log p_i(x, y), \quad (15)$$

where $p_i(x, y)$ are polynomials in x and y , and M_i are constant matrices. We use the Mathematica package *Libra* [5] which implements the algorithm of [6], and obtain

$$\begin{aligned} T &= \begin{pmatrix} 1 & 0 & l_1 \frac{1+x^2-2xy}{1-x^2} \\ 0 & 1 & l_2 \frac{1+x^2-2xy^{-1}}{1-x^2} \\ 0 & 0 & -\frac{1+2(l_1+l_2)\epsilon}{\epsilon} \frac{x}{1-x^2} \end{pmatrix}, \\ T^{-1} &= \begin{pmatrix} 1 & 0 & \frac{l_1 \epsilon}{1+2(l_1+l_2)\epsilon} \frac{1+x^2-2xy}{x} \\ 0 & 1 & \frac{l_2 \epsilon}{1+2(l_1+l_2)\epsilon} \frac{1+x^2-2xy^{-1}}{x} \\ 0 & 0 & -\frac{\epsilon}{1+2(l_1+l_2)\epsilon} \frac{1-x^2}{x} \end{pmatrix}. \end{aligned} \quad (16)$$

The symmetry properties of the canonical master integrals are

$$\begin{aligned} F(x^{-1}, y) &= S_x F(x, y), \quad S_x = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \\ F(x, y^{-1}) &= S_y [F(x, y)]_{l_1 \leftrightarrow l_2}. \end{aligned} \quad (17)$$

The initial conditions for the differential equations (15) are

$$F(1, 1) = T^{-1}(1, 1) f(1, 1) = (1, 1, 0)^T. \quad (18)$$

The matrix $M(x, y)$ is

$$\begin{aligned} M &= M_1 \log x + M_2 [\log(1+x) + \log(1-x)] + M_3 \log y \\ &\quad + M_4 \log(x-y) + M_5 \log(1-xy), \\ M_1 &= \begin{pmatrix} l_1 & -l_1 & l_1(l_1-l_2+l_0) \\ -l_2 & l_2 & l_2(-l_1+l_2+l_0) \\ 1 & 1 & l_1+l_2-l_0 \end{pmatrix}, \end{aligned} \quad (19)$$

$$\begin{aligned} M_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2l_0 \end{pmatrix}, \quad M_3 = \begin{pmatrix} l & 0 & 0 \\ -2l_2 & -l_1+l_2-l_0 & 0 \\ 0 & 0 & l \end{pmatrix}, \\ M_4 &= \begin{pmatrix} -l_1 & l_1 & l_1 l \\ l_2 & -l_2 & -l_2 l \\ 1 & -1 & -l \end{pmatrix}, \quad M_5 = \begin{pmatrix} -l_1 & l_1 & -l_1 l \\ l_2 & -l_2 & l_2 l \\ -1 & 1 & -l \end{pmatrix} \end{aligned}$$

(only derivatives of M matter, and hence we may freely substitute $\log(y-x) \rightarrow \log(x-y)$, etc.). This matrix has symmetry properties

$$\begin{aligned} M(x^{-1}, y) &= S_x M(x, y) S_x, \\ M(x, y^{-1}) &= S_y [M(x, y)]_{l_1 \leftrightarrow l_2} S_y \end{aligned} \quad (20)$$

(again, up to inessential additive constants).

If $l_1 = 0$ then $F_1(x, y) = y^{l\epsilon}$. The first equation decouples, and this trivial function satisfies this equation. The two non-trivial master integrals $F_{2,3}$ are determined by coupled equations. The case $l_2 = 0$ is similar. If $l_1 = l_2 = 0$ then $F_{1,2}(x, y) = y^{\pm l\epsilon}$; the only non-trivial master integral F_3 is determined by the third equation.

First we consider the single-scale case $y = 1$. The differential equations for $x < 1$ are

$$\begin{aligned} \frac{dF(x, 1)}{dx} &= \epsilon \left[\frac{M_1}{x} + \frac{M_2}{x+1} + \frac{M_2+M_4+M_5}{x-1} \right] F(x, 1), \\ M_2+M_4+M_5 &= 2 \begin{pmatrix} -l_1 & l_1 & 0 \\ l_2 & -l_2 & 0 \\ 0 & 0 & -l_1-l_2 \end{pmatrix}. \end{aligned} \quad (21)$$

For $x > 1$ we have $F(x, 1) = S_x F(x^{-1}, 1)$; these functions satisfy the equations

$$\begin{aligned} \frac{dF(x, 1)}{dx^{-1}} &= \epsilon \left[-\frac{M_1+2M_2+M_4+M_5}{x^{-1}} \right. \\ &\quad \left. + \frac{M_2}{x^{-1}+1} + \frac{M_2+M_4+M_5}{x^{-1}-1} \right] F(x, 1) \end{aligned}$$

because $-S_x(M_1+2M_2+M_4+M_5)S_x = M_1$, $S_x M_2 S_x = M_2$, $S_x(M_2+M_4+M_5)S_x = M_2+M_4+M_5$ (this follows from (20)).

The solution of the differential equations (21) with the initial conditions (18) as a series in ϵ can be obtained using *Libra*. The coefficients are uniform-weight combinations of harmonic polylogarithms [7] (we use HPL [8, 9] to reduce them to a minimal set):

$$\begin{aligned} F_1(x, 1) &= 1 + l_1(l_1-l_2+l_0)H_0^2(x)\epsilon^2 \\ &+ 2l_1 \left\{ (l_1-l_2+l_0) \left[-4l_0 H_{0,0,-1}(x) + 2l_0 H_0(x)H_{0,-1}(x) + (2l-l_0) \frac{\pi^2}{6} H_0(x) \right] - 2(l_1^2-l_2^2+(l_1+3l_2)l_0)H_{0,0,1}(x) \right. \\ &\quad \left. + 4l_2 l_0 H_0(x)H_{0,1}(x) + (l_1-l_2)l H_0^2(x)H_1(x) + \frac{1}{6}(2(l_1^2-l_2^2)+(l_1+l_2)l_0-l_0^2)H_0^3(x) + (2(l_1^2-l_2^2)+(5l_1+3l_2)l_0+3l_0^2)\zeta_3 \right\} \epsilon^3 \\ &+ 2l_1 \left\{ (l_1-l_2)l \left[2l_0(4H_{0,0,1,-1}(x) + 4H_{0,0,-1,1}(x) + 2H_{0,1,0,-1}(x) - 3H_{0,0,0,-1}(x) - 4H_1(x)H_{0,0,-1}(x) + 2H_0(x)H_1(x)H_{0,-1}(x)) \right. \right. \\ &\quad \left. \left. - (l_1+l_2)(2H_{0,1,0,1}(x) + 4H_1(x)H_{0,0,1}(x) - H_0^2(x)H_1^2(x)) - (2(l_1+l_2)-l_0) \left(2H_{0,0,0,1}(x) - \frac{1}{3}H_0^3(x)H_1(x) \right) \right] \right\} \epsilon^4 \end{aligned}$$

$$\begin{aligned}
& + (2l - l_0) \frac{\pi^2}{3} H_0(x) H_1(x) + 2(2l + l_0) \zeta_3 H_1(x) \Big] \\
& + (l_1 - l_2 + l_0) \Big[4l_0^2 H_0(x) H_{0,-1,-1}(x) - 2l_0^2 H_{0,-1}^2(x) - 4(l_1 + l_2) l_0 H_{0,1}(x) H_{0,-1}(x) + (l_1 + l_2 - l_0) l_0 H_0^2(x) H_{0,-1}(x) \\
& + (2l - l_0) l_0 \frac{\pi^2}{3} H_{0,-1}(x) \Big] + (l_1 + l_2) [8l_2 l_0 H_0(x) H_{0,1,1}(x) + (l_1^2 - l_2^2 + (l_1 - 5l_2) l_0) H_{0,1}^2(x)] \\
& + 8l_2 l_0^2 H_0(x) (H_{0,1,-1}(x) + H_{0,-1,1}(x)) - 2l_2 (l_1 + l_2 - l_0) l_0 H_0(x) (2H_{0,0,1}(x) - H_0(x) H_{0,1}(x)) \\
& + 2(l_1 - 3l_2 + l_0) l_0^2 H_0(x) H_{0,0,-1}(x) + \frac{1}{24} (4(l_1^3 - l_2^3) + 2(l_1^2 + l_2^2) l_0 - (l_1 + l_2) l_0^2 + l_0^3) H_0^4(x) \\
& + (2l - l_0) \frac{\pi^2}{12} \Big[8l_2 l_0 H_{0,1}(x) + (2(l_1^2 - l_2^2) + (l_1 + l_2) l_0 - l_0^2) H_0^2(x) + (6(l_1^2 - l_2^2) + (l_1 - 21l_2) l_0 - 5l_0^2) \frac{\pi^2}{30} \Big] \\
& - 2(2l_2 l + l_1 l_0 + l_0^2) l_0 \zeta_3 H_0(x) \Big\} \varepsilon^4 + \mathcal{O}(\varepsilon^5),
\end{aligned}$$

$$F_2(x, 1) = [F_1(x, 1)]_{l_1 \leftrightarrow l_2},$$

$$F_3(x, 1) = [F_3(x, 1)]_{l_1 \leftrightarrow l_2} = 2H_0(x) \varepsilon$$

$$\begin{aligned}
& + \left[-4(l_1 + l_2) (H_{0,1}(x) - H_0(x) H_1(x)) - 4l_0 (H_{0,-1}(x) - H_0(x) H_{-1}(x)) + (l_1 + l_2 - l_0) H_0^2(x) + (2l - l_0) \frac{\pi^2}{3} \right] \varepsilon^2 \\
& + \left\{ 4(l_1 + l_2)^2 (2H_{0,1,1}(x) - 2H_1(x) H_{0,1}(x) + H_0(x) H_1^2(x)) \right. \\
& + 2(l_1 + l_2) \left[4l_0 (H_{0,1,-1}(x) + H_{0,-1,1}(x) - H_{-1}(x) H_{0,1}(x) - H_1(x) H_{0,-1}(x) + H_0(x) H_1(x) H_{-1}(x)) \right. \\
& \quad \left. - (l_1 + l_2 - l_0) (2H_{0,0,1}(x) - H_0^2(x) H_1(x)) + (2l - l_0) \frac{\pi^2}{3} H_1(x) \right] \\
& + 4l_0^2 (2H_{0,-1,-1}(x) - 2H_{-1}(x) H_{0,-1}(x) + H_0(x) H_{-1}^2(x)) - (l_1 + l_2 - l_0) \left[4l_0 H_{0,0,-1}(x) - 2l_0 H_0^2(x) H_{-1}(x) - (2l - l_0) \frac{\pi^2}{3} H_0(x) \right] \\
& \quad \left. + \frac{1}{3} (2(l_1^2 + l_2^2) - (l_1 + l_2) l_0 + l_0^2) H_0^3(x) + 2(2l - l_0) l_0 \frac{\pi^2}{3} H_{-1}(x) - 2(2(l_1 + l_2)^2 + 3(l_1 + l_2) l_0 + 2l_0^2) \zeta_3 \right\} \varepsilon^3 \\
& + \left\{ 8(l_1 + l_2)^3 \left(-2H_{0,1,1,1}(x) + 2H_1(x) H_{0,1,1}(x) - H_1^2(x) H_{0,1}(x) + \frac{1}{3} H_0(x) H_1^3(x) \right) \right. \\
& - 2(l_1 + l_2)^2 \left[(l_1 + l_2 - l_0) (2H_{0,1,0,1}(x) + 4H_1(x) H_{0,0,1}(x) - H_{0,1}^2(x) - H_0^2(x) H_1^2(x)) \right. \\
& \quad + 4l_0 (2H_{0,1,1,-1}(x) + 2H_{0,1,-1,1}(x) + 2H_{0,-1,1,1}(x) - 2H_{-1}(x) H_{0,1,1}(x) - 2H_1(x) H_{0,1,-1}(x) - 2H_1(x) H_{0,-1,1}(x) \\
& \quad \left. + 2H_1(x) H_{-1}(x) H_{0,1}(x) + H_1^2(x) H_{0,-1}(x) - H_0(x) H_{-1}(x) H_1^2(x)) - (2l - l_0) \frac{\pi^2}{3} H_1^2(x) \right] \\
& \quad \left. - 8l_0^3 \left(2H_{0,-1,-1,-1}(x) - 2H_{-1}(x) H_{0,-1,-1}(x) + H_{-1}^2(x) H_{0,-1}(x) - \frac{1}{3} H_0(x) H_{-1}^3(x) \right) \right. \\
& + 2(l_1 + l_2 - l_0) \left[(l_1 + l_2) \left(2l_0 (2H_{0,0,1,-1}(x) + 2H_{0,0,-1,1}(x) - 2H_{-1}(x) H_{0,0,1}(x) - 2H_1(x) H_{0,0,-1}(x) + H_0^2(x) H_1(x) H_{-1}(x)) \right. \right. \\
& \quad \left. \left. + (2l - l_0) \frac{\pi^2}{3} H_0(x) H_1(x) \right) + l_0^2 (4H_{0,0,-1,-1}(x) - 4H_{-1}(x) H_{0,0,-1}(x) + H_0^2(x) H_{-1}^2(x)) + (2l - l_0) l_0 \frac{\pi^2}{3} H_0(x) H_{-1}(x) \right] \\
& - 2(l_1 + l_2) \left[4l_0^2 (2H_{0,1,-1,-1}(x) + 2H_{0,-1,1,-1}(x) + 2H_{0,-1,-1,1}(x) - 2H_{-1}(x) H_{0,1,-1}(x) - 2H_{-1}(x) H_{0,-1,1}(x) \right. \\
& \quad \left. - 2H_1(x) H_{0,-1,-1}(x) + H_{-1}^2(x) H_{0,1}(x) + 2H_1(x) H_{-1}(x) H_{0,-1}(x) - H_0(x) H_1(x) H_{-1}^2(x)) - 2(2l - l_0) l_0 \frac{\pi^2}{3} H_1(x) H_{-1}(x) \right] \\
& - 4(2(l_1 + l_2) (l_1^2 + l_2^2) - (l_1^2 + l_2^2 - 10l_1 l_2) l_0 + (l_1 + l_2) l_0^2) H_{0,0,0,1}(x) - 4(5(l_1^2 + l_2^2) - 6l_1 l_2 + 2(l_1 + l_2) l_0 + l_0^2) l_0 H_{0,0,0,-1}(x) \\
& \quad \left. + 8l_1 l_2 l_0 H_0(x) (4H_{0,0,1}(x) - H_0(x) H_{0,1}(x)) + 2((l_1 - l_2)^2 + (l_1 + l_2) l_0) l_0 H_0(x) (4H_{0,0,-1}(x) - H_0(x) H_{0,-1}(x)) \right\} \varepsilon^4
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{3} (2(l_1^2 + l_2^2) - (l_1 + l_2)l_0 + l_0^2) H_0^2(x) \left[2(l_1 + l_2) H_0(x) H_1(x) + 2l_0 H_0(x) H_{-1}(x) + (2l - l_0) \frac{\pi^2}{2} \right] \\
& + \frac{1}{12} (4(l_1^3 + l_2^3) - 2(l_1^2 + l_2^2)l_0 + (l_1 + l_2)l_0^2 - l_0^3) H_0^4(x) \\
& + (2l - l_0) \frac{\pi^2}{3} \left[2l_0^2 H_{-1}^2(x) + (22(l_1^2 + l_2^2) + 28l_1 l_2 + 13(l_1 + l_2)l_0 + 9l_0^2) \frac{\pi^2}{60} \right] \\
& - 4(2(l_1 + l_2)^2 + 3(l_1 + l_2)l_0 + 2l_0^2) \zeta_3 [(l_1 + l_2) H_1(x) + l_0 H_{-1}(x)] \\
& - 4(4l_1 l_2 (l_1 + l_2) - 2(l_1^2 + l_2^2 + l_1 l_2)l_0 - 2(l_1 + l_2)l_0^2 - l_0^3) \zeta_3 H_0(x) \Big\} \varepsilon^4 + \mathcal{O}(\varepsilon^5). \tag{22}
\end{aligned}$$

This expansion can be straightforwardly extended to any order in ε . We have also expanded the exact hypergeometric representations of $F_{1,3}(x, 1)$ which follow from (6) up to ε^3 using HypExp [10, 11]. The results can be expressed via ordinary polylogarithms up to Li_3 , and agree with (22). They also agree with the expansions up to ε^3 obtained in [1] (also using (6) and HypExp). When $l_1 = l_2 = 0$, the only non-trivial master integral F_3 is expressed (6) via the ${}_2F_1$ function whose ε expansion is known to all orders [12]. The expansion in euclidean case is given there (there is a typo in the journal version corrected in the version 4 in arXiv); the Minkowski case is given by the formula (41) in [13]. Our result (22) at $l_1 = l_2 = 0$ agrees with the formula (B.10) in [13] (it contains 3 further expansion terms).

Any finite number of terms in the expansion of $F(x, 1)$ in $\bar{x} = 1 - x$ can be straightforwardly obtained from (6):

$$\begin{aligned}
F_1(x, 1) &= 1 + \frac{\varepsilon^2 l_1 \bar{x}^2}{(1 + (l_1 + l_2)\varepsilon)(1 + 2(l_1 + l_2)\varepsilon)} \\
&\quad \{ (l_1 - l_2 + l_0 + 2l_2 l_0 \varepsilon)(1 + \bar{x}) + \mathcal{O}(\bar{x}^2) \}, \\
F_2(x, 1) &= [F_1(x, 1)]_{l_1 \leftrightarrow l_2}, \\
F_3(x, 1) &= -\frac{\varepsilon \bar{x}}{1 + 2(l_1 + l_2)\varepsilon} [2 + \bar{x} + \mathcal{O}(\bar{x}^2)] \tag{23}
\end{aligned}$$

(we have obtained them up to $\bar{x}^{20}$). The coefficients are exact functions of ε . This expansion satisfies the differential equation (21) with the initial condition (18). Expanding each coefficient of (23) in ε , and each coefficient of (22) in $\bar{x}$, we obtain two identical double expansions up to ε^4 and $\bar{x}^{20}$; this is a strong check of our result (22).

Next we consider the straight-line case $x = 1$. From the form of the matrix T^{-1} (16) at $x = 1$ we see that $F_3(1, y) = 0$. The differential equations for $y < 1$ are

$$\begin{aligned}
\frac{dF(1, y)}{dy} &= \varepsilon \left[\frac{M_3}{y} + \frac{M_4 + M_5}{y - 1} \right] F(1, y), \\
M_4 + M_5 &= 2 \begin{pmatrix} -l_1 & l_1 & 0 \\ l_2 & -l_2 & 0 \\ 0 & 0 & -l \end{pmatrix} \tag{24}
\end{aligned}$$

(they are, of course, consistent with $F_3 = 0$). For $y > 1$ we have $F(1, y) = S_y [F(1, y^{-1})]_{l_1 \leftrightarrow l_2}$; these functions satisfy the equations

$$\begin{aligned}
\frac{dF(1, y)}{dy^{-1}} &= \varepsilon \left[-\frac{M_3 + M_4 + M_5}{y^{-1}} + \frac{M_4 + M_5}{y^{-1} - 1} \right] F(1, y) \\
&\text{because } -S_y [M_3 + M_4 + M_5]_{l_1 \leftrightarrow l_2} S_y = M_3, \\
&S_y [M_4 + M_5]_{l_1 \leftrightarrow l_2} S_y = M_4 + M_5 \text{ (this follows from (20)).} \\
&\text{Solving the differential equations (24) with the initial conditions (18) we obtain}
\end{aligned}$$

$$\begin{aligned}
y^{-l\varepsilon} F_1(1, y) &= 1 - 4l_1 l \left(H_{0,1}(y) - H_0(y) H_1(y) - \frac{\pi^2}{6} \right) \varepsilon^2 \\
&+ 4l_1 l \left[(l_1 + l_2) \left(2H_{0,1,1}(y) - 2H_1(y) H_{0,1}(y) + H_0(y) H_1^2(y) + \frac{\pi^2}{3} H_1(y) \right) \right. \\
&\quad \left. - (l_1 + l_0) (2H_{0,0,1}(y) - 2H_0(y) H_{0,1}(y) + H_0^2(y) H_1(y)) - 2(l_2 - l_0) \zeta_3 \right] \varepsilon^3 \\
&- 4l_1 l \left[2(l_1 + l_2)^2 \left(2H_{0,1,1,1}(y) - 2H_1(y) H_{0,1,1}(y) + H_1^2(y) H_{0,1}(y) - \frac{1}{3} H_0(y) H_1^3(y) - \frac{\pi^2}{6} H_1^2(y) \right) \right. \\
&\quad \left. + 2(l_1 + l_0)^2 \left(2H_{0,0,0,1}(y) - 2H_0(y) H_{0,0,1}(y) + H_0^2(y) H_{0,1}(y) - \frac{1}{3} H_0^3(y) H_1(y) \right) \right. \\
&\quad \left. + 2(l_1 l - l_2 l_0) \left(H_{0,1,0,1}(y) + 2H_0(y) H_{0,1,1}(y) + 2H_1(y) H_{0,0,1}(y) \right) - (l_1 l - 3l_2 l_0) H_{0,1}^2(y) - 4l_1 l H_0(y) H_1(y) H_{0,1}(y) \right. \\
&\quad \left. + (l_1 + l_2)(l_1 + l_0) H_0^2(y) H_1^2(y) - 2l_2 l_0 \frac{\pi^2}{3} (H_{0,1}(y) - H_0(y) H_1(y)) + 4(l_2(l_1 + l_2) - (l_1 - l_2)l_0) \zeta_3 H_1(y) \right] \varepsilon^4
\end{aligned}$$

$$\begin{aligned}
& - (7l_1l + 4(l_2^2 - l_2l_0 + l_0^2)) \frac{\pi^4}{90} \Big] \varepsilon^4 + \mathcal{O}(\varepsilon^5), \\
y^l F_2(1, y) = & 1 + 2l_2l \left(2H_{0,1}(y) - 2H_0(y)H_1(y) - H_0^2(y) - \frac{\pi^2}{3} \right) \varepsilon^2 \\
& - 4l_2l \left[(l_1 + l_2) \left(2H_{0,1,1}(y) - 2H_1(y)H_{0,1}(y) + H_0(y)H_1^2(y) + \frac{\pi^2}{3} (H_0(y) + H_1(y)) \right) - 2(l_1 - l_0)H_{0,0,1}(y) \right. \\
& \left. - 2(l_2 + l_0)(H_0(y)H_{0,1}(y) + \zeta_3) + (l_2 + l)H_0^2(y) \left(H_1(y) + \frac{1}{3}H_0(y) \right) \right] \varepsilon^3 \\
& + 4l_2l \left[2(l_1 + l_2)^2 \left(2H_{0,1,1,1}(y) - 2H_1(y)H_{0,1,1}(y) + H_1^2(y)H_{0,1}(y) - \frac{1}{3}H_0(y)H_1^3(y) - \frac{\pi^2}{6}H_1^2(y) \right) \right. \\
& + 2(l_1l - l_2l_0)(H_{0,1,0,1}(y) + 2H_1(y)H_{0,0,1}(y)) + 4(l_1 + l_0)^2H_{0,0,0,1}(y) \\
& - 4(l_2(l_1 + l_2) - (l_1 - l_2)l_0)(H_0(y)H_{0,1,1}(y) - \zeta_3(H_0(y) + H_1(y))) + 4(l_1l_2 - ll_0)H_0(y)H_{0,0,1}(y) \\
& - (l_1(l_1 + l_2 + 3l_0) - l_2l_0)H_{0,1}^2(y) + 2((l_2 + l_0)^2 + l_1l_0)H_0^2(y)H_{0,1}(y) + 4l_2lH_0(y)H_1(y)H_{0,1}(y) - (l_1 + l_2)(l_2 + l)H_0^2(y)H_1^2(y) \\
& \left. - \frac{1}{6}((l_1 + l_0)^2 + 3l_2l)H_0^3(y)(4H_1(y) + H_0(y)) + \frac{\pi^2}{3}(2l_1l_0H_{0,1}(y) - ((l_1 + l_2)^2 + l_1l_0)H_0(y)(2H_1(y) + H_0(y))) \right. \\
& \left. - (7l_1(l_1 + l_2) + 4l_2^2 + (12l_1 + l_2)l_0 + 4l_0^2) \frac{\pi^4}{90} \right] \varepsilon^4 + \mathcal{O}(\varepsilon^5). \tag{25}
\end{aligned}$$




FIG. 2. Paths from $(1, 1)$ to (x, y) .

This expansion can be straightforwardly extended to any order in ε . We have also expanded the exact hypergeometric representations of $F_{1,2}(1, y)$ which follow from (7) up to ε^3 using HypExp [10, 11]. The results can be expressed via ordinary polylogarithms up to Li_3 , and agree with (25).

Any finite number of terms in the expansion of $F(1, y)$ in $\bar{y} = 1 - y$ can be straightforwardly obtained from (7):

$$F_1(1, y) = 1 - \frac{\varepsilon l \bar{y}}{1 + 2(l_1 + l_2)\varepsilon} [1 - 2(l_1 - l_2)\varepsilon + \mathcal{O}(\bar{y})],$$

$$F_2(1, y) = 1 + \frac{\varepsilon l \bar{y}}{1 + 2(l_1 + l_2)\varepsilon} [1 + 2(l_1 - l_2)\varepsilon + \mathcal{O}(\bar{y})] \tag{26}$$

(we have obtained them up to $\bar{y}^{20}$). This expansion satisfies the differential equations (24) with the initial conditions (18). Expanding each coefficient of (26) in ε , and each coefficient of (25) in $\bar{y}$, we obtain two identical double expansions up to ε^4 and $\bar{y}^{20}$; this is a strong check of our result (25).

Finally, we discuss the general case. Due to the symmetry relations (17) it is sufficient to consider the region $x \leq 1, y \leq 1$. We can solve the differential equations (15) along one of the two paths in Fig. 2. The result is a combination of products of Goncharov polylogarithms [14]

$$G_{\underbrace{0, \dots, 0}_n}(x) = \frac{1}{n!} \log^n x, \quad G_{a, \dots}(x) = \int_0^x \frac{dt}{t-a} G_{\dots}(t)$$

of $\bar{x} = 1 - x$ and $\bar{y} = 1 - y$. Numerical evaluation of Goncharov polylogarithms is available [15] in GiNaC [16]. We make no efforts to express some of them via harmonic polylogarithms of x and y because some Goncharov polylogarithms are bound to remain. Using Libra we obtain

$$\begin{aligned}
y^{-l\varepsilon} F_1(x, y) = & 1 + 2l_1 \{ l [G_{\bar{y},1}(\bar{x}) - G_{\bar{y},1}(\bar{x}) - 2G_{0,1}(\bar{y}) + G_1(\bar{y})(G_1(\bar{x}) - G_{\bar{y}}(\bar{x}) - G_{\bar{y}}(\bar{x}))] + (l_1 - l_2 + l_0)G_{1,1}(\bar{x}) \} \varepsilon^2 \\
& - 2l_1 \{ l [l_0(2(G_{\bar{y},0,1}(\bar{x}) - G_{\bar{y},0,1}(\bar{x}) + G_{\bar{y},2,1}(\bar{x}) - G_{\bar{y},2,1}(\bar{x})) + G_{\bar{y},\bar{y},1}(\bar{x}) - G_{\bar{y},\bar{y},1}(\bar{x}) + G_1(\bar{y})(G_{\bar{y},\bar{y}}(\bar{x}) + G_{\bar{y},\bar{y}}(\bar{x}) - G_{\bar{y},1}(\bar{x}))) \\
& + (2l - l_0)(G_{\bar{y},\bar{y},1}(\bar{x}) - G_{\bar{y},\bar{y},1}(\bar{x}) - G_1(\bar{y})(G_{\bar{y},\bar{y}}(\bar{x}) + G_{\bar{y},\bar{y}}(\bar{x}) - G_{\bar{y},1}(\bar{x}))) \\
& + 2(l_1 + l_2)(G_{0,1}(\bar{y})(G_1(\bar{x}) - G_{\bar{y}}(\bar{x}) - G_{\bar{y}}(\bar{x})) - 2G_{0,0,1}(\bar{y})) + (2l_1 + l_0)(G_{1,\bar{y},1}(\bar{x}) + G_1(\bar{y})G_{1,\bar{y}}(\bar{x})) + (2l_1 - l_0)G_{\bar{y},1,1}(\bar{x}) \\
& - (2l_2 - l_0)(G_{\bar{y},1,1}(\bar{x}) + G_{1,\bar{y},1}(\bar{x}) - G_1(\bar{y})(G_{1,\bar{y}}(\bar{x}) - G_{1,1}(\bar{x}))) + 2(l_1 + l_0)(G_{1,1}(\bar{y})(G_1(\bar{x}) - G_{\bar{y}}(\bar{x}) - G_{\bar{y}}(\bar{x})) - 2G_{0,1,1}(\bar{y}))] \\
& - 2l_0(l_1 - l_2 + l_0)(G_{1,0,1}(\bar{x}) + G_{1,2,1}(\bar{x})) - (2(l_1^2 - l_2^2) + (l_1 + l_2)l_0 - l_0^2)G_{1,1,1}(\bar{x}) \} \varepsilon^3 + \mathcal{O}(\varepsilon^4) \\
= & 1 + 2l_1 \{ l [G_1(\bar{x})(G_{\bar{x}}(\bar{y}) - G_{\bar{x}}(\bar{y})) - G_{\bar{x},1}(\bar{y}) - G_{\bar{x},1}(\bar{y})] + (l_1 - l_2 + l_0)G_{1,1}(\bar{x}) \} \varepsilon^2
\end{aligned}$$

$$\begin{aligned}
& -2l_1 \{ l [l_0 (2G_{2,1}(\bar{x})(G_{\hat{x}}(\bar{y}) - G_{\bar{x}}(\bar{y})) + G_1(\bar{x})(G_{\bar{x},\hat{x}}(\bar{y}) - G_{\hat{x},\bar{x}}(\bar{y})) + G_{\bar{x},\hat{x},1}(\bar{y}) + G_{\hat{x},\bar{x},1}(\bar{y})) \\
& \quad + (2l - l_0)(G_1(\bar{x})(G_{\bar{x},\bar{x}}(\bar{y}) - G_{\hat{x},\bar{x}}(\bar{y})) - G_{\bar{x},\hat{x},1}(\bar{y}) - G_{\hat{x},\bar{x},1}(\bar{y})) + 2(l_1 + l_2)G_{0,1}(\bar{x})(G_{\bar{x}}(\bar{y}) - G_{\hat{x}}(\bar{y})) + 2(l_1 - l_2)G_{0,1,1}(\bar{x}) \\
& \quad + (2l_1 - l_0)G_{1,1}(\bar{x})G_{\hat{x}}(\bar{y}) - (2l_2 - l_0)G_{1,1}(\bar{x})G_{\bar{x}}(\bar{y}) - 2(l_1 + l_0)(G_{\bar{x},1,1}(\bar{y}) + G_{\hat{x},1,1}(\bar{y}))] \\
& \quad - 2(l_1 - l_2 + l_0)[l_0G_{1,2,1}(\bar{x}) - (l_1 + l_2)G_{1,0,1}(\bar{x})] - (2(l_1^2 - l_2^2) + (l_1 + l_2)l_0 - l_0^2)G_{1,1,1}(\bar{x})\} \varepsilon^3 + \mathcal{O}(\varepsilon^4), \\
y^{le} F_2(x, y) &= 1 + 2l_2 \{ l [2(G_{0,1}(\bar{y}) - G_{1,1}(\bar{y})) - G_{\bar{y},1}(\bar{x}) + G_{\hat{y},1}(\bar{x}) - G_1(\bar{y})(G_1(\bar{x}) - G_{\hat{y}}(\bar{x}) - G_{\bar{y}}(\bar{x}))] - (l_1 - l_2 - l_0)G_{1,1}(\bar{x})\} \varepsilon^2 \\
& - 2l_2 \{ l [l_0(G_1(\bar{y})(G_{\bar{y},1}(\bar{x}) - G_{\bar{y},\hat{y}}(\bar{x}) - G_{\hat{y},\bar{y}}(\bar{x})) + 2(G_{\bar{y},0,1}(\bar{x}) - G_{\hat{y},0,1}(\bar{x}) + G_{\bar{y},2,1}(\bar{x}) - G_{\hat{y},2,1}(\bar{x})) - G_{\bar{y},\hat{y},1}(\bar{x}) + G_{\hat{y},\bar{y},1}(\bar{x})) \\
& \quad + (2l - l_0)(G_1(\bar{y})(G_{\bar{y},\bar{y}}(\bar{x}) + G_{\hat{y},\bar{y}}(\bar{x}) - G_{\bar{y},1}(\bar{x})) - G_{\bar{y},\bar{y},1}(\bar{x}) + G_{\hat{y},\bar{y},1}(\bar{x})) \\
& \quad + (l_1 + l_2)(4(G_{0,0,1}(\bar{y}) - G_{1,0,1}(\bar{y})) - 2G_{0,1}(\bar{y})(G_1(\bar{x}) - G_{\bar{y}}(\bar{x}) - G_{\hat{y}}(\bar{x}))) \\
& \quad - (2l_1 - l_0)(G_1(\bar{y})(G_{1,\hat{y}}(\bar{x}) - G_{1,1}(\bar{x})) + G_{\hat{y},1,1}(\bar{x}) + G_{1,\hat{y},1}(\bar{x})) - (2l_2 + l_0)(G_1(\bar{y})G_{1,\bar{y}}(\bar{x}) - G_{1,\bar{y},1}(\bar{x})) \\
& \quad + (2l_2 - l_0)G_{\bar{y},1,1}(\bar{x}) + 2(l_2 + l)(2(G_{1,1,1}(\bar{y}) - G_{0,1,1}(\bar{y})) + G_{1,1}(\bar{y})(G_1(\bar{x}) - G_{\bar{y}}(\bar{x}) - G_{\hat{y}}(\bar{x})))] \\
& \quad + 2l_0(l_1 - l_2 - l_0)(G_{1,0,1}(\bar{x}) + G_{1,2,1}(\bar{x})) + (2(l_1^2 - l_2^2) - (l_1 + l_2)l_0 + l_0^2)G_{1,1,1}(\bar{x})\} \varepsilon^3 + \mathcal{O}(\varepsilon^4) \\
&= 1 + 2l_2 \{ l [G_1(\bar{x})(G_{\hat{x}}(\bar{y}) - G_{\bar{x}}(\bar{y})) - 2G_{1,1}(\bar{y}) + G_{\bar{x},1}(\bar{y}) + G_{\hat{x},1}(\bar{y})] - (l_1 - l_2 - l_0)G_{1,1}(\bar{x})\} \varepsilon^2 \\
& - 2l_2 \{ 2l^2 G_1(\bar{x})(G_{\bar{x},1}(\bar{y}) - G_{\hat{x},1}(\bar{y})) + l [l_0(2G_{2,1}(\bar{x})(G_{\bar{x}}(\bar{y}) - G_{\hat{x}}(\bar{y})) + G_1(\bar{x})(G_{\bar{x},\bar{x}}(\bar{y}) - G_{\hat{x},\bar{x}}(\bar{y})) - G_{\bar{x},\hat{x},1}(\bar{y}) - G_{\hat{x},\bar{x},1}(\bar{y})) \\
& \quad + (2l - l_0)(G_1(\bar{x})(G_{\hat{x},\hat{x}}(\bar{y}) - G_{\bar{x},\bar{x}}(\bar{y})) + G_{\bar{x},\bar{x},1}(\bar{y}) + G_{\hat{x},\hat{x},1}(\bar{y})) \\
& \quad + 2(l_1 + l_2)(G_{0,1}(\bar{x})(G_{\hat{x}}(\bar{y}) - G_{\bar{x}}(\bar{y})) + G_1(\bar{x})(G_{1,\bar{x}}(\bar{y}) - G_{1,\hat{x}}(\bar{y})) - G_{1,\bar{x},1}(\bar{y}) - G_{1,\hat{x},1}(\bar{y})) \\
& \quad + 2(l_1 - l_2)(G_{1,1}(\bar{x})G_1(\bar{y}) - G_{0,1,1}(\bar{x})) - (2l_1 - l_0)G_{1,1}(\bar{x})G_{\hat{x}}(\bar{y}) + (2l_2 - l_0)G_{\bar{x}}(\bar{y})G_{1,1}(\bar{x}) \\
& \quad + 2(l_2 + l)(2G_{1,1,1}(\bar{y}) - G_{\bar{x},1,1}(\bar{y}) - G_{\hat{x},1,1}(\bar{y}))] \\
& \quad + 2(l_1 - l_2 - l_0)[l_0G_{1,2,1}(\bar{x}) - (l_1 + l_2)G_{1,0,1}(\bar{x})] + (2(l_1^2 - l_2^2) - (l_1 + l_2)l_0 + l_0^2)G_{1,1,1}(\bar{x})\} \varepsilon^3 + \mathcal{O}(\varepsilon^4), \\
F_3(x, y) &= 2G_1(\bar{x})\varepsilon + 2 \{ l [G_1(\bar{y})(G_{\bar{y}}(\bar{x}) - G_{\hat{y}}(\bar{x})) - G_{\bar{y},1}(\bar{x}) - G_{\hat{y},1}(\bar{x})] + 2l_0(G_{0,1}(\bar{x}) + G_{2,1}(\bar{x})) + (l_1 + l_2 - l_0)G_{1,1}(\bar{x})\} \varepsilon^2 \\
& - 2 \{ l [l_0(G_1(\bar{y})(2(G_{0,\hat{y}}(\bar{x}) - G_{0,\bar{y}}(\bar{x}) + G_{2,\hat{y}}(\bar{x}) - G_{2,\bar{y}}(\bar{x})) - G_{\bar{y},\hat{y}}(\bar{x}) + G_{\hat{y},\bar{y}}(\bar{x})) \\
& \quad + 2(G_{0,\bar{y},1}(\bar{x}) + G_{0,\hat{y},1}(\bar{x}) + G_{2,\bar{y},1}(\bar{x}) + G_{2,\hat{y},1}(\bar{x}) + G_{\bar{y},0,1}(\bar{x}) + G_{\hat{y},0,1}(\bar{x}) + G_{\bar{y},2,1}(\bar{x}) + G_{\hat{y},2,1}(\bar{x})) - G_{\bar{y},\hat{y},1}(\bar{x}) - G_{\hat{y},\bar{y},1}(\bar{x})) \\
& \quad + (2l - l_0)(G_1(\bar{y})(G_{\bar{y},\bar{y}}(\bar{x}) - G_{\hat{y},\bar{y}}(\bar{x})) - G_{\bar{y},\bar{y},1}(\bar{x}) - G_{\hat{y},\bar{y},1}(\bar{x})) \\
& \quad + (l_1 + l_2)(2G_{0,1}(\bar{y})(G_{\bar{y}}(\bar{x}) - G_{\hat{y}}(\bar{x})) - G_1(\bar{y})(G_{\bar{y},1}(\bar{x}) - G_{\hat{y},1}(\bar{x}))) + (l_1 - l_2)(2G_{0,1}(\bar{y})G_1(\bar{x}) - G_1(\bar{y})G_{1,1}(\bar{x})) \\
& \quad + (2l_1 - l_0)(G_1(\bar{y})G_{1,\hat{y}}(\bar{x}) + G_{\hat{y},1,1}(\bar{x}) + G_{1,\hat{y},1}(\bar{x})) - (2l_2 - l_0)(G_1(\bar{y})G_{1,\bar{y}}(\bar{x}) - G_{1,\bar{y},1}(\bar{x}) - G_{\bar{y},1,1}(\bar{x})) \\
& \quad - (l_1 - l_2 + l_0)G_1(\bar{x})G_{1,1}(\bar{y}) + 2l_2 G_{1,1}(\bar{y})(G_{\hat{y}}(\bar{x}) - G_{\bar{y}}(\bar{x}))] \\
& \quad - 4l_0^2(G_{0,0,1}(\bar{x}) + G_{0,2,1}(\bar{x}) + G_{2,0,1}(\bar{x}) + G_{2,2,1}(\bar{x})) - 2l_0(l_1 + l_2 - l_0)(G_{0,1,1}(\bar{x}) + G_{1,0,1}(\bar{x}) + G_{1,2,1}(\bar{x}) + G_{2,1,1}(\bar{x})) \\
& \quad - (2(l_1^2 + l_2^2) - (l_1 + l_2)l_0 + l_0^2)G_{1,1,1}(\bar{x})\} \varepsilon^3 + \mathcal{O}(\varepsilon^4) \\
&= 2G_1(\bar{x})\varepsilon + 2 \{ l [G_1(\bar{x})(G_1(\bar{y}) - G_{\bar{x}}(\bar{y}) - G_{\hat{x}}(\bar{y})) + G_{\bar{x},1}(\bar{y}) - G_{\hat{x},1}(\bar{y})] + (l_1 + l_2 - l_0)G_{1,1}(\bar{x}) - 2(l_1 + l_2)G_{0,1}(\bar{x}) + 2l_0G_{2,1}(\bar{x})\} \varepsilon^2 \\
& - 2 \{ l^2 [G_1(\bar{x})(G_{1,\bar{x}}(\bar{y}) + G_{1,\hat{x}}(\bar{y}) + G_{\bar{x},1}(\bar{y}) + G_{\hat{x},1}(\bar{y}) - G_{1,1}(\bar{y})) - G_{1,\bar{x},1}(\bar{y}) + G_{1,\hat{x},1}(\bar{y})] \\
& \quad + l [l_0(2G_{2,1}(\bar{x})(G_{\bar{x}}(\bar{y}) + G_{\hat{x}}(\bar{y}) - G_1(\bar{y})) - G_1(\bar{x})(G_{\bar{x},\bar{x}}(\bar{y}) + G_{\hat{x},\bar{x}}(\bar{y})) - G_{\bar{x},\hat{x},1}(\bar{y}) + G_{\hat{x},\bar{x},1}(\bar{y})) \\
& \quad - (2l - l_0)[G_1(\bar{x})(G_{\bar{x},\bar{x}}(\bar{y}) + G_{\hat{x},\bar{x}}(\bar{y})) - G_{\bar{x},\bar{x},1}(\bar{y}) + G_{\hat{x},\bar{x},1}(\bar{y})] + 2(l_1 + l_2)[G_{0,1}(\bar{x})(G_1(\bar{y}) - G_{\hat{x}}(\bar{y})) - G_{0,1}(\bar{x})G_{\bar{x}}(\bar{y})] \\
& \quad + G_{1,1}(\bar{x})[(2l_1 - l_0)G_{\hat{x}}(\bar{y}) + (2l_2 - l_0)G_{\bar{x}}(\bar{y}) - (l_1 + l_2 - l_0)G_1(\bar{y})] - 2l_2(G_{\bar{x},1,1}(\bar{y}) - G_{\hat{x},1,1}(\bar{y}))] \\
& \quad - 4(l_1 + l_2)^2 G_{0,0,1}(\bar{x}) - 2(l_1 + l_2 - l_0)[l_0(G_{2,1,1}(\bar{x}) + G_{1,2,1}(\bar{x})) - (l_1 + l_2)(G_{1,0,1}(\bar{x}) + G_{0,1,1}(\bar{x}))] \\
& \quad + 4l_0(l_1 + l_2)(G_{2,0,1}(\bar{x}) + G_{0,2,1}(\bar{x})) - 4l_0^2 G_{2,2,1}(\bar{x}) - (2(l_1^2 + l_2^2) - (l_1 + l_2)l_0 + l_0^2)G_{1,1,1}(\bar{x})\} \varepsilon^3 + \mathcal{O}(\varepsilon^4), \tag{27}
\end{aligned}$$

where $\hat{x} = 1 - x^{-1}$, $\hat{y} = 1 - y^{-1}$. All Goncharov polylogarithms up to weight 2 can be expressed via Li_2 and logarithms. This expansion can be straightforwardly extended to any order in ε .

I am grateful to R. N. Lee for numerous consultations on Libra. The work was supported by the Russian ministry of science and higher education.

- (1993), arXiv:hep-ph/9209277.
- [3] A. V. Kotikov, Phys. Lett. **B254**, 158 (1991).
 - [4] J. M. Henn, Phys. Rev. Lett. **110**, 251601 (2013), arXiv:1304.1806 [hep-th].
 - [5] R. N. Lee, “Libra,” (2018–2020), available from the author by request.
 - [6] R. N. Lee, JHEP **04**, 108 (2015), arXiv:1411.0911 [hep-ph].
 - [7] E. Remiddi and J. Vermaseren, Int. J. Mod. Phys. A **15**, 725 (2000), arXiv:hep-ph/9905237.
 - [8] D. Maître, Comput. Phys. Commun. **174**, 222 (2006), arXiv:hep-ph/0507152 [hep-ph].
 - [9] D. Maître, Comput. Phys. Commun. **183**, 846 (2012), arXiv:hep-ph/0703052 [hep-ph].
 - [10] T. Huber and D. Maître, Comput. Phys. Commun. **175**, 122 (2006), arXiv:hep-ph/0507094 [hep-ph].
 - [11] T. Huber and D. Maître, Comput. Phys. Commun. **178**, 755 (2008), arXiv:0708.2443 [hep-ph].
 - [12] A. I. Davydychev and M. Kalmykov, Nucl. Phys. B **605**, 266 (2001), arXiv:hep-th/0012189.
 - [13] A. Grozin, Eur. Phys. J. C **77**, 453 (2017), arXiv:1704.07968 [hep-ph].
 - [14] A. B. Goncharov, Math. Res. Lett. **5**, 497 (1998), arXiv:1105.2076 [math.AG].
 - [15] J. Vollinga and S. Weinzierl, Comput. Phys. Commun. **167**, 177 (2005), arXiv:hep-ph/0410259.
 - [16] C. W. Bauer, A. Frink, and R. Kreckel, J. Symb. Comput. **33**, 1 (2002), <https://ginac.de/>, arXiv:cs/0004015.