

Hypergeometric SLE with $\kappa = 8$: Convergence of UST and LERW in Topological Rectangles

Yong Han¹, Mingchang Liu², and Hao Wu^{*3}

¹Shenzhen University, China. Email: hanyongprobability@gmail.com

²Tsinghua University, China. Email: liumc_prob@163.com

³Tsinghua University, China. Email: hao.wu.proba@gmail.com

Abstract

We consider uniform spanning tree (UST) in topological rectangles with alternating boundary conditions. The Peano curves associated to the UST converge weakly to hypergeometric SLE₈, denoted by hSLE₈. From the convergence result, we obtain the continuity and reversibility of hSLE₈ as well as an interesting connection between SLE₈ and hSLE₈. The loop-erased random walk (LERW) branch in the UST converges weakly to SLE₂(-1, -1; -1, -1). We also obtain the limiting joint distribution of the two end points of the LERW branch.

Keywords: uniform spanning tree, loop-erased random walk, Schramm Loewner evolution.
MSC: 60J67

Contents

1	Introduction	2
1.1	Hypergeometric SLE with $\kappa = 8$	2
1.2	Uniform spanning tree (UST)	3
1.3	Loop-erased random walk (LERW)	4
2	Preliminaries on SLE	5
3	Hypergeometric SLE with $\kappa \geq 8$	7
3.1	Continuity of hSLE with $\kappa \geq 8$	7
3.2	Discussion on time-reversal of hSLE	10
3.3	Continuity and reversibility of hSLE with $\kappa = 8$	10
4	Convergence of UST in quads	12
4.1	UST with Dobrushin boundary conditions	13
4.2	UST with alternating boundary conditions in quads	16
4.3	Discrete observable for UST in quads	19
4.4	Proof of Theorem 4.2	20
4.5	Proof of Theorem 1.5	25
5	Convergence of LERW in quads	27
5.1	The pair of random points (X^M, Y^M)	27
5.2	Proof of Theorem 1.6	32
5.3	Consequences	39
A	Hypergeometric function and elliptic integral	40
B	Convergence of discrete harmonic functions	41

*Supported by Chinese Thousand Talents Plan for Young Professionals and Beijing Natural Science Foundation (Z180003).

1 Introduction

In [Sch00], O. Schramm introduced a random process—Schramm Loewner Evolution (SLE)—as a candidate for the scaling limit of interfaces in two-dimensional critical lattice models. The setup is as follows. A Dobrushin domain $(\Omega; x, y)$ is a bounded simply connected domain $\Omega \subsetneq \mathbb{C}$ with two boundary points x, y such that $\partial\Omega$ is locally connected. We denote by (xy) the boundary arc going from x to y in counterclockwise order. Suppose $(\Omega_\delta; x_\delta, y_\delta)$ is an approximation of $(\Omega; x, y)$ on $\delta\mathbb{Z}^2$. Consider a critical lattice model on Ω_δ with certain Dobrushin boundary conditions, for instance, Ising model, percolation, uniform spanning tree etc. In these examples, it comes naturally an interface in Ω_δ connecting x_δ to y_δ . It is conjectured that such interface has a conformally invariant scaling limit which can be identified by SLE_κ where the parameter κ varies for different models. Since the introduction of SLE, there are several models for which the conjecture is proved: the Peano curve in uniform spanning tree converges to SLE_8 and the loop-erased random walk converges to SLE_2 [LSW04], the interface in percolation converges to SLE_6 [Smi01], the level line of discrete Gaussian free field converges to SLE_4 [SS09], the interface in Ising model converges to SLE_3 and the interface in FK-Ising model converges to $\text{SLE}_{16/3}$ [CDCH⁺14]. The proof requires two inputs: 1st. the tightness of interfaces; 2nd. discrete martingale observable. With the tightness, there is always subsequential limits of interfaces; and then one uses the observable to identify the subsequential limits.

Dobrushin boundary conditions are the simplest. It is then natural to consider critical lattice models with more complicated boundary conditions. One possibility is to consider the model in polygons with alternating boundary conditions. In general, a polygon $(\Omega; x_1, \dots, x_p)$ is a bounded simply connected domain $\Omega \subset \mathbb{C}$ with distinct boundary points x_1, \dots, x_p in counterclockwise order, such that $\partial\Omega$ is locally connected. In this article, we focus on topological rectangles $(\Omega; a, b, c, d)$, i.e. a polygon with four marked points on the boundary, and we call it a quad. Suppose $(\Omega_\delta; a_\delta, b_\delta, c_\delta, d_\delta)$ is an approximation of $(\Omega; a, b, c, d)$ on $\delta\mathbb{Z}^2$. Consider a critical lattice model on Ω_δ with alternating boundary conditions, it turns out that the scaling limit of interfaces in this case becomes hypergeometric SLE, denoted by hSLE, which is a variant of SLE process. For instance, the interface in critical Ising model in quad converges to hSLE₃, see [Izy15] and [Wu20], the interface in critical FK-Ising model in quad converges to hSLE_{16/3}, see [KS18] and [BPW18]. In this article, we focus on uniform spanning tree in quad with alternating boundary conditions. Not surprisingly, the associated Peano curve converges to hSLE₈ process. When the third author of this article introduced hSLE process in [Wu20], she only treated the process with $\kappa \in (0, 8)$ due to technical difficulty. The first goal of this article is to address hSLE process with $\kappa = 8$.

1.1 Hypergeometric SLE with $\kappa = 8$

Hypergeometric SLE is a two-parameter family of random curves in quad. The two parameters are $\kappa > 0$ and $\nu \in \mathbb{R}$, and we denote it by $\text{hSLE}_\kappa(\nu)$. The continuity and reversibility of such process are addressed in [Wu20] for $\kappa \in (0, 8)$. Our first main result is about the continuity and reversibility of $\text{hSLE}_\kappa(\nu)$ with $\kappa = 8$.

Theorem 1.1. *Fix $\nu \geq 0$ and $x_1 < x_2 < x_3 < x_4$. The process $\text{hSLE}_8(\nu)$ in the upper half-plane \mathbb{H} from x_1 to x_4 with marked points (x_2, x_3) is almost surely generated by a continuous curve denoted by η . Furthermore, the process η enjoys reversibility: the time-reversal of η has the law of $\text{hSLE}_8(\nu)$ in \mathbb{H} from x_4 to x_1 with marked points (x_3, x_2) .*

We denote $\text{hSLE}_8(\nu)$ by hSLE_8 when $\nu = 0$, and the rest of the introduction will focus on hSLE_8 . We will discuss the connection between SLE_8 and hSLE_8 . To this end, we first generalize the definition of hSLE_8 for general quad. For a quad $(\Omega; a, b, c, d)$, let ϕ be any conformal map from Ω onto \mathbb{H} such that $\phi(a) < \phi(b) < \phi(c) < \phi(d)$. We define hSLE_8 in Ω from a to d with marked points (b, c) to be $\phi^{-1}(\eta)$ where η is an hSLE_8 in \mathbb{H} from $\phi(a)$ to $\phi(d)$ with marked points $(\phi(b), \phi(c))$. Let us discuss the connection between SLE_8 and hSLE_8 . On the one hand, hSLE_8 degenerates to SLE_8 when the marked points collapse, see Lemma 1.2. On the other hand, we may also find hSLE_8 inside SLE_8 , see Proposition 1.3.

Lemma 1.2. *Suppose $x_1 < x_2 < x_3 < x_4$. Suppose $\eta \sim \text{hSLE}_8$ in \mathbb{H} from x_1 to x_4 with marked points (x_2, x_3) . Then, let $x_3 \rightarrow x_4$, the law of η converges weakly to $\text{SLE}_8(2)$ in \mathbb{H} from x_1 to x_4 with force point x_2 . If we further let $x_2 \rightarrow x_4$, the law of $\text{SLE}_8(2)$ converges weakly to SLE_8 in \mathbb{H} from x_1 to x_4 .*

Proposition 1.3. *Fix $x < y$ and suppose $\eta \sim \text{SLE}_8$ in \mathbb{H} from x to ∞ . Let T_y be the first time that η swallows y , and denote by γ the right boundary of $\eta[0, T_y]$. Note that γ is a continuous simple curve starting from y and terminating at some point in $(-\infty, x)$. Let τ be any stopping time for γ before the terminating time. Then the conditional law of $(\eta(t), 0 \leq t \leq T_y)$ given $\gamma[0, \tau]$ is hSLE_8 in $\mathbb{H} \setminus \gamma[0, \tau]$ from x to y^- with marked points (y^+, ∞) conditional that its first hitting point on $\gamma[0, \tau]$ is given by $\gamma(\tau)$.*

In Section 2, we will give preliminaries on SLE; and in Section 3, we will give definition of $\text{hSLE}_\kappa(\nu)$. In fact, we will address $\text{hSLE}_\kappa(\nu)$ for $\kappa \geq 8$ in Section 3. As the case of $\kappa > 8$ is less relevant, we omit the corresponding conclusion in the introduction. We will derive some partial result towards continuity of hSLE_8 in Section 3. The full continuity result will be based on the convergence of UST, see Section 1.2. Note also that Lemma 1.2 holds for general $\kappa > 0$. However, Proposition 1.3 only holds for $\kappa = 8$ and its proof is based on an interesting observation for UST which will be given in Section 5. Moreover, the calculation in Section 5 also gives the following consequence.

Proposition 1.4. *Fix a quad $(\Omega; a, b, c, d)$. Let $K > 0$ be the conformal modulus of the quad $(\Omega; a, b, c, d)$, and let f be the conformal map from Ω onto the rectangle $(0, 1) \times (0, iK)$ which sends (a, b, c, d) to $(0, 1, 1 + iK, iK)$. Suppose $\eta \sim \text{hSLE}_8$ in Ω from a to d with marked points (b, c) . Then we have*

$$\mathbb{P}[z \notin \eta] = \text{Re}f(z), \quad \forall z \in \Omega.$$

1.2 Uniform spanning tree (UST)

Let us come back to uniform spanning tree. The square lattice \mathbb{Z}^2 is the graph with vertex set $V(\mathbb{Z}^2) := \{(m, n) : m, n \in \mathbb{Z}\}$ and edge set $E(\mathbb{Z}^2)$ given by edges between nearest neighbors. This is our primal lattice. Its dual lattice is denoted by $(\mathbb{Z}^2)^*$. The medial lattice $(\mathbb{Z}^2)^\diamond$ is the graph with centers of edges of \mathbb{Z}^2 as vertex set and edges connecting nearest vertices. Suppose a sequence of medial quads $(\Omega_\delta^\diamond; a_\delta^\diamond, b_\delta^\diamond, c_\delta^\diamond, d_\delta^\diamond)$ approximates some quad $(\Omega; a, b, c, d)$, and let $\Omega_\delta \subset \mathbb{Z}^2$ be the corresponding graph on the primal lattice, see details in Section 4.2. We consider uniform spanning tree (UST) on Ω_δ with alternating boundary conditions: the edges in the boundary arcs $(a_\delta b_\delta)$ and $(c_\delta d_\delta)$ are forced to be contained in the UST. Let \mathcal{T}_δ be the UST on Ω_δ with such alternating boundary conditions. Then there exists a triple of curves $(\eta_\delta^L; \gamma_\delta^M; \eta_\delta^R)$ such that η_δ^L runs along the tree \mathcal{T}_δ from a_δ to d_δ , and γ_δ^M is the unique branch in \mathcal{T}_δ connecting $(a_\delta b_\delta)$ to $(c_\delta d_\delta)$, and η_δ^R runs along the tree \mathcal{T}_δ from b_δ to c_δ , see Figure 1.1 and see detail in Section 4.2. We have the following convergence of the triple $(\eta_\delta^L; \gamma_\delta^M; \eta_\delta^R)$.

Theorem 1.5. *Fix a quad $(\Omega; a, b, c, d)$ such that $\partial\Omega$ is C^1 and simple. Suppose that a sequence of medial quads $(\Omega_\delta^\diamond; a_\delta^\diamond, b_\delta^\diamond, c_\delta^\diamond, d_\delta^\diamond)$ converges to $(\Omega; a, b, c, d)$ in the following sense (see (4.2)):*

$$(a_\delta^\diamond b_\delta^\diamond) \rightarrow (ab), \quad (b_\delta^\diamond c_\delta^\diamond) \rightarrow (bc), \quad (c_\delta^\diamond d_\delta^\diamond) \rightarrow (cd), \quad (d_\delta^\diamond a_\delta^\diamond) \rightarrow (da) \quad \text{as curves, as } \delta \rightarrow 0.$$

Consider the UST in $(\Omega_\delta; a_\delta, b_\delta, c_\delta, d_\delta)$ with alternating boundary conditions and consider the triple of curves $(\eta_\delta^L; \gamma_\delta^M; \eta_\delta^R)$ as described above. Then the triple $(\eta_\delta^L; \gamma_\delta^M; \eta_\delta^R)$ converges weakly to a triple of continuous curves $(\eta^L; \gamma^M; \eta^R)$ whose law is characterized by the following properties: the marginal law of η^L is hSLE_8 in Ω from a to d with marked points (b, c) ; given η^L , the conditional law of η^R is SLE_8 in $\Omega \setminus \eta^L$ from b to c ; and $\gamma^M = \eta^L \cap \eta^R$.

Furthermore, given γ^M , denote by Ω^L and Ω^R the two connected components of $\Omega \setminus \gamma^M$ such that Ω^L has a, d on the boundary and Ω^R has b, c on the boundary, then the conditional law of η^L is SLE_8 in Ω^L from a to d and the conditional law of η^R is SLE_8 in Ω^R from b to c , and η^L and η^R are conditionally independent given γ^M .




Figure 1.1: The solid edges in black are wired boundary arcs (ab) and (cd) , the solid edges in red are dual-wired boundary arcs (b^*c^*) and (d^*a^*) . The thin edges are in the tree \mathcal{T} . The solid edges in green are in the branch γ^M in \mathcal{T} . This branch intersects (ab) at X^M and intersects (cd) at Y^M . The two curves in orange are the Peano curves η^L (from a to d) and η^R (from b to c).

In Section 4, we introduce UST and prove the convergence of the Peano curve and complete the proof of Theorem 1.5. The strategy is similar as for Dobrushin boundary conditions: we first derive the tightness of the Peano curves and construct a discrete martingale observable, and then identify the subsequential limits through the observable. This part is a generalization of the work in [Sch00] and [LSW04], and we proceed following a simplification advocated by Smirnov [Smi06] and [DCS12]. As a byproduct, we obtain the continuity and reversibility of hSLE₈ and complete the proof of Theorem 1.1.

1.3 Loop-erased random walk (LERW)

From Theorem 1.5, we see that the Peano curve η_δ^L converges weakly to hSLE₈ and the limit of γ_δ^M is part of the boundary of hSLE₈. In the following theorem, we provide an explicit characterization of the limiting distribution of γ_δ^M .

Theorem 1.6. *Fix a quad $(\Omega; a, b, c, d)$ such that $\partial\Omega$ is C^1 and simple. Let $K > 0$ be the conformal modulus of the quad $(\Omega; a, b, c, d)$, and let f be the conformal map from Ω onto the rectangle $(0, 1) \times (0, iK)$ which sends (a, b, c, d) to $(0, 1, 1 + iK, iK)$. Assume the same setup as in Theorem 1.5. Then the law of γ_δ^M converges weakly to a continuous curve γ^M whose law is characterized by the following properties. Denote by $X^M = \gamma^M \cap (ab)$ and by $Y^M = \gamma^M \cap (cd)$.*

- (1) *The law of the point $f(X^M)$ is uniform in $(0, 1)$.*
- (2) *Given X^M , the conditional law of γ^M is an SLE₂ $(-1, -1; -1, -1)$ in Ω from X^M to (cd) with force points $(d, a; b, c)$ stopped at the first hitting time of (cd) .*

Furthermore, denote by $x^M = f(X^M)$ and $y^M = \text{Re}f(Y^M)$, the joint density of (x^M, y^M) is given by

$$\rho_K(x, y) = \frac{\pi}{4K} \sum_{n \in \mathbb{Z}} \left(\frac{1}{\cosh^2\left(\frac{\pi}{2K}(x - y - 2n)\right)} + \frac{1}{\cosh^2\left(\frac{\pi}{2K}(x + y - 2n)\right)} \right), \quad \forall x, y \in (0, 1). \quad (1.1)$$

In Section 5, we work on the LERW branch and complete the proof of Theorem 1.6. We first derive the joint distribution of the pair (X^M, Y^M) in Section 5.1. We derive the formula (1.1) through discrete observable. The analysis on discrete harmonic function from [CW19] plays an important role. We then derive the conditional law of γ^M given X^M in Section 5.2. The derivation follows the standard strategy: showing the tightness and constructing discrete martingale observable. In fact, this part of the conclusion is already solved in [Zha08c] in an implicit form for more generality. We provide detail of the proof in a self-contained way for our particular setting and derive the explicit answer in Section 5.2. In Section 5.3, we provide an interesting observation for UST and complete the proof of Proposition 1.3. We also remark that the proof for the law of γ^M in Theorem 1.6 also provides an alternative proof for the duality result of SLE₈, see Corollary 1.7. Such duality relation was previously proved in [Zha08a] and [MS16a] in the continuous setting for general κ . Our proof in Section 5.3 is specific for $\kappa = 8$ because we use the convergence of UST and LERW.

Corollary 1.7. *Fix $x < y$ and suppose η is an SLE₈ in \mathbb{H} from x to ∞ . Let T_y be the first time that η swallows y , and denote by γ the right boundary of $\eta[0, T_y]$. Then the law of γ is the same as SLE₂(-1, -1; -1, -1) in \mathbb{H} from y to $(-\infty, x)$ with force points $(x, y^-; y^+, \infty)$.*

Remark 1.8. *The C^1 -regularity on $\partial\Omega$ in Theorem 1.5 plays a crucial role when we derive the tightness of the Peano curves, see the discussion in the proof of Theorem 4.1. In contrast, the C^1 -regularity on $\partial\Omega$ in Theorem 1.6 is not necessary for the conclusion to be true. But we still keep the assumption as it simplifies the proof.*

Acknowledgment. We thank Yijun Wan and Dapeng Zhan for helpful discussion on UST and LERW.

2 Preliminaries on SLE

Notations

For $z \in \mathbb{C}$ and $r > 0$, we denote by $B(z, r)$ the ball with center x and radius r . In particular, we denote $B(0, 1)$ by \mathbb{U} .

Loewner chain

An \mathbb{H} -hull is a compact subset K of $\overline{\mathbb{H}}$ such that $\mathbb{H} \setminus K$ is simply connected. By Riemann's mapping theorem, there exists a unique conformal map g_K from $\mathbb{H} \setminus K$ onto \mathbb{H} with normalization $\lim_{z \rightarrow \infty} |g_K(z) - z| = 0$, and we call $a(K) := \lim_{z \rightarrow \infty} z(g_K(z) - z)$ the half-plane capacity of K seen from ∞ . Loewner chain is a collection of \mathbb{H} -hulls $(K_t, t \geq 0)$ associated to the family of conformal maps $(g_t, t \geq 0)$ which solves the following Loewner equation: for each $z \in \mathbb{H}$,

$$\partial_t g_t(z) = \frac{2}{g_t(z) - W_t}, \quad g_0(z) = z,$$

where $(W_t, t \geq 0)$ is a one-dimensional continuous function which we call the driving function. For $z \in \overline{\mathbb{H}}$, the swallowing time of z is defined to be $\sup \{t \geq 0 : \min_{s \in [0, t]} |g_s(z) - W_s| > 0\}$. Let K_t be the closure of $\{z \in \mathbb{H} : T_z \leq t\}$. It turns out that g_t is the unique conformal map from $\mathbb{H} \setminus K_t$ onto \mathbb{H} with normalization $\lim_{z \rightarrow \infty} |g_t(z) - z| = 0$. Since the half-plane capacity of K_t is $\lim_{z \rightarrow \infty} z(g_t(z) - z) = 2t$, we say that the process $(K_t, t \geq 0)$ is parameterized by the half-plane capacity. We say that $(K_t, t \geq 0)$ can be generated by continuous curve $(\eta(t), t \geq 0)$ if, for any t , the unbounded connected component of $\mathbb{H} \setminus \eta[0, t]$ is the same as $\mathbb{H} \setminus K_t$.

Schramm Loewner evolution

Schramm Loewner evolution SLE_κ is the random Loewner chain driven by $W_t = \sqrt{\kappa}B_t$ where $\kappa > 0$ and $(B_t, t \geq 0)$ is one-dimension Brownian motion. SLE_κ process is almost surely generated by continuous curve η . The continuity for $\kappa \neq 8$ is proved in [RS05], and the continuity for $\kappa = 8$ is proved in [LSW04]. Moreover, the curve η is almost surely transient: $\lim_{t \rightarrow \infty} |\eta(t)| = \infty$. When $\kappa \in (0, 4]$, the curve is simple; when $\kappa \in (4, 8)$, the curve is self-touching; when $\kappa \geq 8$, the curve is space-filling.

In the above, SLE_κ is in \mathbb{H} from 0 to ∞ , we may define it in any Dobrushin domain $(\Omega; x, y)$ via conformal image: let ϕ be any conformal map from Ω onto \mathbb{H} such that $\phi(x) = 0$ and $\phi(y) = \infty$. We define SLE_κ in Ω from x to y to be $\phi^{-1}(\eta)$ where η is an SLE_κ in \mathbb{H} from 0 to ∞ . When $\kappa \in (0, 8]$, SLE_κ enjoys reversibility: suppose η is an SLE_κ in Ω from x to y , the time-reversal of η has the same law as SLE_κ in Ω from y to x , proved in [Zha08b], [MS16b], [MS16c].

$\text{SLE}_\kappa(\underline{\rho})$ process

$\text{SLE}_\kappa(\underline{\rho})$ process is a variant of SLE_κ where one keeps track of multiple marked points. Suppose $\underline{y}^L = (y^{L,l} < \dots < y^{L,1} \leq 0)$, $\underline{y}^R = (0 \leq y^{R,1} < y^{R,2} < \dots < y^{R,r})$ and $\underline{\rho}^L = (\rho^{L,l}, \dots, \rho^{L,1})$, $\underline{\rho}^R = (\rho^{R,1}, \dots, \rho^{R,r})$ with $\rho^{L,i}, \rho^{R,i} \in \mathbb{R}$. An $\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ process with force points $(\underline{y}^L; \underline{y}^R)$ is the Loewner chain driven by W_t which is the solution to the following system of SDEs:

$$\begin{cases} dW_t = \sqrt{\kappa}dB_t + \sum_{i=1}^l \frac{\rho^{L,i}dt}{W_t - V_t^{L,i}} + \sum_{i=1}^r \frac{\rho^{R,i}dt}{W_t - V_t^{R,i}}, & W_0 = 0; \\ dV_t^{L,i} = \frac{2dt}{V_t^{L,i} - W_t}, & V_0^{L,i} = y^{L,i}, \quad \text{for } 1 \leq i \leq l; \\ dV_t^{R,i} = \frac{2dt}{V_t^{R,i} - W_t}, & V_0^{R,i} = y^{R,i}, \quad \text{for } 1 \leq i \leq r; \end{cases}$$

where $(B_t, t \geq 0)$ is one-dimensional Brownian motion. We define the continuation threshold of $\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ to be the infimum of the time t for which

$$\text{either } \sum_{i: V_t^{L,i} = W_t} \rho^{L,i} \leq -2, \quad \text{or } \sum_{i: V_t^{R,i} = W_t} \rho^{R,i} \leq -2.$$

$\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ process is well-defined up to the continuation threshold and it is almost surely generated by continuous curve up to and including the continuation threshold, see [MS16a].

The law of $\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ is absolutely continuous with respect to SLE_κ , and we will give the Radon-Nikodym derivative below, see also [SW05]. To simplify the notation for the Radon-Nikodym derivative, we focus on $\text{SLE}_\kappa(\underline{\rho})$ process when all force points are located to the same side of the process. Consider $\text{SLE}_\kappa(\underline{\rho})$ with force points \underline{y} where $\underline{\rho} = (\rho_1, \dots, \rho_n) \in \mathbb{R}^n$ and $\underline{y} = (0 \leq y_1 < \dots < y_n)$. The law of $\text{SLE}_\kappa(\underline{\rho})$ with force points \underline{y} is absolutely continuous with respect to SLE_κ up to the first time that y_1 is swallowed, and the Radon-Nikodym derivative is M_t/M_0 where

$$M_t = \prod_{1 \leq i \leq n} \left(g'_t(y_i)^{\rho_i(\rho_i + 4 - \kappa)/(4\kappa)} (g_t(y_i) - W_t)^{\rho_i/\kappa} \right) \times \prod_{1 \leq i < j \leq n} (g_t(y_j) - g_t(y_i))^{\rho_i \rho_j / (2\kappa)}. \quad (2.1)$$

$\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ process can be defined in general polygons. Suppose $(\Omega; y^{L,l}, \dots, y^{L,1}, x, y^{R,1}, \dots, y^{R,r}, y)$ is a polygon with $l+r+2$ marked points. Let ϕ be any conformal map from Ω onto \mathbb{H} such that $\phi(x) = 0$ and $\phi(y) = \infty$. We define $\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ in Ω from x to y with force points $(y^{L,l}, \dots, y^{L,1}; y^{R,1}, \dots, y^{R,r})$ to be $\phi^{-1}(\eta)$ where η is an $\text{SLE}_\kappa(\underline{\rho}^L; \underline{\rho}^R)$ in \mathbb{H} from 0 to ∞ with force points $(\phi(y^{L,l}), \dots, \phi(y^{L,1}); \phi(y^{R,1}), \dots, \phi(y^{R,r}))$.

$\text{SLE}_2(-1, -1; -1, -1)$ process

Let us discuss $\text{SLE}_2(-1, -1; -1, -1)$ mentioned in Theorem 1.6. Suppose $(\Omega; d, a, x, b, c, y)$ is a polygon with six marked points and consider $\text{SLE}_2(-1, -1; -1, -1)$ in Ω from x to y with force points $(d, a; b, c)$.

Note that the total of the force point weights is -4 which is $2-6$. Such process is target independent in the following sense: for distinct $y_1, y_2 \in (cd)$, let η_i be the $\text{SLE}_2(-1, -1; -1, -1)$ in Ω from x to y_i with force points $(d, a; b, c)$, and let T_i be the first time that η_i hits (cd) (in fact, T_i is the continuation threshold of η_i) for $i = 1, 2$. Then the law of $(\eta_1(t), 0 \leq t \leq T_1)$ is the same as the law of $(\eta_2(t), 0 \leq t \leq T_2)$. See [SW05] for the target-independence for a general setup. As the law of $(\eta_i(t), 0 \leq t \leq T_i)$ does not depend on the location of the target point, we say that it is an $\text{SLE}_2(-1, -1; -1, -1)$ in Ω from x to (cd) with force points $(d, a; b, c)$.

3 Hypergeometric SLE with $\kappa \geq 8$

Define the hypergeometric function (see Appendix A):

$$F(z) := {}_2F_1\left(\frac{2\nu+4}{\kappa}, 1 - \frac{4}{\kappa}, \frac{2\nu+8}{\kappa}; z\right). \quad (3.1)$$

Note that it is a solution to the Euler's hypergeometric equation (A.2). Set

$$h = \frac{6-\kappa}{2\kappa}, \quad a = \frac{\nu+2}{\kappa}, \quad b = \frac{(\nu+2)(\nu+6-\kappa)}{4\kappa}. \quad (3.2)$$

For $x_1 < x_2 < x_3 < x_4$, define partition function

$$\mathcal{Z}_{\kappa, \nu}(x_1, x_2, x_3, x_4) = (x_4 - x_1)^{-2h} (x_3 - x_2)^{-2b} z^a F(z), \quad \text{where } z = \frac{(x_2 - x_1)(x_4 - x_3)}{(x_3 - x_1)(x_4 - x_2)}. \quad (3.3)$$

The process $\text{hSLE}_\kappa(\nu)$ in \mathbb{H} from x_1 to x_4 with marked points (x_2, x_3) is the Loewner chain driven by W_t which is the solution to the following SDEs:

$$\begin{cases} dW_t = \sqrt{\kappa} dB_t + \kappa(\partial_1 \log \mathcal{Z}_{\kappa, \nu})(W_t, V_t^2, V_t^3, V_t^4) dt, & W_0 = x_1; \\ dV_t^i = \frac{2dt}{V_t^i - W_t}, & V_0^i = x_i, \quad \text{for } i = 2, 3, 4; \end{cases} \quad (3.4)$$

where $(B_t, t \geq 0)$ is one-dimensional Brownian motion. It is clear that the process is well-defined up to the swallowing time of x_2 . Combining (3.4) and (A.2), the law of $\text{hSLE}_\kappa(\nu)$ is the same as SLE_κ in \mathbb{H} from x_1 to ∞ weighted by the following local martingale:

$$M_t = g_t'(x_2)^b g_t'(x_3)^b g_t'(x_4)^h \mathcal{Z}_{\kappa, \nu}(W_t, g_t(x_2), g_t(x_3), g_t(x_4)). \quad (3.5)$$

In [Wu20, Section 3], the author proved the continuity of $\text{hSLE}_\kappa(\nu)$ with $\kappa \in (0, 8)$. In this section, we will discuss the case for $\kappa \geq 8$: We will show that the process is well-defined up to the swallowing time of x_3 which is denoted by T_{x_3} ; moreover, it is generated by a continuous curve η up to and including T_{x_3} . After T_{x_3} , we continue the process as a standard SLE_κ from $\eta(T_{x_3})$ towards x_4 in the remaining domain.

In the above, we have defined hSLE in \mathbb{H} and we may extend the definition to general quad via conformal image: For a quad $(\Omega; a, b, c, d)$, let ϕ be any conformal map from Ω onto \mathbb{H} such that $\phi(a) < \phi(b) < \phi(c) < \phi(d)$. We define $\text{hSLE}_\kappa(\nu)$ in Ω from a to d with marked points (b, c) to be $\phi^{-1}(\eta)$ where η is an $\text{hSLE}_\kappa(\nu)$ in \mathbb{H} from $\phi(a)$ to $\phi(d)$ with marked points $(\phi(b), \phi(c))$.

3.1 Continuity of hSLE with $\kappa \geq 8$

To derive the continuity of $\text{hSLE}_\kappa(\nu)$, it is more convenient to work in \mathbb{H} with $x_1 = 0$ and $x_4 = \infty$. Consider $\text{hSLE}_\kappa(\nu)$ in \mathbb{H} from 0 to ∞ with marked points (x, y) where $0 < x < y$. In this case, the SDEs (3.4) becomes the following:

$$\begin{cases} dW_t = \sqrt{\kappa} dB_t + \frac{(\nu+2)dt}{W_t - V_t^x} + \frac{-(\nu+2)dt}{W_t - V_t^y} - \kappa \frac{F'(Z_t)}{F(Z_t)} \left(\frac{1-Z_t}{V_t^y - W_t} \right) dt, & W_0 = 0; \\ dV_t^x = \frac{2dt}{V_t^x - W_t}, & dV_t^y = \frac{2dt}{V_t^y - W_t}, \quad V_0^x = x, V_0^y = y; \quad \text{where } Z_t = \frac{V_t^x - W_t}{V_t^y - W_t}. \end{cases} \quad (3.6)$$

We denote by T_x the swallowing time of x and by T_y the swallowing time of y . The main result of this section is the continuity of the process up to and including T_y .

From (3.6), it is clear that the Loewner chain is well-defined up to T_x . As in (3.5), the process has the same law as SLE_κ in \mathbb{H} from 0 to ∞ weighted by the following local martingale:

$$M_t = g'_t(x)^b g'_t(y)^b (g_t(y) - g_t(x))^{-2b} Z_t^a F(Z_t). \quad (3.7)$$

In particular, it is generated by continuous curve up to T_x . However, the continuity of the process around T_x or T_y can be problematic in general. To understand the behavior of the process near T_x or T_y , we first need to understand the asymptotic of the hypergeometric function F in the definition of the driving function.

Lemma 3.1. *Fix $\kappa \geq 8$ and $\nu > -2$. Denote by Γ the Gamma function. The function F defined in (3.1) is increasing on $[0, 1)$ with $F(0) = 1$. Moreover, we have the following asymptotic.*

- When $\kappa > 8$ and $\nu > -2$, we have

$$\lim_{z \rightarrow 1^-} (1-z)^{1-8/\kappa} F(z) = {}_2F_1 \left(\frac{4}{\kappa}, \frac{2\nu+12}{\kappa} - 1, \frac{2\nu+8}{\kappa}; 1 \right) = \frac{\Gamma(\frac{2\nu+8}{\kappa})\Gamma(1-\frac{8}{\kappa})}{\Gamma(\frac{2\nu+4}{\kappa})\Gamma(1-\frac{4}{\kappa})} \in (0, \infty). \quad (3.8)$$

- When $\kappa = 8$ and $\nu > -2$, we have

$$\lim_{z \rightarrow 1^-} \frac{F(z)}{\log \frac{1}{1-z}} = \frac{1}{\sqrt{\pi}} \frac{(\nu+2)\Gamma(2+\frac{\nu}{4})}{(\nu+4)\Gamma(\frac{3}{2}+\frac{\nu}{4})} \in (0, \infty). \quad (3.9)$$

Proof. In this lemma, we set

$$A = \frac{2\nu+4}{\kappa}, \quad B = 1 - \frac{4}{\kappa}, \quad C = \frac{2\nu+8}{\kappa}.$$

When $\kappa \geq 8$ and $\nu > -2$, we have $A > 0, B > 0, C > 0$, thus $F(z)$ is increasing on $[0, 1)$. It remains to derive the asymptotic.

When $\kappa > 8$ and $\nu > -2$, by (A.3), we have

$$F(z) = (1-z)^{\frac{8}{\kappa}-1} {}_2F_1 \left(\frac{4}{\kappa}, \frac{2\nu+12}{\kappa} - 1, \frac{2\nu+8}{\kappa}; z \right).$$

By (A.5), we have

$${}_2F_1 \left(\frac{4}{\kappa}, \frac{2\nu+12}{\kappa} - 1, \frac{2\nu+8}{\kappa}; 1 \right) = \frac{\Gamma(\frac{2\nu+8}{\kappa})\Gamma(1-\frac{8}{\kappa})}{\Gamma(\frac{2\nu+4}{\kappa})\Gamma(1-\frac{4}{\kappa})} \in (0, \infty).$$

This gives (3.8).

When $\kappa = 8$ and $\nu > -2$, we have

$$\begin{aligned} \lim_{z \rightarrow 1^-} \frac{F(z)}{\log \frac{1}{1-z}} &= \lim_{z \rightarrow 1^-} (1-z)F'(z) && \text{(by L'Hospital rule)} \\ &= \lim_{z \rightarrow 1^-} \frac{AB}{C} (1-z) {}_2F_1(A+1, B+1, C+1; z) && \text{(by (A.4))} \\ &= \lim_{z \rightarrow 1^-} \frac{AB}{C} {}_2F_1(C-A, C-B, C+1; z) && \text{(by (A.3))} \\ &= \frac{AB}{C} \frac{\Gamma(C+1)\Gamma(A+B+1-C)}{\Gamma(1+A)\Gamma(1+B)}, && \text{(by (A.5))} \end{aligned}$$

as desired in (3.9). □

Lemma 3.2. Fix $\kappa \geq 8, \nu > -2$ and $0 < x < y$. Suppose $\eta \sim \text{hSLE}_\kappa(\nu)$ in \mathbb{H} from 0 to ∞ with marked points (x, y) . Then η is generated by a continuous curve up to T_y .

Proof. We compare the law of η with $\text{SLE}_\kappa(\nu + 2, \kappa - 6 - \nu)$ in \mathbb{H} from 0 to ∞ with force points (x, y) . By (2.1) and (3.7), the Radon-Nikodym derivative is given by R_t/R_0 where

$$R_t = F(Z_t)(V_t^y - W_t)^{4/\kappa-1}.$$

When $\kappa > 8$, we write

$$R_t = F(Z_t)(1 - Z_t)^{1-8/\kappa}(V_t^y - V_t^x)^{8/\kappa-1}(V_t^y - W_t)^{-4/\kappa}.$$

By (3.8), the function $F(z)(1 - z)^{1-8/\kappa}$ is uniformly bounded for $z \in [0, 1]$. Define, for $n \geq 1$,

$$S_n = \inf\{t : V_t^y - V_t^x \leq 1/n\}.$$

Then R_t is bounded up to S_n . Since $\text{SLE}_\kappa(\nu + 2, \kappa - 6 - \nu)$ is generated by a continuous curve, the process η is generated by a continuous curve up to S_n . This holds for any n , thus η is generated by a continuous curve up to $T_y = \lim_n S_n$.

When $\kappa = 8$, we write

$$R_t = \frac{F(Z_t)}{\log \frac{1}{1-Z_t}} \left(\log \frac{1}{1-Z_t} \right) (V_t^y - W_t)^{4/\kappa-1}.$$

By (3.9), we know that $F(z)/\left(\log \frac{1}{1-z}\right)$ is uniformly bounded for $z \in [0, 1]$. Define S_n in the same way as before. Then R_t is bounded up to S_n . Similarly, the process η is continuous up to $T_y = \lim_n S_n$. \square

In Lemma 3.2, we obtain the continuity of $\text{hSLE}_\kappa(\nu)$ up to T_y by showing that the process is absolutely continuous with respect to $\text{SLE}_\kappa(\nu + 2, \kappa - 6 - \nu)$. However, the absolute continuity is no longer true when the process approaches T_y . In the following, we will derive the continuity of the process as $t \rightarrow T_y$. From Lemma 3.1, we see that the asymptotic of $F(z)$ as $z \rightarrow 1$ is very different between $\kappa > 8$ and $\kappa = 8$. We will treat the two cases separately: we prove the continuity of hSLE with $\kappa > 8$ in Lemma 3.3, and the continuity with $\kappa = 8$ in Corollary 4.8.

Lemma 3.3. Fix $\kappa > 8, \nu \geq \kappa/2 - 6$ and $0 < x < y$. Suppose $\eta \sim \text{hSLE}_\kappa(\nu)$ in \mathbb{H} from 0 to ∞ with marked points (x, y) . Then η is generated by a continuous curve up to and including T_y .

Proof. Since $\text{hSLE}_\kappa(\nu)$ is scaling invariant, we may assume $y = 1$ and $x \in (0, 1)$. We denote T_y by T . In this lemma, we discuss the continuity of the process $(K_t, 0 \leq t \leq T)$ as $t \rightarrow T$. We need to zoom in around the point 1. To this end, we perform a standard change of coordinate and parameterize the process according to the capacity seen from the point 1. See [SW05, Theorem 3] or [Wu20, Proof of Proposition 3.2].

Set $\varphi(z) = z/(1-z)$, this is the Möbius transformation of \mathbb{H} that sends the triple $(0, 1, \infty)$ to $(0, \infty, -1)$. Denote by $\tilde{x} = \varphi(x) = x/(1-x) > 0$. Denote the image of $(K_t, 0 \leq t \leq T)$ under φ by $(\tilde{K}_s, 0 \leq s \leq \tilde{S})$ where we parameterize this process by its capacity seen from ∞ . Let $(\tilde{g}_s, s \geq 0)$ be the corresponding family of conformal maps and $(\tilde{W}_s, s \geq 0)$ be the driving function. Then we know that the law of \tilde{K}_s is the same as SLE_κ in \mathbb{H} from 0 to ∞ weighted by the following local martingale:

$$\tilde{M}_s = \tilde{g}'_s(-1)^h \tilde{g}'_s(\tilde{x})^b (\tilde{W}_s - \tilde{g}_s(-1))^{-2h} \tilde{Z}_s^a F(\tilde{Z}_s), \quad \text{where } \tilde{Z}_s = \frac{\tilde{g}_s(\tilde{x}) - \tilde{W}_s}{\tilde{g}_s(\tilde{x}) - \tilde{g}_s(-1)}.$$

Compare the law of \tilde{K} with respect to $\text{SLE}_\kappa(2; \nu + 2)$ in \mathbb{H} from 0 to ∞ with force points $(-1; \tilde{x})$. The Radon-Nikodym derivative is given by \tilde{R}_s/\tilde{R}_0 where

$$\tilde{R}_s = (1 - \tilde{Z}_s)^{1-8/\kappa} F(\tilde{Z}_s) (\tilde{g}_s(\tilde{x}) - \tilde{g}_s(-1))^{1-8/\kappa-2a}.$$

When $\kappa > 8$ and $\nu \geq \kappa/2 - 6 > -2$, the function $(1-z)^{1-8/\kappa}F(z)$ is uniformly bounded on $z \in [0, 1]$ due to (3.8). The process $\tilde{g}_s(\tilde{x}) - \tilde{g}_s(-1)$ is increasing in s , thus $\tilde{g}_s(\tilde{x}) - \tilde{g}_s(-1) \geq 1/(1-x)$. Since $\nu \geq \kappa/2 - 6$, the exponent of the term $\tilde{g}_s(\tilde{x}) - \tilde{g}_s(-1)$ is $1 - 8/\kappa - 2a \leq 0$. Therefore, \tilde{R}_s is bounded. This implies that the law of \tilde{K}_s is absolutely continuous with respect to the law of $\text{SLE}_\kappa(2; \nu + 2)$ up to and including the swallowing time of -1 . Hence $(\tilde{K}_s, 0 \leq s \leq \tilde{S})$ is generated by a continuous curve up to and including \tilde{S} . In particular, this implies that the original process $(K_t, 0 \leq t \leq T)$ is generated by a continuous curve up to and including T . \square

To sum up the results in this section, we have the following continuity of $\text{hSLE}_\kappa(\nu)$ when $\kappa > 8$.

Proposition 3.4. *Fix $\kappa > 8, \nu \geq \kappa/2 - 6$ and $x_1 < x_2 < x_3 < x_4$. The process $\text{hSLE}_\kappa(\nu)$ in \mathbb{H} from x_1 to x_4 is almost surely generated by a continuous curve.*

Proof. Lemma 3.3 gives the continuity of $\text{hSLE}_\kappa(\nu)$ up to and including T_{x_3} . After T_{x_3} , we continue the process by standard SLE_κ in the remaining domain from $\eta(T_{x_3})$ to x_4 . Thus, the process $\text{hSLE}_\kappa(\nu)$ is continuous for all time. \square

3.2 Discussion on time-reversal of hSLE

The time-reversal of SLE_κ with $\kappa > 8$ was fully addressed in [MS17, Theorem 1.19]: consider $\text{SLE}_\kappa(\rho_1; \rho_2)$ with force points next to the starting point for $\rho_1, \rho_2 \in (-2, \kappa/2 - 2)$, its time-reversal is an $\text{SLE}_\kappa(\kappa/2 - 4 - \rho_2; \kappa/2 - 4 - \rho_1)$ process with force points next to the starting point. In particular, the time-reversal of SLE_κ is an $\text{SLE}_\kappa(\kappa/2 - 4; \kappa/2 - 4)$ process. This indicates that the time-reversal of $\text{hSLE}_\kappa(\nu)$ with $\kappa > 8$ is a variant of SLE_κ where one has four extra marked points. In particular, the time-reversal is no longer in the family of hSLE which is a variant of SLE with two extra marked points, see Lemma 3.5. Therefore, it is only reasonable to talk about reversibility of $\text{hSLE}_\kappa(\nu)$ with $\kappa \leq 8$. The reversibility of $\text{hSLE}_\kappa(\nu)$ with $\kappa < 8$ was addressed in [Wu20, Section 3.3]. We will discuss the reversibility of $\text{hSLE}_\kappa(\nu)$ with $\kappa = 8$ in Section 3.3.

Lemma 3.5. *When $\kappa > 8$ and $\nu \geq \kappa/2 - 6$. The time-reversal of $\text{hSLE}_\kappa(\nu)$ is not an $\text{hSLE}_\kappa(\tilde{\nu})$ for any value of $\tilde{\nu}$.*

Proof. Fix $x_1 < x_2 < x_3 < x_4$. Suppose η is an $\text{hSLE}_\kappa(\nu)$ in \mathbb{H} from x_1 to x_4 and let $\hat{\eta}$ be its time-reversal. Let γ be an $\text{hSLE}_\kappa(\tilde{\nu})$ in \mathbb{H} from x_4 to x_1 with marked points (x_3, x_2) . We will compare the laws of $\hat{\eta}$ and γ in small neighborhood of x_4 .

- In the construction of η , we know that the process almost surely hits the interval (x_3, x_4) and after the hitting time, we continue the process as a standard SLE_κ in the remaining domain. Therefore, the initial segment of $\hat{\eta}$ is the time-reversal of a standard SLE_κ . By [MS17, Theorem 1.19], we know that, in small neighborhood of x_4 , the law of $\hat{\eta}$ and the law of $\text{SLE}_\kappa(\kappa/2 - 4; \kappa/2 - 4)$ are absolutely continuous with respect to each other.
- From the definition of $\text{hSLE}_\kappa(\tilde{\nu})$, we know that, in small neighborhood of x_4 , the law of γ and the law of SLE_κ are absolutely continuous with respect to each other.

Combining the above two observations, we see that $\hat{\eta}$ can not have the same law as γ , because $\text{SLE}_\kappa(\kappa/2 - 4; \kappa/2 - 4)$ and SLE_κ are not absolutely continuous with respect to each other. \square

3.3 Continuity and reversibility of hSLE with $\kappa = 8$

The continuity and reversibility of hSLE_8 will be given in Corollary 4.8 in Section 4.4. The proof there is based on the convergence of the Peano curve for UST. Assuming this is true, we are able to complete the proof of Theorem 1.1.

Proof of Theorem 1.1. We may assume $x_1 = 0 < x_2 = x < x_3 = y < x_4 = \infty$. Suppose $\eta \sim \text{hSLE}_8(\nu)$ in \mathbb{H} from 0 to ∞ with marked points (x, y) . Recall that η has the same law as SLE_8 in \mathbb{H} from 0 to ∞ weighted by the following local martingale:

$$M_t = g'_t(x)^b g'_t(y)^b (g_t(y) - g_t(x))^{-2b} Z_t^a F(Z_t), \quad (3.10)$$

where

$$h = \frac{-1}{8}, \quad a = \frac{\nu + 2}{8}, \quad b = \frac{(\nu + 2)(\nu - 2)}{32}, \quad F(z) = {}_2F_1\left(2a, \frac{1}{2}, 2a + \frac{1}{2}; z\right).$$

Suppose $\gamma \sim \text{hSLE}_8$ in \mathbb{H} from 0 to ∞ with marked points (x, y) . Then γ has the same law as SLE_8 in \mathbb{H} from 0 to ∞ weighted by the following local martingale:

$$N_t = g'_t(x)^h g'_t(y)^h (g_t(y) - g_t(x))^{-2h} Z_t^{1/4} G(Z_t), \quad \text{where } G(z) = {}_2F_1\left(\frac{1}{2}, \frac{1}{2}, 1; z\right). \quad (3.11)$$

Combining (3.10) and (3.11), we see that the law of η is the same as the law of γ weighted by the following local martingale:

$$R_t = \frac{M_t}{N_t} = \left(\frac{g'_t(x)g'_t(y)}{(g_t(y) - g_t(x))^2} \right)^{\nu^2/32} \times Z_t^{\nu/8} \times \frac{F(Z_t)}{G(Z_t)}.$$

The term Z_t takes values in $[0, 1]$, the term $F(Z_t)/G(Z_t)$ is uniformly bounded due to (3.9). The term

$$\frac{g'_t(x)g'_t(y)}{(g_t(y) - g_t(x))^2}$$

is the boundary Poisson kernel of the domain $\mathbb{H} \setminus \gamma[0, t]$ and it is positive and bounded from above by $(y - x)^{-2}$. Thus R_t is uniformly bounded and the law of η is absolutely continuous with respect to the law of γ up to and including T_y .

By Corollary 4.8, the process γ is continuous up to and including T_y and $\gamma \cap [x, y] = \emptyset$. By the absolute continuity, the process η is continuous up to and including T_y and $\eta \cap [x, y] = \emptyset$. After T_y , we continue the process by standard SLE_8 in the remaining domain from $\eta(T_y)$ to ∞ . Thus η is a continuous curve for all time.

It remains to show the reversibility. We denote by D the connected component of $\mathbb{H} \setminus \gamma$ with $[x, y]$ on the boundary. Since γ is continuous and $\gamma \cap [x, y] = \emptyset$, we have $Z_t \rightarrow 1$ as $t \rightarrow T_y$. Thus, from (3.9),

$$R_t \rightarrow H_D(x, y)^{\nu^2/32} \times \sqrt{\pi} \frac{(\nu + 2)\Gamma(2 + \frac{\nu}{4})}{(\nu + 4)\Gamma(\frac{3}{2} + \frac{\nu}{4})}, \quad \text{as } t \rightarrow T_y.$$

From this, we find that the law of η is the same as the law of γ weighted by the boundary Poisson kernel $H_D(x, y)^{\nu^2/32}$. Since the law of γ is reversible due to Corollary 4.8, and the boundary Poisson kernel is conformally invariant, we obtain the reversibility of the law of η . This completes the proof. \square

Proof of Lemma 1.2. We may assume $x_1 = 0 < x_2 = x < x_3 = y < x_4 = \infty$. Suppose $\gamma \sim \text{hSLE}_8$ in \mathbb{H} from 0 to ∞ with marked points (x, y) . The law of γ is the same as SLE_8 in \mathbb{H} from 0 to ∞ weighted by N_t/N_0 where N_t is the local martingale in (3.11). Note that

$$\frac{N_t}{N_0} = g'_t(x)^h g'_t(y)^h \left(\frac{g_t(y) - g_t(x)}{y - x} \right)^{-2h} \left(\frac{Z_t}{Z_0} \right)^{1/4} \frac{G(Z_t)}{G(Z_0)}, \quad \text{where } Z_0 = \frac{x}{y}, \quad Z_t = \frac{g_t(x) - W_t}{g_t(y) - W_t}.$$

Fix t , we first let $y \rightarrow \infty$, then

$$\frac{N_t}{N_0} \rightarrow g'_t(x)^h \left(\frac{g_t(x) - W_t}{x} \right)^{1/4}.$$

This implies that the law of γ converges weakly to $\text{SLE}_8(2)$ from 0 to ∞ with marked point x . We then let $x \rightarrow \infty$, then $N_t/N_0 \rightarrow 1$. This implies that the law of $\text{SLE}_8(2)$ converges weakly to SLE_8 , and completes the proof. \square

4 Convergence of UST in quads

For a finite subgraph $G = (V(G), E(G)) \subset \mathbb{Z}^2$, we denote by ∂G the inner boundary of G : $\partial G = \{x \in V(G) : \exists y \notin V(G) \text{ such that } \{x, y\} \in E(\mathbb{Z}^2)\}$. In this article, when we add the subscript or superscript δ , we mean scaling subgraphs of the lattices $\mathbb{Z}^2, (\mathbb{Z}^2)^*, (\mathbb{Z}^2)^\diamond$ by δ .

Uniform spanning tree

Suppose that $G = (V, E)$ is a finite connected graph, a tree T is a subgraph of G that has no loops. A spanning tree is a tree that covers all the vertices. A uniform spanning tree on G is a probability measure on the set of all spanning trees of G in which every tree is chosen with equal probability. Given a disjoint sequence $(\alpha_k : 1 \leq k \leq N)$ of trees of G , a spanning tree with $(\alpha_k : 1 \leq k \leq N)$ wired is a spanning tree T such that $\alpha_k \subset T$ for $1 \leq k \leq N$. A uniform spanning tree with $(\alpha_k : 1 \leq k \leq N)$ wired is a probability measure on the set of all spanning trees of G with $(\alpha_k : 1 \leq k \leq N)$ wired in which every tree is chosen with equal probability.

Space of curves

A path is defined by a continuous map from $[0, 1]$ to \mathbb{C} . Let \mathcal{C} be the space of unparameterized paths in \mathbb{C} . Define the metric on \mathcal{C} as follows:

$$d(\gamma_1, \gamma_2) := \inf \sup_{t \in [0, 1]} |\hat{\gamma}_1(t) - \hat{\gamma}_2(t)|, \quad (4.1)$$

where the infimum is taken over all the choices of parameterizations $\hat{\gamma}_1$ and $\hat{\gamma}_2$ of γ_1 and γ_2 . The metric space (\mathcal{C}, d) is complete and separable.

Let \mathcal{P} be a family of probability measures on \mathcal{C} . We say \mathcal{P} is tight if for any $\epsilon > 0$, there exists a compact set K_ϵ such that $\mathbb{P}[K_\epsilon] \geq 1 - \epsilon$ for any $\mathbb{P} \in \mathcal{P}$. We say \mathcal{P} is relatively compact if every sequence of elements in \mathcal{P} has a weakly convergent subsequence. As the metric space is complete and separable, relative compactness is equivalent to tightness.

Convergence of discrete polygons

Recall that a polygon $(\Omega; x_1, \dots, x_p)$ is a bounded simply connected domain $\Omega \subset \mathbb{C}$ with distinct boundary points x_1, \dots, x_p in counterclockwise order, such that $\partial\Omega$ is locally connected. Let ϕ be any conformal map from \mathbb{U} onto Ω . Note that the local connectedness of $\partial\Omega$ is equivalent to that ϕ can be extended continuously to $\bar{\mathbb{U}}$ and is equivalent to that $\partial\Omega$ is a curve. See [Pom92, Theorem 2.1].

A sequence of discrete polygons $(\Omega_\delta; x_1^\delta, \dots, x_p^\delta)$ on $\delta\mathbb{Z}^2$ is said to converge to a polygon $(\Omega; x_1, \dots, x_p)$ in the Carathéodory sense if there exist conformal maps ϕ_δ from \mathbb{U} onto Ω_δ and conformal map ϕ from \mathbb{U} onto Ω such that $\phi_\delta \rightarrow \phi$ as $\delta \rightarrow 0$ uniformly on compact subsets of \mathbb{U} and $\phi_\delta^{-1}(x_j^\delta) \rightarrow \phi^{-1}(x_j)$ for $1 \leq j \leq p$.

We will encounter another type of convergence of polygons: consider a sequence of discrete polygons $(\Omega_\delta; x_1^\delta, \dots, x_p^\delta)$ on $\delta\mathbb{Z}^2$ converge to a polygon $(\Omega; x_1, \dots, x_p)$ in the following sense:

$$(x_i^\delta x_{i+1}^\delta) \rightarrow (x_i x_{i+1}) \quad \text{as curves in metric (4.1),} \quad \text{as } \delta \rightarrow 0, \quad \text{for } i = 1, \dots, p, \quad (4.2)$$

where we use the convention that $x_{p+1} = x_1$. Note that such convergence implies the convergence in the Carathéodory sense. These two types of convergence apply to polygons in the medial lattice in a similar way.

4.1 UST with Dobrushin boundary conditions

Consider the medial lattice $(\mathbb{Z}^2)^\diamond$. We orient the edges of the medial lattice such that edges of a face containing a vertex in \mathbb{Z}^2 are oriented counterclockwise and edges of a face containing a vertex in $(\mathbb{Z}^2)^*$ are oriented clockwise. Let a^\diamond, b^\diamond be two distinct medial vertices, and $(a^\diamond b^\diamond) = \{a^\diamond = v_0 \sim v_1 \sim \dots \sim v_n = b^\diamond\}$ and $(b^\diamond a^\diamond) = \{b^\diamond = q_0 \sim q_1 \sim \dots \sim q_s = a^\diamond\}$ be two paths of neighboring medial vertices satisfying the following conditions: (1) the edges along $(a^\diamond b^\diamond)$ point in clockwise way with the orientation inherited from the medial lattice; (2) the edges along $(b^\diamond a^\diamond)$ point in counterclockwise way with the orientation inherited from the medial lattice; (3) the two paths are edge-avoiding and $(a^\diamond b^\diamond) \cap (b^\diamond a^\diamond) = \{a^\diamond, b^\diamond\}$. See Figure 4.1.

Given $(a^\diamond b^\diamond)$ and $(b^\diamond a^\diamond)$, the medial Dobrushin domain $(\Omega^\diamond; a^\diamond, b^\diamond)$ is defined as the subgraph of $(\mathbb{Z}^2)^\diamond$ induced by the vertices enclosed by or on the path $(a^\diamond b^\diamond) \cup (b^\diamond a^\diamond)$. Let $\Omega \subset \mathbb{Z}^2$ be the graph with edge set consisting of edges passing through end-points of medial edges in $E(\Omega^\diamond) \setminus (b^\diamond a^\diamond)$ and with vertex set given by the endpoints of these edges. The vertices of Ω nearest to a^\diamond, b^\diamond are denoted by a, b and we call $(\Omega; a, b)$ primal Dobrushin domain. Let (ab) be the set of edges corresponding to medial vertices in $(a^\diamond b^\diamond) \cap \partial\Omega^\diamond$. Let $\Omega^* \subset \mathbb{Z}^2$ be the graph with edge set consisting of edges passing through end-points of medial edges in $E(\Omega^\diamond) \setminus (a^\diamond b^\diamond)$ and with vertex set given by the endpoints of these edges. The vertices of Ω^* nearest to a^\diamond, b^\diamond are denoted by a^*, b^* . Let $(b^* a^*)$ be the set of edges corresponding to medial vertices in $(b^\diamond a^\diamond) \cap \partial\Omega^\diamond$. Note that a is the vertex of Ω that is nearest to a^\diamond and a^* is the vertex of Ω^* that is nearest to a^\diamond . See Figure 4.1.




Figure 4.1: In the left panel, the solid edges in black are wired boundary arc (ab) , the solid edges in red are dual-wired boundary arc (b^*a^*) . The edges in blue are the boundary arcs $(a^\diamond b^\diamond)$ and $(b^\diamond a^\diamond)$ on the medial lattice. The thin edges are in the tree \mathcal{T} . In the right panel, the curve in orange is the Peano curve associated to \mathcal{T} .

Suppose that \mathcal{T} is a spanning tree on some primal Dobrushin domain $(\Omega; a, b)$ with (ab) wired, then there exists a unique path, called Peano curve, on $(\mathbb{Z}^2)^\diamond$ running along the tree from a^\diamond to b^\diamond . The following theorem concerns the convergence of the Peano curve of UST.

Theorem 4.1. *Fix a Dobrushin domain $(\Omega; a, b)$ such that $\partial\Omega$ is C^1 and simple. Suppose that a sequence of medial Dobrushin domains $(\Omega_\delta^\diamond; a_\delta^\diamond, b_\delta^\diamond)$ converges to $(\Omega; a, b)$ as in (4.2). Consider UST on the primal domain Ω_δ with $(a_\delta b_\delta)$ wired. Denote by η_δ the induced Peano curve. Then the law of η_δ converges weakly to SLE_8 in Ω from a to b .*

Proof. This statement is proved in [LSW04, Theorems 4.7 and 4.8]. Let us summarize the proof and we will adjust it for our setting later. The proof is divided into two steps. The first step is to show the convergence of driving function by martingale observable [LSW04, Theorem 4.4]. This step only requires $\Omega_\delta^\diamond \rightarrow \Omega$ in the Carathéodory sense and does not need any regularity assumption on $\partial\Omega$. The second

step is to show the tightness of the discrete curves. We restate the proof for the tightness: (4.3), (4.4) and (4.9) and point out why we need the C^1 -regularity on $\partial\Omega$.

We choose conformal maps $\phi_\delta : \Omega_\delta^\circ \rightarrow \mathbb{H}$ with $\phi_\delta(a_\delta^\circ) = 0$ and $\phi_\delta(b_\delta^\circ) = \infty$ such that ϕ_δ^{-1} converges locally uniformly. We define $\hat{\eta}_\delta := \phi_\delta(\eta_\delta)$ and we parameterize $\hat{\eta}_\delta$ by the half-plane capacity and let $(g_t^\delta, t \geq 0)$ be the corresponding conformal maps. For $0 < t_1 < t_2 < +\infty$, let $Y_\delta(t_1, t_2) := \text{diam}(g_{t_1}^\delta(\hat{\eta}_\delta[t_1, t_2]))$. By [LSW04, Lemma 4.6], for every $\epsilon > 0$, there exist $\epsilon_0 > 0$ and $\delta_0 > 0$ such that, for all $\delta < \delta_0$,

$$\mathbb{P}[\sup\{|\hat{\eta}_\delta(t_2) - \hat{\eta}_\delta(t_1)| : 0 \leq t_1 < t_2 \leq \tau_\delta, Y_\delta(t_1, t_2) \leq \epsilon_0\} \geq \epsilon] < \epsilon \quad (4.3)$$

where $\tau_\delta := \inf\{t \geq 0 : |\eta_\delta(t)| = \epsilon^{-1}\}$. Combining with [LSW04, Lemma 2.1], we obtain the following tightness [LSW04, Proposition 4.5]: For every $t > 0, \epsilon > 0$, there exists $\epsilon_0 > 0$ such that

$$\mathbb{P}[\sup\{|\hat{\eta}_\delta(t_2) - \hat{\eta}_\delta(t_1)| : 0 \leq t_1 < t_2 \leq t, |t_2 - t_1| \leq \epsilon_0\} \geq \epsilon] < \epsilon. \quad (4.4)$$

The key in the proof of the tightness is (4.3) and we will summarize its proof below. For $\delta, \epsilon, \epsilon_0 > 0$, define

$$\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} = \{\exists 0 \leq t_1 < t_2 \leq \tau_\delta \text{ such that } |\hat{\eta}_\delta(t_2) - \hat{\eta}_\delta(t_1)| \geq \epsilon, \text{ but } Y_\delta(t_1, t_2) \leq \epsilon_0\}.$$

It suffices to show that, for a well-chosen positive function $\epsilon_0(\delta)$ such that $\epsilon_0(\delta) \rightarrow 0$ as $\delta \rightarrow 0$, we have

$$\lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \sup_{\epsilon_0 \leq \epsilon_0(\delta)} \mathbb{P}[\mathcal{A}_{\epsilon_0}^{\delta, \epsilon}] = 0. \quad (4.5)$$

We denote by Z the semicircle $2\epsilon^{-1}\partial\mathbb{U} \cap \mathbb{H}$. On the one hand, there is a constant $C > 0$ such that $\text{dist}(g_t^\delta(Z), g_t^\delta(\hat{\eta}_\delta[t, \tau_\delta])) \geq C$ for all $t \leq \tau_\delta$. On the other hand, on the event $\mathcal{A}_{\epsilon_0}^{\delta, \epsilon}$, we have $Y_\delta(t_1, t_2) \leq \epsilon_0 \rightarrow 0$ as $\epsilon_0 \rightarrow 0$. These two facts guarantee that the extremal length of simple arcs in $\mathbb{H} \setminus g_{t_1}^\delta(\hat{\eta}_\delta[t_1, t_2])$ which separate $g_{t_1}^\delta(Z)$ from $g_{t_1}^\delta(\hat{\eta}_\delta[t_1, t_2])$ tends to 0 as $\epsilon_0 \rightarrow 0$. By the conformal invariance of extremal length, there exists a simple curve $\gamma_{\delta, \epsilon_0}$ in $\mathbb{H} \setminus \hat{\eta}_\delta[0, t_1]$ separating $\hat{\eta}_\delta[t_1, t_2]$ and Z such that the length of $\gamma_{\delta, \epsilon_0}$ tends to 0 as $\epsilon_0 \rightarrow 0$.

For $s > 0$, we denote by $\chi_{0, \epsilon_0}^\delta(s)$ the event $\{\text{dist}(0, \gamma_{\delta, \epsilon_0}) < s\}$ and we denote by $\chi_{1, \epsilon_0}^\delta(s)$ the event $\{\text{dist}(\mathbb{R}, \gamma_{\delta, \epsilon_0}) < s\}$. We choose two function $s_1(\epsilon)$ and $s_0(\epsilon)$ such that $s_1(\epsilon) < s_0(\epsilon)$ and $\lim_{\epsilon \rightarrow 0} \frac{s_0(\epsilon)}{\epsilon} = 0$. Now, we divide $\mathcal{A}_{\epsilon_0}^{\delta, \epsilon}$ into three events

$$\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} = \{\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \setminus \chi_{1, \epsilon_0}^\delta(s_1(\epsilon))\} \cup \{\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{1, \epsilon_0}^\delta(s_1(\epsilon)) \setminus \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))\} \cup \{\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))\}.$$

On the event $\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \setminus \chi_{1, \epsilon_0}^\delta(s_1(\epsilon))$, let B_0 be the disk such that ∂B_0 is the smallest circle surrounding $\gamma_{\delta, \epsilon_0}$. Moreover, suppose B_1 is the disk with the same center as B_0 and its radius is $\frac{s_1(\epsilon)}{4}$. We define $A := B_1 \setminus B_0$. Then, there will be a $t^* \in (t_1, t_2)$ such that $\hat{\eta}_\delta(t^*) \notin B_1$. This implies that there exists three segments $\hat{\eta}_{\delta, 1}, \hat{\eta}_{\delta, 3} \subset \hat{\eta}_\delta[0, t^*]$ and $\hat{\eta}_{\delta, 2} \subset \hat{\eta}_\delta[t^*, \infty)$ such that each of them hits B_0 and $\hat{\eta}_{\delta, 2}$ separates $\hat{\eta}_{\delta, 1}$ and $\hat{\eta}_{\delta, 3}$ in B_1 . By our assumption, ϕ_δ^{-1} converges to ϕ^{-1} uniformly on B_1 . By definition, one side of η_δ is adjacent to the UST \mathcal{T}_δ and the other side of η_δ is adjacent to the dual tree \mathcal{T}_δ^* . Then, there are two simple paths $\xi_{1, \delta} \subset \mathcal{T}_\delta$ and $\xi_{2, \delta} \subset \mathcal{T}_\delta^*$ such that their endpoints are in $\phi_\delta^{-1}(\partial B_1)$ and each is between $\phi_\delta^{-1}(\hat{\eta}_{\delta, 1})$ and $\phi_\delta^{-1}(\hat{\eta}_{\delta, 2})$ or $\phi_\delta^{-1}(\hat{\eta}_{\delta, 2})$ and $\phi_\delta^{-1}(\hat{\eta}_{\delta, 3})$. This implies that $\text{dist}(\xi_{1, \delta}, \xi_{2, \delta}) \rightarrow 0$ as $\epsilon_0 \rightarrow 0$, but $\text{diam}(\xi_{1, \delta})$ and $\text{diam}(\xi_{2, \delta})$ is bounded from below by a constant depending on ϵ and Ω .

Now, we introduce the conception of trunk and dual trunk. Suppose $\{\Omega_\delta^\circ\}_{\delta > 0}$ and Ω satisfy our assumption. For the UST \mathcal{T}_δ in the primal graph, we define its $\frac{1}{n}$ -trunk as follows. For any $x, y \in \mathcal{T}_\delta$, there is a unique path on \mathcal{T}_δ from x to y , which we denote by $\eta_\delta^{x, y}$. We denote by x' the first point at which $\eta_\delta^{x, y}$ hits $\partial B(x, \frac{1}{n})$ and y' the last point at which $\eta_\delta^{x, y}$ hits $\partial B(y, \frac{1}{n})$. We denote by $\mathcal{I}(x, y)$ the unique path on \mathcal{T}_δ connecting x' and y' . If we can not find x' or y' in this way, we define $\mathcal{I}(x, y) = \emptyset$. Then, the $\frac{1}{n}$ -trunk is defined as $\mathbf{trunk}_\delta(\frac{1}{n}) := \cup_{x, y \in \mathcal{T}_\delta} \mathcal{I}(x, y)$. Note that we can choose $\{\delta_m\}_{m > 0}$ such that $\mathbf{trunk}_{\delta_m}(\frac{1}{n})$ converges in Hausdorff distance for every n as $m \rightarrow \infty$. We define $\mathbf{trunk}_0(\frac{1}{n}) := \lim_{m \rightarrow \infty} \mathbf{trunk}_{\delta_m}(\frac{1}{n})$. The trunk is defined by $\mathbf{trunk} = \cup_{n > 0} \mathbf{trunk}_0(\frac{1}{n})$. The dual trunk \mathbf{trunk}^* is defined similarly. We define

the event $E := \{\mathbf{trunk} \cap \mathbf{trunk}^* \neq \emptyset\}$. In [Sch00, Theorem 11.1], the author proves that $\mathbb{P}[E] = 0$. This is the place that we need the C^1 -regularity on $\partial\Omega$. Thus, we have $\cap_{\delta>0} \cup_{\epsilon_0 \leq \epsilon_0(\delta)} \{\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \setminus \chi_{1, \epsilon_0}^\delta(s_1(\epsilon))\} \subset E$ for every $\epsilon > 0$, which implies that

$$\lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \sup_{\epsilon_0 \leq \epsilon_0(\delta)} \mathbb{P}[\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \setminus \chi_{1, \epsilon_0}^\delta(s_1(\epsilon))] = 0. \quad (4.6)$$

On the event $\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{1, \epsilon_0}^\delta(s_1(\epsilon)) \setminus \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))$, there are two possibilities. If the two endpoints of $\gamma_{\delta, \epsilon_0}$ lie on $\hat{\eta}_\delta[0, t_1]$, the argument is the same as above. If there is one endpoint which lies on \mathbb{R} , we assume it is on \mathbb{R}^+ . Then, the $\text{dist}(0, \gamma_{\delta, \epsilon_0}) \geq s_0(\epsilon)$. By a similar argument as above, we know that there exists $\xi_{2, \delta} \in \mathcal{T}_\delta^*$ such that $\text{dist}(\xi_{2, \delta}, (a_\delta b_\delta)) \rightarrow 0$ as $\epsilon_0 \rightarrow 0$ and $\text{diam}(\xi_{2, \delta})$ is bounded from below by a constant depending on ϵ and Ω . Note that the boundary arc $(ab) \subset \mathbf{trunk}$. This implies that

$$\cap_{\delta>0} \cup_{\epsilon_0 \leq \epsilon_0(\delta)} \{\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{1, \epsilon_0}^\delta(s_1(\epsilon)) \setminus \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))\} \subset E.$$

Thus, we have

$$\lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \sup_{\epsilon_0 \leq \epsilon_0(\delta)} \mathbb{P}[\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{1, \epsilon_0}^\delta(s_1(\epsilon)) \setminus \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))] = 0. \quad (4.7)$$

We still need to prove

$$\lim_{\epsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \sup_{\epsilon_0 \leq \epsilon_0(\delta)} \mathbb{P}[\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))] = 0. \quad (4.8)$$

We fix an $s > 0$ and choose $v \in V(\Omega_\delta)$ which is closest to $\phi_\delta^{-1}(is)$. Let $v^* \in V(\Omega_\delta^*)$ be the vertex which is closest to v . Denote by χ_v the LERW which starts from v and ends at $(a_\delta b_\delta)$. Similarly, we denote by χ_{v^*} the LERW which starts from v^* . We define $A_1 := \{\text{diam}(\phi_\delta(\chi_v)) < \frac{\epsilon}{10}\}$ and define $A_2 := \{\text{dist}(0, \phi_\delta(\chi_v)) > s_0(\epsilon)\}$. We also define A_1^* and A_2^* similarly by replacing χ_v with χ_{v^*} . By the same method as [LSW04, Proposition 4.2], for any ϵ small enough, we can choose s which is much smaller than ϵ and is much larger than $s_0(\epsilon)$, such that

$$\mathbb{P}[A_1] \geq 1 - \frac{1}{10}\epsilon, \quad \mathbb{P}[A_2] \geq 1 - \frac{1}{10}\epsilon, \quad \mathbb{P}[A_1^*] \geq 1 - \frac{1}{10}\epsilon, \quad \mathbb{P}[A_2^*] \geq 1 - \frac{1}{10}\epsilon.$$

We denote by D' the domain bounded by $\chi_v \cup [vv^*] \cup \chi_{v^*} \cup \partial\Omega_\delta^*$, where $[vv^*]$ is the straight line segment connecting v and v^* . Then, we have that η_δ will exit D' through $[vv^*]$. Note that on $A_1 \cap A_2 \cap A_1^* \cap A_2^* \cap \{\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))\}$, we have that $\text{diam}(\phi_\delta(D')) < \epsilon$ and $\gamma_{\delta, \epsilon_0} \subset \phi_\delta(D')$. This implies that $\hat{\eta}_\delta[t_1, t_2] \subset \phi_\delta(D')$. But this contradicts with the assumption that $|\hat{\eta}_\delta(t_2) - \hat{\eta}_\delta(t_1)| \geq \epsilon$. Thus, we have $\{\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))\} \subset (A_1 \cap A_2 \cap A_1^* \cap A_2^*)^c$. Therefore,

$$\mathbb{P}[\mathcal{A}_{\epsilon_0}^{\delta, \epsilon} \cap \chi_{0, \epsilon_0}^\delta(s_0(\epsilon))] < \epsilon.$$

This implies (4.8). Note that (4.6)–(4.8) imply (4.5) which gives (4.3).

Finally, we will show the transience of curves at b : for any $\epsilon > 0$, there exist $\epsilon' \leq \epsilon$ and $\delta_0 > 0$ such that, for all $\delta \leq \delta_0$,

$$\mathbb{P}[\eta_\delta \text{ hits } \partial B(b, \epsilon) \text{ after hitting } \partial B(b, \epsilon')] < \epsilon. \quad (4.9)$$

The proof is similar to the proof of (4.8). Fix two constants $r < \epsilon$ and $C > 0$. We choose $v \in V(\Omega_\delta)$ which is adjacent to $\partial B(b, r)$. Suppose $v^* \in V(\Omega_\delta^*)$ which is adjacent to v . We denote by χ_v the LERW which starts from v and ends at $(a_\delta b_\delta)$. Similarly, we denote by χ_{v^*} the LERW which starts from v^* and ends at $(b_\delta^* a_\delta^*)$. We define $A_1 := \{\text{diam}(\chi_v) < Cr\}$ and $A_2 := \{\text{dist}(b, \chi_v) > \frac{1}{C}r\}$. We also define A_1^* and A_2^* similarly by replacing χ_v with χ_{v^*} . We choose C such that

$$\mathbb{P}[A_1 \cap A_2 \cap A_1^* \cap A_2^*] \geq 1 - \epsilon.$$

We choose r such that $Cr < \epsilon$ and we choose $\epsilon' < \frac{1}{C}r$. Note that on the event $A_1 \cap A_2 \cap A_1^* \cap A_2^*$, we have that η_δ can not hit $\partial B(b, \epsilon)$ after hitting $\partial B(b, \epsilon')$. Since η_δ can only go through $[vv^*]$ once. This implies (4.9). Together with (4.3) and (4.4), we complete the proof of tightness. \square

4.2 UST with alternating boundary conditions in quads

Now we introduce the discrete quad in Theorem 1.5. Let $a^\diamond, b^\diamond, c^\diamond, d^\diamond$ be four distinct medial vertices, and $(a^\diamond b^\diamond) = \{a^\diamond = v_0 \sim v_1 \sim \dots \sim v_n = b^\diamond\}$, $(b^\diamond c^\diamond) = \{b^\diamond = w_0 \sim w_1 \sim \dots \sim w_m = c^\diamond\}$, $(c^\diamond d^\diamond) = \{c^\diamond = p_0 \sim p_1 \sim \dots \sim p_l = d^\diamond\}$ and $(d^\diamond a^\diamond) = \{d^\diamond = q_0 \sim q_1 \sim \dots \sim q_s = a^\diamond\}$ be four paths of neighboring medial vertices satisfying the following conditions: (1) the edges along $(a^\diamond b^\diamond)$ and $(c^\diamond d^\diamond)$ point in clockwise way with the orientation inherited from the medial lattice; (2) the edges along $(b^\diamond c^\diamond)$ and $(d^\diamond a^\diamond)$ point in counterclockwise way with the orientation inherited from the medial lattice; (3) all the paths are edge-avoiding and $(a^\diamond b^\diamond) \cap (b^\diamond c^\diamond) = \{b^\diamond\}$, $(b^\diamond c^\diamond) \cap (c^\diamond d^\diamond) = \{c^\diamond\}$, $(c^\diamond d^\diamond) \cap (d^\diamond a^\diamond) = \{d^\diamond\}$, $(d^\diamond a^\diamond) \cap (a^\diamond b^\diamond) = \{a^\diamond\}$. See Figure 4.2.

Given $(a^\diamond b^\diamond), (b^\diamond c^\diamond), (c^\diamond d^\diamond)$ and $(d^\diamond a^\diamond)$, the medial quad $(\Omega^\diamond; a^\diamond, b^\diamond, c^\diamond, d^\diamond)$ is defined as the subgraph of $(\mathbb{Z}^2)^\diamond$ induced by the vertices enclosed by or on the path $(a^\diamond b^\diamond) \cup (b^\diamond c^\diamond) \cup (c^\diamond d^\diamond) \cup (d^\diamond a^\diamond)$. Recall that the inner boundary $\partial\Omega^\diamond$ is the set of vertices of Ω^\diamond with strictly fewer than four incident edges in $E(\Omega^\diamond)$. Let $\Omega \subset \mathbb{Z}^2$ be the graph with edge set consisting of edges passing through end-points of medial edges in $E(\Omega^\diamond) \setminus ((b^\diamond c^\diamond) \cup (d^\diamond a^\diamond))$ and with vertex set given by the endpoints of these edges. The vertices of Ω nearest to $a^\diamond, b^\diamond, c^\diamond, d^\diamond$ are denoted by a, b, c, d and we call $(\Omega; a, b, c, d)$ the primal quad. Let (ab) and (cd) be the set of edges corresponding to medial vertices in $\partial\Omega^\diamond$, which are also endpoints of medial edges in $(a^\diamond b^\diamond)$ and $(c^\diamond d^\diamond)$ respectively. One can define (bc) and (da) to be the two components of $\partial\Omega \setminus ((ab) \cup (cd))$. See Figure 4.2.

Suppose that \mathcal{T} is a spanning tree on some primal quad $(\Omega; a, b, c, d)$ with (ab) wired and (cd) wired respectively, then there exist two paths on $(\mathbb{Z}^2)^\diamond$ running along \mathcal{T} from a^\diamond to d^\diamond and from b^\diamond to c^\diamond respectively, see Figure 1.1. We still call them Peano curves as before. With the same notations as in Section 1, we denote by η^L the Peano curve from a^\diamond to d^\diamond , by η^R the Peano curve from b^\diamond to c^\diamond and by γ^M the unique path in \mathcal{T} from (ab) to (cd) .




Figure 4.2: The solid edges in black are wired boundary arcs (ab) and (cd) , the solid edges in red are dual-wired boundary arcs (b^*c^*) and (d^*a^*) . The edges in blue are the boundary arcs $(a^\diamond b^\diamond)$, $(b^\diamond c^\diamond)$, $(c^\diamond d^\diamond)$, and $(d^\diamond a^\diamond)$ on the medial lattice. The thin edges are in the tree \mathcal{T} .

Theorem 4.2. Fix a quad $(\Omega; a, b, c, d)$ such that $\partial\Omega$ is C^1 and simple. Suppose that a sequence of medial quads $(\Omega_\delta^\diamond; a_\delta^\diamond, b_\delta^\diamond, c_\delta^\diamond, d_\delta^\diamond)$ converges to $(\Omega; a, b, c, d)$ as in (4.2). Consider UST on the primal domain Ω_δ with $(a_\delta b_\delta)$ wired and $(c_\delta d_\delta)$ wired. Denote by η_δ^L the Peano curve connecting a_δ^\diamond and d_δ^\diamond . Then the law of

η_δ^L converges weakly to hSLE_8 in Ω from a to d with marked points (b, c) as $\delta \rightarrow 0$.

In [Dub06], the author considers the UST in general polygon with alternating boundary conditions. Theorem 4.2 is its special case for quads. The proof in [Dub06] lacks certain essential technical detail. To be self-contained, we include a complete proof for Theorem 4.2 below.

The proof of Theorem 4.2 consists of two steps: the first step is the tightness of the Peano curves, see Lemma 4.3; the second step is constructing an holomorphic observable, see Lemmas 4.4 and 4.5. With these two at hand, we complete the proof of Theorem 4.2 in Section 4.4. Although the steps are standard, the proof involves a non-trivial calculation, see Lemma 4.7.

Lemma 4.3. *Assume the same setup as in Theorem 4.2. The family of Peano curves $\{\eta_\delta^L\}_{\delta>0}$ is tight. Furthermore, for any subsequential limit η^L , we have $\mathbb{P}[\eta^L \cap [bc] = \emptyset] = 1$.*

Proof. The proof of the tightness is similar as the proof of Theorem 4.1. The only difference is the proof of (4.7). Recall that, in the proof of Theorem 4.1, we define the event $E = \{\mathbf{trunk} \cap \mathbf{trunk}^* \neq \emptyset\}$ and show that $\mathbb{P}[E] = 0$ which gives (4.7). In the setup of Theorem 4.2, the event E does not exclude the possibility that the \mathbf{trunk} hits b or c . Thus, to get the tightness, it remains to prove

$$\mathbb{P}[\{b, c\} \cap \mathbf{trunk} \neq \emptyset] = 0. \quad (4.10)$$

It suffices to show

$$\lim_{\epsilon \rightarrow 0} \overline{\lim}_{\delta \rightarrow 0} \mathbb{P}[\eta_\delta^L \text{ hits } \partial B(c_\delta, \epsilon)] = 0. \quad (4.11)$$

The proof for b_δ is similar where one considers the reversal of η_δ^L .

Suppose $r_0 > 0$ is small enough such that $d(c, (ab)) > r_0$. We denote by l the component of $\partial B(c, r_0) \cap \Omega$ which separates (ab) and c , denote by x the intersection point of l with (cd) and by y the intersection point of l with (bc) . Moreover, we take $\epsilon_0 < r_0$ small enough. Define

$$l_{\epsilon_0} := l \setminus (B(x, \epsilon_0) \cup B(y, \epsilon_0)), \quad l_{x, \epsilon_0} := B(x, \epsilon_0) \cap l, \quad l_{y, \epsilon_0} := B(y, \epsilon_0) \cap l.$$

We choose the discrete approximation $l_{\epsilon_0}^\delta$, l_{x, ϵ_0}^δ and l_{y, ϵ_0}^δ of these three arcs on the primal graph Ω_δ . Recall that γ_δ^M is the unique path in the uniform spanning tree \mathcal{T}_δ from $(a_\delta b_\delta)$ to $(c_\delta d_\delta)$. Suppose $\epsilon < \epsilon_0$ is small enough. We define the following four events:

$$\begin{aligned} A_\delta &:= \{\gamma_\delta^M \text{ hits } B(c_\delta, \epsilon) \text{ before } (c_\delta d_\delta)\}, & A_{1, \delta} &:= \{\gamma_\delta^M \text{ hits } l_{\epsilon_0}^\delta \cup l_{x, \epsilon_0}^\delta \cup l_{y, \epsilon_0}^\delta \text{ at } l_{x, \epsilon_0}^\delta\}, \\ A_{2, \delta} &:= \{\gamma_\delta^M \text{ hits } l_{\epsilon_0}^\delta \cup l_{x, \epsilon_0}^\delta \cup l_{y, \epsilon_0}^\delta \text{ at } l_{\epsilon_0}^\delta\}, & A_{3, \delta} &:= \{\gamma_\delta^M \text{ hits } l_{\epsilon_0}^\delta \cup l_{x, \epsilon_0}^\delta \cup l_{y, \epsilon_0}^\delta \text{ at } l_{y, \epsilon_0}^\delta\}. \end{aligned}$$

Note that

$$\begin{aligned} \mathbb{P}[\eta_\delta^L \text{ hits } B(c_\delta, \epsilon)] &\leq \mathbb{P}[\{\gamma_\delta^M \text{ hits } B(c_\delta, \epsilon)\} \cap \{\gamma_\delta^M \text{ hits } l_{\epsilon_0}^\delta \cup l_{x, \epsilon_0}^\delta \cup l_{y, \epsilon_0}^\delta \text{ at } l_{\epsilon_0}^\delta\}] + \mathbb{P}[A_{1, \delta} \cup A_{3, \delta}] \\ &\leq \mathbb{P}[A_\delta \cap A_{2, \delta}] + \mathbb{P}[A_{1, \delta} \cup A_{3, \delta}]. \end{aligned}$$

We first estimate $\mathbb{P}[A_\delta \cap A_{2, \delta}]$. Denote by $\tilde{\Omega}_\delta$ the graph obtained from Ω_δ by regarding $(a_\delta b_\delta)$ as a single vertex and regarding $(c_\delta d_\delta)$ as a single vertex. We denote by $\tilde{\mathcal{R}}$ simple random walk on $\tilde{\Omega}_\delta$ and by \mathcal{R} simple random walk on Ω_δ . Wilson's algorithm tells that γ_δ^M has the same law as the loop-erased simple random walk on $\tilde{\Omega}_\delta$ from $(a_\delta b_\delta)$ to $(c_\delta d_\delta)$. Thus, we have

$$\begin{aligned} \mathbb{P}[A_\delta \cap A_{2, \delta}] &\leq \mathbb{P}[\tilde{\mathcal{R}} \text{ starting from } (a_\delta b_\delta) \text{ hits } l_{\epsilon_0}^\delta \text{ first and then hits } B(c_\delta, \epsilon) \text{ before } (c_\delta d_\delta)] \\ &\leq \sum_{z_\delta \in l_{\epsilon_0}^\delta} \mathbb{P}[\tilde{\mathcal{R}} \text{ starting from } (a_\delta b_\delta) \text{ hits } l_{\epsilon_0}^\delta \text{ at } z_\delta] \mathbb{P}[\tilde{\mathcal{R}} \text{ starting from } z_\delta \text{ hits } B(c_\delta, \epsilon) \text{ before } (c_\delta d_\delta)] \\ &\leq \max_{z_\delta \in l_{\epsilon_0}^\delta} \mathbb{P}[\tilde{\mathcal{R}} \text{ starting from } z_\delta \text{ hits } B(c_\delta, \epsilon) \text{ before } (c_\delta d_\delta)] \\ &\leq \max_{z_\delta \in l_{\epsilon_0}^\delta} \mathbb{P}[\tilde{\mathcal{R}} \text{ starting from } z_\delta \text{ hits } B(c_\delta, \epsilon) \cup (a_\delta b_\delta) \text{ before } (c_\delta d_\delta)] \\ &= \max_{z_\delta \in l_{\epsilon_0}^\delta} \mathbb{P}[\mathcal{R} \text{ starting from } z_\delta \text{ hits } B(c_\delta, \epsilon) \cup (a_\delta b_\delta) \text{ before } (c_\delta d_\delta)]. \end{aligned}$$

The last equality is because $\tilde{\mathcal{R}}$ starting from z_δ and \mathcal{R} starting from z_δ have the same law before hitting $(a_\delta b_\delta) \cup (c_\delta d_\delta)$. We define, for every $w_\delta \in \Omega_\delta$,

$$u_{\delta,\epsilon}(w_\delta) := \mathbb{P}[\mathcal{R} \text{ starting from } w_\delta \text{ hits } B(c_\delta, \epsilon) \cup (a_\delta b_\delta) \text{ before } (c_\delta d_\delta)].$$

We denote by Ω_ϵ the connected component of $\Omega \setminus B(c, \epsilon)$ with (ab) on its boundary and denote $\partial\Omega_\epsilon \cap \partial B(c, \epsilon)$ by ∂_ϵ . By a similar argument as in the proof of Lemma B.1, the sequence $u_{\delta,\epsilon}$ converges to u_ϵ uniformly on every compact set $K \subset \Omega_\epsilon$ as $\delta \rightarrow 0$, where u_ϵ is the harmonic function in Ω_ϵ with the following boundary data: u_ϵ equals 1 on $\partial B(c, \epsilon) \cup (ab)$ and equals 0 on $(cd) \setminus B(c, \epsilon)$ and $\frac{\partial}{\partial \vec{n}} u_\epsilon = 0$ on $(bc) \cup (da) \setminus B(c, \epsilon)$, where \vec{n} is the outer normal vector. We will prove that

$$\lim_{r_0 \rightarrow 0} \lim_{\epsilon \rightarrow 0} \max_{z \in l_{\epsilon_0}} u_\epsilon(z) = 0.$$

Let ϕ be the conformal map from Ω onto $(0, 1) \times (0, iK)$ sending (a, b, c, d) to $(0, 1, 1 + iK, iK)$. We denote by $\text{hm}(w, S)$ the harmonic measure of S from w in the strip $T := (-\infty, +\infty) \times (0, iK)$ for every $w \in T$ and $S \subset T$. We define $B_{j,\epsilon} := (\phi(\partial_\epsilon) + 2j) \cup (2j + 2 - \phi(\partial_\epsilon))$. Then, by the boundary data of u_ϵ , we have $u_\epsilon(z) = \text{hm}(\phi(z), \cup_{j \in \mathbb{Z}} B_{j,\epsilon} \cup \mathbb{R})$. Thus, we have

$$\begin{aligned} u_\epsilon(z) &\leq \mathbb{P}[\text{Brownian motion starting from } \phi(z) \text{ exits } T \text{ from } \mathbb{R}] \\ &\quad + \mathbb{P}[\text{Brownian motion starting from } \phi(z) \text{ exits } T \text{ from } \cup_{j \in \mathbb{Z}} B_{j,\epsilon}]. \end{aligned}$$

Note that

$$\mathbb{P}[\text{Brownian motion starting from } \phi(z) \text{ exits } T \text{ from } \mathbb{R}] = 1 - \frac{\text{Im}\phi(z)}{K}.$$

By Beurling estimate, there exist $C > 0$ and $c > 0$ such that

$$\max_{w \in \phi(l_{\epsilon_0})} \mathbb{P}[\text{Brownian motion starting from } w \text{ exits } T \text{ from } \cup_{j \notin [-N, N]} B_{j,\epsilon}] \leq C \left(\frac{K}{N} \right)^c$$

for every $\epsilon > 0$ and for every N . Furthermore, since ϕ is continuous on $\bar{\Omega}$ by the fact that $\partial\Omega$ is C^1 , we have $\lim_{\epsilon \rightarrow 0} \text{diam}(B_{j,\epsilon}) = 0$. This implies

$$\lim_{\epsilon \rightarrow 0} \max_{w \in \phi(l_{\epsilon_0})} \mathbb{P}[\text{Brownian motion starting from } w \text{ exits } T \text{ from } \cup_{j \in [-N, N]} B_{j,\epsilon}] = 0.$$

Thus, by letting $\epsilon \rightarrow 0$, $N \rightarrow \infty$ and then $r_0 \rightarrow 0$, we have

$$\lim_{r_0 \rightarrow 0} \overline{\lim}_{\epsilon \rightarrow 0} \overline{\lim}_{\delta \rightarrow 0} \mathbb{P}[A_\delta \cap A_{2,\delta}] = 0. \quad (4.12)$$

Next, we estimate $\mathbb{P}[A_{1,\delta} \cup A_{3,\delta}]$. Note that we can choose n large enough such that

$$\cap_{\delta_0 > 0} \cup_{\delta \leq \delta_0} (A_{1,\delta} \cup A_{3,\delta}) \subset \left\{ d \left(\mathbf{trunk}_0 \left(\frac{1}{n} \right), \mathbf{trunk}_0^* \left(\frac{1}{n} \right) \right) < \epsilon_0 \right\}.$$

Then, we have

$$\lim_{\epsilon_0 \rightarrow 0} \overline{\lim}_{\delta \rightarrow 0} \mathbb{P}[A_{1,\delta} \cup A_{3,\delta}] = 0. \quad (4.13)$$

Combining (4.12) and (4.13), we obtain (4.11) and completes the proof of the tightness.

For the second statement, we already have $\eta^L \cap \{b, c\} = \emptyset$ almost surely due to (4.10), it remains to prove $\mathbb{P}[\eta^L \cap (bc) = \emptyset] = 1$. Note that the event $\{\eta^L \cap (bc) \neq \emptyset\}$ implies $\{\mathbf{trunk} \cap \mathbf{trunk}^* \neq \emptyset\}$ which has zero probability. Thus, we have $\mathbb{P}[\eta^L \cap (bc) = \emptyset] = 1$ and this completes the proof. \square

4.3 Discrete observable for UST in quads

A function $u : \mathbb{Z}^2 \rightarrow \mathbb{C}$ is called (discrete) harmonic at a vertex $x \in \mathbb{Z}^2$ if $\sum_{i=1}^4 u(x_i) = 4u(x)$, where $(x_i : i = 1, 2, 3, 4)$ are the four neighbors of x in \mathbb{Z}^2 . We say a function u is harmonic on a subgraph of \mathbb{Z}^2 if it is harmonic at all vertices in the subgraph. A function $f : \mathbb{Z}^2 \cup (\mathbb{Z}^2)^* \rightarrow \mathbb{C}$ is said to be (discrete) holomorphic around a medial vertex x^\diamond if one has $f(n) - f(s) = i(f(e) - f(w))$, where n, s, w, e are the vertices incident to x^\diamond in counterclockwise order. We say a function f is holomorphic on a subgraph of $\mathbb{Z}^2 \cup (\mathbb{Z}^2)^*$ if it is holomorphic at all vertices in the subgraph. Note that, for a discrete holomorphic function f on a subgraph of $\mathbb{Z}^2 \cup (\mathbb{Z}^2)^*$, its restriction on \mathbb{Z}^2 and its restriction on $(\mathbb{Z}^2)^*$ are both harmonic (see [DC13, Proposition 8.15]). We summarize the setup for discrete observable below.

- Consider the set of spanning trees in the primal domain Ω_δ with $(a_\delta b_\delta)$ wired and $(c_\delta d_\delta)$ wired. Denote this set by $\text{ST}(\delta)$ and denote its cardinality by $|\text{ST}(\delta)|$. Let \mathcal{T}_δ be chosen uniformly among these trees. Recall that η_δ^L is the Peano curve along \mathcal{T}_δ from a_δ^\diamond to d_δ^\diamond , and η_δ^R is the Peano curve along \mathcal{T}_δ from b_δ^\diamond to c_δ^\diamond , and γ_δ^M is the path in \mathcal{T}_δ connecting $(a_\delta b_\delta)$ to $(c_\delta d_\delta)$. For a vertex z^* in Ω_δ^* , define $u_\delta(z^*)$ to be the probability that z^* lies to the right of η_δ^L , i.e. z^* lies in the component of $\Omega_\delta^* \setminus \eta_\delta^L$ with b_δ^* and c_δ^* on the boundary.
- Consider the set of spanning forests in the primal domain Ω_δ with $(a_\delta b_\delta)$ wired and $(c_\delta d_\delta)$ wired such that it has only two trees: one of them contains the wired arc $(a_\delta b_\delta)$ and the other one contains the wired arc $(c_\delta d_\delta)$. Denote this set by $\text{SF}_2(\delta)$ and denote its cardinality by $|\text{SF}_2(\delta)|$. Let \mathcal{F}_δ be chosen uniformly among these forests. For $z \in \Omega_\delta$, define $v_\delta(z)$ to be the probability that z lies in the same tree as the wired arc $(c_\delta d_\delta)$ in \mathcal{F}_δ .

Lemma 4.4. *Define*

$$f_\delta(\cdot) := u_\delta(\cdot) + i \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} v_\delta(\cdot).$$

We view it as a function on $\Omega_\delta \cup \Omega_\delta^$: it equals u_δ on Ω_δ^* and it equals $i \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} v_\delta$ on Ω_δ . Then f_δ is discrete holomorphic on $(\Omega_\delta \cup \Omega_\delta^*) \setminus ((a_\delta b_\delta) \cup (c_\delta d_\delta) \cup (b_\delta^* c_\delta^*) \cup (d_\delta^* a_\delta^*))$. Moreover, it has the following boundary data:*

$$\begin{cases} \text{Re} f_\delta = 0, & \text{on } (d_\delta^* a_\delta^*), \\ \text{Re} f_\delta = 1, & \text{on } (b_\delta^* c_\delta^*), \\ \text{Im} f_\delta = 0, & \text{on } (a_\delta b_\delta), \\ \text{Im} f_\delta = \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|}, & \text{on } (c_\delta d_\delta). \end{cases}$$

Proof. For $z^* \in \Omega_\delta^*$, denote by $E(\mathcal{T}_\delta; z^*)$ the event that z^* lies to the right of η_δ^L . For $x \in \Omega_\delta$, denote by $E(\mathcal{F}_\delta; x)$ the event that x lies in the same tree as the wired arc $(c_\delta d_\delta)$ in \mathcal{F}_δ . Assume $\{n, s\}$ is a primal edge of Ω_δ , and the corresponding dual edge is denoted by $\{w, e\}$ such that w, s, e, n are in counterclockwise order (see Figure 4.3). Then we have

$$\begin{aligned} u_\delta(e) - u_\delta(w) &= \mathbb{P}[E(\mathcal{T}_\delta; e)] - \mathbb{P}[E(\mathcal{T}_\delta; w)] \\ &= \mathbb{P}[E(\mathcal{T}_\delta; e) \cap E(\mathcal{T}_\delta; w)^c] - \mathbb{P}[E(\mathcal{T}_\delta; w) \cap E(\mathcal{T}_\delta; e)^c] \\ &= \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} \mathbb{P}[E(\mathcal{F}_\delta; n) \cap E(\mathcal{F}_\delta; s)^c] - \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} \mathbb{P}[E(\mathcal{F}_\delta; s) \cap E(\mathcal{F}_\delta; n)^c] \\ &= \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} (\mathbb{P}[E(\mathcal{F}_\delta; n)] - \mathbb{P}[E(\mathcal{F}_\delta; s)]) \\ &= \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} (v_\delta(n) - v_\delta(s)). \end{aligned}$$

The third equal sign is due to the observation explained in Figure 4.3. This gives the discrete holomorphicity of f_δ . The boundary data is clear from the construction. \square




Figure 4.3: We denote by $\text{ST}(\delta; w, e)$ the subset of $\text{ST}(\delta)$ consisting of spanning trees such that w lies to the left of γ_δ^M and e lies to the right of γ_δ^M . We denote by $\text{SF}_2(\delta; n, s)$ the subset of $\text{SF}_2(\delta)$ consisting of spanning forests such that n lies in the same tree as $(c_\delta d_\delta)$ and s lies in the same tree as $(a_\delta b_\delta)$. Deleting the edge $\{n, s\}$ induces a bijection from $\text{ST}(\delta; w, e)$ (left) to $\text{SF}_2(\delta; n, s)$ (right).

Lemma 4.5. *Fix a quad $(\Omega; a, b, c, d)$. Suppose a sequence of medial domains $(\Omega_\delta^\circ; a_\delta^\circ, b_\delta^\circ, c_\delta^\circ, d_\delta^\circ)$ converges to $(\Omega; a, b, c, d)$ in the Carathéodory sense as $\delta \rightarrow 0$. Let $K > 0$ be the conformal modulus of the quad $(\Omega; a, b, c, d)$, and let f be the conformal map from Ω to the rectangle $(0, 1) \times (0, iK)$ which sends (a, b, c, d) to $(0, 1, 1 + iK, iK)$. Then the discrete holomorphic function f_δ in Lemma 4.4 (regarded as a function on Ω_δ by interpolating among vertices) converges to f locally uniformly as $\delta \rightarrow 0$.*

We emphasize that, in Lemma 4.5, we do not require extra regularity on $\partial\Omega$ and we only require the convergence of polygons in the Carathéodory sense.

Proof. We claim that $\left\{ \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} \right\}_{\delta > 0}$ is uniformly bounded. Assume this is true, for any sequence $\delta_n \rightarrow 0$, there exists a subsequence, still denoted by δ_n , and a constant \tilde{K} such that

$$u_{\delta_n} \rightarrow u, \quad v_{\delta_n} \rightarrow v, \quad \text{locally uniformly; and} \quad \frac{|\text{SF}_2(\delta_n)|}{|\text{ST}(\delta_n)|} \rightarrow \tilde{K}, \quad \text{as } n \rightarrow \infty.$$

From the definition of u_δ and v_δ and Beurling estimate, it is clear that $u = 1$ on (bc) and $u = 0$ on (da) , and that $v = 1$ on (cd) and $v = 0$ on (ab) . By a similar argument as in the proof of Lemma B.1, the function $g := u + i\tilde{K}v$ is conformal on Ω as $f_{(\Omega_\delta; a_\delta, b_\delta, c_\delta, d_\delta)}$ is discrete holomorphic. Moreover, if we fix a conformal map ξ from Ω onto \mathbb{U} , then the boundary data of $u \circ \xi^{-1}$ is as follows: $u \circ \xi^{-1} = 1$ on $(\xi(b)\xi(c))$, and $u = 0$ on $(\xi(d)\xi(a))$, and $\partial_n u \circ \xi^{-1} = 0$ on $(\xi(a)\xi(b)) \cup (\xi(c)\xi(d))$. Note that, the boundary data of $u \circ \xi^{-1}$ and its harmonicity uniquely determine $u \circ \xi^{-1}$. Thus, they uniquely determine u which in turn uniquely determine g . Therefore, the function g is the conformal map from Ω onto the rectangle $(0, 1) \times (0, i\tilde{K})$. This implies that $\tilde{K} = K$ and $g = f$.

It remains to show that $\left\{ \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} \right\}_{\delta > 0}$ is uniformly bounded. If this is not the case, there exists a sequence $\delta_n \rightarrow 0$ such that $\frac{|\text{SF}_2(\delta_n)|}{|\text{ST}(\delta_n)|} \rightarrow \infty$. Then, by the same argument as above, the function $\frac{|\text{ST}(\delta_n)|}{|\text{SF}_2(\delta_n)|} f_{(\Omega_{\delta_n}; a_{\delta_n}, b_{\delta_n}, c_{\delta_n}, d_{\delta_n})}$ converges to a conformal map h on Ω locally uniformly and h extends continuously to $\bar{\Omega}$. In such case, $\text{Re}h = 0$ on Ω , thus h has to be constant. But $\text{Im}h = 1$ on (cd) and $\text{Im}h = 0$ on (ab) . This is a contradiction. Therefore, $\left\{ \frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} \right\}_{\delta > 0}$ is uniformly bounded and we complete the proof. \square

As a consequence of Lemmas 4.4 and 4.5, we see that $\frac{|\text{SF}_2(\delta)|}{|\text{ST}(\delta)|} \rightarrow K$ as $\delta \rightarrow 0$. This is a special case of [KW11, Theorem 1.1] for the grove with two nodes.

4.4 Proof of Theorem 4.2

Fix a quad $(\Omega; a, b, c, d)$. Let K be its conformal modulus and denote by $f_{(\Omega; a, b, c, d)}$ the conformal map from Ω onto $(0, 1) \times (0, iK)$ sending (a, b, c, d) to $(0, 1, 1 + iK, iK)$. Suppose a sequence of medial quads

$(\Omega_\delta^\circ; a_\delta^\circ, b_\delta^\circ, c_\delta^\circ, d_\delta^\circ)$ and $(\Omega; a, b, c, d)$ satisfy the assumptions in Theorem 4.2. We choose conformal maps $\phi_\delta : \Omega_\delta^\circ \rightarrow \mathbb{H}$ with $\phi_\delta(a_\delta^\circ) = 0, \phi_\delta(d_\delta^\circ) = \infty$ and $\phi : \Omega \rightarrow \mathbb{H}$ with $\phi(a) = 0, \phi(d) = \infty$ such that ϕ_δ^{-1} converges to ϕ^{-1} uniformly on $\overline{\mathbb{H}}$. The uniform spanning tree \mathcal{T}_δ and the Peano curve η_δ^L are defined in the same way as in Section 4.2. Define $\hat{\eta}_\delta^L := \phi_\delta(\eta_\delta^L)$ and parameterize $\hat{\eta}_\delta^L$ by the half-plane capacity and parameterize η_δ^L so that $\hat{\eta}_\delta^L(t) = \phi_\delta(\eta_\delta^L(t))$.

For $\hat{\eta}_\delta^L$, denote by $(W_t^\delta, t \geq 0)$ the driving function of $\hat{\eta}_\delta^L$ and by $(g_t^\delta, t \geq 0)$ the corresponding conformal maps. Define $X_t^\delta := g_t^\delta(\phi_\delta(b_\delta)), Y_t^\delta := g_t^\delta(\phi_\delta(c_\delta))$. Denote by \mathcal{F}_t^δ the filtration generated by $(W_s^\delta, 0 \leq s \leq t)$. Let K_δ be the modulus of the quad $(\mathbb{H}; W_t^\delta, X_t^\delta, Y_t^\delta, \infty)$ and define f_t^δ to be the conformal map from \mathbb{H} onto $(0, 1) \times (0, iK_\delta)$ sending $(W_t^\delta, X_t^\delta, Y_t^\delta, \infty)$ to $(0, 1, 1 + iK_\delta, iK_\delta)$.

For η_δ^L , let τ_δ be the first time that η_δ^L hits $(c_\delta^\circ, d_\delta^\circ)$. For every $t < \tau_\delta$, the slit domain $\Omega_\delta^\circ(t)$ is defined as the component of $\Omega_\delta^\circ \setminus \eta_\delta^L[0, t]$ that contains c_δ° and d_δ° on the boundary. We define $(\Omega_\delta(t); a_\delta(t), b_\delta(t), c_\delta, d_\delta)$ to be the primal discrete quad as follows: The domain $\Omega_\delta(t)$ is the primal domain associated to $\Omega_\delta^\circ(t)$. The point $a_\delta(t)$ is the primal vertex nearest to $\eta_\delta^L(t)$. The definition of the point $b_\delta(t)$ is a little bit complicated: if $\eta_\delta^L[0, t]$ does not hit $(b_\delta^\circ, c_\delta^\circ)$, then $b_\delta(t) = b_\delta$; if $\eta_\delta^L[0, t]$ hits $(b_\delta^\circ, c_\delta^\circ)$, then $b_\delta(t)$ is the last primal vertex in $\eta_\delta^L[0, t] \cap (b_\delta^\circ, c_\delta^\circ)$. The boundary conditions for $(\Omega_\delta(t); a_\delta(t), b_\delta(t), c_\delta, d_\delta)$ are inherited from $(\Omega_\delta; a_\delta, b_\delta, c_\delta, d_\delta)$ and $\eta_\delta^L[0, t]$: the boundary arc $(a_\delta(t), b_\delta(t))$ is wired and the boundary arc (c_δ, d_δ) is wired.

Lemma 4.6. *For $z \in \Omega_\delta$, denote by $\tau_z^\delta := \inf\{t : z \notin \Omega_\delta^\circ(t)\}$. Then, the process*

$$(f_{(\Omega_\delta(t); a_\delta(t), b_\delta(t), c_\delta, d_\delta)}(z), t \geq 0)$$

is a martingale up to $\tau_z^\delta \wedge \tau_\delta$ with respect to the filtration $(\mathcal{F}_t^\delta, t \geq 0)$. Moreover, for every $\epsilon > 0$ and for any compact subset K on Ω , one has

$$\mathbb{E} \left[f_{\tau_2^\delta}^\delta \left(g_{\tau_2^\delta}^\delta(\phi_\delta(z)) \right) - f_{\tau_1^\delta}^\delta \left(g_{\tau_1^\delta}^\delta(\phi_\delta(z)) \right) \mid \mathcal{F}_{\tau_1}^\delta \right] \xrightarrow{\delta \rightarrow 0} 0. \quad (4.14)$$

uniformly on K and uniformly for any stopping times $0 < \tau_1^\delta < \tau_2^\delta < \tau_\epsilon^\delta \wedge \tau_{K, \epsilon}^\delta$, where $\tau_{K, \epsilon}^\delta := \inf\{t : d(\eta_\delta^L([0, t]), K) = \epsilon\}$ and τ_ϵ^δ is the first time that η_δ^L hits the ϵ -neighborhood of $(b_\delta^\circ, c_\delta^\circ)$.

Proof. The first conclusion that $(f_{(\Omega_\delta(t); a_\delta(t), b_\delta(t), c_\delta, d_\delta)}(z), t \geq 0)$ is a martingale comes from the construction. It remains to show (4.14). This suffices to prove that $f_{\tau_i^\delta}^\delta \circ g_{\tau_i^\delta}^\delta \circ \phi_\delta - f_{(\Omega_\delta(\tau_i^\delta); a_\delta(\tau_i^\delta), b_\delta, c_\delta, d_\delta)}$ converges to 0 uniformly on K for $i = 1, 2$. If this is not the case, there exists a sequence $\delta_n \rightarrow 0$ such that the convergence does not hold. Since $\{(\Omega_{\delta_n}(\tau_i^{\delta_n}); a_{\delta_n}(\tau_i^{\delta_n}), b_{\delta_n}, c_{\delta_n}, d_{\delta_n})\}_n$ is tight in the Carathéodory sense, there exists a subsequence, still denoted by δ_n , such that $(\Omega_{\delta_n}(\tau_i^{\delta_n}); a_{\delta_n}(\tau_i^{\delta_n}), b_{\delta_n}, c_{\delta_n}, d_{\delta_n})$ converges to a quad $(\Omega_i; a_i, b_i, c_i, d_i)$ in the Carathéodory sense for $i = 1, 2$. By Lemma 4.5, the sequence $f_{(\Omega_{\delta_n}(\tau_i^{\delta_n}); a_{\delta_n}(\tau_i^{\delta_n}), b_{\delta_n}, c_{\delta_n}, d_{\delta_n})}$ converges to $f_{(\Omega_i; a_i, b_i, c_i, d_i)}$ uniformly on K . Note that $f_{\tau_i^{\delta_n}}^{\delta_n} \circ g_{\tau_i^{\delta_n}}^{\delta_n} \circ \phi_{\delta_n}$ is the conformal map from $\Omega_{\delta_n}^\circ(\tau_i^{\delta_n})$ to a rectangle which maps $(\eta_{\delta_n}^L(\tau_i^{\delta_n}), b_{\delta_n}^\circ, c_{\delta_n}^\circ, d_{\delta_n}^\circ)$ to the four corners. By the Carathéodory convergence and the description of $f_{(\Omega_i; a_i, b_i, c_i, d_i)}$ in Lemma 4.5, the sequence $f_{\tau_i^{\delta_n}}^{\delta_n} \circ g_{\tau_i^{\delta_n}}^{\delta_n} \circ \phi_{\delta_n}$ converges to $f_{(\Omega_i; a_i, b_i, c_i, d_i)}$ on K as well. This is a contradiction. This completes the proof. \square

For $x < y < w$, define

$$\Theta(x, y, w) := \frac{2}{w-x} + \frac{-2}{w-y} - 8 \frac{y-x}{(y-w)^2} \frac{F' \left(\frac{x-w}{y-w} \right)}{F \left(\frac{x-w}{y-w} \right)}, \quad \text{where } F(z) = {}_2F_1 \left(\frac{1}{2}, \frac{1}{2}, 1; z \right). \quad (4.15)$$

Note that F is the hypergeometric function in (3.1) with $\kappa = 8, \nu = 0$.

Lemma 4.7. *Suppose $\hat{\eta}_\delta^L$ converges to $\hat{\eta}^L$ locally uniformly as $\delta \rightarrow 0$ almost surely. For every $\epsilon > 0$, we denote by τ_ϵ the first time that $\phi^{-1}(\hat{\eta}^L)$ hits the ϵ -neighbourhood of (bd) . Then, the law of the driving*

function of $\hat{\eta}^L$, denoted by W_t , is given by the following SDEs up to τ_ϵ :

$$\begin{cases} dW_t = \sqrt{8}dB_t + \Theta(X_t, Y_t, W_t)dt, & W_0 = 0; \\ dX_t = \frac{2dt}{X_t - W_t}, & X_0 = \phi(b); \\ dY_t = \frac{2dt}{Y_t - W_t}, & Y_0 = \phi(c); \end{cases} \quad (4.16)$$

where $(B_t, t \geq 0)$ is one-dimensional Brownian motion and Θ is defined in (4.15).

Proof. We define $T_M := \inf\{t : \hat{\eta}^L \text{ hits } \partial B(0, M)\}$ for every $M > 0$ and $\tau_{\epsilon', z} := \inf\{t : \hat{\eta}^L \text{ hits } \partial B(z, \epsilon')\}$ for every $z \in \mathbb{H}$ and $\epsilon' > 0$. It suffices to prove that (4.16) holds up to $\tau_\epsilon \wedge T_M$. Then, by letting $M \rightarrow \infty$, we get the result. We define T_M^δ and $\tau_{\epsilon', z}^\delta$ similarly for $\hat{\eta}_\delta^L$. We may assume $T_M^\delta \rightarrow T_M$, $\tau_{\epsilon', z}^\delta \rightarrow \tau_{\epsilon', z}$ and $\tau_\epsilon^\delta \rightarrow \tau_\epsilon$ by considering a continuous modification, see details in [Kar19] and [Kar20]. Then, Lemma 4.6 implies that

$$M_t(z) := f_{(\mathbb{H}; 0, X_t - W_t, Y_t - W_t, \infty)}(g_t(z) - W_t)$$

is a martingale up to $\tau_\epsilon \wedge T_M \wedge \tau_{\epsilon', z}$. By the same argument in [Kar20, Lemma 5.3], we can deduce that W_t is a semimartingale. We will calculate the drift term of $M_t(z)$ in terms of W_t, X_t, Y_t and $g_t(z)$.

Due to Schwarz-Christoffel formula (see e.g. [Ahl78, Chapter 6-Section 2.2]), we have, for $0 < x < y$ and for $z \in \mathbb{H}$,

$$f_{(\mathbb{H}; 0, x, y, \infty)}(z) = \frac{\int_0^{z/x} (s(s-1)(s-\frac{y}{x}))^{-1/2} ds}{\int_0^1 (s(s-1)(s-\frac{y}{x}))^{-1/2} ds}.$$

Denote by \mathcal{K} the elliptic integral of the first kind (A.6). By changing of variable $s = \sin^2 \theta$, we have

$$f_{(\mathbb{H}; 0, x, y, \infty)}(z) = \frac{\mathcal{K}(\arcsin \sqrt{z/x}, x/y)}{\mathcal{K}(x/y)}. \quad (4.17)$$

Therefore,

$$M_t(z) = \frac{\mathcal{K}(S_t, U_t)}{\mathcal{K}(U_t)}, \quad \text{where } S_t = \arcsin \sqrt{\frac{g_t(z) - W_t}{X_t - W_t}}, \quad U_t = \frac{X_t - W_t}{Y_t - W_t}. \quad (4.18)$$

We first calculate dU_t and dS_t :

$$\begin{aligned} dU_t &= \frac{1}{(Y_t - W_t)^2} \left(\left(\frac{2}{U_t} - 2U_t \right) dt + (X_t - Y_t)dW_t - (1 - U_t)d\langle W \rangle_t \right), \\ dS_t &= \frac{\cot S_t}{(X_t - W_t)^2} \left((2 + \cot^2 S_t)dt - \frac{1}{2}(X_t - W_t)dW_t - \frac{1}{8}(3 + \cot^2 S_t) d\langle W \rangle_t \right). \end{aligned} \quad (4.19)$$

Applying Itô's formula in (4.18), we have

$$dM_t(z) = \frac{1}{\mathcal{K}(U_t)} d\mathcal{K}(S_t, U_t) - \frac{\mathcal{K}(S_t, U_t)}{\mathcal{K}^2(U_t)} d\mathcal{K}(U_t) - \frac{1}{\mathcal{K}^2(U_t)} d\langle \mathcal{K}(S, U), \mathcal{K}(U) \rangle_t + \frac{\mathcal{K}(S_t, U_t)}{\mathcal{K}^3(U_t)} d\langle \mathcal{K}(U) \rangle_t. \quad (4.20)$$

We denote by L_t the drift term of W_t . Since the drift term of $M_t(z)$ is zero, plugging (4.19) into (4.20),

we have

$$\begin{aligned}
0 = & \frac{\partial_\varphi \mathcal{K}(S_t, U_t)}{\mathcal{K}(U_t)} \frac{\cot S_t}{(X_t - W_t)^2} \left((2 + \cot^2 S_t) dt - \frac{1}{2} (X_t - W_t) dL_t - \frac{1}{8} (3 + \cot^2 S_t) d\langle W \rangle_t \right) \\
& + \frac{\partial_x \mathcal{K}(S_t, U_t)}{\mathcal{K}(U_t)} \frac{1}{(Y_t - W_t)^2} \left(\left(\frac{2}{U_t} - 2U_t \right) dt + (X_t - Y_t) dL_t - (1 - U_t) d\langle W \rangle_t \right) \\
& + \frac{1}{8} \frac{\partial_\varphi^2 \mathcal{K}(S_t, U_t)}{\mathcal{K}(U_t)} \frac{\cot^2 S_t}{(X_t - W_t)^2} d\langle W \rangle_t + \frac{1}{2} \frac{\partial_x^2 \mathcal{K}(S_t, U_t)}{\mathcal{K}(U_t)} \frac{(X_t - Y_t)^2}{(Y_t - W_t)^4} d\langle W \rangle_t \\
& - \frac{1}{2} \frac{\partial_x \partial_\varphi \mathcal{K}(S_t, U_t)}{\mathcal{K}(U_t)} \frac{(X_t - Y_t) \cot S_t}{(Y_t - W_t)^2 (X_t - W_t)} d\langle W \rangle_t \\
& - \frac{\mathcal{K}(S_t, U_t)}{\mathcal{K}^2(U_t)} \frac{\partial_x \mathcal{K}(U_t)}{(Y_t - W_t)^2} \left(\left(\frac{2}{U_t} - 2U_t \right) dt + (X_t - Y_t) dL_t - (1 - U_t) d\langle W \rangle_t \right) \\
& - \frac{1}{2} \frac{\mathcal{K}(S_t, U_t)}{\mathcal{K}^2(U_t)} \frac{(X_t - Y_t)^2}{(Y_t - W_t)^4} \partial_x^2 \mathcal{K}(U_t) d\langle W \rangle_t - \frac{\partial_x \mathcal{K}(U_t) \partial_x \mathcal{K}(S_t, U_t)}{\mathcal{K}^2(U_t)} \frac{(X_t - Y_t)^2}{(Y_t - W_t)^4} d\langle W \rangle_t \\
& + \frac{\partial_x \mathcal{K}(U_t) \partial_\varphi \mathcal{K}(S_t, U_t)}{\mathcal{K}^2(U_t)} \frac{(X_t - Y_t) \cot S_t}{2(Y_t - W_t)^2 (X_t - W_t)} d\langle W \rangle_t \\
& + \frac{\mathcal{K}(S_t, U_t)}{\mathcal{K}^3(U_t)} \frac{(X_t - Y_t)^2}{(Y_t - W_t)^4} (\partial_x \mathcal{K}(U_t))^2 d\langle W \rangle_t.
\end{aligned} \tag{4.21}$$

Note that

$$g_t(z) - W_t \rightarrow 0, \quad S_t \rightarrow 0, \quad \sqrt{g_t(z) - W_t} \cot S_t \rightarrow \sqrt{X_t - W_t}, \quad \text{as } z \rightarrow \hat{\eta}^L(t).$$

Combining with (A.8) and the trivial facts $\partial_x \mathcal{K}(\varphi, x), \partial_x^2 \mathcal{K}(\varphi, x) \rightarrow 0$ as $\varphi \rightarrow 0$, we have

$$\begin{aligned}
\mathcal{K}(S_t, U_t) & \rightarrow 0, \quad \partial_\varphi \mathcal{K}(S_t, U_t) \rightarrow 1, \quad \partial_x \mathcal{K}(S_t, U_t) \rightarrow 0, \\
\frac{\partial_\varphi^2 \mathcal{K}(S_t, U_t)}{\sqrt{g_t(z) - W_t}} & \rightarrow \frac{U_t}{\sqrt{X_t - W_t}}, \quad \frac{\partial_x \partial_\varphi \mathcal{K}(S_t, U_t)}{g_t(z) - W_t} \rightarrow \frac{1}{2(X_t - W_t)}, \quad \partial_x^2 \mathcal{K}(S_t, U_t) \rightarrow 0, \quad \text{as } z \rightarrow \hat{\eta}^L(t).
\end{aligned}$$

In the right hand-side of (4.21), the leading term is of order $\cot^3 S_t$. Dividing (4.21) by $\cot^3 S_t$ and letting $z \rightarrow \hat{\eta}^L(t)$, we have

$$d\langle W \rangle_t = 8dt. \tag{4.22}$$

Plugging (4.22) into (4.21), the leading term now is of order $\cot S_t$. Dividing (4.21) by $\cot S_t$ and letting $z \rightarrow \hat{\eta}^L(t)$, we have

$$-\frac{1}{X_t - W_t} \left(dt + \frac{1}{2} (X_t - W_t) dL_t \right) + \frac{U_t}{X_t - W_t} dt + \frac{4\partial_x \mathcal{K}(U_t)}{\mathcal{K}(U_t)} \frac{X_t - Y_t}{(Y_t - W_t)^2} dt = 0.$$

By (A.7), we have

$$dL_t = \Theta(X_t, Y_t, W_t) dt. \tag{4.23}$$

Combining (4.22) and (4.23), we obtain the conclusion. \square

Proof of Theorem 4.2. By Lemma 4.3, we may choose a subsequence $\delta_n \rightarrow 0$ such that $\eta_{\delta_n}^L$ converges in law as $n \rightarrow \infty$. Denote by η^L the limit and define $\hat{\eta}^L := \phi(\eta^L)$. By Lemma 4.3, the sequence $\{\hat{\eta}_{\delta_n}^L|_{[0,t]}\}_n$ is tight. Thus, by the diagonal method and Skorokhod's representation theorem, we can choose a subsequence, still denoted by δ_n , such that $\hat{\eta}_{\delta_n}^L$ converges to $\hat{\eta}^L$ locally uniformly as $n \rightarrow \infty$ almost surely. We may assume that (by subtracting a further subsequence), in such coupling, the trunk $\{\mathbf{trunk}_{\delta_n}^* (\frac{1}{m})\}_n$, defined in the proof of Theorem 4.1, converges to $\mathbf{trunk}_0^* (\frac{1}{m})$ in Hausdorff distance for every $m \in \mathbb{N}$. It suffices to prove that the driving function of $\hat{\eta}^L$ has the same law as the one for hSLE₈.

Define

$$\tau_{\delta_n, \epsilon} := \inf\{t > 0 : d(\eta_{\delta_n}^L(t), (c_{\delta_n}^\diamond d_{\delta_n}^\diamond)) = \epsilon\}, \quad \tau := \inf\{t > 0 : \eta^L(t) \text{ hits } (cd)\}.$$

Recall that τ_{δ_n} is the first time that $\eta_{\delta_n}^L$ hits $(c_{\delta_n}^\diamond d_{\delta_n}^\diamond)$. First, we will show

$$\lim_{n \rightarrow \infty} \tau_{\delta_n} = \tau \quad \text{almost surely.} \quad (4.24)$$

It is clear that

$$\lim_{\epsilon \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} \tau_{\delta_n, \epsilon} \leq \tau \leq \underline{\lim}_{n \rightarrow \infty} \tau_{\delta_n} \leq \overline{\lim}_{n \rightarrow \infty} \tau_{\delta_n} \quad \text{almost surely.}$$

Denote by $T = \lim_{\epsilon \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} \tau_{\delta_n, \epsilon}$. By Lemma 4.3, we have $\hat{\eta}^L(T) \in (cd)$. If (4.24) fails, there exists t between T and $\overline{\lim}_{n \rightarrow \infty} \tau_{\delta_n}$ such that $d := d(\eta^L(t), \partial\Omega) > 0$. By the locally uniformly convergence, we have $d(\eta_{\delta_n}^L(t), \partial\Omega_{\delta_n}) > \frac{d}{2}$ and $t \in (\tau_{\delta_n, \epsilon}, \tau_{\delta_n})$ for a subsequence of $\{\delta_n\}$, still denoted by $\{\delta_n\}$, for some ϵ . In such case, we can choose $v \in \Omega_{\delta_n}^*$ adjacent to the right-side of $\eta_{\delta_n}^L$ such that it connects to $(b_{\delta_n}^* c_{\delta_n}^*)$ in the dual tree by a unique path which we denote by $T_{v,n}$. Then, we have $\text{diam}(T_{v,n}) > \frac{d}{2}$. Since the trunk $\{\mathbf{trunk}_{\delta_n}^* (\frac{1}{m})\}_n$ converges to $\mathbf{trunk}_0^* (\frac{1}{m})$ in Hausdorff distance for every $m \in \mathbb{N}$, this implies $\{\mathbf{trunk} \cap \mathbf{trunk}^* \neq \emptyset\}$. Note that $\{\mathbf{trunk} \cap \mathbf{trunk}^* \neq \emptyset\}$ has zero probability. This implies (4.24).

Second, we see that the driving function of $\hat{\eta}^L$, denoted by W , solves (4.16) up to τ due to Lemma 4.7. By Lemma 4.3, the curve η^L does not hit $[bc]$. We define $x := \phi(b)$ and $y := \phi(c)$. By Lemma 4.7, we can couple W and a Brownian motion B together such that, for $t < \tau$,

$$W_t = \sqrt{8}B_t + \int_0^t \frac{2ds}{W_s - V_s^x} + \int_0^t \frac{-2ds}{W_s - V_s^y} - 8 \int_0^t \frac{F'(Z_s)}{F(Z_s)} \left(\frac{1 - Z_s}{V_s^y - W_s} \right) ds.$$

Third, we prove that W solves (4.16) up to and including τ . Note that, for any $t < \tau$,

$$W_t = \sqrt{8}B_t - 2 \int_0^t \frac{V_s^y - V_s^x}{(W_s - V_s^x)(W_s - V_s^y)} ds - 8 \int_0^t \frac{F'(Z_s)}{F(Z_s)} \left(\frac{1 - Z_s}{V_s^y - W_s} \right) ds.$$

Moreover, since $\hat{\eta}_{\delta_n}^L$ converges to $\hat{\eta}^L$ locally uniformly as $n \rightarrow \infty$, the driving function of $\hat{\eta}_{\delta_n}^L$ converges to W locally uniformly as $n \rightarrow \infty$. This implies $W : [0, \infty) \rightarrow \mathbb{R}$ is a continuous function. Thus, we have

$$8 \int_0^t \frac{F'(Z_s)}{F(Z_s)} \left(\frac{1 - Z_s}{V_s^y - W_s} \right) ds \leq \max_{t \in [0, \tau]} |W_t - \sqrt{8}B_t| < \infty.$$

Then, by monotone convergence theorem, we have

$$\lim_{t \rightarrow \tau} \int_0^t \frac{F'(Z_s)}{F(Z_s)} \left(\frac{1 - Z_s}{V_s^y - W_s} \right) ds = \int_0^\tau \frac{F'(Z_s)}{F(Z_s)} \left(\frac{1 - Z_s}{V_s^y - W_s} \right) ds < \infty.$$

We take $z \in \mathbb{R}$ such that $z > \hat{\eta}^L(\tau)$. For any $t < \tau$, we have $g_t(y) \leq g_t(z)$. This implies

$$\int_0^t \frac{2ds}{V_s^y - W_s} \leq g_t(z).$$

By monotone convergence theorem, we have

$$\lim_{t \rightarrow \tau} \int_0^t \frac{2ds}{V_s^x - W_s} = \int_0^\tau \frac{2ds}{V_s^x - W_s} \leq \lim_{t \rightarrow \tau} \int_0^t \frac{2ds}{V_s^y - W_s} = \int_0^\tau \frac{2ds}{V_s^y - W_s} \leq g_\tau(z) < \infty.$$

Therefore, letting $t \rightarrow \tau$, we have

$$W_\tau = \sqrt{8}B_\tau + \int_0^\tau \frac{2ds}{W_s - V_s^x} + \int_0^\tau \frac{-2ds}{W_s - V_s^y} - 8 \int_0^\tau \frac{F'(Z_s)}{F(Z_s)} \left(\frac{1 - Z_s}{V_s^y - W_s} \right) ds.$$

In other words, the driving function W of $\hat{\eta}^L$ has the same law as the one for hSLE₈ up to and including τ . In this step, it is important that W is continuous up to and including τ which is a consequence of Lemma 4.3.

Finally, we will show that the driving function of $\eta^L[\tau, \infty]$ given $\eta^L[0, \tau]$ is $\sqrt{8}$ times Brownian motion. Denote by $\Omega(\tau)$ the connected component of $\Omega \setminus \eta^L[0, \tau]$ having d on its boundary. From above, we know that $\Omega_{\delta_n}(\tau_{\delta_n})$ converges to $\Omega(\tau)$ in the Carathéodory sense by Carathéodory kernel theorem. By the domain Markov property, conditioning on $\eta_{\delta_n}[0, \tau_{\delta_n}]$, the remaining curve $\eta_{\delta_n}[\tau_{\delta_n}, \infty]$ has the same law as the Peano curve from $\eta_{\delta_n}(\tau_{\delta_n})$ to $d_{\delta_n}^\circ$ in $\Omega_{\delta_n}(\tau_{\delta_n})$ with Dobrushin boundary conditions. By [LSW04, Theorem 4.4](see also the proof of Theorem 4.1), the driving function of $\hat{\eta}^L[\tau, \infty]$ has the same law as the driving function of SLE₈ in Ω_τ from $\eta(\tau)$ to d . It is important that the convergence of driving function only requires the convergence of domains in the Carathéodory sense and there is no regularity requirement on the boundary of the limiting domain. Thus, the driving function of $\hat{\eta}^L[\tau, \infty]$ given $\hat{\eta}^L[0, \tau]$ is $\sqrt{8}$ times Brownian motion.

In summary, the driving function of $\hat{\eta}^L$ is the same as the one for hSLE₈ as a whole process. This completes the proof. \square

As a consequence of Theorem 4.2, we have the following.

Corollary 4.8. *Fix a quad $(\Omega; a, b, c, d)$. The process $\eta \sim \text{hSLE}_8$ in Ω from a to d with marked points (b, c) has the following properties: It is almost surely generated by continuous curve and $\eta \cap [b, c] = \emptyset$ almost surely. Moreover, it is reversible: the time-reversal of η has the law of hSLE₈ in Ω from d to a with marked points (c, b) .*

Proof. We may assume that $\partial\Omega$ is C^1 and simple. Note that, if the conclusion holds under such assumption, the conclusion would also hold for a general quad with locally connected boundary via conformal image.

First of all, we argue that there exists a unique solution in law to the SDE (4.16) up to and including τ —the first time that $\phi(b)$ is swallowed. From the proof of Theorem 4.2, we see that there exists a version of solution W to the SDE (4.16) up to τ . Moreover, it is generated by a continuous curve η up to and including τ and $\eta \cap [\phi(b), \phi(c)] = \emptyset$. Suppose \tilde{W} is another solution. Denote by τ_ϵ the first time that the process gets within ϵ -neighborhood of $(\phi(b), \infty)$. As the SDE (4.16) has a unique solution up to τ_ϵ , the two processes W and \tilde{W} have the same law up to τ_ϵ . We may couple them so that $W_t = \tilde{W}_t$ for $t \leq \tau_\epsilon$. As the family of the laws of $\{W|_{[0, \tau_\epsilon]}, \tilde{W}|_{[0, \tau_\epsilon]}\}_{\epsilon > 0}$ is tight, there exists subsequence $\epsilon_n \rightarrow 0$ along which W and \tilde{W} converge. Therefore, $W_t = \tilde{W}_t$ for all $t < T$ where T is the first hitting time of $[\phi(b), \infty)$. Since $\eta \cap [\phi(b), \phi(c)] = \emptyset$, we have $\tau = T$. This implies that the SDE (4.16) has a unique solution up to and including τ , which is given by the limit of the Peano curve in Theorem 4.2. Then, the continuity of hSLE₈ is a consequence of Lemma 4.3. For the reversibility, we denote by $\mathcal{R}(\eta_\delta^L)$ the time-reversal of η_δ^L . By Theorem 4.2, the law of $\mathcal{R}(\eta_\delta^L)$ converges to hSLE₈ in Ω from d to a with marked points (c, b) as $\delta \rightarrow 0$. This implies the reversibility and completes the proof. \square

4.5 Proof of Theorem 1.5

In this section, we will complete the proof of Theorem 1.5. Before that, we first show the tightness of the LERW branch.

Lemma 4.9. *Assume the same setup as in Theorem 1.5. Then $\{\gamma_\delta^M\}_{\delta > 0}$ is tight. Moreover, any subsequential limit is a simple curve in $\bar{\Omega}$ which intersects $\partial\Omega$ only at two ends. Furthermore, the two ends are one in (ab) and one in (cd) .*

Proof. The proof is similar as the proof of [LSW04, Theorem 1.1], we summarize it here for concreteness. First, we prove the tightness. Suppose $\Upsilon : (0, \infty) \rightarrow (0, 1]$ is an increasing function. Denote by $\chi_\Upsilon(\Omega)$ the

space of simple curves $\gamma : [0, 1] \rightarrow \bar{\Omega}$, such that for every $0 \leq s_1 < s_2 \leq 1$,

$$d(\gamma[0, s_1], \gamma[s_2, 1]) \geq \Upsilon(\text{diam}(\gamma[s_1, s_2])).$$

We claim that for $\epsilon > 0$, there exists Υ such that

$$\mathbb{P}[\gamma_\delta^M \in \chi_\Upsilon(\Omega_\delta)] \geq 1 - \epsilon, \quad \text{for all } \delta. \quad (4.25)$$

Assuming this is true. We choose $R > 0$ such that $\Omega_\delta \subset B(0, R)$ for all δ . Then, we have

$$\mathbb{P}[\gamma_\delta^M \in \chi_\Upsilon(B(0, R))] \geq 1 - \epsilon.$$

By the same argument as in [LSW04, Lemma 3.10], the set $\chi_\Upsilon(B(0, R))$ is a compact set of curves. This completes the proof of tightness.

Second, we prove that any subsequential limit is a simple curve in $\bar{\Omega}$ which intersects $\partial\Omega$ only at two ends such that one in (ab) and one in (cd) . Denote by S_ϵ the ϵ -neighbourhood of Ω . By the convergence of $\partial\Omega_\delta$, it is clear that any subsequential limit is a simple curve on S_ϵ for every $\epsilon > 0$. This implies that any subsequential limit is in $\bar{\Omega}$. Suppose γ^M is a subsequential limit and $\gamma_{\delta_n}^M \rightarrow \gamma^M$. We may couple $\{(\gamma_{\delta_n}^M, \eta_{\delta_n}^L)\}$ and (γ^M, η^L) together such that $\gamma_{\delta_n}^M \rightarrow \gamma^M$ and $\eta_{\delta_n}^L \rightarrow \eta^L$ almost surely. Recall that τ_{δ_n} is the first time that $\eta_{\delta_n}^L$ hits $(c_{\delta_n}^\diamond, d_{\delta_n}^\diamond)$ and τ is the first time that η^L hits (cd) . As explained in the proof of Theorem 4.2, we can modify this coupling such that $\tau_{\delta_n} \rightarrow \tau$ almost surely. Note that if γ^M intersects $[cd]$ at more than two points, we have $\tau < \liminf_{n \rightarrow \infty} \tau_{\delta_n}$. This is a contradiction. Define $Y^M := \gamma^M \cap [cd]$ and $Y_{\delta_n}^M = \gamma_{\delta_n}^M \cap [c_{\delta_n}^\diamond, d_{\delta_n}^\diamond]$. Note that $d(\eta_{\delta_n}^L(\tau_{\delta_n}), Y_{\delta_n}^M) \leq \delta_n$. This implies that $Y^M = \eta^L(\tau)$. Since $\eta^L \cap [bc] = \emptyset$, we have $Y^M \neq c$. Similarly, we have $Y^M \neq d$. Thus γ^M intersects $[cd]$ only at one point in (cd) . Similarly, we have that γ^M intersects $[ab]$ only at one point in (ab) .

Finally, it remains to prove (4.25). It suffices to show it for any sequence $\delta_n \rightarrow 0$. For $\alpha, \beta > 0$, denote by $\mathcal{A}_{\delta_n}(\beta, \alpha)$ the event that there exists $0 \leq s_1 < s_2 \leq 1$ such that $d(\gamma_{\delta_n}^M[0, s_1], \gamma_{\delta_n}^M[s_2, 1]) \leq \alpha$ but $\text{diam}(\gamma_{\delta_n}^M[s_1, s_2]) \geq \beta$. Note that $\cap_{\alpha > 0} \cap_{i=1}^\infty \cup_{n=i}^\infty \mathcal{A}_{\delta_n}(2^{-m}, \alpha) \subset \{\mathbf{trunk} \cap \mathbf{trunk}^* \neq \emptyset\}$ which has zero probability. Thus, we can choose α_m such that

$$\mathbb{P}[\mathcal{A}_{\delta_n}(2^{-m}, \alpha_m)] \leq \frac{\epsilon}{2^m}, \quad \text{for all } n.$$

We choose Υ such that $\Upsilon(t) < \alpha_m$ for every $t \leq 2^{1-m}$. Then, we have

$$\mathbb{P}[\gamma_{\delta_n}^M \in \chi_\Upsilon(\Omega_{\delta_n})] \geq \mathbb{P}[(\cup_{m=1}^\infty \mathcal{A}_{\delta_n}(2^{-m}, \alpha_m))^c] \geq 1 - \epsilon.$$

This gives (4.25) and completes the proof. \square

Proof of Theorem 1.5. Note that, the families $\{\eta_\delta^L\}_{\delta > 0}$ and $\{\eta_\delta^R\}_{\delta > 0}$ are tight due to Lemma 4.3; and the family $\{\gamma_\delta^M\}_{\delta > 0}$ is tight due to Lemma 4.9. For any sequence $\delta_n \rightarrow 0$, there exists a subsequence, still denoted by δ_n , such that $\{(\eta_{\delta_n}^L; \gamma_{\delta_n}^M; \eta_{\delta_n}^R)\}$ converges in law as $n \rightarrow \infty$. We couple $\{(\eta_{\delta_n}^L; \gamma_{\delta_n}^M; \eta_{\delta_n}^R)\}$ together such that $\eta_{\delta_n}^L \rightarrow \eta^L$ and $\gamma_{\delta_n}^M \rightarrow \gamma^M$ and $\eta_{\delta_n}^R \rightarrow \eta^R$ as curves almost surely as $n \rightarrow \infty$. We will prove that the law of the triple $(\eta^L; \gamma^M; \eta^R)$ is the one in Theorem 1.5.

First of all, the law of η^L is hSLE $_\delta$ in Ω from a to d with marked points (b, c) due to Theorem 4.2.

Next, we derive the conditional law of η^R given γ^M . Denote by $\Omega_{\delta_n}^R$ the connected component of $\Omega_{\delta_n} \setminus \gamma_{\delta_n}^M$ which contains $[b_{\delta_n}, c_{\delta_n}]$ on its boundary. Denote by $\Omega_{\delta_n}^{\diamond, R}$ the medial graph associated with $\Omega_{\delta_n}^R$. Recall that we denote by Ω^R the connected component of $\Omega \setminus \gamma^M$ which contains $[bc]$ on its boundary. This is well-defined since we have $\gamma^M \cap [bc] = \emptyset$ almost surely due to Lemma 4.9. Since $\gamma_{\delta_n}^M \rightarrow \gamma^M$ as curves, the medial Dobrushin domains $(\Omega_{\delta_n}^{\diamond, R}; b_{\delta_n}^\diamond, c_{\delta_n}^\diamond)$ converges to $(\Omega^R; b, c)$ as in (4.2). Note that in the proof of Theorem 4.1, we use the condition that $\partial\Omega$ is C^1 to ensure that $\mathbf{trunk} \cap \mathbf{trunk}^* = \emptyset$ almost surely. Although we do not have C^1 regularity on $\partial\Omega^R$, but we already have $\mathbf{trunk} \cap \mathbf{trunk}^* = \emptyset$ almost surely.

This is because $\mathbf{trunk}_{\delta_n}$ in the primal graph associated with the medial Dobrushin domain $(\Omega_{\delta_n}^{\diamond,R}; b_{\delta_n}^{\diamond}, c_{\delta_n}^{\diamond})$ is a subset of $\mathbf{trunk}_{\delta_n}$ in $(\Omega_{\delta_n}; a_{\delta_n}, b_{\delta_n}, c_{\delta_n}, d_{\delta_n})$ and $\mathbf{trunk}_{\delta_n}^*$ in the dual graph associated with the medial Dobrushin domain $(\Omega_{\delta_n}^{\diamond,R}; b_{\delta_n}^{\diamond}, c_{\delta_n}^{\diamond})$ is a subset of $\mathbf{trunk}_{\delta_n}^*$ in $(\Omega_{\delta_n}^*; a_{\delta_n}^*, b_{\delta_n}^*, c_{\delta_n}^*, d_{\delta_n}^*)$. Thus, by the same proof of Theorem 4.1, we have that η^R has the same law as SLE_8 in Ω^R from b to c . Similarly, we have η^L is SLE_8 in Ω^L from a to d .

From the above argument, we see that, the conditional law of $\eta_{\delta_n}^R$ given $\gamma_{\delta_n}^M$ converges to SLE_8 , and the conditional law of $\eta_{\delta_n}^L$ given $\gamma_{\delta_n}^M$ converges to SLE_8 as well. These imply that η^R and η^L are conditionally independent given γ^M . Since SLE_8 is space-filling, we have $\gamma^M = \eta^L \cap \eta^R$. This completes the proof. \square

Corollary 4.10. *Consider the continuous curve γ^M in the triple of Theorem 1.5. We have $\mathbb{P}[z \in \gamma^M] = 0$ for any $z \in \Omega$ and $\text{Leb}(\gamma^M) = 0$ almost surely.*

Proof. From Theorem 4.2, the law of η^L is hSLE_8 in Ω from a to d with marked points (b, c) . From Theorem 1.5, the curve γ^M is the part of the boundary of η^L inside Ω . We parameterize γ^M so that $\gamma^M(0) = X^M$ and $\gamma^M(1) = Y^M$. Let η be an SLE_8 in Ω from a to d and denote by τ the first time that η swallows b . From (3.11), the law of η^L is absolutely continuous with respect to η up to τ . As the frontier of SLE_8 has zero Lebesgue measure, we have $\text{Leb}(\gamma^M[0, t]) = 0$ almost surely for any $t < 1$. As $\gamma^M = \cup_n \gamma^M[0, 1 - 1/n]$, we have $\text{Leb}(\gamma^M) = 0$ almost surely as desired. \square

5 Convergence of LERW in quads

5.1 The pair of random points (X^M, Y^M)

The goal of this section is to derive the limiting distribution of the pair of random points (X^M, Y^M) in Theorem 1.6. We summarize the setup for the conclusion below.

- Fix a quad $(\Omega; a, b, c, d)$ such that $\partial\Omega$ is C^1 and simple. Suppose that a sequence of medial quads $(\Omega_{\delta}; a_{\delta}^{\diamond}, b_{\delta}^{\diamond}, c_{\delta}^{\diamond}, d_{\delta}^{\diamond})$ converges to $(\Omega; a, b, c, d)$ as in (4.2). Assume the same setup as in Section 4.2. We consider the UST \mathcal{T}_{δ} in Ω_{δ} with $(a_{\delta}b_{\delta})$ wired and $(c_{\delta}d_{\delta})$ wired. There are two Peano curves running along \mathcal{T}_{δ} , and we denote by η_{δ}^L the one from a_{δ}^{\diamond} to d_{δ}^{\diamond} and by η_{δ}^R the one from b_{δ}^{\diamond} to c_{δ}^{\diamond} . There exists a unique branch in \mathcal{T}_{δ} , denoted by γ_{δ}^M , connecting $(a_{\delta}b_{\delta})$ to $(c_{\delta}d_{\delta})$. Recall that $X_{\delta}^M := \gamma_{\delta}^M \cap (a_{\delta}b_{\delta})$ and $Y_{\delta}^M := \gamma_{\delta}^M \cap (c_{\delta}d_{\delta})$.
- For the quad $(\Omega; a, b, c, d)$, denote by $f = f_{(\Omega; a, b, c, d)}$ the conformal map from Ω onto $(0, 1) \times (0, iK)$ which sends (a, b, c, d) to $(0, 1, 1 + iK, iK)$ and extend its definition continuously to the boundary.
- Consider the Poisson kernel for the rectangle $f(\Omega) = (0, 1) \times (0, iK)$. Define, for all $r \in (f(a)f(b)) \cup (f(c)f(d))$ and for all $z \in [0, 1] \times [0, iK] \setminus \{r\}$,

$$P_K(z, r) = \text{Im} \sum_{n \in \mathbb{Z}} \left(\frac{1}{\exp(\frac{\pi}{K}(2n - r + z)) - 1} + \frac{1}{\exp(\frac{\pi}{K}(2n - r - \bar{z})) - 1} \right). \quad (5.1)$$

Note that $P_K(\cdot, r)$ is continuous on $[0, 1] \times [0, iK] \setminus \{r\}$, and it is harmonic on $(0, 1) \times (0, iK)$ with the following boundary data:

$$P_K(\cdot, r) = 0 \text{ on } (f(a)f(b)) \cup (f(c)f(d)) \setminus \{r\}, \quad \partial_n P_K(\cdot, r) = 0 \text{ on } (f(b)f(c)) \cup (f(d)f(a)), \quad (5.2)$$

where n is the outer normal vector.

Proposition 5.1. *The pair $(X_{\delta}^M, Y_{\delta}^M)$ converges weakly to a random pair of points (X^M, Y^M) as $\delta \rightarrow 0$. Denote by $x^M := f(X^M)$ and $y^M := \text{Ref}(Y^M)$. The law of the pair (x^M, y^M) is characterized by the following.*

(1) The law of x^M is uniform on $(0, 1)$.

(2) The conditional density of y^M given $x^M \in (0, 1)$ is the following:

$$\rho_K(x^M, y) = \partial_n P_K(z, y + iK)|_{z=x^M}, \quad \forall y \in (0, 1). \quad (5.3)$$

In particular, the joint density of (x^M, y^M) is given by (1.1).

The proof of Proposition 5.1 is split into three lemmas.

Lemma 5.2. *The pair (X_δ^M, Y_δ^M) converges weakly to a random pair of points (X^M, Y^M) as $\delta \rightarrow 0$. Moreover, the law of $x^M = f(X^M)$ is uniform on $(0, 1)$.*

Proof. From Theorem 1.5, the curve γ_δ^M converges weakly to γ^M as $\delta \rightarrow 0$. This implies that (X_δ^M, Y_δ^M) converges weakly to (X^M, Y^M) as $\delta \rightarrow 0$, where $X^M = \gamma^M \cap (ab)$ and $Y^M = \gamma^M \cap (cd)$. It remains to show that $f(X^M)$ is uniform in $(0, 1)$.

Recall from Lemma 4.4 that $u_\delta(z^*)$ is the probability that z^* lies to the right of η_δ^L for every $z^* \in \Omega_\delta^*$ and that u_δ converges to Ref locally uniformly due to Lemma 4.5. We denote by Ω^R the connected component of $\Omega \setminus \gamma^M$ which contains $[bc]$ on its boundary. It is same as the the connected component of $\Omega \setminus \eta^L$ which contains $[bc]$ on its boundary. For every $z \in \Omega$ and $z_{\delta_n}^* \in \Omega_{\delta_n}^*$ such that $z_{\delta_n}^* \rightarrow z$, we have

$$\begin{aligned} \{z \in \Omega^R\} &\subset \bigcup_{i=1}^{\infty} \bigcap_{n=i}^{\infty} \{z_{\delta_n}^* \text{ lies to the right of } \eta_{\delta_n}^L\} \\ &\subset \bigcap_{i=1}^{\infty} \bigcup_{n=i}^{\infty} \{z_{\delta_n}^* \text{ lies to the right of } \eta_{\delta_n}^L\} \subset \{z \in \overline{\Omega}^R\}. \end{aligned}$$

This implies that

$$\mathbb{P}[z \in \Omega^R] \leq \varliminf_{n \rightarrow \infty} u_{\delta_n}(z_{\delta_n}^*) = \text{Ref}(z) = \overline{\lim}_{n \rightarrow \infty} u_{\delta_n}(z_{\delta_n}^*) \leq \mathbb{P}[z \in \overline{\Omega}^R].$$

Note that $\mathbb{P}[z \in \Omega^R] = \mathbb{P}[z \in \overline{\Omega}^R]$ as $\mathbb{P}[z \in \gamma^M] = 0$ due to Corollary 4.10. Therefore,

$$\mathbb{P}[z \in \Omega^R] = \text{Ref}(z), \quad \forall z \in \Omega. \quad (5.4)$$

For every $\theta \in (ab)$, we choose $\{w_n\} \subset \Omega$ such that $w_n \rightarrow \theta$ as $n \rightarrow \infty$. Then, we have

$$\{X^M \in (a\theta)\} \subset \bigcup_{i=1}^{\infty} \bigcap_{n=i}^{\infty} \{w_n \in \Omega^R\} \subset \bigcap_{i=1}^{\infty} \bigcup_{n=i}^{\infty} \{w_n \in \Omega^R\} \subset \{X^M \in (a\theta)\}.$$

Since f is continuous on $\overline{\Omega}$, we have

$$\mathbb{P}[X^M \in (a\theta)] \leq \varliminf_{n \rightarrow \infty} \text{Ref}(w_n) = \text{Ref}(\theta) \leq \overline{\lim}_{n \rightarrow \infty} \text{Ref}(w_n) \leq \mathbb{P}[X^M \in (a\theta)].$$

Furthermore, for every $\tilde{\theta} \in (\theta b)$, we have

$$\text{Ref}(\theta) \leq \mathbb{P}[X^M \in (a\theta)] \leq \mathbb{P}[X^M \in (a\tilde{\theta})] \leq \text{Ref}(\tilde{\theta}).$$

By letting $\tilde{\theta} \rightarrow \theta$, we have

$$\mathbb{P}[X^M \in (a, \theta]] = \text{Ref}(\theta).$$

This implies that $\mathbb{P}[f(X^M) \in (0, \text{Ref}(\theta))] = \mathbb{P}[X^M \in (a, \theta]] = \text{Ref}(\theta)$ as desired. \square

Next, we will derive the conditional law of y^M given x^M . To this end, we analyze the conditional probability in discrete and use good control on discrete harmonic functions.




Figure 5.1: The function $\lambda = \lambda_{\tilde{c}, \tilde{d}}$ is the unique bounded harmonic function on $(0, 1) \times (0, iK)$ with the following boundary data: $\lambda = 0$ on $(f(a)f(b)) \cup (f(c)f(\tilde{c})) \cup (f(\tilde{d})f(d))$; $\lambda = 1$ on $(f(\tilde{c})f(\tilde{d}))$; and $\partial_n \lambda = 0$ on $(f(b)f(c)) \cup (f(d)f(a))$ where n is the outer normal vector.

Lemma 5.3. *Fix a polygon $(\Omega; a, \tilde{a}, \tilde{b}, b, c, \tilde{c}, \tilde{d}, d)$ with eight marked points. Suppose that a sequence of medial polygons $(\Omega_\delta; a_\delta^\circ, \tilde{a}_\delta^\circ, \tilde{b}_\delta^\circ, b_\delta^\circ, c_\delta^\circ, \tilde{c}_\delta^\circ, \tilde{d}_\delta^\circ, d_\delta^\circ)$ converges to $(\Omega; a, \tilde{a}, \tilde{b}, b, c, \tilde{c}, \tilde{d}, d)$ in the sense of (4.2). Denote by $\lambda_{\tilde{c}, \tilde{d}}$ the unique bounded harmonic function on $(0, 1) \times (0, iK)$ with the boundary data as shown in Figure 5.1. Denote by $\lambda_{c,d}$ when $\tilde{c} = c$ and $\tilde{d} = d$. Then, for every $\epsilon > 0$, there exist $\delta_0 > 0$ and $s > 0$ such that for all $\delta \leq \delta_0$ and $x_\delta \in (\tilde{a}_\delta \tilde{b}_\delta)$ and $x \in (\tilde{a} \tilde{b})$ with $d(x_\delta, x) \leq s$, we have*

$$\left| \mathbb{P} \left[Y_\delta^M \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M = x_\delta \right] - \frac{\partial_n \lambda_{\tilde{c}, \tilde{d}}(f(x))}{\partial_n \lambda_{c,d}(f(x))} \right| \leq \epsilon. \quad (5.5)$$

Proof. First, we show that $\partial_n \lambda_{c,d}(f(x)) > 0$ for all $x \in (ab)$. Let g be the bounded harmonic function on $(0, 1) \times (0, iK)$ with the following boundary data: $g = 0$ on $(f(a)f(b))$ and $g = 1$ on $(f(b)f(a))$. By maximum principle, we have $\lambda_{c,d}(y) \leq g(y)$ for every $y \in [0, 1] \times [0, iK]$. Thus, we have

$$\partial_n \lambda_{c,d}(f(x)) \geq \partial_n g(f(x)) > 0, \quad \text{for all } x \in (ab).$$

Next, we prove (5.5). For any $w_\delta \in \Omega_\delta$, denote by \mathbb{P}^{w_δ} the law of random walk in Ω_δ starting from w_δ . Define

$$\begin{aligned} u_\delta(w_\delta) &:= \mathbb{P}^{w_\delta} [\text{RW hits } (a_\delta b_\delta) \cup (c_\delta d_\delta) \text{ at } (c_\delta d_\delta)], \\ \tilde{u}_\delta(w_\delta) &:= \mathbb{P}^{w_\delta} [\text{RW hits } (a_\delta b_\delta) \cup (c_\delta d_\delta) \text{ at } (\tilde{c}_\delta \tilde{d}_\delta)]. \end{aligned}$$

Note that

$$\mathbb{P} \left[Y_\delta^M \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M = x_\delta \right] = \frac{\sum_{v_\delta \sim x_\delta} \mathbb{P}^{v_\delta} [\text{RW hits } (a_\delta b_\delta) \cup (c_\delta d_\delta) \text{ at } (\tilde{c}_\delta \tilde{d}_\delta)]}{\sum_{v_\delta \sim x_\delta} \mathbb{P}^{v_\delta} [\text{RW hits } (a_\delta b_\delta) \cup (c_\delta d_\delta) \text{ at } (c_\delta d_\delta)]} = \frac{\sum_{v_\delta \sim x_\delta} \tilde{u}_\delta(v_\delta)}{\sum_{v_\delta \sim x_\delta} u_\delta(v_\delta)}.$$

The function \tilde{u}_δ is a discrete harmonic function on $\Omega_\delta \setminus (a_\delta b_\delta) \cup (c_\delta d_\delta)$ with the following boundary data: $\tilde{u}_\delta = 0$ on $(a_\delta b_\delta) \cup (c_\delta \tilde{c}_\delta) \cup (\tilde{d}_\delta d_\delta)$ and $\tilde{u}_\delta = 1$ on $(\tilde{c}_\delta \tilde{d}_\delta)$. Similarly, u_δ is a discrete harmonic function on $\Omega_\delta \setminus (a_\delta b_\delta) \cup (c_\delta d_\delta)$ with the following boundary data: $u_\delta = 0$ on $(a_\delta b_\delta)$ and $u_\delta = 1$ on $(c_\delta d_\delta)$. By [CW19, Corollary 3.8], for every $\epsilon > 0$, there exists $s_1 > 0$ such that

$$1 - \epsilon \leq \frac{\tilde{u}_\delta(v_\delta)}{u_\delta(v_\delta)} \times \frac{u_\delta(y_\delta)}{\tilde{u}_\delta(y_\delta)} \leq 1 + \epsilon, \quad \text{for all } y_\delta \in \Omega_\delta \cap B(x_\delta, s_1) \setminus \partial\Omega_\delta \text{ and all } x_\delta \in (\tilde{a}_\delta \tilde{b}_\delta).$$

This implies that

$$1 - \epsilon \leq \mathbb{P} \left[Y_\delta^M \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M = x_\delta \right] \times \frac{u_\delta(y_\delta)}{\tilde{u}_\delta(y_\delta)} \leq 1 + \epsilon.$$

We choose $s < \frac{s_1}{4}$. Since $\partial\Omega$ is locally connected (which means it is a curve), we can choose a simple curve L such that $\frac{s}{4} < d(L, (\tilde{a}\tilde{b})) < \frac{s}{2}$. Note that there exists $\delta_1 > 0$, for all $\delta < \delta_1$, we can choose a discrete simple curve $L_\delta \subset \Omega_\delta$ with $\frac{s}{4} < d(L_\delta, (\tilde{a}_\delta\tilde{b}_\delta)) < \frac{s}{2}$ such that $L_\delta \rightarrow L$ as curves. By the same argument as in the proof of Lemma B.1, we have that $\tilde{u}_\delta \rightarrow \lambda_{\tilde{c},\tilde{d}} \circ f$ and $u_\delta \rightarrow \lambda_{c,d} \circ f$ locally uniformly in Ω . Then, there exists $\delta_2 > 0$, if $\delta < \delta_2$ and $d(x_\delta, x) < s$, for every $y_\delta \in B(x_\delta, s) \cap L_\delta$, there exists $y \in B(x, s) \cap L$ such that

$$\left| \frac{\tilde{u}_\delta(y_\delta)}{u_\delta(y_\delta)} - \frac{\lambda_{\tilde{c},\tilde{d}}(f(y))}{\lambda_{c,d}(f(y))} \right| < \epsilon.$$

Since f is continuous on $\bar{\Omega}$, we have $\text{diam}(f(B(x, s))) \rightarrow 0$ as $s \rightarrow 0$. By Taylor expansion, we can choose s small enough such that

$$\left| \frac{\lambda_{\tilde{c},\tilde{d}}(f(y))}{\lambda_{c,d}(f(y))} - \frac{\partial_n \lambda_{\tilde{c},\tilde{d}}(f(x))}{\partial_n \lambda_{c,d}(f(x))} \right| \leq \epsilon, \quad \text{for all } x \in (\tilde{a}\tilde{b}) \text{ and } y \in B(x, s) \cap L.$$

This implies that, if $\delta < \delta_1 \wedge \delta_2$ and $d(x_\delta, x) < s$, we have

$$(1 - 3\epsilon) \frac{\partial_n \lambda_{\tilde{c},\tilde{d}}(f(x))}{\partial_n \lambda_{c,d}(f(x))} \leq \mathbb{P} \left[Y_\delta^M \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M = x_\delta \right] \leq (1 + 3\epsilon) \frac{\partial_n \lambda_{\tilde{c},\tilde{d}}(f(x))}{\partial_n \lambda_{c,d}(f(x))}.$$

This completes the proof. \square

Lemma 5.4. *The conditional law of Y^M given X^M is given by*

$$\mathbb{P}[Y^M \in (\tilde{c}\tilde{d}) \mid X^M] = \frac{\partial_n \lambda_{\tilde{c},\tilde{d}}(f(X^M))}{\partial_n \lambda_{c,d}(f(X^M))}. \quad (5.6)$$

Proof. By the conformal invariance, we may assume $\Omega = (0, 1) \times (0, iK)$. We couple (X_δ^M, Y_δ^M) and (X^M, Y^M) together such that $X_\delta^M \rightarrow X^M$ and $Y_\delta^M \rightarrow Y^M$ almost surely. Fix a polygon $(\Omega; a, \tilde{a}, \tilde{b}, b, c, \tilde{c}, \tilde{d}, d)$ with eight marked points. Suppose that a sequence of medial polygons $(\Omega_\delta^\circ; a_\delta^\circ, \tilde{a}_\delta^\circ, \tilde{b}_\delta^\circ, b_\delta^\circ, c_\delta^\circ, \tilde{c}_\delta^\circ, \tilde{d}_\delta^\circ, d_\delta^\circ)$ converges to $(\Omega; a, \tilde{a}, \tilde{b}, b, c, \tilde{c}, \tilde{d}, d)$ in the sense of (4.2). For any $\delta_n \rightarrow 0$, we have

$$\begin{aligned} \{X^M \in (\tilde{a}\tilde{b}), Y^M \in (\tilde{c}\tilde{d})\} &\subset \bigcup_{j=1}^\infty \bigcap_{n=j}^\infty \{X_{\delta_n}^M \in (\tilde{a}_{\delta_n} \tilde{b}_{\delta_n}), Y_{\delta_n}^M \in (\tilde{c}_{\delta_n} \tilde{d}_{\delta_n})\} \\ &\subset \bigcap_{j=1}^\infty \bigcup_{n=j}^\infty \{X_{\delta_n}^M \in (\tilde{a}_{\delta_n} \tilde{b}_{\delta_n}), Y_{\delta_n}^M \in (\tilde{c}_{\delta_n} \tilde{d}_{\delta_n})\} \subset \{X^M \in [\tilde{a}\tilde{b}], Y^M \in [\tilde{c}\tilde{d}]\}. \end{aligned}$$

Thus,

$$\begin{aligned} \mathbb{P}[X^M \in (\tilde{a}\tilde{b}), Y^M \in (\tilde{c}\tilde{d})] &\leq \varliminf_{n \rightarrow \infty} \mathbb{P}[X_{\delta_n}^M \in (\tilde{a}_{\delta_n} \tilde{b}_{\delta_n}), Y_{\delta_n}^M \in (\tilde{c}_{\delta_n} \tilde{d}_{\delta_n})] \\ &\leq \varlimsup_{n \rightarrow \infty} \mathbb{P}[X_{\delta_n}^M \in (\tilde{a}_{\delta_n} \tilde{b}_{\delta_n}), Y_{\delta_n}^M \in (\tilde{c}_{\delta_n} \tilde{d}_{\delta_n})] \leq \mathbb{P}[X^M \in [\tilde{a}\tilde{b}], Y^M \in [\tilde{c}\tilde{d}]]. \end{aligned} \quad (5.7)$$

For every $\epsilon > 0$, we choose s the same as in Lemma 5.3. Divide $(\tilde{a}\tilde{b})$ into $\bigcup_{j=0}^m [x^j x^{j+1}]$ with $x^0 = \tilde{a}$ and $x^{m+1} = \tilde{b}$ such that the length of $[x^j x^{j+1}]$ is less than s . Denote by $\{x_{\delta_n}^0 = \tilde{a}_{\delta_n}, x_{\delta_n}^1, \dots, x_{\delta_n}^{m+1} = \tilde{b}_{\delta_n}\} \subset (\tilde{a}_{\delta_n} \tilde{b}_{\delta_n})$ the discrete approximation. By Lemma 5.3, for n large enough, we have

$$\left| \mathbb{P}[X_{\delta_n}^M \in (\tilde{a}_{\delta_n} \tilde{b}_{\delta_n}), Y_{\delta_n}^M \in (\tilde{c}_{\delta_n} \tilde{d}_{\delta_n})] - \sum_{j=0}^m \frac{\partial_n \lambda_{\tilde{c},\tilde{d}}(x^j)}{\partial_n \lambda_{c,d}(x^j)} \mathbb{P}[X_{\delta_n}^M \in [x_{\delta_n}^j x_{\delta_n}^{j+1}]] \right| \leq \epsilon.$$

By Lemma 5.2, the law of X^M is uniform on (ab) . This implies that

$$\lim_{n \rightarrow \infty} \mathbb{P}[X_{\delta_n}^M \in [x_{\delta_n}^j x_{\delta_n}^{j+1}]] = \mathbb{P}[X^M \in [x^j x^{j+1}]], \quad \text{for } 0 \leq j \leq m.$$

Thus, we have

$$\overline{\lim}_{n \rightarrow \infty} \left| \mathbb{P}[X_{\delta_n}^M \in (\tilde{a}_{\delta_n}, \tilde{b}_{\delta_n}), Y_{\delta_n}^M \in (\tilde{c}_{\delta_n}, \tilde{d}_{\delta_n})] - \sum_{j=0}^m \frac{\partial_n \lambda_{\tilde{c}, \tilde{d}}(x^j)}{\partial_n \lambda_{c, d}(x^j)} \mathbb{P}[X^M \in [x^j, x^{j+1}]] \right| \leq \epsilon.$$

By letting $s \rightarrow 0$ ($m \rightarrow \infty$) and $\epsilon \rightarrow 0$, we have

$$\lim_{n \rightarrow \infty} \mathbb{P}[X_{\delta_n}^M \in (\tilde{a}_{\delta_n}, \tilde{b}_{\delta_n}), Y_{\delta_n}^M \in (\tilde{c}_{\delta_n}, \tilde{d}_{\delta_n})] = \int_{\tilde{a}}^{\tilde{b}} \frac{\partial_n \lambda_{\tilde{c}, \tilde{d}}(x)}{\partial_n \lambda_{c, d}(x)} dx.$$

Plugging into (5.7), we have

$$\mathbb{P}[X^M \in (\tilde{a}\tilde{b}), Y^M \in (\tilde{c}\tilde{d})] \leq \int_{\tilde{a}}^{\tilde{b}} \frac{\partial_n \lambda_{\tilde{c}, \tilde{d}}(x)}{\partial_n \lambda_{c, d}(x)} dx \leq \mathbb{P}[X^M \in [\tilde{a}\tilde{b}], Y^M \in [\tilde{c}\tilde{d}]].$$

Note that the marginal law of X^M is uniform on (ab) and the marginal law of Y^M is uniform on (cd) , we have

$$\mathbb{P}[X^M \in (\tilde{a}\tilde{b}), Y^M \in (\tilde{c}\tilde{d})] = \mathbb{P}[X^M \in [\tilde{a}\tilde{b}], Y^M \in [\tilde{c}\tilde{d}]].$$

Therefore,

$$\mathbb{P}[X^M \in (\tilde{a}\tilde{b}), Y^M \in (\tilde{c}\tilde{d})] = \int_{\tilde{a}}^{\tilde{b}} \frac{\partial_n \lambda_{\tilde{c}, \tilde{d}}(x)}{\partial_n \lambda_{c, d}(x)} dx.$$

This gives (5.6) and completes the proof. \square

Proof of Proposition 5.1. The convergence of $(X_{\delta}^M, Y_{\delta}^M)$ and the law of $x^M = f(X^M)$ is derived Lemma 5.2. The conditional law of Y^M given X^M is derived in Lemma 5.4. We only need to explain that (5.6) is equivalent to (5.3). Consider the following two functions:

$$\lambda_{\tilde{c}, \tilde{d}}(\cdot) \quad \text{and} \quad \int_{f(\tilde{c})}^{f(\tilde{d})} P_K(\cdot, r) dr.$$

Both of them are harmonic functions on $(0, 1) \times (0, iK)$. Their boundary data is the same except at $(f(\tilde{c}), f(\tilde{d}))$ along which both of them are constant. Both of them are bounded on $[0, 1] \times [0, iK]$. Thus, they are the same up to a multiplicative constant. This explains the equivalence between (5.6) and (5.3).

Let us calculate the outer normal derivative of P_K : for $x, y \in (0, 1)$,

$$\partial_n P_K(z, y + iK)|_{z=x} = \frac{\pi}{4K} \sum_{n \in \mathbb{Z}} \left(\frac{1}{\cosh^2\left(\frac{\pi}{2K}(x - y - 2n)\right)} + \frac{1}{\cosh^2\left(\frac{\pi}{2K}(x + y - 2n)\right)} \right), \quad (5.8)$$

where the right-hand side is the same as $\rho_K(x, y)$ defined in (1.1). From here, we have

$$\begin{aligned} & \int_0^1 (\partial_n P_K(z, y + iK)|_{z=x}) dy \\ &= \frac{\pi}{4K} \sum_{n \in \mathbb{Z}} \left(\int_0^1 \frac{dy}{\cosh^2\left(\frac{\pi}{2K}(x - y - 2n)\right)} + \int_0^1 \frac{dy}{\cosh^2\left(\frac{\pi}{2K}(x + y - 2n)\right)} \right) \\ &= \frac{\pi}{4K} \sum_{n \in \mathbb{Z}} \left(\int_{2n}^{2n+1} \frac{dr}{\cosh^2\left(\frac{\pi}{2K}(x - r)\right)} + \int_{2n-1}^{2n} \frac{dr}{\cosh^2\left(\frac{\pi}{2K}(x - r)\right)} \right) \\ &= \frac{\pi}{4K} \int_{-\infty}^{+\infty} \frac{dr}{\cosh^2\left(\frac{\pi}{2K}(x - r)\right)} = \frac{1}{2} \int_{-\infty}^{+\infty} \frac{dr}{\cosh^2(r)} = 1. \end{aligned}$$

Therefore,

$$\frac{\partial_n \lambda_{\tilde{c}, \tilde{d}}(f(X^M))}{\partial_n \lambda_{c, d}(f(X^M))} = \int_{f(\tilde{c})}^{f(\tilde{d})} \rho_K(x^M, \operatorname{Re} y) dy. \quad (5.9)$$

This gives the density in (1.1) and completes the proof. \square

Corollary 5.5. *Fix a polygon $(\Omega; a, x, b, c, \tilde{c}, \tilde{d}, d)$ with seven marked points. Suppose that a sequence of medial polygons $(\Omega_\delta^\circ; a_\delta^\circ, x_\delta^\circ, b_\delta^\circ, c_\delta^\circ, \tilde{c}_\delta^\circ, \tilde{d}_\delta^\circ, d_\delta^\circ)$ converges to $(\Omega; a, x, b, c, \tilde{c}, \tilde{d}, d)$ in the sense of (4.2). Denote by $f = f_{(\Omega; a, b, c, d)}$ the conformal map from Ω onto $(0, 1) \times (0, iK)$ which sends (a, b, c, d) to $(0, 1, 1 + iK, iK)$ and extend its definition continuously to the boundary. Then, we have*

$$\lim_{\delta \rightarrow 0} \mathbb{P} \left[Y_\delta^M \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M = x_\delta \right] = \int_{f(\tilde{c})}^{f(\tilde{d})} \rho_K(f(x), \operatorname{Re} y) dy.$$

Proof. First let $\delta \rightarrow 0$ and then let $\epsilon \rightarrow 0$ in (5.5), combining with (5.9), we obtain the conclusion. \square

We emphasize that the assumption on $\partial\Omega$ in Corollary 5.5 is locally connected, and we do not require extra regularity.

5.2 Proof of Theorem 1.6

The joint law of (X^M, Y^M) in Theorem 1.6 is given in Proposition 5.1, and to complete the proof of Theorem 1.6, it remains to show that the conditional law of γ^M given X^M is $\operatorname{SLE}_2(-1, -1; -1, -1)$. We follow the strategy in [Zha08c] and transform the notations into an amenable way. We fix the following notation in this section.

- Fix $d = -\infty < a < b < c$. Denote by K the conformal modulus of the quad $(\mathbb{H}; a, b, c, \infty)$ and by $f(\cdot; a, b, c)$ the conformal map from \mathbb{H} onto $(0, 1) \times (0, iK)$ sending (a, b, c, ∞) to $(0, 1, 1 + iK, iK)$.
- Define

$$P(z; a, w, b, c) := P_K(f(z; a, b, c), f(w; a, b, c)), \quad \forall z \in \mathbb{H}, \quad (5.10)$$

where P_K is given in (5.1). Note that $P(\cdot; \cdot, \cdot, \cdot, \cdot)$ is smooth on $\mathbb{H} \times \{(a, w, b, c) \in \mathbb{R}^4 : a < w < b < c\}$. Fix $a < w < b < c$, it is the Poisson kernel on \mathbb{H} with the boundary data:

$$P(\cdot; a, w, b, c) = 0, \text{ on } (a, w) \cup (w, b) \cup (c, \infty); \quad \partial_n P(\cdot; a, w, b, c) = 0, \text{ on } (-\infty, a) \cup (b, c); \quad (5.11)$$

and the normalization:

$$\int_c^\infty (\partial_n P(z; a, w, b, c)|_{z=x}) dx = 1. \quad (5.12)$$

First of all, we show that the Poisson kernel satisfies a certain PDE.

Lemma 5.6. *For $a < w < b < c$ and $z \in \mathbb{H}$, consider the function $P(z; a, w, b, c)$ in (5.10). Denote by ∂_x the partial derivative with respect to the real part of the first (complex) variable and by ∂_y the partial derivative with respect to the imaginary part of the first (complex) variable. Define*

$$\mathcal{D} := \frac{2}{a-w} \partial_a + \frac{2}{b-w} \partial_b + \frac{2}{c-w} \partial_c + 2 \frac{f''(w; a, b, c)}{f'(w; a, b, c)} \partial_w + \partial_w^2 + \operatorname{Re} \left(\frac{2}{z-w} \right) \partial_x + \operatorname{Im} \left(\frac{2}{z-w} \right) \partial_y.$$

Then, we have $\mathcal{D}P(z; a, w, b, c) = 0$.

Before the proof of Lemma 5.6, we first show that the Poisson kernel (5.10) gives a martingale observable for $\operatorname{SLE}_2(-1, -1; -1, -1)$ in the following lemma. In fact, this lemma is not needed in this article, but it explains the origin of the PDE in Lemma 5.6, thus we still keep it.

Lemma 5.7. *Suppose $d = -\infty < a < w < b < c$. Let $\gamma \sim \text{SLE}_2(-1; -1, -1)$ in \mathbb{H} from w to ∞ with force points $(a; b, c)$. Denote by $(W_t, t \geq 0)$ be the driving function and by $(g_t, t \geq 0)$ the corresponding conformal maps. Denote by $(\mathcal{F}_t, t \geq 0)$ the filtration generated by γ . Fix $z \in \mathbb{H}$, the process*

$$(P(g_t(z); g_t(a), W_t, g_t(b), g_t(c)), t \geq 0)$$

is a local martingale with respect to the filtration $(\mathcal{F}_t, t \geq 0)$.

Proof. From Itô's formula, we have

$$\begin{aligned} & dP(g_t(z); g_t(a), W_t, g_t(b), g_t(c)) \\ &= \mathcal{D}P(g_t(z); g_t(a), W_t, g_t(b), g_t(c))dt + \partial_w P(g_t(z); g_t(a), W_t, g_t(b), g_t(c)) \left(dW_t - 2 \frac{f''(W_t; g_t(a), g_t(b), g_t(c))}{f'(W_t; g_t(a), g_t(b), g_t(c))} dt \right) \\ &= \partial_w P(g_t(z); g_t(a), W_t, g_t(b), g_t(c)) \left(dW_t - 2 \frac{f''(W_t; g_t(a), g_t(b), g_t(c))}{f'(W_t; g_t(a), g_t(b), g_t(c))} dt \right). \end{aligned} \quad (\text{by Lemma 5.6})$$

From (4.17), we have

$$f(z; a, b, c) = \mathcal{K} \left(\arcsin \sqrt{\frac{z-a}{b-a}, \frac{b-a}{c-a}} \right) / \mathcal{K} \left(\frac{b-a}{c-a} \right).$$

From (A.8), we have

$$2 \frac{f''(w; a, b, c)}{f'(w; a, b, c)} = \frac{1}{a-w} + \frac{1}{b-w} + \frac{1}{c-w}. \quad (5.13)$$

Therefore,

$$dW_t = \sqrt{2}dB_t + 2 \frac{f''(W_t; g_t(a), g_t(b), g_t(c))}{f'(W_t; g_t(a), g_t(b), g_t(c))} dt,$$

and

$$dP(g_t(z); g_t(a), W_t, g_t(b), g_t(c)) = \partial_w P(g_t(z); g_t(a), W_t, g_t(b), g_t(c)) \sqrt{2}dB_t.$$

This completes the proof. \square

Proof of Lemma 5.6. When there is no ambiguity, we write $\partial_n P(z; a, w, b, c)|_{z=x}$ as $\partial_n P(x; a, w, b, c)$. Define

$$\mathcal{V}(z) := \mathcal{D}P(z; a, w, b, c). \quad (5.14)$$

First, we show that $\mathcal{V}(\cdot)$ has the same boundary data as $P(\cdot; a, w, b, c)$. From (5.11), we have $P(z; a, w, b, c) = 0$ for all $a < z \neq w < b < c$. Thus

$$\ell P(z; a, w, b, c) = 0, \quad \forall a < z \neq w < b < c, \quad \text{for all } \ell = \partial_a, \partial_b, \partial_c, \partial_w, \partial_w^2, \partial_x.$$

Therefore, $\mathcal{V}(\cdot) = 0$ on $(a, w) \cup (w, b)$. Similarly, since $P(z; a, w, b, c) = 0$ for all $a < w < b < c < z$, we have $\mathcal{V}(\cdot) = 0$ on $(c, +\infty)$. Since $\partial_n P(x; a, w, b, c) = 0$ for all $a < w < b < x < c$ or $x < a < w < b < c$, we have

$$\begin{aligned} & \partial_n \ell P(x; a, w, b, c) = \ell \partial_n P(x; a, w, b, c) = 0, \quad \text{for all } \ell = \partial_a, \partial_b, \partial_c, \partial_w, \partial_w^2, \\ & \partial_n \left(\text{Re} \frac{2}{z-w} \partial_x P(z; a, w, b, c) \right) \Big|_{z=x} = \frac{2}{x-w} \partial_x \partial_n P(x; a, w, b, c) = 0, \\ & \partial_n \left(\text{Im} \frac{2}{z-w} \partial_y P(z; a, w, b, c) \right) \Big|_{z=x} = \frac{-2}{(x-w)^2} \partial_n P(x; a, w, b, c) = 0, \\ & \quad \forall a < w < b < x < c \text{ or } x < a < w < b < c. \end{aligned}$$

Here the interchange of ∂_n and $\partial_a, \partial_b, \partial_c, \partial_w, \partial_x$ is legal due to the explicit form of $P(z; a, w, b, c)$ in (5.10). Thus, $\partial_n \mathcal{V}(\cdot) = 0$ on $(b, c) \cup (-\infty, a)$.

Second, we show that $\mathcal{V}(\cdot)$ is harmonic in \mathbb{H} . Note that by the explicit form of $P(z; a, w, b, c)$ in (5.10), there exists a smooth function $G(\cdot; \cdot, \cdot, \cdot, \cdot)$ on $\overline{\mathbb{H}} \times \{(a, w, b, c) \in \mathbb{R}^4 : a < w < b < c\}$ which is analytic on $\overline{\mathbb{H}} \setminus \{a, b, c\}$ when $a < w < b < c$ is fixed, such that

$$P(z; a, w, b, c) = \operatorname{Im} \left(G(z; a, w, b, c) + \frac{K}{\pi f'(w; a, b, c)} \frac{1}{z - w} \right), \quad \forall z \in \mathbb{H}, w \in (a, b).$$

Then, we calculate

$$\begin{aligned} & \left(\operatorname{Re} \left(\frac{2}{z - w} \right) \partial_x + \operatorname{Im} \left(\frac{2}{z - w} \right) \partial_y \right) P(z; a, w, b, c) \\ &= \operatorname{Im} \left(\frac{2G'(z; a, w, b, c)}{z - w} - \frac{2K}{\pi f'(w; a, b, c)(z - w)^3} \right) \\ & \quad \partial_w P(z; a, w, b, c) \\ &= \operatorname{Im} \left(\partial_w G(z; a, w, b, c) + \frac{K}{\pi f'(w; a, b, c)} \frac{1}{(z - w)^2} - \frac{K}{\pi} \frac{f''(w; a, b, c)}{f'(w; a, b, c)^2} \frac{1}{z - w} \right) \\ & \quad \partial_w^2 P(z; a, w, b, c) \\ &= \operatorname{Im} \left(\partial_w^2 G(z; a, w, b, c) - \frac{K}{\pi} \frac{f'''(w; a, b, c)f'(w; a, b, c) - 2f''(w; a, b, c)^2}{f'(w; a, b, c)^3} \frac{1}{z - w} \right) \\ & \quad + \operatorname{Im} \left(\frac{2K}{\pi f'(w; a, b, c)} \frac{1}{(z - w)^3} - \frac{2K}{\pi} \frac{f''(w; a, b, c)}{f'(w; a, b, c)^2} \frac{1}{(z - w)^2} \right). \end{aligned}$$

Therefore, we have

$$\mathcal{V}(z) = \operatorname{Im} \left(G_1(z) + \frac{G_2(z)}{z - w} \right),$$

where G_1 and G_2 are analytic functions on $\overline{\mathbb{H}} \setminus \{a, b, c\}$. This implies that \mathcal{V} is harmonic.

Finally, we prove that $\mathcal{V}(z) = 0$ for every $z \in \mathbb{H}$. To this end, we evaluate the value of $\int_c^\infty \partial_n \mathcal{V}(x) dx$. On the one hand, consider the following function:

$$\mathcal{V}(\cdot) - \frac{\pi}{K} G_2(w) f'(w; a, b, c) P(\cdot; a, w, b, c).$$

It is harmonic on \mathbb{H} , and it has the same boundary data as $P(\cdot; a, w, b, c)$. Moreover, it is bounded near w from the construction and it is bounded near a, b, c, ∞ due to Lemma 5.8. Thus, it is bounded in $\overline{\mathbb{H}}$. Therefore, it has to vanish. This implies that

$$\int_c^{+\infty} \partial_n \mathcal{V}(x) dx = \frac{\pi G_2(w) f'(w; a, b, c)}{K} \int_c^{+\infty} \partial_n P(x; a, w, b, c) dx = \frac{\pi}{K} G_2(w) f'(w; a, b, c), \quad (5.15)$$

where the second equal sign is due to (5.12). On the other hand, we have

$$\begin{aligned} \int_c^\infty \partial_n \ell P(x; a, w, b, c) dx &= \ell \int_c^\infty \partial_n P(x; a, w, b, c) dx = 0, \quad \text{for } \ell = \partial_a, \partial_b, \partial_w, \partial_w^2; & \text{(by (5.12))} \\ \int_c^\infty \partial_n \frac{2}{c - w} \partial_c P(x; a, w, b, c) dx &= \frac{2}{c - w} \int_c^\infty \partial_c \partial_n P(x; a, w, b, c) dx = \frac{2}{c - w} \partial_n P(c; a, w, b, c); & \text{(by (5.12))} \end{aligned}$$

$$\begin{aligned} & \int_c^\infty \left(\partial_n \left(\operatorname{Re} \frac{2}{z - w} \partial_x P(z; a, w, b, c) \right) \right) \Big|_{z=x} dx \\ &= \int_c^\infty \frac{2}{x - w} \partial_x \partial_n P(x; a, w, b, c) dx = \frac{-2}{c - w} \partial_n P(c; a, w, b, c) + \int_c^\infty \frac{2}{(x - w)^2} \partial_n P(x; a, w, b, c) dx; \\ & \int_c^\infty \left(\partial_n \left(\operatorname{Im} \frac{2}{z - w} \partial_y P(z; a, w, b, c) \right) \right) \Big|_{z=x} dx = \int_c^\infty \frac{-2}{(x - w)^2} \partial_n P(x; a, w, b, c) dx. \end{aligned}$$

Therefore,

$$\int_c^{+\infty} \partial_n \mathcal{V}(x) dx = \int_c^{+\infty} \partial_n \mathcal{D}P(z; a, w, b, c) \Big|_{z=x} dx = 0.$$

Comparing with (5.15), we have $G_2(w) = 0$ and $\mathcal{V} = 0$. \square

Lemma 5.8. *The function \mathcal{V} in (5.14) is bounded near a, b, c, ∞ .*

Proof. We first investigate its behavior around a . From (4.17), we have

$$f(z; a, b, c) = \mathcal{K} \left(\arcsin \sqrt{\frac{z-a}{b-a}, \frac{b-a}{c-a}} \right) / \mathcal{K} \left(\frac{b-a}{c-a} \right),$$

and

$$K = K(a, b, c) = \text{Im} \mathcal{K} \left(\arcsin \sqrt{\frac{c-a}{b-a}, \frac{b-a}{c-a}} \right) / \mathcal{K} \left(\frac{b-a}{c-a} \right).$$

Note that $f(\cdot; \cdot, \cdot, \cdot)$ is smooth on $\overline{\mathbb{H}} \setminus \{a, b, c\} \times \{(a, b, c) \in \mathbb{R}^3 : a < b < c\}$ and $K(\cdot, \cdot, \cdot)$ is smooth on $\{(a, b, c) \in \mathbb{R}^3 : a < b < c\}$. This implies that $\partial_w P$ and $\partial_w^2 P$ are continuous at a . Moreover, for all $\ell \in \{\partial_a, \partial_b, \partial_c, \partial_x, \partial_y\}$, we have

$$\begin{aligned} \ell P(z; a, w, b, c) = & \text{Im} \sum_{n \in \mathbb{Z}} \frac{1}{\sinh^2 \left(\frac{\pi}{2K} (2n - f(w; a, b, c) + f(z; a, b, c)) \right)} \ell \left(\frac{\pi}{4K} (f(z; a, b, c) - f(w; a, b, c)) \right) \\ & - \text{Im} \sum_{n \in \mathbb{Z}} \frac{1}{\sinh^2 \left(\frac{\pi}{2K} (2n - f(w; a, b, c) - \bar{f}(z; a, b, c)) \right)} \ell \left(\frac{\pi}{4K} (\bar{f}(z; a, b, c) + f(w; a, b, c)) \right). \end{aligned}$$

Denote by $\tilde{z} := \arcsin \sqrt{(z-a)/(b-a)}$ and $s := (b-a)/(c-a)$. We have

$$\begin{aligned} \partial_b f(z; a, b, c) &= -\frac{\sqrt{(c-a)(z-a)}}{2(b-a)\mathcal{K}(s)\sqrt{(c-z)(b-z)}} + \frac{\partial_x \mathcal{K}(\tilde{z}, s)}{(c-a)\mathcal{K}(s)} - \frac{\mathcal{K}(\tilde{z}, s)\mathcal{K}'(s)}{(c-a)\mathcal{K}^2(s)}, \\ \partial_c f(z; a, b, c) &= -\frac{(b-a)\partial_x \mathcal{K}(\tilde{z}, s)}{(c-a)^2\mathcal{K}(s)} + \frac{(b-a)\mathcal{K}(\tilde{z}, s)\mathcal{K}'(s)}{(c-a)^2\mathcal{K}^2(s)}. \end{aligned}$$

This implies that

$$\partial_b P(z; a, w, b, c) \rightarrow 0, \quad \partial_c P(z; a, w, b, c) \rightarrow 0, \quad \text{as } z \rightarrow a. \quad (5.16)$$

We have

$$\begin{aligned} \left| \partial_y P(z; a, w, b, c) \text{Im} \left(\frac{2}{z-w} \right) \right| &= \left| \frac{\sqrt{c-a}}{2\sqrt{(b-z)(c-z)(z-a)}\mathcal{K}(s)} \text{Im} \frac{2}{z-w} \right| \\ &= \frac{\sqrt{c-a} \times \text{Im} z}{|\sqrt{(b-z)(c-z)(z-a)}\mathcal{K}(s)| |z-w|^2}. \end{aligned}$$

Thus,

$$\partial_y P(z; a, w, b, c) \text{Im} \left(\frac{2}{z-w} \right) \rightarrow 0, \quad \text{as } z \rightarrow a. \quad (5.17)$$

We have

$$\begin{aligned} & \left(\frac{2}{a-w} \partial_a + \text{Re} \left(\frac{2}{z-w} \right) \partial_x \right) f(z; a, w, b, c) \\ &= \frac{\sqrt{c-a}}{\sqrt{(c-z)(b-z)(z-a)}\mathcal{K}(s)} \left(\frac{z-a}{(b-a)(a-w)} - \text{Re} \frac{z-a}{(z-w)(a-w)} \right) \\ & \quad + \frac{(b-c)\partial_x \mathcal{K}(\tilde{z}, s)}{(c-a)^2\mathcal{K}(s)} - \frac{(b-c)\mathcal{K}(\tilde{z}, s)\mathcal{K}'(s)}{(c-a)^2\mathcal{K}^2(s)}. \end{aligned}$$

Thus,

$$\left(\frac{2}{a-w} \partial_a + \operatorname{Re} \left(\frac{2}{z-w} \partial_x \right) \right) P(z; a, w, b, c) \rightarrow 0, \quad \text{as } z \rightarrow a. \quad (5.18)$$

Recall that $\partial_w P$ and $\partial_w^2 P$ are continuous at a , combining with (5.16), (5.17) and (5.18), we see that $\mathcal{V}(z)$ remains bounded as $z \rightarrow a$. We may show that it is also bounded near b, c, ∞ similarly. \square

Corollary 5.9. *For $a < w < b < c < x$, define*

$$F(x; a, w, b, c) = \partial_n P(z; a, w, b, c)|_{z=x}. \quad (5.19)$$

Then we have

$$\left(\frac{2}{a-w} \partial_a + \frac{2}{b-w} \partial_b + \frac{2}{c-w} \partial_c + 2 \frac{f''(w; a, b, c)}{f'(w; a, b, c)} \partial_w + \partial_w^2 + \frac{2}{x-w} \partial_x + \frac{-2}{(x-w)^2} \right) F = 0. \quad (5.20)$$

Proof. The PDE (5.20) can be obtained by taking ∂_n in $\mathcal{D}P = 0$ from Lemma 5.6. \square

Now, we come back to derive the conditional law of γ^M given X^M in Theorem 1.6. We will argue that the conditional density in (5.3) gives a martingale observable for γ^M .

Lemma 5.10. *Assume the same setup as in Theorem 1.6. Choose a conformal map ϕ from Ω onto \mathbb{H} such that $\phi(d) = \infty$ and $\phi(a) < \phi(b) < \phi(c)$. Denote by $(W_t, t \geq 0)$ the driving function of $\phi(\gamma^M)$ and by $(g_t, t \geq 0)$ the corresponding conformal maps. For any $x \in (\phi(c), +\infty)$, then the process*

$$(g'_t(x)F(g_t(x); g_t(\phi(a)), W_t, g_t(\phi(b)), g_t(\phi(c))), t \geq 0)$$

is a martingale up to the first time that γ^M hits (cd) where F is defined in (5.19).

Proof. Fix two boundary points \tilde{c}, \tilde{d} such that $a, b, c, \tilde{c}, \tilde{d}, d$ are in counterclockwise order. Choose a sequence of medial polygons $(\Omega_\delta^\circ; a_\delta^\circ, b_\delta^\circ, c_\delta^\circ, \tilde{c}_\delta^\circ, \tilde{d}_\delta^\circ, d_\delta^\circ)$ converges to $(\Omega; a, b, c, \tilde{c}, \tilde{d}, d)$ in the sense of (4.2) and choose a sequence of conformal maps $\phi_\delta : \Omega_\delta \rightarrow \mathbb{H}$ with $\phi_\delta(d_\delta) = \infty$ such that ϕ_δ^{-1} converges to ϕ^{-1} uniformly on $\overline{\mathbb{H}}$ as $\delta \rightarrow 0$. By Theorem 1.5, we have that $\gamma_\delta^M \rightarrow \gamma^M$ in law as $\delta \rightarrow 0$. Couple $\{\gamma_\delta^M\}_{\delta>0}$ and γ^M together such that $\gamma_\delta^M \rightarrow \gamma^M$ almost surely as $\delta \rightarrow 0$. Recall that $X^M = \gamma^M \cap (ab)$, $Y^M = \gamma^M \cap (cd)$ and $X_\delta^M = \gamma_\delta^M \cap (a_\delta b_\delta)$, $Y_\delta^M = \gamma_\delta^M \cap (c_\delta d_\delta)$.

We parameterize $\phi(\gamma^M)$ by the half-plane capacity and parameterize γ^M such that $\phi(\gamma^M(t)) = \phi(\gamma^M(t))$. Denote by T the first time that γ^M hits (cd) . For $\epsilon > 0$, define $T_\epsilon = \inf\{t : d(\gamma^M(t), (ba)) = \epsilon\}$. For $t < T$, denote by K_t the conformal modulus of the quad $(\mathbb{H}; g_t(\phi(a)), g_t(\phi(b)), g_t(\phi(c)), \infty)$ and by f_t the conformal map from \mathbb{H} onto $(0, 1) \times (0, iK_t)$ sending $(g_t(\phi(a)), g_t(\phi(b)), g_t(\phi(c)), \infty)$ to $(0, 1, 1 + iK_t, iK_t)$.

We parameterize γ_δ^M similarly, define $T_\epsilon^\delta = \inf\{t : d(\gamma_\delta^M, (b_\delta a_\delta)) = \epsilon\}$. We may assume $T_\epsilon^\delta \rightarrow T_\epsilon$ almost surely as $\delta \rightarrow 0$ by considering the continuous modification, see details in [Kar19] and [Kar20]. For every $t < T_\epsilon^\delta$, define $\Omega_\delta(t) := \Omega_\delta \setminus \gamma_\delta^M[0, t]$. The boundary conditions of $\Omega_\delta(t)$ are inherited from $(\Omega_\delta; a_\delta, b_\delta, c_\delta, d_\delta)$ and $\gamma_\delta^M[0, t]$ as follows: $(a_\delta b_\delta) \cup \gamma_\delta^M[0, t]$ is wired and $(c_\delta d_\delta)$ is wired. Consider the UST in the quad $(\Omega_\delta(t); a_\delta, b_\delta, c_\delta, d_\delta)$ with such boundary condition. Let $\gamma_{\delta,t}^M$ be the unique branch from $(a_\delta b_\delta) \cup \gamma_\delta^M[0, t]$ to $(c_\delta d_\delta)$ in the UST on $\Omega_\delta(t)$. Denote by $X_\delta^M(t)$ the starting point of $\gamma_{\delta,t}^M$ on $(a_\delta b_\delta) \cup \gamma_\delta^M[0, t]$ and by $Y_\delta^M(t)$ the ending point of $\gamma_{\delta,t}^M$ on $(c_\delta d_\delta)$. For any bounded continuous function R on curves, we have

$$\begin{aligned} & \mathbb{E} \left[\mathbb{1}_{\{Y_\delta^M \in (\tilde{c}_\delta \tilde{d}_\delta)\}} R(\gamma_\delta^M[0, t \wedge T_\epsilon^\delta]) \right] \\ &= \mathbb{E} \left[\mathbb{P} \left[Y_\delta^M(t \wedge T_\epsilon^\delta) \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M(t \wedge T_\epsilon^\delta) = \gamma_\delta^M(t \wedge T_\epsilon^\delta) \right] R(\gamma_\delta^M[0, t \wedge T_\epsilon^\delta]) \right]. \end{aligned}$$

From the convergence of γ_δ^M to γ^M , we have

$$\mathbb{E} \left[\mathbb{1}_{\{Y_\delta^M \in (\tilde{c}_\delta \tilde{d}_\delta)\}} R(\gamma_\delta^M[0, t \wedge T_\epsilon^\delta]) \right] \rightarrow \mathbb{E} \left[\mathbb{1}_{\{Y^M \in (\tilde{c} \tilde{d})\}} R(\gamma^M[0, t \wedge T_\epsilon]) \right], \quad \text{as } \delta \rightarrow 0. \quad (5.21)$$

From Corollary 5.5, we have

$$\mathbb{P} \left[Y_\delta^M(t \wedge T_\epsilon^\delta) \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M(t \wedge T_\epsilon^\delta) = \gamma_\delta^M(t \wedge T_\epsilon^\delta) \right] \rightarrow \int_{f_{t \wedge T_\epsilon}(g_{t \wedge T_\epsilon}(\phi(\tilde{c})))}^{f_{t \wedge T_\epsilon}(g_{t \wedge T_\epsilon}(\phi(\tilde{d})))} \rho_{K_{t \wedge T_\epsilon}}(f_{t \wedge T_\epsilon}(W_{t \wedge T_\epsilon}), \text{Re}y) dy,$$

where ρ_K is defined in (1.1). Thus, by bounded convergence theorem, we have

$$\begin{aligned} & \mathbb{E} \left[\mathbb{E} \left[Y_\delta^M(t \wedge T_\epsilon^\delta) \in (\tilde{c}_\delta \tilde{d}_\delta) \mid X_\delta^M(t \wedge T_\epsilon^\delta) = \gamma_\delta^M(t \wedge T_\epsilon^\delta) \right] R(\gamma_\delta^M[0, t \wedge T_\epsilon^\delta]) \right] \\ & \rightarrow \mathbb{E} \left[R(\gamma^M[0, t \wedge T_\epsilon]) \int_{f_{t \wedge T_\epsilon}(g_{t \wedge T_\epsilon}(\phi(\tilde{c})))}^{f_{t \wedge T_\epsilon}(g_{t \wedge T_\epsilon}(\phi(\tilde{d})))} \rho_{K_{t \wedge T_\epsilon}}(f_{t \wedge T_\epsilon}(W_{t \wedge T_\epsilon}), \text{Re}y) dy \right], \quad \text{as } \delta \rightarrow 0. \end{aligned} \quad (5.22)$$

Combining (5.21) and (5.22), we have

$$\begin{aligned} & \mathbb{E} \left[\mathbb{1}_{\{Y^M \in (\tilde{c} \tilde{d})\}} R(\gamma^M([0, t \wedge T_\epsilon]) \right] \\ & = \mathbb{E} \left[R(\gamma^M([0, t \wedge T_\epsilon]) \int_{f_{t \wedge T_\epsilon}(g_{t \wedge T_\epsilon}(\phi(\tilde{c})))}^{f_{t \wedge T_\epsilon}(g_{t \wedge T_\epsilon}(\phi(\tilde{d})))} \rho_{K_{t \wedge T_\epsilon}}(f_{t \wedge T_\epsilon}(W_{t \wedge T_\epsilon}), \text{Re}y) dy \right]. \end{aligned}$$

This implies that the process

$$\left(\int_{f_t(g_t(\phi(\tilde{c})))}^{f_t(g_t(\phi(\tilde{d})))} \rho_{K_t}(f_t(W_t), \text{Re}y) dy, t \geq 0 \right)$$

is a martingale up to T_ϵ . Thus, the process

$$((f_t \circ g_t)'(x) \rho_{K_t}(f_t(W_t), \text{Re}f_t(g_t(x))), t \geq 0)$$

is a martingale up to T_ϵ for every $x \in (\phi(c), +\infty)$. Combining (1.1), (5.8), (5.10) and (5.19), we have

$$F(x; \phi(a), W_0, \phi(b), \phi(c)) = f'(x) \rho_K(f(W_0), \text{Re}f(x)).$$

Thus, the process

$$(g_t'(x) F(g_t(x); g_t(\phi(a)), W_t, g_t(\phi(b)), g_t(\phi(c))), t \geq 0)$$

is a martingale up to T_ϵ for every $x \in (\phi(c), +\infty)$. From Lemma 4.9, the curve γ^M intersects $\partial\Omega$ only at two ends almost surely. Thus $T_\epsilon \rightarrow T$ as $\epsilon \rightarrow 0$. This completes the proof. \square

Proof of Theorem 1.6. The joint law of (X^M, Y^M) is derived in Proposition 5.1. It remains to show that the conditional law of γ^M given X^M is $\text{SLE}_2(-1, -1; -1, -1)$. To this end, we may assume $\Omega = \mathbb{H}$ with $d = \infty$ and $a < b < c$ and parameterize γ^M by the half-plane capacity. Denote by T the first time that γ^M hits (c, ∞) . Denote by $(W_t, t \geq 0)$ the driving function of γ^M and by $(g_t, t \geq 0)$ the corresponding conformal maps. For $a < w < b < c < x$, define $F(x; a, w, b, c)$ as in (5.19). Lemma 5.10 tells that the process

$$(g_t'(x) F(g_t(x); g_t(a), W_t, g_t(b), g_t(c)), t \geq 0)$$

is a martingale up to T . By the same argument in [Kar20, Lemma 5.3], we can deduce that $(W_t, t \geq 0)$ is a semimartingale. Denote by L_t the drift term of W_t . By Itô's formula, we have

$$\begin{aligned} & \left(\frac{2}{g_t(x) - W_t} \partial_x + \frac{2}{g_t(a) - W_t} \partial_a + \frac{2}{g_t(b) - W_t} \partial_b + \frac{2}{g_t(c) - W_t} \partial_c + \frac{-2}{(g_t(x) - W_t)^2} \right) F dt \\ & \quad + \partial_w F dL_t + \frac{1}{2} \partial_w^2 F d\langle W \rangle_t = 0. \end{aligned}$$

Combining with (5.20), we have

$$\partial_w F \left(dL_t - 2 \frac{f''(W_t; g_t(a), g_t(b), g_t(c))}{f'(W_t; g_t(a), g_t(b), g_t(c))} dt \right) + \frac{1}{2} \partial_w^2 F (d\langle W \rangle_t - 2dt) = 0.$$

From (5.13), it simplifies as

$$\partial_w F \left(dL_t - \left(\frac{1}{g_t(a) - W_t} + \frac{1}{g_t(b) - W_t} + \frac{1}{g_t(c) - W_t} \right) dt \right) + \frac{1}{2} \partial_w^2 F (d\langle W \rangle_t - 2dt) = 0. \quad (5.23)$$

Note that (5.23) holds for all $x \in \mathbb{Q} \cap (c, +\infty)$ almost surely. By the continuity, we have that (5.23) holds for all $x \in (c, +\infty)$ almost surely. Now we fix (a, b, c) and t . Define $S_1(x) := \partial_w F(g_t(x); g_t(a), W_t, g_t(b), g_t(c))$ and $S_2(x) := \partial_w^2 F(g_t(x); g_t(a), W_t, g_t(b), g_t(c))$. It suffices to prove

$$\exists x, x' \in (c, \infty) \text{ such that } S_1(x)S_2(x') \neq S_2(x)S_1(x'). \quad (5.24)$$

Assume this is true, then we have

$$dL_t = \left(\frac{1}{g_t(a) - W_t} + \frac{1}{g_t(b) - W_t} + \frac{1}{g_t(c) - W_t} \right) dt \quad \text{and} \quad d\langle W \rangle_t = 2dt.$$

This shows that γ^M is $\text{SLE}_2(-1; -1, -1)$ as desired.

It remains to show (5.24). Denote $f(\cdot) = f(\cdot; a, b, c)$. Note that

$$\begin{aligned} S_1(x) &= -2f(g_t(x)) \frac{\pi}{4K} \times \frac{\pi}{2K} f'(W_t) \times \sum_{n \in \mathbb{Z}} \left(\frac{\sinh\left(\frac{\pi}{2K}(f(W_t) - \text{Re}f(g_t(x)) - 2n)\right)}{\cosh^3\left(\frac{\pi}{2K}(f(W_t) - \text{Re}f(g_t(x)) - 2n)\right)} + \frac{\sinh\left(\frac{\pi}{2K}(f(W_t) + \text{Re}f(g_t(x)) - 2n)\right)}{\cosh^3\left(\frac{\pi}{2K}(f(W_t) + \text{Re}f(g_t(x)) - 2n)\right)} \right), \\ S_2(x) &= \frac{f''(W_t)}{f'(W_t)} S_1(x) \\ & \quad + 2f'(g_t(x)) \frac{\pi}{4K} \times \left(\frac{\pi}{2K} f'(W_t) \right)^2 \times \sum_{n \in \mathbb{Z}} \left(\frac{2 \sinh^2\left(\frac{\pi}{2K}(f(W_t) - \text{Re}f(g_t(x)) - 2n)\right) - 1}{\cosh^4\left(\frac{\pi}{2K}(f(W_t) - \text{Re}f(g_t(x)) - 2n)\right)} + \frac{2 \sinh^2\left(\frac{\pi}{2K}(f(W_t) + \text{Re}f(g_t(x)) - 2n)\right) - 1}{\cosh^4\left(\frac{\pi}{2K}(f(W_t) + \text{Re}f(g_t(x)) - 2n)\right)} \right). \end{aligned}$$

Define

$$R_1(z) := \sum_{n \in \mathbb{Z}} \left(\frac{\cosh\left(\frac{\pi}{2K}(f(W_t) - z - 2n)\right)}{\sinh^3\left(\frac{\pi}{2K}(f(W_t) - z - 2n)\right)} + \frac{\cosh\left(\frac{\pi}{2K}(f(W_t) + z - 2n)\right)}{\sinh^3\left(\frac{\pi}{2K}(f(W_t) + z - 2n)\right)} \right)$$

and

$$R_2(z) := - \sum_{n \in \mathbb{Z}} \left(\frac{2 \cosh^2\left(\frac{\pi}{2K}(f(W_t) - z - 2n)\right) + 1}{\sinh^4\left(\frac{\pi}{2K}(f(W_t) - z - 2n)\right)} + \frac{2 \cosh^2\left(\frac{\pi}{2K}(f(W_t) + z - 2n)\right) + 1}{\sinh^4\left(\frac{\pi}{2K}(f(W_t) + z - 2n)\right)} \right).$$

If (5.24) is false, then $S_1(x)/S_2(x)$ is constant for $x > c$. Thus, there exists λ (which is random) such that $(R_1 - \lambda R_2)(z)|_{z \in (iK, 1+iK)} = 0$. Since R_1 and R_2 are analytic functions in $(0, 1) \times (0, iK)$, this implies that $R_1 = \lambda R_2$ in $(0, 1) \times (0, iK)$. This is a contradiction by considering the asymptotic of R_1 and R_2 when $z \rightarrow f(W_t)$:

$$\lim_{z \rightarrow f(W_t)} R_1(z)(z - f(W_t))^3 = \left(\frac{-2K}{\pi} \right)^3, \quad \lim_{z \rightarrow f(W_t)} R_2(z)(z - f(W_t))^4 = -3 \left(\frac{2K}{\pi} \right)^4.$$

This completes the proof. \square

5.3 Consequences

In this section, we complete the proof for Propositions 1.3 and 1.4 and Corollary 1.7.

Proof of Proposition 1.4. The conclusion is immediate from Theorem 1.5 and (5.4). \square

The proof for Proposition 1.3 and Corollary 1.7 bases on the following observation in the discrete for UST. Fix a Dobrushin domain $(\Omega; c, d)$ such that $\partial\Omega$ is C^1 and simple. Suppose $(\Omega_\delta; c_\delta, d_\delta)$ is an approximation of $(\Omega; c, d)$ on $\delta\mathbb{Z}^2$ as in Section 4.1. Let \mathcal{T}_δ be the UST in $(\Omega_\delta; c_\delta, d_\delta)$ with $(c_\delta d_\delta)$ wired. Denote by η_δ the associated Peano curve along \mathcal{T}_δ from d_δ^\diamond to c_δ^\diamond . Fix $a \in (dc)$ and let a_δ^\diamond be the medial vertex along $(c_\delta^\diamond d_\delta^\diamond)$ nearest to a . Let $a_\delta \in V(\Omega_\delta)$ be the primal vertex in Ω_δ that is nearest to a_δ^\diamond . Let a_δ^* (resp. b_δ^*) be the dual vertex along $(d_\delta^* c_\delta^*)$ that is nearest to a_δ^\diamond and is closer to d_δ^* (resp. closer to c_δ^*) along $(d_\delta^* c_\delta^*)$. See Figure 5.2. We divide the Peano curve η_δ into two parts: denote by $\tilde{\eta}_\delta^L$ the part of η_δ from d_δ^\diamond to a_δ^\diamond , and denote by η_δ^R the part of η_δ from a_δ^\diamond to c_δ^\diamond . Denote by η_δ^L the time-reversal of $\tilde{\eta}_\delta^L$. There is a branch in \mathcal{T}_δ connecting a_δ to $(c_\delta d_\delta)$ and we denote it by γ_δ . We parameterize γ_δ so that it starts from a_δ and terminates when it hits $(c_\delta d_\delta)$. We have the convergence of the triple $(\eta_\delta^L; \gamma_\delta; \eta_\delta^R)$.




Figure 5.2: In the left panel, the solid edges in black are wired boundary arc (cd) , and the solid edges in red are dual-wired boundary arc (d^*c^*) . The thin edges are in the UST. The thin edges in red are the branch γ in the tree connecting a to (cd) . In the right panel, the solid edges in blue are $\gamma[0, \tau]$ and the solid edges in green are $\gamma[\tau, T]$. The two orange curves are η^L and η^R .

Lemma 5.11. *Fix a polygon $(\Omega; d, a, c)$ with three marked points such that $\partial\Omega$ is C^1 and simple. Suppose that a sequence of medial polygons $(\Omega_\delta^\diamond; d_\delta^\diamond, a_\delta^\diamond, c_\delta^\diamond)$ converges to $(\Omega; d, a, c)$ as in (4.2). Then the triple $(\eta_\delta^L; \gamma_\delta; \eta_\delta^R)$ converges weakly to a triple of continuous curves $(\eta^L; \gamma; \eta^R)$ whose law is characterized as follows. Let η be an SLE_8 in Ω from d to c and let T_a be the first time that it swallows a . Then, the joint distribution of $(\eta^L; \eta^R)$ is the same as $(\eta(T_a - t), 0 \leq t \leq T_a; \eta(t), t \geq T_a)$; and $\gamma = \eta^L \cap \eta^R$.*

Proof. First of all, we prove the tightness of $\{(\eta_\delta^L; \gamma_\delta; \eta_\delta^R)\}$. The tightness of $\{\eta_\delta^L\}$ and $\{\eta_\delta^R\}$ is given in the proof of Theorem 4.1. The tightness of $\{\gamma_\delta\}$ can be proved in the same way as in Lemma 4.9. Therefore, the triple $\{(\eta_\delta^L; \gamma_\delta; \eta_\delta^R)\}$ is tight.

Next, we determine the law of subsequential limits. Suppose $(\eta^L; \gamma; \eta^R)$ is any subsequential limit. There exists $\{\delta_n\}$ with $\delta_n \rightarrow 0$ as $n \rightarrow \infty$, such that $\eta_{\delta_n}^L \rightarrow \eta^L$ and $\gamma_{\delta_n} \rightarrow \gamma$ and $\eta_{\delta_n}^R \rightarrow \eta^R$ in law as $n \rightarrow \infty$. By Theorem 4.1, $\{\eta_{\delta_n}\}_n$ converges weakly to η as $n \rightarrow \infty$. Thus, $(\eta^L; \eta^R)$ has the same law as $(\eta(T_a - t), 0 \leq t \leq T_a; \eta(t), t \geq T_a)$. Since SLE_8 is space filling, we have $\gamma = \eta^L \cap \eta^R$. This completes the proof. \square

From the observation in Lemma 5.11, we arrive at the following lemma.

Lemma 5.12. *Fix a polygon $(\Omega; d, a, c)$ with three marked points such that $\partial\Omega$ is C^1 and simple. Let η be an SLE_8 in Ω from d to c and let T_a be the first time that it swallows a . Denote by γ the right boundary of $\eta[0, T_a]$. Denote by Ω^L and Ω^R the two connected components of $\Omega \setminus \gamma$ such that Ω^L has d on the boundary and Ω^R has c on the boundary. The joint law of the triple*

$$(\eta(T_a - t), 0 \leq t \leq T_a; \quad \gamma; \quad \eta(t), t \geq T_a)$$

can be characterized as follows: γ is $\text{SLE}_2(-1, -1; -1, -1)$ in Ω from a to (cd) with force points $(d, a^-; a^+, c)$; given γ , the conditional law of $(\eta(T_a - t), 0 \leq t \leq T_a)$ is SLE_8 in Ω^L from a^- to d and the conditional law of $(\eta(t), t \geq T_a)$ is SLE_8 in Ω^R from a^+ to c , and $(\eta(T_a - t), 0 \leq t \leq T_a)$ and $(\eta(t), t \geq T_a)$ are conditionally independent given γ .

Proof. First, we derive the marginal law of γ . Choose a conformal map ϕ from Ω onto \mathbb{H} such that $\phi(a) = 0$ and $\phi(d) = \infty$. Denote by $(W_t, t \geq 0)$ the driving function of $\phi(\gamma)$ and by $(g_t, t \geq 0)$ the corresponding conformal maps. By the argument in the proof of Lemma 5.10, for any $x \in (\phi(c), +\infty)$, the process

$$(g'_t(x)F(g_t(x); g_t(0^-), W_t, g_t(0^+), g_t(\phi(c))), t > 0)$$

is a martingale up to the first time that γ hits (cd) where F is defined in (5.19). For $\epsilon > 0$, define $\tau_\epsilon := \inf\{t : d(\phi(\gamma(t)), 0) \geq \epsilon\}$. By the argument in the proof of Theorem 1.6, the conditional law of $(\gamma(t), t \geq \tau_\epsilon)$ given $\gamma[0, \tau_\epsilon]$ is $\text{SLE}_2(-1, -1; -1, -1)$ in $\Omega \setminus \gamma[0, \tau_\epsilon]$ from $\gamma(\tau_\epsilon)$ to (cd) with force points $(d, a^-; a^+, c)$. Let $\epsilon \rightarrow 0$, the law of γ is $\text{SLE}_2(-1, -1; -1, -1)$ in Ω from a to (cd) with force points $(d, a^-; a^+, c)$.

Second, the conditional law of $(\eta(T_a - t), 0 \leq t \leq T_a; \eta(t), t \geq T_a)$ given γ can be proved in the same way as in Theorem 1.5, thanks to the observation in Lemma 5.11 and Figure 5.2. \square

Proof of Corollary 1.7. The conclusion is immediate from Lemma 5.12. \square

Proof of Proposition 1.3. Assume the same notation as in Lemma 5.12. We parameterize γ so that it starts from a and terminates when it hits (cd) at time T . Let τ be any stopping time of γ before T . Consider the conditional law of the following triple given $\gamma[0, \tau]$:

$$(\eta(T_a - t), 0 \leq t \leq T_a; \quad \gamma(t), \tau \leq t \leq T; \quad \eta(t), t \geq T_a).$$

From Lemma 5.12, the law of $(\gamma(t), \tau \leq t \leq T)$ is $\text{SLE}_2(-1, -1; -1, -1)$ in $\mathbb{H} \setminus \gamma[0, \tau]$ from $\gamma(\tau)$ to $(-\infty, x)$ with force points $(d, a^-; a^+, c)$; the conditional law of $(\eta(T_a - t), 0 \leq t \leq T_a)$ given $\gamma[\tau, T]$ is SLE_8 in Ω^L from a^- to d , the conditional law of $(\eta(t), t \geq T_a)$ given $\gamma[\tau, T]$ is SLE_8 in Ω^R from a^+ to c , and $(\eta(T_a - t), 0 \leq t \leq T_a)$ and $(\eta(t), t \geq T_a)$ are conditionally independent given $\gamma[\tau, T]$. Comparing with Theorems 1.5 and 1.6, we see that the triple has the same law as the triple $(\eta^L; \gamma^M; \eta^R)$ in Theorem 1.5 in the quad $(\Omega \setminus \gamma[0, \tau]; d, a^-, a^+, c)$ conditional on $X^M = \gamma(\tau)$. In particular, the law of $(\eta(T_a - t), 0 \leq t \leq T_a)$ is hSLE_8 in $\Omega \setminus \gamma[0, \tau]$ from a^- to d conditional that its last hitting point of $\gamma[0, \tau]$ is $\gamma(\tau)$. Combining with reversibility of hSLE_8 , we obtain the conclusion. \square

A Hypergeometric function and elliptic integral

For $A, B, C \in \mathbb{R}$, the hypergeometric function is defined for $|z| < 1$ by the power series:

$$F(z) = {}_2F_1(A, B, C; z) = \sum_{n=0}^{\infty} \frac{(A)_n (B)_n}{(C)_n} \frac{z^n}{n!}, \quad (\text{A.1})$$

where $(x)_n := x(x+1)\cdots(x+n-1)$ for $n \geq 1$ and $(x)_n = 1$ for $n = 0$. The power series is well-defined when $C \notin \{0, -1, -2, -3, \dots\}$. The hypergeometric function is a solution of Euler's hypergeometric differential equation:

$$z(1-z)F''(z) + \left(\frac{2\nu+8}{\kappa} - \frac{2\nu+2\kappa}{\kappa}z \right) F'(z) - \frac{2(\nu+2)(\kappa-4)}{\kappa^2} F(z) = 0. \quad (\text{A.2})$$

We collect some properties for hypergeometric functions here. For $z \in (-1, 1)$, we have (see [AS92, Eq. 15.2.1 and Eq. 15.3.3])

$${}_2F_1(A, B, C; z) = (1-z)^{C-A-B} {}_2F_1(C-A, C-B, C; z), \quad (\text{A.3})$$

$$\frac{d}{dz} {}_2F_1(A, B, C; z) = \frac{AB}{C} {}_2F_1(A+1, B+1, C+1; z). \quad (\text{A.4})$$

The series (A.1) is absolutely convergent on $z \in [0, 1]$ when $C > A+B$ and $C \notin \{0, -1, -2, \dots\}$. In this case, we have (see [AS92, Eq. 15.1.20])

$${}_2F_1(A, B, C; 1) = \frac{\Gamma(C)\Gamma(C-A-B)}{\Gamma(C-A)\Gamma(C-B)}, \quad (\text{A.5})$$

where Γ is Gamma Function.

Denote by \mathcal{K} the elliptic integral of the first kind (see [AS92, Eq. 17.2.6]): for $\varphi \in \overline{\mathbb{H}}$ and $x \in (0, 1)$,

$$\mathcal{K}(\varphi, x) := \int_0^\varphi \frac{d\theta}{\sqrt{1-x\sin^2\theta}}, \quad \mathcal{K}(x) := \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-x\sin^2\theta}}. \quad (\text{A.6})$$

There is a relation between complete elliptic integral and hypergeometric function (see [AS92, Eq. 17.3.9]):

$$\mathcal{K}(x) = \frac{\pi}{2} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}, 1; x\right), \quad \forall x \in (0, 1). \quad (\text{A.7})$$

Let us calculate the derivatives of \mathcal{K} :

$$\partial_\varphi \mathcal{K}(\varphi, x) = \frac{1}{\sqrt{1-x\sin^2\varphi}}, \quad \partial_\varphi^2 \mathcal{K}(\varphi, x) = \frac{x \sin \varphi \cos \varphi}{\sqrt{1-x\sin^2\varphi}^3}, \quad \partial_\varphi \partial_x \mathcal{K}(\varphi, x) = \frac{\sin^2 \varphi}{2\sqrt{1-x\sin^2\varphi}^3}. \quad (\text{A.8})$$

The derivatives $\partial_x \mathcal{K}$ and $\partial_x^2 \mathcal{K}$ involve the elliptic integral of the second kind, but luckily, we do not need them.

B Convergence of discrete harmonic functions

Lemma B.1. *Fix a quad $(\Omega; a, b, c, d)$ and fix a conformal map ξ from Ω onto \mathbb{U} and extend its definition continuously to the boundary. Suppose a sequence of domains $(\Omega_\delta; a_\delta, b_\delta, c_\delta, d_\delta)$ converges to $(\Omega; a, b, c, d)$ in the Carathéodory sense as $\delta \rightarrow 0$. For $z \in V(\Omega_\delta)$, let $u_\delta(z)$ be the probability that a simple random walk in Ω_δ starting from z hits $(a_\delta b_\delta)$ before $(c_\delta d_\delta)$. Let $u(z)$ be the bounded harmonic function on Ω such that $u \circ \xi^{-1}$ satisfies the following boundary data: $u \circ \xi^{-1} = 1$ on $(\xi(a)\xi(b))$, and $u \circ \xi^{-1} = 0$ on $(\xi(c)\xi(d))$, and $\partial_n u \circ \xi^{-1} = 0$ on $(\xi(b)\xi(c)) \cup (\xi(d)\xi(a))$ where n is the outer normal vector. Then u_δ converges to u locally uniformly as $\delta \rightarrow 0$.*

Proof. This is a consequence of [LSW04, Proposition 4.2]. We summarize the proof here for concreteness. For any function g_δ on $V(\Omega_\delta)$ or $V(\Omega_\delta^*)$, define its discrete derivatives as

$$\partial_x^\delta g_\delta := \delta^{-1}(g_\delta(v+\delta) - g_\delta(v)), \quad \partial_y^\delta g_\delta := \delta^{-1}(g_\delta(v+i\delta) - g_\delta(v)).$$

We extend u_δ and all its derivatives to functions on Ω_δ by linear interpolation.

First, we show that any subsequential limit of u_δ is harmonic on Ω . Since u_δ is discrete harmonic and $0 \leq u_\delta \leq 1$, by [LSW04, Lemma 5.2], for any compact set $S \subset \Omega$ and $k \in \mathbb{N}$, there exists a constant $C > 0$ which depends on S and k such that

$$|\partial_{a_1}^\delta \partial_{a_2}^\delta \cdots \partial_{a_k}^\delta u_\delta(v)| \leq C, \quad \text{for any } \partial_{a_1}^\delta, \dots, \partial_{a_k}^\delta \in \{\partial_x^\delta, \partial_y^\delta\}, \text{ and } v \in S \cap \Omega_\delta.$$

By Arzela-Ascoli theorem, for any sequence $\delta_n \rightarrow 0$, there exist a subsequence, still denoted by $\{\delta_n\}$, and continuous functions $u, u_x, u_y, u_{xx}, u_{yy}$ such that

$$u_\delta \rightarrow u, \quad \partial_x^{\delta_n} u_{\delta_n} \rightarrow u_x, \quad \partial_y^{\delta_n} u_{\delta_n} \rightarrow u_y, \quad \partial_x^{\delta_n} \partial_x^{\delta_n} u_{\delta_n} \rightarrow u_{xx}, \quad \partial_y^{\delta_n} \partial_y^{\delta_n} u_{\delta_n} \rightarrow u_{yy}, \quad \text{locally uniformly.}$$

This implies

$$u_x = \partial_x u, \quad u_y = \partial_y u, \quad u_{xx} = \partial_x^2 u, \quad u_{yy} = \partial_y^2 u.$$

Since u_{δ_n} is discrete harmonic, the function u is harmonic. This shows that any subsequential limit of u_δ is harmonic.

Next, we show that any subsequential limit u has the boundary data given in the statement. Note that such boundary data and harmonicity and boundedness uniquely determines the subsequential limit, hence gives the convergence of sequence u_δ . From the definition of u_δ and Beurling estimate, it is clear that $u = 1$ on (ab) and $u = 0$ on (cd) . To derive the boundary data along $(bc) \cup (ad)$, we need to introduce the discrete harmonic conjugate function v_δ^* of u_δ .

Define $v_\delta^* = 0$ on $(b_\delta^* c_\delta^*)$. For every oriented edge $e_\delta^* = \{x_\delta^*, y_\delta^*\} \in E(\Omega_\delta^*)$, there is a unique oriented edge $e = \{x_\delta, y_\delta\} \in E(\Omega_\delta)$ which crosses e_δ^* from its right-side. Define $v_\delta^*(y_\delta^*) - v_\delta^*(x_\delta^*) := u_\delta(y_\delta) - u_\delta(x_\delta)$. Since u_δ is discrete harmonic, this is well-defined and v_δ^* is constant on $(d_\delta^* a_\delta^*)$. Moreover, v_δ^* takes its maximum, denoted by L_δ , on $(d_\delta^* a_\delta^*)$.

We claim that $\{L_\delta\}_{\delta>0}$ is uniformly bounded. Assume this is true, by the same argument as above, for any sequence $\delta_n \rightarrow 0$, there exists a subsequence, still denoted by δ_n , and a constant \tilde{K} and a harmonic function v such that $v_{\delta_n}^* \rightarrow v$ and the other related derivatives also converge locally uniformly and that $L_n \rightarrow \tilde{K}$ as $n \rightarrow \infty$. From the construction and Beurling estimate, we have $v = \tilde{K}$ on (da) and $v = 0$ on (bc) . By the definition of v_δ^* , the function $f_\delta := u_\delta + i v_\delta^*$ is discrete holomorphic. Then, the convergence of discrete derivatives implies that $f := u + i v$ is conformal on Ω . By Schwartz reflection principle, we can extend $f \circ \xi^{-1}$ to $\partial\mathbb{U} \setminus \{\xi(a), \xi(b), \xi(c), \xi(d)\}$ analytically. By Cauchy-Riemann equation, we have

$$\partial_n(u \circ \xi^{-1}) = \partial_t(v \circ \xi^{-1}) = 0$$

on $(\xi(b)\xi(c))$ and $(\xi(d)\xi(a))$ where t is the tangential vector. This gives the required boundary data of u along $(bc) \cup (da)$.

Finally, it remains to show that $\{L_\delta\}_{\delta>0}$ is uniformly bounded. If this is not the case, there exists a sequence $\delta_n \rightarrow 0$ such that $L_{\delta_n} \rightarrow \infty$ as $n \rightarrow \infty$. By the same argument as above, the sequence $\frac{1}{L_n} f_{(\Omega_{\delta_n}; a_{\delta_n}, b_{\delta_n}, c_{\delta_n}, d_{\delta_n})}$ converges to a conformal map h locally uniformly. In such case, we have $\text{Re} h = 0$ on Ω , thus h is constant. But $\text{Im} h = 1$ on (da) and $\text{Im} h = 0$ on (bc) , this gives a contradiction. Thus, $\{L_\delta\}_{\delta>0}$ is uniformly bounded and we complete the proof. \square

References

- [Ahl78] Lars V. Ahlfors. Complex analysis. *McGraw-Hill Book Co., New York, third edition, 1978.*
- [AS92] Milton Abramowitz and Irene A. Stegun, editors. Handbook of mathematical functions with formulas, graphs, and mathematical tables. *Dover Publications, Inc., New York, 1992.*
- [BPW18] Vincent Beffara, Eveliina Peltola, and Hao Wu. On the uniqueness of global multiple SLEs, *arXiv:1801.07699*, 2018.

- [CDCH⁺14] Dmitry Chelkak, Hugo Duminil-Copin, Clément Hongler, Antti Kemppainen, and Stanislav Smirnov. Convergence of Ising interfaces to Schramm’s SLE curves. *C. R. Math. Acad. Sci. Paris*, 352(2):157–161, 2014.
- [CW19] Dmitry Chelkak and Yijun Wan. On the convergence of massive loop-erased random walks to massive SLE(2) curves. *arXiv:1903.08045*, 2019.
- [DC13] Hugo Duminil-Copin. Parafermionic observables and their applications to planar statistical physics models. *Ensaïos Matemáticos*, 25:1–371, 2013.
- [DCS12] Hugo Duminil-Copin and Stanislav Smirnov. Conformal invariance of lattice models. In *Probability and statistical physics in two and more dimensions*, volume 15 of *Clay Math. Proc.*, pages 213–276. Amer. Math. Soc., Providence, RI, 2012.
- [Dub06] Julien Dubédat. Euler integrals for commuting SLEs. *J. Stat. Phys.*, 123(6):1183–1218, 2006.
- [Izy15] Konstantin Izyurov. Smirnov’s observable for free boundary conditions, interfaces and crossing probabilities. *Comm. Math. Phys.*, 337(1):225–252, 2015.
- [Kar19] Alex Karrila. Multiple SLE type scaling limits: from local to global, *arXiv:1903.10354*, 2019.
- [Kar20] Alex Karrila. UST branches, martingales, and multiple SLE(2), *arXiv:2002.07103*, 2020.
- [KS18] Antti Kemppainen and Stanislav Smirnov. Configurations of FK Ising interfaces and hypergeometric SLE. *Math. Res. Lett.*, 25(3):875–889, 2018.
- [KW11] Richard W. Kenyon and David B. Wilson. Boundary partitions in trees and dimers. *Trans. Amer. Math. Soc.*, 363(3):1325–1364, 2011.
- [LSW04] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Ann. Probab.*, 32(1B):939–995, 2004.
- [MS16a] Jason Miller and Scott Sheffield. Imaginary geometry I: Interacting SLEs. *Probab. Theory Related Fields*, 164(3-4):553–705, 2016.
- [MS16b] Jason Miller and Scott Sheffield. Imaginary geometry II: Reversibility of $SLE_{\kappa}(\rho_1; \rho_2)$ for $\kappa \in (0, 4)$. *Ann. Probab.*, 44(3):1647–1722, 2016.
- [MS16c] Jason Miller and Scott Sheffield. Imaginary geometry III: Reversibility of SLE_{κ} for $\kappa \in (4, 8)$. *Ann. of Math. (2)*, 184(2):455–486, 2016.
- [MS17] Jason Miller and Scott Sheffield. Imaginary geometry IV: interior rays, whole-plane reversibility, and space-filling trees. *Probab. Theory Related Fields*, 169(3-4):729–869, 2017.
- [Pom92] Ch. Pommerenke. Boundary behaviour of conformal maps, volume 299 of *Grundlehren der Mathematischen Wissenschaften*, Springer-Verlag, Berlin, 1992.
- [RS05] Steffen Rohde and Oded Schramm. Basic properties of SLE. *Ann. of Math. (2)*, 161(2):883–924, 2005.
- [Sch00] Oded Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118:221–288, 2000.
- [Smi01] Stanislav Smirnov. Critical percolation in the plane: conformal invariance, Cardy’s formula, scaling limits. *C. R. Acad. Sci. Paris Sér. I Math.*, 333(3):239–244, 2001.
- [Smi06] Stanislav Smirnov. Towards conformal invariance of 2D lattice models. In *International Congress of Mathematicians. Vol. II*, pages 1421–1451. Eur. Math. Soc., Zürich, 2006.
- [SS09] Oded Schramm and Scott Sheffield. Contour lines of the two-dimensional discrete Gaussian free field. *Acta Math.*, 202(1):21–137, 2009.
- [SW05] Oded Schramm and David B. Wilson. SLE coordinate changes. *New York J. Math.*, 11:659–669, 2005.
- [Wu20] Hao Wu. Hypergeometric SLE: conformal Markov characterization and applications. *Comm. Math. Phys.*, 374(2):433–484, 2020.
- [Zha08a] Dapeng Zhan. Duality of chordal SLE. *Invent. Math.*, 174(2):309–353, 2008.
- [Zha08b] Dapeng Zhan. Reversibility of chordal SLE. *Ann. Probab.*, 36(4):1472–1494, 2008.
- [Zha08c] Dapeng Zhan. The scaling limits of planar LERW in finitely connected domains. *Ann. Probab.*, 36(2):467–529, 2008.