Preprint typeset in JHEP style - HYPER VERSION

arXiv:2008.00659v2 [gr-qc] 18 Aug 2020

Covariant Hamiltonian Formalism for $F(R)-$Gravity

J. Klusoň

Department of Theoretical Physics and Astrophysics
Faculty of Science, Masaryk University
Kotlářská 2, 611 37, Brno
Czech Republic
E-mail: klu@physics.muni.cz

B. Matouš

Department of Theoretical Physics and Astrophysics
Faculty of Science, Masaryk University
Kotlářská 2, 611 37, Brno
Czech Republic
E-mail: 426945@mail.muni.cz

Abstract: In this short note we perform covariant Hamiltonian analysis of $F(R)-$gravity.

Contents

1. Introduction and Summary **1**

2. Introduction of $F(R)$-Gravity **2**

3. Covariant Hamiltonian Formalism with (g_{ab}, M^{abc}) as canonical variables **4**

4. Hamiltonian Analysis in New Variables (f^{ab}, N^c_{ab}) **5**

1. Introduction and Summary

The fundamental concept of modern field theories is their formulation with the help of action and corresponding Lagrangian density. This formulation is manifestly covariant which reflects basic concept of general relativity. Dynamics of these fields is then governed by Lagrange equations of motion which are again manifestly covariant. In other words, in the Lagrangian formulation all space-time coordinates are treated on equal footing.

The situation is different when we switch to the Hamiltonian formalism when one direction known as time direction is selected and Hamiltonian equations of motion basically determine evolution of the system along this direction. Clearly such a splitting of space-time into space and time breaks manifest covariance of the theory. On the other hand there exists covariant Hamiltonian formulation of field theory known as Weyl-De Donder theory [1, 2]. Let us demonstrate its principle on the simple example of the Lagrangian for scalar field ϕ that clearly depend on derivative $\partial_a\phi$. In conventional canonical formalism we consider time derivative as special one and define conjugate momentum as derivative of Lagrangian density with respect to $\partial_t\phi$. In case of Weyl-DeDonder theory we treat all partial derivatives on the equal footing which clearly preserves diffeomorphism invariance. Then covariant canonical Hamiltonian density depends on conjugate momenta p^a_M which are variables conjugate to $\partial_a\phi^M$. This approach is also known as multisymplectic field theory, see for example [3, 4, 5], for review, see [6] and for recent interesting application of this formalism in string theory, see [7, 8].

Intuitively it is clear that such a covariant Hamiltonian approach could be very convenient for all covariant theories and especially it would be very appropriate in case of general relativity. In fact, covariant Hamiltonian formulation of gravity was presented by P. Horava in his paper [13]. It was shown there that the analysis simplifies considerably when instead of conventional canonical variable g_{ab} we use $f^{ab} \equiv \sqrt{-g}g^{ab}$. In fact, an importance of this variable in the holographic formulation of gravity was stressed recently by T. Padmanabhan in his works, for very clear and detailed discussion of properties of gravity formulated in f^{ab} variables we recommend his paper [11].

It is natural to ask the question whether covariant Hamiltonian analysis can be performed with more general theories of gravity, as for example $F(R)-$gravity, for review and extensive list of references see [15, 16, 14]. These theories can be considered as simplest modification of general relativity that have a potential to explain some aspects of inflation or dark energy. On the other hand considering $F(R)-$gravity as generalization of general relativity, it is natural to study whether it is possible to find its covariant Hamiltonian formulation. In fact, this is the goal of this paper. As the first step we introduce auxiliary fields in order to be able to deal with $F(R)-$gravity and then we carefully split Lagrangian density for $F(R)-$gravity into bulk and boundary part. We find that there is an additional contribution to the bulk part with respect to ordinary gravity that is proportional to the partial derivative of the auxiliary field. Then we can straightforwardly proceed to the covariant Hamiltonian formulation when we firstly determine momenta conjugate to g_{ab}. We find that they are rather complicated which has been previously stressed in [11]. Then, following discussion presented in that paper, we proceed to the canonical variable $f^{ab} = \sqrt{-g}g^{ab}$ that are related to g_{ab} through point transformations. We determine momenta N^c_{ab} conjugate to f^{ab} and p^c_B conjugate to auxiliary field B. Finally we invert relations between conjugate momenta N^c_{ab} and Christoffel symbols and also between p^c_B and derivative of B and obtain corresponding Hamiltonian. We also determine canonical equations of motion.

Let us outline our results. We find covariant Hamiltonian for $F(R)-$gravity which is the first step in the application of this formalism to more general form of gravity. We also determine corresponding equations of motion. It would be certainly nice to extend this result along directions that are related to Padmanabhan's work. It would be also nice to apply covariant Hamiltonian analysis in case of more general theories of gravity, as for example curvature-squared gravity [17]. Finally, it would be interesting to find covariant Hamiltonian formalism for $F(R)-$gravity formulated in Jordan frame and compare it with the result derived in this paper and perform similar analysis as in [18, 19, 20]. We hope to return to these problems in future.

This paper is organized as follows. In the next section (2) we introduce Lagrangian for $F(R)-$gravity and split it into bulk and boundary parts. Then in section (3) we present canonical formalism in g_{ab} and M^{abc} variables. Finally in section (4) we proceed to the Hamiltonian analysis in f^{ab}, N^c_{ab} variables and determine corresponding Hamiltonian density and canonical equations of motion.

2. Introduction of $F(R)$-Gravity

In this section we introduce an action for $F(R)-$gravity. This theory is generalization of Einstein-Hilbert action when term linear in curvature R is replaced by arbitrary function $F(R)$. Explicitly, the action for $F(R)-$gravity has the form

$$S = \int d^4x \mathcal{L}^F \ , \quad \mathcal{L}^F = \frac{1}{16\pi}\sqrt{-g}F(R) \ , \tag{2.1}$$

where we set gravitational constant and speed of light to be equal one and where F is an arbitrary function of Ricci scalar R that is defined as [1]

$$R = g^{mn}R_{mn} = g^{mn}R^{\ k}_{mkn} \ ,$$
$$R^k_{\ lmn} = \partial_m\Gamma^k_{nl} - \partial_n\Gamma^k_{ml} + \Gamma^k_{mp}\Gamma^p_{nl} - \Gamma^k_{np}\Gamma^p_{ml} \ . \tag{2.2}$$

In order to find Hamiltonian formulation of this theory it is convenient to introduce two scalar fields A and B and write the Lagrangian density in the form

$$\mathcal{L} = \frac{1}{16\pi}\sqrt{-g}[F(B) + A(R-B)] \ , \tag{2.3}$$

where the equations of motion for A implies $R = B$ while equation of motion for B implies $F'(B) - A = 0$. Inserting the first result into (2.3) we obtain original action. Instead it is useful to use second equation to express $A = F'(B)$ and write the Lagrangian density (2.3) in the form

$$\mathcal{L} = \frac{1}{16\pi}\sqrt{-g}[F(B) + F'(B)(R-B)] \ . \tag{2.4}$$

In order to check consistency of this action let us solve equations of motion for B that follow from (2.4)

$$F'(B) + F''(B)(R-B) - F'(B) = F''(B)(R-B) = 0 \tag{2.5}$$

which again implies $R = B$ on condition $F''(B) \neq 0$. Inserting this result into (2.4) we obtain original Lagrangian density (2.1). In fact, for $F''(B) = 0$ we find that $F'(B) = K$ where $K = \text{const}$ and hence $F(B) = KB + C$ where C is another constant. Inserting this result into (2.4) we obtain Einstein-Hilbert action with cosmological constant proportional to C after appropriate rescaling. In what follows we will presume condition $F''(B) \neq 0$ that leads to $F(R)-$gravity.

Following [11] we write Ricci scalar as

$$R = Q_k^{\ mnl}R^k_{\ mnl} \ , \tag{2.6}$$

where

$$Q_k^{\ mnl} = \frac{1}{2}(g^{ml}\delta^n_k - g^{mn}\delta^l_k) \ , \quad Q_k^{\ mnl} = -Q_k^{\ mln} \ . \tag{2.7}$$

As the next step we split Lagrangian density (2.4) into bulk and boundary part [9, 10], for recent careful discussion, see also [11][2]. This procedure is trivial in case of terms quadratic in Christoffel symbols in Riemann tensor however we should be more careful with terms that contain partial derivatives of Christoffel tensor since for example

$$\sqrt{-g}F'(B)Q_k^{\ mnl}\partial_m\Gamma^k_{nl} = \partial_m[\sqrt{-g}F'(B)Q_k^{\ mnl}\Gamma^k_{nl}] -$$
$$-\partial_m[\sqrt{-g}Q_k^{\ mnl}]F'(B)\Gamma^k_{nl} - \sqrt{-g}Q_k^{\ mnl}F''(B)\partial_m B\Gamma^k_{nl} \ . \tag{2.8}$$

[1] We work with the metric g_{ab} with signature $(-1,1,1,1)$ where $a,b,c,\ldots = 0,\ldots,3$.

[2] It is important to stress that the variation of the bulk term alone leads to the equations of motion while the surface term, when integrated over a horizon, is related to the entropy of the horizon, as was shown for example in [12].

The second term on the second line gives bulk contribution corresponding to the kinetic term for auxiliary field B. Further, the first term gives also bulk contribution as follows from the fact that

$$\partial_c(\sqrt{-g}Q_a{}^{bcd}) = -\sqrt{-g}\Gamma^b_{cm}Q_a{}^{mcd} + \Gamma^m_{ca}Q_m{}^{bcd}\sqrt{-g} \tag{2.9}$$

which is consequence of the condition $\nabla_c(\sqrt{-g}Q_a{}^{bcd}) = 0$. Collecting these terms together we obtain that surface term has explicit form

$$\begin{aligned} &16\pi\mathcal{L}^F_{surf} = 2\partial_c[\sqrt{-g}F'(B)Q_a{}^{bcd}\Gamma^a_{bd}] = \\ &= 2F''(B)\partial_c B Q_a{}^{bcd}\Gamma^a{}_{bd} - 4F'(B)\sqrt{-g}Q_a{}^{bcd}\Gamma^a_{dm}\Gamma^m_{bc} + 2F'(B)\sqrt{-g}Q_a{}^{bcd}\partial_c\Gamma^a_{bd} \end{aligned} \tag{2.10}$$

and hence we find that decomposition of the Lagrangian density of $F(R)-$gravity into surface and bulk terms has the form

$$\begin{aligned} &16\pi\mathcal{L}^F = 16\pi\mathcal{L}^F_{quadr} + 16\pi\mathcal{L}^F_{surf}\ , \\ &16\pi\mathcal{L}^F_{quadr} = 2F'(B)Q_a{}^{bcd}\Gamma^a_{dm}\Gamma^m_{bc} - 2F''(B)\partial_c B Q_a{}^{bcd}\Gamma^a{}_{bcd} + \sqrt{-g}[F(B) - F'(B)B]\ , \\ &16\pi\mathcal{L}^F_{surf} = 2\partial_c[\sqrt{-g}F'(B)Q_a{}^{bcd}\Gamma^a_{bd}]\ . \end{aligned} \tag{2.11}$$

The Lagrangian density (2.11) is our starting point for the covariant Hamiltonian formulation of $F(R)-$gravity which will be analysed in the next section.

3. Covariant Hamiltonian Formalism with (g_{ab}, M^{abc}) as canonical variables

In this section we proceed to the covariant Hamiltonian formalism of $F(R)-$gravity when we consider g_{bc} as basic dynamical variable. Then, according to the basic principle of covariant field theory, the conjugate momentum M^{abc} is defined as variation of the bulk part of the action with respect to $\partial_a g_{bc}$. Explicitly, we know that quadratic part of Lagrangian density for $F(R)-$gravity has the form (2.11), or alternatively, using explicit form of $Q_a{}^{bcd}$ we have

$$\begin{aligned} &16\pi\mathcal{L}^F_{quad} = \sqrt{-g}F'(B)[\Gamma^h_{dk}\Gamma^k_{gh}g^{gd} - \Gamma^f_{fk}\Gamma^k_{gh}g^{gh}] - \\ &-F''(B)\sqrt{-g}\partial_c B[\Gamma^c_{bd}g^{bd} - g^{cb}\Gamma^d_{bd}] + \sqrt{-g}[F(B) - F'(B)B]\ . \end{aligned} \tag{3.1}$$

In order to define conjugate momenta M^{abc} we need following expression

$$\frac{\delta\Gamma^k_{bc}}{\delta\partial_p g_{rs}} = \frac{1}{4}\delta^p_b(g^{kr}\delta^s_c + g^{ks}\delta^r_c) + \frac{1}{4}\delta^p_c(g^{kr}\delta^s_b + g^{ks}\delta^r_b) - \frac{1}{4}g^{kp}(\delta^r_b\delta^s_c + \delta^r_c\delta^s_b)\ . \tag{3.2}$$

Using this result we get

$$16\pi M^{abc} = \frac{\partial\mathcal{L}^F_{quad}}{\partial(\partial_a g_{bc})} = \frac{1}{2}F'(B)\sqrt{-g}[g^{bd}\Gamma^a_{dk}g^{kc} + g^{cd}\Gamma^a_{dk}g^{kb}]$$

$$-\frac{1}{2}F'(B)\sqrt{-g}\Gamma^k_{fk}(g^{fb}g^{ac}+g^{fc}g^{ab})-\frac{1}{2}F'(B)\sqrt{-g}g^{bc}\Gamma^a_{gh}g^{gh}+\frac{1}{2}F'(B)\sqrt{-g}\Gamma^k_{fk}g^{fa}g^{bc}-$$
$$-\frac{1}{2}F''(B)\sqrt{-g}\partial_f B[g^{fb}g^{ac}+g^{fc}g^{ab}]+F''(B)\sqrt{-g}\partial_f B g^{fa}g^{bc} \tag{3.3}$$

and momentum conjugate to B as

$$16\pi p^c_B=\frac{\partial\mathcal{L}^F_{quadr}}{\partial(\partial_c B)}=-F''(B)\sqrt{-g}[\Gamma^c_{bd}g^{bd}-g^{cb}\Gamma^d_{bd}]\ . \tag{3.4}$$

To proceed further let us introduce $V^a\equiv -g_{bc}M^{abc}$ that, using (3.3) has the form

$$16\pi V^a=\sqrt{-g}F'(B)\Gamma^a_{gh}g^{gh}-F'(B)\sqrt{-g}\Gamma^k_{kf}g^{fa}-3F''(B)\sqrt{-g}\partial_f B g^{fa}\ . \tag{3.5}$$

From V^a we can express $\partial_f B$ as

$$\partial_f B=-\frac{16\pi g_{fa}}{3F''(B)\sqrt{-g}}\left(g_{bc}M^{abc}+\frac{F'(B)}{F''(B)}p^a_B\right)\ . \tag{3.6}$$

Then we can insert this result into definition of M^{abc} and find relation between M^{abc} and Γ^a_{bc}. However resulting expression is rather complicated and corresponding Hamiltonian as well. For that reason we rather focus on set of variables that were used in [13].

4. Hamiltonian Analysis in New Variables (f^{ab}, N^c_{ab})

We define new metric variable as

$$f^{ab}=\sqrt{-g}g^{ab}\ . \tag{4.1}$$

Even if we call it as a new one we should stress that it was used in classical literature long time ago [9, 10] and it was also used by P. Horava for covariant Hamiltonian formulation of gravity [13]. An importance of these variables was also stressed many times in works by Padmanabhan, see for example [11]. From definition of f^{ab} given in (4.1) we find that

$$f\equiv\det f^{ab}=\det g \tag{4.2}$$

and also we have following relation between variation of g^{ab} and f^{ab}

$$\delta g^{ab}=\frac{\delta f^{ab}}{\sqrt{-f}}-\frac{1}{2\sqrt{-f}}f^{ab}f_{mn}\delta f^{mn}\equiv\frac{1}{\sqrt{-f}}B^{ab}_{mn}\delta f^{mn}\ , \tag{4.3}$$

where

$$B^{ab}_{mn}=\frac{1}{2}(\delta^a_m\delta^b_n+\delta^b_m\delta^a_n)-\frac{1}{2}f^{ab}f_{mn}=\frac{1}{2}(\delta^a_m\delta^b_n+\delta^b_m\delta^a_n-g^{ab}g_{mn})\ . \tag{4.4}$$

Note that in our convention f_{ab} is inverse to f^{ab} and it has explicit form $f_{ab}=\frac{1}{\sqrt{-g}}g_{ab}$.

Now let us consider f^{ab} as canonical variable and introduce corresponding conjugate momenta N^c_{ab} as variation of the action with respect to $\partial_c f^{ab}$. Since boundary term does not contribute to this definition we have

$$N^c{}_{ab} = \frac{\partial L^F_{quadr}}{\partial(\partial_c f^{ab})} = \frac{\partial L^F_{quadr}}{\partial(\partial_d g_{mn})}\frac{\partial(\partial_d g_{mn})}{\partial(\partial_c f^{ab})} \ . \tag{4.5}$$

Now f^{ab} and g_{mn} are related by point transformations so that $f^{ab} = f^{ab}(g_{mn})$ or inverse $g_{mn} = g_{mn}(f^{ab})$. Then

$$\partial_d g_{mn} = \frac{\delta g_{mn}}{\delta f^{ab}}\partial_d f^{ab} \tag{4.6}$$

so that $\partial_d g_{mn} = \partial_d g_{mn}(f^{ab}, \partial_c f^{ab})$. Then clearly we have

$$\frac{\delta(\partial_d g_{mn})}{\delta(\partial_c f^{ab})} = \frac{\delta g_{mn}}{\delta f^{ab}}\delta^c_d \ . \tag{4.7}$$

Returning to $N^c{}_{ab}$ we obtain

$$N^c{}_{ab} = \frac{\partial \mathcal{L}^F_{quadr}}{\partial(\partial_c g_{mn})}(-g_{mk}\frac{1}{\sqrt{-f}}B^{kl}{}_{ab}g_{ln}) = -M^{cmn}\frac{1}{\sqrt{-f}}B_{mn,ab} \ , \tag{4.8}$$

where we used

$$\frac{\delta g_{mn}}{\delta f^{ab}} = -\frac{1}{\sqrt{-f}}g_{mk}B^{kl}{}_{ab}g_{ln} \tag{4.9}$$

and also we defined $B_{mn,ab}$ as

$$B_{mn,ab} = g_{mk}B^{kl}_{ab}g_{ln} = \frac{1}{2}(g_{ma}g_{nb} + g_{mb}g_{na} - g_{mn}g_{ab}) \ . \tag{4.10}$$

Since

$$\begin{aligned}
&16\pi M^{abc} = \frac{1}{2}F'(B)\sqrt{-g}[g^{bd}\Gamma^a_{dk}g^{kc} + g^{cd}\Gamma^a_{dk}g^{kb}] \\
&-\frac{1}{2}F'(B)\sqrt{-g}\Gamma^k_{fk}(g^{fb}g^{ac} + g^{fc}g^{ab}) - \frac{1}{2}F'(B)\sqrt{-g}g^{bc}\Gamma^a_{gh}g^{gh} + \frac{1}{2}F'(B)\sqrt{-g}\Gamma^k_{fk}g^{fa}g^{bc} - \\
&-\frac{1}{2}F''(B)\sqrt{-g}\partial_f B[g^{fb}g^{ac} + g^{fc}g^{ab}] + F''(B)\sqrt{-g}\partial_f B g^{fa}g^{bc}
\end{aligned} \tag{4.11}$$

we obtain from (4.8) following conjugate momenta N^c_{ab}

$$\begin{aligned}
&16\pi N^c_{ab} = -F'(B)\Gamma^c_{ab} + \frac{1}{2}F'(B)(\Gamma^k_{ak}\delta^c_b + \Gamma^k_{bk}\delta^c_a) + \\
&+\frac{1}{2}F''(B)(\partial_a B\delta^c_b + \partial_b B\delta^c_a) + \frac{1}{2}F''(B)\partial_g B f^{gc}f_{ab}
\end{aligned} \tag{4.12}$$

and

$$16\pi p^c_B = -F''(B)[\Gamma^c_{bd}f^{bd} - f^{cb}\Gamma^d_{bd}] \ . \tag{4.13}$$

It is crucial that for $F(B) = B, F'(B) = 1\ , F''(B) = 0$ we get that N^c_{ab} has the same form as in case of pure gravity and that $p^c_B = 0$. Further, if we take the trace $f^{ab}N^c_{ab}$ we obtain

$$16\pi f^{ab}N^c_{ab} = -F'(B)\Gamma^c_{ab}f^{ab} + F'(B)\Gamma^k_{kb}f^{bc} + 3F''(B)\partial_g B f^{gc}\ . \tag{4.14}$$

In case when $F'(B) = 1$ we get trivial identity while for $F''(B) \neq 0$ we can use previous equation and also definition of canonical momenta p^c_B given in (4.13) to express $\partial_a B$ as

$$16\pi f^{ab}N^c_{ab} = 16\pi\frac{F'(B)}{F''(B)}p^c_B + 3F''(B)\partial_g B f^{gc} \tag{4.15}$$

so that

$$\partial_g B = \frac{16\pi}{3F''(B)}f_{gc}\left(f^{ab}N^c_{ab} - \frac{F'(B)}{F''(B)}p^c_B\right)\ . \tag{4.16}$$

Inserting this result into (4.12) we obtain

$$\begin{aligned}
16\pi N^c_{ab} &= -F'(B)\Gamma^c_{ab} + \frac{1}{2}F'(B)(\Gamma^k_{ak}\delta^c_b + \Gamma^k_{bk}\delta^c_a) + \\
&+\frac{16\pi}{6}(f_{ad}f^{mn}N^d_{mn}\delta^c_b + f_{bd}f^{mn}N^d_{mn}\delta^c_a + f^{mn}N^c_{mn}f_{ab}) - \\
&-\frac{16\pi}{6}\frac{F'(B)}{F''(B)}(f_{ad}p^d_B + f_{bd}p^d_B + p^c_B f_{ab})
\end{aligned} \tag{4.17}$$

that allows us to write

$$16\pi\mathbf{N}^c_{ab} = -F'(B)\Gamma^c_{ab} + \frac{1}{2}F'(B)(\Gamma^k_{ak}\delta^c_b + \Gamma^k_{bk}\delta^c_a)\ , \tag{4.18}$$

where

$$\begin{aligned}
\mathbf{N}^c_{ab} &= N^c_{ab} - \frac{1}{6}(f_{ad}f^{mn}N^d_{mn}\delta^c_b + f_{bd}f^{mn}N^d_{mn}\delta^c_a + f^{mn}N^c_{mn}f_{ab}) + \\
&+\frac{1}{6}\frac{F'(B)}{F''(B)}(f_{ad}p^d_B\delta^c_b + f_{bd}p^d_B\delta^c_a + p^c_B f_{ab})\ .
\end{aligned} \tag{4.19}$$

As the next step we have to find inverse relation between Γ^c_{ab} and $\mathbf{N}^c_{ab}$. Following [11] we presume that it has the form

$$\Gamma^c_{ab} = a\mathbf{N}^c_{ab} + b(\mathbf{N}^d_{ad}\delta^c_b + \mathbf{N}^d_{bd}\delta^c_a)\ , \tag{4.20}$$

where a, b are unknown coefficients which we determine after inserting this ansatz into (4.18). Explicitly we get

$$\begin{aligned}
\frac{16\pi}{F'(B)}\mathbf{N}^c_{ab} &= -(a\mathbf{N}^c_{ab} + b(\mathbf{N}^d_{ad}\delta^c_b + \mathbf{N}^d_{bd}\delta^c_a)) \\
&+\frac{1}{2}[(a\mathbf{N}^d_{ad} + b(4\mathbf{N}^k_{ak} + \mathbf{N}^k_{ak}))\delta^c_b + (a\mathbf{N}^d_{bd} + b(\mathbf{N}^k_{bk} + 4\mathbf{N}^k_{bk})\delta^c_b)]\ .
\end{aligned} \tag{4.21}$$

Comparing left and right side of this expression we obtain following equations

$$a = -\frac{16\pi}{F'(B)} \ , \quad -b + \frac{(a+5b)}{2} = 0 \tag{4.22}$$

that has solution

$$b = -\frac{a}{3} \ . \tag{4.23}$$

Using these values of a and b we obtain inverse relation

$$\Gamma^c_{ba} = -\frac{16\pi}{F'(B)}\mathbf{N}^c_{ab} + \frac{16\pi}{3F'(B)}(\mathbf{N}^d_{ad}\delta^c_b + \mathbf{N}^d_{bd}\delta^c_a) \ . \tag{4.24}$$

Now the Hamiltonian density has the form

$$\begin{aligned}
&\mathcal{H}^F = \partial_c f^{ab} N^c_{ab} + p^c_B \partial_c B - \mathcal{L}^F_{quadr} = (\Gamma^d_{dc} f^{ab} - \Gamma^a_{cd} f^{db} - \Gamma^b_{dc} f^{da}) N^c_{ab} + p^c_B \partial_c B - \mathcal{L}^F_{quadr} = \\
&= \frac{1}{16\pi} F'(B)[\Gamma^h_{dk}\Gamma^k_{gh} f^{gd} - \Gamma^f_{fk}\Gamma^k_{gh} f^{gh}] + \\
&+ \frac{1}{16\pi} F''(B)\partial_f B[f^{fa}\Gamma^b_{ab} - \Gamma^f_{ab} f^{ab}] - \frac{1}{16\pi}\sqrt{-f}[F(B) - F'(B)B] \ ,
\end{aligned} \tag{4.25}$$

using

$$\partial_c f^{ab} = \partial_c \sqrt{-g} g^{ab} + \sqrt{-g}\partial_c g^{ab} = \Gamma^d_{dc} f^{ab} - \Gamma^a_{cd} f^{db} - \Gamma^b_{dc} f^{da} \ , \tag{4.26}$$

where we used the fact that $\nabla_c \sqrt{-g} = 0$ implies

$$\partial_c \sqrt{-g} = \Gamma^d_{dc}\sqrt{-g} \ . \tag{4.27}$$

In the same way the condition $\nabla_c g^{ab} = 0$ implies

$$\partial_c g^{ab} = -(\Gamma^a_{cd} g^{db} + \Gamma^b_{cd} g^{da}) \ . \tag{4.28}$$

Then inserting (4.24) into (4.25) we obtain Hamiltonian density in the form

$$\begin{aligned}
&\mathcal{H}_f = \frac{16\pi}{F'(B)}\mathbf{N}^h_{dk} f^{gd} \mathbf{N}^k_{gh} - \frac{16\pi}{3F'(B)}\mathbf{N}^m_{mk} f^{kg} \mathbf{N}^d_{dg} + \\
&+ \frac{16\pi}{3F''(B)} p^g_B f_{gc}(f^{ab} N^c_{ab} - \frac{F'(B)}{F''(B)} p^c_B) - \frac{1}{16\pi}\sqrt{-f}[F(B) - F'(B)B] \ .
\end{aligned} \tag{4.29}$$

Finally in order to complete Hamiltonian analysis we have to insert explicit form of $\mathbf{N}^c_{ab}$ given in (4.19) to (4.29). In fact, after some tedious calculations we find that the Hamiltonian density of $F(R)$ gravity has the form

$$\begin{aligned}
&\mathcal{H}^F = \frac{16\pi}{F'}[N^h_{kd} f^{dg} N^k_{gh} - \frac{1}{3} N^m_{mk} f^{kg} N^n_{ng}] + \\
&-.\frac{16\pi}{6} F'[\frac{1}{F''}p^a + \frac{1}{F'} N^a_{mn} f^{mn}] f_{ab} [\frac{1}{F''} p^b + \frac{1}{F'} f^{rs} N^b_{rs}] - \frac{1}{16\pi}\sqrt{-f}[F(B) - F'(B)B] \ .
\end{aligned} \tag{4.30}$$

This is the final form of the covariant Hamiltonian density for $F(R)-$gravity. In order to find corresponding equations of motion we begin with the canonical form of the action

$$S = \int d^4x (N^c_{ab}\partial_c f^{ab} + p^c_B \partial_c B - \mathcal{H}^F) \tag{4.31}$$

so that its variation has the form

$$\begin{aligned}\delta S = \int d^4x \Big(&\delta N^c_{ab}\partial_c f^{ab} + N^c_{ab}\partial_c \delta f^{ab} + \delta p^c_B \partial_c B + p^c_B \partial_c \delta B - \\ &- \frac{\delta \mathcal{H}^F}{\delta N^c_{ab}}\delta N^c_{ab} - \frac{\delta \mathcal{H}^F}{\delta f^{ab}}\delta f^{ab} - \frac{\delta \mathcal{H}^F}{\delta p^c_B}\delta p^c_B - \frac{\delta \mathcal{H}^F}{\delta B}\delta B\Big) = 0\end{aligned} \tag{4.32}$$

that implies following equations of motion

$$\begin{aligned}
&\partial_c f^{ab} = \frac{\delta \mathcal{H}^F}{\delta N^c_{ab}} = \frac{16\pi}{F'}\left[f^{bg}N^a_{gc} + f^{ag}N^b_{gc}\right] - \frac{16\pi}{3F'}\left[f^{bg}N^n_{ng}\delta^c_a + f^{ag}N^n_{ng}\delta^b_c\right] - \\
&-\frac{32\pi}{3}f^{ab}f_{ch}\left[\sqrt{-f}\frac{1}{F''}p^h + \frac{1}{F'}N^h_{rs}f^{rs}\right] \ , \\
&\partial_c N^c_{ab} = -\frac{\delta \mathcal{H}^F}{\delta f^{ab}} = -\frac{8\pi}{F'}\left[N^h_{ka}N^k_{bh} + N^h_{kb}N^h_{ad}\right] + \frac{16\pi}{3F'}N^m_{ma}N^n_{nb} + \\
&-\frac{8\pi F'}{3}\left[\frac{1}{F''}p^x + \frac{1}{F'}N^x_{mn}f^{mn}\right]f_{xa}\left[\frac{1}{F''}p^y + \frac{1}{F'}f^{ys}N^b_{rs}\right]f_{yb} + \\
&+\frac{16\pi}{3}\frac{F'}{F''}N^x_{ab}f_{xy}\left[\frac{1}{F''}p^y + \frac{1}{F'}N^y_{mn}f^{mn}\right] + \frac{1}{32\pi}f_{ab}\sqrt{-f}\left[F(B) - F'(B)B\right] \ , \\
&\partial_c B = \frac{\delta \mathcal{H}^F}{\delta p^c_B} = -\frac{8\pi}{3}\frac{F'}{F''}f_{ab}\left[\frac{1}{F''}p^b + \frac{1}{F'}N^b_{mn}f^{mn}\right] \ , \\
&\partial_c p^c_B = -\frac{\delta \mathcal{H}^F}{\delta B} = \frac{16\pi F''}{F'^2}\left[N^h_{kd}f^{dg}N^k_{gh} - \frac{1}{3}N^m_{mk}f^{kg}N^n_{ng}\right] + \\
&+\frac{8\pi}{3}F''\left[\frac{1}{F''}p^a + \frac{1}{F'}N^a_{mn}f^{mn}\right]f_{ab}\left[\frac{1}{F''}p^b + \frac{1}{F'}f^{rs}N^b_{rs}\right] + \\
&-\frac{16\pi}{3}\left[\frac{F'''}{F''^2}p^a + \frac{F''}{F'^2 N^a_{mn}f^{mn}}\right]f_{ab}\left[\frac{1}{F''}p^b + \frac{1}{F'}N^b_{rs}f^{rs}\right] - \frac{1}{16\pi}\sqrt{-f}F''(B)B \ .
\end{aligned} \tag{4.33}$$

Further, the boundary term for $F(R)-$gravity has the form

$$\mathcal{L}^F_{surf} = -\partial_c\left[\frac{F'}{F''}p^c_B\right] \tag{4.34}$$

which shows an importance of the field B and corresponding conjugate momenta p^c_B. On the other hand if we presume that the solution of the equation of motion for B is $B = \text{const}$ we find that p^b_B is equal $-\frac{F''}{F'}N^b_{mn}f^{mn}$. Inserting this result into (4.34) we find that it is equal to

$$\mathcal{L}^F_{surf} = \partial_c[N^c_{mn}f^{mn}] \tag{4.35}$$

which is the same as in case of pure gravity. Certainly it would be nice to study consequence of this result for thermodynamics aspects of $F(R)-$gravity.

Acknowledgement:
The work of JK is supported by the grant Integrable Deformations (GA20-04800S) from the Czech Science Foundation (GACR).

References

[1] Th. De Donder, *"Thorie Invariantive Du Calcul des Variations"*, (Gaulthier-Villars and Cie., Paris, 1930)

[2] H. Weyl, *"Geodesic Fields in the Calculus of Variation for Multiple Integrals"* Annals of Mathematics, **36** , p.607

[3] J. Struckmeier and A. Redelbach, *"Covariant Hamiltonian field theory,"* Int. J. Mod. Phys. E **17** (2008), 435-491 doi:10.1142/S0218301308009458 [arXiv:0811.0508 [math-ph]].

[4] I. V. Kanatchikov, *"Canonical structure of classical field theory in the polymomentum phase space,"* Rept. Math. Phys. **41** (1998), 49-90 doi:10.1016/S0034-4877(98)80182-1 [arXiv:hep-th/9709229 [hep-th]].

[5] M. Forger, C. Paufler and H. Roemer, *"The Poisson bracket for Poisson forms in multisymplectic field theory,"* Rev. Math. Phys. **15** (2003), 705-744 doi:10.1142/S0129055X03001734 [arXiv:math-ph/0202043 [math-ph]].

[6] H. Kastrup, *"Canonical Theories of Dynamical Systems in Physics,"* Phys. Rept. **101** (1983), 1 doi:10.1016/0370-1573(83)90037-6

[7] U. Lindstrm, *"Covariant Hamiltonians, sigma models and supersymmetry,"* [arXiv:2004.01073 [hep-th]].

[8] J. Klusoň, *"Note About Covariant Hamiltonian Formalism for Strings, p-Branes and Unstable Dp-Branes,"* [arXiv:2004.14654 [hep-th]].

[9] A. Eddington, *"The Mathematical Theory of Relativity:*, Cambridge University Press, Cambridge, UK, 2 ed., 1924.

[10] E. Schrodinger, *"Space-time Structure"*, Cambridge Science Classics. Cambridge University Press, Cambridge, UK, 1950.

[11] K. Parattu, B. R. Majhi and T. Padmanabhan, *""Structure of the gravitational action and its relation with horizon thermodynamics and emergent gravity paradigm,"* Phys. Rev. D **87** (2013) no.12, 124011 doi:10.1103/PhysRevD.87.124011 [arXiv:1303.1535 [gr-qc]].

[12] T.Padmanabhan, *"Gravitation: Foundations and Frontiers,"* Cambridge University Press, Cambridge, UK, 2010.

[13] P. Horava, *"On a covariant Hamilton-Jacobi framework for the Einstein-Maxwell theory,"* Class. Quant. Grav. **8** (1991), 2069-2084 doi:10.1088/0264-9381/8/11/016

[14] A. De Felice and S. Tsujikawa, *"f(R) theories,"* Living Rev. Rel. **13** (2010), 3 doi:10.12942/lrr-2010-3 [arXiv:1002.4928 [gr-qc]].

[15] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, *"Modified Gravity Theories on a Nutshell: Inflation, Bounce and Late-time Evolution,"* Phys. Rept. **692** (2017) 1 doi:10.1016/j.physrep.2017.06.001 [arXiv:1705.11098 [gr-qc]].

[16] S. Nojiri and S. D. Odintsov, *"Unified cosmic history in modified gravity: from F(R) theory to Lorentz non-invariant models,"* Phys. Rept. **505**, 59 (2011) doi:10.1016/j.physrep.2011.04.001 [arXiv:1011.0544 [gr-qc]].

[17] J. Klusoň, M. Oksanen and A. Tureanu, *"Hamiltonian analysis of curvature-squared gravity with or without conformal invariance,"* Phys. Rev. D **89** (2014) no.6, 064043 doi:10.1103/PhysRevD.89.064043 [arXiv:1311.4141 [hep-th]].

[18] S. Bahamonde, S. D. Odintsov, V. K. Oikonomou and M. Wright, *"Correspondence of $F(R)$ Gravity Singularities in Jordan and Einstein Frames,"* Annals Phys. **373** (2016) 96 doi:10.1016/j.aop.2016.06.020 [arXiv:1603.05113 [gr-qc]].

[19] S. Bahamonde, S. D. Odintsov, V. K. Oikonomou and P. V. Tretyakov, *"Deceleration versus acceleration universe in different frames of $F(R)$ gravity,"* Phys. Lett. B **766** (2017) 225 doi:10.1016/j.physletb.2017.01.012 [arXiv:1701.02381 [gr-qc]].

[20] G. G. L. Nashed, W. El Hanafy, S. D. Odintsov and V. K. Oikonomou, *"Thermodynamical correspondence of $f(R)$ gravity in Jordan and Einstein frames,"* arXiv:1912.03897 [gr-qc].