

Double-heavy tetraquark states with heavy diquark-antiquark symmetry

Jian-Bo Cheng¹, Shi-Yuan Li¹, Yan-Rui Liu¹, Zong-Guo Si¹, Tao Yao¹

¹*School of Physics, Shandong University, Jinan, Shandong, 250100, China*

(Dated: February 10, 2022)

We calculate the masses of the $QQ\bar{q}\bar{q}$ ($Q = c, b$; $q = u, d, s$) tetraquark states with the aid of heavy diquark-antiquark symmetry (HDAS) and the chromomagnetic interaction (CMI) model. The masses of the highest-spin ($J = 2$) tetraquarks that have only the $(QQ)_{\bar{3}_c}(\bar{q}\bar{q})_{3_c}$ color structure are related with those of conventional hadrons using HDAS. Thereafter, the masses of their partner states are determined with the mass splittings in the CMI model. Our numerical results reveal that: (i) the lightest $cc\bar{n}\bar{n}$ ($n = u, d$) is an $I(J^P) = 0(1^+)$ state around 3929 MeV (53 MeV above the DD^* threshold) and none of the double-charm tetraquarks are stable; (ii) the stable double-bottom tetraquarks are the lowest $0(1^+)$ $bb\bar{n}\bar{n}$ around 10488 MeV (≈ 116 MeV below the BB^* threshold) and the lowest $1/2(1^+)$ $bb\bar{n}\bar{s}$ around 10671 MeV (≈ 20 MeV below the BB_s^*/B_sB^* threshold); and (iii) the two lowest $bc\bar{n}\bar{n}$ tetraquarks, namely the lowest $0(0^+)$ around 7167 MeV and the lowest $0(1^+)$ around 7223 MeV, are near-threshold states. Moreover, we discuss the constraints on the masses of double-heavy hadrons. Specifically, for the lowest nonstrange tetraquarks, we obtain $T_{cc} < 3965$ MeV, $T_{bb} < 10627$ MeV, and $T_{bc} < 7199$ MeV.

I. INTRODUCTION

Recently, the LHCb Collaboration [1] observed the doubly charmed baryon Ξ_{cc}^{++} in the $\Lambda_c^+ K^- \pi^+ \pi^+$ mass distribution. Its mass was determined to be $3621.40 \pm 0.72(\text{stat.}) \pm 0.27(\text{syst.}) \pm 0.14(\Lambda_c^+)$ MeV/c². This value is 100 MeV higher than the mass of Ξ_{cc}^+ , which was determined in the channels $\Lambda_c^+ K^- \pi^+$ and $p D^+ K^-$ by the SELEX Collaboration [2, 3] more than fifteen years ago. The doubly heavy baryons Ξ_{cc}^{++} and Ξ_{cc}^+ have also been searched for in the FOCUS [4], BABAR [5], and Belle [6] detectors, with negative results. Thus far, the LHCb Collaboration has still not been able to confirm the Ξ_{cc}^+ baryon [7].

The confirmation of Ξ_{cc}^{++} has important implications as it indicates that two identical charm quarks can exist in a hadronic state. The observation of this baryon has motivated several theoretical discussions regarding the possible double-charm tetraquark T_{cc} and its partner states¹. In the literature, various approaches have been applied to the double-heavy tetraquark structures $QQ\bar{q}\bar{q}$ ($Q = c, b$; $q = u, d, s$), including the color-magnetic interaction (CMI) model [8–12], quark-level models [13–32], QCD sum rule method [33–40], lattice QCD simulation [41–48], and holographic model [49]. One may consult Ref. [50] for further discussions on such exotic states and related methods.

In Ref. [11], the $QQ\bar{q}\bar{q}$ states were systematically studied using a CMI model in which the color mixing effects between $(QQ)_{\bar{3}_c}(\bar{q}\bar{q})_{3_c}$ and $(QQ)_{6_c}(\bar{q}\bar{q})_{\bar{6}_c}$ structures were considered, and the thresholds of the meson-meson channels were treated as reference scales to estimate the tetraquark masses. According to a series of studies on multiquark states that used the above model [11, 51–59], it appears that the method that uses thresholds usually yields underestimated masses [50]. A possible reason for the underestimation is that the color-electric contribution to the two heavy quarks in the tetraquarks was not explicitly considered [19, 60]. Considering the color-Coulomb interaction, the binding energy of two heavy quarks exhibits a positive correlation with their reduced mass. When the two heavy quarks are separated by a large distance, the $QQ\bar{q}\bar{q}$ state will form a mixed $(Q\bar{q})_{1_c}(Q\bar{q})_{1_c}-(Q\bar{q})_{8_c}(Q\bar{q})_{8_c}$ meson-meson type structure. In contrast, if the two heavy quarks move in a small spatial region because of the attraction, they may form a $\bar{3}_c$ substructure and the tetraquark can be treated as a diquark-antidiquark state. In this case, it is not necessary for the distance between the two light antiquarks to be small due to the considerable relativistic effect and small color-Coulomb potential. In this study, we aimed to perform a further investigation of double-heavy tetraquark systems, particularly the nonstrange double-charm system.

The heavy quark flavor-spin symmetry appears in the limit $m_Q \rightarrow \infty$, and it is used extensively to study the properties of heavy quark hadrons. For states containing two heavy quarks, the heavy diquark-antiquark symmetry (HDAS) can also be considered [61–67]. According to this symmetry, the mass splittings between QQq baryons and those between $\bar{Q}'q$ mesons can be related with the correspondence $QQ \leftrightarrow \bar{Q}'$. The consideration is based on the observations that i) the size of QQ is small in the heavy quark limit, ii) the color representations of QQ and \bar{Q}' are both $\bar{3}_c$, and iii) the interaction between light and heavy components is suppressed, even though QQ and \bar{Q}' have different spins. Similarly, one can relate the double-heavy tetraquarks $QQ\bar{q}\bar{q}$, with the color structure $(QQ)_{3_c}$, to the

¹ We use T_{cc} to specifically denote the lowest $cc\bar{u}\bar{d}$ tetraquark state with $I(J^P) = 0(1^+)$ in this article. Similarly, T_{bb} represents the lowest $bb\bar{u}\bar{d}$ with $I(J^P) = 0(1^+)$. However, T_{bc} represents the lowest $bc\bar{u}\bar{d}$ with $I(J^P) = 0(0^+)$.

singly heavy antibaryons $\bar{Q}'\bar{q}\bar{q}$. According to the HDAS, for example, we can estimate the mass of a double-charm tetraquark with the relation $T_{cc\bar{n}\bar{n}} - \Xi_{cc}^* = \Sigma_b^* - \bar{B}^*$, where $n = u, d$, and the hadron symbol represents its mass. Obviously, the required unknown input is only the mass of Ξ_{cc}^* , which can be estimated with the experimental mass of Ξ_{cc}^{++} in the CMI model. Although the quantum numbers of LHCb Ξ_{cc}^{++} have not been measured, its mass is very close to the theoretical value of the ground state predicted by Karliner and Rosner in Ref. [68]. In the following calculations, we will use $m_{\Xi_{cc}} = 3621$ MeV as the input to estimate the masses of the double-charm tetraquark states. Other double-heavy tetraquarks will also be systematically investigated. If the LHCb Ξ_{cc} is actually the Ξ_{cc}^* state with spin=3/2, $m_{\Xi_{cc}^*} - m_{\Xi_{cc}} \approx 70$ MeV should be subtracted from the obtained masses of relevant tetraquarks.

Unlike the conventional hadrons, the QQ diquark in tetraquarks may also be in the color 6_c representation. In Ref. [9], the mass of $I(J^P) = 0(1^+) T_{cc}$ with $6_c cc$ was estimated. To include the mixing effects between the $(cc)_{\bar{3}_c}(\bar{n}\bar{n})_{3_c}$ and $(cc)_{6_c}(\bar{n}\bar{n})_{\bar{6}_c}$ configurations and to estimate the masses of all the $(cc\bar{n}\bar{n})$ states, in the current study, we first need to identify the position of the $(cc)_{\bar{3}_c}(\bar{n}\bar{n})_{3_c}$ state determined with HDAS. Observing that the $J = 2$ tetraquark is the pure $(cc)_{\bar{3}_c}(\bar{n}\bar{n})_{3_c}$ state because of the constraint from the Pauli principle, we directly relate its mass to that of Ξ_{cc}^* . Thereafter, we determine the masses of the lower tetraquark states from the mass splittings within the CMI model. Other double-heavy tetraquark states will be studied similarly. This concept is contrary to the estimation strategy adopted in our recent works [11, 51–59, 69] where the multiquark masses were determined from lower mass scales.

The remainder of this paper is organized as follows. In Sec. II, we present the method and formalism for the study. Thereafter, we provide our analysis and numerical results for the $cc\bar{u}\bar{d}$ states in Sec. III and the predictions on their partners in Sec. IV. In Sec. V, we discuss the constraints on the masses of the involved heavy quark hadrons. Finally, Sec. VI presents our discussions and a summary.

II. MODEL AND METHOD

A. CMI model

For ground state hadrons, the mass splittings of different spin states with the same quark content are mainly determined by the color-spin (color-magnetic) interaction in the quark model [70],

$$H_{CM} = - \sum_{i < j} C_{ij} \vec{\lambda}_i \cdot \vec{\lambda}_j \vec{\sigma}_i \cdot \vec{\sigma}_j. \quad (1)$$

Here, $i(j)$ represents the i th (j th) quark component of the tetraquark state, $\vec{\lambda}_i$ ($\vec{\lambda}_j$) is the vector containing the eight Gell-Mann matrices for the i th (j th) quark component, and $\vec{\sigma}_i$ ($\vec{\sigma}_j$) is the vector containing the three Pauli matrices for the i th (j th) quark component. It should be noted that $\vec{\lambda}_i$ ($\vec{\lambda}_j$) should be replaced with $-\vec{\lambda}_i^*$ ($-\vec{\lambda}_j^*$) if the quark component is an antiquark. The effective coupling parameters C_{ij} , which actually depend on the systems, include effects from the spatial wave function and the constituent quark masses. Thus, the mass formula in the CMI model is

$$M = \sum_i m_i + \langle H_{CM} \rangle, \quad (2)$$

where the effective mass of the i th quark m_i includes the constituent quark mass and contributions from other terms such as the color-Coulomb interaction and color confinement. In the following calculations, we will adopt the values of the parameters presented in Table I, which are determined from the masses of conventional hadrons.

TABLE I: Coupling parameters (units: MeV) extracted from conventional hadrons. The value of $C_{c\bar{b}}$ is estimated using the mass splitting in the Godfrey-Isgur model [71]. Approximations $C_{cc} = kC_{c\bar{c}}$, $C_{bb} = kC_{b\bar{b}}$, $C_{cb} = kC_{c\bar{b}}$, and $C_{s\bar{s}} = C_{ss}/k$ are adopted [72, 73], where $k \equiv C_{nn}/C_{n\bar{n}} \approx 2/3$. The effective quark masses determined from the masses of the ground hadrons [74] are $m_n = 361.8$ MeV, $m_s = 542.4$ MeV, $m_c = 1724.1$ MeV, and $m_b = 5054.4$ MeV.

$C_{nn} = 18.3$	$C_{ns} = 12.0$	$C_{nc} = 4.0$	$C_{nb} = 1.3$	$C_{ss} = 5.7$	$C_{sc} = 4.4$	$C_{sb} = 0.9$	$C_{cc} = 3.2$	$C_{bb} = 1.8$	$C_{cb} = 2.0$
$C_{n\bar{n}} = 29.9$	$C_{n\bar{s}} = 18.7$	$C_{n\bar{c}} = 6.6$	$C_{n\bar{b}} = 2.1$	$C_{s\bar{s}} = 9.3$	$C_{s\bar{c}} = 6.7$	$C_{s\bar{b}} = 2.3$	$C_{c\bar{c}} = 5.3$	$C_{b\bar{b}} = 2.9$	$C_{c\bar{b}} = 3.3$

The CMI model can provide relatively reasonable predictions for the mass splittings for various hadronic systems, but it is not good enough to estimate hadron masses because the effective quark masses have large uncertainties. Ref. [11] presented two methods for estimating the double-heavy tetraquark masses: one employs the mass formula (2) and the other uses the modified formula

$$M = (M_{threshold} - \langle H_{CM} \rangle_{threshold}) + \langle H_{CM} \rangle. \quad (3)$$

The first method, which uses the parameters in Table I, provides theoretical upper limits for the masses. The differences between these upper limits and the “realistic” masses would be very large for heavy quark multiquark states. This can be observed, for example, in the results for conventional hadrons and the $c\bar{c}s$ tetraquark states [51]. A possible means of remedying the deviations is to include a color-electric term appropriately in the Hamiltonian of the CMI model [68, 75]. The second method yields more reasonable results than the first one, but it suffers from the problem of selecting the reference scale. For example, the threshold of the $J/\psi\phi$ channel leads to a lower mass $X(4140)$ than that of $D_s^+D_s^-$ does [51]. If the state has a mixed structure of $(c\bar{s})_{8_c}(\bar{c}s)_{8_c}$ and $(c\bar{s})_{1_c}(\bar{c}s)_{1_c}$, where the separation between c and \bar{s} is small and the distance between $c\bar{s}$ and $\bar{c}s$ is large, the choice to use $D_s^{(*)+}D_s^{(*)-}$ as a reference system to estimate the tetraquark masses is more natural, even though the resulting tetraquark mass is probably still lower than the measured one. In the $cc\bar{n}\bar{n}$ case, the threshold that may be used is only for the $D^{(*)}D^{(*)}$ channel. However, when cc can be considered as a diquark with a small spatial separation, using such a threshold as a reference scale appears to not be a good choice. The T_{cc} mass (approximately 100 MeV below the DD^* threshold) will probably also be underestimated.

In the second method, better choices for estimating the tetraquark masses than hadron-hadron thresholds should exist. When estimating the tetraquark masses of $Qq\bar{Q}\bar{q}$ and $q_1q_2\bar{q}_3\bar{q}_4$ in Refs. [56, 69], we attempted to relate the reference scales to the $X(4140)$ mass. The obtained masses were higher than those with the meson-meson thresholds. In the current study, we examine the results with the aid of the heavy diquark-antiquark symmetry. At present, it is not clear which choice yields more realistic results. Hopefully, future measurements regarding the predicted tetraquark states can provide an answer to this. In the following discussions, to compare the results, we will refer to the methods corresponding to these three reference choices as the threshold, $X(4140)$, and HDAS approaches.

B. Diquark-antiquark symmetry

A diquark is generally assumed to be a color- $\bar{3}$ correlated quark-quark subsystem of a bound or resonant state. It shares certain similar properties with an antiquark, and diquark-antiquark symmetry (DAS) may exist, even though the diquark and antiquark have different masses and spins, and their dynamics are not necessarily the same. This symmetry works for hadrons that contain a heavy quark with a mass that is much larger than Λ_{QCD} . In this case, the heavy quark can be treated as a static color source, and many properties are independent of the quark mass. For example, if we treat the $\bar{n}\bar{n}$ antidiquark as a heavier light quark n' , a $cc\bar{n}\bar{n}$ state will become a ccn' state, which resembles a baryon structure. Subsequently, the mass difference between the ccn' tetraquark and the ccn baryon is independent of the heavy quark mass in the heavy quark limit. This similarity based on DAS provides a method for estimating the masses of unknown states from those of known baryons. Likewise, as discussed by Savage and Wise in [64], if the cc heavy diquark is treated as an antiquark \bar{Q}' , the masses of various $\bar{Q}'\bar{n}\bar{n}$ tetraquarks can be related to those of Qnn baryons. As DAS is an approximate symmetry, whether or not the predictions based on it are correct needs to be tested experimentally. Before estimating the tetraquark masses, we provide further explanations of the diquark-antiquark symmetry.

It is known that there are three types of light quarks (u , d , and s) which form the base representation of flavor $SU(3)$. As quarks have spin, the flavor-spin $SU(6)$ can conventionally be used to classify various quark states. If a light diquark is a stable object, it has been argued that a symmetry exists between the baryons and mesons (diquarks and antiquarks) [76–78]. Considering the color- $\bar{3}$ diquark and antiquark together, we obtain a flavor-spin 27-plet. Its $SU(3) \otimes SU(2)$ decomposition reads $27 = (6_f, 3_s) + (\bar{3}_f, 1_s) + (\bar{3}_f, 2_s)$. The substructure $(6_f, 3_s)$ represents an 18-plet with flavor-symmetric and spin-symmetric diquarks. Similarly, $(\bar{3}_f, 1_s)$ represents a triplet with flavour-antisymmetric and spin-antisymmetric diquarks. The last substructure $(\bar{3}_f, 2_s)$ represents the antiquark sextet. The decomposition indicates that the diquark and antiquark can be combined into the $SU(6/21)$ symmetry algebra. We present the group structures in Fig. 1.

Lattice QCD simulations have indicated such a diquark-antiquark symmetry [79, 80]: the static quark-diquark potential is almost equal to the static quark-antiquark potential, and a quark-antiquark pair and a quark-diquark pair have similar wave functions. However, according to Ref. [61], the diquark-antiquark symmetry in light quark sector is broken for at least three reasons: (1) a diquark and an antiquark have different masses, lead to kinematical differences; (2) the diquark and antiquark have different spin-dependent and velocity-dependent terms; and (3) the diquark is not a point particle, and its finite size must affect its interactions. Thus, the breaking effects for the diquark-antiquark symmetry between the light diquarks and the light antiquarks are significant.

In contrast, for hadrons containing one heavy quark, the aforementioned symmetry breaking effects will be largely suppressed [61]. According to the heavy quark effective theory (HQET) [81], the kinematic and spin-dependent terms are inversely proportional to the heavy quark mass and make small contributions to the hadron mass. The size of the diquark is not a major concern, because the constituent quark model is still successful in handling the properties of conventional hadrons, even though the constituent quark has a comparable size to the diquark [61]. Thus, the size




FIG. 1: 27-multiplet in $SU(6)$ group with diquark-antiquark symmetry. The first number in the parentheses denotes the flavor $SU(3)$ representation and the second the spin $SU(2)$ representation.

of diquark has no significant impact on the DAS. Therefore, in general, there is possibly a better diquark-antiquark symmetry for hadrons containing one heavy quark. In Ref. [82], Lichtenberg, Roncaglia, and Predazzi analyzed relations for the masses of heavy quark hadrons using the Feynman-Hellmann theorem and semiempirical formulas. They obtained some mass sum rules for heavy quark hadrons. We have listed a selection of these as follows:

$$D_s^* - D^* = \bar{B}_s^* - \bar{B}^*, \quad (4)$$

$$\Sigma_b^* - \Sigma_c^* = \bar{B}^* - D^*, \quad (5)$$

$$\Xi_b^* - \Xi_c^* = \bar{B}^* - D^*, \quad (6)$$

$$\Omega_b^* - \Omega_c^* = \bar{B}^* - D^*, \quad (7)$$

which are adopted in the following discussions. These also follow from the heavy quark flavor symmetry. These relations can be confirmed by observing that the values of (*l.h.s.* - *r.h.s.*) are 12.9 MeV, -0.6 MeV, -9.5 MeV, and -21.5 MeV ($\Omega_b^* - \Omega_b = 14.4$ MeV is used in the CMI model), respectively. The common feature of these four relations is that only highest spins are involved. In fact, the final three relations also satisfy the diquark-antiquark symmetry for light quarks in which the diquark spin is 1. It is true that a better DAS exists for hadrons containing one heavy quark than for those without heavy quarks. Although more relations can be found in Ref. [82], it is not necessary to consider them in this work.

Our strategy for estimating the masses of double-heavy tetraquarks is to combine HDAS and the aforementioned four mass sum rules. To illustrate the concept, we temporarily focus only on Eq. (4). If we consider the heavy diquark-antiquark symmetry for the cc diquark, we obtain

$$\Omega_{cc}^* - \Xi_{cc}^* = \bar{B}_s^* - \bar{B}^* (= D_s^* - D^*), \quad (8)$$

which can be used to estimate the mass of Ω_{cc}^* with that of Ξ_{cc}^* . As a better heavy quark symmetry exists in bottom systems than in charmed systems, we use the masses of the bottom mesons. Here and in the following discussions, we only consider the highest spin hadrons while adopting the diquark-antiquark symmetry, because then, the possible contributions from other color or spin structures will be avoided. As explained in Sec. I, we assume that the spin of the LHCb Ξ_{cc} is 1/2. We can evaluate the mass of Ξ_{cc}^* with the CMI model $\Xi_{cc}^* = \Xi_{cc} + 16C_{cn} = 3685$ MeV. Thereafter, the mass of Ω_{cc}^* , namely 3776 MeV, is obtained. By using the CMI model again, we can further obtain $\Omega_{cc} = 3706$ MeV.

At present, doubly heavy baryons other than Ξ_{cc} have not been observed, and the accuracy of Eq. (8) cannot be verified. However, in theory, the reasonability of treating a QQ diquark as a heavy antiquark \bar{Q}' can be argued. Because the heavy diquark has lower kinematic energy and less spin-dependent interaction, the heavy quark approximation works better than the single Q case. Furthermore, the heavy diquark (with a light quark spectator) actually has better symmetry properties than the light diquark (with a heavy quark spectator) because a heavy diquark has small spatial separation, which means that its interaction with light quarks is not affected significantly by its size. In the following parts of this paper, we first focus on the double-charm tetraquark $cc\bar{n}\bar{n}$ states by treating them as systems that are composed of a small-size double-charm diquark and a light antidiquark. According to HDAS, the study of $cc\bar{n}\bar{n}$ states becomes that of heavy $Q'nn$ “baryons”. This approximate symmetry, together with the above mass sum rules, is evidently convenient for us to relate $cc\bar{n}\bar{n}$ to Ξ_{cc} . Focusing on the $cc\bar{n}\bar{n}$ states, we perform similar studies on other double-heavy tetraquarks. Whether or not the adopted approximation is effective should be tested in future experiments.

III. $cc\bar{n}\bar{n}$ SPECTRUM IN TERMS OF KNOWN Ξ_{cc}

Based on the symmetry consideration, we obtain a good flavor-spin supermultiplet [62] that contains both tetraquark mesons $cc\bar{q}\bar{q}$ and three-quark baryons ccq . This double-charm supermultiplet is classified into three types of states,

$$6_{fs} : ccu, \quad ccd, \quad ccs; \quad (9)$$

$$18_{fs} : cc\bar{u}\bar{u}, \quad cc\bar{u}\bar{d}, \quad cc\bar{d}\bar{d}, \quad cc\bar{u}\bar{s}, \quad cc\bar{d}\bar{s}, \quad cc\bar{s}\bar{s}; \quad (10)$$

$$3_{fs} : cc\bar{u}\bar{d}, \quad cc\bar{u}\bar{s}, \quad cc\bar{d}\bar{s}. \quad (11)$$

The members in Eq. (10) contain a light antiquark with spin 1, and those in (11) have a light antiquark with spin 0. We treat the cc diquark as a heavy $\bar{3}_c$ “antiquark” \bar{Q}' . Following the replacement, $cc \rightarrow \bar{Q}'$, the Ξ_{cc}^* with spin 3/2 and the $cc\bar{n}\bar{n}$ with spin 2 are transformed into $\bar{M}_{Q'}^* = (\bar{Q}'n)^{J=3/2}$ and $\bar{\Sigma}_{Q'}^* = (\bar{Q}'\bar{n}\bar{n})^{J=2}$, respectively, i.e.

$$\bar{\Xi}_{cc}^*(\bar{c}\bar{c}\bar{n}) \rightarrow \bar{M}_{Q'}^*(Q'\bar{n}), \quad (12)$$

$$T_{\bar{c}\bar{c}nn}^{I=1, J=2} \rightarrow \bar{\Sigma}_{Q'}^*(Q'nn). \quad (13)$$

Note that only the highest spins are involved, and the cc diquark has unique quantum numbers $color = \bar{3}_c, spin = 1$. The forms on the right hand side remind us of the relation in Eq. (5), where the light diquark-antiquark symmetry is used. With that equation, one can naturally obtain

$$T_{cc\bar{n}\bar{n}}^{I=1, J=2} - \bar{\Xi}_{cc}^* = \Sigma_c^* - D^* = \Sigma_b^* - \bar{B}^*. \quad (14)$$

The obtained mass of $cc\bar{n}\bar{n}$ with $I = 1, J = 2$ is, thus, 4195 MeV (with Σ_c^* and D^*) or 4194 MeV (with Σ_b^* and \bar{B}^*). We select the later value owing to the better heavy quark symmetry for the bottom hadrons. Subsequently, we have a good reference hadron $T_{cc\bar{n}\bar{n}}^{I=1, J=2}$ and can estimate the masses of other $cc\bar{n}\bar{n}$ states by considering the CMI differences. When only $\bar{3}_c$ cc is considered, the masses of the other three double-charm tetraquarks are $T_{cc\bar{n}\bar{n}}^{I=1, J=0} = 4087$ MeV, $T_{cc\bar{n}\bar{n}}^{I=1, J=1} = 4122$ MeV, and $T_{cc\bar{n}\bar{n}}^{I=0, J=1} = 3961$ MeV. It is obvious that these tetraquarks are all above the DD^* threshold (3876 MeV).

TABLE II: Color-spin bases for $cc\bar{q}\bar{q}$ ($q = u, d, s$) states [11]. The superscripts indicate the spin, and the subscripts indicate the color representations.

States	$I(J^P)$	Bases
$(cc\bar{n}\bar{n})$	$1(2^+)$	$[(cc)_{\frac{1}{3}}(\bar{n}\bar{n})_{\frac{1}{3}}]^2$
	$1(1^+)$	$[(cc)_{\frac{1}{3}}(\bar{n}\bar{n})_{\frac{1}{3}}]^1$
	$1(0^+)$	$[(cc)_{\frac{1}{3}}(\bar{n}\bar{n})_{\frac{1}{3}}]^0 \quad [(cc)_6^0(\bar{n}\bar{n})_6^0]$
	$0(1^+)$	$[(cc)_{\frac{1}{3}}(\bar{n}\bar{n})_{\frac{1}{3}}]^1 \quad [(cc)_6^0(\bar{n}\bar{n})_6^1]$
$(cc\bar{n}\bar{s})$	(2^+)	$[(cc)_{\frac{1}{3}}(\bar{n}\bar{s})_{\frac{1}{3}}]^2$
	(1^+)	$[(cc)_{\frac{1}{3}}(\bar{n}\bar{s})_{\frac{1}{3}}]^1 \quad [(cc)_{\frac{1}{3}}(\bar{n}\bar{s})_{\frac{1}{3}}]^0 \quad [(cc)_6^0(\bar{n}\bar{s})_6^1]$
	(0^+)	$[(cc)_{\frac{1}{3}}(\bar{n}\bar{s})_{\frac{1}{3}}]^0 \quad [(cc)_6^0(\bar{n}\bar{s})_6^0]$
$(cc\bar{s}\bar{s})$	(2^+)	$[(cc)_{\frac{1}{3}}(\bar{s}\bar{s})_{\frac{1}{3}}]^2$
	(1^+)	$[(cc)_{\frac{1}{3}}(\bar{s}\bar{s})_{\frac{1}{3}}]^1$
	(0^+)	$[(cc)_{\frac{1}{3}}(\bar{s}\bar{s})_{\frac{1}{3}}]^0 \quad [(cc)_6^0(\bar{s}\bar{s})_6^0]$

Although the color structure of Ξ_{cc} is unique, that of the exotic $cc\bar{n}\bar{n}$ states is not. The mixing or channel coupling effects from the $(cc)_{6_c}(\bar{n}\bar{n})_{\bar{6}_c}$ color structure may significantly change the tetraquark masses. When such contributions are considered, two more tetraquarks appear. We collect all the color-spin bases [11] for the $cc\bar{n}\bar{n}$ states, which are displayed in Table II. With these wave functions, we finally obtain the numerical results listed in the fourth column of Table III and illustrated in Fig. 2(a). Moreover, we provide the masses estimated using Eq. (3) ($D^{(*)}D^{(*)}$ as the reference state) and those using Eq. (2) in the table. They can be viewed as the theoretical lower (fifth column) and upper (sixth column) limits, respectively, in the present framework. Comparing these three results from different considerations, it is found that the new masses fall within the range constrained by the lower and upper limits, and the values are slightly larger than the averages of the two limits. The masses of the lowest $1(0^+)$ and $0(1^+)$ states in Table III are evidently smaller than the above values without the $(cc)_{6_c}(\bar{n}\bar{n})_{\bar{6}_c}$ contributions. However, the T_{cc} state, the mass of which in the HDAS approach is 155 MeV higher than the lower limit (3774 MeV), is still above the DD^* threshold. The other $cc\bar{n}\bar{n}$ states are also above the respective fall-apart thresholds. Therefore, the HDAS approach results in an unstable T_{cc} , which is consistent with the conclusion obtained in Refs. [19, 20, 30, 31, 33, 39, 44, 45, 63, 83, 84].

Because the mass of Ω_b^* has not yet been measured, we opt to use the masses of Ω_c^* and D_s^* . With $\Xi_{cc}^* = 3685$ MeV and $\Omega_{cc}^* = 3776$ MeV, estimated using Eq. (8), we obtain $T_{cc\bar{n}\bar{s}}^{J=2} = 4313$ MeV and $T_{cc\bar{s}\bar{s}}^{J=2} = 4430$ MeV. These will be treated as reference scales to determine the masses of other double-charm strange tetraquarks.

Before proceeding further, we investigate the masses of $T_{cc\bar{n}\bar{n}}^{J=2}$, $T_{cc\bar{n}\bar{s}}^{J=2}$, and $T_{cc\bar{s}\bar{s}}^{J=2}$ in different approaches. The current estimation yields 177 MeV, 193 MeV, and 205 MeV for the mass distances measured from the D^*D^* , $D^*D_s^*$, and $D_s^*D_s^*$ thresholds, respectively, which are gradually increasing numbers. Those in Ref. [11] are gradually decreasing numbers, 23 MeV, 8 MeV, and -6 MeV. Therefore, no stable $J = 2$ tetraquarks are obtained in this study, whereas the states in Ref. [11] are around their fall-apart thresholds. This feature is an apparent difference between the HDAS approach and the threshold approach.

With the above reference states, $T_{cc\bar{n}\bar{s}}^{J=2}$ and $T_{cc\bar{s}\bar{s}}^{J=2}$, and the mass splittings in the CMI model, we can estimate the masses of strange partners of $cc\bar{n}\bar{n}$ states. The base structures for the calculation are presented in Table II. We list the numerical results for all of the $(cc\bar{n}\bar{s})$ and $(cc\bar{s}\bar{s})$ states in Table IV, where we also display the theoretical lower and upper limits for the tetraquark masses. Comparing the values in 4th, 5th, and 6th columns, it is obvious that our results with diquark-antiquark symmetry are slightly larger than the averages of the two limits, which is the same feature as in the $cc\bar{n}\bar{n}$ case. The relative positions for the $(cc\bar{n}\bar{s})$ and $(cc\bar{s}\bar{s})$ tetraquark states are illustrated in Fig. 2(b) and Fig. 2(c), respectively. According to the figure, similar to the $cc\bar{n}\bar{n}$ case, all the obtained double-charm strange tetraquarks can decay through rearrangement mechanisms, and no such stable states exist. This observation is different from that in Ref. [11], where the lowest 1^+ $cc\bar{n}\bar{s}$ is stable.

TABLE IV: Results for $cc\bar{n}\bar{s}$ and $cc\bar{s}\bar{s}$ states (unit: MeV). The second and third columns provide the numerical values of the CMI matrices and their eigenvalues, respectively. The fifth and sixth columns list the masses estimated using Eq. (3) (with $D^{(*)}D_s^{(*)}/D_s^{(*)}D_s^{(*)}$ as the reference state) and Eq. (2) (parameters provided in Table I), respectively. These can be viewed as the theoretical lower limits (low.) and upper limits (up.), respectively, for the tetraquark masses in the current framework. The fourth column displays our predictions with the heavy diquark-antiquark symmetry (HDAS) consideration.

$(cc\bar{n}\bar{s})$ J^P	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $D^{(*)}D_s^{(*)}$	Mass (up.) Eq. (2)
2^+	$\begin{pmatrix} 76.1 \end{pmatrix}$	$\begin{pmatrix} 76.1 \end{pmatrix}$	$\begin{pmatrix} 4313.3 \end{pmatrix}$	$\begin{pmatrix} 4125.4 \end{pmatrix}$	$\begin{pmatrix} 4428.5 \end{pmatrix}$
1^+	$\begin{pmatrix} 5.2 & 0.0 & 0.0 \\ 0.0 & -87.3 & -75.2 \\ 0.0 & -75.2 & -3.0 \end{pmatrix}$	$\begin{pmatrix} -131.4 \\ 5.2 \\ 41.1 \end{pmatrix}$	$\begin{pmatrix} 4105.8 \\ 4242.4 \\ 4278.3 \end{pmatrix}$	$\begin{pmatrix} 3917.9 \\ 4054.5 \\ 4090.4 \end{pmatrix}$	$\begin{pmatrix} 4221.0 \\ 4357.6 \\ 4393.5 \end{pmatrix}$
0^+	$\begin{pmatrix} -30.3 & 130.3 \\ 130.3 & 61.0 \end{pmatrix}$	$\begin{pmatrix} -122.7 \\ 153.4 \end{pmatrix}$	$\begin{pmatrix} 4114.5 \\ 4390.6 \end{pmatrix}$	$\begin{pmatrix} 3926.6 \\ 4202.7 \end{pmatrix}$	$\begin{pmatrix} 4229.7 \\ 4505.8 \end{pmatrix}$
$(cc\bar{s}\bar{s})$ J^P	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $D_s^{(*)}D_s^{(*)}$	Mass (up.) Eq. (2)
2^+	$\begin{pmatrix} 59.6 \end{pmatrix}$	$\begin{pmatrix} 59.6 \end{pmatrix}$	$\begin{pmatrix} 4429.9 \end{pmatrix}$	$\begin{pmatrix} 4211.6 \end{pmatrix}$	$\begin{pmatrix} 4592.6 \end{pmatrix}$
1^+	$\begin{pmatrix} -11.9 \end{pmatrix}$	$\begin{pmatrix} -11.9 \end{pmatrix}$	$\begin{pmatrix} 4358.4 \end{pmatrix}$	$\begin{pmatrix} 4140.1 \end{pmatrix}$	$\begin{pmatrix} 4521.1 \end{pmatrix}$
0^+	$\begin{pmatrix} -47.6 & 131.3 \\ 131.3 & 35.8 \end{pmatrix}$	$\begin{pmatrix} -143.7 \\ 131.8 \end{pmatrix}$	$\begin{pmatrix} 4226.6 \\ 4502.1 \end{pmatrix}$	$\begin{pmatrix} 4008.3 \\ 4283.8 \end{pmatrix}$	$\begin{pmatrix} 4389.3 \\ 4664.8 \end{pmatrix}$

B. $bb\bar{q}\bar{q}$ and $bc\bar{q}\bar{q}$ states

According to the diquark-antiquark symmetry, no stable double-charm tetraquark states exist. In the bottom case, the attractive color-Coulomb interaction between the two heavy quarks may be strong enough to aid in the formation of stable tetraquarks. Next, we investigate the $bb\bar{q}\bar{q}$ and $bc\bar{q}\bar{q}$ systems, where $q = u, d$, or s . First, we focus on the double-bottom tetraquarks, which have exactly the same group structure as the double-charm states. For the HDAS relations and the wave function bases, we simply need to perform a simple substitution of (bb) for (cc) in Eqs. (14)-(16) and Table II. However, considerable difficulty arises in applying the formulas, as the masses of Ξ_{bb}^* and Ω_{bb}^* have not been measured. In this study, we need to select appropriate predictions for their values from various investigations.

In the literature, numerous analyses on the masses of Ξ_{bb} and Ξ_{bb}^* have been performed (see Table I of Ref. [85] for a collection). We list several of the results in Table V, where the involved approaches include lattice QCD [86], chromomagnetic models [60, 68, 87], relativistic quark model [88], nonrelativistic quark model [89–91], bag model [92], and Bethe-Salpeter equation [93]. To select an appropriate value for the mass of Ξ_{bb} , we adopt the following

TABLE V: Theoretical predictions for the masses of doubly heavy baryons (unit: MeV) in various approaches: lattice QCD [86], chromomagnetic models [60, 68, 87], relativistic quark model [88], nonrelativistic quark model [89–91], bag model [92], Bethe-Salpeter equation [93], and QCD sum rules [109, 110].

	Ref. [86]	Ref. [60]	Ref. [88]	Ref. [89]	Ref. [90]	Ref. [91]	Refs. [68, 87]	Ref. [92]	Ref. [93]	Ref. [109, 110]
Ξ_{bb}	10143(30)(23)	10168.9 \pm 9.2	10202	10340	10197 $^{+10}_{-17}$	10204	10162 \pm 12	10272	10090 \pm 10	10170 \pm 140
Ξ_{bb}^*	10178(30)(24)	10188 \pm 7.1	10237	10367	10236 $^{+9}_{-17}$	-	10184 \pm 12	-	10337	10220 \pm 150
Ω_{bb}	10273(27)(20)	10259.0 \pm 15.5	10359	10454	10260 $^{+14}_{-34}$	10258	10208 \pm 18	10369	10180 \pm 5	10320 \pm 140
Ω_{bb}^*	10308(27)(21)	10267.5 \pm 12.1	10389	10486	10297 $^{+5}_{-28}$	-	-	10429	-	10380 \pm 140
Ξ_{bc}	6943(33)(28)	6922.3 \pm 6.9	6933	7011	6919 $^{+17}_{-7}$	6932	6914 \pm 13	6838	6840 \pm 10	-
Ξ'_{bc}	6959(36)(28)	6947.9 \pm 6.9	6963	7047	6948 $^{+17}_{-6}$	-	6933 \pm 12	7028	-	-
Ξ_{bc}^*	6985(36)(28)	6973.2 \pm 5.5	6980	7074	6986 $^{+14}_{-5}$	-	6960 \pm 14	6986	-	-
Ω_{bc}	6998(27)(20)	7010.7 \pm 9.3	7088	7136	6986 $^{+27}_{-17}$	6996	6968 \pm 19	6941	6945 \pm 5	-
Ω'_{bc}	7032(28)(20)	7047.0 \pm 9.3	7116	7165	7009 $^{+24}_{-15}$	-	6984 \pm 19	7116	-	-
Ω_{bc}^*	7059(28)(21)	7065.7 \pm 7.5	7130	7187	7046 $^{+11}_{-9}$	-	-	7077	-	-

criteria: 1) the baryon masses satisfy the light-flavor symmetry $\Omega_{bb}^* - \Omega_{bb} = \Xi_{bb}^* - \Xi_{bb}$ in the heavy quark limit; 2) the HDAS relation $\Omega_{bb}^* - \Xi_{bb}^* = \bar{B}_s^* - \bar{B}^* \approx 91$ MeV holds; and 3) the inequality $\Xi_{bb} - (\Xi_{bb})_{CMI} - 2(\bar{B} - \bar{B}_{CMI}) < \Xi_{cc} - (\Xi_{cc})_{CMI} - 2(D - D_{CMI})$ for the mass of Ξ_{bb} is required. The final criterion leading to $\Xi_{bb} < 10327$ MeV means that the color-Coulomb contribution to the bottom diquark is larger than that of the charm case once the contributions from the effective quark masses and color-magnetic interactions have been subtracted. It can be confirmed that a similar inequality holds for heavy quarkonia, $\eta_c - (\eta_c)_{CMI} - 2(D - D_{CMI}) > \eta_b - (\eta_b)_{CMI} - 2(\bar{B} - \bar{B}_{CMI})$ (numerically, -882 MeV $>$ -1181 MeV). The lattice results in Ref. [86] meet the first and third criteria but not the second because $\Omega_{bb}^* - \Xi_{bb}^* \approx 130$ MeV $>$ 91 MeV. In the chromomagnetic models [60, 68, 87], the results are compatible with all the criteria. In this case, we use the ground baryon mass $\Xi_{bb} = 10169$ MeV from Ref. [60], whereas the mass of Ξ_{bb}^* is evaluated to be $\Xi_{bb}^* = \Xi_{bb} + 16C_{bn} = 10190$ MeV. The mass of Ω_{bb}^* is, subsequently, further determined to be $\Omega_{bb}^* = \Xi_{bb}^* + \bar{B}_s^* - \bar{B}^* = 10280$ MeV and that of Ω_{bb} is $\Omega_{bb}^* - 16C_{bs} = 10266$ MeV.

By repeating the procedure for studying the double-charm tetraquarks, we similarly obtain the masses of the highest-spin double-bottom tetraquark states,

$$\begin{aligned}
T_{bb\bar{n}\bar{n}}^{I=1, J=2} &= \Xi_{bb}^* + \Sigma_b^* - \bar{B}^* = 10699 \text{ MeV}, \\
T_{bb\bar{n}\bar{s}}^{J=2} &= \Xi_{bb}^* + \Xi_b^* - \bar{B}^* = 10818 \text{ MeV}, \\
T_{bb\bar{s}\bar{s}}^{J=2} &= \Omega_{bb}^* + \Omega_b^* - \bar{B}_s^* = 10926 \text{ MeV}.
\end{aligned} \tag{17}$$

These values are 49 MeV, 78 MeV, and 95 MeV (increasing numbers) higher than the $\bar{B}^*\bar{B}^*$, $\bar{B}^*\bar{B}_s^*$, and $\bar{B}_s^*\bar{B}_s^*$ thresholds, respectively. Such mass distances from the corresponding thresholds in Ref. [11] are approximately 45 MeV, 29 MeV, and 13 MeV (decreasing numbers). With the newly obtained masses of the spin-2 tetraquarks, we can further estimate those of the lower double-bottom states in the CMI model. We list all the results in Table VI and plot the relative positions for the $bb\bar{q}\bar{q}$ tetraquarks in Fig. 3.




FIG. 3: Relative positions for double-bottom tetraquark states (solid and dashed lines) and relevant meson-meson thresholds (dotted lines). The masses are given in MeV. In (a), the solid (dashed) lines denote the $I = 1$ ($I = 0$) states.

By comparing the current results with those in Ref. [11], a similar $bb\bar{n}\bar{n}$ spectrum can be found. According to Fig. 3(a), the T_{bb} state with a mass of 10488 MeV is approximately 116 MeV below the $\bar{B}\bar{B}^*$ threshold (10604 MeV), and it should be rather stable, but other $bb\bar{n}\bar{n}$ states are not. This observation is the same as that in Ref. [11]. For the $bb\bar{n}\bar{s}$

TABLE VI: Results for the $bb\bar{q}\bar{q}$ ($q = u, d, s$) states (unit: MeV). The second and third columns provide the numerical values of the CMI matrices and their eigenvalues, respectively. The fifth and sixth columns list the masses estimated using Eq. (3) (with $\bar{B}^{(*)}\bar{B}^{(*)}/\bar{B}^{(*)}\bar{B}_s^{(*)}/\bar{B}_s^{(*)}\bar{B}_s^{(*)}$ as the reference state) and Eq. (2) (parameters provided in Table I), respectively. They can be viewed as the theoretical lower limits (low.) and upper limits (up.), respectively, for the tetraquark masses in the current framework. The fourth column displays our predictions with the heavy diquark-antiquark symmetry (HDAS) consideration.

$(bb\bar{n}\bar{n})$ $I(J^P)$	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $\bar{B}^{(*)}\bar{B}^{(*)}$	Mass (up.) Eq. (2)
$1(2^+)$	$\begin{pmatrix} 64.7 \end{pmatrix}$	$\begin{pmatrix} 64.7 \end{pmatrix}$	$\begin{pmatrix} 10698.7 \end{pmatrix}$	$\begin{pmatrix} 10691.3 \end{pmatrix}$	$\begin{pmatrix} 10897.1 \end{pmatrix}$
$1(1^+)$	$\begin{pmatrix} 42.3 \end{pmatrix}$	$\begin{pmatrix} 42.3 \end{pmatrix}$	$\begin{pmatrix} 10676.3 \end{pmatrix}$	$\begin{pmatrix} 10668.9 \end{pmatrix}$	$\begin{pmatrix} 10874.7 \end{pmatrix}$
$1(0^+)$	$\begin{pmatrix} 31.1 & 41.2 \\ 41.2 & 80.3 \end{pmatrix}$	$\begin{pmatrix} 7.8 \\ 103.7 \end{pmatrix}$	$\begin{pmatrix} 10641.7 \\ 10737.6 \end{pmatrix}$	$\begin{pmatrix} 10634.3 \\ 10730.2 \end{pmatrix}$	$\begin{pmatrix} 10840.2 \\ 10936.1 \end{pmatrix}$
$0(1^+)$	$\begin{pmatrix} -141.7 & -23.8 \\ -23.8 & -17.3 \end{pmatrix}$	$\begin{pmatrix} -146.1 \\ -12.9 \end{pmatrix}$	$\begin{pmatrix} 10487.9 \\ 10621.0 \end{pmatrix}$	$\begin{pmatrix} 10480.5 \\ 10613.6 \end{pmatrix}$	$\begin{pmatrix} 10686.3 \\ 10819.5 \end{pmatrix}$
$(bb\bar{n}\bar{s})$ J^P	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $\bar{B}^{(*)}\bar{B}_s^{(*)}$	Mass (up.) Eq. (2)
2^+	$\begin{pmatrix} 48.5 \end{pmatrix}$	$\begin{pmatrix} 48.5 \end{pmatrix}$	$\begin{pmatrix} 10817.6 \end{pmatrix}$	$\begin{pmatrix} 10764.7 \end{pmatrix}$	$\begin{pmatrix} 11061.5 \end{pmatrix}$
1^+	$\begin{pmatrix} 25.0 & 0.0 & 0.0 \\ 0.0 & -91.3 & -24.9 \\ 0.0 & -24.9 & -8.9 \end{pmatrix}$	$\begin{pmatrix} -98.2 \\ -2.0 \\ 25.0 \end{pmatrix}$	$\begin{pmatrix} 10670.9 \\ 10767.2 \\ 10794.1 \end{pmatrix}$	$\begin{pmatrix} 10618.0 \\ 10714.2 \\ 10741.2 \end{pmatrix}$	$\begin{pmatrix} 10914.8 \\ 11011.0 \\ 11038.0 \end{pmatrix}$
0^+	$\begin{pmatrix} 13.3 & 43.1 \\ 43.1 & 55.1 \end{pmatrix}$	$\begin{pmatrix} -13.7 \\ 82.1 \end{pmatrix}$	$\begin{pmatrix} 10755.4 \\ 10851.2 \end{pmatrix}$	$\begin{pmatrix} 10702.5 \\ 10798.3 \end{pmatrix}$	$\begin{pmatrix} 10999.3 \\ 11095.1 \end{pmatrix}$
$(bb\bar{s}\bar{s})$ J^P	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $\bar{B}_s^{(*)}\bar{B}_s^{(*)}$	Mass (up.) Eq. (2)
2^+	$\begin{pmatrix} 32.2 \end{pmatrix}$	$\begin{pmatrix} 32.2 \end{pmatrix}$	$\begin{pmatrix} 10925.6 \end{pmatrix}$	$\begin{pmatrix} 10838.1 \end{pmatrix}$	$\begin{pmatrix} 11225.8 \end{pmatrix}$
1^+	$\begin{pmatrix} 7.7 \end{pmatrix}$	$\begin{pmatrix} 7.7 \end{pmatrix}$	$\begin{pmatrix} 10901.0 \end{pmatrix}$	$\begin{pmatrix} 10813.6 \end{pmatrix}$	$\begin{pmatrix} 11201.3 \end{pmatrix}$
0^+	$\begin{pmatrix} -4.6 & 45.1 \\ 45.1 & 29.9 \end{pmatrix}$	$\begin{pmatrix} -35.6 \\ 60.9 \end{pmatrix}$	$\begin{pmatrix} 10857.7 \\ 10954.3 \end{pmatrix}$	$\begin{pmatrix} 10770.3 \\ 10866.8 \end{pmatrix}$	$\begin{pmatrix} 11158.0 \\ 11254.5 \end{pmatrix}$

states, the masses in the current work are approximately 50 MeV higher than those in Ref. [11]. According to Fig. 3(b), only the lowest $T_{bb\bar{n}\bar{s}}^{J=1}$, which is slightly (~ 20 MeV) below the $\bar{B}\bar{B}_s^*$ threshold, is possibly a stable tetraquark. This conclusion is similar to that of Ref. [11]. For the heaviest $bb\bar{s}\bar{s}$ states, the masses in the current work are approximately 80 MeV higher than those in Ref. [11], and no stable state is found in both approaches. Therefore, the HDAS and threshold approaches yield similar conclusions regarding the state stabilities for double-bottom tetraquarks.

It is interesting that the masses of the $bb\bar{n}\bar{n}$ states from HDAS consideration coincide with those from the threshold approach. This coincidence may mean that the $(bb)(\bar{n}\bar{n})$ diquark-antidiquark structure and the $(b\bar{n})(b\bar{n})$ molecule-like structure have similar effects on the mass spectrum. It probably also implies that the four quark components have an almost equal spatial distance. If this is true, it is also possible that more than one structure exists near the $\bar{B}\bar{B}^*$ threshold [27]. Of course, future experimental data are required to evaluate which approach provides more reasonable results and how large the effects from mass uncertainty of Ξ_{bb} would be.

HDAS relations similar to Eqs. (14), (15), and (16) can also be applied to bottom-charm tetraquark systems with the replacement $cc \rightarrow bc$. It should be noted that there are two bases for the highest-spin $bc\bar{n}\bar{n}$ and $bc\bar{n}\bar{s}$ tetraquarks. For the former case, the color-triplet (color-sextet) diquark exists only in the isovector (isoscalar) state. For the latter case, the two bases are nearly uncoupled, and the color-triplet diquark exists mainly in the higher state. The color-sextet contributions to the higher $T_{bc\bar{n}\bar{s}}^{J=2}$ are not a concern.

Using the mass $\Xi_{bc} = 6922.3 \pm 6.9$ MeV from Ref. [60] and our CMI model, we obtain $\Xi_{bc}^* = 6974$ MeV [50] and $\Omega_{bc}^* = \Xi_{bc}^* + \bar{B}_s^* - \bar{B}^* = 7065$ MeV. Thereafter,

$$\begin{aligned}
T_{bc\bar{n}\bar{n}}^{I=1, J=2} &= \Xi_{bc}^* + \Sigma_b^* - \bar{B}^* = 7483 \text{ MeV}, \\
T_{bc\bar{n}\bar{s}}^{J=2, \text{higher}} &= \Xi_{bc}^* + \Xi_b^* - \bar{B}^* = 7602 \text{ MeV}, \\
T_{bc\bar{s}\bar{s}}^{J=2} &= \Omega_{bc}^* + \Omega_b^* - \bar{B}_s^* = 7710 \text{ MeV},
\end{aligned} \tag{18}$$

are obtained. These values are 150 MeV, 165 MeV, and 183 MeV (increasing numbers) higher than the \bar{B}^*D^* , $\bar{B}^*D_s^*$, and $\bar{B}_s^*D_s^*$ thresholds, respectively. In Ref. [11], such states are approximately 34 MeV, 5 MeV, and 1 MeV (decreasing numbers) higher than the corresponding thresholds. With the reference scales in Eq. (18), we obtain the numerical results for the $bc\bar{q}\bar{q}$ tetraquark states in the CMI model. These are listed in Table VII, and the spectra are plotted in Fig. 4.

TABLE VII: Results for the $bc\bar{q}\bar{q}$ ($q = u, d, s$) states (unit: MeV). The second and third columns provide the numerical values of the CMI matrices and their eigenvalues, respectively. The fifth and sixth columns list the masses estimated using Eq. (3) (with $\bar{B}^{(*)}D^{(*)}/\bar{B}^{(*)}D_s^{(*)}/\bar{B}_s^{(*)}D_s^{(*)}$ as the reference state) and Eq. (2) (parameters provided in Table I), respectively. These can be viewed as the theoretical lower limits (low.) and upper limits (up.), respectively, for the tetraquark masses in the current framework. The fourth column displays our predictions with the heavy diquark-antiquark symmetry (HDAS) consideration.

$(bc\bar{n}\bar{n})$ $I(J^P)$	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $\bar{B}^{(*)}D^{(*)}$	Mass (up.) Eq. (2)
$1(2^+)$	$\begin{pmatrix} 77.4 \end{pmatrix}$	$\begin{pmatrix} 77.4 \end{pmatrix}$	$\begin{pmatrix} 7483.2 \end{pmatrix}$	$\begin{pmatrix} 7363.8 \end{pmatrix}$	$\begin{pmatrix} 7579.5 \end{pmatrix}$
$1(1^+)$	$\begin{pmatrix} 31.0 & 17.0 & 36.0 \\ 17.0 & 32.6 & -49.2 \\ 36.0 & -49.2 & 70.5 \end{pmatrix}$	$\begin{pmatrix} -24.0 \\ 47.9 \\ 110.2 \end{pmatrix}$	$\begin{pmatrix} 7381.9 \\ 7453.7 \\ 7516.0 \end{pmatrix}$	$\begin{pmatrix} 7262.5 \\ 7334.3 \\ 7396.6 \end{pmatrix}$	$\begin{pmatrix} 7478.1 \\ 7550.0 \\ 7612.3 \end{pmatrix}$
$1(0^+)$	$\begin{pmatrix} 7.8 & 85.2 \\ 85.2 & 81.3 \end{pmatrix}$	$\begin{pmatrix} -48.3 \\ 137.4 \end{pmatrix}$	$\begin{pmatrix} 7357.5 \\ 7543.2 \end{pmatrix}$	$\begin{pmatrix} 7238.1 \\ 7423.8 \end{pmatrix}$	$\begin{pmatrix} 7453.8 \\ 7639.5 \end{pmatrix}$
$0(2^+)$	$\begin{pmatrix} 30.9 \end{pmatrix}$	$\begin{pmatrix} 30.9 \end{pmatrix}$	$\begin{pmatrix} 7436.7 \end{pmatrix}$	$\begin{pmatrix} 7317.3 \end{pmatrix}$	$\begin{pmatrix} 7533.0 \end{pmatrix}$
$0(1^+)$	$\begin{pmatrix} -85.1 & 36.0 & 42.4 \\ 36.0 & -141.0 & -49.2 \\ 42.4 & -49.2 & -16.3 \end{pmatrix}$	$\begin{pmatrix} -182.7 \\ -70.2 \\ 10.5 \end{pmatrix}$	$\begin{pmatrix} 7223.1 \\ 7335.6 \\ 7416.3 \end{pmatrix}$	$\begin{pmatrix} 7103.7 \\ 7216.2 \\ 7296.9 \end{pmatrix}$	$\begin{pmatrix} 7319.4 \\ 7431.9 \\ 7512.6 \end{pmatrix}$
$0(0^+)$	$\begin{pmatrix} -143.1 & 85.2 \\ 85.2 & -162.6 \end{pmatrix}$	$\begin{pmatrix} -238.6 \\ -67.0 \end{pmatrix}$	$\begin{pmatrix} 7167.2 \\ 7338.8 \end{pmatrix}$	$\begin{pmatrix} 7047.8 \\ 7219.4 \end{pmatrix}$	$\begin{pmatrix} 7263.5 \\ 7435.1 \end{pmatrix}$
$(bc\bar{n}\bar{s})$ J^P	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $\bar{B}^{(*)}D_s^{(*)}$	Mass (up.) Eq. (2)
2^+	$\begin{pmatrix} 61.0 & 0.3 \\ 0.3 & 40.3 \end{pmatrix}$	$\begin{pmatrix} 40.3 \\ 61.0 \end{pmatrix}$	$\begin{pmatrix} 7581.5 \\ 7602.2 \end{pmatrix}$	$\begin{pmatrix} 7429.8 \\ 7450.5 \end{pmatrix}$	$\begin{pmatrix} 7723.0 \\ 7743.7 \end{pmatrix}$
1^+	$\begin{pmatrix} 13.8 & -0.6 & 16.8 & -0.3 & 35.6 & -1.2 \\ -0.6 & -90.6 & -0.1 & 35.6 & 0.0 & -50.1 \\ 16.8 & -0.1 & 15.8 & -1.2 & -50.1 & 0.0 \\ -0.3 & 35.6 & -1.2 & -77.7 & -1.4 & 42.0 \\ 35.6 & 0.0 & -50.1 & -1.4 & 45.3 & -0.3 \\ -1.2 & -50.1 & 0.0 & 42.0 & -0.3 & -7.9 \end{pmatrix}$	$\begin{pmatrix} -150.3 \\ -48.2 \\ -43.5 \\ 21.9 \\ 30.6 \\ 88.1 \end{pmatrix}$	$\begin{pmatrix} 7390.9 \\ 7493.0 \\ 7497.7 \\ 7563.1 \\ 7571.8 \\ 7629.3 \end{pmatrix}$	$\begin{pmatrix} 7239.2 \\ 7341.4 \\ 7346.1 \\ 7411.5 \\ 7420.2 \\ 7477.7 \end{pmatrix}$	$\begin{pmatrix} 7532.4 \\ 7634.5 \\ 7639.2 \\ 7704.6 \\ 7713.3 \\ 7770.8 \end{pmatrix}$
0^+	$\begin{pmatrix} -9.8 & 0.2 & -0.6 & 86.7 \\ 0.2 & -112.2 & 86.7 & 0.0 \\ -0.6 & 86.7 & -136.7 & 0.6 \\ 86.7 & 0.0 & 0.6 & 56.1 \end{pmatrix}$	$\begin{pmatrix} -212.0 \\ -69.6 \\ -36.8 \\ 115.9 \end{pmatrix}$	$\begin{pmatrix} 7329.2 \\ 7471.6 \\ 7504.3 \\ 7657.1 \end{pmatrix}$	$\begin{pmatrix} 7177.5 \\ 7319.9 \\ 7352.7 \\ 7505.4 \end{pmatrix}$	$\begin{pmatrix} 7470.7 \\ 7613.1 \\ 7645.9 \\ 7798.6 \end{pmatrix}$
$(bc\bar{s}\bar{s})$ J^P	$\langle H_{CM} \rangle$	Eigenvalues	Mass (our) HDAS	Mass (low.) $\bar{B}_s^{(*)}D_s^{(*)}$	Mass (up.) Eq. (2)
2^+	$\begin{pmatrix} 44.6 \end{pmatrix}$	$\begin{pmatrix} 44.6 \end{pmatrix}$	$\begin{pmatrix} 7710.1 \end{pmatrix}$	$\begin{pmatrix} 7523.9 \end{pmatrix}$	$\begin{pmatrix} 7907.9 \end{pmatrix}$
1^+	$\begin{pmatrix} -3.4 & 16.6 & 35.2 \\ 16.6 & -1.0 & -50.9 \\ 35.2 & -50.9 & 20.1 \end{pmatrix}$	$\begin{pmatrix} -63.7 \\ 13.0 \\ 66.4 \end{pmatrix}$	$\begin{pmatrix} 7601.9 \\ 7678.5 \\ 7731.9 \end{pmatrix}$	$\begin{pmatrix} 7415.6 \\ 7492.3 \\ 7545.7 \end{pmatrix}$	$\begin{pmatrix} 7799.6 \\ 7876.3 \\ 7929.7 \end{pmatrix}$
0^+	$\begin{pmatrix} -27.4 & 88.2 \\ 88.2 & 30.9 \end{pmatrix}$	$\begin{pmatrix} -91.1 \\ 94.6 \end{pmatrix}$	$\begin{pmatrix} 7574.4 \\ 7760.1 \end{pmatrix}$	$\begin{pmatrix} 7388.1 \\ 7573.9 \end{pmatrix}$	$\begin{pmatrix} 7772.2 \\ 7957.9 \end{pmatrix}$

For the $bc\bar{n}\bar{n}$ system, stable states are not found from our results. However, if the errors in the adopted approach are considered, the lowest $I(J^P) = 0(0^+)$ and $0(1^+)$ tetraquarks may be around the $\bar{B}D$ and \bar{B}^*D thresholds, respectively. This conclusion differs from that in Ref. [11], where these two states and the isoscalar spin-2 state are all stable. In

Ref. [69], we investigated the $bc\bar{n}\bar{n}$ spectrum with a reference scale related to the $X(4140)$ by assuming it to be a $c\bar{s}\bar{s}$ tetraquark. It is interesting that the masses of the $bc\bar{n}\bar{n}$ states in that approach are consistent with the present results. The conclusion that the lowest 0^+ T_{bc} may be around the $\bar{B}D$ threshold is also consistent with the findings in Ref. [19]. For the $(bc\bar{n}\bar{s})$ and $(bc\bar{s}\bar{s})$ systems, according to Fig. 4, no stable tetraquarks can be found, which is consistent with the conclusion in Ref. [69] but different from that in Ref. [11], where stable $bc\bar{n}\bar{s}$ is still possible. Although the masses of $bc\bar{n}\bar{n}$ agree with those in Ref. [69], those of the $bc\bar{n}\bar{s}$ and $bc\bar{s}\bar{s}$ states are higher. Future experiments will be required to evaluate which approach, threshold, $X(4140)$, or HDAS, is better.




FIG. 4: Relative positions for bottom-charm tetraquark states (solid and dashed lines) and relevant meson-meson thresholds (dotted lines). The masses are given in MeV. In (a), the solid (dashed) lines denote the $I = 1$ ($I = 0$) states.

V. CONSTRAINTS ON MASSES OF QQq , QQQ , AND $QQ\bar{q}\bar{q}$ STATES

We have obtained the masses of the double-heavy tetraquark states with the aid of heavy diquark-antiquark symmetry. The values are all larger than those in the threshold approach in Ref. [11], in which reference meson-meson thresholds were adopted. For the $bc\bar{n}\bar{n}$ and $bc\bar{n}\bar{s}$ states, the current masses are also heavier than those in the $X(4140)$ approach in Ref. [69], in which the reference scales were related to the mass of $X(4140)$. At present, it cannot be judged where such states are actually located, as there is still no observed double-heavy tetraquark. Even for conventional baryons, a state containing two heavy quarks that are heavier than Ξ_{cc} has not been reported. It is helpful to make a theoretical estimation on the range of their masses to understand the possible structures of an observed state. In Tables III, IV, VI, and VII, we treat the masses obtained using Eq. (2) as the upper limits. In fact, the range may be narrowed further from another perspective. We now investigate the constraints on the masses of the QQq and QQQ baryons first, and then those of the double-heavy tetraquarks. If the experiments reveal a state with a larger mass than the obtained limit, that hadron should not be a ground state with high probability.

Suppose that we are estimating the mass of a multiquark state X using Eq. (3). In principle, several reference hadron-hadron systems can be adopted. Their thresholds will result in different values for the mass of the X state. Considering two reference systems $\mathcal{A} + \mathcal{D}$ and $\mathcal{B} + \mathcal{C}$, where $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ are four hadrons, and $\mathcal{A} + \mathcal{D}$ and $\mathcal{B} + \mathcal{C}$ have the same quark content as X , two values for the mass of X ($M_{\mathcal{AD}}$ and $M_{\mathcal{BC}}$) can be obtained. An inequality between these must exist. For convenience, we further assume that they satisfy $M_{\mathcal{AD}} < M_{\mathcal{BC}}$. Thus, according to the estimation formula (3), we obtain

$$[M_{\mathcal{A}} - \langle H_{CMI} \rangle_{\mathcal{A}}] + [M_{\mathcal{D}} - \langle H_{CMI} \rangle_{\mathcal{D}}] < [M_{\mathcal{B}} - \langle H_{CMI} \rangle_{\mathcal{B}}] + [M_{\mathcal{C}} - \langle H_{CMI} \rangle_{\mathcal{C}}], \quad (19)$$

where $M_{\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}}$ should be the measured masses and no longer the theoretical masses obtained using Eq. (2) or (3). This formula means that the color-electric interactions in the hadrons have different effects on the two reference hadron-hadron systems. In many cases, it has been found that systems with $M_{\mathcal{A}} > M_{\mathcal{B}}/M_{\mathcal{C}} > M_{\mathcal{D}}$ satisfy the inequality [51, 54, 55]. If the assignment for the four hadrons can be provided, and the masses of three of them have been measured, a constraint on the mass of the fourth hadron will be obtained. In this case, we do not demand that the multiquark state X must exist. For the involved states in the present study, \mathcal{A} has two heavy quarks, whereas \mathcal{B}/\mathcal{C} has one heavy quark. Because the binding between two heavy quarks owing to the color-Coulomb potential is positively associated with their reduced mass, the QQ attraction inside \mathcal{A} is large. When the quark structure in $\mathcal{A} + \mathcal{D}$

is changed to $\mathcal{B} + \mathcal{C}$, the two heavy quarks require additional energy so that they can be separated and recombined with other light quarks into two single-heavy hadrons. Subsequently, an inequality (19) naturally follows. It should be noted that it is different from the Hall-Post inequalities, which link N -body energies to N' -body energies ($N' < N$) [94].

As an example, we consider the ground Ξ_{cc} state. When we discuss the case whereby $\mathcal{A} = \Xi_{cc}$ and \mathcal{D} is a light meson, we can select the following four sets of reference meson-baryon systems,

$$\text{I: } \Xi_{cc} + \pi \leftrightarrow \Sigma_c + D; \quad \text{II: } \Xi_{cc} + K \leftrightarrow \Sigma_c + D_s; \quad \text{III: } \Xi_{cc} + \bar{K} \leftrightarrow \Xi'_c + D; \quad \text{IV: } \Xi_{cc} + \phi \leftrightarrow \Xi'_c + D_s.$$

Their corresponding X states are $ccnn\bar{n}$, $ccnn\bar{s}$, $ccns\bar{n}$, and $ccns\bar{s}$, respectively. When we adopt the inequality $M_{\mathcal{AD}} < M_{\mathcal{BC}}$, we can obtain the upper limits for the mass of Ξ_{cc} : 3770 MeV, 3694 MeV, 3735 MeV, and 3663 MeV, respectively. Their difference may be a considerable 100 MeV, but the lowest value 3663 MeV should be used as the constraint. This value is approximately 40 MeV larger than the LHCb result $\Xi_{cc}^{++} = 3621$ MeV [1]. To obtain the mass constraint, we can also use higher spin states herein, such as $\Xi_{cc} + \rho \rightarrow \Sigma_c + D^*$, but no new results are obtained. Another case that can be discussed is that in which \mathcal{D} is a light baryon. Five sets of reference baryon-baryon systems can be considered:

$$\text{I: } \Xi_{cc} + N \leftrightarrow \Sigma_c + \Sigma_c; \quad \text{II: } \Xi_{cc} + \Sigma \leftrightarrow \Sigma_c + \Xi'_c; \quad \text{III: } \Xi_{cc} + \Xi \leftrightarrow \Xi'_c + \Xi'_c; \quad \text{IV: } \Xi_{cc} + \Xi \leftrightarrow \Sigma_c + \Omega_c; \quad \text{V: } \Xi_{cc} + \Omega \leftrightarrow \Xi'_c + \Omega_c.$$

Their corresponding X states are $ccnnnn$, $ccnnns$, $ccnnss$, $ccnnss$, and $ccnsss$, respectively. The upper limits that are obtained are 3776 MeV, 3732 MeV, 3717 MeV, 3709 MeV, and 3657 MeV, respectively. The situation is very similar to the above case. Combining the analyses in these two cases, we obtain $\Xi_{cc} < 3657$ MeV, which is approximately 30 MeV higher than the measured value.

Extending the discussions to Ξ_{bb} and Ξ_{bc} , we can similarly determine the minimum upper limits for their masses. We display the relevant reference states and results in Table VIII. The obtained constraints are $\Xi_{bb} < 10319$ MeV and $\Xi_{bc} < 6972$ MeV. Similar constraints can also be found in Ref. [50]. In fact, more stringent constraints are possible.

TABLE VIII: \mathcal{DBC} states in constraining the upper limits for the masses of the QQn baryons and the obtained limits (Up.) in MeV. Here, “Meson” (“Baryon”) means that \mathcal{D} is a light-quark meson (baryon).

Ξ_{cc}	Meson	$\pi\Sigma_c D$	$K\Sigma_c D_s$	$K\Xi'_c D$	$\phi\Xi'_c D_s$	
	Up.	3770	3694	3735	3663	
	Baryon	$N\Sigma_c \Sigma_c$	$\Sigma\Sigma_c \Xi'_c$	$\Xi\Xi'_c \Xi'_c$	$\Xi\Sigma_c \Omega_c$	$\Omega\Xi'_c \Omega_c$
	Up.	3775	3732	3717	3709	3657
Ξ_{bb}	Meson	$\pi\Sigma_b \bar{B}$	$K\Sigma_b \bar{B}_s$	$K\Xi'_b \bar{B}$	$\phi\Xi'_b \bar{B}_s$	
	Up.	10466	10377	10423	10339	
	Baryon	$N\Sigma_b \Sigma_b$	$\Sigma\Sigma_b \Xi'_b$	$\Xi\Xi'_b \Xi'_b$	$\Xi\Sigma_b \Omega_b$	$\Omega\Xi'_b \Omega_b$
	Up.	10462	10412	10389	10379	10319
Ξ_{bc}	Meson	$\pi\Sigma_b D$	$\pi\Sigma_c B$	$K\Sigma_b D_s$	$K\Sigma_c \bar{B}_s$	$\phi\Xi'_b D_s$ $\phi\Xi'_c \bar{B}_s$
	Up.	7102	7111	7027	7022	6988 6991
	Baryon	$N\Sigma_b \Sigma_c$	$\Sigma\Sigma_b \Xi'_c$	$\Sigma\Sigma_c \Xi'_b$	$\Xi\Sigma_b \Omega_c$	$\Xi\Sigma_c \Omega_b$ $\Xi\Xi'_b \Xi'_c$ $\Omega\Xi'_b \Omega_c$ $\Omega\Xi'_c \Omega_b$
	Up.	7108	7065	7057	7042	7024 7042 6982 6972

In Sec. IV B, we obtained the constraint $\Xi_{bb} < 10327$ MeV with the inequality $\Xi_{bb} - (\Xi_{bb})_{CMI} - 2(\bar{B} - \bar{B}_{CMI}) < \Xi_{cc} - (\Xi_{cc})_{CMI} - 2(D - D_{CMI})$ while determining an appropriate mass of Ξ_{bb} . This inequality, similar to (19), also arises from the color-Coulomb interaction between two heavy quarks. Naturally, the mass of Ξ_{bc} can also be taken into consideration, and we obtain

$$\begin{aligned} & \Xi_{bb} - (\Xi_{bb})_{CMI} - 2(\bar{B} - \bar{B}_{CMI}) \\ & < \Xi_{bc} - (\Xi_{bc})_{CMI} - (\bar{B} - \bar{B}_{CMI}) - (D - D_{CMI}) \\ & < \Xi_{cc} - (\Xi_{cc})_{CMI} - 2(D - D_{CMI}). \end{aligned} \tag{20}$$

The constraint $\Xi_{bc} < 6963$ MeV subsequently follows. Replacing the reference mesons with reference baryons, we similarly obtain

$$\begin{aligned} & \Xi_{bb} - (\Xi_{bb})_{CMI} - 2(\Sigma_b - (\Sigma_b)_{CMI}) \\ & < \Xi_{bc} - (\Xi_{bc})_{CMI} - (\Sigma_b - (\Sigma_b)_{CMI}) - (\Sigma_c - (\Sigma_c)_{CMI}) \\ & < \Xi_{cc} - (\Xi_{cc})_{CMI} - 2(\Sigma_c - (\Sigma_c)_{CMI}). \end{aligned} \tag{21}$$

Now, slightly smaller numbers ($\Xi_{bb} < 10308$ MeV and $\Xi_{bc} < 6954$ MeV) than those shown in Table VIII are obtained.

According to the above discussions, our short summary on the mass constraints for the doubly heavy baryons is as follows: $\Xi_{cc} < 3657$ MeV, $\Xi_{bb} < 10308$ MeV, and $\Xi_{bc} < 6954$ MeV. Of course, the constraint on Ξ_{bb} may be updated once the mass of Ξ_{bc} is measured, or vice versa. If an observed Ξ_{bb} or Ξ_{bc} has a larger mass than the limit provided here, it should not be the ground state. In the previous sections, the masses adopted were $\Xi_{cc} = 3621$ MeV, $\Xi_{bb} = 10169$ MeV, and $\Xi_{bc} = 6922$ MeV, which satisfy the obtained constraints. We do not discuss the QQs case. At present, without experimental data regarding the QQs baryons, we cannot obtain more stringent constraints than those provided in Ref. [50].

Using the same concept, we can estimate the upper limits for the masses of triply heavy baryons. These rely on the masses of doubly heavy baryons. We use $\Xi_{cc} = 3621$ MeV, $\Xi_{bc} < 6954$ MeV, and $\Xi_{bb} < 10308$ MeV as the inputs and list our results for all of the ground triply heavy baryons in Table IX, from which $\Omega_{ccc} < 4962$ MeV, $\Omega_{ccb} < 8250$ MeV, $\Omega_{bbc} < 11578$ MeV, and $\Omega_{bbb} < 14939$ MeV can be obtained. Similar constraints can also be found in Ref. [50]. The upper limits for the Ω_{ccb} , Ω_{bbc} , and Ω_{bbb} states could be changed to lower values if the masses of Ξ_{bc} and Ξ_{bb} are measured experimentally.

TABLE IX: \mathcal{DBC} states in constraining the upper limits for the masses of the QQQ baryons and the obtained limits (Up.) in MeV. Here, “Meson” (“Baryon”) means that \mathcal{D} is a light-quark meson (baryon).

Ω_{ccc}	Meson	$\pi\Xi_{cc}D$	$K\Xi_{cc}D_s$			
	Up.	5038	4962			
	Baryon	$N\Xi_{cc}\Sigma_c$	$\Sigma\Xi_{cc}\Xi'_c$	$\Xi\Xi_{cc}\Omega_c$		
	Up.	5043	5000	4977		
Ω_{ccb}	Meson	$\pi\Xi_{cc}B$	$K\Xi_{cc}B_s$	$\pi\Xi_{bc}D$	$K\Xi_{bc}D_s$	
	Up.	8339	8250	8298	8254	
	Baryon	$N\Xi_{cc}\Sigma_b$	$\Sigma\Xi_{cc}\Xi'_b$	$\Xi\Xi_{cc}\Omega_b$	$N\Xi_{bc}\Sigma_c$	$\Sigma\Xi_{bc}\Xi'_c$
	Up.	8336	8285	8252	8336	8292
Ω_{bbc}	Meson	$\pi\Xi_{bc}\bar{B}$	$K\Xi_{bc}\bar{B}_s$	$\pi\Xi_{bb}D$	$K\Xi_{bb}D_s$	
	Up.	11666	11578	11657	11581	
	Baryon	$N\Xi_{bc}\Sigma_b$	$\Sigma\Xi_{bc}\Xi'_b$	$\Xi\Xi_{bc}\Omega_b$	$N\Xi_{bb}\Sigma_c$	$\Sigma\Xi_{bb}\Xi'_c$
	Up.	11663	11612	11579	11662	11619
Ω_{bbb}	Meson	$\pi\Xi_{bb}\bar{B}$	$K\Xi_{bb}\bar{B}_s$			
	Up.	16028	14939			
	Baryon	$N\Xi_{bb}\Sigma_b$	$\Sigma\Xi_{bb}\Xi'_b$	$\Xi\Xi_{bb}\Omega_b$		
	Up.	15025	14974	14941		

Now, we investigate the $QQ\bar{q}\bar{q}$ case. When we consider a double-heavy tetraquark state in terms of the diquark-antiquark symmetry, its mass is linearly dependent on the mass of a related double-heavy baryon, which is treated as an input, see, e.g., Eq. (14). If we use Δ^{up} to denote the difference between the upper limit for the mass of this QQq baryon and the mass that we adopt, we can set the upper limit for the mass of any tetraquark state by adding Δ^{up} to the obtained tetraquark mass, so that the HDAS relation still holds. Explicitly, we need to add 36 MeV, 139 MeV, and 32 MeV for $cc\bar{q}\bar{q}$, $bb\bar{q}\bar{q}$, and $bc\bar{q}\bar{q}$, respectively. For the lowest $QQ\bar{n}\bar{n}$ tetraquarks, we have $T_{cc} < 3965$ MeV, $T_{bb} < 10627$ MeV ($\approx BB^*$ threshold+17 MeV), and $T_{bc} < 7199$ MeV. However, because the symmetry relations are only approximately correct, the measured tetraquark masses in future experiments may exceed such limits.

Let us return to the upper limits using the inequality (19). Naturally, the involved X systems are $QQq\bar{q}\bar{q}$ and $QQqq\bar{q}\bar{q}$. In the latter case, tetraquarks are always involved in the reference channels ($\mathcal{A} + \mathcal{D}$ or $\mathcal{B} + \mathcal{C}$), and we cannot obtain useful information, at least presently. In the former case, we, unfortunately, cannot obtain reliable constraints, either. If $X = cc\bar{n}\bar{n}\bar{s}$, for example, the reference system can be $(cc\bar{n}\bar{n})(s\bar{s})$, $(cs\bar{n}\bar{n})(c\bar{s})$, or $(ccs)(\bar{n}\bar{n}\bar{s})$. However, in constraining the mass of $cc\bar{n}\bar{n}$, neither $(cs\bar{n}\bar{n})(c\bar{s})$ nor $(ccs)(\bar{n}\bar{n}\bar{s})$ can be adopted. The former system involves another tetraquark state, whereas the latter does not meet the requirement to use the inequality (19) that the two heavy quarks should be separated into two hadrons, which guarantees the difference caused by the color-Coulomb potential. Another reason is that the mass of ccs baryon has not been measured. If we neglect the requirement to use (19) and consider the case where $X = ccn\bar{n}\bar{n}$, the reference state $(ccn)(\bar{n}\bar{n})$ can be adopted and a mass constraint $T_{cc} < 3952$ MeV is obtained. This value appears to be the upper limit of the mass, but this is simply conjecture and not a conclusion. Therefore, we could not get more information from the case that \mathcal{D} is a light hadron. One may wonder whether we can estimate the lower limit of the $QQ\bar{q}\bar{q}$ mass with the minimum theoretical mass of QQQ baryon by considering the case $X = QQQ\bar{q}\bar{q}$. In fact, obtaining the limit is possible, but the constraint is probably

not useful. For example, one could get $T_{cc} > 3704$ MeV with $\Omega_{ccc} = 4790$ MeV [95]. This value is approximately 70 MeV smaller than the lower limit given in Table III, and no meaningful constraint is obtained.

Similar to (20), another inequality exists for tetraquarks. In this case, we only consider the case without strange quarks. Subsequently, we obtain

$$\begin{aligned} & T_{bb} - (T_{bb})_{CMI} - 2(\bar{B} - \bar{B}_{CMI}) \\ & < T_{bc} - (T_{bc})_{CMI} - (\bar{B} + D - \bar{B}_{CMI} - D_{CMI}) \\ & < T_{cc} - (T_{cc})_{CMI} - 2(D - D_{CMI}), \end{aligned} \quad (22)$$

which can be employed to verify the results obtained in the previous sections. As $T_{bb} - (T_{bb})_{CMI} - 2(\bar{B} - \bar{B}_{CMI}) = 8$ MeV, $T_{bc} - (T_{bc})_{CMI} - (\bar{B} + D - \bar{B}_{CMI} - D_{CMI}) = 120$ MeV, and $T_{cc} - (T_{cc})_{CMI} - 2(D - D_{CMI}) < 156$ MeV, the inequalities are certainly satisfied. We can also consider the inequality to be similar to (21), but the obtained relations do not change. Thus, the inequality (22) is sufficient for the purpose of conducting a simple check on the obtained tetraquark masses.

VI. DISCUSSIONS AND SUMMARY

It is known that the mass splittings of conventional hadrons are mainly determined by the chromomagnetic interactions. However, while applying the CMI model (2) to hadron masses, the deviations from the experimental data may be large (e.g. Table 2 of Ref. [50]). After all, the model is a simplified version of potential quark models. Contributions from color-Coulomb interaction, color confinement, and others are simply effectively absorbed into the masses of the quarks and coupling parameters. In principle, it is unrealistic to determine all the hadron masses with only one set of parameters. In the multiquark case, we tend to adopt a method to compensate for the above effects partially by selecting a suitable reference system, instead of using Eq. (2) directly. This appears to be more reasonable than simply taking a set of effective quark masses as input, but the details of the kinematic and dynamic effects may still lead to a significant shift in the spectrum. To fix the deviation, we can take the other effects into account explicitly by sacrificing the concision and simplicity of calculation. However, we can also balance simplicity and rationality in certain peculiar cases, as in the double-heavy tetraquark systems explored here.

When the $\bar{3}_c$ QQ diquark is regarded as a heavier antiquark \bar{Q}' , the double-heavy tetraquark $QQ\bar{q}\bar{q}$ can be viewed as a single-heavy “antibaryon” $\bar{Q}'\bar{q}\bar{q}$ in the sense that they have the same color configuration. The mass relations among the $QQ\bar{q}\bar{q}$, QQq , Qqq , and $Q\bar{q}$ states follow such a heavy diquark-antiquark symmetry. As only the highest-spin $QQ\bar{q}\bar{q}$ states may contain the pure $\bar{3}_c$ QQ diquark, and no mixing effects are involved, their masses are determined with the symmetry relations, and they are selected as the reference states to obtain the tetraquark spectra. Another consideration for using the highest-spin states is that the spin-dependent terms between the light quarks will be cancelled, and those between the heavy and light quarks can be ignored.

Once the masses of all the double-heavy tetraquarks are obtained, it is easy to determine whether or not stable tetraquarks exist from Figs. 2-4. In Table X, we present our answers to the question. In fact, numerous discussions on double-heavy tetraquarks can be found in the literature [50]. For example, Carlson et. [63] discussed non-strange $QQ\bar{q}\bar{q}$ systems and found that T_{bb} is sufficiently stable against strong decay, T_{cc} is unstable, and T_{bc} is uncertain. For comparison, we have also displayed the results obtained in some reference studies in the table. In general, all the studies support the stable double-bottom tetraquark T_{bb} . The results indicate that the double-charm T_{cc} state is probably unstable, whereas the stability of T_{bc} remains controversial.

The consistency between our results and others indicates that the estimation method with HDAS is reasonable. However, how reliable the numerical results are is not clear because they are affected by several factors. First, the accuracy of the approximate HDAS relations and errors of the input QQq masses determine the location of the tetraquark spectra. Second, the existence of $(QQ)_{6_c}(\bar{q}\bar{q})_{\bar{6}_c}$ configuration may significantly affect the mass splittings if the color-electric contributions are considered explicitly. Third, the values of C_{ij} determining the mass splittings are extracted from conventional hadrons. Whether they can be applied to multiquark states remains an open question.

The spacial structure of the tetraquark states was not considered in the above discussions. An observed double-heavy state can also be a meson-meson molecule, the spatial structure of which differs from the compact tetraquark. At present, it is generally difficult to determine a criterion to distinguish a compact multiquark state from a molecular state, but there are cases where this is possible. In the $b\bar{b}\bar{n}\bar{n}$ case, both compact tetraquark and molecules [96–100] are possible, but the binding energies in these two configurations differ. It is possible to identify the inner structure of the observed state: a large (small) binding energy corresponds to a compact (molecular) state. However, in the $c\bar{c}\bar{n}\bar{n}$ case, the observed state should be a molecule [96, 97, 99] if it is below the related meson-meson threshold. In the $b\bar{c}\bar{n}\bar{n}$ case, both molecules [97, 98] and compact tetraquarks are around the related meson-meson thresholds. The situation is complicated, and further discussions are required.

TABLE X: Stability of the double-heavy tetraquarks in various studies. The meanings of “S”, “US”, and “ND” are “stable”, “unstable”, and “not determined”, respectively.

Reference	$(cc\bar{n}\bar{n})$	$(cc\bar{n}\bar{s})$	$(cc\bar{s}\bar{s})$	$(bb\bar{n}\bar{n})$	$(bb\bar{n}\bar{s})$	$(bb\bar{s}\bar{s})$	$(bc\bar{n}\bar{n})$	$(bc\bar{n}\bar{s})$	$(bc\bar{s}\bar{s})$
This work	US	US	US	S	S	US	ND	US	US
[8]	S	S		S	S		S	US	
[11]	S	S	US	S	S	US	S	S	US
[16]	S			S					
[18]	S			S			S		
[19]	US			S			S		
[20]	US			S	S		US	US	
[24]	S			S			S		
[28]	S	US	US	S	S	US	S	US	US
[29]	S			S			S		
[30]	US	US	US	S	US	US	US	US	US
[31]	US	US	US	S	US	US	US	US	US
[32]			US			US			US
[33]	US	US	US	S	S	S			
[34]								S	S
[39]	US			S					
[44, 45]	US	US		S	S		S	US	
[47]							S		
[48]				S	S		US	US	
[63]	US			S			ND		
[69]							ND	US	
[83]	US	US	US	S	S	US	US	US	US
[84]	US	US	US	S	S	US	US	US	US

TABLE XI: Strong and electromagnetic decay patterns for the lowest tetraquark states.

System	Mass (MeV)	Strong Decay		Electromagnetic Decay
		2 body	3 body	3 body
$(cc\bar{n}\bar{n})_{I=0}^{J=1}$	3929	DD^*	$DD\pi$	$DD\gamma/DD^*\gamma$
$(cc\bar{n}\bar{s})_{I=0}^{J=1}$	4106	D^*D_s/DD_s^*	$DD_s\pi$	$DD_s\gamma/D^*D_s\gamma/DD_s^*\gamma$
$(cc\bar{s}\bar{s})_{I=0}^{J=0}$	4227	D_sD_s	$D_sD_s^*\pi$	$D_sD_s\gamma/D_sD_s^*\gamma$
$(bb\bar{n}\bar{n})_{I=0}^{J=1}$	10488	—	—	—
$(bb\bar{n}\bar{s})_{I=0}^{J=1}$	10671	—	—	$\bar{B}\bar{B}_s\gamma$
$(bb\bar{s}\bar{s})_{I=0}^{J=0}$	10858	$\bar{B}_s\bar{B}_s/\bar{B}_s^*\bar{B}_s^*$	—	$\bar{B}_s\bar{B}_s\gamma/\bar{B}_s\bar{B}_s^*\gamma/\bar{B}_s^*\bar{B}_s^*\gamma$
$(bc\bar{n}\bar{n})_{I=0}^{J=0}$	7167	$\bar{B}D$	—	$\bar{B}D\gamma$
$(bc\bar{n}\bar{s})_{I=0}^{J=0}$	7329	$\bar{B}_sD/\bar{B}D_s$	—	$\bar{B}_sD\gamma/\bar{B}D_s\gamma/\bar{B}_s^*D\gamma/\bar{B}^*D_s\gamma$
$(bc\bar{s}\bar{s})_{I=0}^{J=0}$	7574	$\bar{B}_sD_s/\bar{B}_s^*D_s^*$	$\bar{B}_s^*D_s\pi$	$\bar{B}_sD_s\gamma/\bar{B}_s^*D_s\gamma/\bar{B}_sD_s^*\gamma/\bar{B}_s^*D_s^*\gamma$

The detailed partial widths for the studied tetraquarks depend on the Hamiltonian and specific processes, and a quantitative calculation will be discussed in future work. Here, we present a brief analysis on their dominant decay patterns. In Table XI, the strong and electromagnetic decay patterns for the lowest state in each system are provided. For higher states, we simply mention the rearrangement decay modes. The thresholds of such meson-meson channels are illustrated in Figs. 2-4. Whether or not the decays can occur is determined mainly by the kinematics and quantum number conservations. In the $cc\bar{n}\bar{n}$ case, the allowed channels for the $I(J^P) = 0(1^+)$ state are DD^* and D^*D^* , those for the $1(0^+)$ states are DD and D^*D^* , those for the $1(1^+)$ state are DD^* and D^*D^* , whereas DD , DD^* , and D^*D^* are all allowed channels for the $1(2^+)$ state. The channels in the case $bb\bar{n}\bar{n}$ are similar. In the $cc\bar{s}\bar{s}$ and $bb\bar{s}\bar{s}$ cases, the channels can be obtained with reference to the $I = 1$ cases. In the $cc\bar{n}\bar{s}$ case, the 0^+ states can decay into DD_s and $D^*D_s^*$, the 1^+ states can decay into DD_s^*/D^*D_s and $D^*D_s^*$, and the 2^+ state can decay into all these channels. The case of $bb\bar{n}\bar{s}$ is similar. In the case of $bc\bar{n}\bar{n}$, the allowed channels for the 0^+ states are $\bar{B}D$ and \bar{B}^*D^* , those for

the 1^+ states are $\bar{B}^*D/\bar{B}D^*$ and \bar{B}^*D^* , and all these decay channels are allowed for the 2^+ state. The channels in the case of $bc\bar{s}\bar{s}$ can be obtained with the replacement $n \rightarrow s$. In the case of $bc\bar{n}\bar{s}$, the allowed decay channels for the 0^+ states are $\bar{B}_sD/\bar{B}D_s$ and $\bar{B}^*D_s/\bar{B}_s^*D^*$, those for the 1^+ states are $\bar{B}_s^*D/\bar{B}^*D_s$ and $\bar{B}_s^*D^*/B^*D_s^*$, and those for the 2^+ states are $\bar{B}_sD/\bar{B}D_s$, $\bar{B}_s^*D/\bar{B}^*D_s$ and $\bar{B}_s^*D^*/B^*D_s^*$.

If the studied compact double-heavy tetraquarks exist, one may wonder where and how to search for them according to the decay channels. In principle, they can be produced at any collider if the collision energy is sufficiently high. For example, they may be produced in the Z boson decay [101], hadron decays [102], pp collision [103], heavy-ion collisions [104–107], and e^+e^- annihilation process [9, 10, 108]. A low production rate and small signal/noise ratio should be the main reasons that double-heavy tetraquarks are not observed. Owing to the clean background, the e^+e^- annihilation process offers its own advantage in searching for a double-charm state. As a signal of double-charm tetraquarks has not been observed in such a process, the detection efficiency should be increased with improved analysis methods, such as that proposed in Ref. [108].

In summary, we have determined the masses of the highest-spin double-heavy tetraquark states $QQ\bar{q}\bar{q}$ with the aid of the heavy diquark-antiquark symmetry. Thereafter, such reference states were used to derive the masses of their partners with mass splittings in the CMI model. We presented the results in Tables III, IV, VI, and VII as well as Figs. 2-4. The double-charm tetraquarks $cc\bar{q}\bar{q}$ ($q = u, d, s$) were significantly higher than their rearrangement decay channels, and we did not find a bound state in the systems. In the double-bottom systems $bb\bar{q}\bar{q}$, we did not obtain stable $bb\bar{s}\bar{s}$ states, but observed a deep $bb\bar{n}\bar{n}$ bound state (T_{bb}) and a shallow $bb\bar{n}\bar{s}$ bound state. Their $I(J^P)$ were both $0(1^+)$, and their masses were 10488 MeV (≈ 116 MeV below the $\bar{B}\bar{B}^*$ threshold) and 10671 MeV (≈ 20 MeV below the $\bar{B}\bar{B}_s^*/\bar{B}_s\bar{B}^*$ threshold), respectively. The T_{bb} mass is very close to that obtained in Ref. [11] when $\bar{B}\bar{B}^*$ was used as the reference state. For the bottom-charm systems ($bc\bar{q}\bar{q}$), no stable $bc\bar{n}\bar{s}$ or $bc\bar{s}\bar{s}$ was found, but we could obtain two near-threshold $bc\bar{n}\bar{n}$ states. These were the lowest $0(0^+)$ state T_{bc} , with a mass of 7167 MeV, and the lowest $0(1^+)$ state, with a mass of 7223 MeV. Considering the model uncertainties, it was difficult to draw a conclusion whether or not they were stable. From our results, it can be concluded that the order of possibility for finding a bound $QQ\bar{q}\bar{q}$ tetraquark should be $(cc) < (bc) < (bb)$ and that a bound tetraquark becomes more difficult to form with the increasing number of strange quarks. Because not all of the input masses were measured, we also discussed the constraints on the masses of heavy quark hadrons, such as $\Xi_{bb} < 10308$ MeV and $\Omega_{ccc} < 4962$ MeV. We obtained $T_{cc} < 3965$ MeV, $T_{bb} < 10627$ MeV, and $T_{bc} < 7199$ MeV for the lowest tetraquark states. Of course, whether or not this is true requires future experimental tests. We hope that our predictions for double-heavy states will be helpful for future investigations.

Acknowledgments

This project is supported by National Natural Science Foundation of China (Grant Nos. 11775130, 11775132, 11635009, 11325525, 11875179) and by Natural Science Foundation of Shandong Province (Grant Nos. ZR2016AM16, ZR2017MA002).

-
- [1] R. Aaij *et al.* [LHCb Collaboration], “Observation of the doubly charmed baryon Ξ_{cc}^{++} ,” Phys. Rev. Lett. **119**, no. 11, 112001 (2017) [arXiv:1707.01621 [hep-ex]].
 - [2] M. Mattson *et al.* [SELEX Collaboration], “First Observation of the Doubly Charmed Baryon Ξ_{cc}^+ ,” Phys. Rev. Lett. **89**, 112001 (2002) [hep-ex/0208014].
 - [3] A. Ocherashvili *et al.* [SELEX], “Confirmation of the double charm baryon $\Xi_{cc}^+(3520)$ via its decay to pD^+K^- ,” Phys. Lett. B **628**, 18-24 (2005) [arXiv:hep-ex/0406033 [hep-ex]].
 - [4] S. P. Ratti, “New results on c-baryons and a search for cc-baryons in FOCUS,” Nucl. Phys. Proc. Suppl. **115**, 33 (2003).
 - [5] B. Aubert *et al.* [BaBar], “Search for doubly charmed baryons Ξ_{cc}^+ and Ξ_{cc}^{++} in BABAR,” Phys. Rev. D **74**, 011103 (2006) [arXiv:hep-ex/0605075 [hep-ex]].
 - [6] Y. Kato *et al.* [Belle Collaboration], “Search for doubly charmed baryons and study of charmed strange baryons at Belle,” Phys. Rev. D **89**, no. 5, 052003 (2014) [arXiv:1312.1026 [hep-ex]].
 - [7] R. Aaij *et al.* [LHCb Collaboration], “Search for the doubly charmed baryon Ξ_{cc}^+ ,” Sci. China Phys. Mech. Astron. **63**, no. 2, 221062 (2020) [arXiv:1909.12273 [hep-ex]].
 - [8] S. H. Lee and S. Yasui, “Stable multiquark states with heavy quarks in a diquark model,” Eur. Phys. J. C **64**, 283 (2009) [arXiv:0901.2977 [hep-ph]].
 - [9] T. Hyodo, Y. R. Liu, M. Oka, K. Sudoh and S. Yasui, “Production of doubly charmed tetraquarks with exotic color configurations in electron-positron collisions,” Phys. Lett. B **721**, 56 (2013) [arXiv:1209.6207 [hep-ph]].
 - [10] T. Hyodo, Y. R. Liu, M. Oka and S. Yasui, arXiv:1708.05169 [hep-ph].

- [11] S. Q. Luo, K. Chen, X. Liu, Y. R. Liu and S. L. Zhu, “Exotic tetraquark states with the $qq\bar{Q}\bar{Q}$ configuration,” *Eur. Phys. J. C* **77**, no. 10, 709 (2017) [arXiv:1707.01180 [hep-ph]].
- [12] X. Yan, B. Zhong and R. Zhu, “Doubly charmed tetraquarks in a diquark-antidiquark model,” *Int. J. Mod. Phys. A* **33**, no. 16, 1850096 (2018) [arXiv:1804.06761 [hep-ph]].
- [13] H. J. Lipkin, *Phys. Lett. B* **172**, 242 (1986).
- [14] S. Zouzou, B. Silvestre-Brac, C. Gignoux and J. M. Richard, *Z. Phys. C* **30**, 457 (1986).
- [15] C. Semay and B. Silvestre-Brac, “Diquonia and potential models,” *Z. Phys. C* **61**, 271 (1994).
- [16] S. Pepin, F. Stancu, M. Genovese and J. M. Richard, “Tetraquarks with color blind forces in chiral quark models,” *Phys. Lett. B* **393**, 119 (1997) [hep-ph/9609348].
- [17] D. M. Brink and F. Stancu, “Tetraquarks with heavy flavors,” *Phys. Rev. D* **57**, 6778 (1998).
- [18] G. Q. Feng, X. H. Guo and B. S. Zou, “ $QQ'\bar{u}\bar{d}$ bound state in the Bethe-Salpeter equation approach,” [arXiv:1309.7813 [hep-ph]].
- [19] M. Karliner and J. L. Rosner, “Discovery of doubly-charmed Ξ_{cc} baryon implies a stable $(bb\bar{u}\bar{d})$ tetraquark,” *Phys. Rev. Lett.* **119**, no. 20, 202001 (2017) [arXiv:1707.07666 [hep-ph]].
- [20] W. Park, S. Noh and S. H. Lee, “Masses of the doubly heavy tetraquarks in a constituent quark model,” *Nucl. Phys. A* **983**, 1 (2019) [arXiv:1809.05257 [nucl-th]].
- [21] T. F. Carames, J. Vijande and A. Valcarce, “Exotic $bc\bar{q}\bar{q}$ four-quark states,” *Phys. Rev. D* **99**, no. 1, 014006 (2019) [arXiv:1812.08991 [hep-ph]].
- [22] E. Hernandez, J. Vijande, A. Valcarce and J. M. Richard, “Spectroscopy, lifetime and decay modes of the T_{bb}^- tetraquark,” *Phys. Lett. B* **800**, 135073 (2020) [arXiv:1910.13394 [hep-ph]].
- [23] Y. Yang, C. Deng, J. Ping and T. Goldman, “S-wave $QQ\bar{q}\bar{q}$ state in the constituent quark model,” *Phys. Rev. D* **80**, 114023 (2009).
- [24] G. Yang, J. Ping and J. Segovia, “Doubly-heavy tetraquarks,” *Phys. Rev. D* **101**, no. 1, 014001 (2020) [arXiv:1911.00215 [hep-ph]].
- [25] M. A. Bedolla, J. Ferretti, C. D. Roberts and E. Santopinto, “Spectrum of fully-heavy tetraquarks from a diquark+antidiquark perspective,” arXiv:1911.00960 [hep-ph].
- [26] P. C. Wallbott, G. Eichmann and C. S. Fischer, [arXiv:2003.12407 [hep-ph]].
- [27] M. T. Yu, Z. Y. Zhou, D. Y. Chen and Z. Xiao, “Possible molecular states in $B^{(*)}B^{(*)}$ scatterings,” *Phys. Rev. D* **101**, no.7, 074027 (2020) [arXiv:1912.07348 [hep-ph]].
- [28] C. Deng, H. Chen and J. Ping, “Systematical investigation on the stability of doubly heavy tetraquark states,” *Eur. Phys. J. A* **56**, no.1, 9 (2020) [arXiv:1811.06462 [hep-ph]].
- [29] Y. Tan, W. Lu and J. Ping, “ $QQ\bar{q}\bar{q}$ in a chiral constituent quark model,” [arXiv:2004.02106 [hep-ph]].
- [30] Q. F. L, D. Y. Chen and Y. B. Dong, “Masses of doubly heavy tetraquarks $T_{QQ'}$ in a relativized quark model,” [arXiv:2006.08087 [hep-ph]].
- [31] D. Ebert, R. N. Faustov, V. O. Galkin and W. Lucha, “Masses of tetraquarks with two heavy quarks in the relativistic quark model,” *Phys. Rev. D* **76**, 114015 (2007) [arXiv:0706.3853 [hep-ph]].
- [32] G. Yang, J. Ping and J. Segovia, “ $QQ\bar{s}\bar{s}$ tetraquarks in the chiral quark model,” [arXiv:2007.05190 [hep-ph]].
- [33] M. L. Du, W. Chen, X. L. Chen and S. L. Zhu, “Exotic $QQ\bar{q}\bar{q}$, $QQ\bar{q}\bar{s}$ and $QQ\bar{s}\bar{s}$ states,” *Phys. Rev. D* **87**, no. 1, 014003 (2013) [arXiv:1209.5134 [hep-ph]].
- [34] W. Chen, T. G. Steele and S. L. Zhu, “Exotic open-flavor $bc\bar{q}\bar{q}$, $bc\bar{s}\bar{s}$ and $qc\bar{q}\bar{b}$, $sc\bar{s}\bar{b}$ tetraquark states,” *Phys. Rev. D* **89**, no. 5, 054037 (2014) [arXiv:1310.8337 [hep-ph]].
- [35] Z. G. Wang and Z. H. Yan, “Analysis of the scalar, axialvector, vector, tensor doubly charmed tetraquark states with QCD sum rules,” *Eur. Phys. J. C* **78**, no. 1, 19 (2018) [arXiv:1710.02810 [hep-ph]].
- [36] S. S. Agaev, K. Azizi and H. Sundu, “Double-heavy axial-vector tetraquark $T_{bc;\bar{u}\bar{d}}^0$,” *Nucl. Phys. B* **951**, 114890 (2020) [arXiv:1905.07591 [hep-ph]].
- [37] L. Tang, B. D. Wan, K. Maltman and C. F. Qiao, “Doubly Heavy Tetraquarks in QCD Sum Rules,” *Phys. Rev. D* **101**, no.9, 094032 (2020) [arXiv:1911.10951 [hep-ph]].
- [38] S. S. Agaev, K. Azizi, B. Barsbay and H. Sundu, “Heavy exotic scalar meson $T_{bb;\bar{u}\bar{s}}^-$,” *Phys. Rev. D* **101**, no.9, 094026 (2020) [arXiv:1912.07656 [hep-ph]].
- [39] F. S. Navarra, M. Nielsen and S. H. Lee, “QCD sum rules study of $QQ - \bar{u}\bar{d}$ mesons,” *Phys. Lett. B* **649**, 166-172 (2007) [arXiv:hep-ph/0703071 [hep-ph]].
- [40] D. Gao, D. Jia, Y. J. Sun, Z. Zhang, W. N. Liu and Q. Mei, “Masses of doubly heavy tetraquark states with isospin $= \frac{1}{2}$ and 1 and spin-parity 1^{++} ,” [arXiv:2007.15213 [hep-ph]].
- [41] Z. S. Brown and K. Orginos, “Tetraquark bound states in the heavy-light heavy-light system,” *Phys. Rev. D* **86**, 114506 (2012) [arXiv:1210.1953 [hep-lat]].
- [42] Y. Ikeda *et al.*, “Charmed tetraquarks T_{cc} and T_{cs} from dynamical lattice QCD simulations,” *Phys. Lett. B* **729**, 85 (2014) [arXiv:1311.6214 [hep-lat]].
- [43] P. Bicudo, K. Cichy, A. Peters, B. Wagenbach and M. Wagner, “Evidence for the existence of $ud\bar{b}\bar{b}$ and the non-existence of $ss\bar{b}\bar{b}$ and $ccb\bar{b}$ tetraquarks from lattice QCD,” *Phys. Rev. D* **92**, no. 1, 014507 (2015) [arXiv:1505.00613 [hep-lat]].
- [44] A. Francis, R. J. Hudspith, R. Lewis and K. Maltman, “Lattice Prediction for Deeply Bound Doubly Heavy Tetraquarks,” *Phys. Rev. Lett.* **118**, no. 14, 142001 (2017) [arXiv:1607.05214 [hep-lat]].
- [45] A. Francis, R. J. Hudspith, R. Lewis and K. Maltman, “Evidence for charm-bottom tetraquarks and the mass dependence of heavy-light tetraquark states from lattice QCD,” *Phys. Rev. D* **99**, no. 5, 054505 (2019) [arXiv:1810.10550 [hep-lat]].

- [46] P. Junnarkar, N. Mathur and M. Padmanath, “Study of doubly heavy tetraquarks in Lattice QCD,” *Phys. Rev. D* **99**, no.3, 034507 (2019) [arXiv:1810.12285 [hep-lat]].
- [47] L. Leskovec, S. Meinel, M. Pflaumer and M. Wagner, “Lattice QCD investigation of a doubly-bottom $\bar{b}\bar{b}ud$ tetraquark with quantum numbers $I(J^P) = 0(1^+)$,” *Phys. Rev. D* **100**, no. 1, 014503 (2019) [arXiv:1904.04197 [hep-lat]].
- [48] R. J. Hudspith, B. Colquhoun, A. Francis, R. Lewis and K. Maltman, “A lattice investigation of exotic tetraquark channels,” [arXiv:2006.14294 [hep-lat]].
- [49] Y. Liu, M. A. Nowak and I. Zahed, “Heavy tetraquark $QQ\bar{q}\bar{q}$ as a hadronic Efimov state,” [arXiv:1909.02497 [hep-ph]].
- [50] Y. R. Liu, H. X. Chen, W. Chen, X. Liu and S. L. Zhu, “Pentaquark and Tetraquark states,” *Prog. Part. Nucl. Phys.* **107**, 237 (2019) [arXiv:1903.11976 [hep-ph]].
- [51] J. Wu, Y. R. Liu, K. Chen, X. Liu and S. L. Zhu, “ $X(4140)$, $X(4270)$, $X(4500)$ and $X(4700)$ and their $cs\bar{c}\bar{s}$ tetraquark partners,” *Phys. Rev. D* **94**, no. 9, 094031 (2016) [arXiv:1608.07900 [hep-ph]].
- [52] K. Chen, X. Liu, J. Wu, Y. R. Liu and S. L. Zhu, “Triply heavy tetraquark states with the $QQ\bar{Q}\bar{q}$ configuration,” *Eur. Phys. J. A* **53**, no. 1, 5 (2017) [arXiv:1609.06117 [hep-ph]].
- [53] J. Wu, Y. R. Liu, K. Chen, X. Liu and S. L. Zhu, “Heavy-flavored tetraquark states with the $QQ\bar{Q}\bar{Q}$ configuration,” *Phys. Rev. D* **97**, no. 9, 094015 (2018) [arXiv:1605.01134 [hep-ph]].
- [54] J. Wu, Y. R. Liu, K. Chen, X. Liu and S. L. Zhu, “Hidden-charm pentaquarks and their hidden-bottom and B_c -like partner states,” *Phys. Rev. D* **95**, no. 3, 034002 (2017) [arXiv:1701.03873 [hep-ph]].
- [55] Q. S. Zhou, K. Chen, X. Liu, Y. R. Liu and S. L. Zhu, “Surveying exotic pentaquarks with the typical $QQqq\bar{q}$ configuration,” *Phys. Rev. C* **98**, no. 4, 045204 (2018) [arXiv:1801.04557 [hep-ph]].
- [56] J. Wu, X. Liu, Y. R. Liu and S. L. Zhu, “Systematic studies of charmonium-, bottomonium-, and B_c -like tetraquark states,” *Phys. Rev. D* **99**, no. 1, 014037 (2019) [arXiv:1810.06886 [hep-ph]].
- [57] H. T. An, Q. S. Zhou, Z. W. Liu, Y. R. Liu and X. Liu, “Exotic pentaquark states with the $qqQQ\bar{Q}$ configuration,” *Phys. Rev. D* **100**, no. 5, 056004 (2019) [arXiv:1905.07858 [hep-ph]].
- [58] J. B. Cheng and Y. R. Liu, “ $P_c(4457)^+$, $P_c(4440)^+$, and $P_c(4312)^+$: molecules or compact pentaquarks?,” *Phys. Rev. D* **100**, no. 5, 054002 (2019) [arXiv:1905.08605 [hep-ph]].
- [59] S. Y. Li, Y. R. Liu, Y. N. Liu, Z. G. Si and J. Wu, “Pentaquark states with the $QQQq\bar{q}$ configuration in a simple model,” *Eur. Phys. J. C* **79**, 87 (2019) [arXiv:1809.08072 [hep-ph]].
- [60] X. Z. Weng, X. L. Chen and W. Z. Deng, “Masses of doubly heavy-quark baryons in an extended chromomagnetic model,” *Phys. Rev. D* **97**, no. 5, 054008 (2018) [arXiv:1801.08644 [hep-ph]].
- [61] M. Anselmino, E. Predazzi, S. Ekelin, S. Fredriksson and D. B. Lichtenberg, “Diquarks,” *Rev. Mod. Phys.* **65**, 1199 (1993).
- [62] D. B. Lichtenberg, “Symmetry and Supersymmetry in Hadrons Containing Both Heavy and Light Quarks,” *J. Phys. G* **16**, 1599 (1990).
- [63] J. Carlson, L. Heller and J. A. Tjon, “Stability of Dimesons,” *Phys. Rev. D* **37**, 744 (1988).
- [64] M. J. Savage and M. B. Wise, “Spectrum of baryons with two heavy quarks,” *Phys. Lett. B* **248**, 177 (1990).
- [65] N. Brambilla, A. Vairo and T. Rosch, “Effective field theory Lagrangians for baryons with two and three heavy quarks,” *Phys. Rev. D* **72**, 034021 (2005) [hep-ph/0506065].
- [66] S. Fleming and T. Mehen, “Doubly heavy baryons, heavy quark-diquark symmetry and NRQCD,” *Phys. Rev. D* **73**, 034502 (2006) [hep-ph/0509313].
- [67] T. D. Cohen and P. M. Hohler, “Doubly heavy hadrons and the domain of validity of doubly heavy diquark-anti-quark symmetry,” *Phys. Rev. D* **74**, 094003 (2006) [hep-ph/0606084].
- [68] M. Karliner and J. L. Rosner, “Baryons with two heavy quarks: Masses, production, decays, and detection,” *Phys. Rev. D* **90**, no. 9, 094007 (2014) [arXiv:1408.5877 [hep-ph]].
- [69] J. B. Cheng, S. Y. Li, Y. R. Liu, Y. N. Liu, Z. G. Si and T. Yao, “Spectrum and rearrangement decays of tetraquark states with four different flavors,” *Phys. Rev. D* **101**, no.11, 114017 (2020) [arXiv:2001.05287 [hep-ph]].
- [70] A. De Rujula, H. Georgi and S. L. Glashow, “Hadron Masses in a Gauge Theory,” *Phys. Rev. D* **12**, 147 (1975).
- [71] S. Godfrey and N. Isgur, “Mesons in a Relativized Quark Model with Chromodynamics,” *Phys. Rev. D* **32**, 189 (1985).
- [72] B. Keren-Zur, “Testing confining potentials through meson/baryon hyperfine splitting ratio,” *Annals Phys.* **323**, 631 (2008) [hep-ph/0703011 [HEP-PH]].
- [73] H. J. Lipkin, “Relations Between Meson and Baryon Hyperfine Splittings,” *Phys. Lett. B* **171**, 293 (1986).
- [74] M. Tanabashi *et al.*, “Review of Particle Physics,” *Phys. Rev. D* **98**, no. 3, 030001 (2018).
- [75] X. Z. Weng, X. L. Chen, W. Z. Deng and S. L. Zhu, “Hidden-charm pentaquarks and P_c states,” *Phys. Rev. D* **100**, no. 1, 016014 (2019) [arXiv:1904.09891 [hep-ph]].
- [76] H. Miyazawa, “Baryon Number Changing Currents,” *Prog. Theor. Phys.* **36**, no. 6, 1266 (1966).
- [77] H. Miyazawa, “Spinor Currents and Symmetries of Baryons and Mesons,” *Phys. Rev.* **170**, 1586 (1968).
- [78] C. s. Gao and T. h. Ho, *Commun. Theor. Phys.* **1**, 761 (1982).
- [79] H. B. Thacker, E. Eichten and J. C. Sexton, “The Three-Body Potential for Heavy Quark Baryons in Lattice QCD,” *Nucl. Phys. Proc. Suppl.* **4**, 234 (1988).
- [80] D. W. Duke and J. F. Owens, Singapore, Singapore: World Scientific (1985) 441p
- [81] A. V. Manohar and M. B. Wise, “Heavy quark physics,” *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **10**, 1 (2000).
- [82] D. B. Lichtenberg, R. Roncaglia and E. Predazzi, “Mass sum rules for singly and doubly heavy flavored hadrons,” *Phys. Rev. D* **53**, 6678 (1996) [hep-ph/9511461].
- [83] E. J. Eichten and C. Quigg, “Heavy-quark symmetry implies stable heavy tetraquark mesons $Q_i Q_j \bar{q}_k \bar{q}_l$,” *Phys. Rev. Lett.* **119**, no. 20, 202002 (2017) [arXiv:1707.09575 [hep-ph]].

- [84] E. Braaten, L. P. He and A. Mohapatra, “Masses of Doubly Heavy Tetraquarks with Error Bars,” [arXiv:2006.08650 [hep-ph]].
- [85] K. W. Wei, B. Chen, N. Liu, Q. Q. Wang and X. H. Guo, “Spectroscopy of singly, doubly, and triply bottom baryons,” Phys. Rev. D **95**, no. 11, 116005 (2017) [arXiv:1609.02512 [hep-ph]].
- [86] Z. S. Brown, W. Detmold, S. Meinel and K. Orginos, “Charmed bottom baryon spectroscopy from lattice QCD,” Phys. Rev. D **90**, no. 9, 094507 (2014) [arXiv:1409.0497 [hep-lat]].
- [87] M. Karliner and J. L. Rosner, “Strange baryons with two heavy quarks,” Phys. Rev. D **97**, no. 9, 094006 (2018) [arXiv:1803.01657 [hep-ph]].
- [88] D. Ebert, R. N. Faustov, V. O. Galkin and A. P. Martynenko, “Mass spectra of doubly heavy baryons in the relativistic quark model,” Phys. Rev. D **66**, 014008 (2002) [hep-ph/0201217].
- [89] W. Roberts and M. Pervin, “Heavy baryons in a quark model,” Int. J. Mod. Phys. A **23**, 2817 (2008) [arXiv:0711.2492 [nucl-th]].
- [90] C. Albertus, E. Hernandez, J. Nieves and J. M. Verde-Velasco, “Static properties and semileptonic decays of doubly heavy baryons in a nonrelativistic quark model,” Eur. Phys. J. A **32**, 183 (2007) Erratum: [Eur. Phys. J. A **36**, 119 (2008)] [hep-ph/0610030].
- [91] B. Silvestre-Brac, “Spectrum and static properties of heavy baryons,” Few Body Syst. **20**, 1 (1996).
- [92] D. H. He, K. Qian, Y. B. Ding, X. Q. Li and P. N. Shen, “Evaluation of spectra of baryons containing two heavy quarks in bag model,” Phys. Rev. D **70**, 094004 (2004) [hep-ph/0403301].
- [93] M.-H. Weng, X.-H. Guo and A. W. Thomas, “Bethe-Salpeter equation for doubly heavy baryons in the covariant instantaneous approximation,” Phys. Rev. D **83**, 056006 (2011) [arXiv:1012.0082 [hep-ph]].
- [94] J. M. Richard, A. Valcarce and J. Vijande, Annals Phys. **412**, 168009 (2020) [arXiv:1910.08295 [nucl-th]].
- [95] P. Hasenfratz, R. R. Horgan, J. Kuti and J. M. Richard, “Heavy Baryon Spectroscopy in the QCD Bag Model,” Phys. Lett. B **94**, 401-404 (1980)
- [96] S. Ohkoda, Y. Yamaguchi, S. Yasui, K. Sudoh and A. Hosaka, “Exotic mesons with double charm and bottom flavor,” Phys. Rev. D **86**, 034019 (2012) [arXiv:1202.0760 [hep-ph]].
- [97] N. Li, Z. F. Sun, X. Liu and S. L. Zhu, “Coupled-channel analysis of the possible $D^{(*)}D^{(*)}$, $\bar{B}^{(*)}\bar{B}^{(*)}$ and $D^{(*)}\bar{B}^{(*)}$ molecular states,” Phys. Rev. D **88**, no.11, 114008 (2013) [arXiv:1211.5007 [hep-ph]].
- [98] S. Sakai, L. Roca and E. Oset, “Charm-beauty meson bound states from $B(B^*)D(D^*)$ and $B(B^*)\bar{D}(\bar{D}^*)$ interaction,” Phys. Rev. D **96**, no.5, 054023 (2017) [arXiv:1704.02196 [hep-ph]].
- [99] H. Xu, B. Wang, Z. W. Liu and X. Liu, “ DD^* potentials in chiral perturbation theory and possible molecular states,” Phys. Rev. D **99**, no. 1, 014027 (2019) [arXiv:1708.06918 [hep-ph]].
- [100] B. Wang, Z. W. Liu and X. Liu, “ $\bar{B}^{(*)}\bar{B}^{(*)}$ interactions in chiral effective field theory,” Phys. Rev. D **99**, no.3, 036007 (2019) [arXiv:1812.04457 [hep-ph]].
- [101] A. Ali, A. Y. Parkhomenko, Q. Qin and W. Wang, “Prospects of discovering stable double-heavy tetraquarks at a Tera-Z factory,” Phys. Lett. B **782**, 412-420 (2018) [arXiv:1805.02535 [hep-ph]].
- [102] A. Esposito, M. Papinutto, A. Pilloni, A. D. Polosa and N. Tantalo, “Doubly charmed tetraquarks in B_c and Ξ_{bc} decays,” Phys. Rev. D **88**, no.5, 054029 (2013) [arXiv:1307.2873 [hep-ph]].
- [103] A. Ali, Q. Qin and W. Wang, “Discovery potential of stable and near-threshold doubly heavy tetraquarks at the LHC,” Phys. Lett. B **785**, 605-609 (2018) [arXiv:1806.09288 [hep-ph]].
- [104] C. E. Fontoura, G. Krein, A. Valcarce and J. Vijande, “Production of exotic tetraquarks $QQ\bar{q}\bar{q}$ in heavy-ion collisions at the LHC,” Phys. Rev. D **99**, no.9, 094037 (2019) [arXiv:1905.03877 [hep-ph]].
- [105] S. Cho *et al.* [ExHIC Collaboration], “Multi-quark hadrons from Heavy Ion Collisions,” Phys. Rev. Lett. **106**, 212001 (2011) [arXiv:1011.0852 [nucl-th]].
- [106] S. Cho *et al.* [ExHIC Collaboration], “Studying Exotic Hadrons in Heavy Ion Collisions,” Phys. Rev. C **84**, 064910 (2011) [arXiv:1107.1302 [nucl-th]].
- [107] J. Hong, S. Cho, T. Song and S. H. Lee, “Hadronic effects on the $cc\bar{q}\bar{q}$ tetraquark state in relativistic heavy ion collisions,” Phys. Rev. C **98**, no.1, 014913 (2018) [arXiv:1804.05336 [nucl-th]].
- [108] Y. Jin, S. Y. Li, Y. R. Liu, Z. G. Si and T. Yao, “Search for a doubly charmed hadron at B factories,” Phys. Rev. D **89**, no. 9, 094006 (2014) [arXiv:1401.6652 [hep-ph]].
- [109] Z. G. Wang, “Analysis of the $\frac{1}{2}^+$ doubly heavy baryon states with QCD sum rules,” Eur. Phys. J. A **45**, 267-274 (2010) [arXiv:1001.4693 [hep-ph]].
- [110] Z. G. Wang, “Analysis of the $\frac{3}{2}^+$ heavy and doubly heavy baryon states with QCD sum rules,” Eur. Phys. J. C **68**, 459-472 (2010) [arXiv:1002.2471 [hep-ph]].