

GELFAND PROBLEM ON A LARGE SPHERICAL CAP

YOSHITSUGU KABEYA AND VITALY MOROZ

ABSTRACT. We study the behaviour of the minimal solution to the Gelfand problem on a spherical cap under the Dirichlet boundary conditions. The asymptotic behaviour of the solution is discussed as the cap approaches the whole sphere. The results are based on the sharp estimate of the torsion function of the spherical cap in terms of the principle eigenvalue which we derive in this work.

1. INTRODUCTION

We consider the nonlinear problem

$$(1.1) \quad \begin{cases} -\Delta_{\mathbb{S}^N} u = \lambda f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\lambda > 0$ is a parameter, $\Delta_{\mathbb{S}^N}$ denotes the Laplace-Beltrami operator on the unit sphere $\mathbb{S}^N \subset \mathbb{R}^{N+1}$ ($N \geq 1$) and $\Omega \subset \mathbb{S}^N$ is a sub-domain in \mathbb{S}^N with a smooth boundary $\partial\Omega \neq \emptyset$. The principal Dirichlet eigenvalue of $-\Delta_{\mathbb{S}^N}$ in Ω is denoted by $\lambda_1(\Omega) > 0$, and $\varphi_{1,\Omega}$ denotes the corresponding positive Dirichlet eigenfunction normalized as $\|\varphi_{1,\Omega}\|_2 = 1$. By w_Ω we denote the torsion function of Ω , that is the unique solution of the Dirichlet problem

$$(1.2) \quad \begin{cases} -\Delta w_\Omega = 1 & \text{in } \Omega, \\ w_\Omega = 0 & \text{on } \partial\Omega. \end{cases}$$

By the standard elliptic regularity, $w_\Omega \in C^2(\Omega)$.

We shall assume that the nonlinearity $f \in C^2(\mathbb{R})$ is a convex monotone increasing function with $f(0) > 0$ that satisfies the Keller-Osserman assumption

$$\int_0^\infty \frac{ds}{f(s)} < \infty.$$

Typical examples include

$$f(s) = \exp(s), \quad f(s) = (1+s)^p \quad (p > 1).$$

We denote

$$(1.3) \quad a_* := \min_{s>0} \frac{f(s)}{s}, \quad s_* := \min \left\{ s > 0 : \frac{f(s)}{s} = a_* \right\}.$$

By convexity and since $f(0) > 0$, we observe that $a_* > 0$ and $s_* > 0$. Denote

$$\lambda_\Omega^* := \sup \{ \lambda > 0 : (1.1) \text{ has a classical positive solution} \}.$$

The following proposition is standard.

Date: August 4, 2020.

2010 Mathematics Subject Classification. 35J60; 33C55, 35R01.

Key words and phrases. Gelfand problem, spherical cap, torsion function, eigenvalues, spherical harmonics.

Proposition 1.1. *For each $\lambda \in (0, \lambda_\Omega^*)$, problem (1.1) admits a unique minimal classical positive solution u_λ . Moreover,*

i) the following estimate holds,

$$(1.4) \quad \frac{1}{a_* \|w_\Omega\|_\infty} \leq \lambda_\Omega^* \leq \frac{\lambda_1(\Omega)}{a_*}.$$

ii) Problem (1.1) admits a unique extremal solution $u^ \in H_0^1(\Omega)$ defined as*

$$(1.5) \quad u^*(x) := \lim_{\lambda \rightarrow \lambda_\Omega^*} u_\lambda(x).$$

Remark 1.1. The bound (1.4) implies implicitly that

$$(1.6) \quad \|w_\Omega\|_\infty > \frac{1}{\lambda_1(\Omega)}.$$

It is easy to prove this directly. Indeed, let $\varphi \in C_0^\infty(\Omega)$. Testing (1.2) against $\frac{\varphi^2}{w_\Omega}$ and using Picone's identity, we conclude that

$$(1.7) \quad \int_\Omega |\nabla \varphi|^2 dS \geq \int_\Omega \frac{\varphi^2}{w_\Omega} dS,$$

cf. [1, Theorem 3.3] for Riemannian manifolds setting, or [7, Lemma A.9]. By density, (1.7) is also valid for all $\varphi \in H_0^1(\Omega)$. Then taking $\varphi_{1,\Omega}$ as a test function in (1.7) and using $\|\varphi_{1,\Omega}\|_2 = 1$ and the fact that w_Ω is non-constant, we conclude that

$$\lambda_1(\Omega) = \int_\Omega |\nabla \varphi_{1,\Omega}|^2 dS \geq \int_\Omega \frac{\varphi_{1,\Omega}^2}{w_\Omega} dS > \frac{1}{\|w_\Omega\|_\infty}.$$

Remark 1.2. As a bi-product of the proof of (1.4), we actually establish that the following pointwise estimate holds

$$(1.8) \quad \lambda f(0) w_\Omega \leq u_\lambda \leq s_* \frac{w_\Omega}{\|w_\Omega\|_\infty},$$

for all $\lambda \leq \frac{1}{a_* \|w_\Omega\|_\infty}$.

In this work we are primarily interested in the special case when

$$(1.9) \quad \Omega_\varepsilon := \left\{ (x_1, x_2, \dots, x_{N+1}) \in \mathbb{R}^{N+1} \mid \sum_{i=1}^{N+1} x_i^2 = 1, \cos((1-\varepsilon)\pi) < x_{N+1} \leq 1 \right\}.$$

is a geodesic ball, also called a ‘‘spherical cap’’, centred at the North Pole of \mathbb{S}^N , and where $\varepsilon \in (0, 1)$. Throughout the paper, for $\varepsilon \ll 1$ and $f(\varepsilon), g(\varepsilon) \geq 0$, we use the following asymptotic notation:

$f(\varepsilon) \lesssim g(\varepsilon)$ if there exists $C > 0$ independent of ε such that $f(\varepsilon) \leq Cg(\varepsilon)$;

$f(\varepsilon) \simeq g(\varepsilon)$ if $f(\varepsilon) \lesssim g(\varepsilon)$ and $g(\varepsilon) \lesssim f(\varepsilon)$;

We also use the standard Landau symbols $f = O(g)$ and $f = o(g)$, with the understanding that $f \geq 0$ and $g \geq 0$. As usual, C, c, c_1 , etc., denote generic positive constants independent of ε .

It is known by Bandle, Kabeya and Ninomiya [2], Kabeya, Kawakami, Kosaka and Ninomiya [5], Kosaka [6] and Macdonald [8] (we will review this in Sections 3), that the principle eigenvalue of Ω_ε satisfy the following asymptotic bound,

$$(1.10) \quad \lambda_1(\Omega_\varepsilon) = \begin{cases} c_N \varepsilon^{N-2} (1 + o(1)) & \text{if } N \geq 4, \\ 2\varepsilon (1 + o(1)) & \text{if } N = 3, \\ \frac{1}{2} \log^{-1} \left(\frac{2}{\varepsilon} \right) (1 + o(1)) & \text{if } N = 2, \end{cases}$$

where coefficients $c_N > 0$ do not depend on $\varepsilon > 0$ and could be computed explicitly as in [2].

In this work we establish the following new estimate on $\|w_{\Omega_\varepsilon}\|_\infty$ which shows that the lower bound (1.6) is asymptotically sharp as $\varepsilon \rightarrow 0$, and which we believe is of independent interest.

Proposition 1.2. *Let $N \geq 2$. As $\varepsilon \rightarrow 0$, the torsion function w_{Ω_ε} of the spherical cap Ω_ε satisfies*

$$(1.11) \quad \|w_{\Omega_\varepsilon}\|_\infty = \frac{1}{\lambda_1(\Omega_\varepsilon)} + O(1).$$

Using (1.11), from (1.4) we deduce a sharp asymptotic estimate on the critical parameter $\lambda_{\Omega_\varepsilon}^*$ in the Gelfand problem (1.1).

Theorem 1.1. *Let $N \geq 2$. As $\varepsilon \rightarrow 0$,*

$$(1.12) \quad \lambda_{\Omega_\varepsilon}^* = \frac{\lambda_1(\Omega_\varepsilon)}{a_*} (1 + o(1)).$$

The paper is organised as follows. In Section 2 we sketch the proof of Proposition 1.1. In Section 3 we review several fundamental properties of the special functions and outline the derivation of the torsion function estimates which lead to the proof of Proposition 1.2. In Section 4 we prove a technical Lemma 3.2. Finally, in Sections 5, 6 we discuss several technical properties of the special functions which were used in the earlier proofs.

2. PROOF OF PROPOSITION 1.1

The proof of Proposition 1.1 is standard. We present a sketch of the proof for completeness.

Lemma 2.1. *Assume (1.1) admits positive, classical super-solution $\bar{u} > 0$. Then (1.1) admits a unique minimal, positive classical solution $u_\lambda \in C^2(\Omega)$, such that*

$$(2.1) \quad \lambda f(0) w_\Omega \leq u_\lambda \leq \bar{u}.$$

Proof. The minimal solution u_λ of (1.1) could be obtained by a monotone iteration arguments. Namely, for $n \in \mathbb{N}$ consider a sequence of functions $\{u_n\}_{n=0}^\infty$ with $u_0 = 0$ and u_n defined as

$$(2.2) \quad \begin{cases} -\Delta_{\mathbb{S}^N} u_n = \lambda f(u_{n-1}) & \text{in } \Omega, \\ u_n = 0 & \text{on } \partial\Omega. \end{cases}$$

For each n problem (2.2) admits a unique solution $u_n \in C^2(\bar{\Omega})$ and $u_n \geq u_{n-1}$ in Ω for every $n \in \mathbb{N}$. In particular,

$$u_n \geq u_1 = \lambda f(0) w_\Omega.$$

We now define the minimal solution of (1.1) as

$$(2.3) \quad u_\lambda(x) := \lim_{n \rightarrow \infty} u_n(x).$$

By the arguments similar to [10, Theorem 2.1] we conclude that u_λ defined by (2.3) satisfies $\bar{u} \geq u_\lambda > 0$ in Ω , belongs to $C^2(\bar{\Omega})$ and solves (1.1) classically. \square

The next lemma uses the notion of a weak solution. Similarly to [4], we define a weak solution of (1.1) as a non-negative function $u \in L^1(\Omega)$ such that $f(u)\delta_{\partial\Omega} \in L^1(\Omega)$, where $\delta_{\partial\Omega}(x) := \text{dist}(x, \partial\Omega)$ is the distance from x to the boundary of Ω and

$$(2.4) \quad - \int_{\Omega} u \Delta_{\mathbb{S}^N} \varphi dS = \lambda \int_{\Omega} f(u) \varphi dS,$$

for all $\varphi \in C^2(\bar{\Omega})$ with $\varphi = 0$ on $\partial\Omega$.

Lemma 2.2. *Problem (1.1) admits a minimal classical solution u_λ for $0 < \lambda < \lambda_\Omega^* < \infty$. Moreover, the extremal solution*

$$u^*(x) := \lim_{\lambda \rightarrow \lambda_\Omega^*} u_\lambda(x),$$

is a weak solution of (1.1).

Proof. First observe that u_λ is a non-decreasing function of λ . This follows from the fact that $u_{\lambda'}$ is a super-solution for problem (1.1) with $\lambda < \lambda'$. Hence, if (1.1) with $\lambda = \lambda'$ admits a classical solution, then (1.1) admits a classical solution for $\lambda \in (0, \lambda']$.

Next, it is easy to see that a multiple of the torsion function w_Ω is a super-solution for (1.1) for small enough λ . Let

$$s_* := \min \left\{ s > 0 : \frac{f(s)}{s} = a_* \right\} > 0.$$

Set

$$t^* := \frac{s_*}{\|w_\Omega\|_\infty}.$$

and take $\bar{u} := t^* w_\Omega$ and $\lambda' := \frac{1}{a_* \|w_\Omega\|_\infty}$. Then

$$(2.5) \quad - \Delta_{\mathbb{S}^N} \bar{u} = t^* = t^* \frac{f(s_*)}{a_* s_*} = \frac{f(t^* \|w_\Omega\|_\infty)}{a_* \|w_\Omega\|_\infty} \geq \lambda' f(\bar{u}),$$

i.e. \bar{u} is a positive supersolution for (1.1) for all $\lambda \leq \lambda'$. In particular, this establishes the lower bound in (1.4).

Now let us show that $\lambda_\Omega^* < \infty$. Similarly to [4, Lemma 5], we test (1.1) against the $\varphi_{1,\Omega}$. Then,

$$(2.6) \quad \lambda_1(\Omega) \int_{\Omega} u_\lambda \varphi_{1,\Omega} dS = - \int_{\Omega} u_\lambda \Delta_{\mathbb{S}^N} \varphi_{1,\Omega} dS = \lambda \int_{\Omega} f(u_\lambda) \varphi_{1,\Omega} dS \geq \lambda a_* \int_{\Omega} u_\lambda \varphi_{1,\Omega} dS.$$

We conclude that $\lambda_1(\Omega) \geq \lambda a_* > 0$, which establishes the upper bound in (1.4).

Finally, proceeding as in the proof of [4, Lemma 5], we recover that u^* is a weak solution of (1.1). \square

Lemma 2.3. *Problem (1.1) admits neither classical, nor weak solutions for $\lambda > \lambda_\Omega^*$.*

Proof. Similar to [4, Theorem 3] – we omit the details. \square

3. TORSION FUNCTION ESTIMATES

In the rest of the paper we always work on the spherical cap Ω_ε . When there is no ambiguity, we suppress the dependence on ε and write λ_1, φ_1 instead of $\lambda_1(\Omega_\varepsilon), \varphi_{1,\Omega_\varepsilon}$. We always assume that $\varepsilon \in (0, 1)$ and is close to 0.

3.1. General Setting. Let

$$0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \leq \lambda_\ell \leq \dots \quad (\ell \in \mathbb{N})$$

be the Dirichlet eigenvalues of $-\Delta_{\mathbb{S}^N}$ on the spherical cap Ω_ε , and φ_ℓ be the corresponding eigenfunctions normalized as $\|\varphi_\ell\|_2 = 1$. That is, φ_ℓ is a regular solution to the Dirichlet problem

$$(3.1) \quad -\Delta\varphi_\ell = \lambda_\ell\varphi_\ell, \quad \varphi_\ell \in H_0^1(\Omega_\varepsilon).$$

It is well-known that the sequence $\{\varphi_\ell\}$ forms the complete ortho-normal system in $L^2(\Omega)$. Then we see that the torsion function w_{Ω_ε} defined as the unique solution to (1.2) is expressed as

$$(3.2) \quad w_{\Omega_\varepsilon} = \sum_{\ell=1}^{\infty} \frac{(1, \varphi_\ell)}{\lambda_\ell} \varphi_\ell,$$

where

$$(\Phi, \Psi) = \int_{\Omega_\varepsilon} \Phi\Psi \, dS.$$

Indeed, due to the Fourier expansion of 1 and the definition of the eigenfunctions, we have

$$-\Delta w_{\Omega_\varepsilon} = \sum_{\ell=1}^{\infty} \frac{(1, \varphi_\ell)}{\lambda_\ell} (-\Delta\varphi_\ell) = \sum_{\ell=1}^{\infty} \frac{(1, \varphi_\ell)}{\lambda_\ell} (\lambda_\ell\varphi_\ell) = \sum_{\ell=1}^{\infty} (1, \varphi_\ell)\varphi_\ell = 1.$$

We are going to estimate the right-hand side of (3.2).

3.2. Three dimensional case, “radial” eigenfunctions. For simplicity, we first consider the case $N = 3$. Then

$$\Omega_\varepsilon = \{(\sin\theta \sin\phi \cos\psi, \sin\theta \sin\phi \sin\psi, \sin\theta \cos\phi, \cos\theta) \mid 0 \leq \theta < (1-\varepsilon)\pi, 0 \leq \phi \leq \pi, 0 \leq \psi < 2\pi\}.$$

First, we consider the solutions to (1.2) depending only on θ . In this case, the Laplacian yields to

$$\Delta_{\mathbb{S}^3}\varphi = \frac{1}{\sin^2\theta}(\varphi' \sin^2\theta)'$$

Hence by the direct calculation, we see that the eigenfunctions depending only on θ are of the form

$$(3.3) \quad \varphi_j(\theta) = K_j \frac{\sin \frac{j}{1-\varepsilon}\theta}{\sin\theta}$$

where K_j are normalization constants and the corresponding eigenvalues λ_j are

$$(3.4) \quad \lambda_j = \frac{j^2}{(1-\varepsilon)^2} - 1 = \frac{j^2 - 1 + 2\varepsilon - \varepsilon^2}{(1-\varepsilon)^2} = 2\varepsilon j^2 + O(\varepsilon^2).$$

Note that $\lim_{\varepsilon \rightarrow 0} \lambda_j = 0$ only when $j > 1$, and that

$$(3.5) \quad \|\varphi_j\|_\infty = \varphi_j(0) = K_j \frac{j}{1-\varepsilon} + o(\varepsilon).$$

The inner product in our case is given by

$$(\Phi, \Psi) = \int_{\Omega} \Phi \Psi \, dS = 4\pi \int_0^{(1-\varepsilon)\pi} \Phi(\theta) \Psi(\theta) \sin^2 \theta \, d\theta.$$

Then we have

$$\begin{aligned} (1, \varphi_j) &= 4\pi K_j \int_0^{(1-\varepsilon)\pi} \sin \frac{j}{1-\varepsilon} \theta \sin \theta \, d\theta \\ &= 2\pi K_j \int_0^{(1-\varepsilon)\pi} \left(\cos \frac{j-1+\varepsilon}{1-\varepsilon} \theta - \cos \frac{j+1-\varepsilon}{1-\varepsilon} \theta \right) d\theta. \end{aligned}$$

Hence we obtain

$$\begin{aligned} (1, \varphi_j) &= (-1)^j 2\pi K_j (1-\varepsilon) \left\{ \frac{\sin(j-2)\varepsilon\pi + O(\varepsilon^3)}{j-1+\varepsilon} - \frac{\sin j\varepsilon\pi + O(\varepsilon^3)}{j+1-\varepsilon} \right\} \\ &= -(-1)^j 2\pi K_j (1-\varepsilon) \frac{2\varepsilon\pi + O(\varepsilon^3)}{(j-1+\varepsilon)(j+1-\varepsilon)}. \end{aligned}$$

As for K_j , we have

$$(\varphi_j, \varphi_j) = 4\pi K_j^2 \int_0^{(1-\varepsilon)\pi} \sin^2 \frac{j}{1-\varepsilon} \theta \, d\theta = 2\pi K_j^2 \int_0^{(1-\varepsilon)\pi} \left(1 - 2\cos \frac{2j}{1-\varepsilon} \theta \right) d\theta$$

and thus, from

$$1 = (\varphi_j, \varphi_j) = 2\pi K_j^2 (1-\varepsilon)\pi,$$

we obtain

$$(3.6) \quad K_j = \frac{1}{\sqrt{2\pi}} + O(\varepsilon).$$

We hence conclude that

$$(3.7) \quad (1, \varphi_j) = \begin{cases} \sqrt{2\pi} + O(\varepsilon), & j = 1, \\ \frac{(-1)^{j+1} 2\sqrt{2\pi}}{(j+1)(j-1)} \varepsilon + O(\varepsilon^2), & j \geq 2. \end{cases}$$

From (3.2) this implies that

$$(3.8) \quad w_{\Omega_\varepsilon} = \frac{\sqrt{2\pi}}{\lambda_1} \varphi_1 + \sum_{j=2}^{\infty} \frac{(1, \varphi_j)}{\lambda_j} \varphi_j,$$

and taking into account (3.7), (3.4), (3.5) and (3.6) we conclude that

$$(3.9) \quad \left| \sum_{j=2}^{\infty} \frac{(1, \varphi_j)}{\lambda_j} \varphi_j \right| \leq \sum_{j=2}^{\infty} O(j^{-4}) \|\varphi_j\|_\infty \leq C.$$

Since $\|\varphi_1\|_\infty = K_1(1-\varepsilon)^{-1}$, using (3.6) we estimate

$$\|w_{\Omega_\varepsilon}\|_\infty \leq \frac{\sqrt{2\pi}}{\lambda_1} \|\varphi_1\|_\infty + O(1) = \frac{1}{\lambda_1} + O(1).$$

Combined with (1.6) and (3.4) this leads to the two sided bound

$$(3.10) \quad \|w_{\Omega_\varepsilon}\|_\infty = \frac{1}{\lambda_1} + O(1) = \frac{1}{2\varepsilon} + O(1).$$

Remark 3.1. We will see in Lemma 3.1 that

$$\lim_{\varepsilon \rightarrow 0} \|\varphi_1\|_\infty = \lim_{\varepsilon \rightarrow 0} \varphi_1(0)$$

holds in all dimensions $N \geq 2$. Thus, we obtain

$$\lim_{\varepsilon \rightarrow 0} (1, \varphi_1) \|\varphi_1\|_\infty = 1$$

and then

$$(3.11) \quad \|w_{\Omega_\varepsilon}\|_\infty = \frac{1}{\lambda_1} + O(1),$$

as in the case $N = 3$ above.

3.3. General dimensions, “radial” eigenfunctions. In the general case $N \geq 3$ (the case $N = 2$ is exceptional and is discussed in Subsection 3.4), the spherical cap we consider is represented as follows:

$$\Omega_\varepsilon = \left\{ (x_1, x_2, \dots, x_{N+1}) \mid \sum_{i=1}^{N+1} x_i^2 = 1, \cos(1 - \varepsilon)\pi < x_{N+1} \leq 1 \right\}.$$

We introduce the polar coordinates

$$\left\{ \begin{array}{l} x_1 = \left(\prod_{j=1}^{N-1} \sin \theta_j \right) \sin \phi, \\ x_2 = \left(\prod_{j=1}^{N-1} \sin \theta_j \right) \cos \phi, \\ x_{N-k+1} = \left(\prod_{j=1}^k \sin \theta_j \right) \cos \theta_{k+1}, \quad k = 1, 2, \dots, N-2, \\ x_{N+1} = \cos \theta_1. \end{array} \right.$$

Then the Laplace-Beltrami operator is expressed as

$$\begin{aligned} \Delta_{\mathbb{S}^N} &= \frac{1}{\sin^{N-1} \theta_1} \frac{\partial}{\partial \theta_1} \left[\sin^{N-1} \theta_1 \frac{\partial}{\partial \theta_1} \right] \\ &+ \frac{1}{\sin^2 \theta_1 \sin^{N-2} \theta_2} \frac{\partial}{\partial \theta_2} \left[\sin^{N-2} \theta_2 \frac{\partial}{\partial \theta_2} \right] \\ &+ \dots \\ &+ \frac{1}{\sin^2 \theta_1 \sin^2 \theta_2 \dots \sin^2 \theta_{N-2} \sin \theta_{N-1}} \frac{\partial}{\partial \theta_{N-1}} \left[\sin \theta_{N-1} \frac{\partial}{\partial \theta_{N-1}} \right] \\ &+ \frac{1}{\sin^2 \theta_1 \dots \sin^2 \theta_{N-1}} \frac{\partial^2}{\partial \phi^2}. \end{aligned}$$

Hence

$$\Delta_{\mathbb{S}^N} = \frac{1}{\sin^{N-1} \theta_1} \frac{\partial}{\partial \theta_1} \left[\sin^{N-1} \theta_1 \frac{\partial}{\partial \theta_1} \right] + \frac{1}{\sin^2 \theta_1} \Delta_{S^{N-1}}, \quad \Delta_{\mathbb{S}^1} = \frac{\partial^2}{\partial \phi^2}.$$

Here we note that $\Delta_{\mathbb{S}^{N-1}}$ is a Laplace-Beltrami operator on \mathbb{S}^{N-1} with the coordinates $\{\theta_2, \dots, \theta_{N-1}, \phi\}$.

It is known (cf. Bandle, Kabeya and Ninomiya [2]) that the Dirichlet eigenfunctions for $-\Delta_{\mathbb{S}^N}$ on Ω_ε depending only on $\theta \in [0, (1-\varepsilon)\pi]$, which are regular solutions to

$$(3.12) \quad \frac{d}{d\theta} \left(\sin^{N-1} \theta \frac{d}{d\theta} \varphi \right) + \lambda \sin^{N-1} \theta \varphi = 0$$

are expressed as a constant multiple of

$$(3.13) \quad \varphi_j(\theta_1) = K_{j,N} \frac{P_{\nu_j}^\mu(\cos \theta_1)}{(\sin \theta_1)^{(N-2)/2}}, \quad j = 1, 2, 3, \dots$$

with

$$\mu = (-1)^N \frac{N-2}{2}$$

and the normalizing constant $K_{j,N}$ so that $\|\varphi_j\|_2 = 1$. Here, P_ν^μ is the associated Legendre function defined in terms of the Gauss hypergeometric function $F(\alpha, \beta, \gamma; x)$ as follows:

$$(3.14) \quad P_\nu^\mu(t) = \frac{1}{\Gamma(1-\mu)} \left(\frac{1+t}{1-t} \right)^{\mu/2} F \left(-\nu, \nu+1, 1-\mu; \frac{1-t}{2} \right), \quad t \in (-1, 1),$$

when $\mu \notin \mathbb{N}$ with $F(\alpha, \beta, \gamma; x)$ being defined as

$$F(\alpha, \beta, \gamma; x) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\beta)} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+n)\Gamma(\beta+n)}{\Gamma(\gamma+n)} \frac{x^n}{n!}.$$

If μ is a natural number, $P_\nu^\mu(t)$ is defined as

$$(3.15) \quad P_\nu^\mu(t) = \frac{\Gamma(\mu+\nu+1)}{\mu! 2^\mu \Gamma(1+\nu-\mu)} (1-t^2)^{\mu/2} F \left(\mu-\nu, \mu+\nu+1, \mu+1; \frac{1-t}{2} \right), \quad t \in (-1, 1).$$

Also, it is known that $P_\nu^\mu(t)$ is a solution to the associated Legendre differential equation

$$(3.16) \quad (1-t^2) \frac{d^2 u}{dt^2} - 2t \frac{du}{dt} + \left\{ \nu(\nu+1) - \frac{\mu^2}{1-t^2} \right\} u = 0.$$

Also, $P_\nu^\mu(t)/(1-t^2)^{-\mu/2}$ is a solution to the ultra-sphere differential equation

$$(3.17) \quad (1-t^2) \frac{d^2 u}{dt^2} - 2(\mu+1)t \frac{du}{dt} + (\nu-\mu)(\nu+\mu+1)u = 0.$$

If we consider all the eigenfunctions, which depend on more than two variables (we discuss this in details later in Sections 3.5, 3.6), those of $-\Delta$ in Ω_ε are represented in terms of the polar coordinate as

$$(3.18) \quad \tilde{\Phi}_\ell = K_\ell \frac{\hat{P}_\nu^{k_2+(N-2)/2}(\cos \theta_1)}{(\sin \theta_1)^{(N-2)/2}} \prod_{j=1}^{N-2} \frac{\hat{P}_{k_{N-j+1}+(j-1)/2}^{k_{N-j}+(j-1)/2}(\cos \theta_{N-j})}{(\sin \theta_{N-j})^{(j-1)/2}} \times \begin{cases} \cos k_N \phi, \\ \sin k_N \phi, \end{cases}$$

where ℓ is a nonnegative integer and where $\{k_s\}$ is a sequence of integers such that

$$(3.19) \quad 0 \leq k_N \leq k_{N-1} \leq \dots \leq k_{j+1} \leq k_j \leq \dots \leq k_2 \leq \ell.$$

The real number ν must be chosen so that $\hat{P}_\nu^{k_2+(N-2)/2}$ solves (3.1). Here we define

$$\hat{P}_\nu^\mu(\cos \theta) = \begin{cases} P_\nu^\mu(\cos \theta_1) & \text{if } \mu \text{ is an integer,} \\ P_\nu^{-\mu}(\cos \theta_1) & \text{if } \mu \text{ is a half integer,} \end{cases}$$

where θ_1 varies in $[0, (1 - \varepsilon)\pi]$, $\theta_i \in [0, \pi]$ ($i = 2, 3, \dots, N - 1$) and $\phi \in [0, 2\pi]$.

Note that $\ell = 0$ corresponds to the first eigenvalue and that the corresponding eigenfunction depends only on θ_1 according to (3.19). First, we study the case when all the eigenfunctions depend only on θ_1 . Then, for the Dirichlet problem, by Bandle, Kabeya and Ninomiya [2] or Macdonald [8], there holds

$$(3.20) \quad \nu_j = (j - 1) + \frac{N - 2}{2} + C_j \varepsilon^{N-2} + o(\varepsilon^{N-2}),$$

where C_j is a constant explicitly determined in [2], which has a complicated expression.

The relation between ν_j and the eigenvalue λ_j is expanded as

$$(3.21) \quad \lambda_j = \nu_j(\nu_j + 1) - \frac{N(N - 2)}{4} = (j - 1)(j + N - 2) + (2j + N - 3)C_j \varepsilon^{N-2} + o(\varepsilon^{N-2}).$$

Note that the first eigenvalue

$$(3.22) \quad \lambda_1 = (N - 1)C_1 \varepsilon^{N-2} + o(\varepsilon^{N-2})$$

is close to zero.

In order to estimate the maximum of the torsion function $\|w_{\Omega_\varepsilon}\|_\infty$, we need the following formula on the definite integral formula, which is valid for any dimension, and which is written on p. 129 in Moriguchi, Udagawa and Hitotsumatsu [9] as

$$(3.23) \quad \int_0^\pi P_\nu^\mu(\cos \theta) \sin^{\alpha-1} \theta d\theta = \frac{2^\mu \pi \Gamma((\alpha + \mu)/2) \Gamma((\alpha - \mu)/2)}{\Gamma((\alpha - \nu)/2) \Gamma((\nu + \alpha + 1)/2) \Gamma((- \mu - \nu + 1)/2) \Gamma((\nu - \mu + 2)/2)}.$$

The expression is valid for any μ , $\alpha \in \mathbb{R}$ such that $|\mu| < \alpha$. In this case, $|\mu| = (N - 2)/2 < N/2 + 1$. Thus, (3.23) is applicable.

3.4. Two dimensional case. Due to Macdonald [8] (see also Section 3 of Kabeya, Kawakami, Kosaka and Ninomiya [5] in the case of the Dirichlet boundary condition or Kosaka [6]), as $\varepsilon \rightarrow 0$, we see that

$$(3.24) \quad \nu_j = j - 1 + \frac{1}{2 \log \left(\frac{2}{\varepsilon}\right)} + o((\log \varepsilon^{-1})^{-1})$$

and

$$(3.25) \quad \lambda_j = \nu_j(\nu_j + 1) = j(j - 1) + \frac{2j - 1}{2 \log \left(\frac{2}{\varepsilon}\right)} + o((\log \varepsilon^{-1})^{-1})$$

holds. We apply (3.23) with $\nu = \nu_j$ in (3.24) and

$$\mu = 0, \quad \alpha = 2.$$

Then there hold

$$\alpha - \nu = 3 - j - \frac{1}{2 \log \left(\frac{2}{\varepsilon}\right)} + o((\log \varepsilon^{-1})^{-1})$$

and

$$-\mu - \nu + 1 = 2 - j - \frac{1}{2 \log \left(\frac{2}{\varepsilon} \right)} + o((\log \varepsilon^{-1})^{-1}).$$

Thus, when $j = 1$, we see that

$$\begin{aligned} \int_0^\pi P_\nu^0(\cos \theta) \sin \theta \, d\theta &= \frac{(\Gamma(1))^2 \pi}{(\Gamma(1))^2 \Gamma(3/2) \Gamma(1/2)} + O\left(\frac{1}{2 \log \left(\frac{2}{\varepsilon} \right)}\right) \\ &= \frac{\pi}{\Gamma(3/2) \Gamma(1/2)} + O\left(\frac{1}{2 \log \left(\frac{2}{\varepsilon} \right)}\right) = 2 + O\left(\frac{1}{2 \log \left(\frac{2}{\varepsilon} \right)}\right) \neq 0. \end{aligned}$$

When $j \geq 2$, we should be careful. Since $\Gamma(-n) = \infty$ ($n = 0, 1, 2, \dots$) (see (i) of Lemma 6.1 below), or more precisely, $z = -n$ is the pole of order one for $\Gamma(z)$, if $(-\mu - \nu + 1)/2$ or $(\alpha - \nu)/2$ are non-positive integers, we regard

$$\int_0^\pi P_\nu^\mu(\cos \theta) \sin^{\alpha-1} \theta \, d\theta = 0.$$

In our case, $(-\mu - \nu + 1)/2$ or $(\alpha - \nu)/2$ are not exactly non-positive integers, but are close to those. This means that the value of the desired definite integral is small enough.

Indeed, if $j \geq 2$, then from (3.23), we have

$$\int_0^\pi P_\nu^0(\cos \theta) \sin \theta \, d\theta \simeq \frac{(\Gamma(1))^2 \pi}{\Gamma\left(\frac{3-j}{2} - \frac{C_j}{4 \log(2/\varepsilon)}\right) \Gamma\left(\frac{2-j}{2} - \frac{C_j}{4 \log(2/\varepsilon)}\right) \Gamma\left(\frac{j+2}{2}\right) \Gamma\left(\frac{j+1}{2}\right)}.$$

Thus, if $j = 2L$ with $L = 1, 2, \dots$, we see that

$$(3.26) \quad \int_0^\pi P_\nu^0(\cos \theta) \sin \theta \, d\theta \simeq \frac{(-1)^L (L-1)! \pi}{4\Gamma\left(\frac{3}{2} - L\right) \Gamma\left(L + \frac{1}{2}\right) L!} \frac{C_j}{\log\left(\frac{2}{\varepsilon}\right)}$$

and

$$(3.27) \quad \int_0^\pi P_\nu^0(\cos \theta) \sin \theta \, d\theta \simeq \frac{(-1)^{L-1} (L-2)! \pi}{4\Gamma\left(\frac{3}{2} - L\right) \Gamma\left(L + \frac{1}{2}\right) (L-1)!} \frac{C_j}{\log\left(\frac{2}{\varepsilon}\right)}$$

when $j = 2L - 1$ with $L = 2, 3, 4, \dots$

3.5. All the eigenfunctions. Three dimensional case. If we consider the eigenfunctions which depends on θ_i and/or ϕ , first we consider the case $N = 3$. Then according to (3.18), we have

$$(3.28) \quad \Phi = K_j \frac{\hat{P}_\nu^{m+1/2}(\cos \theta_1)}{(\sin \theta_1)^{1/2}} P_m^k(\cos \theta_2) (c_1 \cos k\phi + c_2 \sin k\phi).$$

In the three dimensional case, the sphere element dS is expressed as

$$dS = \sin^2 \theta_1 \sin \theta_2 \, d\theta_1 d\theta_2 d\phi.$$

If k is a natural number, it is very easy to see that

$$\begin{aligned} &\int_{\Omega_\varepsilon} \hat{\Phi} \, dS \\ &= \int_0^{(1-\varepsilon)\pi} \int_0^\pi \int_0^{2\pi} \frac{\hat{P}_\nu^{m+1/2}(\cos \theta_1)}{(\sin \theta_1)^{1/2}} P_m^k(\cos \theta_2) (c_1 \cos k\phi + c_2 \sin k\phi) \sin^2 \theta_1 \sin \theta_2 \, d\phi d\theta_2 d\theta_1 \\ &= 0 \end{aligned}$$

since ϕ varies from 0 to 2π . If $k = 0$ in (3.28), then we consider $P_m^0(\cos \theta_2)$. In this case, we need to calculate

$$\int_0^\pi P_m^0(\cos \theta_2) \sin \theta_2 d\theta_2$$

with some natural number m . In (3.23), we take $\mu = 0$, $\nu = m$, $\alpha = 2$. Hence, we have

$$\frac{\alpha - \nu}{2} = 1 - \frac{m}{2}, \quad \frac{-\mu - \nu + 1}{2} = -\frac{m-1}{2}.$$

If n is even, the former is nonpositive integer and if n is odd, the latter is so. Hence, due to (3.23), we obtain

$$\int_0^\pi P_n^0(\cos \theta_2) \sin \theta_2 d\theta_2 = 0.$$

Hence when $N = 3$, it is sufficient to consider the ‘‘radial’’ case only.

3.6. All the eigenfunctions. Four dimensional case and the general case. Similarly to the three dimensional case, the eigenfunction is of the form

$$(3.29) \quad \hat{\Phi} = K_j \frac{\hat{P}_\nu^{n+1}(\cos \theta_1)}{\sin \theta_1} \frac{\hat{P}_{n+1/2}^{m+1/2}(\cos \theta_2)}{(\sin \theta_2)^{1/2}} P_m^k(\cos \theta_3) (c_1 \cos k\phi + c_2 \sin k\phi).$$

In the four dimensional case, the sphere element dS is expressed as

$$dS = \sin^3 \theta_1 \sin^2 \theta_2 \sin \theta_3 d\theta_1 d\theta_2 d\phi.$$

As in the three dimensional case, if one of k , m is positive, then we have

$$\begin{aligned} & \int_{\Omega_\varepsilon} \hat{\Phi} dS \\ &= \int_0^{(1-\varepsilon)\pi} \int_0^\pi \int_0^\pi \int_0^{2\pi} \frac{\hat{P}_\nu^{n+1}(\cos \theta_1)}{\sin \theta_1} \frac{\hat{P}_{n+1/2}^{m+1/2}(\cos \theta_2)}{(\sin \theta_2)^{1/2}} P_m^k(\cos \theta_3) \begin{Bmatrix} c_1 \cos k\phi \\ c_2 \sin k\phi \end{Bmatrix} \times \\ & \quad \sin^3 \theta_1 \sin^2 \theta_2 \sin \theta_3 d\phi d\theta_3 d\theta_2 d\theta_1 \\ &= 0. \end{aligned}$$

Thus, we consider $k = m = 0$ and calculate

$$\int_0^\pi \hat{P}_{n+1/2}^{1/2}(\cos \theta_2) \sin^{3/2} \theta_2 d\theta_2.$$

In this case, we note that $\hat{P}_{n+1/2}^{1/2}(\cos \theta_2) = P_{n+1/2}^{-1/2}(\cos \theta_2)$. Then we apply (3.23) with

$$\alpha = \frac{5}{2}, \quad \mu = -\frac{1}{2}, \quad \nu = n + \frac{1}{2}.$$

Hence we have

$$\frac{\alpha - \nu}{2} = -\frac{n-2}{2}, \quad \frac{-\mu + \nu - 1}{2} = -\frac{n-1}{2}.$$

If n is an even integer, then $(\alpha - \nu)/2$ is a nonpositive integer and if n is odd, so is $-(\mu + \nu - 1)/2$. Thus, in the four dimensional case also, we see that

$$\int_0^\pi \hat{P}_{n+1/2}^{1/2}(\cos \theta_2) \sin^{3/2} \theta_2 d\theta_2 = 0$$

if $n \geq 1$ and we have only to consider the eigenfunctions depending only on θ_1 .

Inductively, we proved that we have only to consider the “radial” eigenfunctions to estimate the torsion function. Thus in (3.18), we have only to consider the case

$$k_2 = k_3 = \cdots = k_N = 0,$$

that is, we consider the eigenfunctions of the form

$$\frac{P_\nu^{(N-2)/2}(\cos \theta_1)}{(\sin \theta_1)^{(N-2)/2}}$$

as in (3.13).

3.7. Proof of Proposition 1.2 completed. To complete the proof of Proposition 1.2 in general dimensions $N \geq 4$, we use the representation of the torsion function w_{Ω_ε} as in (3.2). To calculate $(1, \varphi_j)$, we note that

$$(1, \varphi_j) = \omega_{N-1} \int_0^{(1-\varepsilon)\pi} \varphi_j(\theta) \sin^{N-1} \theta \, d\theta$$

where ω_{N-1} stands for the area of the unit sphere \mathbb{S}^{N-1} . Hence,

$$(1, \varphi_j) = \omega_{N-1} K_j \int_0^{(1-\varepsilon)\pi} P_\nu^\mu(\cos \theta) (\sin \theta)^{N/2} \, d\theta.$$

We recall (3.23) here and the validity of the formula. The conditions are $\mu, \alpha \in \mathbb{R}$ such that $|\mu| < \alpha$. In this case, $|\mu| = (N-2)/2 < N/2 + 1$. Thus, (3.23) is applicable.

The comments given before (3.26) apply also here. That is, $(-\mu - \nu + 1)/2$ or $(\alpha - \nu)/2$ are not exactly non-positive integers, but are close to those. Indeed, when $N = 2m$ ($m = 2, 3, \dots$), we have

$$\begin{aligned} \mu &= m - 1, \quad \nu \simeq j + m - 2 + C_j \varepsilon^{N-2}, \quad \alpha = m + 1, \\ \nu + \alpha + 1 &\simeq 2m + j + C_j \varepsilon^{N-2}, \quad \nu - \mu + 2 \simeq j + 1 + C_j \varepsilon^{N-2}. \end{aligned}$$

Then

$$\alpha - \nu \simeq 3 - j - C_j \varepsilon^{N-2}, \quad -\mu - \nu + 1 \simeq -j - 2m + 4 - C_j \varepsilon^{N-2}.$$

For $j = 1$, these do not converge to a nonpositive integer. Hence we have

$$\begin{aligned} \int_0^\pi P_\nu^\mu(\cos \theta) \sin^m \theta \, d\theta &= \frac{2^{N/2-1} \pi \Gamma(\frac{N}{2}) \Gamma(1)}{(\Gamma(1))^2 \Gamma(\frac{N+1}{2}) \Gamma(\frac{3-N}{2})} + O(\varepsilon^{N-2}) \\ &= \frac{(-1)^{N/2+1} 2^{N/2} (\frac{N}{2} - 1)!}{N-1} + O(\varepsilon^{N-2}) \neq 0 \end{aligned}$$

and

$$\begin{aligned} \int_0^\pi P_\nu^\mu(\cos \theta) \sin^m \theta \, d\theta &\simeq \frac{2^{m-1} \Gamma(m) \Gamma(1) \pi}{\Gamma(\frac{3-j}{2} - \frac{C_j}{2} \varepsilon^{N-2}) \Gamma(-m - \frac{j-4}{2} - \frac{C_j}{2} \varepsilon^{N-2}) \Gamma(m + \frac{j}{2}) \Gamma(\frac{j+1}{2})} \end{aligned}$$

for $j \geq 2$.

As in the case of $N = 2$, when $j = 2L$, we see that

$$(3.30) \quad \int_0^\pi P_\nu^\mu(\cos \theta) \sin^m \theta \, d\theta \simeq \frac{(-1)^{m+L-1} 2^{m-2} (m-1)! \pi C_j}{(m+L-1) \Gamma(\frac{3}{2} - L) \Gamma(L + \frac{1}{2})} \varepsilon^{N-2}$$

and

$$(3.31) \quad \int_0^\pi P_\nu^\mu(\cos \theta) \sin^m \theta \, d\theta \simeq \frac{(-1)^{L-1} 2^{m-2} (m-1)! \pi C_j}{L \Gamma(m+L-\frac{1}{2}) \Gamma(-m-L+\frac{5}{2})} \varepsilon^{N-2}.$$

When $N = 2m - 1$, then we have

$$\mu = -\frac{N-2}{2} = \frac{3}{2} - m, \quad \nu \simeq j + m - \frac{5}{2} + C_j \varepsilon^{N-2}, \quad \alpha = m + \frac{1}{2}.$$

Then

$$\begin{aligned} \alpha - \nu &\simeq 3 - j - C_j \varepsilon^{N-2}, & -\mu - \nu + 1 &\simeq 2 - j - C_j \varepsilon^{N-2}, \\ \alpha + \nu + 1 &\simeq 2m + j - 1 + C_j \varepsilon^{N-2}, & \nu - \mu + 2 &\simeq j + 2m - 2 + C_j \varepsilon^{N-2}. \end{aligned}$$

For $j = 1$, $-\mu - \nu + 1$ does not converge to a nonpositive integer, either. Indeed, we have

$$\begin{aligned} \int_0^\pi P_\nu^\mu(\cos \theta) \sin^{m-1/2} \theta \, d\theta &= \frac{2^{-N/2+1} \pi \Gamma(N) \Gamma(1)}{\Gamma(1) \Gamma(\frac{N+1}{2}) \Gamma(\frac{1}{2}) \Gamma(N)} + O(\varepsilon^{N-2}) \\ &= \frac{2^{-N/2+1} \pi}{\Gamma((N+1)/2) \Gamma(1/2)} + O(\varepsilon^{N-2}) \neq 0 \end{aligned}$$

and for $j \geq 2$, we get

$$\int_0^\pi P_\nu^\mu(\cos \theta) \sin^{m-1/2} \theta \, d\theta \simeq \frac{2^{-(m-3/2)} \Gamma(m-\frac{1}{2}) \Gamma(1) \pi}{\Gamma(\frac{3-j}{2} - \frac{C_j}{2} \varepsilon^{N-2}) \Gamma(\frac{2-j}{2} - \frac{C_j}{2} \varepsilon^{N-2}) \Gamma(m + \frac{j-1}{2}) \Gamma(m + \frac{j-2}{2})}$$

As in the even dimensional case, if $j = 2L$, we have

$$(3.32) \quad \int_0^\pi P_\nu^\mu(\cos \theta) \sin^{m-1/2} \theta \, d\theta \simeq \frac{(-1)^L 2^{-(m-1/2)} \Gamma(m-\frac{1}{2}) (L-1)! \pi}{\Gamma(\frac{3}{2}-L) \Gamma(m+L-\frac{1}{2}) (n+L-2)!} C_j \varepsilon^{N-2}$$

and

$$(3.33) \quad \int_0^\pi P_\nu^\mu(\cos \theta) \sin^{m-1/2} \theta \, d\theta \simeq \frac{(-1)^{L-1} 2^{-(m-1/2)} \Gamma(m-\frac{1}{2}) (L-2)! \pi}{\Gamma(\frac{3}{2}-L) \Gamma(m+L-\frac{3}{2}) (n+L-2)!} C_j \varepsilon^{N-2}$$

when $j = 2L - 1$.

Thus, for any dimension, if $j = 1$, then we have

$$\int_0^\pi P_\nu^\mu(\cos \theta) \sin^{\alpha-1} \theta \, d\theta \neq 0.$$

Also we conclude that

$$\int_0^{(1-\varepsilon)\pi} P_\nu^\mu(\cos \theta) \sin^{\alpha-1} \theta \, d\theta \neq 0,$$

when $j = 1$ for any small $\varepsilon > 0$.

Moreover, we have the following.

Lemma 3.1. *The first eigenfunction $\varphi_1(\theta)$, which is nonnegative and is monotone decreasing, converges to a constant function as $\varepsilon \rightarrow 0$ compact uniformly on $[0, \pi)$. Moreover, there hold*

$$\|\varphi_1\|_\infty \rightarrow \frac{1}{\sqrt{\omega_N}}, \quad (1, \varphi_1) \rightarrow \sqrt{\omega_N}, \quad (1, \varphi_1) \|\varphi_1\|_\infty = 1 + o(1)$$

as $\varepsilon \rightarrow 0$, where $\omega_N = \frac{2\pi^{(N+1)/2}}{\Gamma((N+1)/2)}$ is the surface area of \mathbb{S}^N .

Proof. As we consider the first eigenfunction, first we note that $\nu \rightarrow (N-2)/2$ and that $\lambda_1 \rightarrow 0$ as $\varepsilon \rightarrow 0$ according to (3.20), (3.21), (3.22) or (3.25). Moreover, the first eigenfunction is monotone decreasing in θ in view of (3.12) since φ_1 is positive due to the characterization of the first eigenfunction. Then putting $\lambda = \lambda_1$ in (3.12), due to the continuous dependence of a solution of an ordinary differential equation on the data, a regular solution to (3.12) converges to a regular solution to

$$(3.34) \quad \frac{d}{d\theta}(\sin^{N-1} \theta \frac{d}{d\theta} \varphi) = 0$$

compact uniformly in $[0, \pi)$. (3.34) implies that

$$\frac{d}{d\theta} \varphi = \frac{K}{\sin^{N-1} \theta}$$

with some constant K . In order to have a regular solution at $\theta = 0$, K must be zero and we see that a regular solution to (3.34) must be a constant.

Thus, as $\varepsilon \rightarrow 0$, $\varphi_1 \rightarrow c_N$ for a constant $c_N > 0$, compactly uniformly on $[0, \pi)$. Hence, we have

$$(1, \varphi_1) \rightarrow c_N \omega_N$$

and

$$\|\varphi_1\|_2^2 \rightarrow c_N^2 \omega_N.$$

Using the normalization assumption $\|\varphi\|_2 = 1$ we conclude that $c_N = 1/\sqrt{\omega_N}$. Hence

$$(1, \varphi_1) \|\varphi_1\|_\infty = (\sqrt{\omega_N} + o(1))(1/\sqrt{\omega_N} + o(1)) = 1 + o(1),$$

so the assertion follows. \square

Only the case $j = 1$ is exceptional. Intuitively, we see that the first eigenfunction converges to a positive constant and this makes the limit of $(1, \varphi_1)$ be away from zero as $\varepsilon \rightarrow +0$. If $j \geq 2$ then $(1, \varphi_j) \rightarrow 0$ as $\varepsilon \rightarrow +0$. More precisely, we have the following lemma which will be proved in the next section.

Lemma 3.2. *For each $\ell \geq 2$, as $\varepsilon \rightarrow 0$ there holds*

$$(3.35) \quad (1, \varphi_\ell) = \begin{cases} M_{\ell, N} \varepsilon^{N-2} + o(\varepsilon^{N-2}) & \text{if } N \geq 3, \\ M_{\ell, 2} \frac{1}{|\log \frac{\varepsilon}{2}|} + o\left(\frac{1}{|\log \frac{\varepsilon}{2}|}\right) & \text{if } N = 2, \end{cases}$$

where $M_{\ell, N}$ is a constant which is independent of ε .

Hence, using Lemmas 3.1 and 3.2, we obtain

$$(3.36) \quad \|w_{\Omega_\varepsilon}\|_\infty = \frac{1}{\lambda_1} + O(1),$$

for $N = 2$ and $N \geq 4$ exactly as in the case $N = 3$ is Section 3.2, and this completes the proof of Proposition 1.2.

4. PROOF OF LEMMA 3.2.

Although the integration in (3.23) is done over the interval $[0, \pi]$, we need to evaluate the integration over $[0, (1 - \varepsilon)\pi]$. Thus, we write

$$\int_0^{(1-\varepsilon)\pi} P_\nu^\mu(\cos \theta) \sin^{N/2} \theta d\theta = \left(\int_0^\pi - \int_{(1-\varepsilon)\pi}^\pi \right) P_\nu^\mu(\cos \theta) \sin^{N/2} \theta d\theta.$$

In order to show that the second term is much smaller than the first term, we use the asymptotic behaviour of the Legendre functions near $\theta = \pi$. To this end, we use the following formulas.

For this purpose, we regard the expression of the associated Legendre function in terms of the Gauss hypergeometric functions. We note that the associated Legendre function is expressed by using the Gauss hypergeometric function as below when μ is not a positive integer:

$$(4.1) \quad P_\nu^\mu(t) = \frac{1}{\Gamma(1-\mu)} \left(\frac{1+t}{1-t} \right)^{\mu/2} F(-\nu, \nu+1, 1-\mu; \frac{1-t}{2}).$$

Note that the right-hand side has a singularity at $t = -1$.

4.1. Odd dimensional case. When N is odd, as we take $\mu = -(N-2)/2$, from (4.1), we have

$$P_\nu^{-(N-2)/2}(\cos \theta) = \frac{1}{\Gamma(N/2)} \left(\frac{1+\cos \theta}{1-\cos \theta} \right)^{-(N-2)/4} F(-\nu, \nu+1, \frac{N}{2}; \frac{1-\cos \theta}{2}).$$

From (5.1) in Section 5, we have

$$\begin{aligned} & F(-\nu, \nu+1, \frac{N}{2}; \frac{1-\cos \theta}{2}) \\ &= \frac{\Gamma(N/2)\Gamma(\frac{N-2}{2})}{\Gamma(\frac{N}{2}+\nu)\Gamma(\frac{N-2}{2}-\nu)} F(-\nu, \nu+1, 2-\frac{N}{2}; \frac{1+\cos \theta}{2}) \\ & \quad + \frac{\Gamma(\frac{N}{2})\Gamma(-\frac{N-2}{2})}{\Gamma(-\nu)\Gamma(\nu+1)} \left(\frac{1+\cos \theta}{2} \right)^{(N-2)/2} F(\frac{N}{2}+\nu, \frac{N-2}{2}-\nu, \frac{N}{2}; \frac{1+\cos \theta}{2}). \end{aligned}$$

In general, there holds $F(a, b, c; x) = 1 + (ab/c)x + O(x^2)$ near $x = 0$ if $a, b, c \notin \{0, -1, -2, \dots\}$. Thus, $F(-\nu, \nu+1, N/2; (1-\cos \theta)/2)$ is close to 1 near $\theta = \pi$.

In the odd dimensional case ($N = 2m - 1$), we have seen in the previous section that $\nu = j + m - \frac{5}{2} + C_j \varepsilon^{N-2} + o(\varepsilon^{N-2})$. Thus, according to (i) of Lemma 6.1, there holds

$$\frac{1}{\Gamma(\frac{N-2}{2}-\nu)} = \frac{1}{\Gamma(1-j-C_j \varepsilon^{N-2} + o(\varepsilon^{N-2}))} \simeq \varepsilon^{N-2}$$

and we have

$$(1+\cos \theta)^{(N-2)/2} \lesssim \varepsilon^{N-2}$$

for $\theta \in [(1-\varepsilon)\pi, \pi]$. We note that

$$\int_{(1-\varepsilon)\pi}^\pi P_\nu^\mu(\cos \theta) \sin^{N/2} \theta d\theta \simeq \int_{(1-\varepsilon)\pi}^\pi (1+\cos \theta)^{-(N-2)/4} \sin^{N/2} \theta d\theta$$

holds. Letting $s = \cos \theta$, we have

$$\begin{aligned} \int_{(1-\varepsilon)\pi}^{\pi} (1 + \cos \theta)^{-(N-2)/4} \sin^{N/2} \theta \, d\theta &\simeq \int_{-1}^{\cos(1-\varepsilon)\pi} (1+s)^{-(N-2)/4} (1-s^2)^{N/4+1/2} \, ds \\ &\simeq \int_{-1}^{\cos(1-\varepsilon)\pi} (1+s) \, ds \simeq \varepsilon^4. \end{aligned}$$

Finally, we see that

$$\int_{(1-\varepsilon)\pi}^{\pi} P_{\nu}^{\mu}(\cos \theta) \sin^{N/2} \theta \, d\theta \lesssim \varepsilon^{N+2}.$$

4.2. Even dimensional case. Let $N = 2m$. Then we have $\alpha = m - 1 - \nu$, $\beta = m + \nu$, $\ell = m$. Thus, (4.1) is not applicable and (3.15) has some difficulty as $\cos(1-\varepsilon)\pi \rightarrow -1$. Thus, in the even dimensional case, the conversion formula (5.1) in Section 5 does not hold. Instead we use the function U defined as (5.2) in Section 5. Concerning the relation U and the Gauss hypergeometric function F , the conclusion in [3] in p. 276 is

$$(4.2) \quad F(\alpha, \beta, \ell; x) = \Gamma(\ell)U(\alpha, \beta, \alpha + \beta + 1 - \ell; 1 - x)$$

provided $\alpha + \beta + 1 - \ell$ is a non-positive integer. Also, we use the other expression as

$$(4.3) \quad P_{\nu}^{m-1}(\cos \theta) = \frac{\Gamma(m + \nu)(m - 1)!}{\Gamma(2 + \nu - m)} (\sin^{m-1} \theta) U(m - 1 - \nu, m + \nu, m; \frac{1 + \cos \theta}{2}).$$

As $\theta \rightarrow \pi - 0$, the leading term of U is $(1 + \cos \theta)^{1-m}$ when $m \geq 2$ and $\log(1 + \cos \theta)$ when $m = 1$. Also note that $\alpha = m - 1 - \nu = -j + 1 - C_j \varepsilon^{N-2} + o(\varepsilon^{N-2}) < 0$ if $j \geq 2$. Thus, $\Gamma(\alpha)$ in (5.2) behaves like ε^{N-2} order. Hence, we see that

$$\int_{(1-\varepsilon)\pi}^{\pi} P_{\nu}^{m-1}(\cos \theta) \sin^m \theta \, d\theta \simeq \varepsilon^{N-2} \int_{(1-\varepsilon)\pi}^{\pi} (1 + \cos \theta)^{1-m} \sin^{2m-1} \theta \, d\theta$$

holds. Letting $s = \cos \theta$, for $N = 2m$ with $m \geq 2$, we have

$$\begin{aligned} \int_{(1-\varepsilon)\pi}^{\pi} (1 + \cos \theta)^{1-m} \sin^{2m-1} \theta \, d\theta &\simeq \int_{-1}^{\cos(1-\varepsilon)\pi} (1+s)^{1-m} (1-s^2)^m \, ds \\ &\simeq \int_{-1}^{\cos(1-\varepsilon)\pi} (1+s) \, ds \asymp \varepsilon^4. \end{aligned}$$

When $N = 2$, we have

$$\alpha + 1 - \ell = -\nu - j - \frac{1}{2 \log(\frac{2}{\varepsilon})} + o\left(\frac{1}{2 \log(\frac{2}{\varepsilon})}\right).$$

Thus as above, we get

$$\int_{(1-\varepsilon)\pi}^{\pi} P_{\nu}^0(\cos \theta) \sin \theta \, d\theta \simeq \frac{1}{\log(\frac{2}{\varepsilon})} \int_{(1-\varepsilon)\pi}^{\pi} \log(1 + \cos \theta) \sin \theta \, d\theta.$$

Again, letting $s = \cos \theta$, we see that

$$\begin{aligned} \int_{(1-\varepsilon)\pi}^{\pi} \log(1 + \cos \theta) \sin \theta \, d\theta &\simeq \int_{-1}^{\cos(1-\varepsilon)\pi} (1+s) \log(1+s) \, ds \\ &\simeq \varepsilon^4 |\log \varepsilon| \end{aligned}$$

holds. Hence, in the even dimensional case also, we see that

$$\int_{(1-\varepsilon)\pi}^{\pi} P_{\nu}^{\mu}(\cos \theta) \sin^{N/2} \theta d\theta$$

is negligible.

Thus, all the coefficients in Lemma 3.2 are determined from (3.26), (3.27), (3.30), (3.31), (3.32) and (3.33) and the proof is now complete.

5. PROPERTIES OF THE GAUSS HYPERGEOMETRIC FUNCTIONS.

For the Gauss hypergeometric function $F(a, b, c; x)$, there is a conversion formula, which is useful for the analysis on an odd dimensional case.

Let a, b, c be real numbers such that none of $c, a+b+1-c$ and $c+1-a-b$ is a non-positive integer. Then for $x \in (-1, 1)$, there holds

$$(5.1) \quad \begin{aligned} F(a, b, c; x) &= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} F(a, b, a+b+1-c; 1-x) \\ &\quad + \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} (1-x)^{c-a-b} F(c-a, c-b, 1+c-a-b; 1-x). \end{aligned}$$

However, this conversion formula is not valid for an even dimensional case. To overcome this difficulty, we introduce the other function which is an independent solution to the Gauss hypergeometric differential equation. The singularity can be controlled according to a device found e.g. in Section 8.4 (pp. 274–276) of the book by Beals and Wong [3] by means of the function $U(\alpha, \beta, \ell; x)$. This function solves the hypergeometric differential equation

$$x(1-x)\frac{d^2F}{dx^2} + \{\ell - (\alpha + \beta + 1)x\}\frac{dF}{dx} - \alpha\beta F = 0$$

and is linearly independent of $F(\alpha, \beta, \ell; x)$.

Let α, β be non-integer values and ℓ be a positive integer. The function $U(\alpha, \beta, \ell, x)$ is defined as follows

$$(5.2) \quad \begin{aligned} &U(\alpha, \beta, \ell; x) \\ &= \frac{(-1)^{\ell}}{\Gamma(\alpha+1-\ell)\Gamma(\beta+1-\ell)(\ell-1)!} \times \\ &\quad \left[F(\alpha, \beta, \ell; x) \log x \right. \\ &\quad \left. + \sum_{i=0}^{\infty} \frac{(\alpha)_i(\beta)_i}{(\ell)_i i!} \{\psi(\alpha+i) + \psi(\beta+i) - \psi(i+1) - \psi(\ell+i)\} x^i \right] \\ &\quad + \frac{(\ell-2)!}{\Gamma(\alpha)\Gamma(\beta)} x^{1-\ell} \sum_{i=0}^{\ell-2} \frac{(\alpha+1-\ell)_i(\beta+1-\ell)_i}{(2-\ell)_i i!} x^i, \end{aligned}$$

where $\psi(z)$ is the psi (or di-Gamma) function defined as

$$\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)},$$

$(z)_0 = 1$ and $(z)_i = z(z+1)\cdots(z+i-1)$ with $i \in \mathbb{N}$. The last term in (5.2) does not exist if $\ell = 0, 1$.

6. PROPERTIES OF THE GAMMA FUNCTION

In this subsection, we recall the definition and several properties of the Gamma function. One of the definitions of the Gamma function $\Gamma(z)$ is

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{(n-1)!n^z}{z(z+1)\cdots(z+n-1)}.$$

By this definition we see that $\Gamma(z)$ has a pole of order 1 at $z = 0, -1, -2, \dots$ and the local behaviour around one of these poles.

The definition of the ψ -function is

$$\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)} = \frac{d}{dz} \log \Gamma(z).$$

The ψ -function has the following property

$$(6.1) \quad \psi(z+1) = \psi(z) + \frac{1}{z}.$$

The limiting behaviour of the Gamma function is as follows.

Lemma 6.1 (cf. Sections 1 and 3 in [9]). *Let n and k be nonnegative integers. Then:*

- (i) $\lim_{\zeta \rightarrow 0} \zeta \Gamma(-n - \zeta) = \frac{(-1)^{n+1}}{n!}$.
- (ii) $\lim_{\zeta \rightarrow 0} \zeta \psi(-n - \zeta) = 1$.
- (iii) $\lim_{\zeta \rightarrow 0} \frac{\psi(-n - \zeta)}{\Gamma(-k - \zeta)} = (-1)^{k+1} k!$.

Acknowledgements. Part of this research was carried out while VM was visiting Osaka Prefecture University. VM thanks the Department of Mathematical Sciences for its support and hospitality. YK is supported in part by JSPS KAKENHI Grant Number 19K03588.

REFERENCES

- [1] S. Agmon, On positivity and decay of solutions of second order elliptic equations on Riemannian manifolds, in *Methods of functional analysis and theory of elliptic equations (Naples, 1982)*, 19–52, Liguori, Naples, 1983.
- [2] C. Bandle, Y. Kabeya, H. Ninomiya, Imperfect bifurcations in nonlinear elliptic equations on spherical caps, *Comm. Pure Appl. Anal.* **10** (2010), 1189–1208.
- [3] R. Beals, R. Wong, *Special Functions, A Graduate Text*, Cambridge Studies in Advanced Mathematics, 126. Cambridge, 2010.
- [4] H. Brezis, T. Cazenave, Y. Martel, A. Ramiandrisoa, Blow up for $u_t - \Delta u = g(u)$ revisited, *Adv. Differential Equations* **1** (1996), 73–90.
- [5] Y. Kabeya, T. Kawakami, A. Kosaka, H. Ninomiya, Eigenvalues of the Laplace-Beltrami operator on a large spherical cap under the Robin problem, *Kodai Math. J.* **37** (2014), 620–645.
- [6] A. Kosaka, Bifurcations of solutions to semilinear elliptic problems on \mathbf{S}^2 with a small hole, *Sci. Math. Jpn.* **78** (2015), 17–42.
- [7] V. Liskevich, S. Lyakhova, V. Moroz, Positive solutions to singular semilinear elliptic equations with critical potential on cone-like domains, *Adv. Differential Equations* **11** (2006), 361–398.

- [8] H. M. MacDonald, Zeros of the spherical harmonics $P_n^m(\mu)$ considered as a function of n , Proc. London Math. Soc. **31** (1899), 264–278.
- [9] S. Moriguchi, K. Udagawa, S. Hitotsumatsu, *Mathematical formula III*, Iwanami Shoten, Tokyo, 1960 (in Japanese).
- [10] D. H. Sattinger, Monotone methods in nonlinear elliptic and parabolic boundary value problems, Indiana Univ. Math. J. **21** (1971/72), 979–1000.

DEPARTMENT OF MATHEMATICAL SCIENCES, OSAKA PREFECTURE UNIVERSITY, GAKUENCHO, SAKAI, 599-8531, JAPAN

E-mail address: `kabeya@ms.osakafu-u.ac.jp`

DEPARTMENT OF MATHEMATICS, SWANSEA UNIVERSITY, FABIAN WAY, SWANSEA SA1 8EN, WALES, UK

E-mail address: `v.moroz@swansea.ac.uk`