

# THE ONE-DIMENSIONAL MARTINGALE HARDY SPACES AND PARTIAL SUMS AND FEJÉR MEANS WITH RESPECT TO WALSH SYSTEM

GEORGE TEPHNADZE

ABSTRACT. In this paper we prove and discuss some new  $(H_p, L_p)$  type inequalities for partial Sums and Fejér means with respect to Walsh system. It is also proved that these results are the best possible in a special sense. As applications, both some well-known and new results are pointed out.

**2010 Mathematics Subject Classification.** 42C10.

**Key words and phrases:** Walsh group, Walsh system,  $L_p$  space, weak- $L_p$  space, modulus of continuity, Walsh-Fourier coefficients, Walsh-Fourier series, partial sums, Lebesgue constants, Fejér means, dyadic martingale, Hardy space, maximal operator, strong convergence.

## 1. PRELIMINARIES

It is well-known that (for details see e.g. [27] and [45]) for every  $p > 1$  there exists an absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|S_n f\|_p \leq c_p \|f\|_p, \quad \text{when } p > 1 \text{ and } f \in H_1(G).$$

Moreover, Watari [75] (see also Gosselin [28] and Young [82]) proved that there exists an absolute constant  $c$  such that, for  $n = 1, 2, \dots$ ,

$$\lambda \mu(|S_n f| > \lambda) \leq c \|f\|_1, \quad f \in L_1(G_m), \quad \lambda > 0.$$

On the other hand, it is also well-known that (for details see e.g. [1] and [45]) Walsh system is not Schauder basis in  $L_1(G)$  space. Moreover, there exists function  $f \in H_1(G)$ , such that partial sums with respect to Walsh system are not uniformly bounded in  $L_1(G)$ .

By applying Lebesgue constants

$$L(n) := \|D_n\|_1$$

we easily obtain that (for details see e.g. [2] and [45]) subsequences of partial sums  $S_{n_k} f$  with respect to Walsh system converge to  $f$  in  $L_1$  norm if and only if

$$(1) \quad \sup_{k \in \mathbb{N}} L(n_k) \leq c < \infty.$$

Since  $n$ -th Lebesgue constant with respect to Walsh system, where

$$n = \sum_{j=0}^{\infty} n_j 2^j, \quad (n_j \in \mathbb{Z}_2)$$

---

The research was supported by Shota Rustaveli National Science Foundation grant no. FR-19-676.

can be estimated by variation of natural number

$$V(n) = n_0 + \sum_{k=1}^{\infty} |n_k - n_{k-1}|$$

and it is also well known that (for details see e.g. [13] and [45]) the following two-sided estimate is true

$$\frac{1}{8}V(n) \leq L(n) \leq V(n)$$

to obtain convergence of subsequences of partial sums  $S_{n_k}f$  with respect to Walsh system of  $f \in L_1$  in  $f \in L_1$ -norm. Condition (1) can be replaced by

$$\sup_{k \in \mathbb{N}} V(n_k) \leq c < \infty$$

It follows that (for details see e.g. [45] and [76]) subsequence of partial sums  $S_{2^n}$  are bounded from  $H_p(G)$  to  $H_p(G)$  for every  $p > 0$ , from which we obtain that

$$(2) \quad \|S_{2^n}f - f\|_{H_p(G)} \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

On the other hand, (see e.g. [59]) there exist a martingale  $f \in H_p(G)$  ( $0 < p < 1$ ), such that

$$\sup_{n \in \mathbb{N}} \|S_{2^{n+1}}f\|_{weak-L_p(G)} = \infty.$$

The main reason of divergence of subsequence  $S_{2^{n+1}}f$  of partial sums is that (for details see [60]) Fourier coefficients of  $f \in H_p(G)$  are not uniformly bounded when  $0 < p < 1$ .

When  $0 < p < 1$  in [72] was investigated boundedness of subsequences of partial sums with respect to Walsh system from  $H_p(G)$  to  $H_p(G)$ . In particular, the following result is true:

**Theorem T1.** Let  $0 < p < 1$  and  $f \in H_p(G)$ . Then there exists a absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|S_{m_k}f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}$$

if and only if the following condition holds

$$(3) \quad \sup_{k \in \mathbb{N}} d(m_k) < c < \infty,$$

where

$$d(m_k) := |m_k| - \langle m_k \rangle.$$

In particular, Theorem T1 immediately follows:

**Theorem T2.** Let  $p > 0$  and  $f \in H_p(G)$ . Then there exists a absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|S_{2^n}f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}$$

and

$$\|S_{2^{n+2^{n-1}}}f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}.$$

On the other hand, we have the following result:

**Theorem T3.** Let  $p > 0$ . Then there exists a martingale  $f \in H_p(G)$ , such that

$$\sup_{n \in \mathbb{N}} \|S_{2^{n+1}}f\|_{H_p(G)} = \infty.$$

Taking into account these results it is interesting to find behaviour of rate of divergence of subsequences of partials sums with respect to Walsh system of martingale  $f \in H_p(G)$  in the martingale Hardy spaces  $H_p(G)$ .

In the second chapter of this thesis (see also [61]) we investigate above mentioned problem. For  $0 < p < 1$  we have the following result:

**Theorem 1.** Let  $f \in H_p(G)$ . Then there exists a absolute constant  $c_p$ , depending only on  $p$ , such that the following inequality is true

$$(4) \quad \|S_n f\|_{H_p(G)} \leq c_p 2^{d(n)(1/p-1)} \|f\|_{H_p(G)}.$$

On the other hand, if  $0 < p < 1$ ,  $\{m_k : k \geq 0\}$  be increasing subsequence of natural numbers, such that

$$(5) \quad \sup_{k \in \mathbb{N}} d(m_k) = \infty$$

and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function satisfying the condition

$$\lim_{k \rightarrow \infty} \frac{2^{d(m_k)(1/p-1)}}{\Phi(m_k)} = \infty,$$

then there exists a martingale  $f \in H_p(G)$ , such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{S_{m_k} f}{\Phi(m_k)} \right\|_{weak-L_p(G)} = \infty.$$

Theorem 1 easily follows the following corollary:

**Corollary 1.** Let  $0 < p < 1$  and  $f \in H_p(G)$ . Then there exists a absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|S_n f\|_{H_p(G)} \leq c_p (n \mu \{\text{supp}(D_n)\})^{1/p-1} \|f\|_{H_p(G)}.$$

On the other hand, if  $0 < p < 1$  and  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers, such that

$$\sup_{k \in \mathbb{N}} m_k \mu \{\text{supp}(D_{m_k})\} = \infty$$

and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function satisfying the condition

$$\lim_{k \rightarrow \infty} \frac{(m_k \mu \{\text{supp}(D_{m_k})\})^{1/p-1}}{\Phi(m_k)} = \infty,$$

then there exists a martingale  $f \in H_p(G)$ , such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{S_{m_k} f}{\Phi(m_k)} \right\|_{weak-L_p(G)} = \infty.$$

In particular, we also get the proofs of Theorem T1 and Theorem T2.

In the second chapter of this thesis we also investigate case  $p = 1$ . In this case the following result is true:

**Theorem 2.** Let  $n \in \mathbb{N}_+$  and  $f \in H_1(G)$ . Then there exists a absolute constant  $c$ , such that

$$(6) \quad \|S_n f\|_{H_1(G)} \leq cV(n) \|f\|_{H_1(G)}.$$

Moreover, if  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers  $\mathbb{N}_+$ , such that

$$\sup_{k \in \mathbb{N}} V(m_k) = \infty$$

and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function satisfying the condition

$$\overline{\lim}_{k \rightarrow \infty} \frac{V(m_k)}{\Phi(m_k)} = \infty.$$

Then there exists a martingale  $f \in H_1(G)$ , such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{S_{m_k} f}{\Phi(m_k)} \right\|_1 = \infty.$$

When  $0 < p < 1$  in [72] was proved boundedness of maximal operators of subsequences of partial sums from  $H_p(G)$  to  $L_p(G)$ . In particular, the following is true:

**Theorem T4.** Let  $0 < p < 1$  and  $f \in H_p(G)$ . Then the maximal operator

$$\sup_{k \in \mathbb{N}} |S_{m_k} f|$$

is bounded from  $H_p(G)$  to  $L_p(G)$ , if and only if condition (3) is fulfilled.

In the special cases we obtain that the following is true:

**Theorem T5.** Let  $p > 0$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$ , such that

$$(7) \quad \left\| \sup_{n \in \mathbb{N}} |S_{2^n} f| \right\|_p \leq c_p \|f\|_{H_p(G)}$$

and

$$\left\| \sup_{n \in \mathbb{N}} |S_{2^n + 2^{n-1}} f| \right\|_p \leq c_p \|f\|_{H_p(G)}.$$

On the other hand we have the following result:

**Theorem T6.** Let  $p > 0$ . Then there exists a martingale  $f \in H_p(G)$ , such that

$$\left\| \sup_{n \in \mathbb{N}} |S_{2^{n+1}} f| \right\|_p = \infty.$$

Above mentioned condition (3) is sufficient condition for the case  $p = 1$  also, but there exist subsequences which do not satisfy this condition, but maximal operators of these subsequences of partial sums with respect to Walsh system are not bounded from  $H_1(G)$  to  $L_1(G)$ .

Such necessary and sufficient conditions which provides boundedness of maximal operators of subsequences of partial sums with respect to Walsh system from  $H_1(G)$  to  $L_1(G)$  is open problem.

In [60] and [72] was investigated boundedness of weighted maximal operators from  $H_p(G)$  to  $L_p(G)$ , when  $0 < p \leq 1$ :

**Theorem T7.** Let  $0 < p \leq 1$ . Then weighted maximal operator

$$\tilde{S}_p^* f := \sup_{n \in \mathbb{N}_+} \frac{|S_n f|}{(n+1)^{1/p-1} \log^{[p]}(n+1)}$$

is bounded from  $H_p(G)$  to  $L_p(G)$ , where  $[p]$  denotes integer part of  $p$ .

Moreover, for any non-decreasing function  $\varphi : \mathbb{N}_+ \rightarrow [1, \infty)$  satisfying the condition

$$\overline{\lim}_{n \rightarrow \infty} \frac{(n+1)^{1/p-1} \log^{[p]}(n+1)}{\varphi(n+1)} = +\infty,$$

there exists a martingale  $f \in H_p(G)$  ( $0 < p \leq 1$ ), such that

$$\sup_{n \in \mathbb{N}} \left\| \frac{S_n f}{\varphi(n)} \right\|_p = \infty.$$

According to negative result for weighted maximal operator of partial sums of Walsh-Fourier series we immediately get the following result:

**Theorem S1.** There exists a martingale  $f \in H_p(G)$ , ( $0 < p \leq 1$ ), such that

$$\sup_{n \in \mathbb{N}} \|S_n f\|_p = \infty.$$

On the other hand, boundedness of weighted maximal operators immediately follows the following estimation:

**Theorem S2.** Let  $0 < p \leq 1$ . Then there exists a absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|S_n f\|_p \leq c_p (n+1)^{1/p-1} \log^{[p]}(n+1) \|f\|_{H_p(G)}, \quad \text{for } 0 < p \leq 1,$$

where  $[p]$  denotes integer part of  $p$ .

By applying this inequality (see [58]) we find necessary and sufficient conditions for martingale  $f \in H_p(G)$  for which partial sums with respect to Walsh system of martingale  $f \in H_p(G)$  converge in  $H_p(G)$  norm.

**Theorem T8.** Let  $0 < p \leq 1$ ,  $[p]$  denotes integer part of  $p$ ,  $f \in H_p(G)$  and

$$\omega_{H_p(G)} \left( \frac{1}{2^N}, f \right) = o \left( \frac{1}{2^{N(1/p-1)} N^{[p]}} \right), \quad \text{as } N \rightarrow \infty.$$

Then

$$\|S_n f - f\|_p \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Moreover, there exists a martingale  $f \in H_p(G)$ , where  $0 < p < 1$ , such that

$$\omega_{H_p(G)} \left( \frac{1}{2^N}, f \right) = O \left( \frac{1}{2^{N(1/p-1)} N^{[p]}} \right), \quad \text{as } N \rightarrow \infty$$

and

$$\|S_n f - f\|_{weak-L_p(G)} \not\rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

By taking these results into account, it is interesting to find necessary and sufficient conditions for modulus of continuity, such that subsequences of partial sums with respect to Walsh system of martingale  $f \in H_p(G)$  converge in  $H_p(G)$  norm.

In the second chapter of this thesis (see also [61]) we investigate this problem. By combining inequalities (4) and (6) we get the following theorem:

**Theorem 3.** Let  $2^k < n \leq 2^{k+1}$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$ , such that

$$(8) \quad \|S_n f - f\|_{H_p(G)} \leq c_p 2^{d(n)(1/p-1)} \omega_{H_p(G)} \left( \frac{1}{2^k}, f \right), \quad (0 < p < 1)$$

and

$$(9) \quad \|S_n f - f\|_{H_1(G)} \leq c_1 V(n) \omega_{H_1(G)} \left( \frac{1}{2^k}, f \right)$$

By applying inequality (8) the following result is proved in the second chapter:

**Theorem 4.** Let  $0 < p < 1$ ,  $f \in H_p(G)$  and  $\{m_k : k \geq 0\}$  be increasing sequence of natural number satisfying the condition

$$\omega_{H_p(G)} \left( \frac{1}{2^{|m_k|}}, f \right) = o \left( \frac{1}{2^{d(m_k)(1/p-1)}} \right) \text{ as } k \rightarrow \infty.$$

Then

$$(10) \quad \|S_{m_k} f - f\|_{H_p(G)} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

On the other hand, if  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers satisfying the condition (5), then there exists a martingale  $f \in H_p(G)$  and subsequence  $\{\alpha_k : k \geq 0\} \subset \{m_k : k \geq 0\}$ , for which

$$\omega_{H_p(G)} \left( \frac{1}{2^{|\alpha_k|}}, f \right) = O \left( \frac{1}{2^{d(\alpha_k)(1/p-1)}} \right) \text{ as } k \rightarrow \infty$$

and

$$(11) \quad \limsup_{k \rightarrow \infty} \|S_{\alpha_k} f - f\|_{weak-L_p(G)} > c_p > 0 \text{ as } k \rightarrow \infty,$$

where  $c_p$  is an absolute constant depending only on  $p$ .

According to this theorem we immediately get that the following result is true:

**Corollary 5.** Let  $0 < p < 1$ ,  $f \in H_p(G)$  and  $\{m_k : k \geq 0\}$  be increasing sequence of natural number, satisfying the condition

$$\omega_{H_p(G)} \left( \frac{1}{2^{|m_k|}}, f \right) = o \left( \frac{1}{(m_k \mu(\text{supp } D_{m_k}))^{1/p-1}} \right), \text{ as } k \rightarrow \infty.$$

Then (10) holds.

On the other hand, if  $\{m_k : k \geq 0\}$  be increasing sequence of natural number, satisfying the condition

$$\lim_{k \rightarrow \infty} \frac{(m_k \mu(\text{supp } (D_{m_k})))^{1/p-1}}{\Phi(m_k)} = \infty,$$

then there exists a martingale  $f \in H_p(G)$  and subsequence  $\{\alpha_k : k \geq 0\} \subset \{m_k : k \geq 0\}$  such that

$$\omega_{H_p(G)} \left( \frac{1}{2^{|\alpha_k|}}, f \right) = O \left( \frac{1}{(\alpha_k \mu(\text{supp } D_{\alpha_k}))^{1/p-1}} \right), \text{ as } k \rightarrow \infty$$

and (11) holds.

By applying (9) we prove that the following is true:

**Theorem 5.** Let  $f \in H_1(G)$  and  $\{m_k : k \geq 0\}$  be increasing sequence of natural number, satisfying the condition

$$\omega_{H_1(G)} \left( \frac{1}{2^{|m_k|}}, f \right) = o \left( \frac{1}{V(m_k)} \right) \text{ as } k \rightarrow \infty.$$

Then

$$\|S_{m_k}f - f\|_{H_1(G)} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Moreover, if  $\{m_k : k \geq 0\}$  be increasing sequence of natural number, satisfying the condition (5), then there exists a martingale  $f \in H_1(G)$  and subsequence  $\{\alpha_k : k \geq 0\} \subset \{m_k : k \geq 0\}$  for which

$$\omega_{H_1(G)}\left(\frac{1}{2^{|\alpha_k|}}, f\right) = O\left(\frac{1}{V(\alpha_k)}\right) \text{ as } k \rightarrow \infty$$

and

$$\limsup_{k \rightarrow \infty} \|S_{\alpha_k}f - f\|_1 > c > 0 \text{ as } k \rightarrow \infty,$$

where  $c$  is an absolute constant.

By applying Theorem 4 and Theorem 5 we immediately get proof of Theorem T8.

Weisz [77] consider convergence in norm of Fejér means of the one-dimensional Walsh-Fourier and proved the following:

**Theorem We1.** Let  $p > 1/2$  and  $f \in H_p(G)$ . Then there exists a absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|\sigma_k f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}.$$

Weisz (for details see e.g. [76]) also consider boundedness of subsequences of Fejér means  $\sigma_{2^n}$  of the one-dimensional Walsh-Fourier series from  $H_p(G)$  to  $H_p(G)$  when  $p > 0$ :

**Theorem We2.** Let  $p > 0$  and  $f \in H_p(G)$ . Then

$$(12) \quad \|\sigma_{2^k} f - f\|_{H_p(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

On the other hand, in [54] was proved the following result:

**Theorem T9.** There exists a martingale  $f \in H_p(G)$  ( $0 < p \leq 1/2$ ) such that

$$\sup_{n \in \mathbb{N}} \|\sigma_{2^{n+1}} f\|_{H_p(G)} = \infty.$$

Goginava [25] (see also [42]) proved that the following result is true:

**Theorem Gog1.** Let  $0 < p \leq 1$ . Then the sequence of operators  $|\sigma_{2^n} f|$  are not bounded from  $H_p(G)$  to  $H_p(G)$ .

When  $0 < p < 1/2$  then in [43] was proved bondedness of subsequences of Fejér means of the one-dimensional Walsh-Fourier from  $H_p(G)$  to  $H_p(G)$ . In particular, the following is true:

**Theorem T10.** Let  $0 < p < 1/2$  and  $f \in H_p(G)$ . Then there exists a absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|\sigma_{m_k} f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}$$

estimation holds if and only if the condition (3) is fulfilled.

Theorem T10 immediately follows theorem of Weisz (see Theorem We2) and also interesting results:

**Theorem T11.** Let  $p > 0$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$ , such that

$$\|\sigma_{2^n} f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}$$

and

$$\|\sigma_{2^n+2^{n-1}}f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}.$$

On the other hand, we have the following result:

**Theorem T12.** Let  $p > 0$ . Then there exists a martingale  $f \in H_p(G)$ , such that

$$\sup_{n \in \mathbb{N}} \|\sigma_{2^n+1}f\|_{H_p(G)} = \infty.$$

According to above mentioned results it is interesting to find rate of divergence of subsequences  $\sigma_{n_k}f$  of Fejér means of the one-dimensional Walsh-Fourier series in the Hardy spaces  $H_p(G)$ .

In the third chapter of this thesis (see also [62]) we find rate of divergence of subsequences of Fejér means of the one-dimensional Walsh-Fourier series on the martingale Hardy spaces  $H_p(G)$ , when  $0 < p \leq 1/2$ .

First, we consider case  $p = 1/2$ :

**Theorem 6.** Let  $n \in \mathbb{N}_+$  and  $f \in H_{1/2}(G)$ . Then there exists an absolute constant  $c$ , such that

$$(13) \quad \|\sigma_n f\|_{H_{1/2}(G)} \leq cV^2(n) \|f\|_{H_{1/2}(G)}.$$

Moreover, if  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers, such that

$$\sup_{k \in \mathbb{N}} V(m_k) = \infty$$

and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty]$  be non-decreasing function satisfying the conditions

$$\overline{\lim}_{k \rightarrow \infty} \frac{V^2(m_k)}{\Phi(m_k)} = \infty.$$

then there exists a martingale  $f \in H_{1/2}(G)$ , such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{\sigma_{m_k} f}{\Phi(m_k)} \right\|_{1/2} = \infty.$$

There was also considered case  $0 < p < 1/2$  and was proved that the following is true:

**Theorem 7.** Let  $0 < p < 1/2$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$  such that

$$(14) \quad \|\sigma_n f\|_{H_p(G)} \leq c_p 2^{d(n)(1/p-2)} \|f\|_{H_p(G)}.$$

On the other hand, if  $0 < p < 1/2$ ,  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers satisfying the condition (5) and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function such that

$$\overline{\lim}_{k \rightarrow \infty} \frac{2^{d(m_k)(1/p-2)}}{\Phi(m_k)} = \infty,$$

then there exists a martingale  $f \in H_p(G)$ , such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{\sigma_{m_k} f}{\Phi(m_k)} \right\|_{weak-L_p(G)} = \infty.$$

From these results also follows proof of Theorem We2.

In 1975 Schipp [44] (see also [2] and [83]) proved that the maximal operator of Fejér means  $\sigma^*$  is of type weak-(1,1):

$$\mu(\sigma^* f > \lambda) \leq \frac{c}{\lambda} \|f\|_1, \quad (\lambda > 0).$$

By using Marcinkiewicz interpolation theorem it follows that  $\sigma^*$  is of strong type-( $p, p$ ), when  $p > 1$ :

$$\|\sigma^* f\|_p \leq c \|f\|_p, \quad (p > 1).$$

The boundedness does not hold for  $p = 1$ , but Fujii [16] (see also [81]) proved that maximal operator of Fejér means is bounded from  $H_1(G)$  to  $L_1(G)$ . Weisz in [78] generalized result of Fujii and proved that maximal operator of Fejér means is bounded from  $H_p(G)$  to  $L_p(G)$ , when  $p > 1/2$ . Simon [46] construct the counterexample, which shows that boundedness does not hold when  $0 < p < 1/2$ . Goginava [21] (see also [9] and [10]) generalized this result for  $0 < p \leq 1/2$  and proved that the following is true:

**Theorem Gog2.** There exists a martingale  $f \in H_p(G)$  ( $0 < p \leq 1/2$ ) such that

$$\sup_{n \in \mathbb{N}} \|\sigma_n f\|_p = \infty.$$

Weisz [79] (see also Goginava [23]) proved that the following is true:

**Theorem We3.** Let  $f \in H_{1/2}(G)$ . Then there exists an absolute constant  $c$ , such that

$$\|\sigma^* f\|_{weak-L_{1/2}(G)} \leq c \|f\|_{H_{1/2}(G)}.$$

In [43] was considered boundedness of maximal operators of subsequences of Fejér means of the one-dimensional Walsh-Fourier series from  $H_p(G)$  to  $L_p(G)$  for  $0 < p < 1/2$ . In particular, the following is true:

**Theorem T13.** Let  $0 < p < 1/2$  and  $f \in H_p(G)$ . Then the maximal operator

$$\tilde{\sigma}^* f := \sup_{k \in \mathbb{N}} |\sigma_{m_k} f|$$

is bounded from  $H_p(G)$  to  $L_p(G)$  if and only if when condition (3) is fulfilled.

As consequences the following results are true:

**Theorem T14.** Let  $p > 0$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$  depending only on  $p$ , such that

$$(15) \quad \left\| \sup_{n \in \mathbb{N}} |\sigma_{2^n} f| \right\|_p \leq c_p \|f\|_{H_p(G)}$$

and

$$\left\| \sup_{n \in \mathbb{N}} |\sigma_{2^{2^n+2^{2^n-1}}} f| \right\|_p \leq c_p \|f\|_{H_p(G)}.$$

On the other hand, we have the following negative result:

**Theorem T15.** Let  $0 < p < 1/2$ . Then there exists a martingale  $f \in H_p(G)$ , such that

$$\left\| \sup_{n \in \mathbb{N}} |\sigma_{2^{2^n+1}} f| \right\|_p = \infty.$$

above mentioned condition is sufficient for the case  $p = 1/2$  also, but there exists subsequences, which do not satisfy condition (3), but maximal operator of subsequences of Fejér means of the one-dimensional Walsh-Fourier series are bounded from  $H_{1/2}(G)$  to  $L_{1/2}(G)$ .

However, it is open problem to find necessary and sufficient conditions on the indexes, which provide boundedness of maximal operator of subsequences of Fejér means of the one-dimensional Walsh-Fourier series from  $H_{1/2}(G)$  to  $L_{1/2}(G)$ .

In [22] and [54] (see also [41], [56], [26] and [53]) is proved that the following is true:

**Theorem GT1.** Let  $0 < p \leq 1/2$  and  $f \in H_p(G)$ . Then the maximal operator

$$\tilde{\sigma}_p^* f := \sup_{n \in \mathbb{N}} \frac{|\sigma_n f|}{(n+1)^{1/p-2} \log^{2[1/2+p]}(n+1)}$$

is bounded from  $H_p(G)$  to  $L_p(G)$ .

Moreover, for any nondecreasing function  $\varphi : \mathbb{N}_+ \rightarrow [1, \infty)$  satisfying the condition

$$\lim_{n \rightarrow \infty} \frac{(n+1)^{1/p-2} \log^{2[1/2+p]}(n+1)}{\varphi(n)} = +\infty,$$

there exists a martingale  $f \in H_p(G)$ , ( $0 < p < 1/2$ ) such that

$$\sup_{n \in \mathbb{N}} \left\| \frac{\sigma_n f}{\varphi(n)} \right\|_p = \infty.$$

From the divergence of weighted maximal operators we immediately get that there exists a martingale  $f \in H_p(G)$  ( $0 < p \leq 1/2$ ), such that

$$\sup_{n \in \mathbb{N}} \|\sigma_n f\|_p = \infty.$$

and from the boundedness results of weighted maximal operators we immediately get that for any  $f \in H_p(G)$  there exists an absolute constant  $c_p$ , such that the following inequality holds true:

$$(16) \quad \|\sigma_n f\|_p \leq c_p n^{1/p-2} \log^{2[1/2+p]}(n+1) \|f\|_{H_p(G)}, \quad \text{as } 0 < p \leq 1/2.$$

By applying inequality (16) in [58] was found necessary and sufficient conditions for modulus of continuity of martingale  $f \in H_p(G)$ , for which Fejér means of the one-dimensional Walsh-Fourier series converge in  $H_p(G)$  norm.

**Theorem T16.** Let  $0 < p \leq 1/2$ ,  $f \in H_p(G)$  and

$$\omega_{H_p(G)} \left( \frac{1}{2^N}, f \right) = o \left( \frac{1}{2^{N(1/p-2)} N^{2[1/2+p]}} \right), \quad \text{as } N \rightarrow \infty.$$

Then

$$\|\sigma_n f - f\|_p \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Moreover, there exists a martingale  $f \in H_p(G)$ , for which

$$\omega_{H_{1/2}(G)} \left( \frac{1}{2^N}, f \right) = O \left( \frac{1}{2^{N(1/p-2)} N^{2[1/2+p]}} \right), \quad \text{as } N \rightarrow \infty$$

and

$$\|\sigma_n f - f\|_p \not\rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

According above mentioned results, it is interesting to find necessary and sufficient conditions for the modulus of continuity, for which subsequences  $\sigma_{n_k} f$  of Fejér means of the one-dimensional Walsh-Fourier series converge in  $H_p(G)$  norm.

In the third chapter of this thesis we find necessary and sufficient conditions for the modulus of continuity, for which subsequences  $\sigma_{n_k}f$  of Fejér means of the one-dimensional Walsh-Fourier series converge in  $H_p(G)$  norm (see also [62]).

By applying inequality (13) for the case  $p = 1/2$  the following necessary and sufficient conditions are found:

**Theorem 8.** Let  $f \in H_{1/2}(G)$  and  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers, such that

$$\omega_{H_{1/2}(G)}\left(\frac{1}{2^{|m_k|}}, f\right) = o\left(\frac{1}{V^2(m_k)}\right) \text{ as } k \rightarrow \infty.$$

Then

$$\|\sigma_{m_k}f - f\|_{H_{1/2}(G)} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Moreover, if  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers, such that (5) holds true, then there exists a martingale  $f \in H_{1/2}(G)$  and subsequence  $\{\alpha_k : k \geq 0\} \subset \{m_k : k \geq 0\}$  such that

$$\omega_{H_{1/2}(G)}\left(\frac{1}{2^{|\alpha_k|}}, f\right) = O\left(\frac{1}{V^2(\alpha_k)}\right) \text{ as } k \rightarrow \infty$$

and

$$\limsup_{k \rightarrow \infty} \|\sigma_{\alpha_k}f - f\|_{1/2} > c > 0 \text{ as } k \rightarrow \infty,$$

where  $c$  is an absolute constant.

By applying inequality (14) we also investigate case  $0 < p < 1/2$ . In the third chapter of this thesis we prove that the following is true:

**Theorem 9.** Let  $0 < p < 1/2$ ,  $f \in H_p(G)$  and  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers, such that

$$\omega_{H_p(G)}\left(\frac{1}{2^{|m_k|}}, f\right) = o\left(\frac{1}{2^{d(m_k)(1/p-2)}}\right), \text{ as } k \rightarrow \infty.$$

Then

$$\|\sigma_{m_k}f - f\|_{H_p(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

On the other hand, if  $\{m_k : k \geq 0\}$  be increasing sequence of natural numbers satisfying the condition (5), then there exists a martingale  $f \in H_p(G)$  and subsequence  $\{\alpha_k : k \geq 0\} \subset \{m_k : k \geq 0\}$ , for which

$$\omega_{H_p(G)}\left(\frac{1}{2^{|\alpha_k|}}, f\right) = O\left(\frac{1}{2^{d(\alpha_k)(1/p-2)}}\right), \text{ as } k \rightarrow \infty$$

and

$$\limsup_{k \rightarrow \infty} \|\sigma_{\alpha_k}f - f\|_{weak-L_p(G)} > c_p > 0, \text{ as } k \rightarrow \infty,$$

where  $c_p$  is constant depending only on  $p$ .

However, Simon in [47] and [49] (see also [48, 14, 50]) consider strong convergence theorems of the one-dimensional Walsh-Fouriere series and proved the following:

**Theorem Si1.** Let  $0 < p \leq 1$  and  $f \in H_1(G)$ . Then there exists an absolute constant  $c_p$  depending only on  $p$ , such that the following inequality is true:

$$\frac{1}{\log^{[p]} n} \sum_{k=1}^n \frac{\|S_k f\|_{H_p(G)}}{k^{2-p}} \leq c_p \|f\|_{H_p(G)},$$

Analogical result for trigonometric system was proved in [51], for unbounded Walsh systems in [18].

In [55] was proved that the following is true:

**Theorem T17.** for any  $0 < p < 1$  and non-decreasing function  $\varphi : \mathbb{N}_+ \rightarrow [1, \infty)$  satisfying the condition

$$\overline{\lim}_{n \rightarrow \infty} \frac{n^{2-p}}{\varphi(n)} = +\infty,$$

there exists a martingale  $f \in H_p(G)$ , such that

$$\sum_{k=1}^{\infty} \frac{\|S_k f\|_{weak-L_p(G)}^p}{\varphi(k)} = \infty, \quad (0 < p < 1).$$

Theorem Si1 follows that if  $f \in H_1(G)$  then the following equalities are true:

$$\lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=1}^n \frac{\|S_k f - f\|_1}{k} = 0$$

and

$$\lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=1}^n \frac{\|S_k f\|_1}{k} = \|f\|_{H_1(G)}.$$

When  $0 < p < 1$  and  $f \in H_p(G)$ , then Theorem Si1 follows that there exists an absolute constant  $c_p$  depending only on  $p$ , such that

$$\frac{1}{n^{1/2-p/2}} \sum_{k=1}^n \frac{\|S_k f\|_{H_p(G)}^p}{k^{3/2-p/2}} \leq c_p \|f\|_{H_p(G)}^p.$$

Moreover,

$$\frac{1}{n^{1/2-p/2}} \sum_{k=1}^n \frac{\|S_k f - f\|_{H_p(G)}^p}{k^{3/2-p}} = 0.$$

It follows the following equality

$$\frac{1}{n^{1/2-p/2}} \sum_{k=1}^n \frac{\|S_k f\|_{H_p(G)}^p}{k^{3/2-p/2}} = \|f\|_{H_p(G)}^p.$$

In the third chapter of this thesis we consider strong convergence results of Fejér means of the one-dimensional Walsh-Fourier series. According to Theorem We1 and Theorem Gog2 we only have to consider case  $0 < p \leq 1/2$  (for details see [57], see also [4], [5], [7], [8], [13]):

**Theorem 10.** Let  $0 < p \leq 1/2$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$  depending only on  $p$ , such that

$$\frac{1}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{\|\sigma_m f\|_{H_p(G)}^p}{m^{2-2p}} \leq c_p \|f\|_{H_p(G)}^p.$$

Moreover, let  $0 < p < 1/2$  and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing, non-negative function, such that  $\Phi(n) \uparrow \infty$  and

$$\overline{\lim}_{k \rightarrow \infty} \frac{k^{2-2p}}{\Phi(k)} = \infty.$$

Then there exists a martingale  $f \in H_p(G)$ , such that

$$\sum_{m=1}^{\infty} \frac{\|\sigma_m f\|_{weak-L_p(G)}^p}{\Phi(m)} = \infty.$$

When  $p = 1/2$  is was also proved that the following is true:

**Theorem 11.** Let  $f \in H_{1/2}(G)$ . Then

$$\sup_{n \in \mathbb{N}_+} \sup_{\|f\|_{H_p(G)} \leq 1} \frac{1}{n} \sum_{m=1}^n \|\sigma_m f\|_{1/2}^{1/2} = \infty.$$

Theorem 10 follows that if  $f \in H_{1/2}(G)$  then the following equalities are true:

$$\lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=1}^n \frac{\|\sigma_k f - f\|_{H_{1/2}(G)}^{1/2}}{k} = 0$$

and

$$\lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=1}^n \frac{\|\sigma_k f\|_{H_{1/2}(G)}^{1/2}}{k} = \|f\|_{H_{1/2}(G)}^{1/2}.$$

When  $0 < p < 1/2$  and  $f \in H_p(G)$ , then Theorem 10 follows that there exists an absolute constant  $c_p$  depending only on  $p$ , such that

$$\frac{1}{n^{1/2-p}} \sum_{k=1}^n \frac{\|\sigma_k f\|_{H_p(G)}^p}{k^{3/2-p}} \leq c_p \|f\|_{H_p(G)}^p.$$

Moreover,

$$\frac{1}{n^{1/2-p}} \sum_{k=1}^n \frac{\|\sigma_k f - f\|_{H_p(G)}^p}{k^{3/2-p}} = 0.$$

It follows that

$$\frac{1}{n^{1/2-p}} \sum_{k=1}^n \frac{\|\sigma_k f\|_{H_p(G)}^p}{k^{3/2-p}} = \|f\|_{H_p(G)}^p.$$

2. PARTIAL SUMS WITH RESPECT TO THE ONE-DIMENSIONAL WALSH-FOURIER SERIES  
ON THE MARTINGALE HARDY SPACES

2.1. **Basic notations.**

Denote by  $\mathbb{N}_+$  the set of the positive integers and by  $\mathbb{N} := \mathbb{N}_+ \cup \{0\}$  the set of non-negative integers. Denote by  $Z_2$  the additive group of integers modulo-2, which contains only two elements  $Z_2 := \{0, 1\}$ , group operation is modulo-2 sum and all sets are open.

Define the group  $G$  as the complete direct product of the groups  $Z_2$  with the product of the discrete topologies  $Z_2$ . The direct product  $\mu$  of the measures  $\mu_n(\{j\}) := 1/2$ , ( $j \in Z_2$ ) is the Haar measure on  $G$  with  $\mu(G) = 1$ .

The elements of  $G$  are represented by sequences

$$x := (x_0, x_1, \dots, x_j, \dots) \quad (x_k = 0, 1).$$

It is easy to give a base for the neighbourhood of  $G$

$$I_0(x) := G,$$

$$I_n(x) := \{y \in G \mid y_0 = x_0, \dots, y_{n-1} = x_{n-1}\} \quad (x \in G, n \in \mathbb{N}).$$

Set  $I_n := I_n(0)$  for any  $n \in \mathbb{N}$  and  $\overline{I}_n := G \setminus I_n$ .

It is evident that

$$(17) \quad \overline{I}_M = \left( \bigcup_{k=0}^{M-2} \bigcup_{l=k+1}^{M-1} I_{l+1}(e_k + e_l) \right) \cup \left( \bigcup_{k=0}^{M-1} I_M(e_k) \right) = \bigcup_{k=0}^{M-1} I_k \setminus I_{k+1}.$$

If  $n \in \mathbb{N}$ , then it can be uniquely expressed as  $n = \sum_{k=0}^{\infty} n_j 2^j$  where  $n_j \in Z_2$  ( $j \in \mathbb{N}$ ) and only a finite number of  $n_j$ s differ from zero. Set

$$\langle n \rangle := \min\{j \in \mathbb{N}, n_j \neq 0\} \quad \text{and} \quad |n| := \max\{j \in \mathbb{N}, n_j \neq 0\},$$

It is evident that  $2^{|n|} \leq n \leq 2^{|n|+1}$ . Let

$$d(n) := |n| - \langle n \rangle, \quad \text{for any } n \in \mathbb{N}.$$

Denote by  $V(n)$  variation of natural number  $n \in \mathbb{N}$

$$V(n) = n_0 + \sum_{k=1}^{\infty} |n_k - n_{k-1}|.$$

Define  $k$ -th Rademacher functions by

$$r_k(x) := (-1)^{x_k} \quad (x \in G, k \in \mathbb{N}).$$

By using Rademacher functions we define Walsh system  $w := (w_n : n \in \mathbb{N})$   $G$  as:

$$w_n(x) := \prod_{k=0}^{\infty} r_k^{n_k}(x) = r_{|n|}(x) (-1)^{\sum_{k=0}^{|n|-1} n_k x_k} \quad (n \in \mathbb{N}).$$

The norm (quasi-norm) of space  $L_p(G)$  and *weak*- $L_p(G)$  for  $(0 < p < \infty)$  are respectively defined as

$$\|f\|_p^p := \left( \int_G |f(x)|^p d\mu(x) \right), \quad \|f\|_{\text{weak-}L_p(G)}^p := \sup_{\lambda > 0} \lambda^p \mu(x \in G : |f| > \lambda).$$

Walsh system is orthonormal and complete in  $L_2(G)$  (see [45]).

For any  $f \in L_1(G)$  the numbers

$$\widehat{f}(n) := \int_G f(x) w_n(x) d\mu(x)$$

are called  $n$ -th Walsh-Fourier coefficient of  $f$ .

$n$ -th partial sum is denoted by

$$S_n(f; x) := \sum_{i=0}^{n-1} \widehat{f}(i) w_i(x).$$

Dirichlet kernels are defined by

$$D_n(x) := \sum_{i=0}^{n-1} w_i(x).$$

We also define the following maximal operators

$$S^* f = \sup_{n \in \mathbb{N}} |S_n f|, \quad \widetilde{S}_{\#}^* f = \sup_{n \in \mathbb{N}} |S_{2^n} f|.$$

The  $\sigma$ -algebra generated by the intervals  $I_n(x)$  with measure  $2^{-n}$  is denoted by  $F_n$  ( $n \in \mathbb{N}$ ). Conditional exponential operator with respect to  $F_n$  ( $n \in \mathbb{N}$ ) is denoted by  $E_n$  and it is given by

$$E_n f(x) = S_{2^n} f(x) = \sum_{k=0}^{2^n-1} \widehat{f}(k) w_k(x) = \frac{1}{|I_n(x)|} \int_{I_n(x)} f(x) d\mu(x),$$

where  $|I_n(x)| = 2^{-n}$  denotes length of set  $I_n(x)$ .

Sequence  $f = (f_n, n \in \mathbb{N})$  of functions  $f_n \in L_1(G)$  is called dyadic martingale (for details see [36], [45]) if

- (i)  $f_n$  is measurable with respect to  $\sigma$ -algebras  $F_n$  for any  $n \in \mathbb{N}$ ,
- (ii)  $E_n f_m = f_n$  for any  $n \leq m$ .

The maximal function of a martingale  $f$  is defined by

$$f^* = \sup_{n \in \mathbb{N}} |f_n|.$$

In the case  $f \in L_1(G)$ , the maximal functions are also be given by:

$$f^*(x) = \sup_{n \in \mathbb{N}} \frac{1}{\mu(I_n(x))} \left| \int_{I_n(x)} f(u) d\mu(u) \right|.$$

For  $0 < p < \infty$  the Hardy martingale space  $H_p(G)$  consists of all martingales, for which

$$\|f\|_{H_p(G)} := \|f^*\|_p < \infty.$$

A bounded measurable function  $a$  is said to be a  $p$ -atom if there exists an dyadic interval  $I$ , such that

$$\int_I a d\mu = 0, \quad \|a\|_\infty \leq \mu(I)^{-1/p}, \quad \text{supp}(a) \subset I.$$

It is easy to show that for martingale  $f = (f_n, n \in \mathbb{N})$  and for any  $k \in \mathbb{N}$  there exists a limit

$$\widehat{f}(k) := \lim_{n \rightarrow \infty} \int_G f_n(x) w_k(x) d\mu(x)$$

and it is called  $k$ -th Walsh-Fourier coefficients of  $f$ .

If  $f_0 \in L_1(G)$  and  $f := (E_n f_0 : n \in \mathbb{N})$  is regular martingale then

$$\widehat{f}(k) = \int_G f(x) w_k(x) d\mu(x) = \widehat{f_0}(k), \quad k \in \mathbb{N}.$$

The modulus of continuity in  $H_p(G)$  space is defined by

$$\omega_{H_p(G)}\left(\frac{1}{2^n}, f\right) := \|f - S_{2^n} f\|_{H_p(G)}.$$

It is important to describe how can be understood difference  $f - S_{2^n} f$ , where  $f$  be martingale  $S_{2^n} f$  is a function:

**Remark 1.** Let  $0 < p \leq 1$ . Since

$$S_{2^n} f = f^{(n)} \in L_1(G), \quad \text{where } f = (f^{(n)} : n \in \mathbb{N}) \in H_p(G)$$

and

$$\begin{aligned} (S_{2^k} f^{(n)} : k \in \mathbb{N}) &= (S_{2^k} S_{2^n}, k \in \mathbb{N}) \\ &= (S_{2^0} f, \dots, S_{2^{n-1}} f, S_{2^n} f, S_{2^n} f, \dots) = (f^{(0)}, \dots, f^{(n-1)}, f^{(n)}, f^{(n)}, \dots). \end{aligned}$$

Under the difference  $f - S_{2^n} f$  we mean the following martingale:

$$f := ((f - S_{2^n} f)^{(k)}, \quad k \in \mathbb{N})$$

where

$$(f - S_{2^n} f)^{(k)} = \begin{cases} 0, & k = 0, \dots, n, \\ f^{(k)} - f^{(n)}, & k \geq n + 1, \end{cases}$$

Consequently, the norm  $\|f - S_{2^n} f\|_{H_p(G)}$  is understood as  $H_p$ -norm of

$$f - S_{2^n} f = ((f - S_{2^n} f)^{(k)}, k \in \mathbb{N})$$

.

Watari [74] showed that there are strong connections between

$$\omega_p\left(\frac{1}{2^n}, f\right), \quad E_{2^n}(L_p, f) \quad \text{and} \quad \|f - S_{2^n} f\|_p, \quad p \geq 1, \quad n \in \mathbb{N}.$$

In particular,

$$\frac{1}{2} \omega_p\left(\frac{1}{2^n}, f\right) \leq \|f - S_{2^n} f\|_p \leq \omega_p\left(\frac{1}{2^n}, f\right)$$

and

$$\frac{1}{2} \|f - S_{2^n} f\|_p \leq E_{2^n}(L_p, f) \leq \|f - S_{2^n} f\|_p.$$

## 2.2. Auxiliary lemmas.

First we present and prove equalities and estimations of Dirichlet kernel and Lebesgue constants with respect to the one-dimensional Walsh-Fourier systems (see Lemmas 1-12).

First equality of the following Lemma is proved in [45] and second identity is proved in [19]:

**Lemma 1.** *Let  $j, n \in \mathbb{N}$ . Then*

$$D_{j+2^n} = D_{2^n} + w_{2^n} D_j, \quad \text{when } j \leq 2^n,$$

and

$$D_{2^n-j} = D_{2^n} - \psi_{2^n-1} D_j, \quad \text{when } j < 2^n.$$

The following estimation of Dirichlet kernel with respect to the one-dimensional Walsh-Fourier systems is proved in [45]:

**Lemma 2.** *Let  $n \in \mathbb{N}$ . Then*

$$D_{2^n}(x) = \begin{cases} 2^n, & \text{if } x \in I_n, \\ 0, & \text{if } x \notin I_n, \end{cases}$$

and

$$D_n = w_n \sum_{k=0}^{\infty} n_k r_k D_{2^k} = w_n \sum_{k=0}^{\infty} n_k (D_{2^{k+1}} - D_{2^k}), \quad \text{for } n = \sum_{i=0}^{\infty} n_i 2^i.$$

The following two-sided estimations of Lebesgue constants with respect to the one-dimensional Walsh-Fourier systems is proved in [45] and second equality is proved in [15]:

**Lemma 3.** *Let  $n \in \mathbb{N}$ . Then*

$$\frac{1}{8} V(n) \leq \|D_n\|_1 \leq V(n)$$

and

$$\frac{1}{n \log n} \sum_{k=1}^n V(k) = \frac{1}{4 \log 2} + o(1).$$

Hardy martingale space  $H_p(G)$  for any  $0 < p \leq 1$  can be characterize by simple functions which are called  $p$ -atoms. The following is true (for details see [48], [76] and [80]):

**Lemma 4.** *A martingale  $f = (f_n, n \in \mathbb{N})$  belongs to  $H_p(G)$  ( $0 < p \leq 1$ ) if and only if there exists a sequence of  $p$ -atoms of  $(a_k, k \in \mathbb{N})$  and sequence of real numbers  $(\mu_k, k \in \mathbb{N})$  such that for all  $n \in \mathbb{N}$ ,*

$$(18) \quad \sum_{k=0}^{\infty} \mu_k S_{2^n} a_k = f_n$$

and

$$\sum_{k=0}^{\infty} |\mu_k|^p < \infty.$$

Moreover,

$$\|f\|_{H_p(G)} \sim \inf \left( \sum_{k=0}^{\infty} |\mu_k|^p \right)^{1/p},$$

where the infimum is taken over all decomposition of  $f$  of the form (18).

The next five Examples of martingales will be used many times to prove sharpness of our main results. Such counterexamples first appear in the papers of Goginava [24] (see also [20, 23]). Such constructions of martingales are also used in the papers [3], [4], [11], [12], [30], [29], [31], [32], [33], [34], [35], [37], [38], [39], [40], [52], [57], [61], [62], [63], [64], [65], [66], [67], [68], [69], [70], [71], [73]. For the one-dimensional case we use martingales which were used in [72]. So, we leave out the details of proof.

**Example 1.** Let  $0 < p \leq 1$ ,  $\{\lambda_k : k \in \mathbb{N}\}$  be sequence of real numbers

$$(19) \quad \sum_{k=0}^{\infty} |\lambda_k|^p \leq c_p < \infty$$

and  $\{a_k : k \in \mathbb{N}\}$  be sequence of  $p$ -atoms, given by

$$a_k(x) := 2^{|\alpha_k|(1/p-1)} (D_{2^{|\alpha_k|+1}}(x) - D_{2^{|\alpha_k|}}(x)),$$

where  $|\alpha_k| := \max \{j \in \mathbb{N} : (\alpha_k)_j \neq 0\}$  and  $(\alpha_k)_j$  denotes  $j$ -th binary coefficients of real number of  $\alpha_k \in \mathbb{N}_+$ . Then  $f = (f_n : n \in \mathbb{N})$ , where

$$f_n(x) := \sum_{\{k: |\alpha_k| < n\}} \lambda_k a_k(x)$$

is martingale, which belongs to  $H_p(G)$  for any  $0 < p \leq 1$ .

It is easy to show that

$$(20) \quad \begin{aligned} & \widehat{f}(j) \\ &= \begin{cases} \lambda_k 2^{(1/p-1)|\alpha_k|}, & j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k \in \mathbb{N}_+, \\ 0, & j \notin \bigcup_{k=1}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases} \end{aligned}$$

Let  $2^{|\alpha_{l-1}|+1} \leq j \leq 2^{|\alpha_l|}$ ,  $l \in \mathbb{N}_+$ . Then

$$(21) \quad S_j f = S_{2^{|\alpha_{l-1}|+1}} = \sum_{\eta=0}^{l-1} \lambda_{\eta} 2^{|\alpha_{\eta}|(1/p-1)} (D_{2^{|\alpha_{\eta}|+1}} - D_{2^{|\alpha_{\eta}|}}).$$

Let  $2^{|\alpha_l|} \leq j < 2^{|\alpha_l|+1}$ ,  $l \in \mathbb{N}_+$ . Then

$$(22) \quad \begin{aligned} S_j f &= S_{2^{|\alpha_l|}} + \lambda_l 2^{(1/p-1)|\alpha_l|} w_{2^{|\alpha_l|}} D_{j-2^{|\alpha_l|}} \\ &= \sum_{\eta=0}^{l-1} \lambda_{\eta} 2^{|\alpha_{\eta}|(1/p-1)} (D_{2^{|\alpha_{\eta}|+1}} - D_{2^{|\alpha_{\eta}|}}) \\ &\quad + \lambda_l 2^{(1/p-1)|\alpha_l|} w_{2^{|\alpha_l|}} D_{j-2^{|\alpha_l|}}. \end{aligned}$$

Moreover, for the modulus of continuity for  $0 < p \leq 1$  we have the following estimation:

$$(23) \quad \omega_{H_p} \left( \frac{1}{2^n}, f \right) = O \left( \sum_{\{k: |\alpha_k| \geq n\}} |\lambda_k|^p \right)^{1/p}, \quad \text{as } n \rightarrow \infty.$$

By applying Lemma 4 we easily obtain that the following is true (see [80]):

**Lemma 5.** *Let  $0 < p \leq 1$  and  $T$  be  $\sigma$ -sub-linear operator, such that, for any  $p$ -atom  $a$ ,*

$$\int_G |Ta(x)|^p d\mu(x) \leq c_p < \infty.$$

Then

$$(24) \quad \|Tf\|_p \leq c_p \|f\|_{H_p(G)}.$$

In addition, if  $T$  is bounded from  $L_\infty(G)$  to  $L_\infty(G)$  then to prove (24) it suffices to show that

$$\int_{\bar{I}} |Ta(x)|^p d\mu(x) \leq c_p < \infty,$$

for every  $p$ -atom  $a$ , where  $I$  denotes support of the atom  $a$ .

In the concrete cases the norm of Hardy martingale spaces can be calculated by simpler formulas (for details see [48], [76] and [77]):

**Lemma 6.** *If  $g \in L_1(G)$  and  $f := (E_n g : n \in \mathbb{N})$  be regular martingale, then  $H_p(G)$  for  $0 < p \leq 1$  norm can be calculated by*

$$\|f\|_{H_p(G)} = \left\| \sup_{n \in \mathbb{N}} |S_{2^n} g| \right\|_p.$$

The following lemmas are proved in [57], [61], [62].

**Lemma 7.** *Let  $0 < p \leq 1$ ,  $2^k \leq n < 2^{k+1}$  and  $S_n f$  be  $n$ -th partial sum with respect to the one-dimensional Walsh-Fourier series, where  $f \in H_p(G)$ . Then for any fixed  $n \in \mathbb{N}$ ,*

$$\|S_n f\|_{H_p(G)}^p \leq \left\| \sup_{0 \leq l \leq k} |S_{2^l} f| \right\|_p^p + \|S_n f\|_p^p \leq \left\| \tilde{S}_{\#}^* f \right\|_p^p + \|S_n f\|_p^p.$$

**Proof:** Let consider the following martingales

$$f_{\#} := (S_{2^k} S_n f, k \in \mathbb{N}_+) = (S_{2^0}, S_{2^k} f, S_n f, \dots, S_n f, \dots),$$

Hence, Lemma 6 immediately follows that

$$\|S_n f\|_{H_p(G)}^p \leq \left\| \sup_{0 \leq l \leq k} |S_{2^l} f| \right\|_p^p + \|S_n f\|_p^p \leq \left\| \tilde{S}_{\#}^* f \right\|_p^p + \|S_n f\|_p^p.$$

Lemma is proved.

### 2.3. Boundedness of subsequences of partial sums with respect to the one-dimensional Walsh-Fourier series on the martingale Hardy spaces.

In this section we consider boundedness of subsequences of partial sums with respect to the one-dimensional Walsh-Fourier series in the martingale Hardy spaces (for details see [61]).

**Theorem 1.** *a) Let  $0 < p < 1$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$  depending only on  $p$ , such that*

$$\|S_n f\|_{H_p(G)} \leq c_p 2^{d(n)(1/p-1)} \|f\|_{H_p(G)}.$$

*b) Let  $0 < p < 1$ ,  $\{m_k : k \in \mathbb{N}_+\}$  be non-negative, increasing sequence of natural numbers such that*

$$(25) \quad \sup_{k \in \mathbb{N}} d(m_k) = \infty$$

and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function satisfying the condition

$$(26) \quad \lim_{k \rightarrow \infty} \frac{2^{d(m_k)(1/p-1)}}{\Phi(m_k)} = \infty.$$

Then there exists a martingale  $f \in H_p(G)$  such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{S_{m_k} f}{\Phi(m_k)} \right\|_{\text{weak-}L_p(G)} = \infty.$$

**Proof:** Suppose that

$$(27) \quad \left\| 2^{(1-1/p)d(n)} S_n f \right\|_p \leq c_p \|f\|_{H_p(G)}.$$

By combining Lemma 7 and inequalities (7) and (27), since  $2^{(1-1/p)d(n)} \leq c_p$  we obtain that

$$(28) \quad \begin{aligned} \left\| 2^{(1-1/p)d(n)} S_n f \right\|_{H_p(G)}^p &\leq \left\| 2^{(1-1/p)d(n)} S_n f \right\|_p^p + 2^{(1-1/p)d(n)} \left\| \tilde{S}_{\#}^* f \right\|_p^p \\ &\leq c_p \|f\|_{H_p(G)}^p + c_p \left\| \tilde{S}_{\#}^* f \right\|_p^p \leq c_p \|f\|_{H_p(G)}^p. \end{aligned}$$

By combining Lemma 5 and (28) it is sufficient to show that

$$(29) \quad \int_G |2^{(1-1/p)d(n)} S_n a|^p d\mu \leq c_p < \infty,$$

for every  $p$ -atom  $a$ , with support  $I$ , such that  $\mu(I) = 2^{-M}$ .

Without loss of generality we may assume that  $p$ -atom  $a$  has support  $I = I_M$ . Then it is easy to see that  $S_n a = 0$ , where  $2^M \geq n$ . So, we may assume that  $2^M < n$ . Since  $\|a\|_\infty \leq 2^{M/p}$  we can conclude that

$$(30) \quad \begin{aligned} |2^{(1-1/p)d(n)} S_n a(x)| &\leq 2^{(1-1/p)d(n)} \|a\|_\infty \int_{I_M} |D_n(x+t)| d\mu(t) \\ &\leq 2^{M/p} 2^{(1-1/p)d(n)} \int_{I_M} |D_n(x+t)| d\mu(t). \end{aligned}$$

Let  $x \in I_M$ . Since  $V(n) \leq 2d(n)$ , by using first estimations of Lemma 3 we can conclude that

$$|2^{(1-1/p)d(n)} S_n a| \leq 2^{M/p} 2^{(1-1/p)d(n)} V(n) \leq 2^{M/p} d(n) 2^{(1-1/p)d(n)}$$

and

$$(31) \quad \int_{I_M} |2^{(1-1/p)d(n)} S_n a|^p d\mu \leq d(n) 2^{(1-1/p)d(n)} < c_p < \infty.$$

Let  $t \in I_M$  and  $x \in I_s \setminus I_{s+1}$ , where  $0 \leq s \leq M-1 < \langle n \rangle$  or  $0 \leq s < \langle n \rangle \leq M-1$ . Then  $x+t \in I_s \setminus I_{s+1}$  and if we use both equality of Lemma 2 we get that  $D_n(x+t) = 0$  and it follows that

$$(32) \quad |2^{(1-1/p)d(n)} S_n a(x)| = 0.$$

Let  $x \in I_s \setminus I_{s+1}$ ,  $\langle n \rangle \leq s \leq M-1$ . Then  $x+t \in I_s \setminus I_{s+1}$ , where  $t \in I_M$ . Then by using again both equality of Lemma 2 we have that

$$|D_n(x+t)| \leq \sum_{j=0}^s n_j 2^j \leq c 2^s.$$

If we apply again (30) we can conclude that

$$(33) \quad \begin{aligned} |2^{(1-1/p)d(n)} S_n a(x)| &\leq 2^{(1-1/p)d(n)} 2^{M/p} \frac{2^s}{2^M} \\ &\leq 2^{\langle n \rangle(1/p-1)} 2^{M(1/p-1)} \frac{2^s}{2^{|\langle n \rangle(1/p-1)|}} \leq 2^{\langle n \rangle(1/p-1)} 2^s. \end{aligned}$$

By identity (17) and inequalities (32) and (33) we find that

$$\begin{aligned} &\int_{I_M} |2^{(1-1/p)d(n)} S_n a(x)|^p d\mu(x) \\ &= \sum_{s=\langle n \rangle}^{M-1} \int_{I_s \setminus I_{s+1}} |2^{\langle n \rangle(1/p-1)} 2^s|^p d\mu(x) \leq c \sum_{s=\langle n \rangle}^{M-1} \frac{2^{\langle n \rangle(1-p)}}{2^{s(1-p)}} \leq c_p < \infty. \end{aligned}$$

Now, we prove part b) of Theorem 1. By using condition (26) there exists sequence of natural numbers  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{m_k : k \in \mathbb{N}_+\}$ , such that

$$(34) \quad \sum_{\eta=0}^{\infty} \frac{\Phi^{p/2}(\alpha_\eta)}{2^{d(\alpha_\eta)(1-p)/2}} < \infty,$$

Let  $f = (f_n, n \in \mathbb{N}_+) \in H_p(G)$  be a martingale from the Example 1, where

$$(35) \quad \lambda_k = \frac{\Phi^{1/2}(\alpha_k)}{2^{d(\alpha_k)(1-p)/2}}.$$

Then, if we use (34) we obtain that condition (19) is fulfilled and it follows that  $f = (f_n, n \in \mathbb{N}_+) \in H_p(G)$ .

If we apply (20) when  $\lambda_k$  are given by the formula (35) then we get that

$$(36) \quad \widehat{f}(j) = \begin{cases} \Phi^{1/2}(\alpha_k) 2^{(|\alpha_k| + \langle \alpha_k \rangle)(1/p-1)/2}, & \text{if } j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k \in \mathbb{N}_+ \\ 0, & \text{if } j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

In the view of (22) when  $\lambda_k$  are given by (35) we get that

$$(37) \quad \frac{S_{\alpha_k} f}{\Phi(\alpha_k)} = \frac{1}{\Phi(\alpha_k)} \sum_{\eta=0}^{k-1} \Phi^{1/2}(\alpha_\eta) 2^{(|\alpha_\eta| + \langle \alpha_\eta \rangle)(1/p-1)/2} (D_{2^{|\alpha_\eta|+1}} - D_{2^{|\alpha_\eta|}}) \\ + \frac{2^{(|\alpha_k| + \langle \alpha_k \rangle)(1/p-1)/2} w_{2^{|\alpha_k|}} D_{\alpha_k - 2^{|\alpha_k|}}}{\Phi^{1/2}(\alpha_k)} := I + II.$$

by using (34) for  $I$  we have that

$$(38) \quad \|I\|_{weak-L_p(G)}^p \\ \leq \frac{1}{\Phi^p(\alpha_k)} \sum_{\eta=0}^{k-1} \frac{\Phi^{p/2}(\alpha_\eta)}{2^{d(\alpha_\eta)(1-p)/2}} \|2^{|\alpha_\eta|(1/p-1)} (D_{2^{|\alpha_\eta|+1}} - D_{2^{|\alpha_\eta|}})\|_{weak-L_p(G)}^p \\ \leq \frac{1}{\Phi^p(\alpha_k)} \sum_{\eta=0}^{\infty} \frac{\Phi^{p/2}(\alpha_\eta)}{2^{d(\alpha_\eta)(1-p)/2}} \leq c < \infty.$$

Let  $x \in I_{\langle \alpha_k \rangle} \setminus I_{\langle \alpha_k \rangle + 1}$ . Since  $|\alpha_k| \neq \langle \alpha_k \rangle$  and  $\langle \alpha_k - 2^{|\alpha_k|} \rangle = \langle \alpha_k \rangle$ .

By using both inequalities of Lemma 2 we get that

$$(39) \quad \left| D_{\alpha_k - 2^{|\alpha_k|}}(x) \right| \\ = \left| (D_{2^{\langle \alpha_k \rangle + 1}}(x) - D_{2^{\langle \alpha_k \rangle}}(x)) + \sum_{j=\langle \alpha_k \rangle + 1}^{|\alpha_k| - 1} (\alpha_k)_j (D_{2^{j+1}}(x) - D_{2^j}(x)) \right| \\ = \left| -D_{2^{\langle \alpha_k \rangle}}(x) \right| = 2^{\langle \alpha_k \rangle}$$

and

$$(40) \quad |II| = \frac{2^{(|\alpha_k| + \langle \alpha_k \rangle)(1/p-1)/2}}{\Phi^{1/2}(\alpha_k)} \left| D_{\alpha_k - 2^{|\alpha_k|}}(x) \right| = \frac{2^{|\alpha_k|(1/p-1)/2} 2^{\langle \alpha_k \rangle(1/p+1)/2}}{\Phi^{1/2}(\alpha_k)}.$$

By combining (38) and (40) we get that

$$\left\| \frac{S_{\alpha_k} f}{\Phi(\alpha_k)} \right\|_{weak-L_p(G)}^p \\ \geq \|II\|_{weak-L_p(G)}^p - \|I\|_{weak-L_p(G)}^p \\ \geq \frac{2^{(|\alpha_k|)(1/p-1)/2} 2^{\langle \alpha_k \rangle(1/p+1)/2}}{\Phi^{1/2}(\alpha_k)} \mu \left\{ x \in G : |II| \geq \frac{2^{(|\alpha_k|)(1/p-1)/2} 2^{\langle \alpha_k \rangle(1/p+1)/2}}{\Phi^{1/2}(\alpha_k)} \right\}^{1/p} \\ \geq \frac{2^{(|\alpha_k|)(1/p-1)/2} 2^{\langle \alpha_k \rangle(1/p+1)/2}}{\Phi^{1/2}(\alpha_k)} (\mu \{I_{\langle \alpha_k \rangle} \setminus I_{\langle \alpha_k \rangle + 1}\})^{1/p} \\ \geq c \frac{2^{d(\alpha_k)(1/p-1)/2}}{\Phi^{1/2}(\alpha_k)} \rightarrow \infty, \quad \text{as } k \rightarrow \infty.$$

The proof of Theorem 1 is complete.

**Corollary 1.** *a) Let  $n \in \mathbb{N}_+$ ,  $0 < p < 1$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$  such that*

$$\|S_n f\|_{H_p(G)} \leq c_p (n \mu \{supp(D_n)\})^{1/p-1} \|f\|_{H_p(G)}.$$

b) Let  $0 < p < 1$ ,  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that

$$(41) \quad \sup_{k \in \mathbb{N}} m_k \mu \{ \text{supp}(D_{m_k}) \} = \infty$$

and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function satisfying the condition

$$(42) \quad \overline{\lim}_{k \rightarrow \infty} \frac{(m_k \mu \{ \text{supp}(D_{m_k}) \})^{1/p-1}}{\Phi(m_k)} = \infty.$$

Then there exists a martingale  $f \in H_p(G)$  such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{S_{m_k} f}{\Phi(m_k)} \right\|_{\text{weak-}L_p(G)} = \infty.$$

**Proof:** By applying both inequalities of Lemma 2 we get that

$$I_{\langle n \rangle} \setminus I_{\langle n \rangle + 1} \subset \text{supp} \{D_n\} \subset I_{\langle n \rangle} \quad \text{and} \quad 2^{-\langle n \rangle - 1} \leq \mu \{ \text{supp}(D_n) \} \leq 2^{-\langle n \rangle}.$$

Hence,

$$\frac{2^{d(n)(1/p-1)}}{4} \leq (n \mu \{ \text{supp}(D_n) \})^{1/p-1} \leq 2^{d(n)(1/p-1)}.$$

Corollary 1 is proved.

**Theorem 2.** a) Let  $n \in \mathbb{N}_+$  and  $f \in H_1(G)$ . Then there exists an absolute constant  $c$ , such that

$$\|S_n f\|_{H_1(G)} \leq cV(n) \|f\|_{H_1(G)}.$$

b) Let  $\{m_k : k \in \mathbb{N}_+\}$  be non-negative increasing sequence of natural numbers such that

$$(43) \quad \sup_{k \in \mathbb{N}} V(m_k) = \infty$$

and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function satisfying the condition

$$(44) \quad \overline{\lim}_{k \rightarrow \infty} \frac{V(m_k)}{\Phi(m_k)} = \infty.$$

Then there exists a martingale  $f \in H_1(G)$ , such that

$$\sup_{k \in \mathbb{N}} \left\| \frac{S_{m_k} f}{\Phi(m_k)} \right\|_1 = \infty.$$

**Proof:** Since

$$(45) \quad \left\| \frac{S_n f}{V(n)} \right\|_1 \leq \|f\|_1 \leq \|f\|_{H_1(G)}$$

by combining Lemmas 7 and (45) we can conclude that

$$(46) \quad \begin{aligned} \left\| \frac{S_n f}{V(n)} \right\|_{H_1(G)} &\leq \left\| \frac{S_n f}{V(n)} \right\|_1 + \frac{1}{V(n)} \left\| \tilde{S}_{\#}^* f \right\|_1 \\ &\leq c \|f\|_{H_1(G)} + c \left\| \tilde{S}_{\#}^* f \right\|_1 \leq c \|f\|_{H_1(G)}. \end{aligned}$$

Now prove second part of Theorem 2. Let  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers and function  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  satisfies conditions (43) and (44). Then there exists non-negative, increasing sequence  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{m_k : k \in \mathbb{N}_+\}$  such that

$$(47) \quad \sum_{k=1}^{\infty} \frac{\Phi^{1/2}(\alpha_k)}{V^{1/2}(\alpha_k)} \leq \beta < \infty.$$

Let  $f = (f_n, n \in \mathbb{N}_+)$  be martingale from Example 1, where

$$(48) \quad \lambda_k = \frac{\Phi^{1/2}(\alpha_k)}{V^{1/2}(\alpha_k)}.$$

By applying condition (47) we can conclude that condition (19) is fulfilled and it follows that  $f = (f_n, n \in \mathbb{N}_+) \in H_1(G)$ .

In the view of (20) when  $\lambda_k$  are given by (48) we get that

$$(49) \quad \widehat{f}(j) = \begin{cases} \frac{\Phi^{1/2}(\alpha_k)}{V^{1/2}(\alpha_k)}, & \text{if } j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k = 0, 1, \dots \\ 0, & \text{if } j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

Analogously to (37) if we apply (22) when  $\lambda_k$  are given by (48) we get that

$$S_{\alpha_k} f = \sum_{\eta=0}^{k-1} \frac{\Phi^{1/2}(\alpha_\eta)}{V^{1/2}(\alpha_\eta)} (D_{2^{|\alpha_\eta|+1}} - D_{2^{|\alpha_\eta|}}) + \frac{\Phi^{1/2}(\alpha_k)}{V^{1/2}(\alpha_k)} w_{2^{|\alpha_k|}} D_{\alpha_k - 2^{|\alpha_k|}}.$$

By applying first estimation of Lemma 3 and (47) we can conclude that

$$\begin{aligned} \left\| \frac{S_{\alpha_k} f}{\Phi(\alpha_k)} \right\|_1 &\geq \frac{\Phi^{1/2}(\alpha_k)}{\Phi(\alpha_k) V^{1/2}(\alpha_k)} \left\| D_{\alpha_k - 2^{|\alpha_k|}} \right\|_1 \\ &\quad - \frac{1}{\Phi(\alpha_k)} \sum_{\eta=0}^{k-1} \frac{\Phi^{1/2}(\alpha_\eta)}{V^{1/2}(\alpha_\eta)} \left\| D_{2^{|\alpha_\eta|+1}} - D_{2^{|\alpha_\eta|}} \right\|_1 \\ &\geq \frac{V(\alpha_k - 2^{|\alpha_k|}) \Phi^{1/2}(\alpha_k)}{8\Phi(\alpha_k) V^{1/2}(\alpha_k)} \\ &\quad - \frac{1}{\Phi(\alpha_k)} \sum_{\eta=0}^{\infty} \frac{\Phi^{1/2}(\alpha_\eta)}{V^{1/2}(\alpha_\eta)} \\ &\geq \frac{cV^{1/2}(\alpha_k)}{\Phi^{1/2}(\alpha_k)} \rightarrow \infty, \text{ as } k \rightarrow \infty. \end{aligned}$$

Theorem 2 is proved.

**Corollary 2.** *Let  $n \in \mathbb{N}$ ,  $0 < p \leq 1$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$ , such that*

$$(50) \quad \|S_{2^n} f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}.$$

**Proof:** To prove Theorem 2 we only have to show that

$$|2^n| = n, \quad \langle 2^n \rangle = n - 1 \text{ and } d(2^n) = 0.$$

By applying first part of Theorem 1 we immediately get that (50) for any  $0 < p \leq 1$  and proof of Corollary 2 is proved.

**Corollary 3.** *Let  $n \in \mathbb{N}$ ,  $0 < p \leq 1$  and  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$  such that*

$$(51) \quad \|S_{2^n+2^{n-1}}f\|_{H_p(G)} \leq c_p \|f\|_{H_p(G)}.$$

**Proof:** Since

$$|2^n + 2^{n-1}| = n, \langle 2^n + 2^{n-1} \rangle = n - 1 \quad \text{and} \quad d(2^n + 2^{n-1}) = 1$$

by first part of Theorem 1 we get that (51) holds, for any  $0 < p \leq 1$  and proof of Corollary 3 is complete.

**Corollary 4.** *Let  $n \in \mathbb{N}$  and  $0 < p < 1$ . Then there exists a martingale  $f \in H_p(G)$ , such that*

$$(52) \quad \sup_{n \in \mathbb{N}} \|S_{2^{n+1}}f\|_{weak-L_p(G)} = \infty.$$

*On the other hand, there exists an absolute constant  $c$ , such that*

$$(53) \quad \|S_{2^{n+1}}f\|_{H_1(G)} \leq c \|f\|_{H_1(G)}.$$

**Proof:** Since

$$(54) \quad |2^n + 1| = n, \langle 2^n + 1 \rangle = 0 \quad \text{and} \quad d(2^n + 1) = n.$$

by applying second part of Theorem 1 we get that there exists a martingale  $f = (f_n, n \in \mathbb{N}_+) \in H_p(G)$ , for  $0 < p < 1$ , such that (52) holds.

On the other hand, proof of (53) follows simple observation that

$$V(2^n + 1) = 4 < \infty.$$

Corollary 4 is proved.

#### 2.4. Modulus of continuity and convergence in norm of subsequences of partial sums with respect to the one-dimensional Walsh-Fourier series on the martingale Hardy spaces.

In this section we apply Theorem 1 and Theorem 2 to find necessary and sufficient conditions for modulus of continuity, for which subsequences of partial sums with respect to the one-dimensional Walsh-Fourier series are bounded in the martingale Hardy spaces.

First, we prove the following estimation:

**Theorem 3.** *Let  $n \in \mathbb{N}_+$  and  $2^k < n \leq 2^{k+1}$ . Then there exists an absolute constant  $c_p$ , depending only on  $p$  such that*

$$(55) \quad \|S_n f - f\|_{H_p(G)} \leq c_p 2^{d(n)(1/p-1)} \omega_{H_p(G)}\left(\frac{1}{2^k}, f\right), \quad (f \in H_p(G)) \quad (0 < p < 1)$$

and

$$(56) \quad \|S_n f - f\|_{H_1(G)} \leq c_1 V(n) \omega_{H_1(G)}\left(\frac{1}{2^k}, f\right), \quad (f \in H_1(G)).$$

**Proof:** Let  $0 < p < 1$  and  $2^k < n \leq 2^{k+1}$ . By applying first part of Theorem 1 we get that

$$\begin{aligned}
(57) \quad & \|S_n f - f\|_{H_p(G)}^p \\
& \leq c_p \|S_n f - S_{2^k} f\|_{H_p(G)}^p + c_p \|S_{2^k} f - f\|_{H_p(G)}^p \\
& = c_p \|S_n (S_{2^k} f - f)\|_{H_p(G)}^p + c_p \|S_{2^k} f - f\|_{H_p(G)}^p \\
& \leq c_p (1 + 2^{d(n)(1-p)}) \omega_{H_p(G)}^p \left( \frac{1}{2^k}, f \right) \\
& \leq c_p 2^{d(n)(1-p)} \omega_{H_p(G)}^p \left( \frac{1}{2^k}, f \right).
\end{aligned}$$

The proof of (56) is analogical to (55). Analogously to (55) we can also prove estimation (56). So, we leave out the details.

Theorem 3 is proved.

**Theorem 4.** a) Let  $k \in \mathbb{N}_+$ ,  $0 < p < 1$ ,  $f \in H_p(G)$  and  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that

$$(58) \quad \omega_{H_p(G)} \left( \frac{1}{2^{|m_k|}}, f \right) = o \left( \frac{1}{2^{d(m_k)(1/p-1)}} \right) \quad \text{as } k \rightarrow \infty.$$

Then

$$(59) \quad \|S_{m_k} f - f\|_{H_p(G)} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

b) Let  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that condition (25) is fulfilled. Then there exists a martingale  $f \in H_p(G)$  and increasing sequence of natural numbers  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{m_k : k \in \mathbb{N}_+\}$ , such that

$$\omega_{H_p(G)} \left( \frac{1}{2^{|\alpha_k|}}, f \right) = O \left( \frac{1}{2^{d(\alpha_k)(1/p-1)}} \right) \quad \text{as } k \rightarrow \infty$$

and

$$(60) \quad \limsup_{k \rightarrow \infty} \|S_{\alpha_k} f - f\|_{weak-L_p(G)} > c_p > 0, \quad \text{as } k \rightarrow \infty,$$

where  $c_p$  is an absolute constant depending only on  $p$ .

**Proof:** Let  $0 < p < 1$ ,  $f \in H_p(G)$  and  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that condition (58) is fulfilled. By combining Theorem 3 and estimation (55) we get that (59) holds true.

Now, prove second part of Theorem 4. In the view of (25) we simply get that there exists sequence  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{m_k : k \in \mathbb{N}_+\}$ , such that

$$(61) \quad 2^{d(\alpha_k)} \uparrow \infty, \quad \text{as } k \rightarrow \infty, \quad 2^{2(1/p-1)d(\alpha_k)} \leq 2^{(1/p-1)d(\alpha_{k+1})}.$$

Let  $f = (f_n, n \in \mathbb{N})$  be a martingale from Example 1, such that

$$(62) \quad \lambda_i = 2^{-(1/p-1)d(\alpha_i)}.$$

By applying (61) we obtain that condition (19) is fulfilled and it follows that  $f \in H_p(G)$ .

By applying (20), where  $\lambda_k$  are given by (62), then

$$(63) \quad \widehat{f}(j) = \begin{cases} 2^{(1/p-1)\langle\alpha_k\rangle}, & \text{if } j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k \in \mathbb{N}_+, \\ 0, & \text{if } j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

By combining (61) and (23) we have that

$$(64) \quad \omega_{H_p(G)}\left(\frac{1}{2^{|\alpha_k|}}, f\right) \leq \sum_{i=k}^{\infty} \frac{1}{2^{(1/p-1)d(\alpha_i)}} = O\left(\frac{1}{2^{(1/p-1)d(\alpha_k)}}\right), \text{ as } k \rightarrow \infty.$$

By using (39) we get that

$$\left|D_{\alpha_k - 2^{\langle\alpha_k\rangle}}\right| \geq 2^{\langle\alpha_k\rangle}, \quad \text{where } I_{\langle\alpha_k\rangle} \setminus I_{\langle\alpha_k\rangle+1}.$$

In the view of (22) we can conclude that

$$S_{\alpha_k} f = S_{2^{|\alpha_k|}} f + 2^{(1/p-1)\langle\alpha_k\rangle} w_{2^{|\alpha_k|}} D_{\alpha_k - 2^{|\alpha_k|}},$$

Since

$$\begin{aligned} \|D_{\alpha_k}\|_{weak-L_p(G)} &\geq 2^{\langle\alpha_k\rangle} \mu \{x \in I_{\langle\alpha_k\rangle} \setminus I_{\langle\alpha_k\rangle+1} : |D_{\alpha_k}| \geq 2^{\langle\alpha_k\rangle}\}^{1/p} \\ &\geq 2^{\langle\alpha_k\rangle} (\mu \{I_{\langle\alpha_k\rangle} \setminus I_{\langle\alpha_k\rangle+1}\})^{1/p} \geq 2^{\langle\alpha_k\rangle(1-1/p)}, \end{aligned}$$

if we apply (2) (see also Theorem T2) we obtain that

$$\begin{aligned} \|f - S_{\alpha_k} f\|_{weak-L_p(G)}^p &\geq 2^{(1-p)\langle\alpha_k\rangle} \|w_{2^{|\alpha_k|}} D_{\alpha_k - 2^{|\alpha_k|}}\|_{weak-L_p(G)}^p \\ &\quad - \|f - S_{2^{|\alpha_k|}} f\|_{weak-L_p(G)}^p \\ &\geq c - o(1) > c > 0, \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Proof of Theorem 4 is complete.

**Corollary 5.** *a) Let  $0 < p < 1$ ,  $f \in H_p(G)$  and  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that*

$$(65) \quad \omega_{H_p(G)}\left(\frac{1}{2^{|m_k|}}, f\right) = o\left(\frac{1}{(m_k \mu(\text{supp} D_{m_k}))^{1/p-1}}\right) \text{ as } k \rightarrow \infty.$$

*Then (59) holds.*

*b) Let  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that*

$$(66) \quad \sup_{k \in \mathbb{N}_+} m_k \mu \{\text{supp}(D_{m_k})\} = \infty.$$

*Then there exist a martingale  $f \in H_p(G)$  and sequence  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{m_k : k \in \mathbb{N}_+\}$ , such that*

$$\omega_{H_p(G)}\left(\frac{1}{2^{|\alpha_k|}}, f\right) = O\left(\frac{1}{(\alpha_k \mu(\text{supp} D_{\alpha_k}))^{1/p-1}}\right) \text{ as } k \rightarrow \infty$$

*and (60) holds.*

**Theorem 5.** a) Let  $f \in H_1(G)$  and  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that

$$(67) \quad \omega_{H_1(G)} \left( \frac{1}{2^{|m_k|}}, f \right) = o \left( \frac{1}{V(m_k)} \right) \quad \text{as } k \rightarrow \infty.$$

Then

$$(68) \quad \|S_{m_k} f - f\|_{H_1(G)} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

b) Let  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that condition (43) is fulfilled. Then there exists a martingale  $f \in H_1(G)$  and increasing sequence of natural numbers  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{m_k : k \in \mathbb{N}_+\}$  such that

$$\omega_{H_1(G)} \left( \frac{1}{2^{|\alpha_k|}}, f \right) = O \left( \frac{1}{V(\alpha_k)} \right) \quad \text{as } k \rightarrow \infty$$

and

$$(69) \quad \limsup_{k \rightarrow \infty} \|S_{\alpha_k} f - f\|_1 > c > 0 \quad \text{as } k \rightarrow \infty,$$

where  $c$  is an absolute constant.

**Proof:** Let  $f \in H_1(G)$  and  $\{m_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that (67). By applying Theorem 3 we get that condition (68) is fulfilled.

Now, we prove second part of Theorem 5. By applying (43) we conclude that there exists sequence  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{m_k : k \in \mathbb{N}_+\}$ , such that

$$(70) \quad V(\alpha_k) \uparrow \infty, \quad \text{as } k \rightarrow \infty \quad \text{and} \quad V^2(\alpha_k) \leq V(\alpha_{k+1}) \quad k \in \mathbb{N}_+.$$

Let  $f = (f_n, n \in \mathbb{N}_+)$  be a martingale from the Example 1, where

$$\lambda_k = \frac{1}{V(\alpha_k)}.$$

By applying (70) we conclude that (19) is fulfilled and we conclude that  $f = (f_n, n \in \mathbb{N}_+) \in H_1(G)$ .

In the view of (20) we have that

$$(71) \quad \widehat{f}(j) = \begin{cases} \frac{1}{V(\alpha_k)}, & \text{if } j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, \quad k = 0, 1, \dots \\ 0, & \text{if } j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

According to (23) we get that

$$\begin{aligned} w_{H_1(G)}(1/2^n, f) &= \|f - S_{2^n} f\|_{H_1(G)} \leq \sum_{i=n+1}^{\infty} \frac{1}{V(\alpha_i)} \\ &= O \left( \frac{1}{V(\alpha_n)} \right), \quad \text{as } n \rightarrow \infty. \end{aligned}$$

By applying (22) we can conclude that

$$S_{\alpha_k} f = S_{2^{|\alpha_k|}} f + \frac{w_{2^{|\alpha_k|}} D_{\alpha_k - 2^{|\alpha_k|}}}{V(\alpha_k)},$$

If we use (2) and Theorem T2 we get that

$$\begin{aligned} \|f - S_{\alpha_k} f\|_1 &\geq \left\| \frac{w_{2^{|\alpha_k|}} D_{\alpha_k - 2^{|\alpha_k|}}}{V(\alpha_k)} \right\|_1 - \|f - S_{2^{|\alpha_k|}} f\|_1 \\ &\geq \frac{V(\alpha_k - 2^{|\alpha_k|})}{8V(\alpha_k)} - o(1) > c > 0, \quad \text{as } k \rightarrow \infty. \end{aligned}$$

The proof of Theorem 5 is proved.

Theorem 9 follows the following corollaries which are [59]:

**Corollary 6.** a) Let  $0 < p < 1$ ,  $f \in H_p(G)$  and

$$\omega_{H_p(G)}(1/2^k, f) = o(1/2^{k(1/p-1)}), \quad \text{as } k \rightarrow \infty.$$

Then

$$\|S_k f - f\|_{H_p(G)} \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

b) There exists a martingale  $f \in H_p(G)$  ( $0 < p < 1$ ), such that

$$\omega_{H_p(G)}(1/2^k, f) = O(1/2^{k(1/p-1)}), \quad \text{as } k \rightarrow \infty$$

and

$$\|S_k f - f\|_{\text{weak-}L_p(G)} \not\rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

**Corollary 7.** a) Let  $f \in H_1(G)$  and

$$\omega_{H_1(G)}(1/2^k, f) = o\left(\frac{1}{k}\right), \quad \text{as } k \rightarrow \infty.$$

Then

$$\|S_k f - f\|_{H_1(G)} \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

b) There exists a martingale  $f \in H_1(G)$ , such that

$$\omega_{H_1(G)}(1/2^k, f) = O\left(\frac{1}{k}\right), \quad \text{as } k \rightarrow \infty$$

and

$$\|S_k f - f\|_1 \not\rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

### 3. FEJÉR MEANS WITH RESPECT TO THE ONE-DIMENSIONAL WALSH-FOURIER SERIES ON THE MARTINGALE HARDY SPACES

#### 3.1. Basic notations.

For the one-dimensional case Fejér means with respect to the one-dimensional Walsh-Fourier series  $\sigma_n$  is defined by:

$$\sigma_n f(x) : = \frac{1}{n} \sum_{k=1}^n S_k f(x) \quad (n \in \mathbb{N}_+).$$

The following equality is true (for details see [2] and [45]):

$$\begin{aligned} \sigma_n f(x) &= \frac{1}{n} \sum_{k=0}^{n-1} (D_k * f)(x) \\ &= (f * K_n)(x) = \int_{G_m} f(t) K_n(x-t) d\mu(t). \end{aligned}$$

where

$$K_n(x) : = \frac{1}{n} \sum_{k=1}^n D_k(x) \quad (n \in \mathbb{N}_+)$$

In the literature  $K_n$  is called  $n$ -th Fejér kernel.

We also define the following maximal operators

$$\begin{aligned} \sigma^* f &= \sup_{n \in \mathbb{N}} |\sigma_n f| \\ \tilde{\sigma}_{\#}^* f &= \sup_{n \in \mathbb{N}} |\sigma_{2^n} f|. \end{aligned}$$

For any natural number  $n \in \mathbb{N}$  we also need the following expression

$$n = \sum_{i=1}^s 2^{n_i}, \quad n_1 < n_2 < \dots < n_s.$$

Set

$$n^{(i)} := 2^{n_1} + \dots + 2^{n_{i-1}}, \quad i = 2, \dots, s$$

and

$$\mathbb{A}_{0,2} := \left\{ n \in \mathbb{N} : n = 2^0 + 2^2 + \sum_{i=3}^{s_n} 2^{n_i} \right\}.$$

Then, for any natural number  $n \in \mathbb{N}$  there exists numbers

$$0 \leq l_1 \leq m_1 \leq l_2 - 2 < l_2 \leq m_2 \leq \dots \leq l_s - 2 < l_s \leq m_s$$

such that it can be written as

$$n = \sum_{i=1}^s \sum_{k=l_i}^{m_i} 2^k,$$

where  $s$  is depending on  $n$ .

It is evident that

$$s \leq V(n) \leq 2s + 1.$$

**3.2. Auxiliary lemmas.** The following equality and estimation of Fejér kernels with respect to the one-dimensional Walsh-Fourier series is proved in [45]:

**Lemma 8.** *Let  $n \in \mathbb{N}$  and  $n = \sum_{i=1}^s 2^{n_i}$ ,  $n_1 < n_2 < \dots < n_s$ . Then*

$$nK_n = \sum_{r=1}^s \left( \prod_{j=r+1}^s w_{2^{n_j}} \right) 2^{n_r} K_{2^{n_r}} + \sum_{t=2}^s \left( \prod_{j=t+1}^s w_{2^{n_j}} \right) n^{(t)} D_{2^{n_t}},$$

and

$$\sup_{n \in \mathbb{N}} \int_G |K_n(x)| d\mu(x) \leq c < \infty,$$

where  $c$  is an absolute constant.

The following equality is proved in [45] (see also [17]):

**Lemma 9.** *Let  $n > t$  and  $t, n \in \mathbb{N}$ . Then we have the following expression for  $2^n$ -th Fejér kernels with respect to the one-dimensional Walsh-Fourier series:*

$$K_{2^n}(x) = \begin{cases} 2^{t-1}, & \text{if } x \in I_n(e_t), \\ \frac{2^n+1}{2}, & \text{if } x \in I_n, \\ 0, & \text{otherwise.} \end{cases}$$

The following estimation is proved by Goginava [22]:

**Lemma 10.** *Let  $x \in I_{l+1}(e_k + e_l)$ ,  $k = 0, \dots, M-2$ ,  $l = 0, \dots, M-1$ . Then*

$$\int_{I_M} |K_n(x+t)| d\mu(t) \leq \frac{c2^{l+k}}{n2^M}, \quad \text{where } n > 2^M.$$

Let  $x \in I_M(e_k)$ ,  $m = 0, \dots, M-1$ . Then

$$\int_{I_M} |K_n(x+t)| d\mu(t) \leq \frac{c2^k}{2^M}, \quad \text{for } n > 2^M,$$

where  $c$  is an absolute constant.

The following estimations of Fejér kernels with respect to the one-dimensional Walsh-Fourier series is proved in [62]:

**Lemma 11.** *Let*

$$n = \sum_{i=1}^r \sum_{k=l_i}^{m_i} 2^k,$$

where

$$m_1 \geq l_1 > l_1 - 2 \geq m_2 \geq l_2 > l_2 - 2 > \dots > m_s \geq l_s \geq 0.$$

Then

$$|nK_n| \leq c \sum_{A=1}^r \left( 2^{l_A} |K_{2^{l_A}}| + 2^{m_A} |K_{2^{m_A}}| + 2^{l_A} \sum_{k=l_A}^{m_A} D_{2^k} \right) + cV(n),$$

where  $c$  is an absolute constant.

**Proof:** Let

$$n = \sum_{i=1}^r 2^{n_i}, n_1 > n_2 > \dots > n_r \geq 0.$$

By using Lemma 8 for  $n$ -th Fejér kernels we can conclude that

$$\begin{aligned} nK_n &= \sum_{A=1}^r \left( \prod_{j=1}^{A-1} w_{2^{n_j}} \right) ((2^{n_A} K_{2^{n_A}} + (2^{n_A} - 1) D_{2^{n_A}})) \\ &- \sum_{A=1}^r \left( \prod_{j=1}^{A-1} w_{2^{n_j}} \right) (2^{n_A} - 1 - n^{(A)}) D_{2^{n_A}} = I_1 - I_2. \end{aligned}$$

For  $I_1$  we have the following equality

$$\begin{aligned} I_1 &= \sum_{v=1}^r \left( \prod_{j=1}^{v-1} \prod_{i=l_j}^{m_j} w_{2^i} \right) \left( \sum_{k=l_v}^{m_v} \left( \prod_{j=k+1}^{m_v} w_{2^j} \right) (2^k K_{2^k} - (2^k - 1) D_{2^k}) \right) \\ &= \sum_{v=1}^r \left( \prod_{j=1}^{v-1} \prod_{i=l_j}^{m_j} w_{2^i} \right) \left( \sum_{k=0}^{m_v} - \sum_{k=0}^{l_v-1} \right) \left( \prod_{j=k+1}^{m_v} w_{2^j} \right) (2^k K_{2^k} - (2^k - 1) D_{2^k}) \\ &= \sum_{v=1}^r \left( \prod_{j=1}^{v-1} \prod_{i=l_j}^{m_j} w_{2^i} \right) \left( \sum_{k=0}^{m_v} \left( \prod_{j=k+1}^{m_v} w_{2^j} \right) (2^k K_{2^k} - (2^k - 1) D_{2^k}) \right) \\ &- \sum_{v=1}^r \left( \prod_{j=1}^v \prod_{i=l_j}^{m_j} w_{2^i} \right) \left( \sum_{k=0}^{l_v-1} \left( \prod_{j=k+1}^{l_v-1} w_{2^j} \right) (2^k K_{2^k} - (2^k - 1) D_{2^k}) \right). \end{aligned}$$

Since

$$2^n - 1 = \sum_{k=0}^{n-1} 2^k$$

and

$$(2^n - 1) K_{2^{n-1}} = \sum_{k=0}^{n-1} \left( \prod_{j=k+1}^{n-1} w_{2^j} \right) (2^k K_{2^k} - (2^k - 1) D_{2^k}),$$

we obtain that

$$\begin{aligned} I_1 &= \sum_{v=1}^r \left( \prod_{j=1}^{v-1} \prod_{i=l_j}^{m_j} w_{2^i} \right) (2^{m_v+1} - 1) K_{2^{m_v+1-1}} \\ &- \sum_{v=1}^r \left( \prod_{j=1}^v \prod_{i=l_j}^{m_j} w_{2^i} \right) (2^{l_v} - 1) K_{2^{l_v-1}}. \end{aligned}$$

If we apply estimations

$$|K_{2^n}| \leq c |K_{2^{n-1}}|$$

and

$$|K_{2^{n-1}}| \leq c |K_{2^n}| + c$$

we get that

$$(72) \quad |I_1| \leq c \sum_{v=1}^r (2^{l_v} |K_{2^{l_v}}| + 2^{m_v} |K_{2^{m_v}}| + cr).$$

Let  $l_j < n_A \leq m_j$ , where  $j = 1, \dots, s$ . Then

$$n^{(A)} \geq \sum_{v=l_j}^{n_A-1} 2^v \geq 2^{n_A} - 2^{l_j}$$

and

$$2^{n_A} - 1 - n^{(A)} \leq 2^{l_j}.$$

If  $l_j = n_A$  where  $j = 1, \dots, s$ , then

$$n^{(A)} \leq 2^{m_{j-1}+1} < 2^{l_j}.$$

By using these estimations we can conclude that

$$(73) \quad |I_2| \leq c \sum_{v=1}^r 2^{l_v} \sum_{k=l_v}^{m_v} D_{2^k}.$$

By combining (72)-(73) we get the proof of Lemma 11.

The following estimations of Fejér kernels with respect to the one-dimensional Walsh-Fourier series is proved in [62]:

**Lemma 12.** *Let*

$$n = \sum_{i=1}^s \sum_{k=l_i}^{m_i} 2^k,$$

where

$$0 \leq l_1 \leq m_1 \leq l_2 - 2 < l_2 \leq m_2 \leq \dots \leq l_s - 2 < l_s \leq m_s.$$

Then

$$n |K_n(x)| \geq \frac{2^{2l_i}}{16}, \quad \text{for } x \in I_{l_i+1}(e_{l_i-1} + e_{l_i}).$$

**Proof:** If we apply Lemma 8 for  $n = \sum_{i=1}^s \sum_{k=l_i}^{m_i} 2^k$  we can write that

$$\begin{aligned} nK_n &= \sum_{r=1}^s \sum_{k=l_r}^{m_r} \left( \prod_{j=r+1}^s \prod_{q=l_j}^{m_j} w_{2^q} \prod_{j=k+1}^{m_r} w_{2^j} \right) 2^k K_{2^k} \\ &+ \sum_{r=1}^s \sum_{k=l_r}^{m_r} \left( \prod_{j=r+1}^s \prod_{q=l_j}^{m_j} w_{2^q} \prod_{j=k+1}^{m_r} w_{2^j} \right) \left( \sum_{t=1}^{r-1} \sum_{q=l_t}^{m_t} 2^q + \sum_{q=l_r}^{k-1} 2^q \right) D_{2^k}. \end{aligned}$$

Let  $x \in I_{l_i+1}(e_{l_i-1} + e_{l_i})$ . Then

$$n |K_n| \geq |2^{l_i} K_{2^{l_i}}| - \sum_{r=1}^{i-1} \sum_{k=l_r}^{m_r} |2^k K_{2^k}| - \sum_{r=1}^{i-1} \sum_{k=l_r}^{m_r} |2^k D_{2^k}| = I - II - III.$$

Lemma 9 follows that

$$(74) \quad I = |2^{l_i} K_{2^{l_i}}(x)| = \frac{2^{2l_i}}{4}.$$

Since  $m_{i-1} \leq l_i - 2$ , we easily obtain that the following estimation is true:

$$(75) \quad \begin{aligned} II &\leq \sum_{n=0}^{l_i-2} |2^n K_{2^n}(x)| \\ &\leq \sum_{n=0}^{l_i-2} 2^n \frac{(2^n + 1)}{2} \\ &\leq \frac{2^{2l_i}}{24} + \frac{2^{l_i}}{4} - \frac{2}{3}. \end{aligned}$$

For  $III$  we get that

$$(76) \quad III \leq \sum_{k=0}^{l_i-2} |2^k D_{2^k}(x)| \leq \sum_{k=0}^{l_i-2} 4^k = \frac{2^{2l_i}}{12} - \frac{1}{3}.$$

By combining (74-76) we can conclude that

$$(77) \quad n |K_n(x)| \geq I - II - III \geq \frac{2^{2l_i}}{8} - \frac{2^{l_i}}{4} + 1.$$

Suppose that  $l_i \geq 2$ . Then

$$n |K_n(x)| \geq \frac{2^{2l_i}}{8} - \frac{2^{2l_i}}{16} \geq \frac{2^{2l_i}}{16}.$$

If  $l_i = 0$  or  $l_i = 1$ , then by applying (77) we get that

$$n |K_n(x)| \geq \frac{7}{8} \geq \frac{2^{2l_i}}{16},$$

Lemma is proved.

The following estimations of Fejér kernels with respect to the one-dimensional Walsh-Fourier series is proved in [62] (see also [72]):

**Lemma 13.** *Let  $0 < p \leq 1$ ,  $2^k \leq n < 2^{k+1}$  and  $\sigma_n f$  be Fejér means with respect to the one-dimensional Walsh-Fourier series, where  $f \in H_p(G)$ . Then, for any fixed  $n \in \mathbb{N}$ ,*

$$\begin{aligned} &\|\sigma_n f\|_{H_p(G)} \\ &\leq \left\| \sup_{0 \leq l \leq k} |\sigma_{2^l} f| \right\|_p + \left\| \sup_{0 \leq l \leq k} |S_{2^l} f| \right\|_p + \|\sigma_n f\|_p \\ &\leq \|\tilde{\sigma}_{\#}^* f\|_p + \|\tilde{S}_{\#}^* f\|_p + \|\sigma_n f\|_p. \end{aligned}$$

**Proof:** Let consider the following martingale

$$\begin{aligned} f_{\#} &= (S_{2^k} \sigma_n f, k \in \mathbb{N}) \\ &= \left( \frac{2^0 \sigma_{2^0}}{n} + \frac{(n - 2^0) S_{2^0} f}{n}, \dots, \frac{2^k \sigma_{2^k} f}{n} + \frac{(n - 2^k) S_{2^k} f}{n}, \sigma_n f, \dots, \sigma_n f, \dots \right). \end{aligned}$$

By using Lemma 6 we immediately get

$$\begin{aligned} & \|\sigma_n f\|_{H_p(G^2)}^p \\ & \leq \left\| \sup_{0 \leq l \leq k} |\sigma_{2^l} f| \right\|_p^p + \left\| \sup_{0 \leq l \leq k} |S_{2^l} f| \right\|_p^p + \|S_n f\|_p^p \\ & \leq \|\tilde{\sigma}_\#^* f\|_p^p + \|\tilde{S}_\#^* f\|_p^p + \|\sigma_n f\|_p^p. \end{aligned}$$

Lemma is proved.

### 3.3. Boundedness of subsequences of Fejér means with respect to the one-dimensional Walsh-Fourier series on the martingale Hardy spaces.

In this section we study boundedness of subsequences of Fejér means with respect to the one-dimensional Walsh-Fourier series in the martingale Hardy spaces (For details see [62]).

First, we consider case  $p = 1/2$ . The following estimation is true:

**Theorem 6.** *a) Let  $f \in H_{1/2}(G)$ . Then there exists an absolute constant  $c$ , such that*

$$\|\sigma_n f\|_{H_{1/2}(G)} \leq cV^2(n) \|f\|_{H_{1/2}(G)}.$$

*b) Let  $\{n_k : k \in \mathbb{N}_+\}$  be increasing sequence of natural numbers, such that  $\sup_{k \in \mathbb{N}_+} V(n_k) = \infty$  and  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function satisfying the conditions  $\Phi(n) \uparrow \infty$  and*

$$(78) \quad \lim_{k \rightarrow \infty} \frac{V^2(n_k)}{\Phi(n_k)} = \infty.$$

*Then there exists a martingale  $f \in H_{1/2}(G)$ , such that*

$$\sup_{k \in \mathbb{N}} \left\| \frac{\sigma_{n_k} f}{\Phi(n_k)} \right\|_{1/2} = \infty.$$

**Proof:** Suppose that

$$(79) \quad \left\| \frac{\sigma_n f}{V^2(n)} \right\|_{1/2} \leq c \|f\|_{H_{1/2}(G)}.$$

By combining estimations (7), (15) and Lemma 13 we can conclude that

$$(80) \quad \begin{aligned} & \left\| \frac{\sigma_n f}{V^2(n)} \right\|_{H_{1/2}(G)}^{1/2} \\ & \leq \left\| \frac{\sigma_n f}{V^2(n)} \right\|_{1/2}^{1/2} + \frac{1}{V^2(n)} \|\sigma_\#^* f\|_{1/2}^{1/2} + \frac{1}{V^2(n)} \|\tilde{S}_\#^*\|_{1/2}^{1/2} \\ & \leq \left\| \frac{\tilde{\sigma}_n f}{V^2(n)} \right\|_{1/2}^{1/2} + \|\tilde{\sigma}_\#^* f\|_{1/2}^{1/2} + \|\tilde{S}_\#^* f\|_{1/2}^{1/2} \leq c \|f\|_{H_{1/2}(G)}^{1/2}. \end{aligned}$$

By combining Lemma 5 and (80), Theorem 6 will be proved if we show that

$$\int_{I_M} \left( \frac{|\sigma_n a|}{V^2(n)} \right)^{1/2} d\mu \leq c < \infty,$$

for any 1/2-atom  $a$ .

Without loss the generality we may assume that  $a$  is  $1/2$ -atom, with support  $I$ , for which  $\mu(I) = 2^{-M}$ ,  $I = I_M$ . It is easy to check that  $\sigma_n(a) = 0$ , when  $n \leq 2^M$ . Therefore, we may assume that  $n > 2^M$ . Set

$$\begin{aligned} II_{\alpha_A}^1(x) &:= 2^M \int_{I_M} 2^{\alpha_A} |K_{2^{\alpha_A}}(x+t)| d\mu(t), \\ II_{l_A}^2(x) &= 2^M \int_{I_M} 2^{l_A} \sum_{k=l_A}^{m_A} D_{2^k}(x+t) d\mu(t). \end{aligned}$$

Let  $x \in I_M$ . Since  $\sigma_n$  is bounded from  $L_\infty(G)$  to  $L_\infty(G)$ , for  $n > 2^M$  and  $\|a\|_\infty \leq 2^{2M}$ , by using Lemma 10 we can conclude that

$$\begin{aligned} & \frac{|\sigma_n a(x)|}{V^2(n)} \\ & \leq \frac{c}{V^2(n)} \int_{I_M} |a(x)| |K_n(x+t)| d\mu(t) \\ & \leq \frac{c \|a\|_\infty}{V^2(n)} \int_{I_M} |K_n(x+t)| d\mu(t) \\ & \leq \frac{c 2^{2M}}{V^2(n)} \int_{I_M} |K_n(x+t)| d\mu(t) \\ & \leq \frac{c 2^M}{V^2(n)} \left\{ \sum_{A=1}^s \int_{I_M} 2^{l_A} |K_{2^{l_A}}(x+t)| d\mu(t) + \int_{I_M} 2^{m_A} |K_{2^{m_A}}(x+t)| d\mu(t) \right\} \\ & + \frac{c 2^M}{V^2(n)} \sum_{A=1}^s \int_{I_M} 2^{l_A} \sum_{k=l_A}^{m_A} D_{2^k}(x+t) d\mu(t) + \frac{c 2^M}{V^2(n)} \int_{I_M} V(n) d\mu(t) \\ & = \frac{c}{V^2(n)} \sum_{A=1}^s (II_{l_A}^1(x) + II_{m_A}^1(x) + II_{l_A}^2(x)) + c. \end{aligned}$$

Hence,

$$\begin{aligned} & \int_{I_M} \left| \frac{\sigma_n a(x)}{V^2(n)} \right|^{1/2} d\mu(x) \\ & \leq \frac{c}{V(n)} \left( \sum_{A=1}^s \int_{I_M} |II_{l_A}^1(x)|^{1/2} d\mu(x) \right. \\ & \quad \left. + \int_{I_M} |II_{m_A}^1(x)|^{1/2} d\mu(x) + \int_{I_M} |II_{l_A}^2(x)|^{1/2} d\mu(x) \right) + c. \end{aligned}$$

Since  $s \leq 4V(n)$  we obtain that Theorem 6 will be proved if we show that

$$(81) \quad \int_{I_M} |II_{\alpha_A}^1(x)|^{1/2} d\mu(x) \leq c < \infty, \quad \int_{I_M} |II_{l_A}^2(x)|^{1/2} d\mu(x) \leq c < \infty,$$

where  $\alpha_A = l_A$  or  $\alpha_A = m_A$ ,  $A = 1, \dots, s$ .

Let  $t \in I_M$  and  $x \in I_{l+1}(e_k + e_l)$ ,  $0 \leq k < l < \alpha_A \leq M$  or  $0 \leq k < l \leq M \leq \alpha_A$ . Since  $x + t \in I_{l+1}(e_k + e_l)$ , by applying Lemma 9 we can conclude that

$$(82) \quad K_{2^{\alpha_A}}(x + t) = 0 \quad \text{and} \quad II_{\alpha_A}^1(x) = 0.$$

Let  $x \in I_{l+1}(e_k + e_l)$ ,  $0 \leq k < \alpha_A \leq l \leq M$ . Then  $x + t \in I_{l+1}(e_k + e_l)$ , where  $t \in I_M$  and if we apply again Lemma 9 we get that

$$(83) \quad 2^{\alpha_A} |K_{2^{\alpha_A}}(x + t)| \leq 2^{\alpha_A + k} \quad \text{and} \quad II_{\alpha_A}^1(x) \leq 2^{\alpha_A + k}.$$

Analogously to (83) for  $0 \leq \alpha_A \leq k < l \leq M$  we can prove that

$$(84) \quad 2^{\alpha_A} |K_{2^{\alpha_A}}(x + t)| \leq 2^{2\alpha_A}, \quad II_{\alpha_A}^1(x) \leq 2^{2\alpha_A}, \quad t \in I_M, \quad x \in I_{l+1}(e_k + e_l).$$

Let  $0 \leq \alpha_A \leq M - 1$ , where  $A = 1, \dots, s$ . According to (17) and (82-84) we find that

$$\begin{aligned} & \int_{I_M} |II_{\alpha_A}^1(x)|^{1/2} d\mu(x) \\ &= \sum_{k=0}^{M-2} \sum_{l=k+1}^{M-1} \int_{I_{l+1}(e_k + e_l)} |II_{\alpha_A}^1(x)|^{1/2} d\mu(x) \\ &+ \sum_{k=0}^{M-1} \int_{I_M(e_k)} |II_{\alpha_A}^1(x)|^{1/2} d\mu(x) \\ &\leq c \sum_{k=0}^{\alpha_A-1} \sum_{l=\alpha_A+1}^{M-1} \int_{I_{l+1}(e_k + e_l)} 2^{(\alpha_A+k)/2} d\mu(x) \\ &+ c \sum_{k=\alpha_A}^{M-2} \sum_{l=k+1}^{M-1} \int_{I_{l+1}(e_k + e_l)} 2^{\alpha_A} d\mu(x) \\ &+ c \sum_{k=0}^{\alpha_A-1} \int_{I_M(e_k)} 2^{(\alpha_A+k)/2} d\mu(x) + c \sum_{k=\alpha_A}^{M-1} \int_{I_M(e_k)} 2^{\alpha_A} d\mu(x) \\ &\leq c \sum_{k=0}^{\alpha_A-1} \sum_{l=\alpha_A+1}^{M-1} \frac{2^{(\alpha_A+k)/2}}{2^l} + c \sum_{k=\alpha_A}^{M-2} \sum_{l=k+1}^{M-1} \frac{2^{\alpha_A}}{2^l} \\ &+ c \sum_{k=0}^{\alpha_A-1} \frac{2^{(\alpha_A+k)/2}}{2^M} + c \sum_{k=\alpha_A}^{M-1} \frac{2^{\alpha_A}}{2^M} \leq c < \infty. \end{aligned}$$

Let  $\alpha_A \geq M$ . Analogously to  $II_{\alpha_A}^1(x)$  we can prove (81), for  $A = 1, \dots, s$ .

Now, prove boundedness of  $II_{l_A}^2$ . Let  $t \in I_M$  and  $x \in I_i \setminus I_{i+1}$ ,  $i \leq l_A - 1$ . Since  $x + t \in I_i \setminus I_{i+1}$ , if we apply first equality of Lemma 2 we get that

$$(85) \quad II_{l_A}^2(x) = 0.$$

Let  $x \in I_i \setminus I_{i+1}$ ,  $l_A \leq i \leq m_A$ . Since  $n \geq 2^M$  and  $t \in I_M$ , if we apply first equality of Lemma 2 we get that

$$(86) \quad II_{l_A}^2(x) \leq 2^M \int_{I_M} 2^{l_A} \sum_{k=l_A}^i D_{2^k}(x + t) d\mu(t) \leq c 2^{l_A + i}.$$

Let  $x \in I_i \setminus I_{i+1}$ ,  $m_A < i \leq M-1$ . Then  $x+t \in I_i \setminus I_{i+1}$ , for any  $t \in I_M$  and by first equality of Lemma 2 we have that

$$(87) \quad II_{l_A}^2(x) \leq c2^M \int_{I_M} 2^{l_A+m_A} \leq c2^{l_A+m_A}.$$

Let  $0 \leq l_A \leq m_A \leq M$ . Then, in the view of (17) and (85-87) we can conclude that

$$\begin{aligned} & \int_{I_M} |II_{l_A}^2(x)|^{1/2} d\mu(x) \\ &= \left( \sum_{i=0}^{l_A-1} + \sum_{i=l_A}^{m_A} + \sum_{i=m_A+1}^{M-1} \right) \int_{I_i \setminus I_{i+1}} |II_{l_A}^2(x)|^{1/2} d\mu(x) \\ &\leq c \sum_{i=l_A}^{m_A} \int_{I_i \setminus I_{i+1}} 2^{(l_A+i)/2} d\mu(x) \\ &+ c \sum_{i=m_A+1}^{M-1} \int_{I_i \setminus I_{i+1}} 2^{(l_A+m_A)/2} d\mu(x) \\ &\leq c \sum_{i=l_A}^{m_A} 2^{(l_A+i)/2} \frac{1}{2^i} \\ &+ c \sum_{i=m_A+1}^{M-1} 2^{(l_A+m_A)/2} \frac{1}{2^i} \leq c < \infty. \end{aligned}$$

Analogously, we can prove same estimations in the cases  $0 \leq l_A \leq M < m_A$  and  $M \leq l_A \leq m_A$ .

Now, we prove part b) of Theorem 6. According to (78), there exists increasing sequence  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{n_k : k \in \mathbb{N}_+\}$  of natural numbers such that

$$(88) \quad \sum_{k=1}^{\infty} \frac{\Phi^{1/4}(\alpha_k)}{V^{1/2}(\alpha_k)} \leq c < \infty.$$

Let  $f = (f_n, n \in \mathbb{N}_+)$  be martingale form Example 1, where

$$\lambda_k := \Phi^{1/2}(\alpha_k) / V(\alpha_k).$$

According to (88) we get that condition (19) is fulfilled and it follows that  $f = (f_n, n \in \mathbb{N}_+)$ .

By applying (20) we get that

$$(89) \quad \widehat{f}(j) = \begin{cases} 2^{|\alpha_k|} \Phi^{1/2}(\alpha_k) / V(\alpha_k), & j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k \in \mathbb{N}_+ \\ 0, & j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

Let  $2^{|\alpha_k|} < j < \alpha_k$ . If we apply (22) we get that

$$(90) \quad S_j f = S_{2^{|\alpha_k|}} f + \frac{w_{2^{|\alpha_k|}} D_{j-2^{|\alpha_k|}} \Phi^{1/2}(\alpha_k)}{V(\alpha_k)}$$

Hence,

$$\begin{aligned}
(91) \quad & \frac{\sigma_{\alpha_k} f}{\Phi(\alpha_k)} \\
&= \frac{1}{\Phi(\alpha_k) \alpha_k} \sum_{j=1}^{2^{|\alpha_k|}} S_j f \\
&+ \frac{1}{\Phi(\alpha_k) \alpha_k} \sum_{j=2^{|\alpha_k|+1}}^{\alpha_k} S_j f \\
&= \frac{\sigma_{2^{|\alpha_k|}} f}{\Phi(\alpha_k) \alpha_k} \\
&+ \frac{(\alpha_k - 2^{|\alpha_k|}) S_{2^{|\alpha_k|}} f}{\Phi(\alpha_k) \alpha_k} \\
&+ \frac{w_{2^{|\alpha_k|}} 2^{|\alpha_k|} \Phi^{1/2}(\alpha_k)}{\Phi(\alpha_k) V(\alpha_k) \alpha_k} \sum_{j=2^{|\alpha_k|+1}}^{\alpha_k} D_{j-2^{|\alpha_k|}} \\
&= III_1 + III_2 + III_3.
\end{aligned}$$

For  $III_3$  we can conclude that

$$\begin{aligned}
(92) \quad & |III_3| \\
&= \frac{2^{|\alpha_k|} \Phi^{1/2}(\alpha_k)}{\Phi(\alpha_k) V(\alpha_k) \alpha_k} \left| \sum_{j=1}^{\alpha_k - 2^{|\alpha_k|}} D_j \right| \\
&= \frac{2^{|\alpha_k|} \Phi^{1/2}(\alpha_k)}{\Phi(\alpha_k) V(\alpha_k) \alpha_k} (\alpha_k - 2^{|\alpha_k|}) \left| K_{\alpha_k - 2^{|\alpha_k|}} \right| \\
&\geq \frac{c(\alpha_k - 2^{|\alpha_k|}) \left| K_{\alpha_k - 2^{|\alpha_k|}} \right|}{\Phi^{1/2}(\alpha_k) V(\alpha_k)}.
\end{aligned}$$

Let

$$\alpha_k = \sum_{i=1}^{r_k} \sum_{k=l_i^k}^{m_i^k} 2^k,$$

where

$$m_1^k \geq l_1^k > l_1^k - 2 \geq m_2^k \geq l_2^k > l_2^k - 2 \geq \dots \geq m_s^k \geq l_s^k \geq 0.$$

Since (see theorems 1 and 6)

$$\|III_1\|_{1/2} \leq c, \|III_2\|_{1/2} \leq c,$$

and

$$\mu \left\{ E_{l_i^k} \right\} \geq 1/2^{l_i^k - 1},$$

By combining (91), (92) and Lemma 12 we get that

$$\begin{aligned}
& \int_G |\sigma_{\alpha_k} f(x) / \Phi(\alpha_k)|^{1/2} d\mu(x) \\
& \geq \|III_3\|_{1/2}^{1/2} - \|III_2\|_{1/2}^{1/2} - \|III_1\|_{1/2}^{1/2} \\
& \geq c \sum_{i=2}^{r_k-2} \int_{E_i^k} \left| 2^{2i^k} / (\Phi^{1/2}(\alpha_k) V(\alpha_k)) \right|^{1/2} d\mu(x) - 2c \\
& \geq c \sum_{i=2}^{r_k-2} 1 / (V^{1/2}(\alpha_k) \Phi^{1/4}(\alpha_k)) - 2c \\
& \geq cr_k / (V^{1/2}(\alpha_k) \Phi^{1/4}(\alpha_k)) \\
& \geq cV^{1/2}(\alpha_k) / \Phi^{1/4}(\alpha_k) \rightarrow \infty, \text{ as } k \rightarrow \infty.
\end{aligned}$$

Theorem 6 is proved.

**Theorem 7.** a) Let  $0 < p < 1/2$ ,  $f \in H_p(G)$ . Then there exists an absolute constant  $c_p$  depending only on  $p$  such that

$$\|\sigma_n f\|_{H_p(G)} \leq c_p 2^{d(n)(1/p-2)} \|f\|_{H_p(G)}.$$

b) Let  $0 < p < 1/2$  and  $\Phi(n) : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function such that

$$(93) \quad \sup_{k \in \mathbb{N}_+} d(n_k) = \infty, \quad \lim_{k \rightarrow \infty} \frac{2^{d(n_k)(1/p-2)}}{\Phi(n_k)} = \infty.$$

Then there exist a martingale  $f \in H_p(G)$ , such that

$$\sup_{k \in \mathbb{N}_+} \left\| \frac{\sigma_{n_k} f}{\Phi(n_k)} \right\|_{weak-L_p(G)} = \infty.$$

**Proof:** Let  $n \in \mathbb{N}$ . Analogously to (80) it is sufficient to prove that

$$\int_{\bar{I}_M} (2^{d(n)(2-1/p)} |\sigma_n(a)|)^p d\mu \leq c_p < \infty,$$

for every  $p$ -atom  $a$ , where  $I$  denotes support of the atom.

Analogously to Theorem 6 we may assume that  $a$  is  $p$ -atom with support  $I = I_M$ ,  $\mu(I_M) = 2^{-M}$  and  $n > 2^M$ . Since  $\|a\|_\infty \leq 2^{M/p}$  we can conclude that

$$\begin{aligned}
& 2^{d(n)(2-1/p)} |\sigma_n a| \\
& \leq 2^{d(n)(2-1/p)} \|a\|_\infty \int_{I_M} |K_n(x+t)| d\mu(t) \\
& \leq 2^{d(n)(2-1/p)} 2^{M/p} \int_{I_M} |K_n(x+t)| d\mu(t).
\end{aligned}$$

Let  $x \in I_{l+1}(e_k + e_l)$ ,  $0 \leq k, l \leq [n] \leq M$ . Then, by applying Lemma 9 we get that  $K_n(x+t) = 0$ , where  $t \in I_M$  and hence,

$$(94) \quad 2^{d(n)(2-1/p)} |\sigma_n a| = 0.$$

Let  $x \in I_{l+1}(e_k + e_l)$ ,  $[n] \leq k, l \leq M$  or  $k \leq [n] \leq l \leq M$ . Then Lemma 11 follows that

$$(95) \quad \begin{aligned} 2^{d(n)(2-1/p)} |\sigma_n a| &\leq 2^{d(n)(2-1/p)} 2^{M(1/p-2)+k+l} \\ &\leq c_p 2^{[n](1/p-2)+k+l}. \end{aligned}$$

By combining (17), (94) and (95) we can conclude that

$$\begin{aligned} &\int_{I_M} |2^{d(n)(2-1/p)} \sigma_n a(x)|^p d\mu(x) \\ &\leq \left( \sum_{k=0}^{[n]-2} \sum_{l=k+1}^{[n]-1} + \sum_{k=0}^{[n]-1} \sum_{l=[n]}^{M-1} + \sum_{k=[n]}^{M-2} \sum_{l=k+1}^{M-1} \right) \int_{I_{l+1}(e_k+e_l)} |2^{d(n)(2-1/p)} \sigma_n a(x)|^p d\mu(x) \\ &+ \sum_{k=0}^{M-1} \int_{I_M(e_k)} |2^{d(n)(2-1/p)} \sigma_n a(x)|^p d\mu(x) \\ &\leq c_p \sum_{k=[n]}^{M-2} \sum_{l=k+1}^{M-1} \frac{1}{2^l} 2^{[n](2p-1)} 2^{p(k+l)} \\ &+ c_p \sum_{k=0}^{[n]} \sum_{l=[n]+1}^{M-1} \frac{1}{2^l} 2^{[n](2p-1)} 2^{p(k+l)} \\ &+ \frac{c_p 2^{[n](2p-1)}}{2^M} \sum_{k=0}^{[n]} 2^{p(k+M)} < c_p < \infty. \end{aligned}$$

Now, we prove part b) of Theorem 7. According to (93) there exists an increasing sequence of natural numbers  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{n_k : k \in \mathbb{N}_+\}$ , such that  $\alpha_0 \geq 3$  and

$$(96) \quad \sum_{\eta=0}^{\infty} u^{-p}(\alpha_\eta) < c_p < \infty, \quad u(\alpha_k) = 2^{d(\alpha_k)(1/p-2)/2} / \Phi^{1/2}(\alpha_k).$$

Let  $f$  be martingale from Example 1, where

$$\lambda_k = u^{-1}(\alpha_k),$$

If we apply (96) we get that (19) is fulfilled and it follows that  $f \in H_p(G)$ . According to (20) we have that

$$(97) \quad \widehat{f}(j) = \begin{cases} 2^{|\alpha_k|(1/p-1)} / u(\alpha_k), & j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k \in \mathbb{N}_+, \\ 0, & j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

Let  $2^{|\alpha_k|} < j < \alpha_k$ . Then, analogously to (90) and (91), if we apply (97) we get that

$$\begin{aligned}
& \frac{\sigma_{\alpha_k} f}{\Phi(\alpha_k)} \\
&= \frac{\sigma_{2^{|\alpha_k|}} f}{\Phi(\alpha_k) \alpha_k} + \frac{(\alpha_k - 2^{|\alpha_k|}) S_{2^{|\alpha_k|}} f}{\Phi(\alpha_k) \alpha_k} \\
&+ \frac{2^{|\alpha_k|(1/p-1)}}{\Phi(\alpha_k) u(\alpha_k) \alpha_k} \sum_{j=2^{|\alpha_k|}}^{\alpha_k-1} (D_j - D_{2^{|\alpha_k|}}) \\
&= IV_1 + IV_2 + IV_3.
\end{aligned}$$

Let  $\alpha_k \in \mathbb{N}$  and  $E_{[\alpha_k]} := I_{[\alpha_k]+1}(e_{[\alpha_k]-1} + e_{[\alpha_k]})$ . Since  $[\alpha_k - 2^{|\alpha_k|}] = [\alpha_k]$ , analogously to (92), if we apply Lemma 12 for  $IV_3$  we have the following estimation

$$\begin{aligned}
|IV_3| &= \frac{2^{|\alpha_k|(1/p-1)}}{\Phi(\alpha_k) u(\alpha_k) \alpha_k} (\alpha_k - 2^{|\alpha_k|}) \left| K_{\alpha_k - 2^{|\alpha_k|}} \right| \\
&= \frac{2^{|\alpha_k|(1/p-1)}}{\Phi(\alpha_k) u(\alpha_k) \alpha_k} \left| 2^{[\alpha_k]} K_{[\alpha_k]} \right| \\
&\geq \frac{2^{|\alpha_k|(1/p-2)} 2^{2^{[\alpha_k]}-4}}{\Phi(\alpha_k) u(\alpha_k)} \\
&\geq 2^{|\alpha_k|(1/p-2)/2} 2^{2^{[\alpha_k]}-4} / \Phi^{1/2}(\alpha_k).
\end{aligned}$$

Hence,

$$\begin{aligned}
& \|IV_3\|_{weak-L_p(G)}^p \\
&\geq \left( \frac{2^{|\alpha_k|(1/p-2)/2} 2^{2^{[\alpha_k]}-4}}{\Phi^{1/2}(\alpha_k)} \right)^p \mu \left\{ x \in G : |IV_3| \geq \frac{2^{|\alpha_k|(1/p-2)/2} 2^{2^{[\alpha_k]}-4}}{\Phi^{1/2}(\alpha_k)} \right\} \\
&\geq c_p \left( 2^{2^{[\alpha_k]}+|\alpha_k|(1/p-2)/2} / \Phi^{1/2}(\alpha_k) \right)^p \mu(E_{[\alpha_k]}) \\
&\geq c_p \left( 2^{(|\alpha_k|-[ \alpha_k ])(1/p-2)} / \Phi(\alpha_k) \right)^{p/2} \\
&= c_p \left( 2^{d(\alpha_k)(1/p-2)} / \Phi(\alpha_k) \right)^{p/2} \rightarrow \infty, \text{ as } k \rightarrow \infty.
\end{aligned}$$

By combining Corollary 2 and first part of Theorem 7 we find that

$$\|IV_1\|_{weak-L_p(G)} \leq c_p < \infty, \quad \|IV_2\|_{weak-L_p(G)} \leq c_p < \infty.$$

On the other hand, for sufficiently large  $n$  we can conclude that

$$\begin{aligned}
& \|\sigma_{\alpha_k} f\|_{weak-L_p(G)}^p \\
&\geq \|IV_3\|_{weak-L_p(G)}^p - \|IV_2\|_{weak-L_p(G)}^p - \|IV_1\|_{weak-L_p(G)}^p \\
&\geq \frac{1}{2} \|IV_3\|_{weak-L_p(G)}^p \rightarrow \infty, \text{ as } k \rightarrow \infty.
\end{aligned}$$

Theorem 7 is proved.

The proofs of Corollaries 8-10 are similar to the proofs of Corollaries 2-4. So, we leave out the details of proofs and just present these results:

**Corollary 8.** *Let  $p > 0$  and  $f \in H_p(G)$ . Then*

$$\|\sigma_{2^k} f - f\|_{H_p(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

**Corollary 9.** *Let  $p > 0$  and  $f \in H_p(G)$ . Then*

$$\|\sigma_{2^k + 2^{k-1}} f - f\|_{H_p(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

**Corollary 10.** *Let  $0 < p < 1/2$ . Then there exists a martingale  $f \in H_p(G)$ , such that*

$$\|\sigma_{2^k + 1} f - f\|_{\text{weak-}L_p(G)} \not\rightarrow 0, \text{ as } k \rightarrow \infty.$$

*On the other hand, for any  $f \in H_{1/2}(G)$  the following is true:*

$$\|\sigma_{2^k + 1} f - f\|_{H_{1/2}(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

### 3.4. Modulus of continuity and convergence in norm of subsequences of Fejér means with respect to the one-dimensional Walsh-Fourier series on the martingale Hardy spaces.

In this section we apply Theorem 6 and Theorem 7 to find necessary and sufficient conditions for modulus of continuity of martingale  $f \in H_p$ , for which subsequences of Fejér means with respect to the one-dimensional Walsh-Fourier series converge in  $H_p$ -norm.

First, we prove the following result:

**Theorem 8.** *a) Let  $f \in H_{1/2}(G)$ ,  $\sup_{k \in \mathbb{N}_+} V(n_k) = \infty$  and*

$$(98) \quad \omega_{H_p(G)}(1/2^{|n_k|}, f) = o(1/V^2(n_k)), \text{ as } k \rightarrow \infty.$$

*Then*

$$\|\sigma_{n_k} f - f\|_{H_{1/2}(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

*b) Let  $\sup_{k \in \mathbb{N}_+} V(n_k) = \infty$ . Then there exists a martingale  $f \in H_{1/2}(G)$ , such that*

$$(99) \quad \omega_{H_{1/2}(G)}(1/2^{|n_k|}, f) = O(1/V^2(n_k)), \text{ as } k \rightarrow \infty$$

*and*

$$(100) \quad \|\sigma_{n_k} f - f\|_{H_{1/2}(G)} \not\rightarrow 0, \text{ as } k \rightarrow \infty.$$

**Proof:** Let  $f \in H_{1/2}(G)$  and  $2^k < n \leq 2^{k+1}$ . Then

$$\begin{aligned} & \|\sigma_n f - f\|_{H_{1/2}(G)}^{1/2} \\ & \leq \|\sigma_n f - \sigma_n S_{2^k} f\|_{H_{1/2}(G)}^{1/2} + \|\sigma_n S_{2^k} f - S_{2^k} f\|_{H_{1/2}(G)}^{1/2} + \|S_{2^k} f - f\|_{H_{1/2}(G)}^{1/2} \\ & = \|\sigma_n (S_{2^k} f - f)\|_{H_{1/2}(G)}^{1/2} + \|S_{2^k} f - f\|_{H_{1/2}(G)}^{1/2} + \|\sigma_n S_{2^k} f - S_{2^k} f\|_{H_{1/2}(G)}^{1/2} \\ & \leq c(V(n) + 1) \omega_{H_{1/2}(G)}^{1/2}(1/2^k, f) + \|\sigma_n S_{2^k} f - S_{2^k} f\|_{H_{1/2}(G)}^{1/2}. \end{aligned}$$

It is evident that

$$\sigma_n S_{2^k} f - S_{2^k} f = \frac{2^k}{n} (S_{2^k} \sigma_{2^k} f - S_{2^k} f) = \frac{2^k}{n} S_{2^k} (\sigma_{2^k} f - f).$$

Let  $p > 0$ . By combining Corollaries 2 and 8 we can conclude that

$$\begin{aligned} \|\sigma_n S_{2^k} f - S_{2^k} f\|_{H_{1/2}(G)}^{1/2} &\leq \frac{2^{k/2}}{n^{1/2}} \|S_{2^k} (\sigma_{2^k} f - f)\|_{H_{1/2}(G)}^{1/2} \\ &\leq \|\sigma_{2^k} f - f\|_{H_{1/2}(G)}^{1/2} \rightarrow 0, \text{ as } k \rightarrow \infty. \end{aligned}$$

Now, we prove part b) of Theorem 8. Since  $\sup_{k \in \mathbb{N}_+} V(\alpha_k) = \infty$ , then there exists a martingale  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{n_k : k \in \mathbb{N}_+\}$  such that  $V(\alpha_k) \uparrow \infty$ , as  $k \rightarrow \infty$  and

$$(101) \quad V^2(\alpha_k) \leq V(\alpha_{k+1}).$$

Let  $f$  be martingale from Example 1, where

$$\lambda_k = V^{-2}(\alpha_k),$$

If we apply (101) we get that condition (19) is fulfilled and it follows that  $f \in H_p(G)$ . By using (20) we find that

$$(102) \quad \widehat{f}(j) = \begin{cases} 2^{|\alpha_k|}/V^2(\alpha_k), & j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k \in \mathbb{N}_+ \\ 0, & j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

By combining (23) and (101) we can conclude that

$$(103) \quad \begin{aligned} w_{H_{1/2}(G)}(1/2^n, f) &= \|f - S_{2^n} f\|_{H_{1/2}(G)} \\ &\leq \sum_{i=n+1}^{\infty} 1/V^2(\alpha_i) = O(1/V^2(\alpha_n)), \text{ as } n \rightarrow \infty. \end{aligned}$$

Let  $2^{|\alpha_k|} < j \leq \alpha_k$ . By using (22) we get that

$$S_j f = S_{2^{|\alpha_k|}} f + \frac{2^{|\alpha_k|} w_{2^{|\alpha_k|}} D_{j-2^{|\alpha_k|}}}{V^2(\alpha_k)}.$$

Hence,

$$(104) \quad \begin{aligned} &\sigma_{\alpha_k} f - f \\ &= \frac{2^{|\alpha_k|}}{\alpha_k} (\sigma_{2^{|\alpha_k|}} f - f) + \frac{\alpha_k - 2^{|\alpha_k|}}{\alpha_k} (S_{2^{|\alpha_k|}} f - f) \\ &+ \frac{2^{|\alpha_k|} w_{2^{|\alpha_k|}} (\alpha_k - 2^{|\alpha_k|}) K_{\alpha_k - 2^{|\alpha_k|}}}{\alpha_k V^2(\alpha_k)}. \end{aligned}$$

According to (2), (12) and (104) we have that

$$(105) \quad \begin{aligned} &\|\sigma_{\alpha_k} f - f\|_{1/2}^{1/2} \\ &\geq \frac{c}{V(\alpha_k)} \|(\alpha_k - 2^{|\alpha_k|}) K_{\alpha_k - 2^{|\alpha_k|}}\|_{1/2}^{1/2} \\ &- \left(\frac{2^{|\alpha_k|}}{\alpha_k}\right)^{1/2} \|\sigma_{2^{|\alpha_k|}} f - f\|_{1/2}^{1/2} \\ &- \left(\frac{\alpha_k - 2^{|\alpha_k|}}{\alpha_k}\right)^{1/2} \|S_{2^{|\alpha_k|}} f - f\|_{1/2}^{1/2}. \end{aligned}$$

Let

$$\alpha_k = \sum_{i=1}^{r_k} \sum_{k=l_i^k}^{m_i^k} 2^k,$$

where

$$m_1^k \geq l_1^k > l_1^k - 2 \geq m_2^k \geq l_2^k > l_2^k - 2 > \dots > m_s^k \geq l_s^k \geq 0$$

and

$$E_{l_i^k} := I_{l_i^k+1} \left( e_{l_i^k-1} + e_{l_i^k} \right).$$

By using Lemma 12 we get that

$$\begin{aligned} (106) \quad & \int_G \left| (\alpha_k - 2^{|\alpha_k|}) K_{\alpha_k - 2^{|\alpha_k|}}(x) \right|^{1/2} d\mu \\ & \geq \frac{1}{16} \sum_{i=2}^{r_k-2} \int_{E_{l_i^k}} \left| (\alpha_k - 2^{|\alpha_k|}) K_{\alpha_k - 2^{|\alpha_k|}}(x) \right|^{1/2} d\mu(x) \\ & \geq \frac{1}{16} \sum_{i=2}^{r_k-2} \frac{1}{2^{l_i^k}} 2^{l_i^k} \geq cr_k \geq cV(\alpha_k). \end{aligned}$$

By combining estimations (105-106), Corollaries 2 and 8 we get that (100) holds true and Theorem 8 is proved.

**Theorem 9.** a) Let  $0 < p < 1/2$ ,  $f \in H_p(G)$ ,  $\sup_{k \in \mathbb{N}_+} d(n_k) = \infty$  and

$$(107) \quad \omega_{H_p(G)}(1/2^{|n_k|}, f) = o(1/2^{d(n_k)(1/p-2)}), \text{ as } k \rightarrow \infty.$$

Then

$$(108) \quad \|\sigma_{n_k} f - f\|_{H_p(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

b) Let  $\sup_{k \in \mathbb{N}_+} d(n_k) = \infty$ . Then there exists a martingale  $f \in H_p(G)$  ( $0 < p < 1/2$ ), such that

$$(109) \quad \omega_{H_p(G)}(1/2^{|n_k|}, f) = O(1/2^{d(n_k)(1/p-2)}), \text{ as } k \rightarrow \infty$$

and

$$(110) \quad \|\sigma_{n_k} f - f\|_{weak-L_p(G)} \not\rightarrow 0, \text{ as } k \rightarrow \infty.$$

**Proof:** Let  $0 < p < 1/2$ . Then under condition (107) if we repeat steps of the proof of Theorem 8, we immediately get that (108) holds.

Let prove part b) of Theorem 9. Since  $\sup_k d(n_k) = \infty$ , there exists  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{n_k : k \in \mathbb{N}_+\}$  such that  $\sup_{k \in \mathbb{N}_+} d(\alpha_k) = \infty$  and

$$(111) \quad 2^{2d(\alpha_k)(1/p-2)} \leq 2^{d(\alpha_{k+1})(1/p-2)}.$$

Let  $f$  be a martingale from Lemma 1, where

$$\lambda_k = 2^{-(1/p-2)d(\alpha_k)}.$$

If we use (111) we conclude that condition (19) is fulfilled and it follows that  $f \in H_p(G)$ .

According to (20) we get that

$$(112) \quad \widehat{f}(j) = \begin{cases} 2^{(1/p-2)[\alpha_k]}, & j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, \quad k \in \mathbb{N}_+ \\ 0, & j \notin \bigcup_{n=0}^{\infty} \{2^{|\alpha_n|}, \dots, 2^{|\alpha_n|+1} - 1\}. \end{cases}$$

By combining (23) and (111) we have that

$$(113) \quad \begin{aligned} \omega_{H_p(G)}(1/2^{|\alpha_k|}, f) &\leq \sum_{i=k}^{\infty} 1/2^{d(\alpha_i)(1/p-2)} \\ &= O(1/2^{d(\alpha_k)(1/p-2)}) \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Analogously to the proof of previous theorem, if we use also Corollaries 2 and 8, for the sufficiently large  $k$  we can conclude that

$$(114) \quad \begin{aligned} &\|\sigma_{\alpha_k} f - f\|_{weak-L_p(G)}^p \\ &\geq 2^{(1-2p)[\alpha_k]} \|(\alpha_k - 2^{|\alpha_k|}) K_{\alpha_k - 2^{|\alpha_k|}}\|_{weak-L_p(G)}^p \\ &\quad - \left(\frac{2^{|\alpha_k|}}{\alpha_k}\right)^p \|\sigma_{2^{|\alpha_k|}} f - f\|_{weak-L_p(G)}^p \\ &\quad - \left(\frac{\alpha_k - 2^{|\alpha_k|}}{\alpha_k}\right)^p \|S_{2^{|\alpha_k|}} f - f\|_{weak-L_p(G)}^p \\ &\geq 2^{(1-2p)[\alpha_k]-1} \|(\alpha_k - 2^{|\alpha_k|}) K_{\alpha_k - 2^{|\alpha_k|}}\|_{weak-L_p(G)}^p \end{aligned}$$

Let  $x \in E_{[\alpha_k]}$ . Lemma 12 follows that

$$\mu \left( x \in G : (\alpha_k - 2^{|\alpha_k|}) \left| K_{\alpha_k - 2^{|\alpha_k|}} \right| \geq 2^{2[\alpha_k]-4} \right) \geq \mu(E_{[\alpha_k]}) \geq 1/2^{[\alpha_k]-4}$$

and

$$(115) \quad 2^{2p[\alpha_k]-4} \mu \left( x \in G : (\alpha_k - 2^{|\alpha_k|}) \left| K_{\alpha_k - 2^{|\alpha_k|}} \right| \geq 2^{2[\alpha_k]-4} \right) \geq 2^{(2p-1)[\alpha_k]-4}.$$

Hence, by combining (2), (12), (114) and (115) we get that

$$\|\sigma_{n_k} f - f\|_{weak-L_p(G)} \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

The proof of Theorem 9 is complete.

By using Theorem 9 we easily get an important result which was proved in [58]:

**Corollary 11.** *a) Let  $f \in H_{1/2}(G)$  and*

$$\omega_{H_{1/2}(G)}(1/2^k, f) = o\left(\frac{1}{k^2}\right), \quad \text{as } k \rightarrow \infty.$$

*Then*

$$\|\sigma_k f - f\|_{H_{1/2}(G)} \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

*b) There exists a martingale  $f \in H_{1/2}(G)$ , for which*

$$\omega_{H_{1/2}(G)}(1/2^k, f) = O\left(\frac{1}{k^2}\right), \quad \text{as } k \rightarrow \infty$$

*and*

$$\|\sigma_k f - f\|_{1/2} \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

**Corollary 12.** *a) Let  $0 < p < 1/2$ ,  $f \in H_p(G)$  and*

$$\omega_{H_p(G)}(1/2^k, f) = o(1/2^{k(1/p-2)}), \text{ as } k \rightarrow \infty.$$

*Then*

$$\|\sigma_k f - f\|_{H_p(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

*b) Then there exists a martingale  $f \in H_p(G)$  ( $0 < p < 1/2$ ), for which*

$$\omega_{H_p(G)}(1/2^k, f) = O(1/2^{k(1/p-2)}), \text{ as } k \rightarrow \infty$$

*and*

$$\|\sigma_k f - f\|_{weak-L_p(G)} \rightarrow 0, \text{ as } k \rightarrow \infty.$$

### 3.5. Strong convergence of Fejér means with respect to the one-dimensional Walsh-Fourier series on the martingale Hardy spaces.

In this section we consider strong convergence results of Fejér means with respect to the one-dimensional Walsh-Fourier series in the martingale Hardy spaces, when  $0 < p \leq 1/2$  (for details see [57]).

The following is true:

**Theorem 10.** *a) Let  $0 < p \leq 1/2$  and  $f \in H_p(G)$ . Then there exists a constant  $c_p$ , depending only on  $p$ , such that*

$$\frac{1}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{\|\sigma_m f\|_{H_p(G)}^p}{m^{2-2p}} \leq c_p \|f\|_{H_p(G)}^p.$$

*b) Let  $0 < p < 1/2$ ,  $\Phi : \mathbb{N}_+ \rightarrow [1, \infty)$  be non-decreasing function, such that  $\Phi(n) \uparrow \infty$  and*

$$\overline{\lim}_{k \rightarrow \infty} \frac{k^{2-2p}}{\Phi(k)} = \infty.$$

*Then there exists a martingale  $f \in H_p(G)$ , such that*

$$\sum_{m=1}^{\infty} \frac{\|\sigma_m f\|_{weak-L_p(G)}^p}{\Phi(m)} = \infty.$$

**Proof:** Suppose that

$$\frac{1}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{\|\sigma_m f\|_p^p}{m^{2-2p}} \leq c_p \|f\|_{H_p(G)}^p.$$

By combining (7), (15) and Lemma 13 we can conclude that

$$\begin{aligned} (116) \quad & \frac{1}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{\|\sigma_m f\|_{H_p(G)}^p}{m^{2-2p}} \\ & \leq \frac{1}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{\|\sigma_m f\|_p^p}{m^{2-2p}} + \|\tilde{\sigma}_{\#}^* f\|_{H_p(G)} + \|\tilde{S}_{\#}^* f\|_{H_p(G)} \\ & \leq c_p \|f\|_{H_p(G)}^p. \end{aligned}$$

According to Lemma 5 and (116) Theorem 10 will be proved if we show that

$$\frac{1}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{\|\sigma_m a\|_p^p}{m^{2-2p}} \leq c < \infty, \quad m = 2, 3, \dots$$

for any  $p$ -atom  $a$ . We may assume that  $a$  is  $p$ -atom, with support  $I$ ,  $\mu(I) = 2^{-M}$  and  $I = I_M$ . It is evident that  $\sigma_n(a) = 0$ , when  $n \leq 2^M$ . Therefore, we may assume that  $n > 2^M$ .

Let  $x \in I_M$ . Since  $\sigma_n$  is bounded from  $L_\infty(G)$  to  $L_\infty(G)$  (The boundedness follows fact that Fejér kernels are uniformly bounded in the space  $L_1(G)$ , which is proved in Lemma 8) and  $\|a\|_\infty \leq 2^{M/p}$  we can conclude that

$$\begin{aligned} \int_{I_M} |\sigma_m a(x)|^p d\mu(x) &\leq \|\sigma_m a\|_\infty^p / 2^M \\ &\leq \|a\|_\infty^p / 2^M \leq c < \infty, \quad 0 < p \leq 1/2. \end{aligned}$$

Let  $0 < p \leq 1/2$ . Then

$$\begin{aligned} \frac{1}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{\int_{I_M} |\sigma_m a(x)|^p d\mu(x)}{m^{2-2p}} \\ \leq \frac{c}{\log^{[1/2+p]} n} \sum_{m=1}^n \frac{1}{m^{2-2p}} \leq c < \infty. \end{aligned}$$

It is evident that

$$\begin{aligned} |\sigma_m a(x)| &\leq \int_{I_M} |a(t)| |K_m(x+t)| d\mu(t) \\ &\leq 2^{M/p} \int_{I_M} |K_m(x+t)| d\mu(t). \end{aligned}$$

Lemma 9 follows that

$$(117) \quad |\sigma_m a(x)| \leq \frac{c 2^{k+l} 2^{M(1/p-1)}}{m}, \quad x \in I_{l+1}(e_k + e_l), \quad 0 \leq k < l < M$$

and

$$(118) \quad |\sigma_m a(x)| \leq c 2^{M(1/p-1)} 2^k, \quad x \in I_M(e_k), \quad 0 \leq k < M.$$

If we use identity (17) and (117-118) we get that

$$\begin{aligned}
(119) \quad & \int_{I_M} |\sigma_m a(x)|^p d\mu(x) \\
&= \sum_{k=0}^{M-2} \sum_{l=k+1}^{M-1} \int_{I_{l+1}(e_k+e_l)} |\sigma_m a(x)|^p d\mu(x) \\
&\quad + \sum_{k=0}^{M-1} \int_{I_M(e_k)} |\sigma_m a(x)|^p d\mu(x) \\
&\leq c \sum_{k=0}^{M-2} \sum_{l=k+1}^{M-1} \frac{1}{2^l} \frac{2^{p(k+l)} 2^{M(1-p)}}{m^p} + c \sum_{k=0}^{M-1} \frac{1}{2^M} 2^{M(1-p)} 2^{pk} \\
&\leq \frac{c 2^{M(1-p)}}{m^p} \sum_{k=0}^{M-2} \sum_{l=k+1}^{M-1} \frac{2^{p(k+l)}}{2^l} + c \sum_{k=0}^{M-1} \frac{2^{pk}}{2^{pM}} \\
&\leq \frac{c 2^{M(1-p)} M^{[1/2+p]}}{m^p} + c.
\end{aligned}$$

Hence,

$$\begin{aligned}
& \frac{1}{\log^{[1/2+p]} n} \sum_{m=2^{M+1}}^n \frac{\int_{I_M} |\sigma_m a(x)|^p d\mu(x)}{m^{2-2p}} \\
&\leq \frac{1}{\log^{[1/2+p]} n} \left( \sum_{m=2^{M+1}}^n \frac{c 2^{M(1-p)} M^{[1/2+p]}}{m^{2-p}} + \sum_{m=2^{M+1}}^n \frac{c}{m^{2-2p}} \right) < c < \infty.
\end{aligned}$$

The proof of part a) of theorem 10 is complete.

Now, we prove part b) of Theorem 10. Let  $\Phi(n)$  non-decreasing function satisfying the condition

$$(120) \quad \lim_{k \rightarrow \infty} \frac{2^{(|n_k|+1)(2-2p)}}{\Phi(2^{|n_k|+1})} = \infty.$$

According to (120), there exists an increasing sequence  $\{\alpha_k : k \in \mathbb{N}_+\} \subset \{n_k : k \in \mathbb{N}_+\}$  such that

$$(121) \quad |\alpha_k| \geq 2, \quad \text{where } k \in \mathbb{N}_+$$

and

$$(122) \quad \sum_{\eta=0}^{\infty} \frac{\Phi^{1/2}(2^{|\alpha_\eta|+1})}{2^{|\alpha_\eta|(1-p)}} = 2^{1-p} \sum_{\eta=0}^{\infty} \frac{\Phi^{1/2}(2^{|\alpha_\eta|+1})}{2^{(|\alpha_\eta|+1)(1-p)}} < c < \infty.$$

Let  $f = (f_n, n \in \mathbb{N}_+) \in H_p(G)$  be a martingale from the Example 1, where

$$\lambda_k = \frac{\Phi^{1/2p}(2^{|\alpha_k|+1})}{2^{(|\alpha_k|)(1/p-1)}}$$

By combining (19) and (122) we get that  $f \in H_p(G)$ . According to (20) we have that

$$(123) \quad \widehat{f}(j) = \begin{cases} \Phi^{1/2p} (2^{|\alpha_k|+1}) & \text{if } j \in \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}, k \in \mathbb{N}_+, \\ 0, & \text{if } j \notin \bigcup_{k=0}^{\infty} \{2^{|\alpha_k|}, \dots, 2^{|\alpha_k|+1} - 1\}. \end{cases}$$

Let  $2^{|\alpha_k|} < n < 2^{|\alpha_k|+1}$ . Then

$$(124) \quad \sigma_n f = \frac{1}{n} \sum_{j=1}^{2^{|\alpha_k|}} S_j f + \frac{1}{n} \sum_{j=2^{|\alpha_k|+1}}^n S_j f = III + IV.$$

It is evident that

$$(125) \quad S_j f = 0, \quad \text{if } 0 \leq j \leq 2^{|\alpha_1|}$$

Let  $2^{|\alpha_s|} < j \leq 2^{|\alpha_s|+1}$ , where  $s = 1, 2, \dots, k$ . If we apply (22) we get that

$$(126) \quad S_j f = \sum_{\eta=0}^{s-1} \Phi^{1/2p} (2^{|\alpha_\eta|+1}) (D_{2^{|\alpha_\eta|+1}} - D_{2^{|\alpha_\eta|}}) + \Phi^{1/2p} (2^{|\alpha_s|+1}) w_{2^{|\alpha_s|}} D_{j-2^{|\alpha_s|}}.$$

Let  $2^{|\alpha_s|+1} \leq j \leq 2^{|\alpha_{s+1}|}$ ,  $s = 0, 1, \dots, k-1$ . Then if we use (21) we can conclude that

$$(127) \quad S_j f = \sum_{\eta=0}^s \Phi^{1/2p} (2^{|\alpha_\eta|+1}) (D_{2^{|\alpha_\eta|+1}} - D_{2^{|\alpha_\eta|}}).$$

Let  $x \in I_2(e_0 + e_1)$ . Since (see Lemmas 2 and 9)

$$(128) \quad D_{2^n}(x) = K_{2^n}(x) = 0, \quad \text{where } n \geq 2$$

by combining (121) and (125-128) we get that

$$(129) \quad \begin{aligned} III &= \frac{1}{n} \sum_{\eta=0}^{k-1} \Phi^{1/2p} (2^{|\alpha_\eta|+1}) \sum_{v=2^{|\alpha_\eta|+1}}^{2^{|\alpha_\eta|+1}} D_v(x) \\ &= \frac{1}{n} \sum_{\eta=0}^{k-1} \Phi^{1/2p} (2^{|\alpha_\eta|+1}) (2^{|\alpha_\eta|+1} K_{2^{|\alpha_\eta|+1}}(x) - 2^{|\alpha_\eta|} K_{2^{|\alpha_\eta|}}(x)) = 0. \end{aligned}$$

If we use (126) when  $s = k$  for  $IV$  we can write that

$$(130) \quad \begin{aligned} IV &= \frac{n - 2^{|\alpha_k|}}{n} \sum_{\eta=0}^{k-1} \Phi^{1/2p} (2^{|\alpha_\eta|+1}) (D_{2^{|\alpha_\eta|+1}} - D_{2^{|\alpha_\eta|}}) \\ &+ \frac{\Phi^{1/2p} (2^{|\alpha_k|+1})}{n} \sum_{j=2^{|\alpha_k|+1}}^n w_{2^{|\alpha_k|}} D_{j-2^{|\alpha_k|}} = IV_1 + IV_2. \end{aligned}$$

By combining (121) and (128) we can conclude that

$$(131) \quad IV_1 = 0, \quad \text{where } x \in I_2(e_0 + e_1).$$

Let  $\alpha_k \in \mathbb{A}_{0,2}$ ,  $2^{|\alpha_k|} < n < 2^{|\alpha_k|+1}$  and  $x \in I_2(e_0 + e_1)$ . Since  $n - 2^{|\alpha_k|} \in \mathbb{A}_{0,2}$ , Lemmas 1 and 8 and (128) follows that

$$\begin{aligned} |IV_2| &= \frac{\Phi^{1/2p}(2^{|\alpha_k|+1})}{n} \left| \sum_{j=1}^{n-2^{|\alpha_k|}} D_j(x) \right| \\ &= \frac{\Phi^{1/2p}(2^{|\alpha_k|+1})}{n} \left| (n - 2^{|\alpha_k|}) K_{n-2^{|\alpha_k|}}(x) \right| \geq \frac{\Phi^{1/2p}(2^{|\alpha_k|+1})}{2^{|\alpha_k|+1}}. \end{aligned}$$

Let  $0 < p < 1/2$  and  $n \in \mathbb{A}_{0,2}$ . By combining (124-132) we get that

$$\begin{aligned} (132) \quad & \|\sigma_n f\|_{weak-L_p(G)}^p \\ & \geq \frac{c_p \Phi^{1/2}(2^{|\alpha_k|+1})}{2^{p(|\alpha_k|+1)}} \mu \left\{ x \in I_2(e_0 + e_1) : |\sigma_n f| \geq \frac{c_p \Phi^{1/2p}(2^{|\alpha_k|+1})}{2^{|\alpha_k|+1}} \right\} \\ & \geq \frac{c_p \Phi^{1/2}(2^{|\alpha_k|+1})}{2^{p(|\alpha_k|+1)}} \mu \{I_2(e_0 + e_1)\} \geq \frac{c_p \Phi^{1/2}(2^{|\alpha_k|+1})}{2^{p(|\alpha_k|+1)}}. \end{aligned}$$

Hence,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\|\sigma_n f\|_{weak-L_p(G)}^p}{\Phi(n)} &\geq \sum_{\{n \in \mathbb{A}_{0,2} : 2^{|\alpha_k|} < n < 2^{|\alpha_k|+1}\}} \frac{\|\sigma_n f\|_{weak-L_p(G)}^p}{\Phi(n)} \\ &\geq \frac{1}{\Phi^{1/2}(2^{|\alpha_k|+1})} \sum_{\{n \in \mathbb{A}_{0,2} : 2^{|\alpha_k|} < n < 2^{|\alpha_k|+1}\}} \frac{1}{2^{p(|\alpha_k|+1)}} \\ &\geq \frac{c_p 2^{(1-p)(|\alpha_k|+1)}}{\Phi^{1/2}(2^{|\alpha_k|+1})} \rightarrow \infty, \quad \text{as } k \rightarrow \infty. \end{aligned}$$

The proof of Theorem 10 is complete.

**Theorem 11.** *Let  $f \in H_{1/2}(G)$ . Then*

$$\sup_{n \in \mathbb{N}_+} \sup_{\|f\|_{H_p} \leq 1} \frac{1}{n} \sum_{m=1}^n \|\sigma_m f\|_{1/2}^{1/2} = \infty.$$

**Proof:** Let  $0 < p \leq 1$  and

$$f_k(x) := 2^k (D_{2^{k+1}}(x) - D_{2^k}(x))$$

Since

$$\text{supp}(f_k) = I_k, \quad \int_{I_k} a_k d\mu = 0$$

and

$$\|f_k\|_{\infty} \leq 2^{2k} = (\text{supp } f_k)^{-2},$$

we conclude that  $f_k$  is 1/2-atom, for every  $k \in \mathbb{N}$

Moreover, if we use orthogonality of Walsh functions we get that

$$S_{2^n}(f_k, x)$$

$$= \begin{cases} 0, & n = 0, \dots, k, \\ (D_{2^{k+1}}(x) - D_{2^k}(x)), & n \geq k + 1, \end{cases}$$

and

$$\begin{aligned} & \sup_{n \in \mathbb{N}} |S_{2^n}(f_k, x)| \\ &= |(D_{2^{k+1}}(x) - D_{2^k}(x))|, \end{aligned}$$

where  $x \in G$ .

By combining first equality of Lemma 1 and Lemma 6 we obtain that

$$\begin{aligned} \|a_k\|_{H_p(G)} &= 2^k \left\| \sup_{n \in \mathbb{N}} |S_{2^n}(D_{2^{k+1}}(x) - D_{2^k}(x))| \right\|_{1/2} \\ &= 2^k \|(D_{2^{k+1}}(x) - D_{2^k}(x))\|_{1/2} = 2^k \|D_{2^k}(x)\|_{1/2} \leq 2^k \cdot 2^{-k} \leq 1. \end{aligned}$$

It is easy to show that

$$(133) \quad \widehat{f}_m(i) = \begin{cases} 2^m, & \text{if } i = 2^m, \dots, 2^{m+1} - 1, \\ 0, & \text{otherwise} \end{cases}$$

and

$$(134) \quad S_i f_m = \begin{cases} 2^m (D_i - D_{2^m}), & \text{if } i = 2^m + 1, \dots, 2^{m+1} - 1, \\ f_m, & \text{if } i \geq 2^{m+1}, \\ 0, & \text{otherwise.} \end{cases}$$

Let  $0 < n < 2^m$ . By using first equality of Lemma 1 we have that

$$\begin{aligned} |\sigma_{n+2^m} f_m| &= \frac{1}{n + 2^m} \left| \sum_{j=2^{m+1}}^{n+2^m} S_j f_m \right| \\ &= \frac{1}{n + 2^m} \left| 2^m \sum_{j=2^{m+1}}^{n+2^m} (D_j - D_{2^m}) \right| \\ &= \frac{1}{n + 2^m} \left| 2^m \sum_{j=1}^n (D_{j+2^m} - D_{2^m}) \right| \\ &= \frac{1}{n + 2^m} \left| 2^m \sum_{j=1}^n D_j \right| = \frac{2^m}{n + 2^m} n |K_n|. \end{aligned}$$

Let

$$n = \sum_{i=1}^s \sum_{k=l_i}^{m_i} 2^k,$$

where

$$0 \leq l_1 \leq m_1 \leq l_2 - 2 < l_2 \leq m_2 \leq \dots \leq l_s - 2 < l_s \leq m_s.$$

By applying Lemma 12 and (135) we find that

$$|\sigma_{n+2^m} f_m(x)| \geq c 2^{2l_i}, \quad \text{where } x \in I_{l_i+1}(e_{l_i-1} + e_{l_i}).$$

Hence,

$$\begin{aligned}
& \int_G |\sigma_{n+2^m} f_m(x)|^{1/2} d\mu(x) \\
& \geq \sum_{i=0}^s \int_{I_{i+1}(e_{i_i-1}+e_{i_i})} |\sigma_{n+2^m} f_m(x)|^{1/2} d\mu(x) \\
& \geq c \sum_{i=0}^s \frac{1}{2^{l_i}} 2^{l_i} \geq cs \geq cV(n).
\end{aligned}$$

According to the second estimation of Lemma 3 we can conclude that

$$\begin{aligned}
\sup_{n \in \mathbb{N}_+} \sup_{\|f\|_{H_p} \leq 1} \frac{1}{n} \sum_{k=1}^n \|\sigma_k f\|_{1/2}^{1/2} & \geq \frac{1}{2^{m+1}} \sum_{k=2^{m+1}}^{2^{m+1}-1} \|\sigma_k f_m\|_{1/2}^{1/2} \\
& \geq \frac{c}{2^{m+1}} \sum_{k=2^{m+1}}^{2^{m+1}-1} V(k-2^m) \\
& \geq \frac{c}{2^{m+1}} \sum_{k=1}^{2^m-1} V(k) \geq c \log m \rightarrow \infty, \text{ as } m \rightarrow \infty.
\end{aligned}$$

The proof is complete.

#### REFERENCES

- [1] *G. N. Agaev, N. Ya. Vilenkin, G. M. Dzhafarly and A. I. Rubinshtein*, Multiplicative systems of functions and harmonic analysis on zero-dimensional groups, Baku, Ehim, 1981 (in Russian).
- [2] *N. K. Bary*, Trigonometric series, Gos. Izd. Fiz. Mat. Lit. Moscow 1961.
- [3] *L. Baramidze, L. E. Persson, G. Tephnadze, P. Wall*, Sharp  $H_p - L_p$  type inequalities of weighted maximal operators of Vilenkin-Nörlund means and its applications, *J. Inequal. Appl.*, 2016, DOI: 10.1186/s13660-016-1182-1.
- [4] *I. Blahota, G. Tephnadze*, A note on maximal operators of Vilenkin-Nörlund means, *Acta Math. Acad. Paed. Nyíreg.*, 32 (2016), 203-213.
- [5] *I. Blahota, G. Tephnadze, R. Toledo*, Strong convergence theorem of  $(C, \alpha)$ -means with respect to the Walsh system, *Tohoku Math. J.*, 67, 4 (2015), 573-584.
- [6] *I. Blahota, L. E. Persson, G. Tephnadze*, On the Nörlund means of Vilenkin-Fourier series, *Czech. Math J.*, 65, 4 (2015), 983-1002.
- [7] *I. Blahota, G. Tephnadze*, On the  $(C, \alpha)$ -means with respect to the Walsh system, *Anal. Math.*, 40 (2014), 161-174.
- [8] *I. Blahota, G. Tephnadze*, Strong convergence theorem for Vilenkin-Fejér means, *Publ. Math. Debrecen*, 85 (1-2) (2014), 181-196.
- [9] *I. Blahota, G. Gät, U. Goginava*, Maximal operators of Fejér means of double Vilenkin-Fourier series, *Colloq. Math.* 107 (2007), no. 2, 287-296.
- [10] *I. Blahota, G. Gät, U. Goginava*, Maximal operators of Fejér means of Vilenkin-Fourier series, *J. Inequal. Pure Appl. Math.* 7 (2006), no. 4, Article 149, 7 pp. (electronic).
- [11] *I. Blahota, K. Nagy, L. E. Persson, G. Tephnadze*, A sharp boundedness result concerning some maximal operators of partial sums with respect to Vilenkin systems, *Georgian Math. J.*, 26, 3 (2019), 351-360.
- [12] *I. Blahota, K. Nagy, G. Tephnadze*, Approximation by Marcinkiewicz  $\Theta$ -means of double Walsh-Fourier series, *Math. Inequal. Appl.*, 22, 3 (2019) 837-853.
- [13] *I. Blahota, L. E. Persson, G. Tephnadze*, Two-sided estimates of the Lebesgue constants with respect to Vilenkin systems and applications, *Glasgow Math. J.*, 60 (2018) 17-34.

- [14] *R. Coifman and G. Weiss*, Extensions of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.* 83 (1977), no. 4, 569-645.
- [15] *N. I. Fine*, On Walsh function, *Trans. Amer. Math. Soc.* 65 (1949), 372-414.
- [16] *N. J. Fujii*, A maximal inequality for  $H^1$ -functions on a generalized Walsh-Paley group, *Proc. Amer. Math. Soc.* 77 (1979), no. 1, 111-116.
- [17] *G. Gát*, Cesàro means of integrable functions with respect to unbounded Vilenkin systems, *J. Approx. Theory* 124 (2003), no. 1, 25-43.
- [18] *G. Gát*, Investigations of certain operators with respect to the Vilenkin system, *Acta Math. Hungar.* 61 (1993), no. 1-2, 131-149.
- [19] *G. Gát, U. Goginava*, Uniform and  $L$ -convergence of logarithmic means of Walsh-Fourier series, *Acta Math. Sin.* 22 (2006), no. 2, 497-506.
- [20] *U. Goginava*, The maximal operator of the  $(C, \alpha)$  means of the Walsh-Fourier series, *Ann. Univ. Sci. Budapest. Sect. Comput.* 26 (2006), 127-135.
- [21] *U. Goginava*, Maximal operators of Fejér means of double Walsh-Fourier series, *Acta Math. Hungar.* 115 (2007), no. 4, 333-340.
- [22] *U. Goginava*, Maximal operators of Fejér-Walsh means, *Acta Sci. Math. (Szeged)* 74 (2008), no. 3-4, 615-624.
- [23] *U. Goginava*, The maximal operator of the Fejér means of the character system of the  $p$ -series field in the Kaczmarr rearrangement, *Publ. Math. Debrecen* 71 (2007), no. 1-2, 43-55.
- [24] *U. Goginava*, The maximal operator of Marcinkiewicz-Fejér means of the  $d$ -dimensional Walsh-Fourier series. *East J. Approx.* 12 (3) (2006), 295-302.
- [25] *U. Goginava*, A note on the Walsh-Fejér means, *Anal. Theory Appl.* 26, 4 (2010), 320-325.
- [26] *U. Goginava, K. Nagy*, On the maximal operator of Walsh-Kaczmarr-Fejér means, *Czechoslovak Math. J.* 61(136) (2011), no. 3, 673-686.
- [27] *B. I. Golubov, A. V. Efimov, V. A. Skvortsov*, Walsh series and transforms, (Russian) Nauka, Moscow, 1987, English transl: Mathematics and its Applications, 64. Kluwer Academic Publishers Group, Dordrecht, 1991.
- [28] *J. A. Gosselin*, Almost everywhere convergence of Vilenkin-Fourier series, *Trans. Amer. Math. Soc.*, 185 (1973), 345-370.
- [29] *D. Lukkassen, L.E. Persson, G. Tephnadze, G. Tutberidze*, Some inequalities related to strong convergence of Riesz logarithmic means of Vilenkin-Fourier series, *J. Inequal. Appl.*, 2020, DOI: <https://doi.org/10.1186/s13660-020-02342-8>.
- [30] *N. Memic, L. E. Persson, G. Tephnadze*, A note on the maximal operators of Vilenkin-Nörlund means with non-increasing coefficients, *Stud. Sci. Math. Hung.*, 53, 4, (2016) 545-556.
- [31] *N. Memic, I. Simon, G. Tephnadze*, Strong convergence of two-dimensional Vilenkin-Fourier series, *Math. Nachr.*, 289, 4 (2016) 485-500.
- [32] *K. Nagy, G. Tephnadze*, Strong convergence theorem for Walsh-Marcinkiewicz means, *Math. Inequal. Appl.*, 19, 1 (2016), 185-195.
- [33] *K. Nagy, G. Tephnadze*, Approximation by Walsh-Marcinkiewicz means on the Hardy space  $H_{2/3}$ , *Kyoto J. Math.*, 54 (3), (2014), 641-652.
- [34] *K. Nagy, G. Tephnadze*, On the Walsh-Marcinkiewicz means on the Hardy space, *Cent. Eur. J. Math.*, 12 (8), (2014), 1214-1228.
- [35] *K. Nagy, G. Tephnadze*, Approximation by Walsh-Kaczmarr-Marcinkiewicz means on the Hardy space  $H_{2/3}$ , *Bulletin of TICMI*, 18, 1 (2014), 110-121.
- [36] *J. Neveu*, Discrete-parameter martingales, North-Holland Mathematical Library, Vol. 10. North-Holland Publishing Co., Amsterdam-Oxford; American Elsevier Publishing Co., Inc., New York, 1975.
- [37] *L. E. Persson, G. Tephnadze, G. Tutberidze*, On the boundedness of subsequences of Vilenkin-Fejér means on the martingale Hardy spaces, operators and matrices, 14, 1 (2020), 283-294.
- [38] *L. E. Persson, G. Tephnadze, P. Wall*, On the Nörlund logarithmic means with respect to Vilenkin system in the martingale Hardy space  $H_1$ , *Acta math. Hungar.*, 154, 2 (2018) 289-301.
- [39] *L. E. Persson, G. Tephnadze, P. Wall*, On an approximation of two-dimensional Walsh-Fourier series in the martingale Hardy spaces, *Ann. Funct. Anal.*, 9, 1 (2018), 137-150.
- [40] *L. E. Persson, G. Tephnadze, P. Wall*, Maximal operators of Vilenkin-Nörlund means, *J. Fourier Anal. Appl.*, 21, (2015), no. 1, 76-94.

- [41] *L. E. Persson, G. Tephnadze, P. Wall*, Some new  $(H_p, L_p)$  type inequalities of maximal operators of Vilenkin-Nörlund means with non-decreasing coefficients, *J. Math. Inequal*, 9, 4 (2015), 1055-1069.
- [42] *L. E. Persson, G. Tephnadze*, A note on Vilenkin-Fejér means on the martingale Hardy space  $H_p$ , *Bulletin of TICMI*, Vol. 18, (2014), no. 1, 55-64.
- [43] *L. E. Persson, G. Tephnadze*, A sharp boundedness result concerning some maximal operators of Vilenkin-Fejér means, *Mediterr. J. Math.*, 13, 4 (2016) 1841-1853.
- [44] *F. Schipp*, Certain rearrangements of series in the Walsh system, 18 (1975), no. 2, 193–201.
- [45] *F. Schipp, W. R. Wade, P. Simon, J. Pál*, *Walsh series. An introduction to dyadic harmonic analysis*, Adam Hilger, Ltd., Bristol, 1990.
- [46] *P. Simon*, Cesaro summability with respect to two-parameter Walsh systems, *Monatsh. Math.* 131 (2000), no. 4, 321–334.
- [47] *P. Simon*, Strong convergence of certain means with respect to the Walsh-Fourier series, *Acta Math. Hungar.* 49 (1987), no. 3-4, 425-431.
- [48] *P. Simon*, A note on the Sunouchi operator with respect to Vilenkin system, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* 43 (2000), 101-116.
- [49] *P. Simon*, Strong convergence theorem for Vilenkin-Fourier series, *J. Math. Anal. Appl.* 245 (2000), no. 1, 52–68.
- [50] *P. Simon, F. Weisz*, Weak inequalities for Cesàro and Riesz summability of Walsh-Fourier series, *J. Approx. Theory* 151 (2008), no. 1, 1-19.
- [51] *B. Smith*, A strong convergence theorem for  $H_1(T)$  Banach spaces, harmonic analysis, and probability theory (Storrs, Conn., 1980/1981), 169-173.
- [52] *G. Tephnadze*, Fejér means of Vilenkin-Fourier series, *Stud. Sci. Math. Hung.*, 49, 1 (2012) 79-90.
- [53] *G. Tephnadze*, On the maximal operators of Vilenkin-Fejér means, *Turkish J. Math.* 37 (2013), no. 2, 308-318.
- [54] *G. Tephnadze*, On the maximal operators of Vilenkin-Fejér means on Hardy spaces, *Math. Inequal. Appl.* 16 (2013), no. 1, 301–312.
- [55] *G. Tephnadze*, A note on the Fourier coefficients and partial sums of Vilenkin-Fourier series, *Acta Math. Acad. Paedagog. Nyházi.*, 28 (2012), no. 2, 167-176.
- [56] *G. Tephnadze*, On the maximal operators of Walsh-Kaczmarz-Fejér means, *Period. Math. Hung.*, 67, 1 (2013), 33-45.
- [57] *G. Tephnadze*, Strong convergence theorems for Walsh-Fejér means, *Acta Math. Hungar.* 142 (2014), no. 1, 244-259.
- [58] *G. Tephnadze*, A note on the norm convergence by Vilenkin-Fejér means, *Georgian Math. J.* 21 (2014), no. 4, 511-517.
- [59] *G. Tephnadze*, On the partial sums of Vilenkin-Fourier series, *J. Contemp. Math. Anal.* 49 (2014), no. 1, 23-32.
- [60] *G. Tephnadze*, On the Vilenkin-Fourier coefficients, *Georgian Math. J.* 20 (2013), no. 1, 169-177.
- [61] *G. Tephnadze*, On the partial sums of Walsh-Fourier series, *Colloquium Mathematicum*, 141, 2 (2015), 227-242.
- [62] *G. Tephnadze*, On the convergence of Fejér means of Walsh-Fourier series in the space  $H_p$ , *J. Contemp. Math. Anal.*, 51, 2 (2016), 51-63.
- [63] *G. Tephnadze*, A note on the strong convergence of two-dimensional Walsh-Fourier series, *Proceedings of A. Razmadze Mathematical Institute*, 162, (2013), 93-97.
- [64] *G. Tephnadze*, Strong convergence of two-dimensional Walsh-Fourier series, *Ukr. Math. J.*, 65, (6), (2013), 822-834.
- [65] *G. Tephnadze*, The maximal operators of logarithmic means of one-dimensional Vilenkin-Fourier series, *Acta Math. Acad. Paed. Nyíreg.*, 27 (2011), 245-256.
- [66] *G. Tephnadze*, On the maximal operators of Riesz logarithmic means of Vilenkin-Fourier series, *Stud. Sci. Math. Hung.*, 51, 1 (2014), 105-120.
- [67] *G. Tephnadze*, Approximation by Walsh-Kaczmarz-Fejér means on the Hardy space, *Acta Math. Sci.*, 34, 5 (2014), 1593-1602.
- [68] *G. Tephnadze*, A note on strong summability of two-dimensional Walsh-Fourier series, *Georgian Math. J.*, (2020), <https://doi.org/10.1515/gmj-2020-2060>.
- [69] *G. Tephnadze*, On the maximal operators of Kaczmarz-Nörlund means, *Acta Math. Acad. Paed. Nyíreg.*, 31 (2015), 259-271.

- [70] *G. Tephnadze*, On the convergence of partial sums with respect to Vilenkin system on the martingale Hardy spaces, *J. Contemp. Math. Anal.*, *J. Contemp. Math. Anal.*, 53, 5, (2018) 294-306.
- [71] *G. Tephnadze*, Convergence and Strong Summability of the Two-dimensional Vilenkin-Fourier Series, *Nonlinear Studies*, 26, 4, (2019) 973-989.
- [72] *G. Tephnadze*, Martingale Hardy spaces and summability of the one dimensional Vilenkin-Fourier series, PhD thesis, Department of Engineering Sciences and Mathematics, Luleå University of Technology, Oct. 2015 (ISSN 1402-1544).
- [73] *G. Tephnadze, G. Tutberidze*, A note on the maximal operators of the Nörlund logarithmic means of Vilenkin-Fourier series, *Proc. Razmadze Math. Inst.*, 174, 1 (2020), 107-112.
- [74] *C. Watari*, Best approximation by Walsh polynomials, *Tôhoku Math. J.* 15 (1963), no. 2, 1-5.
- [75] *C. Watari*, On generalized Walsh-Fourier series, *Tôhoku Math. J.*, (2) 10 (1958), 211-241.
- [76] *F. Weisz*, Martingale Hardy spaces and their applications in Fourier analysis, *Lecture Notes in Mathematics*, 1568. Springer-Verlag, Berlin, 1994.
- [77] *F. Weisz*, Hardy spaces and Cesàro means of two-dimensional Fourier series, *Approximation theory and function series (Budapest, 1995)*, 353-367, *Bolyai Soc. Math. Stud.*, 5, János Bolyai Math. Soc., Budapest, 1996.
- [78] *F. Weisz*, Cesàro summability of one-and two-dimensional Walsh-Fourier series, *Anal. Math.* 22 (1996), no. 3, 229-242.
- [79] *F. Weisz*, Weak type inequalities for the Walsh and bounded Ciesielski systems, *Anal. Math.* 30 (2004), no. 2, 147-160.
- [80] *F. Weisz*, Summability of multi-dimensional Fourier series and Hardy spaces, *Mathematics and its Applications*, 541. Kluwer Academic Publishers, Dordrecht, 2002.
- [81] *S.H. Yano*, Cesàro summability of Walsh-Fourier series, *Tôhoku Math. J.* 9 (1957), no. 2, 267-272.
- [82] *W. S. Young*, Mean convergence of generalized Walsh-Fourier series, *Trans. Amer. Math. Soc.*, 218 (1976), 311-320.
- [83] *A. Zygmund*, *Trigonometric series*. 2nd ed. Vols. I, II, Cambridge University Press, New York 1959.

G. TEPHNADZE, THE UNIVERSITY OF GEORGIA, SCHOOL OF SCIENCE AND TECHNOLOGY, 77A MERAB KOSTAVA ST, TBILISI 0128, GEORGIA.

*E-mail address:* `g.tephnadze@ug.edu.ge`