

Quantum Zeno dynamics induced atomic entanglement in a hybrid atom-cavity-fiber system

Jiaojiao Chen,¹ Baolong Fang,² and Wei Xiong^{2,*}

¹*Hefei Preschool Education College, Hefei 230013, China*

²*Institute of Advanced Manufacturing Engineering, Hefei University, Anhui 230022, China*

(Dated: January 13, 2022)

Quantum entanglement is important quantum resources in quantum information science. Here we propose an approach to preparing atomic quantum entanglement in a hybrid atom-cavity-fiber system. Using quantum Zeno dynamics method, the system evolution states are always split into a series of Zeno invariant subspaces consisting of dark and bright states. By choosing the initial state of the system, the bright states are all neglected and only dark states are kept to build the effective Hamiltonian. By tuning the system parameters, two-atom multiple-dimensional entanglement, two-atom Bell state, and three-atom GHZ state can be realized by one step. Our proposal provides a way to perform quantum information processing with dark states.

PACS numbers: 03.67.Mn, 03.67.-a, 42.50.Pq

I. INTRODUCTION

Quantum entanglement, one of the most intriguing aspects of quantum mechanics, is an essential resource to quantum information science [1, 2]. Therefore, generating a variety of entangled states have been paid much attention in diverse quantum systems [3–6]. For example, GHZ states, first proposed by Greenberger, Horne, and Zeilinger, provide much stronger refutations of local realism and offer possibilities to test quantum mechanics against local hidden theory without inequality [7], and it has been demonstrated that violations of local realism by two entangled high-dimensional systems are stronger than that by two-dimensional systems [8]. Thus, much interest has been focused on the generation of high-dimensional entangled states. Among various quantum systems, the emerging hybrid quantum systems, with the goal of harnessing the advantages of different subsystems to better explore new phenomena and potentially bring about novel quantum technologies (see [9, 10] for a review), can have versatile applications in quantum information, especially for the atom-cavity-fiber hybrid systems. Because photons as flying qubits can transfer quantum information fast and atoms as stationary qubits can store quantum information for a long time. In the hybrid atom-cavity-fiber system, highly reliable swap and entangling gates are investigated via resonant interaction [11]. Then a scheme for deterministically realization of a highly reliable quantum swap gate is proposed [12]. Later, a resonant interaction scheme for implementing quantum phase gates [13] and a conditional phase gate [14] via a nonresonant interaction are studied. More recently, a remote swap gate through virtual photon and virtual atomic excitation is achieved [15]. Also, proposals of Fredkin Gates and high-dimensional entangled states

have been proposed [16]. Experimentally, dressed states of two separated atoms with the atom-cavity-fiber system have been observed [17].

On the other hand, quantum Zeno effect has attracted extensive interest because it hinders the evolution out of the initial state after performing frequent measurements. In 1988, testing the Zeno effect on oscillating (mainly two-level) systems is theoretically proposed [18] and demonstrated experimentally [19] a few years later. In fact, a quantum Zeno evolution does not necessarily freeze the dynamics. By frequently projecting onto a multidimensional subspace, the system will evolve away from its initial state, although it remains in the Zeno subspace defined by the measurement [20, 21]. Moreover, the quantum Zeno effect can be regarded as a continuous measurement [22] without employing projection operators and nonunitary dynamics. Since then, lots of schemes for preparation of quantum states [23–26] and quantum gates [27–30] via quantum Zeno dynamics have been proposed.

Based on these advanced works, we propose a scheme based on quantum Zeno dynamics to realize two-atom multiple-dimensional entangled states, two-atom Bell state and three-atom GHZ state in a hybrid atom-cavity-fiber system. In this setup, the six-level atom is trapped in one cavity and the other two atoms with three-level configuration are trapped in the other cavity. Two cavities are connected by a fiber. We find that the six-level atom and one of the three-level atoms can be in non-maximally three-dimensional entangled states depending on the initial state of the six-level atom. When the coupling between the atom and the cavity is comparable to the coupling between the fiber and cavity modes, the maximally three-dimensional quantum entanglement for two separated atoms can be generated. Also, the Bell state for the six-level atom and one of the three-level atoms can be obtained when the coupling between the atom and the cavity is much smaller than the coupling between the fiber and cavity modes, which indicates that not very strong atom-cavity coupling is enough for pro-

*xiongweiphys@hotmail.com




FIG. 1: (Color online) Schematic diagram of the hybrid atom-cavity-fiber system. The atom a with six-level structure is trapped in the cavity A , and atoms b and c with three-level structure are trapped in the cavity B . Two cavities are connected by an optical fiber.

ducing Bell states. Preparing the six-level atoms in a specially initial state, a six-dimensional entangled state and GHZ state for three atoms can be achieved. Our setup may provide a way to realize more complicated quantum information processing.

The paper is organized as follows. Section II introduces the physical model of our hybrid system and the method of the quantum Zeno dynamics. In Sec. III, we give a detailed description of producing various entangled states. Finally, we give a short conclusion in Sec. IV.

II. MODEL AND QUANTUM ZENO DYNAMICS

A. Physical model

We consider a hybrid quantum system consisting of two optical cavities (labeled as A and B) connected by an optical fiber and three atoms (labeled as atom a , b , c) with different level structures (see Fig. 1), where the atom a is trapped in the cavity A and the atoms b and c are trapped in the cavity B . The transition between the level $|f_i\rangle_j$ and $|e_i\rangle_j$ is driven by a driving field with Rabi frequency Ω_{ij} , where $i = l, r$ and $j = a, b, c$. The transition between $|g_i\rangle_a$ and $|e_i\rangle_a$ of the atom a is coupled to the i mode of the cavity A , and the transition between $|g_i\rangle_{b(c)}$ and $|e_i\rangle_{b(c)}$ of the atom b (c) is coupled to the i mode of the cavity B . In the short fiber regime, i.e., $2L\bar{\nu}/2\pi c \leq 1$, the i mode of the fiber can be only coupled to the i -mode of the cavities A and B . Here, L is the length of the optical fiber, $\bar{\nu}$ is the decay rate of the cavity fields flowing into a continuum of the fiber modes, and c is the speed of light in vacuum. In the interaction representation, the total Hamiltonian of the hybrid quantum system within the rotating wave approximation

can be governed by (setting $\hbar = 1$)

$$H_{tot} = H_{ac} + H_{cf} + H_d, \quad (1)$$

where

$$\begin{aligned} H_{ac} = & g_{al}|e_l\rangle_{aa}\langle g_l|c_{Al} + g_{ar}|e_r\rangle_{aa}\langle g_r|c_{Ar} \\ & + g_{bl}|e_l\rangle_{bb}\langle g_l|c_{Bl} + g_{cr}|e_r\rangle_{cc}\langle g_r|c_{Br} + \text{H.c.}, \\ H_{cf} = & \lambda_l d_l^\dagger (c_{Al} + c_{Bl}) + \lambda_r d_r^\dagger (c_{Ar} + c_{Br}) + \text{H.c.}, \\ H_d = & \Omega_{al}|e_l\rangle_{aa}\langle f_l| + \Omega_{ar}|e_r\rangle_{aa}\langle f_r| + \Omega_{bl}|e_l\rangle_{bb}\langle f_l| \\ & + \Omega_{cr}|e_r\rangle_{cc}\langle f_r| + \text{H.c.}, \end{aligned} \quad (2)$$

where the first term H_{ac} in Eq. (1) describes the interaction between the atoms and cavities with coupling strengths g_{al} , g_{bl} and g_{cr} . The second term H_{cf} in Eq. (1) describes the coupling between the i mode of the optical fiber and the i mode of the cavities A and B with coupling strength λ_i . The last term H_d in Eq. (1) describes the interaction between the atom and the driving fields. The operators c_{Ai} and c_{Bi} are the annihilation operators of the i mode of the cavities A and B , and H.c. in Eq. (2) denotes the hermitian conjugate.

From Eq. (1), atom a initially prepared in the state $|f_i\rangle_a$ will be excited to the state $|e_i\rangle_a$, then a photon arising from i mode is produced due to the coupling between the i mode of the cavity A and the transition $|e_i\rangle_a \leftrightarrow |g_i\rangle_a$ of the atom a . The generated photon can flow into the cavity B via the interaction between the fiber and the cavities A or B . Once the photon generated from the cavity A enters the cavity B , it will excite the state $|g_l\rangle$ to $|e_l\rangle$ of the atom b or the state $|g_r\rangle$ to $|e_r\rangle$ of the atom c , and vice versa.

B. Quantum Zeno dynamics

Now we briefly give a review about the quantum Zeno dynamics induced by continuous coupling. We suppose a dynamical evolution process is governed by

$$H_K = H_S + K H_C, \quad (3)$$

where H_S is the Hamiltonian of the subsystem to be investigated and H_C is an additional interaction Hamiltonian which performs the measurement, K is the coupling strength. For a large K , i.e., $K \rightarrow \infty$, the subsystem of interest is dominated by the following evolution operator

$$\mathcal{V}(t) = \lim_{K \rightarrow \infty} \exp(iK H_C t) V(t). \quad (4)$$

Equivalently,

$$\mathcal{V}(t) = \exp(-iH_Z t) \quad (5)$$

with

$$H_Z = \sum_n P_n H_S P_n \quad (6)$$

being the Zeno Hamiltonian, in which P_n is the eigenprojection of the H_C belonging to the corresponding eigenvalue E_n ,

$$H_C = \sum_n E_n P_n. \quad (7)$$

Thus the whole system is governed by the limiting evolution operator

$$V_K(t) = \exp(-iH_{\text{eff}}t) \quad (8)$$

with

$$H_{\text{eff}} = \sum_n K E_n P_n t + P_n H_S P_n. \quad (9)$$

This equation is the important result governing our following works.

III. THE DYNAMICS OF THE HYBRID QUANTUM SYSTEM

To begin our task, we for convenience assume $g_{al} = g_{ar} = g_{bl} = g_{br} = g$, $\Omega_{al} = \Omega_{ar} = \Omega_1$, $\Omega_{bl} = \Omega_{br} = \Omega_2$, $\Omega_{cr} = \Omega_3$, $\lambda_l = \lambda_r = \lambda$. Also, the single-photon excitation approximation is assumed in the process of the dynamics evolution.

A. Initial state $|f_l\rangle_a |g_l\rangle_b |g_r\rangle_c |0_l 0_r\rangle_A |0_l 0_r\rangle_B |0_l 0_r\rangle_F$

Based on above analysis, we initially prepare the hybrid system in the state

$$|\psi_0\rangle = |f_l\rangle_a |g_l\rangle_b |g_r\rangle_c |0_l 0_r\rangle_A |0_l 0_r\rangle_B |0_l 0_r\rangle_F, \quad (10)$$

which indicates that the atom a is initially prepared in the state $|f_l\rangle_a$, both the atoms b and c are respectively in the states $|g_l\rangle_b$ and $|g_r\rangle_c$, the cavities A , B and the fiber are all in their vacuum states. Under the interaction of the Hamiltonian (1), seven states are generated constructing a evolution subspace spanned by

$$U_1 = \left\{ |\phi_0\rangle, |\phi_1\rangle, |\phi_2\rangle, |\phi_3\rangle, |\phi_4\rangle, |\phi_5\rangle, |\phi_6\rangle \right\}, \quad (11)$$

with

$$\begin{aligned} |\phi_0\rangle &= |\psi_0\rangle, \\ |\phi_1\rangle &= |e_l\rangle_a |g_l\rangle_b |g_r\rangle_c |0_l 0_r\rangle_A |0_l 0_r\rangle_B |0_l 0_r\rangle_F, \\ |\phi_2\rangle &= |g_l\rangle_a |g_l\rangle_b |g_r\rangle_c |1_l 0_r\rangle_A |0_l 0_r\rangle_B |0_l 0_r\rangle_F, \\ |\phi_3\rangle &= |g_l\rangle_a |g_l\rangle_b |g_r\rangle_c |0_l 0_r\rangle_A |0_l 0_r\rangle_B |1_l 0_r\rangle_F, \\ |\phi_4\rangle &= |g_l\rangle_a |g_l\rangle_b |g_r\rangle_c |0_l 0_r\rangle_A |1_l 0_r\rangle_B |0_l 0_r\rangle_F, \\ |\phi_5\rangle &= |g_l\rangle_a |e_l\rangle_b |g_r\rangle_c |0_l 0_r\rangle_A |0_l 0_r\rangle_B |0_l 0_r\rangle_F, \\ |\phi_6\rangle &= |g_l\rangle_a |f_l\rangle_b |g_r\rangle_c |0_l 0_r\rangle_A |0_l 0_r\rangle_B |0_l 0_r\rangle_F. \end{aligned}$$

Using these basis, the interaction Hamiltonian (1) can be rewritten as

$$H_1 = \begin{pmatrix} 0 & \Omega_1 & 0 & 0 & 0 & 0 & 0 \\ \Omega_1 & 0 & g & 0 & 0 & 0 & 0 \\ 0 & g & 0 & \lambda & 0 & 0 & 0 \\ 0 & 0 & \lambda & 0 & \lambda & 0 & 0 \\ 0 & 0 & 0 & \lambda & 0 & g & 0 \\ 0 & 0 & 0 & 0 & g & 0 & \Omega_2 \\ 0 & 0 & 0 & 0 & 0 & \Omega_2 & 0 \end{pmatrix}. \quad (12)$$

The eigenvectors of Hamiltonian (12) within the assumption of $\{\Omega_1, \Omega_2\} \ll \{g, \lambda\}$, consisting of dark states ($|D_0\rangle, |D_1\rangle, |D_2\rangle$) with zero eigenvalues and bright states ($|B_0\rangle, |B_1\rangle, |B_2\rangle, |B_3\rangle$) with non-zero values, are

$$\begin{aligned} |D_0\rangle &= |\phi_0\rangle, \quad |D_1\rangle = |\phi_6\rangle, \\ |D_2\rangle &= \frac{\lambda}{g\chi}(|\phi_1\rangle - \frac{g}{\lambda}|\phi_3\rangle + |\phi_5\rangle), \\ |B_0\rangle &= \frac{1}{2}(-|\phi_1\rangle - |\phi_2\rangle + |\phi_4\rangle + |\phi_5\rangle), \\ |B_1\rangle &= \frac{1}{2}(-|\phi_1\rangle + |\phi_2\rangle - |\phi_4\rangle + |\phi_5\rangle), \\ |B_2\rangle &= 2\chi(|\phi_1\rangle + \chi|\phi_2\rangle + \frac{\lambda}{g}|\phi_3\rangle - \chi|\phi_4\rangle + |\phi_5\rangle), \\ |B_3\rangle &= 2\chi(|\phi_1\rangle - \chi|\phi_2\rangle + \frac{\lambda}{g}|\phi_3\rangle - \chi|\phi_4\rangle + |\phi_5\rangle), \end{aligned} \quad (13)$$

where $\chi = (1 + 2\lambda^2/g^2)^{\frac{1}{2}}$ and the corresponding eigenvalues are $E_0 = 0$, $E_1 = 0$, $E_2 = 0$, $E_3 = g$, $E_4 = -g$, $E_5 = g\chi$, $E_6 = -g\chi$. Note that the whole Hilbert subspace for Hamiltonian (12) under $\{\Omega_1, \Omega_2\} \ll \{g, \lambda\}$ can be divided into five invariant subspaces as following,

$$\begin{aligned} \Gamma_{p_0} &= \{|D_0\rangle, |D_1\rangle, |D_2\rangle\}, \quad \Gamma_{p_1} = \{|B_0\rangle\} \\ \Gamma_{p_2} &= \{|B_1\rangle\}, \quad \Gamma_{p_3} = \{|B_2\rangle\}, \quad \Gamma_{p_4} = \{|B_3\rangle\}. \end{aligned} \quad (14)$$

Here $p_j = |\alpha\rangle\langle\alpha|$ with $j = 0, 1, 2, 3, 4$ is the projector and $\alpha \in \{\Gamma_{p_0}, \Gamma_{p_1}, \Gamma_{p_2}, \Gamma_{p_3}, \Gamma_{p_4}\}$. According to quantum Zeno dynamics method, the Hamiltonian in Eq. (12) reduces to

$$\begin{aligned} \mathcal{H}_1 &= \sum_{j=0} E_j p_j + p_j H_d p_j \\ &= g[|B_0\rangle\langle B_0| - |B_1\rangle\langle B_1| + \chi(|B_2\rangle\langle B_2| - |B_3\rangle\langle B_3|)] \\ &\quad + \frac{\lambda}{g\chi}(\Omega_1|D_0\rangle\langle D_2| + \Omega_2|D_1\rangle\langle D_2| + \text{H.c.}). \end{aligned} \quad (15)$$

Since the system is initially prepared in the state $|D_0\rangle$, the system will always evolve in the Zeno subspace spanned by $\{|D_0\rangle, |D_2\rangle, |B_2\rangle\}$. The corresponding dynamics can be governed by the effective Hamiltonian

$$H_{\text{eff}} = \frac{\lambda}{g\chi}(\Omega_1|D_0\rangle\langle D_2| + \Omega_2|D_1\rangle\langle D_2| + \text{H.c.}), \quad (16)$$

and at time τ , the system state is

$$|\psi_\tau\rangle = e^{-iH_{\text{eff}}\tau}|\psi_0\rangle = A_1|D_0\rangle + A_2|D_2\rangle + A_3|B_2\rangle, \quad (17)$$

where

$$\begin{aligned} A_1 &= \sin^2 \theta + \cos^2 \theta \cos(\Omega\lambda\tau/g\chi), \\ A_2 &= -i \cos \theta \sin(\Omega\lambda\tau/g\chi) \\ A_3 &= \frac{1}{2} \sin(2\theta) [\cos(\Omega\lambda\tau/g\chi) - 1] \end{aligned} \quad (18)$$

with $\Omega_1^2 + \Omega_2^2 = \Omega^2$ and $\tan \theta = \Omega_2/\Omega_1$. By controlling the system parameters $\{\Omega_1, \Omega_2, \lambda, g, \tau\}$, different quantum information task can be achieved. For example, when $\Omega_2 \ll \Omega_1$, $\sin \theta \rightarrow 0$ and $\cos \theta \rightarrow 1$. Thus, $A_1 \rightarrow \cos(\Omega\lambda\tau/g\chi)$, $A_2 \rightarrow -i \sin(\Omega\lambda\tau/g\chi)$ and $A_3 \rightarrow 0$. Then we choose $\Omega\lambda\tau/g\chi = (2k-1)\pi/2$ with $k = 1, 2, 3, \dots$, leading to $\cos(\Omega\lambda\tau/g\chi) = 0$ and $\sin(\Omega\lambda\tau/g\chi) = 1$. This condition can be achieved by tuning the parameters $\{\Omega_1, \Omega_2, \lambda, g, \tau\}$, which greatly reduces the difficulties experimentally. With this condition, Eq. (17) becomes

$$|\psi_\tau\rangle_{\text{ST}} = -i|D_2\rangle, \quad (19)$$

which indicates quantum state transition between two degenerate states is allowed and the state $|D_1\rangle$ is safely neglected. Note that when the photons in the l mode of the optical fiber experience a Hadamard gate, i.e., $|0\rangle \rightarrow \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ and $|1\rangle \rightarrow \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$, and then trace the degrees of the freedom of the atom b and the photons in the cavity and fiber modes out, the quantum state in Eq. (19) further reduces to

$$|\psi_\tau\rangle_{\text{TD}} = \frac{1}{\sqrt{3}}(|e_l\rangle|g_l\rangle - |g_l\rangle|g_l\rangle + |g_l\rangle|e_l\rangle)_{ab} \quad (20)$$

by setting $g = \lambda$, which is a three-dimensional quantum entanglement between atoms a and b . However, when $g \ll \lambda$, Eq. (19) describes a Bell state between atoms a and b , i.e.,

$$|\psi_\tau\rangle_{\text{Bell}} = \frac{1}{\sqrt{2}}(|e_l\rangle|g_l\rangle + |g_l\rangle|e_l\rangle)_{ab}. \quad (21)$$

On the other hand, Eq. (17) can be rewritten as

$$|\psi_\tau\rangle_{\text{SW}} = |D_1\rangle \quad (22)$$

by setting $\Omega_1 = \Omega_2 = \Omega/\sqrt{2}$ and $\Omega\lambda\tau/g\chi = k\pi$ with $k = 1, 2, 3, \dots$. This implies that state swapping ($|f_l\rangle_a|g_l\rangle_b \leftrightarrow |g_l\rangle_a|f_l\rangle_b$) between atoms a and b is achieved.

B. Initial state $|f_r\rangle_a|g_l\rangle_b|g_r\rangle_c|0_l0_r\rangle_A|0_l0_r\rangle_B|0_l0_r\rangle_F$

Below we assume that the initial state of the atom-cavity-fiber hybrid system is

$$|\psi'_0\rangle = |f_r\rangle_a|g_l\rangle_b|g_r\rangle_c|0_l0_r\rangle_A|0_l0_r\rangle_B|0_l0_r\rangle_F, \quad (23)$$

which indicates that the atom a is initially prepared in the state $|f_r\rangle_a$, both the atoms b and c are respectively in the states $|g_l\rangle_b$ and $|g_r\rangle_c$, the cavities A , B and the

fiber are all in their vacuum states. Under the interaction of the Hamiltonian (1), seven states are generated constructing a evolution subspace spanned by

$$U'_1 = \left\{ |\phi'_0\rangle, |\phi'_1\rangle, |\phi'_2\rangle, |\phi'_3\rangle, |\phi'_4\rangle, |\phi'_5\rangle, |\phi'_6\rangle \right\}, \quad (24)$$

with

$$\begin{aligned} |\phi'_0\rangle &= |\psi'_0\rangle, \\ |\phi'_1\rangle &= |e_r\rangle_a|g_l\rangle_b|g_r\rangle_c|0_l0_r\rangle_A|0_l0_r\rangle_B|0_l0_r\rangle_F, \\ |\phi'_2\rangle &= |g_r\rangle_a|g_l\rangle_b|g_r\rangle_c|1_l0_r\rangle_A|0_l0_r\rangle_B|0_l0_r\rangle_F, \\ |\phi'_3\rangle &= |g_r\rangle_a|g_l\rangle_b|g_r\rangle_c|0_l0_r\rangle_A|0_l0_r\rangle_B|0_l1_r\rangle_F, \\ |\phi'_4\rangle &= |g_r\rangle_a|g_l\rangle_b|g_r\rangle_c|0_l0_r\rangle_A|0_l1_r\rangle_B|0_l0_r\rangle_F, \\ |\phi'_5\rangle &= |g_r\rangle_a|g_l\rangle_b|e_r\rangle_c|0_l0_r\rangle_A|0_l0_r\rangle_B|0_l0_r\rangle_F, \\ |\phi'_6\rangle &= |g_r\rangle_a|g_l\rangle_b|f_r\rangle_c|0_l0_r\rangle_A|0_l0_r\rangle_B|0_l0_r\rangle_F. \end{aligned}$$

Using these basis, the interaction Hamiltonian (1) can be rewritten as

$$H'_1 = \begin{pmatrix} 0 & \Omega_1 & 0 & 0 & 0 & 0 & 0 \\ \Omega_1 & 0 & g & 0 & 0 & 0 & 0 \\ 0 & g & 0 & \lambda & 0 & 0 & 0 \\ 0 & 0 & \lambda & 0 & \lambda & 0 & 0 \\ 0 & 0 & 0 & \lambda & 0 & g & 0 \\ 0 & 0 & 0 & 0 & g & 0 & \Omega_3 \\ 0 & 0 & 0 & 0 & 0 & \Omega_3 & 0 \end{pmatrix}. \quad (25)$$

The eigenvectors of Hamiltonian (25) within the assumption of $\{\Omega_1, \Omega_3\} \ll \{g, \lambda\}$, consisting of dark states ($|D'_0\rangle, |D'_1\rangle, |D'_2\rangle$) with zero eigenvalues and bright states ($|B'_0\rangle, |B'_1\rangle, |B'_2\rangle, |B'_3\rangle$) with non-zero values, are

$$\begin{aligned} |D'_0\rangle &= |\phi'_0\rangle, \quad |D'_1\rangle = |\phi'_6\rangle, \\ |D'_2\rangle &= \frac{\lambda}{g\chi}(|\phi'_1\rangle - \frac{g}{\lambda}|\phi'_3\rangle + |\phi'_5\rangle), \\ |B'_0\rangle &= \frac{1}{2}(-|\phi'_1\rangle - |\phi'_2\rangle + |\phi'_4\rangle + |\phi'_5\rangle), \\ |B'_1\rangle &= \frac{1}{2}(-|\phi'_1\rangle + |\phi'_2\rangle - |\phi'_4\rangle + |\phi'_5\rangle), \\ |B'_2\rangle &= 2\chi(|\phi'_1\rangle + \chi|\phi'_2\rangle + \frac{\lambda}{g}|\phi'_3\rangle - \chi|\phi'_4\rangle + |\phi'_5\rangle), \\ |B'_3\rangle &= 2\chi(|\phi'_1\rangle - \chi|\phi'_2\rangle + \frac{\lambda}{g}|\phi'_3\rangle - \chi|\phi'_4\rangle + |\phi'_5\rangle), \end{aligned} \quad (26)$$

where $\chi = (1 + 2\lambda^2/g^2)^{\frac{1}{2}}$ and the corresponding eigenvalues are $E'_0 = 0$, $E'_1 = 0$, $E'_2 = 0$, $E'_3 = g$, $E'_4 = -g$, $E'_5 = g\chi$, $E'_6 = -g\chi$. Note that the whole Hilbert subspace for Hamiltonian (25) under $\{\Omega_1, \Omega_3\} \ll \{g, \lambda\}$ can be divided into five invariant subspaces as following,

$$\begin{aligned} \Gamma_{p'_0} &= \{|D'_0\rangle, |D'_1\rangle, |D'_2\rangle\}, \quad \Gamma_{p'_1} = \{|B'_0\rangle\} \\ \Gamma_{p'_2} &= \{|B'_1\rangle\}, \quad \Gamma_{p'_3} = \{|B'_2\rangle\}, \quad \Gamma_{p'_4} = \{|B'_3\rangle\}. \end{aligned} \quad (27)$$

Here $p'_j = |\alpha'\rangle\langle\alpha'|$ with $j = 0, 1, 2, 3, 4$ is the projector and $\alpha' \in \{\Gamma_{p'_0}, \Gamma_{p'_1}, \Gamma_{p'_2}, \Gamma_{p'_3}, \Gamma_{p'_4}\}$. According to quantum Zeno dynamics method, the Hamiltonian in Eq. (25)

reduces to

$$\begin{aligned}\mathcal{H}'_1 &= \sum_{j=0} E'_j p'_j + p'_j H_d p'_j \\ &= g[|B'_0\rangle\langle B'_0| - |B'_1\rangle\langle B'_1| + \chi(|B'_2\rangle\langle B'_2| - |B'_3\rangle\langle B'_3|)] \\ &\quad + \frac{\lambda}{g\chi}(\Omega_1|D'_0\rangle\langle D'_2| + \Omega_3|D'_1\rangle\langle D'_2| + \text{H.c.}).\end{aligned}\quad (28)$$

Since the system is initially prepared in the state $|D'_0\rangle$, the system will always evolve in the Zeno subspace spanned by $\{|D'_0\rangle, |D'_2\rangle, |B'_2\rangle\}$. The corresponding dynamics can be governed by the effective Hamiltonian

$$H'_{\text{eff}} = \frac{\lambda}{g\chi}(\Omega_1|D'_0\rangle\langle D'_2| + \Omega_3|D'_1\rangle\langle D'_2| + \text{H.c.}), \quad (29)$$

and at time τ' , the system state is

$$|\psi'_{\tau'}\rangle = e^{-iH'_{\text{eff}}\tau'}|\psi'_0\rangle = A'_1|D'_0\rangle + A'_2|D'_2\rangle + A'_3|D'_1\rangle, \quad (30)$$

where

$$\begin{aligned}A'_1 &= \sin^2 \theta' + \cos^2 \theta' \cos(\bar{\Omega}\lambda\tau'/g\chi), \\ A'_2 &= -i \cos \theta' \sin(\bar{\Omega}\lambda\tau'/g\chi) \\ A'_3 &= \frac{1}{2} \sin(2\theta') [\cos(\bar{\Omega}\lambda\tau'/g\chi) - 1]\end{aligned}\quad (31)$$

with $\Omega_1^2 + \Omega_3^2 = \bar{\Omega}^2$ and $\tan \theta' = \Omega_3/\Omega_1$.

By tuning the system parameters $\{\Omega_1, \Omega_3, \lambda, g, \tau'\}$, the similar quantum information tasks can be achieved in Sec. III A. For example, when $\Omega_3 \ll \Omega_1$, $\sin \theta' \rightarrow 0$ and $\cos \theta' \rightarrow 1$. Thus, $A'_1 \rightarrow \cos(\bar{\Omega}\lambda\tau'/g\chi)$, $A'_2 \rightarrow -i \sin(\bar{\Omega}\lambda\tau'/g\chi)$ and $A'_3 \rightarrow 0$. Then we choose $\bar{\Omega}\lambda\tau'/g\chi = (2k-1)\pi/2$ with $k = 1, 2, 3, \dots$, leading to $\cos(\bar{\Omega}\lambda\tau'/g\chi) = 0$ and $\sin(\bar{\Omega}\lambda\tau'/g\chi) = 1$. This condition can be achieved by tuning the parameters $\{\Omega_1, \Omega_3, \lambda, g, \tau'\}$, which greatly reduces the difficulties experimentally. With this condition, Eq. (30) becomes

$$|\psi'_{\tau'}\rangle_{\text{ST}} = -i|D'_2\rangle, \quad (32)$$

which indicates quantum state transition between two degenerate states is allowed and the state $|D'_1\rangle$ is safely neglected. Note that when the photons in the r mode of the optical fiber experience a Hadamard gate, i.e., $|0_r\rangle \rightarrow \frac{1}{\sqrt{2}}(|0_r\rangle + |1_r\rangle)$ and $|1_r\rangle \rightarrow \frac{1}{\sqrt{2}}(|0_r\rangle - |1_r\rangle)$, and then trace the degrees of the freedom of the atom b and the photons in the cavity and fiber modes out, the quantum state in Eq. (32) further reduces to

$$|\psi'_{\tau'}\rangle_{\text{TD}} = \frac{1}{\sqrt{3}}(|e_r\rangle|g_r\rangle - |g_r\rangle|g_r\rangle + |g_r\rangle|e_r\rangle)_{ac} \quad (33)$$

by setting $g = \lambda$, which is a three-dimensional quantum entanglement between atoms a and c . However, when $g \ll \lambda$, Eq. (32) describes a Bell state between atoms a and c , i.e.,

$$|\psi'_{\tau'}\rangle_{\text{Bell}} = \frac{1}{\sqrt{2}}(|e_r\rangle|g_r\rangle + |g_r\rangle|e_r\rangle)_{ac}. \quad (34)$$

However, by setting $\Omega_1 = \Omega_3 = \bar{\Omega}/\sqrt{2}$ and $\bar{\Omega}\lambda\tau'/g\chi = k\pi$ with $k = 1, 2, 3, \dots$, Eq. (30) becomes

$$|\psi'_{\tau'}\rangle_{\text{SW}} = |D'_1\rangle. \quad (35)$$

This implies that state swapping between atoms a and c , i.e., $|f_r\rangle_a|g_r\rangle_c \leftrightarrow |g_r\rangle_a|f_r\rangle_c$, can be achieved.

C. Initial state $\Psi_0 = |\psi_0\rangle + |\psi'_0\rangle$

Based on the analysis in Sec. III A and Sec. III B, we further consider that the atom a is initially prepared in the superposition state $|g_l\rangle_a + |e_r\rangle_a$, atoms b and c are in the states $|g_l\rangle_b$ and $|g_r\rangle_c$, respectively. Cavities A and B , and the fiber are all in their vacuum states. Therefore, the initial state of the considered system can be written as

$$|\Psi_0\rangle = |\psi_0\rangle + |\psi'_0\rangle. \quad (36)$$

According to Eqs. (11) and (24), the evolution space governed by Eq. (1) with the initial state $|\Psi_0\rangle$ is spanned by states in U_1 plus U'_1 , i.e.,

$$U = \{|\Phi_0\rangle, |\Phi_1\rangle, |\Phi_2\rangle, |\Phi_3\rangle, |\Phi_4\rangle, |\Phi_5\rangle, |\Phi_6\rangle\}, \quad (37)$$

where

$$\begin{aligned}|\Phi_0\rangle &= |\Psi_0\rangle, \\ |\Phi_1\rangle &= |\phi_1\rangle + |\phi'_1\rangle, \\ |\Phi_2\rangle &= |\phi_2\rangle + |\phi'_2\rangle, \\ |\Phi_3\rangle &= |\phi_3\rangle + |\phi'_3\rangle, \\ |\Phi_4\rangle &= |\phi_4\rangle + |\phi'_4\rangle, \\ |\Phi_5\rangle &= |\phi_5\rangle + |\phi'_5\rangle, \\ |\Phi_6\rangle &= |\phi_6\rangle + |\phi'_6\rangle.\end{aligned}\quad (38)$$

Using the basis in Eq. (37), the system Hamiltonian in Eq. (1) reduces to

$$H = \begin{pmatrix} 0 & \Omega_1 & 0 & 0 & 0 & 0 & 0 \\ \Omega_1 & 0 & g & 0 & 0 & 0 & 0 \\ 0 & g & 0 & \lambda & 0 & 0 & 0 \\ 0 & 0 & \lambda & 0 & \lambda & 0 & 0 \\ 0 & 0 & 0 & \lambda & 0 & g & 0 \\ 0 & 0 & 0 & 0 & g & 0 & \Omega_2 + \Omega_3 \\ 0 & 0 & 0 & 0 & 0 & \Omega_2 + \Omega_3 & 0 \end{pmatrix}. \quad (39)$$

When we choose $\Omega_1, \Omega_2, \Omega_3 \ll g, \lambda$, the states in Eqs. (13) and (26) can be simultaneously obtained. Thus, the eigenvectors of the Hamiltonian in Eq. (1) are

$$\begin{aligned}|\mathcal{D}_0\rangle &= |D_0\rangle + |D'_0\rangle, \\ |\mathcal{D}_1\rangle &= |D_1\rangle + |D'_1\rangle, \\ |\mathcal{D}_2\rangle &= |D_2\rangle + |D'_2\rangle, \\ |\mathcal{B}_0\rangle &= |B_0\rangle + |B'_0\rangle, \\ |\mathcal{B}_1\rangle &= |B_1\rangle + |B'_1\rangle, \\ |\mathcal{B}_2\rangle &= |B_2\rangle + |B'_2\rangle, \\ |\mathcal{B}_3\rangle &= |B_3\rangle + |B'_3\rangle,\end{aligned}\quad (40)$$

with $\chi = (1+2\lambda^2/g^2)^{\frac{1}{2}}$ and the corresponding eigenvalues are $\mathcal{E}_0 = 0$, $\mathcal{E}_1 = 0$, $\mathcal{E}_2 = 0$, $\mathcal{E}_3 = g$, $\mathcal{E}_4 = -g$, $\mathcal{E}_5 = g\chi$, $\mathcal{E}_6 = -g\chi$. The space consisting of these eigenvectors can be split into five invariant Zeno subspace

$$\begin{aligned} \Gamma_{\mathcal{P}_0} &= \{|\mathcal{D}_0\rangle, |\mathcal{D}_1\rangle, |\mathcal{D}_2\rangle\}, \quad \Gamma_{\mathcal{P}_1} = \{|\mathcal{B}_0\rangle\} \\ \Gamma_{\mathcal{P}_2} &= \{|\mathcal{B}_1\rangle\}, \quad \Gamma_{\mathcal{P}_3} = \{|\mathcal{B}_2\rangle\}, \quad \Gamma_{\mathcal{P}_4} = \{|\mathcal{B}_3\rangle\}. \end{aligned} \quad (41)$$

Here $\mathcal{P}_j = |\bar{\alpha}\rangle\langle\bar{\alpha}|$ with $j = 0, 1, 2, 3, 4$ is the projector and $\bar{\alpha} \in \{\Gamma_{\mathcal{P}_0}, \Gamma_{\mathcal{P}_1}, \Gamma_{\mathcal{P}_2}, \Gamma_{\mathcal{P}_3}, \Gamma_{\mathcal{P}_4}\}$. Combining Eqs. (15) and (28), the Hamiltonian in Eq. (39) reduces to

$$\begin{aligned} \mathcal{H} &= \sum_{j=0} \mathcal{E}_j \mathcal{P}_j + \mathcal{P}_j H_d \mathcal{P}_j \\ &= g[|\mathcal{B}_0\rangle\langle\mathcal{B}_0| - |\mathcal{B}_1\rangle\langle\mathcal{B}_1| + \chi(|\mathcal{B}_2\rangle\langle\mathcal{B}_2| - |\mathcal{B}_3\rangle\langle\mathcal{B}_3|)] \\ &\quad + \frac{\lambda}{g\chi}(\Omega_1|\mathcal{D}_0\rangle\langle\mathcal{D}_2| + (\Omega_2|\mathcal{D}_1\rangle + \Omega_3|\mathcal{D}'_1\rangle)\langle\mathcal{D}_2| + \text{H.c.}). \end{aligned} \quad (42)$$

When the system state is initially prepared in the state $|\mathcal{D}_0\rangle$, Eq. (42) can be rewritten as

$$\mathcal{H}_{\text{eff}} = \frac{\lambda}{g\chi}(\Omega_1|\mathcal{D}_0\rangle\langle\mathcal{D}_2| + (\Omega_2|\mathcal{D}_1\rangle + \Omega_3|\mathcal{D}'_1\rangle)\langle\mathcal{D}_2| + \text{H.c.}). \quad (43)$$

At time T , the system state is

$$\begin{aligned} |\Psi_T\rangle &= \mathcal{A}_1|\mathcal{D}_0\rangle + \mathcal{A}'_1|\mathcal{D}'_0\rangle + \mathcal{A}_2|\mathcal{D}_2\rangle \\ &\quad + \mathcal{A}'_2|\mathcal{D}'_2\rangle + \mathcal{A}_3|\mathcal{D}_1\rangle + \mathcal{A}'_3|\mathcal{D}'_1\rangle, \end{aligned} \quad (44)$$

where

$$\begin{aligned} \mathcal{A}_1 &= A_1, \quad \mathcal{A}'_1 = A'_1, \quad \mathcal{A}_2 = A_2, \\ \mathcal{A}'_2 &= A'_2, \quad \mathcal{A}_3 = A_3, \quad \mathcal{A}'_3 = A'_3. \end{aligned} \quad (45)$$

Obviously, $\Omega_2, \Omega_3 \ll \Omega_1$, the state q becomes

$$|\Psi_T\rangle_{\text{ST}} = -i\mathcal{D}_2, \quad (46)$$

which indicates that quantum state transition between two dark states can be achieved. The similar process

from Eq. (20) to Eq. (22) or Eq. (33) to Eq. (35) is performed, the high-dimensional quantum entangled state among three atoms with setting $g = \lambda$

$$\begin{aligned} |\Psi_T\rangle_{\text{HD}} &= \frac{1}{\sqrt{6}}[(|e_r\rangle|g_l\rangle|g_r\rangle - |g_r\rangle|g_l\rangle|g_r\rangle + |g_r\rangle|g_l\rangle|e_r\rangle)_{abc} \\ &\quad + (|e_l\rangle|g_l\rangle|g_r\rangle - |g_l\rangle|g_l\rangle|g_r\rangle + |g_l\rangle|e_l\rangle|g_r\rangle)_{abc}] \end{aligned} \quad (47)$$

can be obtained, and the three atoms GHZ state

$$|\Psi_T\rangle_{\text{GHZ}} = \frac{1}{\sqrt{2}}\mathcal{D}_1 \quad (48)$$

is also realized when $\Omega_1 = \Omega_2 = \Omega_3$ and $\Omega_3\lambda T/g\chi = k\pi$ are satisfied.

IV. CONCLUSION

In conclusion, we have proposed a hybrid system to entangle remote atoms trapped in two cavities linked by an optical fiber. Employing quantum Zeno dynamics method, bright states of the system are all neglected and dark states are kept to build the effective Hamiltonian. By considering the relative strength between the atom-cavity coupling and cavity-fiber coupling and preparing the six-level atom in different initial state, the Bell state, the two-atom three-dimensional entangled states, the three-atom GHZ state and six-dimensional entangled state are achieved. Our proposal provides a path to perform quantum information processing with dark states.

Acknowledgments

This work is supported by the National Natural Science Foundation of China (Grant No. 11804074), Talent Development Funding of Hefei University (Grant No. 18-19RC60), Candidates of Hefei University Academic Leader(Grant No.2016dtr02), Major Project of Hefei Preschool Education College (Grant No.hyzzd2018003).

-
- [1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, Rev. Mod. Phys. 81, 865 (2009).
 - [2] F. Giraldi and P. Grigolini, Quantum entanglement and entropy, Phys. Rev. A 64, 032310 (2001).
 - [3] I. L. Chuang and Y. Yamamoto, Simple quantum computer, Phys. Rev. A 52, 3489 (1995).
 - [4] J. H. Reina, L. Quiroga, and N. F. Johnson, Quantum entanglement and information processing via excitons in optically driven quantum dots, Phys. Rev. A 62, 012305 (2000).
 - [5] S. B. Zheng and G. C. Guo, Efficient Scheme for Two-Atom Entanglement and Quantum Information Process-

- ing in Cavity QED, Phys. Rev. Lett. 85, 2392 (2000).
- [6] R. G. Unanyan and M. Fleischhauer, Decoherence-Free Generation of Many-Particle Entanglement by Adiabatic Ground-State Transitions, Phys. Rev. Lett. 90, 133601 (2003).
- [7] D. M. Greenberger, M. A. Horne, A. Shimony, and A. Zeilinger, Bells theorem without inequalities, Am. J. Phys. 58, 1131 (1990).
- [8] D. Kaszlikowski, P. Gnacinski, M. Zukowski, W. Miklaszewski, and A. Zeilinger, Violations of Local Realism by Two Entangled N -Dimensional Systems Are Stronger than for Two Qubits, Phys. Rev. Lett. 85, 4418 (2000).
- [9] Z. L. Xiang, S. Ashhab, J. Q. You, and F. Nori, Hybrid

- quantum circuits: Superconducting circuits interacting with other quantum systems, *Rev. Mod. Phys.* **85**, 623 (2013).
- [10] G. Kurizki, P. Bertet, Y. Kubo, K. Mølmer, D. Petrosyan, P. Rabl, and J. Schmiedmayer, Quantum technologies with hybrid systems, *Proc. Natl. Acad. Sci. USA* **112**, 3866 (2015).
 - [11] A. Serafini, S. Mancini, and S. Bose, Distributed Quantum Computation via Optical Fibers, *Phys. Rev. Lett.* **96**, 010503 (2006).
 - [12] Z. Yin and F. Li, Multiatom and resonant interaction scheme for quantum state transfer and logical gates between two remote cavities via an optical fiber, *Phys. Rev. A* **75**, 012324 (2007).
 - [13] Y. H. Chen, Y. Xia, Q. Q. Chen, and J. Song, Fast and noise-resistant implementation of quantum phase gates and creation of quantum entangled states, *Phys. Rev. A* **91**, 012325 (2015).
 - [14] S. B. Zheng, Virtual-photon-induced quantum phase gates for two distant atoms trapped in separate cavities, *Appl. Phys. Lett.* **94**, 154101 (2009).
 - [15] J. L. Wu and S. L. Su, Auxiliary-qubit-driving-induced entanglement and logic gate, *Europhys. Lett.* **126**, 30001 (2019).
 - [16] G. Q. Sun, J. L. Wu, W. Niu, W. R. Yu, and X. Ji, Remote Implementation of a Fredkin Gate via Virtual Excitation of an Atom-Cavity-Fiber System, *Ann. Phys. (Berlin)* **532**, 1900372 (2020).
 - [17] S. Kato, N. Német, K. Senga, S. Mizukami, X. Huang, S. Parkins, and T. Aoki, Observation of dressed states of distant atoms with delocalized photons in coupled-cavities quantum electrodynamics, *Nat. Commun* **10**, 1160 (2019).
 - [18] R. J. Cook, What are Quantum Jumps? *Phys. Scr. T* **21**, 49 (1988).
 - [19] W. M. Itano, D. J. Heinzen, J. J. Bollinger, and D. J. Wineland, Quantum Zeno effect, *Phys. Rev. A* **41**, 2295 (1990).
 - [20] P. Facchi, V. Gorini, G. Marmo, S. Pascazio, and E. C. G. Sudarshan, Quantum Zeno dynamics, *Phys. Lett. A* **275**, 12 (2000).
 - [21] P. Facchi, S. Pascazio, A. Scardicchio, and L. S. Schulman, Zeno dynamics yields ordinary constraints, *Phys. Rev. A* **65**, 012108 (2001).
 - [22] P. Facchi, G. Marmo, and S. Pascazio, Quantum Zeno dynamics and quantum Zeno subspaces, *J. Phys: Conf. Ser.* **196**, 012017 (2009).
 - [23] A. Luis, Quantum-state preparation and control via the Zeno effect, *Phys. Rev. A* **63**, 052112 (2001).
 - [24] X. B. Wang, J. Q. You, and F. Nori, Quantum entanglement via two-qubit quantum Zeno dynamics, *Phys. Rev. A* **77**, 062339 (2008).
 - [25] W. A. Li and G. Y. Huang, Deterministic generation of a three-dimensional entangled state via quantum Zeno dynamics, *Phys. Rev. A* **83**, 022322 (2011).
 - [26] L. T. Shen, H. Z. Wu and R. X. Chen, Robust generation of a four-dimensional entangled state in separate cavities via quantum Zeno dynamics, *J. Phys. B: At. Mol. Opt. Phys.* **44**, 205503 (2011).
 - [27] A. Beige, D. Braun, B. Tregenna, and P. L. Knight, Quantum Computing Using Dissipation to Remain in a Decoherence-Free Subspace, *Phys. Rev. Lett.* **85**, 1762 (2000).
 - [28] J. D. Franson, B. C. Jacobs, and T. B. Pittman, Quantum computing using single photons and the Zeno effect, *Phys. Rev. A* **70**, 062302 (2004).
 - [29] X. Q. Shao, L. Chen, S. Zhang, and K. H. Yeon, Fast CNOT gate via quantum Zeno dynamics, *J. Phys. B: At. Mol. Opt. Phys.* **42**, 165507 (2009).
 - [30] J. D. Franson, T. B. Pittman, and B. C. Jacobs, Zeno logic gates using microcavities, *J. Opt. Soc. Am. B* **24**, 209 (2007).