

Bubbling and extinction for some fast diffusion equations in bounded domains

Tianling Jin*, Jingang Xiong†

August 4, 2020

Abstract

We study a Sobolev critical fast diffusion equation in bounded domains with the Brezis-Nirenberg effect. We obtain extinction profiles of its positive solutions, and show that the convergence rates of the relative error in regular norms are at least polynomial. Exponential decay rates are proved for generic domains. Our proof makes use of its regularity estimates, a curvature type evolution equation, as well as blow up analysis. Results for Sobolev subcritical fast diffusion equations are also obtained.

1 Introduction

Let Ω be a bounded domain in \mathbb{R}^n , $n \geq 3$, with smooth boundary $\partial\Omega$. We consider the Cauchy-Dirichlet problem for the fast diffusion equation with the Sobolev critical exponent

$$\begin{aligned} \frac{\partial}{\partial t} u^{\frac{n+2}{n-2}} &= \Delta u + bu & \text{in } \Omega \times (0, \infty), \\ u &= 0 & \text{on } \partial\Omega \times (0, \infty), \\ u(\cdot, 0) &= u_0 \geq 0 & \text{in } \Omega, \end{aligned} \tag{1}$$

where $\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$ is the Laplace operator, u_0 is not identically zero, and

$$b \in [0, \lambda_1) \text{ is a constant} \tag{2}$$

with λ_1 being the first eigenvalue of $-\Delta$ in Ω with zero Dirichlet boundary condition. Hence, the operator $-\Delta - b$ is coercive on the Sobolev space $H_0^1(\Omega)$. The fast diffusion equations arise in the modelling of gas-kinetics, plasmas, thin liquid film dynamics driven by Van der Waals forces, and etc. If $b = 0$, this Sobolev critical equation (1) can be viewed a unnormalized Yamabe flow with metrics degenerate on the boundary.

The theory of existence and uniqueness of solutions to (1) is well understood, see Vázquez [33, 34]. If $u_0 \in L^q(\Omega)$ for some $q > \frac{2n}{n-2}$, then the solution will become instantaneously positive

*T. Jin is partially supported by Hong Kong RGC grants GRF 16302217 and GRF 16306320.

†J. Xiong is partially supported by NSFC 11922104 and 11631002.

in Ω and globally bounded. Moreover, the solution will vanish in a finite time $T^* > 0$. If we assume that $u_0 \in H_0^1(\Omega) \cap L^q(\Omega)$ for some $q > \frac{2n}{n-2}$, then it follows from the work of Chen-DiBenedetto [11], DiBenedetto-Kwong-Vespri [18] and Jin-Xiong [23] that the solution is of $C_{x,t}^{3,2}(\overline{\Omega} \times (0, T^*))$. In particular, the solutions are classical. Therefore, when we investigate the asymptotic behavior of nonnegative solutions to (1) as t approaching to the extinction time T^* , there is no loss of generality to consider classical (up to the boundary) solutions to (1).

When $\frac{n+2}{n-2}$ is replaced by $p \in (1, \frac{n+2}{n-2})$ if $n \geq 3$, or $p \in (1, \infty)$ if $n = 1, 2$, which is a Sobolev subcritical exponent, the extinction behavior of solutions to the fast diffusion equation

$$\begin{aligned} \frac{\partial}{\partial t} u^p &= \Delta u & \text{in } \Omega \times (0, \infty), \\ u &= 0 & \text{on } \partial\Omega \times (0, \infty) \end{aligned} \quad (3)$$

has been well-studied. By the scaling

$$v(x, t) = \left(\frac{p}{(p-1)(T^* - \tau)} \right)^{\frac{1}{p-1}} u(x, \tau), \quad t = \frac{p}{p-1} \ln \left(\frac{T^*}{T^* - \tau} \right), \quad (4)$$

where T^* is the extinction time, the equation (3) becomes

$$\begin{aligned} \frac{\partial}{\partial t} v^p &= \Delta v + v^p & \text{in } \Omega \times (0, \infty), \\ v &= 0 & \text{on } \partial\Omega \times (0, \infty). \end{aligned} \quad (5)$$

Berryman-Holland [3] proved that the solution of (5) converges to a stationary solution v_∞ in $H_0^1(\Omega)$ along a sequence of times. Feireisl-Simondon [20] proved the full convergence in the $C^0(\overline{\Omega})$ topology. Bonforte-Grillo-Vázquez [5] proved that the relative error $v(\cdot, t)/v_\infty$ converges to 1 in $L^\infty(\Omega)$. Recently, Bonforte-Figalli [4] proved the exponential convergence of the relative error for generic domains Ω , which means that the domains Ω satisfy

$$\text{For every nonnegative } H_0^1 \text{ solution } v \text{ of } -\Delta v - v^p = 0 \text{ in } \Omega, \text{ the linearized operator at } v, \text{ that is } L_v := -\Delta - pv^{p-1}, \text{ has a trivial kernel in } H_0^1(\Omega). \quad (6)$$

The set of smooth domains satisfying (6) has generic properties, see Saut-Temam [28].

The main advantage of the subcritical regime is the upper bound of solutions u to (3) proved in DiBenedetto-Kwong-Vespri [18]

$$u(x, t) \leq Cd(x)(T^* - t)^{\frac{1}{p-1}} \quad \text{for } t < T^*, \quad (7)$$

where $d(x) = \text{dist}(x, \partial\Omega)$. The estimate (7) implies that the function v defined by (4), which satisfies (5), is uniformly bounded as $t \rightarrow \infty$, and consequently, has uniform regularity estimates up to the boundary $\partial\Omega$ by the work of [11, 18, 23].

However, this uniform bound in general does not hold for (5) if $p = \frac{n+2}{n-2}$. For instance, it is the case if Ω is star-shaped, since there is no stationary solution of (5) due to the Pohozaev identity. In this paper, we will show that the uniform boundedness still holds for the equation (1) assuming $b > 0$ and $n \geq 4$. The role of the positivity of b when $n \geq 4$ was first discovered in the seminal paper Brezis-Nirenberg [8], and is similar to the role that the non-vanishing Weyl tensor and the positive

mass theorem play in the resolution of the Yamabe problem on compact manifolds by Aubin [1] and Schoen [29].

Under the scaling

$$v(x, t) = \left(\frac{n+2}{4(T^* - \tau)} \right)^{\frac{n-2}{4}} u(x, \tau), \quad t = \frac{n+2}{4} \ln \left(\frac{T^*}{T^* - \tau} \right), \quad (8)$$

the equation (1) becomes

$$\begin{aligned} \frac{\partial}{\partial t} v^{\frac{n+2}{n-2}} &= \Delta v + bv + v^{\frac{n+2}{n-2}} \quad \text{in } \Omega \times (0, \infty), \\ v &= 0 \quad \text{on } \partial\Omega \times (0, \infty). \end{aligned} \quad (9)$$

We will show that every solution of (9) converges to a stationary solution, that is a solution of

$$\Delta v + bv + v^{\frac{n+2}{n-2}} = 0 \quad \text{in } \Omega, \quad v = 0 \quad \text{on } \partial\Omega, \quad (10)$$

with at least polynomial rates. Moreover, the convergence rate will be exponential if the domain Ω satisfies the following condition:

$$\begin{aligned} \text{For every nonnegative } H_0^1 \text{ solution } v \text{ of } -\Delta v - bv - v^{\frac{n+2}{n-2}} = 0 \text{ in } \Omega, \text{ the linearized} \\ \text{operator at } v, \text{ that is } L_v := -\Delta - b - \frac{n+2}{n-2} v^{\frac{4}{n-2}}, \text{ has a trivial kernel in } H_0^1(\Omega). \end{aligned} \quad (11)$$

The set of smooth domains satisfying (11) also has generic properties, see Saut-Temam [28].

Theorem 1.1. *Let $n \geq 4$, and $b > 0$ satisfy (2). Let u be a classical nonnegative solution of (1) with extinction time $T^* > 0$. Let v be defined by (8). Then there is a nonzero stationary solution v_∞ of (9), and two positive constants θ and C such that*

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{C^2(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t \geq 1.$$

If Ω satisfies (11), then there exist two positive constants γ and C such that

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{C^2(\bar{\Omega})} \leq Ce^{-\gamma t} \quad \text{for all } t \geq 1.$$

When $n = 3$, it was shown in Brezis-Nirenberg [8] that the situation for the stationary equation (10) changes drastically from dimensions $n \geq 4$. The positivity of b is not sufficient to give a minimal energy solution of (10). Druet [19] showed that the necessary and sufficient condition is the positivity of the regular part of the Green's function of $-\Delta - b$ at a diagonal point. There should be similar changes for the parabolic equation (9) as well.

When $b = 0$, Sire-Wei-Zheng [31] recently proved the existence of some initial data such that the solution of (9) blows up at finitely many points with an explicit blow up rate as $t \rightarrow \infty$, using the gluing method for parabolic equations in the spirit of Cortázar-del Pino-Musso [13] and Dávila-del Pino-Wei [15]. This generalizes and provides rigorous proof of a result of Galaktionov-King [21] for the radially symmetric case, where the solution blows up at one point. A class of type II ancient solutions to the Yamabe flow, which are rotationally symmetric and converge to a tower of spheres

as $t \rightarrow -\infty$, was constructed by Daskalopoulos-del Pino-Sesum [14]. Bubble tower solutions for the energy critical heat equation were constructed in del Pino-Monica-Wei [16]. It is conjectured in Sire-Wei-Zheng [31] that bubble tower solutions to (9) with $b = 0$ also exist. Nevertheless, if it is the global case (Ω replaced by \mathbb{R}^n), then it has been proved by del Pino-Sáez [17] that the solution of (9) for $b = 0$ with fast decay initial data will converge to a nontrivial stationary solution, which is in fact a standard bubble.

To prove Theorem 1.1, we will adapt the blow up analysis of Struwe [32], Bahri-Coron [2], Schwetlick-Struwe [30] and Brendle [6]. See also Chen-Xu [12] and Mayer [25] for similar analysis of scalar curvature flows. Here, we define a curvature type quantity \mathcal{R} , and derive its equation along the parabolic equation (9). Due to the lack of information of \mathcal{R} on the boundary $\partial\Omega$, extra work is needed to obtain estimates for \mathcal{R} . Here the optimal boundary regularity proved in our previous paper [23] is crucial. Part of the blow up analysis in this paper remains valid when $b = 0$ or $n = 3$. The condition $n \geq 4$ and $b > 0$ is used in the final step to rule out bubbles.

Our proof of the polynomial decay rates in Theorem 1.1 can be applied to prove the polynomial rate of the convergence of the relative error for the Sobolev subcritical fast diffusion equation (5) in all smooth domains. We also provide an alternative proof the exponential convergence result of Bonforte-Figalli [4] for Ω satisfying (6).

Theorem 1.2. *Suppose $p \in (1, \frac{n+2}{n-2})$ if $n \geq 3$, and $p \in (1, \infty)$ if $n = 1, 2$. Let u be a classical nonnegative solution of (3) with extinction time $T^* > 0$. Let v be defined by (4). Then there is a stationary solution v_∞ of (5), and two positive constants θ and C such that*

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{C^2(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t \geq 1.$$

If Ω satisfies (6), then there exist two positive constants γ and C such that

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{C^2(\bar{\Omega})} \leq Ce^{-\gamma t} \quad \text{for all } t \geq 1.$$

We know from the work of Carlotto-Chodosh-Rubinstein [10] that there exists a Yamabe flow on $\mathbb{S}^1(1/\sqrt{n-2}) \times \mathbb{S}^{n-1}(1)$ such that it converges exactly at a polynomial rate. In a forthcoming paper [24], we will show examples of polynomial rate convergence of the relative error for the solutions of the fast diffusion equations (5) and (9).

This paper is organized as follows. Sections 2–5 deal with the critical equation (9). We first obtain certain integral bounds for solutions of this critical equation in section 2. Section 3 is for the possible concentration phenomenon for its solutions. In section 4, we use blow up analysis to rule out such possible concentration phenomenon. Section 5 is devoted to the proof of the uniform boundedness and convergence results in Theorem 1.1. In section 6, we consider the subcritical equation (5) and prove Theorem 1.2.

Acknowledgement: Part of this work was completed while the second named author was visiting the Hong Kong University of Science and Technology and Rutgers University, to which he is grateful for providing very stimulating research environments and supports. Both authors would like to thank Professors Haïm Brezis and YanYan Li for their interests and useful comments.

2 Integral bounds

For an open set Ω , let $H_0^1(\Omega)$ be the closure of $C_c^\infty(\Omega)$ under the norm

$$\|u\|_{H_0^1(\Omega)} := \left(\int_{\Omega} |\nabla u|^2 dx \right)^{1/2}.$$

For convenience, we define

$$\|u\| := \left(\int_{\Omega} |\nabla u|^2 - bu^2 dx \right)^{1/2} \quad (12)$$

and

$$\langle u, v \rangle = \int_{\Omega} (\nabla u \nabla v - buv) dx \quad (13)$$

be the associated inner product. Since $b < \lambda_1$, there exists a constant $K_b > 0$ such that

$$\|u\|_{L^{\frac{2n}{n-2}}} \leq K_b^{1/2} \|u\| \quad \text{for any } u \in H_0^1(\Omega). \quad (14)$$

Recall that from [23], we know that the solution $u(x, t)$ of (1) is smooth in $t \in (0, T^*)$ for every $x \in \overline{\Omega}$, and $\partial_t^l u(\cdot, t) \in C^{\frac{3n-2}{n-2}}(\overline{\Omega})$ for all $l \geq 0$ and all $t \in (0, T^*)$.

Lemma 2.1. *Let u be a solution of (1), and T^* be the extinction time of u . Then for every $0 < t < T^*$,*

$$\frac{1}{C} (T^* - t)^{\frac{n}{2}} \leq \int_{\Omega} u(x, t)^{\frac{2n}{n-2}} dx \leq (1 - t/T^*)^{\frac{n}{2}} \int_{\Omega} u_0(x)^{\frac{2n}{n-2}} dx,$$

where C is a positive constant depending only on n and Ω .

Proof. If $b = 0$, the lemma was proved by [3] (noting that by our regularity result in [23], the regularity assumptions in [3] are satisfied). The same proof applies if $b \in [0, \lambda_1)$. \square

Let v be as in (8). By Lemma 2.1, we have

$$\frac{1}{C} \leq \int_{\Omega} v(x, t)^{\frac{2n}{n-2}} dx \leq C \quad (15)$$

for some positive constant C depending only on n, Ω and u_0 . Define

$$F(v(t)) = \int_{\Omega} \left(|\nabla v(x, t)|^2 - bv(x, t)^2 - \frac{n-2}{n} v(x, t)^{\frac{2n}{n-2}} \right) dx. \quad (16)$$

It follows that $F(v(t))$ is bounded from below. By the equation of v and integrating by parts,

$$\frac{d}{dt} F(v(t)) = -2 \int_{\Omega} (\Delta v + bv + v^{\frac{n+2}{n-2}}) \partial_t v dx = -\frac{2(n+2)}{n-2} \int_{\Omega} v^{\frac{4}{n-2}} |\partial_t v|^2 dx \leq 0. \quad (17)$$

Hence, $F(v(t))$ is non-increasing in t , and thus, together with (15), we have $\|v(\cdot, t)\|_{H_0^1(\Omega)}$ is uniformly bounded. Moreover, there exists some constant F_∞ such that

$$\lim_{t \rightarrow \infty} F(v(t)) = F_\infty. \quad (18)$$

Define

$$\mathcal{R} = v^{-\frac{n+2}{n-2}}(-\Delta v - bv) \quad (19)$$

and

$$M_q(t) = \int_{\Omega} |\mathcal{R} - 1|^q v^{\frac{2n}{n-2}} dx, \quad \geq 1. \quad (20)$$

In [23], we proved that $\mathcal{R} = 1 - \frac{n+2}{n-2} \frac{\partial_t v}{v}$ is C^2 up to the boundary $\partial\Omega$. However, all the estimates there for solutions of (9) are only locally uniform in $t \in (0, \infty)$. We shall prove some uniform estimates for all $t \in [1, \infty)$ and $M_q(t) \rightarrow \infty$ as $t \rightarrow \infty$.

To do this, we will first use Moser's iteration to obtain a uniform lower bound of \mathcal{R} as an intermediate step. So we need the following evolution equation of \mathcal{R} and integration by parts formula.

Lemma 2.2. *Let $g = u^{\frac{4}{n-2}} g_{flat}$. Then*

(i).

$$\partial_t v^{\frac{2n}{n-2}} = -\frac{2n}{n+2} (\mathcal{R} - 1) v^{\frac{2n}{n-2}}. \quad (21)$$

(ii).

$$\partial_t (\mathcal{R} - 1) = \frac{n-2}{n+2} \Delta_g (\mathcal{R} - 1) + \frac{4}{n+2} (\mathcal{R} - 1)^2 + \frac{4}{n+2} (\mathcal{R} - 1), \quad (22)$$

where Δ_g is the Laplace-Beltrami operator of g .

(iii). For any $f \in H^2(\Omega)$ and $h \in H^1(\Omega)$,

$$\int_{\Omega} h \Delta_g f \, d\text{vol}_g = - \int_{\Omega} \langle \nabla_g f, \nabla_g h \rangle_g \, d\text{vol}_g. \quad (23)$$

Proof. The equation (21) follows immediately from (9) and (19). We also have $\partial_t v = \frac{n-2}{n+2} v(1 - \mathcal{R})$.

By the definition of \mathcal{R} , we have

$$\begin{aligned} \partial_t (\mathcal{R} - 1) &= \frac{n+2}{n-2} v^{-\frac{2n}{n-2}} \partial_t v (\Delta + b)v - v^{-\frac{n+2}{n-2}} (\Delta + b) \partial_t v \\ &= v^{-\frac{n+2}{n-2}} (1 - \mathcal{R})(\Delta + b)v - \frac{n-2}{n+2} v^{-\frac{n+2}{n-2}} (\Delta + b)(v(1 - \mathcal{R})) \\ &= (\mathcal{R} - 1)\mathcal{R} - \frac{n-2}{n+2} v^{-\frac{n+2}{n-2}} (\Delta + b)(v(1 - \mathcal{R})). \end{aligned}$$

Let $L_g = \Delta_g - \frac{n-2}{4(n-1)} R_g$ be the conformal Laplacian of g , where R_g is the the scalar curvature of g . By the conformal transformation law

$$L_g(v^{-1}\varphi) = v^{-\frac{n+2}{n-2}} \Delta \varphi, \quad \forall \varphi \in C^2(\Omega),$$

we have

$$\frac{n-2}{4(n-1)} R_g = -L_g(1) = -v^{-\frac{n+2}{n-2}} \Delta v = \mathcal{R} + bv^{-\frac{4}{n-2}}$$

and

$$\begin{aligned}
v^{-\frac{n+2}{n-2}}(\Delta + b)(v(1 - \mathcal{R})) &= L_g(1 - \mathcal{R}) + bv^{-\frac{4}{n-2}}(1 - \mathcal{R}) \\
&= \Delta_g(1 - \mathcal{R}) - \frac{n-2}{4(n-1)}R_g(1 - \mathcal{R}) + bv^{-\frac{4}{n-2}}(1 - \mathcal{R}) \\
&= \Delta_g(1 - \mathcal{R}) - \mathcal{R}(1 - \mathcal{R}).
\end{aligned}$$

Then, (22) follows.

Finally,

$$\begin{aligned}
\int_{\Omega} h \Delta_g f \, dvol_g &= \int_{\Omega} h v^{-\frac{2n}{n-2}} \partial_i (v^{\frac{2n}{n-2}} v^{-\frac{4}{n-2}} \partial_i f) v^{\frac{2n}{n-2}} \, dx \\
&= \int_{\Omega} h \partial_i (v^2 \partial_i f) \, dx = - \int_{\Omega} v^2 \partial_i f \partial_i h \, dx = - \int_{\Omega} \langle \nabla_g f, \nabla_g h \rangle_g \, dvol_g,
\end{aligned}$$

where we used $v = 0$ on $\partial\Omega$ in the third equality. \square

We have the following Sobolev inequality regarding the metric $g = v^{\frac{4}{n-2}} g_{flat}$:

Lemma 2.3. *There holds*

$$\left(\int_{\Omega} |f|^{\frac{2n}{n-2}} \, dvol_g \right)^{\frac{n-2}{n}} \leq K_b \int_{\Omega} (|\nabla_g f|_g^2 + \mathcal{R}f^2) \, dvol_g$$

for any $f \in H^1(\overline{\Omega})$, where K_b is the constant in (14).

Proof. Note that

$$\begin{aligned}
|\nabla(fv)| &= v^2 |\nabla f|^2 + f^2 |\nabla v|^2 + 2vf \nabla v \cdot \nabla f, \\
\int_{\Omega} (f^2 |\nabla v|^2 + 2v \nabla v f \nabla f) \, dx &= \int_{\Omega} (f^2 |\nabla v|^2 + v \nabla v \nabla f^2) \, dx \\
&= - \int_{\Omega} v f^2 \Delta v \, dx \\
&= \int_{\Omega} (\mathcal{R}f^2 v^{\frac{2n}{n-2}} + bv^2 f^2) \, dx.
\end{aligned}$$

Hence,

$$\int_{\Omega} (|\nabla_g f|_g^2 + \mathcal{R}f^2) \, dvol_g = \int_{\Omega} (v^2 |\nabla f|^2 + \mathcal{R}f^2 v^{\frac{2n}{n-2}}) \, dx = \int_{\Omega} (|\nabla(fv)| - b(fv)^2) \, dx.$$

Therefore, the lemma follows from (14). \square

For any $t_0 \geq 0$ and $T > 0$, let

$$V^1(\Omega \times (t_0, t_0 + T)) = C^0((t_0, t_0 + T); L^2(\Omega)) \cap L^2((t_0, t_0 + T); H^1(\Omega)),$$

equipped with the norm

$$\|f\|_{V^1(\Omega \times (t_0, t_0 + T))}^2 = \sup_{t_0 < t < t_0 + T} \int_{\Omega} f(x, t)^2 \, dvol_g + \int_{t_0}^{t_0 + T} \int_{\Omega} (|\nabla_g f|_g^2 + \mathcal{R}f^2) \, dvol_g \, dt.$$

We have the following parabolic version of Sobolev inequality.

Lemma 2.4. For any $f \in V^1(\Omega \times (t_0, t_0 + T))$, we have

$$\left(\int_{t_0}^{t_0+T} \int_{\Omega} |f|^{\frac{2(n+2)}{n}} \, d\text{vol}_g \, dt \right)^{\frac{n}{n+2}} \leq K_b^{\frac{n}{n+2}} \|f\|_{V^1(\Omega \times (t_0, t_0+T))}^2.$$

Proof. By Hölder's inequality and Lemma 2.3, we have

$$\begin{aligned} \int_{\Omega} |f|^{\frac{2(n+2)}{n}} \, d\text{vol}_g &= \int_{\Omega} |f|^2 |f|^{\frac{4}{n}} \, d\text{vol}_g \\ &\leq \left(\int_{\Omega} |f|^{\frac{2n}{n-2}} \, d\text{vol}_g \right)^{\frac{n-2}{n}} \left(\int_{\Omega} |f|^2 \, d\text{vol}_g \right)^{\frac{2}{n}} \\ &\leq K_b \int_{\Omega} (|\nabla_g f|_g^2 + \mathcal{R}f^2) \, d\text{vol}_g \left(\int_{\Omega} |f|^2 \, d\text{vol}_g \right)^{\frac{2}{n}}. \end{aligned}$$

Hence, by Young's inequality

$$\begin{aligned} &\left(\int_{t_0}^{t_0+T} \int_{\Omega} |f|^{\frac{2(n+2)}{n}} \, d\text{vol}_g \, dt \right)^{\frac{n}{n+2}} \\ &\leq K_b^{\frac{n}{n+2}} \left(\int_{\Omega} (|\nabla_g f|_g^2 + \mathcal{R}f^2) \, d\text{vol}_g \right)^{\frac{n}{n+2}} \left(\sup_{t_0 < t < t_0+T} \int_{\Omega} f(x, t)^2 \, d\text{vol}_g \right)^{\frac{2}{n+2}} \\ &\leq K_b^{\frac{n}{n+2}} \|f\|_{V^1(\Omega \times (t_0, t_0+T))}^2. \end{aligned}$$

Therefore, we complete the proof. \square

With the Sobolev inequality in Lemma 2.4, we will apply Moser's iterations to the equation (22) to obtain a uniform lower bound of \mathcal{R} .

Lemma 2.5. For $t \geq 1$, we have

$$\mathcal{R} - 1 \geq -C,$$

where C is a constant depending only on Ω, n, b and v_0 .

Proof. Let $T > 2$, $\frac{1}{2} \leq T_2 < T_1 \leq 1$, $\eta(t)$ be a smooth cut-off function so that $\eta(t) = 0$ for all $t < T_2$, $0 \leq \eta(t) \leq 1$ for $t \in [T_2, T_1]$, $\eta(t) = 1$ for all $t > T_1$, and $|\eta'(t)| \leq \frac{2}{T_1 - T_2}$. Denote $\phi = (1 - \mathcal{R})^+$. By (22), we have

$$\partial_t(1 - \mathcal{R}) = \frac{n-2}{n+2} \Delta_g(1 - \mathcal{R}) - \frac{4}{n+2}(1 - \mathcal{R})^2 + \frac{4}{n+2}(1 - \mathcal{R}).$$

Let $k \geq \frac{n}{2} - 1$ be a real number. Multiplying both sides of the inequality by $\eta^2 \phi^{1+k}$ and integrating by parts, we see that, for any $0 < s < T$,

$$\begin{aligned} &\frac{1}{2+k} \int_0^s \int_{\Omega} \eta^2 \partial_t \phi^{2+k} \, d\text{vol}_g \, dt + \frac{4(n-2)(k+1)}{(n+2)(k+2)^2} \int_0^s \int_{\Omega} \eta^2 |\nabla_g \phi^{\frac{k+2}{2}}|_g^2 \, d\text{vol}_g \, dt \\ &\leq -\frac{4}{n+2} \int_0^s \int_{\Omega} \phi^{3+k} \eta^2 \, d\text{vol}_g \, dt + \frac{4}{n+2} \int_0^s \int_{\Omega} \phi^{2+k} \eta^2 \, d\text{vol}_g \, dt. \end{aligned}$$

Note that using (21), we have

$$\begin{aligned}
& \frac{1}{2+k} \int_0^s \int_{\Omega} \eta^2 \partial_t \phi^{2+k} \, d\text{vol}_g dt \\
&= \frac{1}{2+k} \int_{\Omega} \phi^{2+k} \eta^2 \, d\text{vol}_g \Big|_{t=s} - \frac{1}{2+k} \int_0^s \int_{\Omega} \phi^{2+k} \left(2\eta \partial_t \eta + \frac{2n}{n+2} (1-\mathcal{R}) \eta^2 \right) \, d\text{vol}_g dt \\
&= \frac{1}{2+k} \int_{\Omega} \phi^{2+k} \eta^2 \, d\text{vol}_g \Big|_{t=s} - \frac{1}{2+k} \int_0^s \int_{\Omega} \left(2\phi^{2+k} \eta \partial_t \eta + \frac{2n}{n+2} \phi^{3+k} \eta^2 \right) \, d\text{vol}_g dt.
\end{aligned}$$

Since $k \geq \frac{n}{2} - 1$, $\frac{1}{2+k} \frac{2n}{n+2} < \frac{4}{n+2}$. Furthermore,

$$\int_{\Omega} \mathcal{R} \phi^{2+k} \eta^2 \, d\text{vol}_g = - \int_{\Omega} (1-\mathcal{R}) \phi^{2+k} \eta^2 \, d\text{vol}_g + \int_{\Omega} \phi^{2+k} \eta^2 \, d\text{vol}_g \leq \int_{\Omega} \phi^{2+k} \eta^2 \, d\text{vol}_g.$$

It follows that

$$\|\eta \phi^{\frac{2+k}{2}}\|_{V^1(\Omega \times (0, T))}^2 \leq C(2+k) \int_0^T \int_{\Omega} \phi^{2+k} (\eta^2 + |\partial_t \eta| \eta) \, d\text{vol}_g dt,$$

where $C > 0$ depends only on n . Making use of Lemma 2.4, we have for all $\gamma := k+2 \geq \frac{n+2}{2}$ that

$$\left(\int_{T_1}^T \int_{\Omega} \phi^{\frac{\gamma(n+2)}{n}} \, d\text{vol}_g dt \right)^{\frac{n}{\gamma(n+2)}} \leq \left(\frac{C\gamma}{T_1 - T_2} \right)^{\frac{1}{\gamma}} \left(\int_{T_2}^T \int_{\Omega} \phi^{\gamma} \, d\text{vol}_g dt \right)^{\frac{1}{\gamma}},$$

By the standard Moser's iteration argument, we have

$$\sup_{\Omega \times [1, T]} \phi \leq C(n, K_b) \left(\int_{1/2}^T \int_{\Omega} \phi^{\frac{n+2}{2}} \, d\text{vol}_g dt \right)^{\frac{2}{n+2}},$$

where $C(n, K_b) > 0$ depending only on n and K_b . Thus,

$$\begin{aligned}
\sup_{\Omega \times [1, T]} \phi &\leq C(n, K_b) \left(\int_{1/2}^1 \int_{\Omega} \phi^{\frac{n+2}{2}} \, d\text{vol}_g dt \right)^{\frac{2}{n+2}} + C(n, K_b) \left(\int_1^T \int_{\Omega} \phi^{\frac{n+2}{2}} \, d\text{vol}_g dt \right)^{\frac{2}{n+2}} \\
&\leq C(n, K_b) \|\mathcal{R} - 1\|_{L^\infty(\Omega \times (1/2, 1))} + C(n, K_b) \left(\int_1^T \int_{\Omega} \phi^{\frac{n+2}{2}} \, d\text{vol}_g dt \right)^{\frac{2}{n+2}},
\end{aligned}$$

By Young's inequality, we have

$$\begin{aligned}
\left(\int_1^T \int_{\Omega} \phi^{\frac{n+2}{2}} \, d\text{vol}_g dt \right)^{\frac{2}{n+2}} &\leq \left(\sup_{\Omega \times [1, T]} \phi \right)^{\frac{n-2}{n+2}} \left(\int_1^T \int_{\Omega} \phi^2 \, d\text{vol}_g dt \right)^{\frac{2}{n+2}} \\
&\leq \varepsilon \sup_{\Omega \times [1, T]} \phi + C(\varepsilon) \left(\int_1^T \int_{\Omega} \phi^2 \, d\text{vol}_g dt \right)^{\frac{1}{2}},
\end{aligned}$$

for any small constant ε . Therefore, by choosing a small ε , we have

$$\sup_{\Omega \times [1, T]} \phi \leq C(n, K_b) \left\{ \|\mathcal{R} - 1\|_{L^\infty(\Omega \times (1/2, 1))} + \left(\int_1^T M_2 \, dt \right)^{\frac{1}{2}} \right\} \quad (24)$$

By (17) and the definition of \mathcal{R} , we have

$$\frac{d}{dt}F(v(t)) = -\frac{2(n-2)}{n+2}M_2(t).$$

It follows that

$$\int_0^\infty M_2(t) dt \leq \frac{n+2}{2(n-2)}(F(v(0)) - F_\infty) < \infty. \quad (25)$$

Moreover, it was proved in [23] that $\|\mathcal{R} - 1\|_{L^\infty(\Omega \times (1/2, 1))} \leq C$. Sending $T \rightarrow \infty$ in (24), we have

$$\sup_{\Omega \times [1, \infty)} (1 - \mathcal{R})^+ \leq C.$$

Therefore, we complete the proof. \square

Using this uniform lower bound of \mathcal{R} , we can derive some useful differential inequalities for M_q defined in (20).

For $q > 1$, using Lemma 2.2, we have

$$\begin{aligned} \frac{dM_q}{dt} &= \int_\Omega q|\mathcal{R} - 1|^{q-2}(\mathcal{R} - 1)\frac{\partial}{\partial t}(\mathcal{R} - 1) dvol_g - \int_\Omega (\mathcal{R} - 1)^q \frac{\partial}{\partial t} v^{\frac{2n}{n-2}} dx \\ &= q\frac{n-2}{n+2} \int_\Omega |\mathcal{R} - 1|^{q-2}(\mathcal{R} - 1)\Delta_g(\mathcal{R} - 1) dvol_g \\ &\quad + \frac{4q}{n+2} \int_\Omega |\mathcal{R} - 1|^q dvol_g + \frac{4}{n+2}\left(q - \frac{n}{2}\right) \int_\Omega |\mathcal{R} - 1|^q(\mathcal{R} - 1) dvol_g. \end{aligned}$$

Using Lemma 2.5, we have for $t \geq 1$ that

$$\left| \int_\Omega |\mathcal{R} - 1|^q(\mathcal{R} - 1) dvol_g - \int_\Omega |\mathcal{R} - 1|^{q+1} dvol_g \right| = 2 \int_\Omega |\mathcal{R} - 1|^q(\mathcal{R} - 1)^- dvol_g \leq CM_q.$$

Using Lemma 2.2, we have

$$\int_\Omega |\mathcal{R} - 1|^{q-2}(\mathcal{R} - 1)\Delta_g(\mathcal{R} - 1) dvol_g = -\frac{4(q-1)}{q^2} \int_\Omega |\nabla_g |\mathcal{R} - 1|^{\frac{q}{2}}|_g^2 dvol_g \leq 0.$$

Therefore, for $q \leq \frac{n}{2}$ we have,

$$\frac{dM_q}{dt} + \frac{4}{n+2} \left(\frac{n}{2} - q\right) M_{q+1} \leq CM_q \quad \text{for } t \geq 1, \quad (26)$$

where $C > 0$ is a constant depending on q .

For $q \geq \frac{n}{2}$, we first obtain from Lemma 2.3 that

$$\begin{aligned} &\int_\Omega |\mathcal{R} - 1|^{q-2}(\mathcal{R} - 1)\Delta_g(\mathcal{R} - 1) dvol_g \\ &= -\frac{4(q-1)}{q^2} \int_\Omega |\nabla_g |\mathcal{R} - 1|^{\frac{q}{2}}|_g^2 dvol_g \\ &\leq -\beta M^{\frac{n-2}{qn}} + \frac{4(q-1)}{q^2} \int_\Omega \mathcal{R}|\mathcal{R} - 1|^q dvol_g \\ &\leq -\beta M^{\frac{n-2}{qn}} + \frac{4(q-1)}{q^2} \int_\Omega (\mathcal{R} - 1)|\mathcal{R} - 1|^q dvol_g + \frac{4(q-1)}{q^2} M_q, \end{aligned}$$

where $\beta > 0$ is a constant depending on K_b and q . Then, we have

$$\frac{dM_q}{dt} + \beta M_q^{\frac{n-2}{n}} \leq \frac{4}{n+2} \left(q - \frac{n}{2} + \frac{(n-2)(q-1)}{q} \right) M_{q+1} + CM_q.$$

By the interpolation inequality and Young's inequality we have

$$M_{q+1} \leq M_q^{\frac{n-2}{2q}} M_q^{\frac{2(q+1)-n}{2q}} \leq \varepsilon M_q^{\frac{n-2}{q(p+1)/2}} + C(\varepsilon) M_q^{\frac{2(q+1)-n}{2q-n}}.$$

By choosing a small ε , we obtain

$$\frac{d}{dt} M_q(t) + \beta M_q^{\frac{n-2}{n}}(t) \leq C \left(M_q(t) + M_q(t)^{1+\frac{2}{2q-n}} \right) \quad \text{for } t \geq 1 \quad (27)$$

for $q > \frac{n}{2}$, where β and C are positive constants depending on q .

The differential inequalities (26) and (27) will be used to prove the decay of M_q .

Proposition 2.6. *For every $1 \leq q < \infty$, we have*

$$\lim_{t \rightarrow \infty} M_q(t) = 0.$$

Proof. By Hölder's inequality and (15), we only need to consider $q \geq 2$.

Let us assume $n \geq 4$ first.

Case 1. $2 \leq q \leq \frac{n}{2}$.

Since $M_2 \in L^1(0, \infty)$, we can pick $t_j \rightarrow \infty$ such that $M_2(t_j) \rightarrow 0$ as $j \rightarrow \infty$. By (26) we have

$$\frac{d}{dt} M_2(t) \leq CM_2(t).$$

Integrating the above inequality we have

$$M_2(t) \leq M_2(t_j) + C \int_{t_j}^{\infty} M_2(s) ds \quad \text{for } t \geq t_j.$$

Hence, $\lim_{t \rightarrow \infty} M_2(t) = 0$. If $2 < q \leq \frac{n}{2}$, (26) we have

$$\int_1^{\infty} M_3(t) dt \leq C \left(\int_1^{\infty} M_2(t) dt + M_2(1) \right) < \infty.$$

For any $2 < q \leq \min\{3, \frac{n}{2}\}$, we have $M_q(t) \leq M_2(t) + M_3(t)$. Hence, $\int_1^{\infty} M_q(t) dt < \infty$. We can repeat the argument for M_2 to show that $M_q(t) \rightarrow 0$ as $t \rightarrow \infty$. If $3 < \frac{n}{2}$, we can show that $\int_1^{\infty} M_4 < \infty$ and $M_q(t) \rightarrow 0$ as $t \rightarrow \infty$ for all $3 < q \leq \min\{4, \frac{n}{2}\}$. Repeating this argument in finite times, we then have $M_q \in L^1(1, \infty)$ and $M_q(t) \rightarrow 0$ as $t \rightarrow \infty$ for all $2 \leq q \leq \frac{n}{2}$.

Case 2. $q > \max\{2, \frac{n}{2}\}$.

By (26) with $q = n/2$, we have

$$\int_1^{\infty} M_q^{\frac{n-2}{2(n-2)}}(t) dt < \infty. \quad (28)$$

Using (27) to have

$$\frac{d}{dt}M_q(t) \leq CM_q(t)^{\frac{n-2}{n}} \left(M_q(t)^{\frac{2}{n}} + M_q(t)^{\frac{2}{n} + \frac{2}{2q-n}} \right) \quad \text{for } t \geq 1$$

Hence,

$$H(M_q(t)) \leq H(M_q(T)) + C \int_T^\infty M_q^{\frac{n-2}{n}} dt \quad \text{for } 1 \leq T < t < \infty$$

where

$$H(\rho) = \int_0^\rho \frac{1}{s^{\frac{2}{n}} + s^{\frac{2}{n} + \frac{2}{2q-n}}} ds.$$

Let

$$q_0 = \frac{n^2}{2(n-2)}, \quad q_k = \frac{n}{n-2}q_{k-1}, \quad k = 1, 2, \dots$$

Note that $\frac{2}{n} + \frac{2}{2q_0-n} = 1$ and $\frac{2}{n} + \frac{2}{2q_k-n} < 1$ for all $k \geq 1$. Hence, starting with (28) that $\int_1^\infty M_{q_0}(t)^{\frac{n-2}{n}} dt < \infty$, using similar arguments to those in Case 1, we can recursively prove in the order of $k = 0, 1, 2, \dots$ that $M_{q_k}(t) \rightarrow 0$ as $t \rightarrow \infty$, $\int_1^\infty \left(M_{q_k}(t) + M_{q_k}(t)^{1 + \frac{2}{2q_k-n}} \right) < \infty$, $\int_1^\infty M_{q_{k+1}}(t)^{\frac{n-2}{n}} dt < \infty$, and $M_{q_{k+1}}(t) \rightarrow 0$ as $t \rightarrow \infty$. Hence, $M_q(t) \rightarrow 0$ for any $q \geq \frac{n^2}{2(n-2)}$.

Finally, let us consider $n = 3$. By (27), we have

$$\frac{d}{dt}M_2(t) \leq CM_2(t)(1 + M_2(t)^2).$$

Using (25), we can pick $t_i \rightarrow \infty$ such that $M_2(t_i) \rightarrow 0$. Hence,

$$\arctan M_2(t) \leq \arctan M_2(t_i) + C \int_{t_i}^\infty M_2(t) dt.$$

It follows that $\lim_{t \rightarrow \infty} \arctan M_2(t) = 0$ and thus $\lim_{t \rightarrow \infty} M_2(t) = 0$. Hence, $\int_1^\infty M_6(t)^{\frac{1}{3}} dt < \infty$. Since $6 > q_0$ when $n = 3$, we can use the argument of those in Case 2 to show that $M_q(t) \rightarrow 0$ for all $q \geq 6$. By Hölder inequality, we conclude that $M_q(t) \rightarrow 0$ for all $q \geq 1$. \square

Corollary 2.7. *We have*

$$\lim_{t \rightarrow \infty} \|\mathcal{R} - 1\|_{L^\infty(\Omega)} = 0.$$

Proof. We know from the proof of Proposition 2.6 that there exists a sufficiently large $q > 1$ such that

$$\int_1^\infty M_q(t) dt < \infty.$$

This means that $\mathcal{R} - 1$ has very high integrability in space-time. Then we can apply the Moser's iteration as in the proof of Lemma 2.5 to obtain

$$\|\mathcal{R} - 1\|_{L^\infty(\Omega \times (T, \infty))} \leq C \left(\int_{T-1}^\infty M_q(t) dt \right)^{\frac{1}{q}}$$

for all large T . Hence, the corollary follows. \square

3 Concentration compactness

The solution of (9) may blow up as $t \rightarrow \infty$ because of the critical exponent $\frac{n+2}{n-2}$. Nevertheless, we also know how the solutions may blow up.

Proposition 3.1. *Let v be a solution of (9). For any $t_\nu \rightarrow \infty$, $\nu \rightarrow \infty$, $v_\nu = v(\cdot, t_\nu)$ is a Palais-Smale sequence of the functional F given by (16) in $H_0^1(\Omega)$.*

Proof. We have already proved that v_ν is bounded in $H_0^1(\Omega)$ and $F(v_\nu) \rightarrow F_\infty$ as $\nu \rightarrow \infty$. It remains to show the derivative of F at v_ν tends to zero. Indeed, for any $\varphi \in H_0^1(\Omega)$, we have

$$\begin{aligned} \langle dF(v_\nu), \varphi \rangle &= 2 \int_{\Omega} (-\Delta v_\nu - b v_\nu - v_\nu^{\frac{n+2}{n-2}}) \varphi \, dx \\ &= 2 \int_{\Omega} (\mathcal{R} - 1) v_\nu^{\frac{n+2}{n-2}} \varphi \, dx \\ &\leq 2 \left(\int_{\Omega} |\mathcal{R} - 1|^{\frac{2n}{n+2}} v_\nu^{\frac{2n}{n-2}} \, dx \right)^{\frac{n+2}{2n}} \left(\int_{\Omega} |\varphi|^{\frac{2n}{n-2}} \, dx \right)^{\frac{n-2}{2n}} \\ &\leq C(n) M_{\frac{2n}{n+2}}(t_\nu)^{\frac{n+2}{2n}} \|\varphi\|_{H_0^1(\Omega)}, \end{aligned}$$

where we used Hölder's inequality and the Sobolev inequality. It follows from Proposition 2.6 that $dF(v_\nu)$ strongly converges to 0 in $H^{-1}(\Omega)$.

Therefore, we complete the proof. \square

The next proposition shows that the blow up points, if exist, will stay uniformly away from the boundary $\partial\Omega$.

Proposition 3.2. *There exist two positive constants δ_0 and C , depending on $v(\cdot, 1)$, such that for all $x \in \Omega$ with $d(x) := \text{dist}(x, \partial\Omega) < \delta_0$ and $t \geq 1$,*

$$v(x, t) \leq C d(x).$$

Proof. We are going to use the moving plane method as Han [22] did for the elliptic case. By the Hopf Lemma, there exist $\rho_0 > 0$ and $\alpha_0 > 0$ such that $v(z - \rho e, 1)$ is nondecreasing for $0 < \rho < \rho_0$, where $z \in \partial\Omega$, $e \in \mathbb{R}^n$ with $|e| = 1$, and $(e, \nu(z)) \geq \alpha_0$ with $\nu(z)$ the unit out normal to $\partial\Omega$ at z . If Ω is strictly convex, using the moving plane method we can conclude that $v(z - \rho e, t)$ is nondecreasing for $0 < \rho < \rho_0$, and for all $t \geq 1$. Therefore, we can find $\gamma > 0$ and $\delta > 0$ such that for any fixed $t \geq 1$, and any $x \in \Omega$ satisfying $0 < d(x) < \delta$, there exists a measurable set Γ_x with (i) $\text{meas}(\Gamma_x) \geq \gamma$, (ii) $\Gamma_x \subset \{z : d(z) \geq \delta/2\}$, and (iii) $v(y, t) \geq v(x, t)$ for any $y \in \Gamma_x$. Actually, Γ_x can be taken to a piece of cone with vertex at x . It follows that for any $x \in \{z : 0 < d(z) < \delta\}$, we have

$$v(x, t) \leq \frac{1}{\text{meas}(\Gamma_x)} \int_{\Gamma_x} v(y, t) \, dy \leq \frac{C}{\gamma},$$

where we used (15) and Hölder's inequality. Namely, $v(x, t) \leq C$ for $(x, t) \in \{z : 0 < d(z) < \delta\} \times [1, \infty)$. By the proof of Theorem 4.1 in [18], we have $v(x, t) \leq C d(x)$ for $(x, t) \in \{z : 0 < d(z) < \delta\} \times [1, \infty)$.

For a general domain, one can first use a Kelvin transform near each boundary point, and then apply the moving plane method. Pick any point $P \in \partial\Omega$ for instance. Since we assume the boundary

of the domain Ω is smooth, we may assume, without loss of generality, that the unit ball B_1 contacts P from the exterior of Ω . Let $w(x, t)$ be the Kelvin transform of v :

$$w(x, t) = |x|^{2-n} v\left(\frac{x}{|x|^2}, t\right).$$

Then

$$\begin{cases} \partial_t w^{\frac{n+2}{n-2}} = \Delta w + b|x|^{-4}w + w^{\frac{n+2}{n-2}} & \text{in } \Omega_P \times (0, \infty) \\ w = 0 & \text{on } \partial\Omega_P \times (0, \infty), \end{cases}$$

where Ω_P is the image of Ω under the Kelvin transform. Since $b \geq 0$, $b|x|^{-4}$ is nondecreasing along the $-P$ direction. Applying the moving plane method we have that $w(\cdot, t)$ is nondecreasing along the $-P$ direction in a neighborhood (uniform in t) of P . Since the $L^{\frac{2n}{n-2}}$ norm is invariant under the Kelvin transform, using the above argument in the case of strictly convex domains, we conclude that $w(\cdot, t)$ is bounded in a neighborhood of P independent of t and so is $v(\cdot, t)$. It follows that $v(x, t) \leq Cd(x)$ for $(x, t) \in \{z : 0 < d(z) < \delta\} \times [1, \infty)$ for some $\delta > 0$.

Therefore, we complete the proof. \square

For $a \in \mathbb{R}^n$ and $\lambda \in (0, \infty)$, let

$$\bar{\xi}_{a,\lambda}(x) = c_0 \left(\frac{\lambda}{1 + \lambda^2|x - a|^2} \right)^{\frac{n-2}{2}} \quad (29)$$

with $c_0 = (n(n-2))^{\frac{n-2}{4}}$. Then we have

$$-\Delta \bar{\xi}_{a,\lambda} = \bar{\xi}_{a,\lambda}^{\frac{n+2}{n-2}} \quad \text{in } \mathbb{R}^n$$

and

$$\int_{\mathbb{R}^n} \bar{\xi}_{a,\lambda}^{\frac{2n}{n-2}} = Y(\mathbb{S}^n)^{\frac{n}{2}},$$

where \mathbb{S}^n is the standard unit sphere in \mathbb{R}^{n+1} ,

$$Y(\mathbb{S}^n) = \frac{n(n-2)}{4} |\mathbb{S}^n|^{\frac{2}{n}} = \inf_{u \in H^1(\mathbb{S}^n)} \frac{\int_{\mathbb{S}^n} |\nabla u|^2 + \frac{n(n-2)}{4} u^2 \, d\text{vol}_{g_{\mathbb{S}^n}}}{\left(\int_{\mathbb{S}^n} |u|^{\frac{2n}{n-2}} \, d\text{vol}_{g_{\mathbb{S}^n}} \right)^{\frac{n-2}{n}}}$$

and $|\mathbb{S}^n|$ is the area of \mathbb{S}^n . Define

$$\xi_{a,\lambda}(x) = \bar{\xi}_{a,\lambda}(x) - h_{a,\lambda}(x), \quad (30)$$

where $\Delta h_{a,\lambda}(x) = 0$ in Ω and $h_{a,\lambda} = \bar{\xi}_{a,\lambda}$ on $\partial\Omega$. By the maximum principle, $\xi_{a,\lambda} > 0$ in Ω and $h_{a,\lambda} > 0$ in $\bar{\Omega}$.

Proposition 3.3. *Let v be a solution of (9). For any $t_\nu \rightarrow \infty$, $\nu \rightarrow \infty$, after passing to a subsequence if necessary, v_ν weakly converges to v_∞ in $H_0^1(\Omega)$ and we can find a non-negative integer m and a sequence of m -tuplets $(x_{k,\nu}^*, \lambda_{k,\nu}^*)_{1 \leq k \leq m}$, $(x_{k,\nu}^*, \lambda_{k,\nu}^*) \in \Omega \times (0, \infty)$, with the following properties.*

- The function $v_\infty \in H_0^1(\Omega)$ satisfies the equation $-\Delta v_\infty - b v_\infty = v_\infty^{\frac{n+2}{n-2}}$ in Ω .

- There hold, for all $i \neq j$,

$$\frac{\lambda_{i,\nu}^*}{\lambda_{j,\nu}^*} + \frac{\lambda_{j,\nu}^*}{\lambda_{i,\nu}^*} + \lambda_{i,\nu}^* \lambda_{j,\nu}^* |x_{i,\nu}^* - x_{j,\nu}^*|^2 \rightarrow \infty,$$

and for all k

$$\lambda_{k,\nu}^* d(x_{k,\nu}^*) \rightarrow \infty$$

as $\nu \rightarrow \infty$.

- We have

$$\left\| v_\nu - v_\infty - \sum_{k=1}^m \xi_{x_{k,\nu}^*, \lambda_{k,\nu}^*} \right\| \rightarrow 0$$

as $\nu \rightarrow \infty$.

- We have

$$F(v_\nu) = F(v_\infty) + \frac{2m}{n} Y(\mathbb{S}^n)^{n/2} + o(1),$$

where $o(1) \rightarrow 0$ as $\nu \rightarrow \infty$.

Proof. In view of Propositions 3.1 and 3.2, this proposition follows from Struwe [32] and Bahri-Coron [2]. \square

A similar result for the harmonic map heat flow was proved by Qing-Tian [27]. The correction term $h_{a,\lambda}$ in (30) is small and can be controlled.

Lemma 3.4. *Let $\xi_{a,\lambda}$ and $h_{a,\lambda}$ be defined as in (30). Suppose $a \in \Omega$ with $d(a) > \delta > 0$ and $\lambda > 1$. Then we have*

$$|h_{a,\lambda}(x)| + |\nabla_a h_{a,\lambda}(x)| + \lambda |\nabla_\lambda h_{a,\lambda}(x)| \leq C(n, \Omega, \delta) \lambda^{-\frac{n-2}{2}}, \quad x \in \Omega,$$

$$\nabla_a \xi_{a,\lambda}(x) = (n-2) \xi_{a,\lambda} \frac{\lambda^2(x-a)}{1 + \lambda^2|x-a|^2} + O(\lambda^{-\frac{n-2}{2}}),$$

and

$$\nabla_\lambda \xi_{a,\lambda}(x) = \frac{(n-2)}{2\lambda} \xi_{a,\lambda} \frac{1 - \lambda^2|x-a|^2}{1 + \lambda^2|x-a|^2} + O(\lambda^{-\frac{n}{2}}),$$

where $|O(\lambda^{-\frac{n-2}{2}})| \leq C \lambda^{-\frac{n-2}{2}}$ for some C depending only on n, Ω and δ .

Proof. It suffices to prove the estimate of $h_{a,\lambda}$, which follows from Poisson formula for the Laplace equation. \square

4 Refined blow up analysis

We continue from Proposition 3.3. By the strong maximum principle, the nonnegative limit v_∞ either is positive in Ω or identically equals to zero. We will adapt the refined blow up analysis in Brendle [6] to the equation (9).

4.1 The case $v_\infty \equiv 0$

For every ν , let \mathcal{A}_ν be the closed set of all m -tuplets $(x_k, \lambda_k, \alpha_k)_{1 \leq k \leq m}$ satisfying $(x_k, \lambda_k, \alpha_k) \in \overline{B}_{\frac{1}{\lambda_{k,\nu}^*}}(x_{k,\nu}^*) \times [\frac{\lambda_{k,\nu}^*}{2}, \frac{3\lambda_{k,\nu}^*}{2}] \times [\frac{1}{2}, \frac{3}{2}]$. Choose an m -tuple $(x_{k,\nu}, \lambda_{k,\nu}, \alpha_{k,\nu})_{1 \leq k \leq m} \in \mathcal{A}_\nu$ such that

$$\left\| v_\nu - \sum_{k=0}^m \alpha_{k,\nu} \xi_{x_{k,\nu}, \lambda_{k,\nu}} \right\| = \inf_{(x_k, \lambda_k, \alpha_k)_{1 \leq k \leq m} \in \mathcal{A}_\nu} \left\| v_\nu - \sum_{k=0}^m \alpha_k \xi_{x_k, \lambda_k} \right\|. \quad (31)$$

By Proposition 3.3, Proposition 3.2 and the definition of $(x_{k,\nu}, \lambda_{k,\nu}, \alpha_{k,\nu})_{1 \leq k \leq m}$, we have, for all $i \neq j$,

$$\frac{\lambda_{i,\nu}}{\lambda_{j,\nu}} + \frac{\lambda_{j,\nu}}{\lambda_{i,\nu}} + \lambda_{i,\nu} \lambda_{j,\nu} |x_{i,\nu} - x_{j,\nu}|^2 \rightarrow \infty, \quad (32)$$

and for all k

$$\lambda_{k,\nu} d(x_{k,\nu}) \rightarrow \infty \quad (33)$$

as $\nu \rightarrow \infty$. In addition, $d(x_{k,\nu}) > \delta/2$, and

$$\left\| v_\nu - \sum_{k=1}^m \alpha_k \xi_{x_{k,\nu}, \lambda_{k,\nu}} \right\| \rightarrow 0 \quad (34)$$

as $\nu \rightarrow \infty$.

By the triangle inequality,

$$\begin{aligned} & \left\| \sum_{k=1}^m \alpha_k \xi_{x_{k,\nu}, \lambda_{k,\nu}} - \sum_{k=1}^m \xi_{x_{k,\nu}^*, \lambda_{k,\nu}^*} \right\| \\ & \leq \left\| v_\nu - \sum_{k=1}^m \alpha_k \xi_{x_{k,\nu}, \lambda_{k,\nu}} \right\| + \left\| v_\nu - \sum_{k=1}^m \xi_{x_{k,\nu}^*, \lambda_{k,\nu}^*} \right\| = o(1). \end{aligned}$$

It follows that, for all $1 \leq k \leq m$,

$$|x_{k,\nu} - x_{k,\nu}^*| = o(1) \frac{1}{\lambda_{k,\nu}^*}, \quad \frac{\lambda_{k,\nu}}{\lambda_{k,\nu}^*} = 1 + o(1), \quad \alpha_{k,\nu} = 1 + o(1). \quad (35)$$

In particular, $(x_{k,\nu}, \lambda_{k,\nu}, \alpha_{k,\nu})_{1 \leq k \leq m}$ is an interior point of \mathcal{A}_ν .

In the sequel, we assume

$$\lambda_{1,\nu} \geq \lambda_{2,\nu} \geq \cdots \geq \lambda_{m,\nu}. \quad (36)$$

Let

$$U_\nu = \sum_{k=1}^m \alpha_{k,\nu} \xi_{x_{k,\nu}, \lambda_{k,\nu}}, \quad w_\nu = v_\nu - U_\nu. \quad (37)$$

Lemma 4.1. *We have for $1 \leq k \leq m$,*

$$\begin{aligned} & \left| \int_\Omega \xi_{x_{k,\nu}, \lambda_{k,\nu}}^{\frac{n+2}{n-2}} w_\nu \, dx \right| + \left| \int_\Omega \xi_{x_{k,\nu}, \lambda_{k,\nu}}^{\frac{n+2}{n-2}} \frac{1 - \lambda^2 |x - x_{k,\nu}|^2}{1 + \lambda^2 |x - x_{k,\nu}|^2} w_\nu \, dx \right| \\ & + \left| \int_\Omega \xi_{x_{k,\nu}, \lambda_{k,\nu}}^{\frac{n+2}{n-2}} \frac{\lambda^2 (x - x_{k,\nu})}{1 + \lambda^2 |x - x_{k,\nu}|^2} w_\nu \, dx \right| \leq o(1) \left(\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \, dx \right)^{\frac{n-2}{2n}} \end{aligned}$$

Proof. By the finite dimensional variational problem (31) and (35), taking derivatives in \mathcal{A}_ν , we have

$$\int_{\Omega} \left[\nabla(\nabla_{a,\lambda} \xi_{x_k,\nu,\lambda_k,\nu}) \nabla w_\nu - b \nabla_{a,\lambda} \xi_{x_k,\nu,\lambda_k,\nu} w_\nu \right] dx = 0$$

and

$$\int_{\Omega} \left[\nabla \xi_{x_k,\nu,\lambda_k,\nu} \nabla w_\nu - b \xi_{x_k,\nu,\lambda_k,\nu} w_\nu \right] dx = 0,$$

where $\nabla_{a,\lambda} \xi_{x_k,\nu,\lambda_k,\nu} = \nabla_{a,\lambda} \xi_{a,\lambda} \Big|_{(a,\lambda)=(x_k,\nu,\lambda_k,\nu)}$. Integrating by parts, using the equation of $\bar{\xi}_{a,\lambda}$, Hölder's inequality and Lemma 3.4, the lemma follows. \square

Note that the bubbles are non-degenerate, since we have the following well known lemma (see (3.14) in Rey [26]).

Lemma 4.2. *Let $\bar{\xi}_{a,\lambda}$ be defined in (29). Then there exists a constant $c_1 > 0$ such that*

$$(1 - c_1) \int_{\mathbb{R}^n} |\nabla \varphi|^2 \geq \frac{n+2}{n-2} \int_{\mathbb{R}^n} \bar{\xi}_{0,1}^{\frac{4}{n-2}} \varphi^2$$

for any $\varphi \in H_0^1(\mathbb{R}^n)$ satisfying

$$\int_{\mathbb{R}^n} \bar{\xi}_{0,1}^{\frac{4}{n-2}} (\nabla_{a,\lambda} \bar{\xi}_{0,1}) \varphi dx = 0.$$

We have the following non-degeneracy estimates of the second variation of F for w_ν .

Lemma 4.3. *For large ν , we have*

$$\frac{n+2}{n-2} \int_{\Omega} \sum_{k=1}^m \xi_{x_k,\nu,\lambda_k,\nu}^{\frac{4}{n-2}} w_\nu^2 \leq (1-c) \int_{\Omega} (|\nabla w_\nu|^2 - b w_\nu^2) dx,$$

where $c > 0$ is independent of ν .

Proof. We assume w_ν is not zero, otherwise there is nothing to prove. Define $\tilde{w}_\nu = \frac{w_\nu}{\|w_\nu\|}$. Suppose the lemma is not true. Then we can find a subsequence of $\{\tilde{w}_\nu\}$ (still denoted by $\{\tilde{w}_\nu\}$) satisfying

$$\lim_{\nu \rightarrow \infty} \frac{n+2}{n-2} \int_{\Omega} \sum_{k=1}^m \xi_{x_k,\nu,\lambda_k,\nu}^{\frac{4}{n-2}} \tilde{w}_\nu^2 \geq 1. \quad (38)$$

By (14),

$$\int_{\Omega} |\tilde{w}_\nu|^{\frac{2n}{n-2}} \leq K_b^{\frac{n}{n-2}} \|\tilde{w}_\nu\| = K_b^{\frac{n}{n-2}}. \quad (39)$$

By (32) and (36), we can find $R_\nu \rightarrow \infty$, $R_\nu \lambda_{j,\nu}^{-1} \rightarrow 0$ for all $1 \leq j \leq m$, and

$$\frac{\lambda_{i,\nu}}{R_\nu} (\lambda_{j,\nu}^{-1} + |x_{i,\nu} - x_{j,\nu}|) \rightarrow \infty \quad (40)$$

for all $i < j$. Set

$$\Omega_{j,\nu} = B_{R_\nu \lambda_{j,\nu}^{-1}}(x_{j,\nu}) \setminus \bigcup_{i=1}^{j-1} B_{R_\nu \lambda_{i,\nu}^{-1}}(x_{i,\nu}).$$

By (38) and $\|\tilde{w}_\nu\| = 1$, we can find $1 \leq j \leq m$ such that

$$\lim_{\nu \rightarrow \infty} \int_{\Omega} \xi_{x_{j,\nu}, \lambda_{j,\nu}}^{\frac{4}{n-2}} \tilde{w}_\nu^2 > 0$$

and

$$\lim_{\nu \rightarrow \infty} \int_{\Omega_{j,\nu}} (|\nabla \tilde{w}_\nu|^2 - b\tilde{w}_\nu^2) \leq \lim_{\nu \rightarrow \infty} \frac{n+2}{n-2} \int_{\Omega} \xi_{x_{j,\nu}, \lambda_{j,\nu}}^{\frac{4}{n-2}} \tilde{w}_\nu^2.$$

Let $\hat{w}_\nu(x) = \lambda_{j,\nu}^{-\frac{n-2}{2}} \tilde{w}_\nu(x_{j,\nu} + \lambda_{j,\nu}^{-1}x)$. Under this scaling, by using (40), we know that either $B_{R_\nu \lambda_{j,\nu}^{-1}}(x_{j,\nu})$ will be disjoint with $\bigcup_{i=1}^{j-1} B_{R_\nu \lambda_{i,\nu}^{-1}}(x_{i,\nu})$, or $B_{R_\nu \lambda_{i,\nu}^{-1}}(x_{i,\nu})$ will shrink to a point for every $1 \leq i \leq j-1$. By passing to a weak limit in $H_{loc}^1(\mathbb{R}^n)$, and using the above two inequalities and Lemma 4.1, we then obtain a contradiction to Lemma 4.2.

Therefore, Lemma 4.3 is proved. \square

Corollary 4.4. *For large ν , we have*

$$\frac{n+2}{n-2} \int_{\Omega} U_\nu^{\frac{4}{n-2}} w_\nu^2 \leq (1-c) \int_{\Omega} (|\nabla w_\nu|^2 - b w_\nu^2) dx,$$

where $c > 0$ is independent of ν .

Proof. It follows from Lemma 4.3, Hölder's inequality, the Sobolev inequality (14) and the fact that

$$\int_{\Omega} \left| U_\nu^{\frac{4}{n-2}} - \sum_{k=1}^m \xi_{x_{k,\nu}, \lambda_{k,\nu}}^{\frac{4}{n-2}} \right|^{\frac{n}{2}} = o(1).$$

\square

Now we can have an expansion of the Hamiltonian F defined in (16).

Proposition 4.5. *When $n \geq 4$ and ν is sufficiently large, we have*

$$F(U_\nu) \leq \sum_{k=1}^m F(\xi_{x_{k,\nu}, \lambda_{k,\nu}}) + o(1) \sum_{k=1}^m \int_{\Omega} \xi_{x_{k,\nu}, \lambda_{k,\nu}}^2 + C \sum_{k=1}^m \lambda_{k,\nu}^{2-n}.$$

Proof. We shall need the following inequality

$$\left(\sum_{k=1}^m a_k \right)^{\frac{2n}{n-2}} \geq \sum_{k=1}^m a_k^{\frac{2n}{n-2}} + \frac{2n}{n-2} \sum_{k < l} a_k^{\frac{n+2}{n-2}} a_l + c_{n,m} \sum_{k < l} (a_k \vee a_l)^{\frac{4}{n-2}} (a_k \wedge a_l)^2 \quad (41)$$

for any $a_1, \dots, a_m \geq 0$, where $c_{n,m} > 0$ is a constant, and $a_k \vee a_l = \max(a_k, a_l)$ and $a_k \wedge a_l = \min(a_k, a_l)$. This inequality can be proved using Lemma A.1 and induction.

Using the inequality (41), we have

$$\begin{aligned}
& \int_{\Omega} (|\nabla U_{\nu}|^2 - bU_{\nu}^2) - \frac{n-2}{n} \int_{\Omega} U_{\nu}^{\frac{2n}{n-2}} \\
& \leq \sum_k \alpha_{k,\nu}^2 \int_{\Omega} (|\nabla \xi_{x_k,\nu,\lambda_{k,\nu}}|^2 - b\xi_{x_k,\nu,\lambda_{k,\nu}}^2) - \sum_k \alpha_{k,\nu}^{\frac{2n}{n-2}} \frac{n-2}{n} \int_{\Omega} \xi_{x_k,\nu,\lambda_{k,\nu}}^{\frac{2n}{n-2}} \\
& \quad + 2 \sum_{i<j} \alpha_{j,\nu} \left[\alpha_{i,\nu} \int_{\Omega} (\nabla \xi_{x_i,\nu,\lambda_{i,\nu}} \nabla \xi_{x_j,\nu,\lambda_{j,\nu}} - b\xi_{x_i,\nu,\lambda_{i,\nu}} \xi_{x_j,\nu,\lambda_{j,\nu}}) - \alpha_{i,\nu}^{\frac{n+2}{n-2}} \int_{\Omega} \xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \xi_{x_j,\nu,\lambda_{j,\nu}} \right] \\
& \quad - c_{n,m} \sum_{i<j} \int_{\Omega} (\xi_{x_i,\nu,\lambda_{i,\nu}} \vee \xi_{x_j,\nu,\lambda_{j,\nu}})^{\frac{4}{n-2}} (\xi_{x_i,\nu,\lambda_{i,\nu}} \wedge \xi_{x_j,\nu,\lambda_{j,\nu}})^2. \tag{42}
\end{aligned}$$

By the equation of $\bar{\xi}_{x_k,\nu,\lambda_{k,\nu}}$ and the definition of $\xi_{x_k,\nu,\lambda_{k,\nu}}$, we have

$$\begin{aligned}
& \alpha_{k,\nu}^2 \int_{\Omega} (|\nabla \xi_{x_k,\nu,\lambda_{k,\nu}}|^2 - b\xi_{x_k,\nu,\lambda_{k,\nu}}^2) - \alpha_{k,\nu}^{\frac{2n}{n-2}} \frac{n-2}{n} \int_{\Omega} \xi_{x_k,\nu,\lambda_{k,\nu}}^{\frac{2n}{n-2}} \\
& \leq \alpha_{k,\nu}^2 \int_{\mathbb{R}^n} |\nabla \bar{\xi}_{x_k,\nu,\lambda_{k,\nu}}|^2 - \alpha_{k,\nu}^{\frac{2n}{n-2}} \frac{n-2}{n} \int_{\mathbb{R}^n} \bar{\xi}_{x_k,\nu,\lambda_{k,\nu}}^{\frac{2n}{n-2}} - b\alpha_{k,\nu}^2 \int_{\Omega} \xi_{x_k,\nu,\lambda_{k,\nu}}^2 + C\lambda_{k,\nu}^{2-n} \\
& \leq \int_{\mathbb{R}^n} |\nabla \bar{\xi}_{x_k,\nu,\lambda_{k,\nu}}|^2 - \frac{n-2}{n} \int_{\mathbb{R}^n} \bar{\xi}_{x_k,\nu,\lambda_{k,\nu}}^{\frac{2n}{n-2}} - \frac{4}{n-2} (\alpha_{k,\nu} - 1)^2 Y(\mathbb{S}^n)^{\frac{n}{2}} \\
& \quad - b\alpha_{k,\nu}^2 \int_{\Omega} \xi_{x_k,\nu,\lambda_{k,\nu}}^2 + C\lambda_{k,\nu}^{2-n} \\
& \leq F(\xi_{x_k,\nu,\lambda_{k,\nu}}) - \frac{4}{n-2} (\alpha_{k,\nu} - 1)^2 Y(\mathbb{S}^n)^{\frac{n}{2}} + o(1) \int_{\Omega} \xi_{x_k,\nu,\lambda_{k,\nu}}^2 + C\lambda_{k,\nu}^{2-n}, \tag{43}
\end{aligned}$$

where we used $\alpha_{k,\nu} = 1 + o(1)$, $\alpha_{k,\nu}^2 - \frac{n-2}{n} \alpha_{k,\nu}^{\frac{2n}{n-2}} \leq \frac{2}{n} - \frac{4}{n-2} (\alpha_{k,\nu} - 1)^2$ and

$$\int_{\mathbb{R}^n} |\nabla \bar{\xi}_{x_k,\nu,\lambda_{k,\nu}}|^2 = \int_{\mathbb{R}^n} \bar{\xi}_{x_k,\nu,\lambda_{k,\nu}}^{\frac{2n}{n-2}} = Y(\mathbb{S}^n)^{\frac{n}{2}}.$$

In addition,

$$\begin{aligned}
& \alpha_{i,\nu} \int_{\Omega} (\nabla \xi_{x_i,\nu,\lambda_{i,\nu}} \nabla \xi_{x_j,\nu,\lambda_{j,\nu}} - b\xi_{x_i,\nu,\lambda_{i,\nu}} \xi_{x_j,\nu,\lambda_{j,\nu}}) - \alpha_{i,\nu}^{\frac{n+2}{n-2}} \int_{\Omega} \xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \xi_{x_j,\nu,\lambda_{j,\nu}} \\
& = \alpha_{i,\nu} \int_{\Omega} (-\Delta \xi_{x_i,\nu,\lambda_{i,\nu}} - b\xi_{x_i,\nu,\lambda_{i,\nu}} - \alpha_{i,\nu}^{\frac{4}{n-2}} \xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}}) \xi_{x_j,\nu,\lambda_{j,\nu}} \\
& \leq C|\alpha_{i,\nu} - 1| \int_{\Omega} \xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \xi_{x_j,\nu,\lambda_{j,\nu}} + \int_{\Omega} \left(\xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}} - \xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \right) \xi_{x_j,\nu,\lambda_{j,\nu}} \\
& \leq C|\alpha_{i,\nu} - 1| \int_{\Omega} \xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \xi_{x_j,\nu,\lambda_{j,\nu}} + O(\lambda_{i,\nu}^{2-n} + \lambda_{j,\nu}^{2-n}) \\
& \leq \frac{2}{n-2} (\alpha_{k,\nu} - 1)^2 Y(\mathbb{S}^n)^{\frac{n}{2}} + C \left(\int_{\Omega} \xi_{x_i,\nu,\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \xi_{x_j,\nu,\lambda_{j,\nu}} \right)^2 + O(\lambda_{i,\nu}^{2-n} + \lambda_{j,\nu}^{2-n}), \tag{44}
\end{aligned}$$

where $C > 0$ is independent of ν , and in the second inequality we used

$$\begin{aligned}
\int_{\Omega} \left(\xi_{x_{i,\nu},\lambda_{i,\nu}}^{\frac{n+2}{n-2}} - \bar{\xi}_{x_{i,\nu},\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \right) \xi_{x_{j,\nu},\lambda_{j,\nu}} &\leq C \int_{\Omega} \bar{\xi}_{x_{i,\nu},\lambda_{i,\nu}}^{\frac{4}{n-2}} |h_{x_{i,\nu},\lambda_{i,\nu}}| \bar{\xi}_{x_{j,\nu},\lambda_{j,\nu}} \\
&\leq C \lambda_{i,\nu}^{\frac{2-n}{2}} \left(\int_{\Omega} \bar{\xi}_{x_{i,\nu},\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \right)^{\frac{4}{n+2}} \left(\int_{\Omega} \bar{\xi}_{x_{j,\nu},\lambda_{j,\nu}}^{\frac{n+2}{n-2}} \right)^{\frac{n-2}{n+2}} \\
&\leq C \lambda_{i,\nu}^{\frac{2-n}{2}} \lambda_{i,\nu}^{\frac{2-n}{2} \frac{4}{n+2}} \lambda_{j,\nu}^{\frac{2-n}{2} \frac{n-2}{n+2}} \\
&\leq C \lambda_{i,\nu}^{\frac{2-n}{2}} (\lambda_{i,\nu}^{\frac{2-n}{2}} + \lambda_{j,\nu}^{\frac{2-n}{2}}) \\
&\leq C(\lambda_{i,\nu}^{2-n} + \lambda_{j,\nu}^{2-n}).
\end{aligned}$$

Combining (42), (43) and (44), we have

$$\begin{aligned}
&F(U_{\nu}) \\
&\leq \sum_{k=1}^m F(\xi_{x_{k,\nu},\lambda_{k,\nu}}) + o(1) \sum_{k=1}^m \int_{\Omega} \xi_{x_{k,\nu},\lambda_{k,\nu}}^2 + C \sum_{k=1}^m \lambda_{k,\nu}^{2-n} \\
&\quad + \sum_{i < j} \left[C \left(\int_{\Omega} \xi_{x_{i,\nu},\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \xi_{x_{j,\nu},\lambda_{j,\nu}} \right)^2 - c_{n,m} \int_{\Omega} (\xi_{x_{i,\nu},\lambda_{i,\nu}} \vee \xi_{x_{j,\nu},\lambda_{j,\nu}})^{\frac{4}{n-2}} (\xi_{x_{i,\nu},\lambda_{i,\nu}} \wedge \xi_{x_{j,\nu},\lambda_{j,\nu}})^2 \right].
\end{aligned}$$

Meanwhile, we have

$$\begin{aligned}
&C \left(\int_{\Omega} \xi_{x_{i,\nu},\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \xi_{x_{j,\nu},\lambda_{j,\nu}} \right)^2 - c_{n,m} \int_{\Omega} (\xi_{x_{i,\nu},\lambda_{i,\nu}} \vee \xi_{x_{j,\nu},\lambda_{j,\nu}})^{\frac{4}{n-2}} (\xi_{x_{i,\nu},\lambda_{i,\nu}} \wedge \xi_{x_{j,\nu},\lambda_{j,\nu}})^2 \\
&\leq C \left(\int_{\mathbb{R}^n} \bar{\xi}_{x_{i,\nu},\lambda_{i,\nu}}^{\frac{n+2}{n-2}} \bar{\xi}_{x_{j,\nu},\lambda_{j,\nu}} \right)^2 - c_{n,m} \int_{\mathbb{R}^n} (\bar{\xi}_{x_{i,\nu},\lambda_{i,\nu}} \vee \bar{\xi}_{x_{j,\nu},\lambda_{j,\nu}})^{\frac{4}{n-2}} (\bar{\xi}_{x_{i,\nu},\lambda_{i,\nu}} \wedge \bar{\xi}_{x_{j,\nu},\lambda_{j,\nu}})^2 \\
&\quad + C(\lambda_{i,\nu}^{2-n} + \lambda_{j,\nu}^{2-n}) \\
&\leq C(\lambda_{i,\nu}^{2-n} + \lambda_{j,\nu}^{2-n}) \quad \text{for all large } \nu,
\end{aligned}$$

where we used (77) in the last inequality.

Therefore, we complete the proof. \square

Corollary 4.6. *If $n \geq 4$ and $b > 0$ satisfying (2), we have, for large ν ,*

$$F(U_{\nu}) \leq \frac{2m}{n} Y(\mathbb{S}^n)^{\frac{n}{2}}.$$

Proof. Note that

$$\int_{\Omega} \xi_{x_{j,\nu},\lambda_{j,\nu}}^2 \geq \begin{cases} \frac{1}{C} \lambda_{j,\nu}^{-\frac{4}{n-2}}, & \text{if } n \neq 4, \\ \frac{1}{C} \lambda_{j,\nu}^{-2} \ln \lambda_{j,\nu}, & \text{if } n = 4. \end{cases}$$

Hence, if $n \geq 4$, for any large constant N we can find $j_N > 0$ such that for all $j \geq j_N$ there holds $\int_{\Omega} \xi_{x_{j,\nu},\lambda_{j,\nu}}^2 \geq N \lambda_{j,\nu}^{2-n}$. The corollary follows immediately from Proposition 4.5. \square

4.2 The case $v_\infty > 0$

Let $\mathcal{L}^2(\Omega) := \{f : \int_\Omega f^2 v_\infty^{\frac{4}{n-2}} < \infty\}$ with the inner product $\langle f, g \rangle = \int_\Omega f g v_\infty^{\frac{4}{n-2}}$. Then the operator

$$f \mapsto \left[v_\infty^{-\frac{4}{n-2}} (-\Delta - b) \right]^{-1} f$$

is a bounded linear compact symmetric operator mapping $\mathcal{L}^2(\Omega)$ into itself. Using the spectral theorem, there exists a sequence of $H_0^1(\Omega)$ functions $\{\phi_l : l \in \mathbb{N}\}$ and a sequence of positive real numbers $\{\mu_l : l \in \mathbb{N}\}$ such that $0 < \mu_1 < \mu_2 \leq \mu_3 \leq \dots \rightarrow \infty$,

$$-\Delta \phi_l - b \phi_l = \mu_l v_\infty^{\frac{4}{n-2}} \phi_l \quad \text{in } \Omega, \quad \phi_l = 0 \quad \text{on } \partial\Omega,$$

and $\{\phi_l : l \in \mathbb{N}\}$ forms an orthonormal basis of $\mathcal{L}^2(\Omega)$. In particular,

$$\int_\Omega v_\infty^{\frac{4}{n-2}} \phi_i \phi_j = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$$

By the regularity theory of linear elliptic equations, $\phi_l \in C^{2+\frac{4}{n-2}}(\bar{\Omega}) \cap C^\infty(\Omega)$ for every l . By the equation of v_∞ and the positivity of v_∞ , we know that $\mu_1 = 1$ and $\phi_1 = v_\infty (\int_\Omega v_\infty^{\frac{2n}{n-2}})^{-1/2}$. It is easy to check that $\{\frac{1}{\sqrt{\mu_l}} \phi_l\}$ is also an orthonormal basis of $H_0^1(\Omega)$ with respect to the inner product (13).

Let L be the largest number such that

$$\mu_l \leq \frac{n+2}{n-2} \quad \text{for all } l \leq L.$$

For $f \in L^p(\Omega)$, $p \geq 1$, we denote by Π the projection operator

$$\Pi f = f - \sum_{i=1}^L \left(\int_\Omega f \phi_i \, dx \right) v_\infty^{\frac{4}{n-2}} \phi_i.$$

It is clear that $\Pi(L^p(\Omega)) = \{f \in L^p(\mathbb{R}^n) : \int_\Omega f \phi_i = 0, i = 1, 2, \dots, L\}$. Hence, $\Pi(L^p(\Omega))$ is a closed subspace of $L^p(\Omega)$, and thus, is a Banach space with the inherited L^p norm.

We have several estimates regarding this projection.

Lemma 4.7. *For every $1 \leq p < \infty$, we can find a constant C such that*

$$\|f\|_{L^p(\Omega)} \leq C \left\| \Delta f + b f + \frac{n+2}{n-2} v_\infty^{\frac{4}{n-2}} f \right\|_{L^p(\Omega)} + C \sup_{1 \leq l \leq L} \left| \int_\Omega v_\infty^{\frac{4}{n-2}} \phi_l f \right|$$

for all $f \in W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)$.

Proof. Suppose that this is not true. Then there exists a sequence of functions $f_k \in W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)$ such that $\|f_k\|_{L^p(\Omega)} = 1$ for all k , and

$$\lim_{k \rightarrow \infty} \left\| \Delta f_k + b f_k + \frac{n+2}{n-2} v_\infty^{\frac{4}{n-2}} f_k \right\|_{L^p(\Omega)} + \lim_{k \rightarrow \infty} \sup_{1 \leq l \leq L} \left| \int_\Omega v_\infty^{\frac{4}{n-2}} \phi_l f_k \right| = 0.$$

If $p > 1$, then by the $W^{2,p}$ estimates, we have $\|f_k\|_{W^{2,p}(\Omega)} \leq C$. If $p = 1$, by the estimates of Brezis-Strauss [9], $\|f_k\|_{W^{1,q}(\Omega)} \leq C$ for some $q > 1$. Therefore, by the compactness, we obtain an f such that $\|f\|_{L^p(\Omega)} = 1$, $\int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \phi_l f_k = 0$ for all $1 \leq l \leq L$, and

$$\Delta f + bf + \frac{n+2}{n-2} v_{\infty}^{\frac{4}{n-2}} f = 0$$

in the distribution sense. Multiplying ϕ_l and integrating by parts, we have

$$\left(\mu_l - \frac{n+2}{n-2} \right) \int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \phi_l f_k = 0.$$

Hence, $\int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \phi_l f_k = 0$ for all $l > L$. Meanwhile, from the elliptic regularity, we know that $f \in L^{\infty}(\Omega)$. Hence, $f \in \mathcal{L}^2(\Omega)$, and thus, $f \equiv 0$, which is a contradiction. \square

Lemma 4.8. *There exists a constant C such that*

(i)

$$\|f\|_{L^{\frac{n+2}{n-2}}(\Omega)} \leq C \left\| \Pi(\Delta f + bf + \frac{n+2}{n-2} v_{\infty}^{\frac{4}{n-2}} f) \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega)} + C \sup_{1 \leq l \leq L} \left| \int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \phi_l f \right|$$

for all $f \in W^{2, \frac{n(n+2)}{n^2+4}}(\Omega) \cap W_0^{1, \frac{n(n+2)}{n^2+4}}(\Omega)$.

(ii)

$$\|f\|_{L^1(\Omega)} \leq C \left\| \Pi(\Delta f + bf + \frac{n+2}{n-2} v_{\infty}^{\frac{4}{n-2}} f) \right\|_{L^1(\Omega)} + C \sup_{1 \leq l \leq L} \left| \int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \phi_l f \right|$$

for all $f \in W^{2,1}(\Omega) \cap W_0^{1,1}(\Omega)$.

Proof. Given Lemma 4.7, the proof is the same as that of Lemma 6.3 in [6]. \square

Lemma 4.9. *There exists $\delta_1 > 0$ such that for every $z = (z_1, \dots, z_L) \in \mathbb{R}^L$ with $|z| \leq \delta_1$, there exists $\xi_z \in C_0^{\frac{3n-2}{n-2}}(\overline{\Omega})$ satisfying $1/2 \leq \xi_z/v_{\infty} \leq 2$ in Ω ,*

$$\int_{\Omega} v_{\infty}^{\frac{4}{n-2}} (\xi_z - v_{\infty}) \phi_l dx = z_l, \quad l = 1, \dots, L,$$

and

$$\Pi(\Delta \xi_z + b \xi_z + \xi_z^{\frac{n+2}{n-2}}) = 0. \quad (45)$$

Furthermore, the map $z \mapsto \xi_z$ is real analytic and $\frac{\partial}{\partial z_1} \xi_z(0) = v_{\infty}$, $\frac{\partial}{\partial z_l} \xi_z(0) = \phi_l$ for $2 \leq l \leq L$.

Proof. Let $\xi_z = (1 + z_1)v_{\infty} + \sum_{l=2}^L z_l \phi_l + h$, where

$$h \in \mathcal{H} := \text{span}\{\phi_1, \dots, \phi_L\}^{\perp}$$

and the “ \perp ” is with respect to the inner product (13). By a direct computation,

$$\begin{aligned}
& \Pi(\Delta\xi_z + b\xi_z + \xi_z^{\frac{n+2}{n-2}}) \\
&= (\Delta + b)\xi_z - \sum_{l=1}^L \left(\int_{\Omega} (\Delta + b)\xi_z \phi_l \right) v_{\infty}^{\frac{4}{n-2}} \phi_l + \xi_z^{\frac{n+2}{n-2}} - \sum_{l=1}^L \left(\int_{\Omega} \xi_z^{\frac{n+2}{n-2}} \phi_l \right) v_{\infty}^{\frac{4}{n-2}} \phi_l \\
&= (\Delta + b)h + \xi_z^{\frac{n+2}{n-2}} - \sum_{l=1}^L \left(\int_{\Omega} \xi_z^{\frac{n+2}{n-2}} \phi_l \right) v_{\infty}^{\frac{4}{n-2}} \phi_l =: G(z, h).
\end{aligned}$$

For any $p > n$, we claim that there exists a small constant $\delta > 0$ such that

$$G : \{|z| < \delta\} \times \{\|h\|_{W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)} < \delta\} \rightarrow \Pi(L^p(\Omega))$$

is analytic. Indeed, let $\Phi(z, h) = \xi_z$, $\mathcal{L}u = \Delta u + bu + u^{\frac{n+2}{n-2}}$. Then we have $G = \Pi \circ \mathcal{L} \circ \Phi$. Obviously, the linear maps Φ and Π are analytic. By Lemma 5.3 of Feireisl-Simondon [20], \mathcal{L} is also analytic in some small neighborhood of v_{∞} in $W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)$.

Note that $G(0, 0) = 0$ and

$$G_h(0, 0)\varphi = (\Delta + b)\varphi + \frac{n+2}{n-2} \left(v_{\infty}^{\frac{4}{n-2}} \varphi - \sum_{l=1}^L \left(\int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \varphi \phi_l \right) v_{\infty}^{\frac{4}{n-2}} \phi_l \right).$$

Since $G_h(0, 0)$ is coercive on $H_0^1(\Omega) \cap \mathcal{H}$, then $G_h(0, 0) : W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega) \cap \mathcal{H} \rightarrow \Pi(L^p(\Omega))$ is invertible, and both $G_h(0, 0)$ and $(G_h(0, 0))^{-1}$ are continuous. By the Implicit Function Theorem we can find $h(z) \in W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega) \cap \mathcal{H}$ such that $G(z, h(z)) = 0$ and h is analytic in z . The regularity of $(h(z))(\cdot)$ follows from linear elliptic equations theory. Since $h(0) = 0$, and $0 = G_z(0, 0) + G_h(0, 0)\partial_z h(0) = G_h(0, 0)\partial_z h(0)$, we have $\partial_z h(0) = 0$. It follows that $\frac{\partial}{\partial z_1} \xi_z(0) = v_{\infty}$ and $\frac{\partial}{\partial z_l} \xi_z(0) = \phi_l$ for $2 \leq l \leq L$. Therefore, we complete the proof. \square

The difference of the energy at ξ_z and v_{∞} can be controlled as follows.

Lemma 4.10. *There exists a real number $0 < \gamma < 1$ such that*

$$F(\xi_z) - F(v_{\infty}) \leq 2 \sup_{1 \leq l \leq L} \left| \int_{\Omega} (\Delta\xi_z + b\xi_z + \xi_z^{\frac{n+2}{n-2}}) \phi_l \, dx \right|^{1+\gamma}$$

if z is sufficiently small.

Proof. By Lemma 4.9 above and Lemma 5.3 of [20], the function $z \mapsto F(\xi_z)$ is real analytic. Using the Lojasiewicz inequality, we have

$$|F(\xi_z) - F(v_{\infty})| \leq \sup_l \left| \frac{\partial}{\partial z_l} F(\xi_z) \right|^{1+\gamma}$$

if z is sufficiently small. By a direct computation, we have

$$\frac{\partial}{\partial z_l} F(\xi_z) = -2 \int_{\Omega} (\Delta\xi_z + b\xi_z + \xi_z^{\frac{n+2}{n-2}}) \frac{\partial \xi_z}{\partial z_l} \, dx = -2 \int_{\Omega} (\Delta\xi_z + b\xi_z + \xi_z^{\frac{n+2}{n-2}}) \phi_l \, dx.$$

Therefore, we complete the proof. \square

For every ν , as in the beginning of Section 4.1, let \mathcal{A}_ν be the closed set of all m -tuplets $(x_k, \lambda_k, \alpha_k)_{1 \leq k \leq m}$ satisfying $(x_k, \lambda_k, \alpha_k) \in \overline{B}_{\frac{1}{\lambda_{k,\nu}^*}}(x_{k,\nu}^*) \times [\frac{\lambda_{k,\nu}}{2}, \frac{3\lambda_{k,\nu}}{2}] \times [\frac{1}{2}, \frac{3}{2}]$. Let $\delta_1 > 0$ be the constant in Lemma 4.9 and $\overline{B}_{\delta_1}^L$ is the open ball in \mathbb{R}^L centered at origin with radius δ_1 . Choose an element $(z_\nu, (x_{k,\nu}, \lambda_{k,\nu}, \alpha_{k,\nu})_{1 \leq k \leq m}) \in \overline{B}_{\delta_1}^L \times \mathcal{A}_\nu$ such that

$$\left\| v_\nu - \xi_{z_\nu} - \sum_{k=0}^m \alpha_{k,\nu} \xi_{x_{k,\nu}, \lambda_{k,\nu}} \right\| = \inf_{(z, (x_k, \lambda_k, \alpha_k)_{1 \leq k \leq m}) \in \overline{\mathcal{A}}_\nu} \left\| v_\nu - \xi_z - \sum_{k=0}^m \alpha_k \xi_{x_k, \lambda_k} \right\|. \quad (46)$$

Similar to (32) - (34), we have

$$\frac{\lambda_{i,\nu}}{\lambda_{j,\nu}} + \frac{\lambda_{j,\nu}}{\lambda_{i,\nu}} + \lambda_{i,\nu} \lambda_{j,\nu} |x_{i,\nu} - x_{j,\nu}|^2 \rightarrow \infty, \quad (47)$$

and for all k

$$\lambda_{k,\nu} d(x_{k,\nu}) \rightarrow \infty \quad (48)$$

as $\nu \rightarrow \infty$. In addition, $d(x_{k,\nu}) > \delta/2$, and

$$\left\| v_\nu - \xi_{z_\nu} - \sum_{k=1}^m \alpha_k \xi_{x_{k,\nu}, \lambda_{k,\nu}} \right\| \rightarrow 0 \quad (49)$$

as $\nu \rightarrow \infty$.

By the triangle inequality,

$$\begin{aligned} & \left\| \xi_{z_\nu} - v_\nu + \sum_{k=1}^m \alpha_k \xi_{x_{k,\nu}, \lambda_{k,\nu}} - \sum_{k=1}^m \xi_{x_{k,\nu}^*, \lambda_{k,\nu}^*} \right\| \\ & \leq \left\| v_\nu - \xi_{z_\nu} - \sum_{k=1}^m \alpha_k \xi_{x_{k,\nu}, \lambda_{k,\nu}} \right\| + \left\| v_\nu - v_\infty - \sum_{k=1}^m \xi_{x_{k,\nu}^*, \lambda_{k,\nu}^*} \right\| = o(1). \end{aligned}$$

It follows that, for all $1 \leq k \leq m$,

$$|z_\nu| = o(1), \quad |x_{k,\nu} - x_{k,\nu}^*| = o(1) \frac{1}{\lambda_{k,\nu}^*}, \quad \frac{\lambda_{k,\nu}}{\lambda_{k,\nu}^*} = 1 + o(1), \quad \alpha_{k,\nu} = 1 + o(1). \quad (50)$$

In particular, $(z_\nu, (x_{k,\nu}, \lambda_{k,\nu}, \alpha_{k,\nu})_{1 \leq k \leq m}) \in \overline{B}_{\delta_1}^L \times \mathcal{A}_\nu$ is an interior point.

In the sequel, we assume

$$\lambda_{1,\nu} \geq \lambda_{2,\nu} \geq \cdots \geq \lambda_{m,\nu}. \quad (51)$$

Let

$$U_\nu = \xi_{z_\nu} + \sum_{k=1}^m \alpha_{k,\nu} \xi_{x_{k,\nu}, \lambda_{k,\nu}}, \quad w_\nu = v_\nu - U_\nu. \quad (52)$$

Lemma 4.11. *We have for $1 \leq l \leq L$,*

$$\left| \int_\Omega v_\infty^{\frac{4}{n-2}} \phi_l w_\nu \, dx \right| \leq o(1) \int_\Omega |w_\nu| \, dx, \quad (53)$$

and for $1 \leq k \leq m$,

$$\begin{aligned} & \left| \int_{\Omega} \xi_{(x_{k,\nu}, \lambda_{k,\nu})}^{\frac{n+2}{n-2}} w_{\nu} \, dx \right| + \left| \int_{\Omega} \xi_{(x_{k,\nu}, \lambda_{k,\nu})}^{\frac{n+2}{n-2}} \frac{1 - \lambda^2 |x - x_{k,\nu}|^2}{1 + \lambda^2 |x - x_{k,\nu}|^2} w_{\nu} \, dx \right| \\ & + \left| \int_{\Omega} \xi_{(x_{k,\nu}, \lambda_{k,\nu})}^{\frac{n+2}{n-2}} \frac{\lambda^2 (x - x_{k,\nu})}{1 + \lambda^2 |x - x_{k,\nu}|^2} w_{\nu} \, dx \right| \leq o(1) \left(\int_{\Omega} |w_{\nu}|^{\frac{2n}{n-2}} \, dx \right)^{\frac{n-2}{2n}}. \end{aligned} \quad (54)$$

Proof. Let $\tilde{\phi}_l = \frac{\partial}{\partial z_l} \xi_z$. By (50), we have $\|\tilde{\phi}_1 - v_{\infty}\|_{C^2(\Omega)} = o(1)$ and $\|\tilde{\phi}_l - \phi_l\|_{C^2(\Omega)} = o(1)$ for $l = 2, \dots, L$. By the definition of $(z_{\nu}, (x_{k,\nu}, \lambda_{k,\nu}, \alpha_{k,\nu})_{1 \leq k \leq m})$, we have

$$\int \nabla \tilde{\phi}_l \nabla w_{\nu} - b \tilde{\phi}_l w_{\nu} = 0.$$

Hence,

$$\begin{aligned} \mu_l \int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \phi_l w_{\nu} \, dx &= \int_{\Omega} \left(-\Delta \phi_l - b \phi_l \right) w_{\nu} \, dx \\ &= \int_{\Omega} \left(\Delta (\tilde{\phi}_l - \phi_l) + b (\tilde{\phi}_l - \phi_l) \right) w_{\nu} \, dx. \end{aligned}$$

Since $\mu_l > 0$, then we can conclude (53). The proof of (54) is the same as that of Lemma 4.1. \square

Now we can show the non-degeneracy estimates of the second variation of F for w_{ν} .

Lemma 4.12. *For large ν , we have*

$$\frac{n+2}{n-2} \int_{\Omega} \left(v_{\infty}^{\frac{4}{n-2}} + \sum_{k=1}^m \xi_{x_{k,\nu}, \lambda_{k,\nu}}^{\frac{4}{n-2}} \right) w_{\nu}^2 \leq (1-c) \int_{\Omega} (|\nabla w_{\nu}|^2 - b w_{\nu}^2) \, dx,$$

where $c > 0$ is independent of ν .

Proof. We assume w_{ν} is not zero, otherwise there is nothing to prove. Define $\tilde{w}_{\nu} = \frac{w_{\nu}}{\|w_{\nu}\|}$. Suppose the lemma is not true. Then we can find a subsequence of $\{\tilde{w}_{\nu}\}$ (still denoted by $\{\tilde{w}_{\nu}\}$) satisfying

$$\lim_{\nu \rightarrow \infty} \frac{n+2}{n-2} \int_{\Omega} \left(v_{\infty}^{\frac{4}{n-2}} + \sum_{k=1}^m \xi_{x_{k,\nu}, \lambda_{k,\nu}}^{\frac{4}{n-2}} \right) \tilde{w}_{\nu}^2 \geq 1. \quad (55)$$

By (14),

$$\int_{\Omega} |\tilde{w}_{\nu}|^{\frac{2n}{n-2}} \leq K_b^{\frac{n}{n-2}} \|\tilde{w}_{\nu}\| = K_b^{\frac{n}{n-2}}. \quad (56)$$

By (47) and (51), we can find $R_{\nu} \rightarrow \infty$, $R_{\nu} \lambda_{j,\nu}^{-1} \rightarrow 0$ for all $1 \leq j \leq m$, and

$$\frac{\lambda_{i,\nu}}{R_{\nu}} (\lambda_{j,\nu}^{-1} + |x_{i,\nu} - x_{j,\nu}|) \rightarrow \infty \quad (57)$$

for all $i < j$. Set

$$\Omega_{j,\nu} = B_{R_{\nu} \lambda_{j,\nu}^{-1}}(x_{j,\nu}) \setminus \bigcup_{i=1}^{j-1} B_{R_{\nu} \lambda_{i,\nu}^{-1}}(x_{i,\nu}).$$

By (55) and $\|\tilde{w}_{\nu}\| = 1$, there are two cases:

(i). We can find $1 \leq j \leq m$ such that

$$\lim_{\nu \rightarrow \infty} \int_{\Omega} \xi_{x_{j,\nu}, \lambda_{j,\nu}}^{\frac{4}{n-2}} \tilde{w}_{\nu}^2 > 0$$

and

$$\lim_{\nu \rightarrow \infty} \int_{\Omega_{j,\nu}} (|\nabla \tilde{w}_{\nu}|^2 - b\tilde{w}_{\nu}^2) \leq \frac{n+2}{n-2} \int_{\Omega} \xi_{x_{j,\nu}, \lambda_{j,\nu}}^{\frac{4}{n-2}} \tilde{w}_{\nu}^2.$$

(ii).

$$\lim_{\nu \rightarrow \infty} \int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \tilde{w}_{\nu}^2 > 0$$

and

$$\lim_{\nu \rightarrow \infty} \int_{\Omega \setminus \cup_j \Omega_{j,\nu}} (|\nabla \tilde{w}_{\nu}|^2 - b\tilde{w}_{\nu}^2) \leq \frac{n+2}{n-2} \int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \tilde{w}_{\nu}^2.$$

In the former case, we can obtain a contradiction similar to that in the proof of Lemma 4.3.

In the later case, after passing to subsequence we suppose $\tilde{w}_{\nu} \rightharpoonup \tilde{w}$ in H_0^1 as $\nu \rightarrow \infty$. It follows that

$$\int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \tilde{w}^2 > 0 \quad (58)$$

and

$$\int_{\Omega} (|\nabla \tilde{w}|^2 - b\tilde{w}^2) \leq \frac{n+2}{n-2} \int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \tilde{w}^2. \quad (59)$$

By (53), we further have

$$\int_{\Omega} v_{\infty}^{\frac{4}{n-2}} \tilde{w} \phi_l = 0 \quad \text{for } l = 1, \dots, L. \quad (60)$$

Combing (59) and (60), \tilde{w} has to be identically zero, which contradicts to (58).

Therefore, Lemma 4.12 is proved. \square

Corollary 4.13. *For large ν , we have*

$$\frac{n+2}{n-2} \int_{\Omega} U_{\nu}^{\frac{4}{n-2}} w_{\nu}^2 \leq (1-c) \int_{\Omega} (|\nabla w_{\nu}|^2 - b w_{\nu}^2) dx,$$

where $c > 0$ is independent of ν .

Proof. It follows from Lemma 4.12, Hölder's inequality, the Sobolev inequality (14) and the fact that

$$\int_{\Omega} \left| U_{\nu}^{\frac{4}{n-2}} - v_{\infty}^{\frac{4}{n-2}} - \sum_{k=1}^m \xi_{x_{k,\nu}, \lambda_{k,\nu}}^{\frac{4}{n-2}} \right|^{\frac{n}{2}} = o(1).$$

\square

The following two lemmas are estimates of $v_{\nu} - \xi_{z_{\nu}}$ in $L^{\frac{n+2}{n-2}}(\Omega)$ and $L^1(\Omega)$, respectively.

Lemma 4.14. *For large ν , we have*

$$\|v_{\nu} - \xi_{z_{\nu}}\|_{L^{\frac{n+2}{n-2}}(\Omega)}^{\frac{n+2}{n-2}} \leq C \|v_{\nu}^{\frac{n+2}{n-2}} (\mathcal{R}(t_{\nu}) - 1)\|_{L^{\frac{2n}{n+2}}(\Omega)}^{\frac{n+2}{n-2}} + C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}},$$

where $C > 0$ is independent of ν .

Proof. From (19), we have

$$\Delta v_\nu + b v_\nu + v_\nu^{\frac{n+2}{n-2}} = (1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}}.$$

Combining with (45), we obtain

$$\begin{aligned} & \Pi \left(\Delta(v_\nu - \xi_{z_\nu}) + b(v_\nu - \xi_{z_\nu}) + \frac{n+2}{n-2} v_\infty^{\frac{4}{n-2}} (v_\nu - \xi_{z_\nu}) \right) \\ &= \Pi \left((1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}} - \frac{n+2}{n-2} (\xi_{z_\nu}^{\frac{4}{n-2}} - v_\infty^{\frac{4}{n-2}}) (v_\nu - \xi_{z_\nu}) \right. \\ & \quad \left. + \xi_{z_\nu}^{\frac{n+2}{n-2}} + \frac{n+2}{n-2} \xi_{z_\nu}^{\frac{4}{n-2}} (v_\nu - \xi_{z_\nu}) - v_\nu^{\frac{n+2}{n-2}} \right). \end{aligned} \quad (61)$$

Apply (i) of Lemma 4.8 to $v_\nu - \xi_{z_\nu}$, we obtain

$$\begin{aligned} & \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)} \\ & \leq C \left\| (1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega)} + C \left\| (\xi_{z_\nu}^{\frac{4}{n-2}} - v_\infty^{\frac{4}{n-2}}) (v_\nu - \xi_{z_\nu}) \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega)} \\ & \quad + C \left\| \xi_{z_\nu}^{\frac{n+2}{n-2}} + \frac{n+2}{n-2} \xi_{z_\nu}^{\frac{4}{n-2}} (v_\nu - \xi_{z_\nu}) - v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega)} + C \sup_{1 \leq l \leq L} \left| \int_\Omega v_\infty^{\frac{4}{n-2}} \phi_l (v_\nu - \xi_{z_\nu}) \right|. \end{aligned}$$

Using the estimates for all $a, b \geq 0$ that

$$\left| a^{\frac{n+2}{n-2}} + \frac{n+2}{n-2} a^{\frac{4}{n-2}} (b-a) - b^{\frac{n+2}{n-2}} \right| \leq C a^{\max(0, \frac{4}{n-2}-1)} |b-a|^{\min(\frac{n+2}{n-2}, 2)} + C |b-a|^{\frac{n+2}{n-2}}, \quad (62)$$

we obtain

$$\begin{aligned} & \left\| \xi_{z_\nu}^{\frac{n+2}{n-2}} + \frac{n+2}{n-2} \xi_{z_\nu}^{\frac{4}{n-2}} (v_\nu - \xi_{z_\nu}) - v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega)} \\ & \leq C \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{n+2}{n-2}, 2)} + |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega)} \\ & \leq C \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{n+2}{n-2}, 2)} + |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\cup_{k=1}^m B_{N/\lambda_{k,\nu}}(x_{k,\nu}))} \\ & \quad + C \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{n+2}{n-2}, 2)} + |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega \setminus \cup_{k=1}^m B_{N/\lambda_{k,\nu}}(x_{k,\nu}))}, \end{aligned}$$

where N is a large real number to be chosen later. Using Hölder's inequality, we have

$$\begin{aligned} & \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{n+2}{n-2}, 2)} + |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\cup_{k=1}^m B_{N/\lambda_{k,\nu}}(x_{k,\nu}))} \\ & \leq C \sum_{k=1}^m (N/\lambda_{k,\nu})^{\frac{(n-2)^2}{2(n+2)}} \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{n+2}{n-2}, 2)} + |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^{\frac{2n}{n+2}}(\Omega)} \\ & \leq C \sum_{k=1}^m (N/\lambda_{k,\nu})^{\frac{(n-2)^2}{2(n+2)}} \end{aligned}$$

and

$$\begin{aligned} & \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{n+2}{n-2}, 2)} + |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega \setminus \cup_{k=1}^m B_{N/\lambda_{k,\nu}}(x_{k,\nu}))} \\ & \leq \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{4}{n-2}, 1)} + |v_\nu - \xi_{z_\nu}|^{\frac{4}{n-2}} \right\|_{L^{\frac{n}{2}}(\Omega \setminus \cup_{k=1}^m B_{N/\lambda_{k,\nu}}(x_{k,\nu}))} \cdot \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}. \end{aligned}$$

Since

$$\begin{aligned} & \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{2n}{n-2}}(\Omega \setminus \cup_{k=1}^m B_{N/\lambda_{k,\nu}}(x_{k,\nu}))} \\ & = \left\| \sum_{k=1}^m \alpha_{k,\nu} \xi_{x_{k,\nu}, \lambda_{k,\nu}} + w_\nu \right\|_{L^{\frac{2n}{n-2}}(\Omega \setminus \cup_{k=1}^m B_{N/\lambda_{k,\nu}}(x_{k,\nu}))} \\ & \leq \sum_{k=1}^m \alpha_{k,\nu} \|\xi_{x_{k,\nu}, \lambda_{k,\nu}}\|_{L^{\frac{2n}{n-2}}(\Omega \setminus B_{N/\lambda_{k,\nu}}(x_{k,\nu}))} + \|w_\nu\|_{L^{\frac{2n}{n-2}}(\Omega)} \\ & \leq CN^{-\frac{n-2}{2}} + o(1), \end{aligned}$$

we have

$$\begin{aligned} & \left\| \xi_{z_\nu}^{\frac{n+2}{n-2}} + \frac{n+2}{n-2} \xi_{z_\nu}^{\frac{4}{n-2}} (v_\nu - \xi_{z_\nu}) - v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{n(n+2)}{n^2+4}}(\Omega)} \\ & \leq C \sum_{k=1}^m (N/\lambda_{k,\nu})^{\frac{(n-2)^2}{2(n+2)}} + C(N^{-\frac{n-2}{2}} + N^{-2} + o(1)) \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}. \end{aligned}$$

Also,

$$\begin{aligned} \sup_{1 \leq l \leq L} \left| \int_\Omega v_\nu^{\frac{4}{n-2}} \phi_l(v_\nu - \xi_{z_\nu}) \right| &= \sup_{1 \leq l \leq L} \left| \int_\Omega v_\nu^{\frac{4}{n-2}} \phi_l \left(\sum_{k=1}^m \alpha_{k,\nu} \xi_{x_{k,\nu}, \lambda_{k,\nu}} + w_\nu \right) \right| \\ &\leq C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}} + o(1) \|w_\nu\|_{L^1(\Omega)} \\ &\leq C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}} + o(1) \left\| v_\nu - \xi_{z_\nu} - \sum_{k=1}^m \alpha_{k,\nu} \xi_{x_{k,\nu}, \lambda_{k,\nu}} \right\|_{L^1(\Omega)} \\ &\leq C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}} + o(1) \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)}. \end{aligned} \tag{63}$$

Putting these facts together, we have

$$\begin{aligned} & \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)} \\ & \leq C \left\| (1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{2n}{n+2}}(\Omega)} + C \sum_{k=1}^m (N/\lambda_{k,\nu})^{\frac{(n-2)^2}{2(n+2)}} \\ & + C(N^{-\frac{n-2}{2}} + N^{-2} + o(1)) \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)} + C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}}. \end{aligned}$$

By choosing N sufficiently large, we obtain

$$\|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)} \leq C \left\| (1 - \mathcal{R}(t_\nu))v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{2n}{n+2}}(\Omega)} + C \sum_{k=1}^m \lambda_{k,\nu}^{-\frac{(n-2)^2}{2(n+2)}},$$

from which the conclusion follows. \square

Lemma 4.15. *For large ν , we have*

$$\|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)} \leq C \|v_\nu^{\frac{n+2}{n-2}} (\mathcal{R}(t_\nu) - 1)\|_{L^{\frac{2n}{n+2}}(\Omega)} + C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}},$$

where $C > 0$ is independent of ν .

Proof. Using (61), and applying (ii) of Lemma 4.8 to $v_\nu - \xi_{z_\nu}$, we obtain

$$\begin{aligned} & \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)} \\ & \leq C \left\| (1 - \mathcal{R}(t_\nu))v_\nu^{\frac{n+2}{n-2}} \right\|_{L^1(\Omega)} + C \left\| (\xi_{z_\nu}^{\frac{4}{n-2}} - v_\infty^{\frac{4}{n-2}})(v_\nu - \xi_{z_\nu}) \right\|_{L^1(\Omega)} \\ & \quad + C \left\| \xi_{z_\nu}^{\frac{n+2}{n-2}} + \frac{n+2}{n-2} \xi_{z_\nu}^{\frac{4}{n-2}} (v_\nu - \xi_{z_\nu}) - v_\nu^{\frac{n+2}{n-2}} \right\|_{L^1(\Omega)} + C \sup_{1 \leq l \leq L} \left| \int_\Omega v_\infty^{\frac{4}{n-2}} \phi_l (v_\nu - \xi_{z_\nu}) \right|. \end{aligned}$$

It follows from (62) that

$$\begin{aligned} & \left\| \xi_{z_\nu}^{\frac{n+2}{n-2}} + \frac{n+2}{n-2} \xi_{z_\nu}^{\frac{4}{n-2}} (v_\nu - \xi_{z_\nu}) - v_\nu^{\frac{n+2}{n-2}} \right\|_{L^1(\Omega)} \\ & \leq C \left\| |v_\nu - \xi_{z_\nu}|^{\min(\frac{n+2}{n-2}, 2)} + |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^1(\Omega)} \\ & \leq C \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)}^{\max(0, 1 - \frac{n-2}{4})} \left\| |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}} \right\|_{L^1(\Omega)}^{\min(1, \frac{n-2}{4})} + C \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}^{\frac{n+2}{n-2}} \\ & \leq C \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)}^{\max(0, 1 - \frac{n-2}{4})} \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}^{\min(1, \frac{n-2}{4})} + C \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}^{\frac{n+2}{n-2}} \\ & \leq \frac{1}{2C} \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)} + C \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}^{\frac{n+2}{n-2}}, \end{aligned}$$

where we used Hölder's inequality in the second inequality and the Young inequality in the last inequality. Combining (63), we have

$$\begin{aligned} & \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)} \\ & \leq C \left\| (1 - \mathcal{R}(t_\nu))v_\nu^{\frac{n+2}{n-2}} \right\|_{L^1(\Omega)} + o(1) \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)} \\ & \quad + \frac{1}{2} \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)} + C \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}^{\frac{n+2}{n-2}} + C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}}. \end{aligned}$$

Then the conclusion follows from Lemma 4.14. \square

Using the above two lemmas, we can continue to estimate $F(\xi_{z_\nu}) - F(v_\infty)$ from Lemma 4.10.

Proposition 4.16. *For all large ν , we have*

$$F(\xi_{z_\nu}) - F(v_\infty) \leq C \left(\int_{\Omega} |\mathcal{R}(t_\nu) - 1| \frac{2n}{n+2} v_\nu^{\frac{2n}{n-2}} \right)^{\frac{n+2}{2n}(1+\gamma)} + C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}(1+\gamma)},$$

where $\gamma \in (0, 1)$ is the one in Lemma 4.10.

Proof. It follows from integration by parts that

$$\begin{aligned} & \int_{\Omega} (\Delta \xi_{z_\nu} + b \xi_{z_\nu} + \xi_{z_\nu}^{\frac{n+2}{n-2}}) \phi_l \, dx \\ &= \int_{\Omega} (\Delta v_\nu + b v_\nu + v_\nu^{\frac{n+2}{n-2}}) \phi_l \, dx + \mu_l \int_{\Omega} v_\nu^{\frac{4}{n-2}} \phi_l (v_\nu - \xi_{z_\nu}) \, dx - \int_{\Omega} \phi_l (v_\nu^{\frac{n+2}{n-2}} - \xi_{z_\nu}^{\frac{n+2}{n-2}}) \, dx \\ &= \int_{\Omega} (1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}} \phi_l \, dx + \mu_l \int_{\Omega} v_\nu^{\frac{4}{n-2}} \phi_l (v_\nu - \xi_{z_\nu}) \, dx - \int_{\Omega} \phi_l (v_\nu^{\frac{n+2}{n-2}} - \xi_{z_\nu}^{\frac{n+2}{n-2}}) \, dx. \end{aligned}$$

Using the pointwise estimate

$$|v_\nu^{\frac{n+2}{n-2}} - \xi_{z_\nu}^{\frac{n+2}{n-2}}| \leq C \xi_{z_\nu}^{\frac{4}{n-2}} |v_\nu - \xi_{z_\nu}| + C |v_\nu - \xi_{z_\nu}|^{\frac{n+2}{n-2}},$$

we have

$$\begin{aligned} & \sup_{1 \leq l \leq L} \left| \int_{\Omega} (\Delta \xi_z + b \xi_z + \xi_z^{\frac{n+2}{n-2}}) \phi_l \, dx \right| \\ & \leq C \left\| (1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{2n}{n+2}}(\Omega)} + C \|v_\nu - \xi_{z_\nu}\|_{L^1(\Omega)} + C \|v_\nu - \xi_{z_\nu}\|_{L^{\frac{n+2}{n-2}}(\Omega)}^{\frac{n+2}{n-2}} \\ & \leq C \left\| (1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{2n}{n+2}}(\Omega)} + C \left\| v_\nu^{\frac{n+2}{n-2}} (\mathcal{R}(t_\nu) - 1) \right\|_{L^{\frac{2n}{n+2}}(\Omega)}^{\frac{n+2}{n-2}} + C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}} \\ & \leq C \left\| (1 - \mathcal{R}(t_\nu)) v_\nu^{\frac{n+2}{n-2}} \right\|_{L^{\frac{2n}{n+2}}(\Omega)} + C \sum_{k=1}^m \lambda_{k,\nu}^{\frac{2-n}{2}}, \end{aligned}$$

where we used Lemma 4.14 and Lemma 4.15 in the second inequality, and Proposition 2.6 in the last inequality.

Then the conclusion follows from Lemma 4.10. \square

Corollary 4.17. *If $n \geq 4$ and $b > 0$ satisfying (2), we have*

$$F(U_\nu) \leq F(v_\infty) + \frac{2m}{n} Y(\mathbb{S}^n)^{n/2} + C \left(\int_{\Omega} |\mathcal{R}(t_\nu) - 1| \frac{2n}{n+2} v_\nu^{\frac{2n}{n-2}} \right)^{\frac{n+2}{2n}(1+\gamma)}.$$

Proof. Let $\tilde{U}_\nu = \sum_{k=1}^m \alpha_k \xi_{x_{k,\nu}, \lambda_{k,\nu}}$.

$$\begin{aligned} F(U_\nu) &= \int_{\Omega} |\nabla(\xi_{z_\nu} + \tilde{U}_\nu)|^2 - b(\xi_{z_\nu} + \tilde{U}_\nu)^2 - \frac{n-2}{n} \int_{\Omega} (\xi_{z_\nu} + \tilde{U}_\nu)^{\frac{2n}{n-2}} \\ &= F(\xi_{z_\nu}) + F(\tilde{U}_\nu) + 2 \int_{\Omega} (\nabla \xi_{z_\nu} \nabla \tilde{U}_\nu - b \xi_{z_\nu} \tilde{U}_\nu - \xi_{z_\nu}^{\frac{n+2}{n-2}} \tilde{U}_\nu) \\ &\quad - \frac{n-2}{n} \int_{\Omega} \left((\xi_{z_\nu} + \tilde{U}_\nu)^{\frac{2n}{n-2}} - \frac{2n}{n-2} \xi_{z_\nu}^{\frac{n+2}{n-2}} \tilde{U}_\nu - \xi_{z_\nu}^{\frac{2n}{n-2}} - \tilde{U}_\nu^{\frac{2n}{n-2}} \right). \end{aligned}$$

We have

$$\begin{aligned}
& \left| \int_{\Omega} (\nabla \xi_{z_\nu} \nabla \tilde{U}_\nu - b \xi_{z_\nu} \tilde{U}_\nu - \xi_{z_\nu}^{\frac{n+2}{n-2}} \tilde{U}_\nu) \right| \\
&= \left| \int_{\Omega} \left(\Delta(\xi_{z_\nu} - v_\infty) + b(\xi_{z_\nu} - v_\infty) + \xi_{z_\nu}^{\frac{n+2}{n-2}} - v_\infty^{\frac{n+2}{n-2}} \right) \tilde{U}_\nu \right| \\
&\leq o(1) \sum_{k=1}^m \lambda_k^{\frac{2-n}{2}}.
\end{aligned}$$

By Lemma A.1, there exists $c > 0$, depending only on n such that

$$(\xi_{z_\nu} + \tilde{U}_\nu)^{\frac{2n}{n-2}} - \frac{2n}{n-2} \xi_{z_\nu}^{\frac{n+2}{n-2}} \tilde{U}_\nu - \xi_{z_\nu}^{\frac{2n}{n-2}} - \tilde{U}_\nu^{\frac{2n}{n-2}} \geq \begin{cases} c \xi_{z_\nu}^{\frac{4}{n-2}} (\tilde{U}_\nu)^2, & \text{if } \xi_{z_\nu} \geq U'_\nu, \\ c \xi_{z_\nu} (\tilde{U}_\nu)^{\frac{n+2}{n-2}}, & \text{if } \xi_{z_\nu} < U'_\nu. \end{cases}$$

Since $v_\infty/2 \leq \xi_{z_\nu} \leq 2v_\infty$, and

$$\int_{|x| < \sqrt{\lambda^{-1}}} \left(\frac{\lambda}{1 + \lambda^2 |x|^2} \right)^{\frac{n+2}{2}} \geq \lambda^{-\frac{n-2}{2}} \int_{|y| < 1} \left(\frac{1}{1 + |y|^2} \right)^{\frac{n+2}{2}} \geq c \lambda^{-\frac{n-2}{2}},$$

we have

$$\int_{\Omega} \left((\xi_{z_\nu} + \tilde{U}_\nu)^{\frac{2n}{n-2}} - \frac{2n}{n-2} \xi_{z_\nu}^{\frac{n+2}{n-2}} \tilde{U}_\nu - \xi_{z_\nu}^{\frac{2n}{n-2}} - \tilde{U}_\nu^{\frac{2n}{n-2}} \right) \geq c \sum_{k=1}^m \lambda_k^{\frac{2-n}{2}}.$$

Then, the conclusion follows from Proposition 4.16 and Corollary 4.6. \square

5 Convergence

Using the estimates in Corollaries 4.4 and 4.13, we have the following estimate of $F(v_\nu) - F_\infty$ for any sequence of times $\{t_\nu : \nu \in \mathbb{N}\}$.

Proposition 5.1. *Let $n \geq 4$, and $b > 0$ satisfy (2). Let $\{t_\nu : \nu \in \mathbb{N}\}$ be a sequence of times such that $t_\nu \rightarrow \infty$ as $\nu \rightarrow \infty$. Then, we can find a real number $\gamma \in (0, 1)$ and a constant $C > 0$ such that, after passing to a subsequence, we have*

$$F(v_\nu) - F_\infty \leq C \left(\int_{\Omega} |\mathcal{R}(t_\nu) - 1|^{\frac{2n}{n+2}} v_\nu^{\frac{2n}{n-2}} \right)^{\frac{n+2}{2n}(1+\gamma)}$$

for all integers ν in that subsequence, where F_∞ is the one defined in (18). Note that γ and C may depend on the sequence $\{t_\nu : \nu \in \mathbb{N}\}$.

Proof. It follows from (18) that $F(v_\infty) = F_\infty$. Recall that $U_\nu = v_\nu - w_\nu$. We have

$$\begin{aligned}
F(v_\nu) - F(U_\nu) &= \frac{n-2}{n} \int_{\Omega} (U_\nu^{\frac{2n}{n-2}} - v_\nu^{\frac{2n}{n-2}}) + 2 \int_{\Omega} (\nabla v_\nu \nabla w_\nu - b v_\nu w_\nu) - \int_{\Omega} (|\nabla w_\nu|^2 - b w_\nu^2) \\
&= \frac{n-2}{n} \int_{\Omega} (U_\nu^{\frac{2n}{n-2}} - v_\nu^{\frac{2n}{n-2}}) + 2 \int_{\Omega} \mathcal{R} v_\nu^{\frac{n+2}{n-2}} w_\nu - \int_{\Omega} (|\nabla w_\nu|^2 - b w_\nu^2) \\
&= \frac{n-2}{n} \int_{\Omega} \left(U_\nu^{\frac{2n}{n-2}} - v_\nu^{\frac{2n}{n-2}} + \frac{2n}{n-2} v_\nu^{\frac{n+2}{n-2}} w_\nu - \frac{n(n+2)}{(n-2)^2} U_\nu^{\frac{4}{n-2}} w_\nu^2 \right) \\
&\quad + 2 \int_{\Omega} (\mathcal{R} - 1) v_\nu^{\frac{n+2}{n-2}} w_\nu - \int_{\Omega} \left(|\nabla w_\nu|^2 - b w_\nu^2 - \frac{n+2}{n-2} U_\nu^{\frac{4}{n-2}} w_\nu^2 \right)
\end{aligned}$$

Using the pointwise estimate

$$\begin{aligned}
& \left| U_\nu^{\frac{2n}{n-2}} - v_\nu^{\frac{2n}{n-2}} + \frac{2n}{n-2} v_\nu^{\frac{n+2}{n-2}} w_\nu - \frac{n(n+2)}{(n-2)^2} U_\nu^{\frac{4}{n-2}} w_\nu^2 \right| \\
&= \left| U_\nu^{\frac{2n}{n-2}} - (w_\nu + U_\nu)^{\frac{2n}{n-2}} + \frac{2n}{n-2} (w_\nu + U_\nu)^{\frac{n+2}{n-2}} w_\nu - \frac{n(n+2)}{(n-2)^2} U_\nu^{\frac{4}{n-2}} w_\nu^2 \right| \\
&\leq C U_\nu^{\max\{0, \frac{4}{n-2}-1\}} |w_\nu|^{\min\{\frac{2n}{n-2}, 3\}} + C |w_\nu|^{\frac{2n}{n-2}},
\end{aligned}$$

it follows that

$$\begin{aligned}
& \int_\Omega \left| U_\nu^{\frac{2n}{n-2}} - v_\nu^{\frac{2n}{n-2}} + \frac{2n}{n-2} v_\nu^{\frac{n+2}{n-2}} w_\nu - \frac{n(n+2)}{(n-2)^2} U_\nu^{\frac{4}{n-2}} w_\nu^2 \right| \\
&\leq C \int_\Omega U_\nu^{\max\{0, \frac{4}{n-2}-1\}} |w_\nu|^{\min\{\frac{2n}{n-2}, 3\}} + C \int_\Omega |w_\nu|^{\frac{2n}{n-2}} \\
&\leq C \left(\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n} \min\{\frac{n}{n-2}, \frac{3}{2}\}}.
\end{aligned}$$

By Hölder inequality and Cauchy inequality, we have

$$\begin{aligned}
\left| \int_\Omega (\mathcal{R} - 1) v_\nu^{\frac{n+2}{n-2}} w_\nu \right| &\leq C \left(\int_\Omega |\mathcal{R} - 1|^{\frac{2n}{n+2}} v_\nu^{\frac{2n}{n-2}} \right)^{\frac{n+2}{2n}} \left(\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{2n}} \\
&\leq \varepsilon \left(\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n}} + C(\varepsilon) \left(\int_\Omega |\mathcal{R} - 1|^{\frac{2n}{n+2}} v_\nu^{\frac{2n}{n-2}} \right)^{\frac{n+2}{n}}.
\end{aligned}$$

Finally, by Corollaries 4.4 and 4.13, we have

$$\int_\Omega |\nabla w_\nu|^2 - b w_\nu^2 - \frac{n+2}{n-2} U_\nu^{\frac{4}{n-2}} w_\nu^2 \geq \frac{1}{C} \left(\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n}}.$$

Since $\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \rightarrow 0$, we have, by choosing ε being small, that

$$\begin{aligned}
F(v_\nu) - F(U_\nu) &\leq C \left(\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n} \min\{\frac{n}{n-2}, \frac{3}{2}\}} + C \left(\int_\Omega |\mathcal{R} - 1|^{\frac{2n}{n+2}} v_\nu^{\frac{2n}{n-2}} \right)^{\frac{n+2}{n}} \\
&\quad - \frac{1}{2C} \left(\int_\Omega |w_\nu|^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n}} \\
&\leq C \left(\int_\Omega |\mathcal{R} - 1|^{\frac{2n}{n+2}} v_\nu^{\frac{2n}{n-2}} \right)^{\frac{n+2}{n}}.
\end{aligned}$$

By Corollary 4.17, we complete the proof. \square

Then we can show the estimate for all large time.

Corollary 5.2. *There exist real numbers $\gamma \in (0, 1)$ and $t_0 > 0$ such that*

$$F(v(t)) - F_\infty \leq \left(\int_\Omega |\mathcal{R} - 1|^{\frac{2n}{n+2}} v(x, t)^{\frac{2n}{n-2}} dx \right)^{\frac{n+2}{2n}(1+\gamma)}$$

for all $t \geq t_0$.

Proof. Suppose this is not true. Then, there exists a sequence of times $\{t_\nu : \nu \in \mathbb{N}\}$ such that $t_\nu > \nu$ and

$$F(v(t_\nu)) - F_\infty \geq \left(\int_\Omega |\mathcal{R}(t_\nu) - 1| \frac{2n}{n+2} v(x, t_\nu)^{\frac{2n}{n-2}} dx \right)^{\frac{n+2}{2n}(1+\frac{1}{\nu})}$$

for all $\nu \in \mathbb{N}$. By applying Proposition 5.1 to this sequence $\{t_\nu : \nu \in \mathbb{N}\}$, there exists an infinite subset $I \subset \mathbb{N}$, a real number $\gamma \in (0, 1)$ and $C > 0$ such that

$$F(v(t_\nu)) - F_\infty \leq C \left(\int_\Omega |\mathcal{R}(t_\nu) - 1| \frac{2n}{n+2} v(x, t_\nu)^{\frac{2n}{n-2}} dx \right)^{\frac{n+2}{2n}(1+\gamma)}$$

for all $\nu \in I$. Thus, we have

$$1 \leq C \left(\int_\Omega |\mathcal{R}(t_\nu) - 1| \frac{2n}{n+2} v(x, t_\nu)^{\frac{2n}{n-2}} dx \right)^{\frac{n+2}{2n}(\gamma-\frac{1}{\nu})}$$

for all $\nu \in I$. However, from Proposition 2.6, we have

$$\lim_{\nu \rightarrow \infty} \left(\int_\Omega |\mathcal{R}(t_\nu) - 1| \frac{2n}{n+2} v(x, t_\nu)^{\frac{2n}{n-2}} dx \right)^{\frac{n+2}{2n}(\gamma-\frac{1}{\nu})} = 0.$$

We have reached a contradiction. □

Now we can use a differential inequality of F to obtain a decay estimate.

Proposition 5.3. *There exist $\theta \in (0, 1)$ and $C > 0$ such that for all $T > 1$, there holds*

$$\int_T^\infty M_2(t)^{1/2} dt \leq CT^{-\theta},$$

where M_2 is defined in (20),

Proof. It follows from Corollary 5.2, Hölder's inequality, and (15) that

$$0 \leq F(v(t)) - F_\infty \leq \left(\int_\Omega |\mathcal{R} - 1| \frac{2n}{n+2} v(x, t)^{\frac{2n}{n-2}} dx \right)^{\frac{n+2}{2n}(1+\gamma)} \leq CM_2(t)^{\frac{1+\gamma}{2}}.$$

It follows that

$$\frac{d}{dt}(F(v(t)) - F_\infty) = -\frac{2(n-2)}{n+2}M_2(t) \leq -C(F(v(t)) - F_\infty)^{\frac{2}{1+\gamma}}.$$

Hence,

$$\frac{d}{dt}(F(v(t)) - F_\infty)^{\frac{\gamma+1}{2}} \geq C \frac{1-\gamma}{1+\gamma} > 0.$$

It follows that

$$F(v(t)) - F_\infty \leq Ct^{-\frac{1+\gamma}{2}}$$

for sufficiently large t . Then we have

$$\begin{aligned}
\left(\int_T^{2T} M_2(s)^{1/2} ds \right)^2 &\leq T \int_T^{2T} M_2(s) ds \\
&\leq \frac{n+2}{n-2} T (F(v(T)) - F(v(2T))) \\
&\leq \frac{n+2}{n-2} T (F(v(T)) - F_\infty) \\
&\leq CT^{-\frac{2\gamma}{1-\gamma}},
\end{aligned}$$

where we used the monotonicity of F . It follows that

$$\int_T^\infty M_2(t)^{1/2} dt = \sum_{k=0}^\infty \int_{2^k T}^{2^{k+1} T} M_2(t)^{1/2} dt \leq CT^{-\frac{\gamma}{1-\gamma}} \sum_{k=1}^\infty 2^{-\frac{\gamma}{1-\gamma} k} \leq CT^{-\frac{\gamma}{1-\gamma}}. \quad (64)$$

This finishes the proof. \square

We are ready to show the uniform boundedness, and uniform higher order estimates.

Proposition 5.4. *For any $\varepsilon > 0$, there exists $T_0 > 0$ such that*

$$\|v(\cdot, t) - v(\cdot, T_0)\|_{L^{\frac{2n}{n-2}}(\Omega)} < \varepsilon \quad \text{for all } t > T_0. \quad (65)$$

Consequently, there exists $C > 0$ such that

$$v(x, t) \leq C \quad \text{in } \Omega \text{ for all } t > 1. \quad (66)$$

Proof. For $b > a > 1$, using the pointwise estimate

$$|v(x, b) - v(x, a)|^{\frac{n}{n-2}} \leq |v(x, b)^{\frac{n}{n-2}} - v(x, a)^{\frac{n}{n-2}}|,$$

we have

$$\begin{aligned}
\left(\int_\Omega |v(x, b) - v(x, a)|^{\frac{2n}{n-2}} dx \right)^{1/2} &\leq \left(\int_\Omega |v(x, b)^{\frac{n}{n-2}} - v(x, a)^{\frac{n}{n-2}}|^2 dx \right)^{1/2} \\
&\leq \left(\int_\Omega \left(\int_a^b |\partial_t(v(x, t)^{\frac{n}{n-2}})| dt \right)^2 dx \right)^{1/2} \\
&\leq \int_a^b \left(\int_\Omega |\partial_t(v(x, t)^{\frac{n}{n-2}})|^2 dx \right)^{1/2} dt \\
&\leq C \int_a^\infty M_2(t)^{1/2} dt \\
&\leq Ca^{-\theta},
\end{aligned} \quad (67)$$

where we used Minkowski's integral inequality in the third inequality, and Proposition 5.3 in the last inequality. Hence, for any $\varepsilon > 0$, there exists $T_0 > 0$ such that (65) holds.

Furthermore, it follows from (19) that

$$-\Delta v - bv = \left[(\mathcal{R} - 1)v^{\frac{4}{n-2}} + v^{\frac{4}{n-2}} \right] v. \quad (68)$$

Using (65) and the fact that $M_{n/2}(t) \rightarrow 0$ as $t \rightarrow \infty$, we can apply the standard Brezis-Kato estimates [7] to the equation (68) to obtain that there exists $p > \frac{2n}{n-2}$ and $C > 0$ such that

$$\|v(t)\|_{L^p(\Omega)} < C \quad \text{for all } t > T_0.$$

Then we can further use Proposition 2.6 with large q and apply the standard Moser iteration to the equation (68) to obtain (66). \square

Theorem 5.5. *There exists $C > 0$ such that*

$$\|v(\cdot, t)\|_{C^{2+\frac{n+2}{n-2}}(\bar{\Omega})} \leq C \quad \text{for all } t > 1. \quad (69)$$

Proof. By using (66), it follows from Proposition 6.2 in [18] (more precisely, its proof) that

$$\frac{1}{C}d(x) \leq v(\cdot, t) \leq Cd(x) \quad \text{for all } x \in \Omega, t > 1,$$

where $d(x) := \text{dist}(x, \partial\Omega)$. Then (69) follows from Theorem 5.1 in [23]. \square

Let us conclude this section with the proof of Theorem 1.1.

Proof of Theorem 1.1. It follows from Proposition 5.4, Theorem 5.5 and (15) that there exists a nonzero stationary solution v_∞ of (9) such that

$$\lim_{t \rightarrow \infty} \|v(\cdot, t) - v_\infty\|_{C^3(\bar{\Omega})} = 0.$$

From (67), we know that there exist $C > 0$ and $\theta > 0$ such that

$$\|v(\cdot, t) - v_\infty\|_{L^{\frac{2n}{n-2}}(\Omega)} \leq Ct^{-\theta} \quad \text{for all } t > 1.$$

Using (69) and interpolation inequalities (Gagliardo–Nirenberg interpolation inequality and Morrey inequality), we have

$$\|v(\cdot, t) - v_\infty\|_{C^1(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t > 1.$$

Since $v(\cdot, t) \equiv v_\infty \equiv 0$ on $\partial\Omega$, we have for all $x \in \Omega$ that

$$\left| \frac{v(x, t) - v_\infty(x)}{v_\infty(x)} \right| \leq \frac{\|v(\cdot, t) - v_\infty\|_{C^1(\bar{\Omega})} d(x)}{d(x)/C} \leq C\|v(\cdot, t) - v_\infty\|_{C^1(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t > 1.$$

That is,

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{L^\infty(\Omega)} \leq Ct^{-\theta} \quad \text{for all } t \geq 1.$$

Using (69) again, we have

$$\left\| \frac{v(\cdot, t)}{v_\infty} \right\|_{C^{1+\frac{n+2}{n-2}}(\bar{\Omega})} \leq C \quad \text{for all } t > 1.$$

Then by interpolation inequalities, we have

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{C^2(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t \geq 1. \quad (70)$$

Now let us assume that Ω satisfies the condition (11). Then there exists $C > 0$ such that for all $\varphi \in H_0^1$ satisfying

$$-\Delta\varphi - b\varphi - \frac{n+2}{n-2}v_\infty^{\frac{4}{n-2}}\varphi = f \quad \text{in } \Omega,$$

there holds

$$\|\varphi\|_{H_0^1(\Omega)} \leq C\|f\|_{L^2(\Omega)}. \quad (71)$$

On one hand, using the equation of v_∞ , we have

$$\begin{aligned} & F(v(t)) - F_\infty \\ &= \int_{\Omega} \left(|\nabla v(\cdot, t)|^2 - bv(\cdot, t)^2 - \frac{n-2}{n}v(\cdot, t)^{\frac{2n}{n-2}} \right) - \int_{\Omega} \left(|\nabla v_\infty|^2 - bv_\infty^2 - \frac{n-2}{n}v_\infty^{\frac{2n}{n-2}} \right) \\ &\quad - 2 \int_{\Omega} \left(\nabla v_\infty \nabla(v(\cdot, t) - v_\infty) - bv_\infty(v(\cdot, t) - v_\infty) - v_\infty^{\frac{n+2}{n-2}}(v(\cdot, t) - v_\infty) \right) \\ &= \int_{\Omega} |\nabla(v(\cdot, t) - v_\infty)|^2 - b(v(\cdot, t) - v_\infty)^2 - \frac{n-2}{n} \left(v(\cdot, t)^{\frac{2n}{n-2}} - v_\infty^{\frac{2n}{n-2}} - \frac{2n}{n-2}(v(\cdot, t) - v_\infty) \right) \\ &\leq C\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)}^2. \end{aligned}$$

On the other hand, using the equation of v_∞ , we have

$$\begin{aligned} & \|\Delta v(\cdot, t) + bv(\cdot, t) + v(\cdot, t)^{\frac{n+2}{n-2}}\|_{L^2(\Omega)} \\ &= \|\Delta(v(\cdot, t) - v_\infty) + b(v(\cdot, t) - v_\infty) + v(\cdot, t)^{\frac{n+2}{n-2}} - v_\infty^{\frac{n+2}{n-2}}\|_{L^2(\Omega)} \\ &\geq \left\| \Delta(v(\cdot, t) - v_\infty) + b(v(\cdot, t) - v_\infty) + \frac{n+2}{n-2}v_\infty^{\frac{4}{n-2}}(v(\cdot, t) - v_\infty) \right\|_{L^2(\Omega)} \\ &\quad - \left\| v(\cdot, t)^{\frac{n+2}{n-2}} - v_\infty^{\frac{n+2}{n-2}} - \frac{n+2}{n-2}v_\infty^{\frac{4}{n-2}}(v(\cdot, t) - v_\infty) \right\|_{L^2(\Omega)} \\ &\geq \frac{1}{C}\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} - C \left\| v_\infty^{\max(0, \frac{4}{n-2}-1)} |v(\cdot, t) - v_\infty|^{\min(2, \frac{n+2}{n-2})} \right\|_{L^2(\Omega)} \\ &\geq \frac{1}{C}\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} - o(1)\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} \quad \text{for large } t \\ &\geq \frac{1}{C}\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} \quad \text{for large } t, \end{aligned}$$

where we used (71) in the second inequality. Combining these two inequalities together, we have

$$F(v(t)) - F_\infty \leq \|\Delta v(\cdot, t) + bv(\cdot, t) + v(\cdot, t)^{\frac{n+2}{n-2}}\|_{L^2(\Omega)}^2 = CM_2(t).$$

It follows that

$$\frac{d}{dt}(F(v(t)) - F_\infty) = -\frac{2(n-2)}{n+2}M_2(t) \leq -C(F(v(t)) - F_\infty),$$

and thus,

$$F(v(t)) - F_\infty \leq Ce^{-\gamma t}$$

for some $C > 0, \gamma > 0$. Hence, the proof of (64) will give

$$\int_T^\infty M_2(t)^{1/2} dt \leq Ce^{-\gamma T}.$$

From (67), we obtain that

$$\|v(\cdot, t) - v_\infty\|_{L^{\frac{2n}{n-2}}(\Omega)} \leq Ce^{-\gamma t}.$$

Then the proof of (70) gives

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{C^2(\bar{\Omega})} \leq Ce^{-\gamma t} \quad \text{for all } t \geq 1. \quad (72)$$

This finishes the proof of Theorem 1.1. \square

6 Subcritical case

In this last section, we consider the Sobolev subcritical case (5), and prove Theorem 1.2. The proof is similar to that of Theorem 1.1.

Proof of Theorem 1.2. First, we know from Proposition 6.2 in [18] that

$$\frac{1}{C}d(x) \leq v(\cdot, t) \leq Cd(x) \quad \text{for all } x \in \Omega, t > 1,$$

Secondly, it follows from Theorem 1.1 in [20] and Theorem 5.1 in [23] that there exists a nonzero stationary solution v_∞ of (5) such that

$$\lim_{t \rightarrow \infty} \|v(\cdot, t) - v_\infty\|_{C^3(\bar{\Omega})} = 0.$$

Let

$$\begin{aligned} F(v(t)) &= \int_{\Omega} \left(|\nabla v(x, t)|^2 - \frac{2}{p+1} v(x, t)^{p+1} \right) dx, \\ \mathcal{R} &= -v^{-p} \Delta v = 1 - \frac{p \partial_t v}{v}, \\ M_q(t) &= \int_{\Omega} |\mathcal{R} - 1|^q v^{p+1} dx, \end{aligned}$$

where $q \geq 1$. Note that

$$\frac{d}{dt} F(v(t)) = -2 \int_{\Omega} (\Delta v + v^p) \partial_t v = -2p \int_{\Omega} |\partial_t v|^2 v^{p-1} = -\frac{2}{p} M_2(t).$$

Hence $F(v(\cdot, t))$ decreases to $F(v_\infty)$ as $t \rightarrow \infty$. Furthermore,

$$dF(v) = -2\Delta v - 2v^p = 2(\mathcal{R} - 1)v^p.$$

Hence,

$$\|dF(v(\cdot, t))\|_{L^2(\Omega)} \leq CM_2(t)^{1/2}.$$

Then it follows from Proposition 6.1 in [20] that there exist $C > 0, T_0 > 0, \gamma > 0$ such that for all $t > T_0$, we have

$$F(v(t)) - F(v_\infty) \leq C\|dF(v(\cdot, t))\|_{L^2(\Omega)}^{1+\gamma} \leq CM_2(t)^{\frac{1+\gamma}{2}}.$$

Therefore, similar to the proof of Proposition 5.3, there exist $\theta \in (0, 1)$ and $C > 0$ such that for all $T > 1$, there holds

$$\int_T^\infty M_2(t)^{1/2} dt \leq CT^{-\theta},$$

Then it follows from the proof of (67) that

$$\|v(\cdot, t) - v_\infty\|_{L^{p+1}(\Omega)} \leq C \int_t^\infty M_2(s)^{1/2} ds \leq Ct^{-\theta}, \quad (73)$$

Using Theorem 5.1 in [23] (which is the regularity estimate) and interpolation inequalities, we have

$$\|v(\cdot, t) - v_\infty\|_{C^1(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t > 1.$$

Since $v(\cdot, t) \equiv v_\infty \equiv 0$ on $\partial\Omega$, we have for all $x \in \Omega$ that

$$\left| \frac{v(x, t) - v_\infty(x)}{v_\infty(x)} \right| \leq C\|v(\cdot, t) - v_\infty\|_{C^1(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t > 1.$$

That is,

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{L^\infty(\Omega)} \leq Ct^{-\theta} \quad \text{for all } t \geq 1.$$

Using Theorem 5.1 in [23] again, we have

$$\left\| \frac{v(\cdot, t)}{v_\infty} \right\|_{C^{1+p}(\bar{\Omega})} \leq C \quad \text{for all } t > 1.$$

Then by interpolation inequalities, we have

$$\left\| \frac{v(\cdot, t)}{v_\infty} - 1 \right\|_{C^2(\bar{\Omega})} \leq Ct^{-\theta} \quad \text{for all } t \geq 1. \quad (74)$$

Now let us assume that Ω satisfies the condition (6). Then there exists $C > 0$ such that for all $\varphi \in H_0^1$ satisfying

$$-\Delta\varphi - pv_\infty^{p-1}\varphi = f \quad \text{in } \Omega,$$

there holds

$$\|\varphi\|_{H_0^1(\Omega)} \leq C\|f\|_{L^2(\Omega)}. \quad (75)$$

As before, on one hand, using the equation of v_∞ , we have

$$F(v(t)) - F_\infty \leq C\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)}^2.$$

On the other hand, using the equation of v_∞ , we have

$$\begin{aligned}
& \|\Delta v(\cdot, t) + v(\cdot, t)^p\|_{L^2(\Omega)} \\
& \geq \|\Delta(v(\cdot, t) - v_\infty) + pv_\infty^{p-1}(v(\cdot, t) - v_\infty)\|_{L^2(\Omega)} \\
& \quad - \|v(\cdot, t)^p - v_\infty^p - pv_\infty^{p-1}(v(\cdot, t) - v_\infty)\|_{L^2(\Omega)} \\
& \geq \frac{1}{C}\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} - C\left\|v_\infty^{\max(0, p-2)}|v(\cdot, t) - v_\infty|^{\min(2, p)}\right\|_{L^2(\Omega)} \\
& \geq \frac{1}{C}\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} - o(1)\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} \quad \text{for large } t \\
& \geq \frac{1}{C}\|v(\cdot, t) - v_\infty\|_{H_0^1(\Omega)} \quad \text{for large } t,
\end{aligned}$$

where we used (75) in the second inequality. Combining these two inequalities together, we have

$$F(v(t)) - F_\infty \leq C\|\Delta v(\cdot, t) + v(\cdot, t)^p\|_{L^2(\Omega)}^2 \leq CM_2(t).$$

It follows that

$$\frac{d}{dt}(F(v(t)) - F_\infty) = -\frac{1}{p}M_2(t) \leq -C(F(v(t)) - F_\infty),$$

and thus,

$$F(v(t)) - F_\infty \leq Ce^{-\gamma t}$$

for some $C > 0, \gamma > 0$. Hence, the proof of (64) will give

$$\int_T^\infty M_2(t)^{1/2} dt \leq Ce^{-\gamma T}.$$

From (73), we obtain that

$$\|v(\cdot, t) - v_\infty\|_{L^{p+1}(\Omega)} \leq Ce^{-\gamma t}.$$

Then the proof of (74) gives

$$\left\|\frac{v(\cdot, t)}{v_\infty} - 1\right\|_{C^2(\bar{\Omega})} \leq Ce^{-\gamma t} \quad \text{for all } t \geq 1. \quad (76)$$

This finishes the proof of Theorem 1.2. \square

A Bubbles interactions

Here, we want to calculate

$$\begin{aligned}
I_1 &= \int_{\mathbb{R}^n} \left(\frac{\lambda_{1,\nu}}{1 + \lambda_{1,\nu}^2|x - x_{1,\nu}|^2}\right)^{\frac{n+2}{2}} \left(\frac{\lambda_{2,\nu}}{1 + \lambda_{2,\nu}^2|x - x_{2,\nu}|^2}\right)^{\frac{n-2}{2}} \\
&= \int_{\mathbb{R}^n} \left(\frac{\lambda_{1,\nu}}{1 + \lambda_{1,\nu}^2|x|^2}\right)^{\frac{n+2}{2}} \left(\frac{\lambda_{2,\nu}}{1 + \lambda_{2,\nu}^2|x - (x_{2,\nu} - x_{1,\nu})|^2}\right)^{\frac{n-2}{2}} \\
&= \int_{\mathbb{R}^n} \left(\frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2|x|^2}\right)^{\frac{n+2}{2}} \left(\frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2|x - e_\nu|^2}\right)^{\frac{n-2}{2}},
\end{aligned}$$

where

$$\tilde{\lambda}_{1,\nu} = \lambda_{1,\nu}|x_{2,\nu} - x_{1,\nu}|, \quad \tilde{\lambda}_{2,\nu} = \lambda_{2,\nu}|x_{2,\nu} - x_{1,\nu}|, \quad e_\nu = \frac{x_{2,\nu} - x_{1,\nu}}{|x_{2,\nu} - x_{1,\nu}|}.$$

By (32), we know that

$$\frac{\tilde{\lambda}_{1,\nu}}{\tilde{\lambda}_{2,\nu}} + \frac{\tilde{\lambda}_{2,\nu}}{\tilde{\lambda}_{1,\nu}} + \tilde{\lambda}_{1,\nu}\tilde{\lambda}_{2,\nu} \rightarrow \infty.$$

Recall that $\lambda_{1,\nu} \geq \lambda_{2,\nu}$ for all $\nu = 1, 2, \dots$. Hence, if $\{\tilde{\lambda}_{1,\nu}\}$ is bounded, then $\tilde{\lambda}_{2,\nu} \rightarrow 0$ and $\frac{\tilde{\lambda}_{1,\nu}}{\tilde{\lambda}_{2,\nu}} \rightarrow \infty$.

Case A: $\tilde{\lambda}_{2,\nu} \geq 1$. Then $\tilde{\lambda}_{1,\nu} \rightarrow \infty$, and thus

$$\begin{aligned} I_1 &= \int_{B_{1/4}} \left(\frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2|x|^2} \right)^{\frac{n+2}{2}} \left(\frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2|x - e_\nu|^2} \right)^{\frac{n-2}{2}} \\ &\quad + \int_{\mathbb{R}^n \setminus B_{1/4}} \left(\frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2|x|^2} \right)^{\frac{n+2}{2}} \left(\frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2|x - e_\nu|^2} \right)^{\frac{n-2}{2}} \\ &\leq C \tilde{\lambda}_{1,\nu}^{\frac{2-n}{2}} \tilde{\lambda}_{2,\nu}^{\frac{2-n}{2}}. \end{aligned}$$

Case B: Otherwise. Then

$$I_1 \leq \tilde{\lambda}_{2,\nu}^{\frac{n-2}{2}} \int_{\mathbb{R}^n} \left(\frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2|x|^2} \right)^{\frac{n+2}{2}} \leq C \tilde{\lambda}_{1,\nu}^{\frac{2-n}{2}} \tilde{\lambda}_{2,\nu}^{\frac{n-2}{2}}$$

We also want to calculate

$$\begin{aligned} I_2 &= \int_{\mathbb{R}^n} \left\{ \left(\frac{\lambda_{1,\nu}}{1 + \lambda_{1,\nu}^2|x - x_{1,\nu}|^2} \right) \vee \left(\frac{\lambda_{2,\nu}}{1 + \lambda_{2,\nu}^2|x - x_{2,\nu}|^2} \right) \right\}^2 \\ &\quad \cdot \left\{ \left(\frac{\lambda_{1,\nu}}{1 + \lambda_{1,\nu}^2|x - x_{1,\nu}|^2} \right) \wedge \left(\frac{\lambda_{2,\nu}}{1 + \lambda_{2,\nu}^2|x - x_{2,\nu}|^2} \right) \right\}^{n-2} dx \\ &= \int_{\mathbb{R}^n} \left\{ \left(\frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2|x|^2} \right) \vee \left(\frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2|x - e_\nu|^2} \right) \right\}^2 \\ &\quad \cdot \left\{ \left(\frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2|x|^2} \right) \wedge \left(\frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2|x - e_\nu|^2} \right) \right\}^{n-2} dx. \end{aligned}$$

Let $\varepsilon > 0$ be sufficiently small. The constant c in the below will be independent of ε .

Case A: $\tilde{\lambda}_{2,\nu} \geq 1$. Then $\tilde{\lambda}_{1,\nu} \rightarrow \infty$. We split it into two cases:

Case A1: $\lambda_{2,\nu} \geq \varepsilon \lambda_{1,\nu}$. Then

$$\begin{aligned} I_2 &\geq c \int_{B_{1/2}(e_\nu)} \left\{ \frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2|x - e_\nu|^2} \right\}^2 \cdot \left\{ \frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2|x|^2} \right\}^{n-2} dx \\ &\geq c \tilde{\lambda}_{2,\nu}^{-2} \tilde{\lambda}_{1,\nu}^{2-n} (1 + (5-n)^+(\ln \lambda_{2,\nu})) \\ &\geq c \tilde{\lambda}_{2,\nu}^{2-n} \tilde{\lambda}_{1,\nu}^{2-n} (1 - (5-n)^+(\ln \lambda_{2,\nu})) \quad (\text{if } n \geq 4). \end{aligned}$$

Case A2: $1 \leq \tilde{\lambda}_{2,\nu} < \varepsilon \tilde{\lambda}_{1,\nu}$. Then

$$\begin{aligned} I_2 &\geq c \int_{B_{1/2} \setminus B_{2\sqrt{\varepsilon}}} \left\{ \frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2 |x - e_\nu|^2} \right\}^2 \cdot \left\{ \frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2 |x|^2} \right\}^{n-2} dx \\ &\geq c \tilde{\lambda}_{2,\nu}^{-2} \tilde{\lambda}_{1,\nu}^{2-n} (1 - (5-n)^+ (\ln \varepsilon)) \\ &\geq c \tilde{\lambda}_{2,\nu}^{2-n} \tilde{\lambda}_{1,\nu}^{2-n} (1 - (5-n)^+ (\ln \varepsilon)) \quad (\text{if } n \geq 4). \end{aligned}$$

Case B: $\tilde{\lambda}_{2,\nu} \leq 1$. Then $\frac{\tilde{\lambda}_{1,\nu}}{\tilde{\lambda}_{2,\nu}} \rightarrow \infty$. We split it into three cases.

Case B1: $\tilde{\lambda}_{1,\nu} \tilde{\lambda}_{2,\nu} \geq \varepsilon^{-1}$. Then $\tilde{\lambda}_{1,\nu} \rightarrow \infty$. We have

$$\begin{aligned} I_2 &\geq c \int_{B_{1/2} \setminus B_{2\sqrt{\varepsilon}}} \tilde{\lambda}_{2,\nu}^2 \cdot \left\{ \frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2 |x|^2} \right\}^{n-2} dx \\ &\geq c \tilde{\lambda}_{2,\nu}^2 \tilde{\lambda}_{1,\nu}^{2-n} (1 - (5-n)^+ \ln(\varepsilon)) \\ &\geq c \tilde{\lambda}_{2,\nu}^{n-2} \tilde{\lambda}_{1,\nu}^{2-n} (1 - (5-n)^+ \ln(\varepsilon)) \quad (\text{if } n \geq 4). \end{aligned}$$

Case B2: $\tilde{\lambda}_{1,\nu} \tilde{\lambda}_{2,\nu} \leq \varepsilon^{-1}$. Then

$$\begin{aligned} I_2 &\geq c \int_B \frac{1}{\sqrt{1/(2\tilde{\lambda}_{1,\nu} \tilde{\lambda}_{2,\nu})}} \left\{ \frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2 |x|^2} \right\}^2 \left\{ \frac{\tilde{\lambda}_{2,\nu}}{1 + \tilde{\lambda}_{2,\nu}^2 |x - e_\nu|^2} \right\}^{n-2} dx \\ &\geq c \tilde{\lambda}_{2,\nu}^{n-2} \int_B \frac{1}{\sqrt{1/(2\tilde{\lambda}_{1,\nu} \tilde{\lambda}_{2,\nu})}} \left\{ \frac{\tilde{\lambda}_{1,\nu}}{1 + \tilde{\lambda}_{1,\nu}^2 |x|^2} \right\}^2 \\ &\geq c \tilde{\lambda}_{1,\nu}^{2-n} \tilde{\lambda}_{2,\nu}^{n-2} \left(1 + (5-n)^+ \ln(\tilde{\lambda}_{1,\nu}/\tilde{\lambda}_{2,\nu}) \right). \end{aligned}$$

Therefore,

$$I_1/\sqrt{I_2} = o(1) \quad \text{as } \nu \rightarrow \infty. \quad (77)$$

The following calculus lemma was used.

Lemma A.1. For $p > 2$, $0 \leq \varepsilon \leq 1$, we have

$$(1 + \varepsilon)^p \geq 1 + \varepsilon^p + p\varepsilon + c_p \varepsilon^2$$

and

$$(1 + \varepsilon)^p \geq 1 + \varepsilon^p + p\varepsilon^{p-1} + c_p \varepsilon$$

for some $c_p > 0$ independent of ε .

References

- [1] T. Aubin, *Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire*. J. Math. Pures Appl. (9) **55** (1976), no. 3, 269–296.
- [2] A. Bahri and J.M. Coron, *On a nonlinear elliptic equation involving the critical Sobolev exponent: The effect of the topology of the domain*. Comm. Pure Appl. Math. **41** (1988), 253–294.

- [3] J. B. Berryman and C. J. Holland, *Stability of the separable solution for fast diffusion*. Arch. Rat. Mech. Anal. **74** (1980), 379–388.
- [4] M. Bonforte and A. Figalli, *Sharp extinction rates for fast diffusion equations on generic bounded domains*. Comm. Pure Appl. Math. to appear.
- [5] M. Bonforte, G. Grillo and J. L. Vázquez, *Behaviour near extinction for the Fast Diffusion Equation on bounded domains*. J. Math. Pures Appl. **97** (2012), 1–38.
- [6] S. Brendle, *Convergence of the Yamabe flow for arbitrary initial energy*. J. Differential Geom. **69** (2005), no. 2, 217–278.
- [7] H. Brezis and T. Kato, *Remarks on the Schrödinger operator with singular complex potentials*. J. Math. Pures Appl. **58** (1979), 137–151.
- [8] H. Brezis and L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*. Comm. Pure Appl. Math. **36** (1983), 437–477.
- [9] H. Brezis and W. A. Strauss, *Semi-linear second-order elliptic equations in L^1* . J. Math. Soc. Japan **25** (1973), 565–590.
- [10] A. Carlotto, O. Chodosh and Y.A. Rubinstein, *Slowly converging Yamabe flows*. Geom. Topol. **19** (2015), no. 3, 1523–1568.
- [11] Y.Z. Chen and E. DiBenedetto, *On the local behavior of solutions of singular parabolic equations*. Arch. Rational Mech. Anal. **103** (1988), no. 4, 319–345.
- [12] X. Chen and X. Xu, *The scalar curvature flow on \mathbb{S}^n -perturbation theorem revisited*. Invent Math. **187** (2012), 395–506.
- [13] C. Cortázar, M. del Pino and M. Musso, *Green’s function and infinite-time bubbling in the critical nonlinear heat equation*. J. Eur. Math. Soc. (JEMS) **22** (2020), no. 1, 283–344.
- [14] P. Daskalopoulos, M. del Pino and N. Sesum, *Type II ancient compact solutions to the Yamabe flow*. J. Reine Angew. Math. **738** (2018), 1–71.
- [15] J. Dávila, M. del Pino and J. Wei, *Singularity formation for the two-dimensional harmonic map flow into \mathbb{S}^2* . Invent. Math. **219** (2020), no. 2, 345–466.
- [16] M. del Pino, M. Monica and J. Wei, *Existence and stability of infinite time bubble towers in the energy critical heat equation*. Anal. PDE. to appear.
- [17] M. del Pino and M. Sáez, *On the extinction profile for solutions of $u_t = \Delta u^{(n-2)/(n+2)}$* . Indiana Univ. Math. J. **50** (2001), no. 1, 611–628.
- [18] E. DiBenedetto, Y.C. Kwong, and V. Vespi, *Local space-analyticity of solutions of certain singular parabolic equations*. Indiana Univ. Math. J. **40** (2) (1991), 741–765.
- [19] O. Druet, *Elliptic equations with critical Sobolev exponents in dimension 3*. Ann. Inst. H. Poincaré Anal. Non Linéaire **19** (2002), no. 2, 125–142.
- [20] E. Feireisl and F. Simondon, *Convergence for semilinear degenerate parabolic equations in several space dimension*. J. Dyn. Diff. Eq. **12** (2000), 647–673.
- [21] V.A. Galaktionov and J.R. King, *Fast diffusion equation with critical Sobolev exponent in a ball*. Nonlinearity **15** (2002), no. 1, 173–188.
- [22] Z. Han, *Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent*. Ann. Inst. H. Poincaré Anal. Non Linéaire **8** (1991), 159–174.
- [23] T. Jin and J. Xiong, *Optimal boundary regularity for fast diffusion equations in bounded domains*. Preprint. arXiv:1910.05160.
- [24] T. Jin and J. Xiong, *Slow convergence for some fast diffusion equations in bounded domains*. In preparation.
- [25] M. Mayer, *A scalar curvature flow in low dimensions*. Calc. Var. Partial Differential Equations **56** (2017), no. 2, Paper No. 24, 41 pp.
- [26] O. Rey, *The role of the Green’s function in a nonlinear elliptic equation involving the critical Sobolev exponent*. J. Funct. Anal. **89** (1990), no. 1, 1–52.
- [27] J. Qing and G. Tian, *Bubbling of the heat flows for harmonic maps from surfaces*. Comm. Pure Appl. Math., **50** (1997), 295–310
- [28] J.-C. Saut and R. Temam, *Generic properties of nonlinear boundary value problems*. Comm. Partial

- Differential Equations **4** (1979), no. 3, 293–319.
- [29] R. Schoen, *Conformal deformation of a Riemannian metric to constant scalar curvature*. J. Differential Geom. **20** (1984), no. 2, 479–495.
 - [30] H. Schwetlick and M. Struwe, *Convergence of the Yamabe flow for “large” energies*. J. Reine Angew. Math. **562** (2003), 59–100.
 - [31] Y. Sire, J. Wei and Y. Zheng, *Extinction behaviour for the fast diffusion equations with critical exponent and Dirichlet boundary conditions*. Preprint. arXiv:2006.01308.
 - [32] M. Struwe, *A global compactness result for elliptic boundary value problems involving limiting nonlinearities*. Math. Z. **187** (1984), no. 4, 511–517.
 - [33] J.L. Vázquez, *Smoothing and decay estimates for nonlinear diffusion equations*. Oxford Lecture Notes in Maths. and its Applications, vol. 33. Oxford University Press, New York, 2006.
 - [34] J.L. Vázquez, *The Porous Medium Equation*. Mathematical Theory. Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, Oxford, 2007.

T. Jin

Department of Mathematics, The Hong Kong University of Science and Technology
Clear Water Bay, Kowloon, Hong Kong

Email: tianlingjin@ust.hk

J. Xiong

School of Mathematical Sciences, Laboratory of Mathematics and Complex Systems, MOE
Beijing Normal University, Beijing 100875, China

Email: jx@bnu.edu.cn