One-generated nilpotent terminal algebras

Ivan Kaygorodov^a, Abror Khudoyberdiyev^{b,c} & Aloberdi Sattarov^c

^a CMCC, Universidade Federal do ABC. Santo André, Brazil

^b National University of Uzbekistan, Tashkent, Uzbekistan

^c Institute of Mathematics Academy of Science of Uzbekistan, Tashkent, Uzbekistan

E-mail addresses:

Ivan Kaygorodov (kaygorodov.ivan@gmail.com) Abror Khudoyberdiyev (khabror@mail.ru) Aloberdi Sattarov (saloberdi90@mail.ru)

Abstract. We give an algebraic classification of complex 5-dimensional one-generated nilpotent terminal algebras.

Keywords: Nilpotent algebra, terminal algebra, algebraic classification, central extension.

MSC2010: 17A30.

INTRODUCTION

Algebraic classification (up to isomorphism) of algebras of small dimension from a certain variety defined by a family of polynomial identities is a classic problem in the theory of non-associative algebras. There are many results related to algebraic classification of small dimensional algebras in varieties of Jordan, Lie, Leibniz, Zinbiel and other algebras. Another interesting approach of studying algebras of a fixed dimension is to study them from a geometric point of view (that is, to study degenerations and deformations of these algebras). The results in which the complete information about degenerations of a certain variety is obtained are generally referred to as the geometric classification of the algebras of these variety. There are many results related to geometric classification of Jordan, Lie, Leibniz, Zinbiel and other algebras [1, 21, 32, 36, 37]. Another interesting direction is a study of one-generated objects. The description of one-generated finite groups is well-known: there is only one one-generated group of order n. In the case of algebras, there are some similar results, such that the description of *n*-dimensional one-generated nilpotent associative [12], noncommutative Jordan [22], Leibniz and Zinbiel algebras [38]. It was proven that there is only one *n*-dimensional one-generated nilpotent algebra in these varieties. But on the other side, as we can see in varieties of Novikov [26], assosymmetric [20], bicommutative [31], commutative [13], and terminal [28] algebras, there are more than one 4-dimensional one-generated nilpotent algebra from these varieties. One-generated Novikov algebras in dimensions 5 and 6 were studied in [5]. In the present paper, we give the algebraic classification of 5-dimensional one-generated nilpotent terminal algebras, which were introduced by Kantor in [23].

Our main results are summarized below.

Theorem A. Up to isomorphism, the variety of 4-dimensional complex one-generated nilpotent terminal algebras has infinitely many isomorphism classes, described explicitly in Table A in terms of 1 two-parametric family, 4 one-parameter families, and 5 additional isomorphism classes.

Theorem B. Up to isomorphism, the variety of complex 5-dimensional one-generated nilpotent terminal algebras has infinitely many isomorphism classes, described explicitly in Table C in terms of 1 threeparametric family, 18 two-parametric families, 38 one-parameter families, and 22 additional isomorphism classes.

In 1972, Kantor introduced the notion a conservative algebra as a generalization of Jordan algebras [23]. Unlike other classes of non-associative algebras, this class is not defined by a set of identities. To introduce the notion of a conservative algebra, we need some notation. Let \mathbb{V} be a vector space, let A be a linear operator on \mathbb{V} , and let B and C be bilinear operators on \mathbb{V} . For all $x, y, z \in \mathbb{V}$, put

Consider an algebra as a vector space \mathbb{V} over a field \mathbb{C} , together with an element μ of $\operatorname{Hom}(\mathbb{V} \otimes \mathbb{V}, \mathbb{V})$, so that $a \cdot b = \mu(a \otimes b)$. For an algebra $(\mathbb{V}, \mathcal{P})$ with a multiplication \mathcal{P} and $x \in \mathbb{V}$ we denote by $L_x^{\mathcal{P}}$ the operator of left multiplication by x. Thus, Kantor defines conservative algebras as follows:

Definition. An algebra $(\mathbb{V}, \mathcal{P})$, where \mathbb{V} is the vector space and \mathcal{P} is the multiplication, is called a conservative algebra if there is a new multiplication $\mathcal{P}^* : \mathbb{V} \times \mathbb{V} \to \mathbb{V}$ such that

(1)
$$[L_b^{\mathcal{P}}, [L_a^{\mathcal{P}}, \mathcal{P}]] = -[L_{\mathcal{P}^*(a,b)}^{\mathcal{P}}, \mathcal{P}], \text{ for all } a, b \in \mathbb{V}.$$

Simple calculations take us to the following identity with an additional multiplication \mathcal{P}^* , which must hold for all $a, b, x, y \in \mathbb{V}$:

(2)
$$b(a(xy) - (ax)y - x(ay)) - a((bx)y) + (a(bx))y + (bx)(ay) - a(x(by)) + (ax)(by) + x(a(by)) = -\mathcal{P}^*(a, b)(xy) + (\mathcal{P}^*(a, b)x)y + x(\mathcal{P}^*(a, b)y).$$

The class of conservative algebras is very vast. It includes all associative algebras, all quasi-associative algebras, all Jordan algebras, all Lie algebras, all (left) Leibniz algebras, all (left) Zinbiel algebras, and many other classes of algebras. On the other side, all conservative algebras are "rigid" (in sense of Cantarini and Kac [7]).

However, this class is very hard to study and for now even the basic general questions about it remain unanswered, so it is a good idea to study its subclasses which are sufficiently wide but easier to deal with. In [24] Kantor, studying the generalized TKK functor, introduced the class of terminal algebras which is a subclass of the class of conservative algebras.

Definition. An algebra $(\mathbb{V}, \mathcal{P})$, where \mathbb{V} is a vector space and \mathcal{P} is a multiplication, is called a terminal algebra if for all $a \in \mathbb{V}$ we have

$$(3) \qquad \qquad [[[\mathcal{P}, a], \mathcal{P}], \mathcal{P}] = 0.$$

Note that we can expand the relation (3), obtaining an identity of degree 4. Therefore, the class of terminal algebras is a variety. Also, about terminal and conservative algebras see, for example [3,25,28,29,33,35,39].

The following remark is obtained by straightforward calculations.

Remark 1. A commutative algebra satisfying (3) is a Jordan algebra.

Aside from Jordan algebras, the class of terminal algebras includes all Lie algebras, all (left) Leibniz algebras and some other types of algebras.

The following characterization of terminal algebras, proved by Kantor [24, Theorem 2], provides a description of this class as a subclass of the class of conservative algebras.

Remark 2. An algebra (\mathbb{V}, \mathbf{P}) is terminal if and only if it is conservative and the multiplication in the associated superalgebra \mathbf{P}^* can be defined by

(4)
$$\mathbf{P}^{*}(x,y) = \frac{2}{3}\mathbf{P}(x,y) + \frac{1}{3}\mathbf{P}(y,x).$$

The algebraic classification of nilpotent algebras will be obtained by the calculation of central extensions of algebras from the same variety which have a smaller dimension. Central extensions of algebras from various varieties were studied, for example, in [2, 30, 40]. Skjelbred and Sund [40] used central extensions of Lie algebras to classify nilpotent Lie algebras. The same method was applied to describe, all non-Lie central extensions of all 4-dimensional Malcev algebras [17], all non-associative central extensions of all 3-dimensional Jordan algebras [15], all anticommutative central extensions of 3-dimensional anticommutative algebras, all central extensions of 2-dimensional algebras and some others were described. One can also look at the classification of 3-dimensional nilpotent algebras [13], 4-dimensional nilpotent Novikov algebras [26], 4-dimensional nilpotent bicommutative algebras [31], 4-dimensional nilpotent commutative algebras in [13], 5-dimensional nilpotent associative commutative algebras [31], 5-dimensional nilpotent associative commutative algebras [31], 6-dimensional nilpotent algebras [16], 5-dimensional nilpotent Lie algebras [13], 5-dimensional nilpotent bicommutative algebras [13], 6-dimensional nilpotent Lie algebras [14], 6-dimensional nilpotent binary Lie algebras [18], 6-dimensional nilpotent Tortkara algebras [14], 6-dimensional nilpotent binary Lie algebras [11], 6-dimensional nilpotent anticommutative algebras [27], 8-dimensional nilpotent binary Lie algebras [6] and so on.

Motivation and contextualization. Given algebras A and B in the same variety, we write $A \rightarrow B$ and say that A *degenerates* to B, or that A is a *deformation* of B, if B is in the Zariski closure of the orbit of A (under the aforementioned base-change action of the general linear group). The study of degenerations of algebras is very related to deformation theory, in the sense of Gerstenhaber. It offers an insightful geometric perspective on the subject and has been studied in various papers. In particular, there are many results concerning degenerations of algebras of small dimensions in a variety defined by a set of identities. One of the main problems of the *geometric classification* of a variety of algebras is a description of its irreducible components. In the case studied in various papers finitely many orbits (i.e., isomorphism classes), the irreducible component are determined by the rigid algebras — algebras whose orbit closure is an irreducible component of the variety under consideration.

Observe that the algebraic classification of complex one-generated 5-dimensional nilpotent terminal algebras presented in this paper is the first step in the full algebraic classification of complex one-generated 5-dimensional nilpotent terminal algebras, which will halp to obtain the geometric classification of all complex 5-dimensional nilpotent terminal algebras.

1. PRELIMINARIES AND NOTATIONS

1.1. **Method of classification of nilpotent algebras.** Throughout this paper we use the notation and methods described in [4, 17] and adapted for the terminal case with some modifications (see also [19] for a discussion of extensions of algebras in an arbitrary nonassociative variety). Therefore, all statements in this subsection are given without proofs, which can be found in the papers cited above.

Let A be a terminal algebra over \mathbb{C} and \mathbb{V} a vector space of dimension *s* over the same base field. Then the space $Z^2_T(\mathbf{A}, \mathbb{V})$ is defined as the set of all maps $\theta : \mathbf{A} \times \mathbf{A} \to \mathbb{V}$ such that

$$\theta(b, a(xy) - (ax)y - x(ay)) - \theta(a, (bx)y) + \theta(a(bx), y) + \theta(bx, ay)$$

$$-\theta(a, x(by)) + \theta(ax, by) + \theta(x, a(by)) = -\theta(\mathbf{P}^*(a, b), xy) + \theta(\mathbf{P}^*(a, b)x, y) + \theta(x, \mathbf{P}^*(a, b)y),$$

where \mathbf{P}^* is given by (4). Its elements will be called *cocycles*. For a linear map $f : \mathbf{A} \to \mathbb{V}$ define $\delta f : \mathbf{A} \times \mathbf{A} \to \mathbb{V}$ by $\delta f(x, y) = f(xy)$. One can check that $\delta f \in Z^2_T(\mathbf{A}, \mathbb{V})$. Therefore, $B^2(\mathbf{A}, \mathbb{V}) = \{\theta = \delta f : f \in \text{Hom}(\mathbf{A}, \mathbb{V})\}$ is a subspace of $Z^2_T(\mathbf{A}, \mathbb{V})$ whose elements are called *coboundaries*. We define the *second cohomology space* $H^2_T(\mathbf{A}, \mathbb{V})$ as the quotient space $Z^2_T(\mathbf{A}, \mathbb{V})/B^2(\mathbf{A}, \mathbb{V})$. The equivalence class of $\theta \in Z^2(\mathbf{A}, \mathbb{V})$ in $H^2(\mathbf{A}, \mathbb{V})$ will be denoted by $[\theta]$.

For a bilinear map $\theta : \mathbf{A} \times \mathbf{A} \to \mathbb{V}$ define on the linear space $\mathbf{A}_{\theta} := \mathbf{A} \oplus \mathbb{V}$ a bilinear product $[-, -]_{\theta}$ by $[x + x', y + y']_{\theta} = xy + \theta(x, y)$ for all $x, y \in \mathbf{A}, x', y' \in \mathbb{V}$.

Then A_{θ} is an algebra called an *s*-dimensional central extension of A by \mathbb{V} . The following statement can be verified directly:

Lemma 1. The algebra \mathbf{A}_{θ} is terminal if and only if $\theta \in Z^2_{\mathbb{T}}(\mathbf{A}, \mathbb{V})$.

Recall that the *annihilator* of A is defined as the ideal Ann(A) = { $x \in A : xA + Ax = 0$ }. Given $\theta \in Z_T^2(A, \mathbb{V})$, we call the set Ann(θ) = { $x \in A : \theta(x, A) + \theta(A, x) = 0$ } the *annihilator* of θ .

Lemma 2. Ann $(\mathbf{A}_{\theta}) = (Ann(\theta) \cap Ann(\mathbf{A})) \oplus \mathbb{V}$.

Therefore, $0 \neq \mathbb{V} \subseteq Ann(\mathbf{A}_{\theta})$. The following lemma shows that every algebra with a nonzero annihilator can be obtained in the way described above:

Lemma 3. Let \mathbf{A} be an *n*-dimensional terminal algebra such that $\dim(\operatorname{Ann}(\mathbf{A})) = m \neq 0$. Then there exists, up to isomorphism, a unique (n - m)-dimensional terminal algebra \mathbf{A}' and a bilinear map $\theta \in Z^2_T(\mathbf{A}, \mathbb{V})$ with $\operatorname{Ann}(\mathbf{A}') \cap \operatorname{Ann}(\theta) = 0$, where \mathbb{V} is a vector space of dimension m, such that $\mathbf{A} \cong \mathbf{A}'_{\theta}$ and $\mathbf{A}/\operatorname{Ann}(\mathbf{A}) \cong \mathbf{A}'$.

In particular, any finite-dimensional nilpotent algebra is a central extension of another nilpotent algebra of strictly smaller dimension. Thus, to classify all nilpotent terminal algebras of a fixed dimension, we need to classify cocycles of nilpotent terminal algebras \mathbf{A}' of smaller dimension (with an additional condition $\operatorname{Ann}(\mathbf{A}') \cap \operatorname{Ann}(\theta) = 0$) and central extensions that arise from them.

We can reduce the class of extensions that we need to consider.

Definition. Let A be an algebra and I be a subspace of Ann(A). If $A = A_0 \oplus I$ then I is called an annihilator component of A. A central extension of an algebra A without an annihilator component is called a non-split central extension.

Clearly, we are only interested in non-split extensions (in the contrary case we can cut off annihilator components lying in \mathbb{V} until we obtain a non-split extension).

Let us fix a basis e_1, \ldots, e_s of \mathbb{V} , and $\theta \in Z^2_T(\mathbf{A}, \mathbb{V})$. Then θ can be uniquely written as $\theta(x, y) =$ $\theta(x,y) = \sum_{i=1}^{s} \theta_i(x,y) e_i$, where $\theta_i \in \mathbb{Z}^2_{\mathbb{T}}(\mathbf{A},\mathbb{C})$. Moreover, $\operatorname{Ann}(\theta) = \operatorname{Ann}(\theta_1) \cap \operatorname{Ann}(\theta_2) \cap \cdots \cap \operatorname{Ann}(\theta_s)$. Further, $\theta \in B^2(\mathbf{A}, \mathbb{V})$ if and only if all $\theta_i \in B^2(\mathbf{A}, \mathbb{C})$. Using this presentation, one can determine whether the extension corresponding to a cocycle θ is split:

Lemma 4. [17, Lemma 13] Let $\theta(x, y) = \sum_{i=1}^{s} \theta_i(x, y) e_i \in \mathbb{Z}^2_T(\mathbf{A}, \mathbb{V})$ be such that $\operatorname{Ann}(\theta) \cap \operatorname{Ann}(\mathbf{A}) = 0$. Then \mathbf{A}_{θ} has an annihilator component if and only if $[\theta_1], [\theta_2], \dots, [\theta_s]$ are linearly dependent in $\mathrm{H}^2_{\mathrm{T}}(\mathbf{A}, \mathbb{C})$.

Some cocycles give rise to isomorphic extensions:

Lemma 5. Let $\theta, \vartheta \in Z^2_T(\mathbf{A}, \mathbb{V})$ be such that $[\theta] = [\vartheta]$. Then $\mathbf{A}_{\theta} \cong \mathbf{A}_{\vartheta}$.

By above, the isomorphism classes of extensions correspond to certain equivalence classes on $H^2_T(\mathbf{A}, \mathbb{V})$. These classes can be given in terms of actions of certain groups on this space. In particular, let Aut(A) be the automorphism group of A, let $\phi \in Aut(A)$, and let $\psi \in GL(\mathbb{V})$. For $\theta \in Z^2_T(A, \mathbb{V})$ define

$$\phi\theta(x,y) = \theta(\phi(x),\phi(y)), \quad \psi\theta(x,y) = \psi(\theta(x,y)).$$

Then $\phi\theta, \psi\theta \in Z^2_T(\mathbf{A}, \mathbb{V})$. Hence, $Aut(\mathbf{A})$ and $GL(\mathbb{V})$ act on $Z^2_T(\mathbf{A}, \mathbb{V})$. It is easy to verify that $B^2(\mathbf{A}, \mathbb{V})$ is invariant under both actions. Therefore, we have induced actions on $H^2_T(\mathbf{A}, \mathbb{V})$.

Lemma 6. Let $\theta, \vartheta \in Z^2_T(\mathbf{A}, \mathbb{V})$ be such that $Ann(\mathbf{A}_{\theta}) = Ann(\mathbf{A}_{\vartheta}) = \mathbb{V}$. Then $\mathbf{A}_{\theta} \cong \mathbf{A}_{\vartheta}$ if and only if there exist $a \phi \in Aut(\mathbf{A}), \psi \in GL(\mathbb{V})$ such that $[\phi \theta] = [\psi \vartheta]$.

Now we rewrite the above lemma in a form more suitable for computations.

Let U be a finite-dimensional vector space over C. The Grassmannian $G_k(U)$ is the set of all kdimensional linear subspaces of V. Let $G_s(\mathrm{H}^2_{\mathrm{T}}(\mathbf{A},\mathbb{C}))$ be the Grassmannian of subspaces of dimension s in $H^2_T(\mathbf{A}, \mathbb{C})$. There is a natural action of $Aut(\mathbf{A})$ on $G_s(H^2_T(\mathbf{A}, \mathbb{C}))$: for $\phi \in Aut(\mathbf{A}), W =$ $\langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle \in G_s(\mathrm{H}^2_{\mathrm{T}}(\mathbf{A}, \mathbb{C})),$ define

$$\phi W = \langle [\phi \theta_1], [\phi \theta_2], \dots, [\phi \theta_s] \rangle.$$

Note that this action is compatible with the action of $Aut(\mathbf{A})$ on $H^2_T(\mathbf{A}, \mathbb{V})$ and the above presentation of a cocycle as a collection of s elements of $H^2_T(\mathbf{A}, \mathbb{C})$. Denote the orbit of W under the action of Aut(A) by Orb(W). It is easy to check that given two bases of a subspace

$$W = \langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle = \langle [\vartheta_1], [\vartheta_2], \dots, [\vartheta_s] \rangle \in G_s(\mathrm{H}^2(\mathbf{A}, \mathbb{C})),$$

we have $\bigcap_{i=1}^{s} \operatorname{Ann}(\theta_i) \cap \operatorname{Ann}(\mathbf{A}) = \bigcap_{i=1}^{s} \operatorname{Ann}(\vartheta_i) \cap \operatorname{Ann}(\mathbf{A})$. Therefore, we can introduce the set

$$T_s(\mathbf{A}) = \left\{ W = \langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle \in G_s(\mathrm{H}^2(\mathbf{A}, \mathbb{C})) : \bigcap_{i=1}^s \mathrm{Ann}(\theta_i) \cap \mathrm{Ann}(\mathbf{A}) = 0 \right\},$$

which is stable under the action of $Aut(\mathbf{A})$.

Let us denote by $E(\mathbf{A}, \mathbb{V})$ the set of all *non-split* central extensions of \mathbf{A} by \mathbb{V} . By Lemmas 2 and 4, we can write

$$E(\mathbf{A}, \mathbb{V}) = \left\{ \mathbf{A}_{\theta} : \theta(x, y) = \sum_{i=1}^{s} \theta_{i}(x, y) e_{i} \text{ and } \langle [\theta_{1}], [\theta_{2}], \dots, [\theta_{s}] \rangle \in T_{s}(\mathbf{A}) \right\}.$$

Also, we have the next result, which can be proved in the same manner as [17, Lemma 17].

Lemma 7. Let $\theta(x, y) = \sum_{i=1}^{s} \theta_i(x, y) e_i$ and $\vartheta(x, y) = \sum_{i=1}^{s} \vartheta_i(x, y) e_i$ be such that $\mathbf{A}_{\theta}, \mathbf{A}_{\vartheta} \in E(\mathbf{A}, \mathbb{V})$. Then the algebras \mathbf{A}_{θ} and \mathbf{A}_{ϑ} are isomorphic if and only if

$$\operatorname{Orb}\langle [\theta_1], [\theta_2], \dots, [\theta_s] \rangle = \operatorname{Orb}\langle [\vartheta_1], [\vartheta_2], \dots, [\vartheta_s] \rangle.$$

Thus, there exists a one-to-one correspondence between the set of $Aut(\mathbf{A})$ -orbits on $T_s(\mathbf{A})$ and the set of isomorphism classes of $E(\mathbf{A}, \mathbb{V})$. Consequently, we have a procedure that allows us, given a terminal algebra \mathbf{A}' of dimension n, to construct all non-split central extensions of \mathbf{A}' . This procedure is as follows:

Procedure

- (1) For a given (nilpotent) terminal algebra \mathbf{A}' of dimension n s, determine $T_s(\mathbf{A}')$ and $\operatorname{Aut}(\mathbf{A}')$.
- (2) Determine the set of $Aut(\mathbf{A}')$ -orbits on $T_s(\mathbf{A}')$.
- (3) For each orbit, construct the terminal algebra corresponding to one of its representatives.

1.2. Notations. Let us introduce the following notations. Let \mathbf{A} be a terminal algebra with a basis e_1, e_2, \ldots, e_n . Then by Δ_{ij} we will denote the bilinear form $\Delta_{ij} : \mathbf{A} \times \mathbf{A} \longrightarrow \mathbb{C}$ with $\Delta_{ij}(e_l, e_m) = \delta_{il}\delta_{jm}$. The set $\{\Delta_{ij} : 1 \le i, j \le n\}$ is a basis for the linear space of bilinear forms on \mathbf{A} , so every $\theta \in \mathbb{Z}^2(\mathbf{A}, \mathbb{V})$ can be uniquely written as $\theta = \sum_{1 \le i, j \le n} c_{ij}\Delta_{ij}$, where $c_{ij} \in \mathbb{C}$. Let us fix the following notations:

 \mathbf{T}_{i}^{i} — *j*th *i*-dimensional one-generated terminal algebra.

1.3. The algebraic classification of 3-dimensional one-generated nilpotent terminal algebras. Observe that a 2-dimensional one-generated nilpotent algebra is isomorphic to

$$\mathbf{T}_{01}^2$$
 : $e_1e_1 = e_2$.

We have the following list of all 3-dimensional one-generated nilpotent terminal algebras with an annihilator of codimension 1 or 2:

(5)
$$\begin{array}{rcl} \mathbf{T}_{01}^{3} & : & e_{1}e_{1} = e_{2} & e_{1}e_{2} = e_{3} \\ \mathbf{T}_{02}^{3}(\lambda) & : & e_{1}e_{1} = e_{2} & e_{1}e_{2} = \lambda e_{3} & e_{2}e_{1} = e_{3} \end{array}$$

1.4. Automorphism and cohomology groups of 3-dimensional one-generated nilpotent terminal algebras. In the following table we give the description of the automorphism groups and the second cohomology spaces of 3-dimensional one-generated nilpotent terminal algebras.

Α	$\operatorname{Aut}\left(\mathbf{A} ight)$	$\mathrm{Z}_{\mathrm{T}}^2(\mathbf{A})$	$\mathrm{H}_{\mathrm{T}}^2(\mathbf{A})$
\mathbf{T}_{01}^3	$\begin{pmatrix} x & 0 & 0 \\ y & x^2 & 0 \\ z & xy & x^3 \end{pmatrix}$	$\left\langle \begin{array}{c} \Delta_{11}, \Delta_{12}, \Delta_{13}, \\ \Delta_{21}, \Delta_{22} - 3\Delta_{31} \end{array} \right\rangle$	$\left\langle \begin{array}{c} [\Delta_{13}], [\Delta_{21}], \\ [\Delta_{22}] - 3[\Delta_{31}] \end{array} \right\rangle$

$\mathbf{T}^3_{02}(\lambda)_{\lambda eq 0}$	$ \begin{pmatrix} x & 0 & 0 \\ y & x^2 & 0 \\ z & (\lambda+1)xy & x^3 \end{pmatrix} $	$\left\langle \begin{array}{c} \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \\ \Delta_{21}, \frac{1-\lambda}{3} \Delta_{22} + \Delta_{31} \end{array} \right\rangle$	$\left\langle \begin{array}{c} [\Delta_{12}], [\Delta_{13}] + [\Delta_{22}], \\ \frac{1-\lambda}{3} [\Delta_{22}] + [\Delta_{31}] \end{array} \right\rangle$
${f T}_{02}^3(0)$	$\begin{pmatrix} x & 0 & 0 \\ y & x^2 & 0 \\ z & xy & x^3 \end{pmatrix}$	$\left\langle \begin{array}{c} \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \\ \Delta_{21}, \frac{1}{3}\Delta_{22} + \Delta_{31}, \Delta_{23} \end{array} \right\rangle$	$\left\langle \begin{array}{c} [\Delta_{12}], [\Delta_{13}] + [\Delta_{22}], \\ \frac{1}{3} [\Delta_{22}] + [\Delta_{31}], [\Delta_{23}] \end{array} \right\rangle$

2. CLASSIFICATION OF 4-DIMENSIONAL ONE-GENERATED NILPOTENT TERMINAL ALGEBRAS

Thanks to [4], we have only one 4-dimensional one-generated nilpotent terminal algebra with 2dimensional annihilator:

$$\mathbf{T}_{01}^4$$
 : $e_1e_1 = e_2$ $e_1e_2 = e_4$ $e_2e_1 = e_3$.

2.1. Central extensions of T_{01}^3 . Let us use the following notations:

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = [\Delta_{21}], \quad \nabla_3 = [\Delta_{22}] - 3[\Delta_{31}].$$

Take $\theta = \sum_{i=1}^{3} \alpha_i \nabla_i \in H^2_T(\mathbf{T}^3_{01})$. For an element

$$\phi = \begin{pmatrix} x & 0 & 0 \\ y & x^2 & 0 \\ z & xy & x^3 \end{pmatrix} \in \operatorname{Aut} (\mathbf{T}_{01}^3),$$

we have

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_1 \\ \alpha_2 & \alpha_3 & 0 \\ -3\alpha_3 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha^{**} & \alpha_1^* \\ \alpha_2^* & \alpha_3^* & 0 \\ -3\alpha_3^* & 0 & 0 \end{pmatrix},$$

where

$$\alpha_1^* = x^4 \alpha_1, \quad \alpha_2^* = x^2 (x \alpha_2 - 2y \alpha_3), \quad \alpha_3^* = x^4 \alpha_3.$$

2.1.1. 1-dimensional central extensions of T_{01}^3 . Since we are interested only in new algebras, we have the following cases:

- (1) if $\alpha_3 \neq 0$, then by choosing $y = \frac{x\alpha_2}{2\alpha_3}$, we have the representative $\langle \alpha \nabla_1 + \nabla_3 \rangle$. (2) if $\alpha_3 = 0$, then $\alpha_1 \neq 0$, and we have the representative $\langle \nabla_1 \rangle$ and $\langle \nabla_1 + \nabla_2 \rangle$ depending on whether $\alpha_2 = 0$ or not.

Hence, we have the following 4-dimensional algebras:

$\mathbf{T}_{02}^4(lpha)$:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = \alpha e_4$	$e_2 e_2 = e_4$	$e_3e_1 = -3e_4$
\mathbf{T}_{03}^4	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$		
\mathbf{T}_{04}^4	:	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	$e_2 e_1 = e_4.$	

2.1.2. 2-dimensional central extensions of T_{01}^3 . Consider the vector space generated by the following two cocycles:

$$\begin{aligned} \theta_1 &= \alpha_1 \nabla_1 + \alpha_2 \nabla_2 + \alpha_3 \nabla_3 \\ \theta_2 &= \beta_1 \nabla_1 + \beta_2 \nabla_2. \end{aligned}$$

If $\alpha_3 = 0$, then we have the representative $\langle \nabla_1, \nabla_2 \rangle$. If $\alpha_3 \neq 0$, then:

- (1) if $\beta_1 = 0$, then we have the family of representatives $\langle \nabla_2, \alpha \nabla_1 + \nabla_3 \rangle$.
- (2) if $\beta_1 \neq 0$, then we have the representatives $\langle \nabla_1, \nabla_3 \rangle$, and $\langle \nabla_1 + \nabla_2, \nabla_3 \rangle$ depending on whether $\beta_2 = 0$ or not.

Hence, we have the following 5-dimensional algebras:

\mathbf{T}_{01}^5	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	$e_2 e_1 = e_5$		
$\mathbf{T}_{02}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = \alpha e_5$	$e_2 e_1 = e_4$	$e_2 e_2 = e_5$	$e_3e_1 = -3e_5$
\mathbf{T}_{03}^5	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	$e_2 e_2 = e_5$	$e_3e_1 = -3e_5$	
\mathbf{T}_{04}^5	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	$e_2e_1 = e_4$	$e_2 e_2 = e_5$	$e_3e_1 = -3e_5.$

2.2. Central extensions of $T^3_{02}(\lambda)_{\lambda\neq 0}$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = \frac{1-\lambda}{3} [\Delta_{22}] + [\Delta_{31}]$$

Take $\theta = \sum_{i=1}^{3} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^3_{02}(\lambda))$. For an element

$$\phi = \begin{pmatrix} x & 0 & 0 \\ y & x^2 & 0 \\ z & (\lambda + 1)xy & x^3 \end{pmatrix} \in \operatorname{Aut} (\mathbf{T}_{01}^3),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 \\ 0 & \alpha_2 + \frac{1-\lambda}{3}\alpha_3 & 0 \\ \alpha_3 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha_1^* + \lambda \alpha^{**} & \alpha_2^* \\ \alpha^{**} & \alpha_2^* + \frac{1-\lambda}{3}\alpha_3^* & 0 \\ \alpha_3^* & 0 & 0 \end{pmatrix},$$

where

$$\alpha_1^* = x^3 \alpha_1 + x^2 y \left(2\alpha_2 - \frac{2\lambda^2 + 5\lambda - 1}{3} \alpha_3 \right), \qquad \alpha_2^* = x^4 \alpha_2, \qquad \alpha_3^* = x^4 \alpha_3.$$

2.2.1. 1-dimensional central extensions of $T^3_{02}(\lambda)_{\lambda\neq 0}$. Since we are interested only in new algebras, we have the following cases:

(1) if $\alpha_3 \neq 0$, then: (a) if $\alpha_2 \neq \frac{2\lambda^2 + 5\lambda - 1}{6}\alpha_3$, then by choosing $y = \frac{3x\alpha_1}{6\alpha_2 - (2\lambda^2 + 5\lambda - 1)\alpha_3}$, we have the representative $\langle \alpha \nabla_2 + \nabla_3 \rangle$, with $\alpha \neq \frac{2\lambda^2 + 5\lambda - 1}{6}$. (b) if $\alpha_2 = \frac{2\lambda^2 + 5\lambda - 1}{6}\alpha_3$, then we have the representatives $\langle \frac{2\lambda^2 + 5\lambda - 1}{6}\nabla_2 + \nabla_3 \rangle$ and $\langle \nabla_1 + \frac{2\lambda^2 + 5\lambda - 1}{6}\nabla_2 + \nabla_3 \rangle$,

depending on whether $\alpha_1 = 0$ or not.

(2) if $\alpha_3 = 0$, then $\alpha_2 \neq 0$, and by choosing $y = -\frac{x\alpha_1}{2\alpha_2}$, we have the representative $\langle \nabla_2 \rangle$. Hence, we have the following distinct orbits:

$$\langle \alpha \nabla_2 + \nabla_3 \rangle, \quad \langle \nabla_1 + \frac{2\lambda^2 + 5\lambda - 1}{6} \nabla_2 + \nabla_3 \rangle, \quad \langle \nabla_2 \rangle$$

2.2.2. 2-dimensional central extensions of $T_{02}^3(\lambda)_{\lambda\neq 0}$. Consider the vector space generated by the following two cocycles:

$$\begin{aligned} \theta_1 &= \alpha_1 \nabla_1 + \alpha_2 \nabla_2 + \alpha_3 \nabla_3 \\ \theta_2 &= \beta_1 \nabla_1 + \beta_2 \nabla_2. \end{aligned}$$

If $\alpha_3 = 0$, then we have the representative $\langle \nabla_1, \nabla_2 \rangle$. If $\alpha_3 \neq 0$, then:

- (1) if $\beta_2 = 0$, then we have the family of representatives $\langle \nabla_1, \alpha \nabla_2 + \nabla_3 \rangle$.
- (2) if $\beta_2 \neq 0$, then choosing $y = -\frac{x\beta_1}{2\beta_2}$, we have the representatives $\langle \nabla_2, \nabla_3 \rangle$ and $\langle \nabla_2, \nabla_1 + \nabla_3 \rangle$ depending on whether $\alpha_1^* = 0$ or not.

Hence, we have the following distinct orbits:

$$\langle \nabla_1, \nabla_2 \rangle, \quad \langle \nabla_1, \alpha \nabla_2 + \nabla_3 \rangle, \quad \langle \nabla_2, \nabla_3 \rangle, \quad \langle \nabla_2, \nabla_1 + \nabla_3 \rangle.$$

2.3. Central extensions of $T_{02}^3(0)$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = \frac{1}{3}[\Delta_{22}] + [\Delta_{31}], \quad \nabla_4 = [\Delta_{23}].$$

Take $\theta = \sum_{i=1}^{4} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^3_{01}(0))$. For an element

$$\phi = \begin{pmatrix} x & 0 & 0 \\ y & x^2 & 0 \\ z & xy & x^3 \end{pmatrix} \in \operatorname{Aut} \left(\mathbf{T}_{01}^3(0) \right),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 \\ 0 & \alpha_2 + \frac{1}{3}\alpha_3 & \alpha_4 \\ \alpha_3 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha_1^* & \alpha_2^* \\ \alpha^{**} & \alpha_2^* + \frac{1}{3}\alpha_3^* & \alpha_4^* \\ \alpha_3^* & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl}
\alpha_1^* &=& x^3\alpha_1 + x^2y(2\alpha_2 + \frac{1}{3}\alpha_3) + xy^2\alpha_4, & \alpha_2^* &=& x^4\alpha_2 + x^3y\alpha_4, \\
\alpha_3^* &=& x^4\alpha_3, & \alpha_4^* &=& x^5\alpha_4.
\end{array}$$

2.3.1. 1-dimensional central extensions of $\mathbf{T}_{02}^3(0)$.

(1) if $\alpha_4 = 0$, then:

- (a) if $\alpha_3 \neq 0, \alpha_2 \neq -\frac{1}{6}\alpha_3$, then by choosing $y = \frac{3x\alpha_1}{6\alpha_2 + \alpha_3}$, we have the representative $\langle \alpha \nabla_2 + \nabla_3 \rangle$, with $\alpha \neq -\frac{1}{6}$.
- (b) if $\alpha_3 \neq 0$, $\alpha_2 = -\frac{1}{6}\alpha_3$, then we have the representatives $\langle -\frac{1}{6}\nabla_2 + \nabla_3 \rangle$ and $\langle \nabla_1 \frac{1}{6}\nabla_2 + \nabla_3 \rangle$, depending on whether $\alpha_1 = 0$ or not.
- (c) if $\alpha_3 = 0$, then $\alpha_2 \neq 0$, and by choosing $y = -\frac{x\alpha_1}{2\alpha_2}$, we have the representative $\langle \nabla_2 \rangle$.

(2) if $\alpha_4 \neq 0$, then by choosing $y = -\frac{x\alpha_2}{\alpha_4}$, we have the representatives

$$\langle
abla_4
angle, \ \langle
abla_1 +
abla_4
angle ext{ and } \langle lpha
abla_1 +
abla_3 +
abla_4
angle$$

depending on whether $\alpha_3 = 0$ and $3\alpha_1\alpha_4 - 3\alpha_2^2 - \alpha_2\alpha_3 = 0$ or not. Hence, we have the following distinct orbits:

$$\langle \alpha \nabla_2 + \nabla_3 \rangle, \quad \langle \nabla_1 - \frac{1}{6} \nabla_2 + \nabla_3 \rangle, \quad \langle \nabla_2 \rangle, \quad \langle \nabla_4 \rangle, \quad \langle \nabla_1 + \nabla_4 \rangle, \quad \langle \alpha \nabla_1 + \nabla_3 + \nabla_4 \rangle.$$

Summarizing the cases $\lambda \neq 0$ and $\lambda = 0$, we have the following new 4-dimensional algebras:

$\mathbf{T}_{05}^4(\lambda, \alpha)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3$	$e_1e_3 = \alpha e_4$	
	$e_2e_1 = e_3$	$e_2 e_2 = \left(\alpha + \frac{1-\lambda}{3}\right) e_4$	$e_3e_1 = e_4$	
$\mathbf{T}_{06}^4(\lambda)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3 + e_4$	$e_1 e_3 = \frac{2\lambda + 5\lambda - 1}{6} e_4$	
	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{(2\lambda+1)(\lambda+1)}{6} e_4$	$e_3e_1 = e_4$	
$\mathbf{T}_{07}^4(\lambda)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3$	$e_1e_3 = e_4$ $e_2e_1 = e_3$ $e_2e_2 = e_4$	
\mathbf{T}_{08}^4	$e_1e_1 = e_2$	$e_2 e_1 = e_3$	$e_2e_3 = e_4$	
\mathbf{T}_{09}^4	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_2 e_1 = e_3$ $e_2 e_3 = e_4$	
$\mathbf{T}_{10}^4(\alpha)$	$e_1e_1 = e_2$	$e_1 e_2 = \alpha e_4$	$e_2e_1 = e_3$ $e_2e_2 = \frac{1}{3}e_4$ $e_2e_3 = e_4$ e_3e_3	$e_1 = e_4$

2.3.2. 2-dimensional central extensions of $T_{02}^3(0)$. Consider the vector space generated by the following two cocycles

$$\begin{aligned} \theta_1 &= \alpha_1 \nabla_1 + \alpha_2 \nabla_2 + \alpha_3 \nabla_3 + \alpha_4 \nabla_4 \\ \theta_2 &= \beta_1 \nabla_1 + \beta_2 \nabla_2 + \beta_3 \nabla_3. \end{aligned}$$

Consider the following cases:

(1) if $\alpha_4 = 0$, then similarly to the case of $\lambda \neq 0$, we have the representatives

$$\langle \nabla_1, \nabla_2 \rangle, \ \langle \nabla_1, \alpha \nabla_2 + \nabla_3 \rangle, \ \langle \nabla_2, \nabla_3 \rangle \text{ and } \langle \nabla_2, \nabla_1 + \nabla_3 \rangle.$$

(2) if $\alpha_4 \neq 0$, then:

(a) if $\beta_3 = \beta_2 = 0$, then choosing $y = -\frac{x\alpha_2}{\alpha_4}$, we have the representatives $\langle \nabla_1, \nabla_4 \rangle, \langle \nabla_1, 3\nabla_3 + \nabla_4 \rangle$ depending on whether $\alpha_3 = 0$ or not.

depending on whether $\alpha_3 = 0$ or not. (b) $\beta_3 = 0, \beta_2 \neq 0$, then choosing $y = -\frac{x\beta_1}{2\beta_2}$, we have the representatives

$$\langle
abla_2,
abla_4
angle, \ \langle
abla_2,
abla_1 +
abla_4
angle \ \text{and} \ \langle
abla_2, 3
abla_3 +
abla_4
angle$$

depending on whether $\alpha_1^*=0,\,\alpha_3^*=0$ or not.

(c) $\beta_3 \neq 0$, then we may suppose $\alpha_3 = 0$ and choosing $y = -\frac{x\alpha_2}{\alpha_4}$, we have the representatives

$$\langle \alpha \nabla_2 + 3 \nabla_3, \nabla_4 \rangle, \ \langle \nabla_1 + \alpha \nabla_2 + 3 \nabla_3, \nabla_4 \rangle, \ \langle \beta \nabla_1 + \alpha \nabla_2 + 3 \nabla_3, \nabla_1 + \nabla_4 \rangle.$$

Summarizing the cases $\lambda \neq 0$ and $\lambda = 0$, we have the following 5-dimensional algebras:

$\mathbf{T}_{05}^5(\lambda)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_4$	$e_1e_3 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = e_5$
$\mathbf{T}_{06}^5(\lambda,lpha)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_4$	$e_1e_3 = \alpha e_5$		
	$e_2 e_1 = e_3$	$e_2 e_2 = \left(\alpha + \frac{1-\lambda}{3}\right) e_5$	$e_3e_1 = e_5$		

$\mathbf{T}_{07}^5(\lambda)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3$	$e_1e_3 = e_4$		
	$e_2 e_1 = e_3$	$e_2 e_2 = e_4 + \frac{1-\lambda}{3} e_5$	$e_3e_1 = e_5$		
$\mathbf{T}_{08}^5(\lambda)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_5$	$e_1e_3 = e_4$		
	$e_2 e_1 = e_3$	$e_2 e_2 = e_4 + \frac{1-\lambda}{3}e_5$	$e_3e_1 = e_5$		
T_{09}^5	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_2e_1 = e_3$	$e_2e_3 = e_5$	
\mathbf{T}_{10}^5	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_2e_1 = e_3$	$e_2 e_2 = e_5$	
	$e_2 e_3 = e_5$	$e_3e_1 = 3e_5$			
\mathbf{T}_{11}^5	$e_1e_1 = e_2$	$e_1e_3 = e_4$	$e_2e_1 = e_3$	$e_2 e_2 = e_4$	$e_2e_3 = e_5$
\mathbf{T}_{12}^5	$e_1e_1 = e_2$	$e_1e_2 = e_5$	$e_1e_3 = e_4$	$e_2e_1 = e_3$	
	$e_2 e_2 = e_4$	$e_2e_3=e_5$			
$\mathbf{T}_{13}^5(lpha)$	$e_1e_1 = e_2$	$e_1 e_2 = \alpha e_5$	$e_1e_3 = e_4$	$e_2e_1 = e_3$	
	$e_2 e_2 = e_4 + e_5$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_4$		
$\mathbf{T}_{14}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3$		
	$e_2 e_2 = (\alpha + 1)e_4$	$e_2e_3=e_5$	$e_3e_1 = 3e_4$		
$\mathbf{T}_{15}^5(lpha)$	$e_1e_1 = e_2$	$e_1 e_2 = e_4$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3$	
	$e_2 e_2 = (\alpha + 1)e_4$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_4$		
$T_{16}^{\overline{5}}(\alpha, \overline{\beta})$	$e_1 e_1 = e_2$	$e_1 e_2 = \beta e_4 + e_5$	$e_1\overline{e_3} = \alpha \overline{e_4}$		
	$e_2e_1 = e_3$	$e_2 e_2 = (\alpha + 1)e_4$	$e_2e_3 = e_5$	$e_3e_1 = 3e_4$	

2.4. Classification theorem. Summarizing results of the present sections, we have the following theorem.

Theorem A. Let A be a complex 4-dimensional one-generated nilpotent terminal algebra. Then A is isomorphic to one of the algebras $\mathbf{T}_{01}^4 - \mathbf{T}_{10}^4$ found in Table A (See, Appendix).

3. The algebraic classification of 5-dimensional one-generated nilpotent terminal algebras

To complete the algebraic classification of 5-dimensional one-generated nilpotent terminal algebras, we need to describe all 1-dimensional extensions of the algebras from the table 5. This is done in Subsections 3.1–3.8 and summarized in the main theorem of the paper.

3.1. 1-dimensional central extensions of T_{01}^4 . Let us use the following notations:

$$\nabla_{1} = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_{2} = [\Delta_{14}], \quad \nabla_{3} = [\Delta_{22}] + 3[\Delta_{31}], \quad \nabla_{4} = [\Delta_{23}], \quad \nabla_{5} = [\Delta_{31}] + [\Delta_{41}].$$

Take $\theta = \sum_{i=1}^{5} \alpha_{i} \nabla_{i} \in \mathrm{H}^{2}_{\mathrm{T}}(\mathbf{T}^{4}_{01}).$ For an element
 $\begin{pmatrix} x & 0 & 0 & 0 \end{pmatrix}$

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & xy & x^3 & 0 \\ t & xy & 0 & x^3 \end{pmatrix} \in \operatorname{Aut} (\mathbf{T}_{01}^4),$$

.

12

we have

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_1 & \alpha_2 \\ 0 & \alpha_1 + \alpha_3 & \alpha_4 & 0 \\ 3\alpha_3 + \alpha_5 & 0 & 0 & 0 \\ \alpha_5 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha^{**} & \alpha_1^* & \alpha_2^* \\ \alpha^{***} & \alpha_1^* + \alpha_3^* & \alpha_4^* & 0 \\ 3\alpha_3^* + \alpha_5^* & 0 & 0 & 0 \\ \alpha_5^* & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl} \alpha_1^* &=& x^4 \alpha_1 + x^3 y \alpha_4, & \alpha_2^* &=& x^4 \alpha_2, & \alpha_3^* &=& x^4 \alpha_3, \\ \alpha_4^* &=& x^5 \alpha_4, & \alpha_5^* &=& x^4 \alpha_5. \end{array}$$

Since we are interested only in new algebras, we have $(\alpha_2, \alpha_5) \neq (0, 0)$ and the following cases:

(1) if $\alpha_4 \neq 0$, then choosing $y = -\frac{x\alpha_1}{\alpha_4}$, we have the family of representatives $\langle \alpha_2 \nabla_2 + \alpha_3 \nabla_3 + \alpha_4 \nabla_4 + \alpha_5 \nabla_5 \rangle$. It two three distinct orbits with representatives:

$$\langle \nabla_2 + \alpha \nabla_3 + \nabla_4 + \beta \nabla_5 \rangle$$
 and $\langle \alpha \nabla_3 + \nabla_4 + \nabla_5 \rangle$.

(2) if $\alpha_4 = 0$, then we have the family of representatives $\langle \alpha_1 \nabla_1 + \alpha_2 \nabla_2 + \alpha_3 \nabla_3 + \alpha_5 \nabla_5 \rangle$. It gives the following representatives of distinct orbits:

$$\langle \alpha \nabla_1 + \beta \nabla_2 + \gamma \nabla_3 + \nabla_5 \rangle_{(\alpha;\gamma) \neq (0;-\frac{1}{3})} \text{ and } \langle \alpha \nabla_1 + \nabla_2 + \beta \nabla_3 \rangle_{(\alpha;\beta) \neq (0;0)}.$$

Hence, we have the following 5-dimensional algebras:

$\mathbf{T}_{17}^5(lpha,eta)$	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_1e_4 = e_5$	$e_2 e_1 = e_3$
	$e_2 e_2 = \alpha e_5$	$e_2e_3=e_5$	$e_3e_1 = (3\alpha + \beta)e_5$	$e_4 e_1 = \beta e_5$
$\mathbf{T}_{18}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_2 e_1 = e_3$	$e_2 e_2 = \alpha e_5$
	$e_2 e_3 = e_5$	$e_3e_1 = (3\alpha + 1)e_5$	$e_4 e_1 = e_5$	
$\mathbf{T}_{19}^5(\alpha,\beta,\gamma)_{(\alpha;\gamma)\neq(0;-\frac{1}{3})}$	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_5$	$e_1e_4 = \beta e_5$
Ŭ	$e_2e_1 = e_3$	$e_2 e_2 = (\alpha + \gamma) e_5$	$e_3e_1 = (3\gamma + 1)e_5$	$e_4 e_1 = e_5$
$\mathbf{T}_{20}^5(\alpha,\beta)_{(\alpha;\beta)\neq(0;0)}$	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_5$	$e_1e_4 = e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = (\alpha + \beta) e_5$	$e_3e_1 = 3\beta e_5$	

3.2. 1-dimensional central extensions of $\mathbf{T}_{02}^4(\alpha)$.

(1) $\alpha \neq 6$. Let us use the following notations:

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = [\Delta_{14}] + 2[\Delta_{23}] - 3[\Delta_{32}], \quad \nabla_3 = [\Delta_{21}].$$

Take $\theta = \sum_{i=1}^{3} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{02}(\alpha))$. For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0\\ 0 & x^2 & 0 & 0\\ y & 0 & x^3 & 0\\ z & xy(\alpha - 3) & 0 & x^4 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{02}^4(\alpha)\right),$$

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_1 & \alpha_2 \\ \alpha_3 & 0 & 2\alpha_2 & 0 \\ 0 & -3\alpha_2 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & \alpha^* & \alpha_1^* & \alpha_2^* \\ \alpha_3^* & 0 & 2\alpha_2^* & 0 \\ 0 & -3\alpha_2^* & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$\alpha_1^* = x^4 \alpha_1, \quad \alpha_2^* = x^5 \alpha_2, \quad \alpha_3^* = x^3 \alpha_3 + 2x^2 y \alpha_2.$$

Since $\alpha_2 \neq 0$, then choosing $y = -\frac{x\alpha_3}{2\alpha_2}$, we have the representatives $\langle \nabla_2 \rangle_{\alpha \neq 6}$ and $\langle \nabla_1 + \nabla_2 \rangle_{\alpha \neq 6}$ depending on whether $\alpha_1 = 0$ or not.

(2) $\alpha = 6$. Let us use the following notations:

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = [\Delta_{14}] + 2[\Delta_{23}] - 3[\Delta_{32}], \quad \nabla_3 = [\Delta_{21}], \quad \nabla_4 = [\Delta_{41}] + [\Delta_{23}] - 3[\Delta_{32}].$$

Take $\theta = \sum_{i=1}^{4} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{02}(6))$. For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & 3xy & 0 & x^4 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{02}^4(6)\right),$$

we have

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_1 & \alpha_2 \\ \alpha_3 & 0 & 2\alpha_2 + \alpha_4 & 0 \\ 0 & -3(\alpha_2 + \alpha_4) & 0 & 0 \\ \alpha_4 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & \alpha^* & \alpha_1^* & \alpha_2^* \\ \alpha_3^* & 0 & 2\alpha_2^* + \alpha_4^* & 0 \\ 0 & -3(\alpha_2^* + \alpha_4^*) & 0 & 0 \\ \alpha_4^* & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rclrcl} \alpha_1^* &=& x^4 \alpha_1, & & \alpha_2^* &=& x^5 \alpha_2, \\ \alpha_3^* &=& x^3 \alpha_3 + 2x^2 y (\alpha_2 + 2\alpha_4), & & \alpha_4^* &=& x^5 \alpha_4. \end{array}$$

Since we are interested only in new algebras, we have $(\alpha_2, \alpha_4) \neq (0, 0)$ and the following cases: (a) if $\alpha_4 = 0$, then $\alpha_2 \neq 0$, and choosing $y = -\frac{x\alpha_3}{2\alpha_2}$, we have the representatives $\langle \nabla_2 \rangle$ and $\langle \nabla_1 + \nabla_2 \rangle$ depending on whether $\alpha_1 = 0$ or not.

(b) if $\alpha_4 \neq 0$, then:

(i) if $\alpha_2 \neq -2\alpha_4$, then choosing $y = -\frac{x\alpha_3}{2(\alpha_2+2\alpha_4)}$, we have the representatives $\langle \beta \nabla_2 + \nabla_4 \rangle_{\beta \neq -2}$ and $\langle \nabla_1 + \beta \nabla_2 + \nabla_4 \rangle_{\beta \neq -2}$

depending on whether $\alpha_1 = 0$ or not.

(ii) if $\alpha_2 = -2\alpha_4$, then we have the family of representatives

$$\langle \alpha_1 \nabla_1 - 2\alpha_4 \nabla_2 + \alpha_3 \nabla_3 + \alpha_4 \nabla_4 \rangle_{\pm}$$

which gives three distinct orbits with representatives

$$\langle -2\nabla_2 + \nabla_4 \rangle, \ \langle \nabla_1 - 2\nabla_2 + \nabla_4 \rangle \text{ and } \langle \beta \nabla_1 - 2\nabla_2 + \nabla_3 + \nabla_4 \rangle.$$

Summarizing, all cases, we have the following representatives of distinct orbits:

 $\langle \nabla_2 \rangle, \quad \langle \nabla_1 + \nabla_2 \rangle, \quad \langle \beta \nabla_2 + \nabla_4 \rangle_{\alpha=6}, \quad \langle \nabla_1 + \beta \nabla_2 + \nabla_4 \rangle_{\alpha=6}, \quad \langle \beta \nabla_1 - 2\nabla_2 + \nabla_3 + \nabla_4 \rangle_{\alpha=6}.$ Hence, we have following 5-dimensional algebras:

$\mathbf{T}_{21}^5(\alpha)$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = \alpha e_4$	$e_1e_4 = e_5$	
	$e_2 e_2 = e_4$	$e_2e_3 = 2e_5$	$e_3e_1 = -3e_4$	$e_3e_2 = -3e_5$	
$\mathbf{T}_{22}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = \alpha e_4 + e_5$	$e_1e_4 = e_5$	
	$e_2 e_2 = e_4$	$e_2 e_3 = 2e_5$	$e_3e_1 = -3e_4$	$e_3e_2 = -3e_5$	
$\mathbf{T}_{23}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = 6e_4$	$e_1e_4 = \beta e_5$	$e_2 e_2 = e_4$
	$e_2 e_3 = (2\beta + 1)e_5$	$e_3e_1 = -3e_4$	$e_3 e_2 = -3(\beta + 1)e_5$	$e_4 e_1 = e_5$	
$\mathbf{T}_{24}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = 6e_4 + e_5$	$e_1e_4 = \beta e_5$	$e_2 e_2 = e_4$
	$e_2 e_3 = (2\beta + 1)e_5$	$e_3e_1 = -3e_4$	$e_3 e_2 = -3(\beta + 1)e_5$	$e_4 e_1 = e_5$	
$\mathbf{T}_{25}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = 6e_4 + \beta e_5$	$e_1e_4 = -2e_5$	$e_2 e_1 = e_5$
	$e_2 e_2 = e_4$	$e_2e_3 = -3e_5$	$e_3e_1 = -3e_4$	$e_3e_2 = 3e_5$	$e_4e_1 = e_5$

3.3. 1-dimensional central extensions of T_{03}^4 . Let us use the following notations:

$$\nabla_1 = [\Delta_{14}], \quad \nabla_2 = [\Delta_{21}], \quad \nabla_3 = [\Delta_{22}] - 3[\Delta_{31}].$$

Take $\theta = \sum_{i=1}^{3} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{03}).$ For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & xy & x^3 & 0 \\ t & xz & x^2y & x^4 \end{pmatrix} \in \operatorname{Aut}(\mathbf{T}_{03}^4),$$

we have

$$\phi^T \begin{pmatrix} 0 & 0 & 0 & \alpha_1 \\ \alpha_2 & \alpha_3 & 0 & 0 \\ -3\alpha_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha^{**} & \alpha^* & \alpha_1^* \\ \alpha_2^* & \alpha_3^* & 0 & 0 \\ -3\alpha_3^* & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$\alpha_1^* = x^5 \alpha_1, \quad \alpha_2^* = x^3 \alpha_2 - 2x^2 y \alpha_3, \quad \alpha_3^* = x^4 \alpha_3.$$

Since $\alpha_1 \neq 0$, we have the following cases:

- (1) if α₃ ≠ 0, then choosing x = α₃/α₁, y = xα₂/2α₃, we have the representatives (∇₁ + ∇₃).
 (2) if α₃ = 0, then we have the representatives (∇₁) and (∇₁ + ∇₂) depending on whether α₂ = 0 or not.

Hence we have the following 5-dimensional algebras:

\mathbf{T}_{26}^5	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1 e_3 = e_4$	$e_1e_4 = e_5$	$e_2 e_2 = e_5$	$e_3e_1 = -3e_5$
\mathbf{T}_{27}^5	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1 e_3 = e_4$	$e_1e_4 = e_5$		
\mathbf{T}_{28}^5	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1 e_3 = e_4$	$e_1e_4 = e_5$	$e_2 e_1 = e_5$	

3.4. 1-dimensional central extensions of T_{04}^4 . Let us use the following notations:

$$\nabla_1 = [\Delta_{13}], \quad \nabla_2 = [\Delta_{14}] + 3[\Delta_{31}], \quad \nabla_3 = [\Delta_{22}] - 3[\Delta_{31}].$$

Take $\theta = \sum_{i=1}^{3} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{04})$. For the element

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ x & 1 & 0 & 0 \\ y & x & 1 & 0 \\ z & y + z & x & 1 \end{pmatrix} \in \operatorname{Aut} (\mathbf{T}_{04}^4).$$

we have

$$\phi^T \begin{pmatrix} 0 & 0 & \alpha_1 & \alpha_2 \\ 0 & \alpha_3 & 0 & 0 \\ 3(\alpha_2 - \alpha_3) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha^{**} & \alpha^* + \alpha_1^* & \alpha_2^* \\ \alpha^* & \alpha_3^* & 0 & 0 \\ 3(\alpha_2^* - \alpha_3^*) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$\alpha_1^* = \alpha_1 + 2x(\alpha_3 - \alpha_2), \quad \alpha_2^* = \alpha_2, \quad \alpha_3^* = \alpha_3$$

If $\alpha_2 \neq \alpha_3$, then choosing $x = \frac{\alpha_1}{2(\alpha_2 - \alpha_3)}$, we have the representatives $\langle \nabla_2 + \alpha \nabla_3 \rangle_{\alpha \neq 1}$. If $\alpha_2 = \alpha_3$, then we have the representatives $\langle \alpha \nabla_1 + \nabla_2 + \nabla_3 \rangle$.

Hence, we have following 5-dimensional algebras:

3.5. 1-dimensional central extensions of $\mathbf{T}_{05}^4(\lambda, \alpha)$.

(1) $\lambda = 1, \alpha = 0$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = [\Delta_{14}] + [\Delta_{32}], \quad \nabla_4 = [\Delta_{34}].$$

Take $\theta = \sum_{i=1}^{4} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{05}(1,0))$. For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & xy & 0 & x^4 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{05}^4(1,0)\right),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 & \alpha_3 \\ 0 & \alpha_2 & 0 & 0 \\ 0 & \alpha_3 & 0 & \alpha_4 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha_1^* & \alpha_2^* & \alpha_3^* \\ 0 & \alpha_2^* & 0 & 0 \\ \alpha^{**} & \alpha_3^* & 0 & \alpha_4^* \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl} \alpha_1^* &=& x^3 \alpha_1 + 2x^2 y \alpha_3 + x y^2 \alpha_4, & \alpha_2^* &=& x^4 \alpha_2, \\ \alpha_3^* &=& x^5 \alpha_3 + x^4 y \alpha_4, & \alpha_4^* &=& x^7 \alpha_4. \end{array}$$

Since we are interested only in new algebras, we have $(\alpha_3, \alpha_4) \neq (0, 0)$ and the following cases: (a) if $\alpha_4 = 0$, then:

- (i) if $\alpha_2 = 0$, then choosing $y = -\frac{x\alpha_1}{2\alpha_3}$, we have the representatives $\langle \nabla_3 \rangle$. (ii) if $\alpha_2 \neq 0$, then choosing $x = \frac{\alpha_2}{\alpha_3}$, $y = -\frac{x\alpha_1}{2\alpha_3}$, we have the representatives $\langle \nabla_2 + \nabla_3 \rangle$. (b) if $\alpha_4 \neq 0$, then:
 - (i) if $\alpha_2 = 0$, $\alpha_1 \alpha_4 \alpha_3^2 = 0$, then choosing $y = -\frac{x\alpha_3}{\alpha_4}$, we have the representatives $\langle \nabla_4 \rangle$.
 - (ii) if $\alpha_2 = 0$, $\alpha_1 \alpha_4 \alpha_3^2 \neq 0$, then choosing $x = \sqrt[4]{\frac{\alpha_1 \alpha_4 \alpha_3^2}{\alpha_4^2}}$, $y = -\frac{x\alpha_3}{\alpha_4}$, we have the representatives $\langle \nabla_1 + \nabla_4 \rangle$.

(iii) if
$$\alpha_2 \neq 0$$
, then choosing $x = \sqrt[3]{\frac{\alpha_2}{\alpha_4}}$, $y = -\frac{x\alpha_3}{\alpha_4}$, we have the representatives

$$\langle \beta \nabla_1 + \nabla_2 + \nabla_4 \rangle$$

(2) $\lambda = 0, \alpha = \frac{1}{3}$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = [\Delta_{23}], \quad \nabla_4 = [\Delta_{24}] + [\Delta_{33}].$$

Take
$$\theta = \sum_{i=1}^{n} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{05}(0, \frac{1}{3}))$$
. For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & \frac{4}{3}xy & 0 & x^4 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{05}^4(0, \frac{1}{3})\right),$$

we have

$$\phi^{T} \begin{pmatrix} 0 & \alpha_{1} & \alpha_{2} & 0 \\ 0 & \alpha_{2} & \alpha_{3} & \alpha_{4} \\ 0 & 0 & \alpha_{4} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha_{1}^{*} & \alpha^{*} + \alpha_{2}^{*} & 0 \\ \alpha^{**} & 2\alpha^{*} + \alpha_{2}^{*} & \alpha_{3}^{*} & \alpha_{4}^{*} \\ 3\alpha^{*} & 0 & \alpha_{4}^{*} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

where

$$\begin{array}{rcl} \alpha_1^* &=& x^3 \alpha_1, & \alpha_2^* &=& x^4 \alpha_2 + \frac{2}{3} x^3 y \alpha_4, \\ \alpha_3^* &=& x^5 \alpha_3, & \alpha_4^* &=& x^6 \alpha_4. \end{array}$$

Since $\alpha_4 \neq 0$, choosing $y = -\frac{3x\alpha_2}{2\alpha_4}$, we have the representatives

$$\langle \nabla_4 \rangle, \ \langle \nabla_1 + \nabla_4 \rangle \text{ and } \langle \beta \nabla_1 + \nabla_3 + \nabla_4 \rangle$$

depending on whether $\alpha_1 = 0$, $\alpha_3 = 0$ or not.

(3) $\lambda = 0, \alpha = -\frac{2}{3}$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = [\Delta_{23}], \quad \nabla_4 = [\Delta_{14}] + [\Delta_{32}],$$

Take
$$\theta = \sum_{i=1}^{4} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{05}(0, -\frac{2}{3})).$$
 For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & \frac{1}{3}xy & 0 & x^4 \end{pmatrix} \in \mathrm{Aut}\,(\mathbf{T}^4_{05}(0, -\frac{2}{3})),$$

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 & \alpha_4 \\ 0 & \alpha_2 & \alpha_3 & 0 \\ 0 & \alpha_4 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^* & \alpha_1^* & \alpha_2^* & \alpha_4^* \\ 0 & \alpha_2^* & \alpha_3^* & 0 \\ 0 & \alpha_4^* & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl} \alpha_1^* &=& x^3 \alpha_1 + \frac{4}{3} x^2 y \alpha_4, & \alpha_2^* &=& x^4 \alpha_2, \\ \alpha_3^* &=& x^5 \alpha_3, & \alpha_4^* &=& x^5 \alpha_4. \end{array}$$

Since $\alpha_4 \neq 0$, choosing $y = -\frac{3x\alpha_1}{4\alpha_4}$, we have the representatives $\langle \beta \nabla_3 + \nabla_4 \rangle$ and $\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle$, depending on whether $\alpha_2 = 0$ or not. (4) $\lambda = 0, \alpha = -\frac{1}{5}$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = [\Delta_{23}], \quad \nabla_4 = [\Delta_{14}] - \frac{2}{5}[\Delta_{32}] - \frac{7}{2}[\Delta_{41}].$$

Take
$$\theta = \sum_{i=1}^{4} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{05}(0, -\frac{1}{5}))$$
. For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & \frac{4}{5}xy & 0 & x^4 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{05}^4(0, -\frac{1}{5})\right),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 & \alpha_4 \\ 0 & \alpha_2 & \alpha_3 & 0 \\ 0 & -\frac{2}{5}\alpha_4 & 0 & 0 \\ -\frac{7}{2}\alpha_4 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & \alpha_1^* & \alpha_2^* & \alpha_4^* \\ \alpha^* & \alpha_2^* & \alpha_3^* & 0 \\ 0 & -\frac{2}{5}\alpha_4^* & 0 & 0 \\ -\frac{7}{2}\alpha_4^* & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl} \alpha_1^* &=& x^3 \alpha_1 + \frac{2}{5} x^2 y \alpha_4, & \alpha_2^* &=& x^4 \alpha_2, \\ \alpha_3^* &=& x^5 \alpha_3, & \alpha_4^* &=& x^5 \alpha_4. \end{array}$$

Since $\alpha_4 \neq 0$, choosing $y = -\frac{5x\alpha_1}{2\alpha_4}$, we have the representatives $\langle \beta \nabla_3 + \nabla_4 \rangle$ or $\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle$ depending on whether $\alpha_2 = 0$ or not.

(5) $\lambda = -1$. Let us use the following notations:

$$\begin{aligned} \nabla_1 &= [\Delta_{12}], & \nabla_3 &= [\Delta_{14}] - \frac{3\alpha + 4}{3} [\Delta_{23}] + [\Delta_{32}], \\ \nabla_2 &= [\Delta_{13}] + [\Delta_{22}], & \nabla_4 &= \frac{3\alpha + 1}{3} [\Delta_{23}] - \frac{3\alpha + 1}{2} [\Delta_{32}] + [\Delta_{41}]. \end{aligned}$$

$$\begin{aligned} \text{Take } \theta &= \sum_{i=1}^{4} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{05}(-1,\alpha)). \text{ For the element} \\ \phi &= \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ z & 0 & x^3 & 0 \\ t & xz(\alpha+1) & 0 & x^4 \end{pmatrix} \in \mathrm{Aut} \, (\mathbf{T}^4_{05}(-1,\alpha)_{\alpha \neq -\frac{2}{3}}), \quad \phi = \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & 0 & x^3 & 0 \\ t & \frac{1}{3}xz & 0 & x^4 \end{pmatrix} \in \mathrm{Aut} \, (\mathbf{T}^4_{05}(-1,-\frac{2}{3})), \end{aligned}$$

$$\phi^{T} \begin{pmatrix} 0 & \alpha_{1} & \alpha_{2} & \alpha_{3} \\ 0 & \alpha_{2} & -\frac{3\alpha+4}{3}\alpha_{3} + \frac{3\alpha+1}{3}\alpha_{4} & 0 \\ 0 & \alpha_{3} - \frac{3\alpha+1}{2}\alpha_{4} & 0 & 0 \\ \alpha_{4} & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & -\alpha^{*} + \alpha_{1}^{*} & \alpha_{2}^{*} & \alpha_{3}^{*} \\ \alpha^{*} & \alpha_{2}^{*} & -\frac{3\alpha+4}{3}\alpha_{3}^{*} + \frac{3\alpha+1}{3}\alpha_{4}^{*} & 0 \\ 0 & \alpha_{3}^{*} - \frac{3\alpha+1}{2}\alpha_{4}^{*} & 0 & 0 \\ \alpha_{4}^{*} & 0 & 0 & 0 \end{pmatrix}$$

where

$$\alpha \neq -\frac{2}{3} : \begin{cases} \alpha_1^* = x^3 \alpha_1 + \frac{4\alpha_3 + (3\alpha + 5)\alpha_4}{6} x^2 z, \\ \alpha_2^* = x^4 \alpha_2, \\ \alpha_3^* = x^5 \alpha_3, \\ \alpha_4^* = x^5 \alpha_4. \end{cases}, \quad \alpha = -\frac{2}{3} : \begin{cases} \alpha_1^* = x^3 \alpha_1 + 2x^2 y \alpha_2 + \frac{4\alpha_3 + 3\alpha_4}{6} x^2 z, \\ \alpha_2^* = x^4 \alpha_2, \\ \alpha_3^* = x^5 \alpha_3, \\ \alpha_4^* = x^5 \alpha_4. \end{cases}$$

- Since $(\alpha_3, \alpha_4) \neq (0, 0)$, we have the following cases: (a) if $\alpha_4 = 0$, then by choosing $z = -\frac{3x\alpha_1}{2\alpha_3}$ (or $z = -\frac{3x\alpha_1+6y\alpha_2}{2\alpha_3}$), we have the representatives $\langle \nabla_3 \rangle$ and $\langle \nabla_2 + \nabla_3 \rangle$ depending on whether $\alpha_2 = 0$ or not.
- (b) if $\alpha_4 \neq 0$, then:
 - (i) if $4\alpha_3 + (3\alpha + 5)\alpha_4 \neq 0$, then by choosing $z = -\frac{6x\alpha_1}{4\alpha_3 + (3\alpha + 5)\alpha_4}$ (or $z = -\frac{6x\alpha_1 + 12y\alpha_2}{4\alpha_3 + 3\alpha_4}$), we have the representatives $\langle \beta \nabla_3 + \nabla_4 \rangle$ and $\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle$ depending on whether $\alpha_2 = 0$ or not.
 - (ii) if $4\alpha_3 + (3\alpha + 5)\alpha_4 = 0$, then in case of $\alpha \neq -\frac{2}{3}$, we have the representatives

$$\langle -\frac{3\alpha+5}{4}\nabla_3+\nabla_4\rangle, \ \langle \nabla_2-\frac{3\alpha+5}{4}\nabla_3+\nabla_4\rangle \text{ and } \langle \nabla_1+\beta\nabla_2-\frac{3\alpha+5}{4}\nabla_3+\nabla_4\rangle$$

depending on whether $\alpha_1 = 0$, $\alpha_2 = 0$ or not. If $\alpha = -\frac{2}{3}$, we have the representatives

$$\langle -\frac{3}{4}\nabla_3 + \nabla_4 \rangle, \ \langle \nabla_1 - \frac{3}{4}\nabla_3 + \nabla_4 \rangle \text{ and } \langle \nabla_2 - \frac{3}{4}\nabla_3 + \nabla_4 \rangle$$

depending on whether $\alpha_1 = 0, \alpha_2 = 0$ or not.

(6) $\alpha = \frac{4\lambda - 1}{5 - 2\lambda}, \lambda \neq 0; -1; \frac{5}{2}$. Let us use the following notations:

$$\nabla_{1} = [\Delta_{12}], \qquad \nabla_{3} = [\Delta_{14}] + \frac{4\lambda^{2} - 2\lambda + 7}{3\lambda(5 - 2\lambda)} [\Delta_{23}] + [\Delta_{32}], \\
\nabla_{2} = [\Delta_{13}] + [\Delta_{22}], \qquad \nabla_{4} = \frac{2(\lambda - 1)^{2}}{3\lambda(5 - 2\lambda)} [\Delta_{23}] - \frac{2(\lambda - 1)}{5 - 2\lambda} [\Delta_{32}] + [\Delta_{41}].$$

$$\begin{aligned} \text{Take } \theta &= \sum_{i=1}^{4} \alpha_i \nabla_i \in \mathcal{H}^2_{\mathcal{T}}(\mathcal{T}^4_{05}(\lambda, \frac{4\lambda - 1}{5 - 2\lambda})). \text{ For the element} \\ \phi &= \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & 2xy & x^3 & 0 \\ t & 2xz + y^2 & 3x^2y & x^4 \end{pmatrix} \in \text{Aut} (\mathcal{T}^4_{05}(1, 1)), \quad \phi = \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & \frac{1}{2}xy & x^3 & 0 \\ t & \frac{1}{2}xz & \frac{1}{2}x^2y & x^4 \end{pmatrix} \in \text{Aut} (\mathcal{T}^4_{05}(-\frac{1}{2}, -\frac{1}{2})), \\ \phi &= \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ z & 0 & x^3 & 0 \\ t & \frac{2(\lambda + 2)}{5 - 2\lambda}xz & 0 & x^4 \end{pmatrix} \in \text{Aut} (\mathcal{T}^4_{05}(\lambda, \frac{4\lambda - 1}{5 - 2\lambda})), \quad \lambda \neq 1; -\frac{1}{2}, \end{aligned}$$

$$\phi^{T} \begin{pmatrix} 0 & \alpha_{1} & \alpha_{2} & \alpha_{3} \\ 0 & \alpha_{2} & \frac{4\lambda^{2}-2\lambda+7}{3\lambda(5-2\lambda)}\alpha_{3} + \frac{2(\lambda-1)^{2}}{3\lambda(5-2\lambda)}\alpha_{4} & 0 \\ 0 & \alpha_{3} - \frac{2(\lambda-1)}{(5-2\lambda)}\alpha_{4} & 0 & 0 \\ \alpha_{4} & 0 & 0 & 0 \end{pmatrix} \phi = \\ \begin{pmatrix} \alpha^{**} & \lambda\alpha^{*} + \alpha_{1}^{*} & \alpha_{2}^{*} & \alpha_{3}^{*} \\ \alpha^{*} & \alpha_{2}^{*} & \frac{4\lambda^{2}-2\lambda+7}{3\lambda(5-2\lambda)}\alpha_{3}^{*} + \frac{2(\lambda-1)^{2}}{3\lambda(5-2\lambda)}\alpha_{4}^{*} & 0 \\ 0 & \alpha_{3}^{*} - \frac{2(\lambda-1)}{(5-2\lambda)}\alpha_{4}^{*} & 0 & 0 \\ \alpha_{4}^{*} & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{split} \lambda &= 1 \quad : \quad \begin{cases} \alpha_1^* = x^3 \alpha_1 + 2x^2 y \alpha_2 + (2x^2 z + xy^2)(\alpha_3 - \alpha_4), \\ \alpha_2^* = x^4 \alpha_2 + 3x^3 y(\alpha_3 - \alpha_4), \\ \alpha_3^* = x^5 \alpha_3, \\ \alpha_4^* = x^5 \alpha_4. \\ \alpha_1^* = x^3 \alpha_1 + 2x^2 y \alpha_2 + (x^2 z - \frac{1}{4} xy^2)(\alpha_3 + \frac{1}{2} \alpha_4), \\ \alpha_2^* = x^4 \alpha_2, \\ \alpha_3^* = x^5 \alpha_3, \\ \alpha_4^* = x^5 \alpha_4. \\ \lambda \neq 1; -\frac{1}{2} \quad : \quad \begin{cases} \alpha_1^* = x^3 \alpha_1 + x^2 z \frac{2(\lambda+2)}{3} (\alpha_3 - \frac{4\lambda-1}{5-2\lambda} \alpha_4), \\ \alpha_2^* = x^4 \alpha_2, \\ \alpha_3^* = x^5 \alpha_3, \\ \alpha_4^* = x^5 \alpha_4. \end{cases} \end{split}$$

Since $(\alpha_3, \alpha_4) \neq (0, 0)$, we have the following cases:

- (a) $\lambda = 1$: (i) if $\alpha_4 = 0$, then choosing $y = -\frac{x\alpha_2}{3\alpha_3}$, $z = -\frac{x^2\alpha_1 + 2xy\alpha_2 + y^2\alpha_3}{2x\alpha_3}$, we have the representatives $\langle \nabla_3 \rangle$. (ii) if $\alpha_4 \neq 0$, then:

- (A) if $\alpha_3 = \alpha_4, \alpha_2 = 0$, then we have the representatives $\langle \nabla_3 + \nabla_4 \rangle$ and $\langle \nabla_1 + \nabla_3 + \nabla_4 \rangle$ depending on whether $\alpha_1 = 0$ or not.
- (B) if $\alpha_3 = \alpha_4, \alpha_2 \neq 0$, then choosing $y = -\frac{x\alpha_1}{2\alpha_2}$, we have the representative $\langle \nabla_2 + \nabla_3 + \nabla_4 \rangle$.
- (C) if $\alpha_3 \neq \alpha_4$, then choosing $y = -\frac{x\alpha_2}{3(\alpha_3 \alpha_4)}$, $z = -\frac{x^2\alpha_1 + 2xy\alpha_2 + y^2(\alpha_3 \alpha_4)}{2x(\alpha_3 \alpha_4)}$, we have the representatives $\langle \beta \nabla_3 + \nabla_4 \rangle$.

(b)
$$\lambda = -\frac{1}{2}$$
:

(i) if $\alpha_4 = 0$, then choosing y = 0, $z = -\frac{x\alpha_1}{\alpha_3}$, we have the representatives $\langle \nabla_3 \rangle$ and $\langle \nabla_2 + \nabla_3 \rangle$ depending on whether $\alpha_2 = 0$ or not.

(ii) if
$$\alpha_4 \neq 0$$
, then:

- (A) if $\alpha_3 = -\frac{1}{2}\alpha_4$, $\alpha_2 = 0$, then we have the representatives $\langle -\frac{1}{2}\nabla_3 + \nabla_4 \rangle$ and $\langle \nabla_1 \frac{1}{2}\nabla_3 + \nabla_4 \rangle$ depending on whether $\alpha_1 = 0$ or not.
- (B) if $\alpha_3 = -\frac{1}{2}\alpha_4$, $\alpha_2 \neq 0$, then choosing $y = -\frac{x\alpha_1}{2\alpha_2}$, we have the representative $\langle \nabla_2 \frac{1}{2}\nabla_3 + \nabla_4 \rangle$.
- (C) if $\alpha_3 \neq -\frac{1}{2}\alpha_4$, then choosing $z = -\frac{8x^2\alpha_1 + 16xy\alpha_2 y^2(2\alpha_3 + \alpha_4)}{4x(2\alpha_3 + \alpha_4)}$, we have the representative $\langle \beta \nabla_3 + \nabla_4 \rangle$ and $\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle$ depending on whether $\alpha_2 = 0$ or not.

(c)
$$\lambda = -2$$
:

(i) if $\alpha_4 = 0$, then we have the representatives

$$\langle
abla_3
angle, \ \langle
abla_1 +
abla_3
angle ext{ and } \langle eta
abla_1 +
abla_2 +
abla_3
angle$$

depending on whether $\alpha_1 = 0$, $\alpha_2 = 0$ or not.

- (ii) if $\alpha_4 \neq 0$, then we have the representatives $\langle \beta \nabla_3 + \nabla_4 \rangle$, $\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle$ and $\langle \nabla_1 + \beta \nabla_2 + \gamma \nabla_3 + \nabla_4 \rangle$ depending on whether $\alpha_1 = 0$, $\alpha_2 = 0$ or not.
- (d) $\lambda \neq 1; -2; -\frac{1}{2}:$
 - (i) if $\alpha_4 = 0$, then choosing $z = -\frac{3x\alpha_1}{2(\lambda+2)\alpha_3}$, we have the representatives $\langle \nabla_3 \rangle$ and $\langle \nabla_2 + \nabla_3 \rangle$ depending on whether $\alpha_2 = 0$ or not.
 - (ii) if α₄ ≠ 0, then:
 (A) if α₃ = 4λ-1/(5-2λ)α₄, then we have the representatives (4λ-1/(5-2λ)∇₃ + ∇₄), (∇₂ + 4λ-1/(5-2λ)∇₃ + ∇₄) and (∇₁ + β∇₂ + 4λ-1/(5-2λ)∇₃ + ∇₄) depending on whether α₁ = 0, α₂ = 0 or not.
 (B) if α₃ ≠ 4λ-1/(5-2λ)α₄, then choosing z = -(3xα₁(5-2λ))/(2(λ+2)((5-2λ)α₄-(4λ-1)α₃)), we have the representative (β∇₃ + ∇₄) and (∇₂ + β∇₃ + ∇₄) depending on whether α₂ = 0 or not.

(7) $\lambda \neq 0; -1, \alpha \neq \frac{4\lambda-1}{5-2\lambda}, (\lambda; \alpha) \neq (1; 0)$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = [\Delta_{14}] + \frac{3\alpha - 2(\lambda - 1)}{3\lambda} [\Delta_{23}] + [\Delta_{32}].$$

$$\begin{split} \text{Take } \theta &= \sum_{i=1}^{3} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{05}(\lambda, \alpha)). \text{ For the element} \\ \phi &= \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ z & 0 & x^3 & 0 \\ t & (\alpha+1)xz & 0 & x^4 \end{pmatrix} \in \mathrm{Aut} \left(\mathbf{T}^4_{05}(\lambda, \alpha)\right), \quad \alpha \neq \frac{2\lambda^2 + 5\lambda - 1}{6}, \\ \phi &= \begin{pmatrix} x & 0 & 0 & 0 \\ y & x^2 & 0 & 0 \\ z & (\lambda+1)xy & x^3 & 0 \\ t & \frac{2\lambda^2 + 5\lambda + 5}{6}xz + \frac{2\lambda^2 + 3\lambda + 1}{6}y^2 & \frac{2\lambda^2 + 9\lambda + 7}{6}x^2y & x^4 \end{pmatrix} \in \mathrm{Aut} \left(\mathbf{T}^4_{04}\left(\lambda, \frac{2\lambda^2 + 5\lambda - 1}{6}\right)\right), \\ \text{we have} \\ \phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 & \alpha_3 \\ 0 & \alpha_2 & \frac{3\alpha - 2(\lambda - 1)}{3\lambda}\alpha_3 & 0 \\ 0 & \alpha_3 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & \lambda \alpha^* + \alpha_1^* & \alpha \alpha^* + \alpha_2^* & \alpha_3^* \\ \alpha^* & \left(\alpha + \frac{1 - \lambda}{3}\right)\alpha^* + \alpha_2^* & \frac{3\alpha - 2(\lambda - 1)}{3\lambda}\alpha_3 & 0 \\ \alpha^* & \alpha_3^* & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \end{split}$$

where

$$\alpha \neq \frac{2\lambda^2 + 5\lambda - 1}{6} : \begin{cases} \alpha_1^* = x^3 \alpha_1 + \frac{2}{3} x^2 y (\lambda + 2) \alpha_3, \\ \alpha_2^* = x^4 \alpha_2, \\ \alpha_3^* = x^5 \alpha_3. \end{cases} \\ \alpha = \frac{2\lambda^2 + 5\lambda - 1}{6} : \begin{cases} \alpha_1^* = x^3 \alpha_1 + 2x^2 y \alpha_2 + \frac{(\lambda + 1)(-2\lambda^2 + 2\lambda + 3)xy^2 + 4\lambda(\lambda + 2)x^2 z}{6\lambda} \alpha_3, \\ \alpha_2^* = x^4 \alpha_2 + \frac{(2\lambda + 1)(\lambda + 1)}{2\lambda} x^3 y \alpha_3, \\ \alpha_3^* = x^5 \alpha_3. \end{cases}$$

- Since $\alpha_3 \neq 0$, we have the following cases: (a) $\lambda = -2$, $\alpha = -\frac{1}{2}$, then choosing $y = \frac{4x\alpha_2}{3\alpha_3}$ we have the representatives $\langle \nabla_3 \rangle$ and $\langle \nabla_1 + \nabla_3 \rangle$ depending on whether $3\alpha_1\alpha_3 + 4\alpha_2^2 = 0$ or not.
- (b) $\lambda = -2, \alpha \neq -\frac{1}{2}$, then we have the representatives

$$\langle
abla_3
angle, \ \langle
abla_1 +
abla_3
angle \text{ and } \langle eta
abla_1 +
abla_2 +
abla_3
angle$$

- depending on whether $\alpha_1 = 0$, $\alpha_2 = 0$ or not. (c) $\lambda \neq -2$, $\alpha = \frac{2\lambda^2 + 5\lambda 1}{6}$, then choosing $y = -\frac{2\lambda x \alpha_2}{(2\lambda + 1)(\lambda + 1)\alpha_3}$, $z = -\frac{6x^2\lambda\alpha_1 + 12xy\lambda\alpha_2 (\lambda + 1)(2\lambda^2 2\lambda 3)xy\alpha_3}{4\lambda\alpha_3}$, we have the representatives $\langle \nabla_3 \rangle$. (d) $\lambda \neq -2$, $\alpha \neq \frac{2\lambda^2 + 5\lambda 1}{6}$, then choosing $y = -\frac{3x\alpha_1}{2(\lambda + 2)\alpha_3}$, we have the representatives $\langle \nabla_3 \rangle$ and $\langle \nabla_3 \rangle = 0$ or not.
- $\langle \nabla_2 + \nabla_3 \rangle$ depending on whether $\alpha_2 = 0$ or not.

Thus, we have the following distinct orbits:

$$\lambda = 1, \ \alpha = 0 \quad : \quad \langle \beta \nabla_1 + \nabla_2 + \nabla_4 \rangle, \qquad \qquad \langle \nabla_1 + \nabla_4 \rangle, \qquad \qquad \langle \nabla_4 \rangle,$$

$\lambda = 0, \ \alpha = \frac{1}{3}$:	$\langle \beta \nabla_1 + \nabla_3 + \nabla_4 \rangle,$	$\langle abla_1 + abla_4 angle,$	$\langle abla_4 angle,$
$\lambda = 0, \ \alpha = -\frac{2}{3}$:	$\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle,$	$\langle \beta \nabla_3 + \nabla_4 \rangle,$	
$\lambda = 0, \ \alpha = -\frac{1}{5}$:	$\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle,$	$\langle \beta \nabla_3 + \nabla_4 \rangle,$	
$\lambda = -1$:	$\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle,$	$\langle \beta \nabla_3 + \nabla_4 \rangle,$	
		$\langle \nabla_1 + \beta \nabla_2 - \frac{3\alpha+5}{4} \nabla_3 + \nabla_4 \rangle_{\alpha \neq -\frac{2}{3}},$	$\langle \nabla_1 - \frac{3}{4} \nabla_3 + \nabla_4 \rangle_{\alpha = -\frac{2}{3}},$	$\langle \nabla_2 + \nabla_3 \rangle_{\alpha = -\frac{2}{3}},$
$\alpha = \frac{4\lambda - 1}{5 - 2\lambda}$:	$\langle \nabla_1 + \beta \nabla_2 + \frac{4\lambda - 1}{5 - 2\lambda} \nabla_3 + \nabla_4 \rangle_{\lambda \neq 1, -2, -\frac{1}{2}},$	$\langle \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle_{\lambda \neq 1},$	$\langle \beta \nabla_3 + \nabla_4 \rangle,$
		$\langle abla_1 + abla_3 + abla_4 angle_{\lambda=1},$	$\langle \nabla_2 + \nabla_3 + \nabla_4 \rangle_{\lambda=1},$	
		$\langle \nabla_1 + \beta \nabla_2 + \gamma \nabla_3 + \nabla_4 \rangle_{\lambda = -2},$	$\langle \beta \nabla_1 + \nabla_2 + \nabla_3 \rangle_{\lambda = -2},$	
		$\langle \mathbf{v}_2 - \frac{1}{2}\mathbf{v}_3 + \mathbf{v}_{4/\lambda = -\frac{1}{2}},$	$\langle \mathbf{v}_2 + \mathbf{v}_3 \rangle_{\lambda = -\frac{1}{2}},$	
$\lambda \neq 0$:	$\langle abla_2 + abla_3 angle_{\lambda eq -2; lpha eq rac{2\lambda^2 + 5\lambda - 1}{6}},$	$\langle abla_3 angle,$	
		$\langle \beta \nabla_1 + \nabla_2 + \nabla_3 \rangle_{\lambda = -2, \alpha = -\frac{1}{2}},$	$\langle abla_1 + abla_3 angle_{\lambda = -2}.$	

The corresponding algebras are:

$\mathbf{T}_{31}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = e_3 + \beta e_5$	$e_1e_3 = e_5$	
	$e_2 e_1 = e_3$	$e_2 e_2 = e_5$	$e_3e_1 = e_4$	$e_3 e_4 = e_5$
\mathbf{T}_{32}^5	$e_1e_1 = e_2$	$e_1 e_2 = e_3 + e_5$	$e_2e_1 = e_3$	
	$e_3 e_1 = e_4$	$e_3e_4 = e_5$		
${f T}_{33}^5$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_2 e_1 = e_3$	
	$e_3e_1 = e_4$	$e_3e_4 = e_5$		
$\mathbf{T}_{34}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = \beta e_5$	$e_1 e_3 = \frac{1}{3} e_4$	
	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{2}{3} e_4$	$e_2 e_3 = e_5$	
	$e_2 e_4 = e_5$	$e_3e_1 = e_4$	$e_3 e_3 = e_5$	
\mathbf{T}_{35}^5	$e_1e_1 = e_2$	$e_1e_2 = e_5$	$e_1 e_3 = \frac{1}{3} e_4$	$e_2e_1 = e_3$
	$e_2 e_2 = \frac{2}{3} e_4$	$e_2 e_4 = e_5$	$e_3 e_1 = e_4$	$e_3 e_3 = e_5$
\mathbf{T}_{36}^5	$e_1e_1 = e_2$	$e_1 e_3 = \frac{1}{3} e_4$	$e_2 e_1 = e_3$	
	$e_2 e_2 = \frac{2}{3} e_4$	$e_{2}e_{4} = e_{5}$	$e_3e_1 = e_4$	$e_3 e_3 = e_5$
$\mathbf{T}_{37}^5(eta)$	$e_1 e_1 = e_2$	$e_1 e_3 = -\frac{2}{3}e_4 + e_5$	$e_2 e_1 = e_3$	$e_1e_4 = e_5$
	$e_2 e_2 = -\frac{1}{3}e_4 + e_5$	$e_2 e_3 = \beta e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$
$\mathbf{T}_{38}^5(eta)$	$e_1e_1 = e_2$	$e_1e_3 = -\frac{2}{3}e_4$	$e_2e_1 = e_3$	$e_1e_4 = e_5$
	$e_2 e_2 = -\frac{1}{3} e_4$	$e_2 e_3 = \beta e_5$	$e_3e_1 = e_4$	$e_3 e_2 = e_5$
$\mathbf{T}_{39}^5(eta)$	$e_1e_1 = e_2$	$e_1e_3 = -\frac{1}{5}e_4 + e_5$	$e_1e_4 = e_5$	
	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{2}{15} e_4 + e_5$	$e_2 e_3 = \beta e_5$	
	$e_3 e_1 = e_4$	$e_3e_2 = -\frac{2}{5}e_5$	$e_4 e_1 = -\frac{7}{2} e_5$	
$\mathbf{T}_{40}^5(eta)$	$e_1e_1 = e_2$	$e_1e_3 = -\frac{1}{5}e_4$	$e_1 e_4 = e_5$	
	$e_2e_1 = e_3$	$e_2 e_2 = \frac{2}{15} e_4$	$e_2 e_3 = \beta e_5$	
	$e_3e_1 = e_4$	$e_3 e_2 = -\frac{2}{5} e_5$	$e_4 e_1 = -\frac{7}{2} e_5$	

$\mathbf{T}_{41}^5(lpha,eta)$	$e_1e_1 = e_2$	$e_1e_2 = -e_3$	$e_1e_3 = \alpha e_4 + e_5$
	$e_1e_4 = \beta e_5$	$e_2 e_1 = e_3$	$e_2e_2 = \left(\alpha + \frac{2}{3}\right)e_4 + e_5$
	$e_2 e_3 = \left(\frac{3\alpha + 1}{3} - \frac{\beta(3\alpha + 1)}{3}\right)$	$\left(\frac{\alpha+4)}{3}\right)e_5$	$e_3 e_1 = e_4$
	$e_3e_2 = \left(\beta - \frac{3\alpha+1}{2}\right)e_3$	95 5	$e_4 e_1 = e_5$
$\mathbf{T}_{42}^5(lpha,eta)$	$e_1 e_1 = e_2$	$e_1e_2 = -e_3$	$e_1e_3 = \alpha e_4$
	$e_1e_4 = \beta e_5$	$e_2 e_1 = e_3$	$e_2e_2 = \left(\alpha + \frac{2}{3}\right)e_4$
	$e_2 e_3 = \left(\frac{3\alpha + 1}{3} - \frac{\beta(3\alpha + 1)}{3}\right)$	$\left(\frac{\alpha+4)}{3}\right)e_5$	$e_3e_1 = e_4$
	$e_3e_2 = \left(\beta - \frac{3\alpha + 1}{2}\right)e_3$	95 5	$e_4 e_1 = e_5$
$\mathbf{T}_{43}^5(lpha,eta)$	$e_1e_1 = e_2$	$e_1 e_2 = -e_3 + e_5$	$e_1e_3 = \alpha e_4 + \beta e_5$
$\alpha \neq -\frac{2}{3}$	$e_1 e_4 = -\frac{3\alpha + 5}{4} e_5$	$e_2 e_1 = e_3$	$e_2e_2 = \left(\alpha + \frac{2}{3}\right)e_4 + \beta e_5$
	$e_2e_3 = \frac{3\alpha^2 + 13\alpha + 8}{4}e_5$		$e_3e_1 = e_4$
	$e_3e_2 = -\frac{9\alpha + 7}{4}e_5$	$e_4 e_1 = e_5$	
\mathbf{T}_{44}^5	$e_1 e_1 = e_2$	$e_1 e_2 = -e_3 + e_5$	$e_1e_3 = -\frac{2}{3}e_4$
	$e_1 e_4 = -\frac{3}{4} e_5$	$e_2 e_1 = e_3$	$e_2 e_3 = \frac{1}{6} e_5$
F	$e_3e_1 = e_4$	$e_3 e_2 = -\frac{1}{4} e_5$	$e_4e_1 = e_5$
\mathbf{T}_{45}^{5}	$e_1e_1 = e_2$	$e_1e_2 = -e_3$	$e_1 e_3 = -\frac{2}{3}e_4 + e_5$
	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = e_5$
	$e_2 e_3 = -\frac{2}{3} e_5$	$e_3 e_1 = e_4$	$e_3 e_2 = e_5$
$\mathbf{T}_{46}^{5}(\lambda,eta)$	$e_1 e_1 = e_2$	$e_1 e_2 = \lambda e_3 + e_5$	$e_1e_3 = \frac{4\lambda - 1}{5 - 2\lambda}e_4 + \beta e_5$
$\lambda \neq 1, -2, -\frac{1}{2}, \frac{5}{2}$	$e_1 e_4 = \frac{4\lambda - 1}{5 - 2\lambda} e_5$	$e_2e_1=e_3$	$e_2 e_2 = \frac{(2\lambda+1)(\lambda+2)}{3(5-2\lambda)} e_4 + \beta e_5$
	$e_2 e_3 = \frac{2(2\lambda+1)(3\lambda^2+1)}{3\lambda(5-2\lambda)^2}$	$(1) e_5$	$e_3e_1 = e_4$
	$e_3 e_2 = \frac{2\lambda + 1}{5 - 2\lambda} e_5$	$e_4 e_1 = e_5$	
$\mathbf{T}_{47}^5(\lambda,eta)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3$	$e_1e_3 = \frac{4\lambda - 1}{5 - 2\lambda}e_4 + e_5$
$\lambda \neq 1, \frac{5}{2}$	$e_1e_4 = \beta e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{(2\lambda+1)(\lambda+2)}{3(5-2\lambda)}e_4 + e_5$
	$e_2 e_3 = \frac{\beta(4\lambda^2 - 2\lambda + 7) + \beta(4\lambda^2 - 2\lambda + 7)}{3\lambda(5 - 2\lambda)}$	$\frac{-2(\lambda-1)^2}{\lambda}e_5$	$e_3e_1 = e_4$
	$e_3e_2 = \left(\beta - \frac{2(\lambda-1)}{5-2\lambda}\right)$	e_5	$e_4 e_1 = e_5$
$\mathbf{T}_{48}^5(\lambda,eta)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1e_3 = \frac{4\lambda - 1}{5 - 2\lambda}e_4$
$\lambda \neq 0, \frac{5}{2}$	$e_1e_4 = \beta e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{\overset{3-2\lambda}{(2\lambda+1)(\lambda+2)}}{\overset{3(5-2\lambda)}{(2\lambda-2\lambda)}} e_4$
2	$e_2e_3 = \frac{\beta(4\lambda^2 - 2\lambda + 7) + \gamma}{2\lambda(5-2)}$	$\frac{-2(\lambda-1)^2}{\lambda}e_5$	$e_3e_1 = e_4$
	$e_3 e_2 = \left(\beta - \frac{2(\lambda - 1)}{5}\right)$	e_5	$e_4 e_1 = e_5$
T_{2}^{5}	$\frac{1}{2} \frac{1}{1} \frac{1}{2} \frac{1}$	$\frac{1}{\rho_1\rho_2} = \rho_2 + \rho_r$	$\rho_1 \rho_2 = \rho_4$
- 49	$e_1e_1 = e_5$	$e_1e_2 = e_3 + e_3$ $e_2e_1 = e_3$	$e_1e_3 = e_4$
	$e_2e_3 = e_5$	$e_3 e_1 = e_4$	$e_3e_2 = e_5$
	$e_4 e_1 = e_5$	0.1 .4	52 -0
\mathbf{T}_{50}^5	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4 + e_5$
00	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = e_4 + e_5$
	-	-	

	$e_2e_3 = e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	$e_4 e_1 = e_5$
$\mathbf{T}_{51}^5(eta,\gamma)$	$e_1e_1 = e_2$	$e_1 e_2 = -2e_3 + e_5$	$e_1e_3 = -e_4 + \beta e_5$	
	$e_1e_4 = \gamma e_5$	$e_2e_1 = e_3$	$e_2 e_2 = \beta e_5$	
	$e_2e_3 = -\frac{3\gamma+2}{6}e_5$	$e_3e_1 = e_4$	$e_3e_2 = \frac{3\gamma+2}{3}e_5$	$e_4 e_1 = e_5$
$\mathbf{T}_{52}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = -2e_3 + \beta e_5$	$e_1e_3 = -e_4 + e_5$	
	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = e_5$	
	$e_2 e_3 = -\frac{1}{2} e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
\mathbf{T}_{53}^5	$e_1e_1 = e_2$	$e_1 e_2 = -\frac{1}{2} e_3$	$e_1 e_3 = -\frac{1}{2}e_4 + e_5$	$e_1 e_4 = -\frac{1}{2} e_5$
	$e_2 e_1 = e_3$	$e_2 e_2 = e_5$	$e_3e_1 = e_4$	$e_4e_1 = e_5$
\mathbf{T}_{54}^5	$e_1e_1 = e_2$	$e_1 e_2 = -\frac{1}{2} e_3$	$e_1 e_3 = -\frac{1}{2}e_4 + e_5$	
	$e_1e_4 = e_5$	$e_2e_1 = e_3$	$e_2 e_2 = e_5$	
	$e_2e_3 = -e_5$	$e_3e_1=e_4$	$e_3e_2 = e_5$	
$\mathbf{T}_{55}^5(\lambda, \alpha)_{\lambda \neq -2,0}$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3$	$e_1e_3 = \alpha e_4 + e_5$	
$\alpha \neq \frac{2\lambda^2 + 5\lambda - 1}{6}$	$e_1e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = \left(\alpha + \frac{1-\lambda}{3}\right) e_4$	$+ e_5$
	$e_2 e_3 = \frac{3\alpha - 2(\lambda - 1)}{3\lambda} e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
$\mathbf{T}_{56}^5(\lambda,lpha)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3$	$e_1e_3 = \alpha e_4$	
$\lambda \neq 0$	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = \left(\alpha + \frac{1-\lambda}{3}\right) e_4$	
	$e_2 e_3 = \frac{3\alpha - 2(\lambda - 1)}{3\lambda} e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
$\mathbf{T}_{57}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = -2e_3$	$e_1 e_3 = -\frac{1}{2}e_4 + e_5$	
	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{1}{2}e_4 + e_5$	
	$e_2 e_3 = -\frac{3}{4} e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
$\mathbf{T}_{58}^5(lpha)$	$e_1e_1 = e_2$	$e_1 e_2 = -2e_3 + e_5$	$e_1e_3 = \alpha e_4$	
	$e_1e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = (\alpha + 1)e_4$	
	$e_3e_1=e_4$	$e_2e_3 = -\frac{\alpha+2}{2}e_5$	$e_3e_2 = e_5$	

3.6. 1-dimensional central extensions of $\mathbf{T}_{06}^4(\lambda).$

(1) $\lambda = 1$. Let us use the following notations:

$$\nabla_{1} = [\Delta_{12}], \quad \nabla_{2} = [\Delta_{31}], \quad \nabla_{3} = [\Delta_{14}] + [\Delta_{23}] + [\Delta_{32}], \quad \nabla_{4} = 3[\Delta_{41}] - [\Delta_{22}].$$

Take $\theta = \sum_{i=1}^{4} \alpha_{i} \nabla_{i} \in \mathrm{H}^{2}_{\mathrm{T}}(\mathbf{T}^{4}_{05}(1)).$ For the element
 $\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ x & 1 & 0 & 0 \\ y & 2x & 1 & 0 \\ z & 2y + x + x^{2} & 3x & 1 \end{pmatrix} \in \mathrm{Aut} (\mathbf{T}^{4}_{05}(1)),$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & 0 & \alpha_3 \\ 0 & -\alpha_4 & \alpha_3 & 0 \\ \alpha_2 & \alpha_3 & 0 & 0 \\ 3\alpha_4 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha^{**} + \alpha^* + \alpha_1^* & \alpha^* & \alpha_3^* \\ \alpha^{**} & \alpha^* - \alpha_4^* & \alpha_3^* & 0 \\ \alpha^* + \alpha_2^* & \alpha_3^* & 0 & 0 \\ 3\alpha_4^* & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl} \alpha_1^* &=& \alpha_1 - x(2\alpha_2 + 3\alpha_3 + 3\alpha_4) + (x^2 + 2y)(\alpha_3 - 3\alpha_4), & \alpha_2^* &=& \alpha_2 - 3x(\alpha_3 - 3\alpha_4), \\ \alpha_3^* &=& \alpha_3, & \alpha_4^* &=& \alpha_4. \end{array}$$

Since $(\alpha_3, \alpha_4) \neq (0, 0)$, we have the following cases:

- (a) if $\alpha_4 = 0$, then choosing $x = \frac{\alpha_2}{3\alpha_3}$, $y = \frac{-\alpha_1 + 2x\alpha_2 (x^2 3x)\alpha_3}{2\alpha_3}$, we have the representative $\langle \nabla_3 \rangle$.
- (b) if $\alpha_4 \neq 0$, then:
 - (i) if $\alpha_3 \neq 3\alpha_4$, then choosing $x = \frac{\alpha_2}{3(\alpha_3 3\alpha_4)}$, $y = \frac{-\alpha_1 + 2x\alpha_2 + (3x x^2)\alpha_3 + 3(x^2 + x)\alpha_4}{2(\alpha_3 3\alpha_4)}$, we have the representative $\langle \beta \nabla_3 + \nabla_4 \rangle_{\beta \neq 3}$.
 - (ii) if $\alpha_3 = 3\alpha_4$, then: (A) if $\alpha_2 \neq -6\alpha_4$, then choosing $y = \frac{\alpha_1}{\alpha_2 + 6\alpha_4}$, we have the representative $\langle \beta \nabla_2 + 3\nabla_3 + \nabla_4 \rangle_{\beta \neq -6}$.
 - (B) if $\alpha_2 = -6\alpha_4$, then we have the representative $\langle \beta \nabla_1 6\nabla_2 + 3\nabla_3 + \nabla_4 \rangle$.

(2)
$$\lambda = -1$$
.

Let us use the following notations:

$$\nabla_{1} = [\Delta_{12}], \quad \nabla_{2} = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_{3} = 3[\Delta_{14}] - 2[\Delta_{23}] + 3[\Delta_{32}], \quad \nabla_{4} = 2[\Delta_{13}] - 3[\Delta_{14}] + 6[\Delta_{41}].$$

Take $\theta = \sum_{i=1}^{4} \alpha_{i} \nabla_{i} \in \mathrm{H}^{2}_{\mathrm{T}}(\mathbf{T}^{4}_{06}(-1)).$ For the element

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ x & 1 & 0 & 0 \\ y & 0 & 1 & 0 \\ z & x + \frac{1}{3}y & 0 & 1 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{06}^{4}(-1)\right),$$

we have

$$\phi^{T} \begin{pmatrix} 0 & \alpha_{1} & \alpha_{2} + 2\alpha_{4} & 3\alpha_{3} - 3\alpha_{4} \\ 0 & \alpha_{2} & -2\alpha_{3} & 0 \\ 0 & 3\alpha_{3} & 0 & 0 \\ 6\alpha_{4} & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & -\alpha^{**} + 3\alpha^{*} + \alpha_{1}^{*} & -2\alpha^{*} + \alpha_{2}^{*} + 2\alpha_{4}^{*} & 3\alpha_{3}^{*} - 3\alpha_{4}^{*} \\ \alpha^{**} & \alpha_{2}^{*} & -2\alpha_{3}^{*} & 0 \\ 3\alpha^{*} & 3\alpha_{3}^{*} & 0 & 0 \\ 6\alpha_{4}^{*} & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl}
\alpha_1^* &=& \alpha_1 + x(2\alpha_2 + 3\alpha_4) + y(2\alpha_3 + \alpha_4), & \alpha_2^* &=& \alpha_2, \\
\alpha_3^* &=& \alpha_3, & & \alpha_4^* &=& \alpha_4.
\end{array}$$

Since $(\alpha_3, \alpha_4) \neq (0, 0)$, we have the following cases:

(a) if $\alpha_4 = 0$, then choosing $y = -\frac{\alpha_1 + 2\alpha_2 x}{2\alpha_3}$, we have the representatives $\langle \beta \nabla_2 + \nabla_3 \rangle$.

(b) if $\alpha_4 \neq 0$, then:

(i) if $\alpha_4 = -2\alpha_3$, $\alpha_2 = 3\alpha_3$, then we have the representative

$$\langle \beta \nabla_1 + 3 \nabla_2 + \nabla_3 - 2 \nabla_4 \rangle.$$

(ii) if
$$\alpha_4 = -2\alpha_3$$
, $\alpha_2 \neq 3\alpha_3$, then choosing $x = -\frac{\alpha_1}{2\alpha_2 - 6\alpha_3}$, we have the representative $\langle \beta \nabla_2 + \nabla_3 - 2\nabla_4 \rangle$.

(iii) if $\alpha_4 \neq -2\alpha_3$, then choosing $y = -\frac{\alpha_1 + x(2\alpha_2 + 3\alpha_4)}{2\alpha_3 + \alpha_4}$, we have the representative $\langle \beta \nabla_2 + \gamma \nabla_3 + \nabla_4 \rangle.$

(3) $\lambda = -\frac{1}{2}$. Let us use the following notations:

$$\nabla_{1} = [\Delta_{12}], \quad \nabla_{2} = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_{3} = [\Delta_{14}] - [\Delta_{23}] + [\Delta_{32}], \quad \nabla_{4} = 2[\Delta_{13}] - 3[\Delta_{14}] + 6[\Delta_{41}].$$

Take $\theta = \sum_{i=1}^{4} \alpha_{i} \nabla_{i} \in \mathrm{H}^{2}_{\mathrm{T}}(\mathbf{T}^{4}_{06}(-\frac{1}{2})).$ For the element

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ x & 1 & 0 & 0 \\ y & \frac{1}{2}x & 1 & 0 \\ z & x + \frac{1}{2}y & \frac{1}{2}x & 1 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{06}^{4}(-\frac{1}{2})\right),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 + 2\alpha_4 & \alpha_3 - 3\alpha_4 \\ 0 & \alpha_2 & -\alpha_3 & 0 \\ 0 & \alpha_3 & 0 & 0 \\ 6\alpha_4 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & -\frac{1}{2}\alpha^{**} + \alpha^* + \alpha_1^* & -\frac{1}{2}\alpha^* + \alpha_2^* + 2\alpha_4^* & \alpha_3^* - 3\alpha_4^* \\ \alpha^{**} & \alpha_2^* & -\alpha_3^* & 0 \\ \alpha^* & \alpha_3^* & 0 & 0 \\ 6\alpha_4^* & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl} \alpha_1^* &=& \alpha_1 + 2x(\alpha_2 - \alpha_4) - (\frac{1}{4}x^2 - y)\alpha_3, & \alpha_2^* &=& \alpha_2, \\ \alpha_3^* &=& \alpha_3, & & \alpha_4^* &=& \alpha_4. \end{array}$$

- Since $(\alpha_3, \alpha_4) \neq (0, 0)$, we have the following cases: (a) if $\alpha_4 = 0$, then choosing $y = -\frac{4\alpha_1 + 8x\alpha_2 x^2\alpha_3}{4\alpha_3}$, we have the representatives $\langle \beta \nabla_2 + \nabla_3 \rangle$.
- (b) if $\alpha_4 \neq 0$, then:

 - (i) if $\alpha_3 = 0$, $\alpha_2 = \alpha_4$, then we have the representative $\langle \beta \nabla_1 + \nabla_2 + \nabla_4 \rangle$. (ii) if $\alpha_3 = 0$, $\alpha_2 \neq \alpha_4$, then choosing $x = -\frac{\alpha_1}{2(\alpha_2 \alpha_4)}$, we have the representative $\langle \beta \nabla_2 + \alpha_4 \rangle$. ∇_4 .
- (iii) if $\alpha_3 \neq 0$, then choosing $y = -\frac{4\alpha_1 + 8x(\alpha_2 \alpha_4) x^2\alpha_3}{4\alpha_3}$, we have the representative $\langle \beta \nabla_2 + \alpha_3 \rangle = -\frac{4\alpha_1 + 8x(\alpha_2 \alpha_4) x^2\alpha_3}{4\alpha_3}$. (4) $\lambda \neq 0; 1; -1; -\frac{1}{2}$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = 6\lambda[\Delta_{14}] + (2\lambda^2 + \lambda + 3)[\Delta_{23}] + 6\lambda[\Delta_{32}].$$

Take
$$\theta = \sum_{i=1}^{3} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{06}(\lambda)).$$
 For the element

$$\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ x & 1 & 0 & 0 \\ y & (\lambda+1)x & 1 & 0 \\ z & \frac{2\lambda^2+3\lambda+1}{6}x^2 + x + \frac{2\lambda^2+5\lambda+5}{6}y & \frac{2\lambda^2+9\lambda+7}{6}x & 1 \end{pmatrix} \in \mathrm{Aut}\,(\mathbf{T}^4_{06}(\lambda)),$$

$$\phi^{T} \begin{pmatrix} 0 & \alpha_{1} & \alpha_{2} & 6\lambda\alpha_{3} \\ 0 & \alpha_{2} & (2\lambda^{2} + \lambda + 3)\alpha_{3} & 0 \\ 0 & 6\lambda\alpha_{3} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \lambda\alpha^{**} + \alpha^{*} + \alpha_{1}^{*} & \frac{2\lambda^{2} + 5\lambda - 1}{6}\alpha^{*} + \alpha_{2}^{*} & 6\lambda\alpha_{3}^{*} \\ \alpha^{**} & \frac{2\lambda^{2} + 3\lambda + 1}{6}\alpha^{*} + \alpha_{2}^{*} & (2\lambda^{2} + \lambda + 3)\alpha_{3}^{*} & 0 \\ \alpha^{*} & 6\lambda\alpha_{3}^{*} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rcl}
\alpha_1^* &=& \alpha_1 + 2x\alpha_2 + \left((-2\lambda^3 + 5\lambda + 3)x^2 + 4y\lambda(\lambda + 2) \right) \alpha_3, \\
\alpha_2^* &=& \alpha_2 + 3x(2\lambda^2 + 3\lambda + 1)\alpha_3, \\
\alpha_3^* &=& \alpha_3.
\end{array}$$

Since $\alpha_3 \neq 0$ we have the following cases: (a) if $\lambda \neq -2$, then choosing

$$x = -\frac{\alpha_2}{3(2\lambda^2 + 3\lambda + 1)\alpha_3}, \quad y = -\frac{1}{4\lambda(\lambda + 2)\alpha_3} \Big(\alpha_1 + 2x\alpha_2 - (2\lambda^3 - 5\lambda - 3)x^2\alpha_3\Big),$$

we have the representatives $\langle \nabla_3 \rangle$. (b) if $\lambda = -2$, then choosing $x = -\frac{\alpha_2}{9\alpha_3}$, we have the representative $\langle \beta \nabla_1 + \nabla_3 \rangle$. Summarizing all cases, we have the following distinct orbits:

$$\begin{split} \lambda &= 1 &: \langle \beta \nabla_3 + \nabla_4 \rangle, \qquad \langle \beta \nabla_2 + 3 \nabla_3 + \nabla_4 \rangle_{\beta \neq 0}, \qquad \langle \beta \nabla_1 - 6 \nabla_2 + 3 \nabla_3 + \nabla_4 \rangle_{\beta \neq 0} \\ \lambda &= -1 &: \langle \beta \nabla_2 + \nabla_3 \rangle_{\beta \neq 0}, \quad \langle \beta \nabla_1 + 3 \nabla_2 + \nabla_3 - 2 \nabla_4 \rangle_{\beta \neq 0}, \qquad \langle \beta \nabla_2 + \gamma \nabla_3 + \nabla_4 \rangle \\ \lambda &= -\frac{1}{2} &: \langle \beta \nabla_2 + \nabla_3 \rangle_{\beta \neq 0}, \quad \langle \beta \nabla_1 + \nabla_2 + \nabla_4 \rangle_{\beta \neq 0}, \qquad \langle \beta \nabla_2 + \gamma \nabla_3 + \nabla_4 \rangle \\ \lambda &= -2 &: \langle \beta \nabla_1 + \nabla_3 \rangle_{\beta \neq 0} \\ \lambda &= 0 &: \langle \nabla_3 \rangle. \end{split}$$

The corresponding algebras are:

$\mathbf{T}_{59}^5(eta)$	$e_1e_1 = e_2$	$e_1 e_2 = e_3 + e_4$	$e_1e_3 = e_4$	$e_1e_4 = \beta e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = e_4 - e_5$	$e_2 e_3 = \beta e_5$	$e_3e_1 = e_4$
	$e_3e_2 = \beta e_5$	$e_4 e_1 = 3 e_5$		
$\mathbf{T}_{60}^5(\beta)_{\beta \neq 0}$	$e_1e_1 = e_2$	$e_1 e_2 = e_3 + e_4$	$e_1e_3 = e_4$	$e_1e_4 = 3e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = e_4 - e_5$	$e_2e_3 = 3e_5$	$e_3e_1 = e_4 + \beta e_5$
	$e_3e_2 = 3e_5$	$e_4 e_1 = 3 e_5$		
$\mathbf{T}_{61}^5(\beta)_{\beta \neq 0}$	$e_1e_1 = e_2$	$e_1 e_2 = e_3 + e_4 + \beta e_5$	$e_1e_3 = e_4$	$e_1e_4 = 3e_5$
	$e_2 e_1 = e_3$	$e_2 e_2 = e_4 - e_5$	$e_2e_3 = 3e_5$	$e_3e_1 = e_4 - 6e_5$
	$e_3e_2 = 3e_5$	$e_4 e_1 = 3 e_5$		
$\mathbf{T}_{62}^5(\beta)_{\beta \neq 0}$	$e_1e_1 = e_2$	$e_1 e_2 = -e_3 + e_4$	$e_1 e_3 = -\frac{2}{3}e_4 + \beta e_5$	$e_1e_4 = 3e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = \beta e_5$	$e_2e_3 = -2e_5$	$e_3e_1 = e_4$
	$e_3e_2 = 3e_5$			

$\mathbf{T}_{63}^5(eta)_{eta eq 0}$	$e_1e_1 = e_2$	$e_1 e_2 = -e_3 + e_4 + \beta e_5$	$e_1 e_3 = -\frac{2}{3}e_4 - e_5$	$e_1e_4 = 9e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = 3e_5$	$e_2e_3 = -2e_5$	$e_3e_1 = e_4$
	$e_3e_2 = 3e_5$	$e_4e_1 = -12e_5$		
$\mathbf{T}_{64}^5(eta,\gamma)$	$e_1e_1 = e_2$	$e_1 e_2 = -e_3 + e_4$	$e_1e_3 = -\frac{2}{3}e_4 + (\beta + 2)e_5$	$e_1e_4 = 3(\gamma - 1)e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = \beta e_5$	$e_2 e_3 = -2\gamma e_5$	$e_3e_1 = e_4$
	$e_3e_2 = 3\gamma e_5$	$e_4 e_1 = 6 e_5$		
$\mathbf{T}_{65}^5(\beta)_{\beta \neq 0}$	$e_1e_1 = e_2$	$e_1 e_2 = -\frac{1}{2}e_3 + e_4$	$e_1 e_3 = -\frac{1}{2} e_4 + \beta e_5$	$e_1e_4 = e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = \beta e_5$	$e_2e_3 = -e_5$	$e_3e_1 = e_4$
	$e_3e_2 = e_5$			
$\mathbf{T}_{66}^5(eta)_{eta eq 0}$	$e_1e_1 = e_2$	$e_1 e_2 = -\frac{1}{2}e_3 + e_4 + \beta e_5$	$e_1 e_3 = -\frac{1}{2}e_4 + 3e_5$	$e_1e_4 = -3e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = e_5$	$e_3e_1 = e_4$	$e_4 e_1 = 6 e_5$
$\mathbf{T}_{67}^5(eta,\gamma)$	$e_1e_1 = e_2$	$e_1 e_2 = -\frac{1}{2}e_3 + e_4$	$e_1e_3 = -\frac{1}{2}e_4 + (\beta + 2)e_5$	$e_1e_4 = (\gamma - 3)e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = \beta e_5$	$e_2 e_3 = -\gamma e_5$	$e_3e_1 = e_4$
	$e_3e_2 = \gamma e_5$	$e_4 e_1 = 6 e_5$		
$\mathbf{T}_{68}^5(eta)_{eta eq 0}$	$e_1e_1 = e_2$	$e_1 e_2 = -2e_3 + e_4 + \beta e_5$	$e_1e_3 = -\frac{1}{2}e_4$	$e_1e_4 = 4e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = \frac{1}{2} e_4$	$e_2e_3 = -3e_5$	$e_3e_1 = e_4$
	$e_3e_2 = 4e_5$			
$\mathbf{T}_{69}^5(\lambda)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3 + e_4$	$e_1e_3 = \frac{2\lambda^2 + 5\lambda - 1}{6}e_4$	$e_1 e_4 = 6\lambda e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = \frac{(2\lambda+1)(\lambda+1)}{6} e_4$	$e_2e_3 = (2\lambda^2 + \lambda + 3)e_5$	$e_3e_1 = e_4$
	$e_3 e_2 = 6\lambda e_5$	-		

3.7. 1-dimensional central extensions of $\mathbf{T}_{07}^4(\lambda).$

(1) $\lambda \neq -1$. Let us use the following notations:

$$\nabla_1 = [\Delta_{12}], \quad \nabla_2 = (1 - \lambda)[\Delta_{22}] + [\Delta_{31}], \quad \nabla_3 = \lambda[\Delta_{14}] + [\Delta_{23}].$$

Take
$$\theta = \sum_{i=1}^{\infty} \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{07}(\lambda))$$
. For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & xy & 0 & x^4 \end{pmatrix} \in \operatorname{Aut} (\mathbf{T}_{07}^4(\lambda)),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & 0 & \lambda \alpha_3 \\ 0 & (1-\lambda)\alpha_2 & \alpha_3 & 0 \\ 3\alpha_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & \alpha_1^* + \alpha^* & 0 & \alpha_3^* \\ \alpha^* & (1-\lambda)\alpha_2^* & \alpha_3^* & 0 \\ 3\alpha_2^* & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$\alpha_1^* = x^3 \alpha_1, \quad \alpha_2^* = x^4 \alpha_2, \quad \alpha_3^* = x^5 \alpha_3.$$

Since $\alpha_3 \neq 0$, then we have the representatives

$$\langle \alpha \nabla_1 + \nabla_2 + \nabla_3 \rangle_{\lambda \neq -1}, \ \langle \nabla_1 + \nabla_3 \rangle_{\lambda \neq -1} \text{ and } \langle \nabla_3 \rangle_{\lambda \neq -1}$$

depending on whether $\alpha_1 = 0$, $\alpha_2 = 0$ or not. (2) $\lambda = -1$. Let us use the following notations:

$$\nabla_{1} = [\Delta_{12}], \quad \nabla_{2} = 2[\Delta_{22}] + 3[\Delta_{31}], \quad \nabla_{3} = [\Delta_{14}] - [\Delta_{23}], \quad \nabla_{4} = [\Delta_{41}] + 2[\Delta_{23}] - 3[\Delta_{32}].$$

Take $\theta = \sum_{i=1}^{4} \alpha_{i} \nabla_{i} \in \mathrm{H}^{2}_{\mathrm{T}}(\mathbf{T}^{4}_{07}(-1)).$ For the element
$$\begin{pmatrix} x & 0 & 0 & 0\\ 0 & x^{2} & 0 & 0 \end{pmatrix}$$

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & xy & 0 & x^4 \end{pmatrix} \in \operatorname{Aut} (\mathbf{T}_{07}^4(-1)),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & 0 & \alpha_3 \\ 0 & 2\alpha_2 & 2\alpha_4 - \alpha_3 & 0 \\ 3\alpha_2 & -3\alpha_4 & 0 & 0 \\ \alpha_4 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{**} & \alpha_1^* + \alpha^* & 0 & \alpha_3^* \\ \alpha^* & 2\alpha_2^* & 2\alpha_4^* - \alpha_3^* & 0 \\ 3\alpha_2^* & -3\alpha_4^* & 0 & 0 \\ \alpha_4^* & 0 & 0 & 0 \end{pmatrix},$$

where

$$\begin{array}{rclrcl}
\alpha_1^* &=& x^3 \alpha_1, & \alpha_2^* &=& x^4 \alpha_2, \\
\alpha_3^* &=& x^5 \alpha_3, & \alpha_4^* &=& x^5 \alpha_4.
\end{array}$$

Since $(\alpha_3, \alpha_4) \neq (0, 0)$, we have the following cases:

(a) if $\alpha_4 = 0$, then and we have the representatives

$$\langle \alpha \nabla_1 + \nabla_2 + \nabla_3 \rangle_{\lambda = -1}, \ \langle \nabla_1 + \nabla_3 \rangle_{\lambda = -1} \text{ and } \langle \nabla_3 \rangle_{\lambda = -1}$$

depending on whether $\alpha_1 = 0$, $\alpha_2 = 0$ or not.

(b) if $\alpha_4 \neq 0$, then we have three distinct orbits with family of representatives

$$\langle \nabla_1 + \alpha \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle, \ \langle \nabla_2 + \alpha \nabla_3 + \nabla_4 \rangle \text{ and } \langle \alpha \nabla_3 + \nabla_4 \rangle$$

Summarizing, we have the following representatives of distinct orbits:

$$\langle \alpha \nabla_1 + \nabla_2 + \nabla_3 \rangle, \quad \langle \nabla_1 + \nabla_3 \rangle, \quad \langle \nabla_2 \rangle, \quad \langle \nabla_1 + \alpha \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle, \quad \langle \nabla_2 + \alpha \nabla_3 + \nabla_4 \rangle, \quad \langle \alpha \nabla_3 + \nabla_4 \rangle.$$

Hence, we have following 5-dimensional algebras:

$\mathbf{T}_{70}^5(\lambda,lpha)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3 + \alpha e_5$	$e_1e_3 = e_4$	$e_1 e_4 = \lambda e_5$
	$e_2 e_1 = e_3$	$e_2 e_2 = e_4 + (1 - \lambda) e_5$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_5$
$\mathbf{T}_{71}^5(\lambda)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_5$	$e_1e_3 = e_4$	$e_1 e_4 = \lambda e_5$
	$e_2 e_1 = e_3$	$e_2 e_2 = e_4$	$e_2 e_3 = e_5$	
$\mathbf{T}_{72}^5(\lambda)$	$e_1e_1 = e_2$	$e_1 e_2 = \lambda e_3$	$e_1e_3 = e_4$	$e_1e_4 = \lambda e_5$
	$e_2e_1 = e_3$	$e_2 e_2 = e_4$	$e_2e_3 = e_5$	

$\mathbf{T}_{73}^5(lpha,eta)$	$e_1e_1 = e_2$	$e_1 e_2 = -e_3 + e_5$	$e_1e_3 = e_4$	$e_1e_4 = \alpha e_5$	$e_2e_1 = e_3$
	$e_2 e_2 = e_4 + 2\beta e_5$	$e_2 e_3 = (2 - \alpha) e_5$	$e_3e_1 = 3\beta e_5$	$e_3e_2 = -3e_5$	$e_4e_1 = e_5$
$\mathbf{T}_{74}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_2 = -e_3$	$e_1e_3 = e_4$	$e_1e_4 = \alpha e_5$	$e_2e_1 = e_3$
	$e_2 e_2 = e_4 + 2e_5$	$e_2 e_3 = (2 - \alpha) e_5$	$e_3e_1 = 3e_5$	$e_3e_2 = -3e_5$	$e_4e_1 = e_5$
$\mathbf{T}_{75}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_2 = -e_3$	$e_1e_3 = e_4$	$e_1e_4 = \alpha e_5$	$e_2e_1 = e_3$
	$e_2 e_2 = e_4$	$e_2 e_3 = (2 - \alpha) e_5$	$e_3e_2 = -3e_5$	$e_4e_1 = e_5$	

3.8. 1-dimensional central extensions of T_{08}^4 . Let us use the following notations:

 $\nabla_1 = [\Delta_{12}], \quad \nabla_2 = [\Delta_{13}] + [\Delta_{22}], \quad \nabla_3 = [\Delta_{22}] + 3[\Delta_{31}], \quad \nabla_4 = [\Delta_{24}].$ Take $\theta = \sum_{i=1}^4 \alpha_i \nabla_i \in \mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{08}).$ For the element

$$\phi = \begin{pmatrix} x & 0 & 0 & 0 \\ 0 & x^2 & 0 & 0 \\ y & 0 & x^3 & 0 \\ z & 0 & x^2 y & x^5 \end{pmatrix} \in \operatorname{Aut}\left(\mathbf{T}_{08}^4\right),$$

we have

$$\phi^T \begin{pmatrix} 0 & \alpha_1 & \alpha_2 & 0 \\ 0 & \alpha_2 + \alpha_3 & 0 & \alpha_4 \\ 3\alpha_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \phi = \begin{pmatrix} \alpha^{***} & \alpha_1^* & \alpha_2^* & 0 \\ \alpha^{**} & \alpha_2^* + \alpha_3^* & \alpha^* & \alpha_4^* \\ 3\alpha_3^* & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

where

$$\begin{array}{rcl} \alpha_1^* &=& x^3 \alpha_1, & \alpha_2^* &=& x^4 \alpha_2, \\ \alpha_3^* &=& x^4 \alpha_3, & \alpha_4^* &=& x^7 \alpha_4. \end{array}$$

Since $\alpha_4 \neq 0$, we have the following distinct orbits with representatives

$$\langle \nabla_1 + \alpha \nabla_2 + \beta \nabla_3 + \nabla_4 \rangle, \ \langle \nabla_2 + \alpha \nabla_3 + \nabla_4 \rangle, \ \langle \nabla_3 + \nabla_4 \rangle, \text{ and } \langle \nabla_4 \rangle.$$

Hence, we have following 5-dimensional algebras:

$\mathbf{T}_{76}^5(lpha,eta)$	$e_1e_1 = e_2$	$e_1e_2 = e_5$	$e_1e_3 = \alpha e_5$	$e_2 e_1 = e_3 e_2 e_2 = (\alpha + \beta) e_5$
	$e_2e_3=e_4$	$e_2 e_4 = e_5$	$e_3e_1 = 3\beta e_5$	
$\mathbf{T}_{77}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_3 = e_5$	$e_2e_1 = e_3$	$e_2 e_2 = (\alpha + 1)e_5$
	$e_2 e_3 = e_4$	$e_2 e_4 = e_5$	$e_3e_1 = 3\alpha e_5$	
\mathbf{T}_{78}^5	$e_1e_1 = e_2$	$e_2e_1 = e_3$	$e_2 e_2 = e_5$	$e_2e_3 = e_4$ $e_2e_4 = e_5$ $e_3e_1 = 3e_5$
${f T}_{79}^5$	$e_1e_1 = e_2$	$e_2e_1 = e_3$	$e_2 e_3 = e_4$	$e_2 e_4 = e_5$

3.9. Classification theorem. Summarizing results of the present sections, we have the following theorem.

Theorem B. Let A be a complex 5-dimensional one-generated nilpotent terminal algebra. Then A is isomorphic to one of the algebras $\mathbf{T}_{01}^5 - \mathbf{T}_{79}^5$ found in Table C (see Appendix).

APPENDIX

\mathbf{T}_{01}^4	:	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_2 e_1 = e_3$			
$\mathbf{T}_{02}^4(\alpha)$:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = \alpha e_4$	$e_2 e_2 = e_4$	$e_3e_1 = -3e_4$	
\mathbf{T}_{03}^4	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$			
\mathbf{T}_{04}^4	:	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_2 e_1 = e_4$		
$\mathbf{T}_{05}^4(\lambda, \alpha)$	$\alpha)$:	$e_1e_1=e_2$	$e_1e_2 = \lambda e_3$	$e_1e_3 = \alpha e_4$	$e_2 e_1 = e_3$	$e_2 e_2 = \left(\alpha + \frac{1-\lambda}{3}\right)e_4$	$e_3e_1 = e_4$
$\mathbf{T}_{06}^4(\lambda)$:	$e_1e_1=e_2$	$e_1e_2 = \lambda e_3 + e_4$	$e_1e_3 = \frac{2\lambda + 5\lambda - 1}{6}e_4$	$e_2 e_1 = e_3$	$e_2e_2 = \frac{(2\lambda+1)(\lambda+1)}{6}e_4$	$e_3e_1 = e_4$
$\mathbf{T}_{07}^4(\lambda)$:	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1 e_3 = e_4$	$e_2 e_1 = e_3$	$e_2 e_2 = e_4$	
\mathbf{T}_{08}^4	:	$e_1e_1 = e_2$	$e_2 e_1 = e_3$	$e_2 e_3 = e_4$			
\mathbf{T}_{09}^4	:	$e_1e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_3$	$e_2 e_3 = e_4$		
$\mathbf{T}_{10}^4(\alpha)$:	$e_1 e_1 = e_2$	$e_1e_2 = \alpha e_4$	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{1}{2} e_4$	$e_2 e_3 = e_4$	$e_3e_1 = e_4$

Table A. The list of 4-dimensional one-generated nilpotent terminal algebras.

Table B. Cohomology groups of 4-dimensional one-generated nilpotent terminal algebras.

$B_{T}^{2}(T_{01}^{4})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{21} \rangle$
$Z_{T}^{2}(T_{01}^{4})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{14}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23}, \Delta_{31} + \Delta_{41} \rangle$
$\mathrm{H}^{2}_{\mathrm{T}}(\mathbf{T}^{4}_{01})$	=	$\langle [\Delta_{13}] + [\Delta_{22}], [\Delta_{14}], [\Delta_{22}] + 3[\Delta_{31}], [\Delta_{23}], [\Delta_{31}] + [\Delta_{41}] \rangle$
$B_{\rm T}^2(\mathbf{T}_{02}^4(\alpha)_{\alpha\neq 6})$	=	$\langle \Delta_{11}, \Delta_{12}, \alpha \Delta_{13} + \Delta_{22} - 3 \Delta_{13} \rangle$
$\mathbf{Z}_{\mathrm{T}}^{2}(\mathbf{T}_{02}^{4}(\alpha)_{\alpha\neq 6})$	=	$\left\langle \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14} + 2\Delta_{23} - 3\Delta_{32}, \Delta_{21}, \Delta_{22} - 3\Delta_{31} \right\rangle$
$\mathrm{H}^2_{\mathrm{T}}(\mathbf{T}^4_{02}(\alpha)_{\alpha\neq 6})$	=	$\langle [\Delta_{13}], [\Delta_{14}] + 2[\Delta_{23}] - 3[\Delta_{32}], [\Delta_{21}] \rangle$
$B_T^2(T_{02}^4(6))$	=	$\langle \Delta_{11}, \Delta_{12}, 6\Delta_{13} + \Delta_{22} - 3\Delta_{13} \rangle$
$Z_T^2(T_{02}^4(6))$	=	$ \langle \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14} + 2\Delta_{23} - 3\Delta_{32}, \Delta_{21}, \Delta_{22} - 3\Delta_{31}, \Delta_{41} + \Delta_{23} - 3\Delta_{32} \rangle $
$H_T^2(T_{02}^4(6))$	=	$\langle [\Delta_{13}], [\Delta_{14}] + 2[\Delta_{23}] - 3[\Delta_{32}], [\Delta_{21}], [\Delta_{41}] + [\Delta_{23}] - 3[\Delta_{32}] \rangle$
$\mathrm{B}^2_\mathrm{T}(\mathbf{T}^4_{03})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} \rangle$
$Z_{T}^{2}(T_{03}^{4})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14}, \Delta_{21}, \Delta_{22} - 3\Delta_{31} \rangle$
$\mathrm{H}^2_\mathrm{T}(\mathbf{T}^4_{03})$	=	$\langle [\Delta_{14}], [\Delta_{21}], [\Delta_{22}] - 3[\Delta_{31}] \rangle$
$\mathrm{B}^2_\mathrm{T}(\mathbf{T}^4_{04})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{21} \rangle$
$\mathrm{Z}_\mathrm{T}^2(\mathbf{T}_{04}^4)$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14} + 3\Delta_{31}, \Delta_{21}, \Delta_{22} - 3\Delta_{31} \rangle$
$\mathrm{H}^2_\mathrm{T}(\mathbf{T}^4_{04})$	=	$\langle [\Delta_{13}], [\Delta_{14}] + 3[\Delta_{31}], [\Delta_{22}] - 3[\Delta_{31}] \rangle$
$B_T^2(T_{05}^4(1,0))$	=	$\langle \Delta_{11}, \Delta_{12} + \Delta_{21}, \Delta_{31} \rangle$
$Z_T^2(\mathbf{T}_{05}^4(1,0))$	=	$\left< \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{14} + \Delta_{32}, \Delta_{31}, \Delta_{34} \right>$
$H_T^2(\mathbf{T}_{05}^4(1,0))$	=	$\langle [\Delta_{12}], [\Delta_{13}] + [\Delta_{22}], [\Delta_{14}] + [\Delta_{32}], [\Delta_{34}] \rangle$
$B_T^2(T_{05}^4(0,\frac{1}{3}))$	=	$\langle \Delta_{11}, \Delta_{21}, \Delta_{13} + 2\Delta_{22} + 3\Delta_{31} \rangle$
$Z_T^2(T_{05}^4(0,\frac{1}{3}))$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23}, \Delta_{24} + \Delta_{33} \rangle$
$H_{T}^{2}(T_{05}^{4}(0,\frac{1}{3}))$	=	$\langle [\Delta_{12}], [\Delta_{13}] + [\Delta_{22}], [\Delta_{23}], [\Delta_{24}] + [\Delta_{33}] \rangle$
$B_{T}^{2}(T_{05}^{4}(0,-\frac{2}{3}))$	=	$\langle \Delta_{11}, \Delta_{21}, 2\Delta_{13} + \Delta_{22} - 3\Delta_{31} \rangle$
$Z_{T}^{2}(T_{05}^{4}(0,-\frac{2}{3}))$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23}, \Delta_{14} + \Delta_{32} \rangle$
$H_T^2(\mathbf{T}_{05}^4(0,-\frac{2}{3}))$	=	$\langle [\Delta_{12}], [\Delta_{13}] + [\Delta_{22}], [\Delta_{23}], [\Delta_{14}] + [\Delta_{32}] \rangle$
$B_{T}^{2}(T_{05}^{4}(0,-\frac{1}{5}))$	=	$\left\langle \Delta_{11}, \Delta_{21}, -\frac{1}{5}\Delta_{13} + \frac{2}{15}\Delta_{22} + \Delta_{31} \right\rangle$
$Z_{T}^{2}(T_{05}^{4}(0, -\frac{1}{5}))$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{23}, \Delta_{22} + 3\Delta_{31}, \Delta_{14} - \frac{2}{5}\Delta_{32} - \frac{7}{2}\Delta_{41} \rangle$
		0 2 1

$$\begin{split} & \frac{\mathrm{H}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{4}(0,-\frac{1}{3})) = \langle [\Delta_{12}], [\Delta_{13}] + [\Delta_{22}], [\Delta_{23}], [\Delta_{14}] - \frac{2}{5}[\Delta_{22}] - \frac{2}{5}[\Delta_{41}] \rangle \\ & \alpha \neq \frac{1}{3}; -\frac{2}{5}; -\frac{1}{5} \\ & \mathrm{B}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{4}(0,\alpha)) = \langle \Delta_{11}, \Delta_{21}, \alpha \Delta_{13} + (\alpha + \frac{2}{3}) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{Z}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(0,\alpha)) = \langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23} \rangle \\ & \mathrm{H}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{4}(0,\alpha)) = \langle (\Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23} \rangle \\ & \mathrm{B}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(-1,\alpha)) = \langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, 2\Delta_{22} + 3\Delta_{31}, \Delta_{23} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{4}(2, \Delta_{13}) + (\Delta_{22}), [\Delta_{13}] + [\Delta_{22}], [\Delta_{14}] - \frac{3\alpha_{24}}{3} \Delta_{23} - \frac{3\alpha_{24}}{3} \Delta_{23} + 2\Delta_{24} \rangle \\ & \mathrm{H}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(-1,\alpha)) = \langle \Delta_{11}, \lambda_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (\Delta_{23}) + [\Delta_{32}] \rangle \\ & \mathrm{L}_{\mathrm{T}}^{2}(0, -1; \frac{5}{2}, \alpha = \frac{4\lambda_{-1}}{3} \rangle \\ & \mathrm{B}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(\lambda, \alpha)) = \langle \Delta_{11}, \lambda_{12}, \Delta_{12} + \Delta_{22}, \Delta_{21}, (1, -\lambda) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{4}(1, \alpha, \alpha) = \langle \Delta_{11}, \lambda_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (1, -\lambda) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{4}(\mathrm{T}_{\mathrm{dc}}^{4}(\lambda, \alpha)) = \langle \Delta_{11}, \lambda_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (1, -\lambda) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{4}(\mathrm{T}_{\mathrm{dc}}^{4}(\lambda, \alpha)) = \langle \Delta_{11}, \lambda_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (1, -\lambda) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(\lambda, \alpha)) = \langle \Delta_{11}, \lambda_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (1, -\lambda) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(\lambda, \alpha)) = \langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (1, -\lambda) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(\lambda, \alpha)) = \langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (1, -\lambda) \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(0)) = \langle \Delta_{11}, \Delta_{12}, \Delta_{14} + \Delta_{22}, \Delta_{14}, \Delta_{22} + \Delta_{31} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(0)) = \langle \Delta_{11}, \Delta_{12}, \Delta_{14} + \Delta_{22}, \Delta_{21}, \Delta_{23} + \Delta_{23} \rangle \\ & \mathrm{L}_{\mathrm{T}}^{2}(\mathrm{T}_{\mathrm{dc}}^{6}(0)) = \langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{13}, \Delta_{14} + \Delta_{22} \rangle \\ & \mathrm{L}_{$$

$\mathrm{H}^2_\mathrm{T}(\mathbf{T}^4_{06}(\lambda))$	=	$\langle [\Delta_{12}], \Delta_{13} + \Delta_{22}, 6\lambda[\Delta_{14}] + (2\lambda^2 + \lambda + 3)[\Delta_{23}] + 6\lambda[\Delta_{32}] \rangle$
${\rm B}_{ m T}^2({f T}_{07}^4(-1))$	=	$\langle \Delta_{11}, -\Delta_{12} + \Delta_{21}, \Delta_{13} + \Delta_{22} \rangle$
$Z_T^2(T_{07}^4(-1))$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, 2\Delta_{22} + 3\Delta_{31}, \Delta_{14} - \Delta_{23}, 2\Delta_{23} - 3\Delta_{32} + \Delta_{41} \rangle$
$H_T^2(T_{07}^4(-1))$	=	$\langle [\Delta_{12}], 2[\Delta_{22}] + 3[\Delta_{31}], [\Delta_{14}] - [\Delta_{23}], 2[\Delta_{23}] - 3[\Delta_{32}] + [\Delta_{41}] \rangle$
$\mathrm{B}_{\mathrm{T}}^{2}(\mathbf{T}_{07}^{4}(\lambda)_{\lambda\neq-1})$	=	$\langle \Delta_{11}, \lambda \Delta_{12} + \Delta_{21}, \Delta_{13} + \Delta_{22} \rangle$
$\mathbf{Z}_{\mathrm{T}}^{2}(\mathbf{T}_{07}^{4}(\lambda)_{\lambda\neq-1})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, (1-\lambda)\Delta_{22} + 3\Delta_{31}, \lambda\Delta_{14} + \Delta_{23} \rangle$
$\mathrm{H}^{2}_{\mathrm{T}}(\mathbf{T}^{4}_{07}(\lambda)_{\lambda\neq-1})$	=	$\langle [\Delta_{12}], (1-\lambda)[\Delta_{22}] + 3[\Delta_{31}], \lambda[\Delta_{14}] + [\Delta_{23}] \rangle$
$\mathrm{B}_{\mathrm{T}}^2(\mathbf{T}_{08}^4)$	=	$\langle \Delta_{11}, \Delta_{21}, \Delta_{23} \rangle$
$Z_{T}^{2}(T_{08}^{4})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23}, \Delta_{24} \rangle$
$\mathrm{H}^2_\mathrm{T}(\mathbf{T}^4_{08})$	=	$\langle [\Delta_{13}] + [\Delta_{22}], [\Delta_{21}], [\Delta_{22}] + 3[\Delta_{31}], [\Delta_{24}] \rangle$
$\mathrm{B}_{\mathrm{T}}^2(\mathbf{T}_{09}^4)$	=	$\langle \Delta_{11}, \Delta_{12} + \Delta_{23}, \Delta_{21} \rangle$
$Z_{T}^{2}(T_{09}^{4})$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} - 3\Delta_{31}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23} \rangle$
${ m H}_{ m T}^2({f T}_{09}^4)$	=	$\langle [\Delta_{12}], [\Delta_{13}] - 3[\Delta_{31}], [\Delta_{22}] + 3[\Delta_{31}] \rangle$
$\mathrm{B}_{\mathrm{T}}^{2}(\mathbf{T}_{10}^{4}(lpha))$	=	$\left\langle \Delta_{11}, \Delta_{21}, \alpha \Delta_{12} + \frac{1}{3} \Delta_{22} + \Delta_{23} + \Delta_{31} \right\rangle$
$\mathrm{Z}_{\mathrm{T}}^{2}(\mathbf{T}_{10}^{4}(lpha))$	=	$\langle \Delta_{11}, \Delta_{12}, \Delta_{13} + \Delta_{22}, \Delta_{21}, \Delta_{22} + 3\Delta_{31}, \Delta_{23} \rangle$
$\mathrm{H}^2_\mathrm{T}(\mathbf{T}^4_{10}(lpha))$	=	$\langle [\Delta_{12}], [\Delta_{22}] + 3[\Delta_{31}], [\Delta_{23}] \rangle$

Table C. The list of 5-dimensional one-generated nilpotent terminal algebras.

T_{01}^{5}	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1 e_3 = e_4$	$e_2e_1 = e_5$
$\mathbf{T}_{02}^5(lpha)$	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = \alpha e_5$	
	$e_2e_1 = e_4$	$e_2e_2 = e_5$	$e_3e_1 = -3e_5$	
T_{03}^{5}	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	
	$e_2e_2 = e_5$	$e_3e_1 = -3e_5$		
T_{04}^{5}	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	
	$e_2e_1 = e_4$	$e_2e_2 = e_5$	$e_3e_1 = -3e_5$	
$\mathbf{T}_{05}^5(\lambda)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_4$	$e_1 e_3 = e_5$	
	$e_2e_1 = e_3$	$e_2e_2 = e_5$		
$\mathbf{T}_{06}^5(\lambda, \alpha)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_4$	$e_1e_3 = \alpha e_5$	
	$e_2e_1 = e_3$	$e_2 e_2 = \left(\alpha + \frac{1-\lambda}{3}\right) e_5$	$e_3 e_1 = e_5$	
$\mathbf{T}_{07}^5(\lambda)$	$e_1 e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1 e_3 = e_4$	
	$e_2e_1 = e_3$	$e_2 e_2 = e_4 + \frac{1-\lambda}{3} e_5$	$e_3 e_1 = e_5$	
$\mathbf{T}_{08}^5(\lambda)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_5$	$e_1 e_3 = e_4$	
	$e_2e_1 = e_3$	$e_2 e_2 = e_4 + \frac{1-\lambda}{3} e_5$	$e_3 e_1 = e_5$	
T_{09}^{5}	$e_1e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_3$	$e_2 e_3 = e_5$
T_{10}^{5}	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_2e_1 = e_3$	
	$e_2e_2 = e_5$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_5$	
T_{11}^5	$e_1e_1 = e_2$	$e_1 e_3 = e_4$	$e_2e_1 = e_3$	
	$e_2e_2 = e_4$	$e_2 e_3 = e_5$		
T_{12}^{5}	$e_1e_1 = e_2$	$e_1e_2 = e_5$	$e_1 e_3 = e_4$	
	$e_2e_1 = e_3$	$e_2e_2 = e_4$	$e_2 e_3 = e_5$	
$\mathbf{T}_{13}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_2 = \alpha e_5$	$e_1 e_3 = e_4$	$e_2 e_1 = e_3$
	$e_2 e_2 = e_4 + e_5$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_4$	
$\mathbf{T}_{14}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3$	
	$e_2e_2 = (\alpha + 1)e_4$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_4$	
$\mathbf{T}_{15}^5(\alpha)$	$e_1e_1 = e_2$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	$e_2 e_1 = e_3$
	$e_2 e_2 = (\alpha + 1)e_4$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_4$	
$\mathbf{T}_{16}^5(\alpha,\beta)$	$e_1e_1 = e_2$	$e_1e_2 = \beta e_4 + e_5$	$e_1e_3 = \overline{\alpha e_4}$	$e_2 e_1 = e_3$
	$e_2 e_2 = (\alpha + 1)e_4$	$e_2 e_3 = e_5$	$e_3e_1 = 3e_4$	

$\mathbf{T}_{17}^{\mathrm{o}}(lpha,eta)$	$e_1e_1 = e_2$	$e_1 e_2 = e_4$	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$
	$e_2 e_2 = \alpha e_5$	$e_2 e_3 = e_5$	$e_3e_1 = (3\alpha + \beta)e_5$	$e_4e_1 = \beta e_5$
$T_{18}^5(\alpha)$	$e_1e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_3$	$e_2 e_2 = \alpha e_5$
	$e_2 e_3 = e_5$	$e_3e_1 = (3\alpha + 1)e_5$	$e_4e_1 = e_5$	
$\mathbf{T}_{10}^5(\alpha,\beta,\gamma)$	$e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_1e_3 = \alpha e_5$	$e_1e_4 = \beta e_5$
$(\alpha; \gamma) \neq (0; -\frac{1}{2})$	$e_2 e_1 = e_3$	$e_2 e_2 = (\alpha + \gamma) e_5$	$e_3e_1 = (3\gamma + 1)e_5$	$e_4 e_1 = e_5$
$\mathbf{T}_{20}^{5}(\alpha,\beta)$	$e_1e_1 = e_2$	$e_1 e_2 = e_4$	$e_1e_3 = \alpha e_5$	$e_1e_4 = e_5$
$(\alpha; \beta) \neq (0; 0)$	$e_2e_1 = e_3$	$e_2e_2 = (\alpha + \beta)e_5$	$e_3e_1 = 3\beta e_5$	114 0
$\mathbf{T}_{2}^{5}(\alpha)$	$e_1e_1 = e_2$	$e_1e_2 = e_2$	$e_1e_2 = \alpha e_4$	$e_1e_4 = e_5$
- 21 (**)	$e_2e_2 = e_4$	$e_2e_2 = 2e_5$	$e_2e_1 = -3e_4$	$e_2e_2 = -3e_5$
$\mathbf{T}_{-}^{5}(\alpha)$	$e_{2}e_{2} = e_{2}$	$e_2 e_3 = e_3$	$e_1e_2 = oe_4 + e_5$	$e_1e_4 = e_5$
- 22(a)	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = ae_4 + e_3$	$e_1e_4 = e_3$
$\mathbf{T}^{5}(\beta)$	$e_2 e_2 = e_4$	$c_2 c_3 = 2 c_5$	$c_3c_1 = 5c_4$	$e_3e_2 = -5e_5$
$1_{23}(p)$	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = 0e_4$	
	$e_1e_4 = \rho e_5$	$e_2e_2 = e_4$	$e_2e_3 = (2\beta + 1)e_5$	
$\mathbf{T}_{5}(\rho)$	$e_3e_1 = -3e_4$	$e_3e_2 = -3(p+1)e_5$	$e_4e_1 = e_5$	
$1_{24}(p)$	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 0e_4 + e_5$	
	$e_1e_4 = \rho e_5$	$e_2 e_2 = e_4$	$e_2e_3 = (2\beta + 1)e_5$	
m 5 (0)	$e_3e_1 = -3e_4$	$e_3e_2 = -3(p+1)e_5$	$e_4 e_1 = e_5$	2
$T_{25}^{0}(\beta)$	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = 6e_4 + \beta e_5$	$e_1e_4 = -2e_5$
	$e_2 e_1 = e_5$	$e_2 e_2 = e_4$	$e_2 e_3 = -3 e_5$	
	$e_3e_1 = -3e_4$	$e_3 e_2 = 3 e_5$	$e_4 e_1 = e_5$	
T_{26}^{5}	$e_1e_1 = e_2$	$e_1e_2 = e_3$	$e_1e_3 = e_4$	
	$e_1e_4 = e_5$	$e_2 e_2 = e_5$	$e_3e_1 = -3e_5$	
T_{27}^5	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_1 e_4 = e_5$
\mathbf{T}_{28}^5	$e_1e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	
	$e_1e_4 = e_5$	$e_2 e_1 = e_5$		
$\mathbf{T}_{29}^5(\alpha)$	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_1 e_4 = e_5$
$\alpha \neq 1$	$e_2 e_1 = e_4$	$e_2 e_2 = \alpha e_5$	$e_3e_1 = 3(1-\alpha)e_5$	
$\mathbf{T}_{30}^5(\alpha)$	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1e_3 = e_4 + \alpha e_5$	
	$e_1 e_4 = e_5$	$e_2 e_1 = e_4$	$e_2 e_2 = e_5$	
$T_{31}^{5}(\beta)$	$e_1 e_1 = e_2$	$e_1e_2 = e_3 + \beta e_5$	$e_1 e_3 = e_5$	$e_2 e_1 = e_3$
01	$e_2 e_2 = e_5$	$e_3e_1 = e_4$	$e_3e_4 = e_5$	
T_{32}^5	$e_1 e_1 = e_2$	$e_1e_2 = e_3 + e_5$	$e_2 e_1 = e_3$	
	$e_3 e_1 = e_4$	$e_3e_4 = e_5$		
T_{33}^{5}	$e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_2 e_1 = e_3$	
00	$e_3 e_1 = e_4$	$e_3e_4 = e_5$		
$\mathbf{T}_{24}^5(\beta)$	$e_1 e_1 = e_2$	$e_1e_2 = \beta e_5$	$e_1e_3 = \frac{1}{2}e_4$	$e_2 e_1 = e_3$
94.	$e_2e_2 = \frac{2}{2}e_4$	$e_2 e_3 = e_5$	$e_2 e_4 = e_5$	
	$e_3e_1 = e_4$	$e_3e_3 = e_5$		
$T_{2^{t}}^{5}$	$e_1e_1 = e_2$	$e_1 e_2 = e_5$	$e_1e_3 = \frac{1}{2}e_4$	$e_2 e_1 = e_3$
30	$e_2 e_2 = \frac{2}{2} e_4$	$e_2 e_4 = e_5$	$e_3e_1 = e_4$	$e_3e_3 = e_5$
T_{22}^{5}	$e_1e_1 = e_2$	$e_1e_2 = \frac{1}{2}e_4$	$e_2e_1 = e_2$	$e_2e_2 = \frac{2}{2}e_4$
30	$e_{2}e_{4} = e_{5}$	$e_3e_1 = e_4$	$e_{3}e_{3} = e_{5}$	304
$\mathbf{T}_{-}^{5}(\beta)$	$e_2 e_4 = e_3$	$e_1e_2 = -\frac{2}{2}e_4 + e_5$	$e_3e_3 = e_3$	$e_1e_4 = e_5$
- 37(P)	$e_1e_1 = e_2$ $e_2e_2 = -\frac{1}{2}e_4 + e_5$	$e_1e_3 = 3e_4 + e_3$	$e_2e_1 = e_3$	$e_1e_4 = e_5$
$\mathbf{T}^{5}(\beta)$	$c_2 c_2 = \frac{1}{3} c_4 + c_5$	$\frac{c_2 c_3 - p c_3}{c_1 c_2 - 2 c_4}$		$e_3e_2 = e_3$
1 38(<i>P</i>)	$e_1e_1 = e_2$	$e_1e_3 = -\frac{1}{3}e_4$	$e_2e_1 = e_3$	$e_1e_4 = e_5$
$\mathbf{T}_{2}^{5}(\boldsymbol{\beta})$	$e_2 e_2 = -\frac{1}{3}e_4$	$e_2e_3 = \rho e_5$		C3C2 - C5
⊥ ₃₉ (<i>p</i>)	$e_1e_1 = e_2$	$e_1e_3 = -\frac{1}{5}e_4 + e_5$	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$
	$e_2e_2 = \frac{1}{15}e_4 + e_5$	$e_2 e_3 = \beta e_5$ 7	$e_3 e_1 = e_4$	
m 5 (a)	$e_3e_2 = -\frac{2}{5}e_5$	$e_4 e_1 = -\frac{1}{2} e_5$		
$T_{40}^{5}(\beta)$	$e_1e_1 = e_2$	$e_1e_3 = -\frac{1}{2}e_4$	$e_1 e_4 = e_5$	
	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{2}{15} e_4$	$e_2e_3 = \beta e_5$	
$e_3e_1 = e_4$	$e_3e_2 = -\frac{2}{5}e_5$	$e_4 e_1 = -\frac{i}{2} e_5$		
$\mathbf{T}_{41}^5(\alpha, \overline{\beta})$	$e_1 e_1 = e_2$	$e_1e_2 = -e_3$	$e_1e_3 = \alpha e_4 + e_5$	$e_1e_4 = \beta \overline{e_5}$
	$e_2 e_1 = e_3$	$e_2e_2 = (\alpha + \frac{2}{2})e_4 + e_5$	$e_2e_3 = \left(\frac{3\alpha+1}{2} - \frac{\beta(3\alpha+4)}{2}\right)$	e_5
		` o/	`J J /	

	$e_3e_1 = e_4$	$e_3e_2 = \left(\beta - \frac{3\alpha + 1}{2}\right)e_5$	$e_4 e_1 = e_5$	
$\mathbf{T}_{40}^5(\alpha,\beta)$	$e_1e_1 = e_2$	$e_1e_2 = -e_3$	$e_1e_3 = \alpha e_4$	$e_1e_4 = \beta e_5$
- 42(***,/*)	$e_{2}e_{1} = e_{2}$	$e_{2}e_{2} = (\alpha + \frac{2}{2})e_{4}$	$e_{2}e_{2} - (\frac{3\alpha+1}{2} - \frac{\beta(3\alpha+4)}{2})e_{2}$	-1-4 /0
	$e_2e_1 = e_3$	$e_2e_2 = (\alpha + \frac{3}{3})e_4$ $e_2e_2 = (\beta - \frac{3\alpha + 1}{2})e_5$	$e_4e_1 = e_5$	-0
$\mathbf{T}_{i\alpha}^{5}(\alpha,\beta)$	$e_1e_1 = e_2$	$e_1e_2 = -e_2 + e_5$	$e_1e_2 = \alpha e_4 + \beta e_5$	$e_1e_4 = -\frac{3\alpha+5}{2}e_5$
$= \frac{43}{43} (\alpha, \beta)$		$c_1 c_2 = (c_1 + c_3)$	$3\alpha^2 + 13\alpha + 8 \alpha^2$	4 05
$\alpha \neq -\frac{1}{3}$	$e_2e_1 = e_3$	$e_2e_2 = (\alpha + \frac{1}{3})e_4 + \beta e_5$	$e_2e_3 = \frac{4}{4}e_5$	
T ⁵	$e_3e_1 = e_4$	$e_3e_2 = -\frac{1}{4}e_5$	$e_4e_1 = e_5$	
4 44	$e_1e_1 = e_2$ $e_1e_4 = -\frac{3}{2}e_7$	$e_1e_2 = -e_3 + e_5$	$e_1e_3 = -\frac{1}{3}e_4$	
	$e_1e_4 = \frac{1}{4}e_5$	$e_2e_1 = e_3$ $e_2e_2 = -\frac{1}{2}e_5$	$e_2e_3 = \frac{1}{6}e_5$	
T ⁵ -	$e_3e_1 = e_4$	$e_1e_2 = -e_2$	$e_4e_1 = e_5$ $e_1e_2 = -\frac{2}{2}e_4 + e_5$	
- 45	$e_1e_4 = e_5$	$e_2e_1 = e_3$	$e_2e_2 = e_5$	
	$e_2e_3 = -\frac{2}{2}e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
$\mathbf{T}_{4e}^{5}(\lambda,\beta)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_5$	$e_1e_3 = \frac{4\lambda - 1}{2}e_4 + \beta e_5$	$e_1e_4 = \frac{4\lambda - 1}{5}e_5$
10(177)		$2\lambda = \frac{(2\lambda+1)(\lambda+2)}{(\lambda+2)} + \beta = 0$	$2(2\lambda+1)(3\lambda^2+1) = 2(2\lambda+1)(3\lambda^2+1) = 2(2\lambda+1)(3\lambda^2+1)(3\lambda^2+1) = 2(2\lambda+1)(3\lambda^2+1)(3\lambda^2+1) = 2(2\lambda+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1) = 2(2\lambda+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1)(3\lambda^2+1$	$5-2\lambda$
$\lambda \neq 1, -2, -\frac{1}{2}, \frac{1}{2}$	$e_2e_1 - e_3$	$e_2e_2 = \frac{1}{3(5-2\lambda)}e_4 + \rho e_5$	$e_2e_3 = \frac{1}{3\lambda(5-2\lambda)^2}e_5$	
	$e_3e_1 = e_4$	$e_3e_2 = \frac{2\lambda + 4}{5 - 2\lambda}e_5$	$e_4e_1 = e_5$	
$\mathbf{T}_{47}^{\scriptscriptstyle 5}(\lambda,eta)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1e_3 = \frac{4\lambda - 1}{5 - 2\lambda}e_4 + e_5$	$e_1e_4 = \beta e_5$
$\lambda \neq 1, \frac{5}{2}$	$e_2 e_1 = e_3$	$e_2e_2 = \frac{(2\lambda+1)(\lambda+2)}{3(5-2\lambda)}e_4 + e_5$	$e_2e_3 = \frac{\beta(4\lambda^2 - 2\lambda + 7) + 2(\lambda - 3\lambda)}{3\lambda(5 - 2\lambda)}$	$\frac{1)^2}{2}e_5$
	$e_3e_1 = e_4$	$e_3e_2 = \left(\beta - \frac{2(\lambda - 1)}{5}\right)e_5$	$e_4e_1 = e_5$	
$T_{48}^5(\lambda,\beta)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1e_3 = \frac{4\lambda - 1}{5\lambda}e_4$	$e_1e_4 = \beta e_5$
$\lambda \neq 0^{\frac{5}{2}}$	$e_0e_1 = e_0$	$e_{2}e_{2} = \frac{(2\lambda+1)(\lambda+2)}{(2\lambda+1)(\lambda+2)}e_{4}$	$e_{0}e_{0} = \frac{\beta(4\lambda^{2} - 2\lambda + 7) + 2(\lambda - 1)}{\beta(4\lambda^{2} - 2\lambda + 7) + 2(\lambda - 1)}$	<u>1)²</u> er
$\pi \neq 0, 2$	$c_2 c_1 = c_3$	$c_2 c_2 = \frac{3(5-2\lambda)}{3(5-2\lambda)} c_4$	$c_2 c_3 = 3\lambda(5-2\lambda)$	65
T ⁵	$e_3e_1 = e_4$	$\frac{c_3c_2}{c_1c_2} = \left(p - \frac{c_2}{5-2\lambda}\right)c_5$	$e_4e_1 = e_5$	0101 - 07
1 49	$e_1e_1 = e_2$	$e_1e_2 = e_3 + e_5$	$e_1e_3 = e_4$	$e_1e_4 - e_5$
	$e_2e_1 = e_3$ $e_2e_1 = e_4$	$e_2e_2 = e_4$ $e_2e_2 = e_5$	$e_2e_3 = e_5$	
T_{20}^5	$e_3e_1 = e_4$ $e_1e_1 = e_2$	$e_{1}e_{2} = e_{3}$	$e_4e_1 = e_3$ $e_1e_3 = e_4 + e_5$	$e_1e_4 = e_5$
- 50	$e_2e_1 = e_3$	$e_2e_2 = e_4 + e_5$	$e_2e_3 = e_5$	-1-4 -5
	$e_3e_1 = e_4$	$e_3e_2 = e_5$	$e_4e_1 = e_5$	
$\mathbf{T}_{51}^5(eta,\gamma)$	$e_1e_1 = e_2$	$e_1e_2 = -2e_3 + e_5$	$e_1e_3 = -e_4 + \beta e_5$	$e_1e_4 = \gamma e_5$
	$e_2 e_1 = e_3$	$e_2e_2 = \beta e_5$	$e_2e_3 = -\frac{3\gamma + 2}{6}e_5$	$e_3e_1 = e_4$
	$e_3e_2 = \frac{3\gamma + 2}{3}e_5$	$e_4 e_1 = e_5$	-	
$\mathbf{T}_{52}^5(eta)$	$e_1e_1 = e_2$	$e_1e_2 = -2e_3 + \beta e_5$	$e_1e_3 = -e_4 + e_5$	
	$e_1 e_4 = e_5$	$e_2e_1 = e_3$	$e_2e_2 = e_5$	
	$e_2e_3 = -\frac{1}{2}e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$,
\mathbf{T}_{53}^5	$e_1e_1 = e_2$	$e_1 e_2 = -\frac{1}{2} e_3$	$e_1 e_3 = -\frac{1}{2}e_4 + e_5$	$e_1 e_4 = -\frac{1}{2} e_5$
5	$e_2 e_1 = e_3$	$e_2 e_2 = e_5$	$e_3e_1 = e_4$	$e_4e_1 = e_5$
T_{54}^{5}	$e_1e_1 = e_2$	$e_1e_2 = -\frac{1}{2}e_3$	$e_1e_3 = -\frac{1}{2}e_4 + e_5$	
	$e_1 e_4 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = e_5$	
\mathbf{T} $(1, \mathbf{n})$	$e_2 e_3 = -e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
$1_{55}(\lambda,\alpha)_{\lambda\neq-2,0}$	$e_1 e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1e_3 = \alpha e_4 + e_5$	
$\alpha \neq \frac{2n+6n+1}{6}$	$e_1e_4 = e_5$	$e_2e_1 = e_3$	$e_2 e_2 = (\alpha + \frac{1}{3})e_4 + e_5$	
	$e_2 e_3 = \frac{3\alpha - 2(\lambda - 1)}{3\lambda} e_5$	$e_3e_1 = e_4$	$e_3 e_2 = e_5$	
$\mathbf{T}_{56}^{3}(\lambda, \alpha)$	$e_1 e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1e_3 = \alpha e_4$	
$\lambda \neq 0$	$e_1e_4 = e_5$ $3\alpha - 2(\lambda - 1)$	$e_2e_1 = e_3$	$e_2e_2 = \left(\alpha + \frac{-1}{3}\right)e_4$	
\mathbf{m}_{2}	$e_2e_3 = \frac{1}{3\lambda}e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
⊥ ₅₇ (<i>P</i>)	$e_1e_1 = e_2$	$e_1e_2 = -2e_3$	$e_1e_3 = -\frac{1}{2}e_4 + e_5$	
	$e_1e_4 - e_5$ $e_2e_2\frac{3}{2}e_7$	$c_2c_1 - c_3$	$e_2e_2 = \frac{1}{2}e_4 + e_5$	
$\mathbf{T}_{2}^{5}(\alpha)$	$c_2 c_3 = -\frac{1}{4} c_5$	$e_3e_1 - e_4$ $e_1e_2 = -2e_2 + e_7$	$c_3c_2 = c_0$ $e_1e_2 = 0e_4$	
- 58(4)	$e_1e_4 = e_5$	$e_1e_2 = 2e_3 + e_5$	$e_1e_3 = ae_4$ $e_2e_2 = (\alpha + 1)e_4$	
	$e_3e_1 = e_4$	$e_2e_3 = -\frac{\alpha+2}{2}e_5$	$e_3e_2 = e_5$	
$\mathbf{T}_{50}^5(\beta)$	$e_1e_1 = e_2$	$e_1e_2 = e_3 + e_4$	$e_1e_3 = e_4$	$e_1e_4 = \beta e_5$
99.4.7	$e_2e_1 = e_3$	$e_2e_2 = e_4 - e_5$	$e_2e_3 = \beta e_5$	

	$e_3e_1 = e_4$	$e_3e_2 = \beta e_5$	$e_4 e_1 = 3 e_5$	
$\mathbf{T}_{60}^5(\beta)_{\beta\neq 0}$	$e_1e_1 = e_2$	$e_1e_2 = e_3 + e_4$	$e_1 e_3 = e_4$	$e_1 e_4 = 3 e_5$
00007773	$e_2 e_1 = e_3$	$e_2 e_2 = e_4 - e_5$	$e_2 e_3 = 3 e_5$	$e_3e_1 = e_4 + \beta e_5$
	$e_3e_2 = 3e_5$	$e_4e_1 = 3e_5$		
$\mathbf{T}_{61}^5(\beta)_{\beta\neq 0}$	$e_1 e_1 = e_2$	$e_1e_2 = e_3 + e_4 + \beta e_5$	$e_1 e_3 = e_4$	$e_1 e_4 = 3 e_5$
01 (), / / -	$e_2 e_1 = e_3$	$e_2 e_2 = e_4 - e_5$	$e_2 e_3 = 3 e_5$	
	$e_3e_1 = e_4 - 6e_5$	$e_3e_2 = 3e_5$	$e_4e_1 = 3e_5$	
$\mathbf{T}_{62}^5(\beta)_{\beta\neq 0}$	$e_1 e_1 = e_2$	$e_1e_2 = -e_3 + e_4$	$e_1e_3 = -\frac{2}{2}e_4 + \beta e_5$	
02 (), / / -	$e_1e_4 = 3e_5$	$e_2 e_1 = e_3$	$e_2e_2 = \beta e_5$	
	$e_2e_3 = -2e_5$	$e_3e_1 = e_4$	$e_3e_2 = 3e_5$	
$\mathbf{T}_{63}^5(\beta)_{\beta\neq 0}$	$e_1 e_1 = e_2$	$e_1e_2 = -e_3 + e_4 + \beta e_5$	$e_1e_3 = -\frac{2}{2}e_4 - e_5$	$e_1 e_4 = 9 e_5$
00 () / / /	$e_2 e_1 = e_3$	$e_2 e_2 = 3 e_5$	$e_2 e_3 = -2 e_5$	
	$e_3 e_1 = e_4$	$e_3e_2 = 3e_5$	$e_4 e_1 = -12 e_5$	
$\mathbf{T}_{64}^5(eta,\gamma)$	$e_1 e_1 = e_2$	$e_1e_2 = -e_3 + e_4$	$e_1e_3 = -\frac{2}{2}e_4 + (\beta + 2)e_5$	$e_1e_4 = 3(\gamma - 1)e_5$
010	$e_2 e_1 = e_3$	$e_2e_2 = \beta e_5$	$e_2e_3 = -2\gamma e_5$	
	$e_3e_1 = e_4$	$e_3e_2 = 3\gamma e_5$	$e_4 e_1 = 6 e_5$	
$\mathbf{T}_{65}^5(\beta)_{\beta\neq 0}$	$e_1 e_1 = e_2$	$e_1e_2 = -\frac{1}{2}e_3 + e_4$	$e_1e_3 = -\frac{1}{2}e_4 + \beta e_5$	
03 () / / ($e_1 e_4 = e_5$	$e_2e_1 = e_3^2$	$e_2e_2 = \beta e_5^2$	
	$e_2e_3 = -e_5$	$e_3e_1 = e_4$	$e_3e_2 = e_5$	
$\mathbf{T}_{66}^5(\beta)_{\beta\neq 0}$	$e_1e_1 = e_2$	$e_1e_2 = -\frac{1}{2}e_3 + e_4 + \beta e_5$	$e_1e_3 = -\frac{1}{2}e_4 + 3e_5$	$e_1e_4 = -3e_5$
000 777 0	$e_2 e_1 = e_3$	$e_2 e_2 = e_5^2$	$e_3e_1 = e_4^2$	$e_4 e_1 = 6 e_5$
$\mathbf{T}_{67}^5(\beta,\gamma)$	$e_1e_1 = e_2$	$e_1e_2 = -\frac{1}{2}e_3 + e_4$	$e_1e_3 = -\frac{1}{2}e_4 + (\beta + 2)e_5$	$e_1e_4 = (\gamma - 3)e_5$
01 (0 / //	$e_2 e_1 = e_3$	$e_2e_2 = \beta e_5^2$	$e_2 e_3 = -\gamma e_5$	$e_3e_1 = e_4$
	$e_3e_2 = \gamma e_5$	$e_4 e_1 = 6 e_5$		
$\mathbf{T}_{68}^5(\beta)_{\beta\neq 0}$	$e_1 e_1 = e_2$	$e_1e_2 = -2e_3 + e_4 + \beta e_5$	$e_1e_3 = -\frac{1}{2}e_4$	
00 (), / / -	$e_1 e_4 = 4 e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = \frac{1}{2} e_4^2$	
	$e_2 e_3 = -3e_5$	$e_3 e_1 = e_4$	$e_3e_2 = \tilde{4}e_5$	
$\mathbf{T}_{60}^5(\lambda)$	$e_1 e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_4$	$e_1e_3 = \frac{2\lambda^2 + 5\lambda - 1}{\epsilon}e_4$	
03()	$e_1e_4 = 6\lambda e_5$	$e_2e_1 = e_3$	$e_2e_2 = \frac{(2\lambda+1)(\lambda+1)}{2}e_4$	
	$e_2e_3 = (2\lambda^2 + \lambda + 3)e_5$	$e_3e_1 = e_4$	$e_3e_2 = 6\lambda e_5$	
$\mathbf{T}_{\mathbf{r}\alpha}^{5}(\lambda,\alpha)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + \alpha e_5$	$e_1e_3 = e_4$	$e_1e_4 = \lambda e_5$
70(00)	$e_2e_1 = e_3$	$e_2e_2 = e_4 + (1 - \lambda)e_5$	$e_2e_3 = e_5$	$e_3e_1 = 3e_5$
$\mathbf{T}_{71}^5(\lambda)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3 + e_5$	$e_1e_3 = e_4$	$e_1e_4 = \lambda e_5$
/1 ()	$e_2e_1 = e_3$	$e_2e_2 = e_4$	$e_2e_3 = e_5$	11 0
$\mathbf{T}_{72}^5(\lambda)$	$e_1e_1 = e_2$	$e_1e_2 = \lambda e_3$	$e_1 e_3 = e_4$	$e_1e_4 = \lambda e_5$
12 ()	$e_2 e_1 = e_3$	$e_2 e_2 = e_4$	$e_2 e_3 = e_5$	
$\mathbf{T}_{73}^5(\alpha,\beta)$	$e_1e_1 = e_2$	$e_1e_2 = -e_3 + e_5$	$e_1 e_3 = e_4$	$e_1e_4 = \alpha e_5$
10($e_2 e_1 = e_3$	$e_2e_2 = e_4 + 2\beta e_5$	$e_2 e_3 = (2 - \alpha) e_5$	$e_3e_1 = 3\beta e_5$
	$e_3e_2 = -3e_5$	$e_4 e_1 = e_5$		
$\mathbf{T}_{74}^5(\alpha)$	$e_1 e_1 = e_2$	$e_1 e_2 = -e_3$	$e_1 e_3 = e_4$	$e_1e_4 = \alpha e_5$
	$e_2 e_1 = e_3$	$e_2e_2 = e_4 + 2e_5$	$e_2 e_3 = (2 - \alpha) e_5$	
	$e_3e_1 = 3e_5$	$e_3e_2 = -3e_5$	$e_4 e_1 = e_5$	
$\mathbf{T}_{75}^5(lpha)$	$e_1e_1 = e_2$	$e_1e_2 = -e_3$	$e_1 e_3 = e_4$	
	$e_1e_4 = \alpha e_5$	$e_2e_1 = e_3$	$e_2 e_2 = e_4$	
	$e_2 e_3 = (2 - \alpha) e_5$	$e_3e_2 = -3e_5$	$e_4 e_1 = e_5$	
$\mathbf{T}_{76}^5(\alpha, \overline{\beta})$	$e_1e_1 = e_2$	$e_1 e_2 = e_5$	$e_1e_3 = \alpha e_5$	$e_2e_1 = e_3$
	$e_2 e_2 = (\alpha + \beta) e_5$	$e_2 e_3 = e_4$	$e_2 e_4 = e_5$	$e_3e_1 = 3\beta e_5$
$\mathbf{T}_{77}^5(\alpha)$	$e_1e_1 = e_2$	$e_1 e_3 = e_5$	$e_2 e_1 = e_3$	$e_2 e_2 = (\alpha + 1)e_5$
5	$e_2 e_3 = e_4$	$e_2 e_4 = e_5$	$e_3e_1 = 3\alpha e_5$	
\mathbf{T}_{78}^5	$e_1e_1 = e_2$	$e_2e_1 = e_3$	$e_2e_2 = e_5$	
	$e_2 e_3 = e_4$	$e_2e_4 = e_5$	$e_3e_1 = 3e_5$	
T^{5}	$e_1e_1 = e_2$	$e_{2}e_{1} = e_{2}$	$e_{2}e_{2} = e_{4}$	$e_{2}e_{4} = e_{5}$

REFERENCES

- [1] Abdelwahab H., Calderón A.J., Kaygorodov I., The algebraic and geometric classification of nilpotent binary Lie algebras, International Journal of Algebra and Computation, 29 (2019), 6, 1113–1129.
- [2] Adashev J., Camacho L., Omirov B., Central extensions of null-filiform and naturally graded filiform non-Lie Leibniz algebras, Journal of Algebra, 479 (2017), 461–486.
- [3] Calderón Martín A., Fernández Ouaridi A., Kaygorodov I., The classification of 2-dimensional rigid algebras, Linear and Multilinear Algebra, 68 (2020), 4, 828–844.
- [4] Calderón Martín A., Fernández Ouaridi A., Kaygorodov I., On the classification of bilinear maps with radical of a fixed codimension, Linear and Multilinear Algebra, 2020, to appear, arXiv:1806.07009
- [5] Camacho L., Karimjanov I., Kaygorodov I., Khudoyberdiyev A., One-generated nilpotent Novikov algebras, Linear and Multilinear Algebra, 2020, DOI: 10.1080/03081087.2020.1725411
- [6] Camacho L., Kaygorodov I., Lopatkin V., Salim M., The variety of dual Mock-Lie algebras, Communications in Mathematics, to appear, arXiv:1910.01484
- [7] Cantarini N., Kac V., Classification of linearly compact simple rigid superalgebras, International Mathematics Research Notices, 17 (2010), 3341–3393.
- [8] Cicalò S., De Graaf W., Schneider C., Six-dimensional nilpotent Lie algebras, Linear Algebra and its Applications, 436 (2012), 1, 163–189.
- [9] Darijani I., Usefi H., The classification of 5-dimensional *p*-nilpotent restricted Lie algebras over perfect fields, I., Journal of Algebra, 464 (2016), 97–140.
- [10] De Graaf W., Classification of nilpotent associative algebras of small dimension, International Journal of Algebra and Computation, 28 (2018), 1, 133–161
- [11] De Graaf W., Classification of 6-dimensional nilpotent Lie algebras over fields of characteristic not 2, Journal of Algebra, 309 (2007), 2, 640–653.
- [12] Dekimpe K., Ongenae V., Filiform left-symmetric algebras, Geometriae Dedicata, 74 (1999), no. 2, 165–199.
- [13] Fernández Ouaridi A., Kaygorodov I., Khrypchenko M., Volkov Yu., Degenerations of nilpotent algebras, arXiv:1905.05361
- [14] Gorshkov I., Kaygorodov I., Khrypchenko M., The algebraic classification of nilpotent Tortkara algebras, Communications in Algebra, 48 (2020), 8, 3608–3623.
- [15] Hegazi A., Abdelwahab H., The classification of *n*-dimensional non-associative Jordan algebras with (n 3)-dimensional annihilator, Communications in Algebra, 46 (2018), 2, 629–643.
- [16] Hegazi A., Abdelwahab H., Classification of five-dimensional nilpotent Jordan algebras, Linear Algebra and its Applications, 494 (2016), 165–218.
- [17] Hegazi A., Abdelwahab H., Calderón Martín A., The classification of *n*-dimensional non-Lie Malcev algebras with (n-4)-dimensional annihilator, Linear Algebra and its Applications, 505 (2016), 32–56.
- [18] Hegazi A., Abdelwahab H., Calderón Martín A., Classification of nilpotent Malcev algebras of small dimensions over arbitrary fields of characteristic not 2, Algebras and Representation Theory, 21 (2018), 1, 19–45.
- [19] Jacobson N., Structure and representations of Jordan algebras, American Mathematical Society Colloquium Publications, Vol. XXXIX American Mathematical Society, Providence, R.I. 1968 x+453 pp.
- [20] Ismailov N., Kaygorodov I., Mashurov F., The algebraic and geometric classification of nilpotent assosymmetric algebras, Algebras and Representation Theory, 2020, DOI: 10.1007/s10468-019-09935-y
- [21] Ismailov N., Kaygorodov I., Volkov Yu., The geometric classification of Leibniz algebras, International Journal of Mathematics, 29 (2018), 5, 1850035.
- [22] Jumaniyozov D., Kaygorodov I., Khudoyberdiyev A., The algebraic and geometric classification of nilpotent noncommutative Jordan algebras, Journal of Algebra and its Applications, 2020, DOI:10.1142/S0219498821502029
- [23] Kantor I., Certain generalizations of Jordan algebras [in Russian], Trudy Sem. Vektor. Tenzor. Anal., 16 (1972), 407–499.
- [24] Kantor I., On an extension of a class of Jordan algebras, Algebra and Logic, 28 (1989), 2, 117–121.
- [25] Kantor I., An universal conservative algebra, Siberian Mathematical Journal, 31 (1990), 3, 388–395.

- [26] Karimjanov I., Kaygorodov I., Khudoyberdiyev A., The algebraic and geometric classification of nilpotent Novikov algebras, Journal of Geometry and Physics, 143 (2019), 11–21.
- [27] Kaygorodov I., Khrypchenko M., Lopes S., The algebraic and geometric classification of nilpotent anticommutative algebras, Journal of Pure and Applied Algebra, 224 (2020), 8, 106337
- [28] Kaygorodov I., Khrypchenko M., Popov Yu., The algebraic and geometric classification of nilpotent terminal algebras, arXiv:1909.00358
- [29] Kaygorodov I., Lopatin A., Popov Yu., Conservative algebras of 2-dimensional algebras, Linear Algebra and its Applications, 486 (2015), 255–274.
- [30] Kaygorodov I., Lopes S., Páez-Guillán P., Non-associative central extensions of null-filiform associative algebras, Journal of Algebra, 560 (2020), 1190–1210.
- [31] Kaygorodov I., Páez-Guillán P., Voronin V., The algebraic and geometric classification of nilpotent bicommutative algebras, Algebras and Representation Theory, 2020, DOI: 10.1007/s10468-019-09944-x
- [32] Kaygorodov I., Popov Yu., Pozhidaev A., Volkov Yu., Degenerations of Zinbiel and nilpotent Leibniz algebras, Linear and Multilinear Algebra, 66 (2018), 4, 704–716.
- [33] Kaygorodov I., Popov Yu., Pozhidaev A., The universal conservative superalgebra, Communications in Algebra, 47 (2019), 10, 4064–4074.
- [34] Kaygorodov I., Rakhimov I., Said Husain Sh. K., The algebraic classification of nilpotent associative commutative algebras, Journal of Algebra and its Applications, 2019, DOI: 10.1142/S0219498820502205
- [35] Kaygorodov I., Volkov Yu., Conservative algebras of 2-dimensional algebras. II, Communications in Algebra, 45 (2017), 8, 3413–3421.
- [36] Kaygorodov I., Volkov Yu., The variety of 2-dimensional algebras over an algebraically closed field, Canadian Journal of Mathematics, 71 (2019), 4, 819–842.
- [37] Kaygorodov I., Volkov Yu., Complete classification of algebras of level two, Moscow Mathematical Journal, 19 (2019), 3, 485–521.
- [38] Masutova K., Omirov B., On some zero-filiform algebras, Ukrainian Mathematical Journal, 66 (2014), 4, 541–552.
- [39] Popov Yu., Conservative algebras and superalgebras: a survey, Communications in Mathematics, to appear.
- [40] Skjelbred T., Sund T., Sur la classification des algebres de Lie nilpotentes, C. R. Acad. Sci. Paris Ser. A-B, 286 (1978), 5, A241–A242.