

EXISTENCE AND LOCAL UNIQUENESS OF NORMALIZED PEAK SOLUTIONS FOR A SCHRÖDINGER-NEWTON SYSTEM

QING GUO, PENG LUO, CHUNHUA WANG AND JING YANG

ABSTRACT. In this paper, we investigate the existence and local uniqueness of normalized peak solutions for a Schrödinger-Newton system under the assumption that the trapping potential is degenerate and has non-isolated critical points.

First we investigate the existence and local uniqueness of normalized single-peak solutions for the Schrödinger-Newton system. Precisely, we give the precise description of the chemical potential μ and the attractive interaction a . Then we apply the finite dimensional reduction method to obtain the existence of single-peak solutions. Furthermore, using various local Pohozaev identities, blow-up analysis and the maximum principle, we prove the local uniqueness of single-peak solutions by precise analysis of the concentrated points and the Lagrange multiplier. Finally, we also prove the nonexistence of multi-peak solutions for the Schrödinger-Newton system, which is markedly different from the corresponding Schrödinger equation. The nonlocal term results in this difference.

The main difficulties come from the estimates on Lagrange multiplier, the different degenerate rates along different directions at the critical point of $P(x)$ and some complicated estimates involved by the nonlocal term. To our best knowledge, it may be the first time to study the existence and local uniqueness of solutions with prescribed L^2 -norm for the Schrödinger-Newton system.

Keywords: Normalized solutions; the Schrödinger-Newton system; Degenerated trapping potential

2010 Mathematics Subject Classification: 35A01 · 35B25 · 35J20 · 35J60

1. INTRODUCTION AND OUR MAIN RESULTS

In this paper, we investigate the following nonlinear Schrödinger-Newton system

$$\begin{cases} -\Delta u + P(x)u = a\psi u + \mu u, & x \in \mathbb{R}^3, \\ -\Delta \psi = \frac{u^2}{2}, & x \in \mathbb{R}^3, \end{cases} \quad (1.1)$$

under the mass constraint $\int_{\mathbb{R}^3} u^2(x)dx = 1$, where $P(x)$ is a degenerate trapping potential with non-isolated critical points, $\mu \in \mathbb{R}$ and $a \geq 0$ denote the chemical potential and the attractive interaction respectively.

The Schrödinger-Newton problem arises from describing the quantum mechanics of a polaron at rest, see [28]. Also it was used to describe an electron trapped in its own hole in a certain approximating to Hartree-Fock theory of one component plasma in [19]. In addition, Penrose in [28] applied it as a model of self-gravitating matter, where quantum state reduction is understood as a gravitational phenomenon. Specifically, if m is the mass of the point, the interaction leads

to the system in \mathbb{R}^3

$$\begin{cases} -\frac{\epsilon^2}{2m}\Delta u + P(x)u = \psi u, & x \in \mathbb{R}^3, \\ -\Delta\psi = 4\pi\tau u^2, & x \in \mathbb{R}^3, \end{cases} \quad (1.2)$$

where u is the wave function, ψ is the gravitational potential energy, $P(x)$ is a given Schrödinger potential, ϵ is the Planck constant, $\tau = Gm^2$ and G is the Newton's constant of gravitation.

Set

$$u(x) \mapsto \frac{u}{4\epsilon\sqrt{\pi\tau m}}, \quad P(x) \mapsto \frac{1}{2m}P(x), \quad \psi(x) \mapsto \frac{1}{2m}\psi(x).$$

Then system (1.2) can be written, maintaining the original notations, as

$$\begin{cases} -\epsilon^2\Delta u + P(x)u = \psi u, & x \in \mathbb{R}^3, \\ -\epsilon^2\Delta\psi = \frac{u^2}{2}, & x \in \mathbb{R}^3. \end{cases} \quad (1.3)$$

From the second equation of (1.3), we know $\psi(x) = \frac{1}{8\pi\epsilon^2} \left(\int_{\mathbb{R}^3} \frac{u^2(y)}{|x-y|} dy \right)$. Then the system (1.3) turns into the following single nonlocal equation

$$-\epsilon^2\Delta u + P(x)u = \frac{1}{8\pi\epsilon^2} \left(\int_{\mathbb{R}^3} \frac{u^2(y)}{|x-y|} dy \right) u, \quad x \in \mathbb{R}^3, \quad (1.4)$$

which also appears in the study of standing waves for the following nonlinear Hartree equations

$$i\epsilon \frac{\partial\varphi}{\partial t} = -\epsilon^2\Delta_x\varphi + (P(x) + E)\varphi - \frac{1}{8\pi\epsilon^2} \left(\int_{\mathbb{R}^3} \frac{\varphi^2(y)}{|x-y|} dy \right) \varphi, \quad (x, t) \in \mathbb{R}^3 \times \mathbb{R}^+,$$

with the form $\varphi(x, t) = e^{-iEt/\epsilon} u(x)$, where i is the imaginary unit and ϵ is the Planck constant.

In recent decades, problem (1.4) has been extensively investigated. When $\epsilon = 1$ and $P(x) = 1$, (1.4) becomes

$$-\Delta u + u = \frac{1}{8\pi} \left(\int_{\mathbb{R}^3} \frac{u^2(y)}{|x-y|} dy \right) u, \quad x \in \mathbb{R}^3. \quad (1.5)$$

The existence and uniqueness of ground states for (1.5) was obtained with variational methods by Lieb [19], Lions [21] and Menzala [25]. The nondegeneracy of the ground states for (1.5) was proved by Tod-Moroz [31] and Wei-Winter [34].

When ϵ is small and $P(x)$ is not a constant, in [22] Lions proved the existence of solutions with ground states for (1.4) under some conditions on $P(x)$ since problem (1.4) has a variational structure. Moreover, the solution with ground states concentrates at certain point. Applying the finite dimensional reduction method, in [34] Wei and Winter proved that (1.4) has a solution concentrating at k points which are the local minimum points of $P(x)$. This also means the existence of multiple solutions. Very recently, in [23] Luo, Peng and Wang proved the uniqueness of the concentrated solutions of (1.4) obtained in [34] by some local Pohozaev identities, blow-up analysis and the maximum principle. For more other results of the existence of solutions with concentration, one can refer to [5, 8, 30, 32] and the references therein. For a very similar nonlocal problem i.e. Schrödinger-Poisson equations, one can refer to [14, 15, 33], where some existence of peak solutions are obtained by the finite dimensional reduction method under various assumptions of the potential function.

Actually, the Schrödinger-Newton problem (1.4) is a special type of following Choquard equation:

$$-\epsilon^2 \Delta u + P(x)u = \frac{a}{\epsilon^2} (I_\alpha * |u|^p) |u|^{p-2} u + \mu u, \quad N \geq 2, \quad p > 1, \quad x \in \mathbb{R}^N, \quad (1.6)$$

where $\mu \in \mathbb{R}$ denotes the chemical potential, $a \geq 0$ denotes the attractive interaction and the Riesz potential $I_\alpha : \mathbb{R}^N \rightarrow \mathbb{R}$ is defined (cf. Ref [29]) as

$$I_\alpha := \frac{\Gamma(\frac{N-\alpha}{2})}{\Gamma(\frac{\alpha}{2}) \pi^{\frac{N}{2}} 2^\alpha} \frac{1}{|x|^{N-\alpha}}, \quad x \in \mathbb{R}^N \setminus \{0\}, \quad \alpha \in (0, N). \quad (1.7)$$

Especially, when $N = 3$ and $\alpha = 2$ in (1.7), then $I_\alpha = \frac{1}{4\pi} \frac{1}{|x|}$, $x \in \mathbb{R}^3 \setminus \{0\}$.

When $\epsilon = 1, N = 3, \alpha = 2, P(x) = \text{constant} > 0, \mu = 0$ and $p \rightarrow 2$, in [35] Xiang proved the uniqueness and non-degeneracy of ground states to (1.14). When $\epsilon = 1, N = 3, 4, 5, P(x) = 1, \alpha = 2$ and $\mu = 0$, In [4] Chen proved the nondegeneracy of ground states of (1.14). As an application of the non-degeneracy result he obtained, he then used a Lyapunov-Schmidt reduction argument to construct multiple semi-classical solutions to (1.14) with an external potential. When $N = 3, \alpha = 2$ and $\mu = 0$ in (1.14), in [26, 27] Moroz obtained some results about the existence and concentration of positive solutions to the Choquard equation. For ground states of (1.14) with $\epsilon = 1$, one can refer to [6, 12, 16–18, 36] under various assumptions of trapping potential $P(x)$, which can be described equivalently by positive L^2 minimizers of the following Hartree-type energy functional (c.f. in [12])

$$E_a(u) = \int_{\mathbb{R}^N} (|\nabla u|^2 + P(x)u^2) dx - \frac{a}{p} \int_{\mathbb{R}^N} (I_\alpha * |u(x)|^p) |u(x)|^p dx.$$

To our best knowledge, up to now there are no results about the existence and the local uniqueness of peak solutions to (1.8), that is,

$$-\Delta u + P(x)u = \frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{u^2(y)}{|x-y|} dy u + \mu u, \quad \text{in } \mathbb{R}^3, \quad (1.8)$$

where $P(x)$ denotes a class of degenerate trapping potential which has non-isolated critical points. So in this paper, we are aimed to investigate these problems. Precisely, we study the existence and the local uniqueness of peak solutions for (1.1) with the following L^2 constraint

$$\int_{\mathbb{R}^3} u^2 = 1. \quad (1.9)$$

We first recall the existence result for the ground state. Denote by $U(x)$ the unique positive solution of

$$-\Delta u + u = \int_{\mathbb{R}^3} \frac{u^2(y)}{|x-y|} dy u(x), \quad u \in H^1(\mathbb{R}^3). \quad (1.10)$$

Form [20, 34], we know that $U(0) = \max_{x \in \mathbb{R}^3} U(x)$ and the solution $U(x)$ is strictly decreasing and

$$\lim_{|x| \rightarrow \infty} U(x) e^{|x|} = \lambda_0 > 0, \quad \lim_{|x| \rightarrow \infty} \frac{U'(x)}{U(x)} = -1,$$

for some constant $\lambda_0 > 0$. Moreover, if $\phi(x) \in H^1(\mathbb{R}^3)$ solves the linearized equation

$$-\Delta \phi(x) + \phi(x) = \frac{1}{8\pi} \left(\int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy \right) \phi(x) + \frac{1}{4\pi} \left(\int_{\mathbb{R}^3} \frac{U(y)\phi(y)}{|x-y|} dy \right) U(x),$$

then $\phi(x)$ is a linear combination of $\frac{\partial U}{\partial x_j}$, $j = 1, 2, 3$.

In this paper, we let $a_* = \int_{\mathbb{R}^3} U^2$. First of all, we study the following problem without constraint:

$$\begin{cases} -\Delta w + (\lambda + P(x))w = \int_{\mathbb{R}^3} \frac{w^2(y)}{|x-y|} dy w, & \text{in } \mathbb{R}^3; \\ w \in H^1(\mathbb{R}^3), \end{cases} \quad (1.11)$$

where $\lambda > 0$ is a large parameter. It is well known that for large $\lambda > 0$, we can construct various positive solutions concentrating at some stable critical points of $P(x)$. Particularly, we can construct positive single-peak solutions for (1.11) in the sense that

$$w_\lambda(x) = \lambda \left(U(\sqrt{\lambda}(x - x_\lambda)) + \varpi_\lambda(x) \right),$$

with $\int_{\mathbb{R}^3} \left[\frac{1}{\lambda} |\nabla \varpi_\lambda|^2 + \varpi_\lambda^2 \right] = o(\lambda^{-\frac{3}{2}})$. Let $u_\lambda = \frac{w_\lambda}{\left(\int_{\mathbb{R}^3} w_\lambda^2 \right)^{1/2}}$. Then $\int_{\mathbb{R}^3} u_\lambda^2 = 1$, and

$$\begin{cases} -\Delta u_\lambda + (\lambda + P(x))u_\lambda = a_\lambda \int_{\mathbb{R}^3} \frac{u_\lambda^2(y)}{|x-y|} dy u_\lambda(x), & \text{in } \mathbb{R}^3; \\ u_\lambda \in H^1(\mathbb{R}^3), \end{cases}$$

with

$$a_\lambda = \int_{\mathbb{R}^3} w_\lambda^2 = \lambda^{2-\frac{3}{2}} \left(\int_{\mathbb{R}^3} U^2 + o(1) \right) = \lambda^{\frac{1}{2}} (a_* + o(1)).$$

Note that $a_\lambda > 0$, and as $\lambda \rightarrow +\infty$, so $a_\lambda \rightarrow +\infty$. Therefore, we obtain a concentrated solution with a single-peak for (1.8)–(1.9) with $\mu = -\lambda$ and suitable a_λ . Now the crucial question is for any $a > 0$ large, whether we can choose a suitable large $\lambda_a > 0$, such that (1.8)–(1.9) holds with

$$\mu = -\lambda_a, \quad u_a = \frac{w_{\lambda_a}}{\left(\int_{\mathbb{R}^3} w_{\lambda_a}^2 \right)^{1/2}}.$$

The above discussions show that the existence of concentrated solutions for (1.8)–(1.9) is closely related to the existence of peaked solutions for the nonlinear Schrödinger equation (1.11). In this paper, we will mainly investigate concentrated solutions u_a of (1.8)–(1.9) in the sense that

$$\max_{x \in B_\vartheta(b_0)} u_a(x) \rightarrow +\infty, \quad \text{while } u_a(x) \rightarrow 0 \text{ uniformly in } \mathbb{R}^3 \setminus B_\vartheta(b_0), \text{ for any } \vartheta > 0,$$

as $a \rightarrow +\infty$, where b_0 is a point in \mathbb{R}^3 . Here, we are concerned with the following three aspects: possible values for μ_a ; the exact location of the concentrated points of the concentrated solutions for (1.8)–(1.9); the existence and the local uniqueness of the concentrated solutions for (1.8)–(1.9).

The first result of our paper is as follows.

Theorem 1.1. *Suppose that u_a is a solution of (1.8)–(1.9) concentrated at some points as $a \rightarrow +\infty$. Then it holds*

$$\mu_a \rightarrow -\infty, \quad \text{as } a \rightarrow +\infty.$$

Moreover, u_a satisfies

$$u_a(x) = \frac{-\mu_a}{\sqrt{a}} \left(U(\sqrt{-\mu_a}(x - x_a)) + \varpi_a(x) \right), \quad (1.12)$$

$$\text{with } \int_{\mathbb{R}^3} \left[-\frac{1}{\mu_a} |\nabla \varpi_a|^2 + \varpi_a^2 \right] = o\left(\frac{1}{(\sqrt{-\mu_a})^3}\right).$$

Throughout this paper, we call u_a a single-peak solution of (1.8)–(1.9) if u_a satisfies (1.15). To the best knowledge of us, if the critical point of $P(x)$ is not isolated, not much is known for the exact location of the concentrated point, nor for the local uniqueness of the solutions. In the paper, we assume that $P(x)$ obtains its local minimum or local maximum at Γ_i ($i = 1, \dots, m$) and Γ_i is a closed 2 dimensional hyper-surface satisfying $\Gamma_i \cap \Gamma_j = \emptyset$ for $i \neq j$. More precisely, we assume that the following conditions hold.

(P). *There exist $\delta > 0$ and some C^2 compact hypersurfaces Γ_i ($i = 1, \dots, m$) without boundary, satisfying*

$$P(x) = P_i, \quad \frac{\partial P(x)}{\partial \nu_i} = 0, \quad \frac{\partial^2 P(x)}{\partial \nu_i^2} \neq 0, \quad \text{for any } x \in \Gamma_i \text{ and } i = 1, \dots, m,$$

where $P_i \in \mathbb{R}$, ν_i is the unit outward normal of Γ_i at $x \in \Gamma_i$. Moreover, $P \in C^4(\bigcup_{i=1}^m W_{\delta,i})$ and $P(x) = O(e^{\alpha|x|})$ for some $\alpha \in (0, 2)$. Here we denote $W_{\delta,i} := \{x \in \mathbb{R}^3, \text{dist}(x, \Gamma_i) < \delta\}$.

We would like to point out that the assumption (P) was first introduced in [24] by Luo, Peng and Yan where they studied the Bose-Einstein condensates. A specific example of $P(x)$ was also given by [24]. Observe that the assumption (P) implies that $P(x)$ obtains its local minimum or local maximum on the hypersurface Γ_i for $i = 1, \dots, m$. It is also easy to see that if $\delta > 0$ is small, the set $\Gamma_{t,i} = \{x : P(x) = t\} \cap W_{\delta,i}$ consists of two compact hypersurfaces in \mathbb{R}^3 without boundary for $t \in [P_i, P_i + \theta]$ (or $t \in [P_i - \theta, P_i]$) provided $\theta > 0$ is small. Moreover, the outward unit normal vector $\nu_{t,i}(x)$ and the j -th principal tangential unit vector $\tau_{t,i,j}(x)$ ($j = 1, 2$) of $\Gamma_{t,i}$ at x are Lip-continuous in $W_{\delta,i}$.

Remark 1.2. *When $m = 1$, for simplicity of notations we denote $P_0 =: P_i, b_0 := b_i$ and omit all the subscript i .*

Applying the local Pohozaev identities, we can easily prove that a single peak solution of (1.8)–(1.9) must concentrate at a critical point of $P(x)$, for which we can also refer to [10]. If the assumption (P) holds and the concentrated points belong to Γ , we ask further where the concentrating points locate on Γ . The following result gives our answer for this question.

Theorem 1.3. *Assume that the assumption (P) holds. If u_a is a single-peak solution of (1.8)–(1.9), concentrating at $\{b_0\}$ with $b_0 \in \Gamma$ as $a \nearrow +\infty$, then*

$$(D_{\tau_j} \Delta V)(b_0) = 0, \quad \text{with } j = 1, 2. \tag{1.13}$$

where τ_j is the j -th principal tangential unit vector of Γ at b_0 .

Theorem 1.3 implies that not every $\{b_0\}$ with $b_0 \in \Gamma$ can generate a single-peak solution for (1.8)–(1.9). In order to investigate the converse of Theorem 1.3, we have to add the following non-degenerate assumption on the critical point of $P(x)$. We say that $x_0 \in \Gamma$ is non-degenerate on Γ if it satisfies:

$$\frac{\partial^2 P(x_0)}{\partial \nu^2} \neq 0 \text{ and } \det\left(\left(\frac{\partial^2 \Delta P(x_0)}{\partial \tau_l \partial \tau_j}\right)_{1 \leq l, j \leq 2}\right) \neq 0.$$

Theorem 1.4. *Under the assumption (P), if $b_0 \in \Gamma$ is a non-degenerate critical point of $P(x)$ on Γ satisfying (1.7), then there exists a large constant $a_0 > 0$, such that (1.8)–(1.9) has a single-peak solution u_a concentrating at b_0 as $a \in [a_0, +\infty)$.*

The existence result in Theorem 1.4 is new even without the L^2 -norm constraint since our potential includes the degenerate case. Also if $P(x)$ does not achieve its global minimum on Γ , any solution concentrating at a point on Γ is not a ground state. So our existence result can not be obtained by the method in [6, 12, 17, 18].

To state the local uniqueness result of single-peak solutions for (1.8)–(1.9), we give another assumption (\tilde{P}) of $P(x)$: if b_0 is non-degenerate and

$$\left(\frac{\partial^2 \Delta P(b_0)}{\partial \tau_l \partial \tau_j}\right)_{1 \leq l, j \leq 2} + \frac{\partial \Delta P(b_0)}{\partial \nu} \text{diag}(\kappa_1, \kappa_2)$$

is non-singular, where κ_j is the j -th principal curvature of Γ at b_0 for $j = 1, 2$.

Our second main result is the following.

Theorem 1.5. *Suppose that the assumptions (P) and (\tilde{P}) hold. Let $u_a^{(1)}(x)$ and $u_a^{(2)}(x)$ be two single-peak solutions of (1.8)–(1.9) concentrating at b_0 with $b_0 \in \Gamma$. Then there exists a large positive number a_0 , such that $u_a^{(1)}(x) \equiv u_a^{(2)}(x)$ for all a with $a_0 \leq a < +\infty$.*

Remark 1.6. *Our method can also be used to study the following Choquard equation with the dimensions $N = 4, 5$*

$$-\epsilon^2 \Delta u + P(x)u = \frac{a}{8\pi\epsilon^2} \left(\int_{\mathbb{R}^N} \frac{u^2(y)}{|x-y|^{N-2}} dy \right) u + \mu u, \quad x \in \mathbb{R}^N. \quad (1.14)$$

We would like to point out that in this case a goes to $\begin{cases} a^* = \int_{\mathbb{R}^4} U^2(x) dx, & N = 4, \\ 0, & N = 5. \end{cases}$

The main difference in the discussion of the local uniqueness for (1.8) and [23] is that the Lagrange multiplier μ_a in (1.8) also depends on the solution u_a . Hence its corresponding linearized operator has changed. For the details, see Lemma B.1. Fortunately, such change does not bring much difficulties.

To our knowledge, except [23], there seems to be no other local uniqueness results for peak (or bubbling) solutions of a Schrödinger-Newton system. Even for Schrödinger equations, there are very few results about the local uniqueness of peak solutions. The classical moving plane method is not still effective to study the uniqueness of concentrating solutions. There are two main tools to investigate the uniqueness of concentration solutions, i.e., the topological degree method and Pohozaev identities. Before the local uniqueness result of multi-bump solutions in [24], other local uniqueness results for peak (or bubbling) solutions of Schrödinger equations are available only for the case when the solutions blow up at x_0 , an isolated critical point of the potential $P(x)$. When x_0 is a non-degenerate critical point of $P(x)$, that is, (D^2P) is non-singular at x_0 , one can prove the local uniqueness of the peak solution concentrating at x_0 either by counting the local degree of the corresponding reduced finite dimensional problem as in [1, 3, 9], or by using Pohozaev type identities as in [2, 7, 10, 11, 13].

Compared with the topological degree method, it can deal with the degenerate potential to use the Pohozaev identities to prove the local uniqueness, one can refer to [2, 7, 13]. We want to point out that in [2, 7, 13], though the critical point x_0 is degenerate, the rate of degeneracy along each direction is the same. Moreover, in [10] Grossi gave an example which shows that local uniqueness may not be true at a degenerate critical point x_0 of $P(x)$. When peak solutions concentrates at a degenerate critical point, it is more subtle to study the local uniqueness of it. One can refer to [24]. Under our assumption (P), the potential $P(x)$ is non-degenerate along

the normal direction ν of Γ . But along each tangential direction of Γ , $P(x)$ is degenerate. Such non-uniform degeneracy causes the estimates more complicated. Moreover, there are two terms involving volume integral in the corresponding local Pohozaev identity, which brings us some difficulties.

Finally, we give a non-existence of multi-peak solutions for (1.8)–(1.9).

Theorem 1.7. *Under the assumption (P), assume that $b_i \in \Gamma_i (i = 1, \dots, m)$ are non-degenerate critical points of $P(x)$ on Γ_i satisfying*

$$(D_{\tau_{i,j}} \Delta P)(b_i) = 0, \text{ with } i = 1, \dots, m \text{ and } j = 1, 2,$$

where $\tau_{i,j}$ is the j -th principal tangential unit vector of Γ at b_i . Then there exists a large constant $a_0 > 0$, such that problem (1.8)–(1.9) has no m -peaks solutions ($m \geq 2$) of this form

$$u_a(x) = \frac{-\mu_a}{\sqrt{a}} \sum_{i=1}^m \left(U(\sqrt{-\mu_a}(x - x_{a,i})) + \varpi_a(x) \right), \quad (1.15)$$

with $\int_{\mathbb{R}^3} \left[-\frac{1}{\mu_a} |\nabla \varpi_a|^2 + \varpi_a^2 \right] = o\left(\frac{1}{(-\mu_a)^{\frac{3}{2}}}\right)$ and some points $x_{a,i} \in \mathbb{R}^3 (i = 1, \dots, m)$ satisfying $x_{a,i} \rightarrow b_i, i = 1, \dots, m$ as $a \rightarrow +\infty$ and $b_i \neq b_j$ for $i \neq j$.

Remark 1.8. *The nonexistence result of multi-peak solutions for (1.8)–(1.9) is very different from the Schrödinger problem studied in [24], which is mainly caused by the nonlocal term.*

To prove Theorem 1.7, we mainly use some local Pohozaev identities and some contradiction argument. We have to obtain some accurate estimates involved by the nonlocal term as [23].

This paper is organized as follows. We prove Theorem 1.1 in section 2. In section 3, we estimate the Lagrange multiplier μ_a in terms of a . In section 4 we prove the results for the location of the peaks and for the existence of single-peak solutions. We study the local uniqueness of single-peak solutions in section 5. In section 6, we prove the nonexistence result of multi-peak solutions. We put some known results and some basic and technical estimates in Appendices A to C.

For simplicity, we use $|u|_q (2 \leq q \leq 6)$ to denote $(\int_{\mathbb{R}^3} |u(x)|^q dx)^{\frac{1}{q}}$ and $\|u\|$ the usual $H^1(\mathbb{R}^3)$ norm. In this paper, we always assume that $b_0 \in \Gamma$.

2. THE PROOF OF THEOREM 1.1

First, we study the following problem:

$$-\Delta u = P_1(x)u, \quad u > 0, \quad \text{in } \mathbb{R}^3, \quad (2.1)$$

where the function $P_1(x)$ satisfies $P_1 > 1$ in $B_R(0) \setminus B_t(0)$ for some fixed $t > 0$ and large $R > 0$.

In order to prove Theorem 1.1, first we give the following result.

Proposition 2.1. *(Proposition 2.1, [24]) Problem (2.1) has no solution.*

With Proposition 2.1 at hand, now we are in a position to prove Theorem 1.1.

Proof of Theorem 1.1. First, we prove that $\mu_a \rightarrow -\infty$. We argue by an indirect method. Suppose that $|\mu_a| \leq M$. Since $\int_{\mathbb{R}^3} u_a^2 = 1$, by the Moser iteration, we can prove that u_a is uniformly bounded. That is, u_a does not blow up.

Suppose that $\mu_a \rightarrow +\infty$. Set $P_1(x) = \mu_a - P(x) + \frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{u_a^2(y)}{|x-y|} dy$. Noting that u_a concentrates at some points, we have for $x \in \mathbb{R}^3 \setminus B_t(0)$ and $t > 0$

$$\begin{aligned} & \int_{\mathbb{R}^3} \frac{u_a^2(y)}{|x-y|} dy \\ &= \int_{|x-y| \leq \frac{t}{2}} \frac{u_a^2(y)}{|x-y|} dy + \int_{|x-y| \geq \frac{t}{2}} \frac{u_a^2(y)}{|x-y|} dy \\ &= o(1) \int_{|x-y| \leq \frac{t}{2}} \frac{1}{|x-y|} dy + \left(\int_{|x-y| \geq \frac{t}{2}} |u_a(y)|^{\frac{20}{7}} dy \right)^{\frac{7}{10}} \left(\frac{1}{|x-y|^{\frac{10}{3}}} dy \right)^{\frac{3}{10}} \\ &= o(1)t^2 + O\left(\frac{1}{t^{\frac{1}{10}}}\right), \end{aligned} \tag{2.2}$$

where $o(1)$ goes to zero as $a \rightarrow \infty$. Hence from (2.2) we may suppose that $\frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{u_a^2(y)}{|x-y|} dy \geq -1$ in $\mathbb{R}^3 \setminus B_t(0)$ for some $t > 0$. Hence for any fixed $R > 0$, we usually have

$$P_1(x) = \mu_a - P(x) + \frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{u_a^2(y)}{|x-y|} dy > 1, \quad x \in B_R(0) \setminus B_t(0).$$

From Proposition 2.1, we obtain a contradiction. So we have proved that $\mu_a \rightarrow -\infty$. Let $\lambda_a = -\mu_a$. Let x_a be the maximum point of u_a . By equation (1.8), we find

$$\frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{u_a^2(y)}{|x-y|} dy u_a(x) \geq (\lambda_a + P(x_a)) u_a(x) > 0,$$

which implies that $a > 0$.

Denote $\bar{u}_a(x) = \frac{1}{\lambda_a} u_a\left(\frac{x}{\sqrt{\lambda_a}}\right)$. Then

$$-\Delta \bar{u}_a + \left(1 + \frac{1}{\lambda_a} P\left(\frac{x}{\sqrt{\lambda_a}}\right)\right) \bar{u}_a = \frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{\bar{u}_a^2(y)}{|x-y|} dy \bar{u}_a(x), \quad \text{in } \mathbb{R}^3, \tag{2.3}$$

and

$$\int_{\mathbb{R}^3} \bar{u}_a^2 = \frac{1}{\lambda_a^{\frac{1}{2}}}. \tag{2.4}$$

By (2.3) and (2.4), applying Moser iteration, we can show that $|u_a| \leq C$ for some constant independent of a . Let \bar{x}_a be a maximum point of \bar{u}_a . Then

$$\frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{\bar{u}_a^2(y)}{|\bar{x}_a - y|} dy \bar{u}_a(\bar{x}_a) \geq \left(1 + \frac{1}{\lambda_a} P\left(\frac{\bar{x}_a}{\sqrt{\lambda_a}}\right)\right) \bar{u}_a(\bar{x}_a),$$

which gives $a \geq 8\pi \left(\int_{\mathbb{R}^3} \frac{\bar{u}_a^2(y)}{|\bar{x}_a - y|} dy\right)^{-1} \geq c_0 > 0$. Applying the standard blow-up argument, by (2.4) we can check that there holds

$$\bar{u}_a = U_a(x - \bar{x}_a) + \bar{\omega}_a(x), \tag{2.5}$$

for some $\bar{x}_a \in \mathbb{R}^3$ with

$$\int_{\mathbb{R}^3} [|\nabla \bar{\omega}_a|^2 + (\bar{\omega}_a)^2] = o(1),$$

and U_a is the unique positive solution of

$$-\Delta u + u = \frac{a}{8\pi} \int_{\mathbb{R}^3} \frac{u^2(x)}{|x-y|} dy u, \quad u \in H^1(\mathbb{R}^3), \quad u(0) = \max_{x \in \mathbb{R}^3} u(x).$$

Observing that $U_a = \frac{1}{\sqrt{a}}U$, it follows from (2.4) and (2.5) that $a \rightarrow a^* \lambda_a^{\frac{1}{2}} \rightarrow +\infty$. \square

3. SOME ESTIMATES FOR GENERAL POTENTIALS

In this section, we shall estimate μ_a respect to a .

Let $\epsilon = \frac{1}{\sqrt{-\mu_a}}$ and $u(x) \mapsto \frac{-\mu_a}{\sqrt{a}}u(x)$. Then (1.8) can be rewritten as

$$-\epsilon^2 \Delta u + (1 + \epsilon^2 P(x))u = \frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{u^2(y)}{|x-y|} dy u(x), \quad u \in H^1(\mathbb{R}^3). \quad (3.1)$$

For any $a \in \mathbb{R}^+$, we define $\|u\|_a := \int_{\mathbb{R}^3} (\epsilon^2 |\nabla u|^2 + u^2)^{\frac{1}{2}}$.

By (1.15), we know that a single-peak solution of (3.1) has the following form

$$\tilde{u}_a(x) = U_{\epsilon, x_a}(x) + \varphi_a(x), \quad \text{with } \|\varphi_a\|_a = o(\epsilon^{\frac{3}{2}}),$$

where $U_{\epsilon, x_a}(x) := (1 + \epsilon^2 P_0)U\left(\frac{\sqrt{1 + \epsilon^2 P_0}(x - x_a)}{\epsilon}\right)$. Then, there holds

$$\begin{aligned} -\epsilon^2 \Delta \varphi_a + \left((1 + \epsilon^2 P(x))\varphi_a - \frac{1}{4\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{U_{\epsilon, x_a}(y)\varphi_a(y)}{|x-y|} dy U_{\epsilon, x_a}(x) \right. \\ \left. - \frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(U_{\epsilon, x_a}(y))^2}{|x-y|} dy \varphi_a(x) \right) = \mathcal{R}_{a, \epsilon}(\varphi_a) + \mathcal{L}_a(x), \end{aligned} \quad (3.2)$$

where

$$\begin{aligned} \mathcal{R}_{a, \epsilon}(\varphi_a) &= \left(\frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(U_{\epsilon, x_a} + \varphi_a)^2(y)}{|x-y|} dy (U_{\epsilon, x_a} + \varphi_a)(x) - \frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(U_{\epsilon, x_a}(y))^2}{|x-y|} dy U_{\epsilon, x_a}(x) \right. \\ &\quad \left. - \frac{1}{4\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(U_{\epsilon, x_a}\varphi_a)(y)}{|x-y|} dy U_{\epsilon, x_a}(x) \right) - \frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(U_{\epsilon, x_a})^2(y)}{|x-y|} dy \varphi_a(x) \\ &= \frac{1}{4\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(U_{\epsilon, x_a}\varphi_a)(y)}{|x-y|} dy \varphi_a(x) + \frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(\varphi_a)^2(y)}{|x-y|} dy U_{\epsilon, x_a}(x) \\ &\quad + \frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \frac{(\varphi_a)^2(y)}{|x-y|} dy \varphi_a(x) \end{aligned} \quad (3.3)$$

and

$$\mathcal{L}_a(x) = -\epsilon^2(P(x) - P_0)U_{\epsilon, x_a}(x).$$

We can move x_a a bit (still denoted by x_a), so that the error term $\varphi_a \in E_{a, x_a}$, where

$$E_{a, x_a} := \left\{ u(x) \in H^1(\mathbb{R}^3) : \left\langle u, \frac{\partial U_{\epsilon, x_a}(x)}{\partial x_j} \right\rangle_a = 0, \quad j = 1, 2, 3 \right\}.$$

Let L_a be the bounded linear operator from $H^1(\mathbb{R}^3)$ to itself, defined by

$$\langle L_a u, v \rangle_a = \int_{\mathbb{R}^3} (\epsilon^2 \nabla u \nabla v + (1 + \epsilon^2 P(x))uv) - \frac{3}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{u^2(x)(uv)(y)}{|x-y|} dx dy.$$

Then, it is standard to prove the following lemma.

Lemma 3.1. *There exist constants $\varrho > 0$ and large $a_0 > 0$ such that for all a with $a_0 \leq a < +\infty$, it holds*

$$\|L_a u\|_a \geq \varrho \|u\|_a, \text{ for all } u \in E_{a, x_a}. \quad (3.4)$$

Lemma 3.2. *There exists a constant $C > 0$ independent of a such that*

$$\|\mathcal{R}_{a, \epsilon}(\varphi_a)\|_a = O(\epsilon^{-3} \|\varphi_a\|_a^3 + \epsilon^{-\frac{3}{2}} \|\varphi_a\|_a^2). \quad (3.5)$$

Proof. From (3.3), for any $v \in H^1(\mathbb{R}^3)$ we have

$$\begin{aligned} & (\mathcal{R}_{a, \epsilon}(\varphi_a), v) \\ &= \frac{1}{4\pi\epsilon^2} \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\varphi_a(x)v(x)(U_{\epsilon, x_a}\varphi_a)(y)}{|x-y|} dx dy}_{:=A_1} + \frac{1}{8\pi\epsilon^2} \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U_{\epsilon, x_a}(x)v(x)(\varphi_a)^2(y)}{|x-y|} dx dy}_{:=A_2} \\ &+ \frac{1}{8\pi\epsilon^2} \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\varphi_a(x)v(x)(\varphi_a)^2(y)}{|x-y|} dx dy}_{:=A_3}. \end{aligned} \quad (3.6)$$

By Lemma A.2, we have

$$A_1 \leq C\epsilon^{-1} \|\varphi_a\|_a^2 \|v\|_a \|U_{\epsilon, x_a}\|_a \leq C\epsilon^{\frac{1}{2}} \|\varphi_a\|_a^2 \|v\|_a, \quad A_2 \leq C\epsilon^{\frac{1}{2}} \|\varphi_a\|_a^2 \|v\|_a,$$

and

$$A_3 \leq C\epsilon^{-1} \|\varphi_a\|_a^3 \|v\|_a. \quad (3.7)$$

It follows from (3.6) to (3.7) that (3.5) holds. \square

Lemma 3.3. *There holds*

$$\|\mathcal{L}_a\|_a = O\left(|(P(x_a) - P_0)|\epsilon^{\frac{7}{2}} + |\nabla P(x_a)|\epsilon^{\frac{9}{2}} + \epsilon^{\frac{11}{2}}\right). \quad (3.8)$$

Proof. For any $v \in H^1(\mathbb{R}^3)$, we have

$$\begin{aligned} (\mathcal{L}_a(x), v) &= -\epsilon^2 \int_{\mathbb{R}^3} (P(x) - P_0) U_{\epsilon, x_a}(x) v(x) dx \\ &= -\epsilon^2 \underbrace{\int_{\mathbb{R}^3} (P(x) - P(x_a)) U_{\epsilon, x_a}(x) v(x) dx}_{:=B_1} - \epsilon^2 \underbrace{\int_{\mathbb{R}^3} (P(x_a) - P_0) U_{\epsilon, x_a}(x) v(x) dx}_{:=B_2}. \end{aligned}$$

By Hölder inequality and the exponential of $U(x)$ at infinity, we can estimate

$$\begin{aligned} B_1 &= \int_{B_\delta(x_a)} (P(x) - P(x_a)) U_{\epsilon, x_a}(x) v(x) dx + \int_{B_\delta^c(x_a)} (P(x) - P(x_a)) U_{\epsilon, x_a}(x) v(x) dx \\ &= \int_{B_\delta(x_a)} (\nabla P(x_a) \cdot (x - x_a) + O(|x - x_a|^2)) U_{\epsilon, x_a}(x) v(x) dx + C \left(\int_{B_\delta^c(x_a)} U_{\epsilon, x_a}^2(x) dx \right)^{\frac{1}{2}} |v|_2 \\ &\leq |\nabla P(x_a)| \left(\int_{B_\delta(x_a)} |x - x_a|^2 U_{\epsilon, x_a}^2(x) dx \right)^{\frac{1}{2}} |v|_2 + C \left(\int_{B_\delta(x_a)} |x - x_a|^4 U_{\epsilon, x_a}^2(x) dx \right)^{\frac{1}{2}} |v|_2 \\ &\quad + O(e^{-\frac{\theta}{\epsilon}}) \|v\|_a \\ &= O(|\nabla P(x_a)| \epsilon^{\frac{5}{2}} \|v\|_a) + O(\epsilon^{\frac{7}{2}} \|v\|_a), \end{aligned}$$

and

$$\begin{aligned} B_2 &= \int_{\mathbb{R}^3} (P(x_a) - P_0) U_{\epsilon, x_a}(x) v(x) dx \\ &\leq |P(x_a) - P_0| \left(\int_{\mathbb{R}^3} U_{\epsilon, x_a}^2(x) dx \right)^{\frac{1}{2}} \|v\|_2 = O(|P(x_a) - P_0| \epsilon^{\frac{3}{2}}) \|v\|_a. \end{aligned}$$

Combining all the estimates above, we know that (3.8) holds. \square

With Lemmas 3.1 to 3.3 at hand, we can prove

Lemma 3.4. *A single-peak solution \tilde{u}_a for (3.1) concentrating at b_0 has the following form*

$$\tilde{u}_a(x) = U_{\epsilon, x_a}(x) + \varphi_a(x), \quad (3.9)$$

with $\varphi_a \in E_{a, x_a}$ and

$$\|\varphi_a\|_a = O\left(|P(x_a) - P_0| \epsilon^{\frac{7}{2}} + |\nabla P(x_a)| \epsilon^{\frac{9}{2}} + \epsilon^{\frac{11}{2}}\right). \quad (3.10)$$

Proof. Then from (3.2), (3.4), (3.5), (3.8) and by applying the contraction mapping theorem, we can get (3.9) and (3.10) using the standard argument. \square

Let $\tilde{\varphi}_a(x) = \varphi_a(\epsilon x + x_a)$. Then, $\tilde{\varphi}_a$ satisfies $\|\tilde{\varphi}_a\|_a = O(\epsilon^2)$. Using the Moser iteration, we can prove $\|\tilde{\varphi}_a\|_{L^\infty(\mathbb{R}^3)} = o(1)$. From this and the comparison theorem, similar to Proposition 2.2 in [23], we can prove the following estimates for $\tilde{u}_a(x)$ away from the concentrated point b_0 .

Proposition 3.5. *Suppose that $\tilde{u}_a(x)$ is a single-peak solution of (3.1) concentrating at b_0 . Then for any fixed $R \gg 1$, there exist some $\theta > 0$ and $C > 0$, such that*

$$|\tilde{u}_a(x)| + |\nabla \tilde{u}_a(x)| \leq C e^{-\theta|x-x_a|/\epsilon}, \quad \text{for } x \in \mathbb{R}^3 \setminus B_{R\epsilon}(x_a). \quad (3.11)$$

Lemma 3.6. *There holds*

$$\int_{\mathbb{R}^3} U^2 = \frac{3}{32\pi} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U^2(y)}{|x-y|} dx dy. \quad (3.12)$$

Proof. It follows directly from the following two identities:

$$\int_{\mathbb{R}^3} (|\nabla U|^2 + U^2) = \frac{1}{8\pi} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U^2(y)}{|x-y|} dx dy$$

and

$$\int_{\mathbb{R}^3} (|\nabla U|^2 + 3U^2) = \frac{5}{16\pi} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U^2(y)}{|x-y|} dx dy,$$

where the last equality can be deduced by multiplying $\langle x, \nabla U \rangle$ on both sides of (1.10) and integrating on \mathbb{R}^3 . \square

Proposition 3.7. *Letting $a \rightarrow +\infty$, there holds*

$$\frac{a}{\sqrt{-\mu_a}} = a_* - \frac{P_0}{2\mu_a} a_* + O\left(|P(x_a) - P_0| \frac{1}{(-\mu_a)} + |\nabla P(x_a)| \frac{1}{(\sqrt{-\mu_a})^3} + \frac{1}{\mu_a^2}\right). \quad (3.13)$$

Proof. From (1.8) and (1.9), we have

$$1 = \int_{\mathbb{R}^3} (u_a(x))^2 = \int_{\mathbb{R}^3} \left(\bar{U}_{a,x_a}(x) + \frac{(-\mu_a)}{\sqrt{a}} \varphi_a(x) \right)^2,$$

where $\bar{U}_{a,x_a}(x) = \frac{-\mu_a + P_0}{\sqrt{a}} U(\sqrt{-\mu_a + P_0}(x - x_a))$. By direct computation, we can obtain

$$a = a_* \sqrt{-\mu_a + P_0} + O\left(|P(x_a) - P_0| \frac{1}{\sqrt{-\mu_a}} + |\nabla P(x_a)| \frac{1}{(-\mu_a)} + \frac{1}{(\sqrt{-\mu_a})^3}\right),$$

which implies that (3.13) is true. \square

4. LOCATING THE PEAK AND THE EXISTENCE OF SINGLE-PEAK SOLUTIONS

First, we locate the peak for a single-peak solution. Let \tilde{u}_a be a single-peak solution of (3.1). Then for any small fixed $\rho > 0$, from (B.4) and (C.1) we find

$$\begin{aligned} & \epsilon^2 \int_{B_\rho(x_a)} \frac{\partial P(x)}{\partial x_j} (\tilde{u}_a)^2 dx \\ &= \underbrace{-2\epsilon^2 \int_{\partial B_\rho(x_a)} \frac{\partial \tilde{u}_a}{\partial \bar{\nu}} \frac{\partial \tilde{u}_a}{\partial x_j} d\sigma}_{:=C_1} + \underbrace{\epsilon^2 \int_{\partial B_\rho(x_a)} |\nabla \tilde{u}_a|^2 \bar{\nu}_j(x) d\sigma}_{:=C_2} \\ &+ \underbrace{\int_{\partial B_\rho(x_a)} (1 + \epsilon^2 P(x)) (\tilde{u}_a)^2 \bar{\nu}_j(x) d\sigma}_{:=C_3} - \underbrace{\frac{1}{8\pi\epsilon^2} \int_{\partial B_\rho(x_a)} \int_{\mathbb{R}^3} \frac{(\tilde{u}_a)^2}{|x-y|} dy (\tilde{u}_a)^2 \bar{\nu}_j(x) d\sigma}_{:=C_4} \\ &- \underbrace{\frac{1}{8\pi\epsilon^2} \int_{B_\rho(x_a)} \int_{\mathbb{R}^3} \frac{(\tilde{u}_a(y))^2 (\tilde{u}_a(x))^2 (x_j - y_j)}{|x-y|^3} dy dx}_{:=C_5} \\ &= O(e^{-\frac{\theta}{\epsilon}}), \text{ with some } \theta > 0, \end{aligned} \quad (4.1)$$

where $j = 1, 2, 3$ and $\bar{\nu}(x) = (\bar{\nu}_1(x), \bar{\nu}_2(x), \bar{\nu}_3(x))$ is the outward unit normal of $\partial B_\rho(x_a)$. And then (4.1) implies the first necessary condition for the concentrated point b_0 :

$$\nabla P(b_0) = 0.$$

Now we are in a position to prove Theorem 1.3.

Proof of Theorem 1.3. Since $x_a \rightarrow b_0 \in \Gamma$, we find that there is a $t_a \in [P_0, P_0 + \sigma]$ if Γ is a local minimum set of $P(x)$, or $t_a \in [P_0 - \sigma, P_0]$ if Γ is a local maximum set of $P(x)$, such that $x_a \in \Gamma_{t_a}$. Let τ_a be the unit tangential vector of Γ_{t_a} at x_a . Then

$$G(x_a) = 0, \text{ where } G(x) = \langle \nabla P(x), \tau_a \rangle.$$

We have the following expansion:

$$G(x) = \langle \nabla G(x_a), x - x_a \rangle + \frac{1}{2} \langle \langle \nabla^2 G(x_a), x - x_a \rangle, x - x_a \rangle + o(|x - x_a|^2), \text{ for } x \in B_\rho(x_a).$$

Then it follows from (3.10), (4.1) and the above expansion that

$$\begin{aligned}
\int_{\mathbb{R}^3} G(x)U_{\epsilon, x_a}^2(x) &= \int_{B_\rho(x_a)} G(x)U_{\epsilon, x_a}^2(x) + O(e^{-\frac{\theta}{\epsilon}}) \\
&= -2 \int_{B_\rho(x_a)} G(x)U_{\epsilon, x_a}(x)\varphi_a - \int_{B_\rho(x_a)} G(x)\varphi_a^2 + O(e^{-\frac{\theta}{\epsilon}}) \\
&= O\left([\epsilon^{\frac{5}{2}}|\nabla G(x_a)| + \epsilon^{\frac{7}{2}}\|\varphi_a\|_a + \epsilon|\nabla G(x_a)| \cdot \|\varphi_a\|_a^2\right) + O(e^{-\frac{\theta}{\epsilon}}) \\
&= O(\epsilon^6).
\end{aligned} \tag{4.2}$$

On the other hand, noting that $G(x_a) = 0$, it is easy to check

$$\int_{\mathbb{R}^3} G(x)U_{\epsilon, x_a}^2(x) = \frac{1}{6}\epsilon^5\Delta G(x_a) \int_{\mathbb{R}^3} |x|^2U^2 + O(\epsilon^7). \tag{4.3}$$

Then it follows from (4.2) and (4.3) that $(\Delta G)(x_a) = O(\epsilon)$. Thus by the assumption (P), we get (1.7). \square

Now, we study the existence of single-peak solutions for (1.11) with $\lambda > 0$ a large parameter. Letting $\eta = \frac{1}{\sqrt{\lambda}}$ and $w(x) \mapsto \lambda w(x)$, then (1.11) can be changed to following problem:

$$-\eta^2\Delta w + (1 + \eta^2P(x))w = \frac{1}{8\pi\eta^2} \int_{\mathbb{R}^3} \frac{w^2}{|x-y|} dy w, \quad w \in H^1(\mathbb{R}^3). \tag{4.4}$$

In the sequel, we denote $\langle u, v \rangle_\eta = \int_{\mathbb{R}^3} (\eta^2\nabla u \nabla v + uv)$ and $\|u\|_\eta = \langle u, u \rangle_\eta^{\frac{1}{2}}$. Now for $\eta > 0$ small, we construct a single-peak solution u_η of (4.4) concentrating at b_0 . Here we can prove the following result in a standard way.

Proposition 4.1. *There is an $\eta_0 > 0$, such that for any $\eta \in (0, \eta_0]$, and z_0 close to b_0 , there exists $v_{\eta, z_0} \in F_{\eta, z_0}$ such that*

$$\int_{\mathbb{R}^3} (\eta^2\nabla w_\eta \nabla \psi + (1 + \eta^2P(x))w_\eta\psi) = \frac{1}{8\pi\eta^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{w_\eta^2(y)}{|x-y|} w_\eta(x)\psi(x) dy dx, \quad \text{for all } \psi \in F_{\eta, z_0},$$

where

$$w_\eta(x) = U_{\eta, z_0}(x) + \varphi_{\eta, z_0}(x) \tag{4.5}$$

and

$$F_{\eta, z_0} = \left\{ u(x) \in H^1(\mathbb{R}^3) : \left\langle u, \frac{\partial U_{\eta, z_0}(x)}{\partial x_j} \right\rangle_\eta = 0, \quad j = 1, 2, 3 \right\}.$$

Moreover, it holds

$$\|\varphi_{\eta, z_0}\|_\eta = O(|P(z_0) - P_0|\eta^{\frac{7}{2}} + |\nabla P(z_0)|\eta^{\frac{9}{2}} + \eta^{\frac{11}{2}}). \tag{4.6}$$

To obtain a true solution for (4.4), we have to choose z_0 such that

$$\int_{B_\rho(x_a)} \left(-\eta^2\Delta w_\eta \frac{\partial w_\eta}{\partial x_j} + (1 + \eta^2P(x))w_\eta \frac{\partial w_\eta}{\partial x_j} - \frac{1}{8\pi\eta^2} \int_{\mathbb{R}^3} \frac{w_\eta^2(y)}{|x-y|} dy w_\eta \frac{\partial w_\eta}{\partial x_j} \right) = 0, \quad j = 1, 2, 3.$$

Similar to (4.1), it is easy to check that the above identities are equivalent to

$$\int_{B_\rho(x_a)} \frac{\partial P(x)}{\partial x_j} w_\eta^2 = O(e^{-\frac{\theta}{\eta}}), \quad \forall j = 1, 2, 3. \tag{4.7}$$

For z_0 close to b_0 , and $z_0 \in \Gamma_t$ for some t close to P_0 , now we use ν to denote the unit normal vector of Γ_t at z_0 , while we use τ_j ($j = 1, 2$) to denote the principal directions of Γ_t at x_a . Then, at z_0 , it holds

$$D_{\tau_j}P(z_0) = 0, \text{ for } j = 1, 2, \text{ and } |\nabla P(z_0)| = |D_\nu P(z_0)|.$$

We first prove the following results.

Lemma 4.2. *Under the assumption (P), $\int_{B_\rho(x_a)} D_\nu P(x) u_\eta^2 = O(e^{-\frac{\rho}{\eta}})$ is equivalent to*

$$D_\nu P(z_0) = O(\eta^2). \quad (4.8)$$

Proof. First, from (4.5) we have

$$\begin{aligned} & \int_{\mathbb{R}^3} D_\nu P(x) U_{\eta, z_0}^2(x) \\ &= -2 \int_{\mathbb{R}^3} D_\nu P(x) U_{\eta, z_0}(x) \varphi_{\eta, z_0} - \int_{\mathbb{R}^3} D_\nu P(x) \varphi_{\eta, z_0}^2 + O(e^{-\frac{\rho}{\eta}}) \\ &= O(|D_\nu P(z_0)| \eta^{\frac{3}{2}} \cdot \|\varphi_{\eta, z_0}\|_\eta + \eta^{\frac{5}{2}} \|\varphi_{\eta, z_0}\|_\eta + \|\varphi_{\eta, z_0}\|_\eta^2) = O(\eta^5). \end{aligned} \quad (4.9)$$

On the other hand, we have

$$\int_{\mathbb{R}^3} D_\nu P(x) U_{\eta, z_0}^2(x) = a_* \eta^3 D_\nu P(z_0) + O(\eta^5), \quad (4.10)$$

Then it follows from (4.9) and (4.10) that (4.8) holds. \square

Lemma 4.3. *Under the assumption (P), $\int_{B_\rho(x_a)} D_\tau P(x) u_\eta^2 = O(e^{-\frac{\rho}{\eta}})$ is equivalent to*

$$(D_\tau \Delta P)(z_0) = O(|P(z_0) - P_0| \eta + \eta^2). \quad (4.11)$$

Proof. Let $G(x) = \langle \nabla P(x), \tau \rangle$. Then, similar to the estimate (4.2), by (4.6) we have

$$\begin{aligned} \int_{\mathbb{R}^3} G(x) U_{\eta, z_0}^2(x) &= -2 \int_{\mathbb{R}^3} G(x) U_{\eta, z_0}(x) \varphi_{\eta, z_0} - \int_{\mathbb{R}^3} G(x) \varphi_{\eta, z_0}^2 + O(e^{-\frac{\rho}{\eta}}) \\ &= O(|P(z_0) - P_0| \eta^5 + \eta^7). \end{aligned} \quad (4.12)$$

On the other hand, in view of $G(z_0) = 0$, it is easy to show

$$\int_{\mathbb{R}^3} G(x) U_{\eta, z_0}^2(x) = \frac{1}{2} \eta^5 \Delta G(z_0) B + O(\eta^7), \quad (4.13)$$

where

$$B = \frac{1}{3} \int_{\mathbb{R}^3} |x|^2 U^2. \quad (4.14)$$

Thus, from (4.12) and (4.13) we can obtain (4.11). \square

Theorem 4.4. *For $\lambda > 0$ large, (1.11) has a solution u_λ satisfying*

$$u_\lambda(x) = \lambda \left(U(\sqrt{\lambda}(x - x_\lambda)) + \varpi_\lambda \right),$$

where $x_\lambda \rightarrow b_0$ and $\int_{\mathbb{R}^3} (|\nabla \varpi_\lambda|^2 + \varpi_\lambda^2) \rightarrow 0$ as $\lambda \rightarrow +\infty$.

Proof. As pointed out earlier, we need to solve (4.7). By Lemmas 4.2 and 4.3, the equation (4.7) is equivalent to

$$D_\nu P(z_0) = O(\eta^2), \quad (D_\tau \Delta P)(z_0) = O(|P(z_0) - P_0|\eta + \eta^2).$$

Let $\bar{z}_0 \in \Gamma$ be the point such that $z_0 - \bar{z}_0 = \alpha_0 \nu$ for some $\alpha_0 \in \mathbb{R}$. Then, we have $D_\nu P(\bar{z}_0) = 0$. As a result,

$$D_\nu P(z_0) = D_\nu P(z_0) - D_\nu P(\bar{z}_0) = D_{\nu\nu}^2 P(\bar{z}_0) \langle z_0 - \bar{z}_0, \nu \rangle + O(|z_0 - \bar{z}_0|^2).$$

By the non-degenerate assumption, we find that $D_\nu P(z_0) = O(\eta^2)$ is equivalent to $\langle z_0 - \bar{z}_0, \nu \rangle = O(\eta^2 + |z_0 - \bar{z}_0|^2)$. This means that $D_\nu P(z_0) = O(\eta^2)$ can be written as

$$|z_0 - \bar{z}_0| = O(\eta^2). \quad (4.15)$$

Let $\bar{\tau}_j$ be the j -th tangential unit vector of Γ at \bar{z}_0 . Now by the assumption (P), we have

$$(D_{\bar{\tau}_j} \Delta P)(z_0) = (D_{\bar{\tau}_j} \Delta P)(\bar{z}_0) + O(|z_0 - \bar{z}_0|) = (D_{\bar{\tau}_j} \Delta P)(\bar{z}_0) + O(\eta^2),$$

and

$$(D_{\bar{\tau}_j} \Delta P)(\bar{z}_0) = (D_{\bar{\tau}_j} \Delta P)(\bar{z}_0) - (D_{\tau_{j,0}} \Delta P)(b_0) = \langle (\nabla_T D_{\tau_{j,0}} \Delta P)(b_0), \bar{z}_0 - b_0 \rangle + O(|\bar{z}_0 - b_0|^2),$$

where ∇_T is the tangential gradient on Γ at $b_0 \in \Gamma$, and $\tau_{j,0}$ is the j -th tangential unit vector of Γ at b_0 . Therefore, $(D_\tau \Delta P)(z_0) = O(|P(z_0) - P_0|\eta + \eta^2)$ can be rewritten as

$$\langle (\nabla_T D_{\tau_{j,0}} \Delta P)(b_0), \bar{z}_0 - b_0 \rangle = O(\eta^2 + |\bar{z}_0 - b_0|^2). \quad (4.16)$$

So we can solve (4.15) and (4.16) to obtain $z_0 = x_{\eta,0}$ with $x_{\eta,0} \rightarrow b_0$ as $\eta \rightarrow 0$. \square

Now we are in a position to prove Theorem 1.4.

Proof of Theorem 1.4. Let w_λ be a single-peak solution as in Theorem 4.4, and we define

$$u_\lambda = \frac{w_\lambda}{\left(\int_{\mathbb{R}^3} w_\lambda^2\right)^{\frac{1}{2}}}.$$

Then $\int_{\mathbb{R}^3} u_\lambda^2 = 1$, and

$$-\Delta u_\lambda + P(x)u_\lambda = \frac{a_\lambda}{8\pi} \int_{\mathbb{R}^3} \frac{u_\lambda^2(y)}{|x-y|} dy u_\lambda(x) - \lambda u_\lambda, \quad \text{in } \mathbb{R}^3,$$

with $a_\lambda = \int_{\mathbb{R}^3} w_\lambda^2$.

Similar to (3.13), we can prove

$$\frac{1}{\sqrt{\lambda}} \int_{\mathbb{R}^3} w_\lambda^2 = a_* + o(1), \quad \text{as } \lambda \rightarrow +\infty.$$

Take $\lambda_0 > 0$ large and let $a_0 = \int_{\mathbb{R}^3} w_{\lambda_0}^2$. For any $a > 0$, let $f(\lambda) = \int_{\mathbb{R}^3} w_\lambda^2 - a$. Then for $a \geq a_0$, we have

$$f(\lambda_0) = a_0 - a \leq 0 \quad \text{and} \quad \lim_{\lambda \rightarrow +\infty} f(\lambda) = \lim_{\lambda \rightarrow +\infty} (\sqrt{\lambda}(a_* + o(1)) - a) = +\infty.$$

Hence by the continuity of the function $f(\lambda)$, for any $a \geq a_0$, there exists $\lambda = \lambda_a > 0$ large such that $f(\lambda_a) = 0$, i.e. $\int_{\mathbb{R}^3} w_\lambda^2 = a$, which yields that there exists $\lambda = \lambda_a > 0$ large such that the solution u_a of (1.11) with $\lambda = \lambda_a$ satisfies $\int_{\mathbb{R}^3} w_\lambda^2 = a$. Thus, for such a , we obtain a single-peak solution for (1.8), where $\mu_a = -\lambda_a$. \square

5. LOCAL UNIQUENESS OF SINGLE PEAK SOLUTIONS

From Lemma 3.4, a single-peak solution \tilde{u}_a to (3.1) can be written as

$$\tilde{u}_a(x) = U_{\epsilon, x_a} + \varphi_a(x), \quad (5.1)$$

with $|x_a - b_0| = o(1)$, $\epsilon = \frac{1}{\sqrt{-\mu_a}}$, $\varphi_a \in E_{a, x_a}$ and

$$\|\varphi_a\|_a = O\left(|P(x_a) - P_0|\epsilon^{\frac{7}{2}} + |\nabla P(x_a)|\epsilon^{\frac{9}{2}} + \epsilon^{\frac{11}{2}}\right). \quad (5.2)$$

Also we know $x_a \in \Gamma_{t_a}$ for some $t_a \rightarrow P_0$. Similar to the last section, we use ν_a to denote the unit normal vector of Γ_{t_a} at x_a , while we use $\tau_{a,j}$ to denote the principal direction of Γ_{t_a} at x_a . Then, at x_a , it holds

$$D_{\tau_{a,j}}P(x_a) = 0, \quad |\nabla P(x_a)| = |D_{\nu_a}P(x_a)|. \quad (5.3)$$

We first prove the following result.

Lemma 5.1. *Under the assumption (P), we have*

$$D_{\nu_a}P(x_a) = O(\epsilon^2). \quad (5.4)$$

Proof. We use (4.1) to obtain

$$\int_{B_\rho(x_a)} D_{\nu_a}P(x)\tilde{u}_a^2 = O(e^{-\frac{\rho}{\epsilon}}). \quad (5.5)$$

Then by (5.1)–(5.3) and (5.5), we get

$$\begin{aligned} & \int_{B_\rho(x_a)} D_{\nu_a}P(x)U_{\epsilon, x_a}^2 \\ &= -2 \int_{B_\rho(x_a)} D_{\nu_a}P(x)U_{\epsilon, x_a}\varphi_a - \int_{B_\rho(x_a)} D_{\nu_a}P(x)\varphi_a^2 + O(e^{-\frac{\rho}{\epsilon}}) \\ &= O(|D_{\nu_a}P(x_a)|\epsilon^{\frac{3}{2}} \cdot \|\varphi_a\|_a + \epsilon^{\frac{5}{2}}\|\varphi_a\|_a + \|\varphi_a\|_a^2) + O(e^{-\frac{\rho}{\epsilon}}) \\ &= O(|P(x_a) - P_0|\epsilon^5 + |D_{\nu_a}P(x_a)|\epsilon^6 + \epsilon^7). \end{aligned} \quad (5.6)$$

On the other hand, by Taylor's expansion, we have

$$\int_{B_\rho(x_a)} D_{\nu_a}P(x)U_{\epsilon, x_a}^2 = \epsilon^3 \left[a_* D_{\nu_a}P(x_a) + \frac{B\epsilon^2}{2} \Delta D_{\nu_a}P(x_a) + O(\epsilon^4) \right], \quad (5.7)$$

where B is the constant in (4.14). And then (5.4) follows from (5.6) and (5.7). \square

Let $\bar{x}_a \in \Gamma$ be the point such that $x_a - \bar{x}_a = \beta_a \nu_a$ for some $\beta_a \in \mathbb{R}$. Then we can prove

Lemma 5.2. *If the assumption (P) holds, then we have*

$$\begin{cases} \bar{x}_a - b_0 = L\epsilon^2 + O(\epsilon^4), \\ x_a - \bar{x}_a = -\frac{B}{2a_*} \frac{\partial \Delta P(b_0)}{\partial \nu} \left(\frac{\partial^2 P(b_0)}{\partial \nu^2} \right)^{-1} \epsilon^2 + O(\epsilon^4), \end{cases} \quad (5.8)$$

where B is the constant in (4.14) and L is a vector depending on b_0 .

Proof. It follows from (5.6) and (5.7) that

$$\begin{aligned} & (a_* + O(\epsilon^2)) D_{\nu_a} P(x_a) + \frac{B\epsilon^2}{2} \Delta D_{\nu_a} P(x_a) \\ & = O(\epsilon^4 + \epsilon^2 |P(x_a) - P_0|) = O(\epsilon^4 + \epsilon^2 |x_a - \bar{x}_a|^2). \end{aligned} \quad (5.9)$$

Since $\frac{\partial^2 P(b_0)}{\partial \nu^2} \neq 0$, the outward unit normal vector $\nu_a(x)$ and the tangential unit vector $\tau_a(x)$ of Γ_{t_a} at x_a are Lip-continuous in W_δ , then from (5.9), we find

$$x_a - \bar{x}_a = -\frac{B}{2a_*} (\Delta D_\nu P(b_0)) \left(\frac{\partial^2 P(b_0)}{\partial \nu^2} \right)^{-1} \epsilon^2 + O(\epsilon^4 + \epsilon^2 |\bar{x}_a - b_0|^2). \quad (5.10)$$

Then (5.2) and (5.10) implies

$$\|\varphi_a\|_a = O\left(|x_a - \bar{x}_a|^2 \epsilon^{\frac{7}{2}} + \epsilon^{\frac{11}{2}}\right) = O(\epsilon^{\frac{11}{2}}). \quad (5.11)$$

Recall $G(x) = \langle \nabla P(x), \tau_a \rangle$. Then $G(x_a) = 0$. Similar to (4.2) and (5.6), we have

$$\begin{aligned} \int_{B_\rho(x_a)} G(x) U_{\epsilon, x_a}^2 &= -2 \int_{B_\rho(x_a, i)} G(x) U_{\epsilon, x_a} \varphi_a - \int_{B_\rho(x_a)} G(x) \varphi_a^2 + O(e^{-\frac{\theta}{\epsilon}}) \\ &= -2 \int_{B_\rho(x_a, i)} G(x) U_{\epsilon, x_a} \varphi_a + O(\|\varphi_a\|_a^2) + O(e^{-\frac{\theta}{\epsilon}}) \\ &= -2 \int_{B_\rho(x_a)} \langle \nabla G(x_a), x - x_a \rangle U_{\epsilon, x_a} \varphi_a + O(\epsilon^9). \end{aligned} \quad (5.12)$$

On the other hand, in view of $\nabla P(x) = 0$, $x \in \Gamma$, we find

$$\nabla G(x_a) = \langle \nabla^2 P(x_a), \tau_a \rangle = \langle \nabla^2 P(\bar{x}_a), \bar{\tau}_a \rangle + O(|x_a - \bar{x}_a|) = O(|x_a - \bar{x}_a|), \quad (5.13)$$

where $\bar{x}_a \in \Gamma$ is the point such that $x_a - \bar{x}_a = \beta_a \nu_a$ for some $\beta_a \in \mathbb{R}$, and $\bar{\tau}_a$ is the tangential vector of Γ at $\bar{x}_a \in \Gamma$. Therefore, from (5.10), (5.11) and (5.13), we know

$$\begin{aligned} & \int_{B_\rho(x_a, i)} \langle \nabla G(x_a), x - x_a \rangle U_{\epsilon, x_a} \varphi_a \\ & = O(\epsilon^{\frac{5}{2}} \|\nabla G(x_a)\| \|\varphi_a\|_a) = O(|x_a - \bar{x}_a| \epsilon^8) = O(\epsilon^{10}). \end{aligned} \quad (5.14)$$

Then by (5.12) and (5.14), we find

$$\int_{B_\rho(x_a)} G(x) U_{\epsilon, x_a}^2 = O(\epsilon^9). \quad (5.15)$$

On the other hand, by the Taylor's expansion, we can prove

$$\int_{B_d(x_a)} G(x) U_{\epsilon, x_a}^2 = \left[\frac{B\epsilon^5}{2} (1 + P_0 \epsilon^2)^{\frac{1}{2}} \right] (D_{\tau_a} \Delta P)(x_a) + \frac{H_\tau \epsilon^7}{24} + O(\epsilon^9), \quad (5.16)$$

where

$$H_{\tau_i} = \sum_{l=1}^2 \sum_{m=1}^2 \frac{\partial^4 G(b_0)}{\partial x_l^2 \partial x_m^2} \int_{\mathbb{R}^N} x_l^2 x_m^2 U^2.$$

So (5.15) and (5.16) give

$$(D_{\tau_a} \Delta P)(x_a) = -\frac{H_{\tau} \epsilon^2}{12B} + O(\epsilon^4). \quad (5.17)$$

We denote by $\bar{\tau}_a$ the tangential vector of Γ at \bar{x}_a . Then by (5.10), we get

$$\begin{aligned} (D_{\tau_a} \Delta P)(x_a) &= (D_{\bar{\tau}_a} \Delta P)(\bar{x}_a) + \langle A_{\tau}, x_a - \bar{x}_a \rangle + O(|x_a - \bar{x}_a|^2) \\ &= (D_{\bar{\tau}_a} \Delta P)(\bar{x}_a) + B_{\tau} \epsilon^2 + O(\epsilon^4), \end{aligned}$$

where A_{τ} is a vector depending on b_0 and B_{τ} is a constant depending on b_0 . Moreover,

$$(D_{\bar{\tau}_a} \Delta P)(\bar{x}_a) = \left(D_{\tau}^2(\Delta P)(b_0) \right) (\bar{x}_a - b_0) + O(|\bar{x}_a - b_0|^2). \quad (5.18)$$

Therefore, from (5.17)–(5.18), we find

$$D_{\tau_i}^2(\Delta P)(b_0)(\bar{x}_a - b_0) = -\left(\frac{H_{\tau}}{12B} + B_{\tau} \right) \epsilon^2 + O(\epsilon^4) + O(|\bar{x}_a - b_0|^2). \quad (5.19)$$

Since $D_{\tau}^2(\Delta P)(b_0)$ is non-singular, we can complete the proofs of (5.8) from (5.10) and (5.19). \square

Let

$$\delta_a := \frac{a_*}{a}.$$

Proposition 5.3. *Under the assumption (P), there holds*

$$-\mu_a \delta_a^2 = 1 + \gamma_1 \delta_a^2 + O(\delta_a^4), \quad (5.20)$$

and

$$x_a - b_0 = \bar{L} \delta_a^2 + O(\delta_a^4), \quad (5.21)$$

where γ_1 and the vector \bar{L} are constants.

Proof. First, (3.13) shows that (5.20) holds. Then we can find (5.21) by (5.8) and (5.20). \square

Let $u(x) \mapsto a^{-\frac{1}{2}} \delta_a^{-2} u(x)$. Then the problem (1.8)–(1.9) can be changed into the following problem

$$-\delta_a^2 \Delta u + (-\mu_a \delta_a^2 + \delta_a^2 P(x)) u = \frac{1}{8\pi \delta_a^2} \int_{\mathbb{R}^3} \frac{u^2(y)}{|x-y|} dy u(x), \quad u \in H^1(\mathbb{R}^3), \quad (5.22)$$

and

$$\int_{\mathbb{R}^3} u^2 = a \delta_a^4. \quad (5.23)$$

Then similar to Lemma 3.4, the single-peak solution of (5.22)–(5.23) concentrating at b_0 can be written as $\tilde{U}_{\delta_a, x_a} + \tilde{\varphi}_a(x)$, with $|x_a - b_0| = o(1)$, $\|\tilde{\varphi}_a\|_{\delta_a} = o(\delta_a^{\frac{3}{2}})$, and

$$\tilde{\varphi}_a \in \tilde{E}_{a, x_a} := \left\{ \varphi \in H^1(\mathbb{R}^3) : \left\langle \varphi, \frac{\partial \tilde{U}_{\delta_a, x_a}}{\partial x_j} \right\rangle_{\delta_a} = 0, \quad j = 1, 2, 3 \right\},$$

where $\tilde{U}_{\delta_a, x_a} := (1 + (\gamma_1 + P_0)\delta_a^2)U\left(\frac{\sqrt{1+(\gamma_1+P_0)\delta_a^2}(x-x_a)}{\delta_a}\right)$, $\|\varphi\|_{\delta_a}^2 := \int_{\mathbb{R}^3} (\delta_a^2 |\nabla \varphi|^2 + \varphi^2)$ and γ_1 is the constant in (5.20). Then we can write the equation (5.22) as follows:

$$\bar{L}_a(\tilde{\varphi}_a) = \mathcal{R}_{a, \delta_a}(\tilde{\varphi}_a) + \bar{\mathcal{L}}_a(x),$$

where $\mathcal{R}_{a, \delta_a}$ is defined by (3.3),

$$\begin{aligned} \bar{L}_a(\tilde{\varphi}_a) := & -\delta_a^2 \Delta \tilde{\varphi}_a + \left((-\mu_a \delta_a^2 + \delta_a^2 P(x)) \tilde{\varphi}_a - \frac{1}{4\pi \delta_a^2} \int_{\mathbb{R}^3} \frac{\tilde{U}_{\delta_a, x_a}(y) \tilde{\varphi}_a(y)}{|x-y|} dy \tilde{U}_{\delta_a, x_a}(x) \right. \\ & \left. - \frac{1}{8\pi \delta_a^2} \int_{\mathbb{R}^3} \frac{(\tilde{U}_{\delta_a, x_a}(y))^2}{|x-y|} dy \tilde{\varphi}_a(x) \right) \end{aligned}$$

and

$$\tilde{\mathcal{L}}_a = - \sum_{i=1}^m \left(-\mu_a \delta_a^2 - (1 + \gamma_1 \delta_a^2) + (P(x) - P_0) \delta_a^2 \right) \tilde{U}_{\delta_a, x_a}.$$

Lemma 5.4. *There holds*

$$\|\tilde{\varphi}_a\|_{\delta_a} = O\left(\delta_a^{\frac{11}{2}}\right). \quad (5.24)$$

Proof. The proofs are similar to that of Lemma 3.4, the difference is

$$\|\tilde{\mathcal{L}}_a\|_{\delta_a} = O\left(|P(x_a) - P_0| \delta_a^{\frac{7}{2}} + |\nabla P(x_a)| \delta_a^{\frac{9}{2}} + \delta_a^{\frac{11}{2}}\right) = O\left(\delta_a^{\frac{11}{2}}\right). \quad (5.25)$$

Similar to Lemma 3.1, we can also check that $\tilde{\mathcal{L}}_a$ is invertible in \tilde{E}_{a, x_a} . Finally, (5.25) and the contradiction mapping theorem imply (5.24). \square

For simplicity of notations, hereafter we denote $\mathbf{p}_0 := 1 + (\gamma_1 + P_0)\delta_a^2$. Hence

$$\tilde{U}_{\delta_a, x_a} := \mathbf{p}_0 U\left(\frac{\sqrt{\mathbf{p}_0}(x-x_a)}{\delta_a}\right).$$

Let $u_a^{(1)}$ and $u_a^{(2)}$ be two single-peak solutions of (5.22)–(5.23) concentrating at some point b_0 , which can be written as

$$u_a^{(l)} = \tilde{U}_{\delta_a, x_a^{(l)}} + \tilde{\varphi}_a^{(l)}(x), \text{ for } l = 1, 2, \text{ and } \tilde{\varphi}_a^{(l)} \in \tilde{E}_{a, x_a^{(l)}}.$$

Now we set $\xi_a(x) = \frac{u_a^{(1)}(x) - u_a^{(2)}(x)}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^3)}}$. Then $\xi_a(x)$ satisfies $\|\xi_a\|_{L^\infty(\mathbb{R}^3)} = 1$. And from (5.22), we find that ξ_a satisfies

$$-\delta_a^2 \Delta \xi_a(x) + C_a(x) \xi_a(x) - D_a(x) \xi_a(x) - E_a(x) = g_a(x),$$

where

$$\begin{aligned} C_a(x) &= \delta_a^2 P(x) - \delta_a^2 \mu_a^{(1)}, \quad D_a(x) = \frac{1}{8\pi \delta_a^2} \int_{\mathbb{R}^3} \frac{(u_a^{(1)})^2(y)}{|x-y|} dy, \\ E_a(x) &= \frac{u_a^{(2)}(x)}{8\pi \delta_a^2} \int_{\mathbb{R}^3} \frac{(u_a^{(1)} + u_a^{(2)}) \xi_a(y)}{|x-y|} dy, \quad g_a(x) = \frac{\delta_a^2 (\mu_a^{(1)} - \mu_a^{(2)})}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^3)}} u_a^{(2)}(x). \end{aligned}$$

Also, similar to (3.11), for any fixed $R \gg 1$, there exist some $\theta > 0$ and $C > 0$, such that

$$|u_a^{(l)}(x)| + |\nabla u_a^{(l)}(x)| \leq C e^{-\theta|x-x_a|/\delta_a}, \text{ for } l = 1, 2, \quad x \in \mathbb{R}^3 \setminus B_{R\delta_a}(x_a). \quad (5.26)$$

Now let $\bar{\xi}_a(x) = \xi_a(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}x + x_a^{(1)})$, we have

$$-\Delta \bar{\xi}_a(x) + \frac{C_a(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}x + x_a^{(1)})}{\mathbf{p}_0} \bar{\xi}_a(x) - \frac{\bar{D}_a(x)}{\mathbf{p}_0} \bar{\xi}_a(x) - \frac{\bar{E}_a(x)}{\mathbf{p}_0} = \frac{g_a(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}x + x_a^{(1)})}{\mathbf{p}_0}, \quad (5.27)$$

where

$$\begin{aligned} \bar{D}_a(x) &= \frac{1}{8\pi\mathbf{p}_0} \int_{\mathbb{R}^3} \frac{(u_a^{(1)})^2(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}y + x_a^{(1)})}{|x-y|} dy, \\ \bar{E}_a(x) &= \frac{u_a^{(2)}(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}x + x_a^{(1)})}{8\pi\mathbf{p}_0} \int_{\mathbb{R}^3} \frac{(u_a^{(1)} + u_a^{(2)})(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}y + x_a^{(1)})\bar{\xi}_a(y)}{|x-y|} dy. \end{aligned}$$

Lemma 5.5. For $x \in B_{\rho\sqrt{\mathbf{p}_0}\delta_a^{-1}}(0)$, it holds

$$\frac{\bar{D}_a(x)}{\mathbf{p}_0} = \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy + O(\delta_a^4),$$

and

$$\frac{\bar{E}_a(x)}{\mathbf{p}_0} = \frac{U(x)}{4\pi} \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy + O\left(\delta_a + \tilde{\varphi}_a^2\left(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)}\right)\right).$$

Proof. By direct computations, from (C.4) we have

$$\begin{aligned} \frac{\bar{D}_a(x)}{\mathbf{p}_0} &= \frac{1}{8\pi\mathbf{p}_0^2} \int_{\mathbb{R}^3} \frac{(u_a^{(1)})^2(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}y + x_a^{(1)})}{|x-y|} dy \\ &= \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy + \underbrace{\frac{1}{4\pi\mathbf{p}_0} \int_{\mathbb{R}^3} \frac{U(y)\tilde{\varphi}_a^{(1)}(\frac{\delta_a y}{\sqrt{\mathbf{p}_0}} + x_a^{(1)})}{|x-y|} dy}_{:=F_1} \\ &\quad + \underbrace{\frac{1}{8\pi\mathbf{p}_0^2} \int_{\mathbb{R}^3} \frac{(\tilde{\varphi}_a^{(1)}(\frac{\delta_a y}{\sqrt{\mathbf{p}_0}} + x_a^{(1)}))^2}{|x-y|} dy}_{:=F_2} \\ &= \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(x)}{|x-y|} dy + O(\delta_a). \end{aligned}$$

Also, it follows from (C.7) and (5.21) that

$$\begin{aligned} \frac{\bar{E}_a(x)}{\mathbf{p}_0} &= \frac{u_a^{(2)}(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)})}{8\pi\mathbf{p}_0} \int_{\mathbb{R}^3} \frac{(u_a^{(1)} + u_a^{(2)})(\frac{\delta_a y}{\sqrt{\mathbf{p}_0}} + x_a^{(1)})\bar{\xi}_a(y)}{|x-y|} dy \\ &= \frac{U(x)}{4\pi} \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy \\ &\quad + \underbrace{\frac{1}{4\pi\mathbf{p}_0} (u_a^{(2)}(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)}) - \mathbf{p}_0 U(x)) \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy}_{:=G_1} \end{aligned}$$

$$\begin{aligned}
& + \underbrace{\frac{u_a^{(2)}(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)})}{8\pi \mathbf{p}_0} \int_{\mathbb{R}^3} \frac{((u_a^{(1)} + u_a^{(2)})(\frac{\delta_a y}{\sqrt{\mathbf{p}_0}} + x_a^{(1)}) - 2\mathbf{p}_0 U(y)) \bar{\xi}_a(y)}{|x-y|} dy}_{:=G_2} \\
& = \frac{U(x)}{4\pi} \int_{\mathbb{R}^3} \frac{U(y) \bar{\xi}_a(y)}{|x-y|} dy + O\left(\delta_a + \tilde{\varphi}_a^2\left(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)}\right)\right).
\end{aligned}$$

□

Lemma 5.6. For $x \in B_{\rho\sqrt{\mathbf{p}_0}\delta_a^{-1}}(0)$, it holds

$$\frac{C_a(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)})}{\mathbf{p}_0} = 1 + O\left(\delta_a^4 + \sum_{l=1}^2 \tilde{\varphi}_a^{(l)}\left(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)}\right)\right), \quad (5.28)$$

and

$$\frac{g_a(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)})}{\mathbf{p}_0} = -\frac{1}{4\pi a_*} U(x) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U(y)\bar{\xi}_a(y)}{|x-y|} dx dy + O\left(\delta_a + \sum_{l=1}^2 \tilde{\varphi}_a^{(l)}\left(\frac{\delta_a x}{\sqrt{\mathbf{p}_0}} + x_a^{(1)}\right)\right). \quad (5.29)$$

Proof. First, (5.28) can be deduced by (5.20) and (5.21) directly. Now we prove (5.29).

From (5.22) and (5.23), for $l = 1, 2$, we find

$$a\mu_a^{(l)}\delta_a^6 = \delta_a^2 \int_{\mathbb{R}^3} (|\nabla u_a^{(l)}|^2 + P(x)(u_a^{(l)})^2) - \frac{1}{8\pi\delta_a^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(u_a^{(l)})^2(x)(u_a^{(l)})^2(y)}{|x-y|} dx dy,$$

which gives

$$\begin{aligned}
& \frac{a\delta_a^6(\mu_a^{(1)} - \mu_a^{(2)})}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^3)}} \\
& = -\mu_a^{(2)}\delta_a^2 \int_{\mathbb{R}^3} (u_a^{(1)} + u_a^{(2)})\xi_a + \delta_a^2 \int_{\mathbb{R}^3} (\nabla(u_a^{(1)} + u_a^{(2)}) \cdot \nabla\xi_a + P(x)(u_a^{(1)} + u_a^{(2)})\xi_a) \\
& \quad - \frac{1}{8\pi\delta_a^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(u_a^{(l)})^2(x)(u_a^{(1)} + u_a^{(2)})\xi_a(y) + (u_a^{(1)} + u_a^{(2)})\xi_a(x)(u_a^{(2)})^2(y)}{|x-y|} dx dy \\
& = -(\mu_a^{(2)} - \mu_a^{(1)})\delta_a^2 \int_{\mathbb{R}^3} u_a^{(1)}\xi_a - \frac{1}{8\pi\delta_a^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{[(u_a^{(1)})^2(x)u_a^{(2)}(y) + (u_a^{(2)})^2(x)u_a^{(1)}(y)]\xi_a(y)}{|x-y|} dx dy, \quad (5.30)
\end{aligned}$$

here we use the following identity:

$$\int_{\mathbb{R}^3} (u_a^{(1)} + u_a^{(2)})\xi_a = \frac{1}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^3)}} \left(\int_{\mathbb{R}^3} (u_a^{(1)})^2 - \int_{\mathbb{R}^3} (u_a^{(2)})^2 \right) = 0.$$

Then from (5.20), (5.21), (5.24) and (5.30), we know

$$\begin{aligned} \frac{\delta_a^2(\mu_a^{(1)} - \mu_a^{(2)})}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^3)}} &= \frac{1}{a_*\delta_a^3}(\mu_a^{(2)} - \mu_a^{(1)})\delta_a^2 \int_{\mathbb{R}^3} u_a^{(1)}\xi_a \\ &\quad - \frac{1}{8\pi a_*\delta_a^5} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{[(u_a^{(1)})^2(x)u_a^{(2)}(y) + (u_a^{(2)})^2(x)u_a^{(1)}(y)]\xi_a(y)}{|x-y|} dx dy. \end{aligned}$$

By (5.20), Hölder inequality and (5.24), we have

$$\begin{aligned} &\frac{1}{a_*\delta_a^3}(\mu_a^{(2)} - \mu_a^{(1)})\delta_a^2 \int_{\mathbb{R}^3} u_a^{(1)}\xi_a \\ &= \frac{-1}{a_*\delta_a^3}(\mu_a^{(2)} - \mu_a^{(1)})\delta_a^2 \left(\int_{\mathbb{R}^3} \mathbf{p}_0 U\left(\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x - x_a^{(1)})\right)\xi_a + \int_{\mathbb{R}^3} \tilde{\varphi}_a^{(1)}\xi_a \right) \\ &= \frac{1}{a_*\delta_a^3}O(\delta_a^4)(C\delta_a^{\frac{3}{2}} + C\delta_a^{\frac{11}{2}}) = O(\delta_a^{\frac{5}{2}}). \end{aligned}$$

From (C.12), we can check that

$$\begin{aligned} &\frac{1}{8\pi a_*\delta_a^5} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{[(u_a^{(1)})^2(x)u_a^{(2)}(y) + (u_a^{(2)})^2(x)u_a^{(1)}(y)]\xi_a(y)}{|x-y|} dx dy \\ &= \frac{1}{8\pi a_*\delta_a^5} \left(2\delta_a^5 \mathbf{P}_0^{\frac{1}{2}} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U(y)\bar{\xi}_a(y)}{|x-y|} dx dy + O(\delta_a^6) \right) \\ &= \frac{1}{4\pi a_*} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U(y)\bar{\xi}_a(y)}{|x-y|} dx dy + O(\delta_a^6). \end{aligned}$$

Noting that in $x \in B_{\rho\sqrt{\mathbf{P}_0}\delta_a^{-1}}(0)$, by the mean value theorem and (5.21) we have

$$\begin{aligned} &u_a^{(2)}\left(\frac{\delta_a x}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right) \\ &= \mathbf{p}_0 U(x) + \mathbf{p}_0 \left(U\left(x + \frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x_a^{(1)} - x_a^{(2)})\right) - U(x) \right) + \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a x}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right) \\ &= \mathbf{p}_0 U(x) + \mathbf{p}_0 \nabla U\left(\zeta x + (1-\zeta)\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x_a^{(1)} - x_a^{(2)})\right) \cdot \frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x_a^{(1)} - x_a^{(2)}) + \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a x}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right) \\ &= \mathbf{p}_0 U(x) + O(\delta_a) + \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a x}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right), \end{aligned}$$

which combining all the estimates above implies that (5.29) holds. \square

Then from Lemmas 5.5 and 5.6, we have the following result.

Lemma 5.7. *From $|\bar{\xi}_a| \leq 1$, we suppose that $\bar{\xi}_a(x) \rightarrow \xi(x)$ in $C_{loc}^1(\mathbb{R}^3)$. Then $\xi(x)$ satisfies following system:*

$$\begin{aligned} &-\Delta\xi(x) + \xi(x) - \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy \xi(x) - \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{U(y)\xi(y)}{|x-y|} dy U(x) \\ &= -\frac{1}{4\pi a_*} U(x) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U(y)\xi(y)}{|x-y|} dx dy. \end{aligned}$$

To prove $\xi = 0$, we write

$$\bar{\xi}_a(x) = \sum_{j=0}^3 \beta_{a,j} \psi_j + \tilde{\xi}_a(x), \text{ in } H^1(\mathbb{R}^3), \quad (5.31)$$

where $\psi_j (j = 0, 1, 2, 3)$ are the functions in (B.3) and $\tilde{\xi}_a(x) \in \tilde{E}$ with

$$\tilde{E} = \{u \in H^1(\mathbb{R}^3), \langle u, \psi_j \rangle = 0, \text{ for } j = 0, 1, 2, 3\}.$$

It is standard to prove the following result.

Lemma 5.8. *For any $u \in \tilde{E}$, there exists $\bar{\gamma} > 0$ such that*

$$\|\tilde{L}(u)\| \geq \bar{\gamma} \|u\|,$$

where \tilde{L} is defined by

$$\begin{aligned} \tilde{L}(u) := & -\Delta u(x) + u(x) - \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy u(x) - \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{U(y)u(y)}{|x-y|} dy U(x) \\ & + \frac{1}{4\pi a_*} U(x) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U(y)u(y)}{|x-y|} dx dy. \end{aligned}$$

Proposition 5.9. *Let $\tilde{\xi}_a(x)$ be as in (5.31). Then*

$$\|\tilde{\xi}_a\| = O(\delta_a). \quad (5.32)$$

Proof. First, Lemma 5.8 gives

$$\|\tilde{\xi}_a\| \leq C \|\tilde{L}(\tilde{\xi}_a)\|. \quad (5.33)$$

On the other hand, from (5.27)–(5.31), we can prove

$$\begin{aligned} \tilde{L}(\tilde{\xi}_a) = & \left(1 - \frac{C_a(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}x + x_a^{(1)})}{\mathbf{p}_0}\right) \bar{\xi}_a(x) + \left(-\frac{\bar{D}_a(x)}{\mathbf{p}_0} \bar{\xi}_a(x) - \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy \bar{\xi}_a(x)\right) \\ & + \left(-\frac{\bar{E}_a(x)}{\mathbf{p}_0} \bar{\xi}_a(x) - \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy U(x)\right) \\ & + \left(\frac{g_a(\frac{\delta_a}{\sqrt{\mathbf{p}_0}}x + x_a^{(1)})}{\mathbf{p}_0} \bar{\xi}_a(x) + \frac{1}{4\pi a_*} U(x) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U(y)\bar{\xi}_a(y)}{|x-y|} dx dy\right). \end{aligned} \quad (5.34)$$

So from (5.24), (5.33), (5.34), Lemma 5.5 and Lemma 5.6, we have for any $v(x) \in H^1(\mathbb{R}^3)$

$$\begin{aligned} \langle \tilde{L}(\tilde{\xi}_a), v \rangle &= O(\delta_a) \|v\| + O\left(\int_{\mathbb{R}^3} \sum_{l=1}^2 \varphi_a^{(l)}(\delta_a x + x_{a,i}^{(1)}) |v|\right) \\ &\leq O(\delta_a) \|v\| + C \delta_a^{-\frac{3}{2}} \|\varphi_a^{(l)}\|_{\delta_a} \|v\| \\ &\leq O(\delta_a) \|v\|, \end{aligned}$$

which implies (5.32). □

Letting $\beta_{a,j}$ be as in (5.31) and using $|\bar{\xi}_a| \leq 1$, we find

$$\beta_{a,j} = \frac{\langle \bar{\xi}_a, \psi_j \rangle}{\|\psi_j\|^2} = O(\|\bar{\xi}_a\|) = O(1), \quad j = 0, 1, 2, 3.$$

Lemma 5.10. *There holds*

$$\beta_{a,0} = o(1). \quad (5.35)$$

Proof. On one hand, from (5.31), (5.32) and (5.21) we get

$$\begin{aligned} & \int_{B_\rho(x_a^{(1)})} (u_a^{(1)} + u_a^{(2)}) \xi_a \\ &= \frac{\delta_a^3}{\mathbf{P}_0^{\frac{3}{2}}} \int_{B_{\frac{\sqrt{\mathbf{P}_0}}{\delta_a} \rho(0)}} \left(2\mathbf{p}_0 U(x) + \nabla U(\zeta x + (1-\zeta) \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)})) \right) \cdot \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)}) \\ & \quad + \sum_{l=1}^2 \tilde{\varphi}^{(l)} \left(\frac{\sqrt{\mathbf{P}_0}}{\delta_a} x + x_a^{(1)} \right) \bar{\xi}_a(x) dx \\ &= 2 \frac{\delta_a^3}{\sqrt{\mathbf{P}_0}} \int_{\mathbb{R}^3} U(x) \bar{\xi}_a(x) - \int_{\mathbb{R}^3 \setminus B_{\frac{\sqrt{\mathbf{P}_0}}{\delta_a} \rho(0)}} U(x) \bar{\xi}_a(x) \\ & \quad + O(\delta_a^2 |x_a^{(1)} - x_a^{(2)}|) \left| \nabla U(\zeta x + (1-\zeta) \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)})) \right|_2 |\bar{\xi}_a(x)|_2 \\ & \quad + \frac{\delta_a^3}{\mathbf{P}_0^{\frac{3}{2}}} \sum_{l=1}^2 \left| \tilde{\varphi}^{(l)} \left(\frac{\sqrt{\mathbf{P}_0}}{\delta_a} x + x_a^{(1)} \right) \right|_2 |\bar{\xi}_a(x)|_2 \\ &= 2 \frac{\delta_a^3}{\sqrt{\mathbf{P}_0}} \left(\int_{\mathbb{R}^3} U(x) \beta_{a,0} (2U(x) + x \cdot \nabla U(x)) + \int_{\mathbb{R}^3} U(x) \tilde{\xi}_a(x) \right) + O(e^{-\frac{\theta}{\delta_a}}) \\ & \quad + O(|x_a^{(1)} - x_a^{(2)}| \delta_a^2) + O(\delta_a^{\frac{3}{2}} \sum_{l=1}^2 \|\tilde{\varphi}_a^{(l)}\|_{\delta_a}) \\ &= \beta_{a,0} \frac{\delta_a^3}{\sqrt{\mathbf{P}_0}} \int_{\mathbb{R}^3} U^2(x) + O(\delta_a^4) + O(|x_a^{(1)} - x_a^{(2)}| \delta_a^2 + O(\delta_a^4) + \delta_a^{\frac{3}{2}} \|\tilde{\varphi}_a^{(2)}\|_{\delta_a}) \\ &= a_* \beta_{a,0} \delta_a^3 + O(\delta_a^4). \end{aligned} \quad (5.36)$$

On the other hand, noting that from (5.26)

$$\begin{aligned} \int_{\mathbb{R}^3 \setminus B_\rho(x_a^{(1)})} (u_a^{(1)} + u_a^{(2)}) \xi_a &= O \left(\int_{\mathbb{R}^3 \setminus B_\rho(x_a^{(1)})} |u_a^{(1)} + u_a^{(2)}| \right) \\ &= O \left(\int_{\mathbb{R}^3 \setminus B_\rho(x_a^{(1)})} |u_a^{(1)}| \right) + O \left(\int_{\mathbb{R}^3 \setminus B_{\frac{\rho}{2}}(x_a^{(2)})} |u_a^{(2)}| \right) \\ &= O(e^{-\frac{\theta}{\delta_a}}), \end{aligned}$$

then we can have

$$\begin{aligned} \int_{B_\rho(x_a^{(1)})} (u_a^{(1)} + u_a^{(2)}) \xi_a &= \int_{\mathbb{R}^3} (u_a^{(1)} + u_a^{(2)}) \xi_a + O(e^{-\frac{\theta}{\delta_a}}) \\ &= \frac{1}{\|u_a^{(1)} - u_a^{(2)}\|_{L^\infty(\mathbb{R}^3)}} \int_{\mathbb{R}^3} [(u_a^{(1)})^2 - (u_a^{(2)})^2] + O(e^{-\frac{\theta}{\delta_a}}) \\ &= O(e^{-\frac{\theta}{\delta_a}}), \end{aligned} \quad (5.37)$$

since

$$\int_{\mathbb{R}^3} (u_a^{(1)})^2 = \int_{\mathbb{R}^3} (u_a^{(2)})^2 = 1.$$

It follows from (5.36) and (5.37) that (5.35) holds. \square

Remark 5.11. We would like to point out that since here we consider the single-peak case and observing this fact $\int_{\mathbb{R}^3} U(x)(2U(x) + x \cdot \nabla U(x))dx = \frac{1}{2} \int_{\mathbb{R}^3} U^2(x)dx \neq 0$, the proof of Proposition 5.10 is much simple.

Proposition 5.12. It holds

$$\beta_{a,j} = o(1), \quad j = 1, 2, 3. \quad (5.38)$$

Proof. Step 1: To prove $\beta_{a,3} = O(\delta_a)$.

Using (4.1), we deduce

$$\int_{B_\rho(x_a^{(1)})} \frac{\partial P(x)}{\partial \nu_a} B_a(x) \xi_a = O(e^{-\frac{\rho}{\delta_a}}), \quad (5.39)$$

where ν_a is the outward unit vector of $\partial B_d(x_a^{(1)})$ at x , $B_a(x) = \sum_{l=1}^2 u_a^{(l)}(x)$.

On the other hand, by (5.21), we have

$$B_a(x) = \left(2 + O(\delta_a^2)\right) \tilde{U}_{\delta_a, x_a^{(1)}}(x) + O\left(\sum_{l=1}^2 |\tilde{\varphi}_a^{(l)}(x)|\right), \quad x \in B_\rho(x_a^{(1)}). \quad (5.40)$$

Also, from (5.8), we find

$$\frac{\partial P(x_a^{(1)})}{\partial \nu_a} = \frac{\partial P(x_a^{(1)})}{\partial \nu_a} - \frac{\partial P(\bar{x}_a^{(1)})}{\partial \nu_a} = O(|x_a^{(1)} - \bar{x}_a^{(1)}|) = O(\delta_a^2),$$

and

$$\frac{\partial^2 P(x_a^{(1)})}{\partial \nu_a \partial \tau_{a,j}} = \frac{\partial^2 P(x_a^{(1)})}{\partial \nu_a \partial \tau_{a,j}} - \frac{\partial^2 P(\bar{x}_a^{(1)})}{\partial \nu_a \partial \tau_{a,j}} = O(|x_a^{(1)} - \bar{x}_a^{(1)}|) = O(\delta_a^2), \quad \text{for } j = 1, 2.$$

From (3.11), (4.1) and (5.40), we get

$$\begin{aligned} & \int_{B_\rho(x_a^{(1)})} \frac{\partial P(x)}{\partial \nu_a} B_a(x) \xi_a \\ &= \int_{\mathbb{R}^3} \frac{\partial P(x_a^{(1)})}{\partial \nu_a} B_a(x) \xi_a + \int_{\mathbb{R}^3} \left\langle \nabla \frac{\partial P(x_a^{(1)})}{\partial \nu_a}, x - x_a^{(1)} \right\rangle B_a(x) \xi_a + O(\delta_a^5) \\ &= -\frac{\partial^2 P(x_a^{(1)})}{\partial \nu_a^2} a_* \beta_{a,3} \delta_a^4 + O(\delta_a^5). \end{aligned} \quad (5.41)$$

Then (5.39) and (5.41) imply $\beta_{a,3} = O(\delta_a)$.

Step 2: To prove $\beta_{a,j} = o(1)$ for $j = 1, 2$.

Similar to (5.39), we have

$$\int_{B_\rho(x_a^{(1)})} \frac{\partial P(y)}{\partial \tau_{a,j}} B_a(y) \xi_a = O(e^{-\frac{\rho}{\delta_a}}), \quad \text{for } j = 1, 2. \quad (5.42)$$

Using suitable rotation, we assume that $\tau_{a,1} = (1, 0, 0)$, $\tau_{a,2} = (0, 1, 0)$ and $\nu_a = (0, 0, 1)$. Under the assumption (P), we obtain

$$\begin{aligned} & \frac{\partial P(\frac{\delta_a}{\sqrt{\mathbf{P}_0}}y + x_a^{(1)})}{\partial \tau_{a,j}} \\ &= \frac{\delta_a}{\sqrt{\mathbf{P}_0}} \sum_{l=1}^3 \frac{\partial^2 P(x_a^{(1)})}{\partial y_l \partial \tau_{a,j}} y_l + \frac{\delta_a^2}{2\mathbf{P}_0} \sum_{k=1}^3 \sum_{l=1}^3 \frac{\partial^3 P(x_a^{(1)})}{\partial y_k \partial y_l \partial \tau_{a,j}} y_k y_l \\ &+ \frac{\delta_a^3}{6\mathbf{P}_0^{\frac{3}{2}}} \sum_{s=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \frac{\partial^4 P(x_a^{(1)})}{\partial y_s \partial y_l \partial y_k \partial \tau_{a,i,j}} y_s y_l y_k + o(\delta_a^3 |y|^3), \text{ in } B_{\frac{\rho}{\delta_a} \sqrt{\mathbf{P}_0}}(0). \end{aligned} \quad (5.43)$$

By (1.7), (5.8), (5.24), (5.31), (5.40) and the symmetry of $\psi_j(x)$, we find, for $j = 1, 2$,

$$\begin{aligned} & \sum_{k=1}^3 \sum_{l=1}^3 \frac{\partial^3 P(x_a^{(1)})}{\partial y_k \partial y_l \partial \tau_{a,j}} \int_{B_{\frac{\rho}{\delta_a} \sqrt{\mathbf{P}_0}}(0)} B_a(\frac{\delta_a}{\sqrt{\mathbf{P}_0}}y + x_a^{(1)}) \bar{\xi}_a y_k y_l \\ &= 2 \sum_{k=1}^3 \sum_{l=1}^3 \frac{\partial^3 P(x_a^{(1)})}{\partial y_k \partial y_l \partial \tau_{a,j}} \int_{B_{\frac{\rho}{\delta_a} \sqrt{\mathbf{P}_0}}(0)} U(y) \bar{\xi}_{a,i} y_k y_l + O(\delta_a^2) \\ &= B \beta_{a,0} \frac{\partial \Delta P(x_a^{(1)})}{\partial \tau_{a,j}} + O(\delta_a^2) = O(|x_a^{(1)} - b_0|) + O(\delta_a^2) = O(\delta_a^2). \end{aligned}$$

Also from (5.38) and (5.40), we get

$$\begin{aligned} & \sum_{s=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \frac{\partial^4 P(x_a^{(1)})}{\partial y_s \partial y_l \partial y_k \partial \tau_{a,j}} \int_{B_{\rho \sqrt{\mathbf{P}_0} \delta_a^{-1}}(0)} B_a(\frac{\delta_a}{\sqrt{\mathbf{P}_0}}y + x_a^{(1)}) \bar{\xi}_a y_s y_l y_m \\ &= 2 \sum_{s=1}^3 \sum_{k=1}^3 \sum_{l=1}^3 \frac{\partial^4 P(x_a^{(1)})}{\partial y_s \partial y_l \partial y_k \partial \tau_{a,j}} \int_{B_{\rho \sqrt{\mathbf{P}_0} \delta_a^{-1}}(0)} U(y) \left(\sum_{h=1}^2 \beta_{a,h} \right) \psi_h(y) y_s y_l y_k + o(1) \\ &= 2 \sum_{h=1}^2 \beta_{a,j} \int_{B_{\rho \sqrt{\mathbf{P}_0} \delta_a^{-1}}(0)} U(y) \psi_h(y) y_h \left[\frac{\partial^4 P(x_a^{(1)})}{\partial y_h^3 \partial \tau_{a,j}} y_h^2 + 3 \sum_{l=1, l \neq h}^3 \frac{\partial^4 P(x_a^{(1)})}{\partial y_h \partial y_l^2 \partial \tau_{a,j}} y_l^2 \right] + o(1) \\ &= -3B \left(\sum_{h=1}^2 \frac{\partial^2 \Delta P(x_a^{(1)})}{\partial \tau_{a,h} \partial \tau_{a,j}} \beta_{a,h} \right) + o(1) = -3B \left(\sum_{h=1}^2 \frac{\partial^2 \Delta P(b_0)}{\partial \tau_{a,h} \partial \tau_{a,j}} \beta_{a,h} \right) + o(1). \end{aligned}$$

By (A.3), we estimate

$$\frac{\partial^2 P(x_a^{(1)})}{\partial y_l \partial \tau_{a,j}} = -\frac{\partial P(x_a^{(1)})}{\partial \nu_a} \kappa_l(x_a^{(1)}) \delta_{lj}, \quad l, j = 1, 2.$$

Since $\frac{\partial P(\bar{x}_a^{(1)})}{\partial \nu_a} = 0$, from (5.8), we find

$$\begin{aligned}
\frac{\partial^2 P(x_a^{(1)})}{\partial y_l \partial \tau_{a,j}} &= - \left(\frac{\partial P(x_a^{(1)})}{\partial \nu_a} - \frac{\partial P(\bar{x}_a^{(1)})}{\partial \nu_a} \right) \kappa_l(x_a^{(1)}) \delta_{lj} \\
&= - \frac{\partial^2 P(\bar{x}_a^{(1)})}{\partial \nu_a^2} (x_a^{(1)} - \bar{x}_a^{(1)}) \cdot \nu_a \kappa_l(x_a^{(1)}) \delta_{lj} + o(\delta_a^2) \\
&= - \frac{\partial^2 P(b_0)}{\partial \nu^2} (x_a^{(1)} - \bar{x}_a^{(1)}) \cdot \nu_a \kappa_l(b_0) \delta_{lj} + o(\delta_a^2) \\
&= \frac{B}{2a_*} \frac{\partial \Delta P(b_0)}{\partial \nu} \delta_a^2 \kappa_l(b_0) \delta_{lj} + o(\delta_a^2).
\end{aligned} \tag{5.44}$$

Therefore from (5.24), (5.31), (5.40) and (5.44), we get

$$\begin{aligned}
&\sum_{l=1}^3 \frac{\partial^2 P(x_a^{(1)})}{\partial y_l \partial \tau_{a,j}} \int_{B_{\frac{\rho}{\delta_a} \sqrt{\mathbf{P}_0}}(0)} B_a \left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)} \right) \bar{\xi}_a y_l \\
&= \frac{B}{2a_*} \frac{\partial \Delta P(b_0)}{\partial \nu} \delta_a^2 \kappa_j(b_0) \int_{B_{\frac{\rho}{\delta_a} \sqrt{\mathbf{P}_0}}(0)} B_a \left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)} \right) \bar{\xi}_a y_j + o(\delta_a^2) \\
&= \frac{B}{a_*} \frac{\partial \Delta P(b_0)}{\partial \nu_i} \delta_a^2 \kappa_j(b_0) \beta_{a,j} \int_{\mathbb{R}^3} U(y) \frac{U(y)}{\partial y_j} y_j + o(\delta_a^2) \\
&= - \frac{B}{2} \frac{\partial \Delta P(b_0)}{\partial \nu_i} \delta_a^2 \kappa_j(b_0) \beta_{a,j} + o(\delta_a^2).
\end{aligned} \tag{5.45}$$

Combining (5.43)–(5.45), we obtain

$$\begin{aligned}
&\int_{B_\rho(x_a^{(1)})} \frac{\partial P(y)}{\partial \tau_{a,j}} B_a(y) \xi_a \\
&= - \frac{B}{2} \frac{\partial \Delta P(b_0)}{\partial \nu} \delta_a^6 \kappa_j(b_0) \beta_{a,j} - \frac{B}{2} \left(\sum_{l=1}^2 \frac{\partial^2 \Delta P(b_0)}{\partial \tau_l \partial \tau_j} \beta_{a,j} \right) \delta_a^6 + o(\delta_a^6).
\end{aligned} \tag{5.46}$$

From (5.42) and (5.46), we find

$$\frac{\partial \Delta P(b_0)}{\partial \nu_i} \kappa_j(b_0) \beta_{a,j} + \left(\sum_{l=1}^2 \frac{\partial^2 \Delta P(b_0)}{\partial \tau_l \partial \tau_j} \beta_{a,l} \right) = o(1),$$

which and the assumption (\tilde{P}) implies $\beta_{a,j} = o(1)$ for $j = 1, 2$. □

Finally, we prove Theorem 1.5.

Proof of Theorem 1.5: First, for large fixed R , (5.20), (5.26), (A.1) and (A.2) give

$$C_a(x) - D_a(x) \geq \frac{1}{2}, \quad |g_a(x) + E_a(x)| \leq C e^{-\hat{\theta} R}, \quad x \in \mathbb{R}^3 \setminus B_{R\delta_a}(x_a^{(1)}),$$

for some $\hat{\theta} > 0$.

Using the comparison principle, we get

$$\xi_a(x) = o(1), \quad \text{in } \mathbb{R}^3 \setminus \bigcup B_{R\delta_a}(x_a^{(1)}).$$

On the other hand, it follows from (5.32), (5.35) and (5.38) that

$$\xi_a(x) = o(1), \text{ in } B_{R\delta_a}(x_a^{(1)}).$$

This is in contradiction with $\|\xi_a\|_{L^\infty(\mathbb{R}^3)} = 1$. So $u_a^{(1)}(x) \equiv u_a^{(2)}(x)$ for $a \nearrow +\infty$. \square

6. THE NON-EXISTENCE OF MULTI-PEAK SOLUTIONS

From (3.1), we know that in order to prove Theorem 1.7, it suffices to prove the following result.

Proposition 6.1. *Under the assumption (P), there exists a small constant $\epsilon_0 > 0$ such that problem (3.1) has no m -peak solutions ($m \geq 2$) of the form*

$$\tilde{u}_a(x) = \sum_{i=1}^m U_{\epsilon, x_{a,i}}(x) + \varphi_a(x)$$

with $\|\varphi_a\|_a = O(\epsilon^{\frac{7}{2}})$, $x_{a,i} \rightarrow b_i$ as $0 < \epsilon \leq \epsilon_0$ for each $i = 1, \dots, m$, and $b_i \neq b_j$ for $i \neq j$, where $U_{\epsilon, x_{a,i}}(x) := (1 + \epsilon^2 P_i)U\left(\frac{\sqrt{1 + \epsilon^2 P_i}(x - x_{a,i})}{\epsilon}\right)$.

Let \tilde{u}_a be a m -peak solution of (3.1). Then for any small fixed $\rho > 0$, from (B.4) we find

$$\begin{aligned} & \epsilon^2 \int_{B_\rho(x_{a,i})} \frac{\partial P(x)}{\partial x_j} (\tilde{u}_a)^2 dx \\ &= \underbrace{-2\epsilon^2 \int_{\partial B_\rho(x_{a,i})} \frac{\partial \tilde{u}_a}{\partial \bar{\nu}} \frac{\partial \tilde{u}_a}{\partial x_j} d\sigma}_{:=\tilde{C}_1} + \underbrace{\epsilon^2 \int_{\partial B_\rho(x_{a,i})} |\nabla \tilde{u}_a|^2 \bar{\nu}_j(x) d\sigma}_{:=\tilde{C}_2} \\ &+ \underbrace{\int_{\partial B_\rho(x_{a,i})} (1 + \epsilon^2 P(x)) (\tilde{u}_a)^2 \bar{\nu}_j(x) d\sigma}_{:=\tilde{C}_3} - \underbrace{\frac{1}{8\pi\epsilon^2} \int_{\partial B_\rho(x_{a,i})} \int_{\mathbb{R}^3} \frac{(\tilde{u}_a)^2}{|x-y|} dy (\tilde{u}_a)^2 \bar{\nu}_j(x) d\sigma}_{:=\tilde{C}_4} \\ &- \underbrace{\frac{1}{8\pi\epsilon^2} \int_{B_\rho(x_{a,i})} \int_{\mathbb{R}^3} \frac{(\tilde{u}_a(y))^2 (\tilde{u}_a(x))^2 (x_j - y_j)}{|x-y|^3} dy dx}_{:=\tilde{C}_5}, \end{aligned} \quad (6.1)$$

where $j = 1, 2, 3$ and $\bar{\nu}(x) = (\bar{\nu}_1(x), \bar{\nu}_2(x), \bar{\nu}_3(x))$ is the outward unit normal of $\partial B_\rho(x_{a,i})$.

Let $\tilde{\varphi}_a(x) = \varphi_a(\epsilon x + x_{a,i})$. Then, $\tilde{\varphi}_a$ satisfies $\|\tilde{\varphi}_a\|_a = O(\epsilon^2)$. Using the Moser iteration, we can prove $\|\tilde{\varphi}_a\|_{L^\infty(\mathbb{R}^3)} = o(1)$. From this fact and the comparison theorem, similar to Proposition 2.2 in [23], we can prove the following estimates for $\tilde{u}_a(x)$ away from the concentrated points b_1, \dots, b_m .

Proposition 6.2. *Suppose that $\tilde{u}_a(x)$ is a m -peak solution of (3.1) concentrated at b_1, \dots, b_m . Then for any fixed $R \gg 1$, there exist some $\theta > 0$ and $C > 0$, such that*

$$|\tilde{u}_a(x)| + |\nabla \tilde{u}_a(x)| \leq C \sum_{i=1}^m e^{-\theta|x-x_{a,i}|/\epsilon}, \text{ for } x \in \mathbb{R}^3 \setminus \bigcup_{i=1}^m B_{R\epsilon}(x_{a,i}). \quad (6.2)$$

From the proof of Proposition 4.2 in [23], we have the following result.

Lemma 6.3. *For the small fixed constant $\bar{\rho} > 0$ and any $\rho \in (\bar{\rho}, 2\bar{\rho})$, then there exist $i_0 \in \{1, \dots, m\}$, $j_0 \in \{1, 2, 3\}$ and $C^* = C(i_0, j_0) \neq 0$ such that*

$$\tilde{C}_5 = C^* \epsilon^4 + o(\epsilon^4). \quad (6.3)$$

Now we are ready to prove Proposition 6.1.

Proof of Proposition 6.1. Here we prove it by contradiction. Assume that (3.1) has a m -peak solution $\tilde{u}_a(x)$. Then taking $i = i_0$ and $j = j_0$ as in Lemma 6.3, it follows from (6.1) and (6.3) that

$$\epsilon^2 \int_{B_\rho(x_{a,i})} \frac{\partial P(x)}{\partial x_j} (\tilde{u}_a)^2 dx = C^* \epsilon^4 + o(\epsilon^4) + O(e^{-\frac{\theta}{\epsilon}}),$$

since similar to (C.2) and by (6.2) we can prove

$$\sum_{i=1}^4 \tilde{C}_i \leq C \int_{\partial B_\rho(x_{a,i})} (\epsilon^2 |\nabla \tilde{u}_a|^2 + \tilde{u}_a^2(x)) d\sigma \leq C \int_{\partial B_\rho(x_{a,i})} \sum_{j=1}^m e^{-\frac{\theta \rho}{\epsilon}} d\sigma = O(e^{-\frac{\theta \rho}{\epsilon}}),$$

where we use the fact that $\{x : \partial B_\rho(x_{a,i})\} \subset \{x : \mathbb{R}^3 \setminus \bigcup_{j=1}^m B_{\frac{\rho}{2}}(x_{a,j})\}$. Since $x_{a,i} \rightarrow b_i \in \Gamma_i$, we find that there is a $t_a \in [P_i, P_i + \sigma]$ if Γ is a local minimum set of $P(x)$, or $t_a \in [P_i - \sigma, P_i]$ if Γ_i is a local maximum set of $P(x)$, such that $x_{a,i} \in \Gamma_{t_a,i}$. Let $\tau_{a,i}$ be the unit tangential vector of $\Gamma_{t_a,i}$ at $x_{a,i}$. Then

$$G(x_{a,i}) = 0, \text{ where } G(x) = \langle \nabla P(x), \tau_{a,i} \rangle.$$

We have the following expansion:

$$G(x) = \langle \nabla G(x_{a,i}), x - x_{a,i} \rangle + \frac{1}{2} \langle \langle \nabla^2 G(x_{a,i}), x - x_{a,i} \rangle, x - x_{a,i} \rangle + o(|x - x_{a,i}|^2), \text{ for } x \in B_\rho(x_{a,i}).$$

Then it follows from (4.1), the above expansion and $\|\varphi_a\|_a = O(\epsilon^{\frac{7}{2}})$ that

$$\begin{aligned} \int_{\mathbb{R}^3} G(x) U_{\epsilon, x_{a,i}}^2(x) &= \int_{B_\rho(x_{a,i})} G(x) U_{\epsilon, x_{a,i}}^2(x) + O(e^{-\frac{\tilde{\theta}}{\epsilon}}) \\ &= -2 \int_{B_\rho(x_{a,i})} G(x) U_{\epsilon, x_{a,i}}(x) \varphi_a - \int_{B_\rho(x_{a,i})} G(x) \varphi_a^2 + C^* \epsilon^2 + o(\epsilon^2) \\ &= O\left([\epsilon^{\frac{5}{2}} |\nabla G(x_{a,i})| + \epsilon^{\frac{7}{2}}] \|\varphi_a\|_a + \epsilon |\nabla G(x_{a,i})| \cdot \|\varphi_a\|_a^2\right) + C^* \epsilon^2 + o(\epsilon^2) \\ &= C^* \epsilon^2 + o(\epsilon^2), \end{aligned} \quad (6.4)$$

where $0 < \tilde{\theta} < \theta$ small.

On the other hand, noting that $G(x_{a,i}) = 0$, it is easy to check

$$\int_{\mathbb{R}^3} G(x) U_{\epsilon, x_{a,i}}^2(x) = \frac{1}{6} \epsilon^5 \Delta G(x_{a,i}) \int_{\mathbb{R}^3} |x|^2 U^2 + O(\epsilon^7) = O(\epsilon^5). \quad (6.5)$$

Then it follows from (6.4) and (6.5) that

$$C^* = o(1),$$

which contradicts with $C^* \neq 0$. Therefore (3.1) has no m -peak solution $\tilde{u}_a(x)$. \square

With Proposition 6.1 at hand, we can prove Theorem 1.7 at once.

Proof of Theorem 1.7. Theorem 1.7 follows directly from Proposition 6.1 and the relation between the solutions of equation (1.8) and equation (3.1). \square

APPENDIX

A. SOME KNOWN RESULTS

In this section, we give some known results which are used before. Denote

$$H_\epsilon = \left\{ u \in H^1(\mathbb{R}^3), \int_{\mathbb{R}^3} (\epsilon^2 |\nabla u|^2 + P(x)u^2(x)) dx < \infty \right\}$$

and the corresponding norm

$$\|u\|_\epsilon = (u(x), u(x))_\epsilon^{\frac{1}{2}} = \left(\int_{\mathbb{R}^3} (\epsilon^2 |\nabla u|^2 + P(x)u^2(x)) dx \right)^{\frac{1}{2}}.$$

Lemma A.1. (Lemma 2.1, [8]) For each $u \in L^q(\mathbb{R}^3)$ ($2 \leq q \leq 6$), we have

$$|u|_q \leq C\delta_a^{3(\frac{1}{q}-\frac{1}{2})} \|u\|_{\delta_a}.$$

Lemma A.2. (Lemma 2.2, [8], Lemma A.5, [23]) For any u_i ($i = 1, 2, 3, 4$) $\in H_\epsilon$, then

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{u_1(x)u_2(x)u_3(y)u_4(y)}{|x-y|} dx dy \leq C\epsilon^{-1} \|u_1\|_\epsilon \|u_2\|_\epsilon \|u_3\|_\epsilon \|u_4\|_\epsilon.$$

Lemma A.3. (Lemma A.6, [23]) For any $u_1, u_2, u_3, u_4 \in H^1(\mathbb{R}^3)$, then

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{u_1(x)u_2(x)u_3(y)u_4(y)}{|x-y|} dx dy \leq C \|u_1\|_{H^1(\mathbb{R}^3)} \|u_2\|_{H^1(\mathbb{R}^3)} \|u_3\|_{H^1(\mathbb{R}^3)} \|u_4\|_{H^1(\mathbb{R}^3)}.$$

Lemma A.4. (Lemma B.2, [23]) For any fixed $R > 0$, it holds

$$D_a(x) = o(1)R + O\left(\frac{1}{R}\right), \text{ for } x \in \mathbb{R}^3 \setminus B_{R\delta_a(x_a)}, \quad (\text{A.1})$$

and

$$E_a(x) = O(e^{-\tilde{\theta}R}), \text{ for } x \in \mathbb{R}^3 \setminus B_{R\delta_a(x_a)}, \text{ and some } \tilde{\theta} > 0. \quad (\text{A.2})$$

Now let $\tilde{\Gamma} \in C^2$ be a closed hypersurface in \mathbb{R}^3 . For $y \in \tilde{\Gamma}$, let $\nu(y)$ and $T(y)$ denote respectively the outward unit normal to $\tilde{\Gamma}$ at y and the tangent hyperplane to $\tilde{\Gamma}$ at y . The curvatures of $\tilde{\Gamma}$ at a fixed point $y_0 \in \tilde{\Gamma}$ are determined as follows. By a rotation of coordinates, we can assume that $y_0 = 0$ and $\nu(0)$ is the x_3 -direction, and x_j -direction is the j -th principal direction.

In some neighborhood $\mathcal{N} = \mathcal{N}(0)$ of 0, we have

$$\tilde{\Gamma} = \{x : x_3 = \phi(x')\},$$

where $x' = (x_1, x_2)$,

$$\phi(x') = \frac{1}{2} \sum_{j=1}^2 \kappa_j x_j^2 + O(|x'|^3),$$

where κ_j , is the j -th principal curvature of $\tilde{\Gamma}$ at 0. The Hessian matrix $[D^2\phi(0)]$ is given by

$$[D^2\phi(0)] = \text{diag}[\kappa_1, \kappa_2].$$

Suppose that W is a smooth function, such that $W(x) = \text{constant}$ for all $x \in \tilde{\Gamma}$. It follows from [24] that

Lemma A.5. (Lemma B.1, [24]) We have

$$\begin{aligned} \frac{\partial W(x)}{\partial x_l} \Big|_{x=0} &= 0, \quad l = 1, 2 \\ \frac{\partial^2 W(x)}{\partial x_m \partial x_l} \Big|_{x=0} &= -\frac{\partial W(x)}{\partial x_3} \Big|_{x=0} \kappa_m \delta_{ml}, \quad \text{for } m, l = 1, 2, \end{aligned} \quad (\text{A.3})$$

where κ_1, κ_2 , are the principal curvatures of $\tilde{\Gamma}$ at 0.

B. LINEARIZATION AND A POHOZAEV IDENTITY

Lemma B.1. Let ξ_0 be the solution of following system:

$$\begin{aligned} -\Delta \xi(x) + \xi(x) - \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy \xi(x) - \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{U(y)\xi(y)}{|x-y|} dy U(x) \\ = -\frac{1}{4\pi a_*} U(x) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x)U(y)\xi(y)}{|x-y|} dx dy. \end{aligned} \quad (\text{B.1})$$

Then it holds

$$\xi(x) = \sum_{j=0}^3 \gamma_j \psi_j, \quad (\text{B.2})$$

where γ_j are some constants,

$$\psi_0 = 2U + x \cdot \nabla U, \quad \psi_j = \frac{\partial U}{\partial x_j}, \quad \text{for } j = 1, 2, 3. \quad (\text{B.3})$$

Proof. We set $\bar{L}(u) := -\Delta u(x) + u(x) - \frac{1}{8\pi} \int_{\mathbb{R}^3} \frac{U^2(y)}{|x-y|} dy u(x) - \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{U(y)u(y)}{|x-y|} dy U(x)$. It is obvious that $\bar{L}(\psi_j) = 0$ and ψ_j is the solution of (B.1), for $j = 1, 2, 3$. Also, then using (3.12), we find that ψ_0 is also the solution of (B.1). And we know that $\psi_0, \psi_1, \psi_2, \psi_3$ are linearly independent. Then we get (B.2). \square

Proposition B.2. Let $\tilde{u}_a(x)$ be the solution of (3.1). Then we have following local Pohozaev identity:

$$\begin{aligned} \epsilon^2 \int_{\Omega} \frac{\partial P(x)}{\partial x_j} \tilde{u}_a^2(x) dx &= -2\epsilon^2 \int_{\partial\Omega} \frac{\partial \tilde{u}_a(x)}{\partial \nu} \frac{\partial \tilde{u}_a(x)}{\partial x_j} d\sigma + \epsilon^2 \int_{\partial\Omega} |\nabla \tilde{u}_a(x)|^2 \nu_j(x) d\sigma \\ &+ \int_{\partial\Omega} (1 + \epsilon^2 P(x)) \tilde{u}_a^2(x) \nu_j(x) d\sigma \\ &- \frac{1}{8\pi\epsilon^2} \int_{\partial\Omega} \int_{\mathbb{R}^3} \frac{\tilde{u}_a^2(y) \tilde{u}_a^2(x)}{|x-y|} \nu_j(x) dy d\sigma - \frac{1}{8\pi\epsilon^2} \int_{\Omega} \int_{\mathbb{R}^3} \tilde{u}_a^2(y) \tilde{u}_a^2(x) \frac{x_j - y_j}{|x-y|^3} dy dx, \end{aligned} \quad (\text{B.4})$$

where Ω is a bounded open domain of \mathbb{R}^3 , $j = 1, 2, 3$, $\nu(x) = (\nu_1(x), \nu_2(x), \nu_3(x))$ is the outward unit normal of $\partial\Omega$ and x_j, y_j are the j -th components of x, y .

Proof. Since (B.4) is obtained just by multiplying $\frac{\partial \tilde{u}_a(x)}{\partial x_j}$ on both sides of (1.4) and integrating on Ω , here we omit the details. \square

C. SOME BASIC AND USEFUL ESTIMATES

In this section, we mainly give some basic and useful estimates which are used before.

Lemma C.1. *There holds*

$$C_1 + C_2 + C_3 + C_4 + C_5 = O(e^{-\frac{\theta}{\epsilon}}). \quad (\text{C.1})$$

Proof. From (A.1) and (3.11), we have

$$C_1 + C_2 + C_3 + C_4 \leq C \int_{\partial B_\rho(x_a)} (\epsilon^2 |\nabla \tilde{u}_a|^2 + \tilde{u}_a^2(x)) d\sigma = O(e^{-\frac{\theta}{\epsilon}}). \quad (\text{C.2})$$

By symmetry and (3.11), we have

$$\begin{aligned} C_5 &= \frac{1}{8\pi\epsilon^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(\tilde{u}_a(y))^2 (\tilde{u}_a(x))^2 (x_j - y_j)}{|x - y|^3} dy dx \\ &\quad - \frac{1}{8\pi\epsilon^2} \int_{B_\rho^C(x_a)} \int_{\mathbb{R}^3} \frac{(\tilde{u}_a(y))^2 (\tilde{u}_a(x))^2 (x_j - y_j)}{|x - y|^3} dy dx \\ &= O(e^{-\frac{\theta}{\epsilon}}). \end{aligned} \quad (\text{C.3})$$

By (C.2) and (C.3), (C.1) is true. \square

Lemma C.2. *For any small and fixed $\rho > 0$, if $x \in B_{\rho\delta_a^{-1}\sqrt{\mathbf{P}_0}}(0)$, then we have*

$$F_1 + F_2 = O(\delta_a^4). \quad (\text{C.4})$$

Proof. From (5.24) and Lemma A.1, by Hölder inequality we have

$$\begin{aligned} F_1 &= \frac{1}{4\pi\mathbf{P}_0} \int_{\mathbb{R}^3} \frac{U(y) \tilde{\varphi}_a^{(1)}(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_a^{(1)})}{|x - y|} dy = \frac{1}{4\pi\mathbf{P}_0 \delta_a^2} \int_{\mathbb{R}^3} \frac{U(\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(y - x_a^{(1)})) \tilde{\varphi}_a^{(1)}(y)}{|y - x_a^{(1)} - \frac{\delta_a}{\sqrt{\mathbf{P}_0}}x|} dy \\ &\leq \frac{1}{4\pi\mathbf{P}_0 \delta_a^2} \left| U(\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(y - x_a^{(1)})) \right|_3 |\tilde{\varphi}_a^{(1)}(y)|_6 \left(\int_{|y - x_a^{(1)} - \frac{\delta_a}{\sqrt{\mathbf{P}_0}}x| \leq c_0 \delta_a} \frac{1}{|y - x_a^{(1)} - \frac{\delta_a}{\sqrt{\mathbf{P}_0}}x|^2} \right)^{\frac{1}{2}} \\ &\quad + \frac{1}{4\pi\mathbf{P}_0 \delta_a^3 c_0} \left\| U(\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(y - x_a^{(1)})) \right\|_{\delta_a} \|\tilde{\varphi}_a^{(1)}(y)\|_{\delta_a} \\ &\leq C \delta_a^{-2} \delta_a (\delta_a^{-1} \|\tilde{\varphi}_a^{(1)}(y)\|_{\delta_a}) \delta_a^{\frac{1}{2}} + C \delta_a^{-3} \delta_a^{\frac{3}{2}} \delta_a^{\frac{11}{2}} = O(\delta_a^4), \end{aligned} \quad (\text{C.5})$$

and

$$\begin{aligned} F_2 &= \frac{1}{8\pi\mathbf{P}_0^2} \int_{\mathbb{R}^3} \frac{(\tilde{\varphi}_a^{(1)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}}y + x_a^{(1)}))^2}{|x - y|} dy = \frac{1}{8\pi\mathbf{P}_0 \delta_a^2} \int_{\mathbb{R}^3} \frac{(\tilde{\varphi}_a^{(1)}(y))^2}{|y - x_a^{(1)} - \frac{\delta_a}{\sqrt{\mathbf{P}_0}}x|} dy \\ &\leq \frac{1}{8\pi\mathbf{P}_0 \delta_a^2} |\tilde{\varphi}_a^{(1)}(y)|_4^2 \left(\int_{|y - x_a^{(1)} - \frac{\delta_a}{\sqrt{\mathbf{P}_0}}x| \leq c_0 \delta_a} \frac{1}{|y - x_a^{(1)} - \frac{\delta_a}{\sqrt{\mathbf{P}_0}}x|^2} \right)^{\frac{1}{2}} \\ &\quad + \frac{1}{4\pi\mathbf{P}_0 \delta_a^3 c_0} \|\tilde{\varphi}_a^{(1)}(y)\|_{\delta_a}^2 \\ &\leq C \delta_a^{-2} (\delta_a^{-\frac{3}{4}} \|\tilde{\varphi}_a^{(1)}(y)\|_{\delta_a})^2 \delta_a^{\frac{1}{2}} + C \delta_a^{-3} \delta_a^{11} = O(\delta_a^8). \end{aligned} \quad (\text{C.6})$$

Hence, it follows from (C.5) to (C.6) that (C.4) holds. \square

Lemma C.3. For any small and fixed $\rho > 0$, if $x \in B_{\rho\delta_a^{-1}\sqrt{\mathbf{P}_0}}(0)$, there holds

$$G_1 + G_2 = O\left(\delta_a + \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}}x + x_a^{(1)}\right)\right). \quad (\text{C.7})$$

Proof. By the mean value theorem and (5.21), we have

$$\begin{aligned} G_1 &= \frac{1}{4\pi\mathbf{P}_0} \left(u_a^{(2)}\left(\frac{\delta_a x}{\sqrt{\mathbf{P}_0}} + x_{a,i}^{(1)}\right) - \mathbf{P}_0 U(x)\right) \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy \\ &= \frac{1}{4\pi\mathbf{P}_0} \left(\nabla U(\zeta x + (1-\zeta)\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x_a^{(1)} - x_a^{(2)})) \cdot \frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x_a^{(1)} - x_a^{(2)})\right. \\ &\quad \left.+ \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a x}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right)\right) \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy \\ &\leq C\left(\delta_a + \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}}x + x_a^{(1)}\right)\right) \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy \\ &= O\left(\delta_a + \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}}x + x_a^{(1)}\right)\right), \text{ for } \zeta \in (0, 1) \end{aligned} \quad (\text{C.8})$$

since

$$\begin{aligned} \int_{\mathbb{R}^3} \frac{U(y)\bar{\xi}_a(y)}{|x-y|} dy &\leq \int_{\mathbb{R}^3} \frac{U(y)}{|x-y|} dy \\ &\leq |U(y)|_2 \left(\int_{|x-y|\leq 2R} \frac{1}{|x-y|^2} dy\right)^{\frac{1}{2}} + \frac{1}{2R} \int_{\mathbb{R}^3} U(y) dy \\ &= O\left(R^{\frac{1}{2}} + \frac{1}{R}\right) = O(1). \end{aligned} \quad (\text{C.9})$$

We can also check

$$\begin{aligned} G_2 &= \frac{u_a^{(2)}\left(\frac{\delta_a x}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right)}{8\pi\mathbf{P}_0} \int_{\mathbb{R}^3} \frac{\left((u_a^{(1)} + u_a^{(2)})\left(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right) - 2\mathbf{P}_0 U(y)\right)\bar{\xi}_a(y)}{|x-y|} dy \\ &\leq C \frac{|x_{a,i}^{(1)} - x_a^{(2)}|}{\delta_a} \int_{\mathbb{R}^3} \frac{|\nabla U(\zeta y + (1-\zeta)\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x_a^{(1)} - x_a^{(2)}))| \cdot |\bar{\xi}_a(y)|}{|x-y|} dy \\ &\quad + C \sum_{l=1}^2 \int_{\mathbb{R}^3} \frac{|\tilde{\varphi}_a^{(l)}\left(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_{a,i}^{(1)}\right)| \cdot |\bar{\xi}_a(y)|}{|x-y|} dy \\ &= O(\delta_a), \end{aligned} \quad (\text{C.10})$$

since similar to (C.9), we have

$$\int_{\mathbb{R}^3} \frac{|\nabla U(\zeta y + (1-\zeta)\frac{\sqrt{\mathbf{P}_0}}{\delta_a}(x_a^{(1)} - x_a^{(2)}))| \cdot |\bar{\xi}_a(y)|}{|x-y|} dy = O(1),$$

and

$$\begin{aligned}
& \int_{\mathbb{R}^3} \frac{\tilde{\varphi}_a^{(l)}\left(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right) \bar{\xi}_a(y)}{|x-y|} dy \\
& \leq \int_{|y| \leq 2\frac{\sqrt{\mathbf{P}_0}}{\delta_a} \rho} \frac{\tilde{\varphi}_a^{(l)}\left(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right) \bar{\xi}_a(y)}{|x-y|} dy + \int_{|y| \geq 2\frac{\sqrt{\mathbf{P}_0}}{\delta_a} \rho} \frac{\tilde{\varphi}_a^{(l)}\left(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right) \bar{\xi}_a(y)}{|x-y|} dy \\
& \leq C |\tilde{\varphi}_a^{(l)}\left(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right)|_3 |\bar{\xi}_a(y)|_6 \left(\int_{|x-y| \leq 3\frac{\sqrt{\mathbf{P}_0}}{\delta_a} \rho} \frac{1}{|x-y|^2} dy \right)^{\frac{1}{2}} + C \delta_a |\tilde{\varphi}_a^{(l)}\left(\frac{\delta_a y}{\sqrt{\mathbf{P}_0}} + x_a^{(1)}\right)|_2 |\bar{\xi}_a(y)|_2 \\
& \leq C \delta_a^{-1} |\tilde{\varphi}_a^{(l)}(y)|_3 \delta_a^{-\frac{1}{2}} + C \delta_a \delta_a^{-\frac{3}{2}} |\tilde{\varphi}_a^{(l)}(y)|_2 \\
& \leq C \delta_a^{-\frac{3}{2}} (\delta_a^{-\frac{1}{2}} \delta_a^{\frac{11}{2}}) + C \delta_a^{-\frac{1}{2}} \delta_a^{\frac{11}{2}} = O(\delta_a^{\frac{7}{2}}).
\end{aligned}$$

Then (C.7) follows from (C.8) and (C.10). \square

Lemma C.4. *There holds*

$$\begin{aligned}
& \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{[(u_a^{(1)})^2(x) u_a^{(2)}(y) + (u_a^{(2)})^2(x) u_a^{(1)}(y)] \xi_a(y)}{|x-y|} dx dy \\
& = 2 \delta_a^5 \mathbf{P}_0^{\frac{1}{2}} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) U(y) \bar{\xi}_a(y)}{|x-y|} dx dy + O(\delta_a^6).
\end{aligned} \tag{C.11}$$

Proof. By direct computations, from (3.12) we have

$$\begin{aligned}
& \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(u_a^{(1)})^2(x) u_a^{(2)}(y) \xi_a(y)}{|x-y|} dx dy \\
& = \delta_a^5 \mathbf{P}_0^{\frac{1}{2}} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) U(y) \bar{\xi}_a(y)}{|x-y|} dx dy \\
& \quad + \frac{\delta_a^5}{\mathbf{P}_0^{\frac{1}{2}}} \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) (u_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}\right) - \mathbf{P}_0 U(y)) \bar{\xi}_a(y)}{|x-y|} dx dy}_{:=H_1} \\
& \quad + \delta_a^5 \mathbf{P}_0^{\frac{1}{2}} \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{\tilde{\varphi}_a^{(1)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}\right) (u_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}\right) + \mathbf{P}_0 U(x)) u_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}\right) \bar{\xi}_a(y)}{|x-y|} dx dy}_{:=H_2} \\
& = \delta_a^5 \mathbf{P}_0^{\frac{1}{2}} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) U(y) \bar{\xi}_a(y)}{|x-y|} dx dy + O(\delta_a^6),
\end{aligned} \tag{C.12}$$

since by Lemmas A.2, A.3 and (5.24), we have

$$\begin{aligned}
H_1 &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) \mathbf{P}_0 (U(y + \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)})) - U(y)) \bar{\xi}_a(y)}{|x - y|} dx dy \\
&\quad + \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) \tilde{\varphi}_a^{(2)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}) \bar{\xi}_a(y)}{|x - y|} dx dy \\
&= O(\delta_a) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) |\nabla U(\zeta y + (1 - \zeta) \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)}))| |\bar{\xi}_a(y)|}{|x - y|} dx dy \\
&\quad + C \delta_a^{-1} \|U\|_{\delta_a}^2 \left\| \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}\right) \right\|_{\delta_a} \|\bar{\xi}_a\|_{\delta_a} \\
&= O(\delta_a) \|U\|^2 \left\| \nabla U\left(\zeta y + (1 - \zeta) \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)})\right) \right\| \|\bar{\xi}_a\| + O(\delta_a^3) \\
&= O(\delta_a), \quad \text{for some } \zeta \in (0, 1)
\end{aligned}$$

and

$$\begin{aligned}
H_2 &= 2\mathbf{P}_0^2 \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U(x) \tilde{\varphi}_a^{(1)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}) U(y + \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)})) \bar{\xi}_a(y)}{|x - y|} dx dy \\
&\quad + 2\mathbf{P}_0 \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U(x) \tilde{\varphi}_a^{(1)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}) \tilde{\varphi}_a^{(2)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}) \bar{\xi}_a(y)}{|x - y|} dx dy \\
&\quad + \mathbf{P}_0 \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(\tilde{\varphi}_a^{(1)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}))^2 U(x + \frac{\sqrt{\mathbf{P}_0}}{\delta_a} (x_a^{(1)} - x_a^{(2)})) \bar{\xi}_a(y)}{|x - y|} dx dy \\
&\quad + \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(\tilde{\varphi}_a^{(1)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}))^2 \tilde{\varphi}_a^{(2)}(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}) \bar{\xi}_a(y)}{|x - y|} dx dy \\
&\leq C \delta_a^{-1} \|U\|_{\delta_a}^2 \left\| \tilde{\varphi}_a^{(1)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}\right) \right\|_{\delta_a} \|\bar{\xi}_a(y)\|_{\delta_a} \\
&\quad + C \delta_a^{-1} \|U\|_{\delta_a} \left\| \tilde{\varphi}_a^{(1)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}\right) \right\|_{\delta_a} \left\| \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}\right) \right\|_{\delta_a} \|\bar{\xi}_a(y)\|_{\delta_a} \\
&\quad + C \delta_a^{-1} \left\| \tilde{\varphi}_a^{(1)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}\right) \right\|_{\delta_a}^2 \|U\|_{\delta_a} \|\bar{\xi}_a(y)\|_{\delta_a} \\
&\quad + C \delta_a^{-1} \left\| \tilde{\varphi}_a^{(1)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} x + x_a^{(1)}\right) \right\|_{\delta_a}^2 \left\| \tilde{\varphi}_a^{(2)}\left(\frac{\delta_a}{\sqrt{\mathbf{P}_0}} y + x_a^{(1)}\right) \right\|_{\delta_a} \|\bar{\xi}_a(y)\|_{\delta_a} \\
&= O(\delta_a^3) + O(\delta_a^7) + O(\delta_a^{11}) = O(\delta_a^3).
\end{aligned}$$

Just by the same argument as that of (C.12), we can also check that

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(u_a^{(2)})^2(x) u_a^{(1)}(y) \xi_a(y)}{|x - y|} dx dy = \delta_a^5 \mathbf{P}_0^{\frac{1}{2}} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{U^2(x) U(y) \bar{\xi}_a(y)}{|x - y|} dx dy + O(\delta_a^6). \quad (\text{C.13})$$

It follows from (C.12) and (C.13) that (C.11) holds. \square

Acknowledgments Part of this work was done while Luo was visiting the Mathematics Department of University of Rome “La Sapienza” whose members he would like to thank for their warm hospitality. Guo was supported by NSFC grants (No.11771469). Luo and Wang were supported by the Fundamental Research Funds for the Central Universities(No.KJ02072020-0319). Luo was supported by the China Scholarship Council and NSFC grants (No.11701204, No.11831009). Wang was supported by NSFC grants (No.11671162). Yang was supported by NSFC grants (No.11601194).

REFERENCES

- [1] D. Cao, H. Heinz, *Uniqueness of positive multi-lump bound states of nonlinear Schrödinger equations*. Math. Z. **243** (2003), 599–642.
- [2] D. Cao, S. Li, P. Luo, *Uniqueness of positive bound states with multi-bump for nonlinear Schrödinger equations*. Calc. Var. Partial Differential Equations **54** (2015), 4037–4063.
- [3] D. Cao, E. Noussair, S. Yan, *Existence and uniqueness results on single-peaked solutions of a semi-linear problem*. Ann. Inst. H. Poincaré Anal. Non Linéaire **15** (1998), 73–111.
- [4] G. Chen, *Nondegeneracy of ground states and multiple semiclassical solutions of the Hartree equation for general dimensions*, arxiv:1610.05503v3[Math.AP], 3Feb 2020.
- [5] S. Cingolani, S. Secchi, M. Squassina, *Semi-classical limit for Schrödinger equations with magnetic field and Hartree-type nonlinearities*. Proc. Roy. Soc. Edinburgh Sect. A **140** (2010), 973–1009.
- [6] Y. Deng, Y. Guo, L. Lu, *Threshold behavior and uniqueness of ground states for mass critical inhomogeneous Schrödinger equations*. J. Math. Phys. **59** (2018), 21 pp.
- [7] Y. Deng, C. Lin, S. Yan, *On the prescribed scalar curvature problem in \mathbb{R}^N , local uniqueness and periodicity*. J. Math. Pures Appl. **104** (2015), 1013–1044.
- [8] B. Gheraibia, C. Wang, *Multi-peak positive solutions of a nonlinear Schrödinger-Newton type system*. Adv. Nonlinear Stud. **20** (2020), 53–75.
- [9] L. Ghanetgas, *Uniqueness of positive solutions of a nonlinear elliptic equation involving the critical exponent*. Nonlinear Anal. **20** (1993), 571–603.
- [10] M. Grossi, *On the number of single-peak solutions of the nonlinear Schrödinger equations*. Ann. Inst. H. Poincaré Anal. Non Linéaire **19** (2002), 261–280.
- [11] Y. Guo, C. Lin, J. Wei, *Local uniqueness and refined spike profiles of ground states for two-dimensional attractive Bose-Einstein condensates*. SIAM J. Math. Anal. **49** (2017), 3671–3715.
- [12] Y. Guo, Y. Luo, Z-Q Wang, *Limit behavior of mass critical Hartree minimization problems with steep potential wells*. J. Math. Phys. **59** (2018), 061504, 19 pp.
- [13] Y. Guo, S. Peng, S. Yan, *Local uniqueness and periodicity induced by concentration*. Proc. Lond. Math. Soc. **114** (2017), 1005–1043.
- [14] G. Li, S. Peng, C. Wang, *Multi-bump solutions for the nonlinear Schrödinger-Poisson system*. J. Math. Phys. **52** (2011), 19 pp.
- [15] G. Li, S. Peng, S. Yan, *Infinitely many positive solutions for the nonlinear Schrödinger-Poisson system*. Commun. Contemp. Math. **12** (2010), 1069–1092.
- [16] G. Li, H. Ye, *The existence of positive solutions with prescribed L^2 -norm for nonlinear Choquard equations*. J. Math. Phys. **55** (2014), 121501, 19 pp.
- [17] S. Li, J. Xiang, X. Zeng, *Ground states of nonlinear Choquard equations with multi-well potentials*. J. Math. Phys. **57** (2016), 081515, 19 pp.
- [18] S. Li, X. Zhu, *Mass concentration and local uniqueness of ground states for L^2 -subcritical nonlinear Schrödinger equations*. Z. Angew. Math. Phys. **70** (2019), 26 pp.
- [19] E. Lieb, *Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation*. Studies in Appl. Math. **57** (1976/77), 93–105.
- [20] E. Lieb, M. Loss, *Analysis, second ed., in: Graduate Studies in Mathematics*, vol. 14, American Mathematical Society, Providence, RI, 2001.

- [21] P. L. Lions, *The Choquard equation and related questions*. Nonlinear Anal. **4** (1980), 1063–1072.
- [22] P. L. Lions, *The concentration-compactness principle in the calculus of variations. The locally compact case. II*. Ann. Inst. H. Poincaré Anal. Non Linéaire **1** (1984), 223–283.
- [23] P. Luo, S. Peng, C. Wang, *Uniqueness of positive solutions with concentration for the Schrödinger-Newton problem*. Calc. Var. Partial Differential Equations **59** (2020), Paper No. 60, 41 pp.
- [24] P. Luo, S. Peng, S. Yan, *Excited states on Bose-Einstein condensates with attractive interactions*. arXiv:1909.08828.
- [25] G. P. Menzala, *On regular solutions of a nonlinear equation of Choquard's type*. Proc. Roy. Soc. Edinburgh Sect. A **86** (1980), 291–301.
- [26] V. Moroz, J. Van Schaftingen, *Ground states of nonlinear Choquard equations: existence, qualitative properties and decay asymptotics*. J. Funct. Anal. **265** (2013), 153–184.
- [27] V. Moroz, J. Van Schaftingen, *Existence of ground states for a class of nonlinear Choquard equations*. Trans. Amer. Math. Soc. **367** (2015), 6557–6579.
- [28] R. Penrose, *Quantum computation, entanglement and state reduction*. R. Soc. Lond. Philos. Trans. Ser. A Math. Phys. Eng. Sci. **356** (1998), 1927–1939.
- [29] M. Riesz, *L'intégrale de Riemann-Liouville et le problème de Cauchy*, Acta Math. **81** (1949) 1–233.
- [30] S. Secchi, *A note on Schrödinger-Newton systems with decaying electric potential*. Nonlinear Anal. **72** (2010), 3842–3856.
- [31] P. Tod, I. M. Moroz, *An analytical approach to the Schrödinger-Newton equations*. Nonlinearity **12** (1999), 201–216.
- [32] G. Vaira, *Existence of bound states for Schrödinger-Newton type systems*. Adv. Nonlinear Stud. **13** (2013), 495–516.
- [33] C. Wang, J. Yang, *Positive solutions for a nonlinear Schrödinger-Poisson system*. Discrete Contin. Dyn. Syst. **38** (2018), 5461–5504.
- [34] J. Wei, M. Winter, *Strongly interacting bumps for the Schrödinger-Newton equations*. J. Math. Phys. **50** (2009), 012905.
- [35] C. Xiang, *Uniqueness and nondegeneracy of ground states for Choquard equations in three dimensions*. Calc. Var. Partial Differential Equations **55** (2016), 55–134.
- [36] H. Ye, *Mass minimizers and concentration for nonlinear Choquard equations in R^N* . Topol. Methods Nonlinear Anal. **48** (2016), 393–417.

(Qing Guo) COLLEGE OF SCIENCE, MINZU UNIVERSITY OF CHINA, BEIJING 100081, CHINA
E-mail address: guoqing0117@163.com

(Peng Luo) SCHOOL OF MATHEMATICS AND STATISTICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN 430079, CHINA
E-mail address: pluo@mail.ccnu.edu.cn

(Chunhua Wang) SCHOOL OF MATHEMATICS AND STATISTICS AND HUBEI KEY LABORATORY OF MATHEMATICAL SCIENCES, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN 430079, CHINA
E-mail address: chunhuawang@mail.ccnu.edu.cn

(Jing Yang) SCHOOL OF SCIENCE, JIANGSU UNIVERSITY OF SCIENCE AND TECHNOLOGY, ZHENJIANG 212003, CHINA
E-mail address: yyangecho@163.com