

Integral formulas for a Riemannian manifold with several orthogonal complementary distributions

Vladimir Rovenski*

Abstract

In the paper we prove integral formulae for a Riemannian manifold endowed with $k > 2$ orthogonal complementary distributions, which generalize well-known formula for $k = 2$, and give applications to splitting and isometric immersions of Riemannian manifolds, in particular, multiply warped products, and to hypersurfaces with $k > 2$ distinct principal curvatures of constant multiplicities.

Keywords: Almost k -product manifold, distribution, foliation, mean curvature vector, mixed scalar curvature, isometric immersion, multiply warped product, Dupin hypersurface

Mathematics Subject Classifications (2010) 53C15; 53C12; 53C40

Introduction

Distributions on a manifold (i.e., subbundles of the tangent bundle) are used to build up notions of integrability, and specifically of a foliated manifold. Distributions and foliations on Riemannian manifolds appear in various situations, e.g., [3, 9, 13], among them warped products are fruitful generalizations of the direct product playing important role in mathematics and physics, e.g., [7].

Integral formulae are useful for many problems in the theory of foliations, e.g. [2, 13, 17]:

- a) the existence and characterizing of foliations, whose leaves have a given geometric property;
- b) prescribing the higher mean curvatures of the leaves of a foliation;
- c) minimizing functionals like volume and energy defined for tensor fields on a manifold.

The first known integral formula for a closed Riemannian manifold endowed with a codimension one foliation tells us that the integral mean curvature of the leaves vanishes, see [12]. The second formula in the series of total σ_k 's – elementary symmetric functions of principal curvatures of the leaves – says that for a codimension one foliation with a unit normal N to the leaves the total σ_2 is a half of the total Ricci curvature in the N -direction, e.g., [2]:

$$\int_M (2\sigma_2 - \text{Ric}_{N,N}) \, d \text{vol} = 0, \quad (1)$$

which is a consequence of Stokes' theorem applied to $\nabla_N N + \sigma_1 N$. One can see directly that (1) implies nonexistence of totally umbilical foliations on a closed manifold of negative curvature: if the integrand is strictly positive, so is the value of the integral. The mixed scalar curvature S_{mix} is one of the simplest curvature invariants of a pseudo-Riemannian almost-product structure, which can be defined as an averaged sum of sectional curvatures of planes that non-trivially intersect with both of the distributions. The following integral formula for a closed Riemannian manifold (M, g) endowed with two complementary orthogonal distributions, see [17] (and [11] for foliations), generalizes (1) and has many interesting global corollaries (e.g., decomposition criteria using the sign of the mixed scalar curvature, [14]):

$$\int_M (S_{\text{mix}} + \|h_1\|^2 + \|h_2\|^2 - \|H_1\|^2 - \|H_2\|^2 - \|T_1\|^2 - \|T_2\|^2) \, d \text{vol}_g = 0. \quad (2)$$

*Mathematical Department, University of Haifa, Mount Carmel, 31905 Haifa, Israel
e-mail: vrovenski@univ.haifa.ac.il

Here and $h_i, H_i = \text{Tr}_g h_i$ and T_i are the second fundamental form, the mean curvature vector field and the integrability tensor of distributions \mathcal{D}_1 and \mathcal{D}_2 on (M, g) . The formula (2) was obtained by calculation of the divergence of the vector field $H_1 + H_2$,

$$\text{div}(H_1 + H_2) = S_{\text{mix}} + \|h_1\|^2 + \|h_2\|^2 - \|H_1\|^2 - \|H_2\|^2 - \|T_1\|^2 - \|T_2\|^2, \quad (3)$$

and then applying Stokes' theorem.

Recently, the notion of warped products was naturally extended to multiply warped products, e.g., [7]. The notion of multiply warped product, in turn, is a special case of a Riemannian almost k -product structure with $k \in \{2, \dots, n\}$ foliations. This structure can be also viewed in the theory of webs composed of foliations of different dimensions, see [1]. A Riemannian almost k -product structure appears also on a proper Dupin hypersurface of a real space-form, i.e., the number k of distinct principal curvatures is constant and each principal curvature function is constant along its corresponding surface of curvature, see [6].

Thus, the problem of generalizing (2) to the case of (M, g) with $k > 2$ distributions is actual. In Section 1, we solve this problem for arbitrary $k > 2$. In Sections 2 and 3 we give applications to splitting and isometric immersions of manifolds, in particular, multiply warped products, and to hypersurfaces with $k > 2$ distinct principal curvatures of constant multiplicities.

1 New integral formulas

Let an n -dimensional Riemannian manifold (M, g) with the Levi-Civita connection ∇ and the curvature tensor R be endowed with $k > 2$ orthogonal n_i -dimensional distributions \mathcal{D}_i ($1 \leq i \leq k$) with $\sum n_i = \dim M$. There exists on M a local adapted orthonormal frame $\{E_1, \dots, E_n\}$, where

$$\{E_1, \dots, E_{n_1}\} \subset \mathcal{D}_1, \quad \{E_{n_{i-1}+1}, \dots, E_{n_i}\} \subset \mathcal{D}_i, \quad 2 \leq i \leq k.$$

Such $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ is called a *Riemannian almost k -product manifold*, see [9] for $k = 2$. A plane in TM spanned by two vectors belonging to different distributions, say, \mathcal{D}_i and \mathcal{D}_j is called *mixed*. We will generalize the well-known concept of almost-product manifolds.

Definition 1. The function on (M, g) with $k \geq 2$ non-degenerate distributions,

$$S_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) = \sum_{i < j} K(\mathcal{D}_i, \mathcal{D}_j) \quad (4)$$

will be called the *mixed scalar curvature* of $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$, where

$$K(\mathcal{D}_i, \mathcal{D}_j) = \sum_{n_{i-1} < a \leq n_i, n_{j-1} < b \leq n_j} \langle R(E_a, E_b) E_a, E_b \rangle, \quad i \neq j.$$

Let $P_i : TM \rightarrow \mathcal{D}_i$ be the orthoprojector, and $\widehat{P}_i = \text{id}_{TM} - P_i$ be the orthoprojector onto \mathcal{D}_i^\perp . The second fundamental form $h_i : \mathcal{D}_i \times \mathcal{D}_i \rightarrow \mathcal{D}_i^\perp$ (symmetric) and the integrability tensor $T_i : \mathcal{D}_i \times \mathcal{D}_i \rightarrow \mathcal{D}_i^\perp$ (skew-symmetric) of \mathcal{D}_i are defined by

$$h_i(X, Y) = \frac{1}{2} \widehat{P}_i(\nabla_X Y + \nabla_Y X), \quad T_i(X, Y) = \frac{1}{2} \widehat{P}_i(\nabla_X Y - \nabla_Y X) = \frac{1}{2} \widehat{P}_i[X, Y].$$

Let $h_{ij}, H_{ij} = \text{Tr}_g h_{ij}, T_{ij}$ be the second fundamental forms, mean curvature vector fields and the integrability tensors of distributions $\mathcal{D}_{i,j} = \mathcal{D}_i \oplus \mathcal{D}_j$ in M , and $P_{ij} : TM \rightarrow \mathcal{D}_{i,j}$ be orthoprojectors, etc. Note that

$$H_i = \sum_{j \neq i} P_j H_i, \quad H_{1\dots r} = P_{r+1\dots k}(H_1 + \dots + H_r), \quad \text{etc.}$$

Recall that a distribution \mathcal{D}_i is called integrable if $T_i = 0$, and \mathcal{D}_i is called totally umbilical, harmonic, or totally geodesic, if $h_i = (H_i/n_i)g$, $H_i = 0$, or $h_i = 0$, respectively.

Let $S(r, k)$ be the set of all r -combinations (i.e., subsets of r distinct elements) of $\{1, \dots, k\}$. The $S(r, k)$ contains $C_k^r = \frac{k!}{r!(k-r)!}$ elements (where C_k^r denote binomial coefficients). For example, $S(k-1, k)$ contains k elements. For any $\mathbf{q} \in S(r, k)$ we may assume $q_1 < \dots < q_r$. Set $h_{\mathbf{q}} = h_{q_1, \dots, q_r}$ and $T_{\mathbf{q}} = T_{q_1, \dots, q_r}$.

Our main goal is the following formula for $k \in \{2, \dots, n\}$, which generalizes (3).

Theorem 1. For a Riemannian almost k -product manifold $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ we have

$$\operatorname{div} X = 2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) + \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r, k)} (\|h_{\mathbf{q}}\|^2 - \|H_{\mathbf{q}}\|^2 - \|T_{\mathbf{q}}\|^2), \quad (5)$$

where $X = \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r, k)} H_{\mathbf{q}}$.

Proof. For $k = 2$ we have (3). To illustrate the proof for $k > 2$, first consider the case of $k = 3$. Using (3) for the distributions \mathcal{D}_1 and $\mathcal{D}_1^\perp = \mathcal{D}_2 \oplus \mathcal{D}_3$, we get

$$\operatorname{div}(H_1 + H_{23}) = 2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \mathcal{D}_1^\perp) + (\|h_1\|^2 - \|H_1\|^2 - \|T_1\|^2) + (\|h_{23}\|^2 - \|H_{23}\|^2 - \|T_{23}\|^2), \quad (6)$$

and similarly for $(\mathcal{D}_2, \mathcal{D}_2^\perp)$ and $(\mathcal{D}_3, \mathcal{D}_3^\perp)$. Summing 3 copies of (6), we obtain (5) for $k = 3$:

$$\begin{aligned} \operatorname{div} \left(\sum_i H_i + \sum_{i < j} H_{ij} \right) &= 2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3) \\ &+ \sum_i (\|h_i\|^2 - \|H_i\|^2 - \|T_i\|^2) + \sum_{i < j} (\|h_{ij}\|^2 - \|H_{ij}\|^2 - \|T_{ij}\|^2). \end{aligned} \quad (7)$$

Next, for arbitrary $k > 2$, we apply (3) for pairs of distributions $(\mathcal{D}_i, \mathcal{D}_i^\perp)$ and get k equalities. Summing these equations and using Lemma 1 below, we get (5). \square

Lemma 1. We have

$$2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) = \sum_i \operatorname{S}_{\text{mix}}(\mathcal{D}_i, \mathcal{D}_i^\perp).$$

Proof. This directly follows from definitions (4). \square

Remark 1. Just the particular case of (5) for $k = 3$, see (7), can find many geometrical applications, because an almost 3-product structure appears naturally in several topics:

- 1) almost para- f -manifolds, e.g., [15].
- 2) lightlike manifolds, i.e., (M, g) with degenerate metric g of constant rank and index, see [8].
- 3) on orientable 3-manifolds, since they admits 3 linearly independent vector fields, i.e., $n_i = 1$.
- 4) the theory of 3-webs composed of three generic foliations of different dimensions, e.g., [16].
- 5) minimal hypersurfaces in the sphere with 3 distinct principal curvatures, see [10].
- 6) tubes over a standard embeddings of a projective plane FP^2 , for $F = \mathbb{R}; \mathbb{C}; \mathbb{H}$ or \mathbb{O} (Cayley numbers), in $S^4; S^7; S^{13}$, and S^{25} , respectively, see [6].

Example 1. For $k = 4$ (next to $k = 2, 3$), formula (5) reads as

$$\begin{aligned} \operatorname{div} \left(\sum_i H_i + \sum_{i < j < s} H_{ijs} \right) &= 2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_4) \\ &+ \sum_i (\|h_i\|^2 - \|H_i\|^2 - \|T_i\|^2) + \sum_{i < j < s} (\|h_{ijs}\|^2 - \|H_{ijs}\|^2 - \|T_{ijs}\|^2), \end{aligned}$$

Using Stokes' theorem for (5) on a closed manifold $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ yields the novel *integral formula* for any $k \in \{2, \dots, n\}$, which for $k = 2$ coincides with (2).

Corollary 1. On a closed Riemannian almost k -product manifold we have

$$\int_M \left(2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) + \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r, k)} (\|h_{\mathbf{q}}\|^2 - \|H_{\mathbf{q}}\|^2 - \|T_{\mathbf{q}}\|^2) \right) d \operatorname{vol}_g = 0. \quad (8)$$

Example 2. For few initial values of k , $k = 3, 4$, the integral formula (8) reads as follows:

$$\begin{aligned} \int_M \left(2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3) + \sum_i (\|h_i\|^2 - \|H_i\|^2 - \|T_i\|^2) + \sum_{i < j} (\|h_{ij}\|^2 - \|H_{ij}\|^2 - \|T_{ij}\|^2) \right) d \operatorname{vol}_g &= 0, \\ \int_M \left(2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_4) + \sum_i (\|h_i\|^2 - \|H_i\|^2 - \|T_i\|^2) + \sum_{i < j < s} (\|h_{ijs}\|^2 - \|H_{ijs}\|^2 - \|T_{ijs}\|^2) \right) d \operatorname{vol}_g &= 0, \end{aligned}$$

In order to simplify the LHS of (5) to the shorter view $\operatorname{div}(\sum_i H_i)$, we reorganize the terms $\operatorname{div}(\sum_{\mathbf{q} \in S(k-1,k)} H_{\mathbf{q}})$ for $r > 1$ and obtain new (auxiliary) integral formulae.

Theorem 2. For $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ and any $r \in \{2, \dots, k-1\}$, we have

$$\operatorname{div} X = \sum_{\mathbf{q} \in S(r,k)} (\|H_{\mathbf{q}}\|^2 + \langle \sum_{i=1}^r H_{q(i)} - r H_{\mathbf{q}}, \sum_{j \notin \mathbf{q}} H_j \rangle), \quad (9)$$

where $X = \sum_{\mathbf{q} \in S(r,k)} H_{\mathbf{q}} - C_{k-2}^{r-1} \sum_i H_i$.

Proof. Using equality $H_{1\dots r} = P_{r+1\dots k}(H_1 + \dots + H_r)$ for $\mathbf{q} = \{1, \dots, r\}$, we find

$$\begin{aligned} \operatorname{div} H_{1\dots r} &= \operatorname{div}_{r+1\dots k} H_{1\dots r} - \|H_{1\dots r}\|^2 \\ &= \operatorname{div}_{r+1\dots k}(H_1 + \dots + H_r) + \langle H_1 + \dots + H_r, H_{r+1\dots k} \rangle - \|H_{1\dots r}\|^2, \end{aligned}$$

and similarly for all C_k^r cases of $\mathbf{q} \in S(r, k)$. Summing the above, we use equalities

$$\operatorname{div}_{r+1\dots k} H_1 = \sum_{j>r} \operatorname{div}_j H_1, \quad \operatorname{div}_{2\dots k} H_1 = \operatorname{div} H_1 + \|H_1\|^2, \quad \text{etc.}$$

We apply (9) with $k-1$ for the distributions $\mathcal{D}_1, \dots, \mathcal{D}_{k-2}$ and $\mathcal{D}_{k-1,k}$, and then obtain similar formulas for other choices of a pair of distributions. Summing these C_k^2 equations, we get equation of the form (9) with k , comparing coefficients yield the claim. \square

Corollary 2. The following integral formulae for $r \in \{2, \dots, k-1\}$ take place on a closed Riemannian almost k -product manifold:

$$\int_M \sum_{\mathbf{q} \in S(r,k)} (\|H_{\mathbf{q}}\|^2 + \langle \sum_{i=1}^r H_{q(i)} - r H_{\mathbf{q}}, \sum_{j \notin \mathbf{q}} H_j \rangle) \operatorname{d vol}_g = 0. \quad (10)$$

Example 3. For $k=3$ and $r=2$, the formula (10) reads as

$$\int_M (\sum_i \|H_i\|^2 + \sum_{i<j} (2\langle H_i, H_j \rangle - \|H_{ij}\|^2)) \operatorname{d vol}_g = 0,$$

and for $k=4$ and $r=2$, the formula (10) reads as

$$\int_M (2\sum_i \|H_i\|^2 + \sum_{i,j<s} \langle H_i, H_{js} \rangle - \sum_{i<j} \|H_{ij}\|^2) \operatorname{d vol}_g = 0.$$

The reader can easily find the integral formula corresponding to (10) with $k=4$ and $r=3$.

From Theorem 1 and Theorem 2 for $r=k-1$, we obtain the following companion of (5).

Theorem 3. For a Riemannian almost k -product manifold $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ we have

$$\begin{aligned} 2 \operatorname{div} (\sum_i H_i) &= 2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) + \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r,k)} (\|h_{\mathbf{q}}\|^2 - \|H_{\mathbf{q}}\|^2 - \|T_{\mathbf{q}}\|^2) \\ &\quad - \sum_{\mathbf{q} \in S(k-1,k)} (\|H_{\mathbf{q}}\|^2 + \langle \sum_{i=1}^{k-1} H_{q(i)} - (k-1) H_{\mathbf{q}}, \sum_{j \notin \mathbf{q}} H_j \rangle). \end{aligned}$$

Corollary 3. The following integral formula takes place on a closed $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$:

$$\begin{aligned} \int_M &\left(2 \operatorname{S}_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) + \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(k-1,k)} (\|h_{\mathbf{q}}\|^2 - \|H_{\mathbf{q}}\|^2 - \|T_{\mathbf{q}}\|^2) \right. \\ &\quad \left. - \sum_{\mathbf{q} \in S(k-1,k)} (\|H_{\mathbf{q}}\|^2 + \langle \sum_{i=1}^{k-1} H_{q(i)} - (k-1) H_{\mathbf{q}}, \sum_{j \notin \mathbf{q}} H_j \rangle) \right) \operatorname{d vol}_g = 0. \end{aligned}$$

Example 4. For particular case of $k=3$, the integral formula of Corollary 3 reads as

$$\begin{aligned} \int_M &\left(\operatorname{S}_{\text{mix}}(\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3) - \sum_i \|H_i\|^2 - \sum_{i<j} \langle H_i, H_j \rangle + \frac{1}{2} \sum_i (\|h_i\|^2 - \|T_i\|^2) \right. \\ &\quad \left. + \frac{1}{2} \sum_{i<j} (\|h_{ij}\|^2 - \|T_{ij}\|^2) \right) \operatorname{d vol}_g = 0. \end{aligned}$$

The above formula was obtained in [4] by long direct calculations of $\operatorname{div}(H_1 + H_2 + H_3)$.

2 Splitting and isometric immersions of manifolds

Here, we use results of Section 1 to prove some splitting and non-existence of immersions results for Riemannian almost k -product manifolds with $k > 2$. We say that $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ splits if all distributions \mathcal{D}_i are integrable and M is locally the direct product $M_1 \times \dots \times M_k$ with foliations tangent to \mathcal{D}_i . Recall that if a simply connected manifold splits then it is the direct product.

We apply the submanifolds theory to Riemannian almost k -product manifolds.

Definition 2. A pair $(\mathcal{D}_i, \mathcal{D}_j)$ with $i \neq j$ of distributions on $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ with $k > 2$ is called

- a) *mixed totally geodesic*, if $h_{ij}(X, Y) = 0$ for all $X \in \mathcal{D}_i$ and $Y \in \mathcal{D}_j$.
- b) *mixed integrable*, if $T_{ij}(X, Y) = 0$ for all $X \in \mathcal{D}_i$ and $Y \in \mathcal{D}_j$.

Lemma 2. If each pair $(\mathcal{D}_i, \mathcal{D}_j)$ on $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ with $k > 2$ is

- a) *mixed totally geodesic*, then $h_{\mathbf{q}}(X, Y) = 0$
 - b) *mixed integrable*, then $T_{\mathbf{q}}(X, Y) = 0$
- for all $\mathbf{q} \in S(r, k)$, $2 < r < k$ and $X \in \mathcal{D}_{q(1)}$, $Y \in \mathcal{D}_{q(2)}$.

Proof. This follows by mathematical induction. □

The next our results generalize [14, Theorem 6] and [17, Theorem 2] on case $k = 2$.

Theorem 4. Let a Riemannian almost k -product manifold $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ with $k > 2$ has integrable harmonic distributions $\mathcal{D}_1, \dots, \mathcal{D}_k$. If $S_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) \geq 0$, and each pair $(\mathcal{D}_i, \mathcal{D}_j)$ is mixed integrable, then M splits.

Proof. From the equality $H_{1\dots r} = P_{r+1\dots k}(H_1 + \dots + H_r)$ follows that $H_i = 0$ for all $i \in \{1, \dots, k\}$, then $H_{\mathbf{q}} = 0$ for all $\mathbf{q} \in S(r, k)$ and $2 \leq r < k$. Similarly (by Lemma 2), if $T_{ij} = 0$ for all $i \in \{1, \dots, k\}$, then $T_{\mathbf{q}} = 0$ for all $\mathbf{q} \in S(r, k)$ and $2 \leq r < k$. By conditions and (5),

$$2S_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) + \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r, k)} (\|h_{\mathbf{q}}\|^2) = 0.$$

Thus, $h_{\mathbf{q}} = 0$ for all $\mathbf{q} \in S(r, k)$ with $r \in \{1, k-1\}$, in particular, $h_i = 0$ ($1 \leq i \leq k$). By well-known de Rham decomposition theorem, (M, g) splits. □

Theorem 5. Let a complete open Riemannian almost k -product manifold $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$ with $k > 2$ has totally umbilical distributions such that each pair $(\mathcal{D}_i, \mathcal{D}_j)$ is mixed totally geodesic and $\langle H_i, H_j \rangle = 0$ for $i \neq j$. If $S_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) \leq 0$ and $\|H_i\| \in L^1(M, g)$ for $1 \leq i \leq k$, then M splits.

Proof. By assumptions, from (5) we get

$$\text{div } \xi = 2S_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) + \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r, k)} (\|h_{\mathbf{q}}\|^2 - \|H_{\mathbf{q}}\|^2 - \|T_{\mathbf{q}}\|^2), \quad (11)$$

where $\xi = \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r, k)} H_{\mathbf{q}}$. By conditions, and since $\|H_{\mathbf{q}}\| \leq \sum_{i=1}^k \|H_{q(i)}\|$, we get $\|H_{\mathbf{q}}\| \in L^1(M, g)$. Since $\|\xi\| \leq \sum_{r \in \{1, k-1\}} \sum_{\mathbf{q} \in S(r, k)} \|H_{\mathbf{q}}\|$, then also $\|\xi\| \in L^1(M, g)$. Next, by conditions, for any $\mathbf{q} = (q(1), \dots, q(r)) \in S(r, k)$ with $r \geq 1$ we have

$$\|h_{\mathbf{q}}\|^2 - \|H_{\mathbf{q}}\|^2 = - \sum_{i=1}^r \frac{n_{q(i)} - 1}{n_{q(i)}} \|\widehat{P}_{\mathbf{q}} H_{q(i)}\|^2 \leq 0,$$

where $\widehat{P}_{\mathbf{q}}$ is the orthoprojector on the distribution $\bigoplus_{j \neq \mathbf{q}} \mathcal{D}_j$. Hence, from $S_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) \leq 0$ and (11) we get $\text{div } \xi \leq 0$. By conditions and Lemma 3 below, $\text{div } \xi = 0$. Thus, see (11), $S_{\text{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) = 0$ and T_i and h_i vanish. By de Rham decomposition theorem, (M, g) splits. □

Modifying Stokes' theorem on a complete open manifold (M, g) yields the following.

Lemma 3 (see Proposition 1 in [5]). *Let (M^n, g) be a complete open Riemannian manifold endowed with a vector field ξ such that $\operatorname{div} \xi \geq 0$. If the norm $\|\xi\|_g \in L^1(M, g)$ then $\operatorname{div} \xi \equiv 0$.*

Example 5. Totally umbilical integrable distributions appear on multiply warped products. Let F_1, \dots, F_k be k Riemannian manifolds and let $M = F_1 \times \dots \times F_k$ be their direct product. A multiply warped product $F_1 \times_{u_2} F_2 \times \dots \times_{u_k} F_k$ is M with the metric $g = g_{F_1} \oplus u_2^2 g_{F_2} \oplus \dots \oplus u_k^2 g_{F_k}$, where $u_i : F_1 \rightarrow (0, \infty)$ for $i = 2, \dots, k$ are smooth functions. Let \mathcal{D}_i be the distribution on M obtained from the vectors tangent to (the horizontal lifts of) F_i , e.g., [7]. The leaves (i.e., tangent to \mathcal{D}_i , $i \geq 2$) are totally umbilical submanifolds, with

$$H_i = -n_i \nabla(\log u_i),$$

and the fibers (i.e., tangent to \mathcal{D}_1) are totally geodesic submanifolds. Since

$$\operatorname{div} H_i = -n_i (\Delta u_i)/u_i - (n_i^2 - n_i) \|P_i \nabla u_i\|^2 / u_i^2,$$

where $\Delta = -\operatorname{div} \circ \nabla$ is the Laplacian on $C^2(F_1)$, and we have

$$S_{\operatorname{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) = \sum_{i \geq 2} n_i (\Delta u_i)/u_i. \quad (12)$$

Corollary 4. *Let a multiply warped product (M, g) be complete open and $\langle H_i, H_j \rangle = 0$ for $i \neq j$. If $S_{\operatorname{mix}}(\mathcal{D}_1, \dots, \mathcal{D}_k) \leq 0$ and $\|H_i\| \in L^1(M, g)$ for $1 \leq i \leq k$, then M is the direct product.*

The following theorem on isometric immersions of multiply warped products, see [7, Theorem 10.2], is related to the question by B.-Y. Chen: “What can we conclude from an arbitrary isometric immersion of a warped product into a Riemannian manifold with arbitrary codimension?”

Theorem A *Let $f : F_1 \times_{u_2} F_2 \times \dots \times_{u_k} F_k \rightarrow \tilde{M}$ be an isometric immersion of a multiply warped product $(M, g) := F_1 \times_{u_2} F_2 \times \dots \times_{u_k} F_k$ into an arbitrary Riemannian manifold. Then*

$$\sum_{i=2}^k n_i (\Delta u_i)/u_i \leq \frac{n^2(k-1)}{2k} H^2 + n_1(n - n_1) \max \tilde{K}, \quad n = \sum_{j=1}^k n_j, \quad (13)$$

where $H^2 = \langle H, H \rangle$ is the squared mean curvature of f , $\max \tilde{K}(x)$ is the maximum of the sectional curvature of \tilde{M} restricted to 2-plane sections of the tangent space $T_x M$ of M at $x = (x_1, \dots, x_k)$. The equality sign of (13) holds identically if and only if the following two statements hold:

- 1) f is a mixed totally geodesic immersion satisfying equalities $\operatorname{Tr}_g h_1 = \dots = \operatorname{Tr}_g h_k$;
- 2) at each point $x \in M$, the sectional curvature function \tilde{K} of \tilde{M} satisfies $\tilde{K}(X, Y) = \max \tilde{K}(x)$ for each unit vector $X \in T_{x_1}(F_1)$ and each unit vector Y in $T(x_2, \dots, x_k)(F_2 \times \dots \times F_k)$.

On a multiply warped product with $k > 2$ each pair of distributions is mixed totally geodesic. Indeed, such manifold is diffeomorphic to the direct product and the Lie bracket does not depend on metric. Thus, using Theorem A and our results above, we obtain the following.

Theorem 6 (Non-existence of immersions). *Let $f : F_1 \times_{u_2} F_2 \times \dots \times_{u_k} F_k \rightarrow \tilde{M}$ be an isometric immersion of a closed multiply warped product $(M, g) := F_1 \times_{u_2} F_2 \times \dots \times_{u_k} F_k$ into an arbitrary Riemannian manifold. Let, in addition, $\langle H_i, H_j \rangle = 0$ for $i \neq j$. Then there are no isometric immersions $f : F_1 \times_{u_2} F_2 \times \dots \times_{u_k} F_k \rightarrow \tilde{M}$ satisfying the inequality*

$$\frac{n^2(k-1)}{2k} H^2 < -n_1(n - n_1) \max \tilde{K}; \quad (14)$$

in particular, there are no minimal isometric immersions f when $\tilde{K} < 0$.

Proof. From (12), (11) and (13), applying the Stokes' Theorem, we get

$$\begin{aligned} 0 &\leq (1/2) \int_M \sum_{r \in \{1, k-1\}} \sum_{q \in S(r, k)} \sum_{i=1}^r \frac{n_{q(i)} - 1}{n_{q(i)}} \|\hat{P}_q H_{q(i)}\|^2 \operatorname{d vol}_g \\ &\leq \frac{n^2(k-1)}{2k} \int_M H^2 \operatorname{d vol}_g + n_1(n - n_1) \max \tilde{K} \cdot \operatorname{Vol}(M, g). \end{aligned} \quad (15)$$

The equality in (15) holds if and only if conclusions 1)–2) of Theorem A are satisfied. If such isometric immersion exists, then the inequality (14) yields a contradiction \square

Remark 2. Theorem 6 can be improved replacing $\max \tilde{K}$ by $\bar{S}(n_1, \dots, n_k)$ (with certain factor) defined below. Let L_i ($i = 1, 2$) be two subspaces of $T_x \bar{M}$ at some point $x \in \bar{M}$, and $\{E'_i\}$ and $\{\mathcal{E}'_j\}$ some orthonormal frames of these subspaces. The following quantity:

$$\bar{S}(L_1, L_2) = \sum_{i,j} \langle \bar{R}(E'_i, \mathcal{E}'_j) E'_i, \mathcal{E}'_j \rangle$$

does not depend on the choice of frames. For any $n_1, \dots, n_k \in \mathbb{N}$ with $\sum_i n_i \leq \dim \bar{M}$, let $\Gamma_x(n_1, \dots, n_k)$ be the set of k -tuples of pairwise orthogonal subspaces (L_1, \dots, L_k) with $\dim L_i = n_i$ ($1 \leq i \leq k$) of $T_x \bar{M}$ at some point $x \in \bar{M}$. Set $\bar{S}(n_1, \dots, n_k) = \sup_{x \in \bar{M}} \bar{S}_x(n_1, \dots, n_k)$, where

$$\bar{S}_x(n_1, \dots, n_k) = \max \left\{ \sum_{i < j} \bar{S}(L_i, L_j) : (L_1, \dots, L_k) \in \Gamma_x(n_1, \dots, n_k) \right\}.$$

3 Hypersurfaces with k distinct principal curvatures

Let M^n be a transversely orientable hypersurface in a Riemannian manifold \bar{M} . Denote by A the shape operator of M with respect to a unit normal vector field N , and let $\lambda_1 \leq \dots \leq \lambda_n$ be the principal curvatures (eigenvalues of A) – continuous functions on M . If at any point of M all principal curvatures are equal, then M is totally umbilical.

Suppose that there exists $k \geq 2$ distinct principal curvatures (μ_i) on M of multiplicities n_i , and let \mathcal{D}_i ($1 \leq i \leq k$) be corresponding eigen-distributions – subbundles of TM .

$$\mu_1 := \lambda_1 = \dots = \lambda_{n_1} < \dots < \mu_k := \lambda_{n-n_k+1} = \dots = \lambda_n.$$

Such hypersurface is a Riemannian almost k -product manifold $(M, g; \mathcal{D}_1, \dots, \mathcal{D}_k)$, see Section 1.

For a hypersurface M in a Riemannian manifold $\bar{M}(c)$ of constant curvature c , it is known the following, see [6]: if $n_i > 1$ for some i , then the function $\mu_i : M \rightarrow \mathbb{R}$ is differentiable and \mathcal{D}_i is integrable and its leaves are totally umbilical in $\bar{M}(c)$.

The curvature tensor \bar{R} of $\bar{M}(c)$ has the well-known view

$$\bar{R}(X, Y)Z = c(\langle X, Z \rangle Y - \langle Y, Z \rangle X).$$

Denote by $X_i \in \mathfrak{X}_i$ local unit vector fields on \mathcal{D}_i ($i \leq k$). For the sectional curvature of M we get

$$K(X_i, X_j) = c + \mu_i \mu_j, \quad i \neq j. \quad (16)$$

Example 6. For a hypersurface M in $\bar{M}(c)$ with $k = 2$, using equalities

$$\|h_i\|^2 - \|H_i\|^2 = n_i(1 - n_i) \frac{\|\nabla \mu_i\|^2}{(\mu_i - \mu_j)^2}, \quad S_{\text{mix}}(\mathcal{D}_1, \mathcal{D}_2) = n_1 n_2 (c + \mu_1 \mu_2),$$

formula (3) can be rewritten in terms of A and μ_i ($i = 1, 2$) as follows, see [17]:

$$\text{div}(H_1 + H_2) = n_1 n_2 (c + \mu_1 \mu_2) + \frac{n_1(1 - n_1) \|\nabla \mu_1\|^2 + n_2(1 - n_2) \|\nabla \mu_2\|^2}{(\mu_2 - \mu_1)^2}, \quad (17)$$

The next **problem** was posed by P. Walczak [17]: “To search for formulae analogous to (17) in the case of a hypersurface in $\bar{M}(c)$ of $k > 2$ distinct principal curvatures of constant multiplicities”. By symmetry of A and the Codazzi equation for a hypersurface in $\bar{M}(c)$,

$$(\nabla_X A)(Y) = (\nabla_Y A)(X), \quad (18)$$

the following tensor \mathcal{A} of rank 3 is totally symmetric:

$$\mathcal{A}(X, Y, Z) = \langle (\nabla_X A)(Y), Z \rangle, \quad X, Y, Z \in \mathfrak{X}_M.$$

Remark 3. A submanifold M in \bar{M} is called *curvature-invariant* if $R(X, Y)Z \in T_x M$ for any $X, Y, Z \in T_x M$ and any $x \in M$, e.g., [13]. Examples are arbitrary submanifolds in $\bar{M}(c)$. For a curvature-invariant hypersurface $M \subset \bar{M}$, Codazzi equation (18) is satisfied, thus, results of this section on a hypersurface in $\bar{M}(c)$ can be extended for curvature-invariant hypersurfaces.

Lemma 4. For a hypersurface M in $\bar{M}(c)$ with $k > 2$ distinct principal curvatures, we have

$$\begin{aligned}\mathcal{A}(X_i, X_j, X_l) &= (\mu_j - \mu_l) \langle \nabla_{X_i} X_j, X_l \rangle, \quad j \neq l, \\ \mathcal{A}(X_i, X_j, X_j) &= X_i(\mu_j), \quad \forall i, j.\end{aligned}\tag{19}$$

Proof. After derivation of equality $AX_j = \mu_j X_j$ and using symmetry of A , we obtain

$$\begin{aligned}\langle \nabla_{X_i}(AX_j), X_l \rangle &= \langle (\nabla_{X_i} A)(X_j), X_l \rangle + \langle A \nabla_{X_i} X_j, X_l \rangle \\ &= \mathcal{A}(X_i, X_j, X_l) + \mu_l \langle \nabla_{X_i} X_j, X_l \rangle, \\ \langle \nabla_{X_i}(\mu_j X_j), X_l \rangle &= \mu_j \langle \nabla_{X_i} X_j, X_l \rangle + X_i(\mu_j) \delta_{jl}.\end{aligned}$$

From this equalities in (19) follow. □

Corollary 5. For a hypersurface M in $\bar{M}(c)$ with $k > 2$ distinct principal curvatures, we have

$$(\mu_j - \mu_l) \langle \nabla_{X_i} X_j, X_l \rangle = (\mu_i - \mu_l) \langle \nabla_{X_j} X_i, X_l \rangle, \quad i \neq j \neq l.\tag{20}$$

Proof. This follows from (19) and symmetry of \mathcal{A} . Alternatively, one can use direct derivation,

$$\begin{aligned}0 &= \langle \bar{R}(X_i, X_j)N, X_l \rangle = \langle \mu_{X_i}(\mu_j X_j) - \nabla_{X_j}(\mu_i X_i) - \langle [X_i, X_j], X_l \rangle \langle \mu_l X_l, X_l \rangle \\ &= (\mu_j - \mu_l) \langle \nabla_{X_i} X_j, X_l \rangle - (\mu_i - \mu_l) \langle \nabla_{X_j} X_i, X_l \rangle,\end{aligned}$$

from which (20) follows. □

Theorem 5.13 in [6], which states that *each point of a hypersurface M^n in $\bar{M}(c)$ with 2, 3, 4 or 6 distinct principal curvatures has a principal coordinate neighborhood if and only if each \mathcal{D}_i^\perp is integrable*. We complete this result by the following.

Proposition 1. Let M^n be a hypersurface in $\bar{M}(c)$ with $k > 2$ distinct principal curvatures at each point. Then each \mathcal{D}_i^\perp is integrable if and only if $\mathcal{A}(X_i, X_j, X_l) = 0$ for $X_i \in \mathcal{D}_i$, $X_j \in \mathcal{D}_j$ and $X_l \in \mathcal{D}_l$ with $i < j < l$ on M^n .

Next, we study the problem (posed in [17]) for $k = 3$ (the case of $k > 3$ is similar).

Theorem 7. For a hypersurface M in $\bar{M}(c)$ with $k = 3$ distinct principal curvatures, we have

$$\begin{aligned}\operatorname{div} \sum_i n_i \left(\frac{P_j \nabla \mu_i}{\mu_i - \mu_j} + \frac{P_l \nabla \mu_i}{\mu_i - \mu_l} \right) \\ = \frac{1}{2} \sum_{i < j} n_i n_j (c + \mu_i \mu_j) + \sum_i n_i (1 - n_i) \left(\frac{\|P_j \nabla \mu_i\|^2}{(\mu_i - \mu_j)^2} + \frac{\|P_l \nabla \mu_i\|^2}{(\mu_i - \mu_l)^2} \right),\end{aligned}\tag{21}$$

where $(j, l) \in \{1, 2, 3\} \setminus \{i\}$ and $j < l$.

Proof. By (7) and $T_i = 0$, we have

$$\begin{aligned}\operatorname{div} \left(\sum_i H_i + \sum_{i < j} H_{ij} \right) &= 2 \operatorname{S}_{\operatorname{mix}}(\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3) + \sum_i (\|h_i\|^2 - \|H_i\|^2) \\ &\quad + \sum_{i < j} (\|h_{ij}\|^2 - \|H_{ij}\|^2 - \|T_{ij}\|^2).\end{aligned}\tag{22}$$

By (16), the mixed scalar curvature of M with $k = 3$ is

$$\operatorname{S}_{\operatorname{mix}}(\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3) = \sum_{i < j} n_i n_j (c + \mu_i \mu_j).$$

Note that $\|h_i\|^2 = \|H_i\|^2/n_i$, where

$$h_i(X, Y) = \langle X, Y \rangle \left(\frac{P_j \nabla \mu_i}{\mu_i - \mu_j} + \frac{P_l \nabla \mu_i}{\mu_i - \mu_l} \right), \quad H_i = n_i \left(\frac{P_j \nabla \mu_i}{\mu_i - \mu_j} + \frac{P_l \nabla \mu_i}{\mu_i - \mu_l} \right).$$

Thus (similarly to Example 6), we obtain

$$\|h_i\|^2 - \|H_i\|^2 = n_i(1 - n_i) \left(\frac{\|P_j \nabla \mu_i\|^2}{(\mu_i - \mu_j)^2} + \frac{\|P_l \nabla \mu_i\|^2}{(\mu_i - \mu_l)^2} \right).$$

Next, we consider $\|h_{ij}\|^2$ and $\|T_{ij}\|^2$ for $i \neq j$ along three subsets of $TM \oplus TM$: $\mathcal{D}_i \oplus \mathcal{D}_i$, $\mathcal{D}_j \oplus \mathcal{D}_j$ and $V_{ij} = (\mathcal{D}_i \oplus \mathcal{D}_j) \cup (\mathcal{D}_j \oplus \mathcal{D}_i)$. Using $h_{ij} = \frac{n_i P_l \nabla \mu_i}{\mu_i - \mu_l} + \frac{n_j P_l \nabla \mu_j}{\mu_j - \mu_l}$ for $i \neq j \neq l$, we obtain

$$\|h_{ij}\|^2 - \|H_{ij}\|^2 = n_i(1 - n_i) \frac{\|P_l \nabla \mu_i\|^2}{(\mu_i - \mu_l)^2} + n_j(1 - n_j) \frac{\|P_l \nabla \mu_j\|^2}{(\mu_j - \mu_l)^2} + \|h_{ij}|_{V_{ij}}\|^2,$$

Since \mathcal{D}_i are integrable, we have $\|T_{ij}\|^2 = \|T_{ij}|_{V_{ij}}\|^2$. Given $i \neq j \neq l$, let (e_α) , (ε_β) , (E_γ) be local orthonormal frames of \mathcal{D}_i , \mathcal{D}_j and \mathcal{D}_l , respectively. Using (19), and

$$\|h_{ij}|_{V_{ij}}\|^2 = \frac{1}{4} \sum_{\alpha, \beta, \gamma} \langle \nabla_{e_\alpha} \varepsilon_\beta + \nabla_{\varepsilon_\beta} e_\alpha, E_\gamma \rangle^2, \quad \|T_{ij}|_{V_{ij}}\|^2 = \frac{1}{4} \sum_{\alpha, \beta, \gamma} \langle \nabla_{e_\alpha} \varepsilon_\beta - \nabla_{\varepsilon_\beta} e_\alpha, E_\gamma \rangle^2,$$

we find for $(i, j) \in \{(1, 2), (1, 3), (2, 3)\}$ the equality

$$\|h_{ij}|_{V_{ij}}\|^2 - \|T_{ij}|_{V_{ij}}\|^2 = \sum_{\alpha, \beta, \gamma} \langle \nabla_{e_\alpha} \varepsilon_\beta, E_\gamma \rangle \langle \nabla_{\varepsilon_\beta} e_\alpha, E_\gamma \rangle = \frac{\sum_{\alpha, \beta, \gamma} \|\mathcal{A}(e_\alpha, \varepsilon_\beta, E_\gamma)\|^2}{(\mu_i - \mu_l)(\mu_j - \mu_l)}.$$

Since

$$\frac{1}{(\mu_2 - \mu_3)(\mu_1 - \mu_3)} + \frac{1}{(\mu_1 - \mu_2)(\mu_3 - \mu_2)} + \frac{1}{(\mu_2 - \mu_1)(\mu_3 - \mu_1)} = 0,$$

the sum $\sum_{i < j} (\|h_{ij}|_{V_{ij}}\|^2 - \|T_{ij}|_{V_{ij}}\|^2)$ vanishes. From the above, we have

$$\begin{aligned} \sum_i (\|h_i\|^2 - \|H_i\|^2) + \sum_{i < j} (\|h_{ij}\|^2 - \|H_{ij}\|^2 - \|T_{ij}\|^2) &= 2n_1(1 - n_1) \left(\frac{\|P_2 \nabla \mu_1\|^2}{(\mu_1 - \mu_2)^2} + \frac{\|P_3 \nabla \mu_1\|^2}{(\mu_1 - \mu_3)^2} \right) \\ &+ 2n_2(1 - n_2) \left(\frac{\|P_1 \nabla \mu_2\|^2}{(\mu_2 - \mu_1)^2} + \frac{\|P_3 \nabla \mu_2\|^2}{(\mu_2 - \mu_3)^2} \right) + 2n_3(1 - n_3) \left(\frac{\|P_1 \nabla \mu_3\|^2}{(\mu_3 - \mu_1)^2} + \frac{\|P_2 \nabla \mu_3\|^2}{(\mu_3 - \mu_2)^2} \right). \end{aligned}$$

Note that the terms under the divergence in (22) are

$$\begin{aligned} \sum_i H_i + \sum_{i < j} H_{ij} &= \sum_i n_i \left(\frac{P_j \nabla \mu_i}{\mu_i - \mu_j} + \frac{P_l \nabla \mu_i}{\mu_i - \mu_l} \right) + \sum_{i < j} \left(n_i \frac{P_l \nabla \mu_i}{\mu_i - \mu_l} + n_j \frac{P_l \nabla \mu_j}{\mu_j - \mu_l} \right) \\ &= 2n_1 \left(\frac{P_2 \nabla \mu_1}{\mu_1 - \mu_2} + \frac{P_3 \nabla \mu_1}{\mu_1 - \mu_3} \right) + 2n_2 \left(\frac{P_1 \nabla \mu_2}{\mu_2 - \mu_1} + \frac{P_3 \nabla \mu_2}{\mu_2 - \mu_3} \right) + 2n_3 \left(\frac{P_1 \nabla \mu_3}{\mu_3 - \mu_1} + \frac{P_2 \nabla \mu_3}{\mu_3 - \mu_2} \right). \end{aligned}$$

From the above (21) follows. \square

Using Stokes' theorem to (21) on a closed M , we obtain the following integral formula.

Corollary 6. *In conditions of Theorem 7, for a closed hypersurface $M \subset \bar{M}(c)$, we have*

$$\int_M \left[\sum_{i < j} n_i n_j (c + \mu_i \mu_j) + 2 \sum_i n_i (1 - n_i) \left(\frac{\|P_j \nabla \mu_i\|^2}{(\mu_i - \mu_j)^2} + \frac{\|P_l \nabla \mu_i\|^2}{(\mu_i - \mu_l)^2} \right) \right] d \text{vol}_g = 0,$$

where $(j, l) \in \{1, 2, 3\} \setminus \{i\}$ and $j < l$.

References

- [1] M.A. Akivis, V.V. Goldberg, Differential geometry of webs. In *Handbook of differential geometry*, Vol. I, pp. 1–152, North-Holland, Amsterdam, 2000
- [2] K. Andrzejewski, V. Rovenski and P. Walczak, *Integral formulas in foliation theory*, 73–82, in “*Geometry and its Applications*”, Springer Proc. in Math. and Statistics, 72, Springer, 2014
- [3] A. Bejancu and H. Farran. *Foliations and geometric structures*, Dordrecht: Springer, 2006
- [4] M. Banaszczyk, R. Majchrzak, An integral formula for a Riemannian manifold with three orthogonal distributions. *Acta Sci. Math.*, 54 (1990), 201–207
- [5] A. Caminha, P. Souza, F. Camargo, Complete foliations of space forms by hypersurfaces, *Bull. Braz. Math. Soc., New Series*, 41:3 (2010), 339–353
- [6] T.E. Cecil, P.J. Ryan, *Geometry of Hypersurfaces*, Springer Monographs in Mathematics. Springer-Verlag, New York, 2015
- [7] B.-Y. Chen, *Differential geometry of warped product manifolds and submanifolds*. World Scientific, 2017
- [8] K.L. Duggal, B. Sahin, *Differential Geometry of Lightlike Submanifolds*, Birkhäuser, 2010
- [9] A. Gray, Pseudo-Riemannian almost product manifolds and submersions, *J. Math. Mech.* 16:7 (1967), 715–737
- [10] T. Otsuki, Minimal hypersurfaces with three principal curvatures in S^{n+1} . *Kodai Math. J.* 1 (1978), 1–29
- [11] A. Ranjan, Structural equations and an integral formula for foliated manifolds, *Geom. Dedicata*, 20 (1986), 85–91
- [12] G. Reeb, Sur la courbure moyenne des variétés intégrales d’une équation de Pfaff $\omega = 0$, *C. R. Acad. Sci. Paris* 231 (1950), 101–102
- [13] V. Rovenski, *Foliations on Riemannian Manifolds and Submanifolds*. Birkhäuser, 1998
- [14] S.E. Stepanov and J. Mikeš, Liouville-type theorems for some classes of Riemannian almost product manifolds and for special mappings of Riemannian manifolds, *Differential Geom. and its Appl.* 54, Part A (2017), 111–121
- [15] A. Tarrío, On certain classes of metric para- ϕ -manifolds with parallelizable kernel, *Tebzir*, N.S. 57 (1996), 258–267
- [16] G.A. Tolstikhina, Algebra and geometry of 3-webs formed by foliations of different dimensions. *J. Math. Sci. (N.Y.)* 143 (2007), no. 6, 3630–3721
- [17] P. Walczak, An integral formula for a Riemannian manifold with two orthogonal complementary distributions. *Colloq. Math.*, 58 (1990), 243–252