

# A fractional parabolic inverse problem involving a time-dependent magnetic potential

Li Li

*Department of Mathematics, University of Washington,  
Seattle, WA 98195, USA*

**ABSTRACT.** We study a class of fractional parabolic equations involving a time-dependent magnetic potential and formulate the corresponding inverse problem. We determine both the magnetic potential and the electric potential from the exterior partial measurements of the Dirichlet-to-Neumann map.

## 1 Introduction

The study of fractional operators has been an active research field in past decades. Differential equations involving fractional derivatives have been introduced to describe anomalous diffusion and random processes with jumps in physics and probability theory. See for instance, [19, 20, 25].

Correspondingly, various kinds of inverse problems associated with fractional operators have been extensively studied so far. The study of the inverse problem for space-fractional operators (very different from the one for time-fractional operators studied in [4, 13]), namely the fractional Calderón problem, was initiated in [12] where the authors considered the exterior Dirichlet problem

$$((-\Delta)^s + q)u = 0 \text{ in } \Omega, \quad u|_{\Omega_e} = f$$

and they showed that the electric potential  $q$  in  $\Omega$  can be determined from the exterior partial measurements of the Dirichlet-to-Neumann map

$$\Lambda_q : f \rightarrow (-\Delta)^s u_f|_{\Omega_e}.$$

See [1, 3, 11, 10, 24] for further studies based on [12].

As variants of the inverse problem introduced in [12], parabolic analogues of the fractional Calderón problem have been studied in recent years (see [2, 5] for results in the local parabolic inverse problem). One related work can be found in [15] where the authors studied the inverse problem for the fractional operator

$$(\partial_t - \Delta)^s + q.$$

Another related work can be found in [23] where the authors established the well-posedness of the initial exterior problem associated with the fractional operator

$$\partial_t + (-\Delta)^s$$

and its Runge approximation property.

In this paper, we study an inverse problem for a fractional operator generalizing  $\partial_t + (-\Delta)^s$ . Our operator contains a time-dependent space-fractional derivative and our inverse problem can be viewed as a parabolic analogue of the the fractional magnetic Calderón problem introduced in [16, 17]. See [9, 14, 21, 26] for results in the local magnetic Calderón problem. Also see [6] for the study of a different fractional magnetic Calderón problem.

More precisely, we consider the time-dependent operator  $\mathcal{R}_{A(t)}^s$ , which is formally defined by

$$\langle \mathcal{R}_{A(t)}^s u, v \rangle := 2 \iint (u(x) - R_{A(t)}(x, y)u(y))v(x)K(x, y) dx dy \quad (1)$$

for each  $t$ . Here  $K$  is a function associated with a heat kernel (see Subsection 2.2 in [16] or Section 2 in [10]) satisfying

$$K(x, y) = K(y, x), \quad K(x, y) \sim \frac{1}{|x - y|^{n+2s}},$$

$A(\cdot, t)$  is a time-dependent real vector-valued magnetic potential and

$$R_{A(t)}(x, y) := \cos((x - y) \cdot A(\frac{x + y}{2}, t)). \quad (2)$$

Clearly, the operator  $\mathcal{R}_{A(t)}^s$  coincides with the fractional Laplacian  $(-\Delta)^s$  when  $A \equiv 0$  and  $K(x, y) = c_{n,s}/|x - y|^{n+2s}$ .

Under appropriate assumptions on  $A$  and the time-dependent electric potential  $q(\cdot, t)$ , the initial exterior problem

$$\begin{cases} \partial_t u + \mathcal{R}_{A(t)}^s u + q(t)u = 0 & \text{in } \Omega \times (-T, T) \\ u = g & \text{in } \Omega_e \times (-T, T) \\ u = 0 & \text{in } \mathbb{R}^n \times \{-T\}. \end{cases} \quad (3)$$

is well-posed so we can define the solution operator  $P_{A,q} : g \rightarrow u_g$  and the Dirichlet-to-Neumann map  $\Lambda_{A,q}$ , which is formally given by

$$\Lambda_{A,q}g := \mathcal{R}_A^s(u_g)|_{\Omega_e \times (-T, T)}. \quad (4)$$

Our goal here is to determine both  $A$  and  $q$  from the exterior partial measurements of  $\Lambda_{A,q}$ . The following theorem is the main result in this paper.

**Theorem 1.1.** *Suppose  $\Omega \subset B_r(0)$  for some constant  $r > 0$ ,  $A_j, q_j \in C^2([-T, T]; L^\infty(\Omega))$  and  $q_j \geq c$  for some constant  $c > 0$ ,  $W_j$  are open sets s.t.  $W_j \cap B_{3r}(0) = \emptyset$  ( $j = 1, 2$ ). Let*

$$W^{(1,2)} = \left\{ \frac{x + y}{2} : x \in W_1, y \in W_2 \right\}.$$

*Also assume  $W^{(1,2)} \setminus \Omega \neq \emptyset$ . If*

$$\Lambda_{A_1, q_1} g|_{W_2 \times (-T, T)} = \Lambda_{A_2, q_2} g|_{W_2 \times (-T, T)} \quad (5)$$

*for any  $g \in C_c^\infty(W_1 \times (-T, T))$ , then  $A_1(t) = \pm A_2(t)$  and  $q_1 = q_2$  in  $\Omega \times (-T, T)$ .*

The rest of this paper is organized in the following way. In Section 2, we summarize the background knowledge. We show the well-posedness of the initial exterior problem (3) in Section 3. We introduce the associated Dirichlet-to-Neumann map, prove the Runge approximation property of our fractional operator and the main theorem in Section 4.

**Acknowledgement.** The author is partly supported by National Science Foundation. The author would like to thank Professor Gunther Uhlmann for helpful discussions.

## 2 Preliminaries

Throughout this paper

- Fix the space dimension  $n \geq 2$  and the fractional power  $0 < s < 1$
- Fix the constant  $T > 0$  and  $t$  denotes the time variable
- $\Omega$  denotes a bounded Lipschitz domain and  $\Omega_e := \mathbb{R}^n \setminus \bar{\Omega}$
- $B_r(0)$  denotes the open ball centered at the origin with radius  $r > 0$  in  $\mathbb{R}^n$
- If  $u(\cdot, t)$  is an  $(n+1)$ -variable function, then  $u(t)$  denotes the corresponding  $n$ -variable function for each  $t$
- $A(\cdot, t)$  denotes a time-dependent  $\mathbb{R}^n$ -valued magnetic potential and  $q(\cdot, t)$  denotes a time-dependent electric potential
- If  $A(t) \in L^\infty(\Omega)$ , then identify  $A(t)$  with its zero extension in  $L^\infty(\mathbb{R}^n)$
- $c, C, C', C_1, \dots$  denote positive constants (which may depend on some parameters)
- $\int \cdots \int = \int_{\mathbb{R}^n} \cdots \int_{\mathbb{R}^n}$
- $X^*$  denotes the continuous dual space of  $X$  and write  $\langle f, u \rangle = f(u)$  for  $u \in X, f \in X^*$  when  $X$  is an  $n$ -variable function space.

### 2.1 Function spaces

Throughout this paper we refer all function spaces to real-valued function spaces.

For  $\alpha \in \mathbb{R}$ ,  $H^\alpha(\mathbb{R}^n)$  denotes the Sobolev space  $W^{\alpha,2}(\mathbb{R}^n)$ .

We have the natural identification

$$H^{-\alpha}(\mathbb{R}^n) = H^\alpha(\mathbb{R}^n)^*.$$

Let  $U$  be an open set and  $F$  be a closed set in  $\mathbb{R}^n$ ,

$$H^\alpha(U) := \{u|_U : u \in H^\alpha(\mathbb{R}^n)\}, \quad H_F^\alpha(\mathbb{R}^n) := \{u \in H^\alpha(\mathbb{R}^n) : \text{supp } u \subset F\},$$

$$\tilde{H}^\alpha(U) := \text{the closure of } C_c^\infty(U) \text{ in } H^\alpha(\mathbb{R}^n).$$

$\Omega$  is Lipschitz bounded implies

$$\tilde{H}^\alpha(\Omega) = H_\Omega^\alpha(\mathbb{R}^n).$$

Let  $X$  be a Banach space. We use  $C^m([-T, T]; X)$  (resp.  $AC([-T, T]; X)$ ) to denote the space consisting of the corresponding Banach space-valued continuously differentiable (resp. absolutely continuous) functions on  $[-T, T]$ .

$L^p(-T, T; X)$  denotes the space consisting of the corresponding Banach space-valued  $L^p$  functions, equipped with the standard norm

$$\|u\|_{L^p(-T, T; X)} := \left( \int_{-T}^T \|u(t)\|_X^p dt \right)^{1/p}.$$

## 2.2 The operator $\mathcal{R}_{A(t)}^s$

Recall that we gave the formal definition of  $\mathcal{R}_{A(t)}^s$  in (1). Note that

$$\iint (u(x) - R_{A(t)}(x, y)u(y))v(x)K(x, y) dx dy = \iint (u(y) - R_{A(t)}(x, y)u(x))v(y)K(x, y) dy dx$$

so it is easy to verify that

$$\langle \mathcal{R}_{A(t)}^s u, v \rangle = \operatorname{Re} \iint (u(x) - e^{i(x-y) \cdot A(\frac{x+y}{2}, t)} u(y))(v(x) - e^{-i(x-y) \cdot A(\frac{x+y}{2}, t)} v(y)) K(x, y) dx dy$$

and

$$\langle \mathcal{R}_{A(t)}^s u, v \rangle = \langle \mathcal{R}_{A(t)}^s v, u \rangle. \quad (6)$$

The following lemma is a time-dependent version of Lemma 3.3 in [16].

**Lemma 2.1.** *Suppose  $A \in C([-T, T], L^\infty(\mathbb{R}^n))$ , then for  $u \in H^s(\mathbb{R}^n)$ , we have*

$$c\|u\|_{H^s} \leq \|u\|_{H_{A(t)}^s} \leq C\|u\|_{H^s}$$

where the magnetic Sobolev norm  $\|\cdot\|_{H_{A(t)}^s}$  is defined by

$$\|u\|_{H_{A(t)}^s} := (\|u\|_{L^2}^2 + \langle \mathcal{R}_{A(t)}^s u, u \rangle)^{1/2}$$

and  $c, C$  depend on  $\sup_{t \in [-T, T]} \|A(t)\|_{L^\infty}$  but do not depend on  $t$ .

**Definition 2.2.** *We define the time-dependent bilinear form associated with  $A, q$  by*

$$B_t[u, v] := \langle \mathcal{R}_{A(t)}^s u, v \rangle + \int_{\Omega} q(t)uv, \quad t \in [-T, T] \quad (7)$$

The symmetry of  $B_t$  follows immediately from the symmetry of  $\mathcal{R}_{A(t)}^s$ .

The following estimates will be useful when we show the well-posedness of the initial exterior problem later.

**Lemma 2.3.** *Suppose  $A, q \in C^2([-T, T]; L^\infty(\Omega))$  and  $q \geq c'$  in  $\Omega \times [-T, T]$  for some  $c'$ . Then*

$$|B_t[u, v]| \leq C_0 \|u\|_{H^s} \|v\|_{H^s}, \quad u, v \in H^s(\mathbb{R}^n) \quad (8)$$

and for  $u, v \in \tilde{H}^s(\Omega)$ , we have

$$|B_t[u, u]| \geq c_0 \|u\|_{H^s}^2, \quad (9)$$

$$|B_{t+h}[u, v] - B_t[u, v]| \leq C_1 h \|u\|_{H^s} \|v\|_{H^s}, \quad (10)$$

$$|B_{t+h}[u, v] + B_{t-h}[u, v] - 2B_t[u, v]| \leq C_2 h^2 \|u\|_{H^s} \|v\|_{H^s} \quad (11)$$

where the constants  $c_0, C_0, C_1, C_2$  do not depend on  $t, h, u, v$ .

*Proof.* (8) and (9) follow from Lemma 2.1 immediately.

Note that for  $u, v \in \tilde{H}^s(\Omega)$ , we have

$$\begin{aligned} \frac{d}{dt} \langle \mathcal{R}_{A(t)}^s u, v \rangle &:= -2 \iint \frac{d}{dt} R_{A(t)}(x, y) u(y) v(x) K(x, y) dx dy \\ &= -2 \iint_{\Omega} \frac{d}{dt} R_{A(t)}(x, y) u(y) v(x) K(x, y) dx dy \\ &= 2 \iint_{\Omega} \sin((x-y) \cdot A(\frac{x+y}{2}, t)) (x-y) \cdot \partial_t A(\frac{x+y}{2}, t) u(y) v(x) K(x, y) dx dy \end{aligned}$$

so we have

$$\begin{aligned} \left| \frac{d}{dt} \langle \mathcal{R}_{A(t)}^s u, v \rangle \right| &\leq C \iint_{\Omega} \frac{|u(y)v(x)|}{|x-y|^{n+2s-2}} dx dy \\ &\leq C \left( \iint_{\Omega} \frac{|u(y)|^2}{|x-y|^{n+2s-2}} dx dy \right)^{\frac{1}{2}} \left( \iint_{\Omega} \frac{|v(x)|^2}{|x-y|^{n+2s-2}} dx dy \right)^{\frac{1}{2}}. \end{aligned}$$

Note that

$$\iint_{\Omega} \frac{|u(y)|^2}{|x-y|^{n+2s-2}} dx dy = \int_{\Omega} \left( \int_{\Omega} \frac{1}{|x-y|^{n+2s-2}} dx \right) |u(y)|^2 dy \leq C' \|u\|_{L^2}^2$$

and similarly

$$\iint_{\Omega} \frac{|v(x)|^2}{|x-y|^{n+2s-2}} dx dy \leq C' \|v\|_{L^2}^2$$

so we have

$$\left| \frac{d}{dt} \langle \mathcal{R}_{A(t)}^s u, v \rangle \right| \leq C'' \|u\|_{L^2} \|v\|_{L^2} \leq C'' \|u\|_{H^s} \|v\|_{H^s},$$

which implies (10) holds.

Also note that

$$\begin{aligned} \frac{d^2}{dt^2} R_{A(t)}(x, y) &= -\sin((x-y) \cdot A(\frac{x+y}{2}, t)) (x-y) \cdot \partial_{tt} A(\frac{x+y}{2}, t) \\ &\quad - \cos((x-y) \cdot A(\frac{x+y}{2}, t)) ((x-y) \cdot \partial_t A(\frac{x+y}{2}, t))^2 \end{aligned}$$

so we have

$$\left| \frac{d^2}{dt^2} R_{A(t)}(x, y) \right| \leq C''' |x-y|^2$$

and we can similarly show that

$$\left| \frac{d^2}{dt^2} \langle \mathcal{R}_{A(t)}^s u, v \rangle \right| \leq C'''' \|u\|_{H^s} \|v\|_{H^s}, \quad u, v \in \tilde{H}^s(\Omega),$$

which implies (11) holds. □

We will use the following proposition to prove the Runge approximation property later.

**Proposition 2.4.** (Proposition 2.4 in [18]) Suppose  $\Omega \cup \text{supp } A(t) \subset B_r(0)$  for some  $r > 0$ ,  $W$  is an open set s.t.  $W \cap B_{3r}(0) \neq \emptyset$ . If

$$u \in \tilde{H}^s(\Omega), \quad \mathcal{R}_{A(t)}^s u|_W = 0$$

then  $u = 0$  in  $\mathbb{R}^n$ .

### 3 Initial Exterior Problem

From now on we always assume  $A, q \in C^2([-T, T]; L^\infty(\Omega))$  and  $q \geq c'$  in  $\Omega \times [-T, T]$  for some  $c'$ .

#### 3.1 Discretization in time

First we study the initial value problem

$$\begin{cases} \partial_t u + \mathcal{R}_{A(t)}^s u + q(t)u = f & \text{in } \Omega \times (-T, T) \\ u = 0 & \text{in } \Omega \times \{-T\}. \end{cases} \quad (12)$$

**Proposition 3.1.** *Suppose  $f \in L^2(\Omega \times (-T, T))$  and*

$$\|f(t+h) - f(t)\|_{L^2(\Omega)} \leq Ch \quad (13)$$

for some  $C$  independent of  $t, h$ , then (12) has a unique (weak) solution satisfying

$$u \in L^2(-T, T; \tilde{H}^s(\Omega)) \cap AC([-T, T]; L^2(\Omega)), \quad \partial_t u \in L^2(\Omega \times (-T, T)).$$

**Remark.** *The initial value problem associated with  $\partial_t + (-\Delta)^s$  has been studied in [23] where the authors used a Galerkin approximation to show the existence of solutions. Here the time-dependent fractional operator  $\mathcal{R}_{A(t)}^s$  makes the problem much more complicated. We will use the method of discretization in time instead to show the existence of solutions, which can be viewed as a nonlocal analogue of the Rothe's method for local parabolic problems (see Chapter 15 in [22]).*

*Proof. Existence:* Divide  $[-T, T]$  into  $p$  subintervals of length  $h = \frac{2T}{p} < \frac{1}{2}$  and let  $t_j = -T + jh$ . Consider the discretization in  $t$

$$\begin{cases} \frac{z_j - z_{j-1}}{h} + \mathcal{R}_{A(t_j)}^s z_j + q(t_j)z_j = f(t_j) & j = 1, \dots, p \\ z_0 = 0. \end{cases} \quad (14)$$

We can iteratively determine  $z_j \in \tilde{H}^s(\Omega)$ , which solves the elliptic equation

$$\mathcal{R}_{A(t_j)}^s z_j + (q(t_j) + \frac{1}{h})z_j = f(t_j) + \frac{z_{j-1}}{h}.$$

((8) and (9) ensure the existence and uniqueness of  $z_j$  by Lax-Milgram Theorem.)

Define  $Z_j := (z_j - z_{j-1})/h$  and  $u^{(1)} : [-T, T] \rightarrow \tilde{H}^s(\Omega)$  given by

$$u^{(1)}(t) := z_{j-1} + Z_j(t - t_{j-1}), \quad t \in [t_{j-1}, t_j].$$

Now divide  $[-T, T]$  into  $2^{m-1}p$  subintervals of length  $h = 2T/(2^{m-1}p)$  and let  $t_j^{(m)} = -T + jh_m$ . Similarly, we consider the discretization, obtain a sequence

$$\{z_0^{(m)} = 0, \dots, z_{2^{m-1}p}^{(m)}\}$$

in  $\tilde{H}^s(\Omega)$ , define  $Z_j^{(m)} := (z_j^{(m)} - z_{j-1}^{(m)})/h_m$  and the poly-line

$$u^{(m)}(t) := z_{j-1}^{(m)} + Z_j^{(m)}(t - t_{j-1}^{(m)}), \quad t \in [t_{j-1}^{(m)}, t_j^{(m)}].$$

Also define the step functions

$$\tilde{u}^{(m)}(t) := z_j^{(m)}, \quad \tilde{U}^{(m)}(t) := Z_j^{(m)}, \quad t \in (t_{j-1}^{(m)}, t_j^{(m)}).$$

Note that the constants in Lemma 3.2 do not depend on  $h$  so for general  $m$ ,

$$\|z_j^{(m)}\|_{H^s} \leq c_2, \quad \|Z_j^{(m)}\|_{L^2} \leq c_3$$

hold. This implies the boundedness of  $\{u^{(m)}\}$  in  $L^2(-T, T; \tilde{H}^s(\Omega))$  and the boundedness of  $\{\tilde{U}^{(m)}\}$  in  $L^2(\Omega \times (-T, T))$ . Hence we can choose weakly convergent sequences s.t.

$$u^{(m_k)} \rightharpoonup u \quad \text{in } L^2(-T, T; \tilde{H}^s(\Omega)), \quad \tilde{U}^{(m_k)} \rightharpoonup \tilde{U} \quad \text{in } L^2(\Omega \times (-T, T)).$$

Note that  $\tilde{U}^{(m)}$  is the weak derivative as well as the pointwise derivative of  $u^{(m)}$ . Let  $k \rightarrow \infty$  in

$$u^{(m_k)}(t) = \int_{-T}^t \tilde{U}^{(m_k)}(\tau) d\tau$$

to obtain

$$u(t) = \int_{-T}^t \tilde{U}(\tau) d\tau.$$

Hence  $u$  is absolutely continuous in  $t$ ,  $u(-T) = 0$  and  $\partial_t u(t) = \tilde{U}(t)$ .

Now we show that this  $u$  satisfies the equation in (12).

Define the step function

$$\tilde{f}^{(m)}(t) := f(t_j^{(m)}), \quad t \in (t_{j-1}^{(m)}, t_j^{(m)})$$

and the step bilinear form

$$B_t^{(m)}[\cdot, \cdot] := B_{t_j^{(m)}}[\cdot, \cdot], \quad t \in (t_{j-1}^{(m)}, t_j^{(m)}).$$

Fix  $v \in L^2(-T, T; \tilde{H}^s(\Omega))$  and let both sides of the discretized equation

$$Z_j^{(m)} + \mathcal{R}_{A(t_j^{(m)})}^s z_j^{(m)} + q(t_j^{(m)}) z_j^{(m)} = f(t_j^{(m)})$$

act on  $v(t)$  for  $t \in (t_{j-1}^{(m)}, t_j^{(m)})$  and integrate from  $-T$  to  $T$ , then we have

$$\int_{-T}^T \langle \tilde{U}^{(m)}(t), v(t) \rangle dt + \int_{-T}^T B_t^{(m)}[\tilde{u}^{(m)}(t), v(t)] dt = \int_{-T}^T \langle \tilde{f}^{(m)}(t), v(t) \rangle dt. \quad (15)$$

(13) ensures that

$$\int_{-T}^T \langle \tilde{f}^{(m)}(t), v(t) \rangle dt \rightarrow \int_{-T}^T \langle f(t), v(t) \rangle dt. \quad (16)$$

The weak convergence of  $\tilde{U}^{(m_k)}$  implies

$$\int_{-T}^T \langle \tilde{U}^{(m_k)}(t), v(t) \rangle dt \rightarrow \int_{-T}^T \langle \partial_t u(t), v(t) \rangle dt. \quad (17)$$

Note that (8) ensures that

$$\int_{-T}^T B_t[\cdot, v(t)] dt$$

is a bounded linear functional on  $L^2(-T, T; \tilde{H}^s(\Omega))$  and by Lemma 3.3 we have the weak convergence of  $\{\tilde{u}^{(m_k)}\}$  so

$$\int_{-T}^T B_t[\tilde{u}^{(m_k)}(t), v(t)] dt \rightarrow \int_{-T}^T B_t[u(t), v(t)] dt \quad (18)$$

Also we can show

$$\int_{-T}^T B_t^{(m_k)}[\tilde{u}^{(m_k)}(t), v(t)] dt - \int_{-T}^T B_t[\tilde{u}^{(m_k)}(t), v(t)] dt \rightarrow 0 \quad (19)$$

In fact, we first assume  $v(t) = 1_{[\alpha, \beta]}(t)v_0$  where  $v_0 \in \tilde{H}^s(\Omega)$  and  $\alpha, \beta$  are endpoints of subintervals in the  $m_k$ -division for some  $k$ . For each large  $m_k$ , we write  $\alpha = t_{j_1}^{(m_k)}, \beta = t_{j_2}^{(m_k)}$  for some  $j_1, j_2$ , then

$$\begin{aligned} & \left| \int_{-T}^T B_t^{(m_k)}[\tilde{u}^{(m_k)}(t), v(t)] dt - \int_{-T}^T B_t[\tilde{u}^{(m_k)}(t), v(t)] dt \right| \\ &= \left| \sum_{j=j_1+1}^{j_2} \int_{t_{j-1}^{(m_k)}}^{t_j^{(m_k)}} B_{t_j^{(m_k)}}[z_j^{(m_k)}, v_0] - B_t[z_j^{(m_k)}, v_0] dt \right| \\ &\leq \sum_{j=j_1+1}^{j_2} \int_{t_{j-1}^{(m_k)}}^{t_j^{(m_k)}} C_1 \|z_j^{(m_k)}\|_{H^s} \|v_0\|_{H^s} h_{m_k} dt \leq 2TC_1 c_2 h_{m_k} \|v_0\|_{H^s} \end{aligned}$$

by using (10) and the boundedness of  $\{z_j^{(m)}\}$ .

Since the set consisting of such  $v$  spans a space dense in  $L^2(-T, T; \tilde{H}^s(\Omega))$ , we know (19) holds for all  $v \in L^2(-T, T; \tilde{H}^s(\Omega))$ .

Combine (16), (17), (18), (19) with (15). We conclude that

$$\int_{-T}^T \langle \partial_t u(t), v(t) \rangle dt + \int_{-T}^T B_t[u(t), v(t)] dt = \int_{-T}^T \langle f(t), v(t) \rangle dt. \quad (20)$$

**Uniqueness:** Let  $v = u$  be the solution constructed in the existence part, then (20) becomes

$$\frac{1}{2} \|u(T)\|_{L^2(\Omega)}^2 + \int_{-T}^T B_t[u(t), u(t)] dt = \int_{-T}^T \langle f(t), u(t) \rangle dt,$$

which implies

$$\|u\|_{L^2(-T, T; \tilde{H}^s(\Omega))} \leq C' \|f\|_{L^2(-T, T; H^{-s}(\Omega))}.$$

By (12) and (8),

$$\begin{aligned} \|\partial_t u\|_{L^2(-T, T; H^{-s}(\Omega))} &\leq \|f\|_{L^2(-T, T; H^{-s}(\Omega))} + \|\mathcal{R}_A^s u + qu\|_{L^2(-T, T; H^{-s}(\Omega))} \\ &\leq \|f\|_{L^2(-T, T; H^{-s}(\Omega))} + C'' \|u\|_{L^2(-T, T; \tilde{H}^s(\Omega))}. \end{aligned}$$

Hence we have

$$\|u\|_{L^2(-T, T; \tilde{H}^s(\Omega))} + \|\partial_t u\|_{L^2(-T, T; H^{-s}(\Omega))} \leq \|f\|_{L^2(-T, T; H^{-s}(\Omega))}. \quad (21)$$

The uniqueness is clear when we let  $f = 0$ .  $\square$



We need the following two lemmas to complete the proof of Proposition 3.1. They are essentially the same as their counterparts in the local parabolic problem (see Page 286-294 in [22] for details). We include the proofs here for completeness.

**Lemma 3.2.**  $\{z_j\}$  and  $\{Z_j\}$  defined in the proof of Proposition 3.1 satisfy

$$\|z_j\|_{H^s} \leq c_2, \quad \|Z_j\|_{L^2} \leq c_3$$

where  $c_2, c_3$  do not depend on  $h$ .

*Proof.* We first show that  $\{z_j\}$  is bounded in  $L^2(\Omega)$ .

In fact, let both sides of the equation in (14) act on  $z_j$ , then we have

$$B_{t_j}[z_j, z_j] + \frac{1}{h}\langle z_j - z_{j-1}, z_j \rangle = \langle f(t_j), z_j \rangle,$$

which implies

$$\|z_j\|_{L^2} \leq \|z_{j-1}\|_{L^2} + h\|f(t_j)\|_{L^2},$$

then iteratively we can show

$$\|z_j\|_{L^2} \leq h \sum_{l=1}^j \|f(t_l)\|_{L^2} \leq jhC_f \leq 2TC_f =: c_1.$$

Next we show that  $\{z_j\}$  is bounded in  $\tilde{H}^s(\Omega)$ .

In fact, let both sides of the equation in (14) act on  $z_j - z_{j-1}$ , then we have

$$B_{t_j}[z_j, z_j - z_{j-1}] + \frac{1}{h}\|z_j - z_{j-1}\|_{L^2}^2 = \langle f(t_j), z_j - z_{j-1} \rangle. \quad (22)$$

Note that

$$B_{t_l}[z_l, z_l - z_{l-1}] = \frac{1}{2}(B_{t_l}[z_l, z_l] + B_{t_l}[z_l - z_{l-1}, z_l - z_{l-1}] - B_{t_l}[z_{l-1}, z_{l-1}]),$$

then sum up all the identities in the form (22) for  $1 \leq l \leq j$  and omit all the non-negative terms

$$\|z_l - z_{l-1}\|_{L^2}, \quad B_{t_l}[z_l - z_{l-1}, z_l - z_{l-1}]$$

to obtain the inequality

$$B_{t_j}[z_j, z_j] - \sum_{l=1}^{j-1} (B_{t_{l+1}}[z_l, z_l] - B_{t_l}[z_l, z_l]) \leq 2\langle f(t_j), z_j \rangle + \sum_{l=1}^{j-1} \langle f(t_l) - f(t_{l+1}), z_l \rangle.$$

By (9), (10) and (13), this inequality implies

$$\begin{aligned} c_0\|z_j\|_{H^s}^2 &\leq C_1h \sum_{l=1}^{j-1} \|z_l\|_{H^s}^2 + 2(\|f(t_j)\|_{L^2}\|z_j\|_{L^2} + \sum_{l=1}^{j-1} \|f(t_l) - f(t_{l+1})\|_{L^2}\|z_l\|_{L^2}) \\ &\leq C_1h \sum_{l=1}^{j-1} \|z_l\|_{H^s}^2 + 2c_1C_f + 2c_1C(j-1)h. \end{aligned}$$

Since  $(j-1)h \leq 2T$ , we have

$$\|z_j\|_{H^s}^2 \leq C'_2 + C'_1 h \sum_{l=1}^{j-1} \|z_l\|_{H^s}^2$$

where  $C'_1, C'_2$  do not depend on  $h$ . This implies

$$\|z_j\|_{H^s}^2 \leq C'_2 e^{C'_1(j-1)h}, \quad \|z_j\|_{H^s} \leq (C'_2 e^{2TC'_1})^{\frac{1}{2}} =: c_2.$$

Now we show  $\{Z_j\}$  is bounded in  $L^2(\Omega)$ .

In fact, combine the consecutive equations in (14) to get

$$Z_j - Z_{j-1} + (\mathcal{R}_{A(t_j)}^s z_j + q(t_j)z_j) - (\mathcal{R}_{A(t_{j-1})}^s z_{j-1} + q(t_{j-1})z_{j-1}) = f(t_j) - f(t_{j-1}).$$

Let both sides act on  $Z_j$ , then we have

$$\langle Z_j - Z_{j-1}, Z_j \rangle + B_{t_j}[z_j, Z_j] - B_{t_{j-1}}[z_{j-1}, Z_j] = \langle f(t_j) - f(t_{j-1}), Z_j \rangle. \quad (23)$$

Note that

$$\begin{aligned} B_{t_j}[z_j, Z_j] &= \frac{1}{2h}(B_{t_j}[z_j, z_j] - B_{t_j}[z_{j-1}, z_{j-1}] + h^2 B_{t_j}[Z_j, Z_j]), \\ B_{t_{j-1}}[z_{j-1}, Z_j] &= \frac{1}{2h}(B_{t_{j-1}}[z_j, z_j] - B_{t_{j-1}}[z_{j-1}, z_{j-1}] - h^2 B_{t_{j-1}}[Z_j, Z_j]) \end{aligned}$$

so we have

$$\begin{aligned} & B_{t_j}[z_j, Z_j] - B_{t_{j-1}}[z_{j-1}, Z_j] \\ &= \frac{h}{2}(B_{t_j}[Z_j, Z_j] + B_{t_{j-1}}[Z_j, Z_j]) + \frac{1}{2h}(B_{t_j}[z_j, z_j] - B_{t_{j-1}}[z_j, z_j] + B_{t_{j-1}}[z_{j-1}, z_{j-1}] - B_{t_j}[z_{j-1}, z_{j-1}]). \end{aligned}$$

Also note that

$$\begin{aligned} \langle Z_j - Z_{j-1}, Z_j \rangle &= \frac{1}{2}(\|Z_j\|_{L^2}^2 + \|Z_j - Z_{j-1}\|_{L^2}^2 - \|Z_{j-1}\|_{L^2}^2), \\ \left| \left\langle \frac{f(t_j) - f(t_{j-1})}{h}, Z_j \right\rangle \right| &\leq \frac{1}{2}(\|Z_j\|_{L^2}^2 + C^2). \end{aligned}$$

Now sum up all the identities in the form (23) for  $2 \leq l \leq j$  and omit all the non-negative terms

$$B_{t_l}[Z_l, Z_l], \quad B_{t_{l-1}}[Z_l, Z_l], \quad \|Z_l - Z_{l-1}\|_{L^2}^2$$

to obtain the inequality

$$\begin{aligned} & \|Z_j\|_{L^2}^2 - \|Z_1\|_{L^2}^2 + \frac{1}{h}(B_{t_j}[z_j, z_j] - B_{t_{j-1}}[z_j, z_j]) \\ &+ \frac{1}{h} \sum_{l=2}^{j-1} (-B_{t_{l+1}}[z_l, z_l] + 2B_{t_l}[z_l, z_l] - B_{t_{l-1}}[z_l, z_l]) + \frac{1}{h}(B_{t_1}[z_1, z_1] - B_{t_2}[z_1, z_1]) \\ &\leq h \sum_{l=2}^j (\|Z_l\|_{L^2}^2 + C^2). \end{aligned} \quad (24)$$

By (10) and (11), we have

$$|B_{t_l}[z_l, z_l] - B_{t_{l-1}}[z_l, z_l]| \leq C_1 h \|z_l\|_{H^s}^2 \leq c_2^2 C_1 h,$$

$$|-B_{t_{l+1}}[z_l, z_l] + 2B_{t_l}[z_l, z_l] - B_{t_{l-1}}[z_l, z_l]| \leq C_2 h^2 \|z_l\|_{H^s}^2 \leq c_2^2 C_2 h^2$$

so (24) implies

$$\|Z_j\|_{L^2}^2 \leq \|Z_1\|_{L^2}^2 + (j-1)hC^2 + 2c_2^2 C_1 + (j-2)c_2^2 C_2 h + h \sum_{l=2}^j \|Z_l\|_{L^2}^2.$$

Since  $\|Z_1\|_{L^2} \leq C_f$  and  $(j-1)h \leq 2T$ , we can write

$$\|Z_j\|_{L^2}^2 \leq C_2'' + h \sum_{l=2}^j \|Z_l\|_{L^2}^2$$

where  $C_2''$  does not depend on  $h$ . Since  $h < 1/2$ , we have the estimate

$$\|Z_j\|_{L^2}^2 \leq 2C_2'' e^{2(j-1)h}, \quad \|Z_j\|_{L^2} \leq (2C_2'' e^{4T})^{\frac{1}{2}} =: c_3.$$

□

**Lemma 3.3.**  $u$  and  $\{\tilde{u}^{(m_k)}\}$  defined in the proof of Proposition 3.1 satisfy

$$\tilde{u}^{(m_k)} \rightharpoonup u \quad \text{in } L^2(-T, T; \tilde{H}^s(\Omega)).$$

*Proof.* Since  $u^{(m_k)} \rightharpoonup u$  in  $L^2(-T, T; \tilde{H}^s(\Omega))$ , we only need to show that

$$u^{(m_k)} - \tilde{u}^{(m_k)} \rightharpoonup 0 \quad \text{in } L^2(-T, T; \tilde{H}^s(\Omega)).$$

Consider  $v(t) = 1_{[\alpha, \beta]}(t)v_0$  where  $v_0 \in H^{-s}(\Omega)$  and  $\alpha, \beta$  are endpoints of subintervals in the  $m_k$ -division for some  $k$ . For each large  $m_k$ , we write  $\alpha = t_{j_1}^{(m_k)}, \beta = t_{j_2}^{(m_k)}$  for some  $j_1, j_2$ , then

$$\begin{aligned} \int_{-T}^T \langle v(t), u^{(m_k)}(t) - \tilde{u}^{(m_k)}(t) \rangle dt &= \sum_{j=j_1+1}^{j_2} \int_{t_{j-1}^{(m_k)}}^{t_j^{(m_k)}} \langle v_0, u^{(m_k)}(t) - \tilde{u}^{(m_k)}(t) \rangle dt \\ &= \sum_{j=j_1+1}^{j_2} \int_{t_{j-1}^{(m_k)}}^{t_j^{(m_k)}} \langle v_0, (z_j^{(m_k)} - z_{j-1}^{(m_k)}) \frac{t - t_j^{(m_k)}}{h_{m_k}} \rangle dt \\ &= \sum_{j=j_1+1}^{j_2} \frac{h_{m_k}}{2} \langle v_0, (z_{j-1}^{(m_k)} - z_j^{(m_k)}) \rangle = \frac{h_{m_k}}{2} \langle v_0, (z_{j_1}^{(m_k)} - z_{j_2}^{(m_k)}) \rangle. \end{aligned}$$

By the boundedness of  $\{z_j^{(m)}\}$ , it converges to zero.

By using a density argument, we can conclude that

$$\int_{-T}^T \langle v(t), u^{(m_k)}(t) - \tilde{u}^{(m_k)}(t) \rangle dt \rightarrow 0$$

for general  $v \in L^2(-T, T; H^{-s}(\Omega))$ .

□

### 3.2 Well-posedness

Now we consider (12) for general  $f \in L^2(\Omega \times (-T, T))$ .

In fact, we can choose  $f_m$  satisfying (13) s.t.  $f_m \rightarrow f$  in  $L^2(\Omega \times (-T, T))$ .

Let  $u_m$  be the solution corresponding to  $f_m$  in (12), then we have

$$\|u_m - u_l\|_{L^2(-T, T; \tilde{H}^s(\Omega))} + \|\partial_t(u_m - u_l)\|_{L^2(-T, T; H^{-s}(\Omega))} \leq \|f_m - f_l\|_{L^2(-T, T; H^{-s}(\Omega))}$$

so

$$u_m \rightarrow u \quad \text{in } L^2(-T, T; \tilde{H}^s(\Omega)), \quad \partial_t u_m \rightarrow v \quad \text{in } L^2(-T, T; H^{-s}(\Omega))$$

for some  $u, v$  and  $\partial_t u = v$ .

This implies the convergence of  $\{u_m\}$  in  $C([-T, T]; L^2(\Omega))$  (see, for instance, Theorem 1 in Section 1.2 in Chapter 18 in [7]) and this  $u$  satisfies the estimate (21).

Hence we reach the following conclusion.

**Corollary 3.4.** *Suppose  $f \in L^2(\Omega \times (-T, T))$ , then (12) has a unique (weak) solution satisfying*

$$u \in L^2(-T, T; \tilde{H}^s(\Omega)) \cap C([-T, T]; L^2(\Omega)), \quad \partial_t u \in L^2(-T, T; H^{-s}(\Omega)).$$

From now on we always assume  $\Omega \subset B_r(0)$  for some constant  $r > 0$  and  $W$  is an open set in  $\mathbb{R}^n$  s.t.  $W \cap B_{3r}(0) = \emptyset$ .

**Proposition 3.5.** *Suppose  $g \in C_c^\infty(W \times (-T, T))$ , then (3) has a unique (weak) solution  $u$  satisfying*

$$w \in L^2(-T, T; \tilde{H}^s(\Omega)) \cap C([-T, T]; L^2(\Omega)), \quad \partial_t w \in L^2(-T, T; H^{-s}(\Omega))$$

where  $w := u - g$ .

*Proof.* By (1), we have

$$\mathcal{R}_{A^s} g|_{\Omega \times (-T, T)} = (-\Delta)^s g|_{\Omega \times (-T, T)}$$

where  $(-\Delta)^s$  acts on space variables. This is because  $(x, y) \in \Omega \times W$  implies  $|(x + y)/2| \geq r$  and thus  $A(t) = 0$ ,  $R_{A(t)} = 1$ . Consider the problem

$$\begin{cases} \partial_t w + \mathcal{R}_{A(t)}^s w + q(t)w = f & \text{in } \Omega \times (-T, T) \\ u = 0 & \text{in } \Omega \times \{-T\} \end{cases} \quad (25)$$

where  $f := -\mathcal{R}_{A^s} g|_{\Omega \times (-T, T)}$ . Since

$$(-\Delta)^s : H^\alpha(\mathbb{R}^n) \rightarrow H^{\alpha-2s}(\mathbb{R}^n), \quad \alpha \in \mathbb{R}$$

(see Lemma 2.1 in [12]), it is clear that  $f \in L^2(\Omega \times (-T, T))$ . Now apply the Corollary above.  $\square$

Consider the substitutions  $\tilde{A}(x, t) := A(x, -t)$ ,  $\tilde{q}(x, t) := q(x, -t)$ ,  $\tilde{g}(x, t) := g(x, -t)$  and  $\tilde{u}(x, t) := u(x, -t)$ . Then we know the proposition above holds for the dual problem

$$\begin{cases} -\partial_t u + \mathcal{R}_{A(t)}^s u + q(t)u = 0 & \text{in } \Omega \times (-T, T) \\ u = g & \text{in } \Omega_e \times (-T, T) \\ u = 0 & \text{in } \mathbb{R}^n \times \{T\}. \end{cases} \quad (26)$$

We denote the solution operator  $g \rightarrow u_g$  associated with (3) (resp. (26)) by  $P_{A, q}$  (resp.  $P_{A, q}^*$ ).

## 4 Inverse Problem

### 4.1 Dirichlet-to-Neumann map

Proposition 3.5 ensures that the Dirichlet-to-Neumann map  $\Lambda_{A,q}$  given by (4) is well-defined at least for  $g \in C_c^\infty(W \times (-T, T))$ .

Now let  $g \in C_c^\infty(W_1 \times (-T, T))$  and  $h \in C_c^\infty(W_2 \times (-T, T))$ .

By the definition of the solution operator  $P_{A,q}$  we have

$$\int_{-T}^T \langle \Lambda_{A,q} g(t), h(t) \rangle dt = \int_{-T}^T B_t[u(t), \tilde{h}(t)] dt + \int_{-T}^T \langle \partial_t u(t), \tilde{h}(t) \rangle_\Omega dt \quad (27)$$

for any  $\tilde{h}$  satisfying  $\tilde{h} - h \in L^2(-T, T; \tilde{H}^s(\Omega))$ . Here  $u := P_{A,q}g$ ,  $w := u - g$  and

$$\langle \partial_t u(t), \tilde{h}(t) \rangle_\Omega := \langle \partial_t w(t), \tilde{h}(t) - h(t) \rangle.$$

Similarly we can define

$$\Lambda_{A,q}^* h := \mathcal{R}_A^s u^*|_{\Omega_e \times (-T, T)}$$

where  $u^* := P_{A,q}^* h$  and we have

$$\int_{-T}^T \langle \Lambda_{A,q}^* h(t), g(t) \rangle dt = \int_{-T}^T B_t[u^*(t), \tilde{g}(t)] dt + \int_{-T}^T \langle -\partial_t u^*(t), \tilde{g}(t) \rangle_\Omega dt \quad (28)$$

for any  $\tilde{g}$  satisfying  $\tilde{g} - g \in L^2(-T, T; \tilde{H}^s(\Omega))$ .

**Proposition 4.1.** *For  $g \in C_c^\infty(W_1 \times (-T, T))$  and  $h \in C_c^\infty(W_2 \times (-T, T))$ , we have*

$$\int_{-T}^T \langle \Lambda_{A,q} g(t), h(t) \rangle dt = \int_{-T}^T \langle \Lambda_{A,q}^* h(t), g(t) \rangle dt. \quad (29)$$

*Proof.* Let  $\tilde{h} = u^*$  in (27) and let  $\tilde{g} = u$  in (28). Since  $u(-T) = u^*(T) = 0$ , we have

$$\begin{aligned} & \int_{-T}^T \langle \Lambda_{A,q} g(t), h(t) \rangle dt - \int_{-T}^T \langle \Lambda_{A,q}^* h(t), g(t) \rangle dt \\ &= \int_{-T}^T \langle \partial_t u(t), u^*(t) \rangle_\Omega + \langle \partial_t u^*(t), u(t) \rangle_\Omega dt = \langle u(t), u^*(t) \rangle_\Omega \Big|_{t=-T}^{t=T} = 0. \end{aligned}$$

□

Now we build the integral identity, which will be used in the proof of the main theorem.

For  $g_j \in C_c^\infty(W_j \times (-T, T))$  ( $j = 1, 2$ ), let  $u_1 = P_{A_1, q_1}(g_1)$  and  $u_2^* = P_{A_2, q_2}^*(g_2)$ , i.e.  $u_1$  is the unique weak solution of

$$\begin{cases} \partial_t u + \mathcal{R}_{A_1(t)}^s u + q_1(t)u = 0 & \text{in } \Omega \times (-T, T) \\ u = g_1 & \text{in } \Omega_e \times (-T, T) \\ u = 0 & \text{in } \mathbb{R}^n \times \{-T\}. \end{cases} \quad (30)$$

and  $u_2^*$  is the unique weak solution of

$$\begin{cases} -\partial_t u + \mathcal{R}_{A_2(t)}^s u + q_2(t)u = 0 & \text{in } \Omega \times (-T, T) \\ u = g_2 & \text{in } \Omega_e \times (-T, T) \\ u = 0 & \text{in } \mathbb{R}^n \times \{T\}. \end{cases} \quad (31)$$

then we have

$$\begin{aligned} & \int_{-T}^T \langle \Lambda_{A_1, q_1} g_1(t), g_2(t) \rangle - \langle \Lambda_{A_2, q_2} g_1(t), g_2(t) \rangle dt \\ &= \int_{-T}^T \langle \Lambda_{A_1, q_1} g_1(t), g_2(t) \rangle dt - \int_{-T}^T \langle \Lambda_{A_2, q_2}^* g_2(t), g_1(t) \rangle dt \\ &= \int_{-T}^T B_t^{(1)}[u_1(t), u_2^*(t)] + \langle \partial_t u_1(t), u_2^*(t) \rangle_\Omega dt - \int_{-T}^T B_t^{(2)}[u_2^*(t), u_1(t)] + \langle -\partial_t u_2^*(t), u_1(t) \rangle_\Omega dt \\ &= \int_{-T}^T B_t^{(1)}[u_1(t), u_2^*(t)] dt - \int_{-T}^T B_t^{(2)}[u_1(t), u_2^*(t)] dt \\ &= \int_{-T}^T \iint G(x, y, t) u_1(y, t) u_2^*(x, t) - \int_{-T}^T \int_\Omega (q_2 - q_1) u_1 u_2^* \end{aligned} \quad (32)$$

where

$$G(x, y, t) := 2(R_{A_2(t)}(x, y) - R_{A_1(t)}(x, y))K(x, y).$$

## 4.2 Runge approximation

**Proposition 4.2.** *Suppose  $\Omega \subset B_r(0)$  for some constant  $r > 0$  and  $W$  is an open set in  $\mathbb{R}^n$  s.t.  $W \cap B_{3r}(0) = \emptyset$ , then*

$$\begin{aligned} S &:= \{P_{A, q} g|_{\Omega \times (-T, T)} : g \in C_c^\infty(W \times (-T, T))\}, \\ S^* &:= \{P_{A, q}^* g|_{\Omega \times (-T, T)} : g \in C_c^\infty(W \times (-T, T))\} \end{aligned}$$

are dense in  $L^2(\Omega \times (-T, T))$ .

*Proof.* By the Hahn-Banach Theorem, it suffices to show that:

If  $v \in L^2(\Omega \times (-T, T))$  and  $\int_{-T}^T \int_\Omega v w = 0$  for all  $w \in S$ , then  $v = 0$  in  $\Omega \times (-T, T)$ .

In fact, for a given  $v \in L^2(\Omega \times (-T, T))$ , let  $\phi \in L^2(-T, T; \tilde{H}^s(\Omega))$  be the solution of

$$\begin{cases} -\partial_t \phi + \mathcal{R}_{A(t)}^s \phi + q(t)\phi = v & \text{in } \Omega \times (-T, T) \\ \phi = 0 & \text{in } \Omega \times \{T\}. \end{cases} \quad (33)$$

For  $g \in C_c^\infty(W \times (-T, T))$ , write  $u_g := P_{A, q} g$ , then we have

$$\begin{aligned} & \int_{-T}^T \int_\Omega v u_g = \int_{-T}^T \langle -\partial_t \phi(t) + \mathcal{R}_{A(t)}^s \phi(t) + q(t)\phi, u_g(t) - g(t) \rangle dt \\ &= \int_{-T}^T \langle \partial_t u_g(t), \phi(t) \rangle + B_t[u_g(t), \phi(t)] dt - \int_{-T}^T \langle \mathcal{R}_{A(t)}^s g(t), \phi(t) \rangle dt = - \int_{-T}^T \langle \mathcal{R}_{A(t)}^s \phi(t), g(t) \rangle dt. \end{aligned}$$

The first equality holds since  $u_g - g \in L^2(-T, T; \tilde{H}^s(\Omega))$ , the second equality holds since  $u_g(-T) = \phi(T) = 0$  and the last equality holds since  $\phi \in L^2(-T, T; \tilde{H}^s(\Omega))$  and  $u_g$  is the solution of (3).

Hence, if  $\int_{-T}^T \int_{\Omega} v w = 0$  for all  $w \in S$ , then for each  $t$  we have

$$\phi(t) \in \tilde{H}^s(\Omega), \quad \mathcal{R}_{A(t)}^s \phi(t)|_W = 0,$$

which implies  $\phi(t) = 0$  in  $\mathbb{R}^n$  for each  $t$  by Proposition 2.4 and thus  $v = 0$  in  $\Omega \times (-T, T)$ .

Similarly we can show  $S^*$  is dense in  $L^2(\Omega \times (-T, T))$ .  $\square$

**Remark.** Proposition 4.2 can be viewed as a generalization of Theorem 2 in [23]. We refer readers to [8, 23] for more approximation properties of solutions of nonlocal evolution problems.

### 4.3 Proof of the main theorem

Now we are ready to prove Theorem 1.1. As in the proof of Theorem 1.1 in [17], we exploit the integral identity and the Runge approximation property associated with our operator.

*Proof.* Write  $u_1 = P_{A_1, q_1}(g_1)$  and  $u_2^* = P_{A_2, q_2}^*(g_2)$  for  $g_j \in C_c^\infty(W_j \times (-T, T))$ .

As in the proof of Theorem 1.1 in [17], the assumptions on  $W_1, W_2, W^{(1,2)}$  ensure that

$$\iint G(x, y, t) u_1(y, t) u_2^*(x, t) dx dy = \iint_{\Omega} G(x, y, t) u_1(y, t) u_2^*(x, t) dx dy$$

for each  $t$  (if we shrink  $W_1, W_2$  when necessary).

By the integral identity (32), (5) implies

$$\int_{-T}^T \int_{\Omega} \int_{\Omega} G(x, y, t) u_1(y, t) u_2^*(x, t) = \int_{-T}^T \int_{\Omega} (q_2 - q_1) u_1 u_2^*. \quad (34)$$

**Determine A:** Fix open sets  $\Omega_j \subset \Omega$  s.t.  $\Omega_1 \cap \Omega_2 = \emptyset$ . Also fix  $\phi_j \in C_c^\infty(\Omega_j)$  and the constants  $a, b \in (-T, T)$  and  $\epsilon > 0$ . Write

$$\tilde{\phi}_j(x, t) := 1_{[a, b]}(t) \phi_j(x).$$

By Proposition 4.2, we can choose  $g_1 \in C_c^\infty(W_1 \times (-T, T))$  s.t.

$$\|u_1 - \tilde{\phi}_1\|_{L^2(\Omega \times (-T, T))} \leq \epsilon$$

and for this chosen  $g_1$ , we can choose  $g_2 \in C_c^\infty(W_2 \times (-T, T))$  s.t.

$$\|u_1\|_{L^2(\Omega \times (-T, T))} \|u_2^* - \tilde{\phi}_2\|_{L^2(\Omega \times (-T, T))} \leq \epsilon.$$

Note that  $\phi_1(x) \phi_2(x) = 0$  for  $x \in \Omega$  so

$$\begin{aligned} \left| \int_{-T}^T \int_{\Omega} (q_2 - q_1) u_1 u_2^* \right| &= \left| \int_{-T}^T \int_{\Omega} (q_2 - q_1) (u_1 - \tilde{\phi}_1) \tilde{\phi}_2 + \int_{-T}^T \int_{\Omega} (q_2 - q_1) u_1 (u_2^* - \tilde{\phi}_2) \right| \\ &\leq \|(q_2 - q_1)\|_{L^\infty} \|\tilde{\phi}_2\|_{L^2} \|u_1 - \tilde{\phi}_1\|_{L^2} + \|(q_2 - q_1)\|_{L^\infty} \|u_1\|_{L^2} \|u_2^* - \tilde{\phi}_2\|_{L^2} \leq C\epsilon. \end{aligned} \quad (35)$$

Also note that

$$|G(x, y, t)| \leq \frac{C'}{|x - y|^{n+2s-2}},$$

which implies

$$\int_{\Omega} |G(x, y, t)| dy \leq C'', \quad x \in \Omega, \quad \int_{\Omega} |G(x, y, t)| dx \leq C'', \quad y \in \Omega$$

where  $C', C''$  do not depend on  $t$ . By the generalized Young's Inequality,

$$\|T_t f\|_{L^2(\Omega)} \leq C'' \|f\|_{L^2(\Omega)}, \quad (T_t f)(x) := \int_{\Omega} |G(x, y, t) f(y)| dy.$$

Now note that

$$\begin{aligned} & \int_{-T}^T \int_{\Omega} \int_{\Omega} G(x, y, t) u_1(y, t) u_2^*(x, t) dx dy dt - \int_a^b \int_{\Omega_1} \int_{\Omega_2} G(x, y, t) \phi_1(y) \phi_2(x) dx dy dt \\ &= \int_{-T}^T \int_{\Omega} \int_{\Omega} G(x, y, t) (u_1(y, t) - \tilde{\phi}_1(y, t)) \tilde{\phi}_2(x, t) dx dy dt \\ &+ \int_{-T}^T \int_{\Omega} \int_{\Omega} G(x, y, t) u_1(y, t) (u_2^*(x, t) - \tilde{\phi}_2(x, t)) dx dy dt. \end{aligned}$$

By Cauchy-Schwarz inequality, we have the estimate

$$\begin{aligned} & \int_{-T}^T \int_{\Omega} \left( \int_{\Omega} |G(x, y, t) u_1(y, t)| dy \right) |u_2^*(x, t) - \tilde{\phi}_2(x, t)| dx dt \\ & \leq \left( \int_{-T}^T \int_{\Omega} |(T_t u_1(t))(x)|^2 dx dt \right)^{\frac{1}{2}} \|u_2^* - \tilde{\phi}_2\|_{L^2(\Omega \times (-T, T))} \\ & \leq C'' \left( \int_{-T}^T \|u_1(t)\|_{L^2(\Omega)}^2 dt \right)^{\frac{1}{2}} \|u_2^* - \tilde{\phi}_2\|_{L^2(\Omega \times (-T, T))} = C'' \|u_1\|_{L^2(\Omega \times (-T, T))} \|u_2^* - \tilde{\phi}_2\|_{L^2(\Omega \times (-T, T))}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} & \int_{-T}^T \int_{\Omega} \left( \int_{\Omega} |G(x, y, t) \tilde{\phi}_2(x, t)| dx \right) |u_1(y, t) - \tilde{\phi}_1(y, t)| dy dt \\ & \leq C'' \|\tilde{\phi}_2\|_{L^2(\Omega \times (-T, T))} \|u_1 - \tilde{\phi}_1\|_{L^2(\Omega \times (-T, T))}. \end{aligned}$$

Hence

$$\left| \int_{-T}^T \int_{\Omega} \int_{\Omega} G(x, y, t) u_1(y, t) u_2^*(x, t) - \int_a^b \int_{\Omega_1} \int_{\Omega_2} G(x, y, t) \phi_1(y) \phi_2(x) \right| \leq C''' \epsilon. \quad (36)$$

Combine (35), (36) with (34).  $\epsilon$  is arbitrary implies

$$\int_a^b \int_{\Omega_1} \int_{\Omega_2} G(x, y, t) \phi_1(y) \phi_2(x) dx dy dt = 0.$$

Then  $[a, b]$  is arbitrary implies

$$\int_{\Omega_1} \int_{\Omega_2} G(x, y, t) \phi_1(y) \phi_2(x) dx dy = 0$$



for each  $t$  and thus  $G(x, y, t) = 0$  in  $\Omega_1 \times \Omega_2$  for each  $t$  since  $\phi_1, \phi_2$  are arbitrary. Now we can conclude that  $G(x, y, t) = 0$  for  $x, y \in \Omega$  whenever  $x \neq y$  since  $\Omega_1, \Omega_2$  are arbitrary. Hence

$$R_{A_1(t)}(x, y) = R_{A_2(t)}(x, y), \quad x, y \in \Omega \quad (37)$$

for each  $t$ , which implies  $A_1(t) = \pm A_2(t)$  as in the proof of Theorem 1.1 in [17].

**Determine  $q$ :** Now (34) becomes

$$\int_{-T}^T \int_{\Omega} (q_2 - q_1) u_1 u_2^* = 0.$$

Fix  $\epsilon > 0$  and  $f \in L^2(\Omega \times (-T, T))$ . Choose  $g_1 \in C_c^\infty(W_1 \times (-T, T))$  s.t.

$$\|u_1 - f\|_{L^2(\Omega \times (-T, T))} \leq \epsilon.$$

For this chosen  $u_1$ , choose  $g_2 \in C_c^\infty(W_2)$  s.t.

$$\|u_1\|_{L^2(\Omega \times (-T, T))} \|u_2^* - 1\|_{L^2(\Omega \times (-T, T))} \leq \epsilon.$$

Now we have

$$\left| \int_{-T}^T \int_{\Omega} (q_1 - q_2) f \right| = \left| \int_{-T}^T \int_{\Omega} (q_1 - q_2) (f - u_1) + \int_{-T}^T \int_{\Omega} (q_1 - q_2) u_1 (1 - u_2^*) \right| \leq C\epsilon.$$

We conclude that  $q_1 = q_2$  since  $\epsilon, f$  are arbitrary. □

## References

- [1] Sombuddha Bhattacharyya, Tuhin Ghosh, and Gunther Uhlmann. Inverse problem for fractional-laplacian with lower order non-local perturbations. *(to appear) Transactions of the American Mathematical Society*, 2020.
- [2] Bruno Canuto and Otared Kavian. Determining coefficients in a class of heat equations via boundary measurements. *SIAM Journal on Mathematical Analysis*, 32(5):963–986, 2001.
- [3] Mihajlo Cekić, Yi-Hsuan Lin, and Angkana Rüland. The Calderón problem for the fractional Schrödinger equation with drift. *arXiv preprint arXiv:1810.04211*, 2018.
- [4] Jin Cheng, Junichi Nakagawa, Masahiro Yamamoto, and Tomohiro Yamazaki. Uniqueness in an inverse problem for a one-dimensional fractional diffusion equation. *Inverse problems*, 25(11):115002, 2009.
- [5] Mourad Choulli and Masahiro Yamamoto. Conditional stability in determining a heat source. *Journal of inverse and ill-posed problems*, 12(3):233–243, 2004.
- [6] Giovanni Covi. An inverse problem for the fractional Schrödinger equation in a magnetic field. *Inverse Problems*, 36(4):045004, 2020.
- [7] Robert Dautary and Jacques-Louis Lions. Mathematical analysis and numerical methods for science and technology, vol. 5, 1992.

- [8] Serena Dipierro, Ovidiu Savin, and Enrico Valdinoci. Local approximation of arbitrary functions by solutions of nonlocal equations. *The Journal of Geometric Analysis*, 29(2):1428–1455, 2019.
- [9] David Dos Santos Ferreira, Carlos E Kenig, Johannes Sjöstrand, and Gunther Uhlmann. Determining a magnetic Schrödinger operator from partial cauchy data. *Communications in mathematical physics*, 271(2):467–488, 2007.
- [10] Tuhin Ghosh, Yi-Hsuan Lin, and Jingni Xiao. The Calderón problem for variable coefficients nonlocal elliptic operators. *Communications in Partial Differential Equations*, 42(12):1923–1961, 2017.
- [11] Tuhin Ghosh, Angkana Rüland, Mikko Salo, and Gunther Uhlmann. Uniqueness and reconstruction for the fractional Calderón problem with a single measurement. *Journal of Functional Analysis*, page 108505, 2020.
- [12] Tuhin Ghosh, Mikko Salo, and Gunther Uhlmann. The Calderón problem for the fractional Schrödinger equation. *Analysis & PDE*, 13(2):455–475, 2020.
- [13] Yavar Kian, Lauri Oksanen, Eric Soccorsi, and Masahiro Yamamoto. Global uniqueness in an inverse problem for time fractional diffusion equations. *Journal of Differential Equations*, 264(2):1146–1170, 2018.
- [14] Katsiaryna Krupchyk and Gunther Uhlmann. Uniqueness in an inverse boundary problem for a magnetic Schrödinger operator with a bounded magnetic potential. *Communications in Mathematical Physics*, 327(3):993–1009, 2014.
- [15] Ru-Yu Lai, Yi-Hsuan Lin, and Angkana Rüland. The Calderón problem for a space-time fractional parabolic equation. *SIAM Journal on Mathematical Analysis*, 52(3):2655–2688, 2020.
- [16] Li Li. The Calderón problem for the fractional magnetic operator. *Inverse Problems*, 36(7):075003, 2020.
- [17] Li Li. Determining the magnetic potential in the fractional magnetic Calderón problem. *arXiv preprint arXiv:2006.10150*, 2020.
- [18] Li Li. A semilinear inverse problem for the fractional magnetic Laplacian. *arXiv preprint arXiv:2005.06714*, 2020.
- [19] Mark M Meerschaert, David A Benson, and Boris Bäumer. Multidimensional advection and fractional dispersion. *Physical Review E*, 59(5):5026, 1999.
- [20] Mark M Meerschaert, David A Benson, Hans-Peter Scheffler, and Peter Becker-Kern. Governing equations and solutions of anomalous random walk limits. *Physical Review E*, 66(6):060102, 2002.
- [21] Gen Nakamura, Ziqi Sun, and Gunther Uhlmann. Global identifiability for an inverse problem for the Schrödinger equation in a magnetic field. *Mathematische Annalen*, 303(1):377–388, 1995.
- [22] Karel Rektorys. *The method of discretization in time and partial differential equations*. D. Reidel Publishing Company, 1982.

- [23] Angkana Rüland and Mikko Salo. Quantitative approximation properties for the fractional heat equation. *arXiv preprint arXiv:1708.06300*, 2017.
- [24] Angkana Rüland and Mikko Salo. The fractional Calderón problem: low regularity and stability. *Nonlinear Analysis*, 193:111529, 2020.
- [25] Alexander I Saichev and George M Zaslavsky. Fractional kinetic equations: solutions and applications. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 7(4):753–764, 1997.
- [26] Zi Qi Sun. An inverse boundary value problem for Schrödinger operators with vector potentials. *Transactions of the American Mathematical Society*, 338(2):953–969, 1993.