Quasi-stationary distribution and metastability for the stochastic Becker-Döring model

Erwan Hingant* Romain Yvinec^{†‡}

March 16, 2022

Abstract

We study a stochastic version of the classical Becker-Döring model, a well-known kinetic model for cluster formation that predicts the existence of a long-lived metastable state before a thermodynamically unfavorable nucleation occurs, leading to a phase transition phenomena. This continuous-time Markov chain model has received little attention, compared to its deterministic differential equations counterpart. We show that the stochastic formulation leads to a precise and quantitative description of stochastic nucleation events thanks to an exponentially ergodic quasi-stationary distribution for the process conditionally on nucleation has not yet occurred.

Keywords: Quasi-stationary distribution; Metastability; Exponential ergodicity; Becker-Döring; Nucleation; Phase transition

AMS MSC 2010: 82C26; 60J27.

1 Introduction

The Becker-Döring model is a kinetic model for phase transition phenomenon represented schematically by the reaction network

$$\emptyset \xrightarrow{a_1 z^2} C_2, \quad \text{and} \quad C_i \xrightarrow{a_i z} C_{i+1}, \quad i = 2, 3, \dots$$

$$(1.1)$$

We assume an infinite reservoir of monomer, cluster of size 1, represented in (1.1) by \emptyset . The parameter z represents the *fixed* concentration of monomer and will play a key role in the sequel. A cluster of size $i \geq 2$, whose population is represented in (1.1) by C_i , lengthen to give rise to a cluster of size i+1 at rate a_iz or shorten to give rise a cluster of size i-1 at rate b_i . The rate of apparition of a new cluster of size 2 is a_1z^2 (without loss of generality). All parameters are positives.

The Becker-Döring (BD) model goes back to the seminal work "Kinetic treatment of nucleation in supersaturated vapors" in [2]. Since then, the model met very different applications ranging from physics to biology. From the mathematical point of view, this model received much more attention in the deterministic literature than the probabilistic one. We refer to our review [6] for historical comments and detailed literature review on theoretical results from the deterministic side. See also [7, 14] for recent results on functional law of large number and central limit theorem.

The model was initially designed to explain critical phase condensation phenomena where macroscopic droplets self-assemble and segregate from an initially supersaturated homogeneous mixture of particles, at a rate that is exponentially small in the

^{*}Departamento de Matemática, Universidad del Bío-Bío, Chile. E-mail: ehingant@ubiobio.cl

[†]PRC, INRA, CNRS, IFCE, Université de Tours, 37380 Nouzilly, France. E-mail: romain.yvinec@inrae.fr

[‡]Associate researcher at Cogitamus laboratory.

excess of particles. This led to important applications in kinetic nucleation theory [13]. Mathematical studies in the 90's showed that (in the deterministic context), departing from certain initial conditions, the size distribution of clusters reaches quickly a metastable configuration composed of "small" clusters, and remains arbitrary close to that state for a very large time, before it converges to the true stationary solution that leads to "infinitely large" clusters (interpreted as droplets) [10, 11].

Our objective in this note is to re-visit the metastability theory in Becker-Döring model in terms of quasi-stationary distribution (QSD) for the associated continuous-time Markov chain. We prove existence, uniqueness and exponential ergodicity of a QSD for the BD model conditioned on the event that large clusters have not yet appeared. We prove furthermore that the convergence rate towards the QSD is faster than the rate of apparition of (sufficiently) large clusters. Quantitative results are obtained thanks to a surprisingly simple analytical formula for the QSD, that provides also an exact rate of apparition of stable large clusters, consistently with the original heuristic development of Becker and Döring.

Outline: We introduce in Sec. 2 the stochastic BD model. In Sec. 3, we gather few (known) results on a related Birth-Death process, and from deterministic modeling of the BD model. Then, we prove exponential decay, in total variation, of the BD process towards its stationary measure in Sec. 4. For any $n \geq 2$, similar results are obtained in Sec. 5 for the process conditionally on no cluster larger than n are formed. An estimate on the time for the first cluster larger than n to appear is provided in Sec. 6. The last, but not least, Sec. 7 gathers our results in a quantitative way, and proves that the QSD can be interpreted as a long-lived metastable state, when n equals the critical nucleus size.

Notation: We denote by \mathbb{N}_0 the set of non-negatives integers $\mathbb{N}_0 = \{0,1,2\ldots\}$, \mathbb{N}_i the set of integers greater or equal to i, $[\![\,]\!]$ for integers interval. For a set A, A^c is its complement, $\mathbf{1}_A$ the indicator function on it, #A its cardinality. For two sets A and B their intersection is denoted by A, B. Also, $\mathbf{1}_x = \mathbf{1}_{\{x\}}$ and $\mathbf{1}$ (resp. 0) is the constant function equal to 1 (resp. 0). For two numbers a,b, their minimum is $a \wedge b$. We denote by $\|\mu - \nu\|$ the total variation distance between two probability measure μ and ν on a countable state space $\mathcal S$ namely

$$\|\mu - \nu\| = \frac{1}{2} \sum_{x \in S} |\mu(x) - \mu(x)| = \inf_{\gamma \in \Gamma} \int_{S \times S} \mathbf{1}_{x \neq y} \gamma(dx, dy),$$

where Γ is the set of probability measures on $\mathcal{S} \times \mathcal{S}$ with marginals μ and ν . \mathbf{E} (resp. \mathbf{E}_{μ}) denotes the expectation with respect to the usual probability measure \mathbf{P} (resp. μ).

Below, we set $\mathcal{E} = \ell^1(\mathbb{N}_2, \mathbb{N}_0)$ the space of summable \mathbb{N}_0 -valued sequences indexed by \mathbb{N}_2 .

2 The model

The stochastic Becker-Döring (BD) process is a continuous-time Markov chain on the countable state space $\mathcal E$ with infinitesimal generator $\mathcal A$, given for all ψ with finite support on $\mathcal E$ and $C\in\mathcal E$, by

$$\mathcal{A}\psi(C) = \sum_{i=1}^{+\infty} \left(a_i z C_i [\psi(C + \Delta_i) - \psi(C)] + b_{i+1} C_{i+1} [\psi(C - \Delta_i) - \psi(C)] \right)$$

with the convention $C_1=z$, $\Delta_1=\mathbf{e}_2$ and $\Delta_i=\mathbf{e}_{i+1}-\mathbf{e}_i$, for each $i\geq 2$, where $\{\mathbf{e}_2,\mathbf{e}_3,\ldots\}$ denotes the canonical basis of $\mathcal E$ namely, $e_{i,k}=1$ if k=i and 0 otherwise.

We shall however use a different approach, modeling explicitly the size of each individual cluster. On a sufficiently large probability space $(\Omega, \mathcal{F}, \mathbf{P})$, we introduce:

- N_1, N_2, \ldots a denumerable family of independent Poisson point measure with intensity the Lebesgue measure dsdu on \mathbb{R}^2_+ .
- T_1, T_2, \ldots a collection of random times such that the $T_k T_{k-1}$ are independent exponential random variable of parameter $a_1 z^2$, independent from the above Poisson point measure as well, with $T_0 = 0$.

Let Π^{in} a probability distribution on $\mathcal E$ and $\mathbf C(0)=(C_2(0),C_3(0),\ldots)$ an $\mathcal E$ -valued random variable distributed according to Π^{in} . We denote by $N^{in}=\sum_{i=2}^\infty C_i(0)$. By construction $N^{in}<\infty$ almost surely (a.s.). Then, given $\mathbf C(0)$, we define $X_1(0),X_2(0),\ldots$ a denumerable collection of random variables on $\mathbb N_2$ such that, a.s. for each $i\geq 2$,

$$C_i(0) = \# \left\{ k \in [1, N^{in}] \mid X_k(0) = i \right\},$$
 (2.1)

and $X_k^{\text{in}}=2$ for all $k>N^{\text{in}}$. Note this construction may be achieved by a bijective *labeling* function¹. Finally, we consider the denumerable collection of stochastic processes X_1, X_2, \ldots on \mathbb{N}_1 solution of the stochastic differential equations, for all $t\geq 0$ and $k\geq 1$,

$$X_k(t) = X_k(0) + \sum_{i=2}^{\infty} \int_0^t \int_{\mathbb{R}^+} \mathbf{1}_{s > T_{k-N^{\text{in}}}} \mathbf{1}_{X_k(s^-)=i} (\mathbf{1}_{u \le a_i z} - \mathbf{1}_{a_i z < u \le a_i z + b_i}) N_k(ds, du), \quad (2.2)$$

where by convention $T_k=0$ if $k\leq 0$. The pathwise construction (2.2) is what we call thereafter the particle description of the BD process. The interpretation is clear: $X_k(t)$ denotes the size of the cluster labelled by k at time t; for $k\leq N^{\mathrm{i}n}$, clusters are initially "actives" while for $k>N^{\mathrm{i}n}$ clusters are initially "inactives" at state 2, and become "activated" at the random arrival times $T_{k-N^{\mathrm{i}n}}$. We ensured $X_k(0)<\infty$ a.s. because the $C_i(0)$'s are integer-valued random variables and belong to \mathcal{E} , the sequence $\mathbf{C}(0)$ is a.s. equally 0 from a certain range. Thus, local existence of \mathbf{c} addiag processes $t\mapsto X_k(t)$ on \mathbb{N}_1 solution to (2.2) can classically be obtained inductively. It is clear from (2.2) that each X_k evolves like a Birth-Death process for $t>T_k$ (that will be detailed in the next section 3) and are mutually independent conditionally to their initial value. The Reuter's criterion gives a well-known necessary and sufficient condition so that each process X_k is non-explosive, namely

$$\sum_{n=2}^{\infty} Q_n z^n \left(\sum_{k=n}^{\infty} \frac{1}{a_k Q_k z^{k+1}} \right) = \infty, \text{ with } Q_1 = 1, \ Q_i = \frac{a_1 a_2 \cdots a_{i-1}}{b_2 \cdots b_i}, \ i \ge 2.$$
 (H0)

It is now convenient to go back to the original description at stake. The number of "active" clusters at time $t \ge 0$ is given by the counting process

$$N(t) = N^{\mathrm{i}n} + \sum_{k \ge 1} \mathbf{1}_{t \ge T_k} \,,$$

while the number of cluster of size $i \geq 2$ is

$$C_i(t) = \# \{k \in [1, N(t)] \mid X_k(t) = i\}$$
.

Now, noticing that $C_i(t) = \sum_{k=1}^{N(t)} \mathbf{1}_i(X_k(t))$, we can prove from standard stochastic calculus that the process \mathbf{C} given by $\mathbf{C}(t) = (C_2(t), C_3(t), \ldots)$ for all $t \geq 0$ has infinitesimal generator \mathcal{A} , and being non-explosive under condition (H0), it is the unique regular jump homogeneous Markov chain on \mathcal{E} with infinitesimal generator \mathcal{A} and initial distribution Π^{in} , say the BD process. The proof is left to the reader and follows from classical

 $^{^1}$ A function (that exists) which associates, to each $c \in \mathcal{E}$ such that $N = \sum_{i=2}^{\infty} c_i < \infty$, a unique sequence (x_1, \ldots, x_N) in \mathbb{N}_2 satisfying $c_i = \#\{k \in [\![1,N]\!] \mid x_k = i\}$.

theory, e.g. [1]. In the sequel, $\mathbf{C}(t)$ always denote a BD process, and $\mathbf{P}_{\Pi^{in}} \{ \mathbf{C} \in \cdot \}$ its (unique) finite dimensional probability distribution given that $\mathbf{C}(0)$ is distributed according to Π^{in} . We also set by convention $\mathbf{P}_C = \mathbf{P}_{\delta_C}$ for a deterministic $C \in \mathcal{E}$ and we recover $\mathbf{P}_{\Pi^{in}} \{ \cdot \} = \sum_{C \in \mathcal{E}} \mathbf{P}_C \{ \cdot \} \Pi^{in}(C)$.

3 Behaviour of one cluster

Let X the continuous-time Markov chain on \mathbb{N}_1 with transition rate matrix $(q_{i,j})_{i,j\geq 1}$ whose nonzero entries are

$$q_{i,i+1} = a_i z, \ q_{i,i-1} = b_i, \ q_{i,i} = -(a_i z + b_i), \qquad i \ge 2.$$
 (3.1)

Remark that i = 1 is absorbing in agreement with (1.1): when a cluster size reaches 1, it "leaves the system". We shall assume standard hypotheses in the BD model [10, 11]:

$$\lim_{i \to \infty} b_i / a_i = z_s > 0, \quad \text{and} \quad \lim_{i \to \infty} b_{i+1} / b_i = 1. \tag{H1}$$

Hypothesis (H1) then guarantee (H0) for $z \neq z_s$ for the following reason. The convergence of both series

$$\sum_{k=2}^{\infty}Q_kz^k \quad \text{and} \quad \sum_{k=2}^{\infty}\frac{1}{a_kQ_kz^{k+1}}\,,$$

depends on the value of z. Indeed, z_s is the radius of convergence of the first series while the second series converges for $z>z_s$ and diverges for $z< z_s$. We have a dichotomy in the long time behavior of X related to this value. The case $z< z_s$ is called the *sub-critical case*, for which absorption at state 1 is certain and the expected time of absorption is finite (also called ergodic absorption). The case $z>z_s$ is called the *super-critical case* and absorption at 1 is not certain (also called transient absorption), and the probability to be absorbed at 1 is, according to [8],

$$\lim_{t \to \infty} p_{i1}(t) = J \sum_{k=i}^{\infty} \frac{1}{a_k Q_k z^{k+1}}, \quad \text{with } J = \left(\sum_{k=1}^{\infty} \frac{1}{a_k Q_k z^{k+1}}\right)^{-1}$$
 (3.2)

where $p_{ij}(t) = \mathbf{P}\{X(t) = j \mid X(0) = i\}$ the probability transition function of X. The limit case $z = z_s$ is somewhat technical and depends more deeply on the shape of the coefficients. It is not considered in this note.

Following [10], a precise long time estimate on transient states can be obtained, under the hypothesis (H1) and

$$\frac{b_{i+1}}{b_i} - 1 = O(i^{-1}), \quad \frac{a_{i+1}}{a_i} - 1 = O(i^{-1}), \quad a_i = O(i) \quad \text{and} \quad \lim_{i \to +\infty} a_i = +\infty. \tag{H2}$$

In such a case, the infinite matrix $(q_{i,j})_{i,j\geq 2}$ in (3.1) is self-adjoint on the Hilbert space H consisting of the real sequences $\mathbf{x}=(x_2,x_3,\ldots)$ whose norm is

$$\|\mathbf{x}\|_H = \sqrt{\sum_{i=2}^{\infty} \frac{x_i^2}{Q_i z^i}} < \infty.$$

We denote by $\langle \cdot, \cdot \rangle_H$ the associated scalar product. It turns that $(q_{i,j})_{i,j\geq 2}$ has a negative maximum eigenvalue $-\lambda$, and the following estimate holds for any $i\geq 2$,

$$\|(p_{ij}(t))_{j\geq 2}\|_{H} \leq e^{-\lambda t} \|(p_{ij}(0))_{j\geq 2}\|_{H} = \frac{e^{-\lambda t}}{\sqrt{Q_{i}z^{i}}}.$$
(3.3)

We will also consider the chain X conditioned to remain below a given state $n+1 \geq 2$. We define the exit time

$$T_n = \inf(t \ge 0 \mid X(t) \notin [1, n]) = \inf(t \ge 0 \mid X(t) \ge n + 1)$$
 (3.4)

Let Y the birth-death process defined by $Y(t) = X(t \wedge T_n)$. Hence, Y is absorbed either in 1 or n+1, and the probability to be absorbed at 1 (without visiting state n+1) is, according to [8, p.387],

$$\lim_{t \to +\infty} p_{i1}^n(t) = J_n \sum_{k=i}^n \frac{1}{a_k Q_k z^{k+1}}, \quad \text{with } J_n = \left(\sum_{k=1}^n \frac{1}{a_k Q_k z^{k+1}}\right)^{-1}, \quad (3.5)$$

where $p_{ij}^n(t) = \mathbf{P}\left(Y(t) = j \mid Y(0) = i\right)$ is the probability transition function of Y and clearly

$$\mathbf{P}\left\{T_{n} > t \mid X(0) = i\right\} \ge \lim_{t \to +\infty} \mathbf{P}\left\{T_{n} > t \mid X(0) = i\right\} = J_{n} \sum_{k=i}^{n} \frac{1}{a_{k} Q_{k} z^{k+1}}.$$
 (3.6)

Again, in [10], the author shows that the truncated matrix $(q_{i,j})_{i,j=2,...,n}$ is similar to a symmetric one and then there exists $\gamma_n > 0$ such that for each i = 2,...,n,

$$\sqrt{\sum_{j=2}^{n} \frac{p_{ij}^{n}(t)^{2}}{Q_{j}z^{j}}} \le \frac{e^{-\gamma_{n}t}}{\sqrt{Q_{i}z^{i}}}.$$
(3.7)

Note the probability to be absorbed in 1 before time t, $p_{i1}^n(t)$, is monotonously increasing and $\lim_{t\to+\infty} p_{i1}^n(t) = 1 - \lim_{t\to+\infty} p_{i(n+1)}^n(t)$, thus we deduce that

$$\sum_{i=2}^{n} \mathbf{P} \left\{ Y(t) = j \mid Y(0) = i, \ T_n > t \right\} = \frac{\sum_{j=2}^{n} p_{ij}^n(t)}{1 - p_{i(n+1)}^n(t)} \le M_{i,n} e^{-\gamma_n t}$$

where the constant $M_{i,n}$, obtained thanks to (3.7), Cauchy-Schwarz inequality and (3.5), is given by

$$M_{i,n} = \left(\frac{1}{Q_i z^i} \sum_{j=2}^n Q_j z^j\right)^{\frac{1}{2}} \frac{1}{J_n \sum_{k=i}^n \frac{1}{a_k Q_k z^{k+1}}}.$$

We end this preliminary section, noticing that X(t) = Y(t) on $\{T_n > t\}$, with

$$\mathbf{P}\{X(t) = 1 \mid X(0) = i, T_n > t\} > 1 - M_{i,n}e^{-\gamma_n t} \wedge 1.$$
(3.8)

4 Long-time behaviour of the BD process

In this section we are concerned with the long-time behaviour of the BD process. Formally the measure Π^{eq} , given by

$$\Pi^{\mathrm{e}q}(C) = \prod_{i=2}^{\infty} e^{-c_i^{\mathrm{e}q}} rac{(c_i^{\mathrm{e}q})^{C_i}}{C_i!} \,, \qquad ext{with } c_i^{\mathrm{e}q} = Q_i z^i$$

for all $C \in \mathcal{E}$, satisfies $\mathbf{E}_{\Pi^{\mathrm{eq}}}[\mathcal{A}\psi(C)] = 0$ for any function ψ on \mathcal{E} with finite support². Actually, Π^{eq} satisfies the detailed balance condition $a_i z C_i \Pi^{\mathrm{eq}}(C) = b_{i+1} (C_{i+1} + 1) \Pi^{\mathrm{eq}}(C + \Delta_i)$, for all $i \geq 1$ and all $C \in \mathcal{E}$ (with the convention that $C_1 = z$), as a consequence of

 $^{^2}$ In the sequel C in expectation formula always refers to the free variable of integration.

the relation $a_iQ_i=b_{i+1}Q_{i+1}$. In the sub-critical case, Π^{eq} is a probability measure on $\mathcal E$ (indeed $\Pi^{eq}(\mathbb N_0^{\mathbb N_2})=1$ with support in $\mathcal E$ because of $\mathbf E_{\Pi^{eq}}\left[\sum_{i=2}^\infty C_i\right]=\sum_{i=2}^\infty c_i^{eq}<\infty$) and we prove exponential ergodicity towards Π^{eq} . In the super-critical case, Π^{eq} is not a limiting distribution (and $\sum_{i=2}^\infty c_i^{eq}=\infty$) but the measure defined by

$$\Pi^{stat}(C) = \prod_{i=2}^{\infty} e^{-f_i} \frac{(f_i)^{C_i}}{C_i!}, \quad \text{with } f_i = JQ_i z^i \sum_{k=i}^{\infty} \frac{1}{a_k Q_k z^k}$$
 (4.1)

for all $C \in \mathcal{E}$, where J is given in (3.2), characterizes the long-time behaviour of any finite-number of marginals. Now on, we note $\mathbf{f} = (f_i)_{i \geq 2}$, with f_i defined in (4.1).

Theorem 4.1. Under hypotheses (H1) and (H2). Let Π^{in} a probability distribution on \mathcal{E} such that

$$\mathbf{E}_{\Pi^{in}} \left[\langle C, \sqrt{\mathbf{Q}} \rangle_H \right] < \infty \tag{4.2}$$

where $\sqrt{\mathbf{Q}} = (\sqrt{Q_i z^i})_{i \geq 2}$. With $\lambda > 0$ introduced in Sec. 3, see (3.3), we have:

• In the sub-critical case ($z < z_s$), for all $t \ge 0$,

$$\|\mathbf{P}_{\Pi^{\mathrm{i}n}} \left\{ \mathbf{C}(t) \in \cdot \right\} - \Pi^{\mathrm{e}q} \| \le R^{\mathrm{i}n} e^{-\lambda t}$$

with
$$R^{\mathrm{i}n} = K(\mathbf{E}_{\Pi^{\mathrm{i}n}} \left[\langle C, \sqrt{\mathbf{Q}} \rangle_H \right] + \mathbf{E}_{\Pi^{\mathrm{eq}}} \left[\langle C, \sqrt{\mathbf{Q}} \rangle_H \right])$$
 and $K = (\sum_{k=2}^{\infty} c_i^{\mathrm{eq}})^{\frac{1}{2}}$;

• In the super-critical case ($z>z_s$), for all $t\geq 0$, and for all $n\geq 2$,

$$\|\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{(C_2(t),\ldots,C_n(t))\in\cdot\right\} - \Pi^{\mathrm{stat}}(\cdot\times\prod_{k=n+1}^{\infty}\mathbb{N}_0)\|\leq R_n^{\mathrm{i}n}e^{-\lambda t}$$

with
$$R_n^{\mathrm{i}n}=K_n\mathbf{E}_{\Pi^{\mathrm{i}n}}\left[\langle C,\sqrt{\mathbf{Q}}\rangle_H
ight]+\|\mathbf{f}\|_H$$
 and $K_n=(\sum_{k=2}^nc_i^{\mathrm{eq}})^{rac{1}{2}}.$

Not least, remark that $\mathbf{E}_{\Pi^{eq}}\left[\left\langle C,\sqrt{\mathbf{Q}}\right\rangle_{H}\right]=\sum_{i=2}^{\infty}\sqrt{Q_{i}z^{i}}<\infty$ for $z< z_{s}$ and that $\mathbf{f}\in H$ for $z>z_{s}$ (see [10]). In the sequel, K_{n} and K always refer to the constants given above. The proof is based on a coupling (described below) to a distribution starting from $\mathbf{0}$, so that the control of the initial particles in $\mathbf{C}(0)$ are a key point.

Lemma 4.2. Under the hypothesis of Theorem 4.1. Let the collection of processes X_1, X_2, \ldots being a particle description of the BD process C. We have, for each $n \geq 2$,

$$\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\forall k\in \llbracket 1,N^{\mathrm{i}n}\rrbracket\,,\; X_k(t)\notin \llbracket 2,n\rrbracket\right\}\geq 1-K_n\mathbf{E}_{\Pi^{\mathrm{i}n}}\left[\langle C,\sqrt{\mathbf{Q}}\rangle_H\right]e^{-\lambda t}\,.$$

In particular, for the sub-critical case,

$$\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\forall k \in \llbracket 1, N^{\mathrm{i}n} \rrbracket, \ X_k(t) = 1\right\} \ge 1 - K\mathbf{E}_{\Pi^{\mathrm{i}n}}\left[\langle C, \sqrt{\mathbf{Q}} \rangle_H\right] e^{-\lambda t}.$$

Proof. Fix $n \geq 2$. Let $C \in \mathcal{E}$ deterministic, define $N = \sum_{i=2}^{\infty} C_i$ and $(i_1,\ldots,i_N) \in \mathbb{N}_2^N$ given by the *labelling function*, e.g. $\mathbf{C}(0) = C$ and $X_1(0) = i_1,\ldots X_N(0) = i_N$ satisfy relation (2.1). Since each processes $X_1,\ldots X_N$ are independent copy of the chain X given in Sec. 3, conditionally on their initial condition, we have

$$\mathbf{P}_{C} \{ \forall k \in [1, N], \ X_{k}(t) \notin [2, n] \} = \prod_{k=1}^{N} \mathbf{P} \{ X(t) \notin [2, n] \mid X(0) = i_{k} \}$$
 (4.3)

for all $t \ge 0$. Thanks to Cauchy–Schwarz inequality and (3.3),

$$\mathbf{P}\{X(t) \in [2, n] \mid X(0) = i\} = \sum_{j=2}^{n} p_{ij}(t) \le \left(\frac{1}{\sqrt{Q_i z^i}}\right) K_n e^{-\lambda t} \wedge 1.$$

Hence, with (4.3), we have

$$\mathbf{P}_{C} \{ \forall k \in [1, N], \ X_{k}(t) \notin [2, n] \} \ge 1 - K_{n} e^{-\lambda t} \sum_{k=1}^{N} \frac{1}{\sqrt{Q_{i_{k}} z^{i_{k}}}},$$

remarking that $\prod_{i=1}^{N}(1-x_i\wedge 1)\geq 1-\sum_{i=1}^{N}x_i$ for any non-negatives x_1,\ldots,x_N . Finally, we conclude that

$$\mathbf{P}_{\Pi^{\text{i}n}} \left\{ \forall k \in [\![1, N^{\text{i}n}]\!], \ X_k(t) \notin [\![2, n]\!] \right\} \ge 1 - K_n e^{-\lambda t} \sum_{C \in \mathcal{E}} \sum_{k=1}^N \frac{1}{\sqrt{Q_{i_k} z^{i_k}}} \Pi^{\text{i}n}(C)$$

$$= 1 - K_n e^{-\lambda t} \mathbf{E}_{\Pi^{\text{i}n}} \left[\sum_{i=2}^{\infty} \frac{\# \left\{ k \in [\![1, N^{\text{i}n}]\!] \mid X_k(0) = i \right\}}{\sqrt{Q_i z^i}} \right],$$

and the proof ends.

We now show, by a coupling argument, that any solution satisfying (4.2) is in total variation exponentially close to the solution that starts with no cluster, namely the deterministic initial condition at 0.

Lemma 4.3. Under the hypothesis of Theorem 4.1. For all $t \geq 0$, we have

• In the subcritical case ($z < z_s$),

$$\|\mathbf{P}_{\Pi^{in}} \left\{ \mathbf{C}(t) \in \cdot \right\} - \mathbf{P_0} \left\{ \mathbf{C}(t) \in \cdot \right\} \| \le K \mathbf{E}_{\Pi^{in}} \left[\langle C, \sqrt{\mathbf{Q}} \rangle_H \right] e^{-\lambda t};$$

• In the super-critical case $(z > z_s)$, for all $n \ge 2$,

$$\|\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\left(C_{2}(t),\ldots,C_{n}(t)\right)\in\cdot\right\}-\mathbf{P}_{\mathbf{0}}\left\{\left(C_{2}(t),\ldots,C_{n}(t)\right)\in\cdot\right\}\|\leq K_{n}\mathbf{E}_{\Pi^{\mathrm{i}n}}\left[\langle C,\sqrt{\mathbf{Q}}\rangle_{H}\right]e^{-\lambda t}$$

Proof. Let the collection of processes X_1,X_2,\ldots (resp. Y_1,Y_2,\ldots) being a particle description of the BD process that starts from the initial distribution Π^{in} (resp. from δ_0). We couple the processes X_1,X_2,\ldots to the processes Y_1,Y_2,\ldots such that all new particle "activates" simultaneously and evolves with the same jumps. Namely, $Y_k(t)=X_{k+N^{in}}(t)$ for all k>1 and all $t\geq 0$, where N^{in} is distributed according to Π^{in} .

In the sub-critical case the proof readily follows from Lemma 4.2 and the definition of the total variation since

$$\|\mathbf{P}_{\Pi^{in}} \{ \mathbf{C}(t) \in \cdot \} - \mathbf{P_0} \{ \mathbf{C}(t) \in \cdot \} \| \le 1 - \mathbf{P}_{\Pi^{in}} \{ \forall k \in [1, N^{in}], X_k(t) = 1 \}$$

because all "active" clusters are equal whenever all initial clusters from Π^{in} have been absorbed. A very similar argument holds in the super-critical case.

Proof of Theorem 4.1. We consider first the sub-critical case. As said, condition (4.2) is satisfied for Π^{eq} since $\mathbf{E}_{\Pi^{eq}}\left[\langle C, \sqrt{\mathbf{Q}}\rangle_H\right] < \infty$, thus Lemma 4.3 applies for Π^{eq} as initial distribution. Because the constructed BD process is regular and Π^{eq} is a stationary distribution (i.e. $\mathbf{P}_{\Pi^{eq}}\left\{\mathbf{C}(t) \in \cdot\right\} = \Pi^{eq}$), we deduce

$$\|\mathbf{P_0}\left\{\mathbf{C}(t)\in\cdot\right\} - \Pi^{eq}\| \le K\mathbf{E}_{\Pi^{eq}}\left[\langle C, \sqrt{\mathbf{Q}}\rangle_H\right]e^{-\lambda t}.$$

Going back to any Π^{in} satisfying condition (4.2), applying Lemma 4.3 again and the triangular inequality yield the desired result.

Consider now the super-critical case. Π^{stat} is a product of Poisson distribution $\mathcal{P}(f_i)$ on \mathbb{N}_0 with mean f_i . According to a classical result on Markov population processes, see

e.g. [9, Sec. 4], the law $\mathbf{P_0}\left\{\mathbf{C}(t) \in \cdot\right\}$ is also a product of Poisson distribution $\mathcal{P}(c_i(t))$ on \mathbb{N}_0 with mean $c_i(t)$ such that $\mathbf{c}(t) = (c_2(t), c_3(t), \ldots)$ solves the deterministic (linear) Becker-Döring equations namely, $\mathbf{c}(t) = A\mathbf{c}(t) + a_1z^2\mathbf{e}_2$ where $A = (q_{j,i})_{i,j \geq 2}$ the matrix with entries in (3.1), and initial condition $\mathbf{c}(0) = \mathbf{0}$. Thanks to [10, Theorem III], we have

$$\|\mathbf{c}(t) - \mathbf{f}\|_{H} \le \|\mathbf{f}\|_{H} e^{-\lambda t} \tag{4.4}$$

and according to [12, Lemma 1], $\|\mathcal{P}(c_i(t)) - \mathcal{P}(f_i)\| \le |c_i(t) - f_i|$. The latter, with independence of the marginals of Π^{stat} and of $\mathbf{P_0}$ { $\mathbf{C}(t) \in \cdot$ }, estimate (4.4) and Cauchy-Schwarz inequality, entail

$$\|\mathbf{P_0}\{(C_2(t),\ldots,C_n(t))\in\cdot\} - \Pi^{\text{stat}}(\cdot \times \prod_{k=n+1}^{\infty} \mathbb{N}_0)\| \le \sum_{i=2}^{n} |d_i(t) - f_i| \le K_n \|\mathbf{f}\|_H e^{-\lambda t}.$$

We conclude again by Lemma 4.3 and the triangular inequality.

5 A quasi-stationary distribution

Let $\mathcal{E}_n = \{C \in \mathcal{E} \mid C_i = 0, i \geq n+1\}$. We define the first exit time from \mathcal{E}_n ,

$$\tau_n = \inf \left\{ t \ge 0 \mid \mathbf{C}(t) \notin \mathcal{E}_n \right\}. \tag{5.1}$$

Remark that $\mathbf{P}_{\Pi^{in}}$ $\{\tau_n > t\} > 0$ for all times t > 0 and for any Π^{in} supported on \mathcal{E}_n . We will give in the next section 6 a tight lower bound on that probability in the super-critical case. In this section, we prove exponential ergodicity towards a unique QSD for the BD process conditioned to $\tau_n > t$. It is remarkable that we have at hand an explicit QSD, given, for all $C \in \mathcal{E}_n$, by

$$\Pi_n^{qsd}(C) = \prod_{i=2}^n \frac{(f_i^n)^{C_i}}{C_i!} e^{-f_i^n}, \quad \text{with } f_i^n(z) = J_n Q_i z^i \sum_{k=i}^n \frac{1}{a_k Q_k z^{k+1}}$$
(5.2)

for i = 2, ..., n and J_n defined in (3.5).

Proposition 5.1. Under assumption (H0). The distribution Π_n^{qsd} is a quasi-stationary distribution for the BD process conditioned to stay on \mathcal{E}_n namely,

$$\mathbf{P}_{\Pi_n^{qsd}} \left\{ \mathbf{C}(t) \in \cdot \mid t < \tau_n \right\} = \Pi_n^{qsd}$$

and moreover

$$\mathbf{P}_{\Pi_n^{qsd}}\left\{t < \tau_n\right\} = \exp\left(-J_n t\right) .$$

Proof. Recall assumption (H0) ensures the BD process is regular. Fix $n \geq 2$. Let the semi-group $P_t^n \psi(C) = \mathbf{E}_C \left[\psi(\mathbf{C}(t)) \mathbf{1}_{t < \tau_n} \right]$ for $t \geq 0$ (i.e. $\mathbf{C}(0)$ is distributed according to δ_C), whose generator is

$$A_n \psi(C) = \sum_{i=1}^{n-1} \left(a_i z C_i [\psi(C + \Delta_i) - \psi(c)] + b_{i+1} C_{i+1} [\psi(C - \Delta_i) - \psi(C)] \right) - a_n z C_n \psi(C)$$

for all $C \in \mathcal{E}_n$ (recall $C_1 = z$) and bounded function ψ on \mathcal{E}_n . Denote by \mathcal{A}_n^* the dual operator for the generator \mathcal{A}_n . Some calculations show that the distribution (5.2) satisfies, for any $C \in \mathcal{E}_n$,

$$\mathcal{A}_{n}^{*}\Pi_{n}^{qsd}(C) = \Pi_{n}^{qsd}(C) \left\{ b_{2}f_{2}^{n} - a_{1}z^{2} + \sum_{i=2}^{n} \frac{C_{i}}{f_{i}^{n}} \left(a_{i-1}zf_{i-1}^{n} - (a_{i}z + b_{i})f_{i}^{n} + \mathbf{1}_{i < n}b_{i+1}f_{i+1}^{n} \right) \right\}$$

with the convention $f_1^n=z$. Since the f_i^n given by (5.2) verifies $a_izf_i^n-\mathbf{1}_{i< n}b_{i+1}f_{i+1}^n=J_n$ for all $i\in [\![1,n]\!]$, all terms but the first cancel in the above expression, so that we obtain $\mathcal{A}_n^*\Pi_n^{\mathrm{q}sd}=-J_n\Pi_n^{\mathrm{q}sd}$ which is the classical spectral criteria of QSD, noticing that $J_n\leq a_1z^2$, see [4, Thm 4.4].

The next theorem shows the QSD is a quasi-limiting distribution for a wide range of initial distribution supported on \mathcal{E}_n , with an exponential rate of convergence and an explicit (non-uniform) pre-factor.

Theorem 5.2. Under assumption (H0). Let Π^{in} a probability distribution on \mathcal{E}_n such that $\mathbf{E}_{\Pi^{in}}\left[\sum_{i=2}^{\infty}C_i\right]<\infty$. We have for all $t\geq 0$,

$$\|\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\mathbf{C}(t)\in\cdot\mid\tau_n>t\right\}-\Pi_n^{\mathrm{q}sd}\|\leq K_n\left(\frac{H_n^{\mathrm{i}n}}{\mathbf{P}\left\{\tau_n>t\right\}}+e^{J_nt}H_n^{\mathrm{q}sd}\right)e^{-\gamma_nt},$$

where τ_n is defined in (5.1), J_n in (3.5), K_n in Theorem (4.1), γ_n in (3.7),

$$H_n^{\mathrm{i}n} = \sum_{i=2}^n \sqrt{Q_i z^i} \frac{\mathbf{E}_{\Pi^{\mathrm{i}n}}\left[C_i\right]}{f_i^n} \quad \text{and} \quad H_n^{\mathrm{q}sd} = \sum_{i=2}^n \sqrt{Q_i z^i} \,.$$

It is clear that $H_n^{\mathrm{i}n}$ is finite because $\mathbf{E}_{\Pi^{\mathrm{i}n}}\left[\sum_{i=2}^{\infty}C_i\right]$ is. The proof of Theorem 5.2 is similar to the proof of Theorem 4.1 and consists in a coupling argument together with a control of the initial clusters in $\Pi^{\mathrm{i}n}$. We start by the later, which is the analogous of Lemma 4.2.

Lemma 5.3. Under the hypothesis of Theorem 5.2. Let the collection of processes X_1, X_2, \ldots being a particle description of the BD process C. We have

$$\mathbf{P}_{\Pi^{\text{in}}} \left\{ \forall k \in [1, N^{\text{in}}], \ X_k(t) = 1 \mid \tau_n > t \right\} \ge 1 - e^{-\gamma_n t} \frac{K_n H_n^{\text{in}}}{\mathbf{P}_{\Pi^{\text{in}}} \left\{ t < \tau_n \right\}}.$$

Proof. We start by observing that the following relation holds true,

$$\tau_n = \min(\tau_n^0, T_n^1, \dots, T_n^{N^{in}}),$$
(5.3)

where $T_n^k = \inf\{t > 0 \mid X_k(t) \ge n + 1\}$ for $k = 1, ..., N^{in}$ and

$$\tau_n^0 = \inf \left\{ t \ge 0 \mid \exists k > N^{\text{i}n}, X_k(t) \ge n + 1 \right\}. \tag{5.4}$$

Let $\mathbf{C}(0) = C \in \mathcal{E}_n$ deterministic, define $N = \sum_{i=2}^{\infty} C_i$ and $(i_1, \dots, i_N) \in [\![2, n]\!]^N$ given by labelling function such that $C_i = \#\{k \in [\![1, N]\!] \mid i_k = i\}$. Conditionally on their initial condition, all clusters $X_k(t)$ (starting at i_k) are independent from each other, thus the event

$$A_t = \{ \forall k \in [1, N], \ X_k(t) = 1 \}$$
 (5.5)

is independent of $(X_k)_{k>N}$ and thus independent of τ_n^0 . Then,

$$\mathbf{P}_C\left\{A_t \mid \tau_n > t\right\} = \mathbf{P}_C\left\{A_t \mid \min(T_n^1, \dots, T_n^N) > t\right\}.$$

Still by independence of the clusters from each other, we claim that

$$\mathbf{P}_{C}\left\{A_{t} \mid \min(T_{n}^{1}, \dots, T_{n}^{N}) > t\right\} = \prod_{k=1}^{N} \mathbf{P}\left\{X(t) = 1 \mid X(0) = i_{k}, T_{n} > t\right\}$$
 (5.6)

where X is defined in Sec. 3 and T_n in (3.4). This equation is clear for N=1, and is easily proved by induction. We do it only for N=2, for the sake of simplicity. Let i_1 , $i_2 \in [2, n]$. By definition of A_t ,

$$\mathbf{P}_{C}\left\{A_{t} \mid \min(T_{n}^{1}, T_{n}^{2}) > t\right\} = \frac{\mathbf{P}_{C}\left\{X_{1}(t) = 1, X_{2}(t) = 1, T_{n}^{1} > t, T_{n}^{2} > t\right\}}{\mathbf{P}_{C}\left\{T_{n}^{1} > t, T_{n}^{2} > t\right\}}.$$

Then, by independence of the X_1 and X_2 conditionally on their initial condition we have

$$\mathbf{P}_{C}\left\{A_{t} \mid \min(T_{n}^{1}, T_{n}^{2}) > t\right\} = \prod_{k=1}^{2} \frac{\mathbf{P}\left\{X_{k}(t) = 1, T_{n}^{k} > t, X_{k}(0) = i_{k}\right\}}{\mathbf{P}\left\{T_{n}^{k} > t, X_{k}(0) = i_{k}\right\}}$$

$$= \prod_{k=1}^{2} \mathbf{P}\left\{X(t) = 1 \mid X(0) = i_{k}, T_{n} > t\right\}.$$

since X_1 and X_2 are independent copy of X, which proves the desired result. Thus, going back to (5.6) and thanks to (3.8) we have

$$\mathbf{P}_{C} \{ A_{t} \mid \tau_{n} > t \} \ge \prod_{k=1}^{N} (1 - M_{i_{k},n} e^{-\gamma_{n} t} \wedge 1) \ge 1 - e^{-\gamma_{n} t} \sum_{k=1}^{N} M_{i_{k},n}.$$

Finally, we obtain

$$\mathbf{P}_{\Pi^{\text{in}}} \left\{ A_{t} \mid \tau_{n} > t \right\} = \sum_{C \in \mathcal{E}_{n}} \mathbf{P}_{C} \left\{ A_{t} \mid \tau_{n} > t \right\} \frac{\mathbf{P}_{C} \left\{ \tau_{n} > t \right\}}{\mathbf{P}_{\Pi^{\text{in}}} \left\{ \tau_{n} > t \right\}} \Pi^{\text{in}}(C)
\geq 1 - e^{-\gamma_{n}t} \sum_{C \in \mathcal{E}_{n}} \sum_{k=1}^{N} M_{i_{k}, n} \frac{\mathbf{P}_{C} \left\{ \tau_{n} > t \right\}}{\mathbf{P}_{\Pi^{\text{in}}} \left\{ \tau_{n} > t \right\}} \Pi^{\text{in}}(C) \geq 1 - e^{-\gamma_{n}t} \frac{K_{n} H_{n}^{\text{in}}}{\mathbf{P}_{\Pi^{\text{in}}} \left\{ \tau_{n} > t \right\}},$$

where in the second line, N and $(i_k)_{k=1..N}$ are given by the *labelling* function for each $C \in \mathcal{E}_n$, and using that $\mathbf{P}_C \{ \tau_n > t \} \leq 1$ in the last inequality. Remark the expression of H_n^{in} is obtained thanks to the definition of $M_{i,n}$ in (3.8) and f_i^n in (5.2).

Proof of the Theorem 5.2. Let the collection of processes X_1, X_2, \ldots (resp. Y_1, Y_2, \ldots) being a particle description of the BD process that starts from the initial distribution Π^{in} (resp. from δ_0). We couple the processes X_1, X_2, \ldots to the processes Y_1, Y_2, \ldots as in the proof of Theorem 5.2, namely, $Y_k(t) = X_{k+N^{in}}(t)$ for all $k \geq 1$ and all $t \geq 0$, where N^{in} is distributed according to Π^{in} .

To avoid notation confusion, we write τ_n^X and τ_n^Y the first exit time from \mathcal{E}_n for the collection of processes $\{X_k\}$ and $\{Y_k\}$, respectively. We define $\tau_n^{0,X}$ and $\tau_n^{0,Y}$, respectively to the processes $\{X_k\}$ and $\{Y_k\}$ likewise τ_n^0 in (5.4). Due to the coupling between the $\{X_k\}$ and $\{Y_k\}$, we have

$$\tau_n^{0,Y} = \tau_n^Y = \inf \{ t \ge 0 \mid \exists k > 0, Y_k(t) \ge n+1 \}$$

$$= \inf \{ t \ge 0 \mid \exists k > N^{\text{i}n}, X_k(t) \ge n+1 \} = \tau_n^{0,X}. \quad (5.7)$$

Remark that each Y_k , for $k \geq 1$, is independent of X_i for $i \leq N^{in}$, thus independent of $T_n^1,\ldots,T_n^{N^{in}}$ the exit times arising in (5.3). Hence, by (5.3) and (5.7), the laws of the collection of processes $\{Y_k\}$ conditioned to $\tau_n^Y > t$ equals to the laws of the collection of processes $\{Y_k\}$ conditioned to $\tau_n^X > t$. Also, we have, for any $i \geq 2$ and t > 0, $\#\{k \mid X_k(t) = i\} = \#\{k \mid Y_k(t) = i\}$ on the event A_t , given in (5.5), since all initial particles being absorbed. Finally, we deduce that

$$\begin{aligned} &\|\mathbf{P}_{\Pi^{\text{i}n}}\left\{\mathbf{C}(t) \in \cdot \mid \tau_n > t\right\} - \mathbf{P_0}\left\{\mathbf{C}(t) \in \cdot \mid \tau_n > t\right\} \| \\ &\leq \mathbf{P}\left\{\exists i \geq 2, \#\left\{k \mid X_k(t) = i\right\} \neq \#\left\{k \mid Y_k(t) = i\right\} \mid \tau_n^X > t\right\} \leq \mathbf{P}_{\Pi^{\text{i}n}}\left\{A_t^c \mid t < \tau_n^X\right\} \; . \end{aligned}$$

The latter, with Lemma 5.3, entails

$$\|\mathbf{P}_{\Pi^{in}} \left\{ \mathbf{C}(t) \in \cdot \mid \tau_n > t \right\} - \mathbf{P_0} \left\{ \mathbf{C}(t) \in \cdot \mid \tau_n > t \right\} \| \le e^{-\gamma_n t} \frac{K_n H_n^{in}}{\mathbf{P}_{\Pi^{in}} \left\{ t < \tau_n \right\}}.$$
 (5.8)

Then, applying estimate (5.8) with the initial distribution Π_n^{qsd} since $\mathbf{E}_{\Pi_n^{qsd}}\left[\sum_{i=2}^{\infty}C_i\right]=\sum_{i=2}^n f_i^n<\infty$ and by Proposition 5.1, we deduce

$$\|\prod_{n}^{\operatorname{qsd}} - \mathbf{P_0} \left\{ \mathbf{C}(t) \in \cdot \mid t < \tau_n \right\} \| \le e^{-\gamma_n t} K_n e^{J_n t} H_n^{\operatorname{qsd}}.$$

We end the proof by triangular inequality.

6 Estimates on τ_n and the largest cluster

In this section we consider the super-critical case $z>z_s$. The analysis of the τ_n in (5.1) leads off the simple observation

$$\tau_n > t \Leftrightarrow \forall s \le t, \max_{1 \le k \le N(s)} X_k(s) \le n,$$

where X_1, X_2, \ldots is the particle description of the BD process. We prove

Theorem 6.1. Under assumption (H0). Let $z>z_s$ and $\Pi^{\mathrm{i}n}$ a probability distribution on \mathcal{E}_n such that $\mathbf{E}_{\Pi^{\mathrm{i}n}}\left[\sum_{i=2}^{\infty}C_i\right]<\infty$. We have

$$\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\tau_n > t\right\} \ge G_n^{\mathrm{i}n} e^{-J_n t}$$

where

$$G_n^{in} = \mathbf{E}_{\Pi^{in}} \left[\prod_{i=1}^n \left(\frac{f_i^n}{Q_i z^i} \right)^{C_i} \right] \ge 1 - \sum_{i=2}^n \mathbf{E}_{\Pi^{in}} \left[C_i \right] \left(1 - \frac{f_i^n}{Q_i z^i} \right).$$

In fact, as in the coupling strategy, (5.3) provides a useful understanding of the statistics of τ_n by decomposing between the initial cluster from the ones that will appear at later times. We start with the statistics of the later, namely of τ_n^0 . The next lemma clearly applies for the initial distribution $\Pi^{in} = \delta_0$ but is more general, and related to a maximum principle in the BD model [3].

Lemma 6.2. Under assumption (H0) and $z > z_s$. For any probability distribution Π^{in} on \mathcal{E}_n such that for all $i \in [2, n]$ and $k \in \mathbb{N}$,

$$\mathbf{P}_{\Pi^{\mathrm{i}n}} \left\{ \sum_{i \le j \le n} C_j \ge k \right\} \le \mathbf{P}_{\Pi^{\mathrm{q}sd}} \left\{ \sum_{i \le j \le n} C_j \ge k \right\}, \tag{6.1}$$

we have

$$\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\tau_n > t\right\} \ge e^{-J_n t}.$$

Proof of Lemma 6.2. It is classical that condition (6.1) ensures there exists randoms $\mathbf{C}^{\mathrm{i}n}$ and $\mathbf{C}^{\mathrm{q}sd}$ distributed according to $\Pi^{\mathrm{i}n}$ and $\Pi^{\mathrm{q}sd}$, respectively, such that for each $i \in [\![2,n]\!]$,

$$\sum_{i \leq j \leq n} C_j^{\mathrm{i}n} \leq \sum_{i \leq j \leq n} C_j^{\mathrm{q}sd} \,, \quad a.s.$$

see e.g. [5, Sec. 4.12]. Then, we may construct the collection of processes X_1, X_2, \ldots (resp. Y_1, Y_2, \ldots) as a particle description of the BD process associated to \mathbf{C}^{in} (resp. to \mathbf{C}^{qsd}) such that, a.s., for all $i \geq 1$, $X_i(0) \leq Y_i(0)$. A standard coupling between two copies of the chain X from Sec. 3 consists in having the same jumps in the two copies as soon as they are equal. Such coupling applied to each couple (X_i, Y_i) then ensures that, for all $i \geq 1$ and $t \geq 0$, we have $X_i(t) \leq Y_i(t)$ a.s. In particular,

$$\inf\{t > 0 \mid \max_{k} X_k(t) > n\} \ge \inf\{t > 0 \mid \max_{k} Y_k(t) > n\} \quad a.s.$$

thus, with Proposition 5.1,

$$\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\tau_{n} > t\right\} \ge \mathbf{P}_{\Pi^{\mathrm{q}sd}}\left\{\tau_{n} > t\right\} = e^{-J_{n}t}.$$

Proof of Theorem 6.1. Let $n \geq 2$ and $i \in [2, n]$. Define $g_{n,i}(t) = \mathbf{P}\{T_n > t \mid X(0) = i\}$ where T_n and X are given in Sec. 3. Thanks to (3.6), we have

$$g_{n,i}(t) \ge \lim_{t \to +\infty} g_{n,i}(t) = J_n \sum_{k=i}^n \frac{1}{a_k Q_k z^{k+1}} = \frac{f_i^n}{Q_i z^i}$$
 (6.2)

Let $\psi_n(t,x)=1$ if $\max_{0\leq \tau\leq t} x(\tau)\leq n$ and 0 otherwise. We have

$$\mathbf{P}_{\Pi^{\mathrm{i}n}}\left\{\tau_n > t\right\} = \sum_{C \in \mathcal{E}_n} \mathbf{P}_C \left\{ \prod_{i=1}^{N(t)} \psi_n(t, X_i) = 1 \right\} \Pi^{\mathrm{i}n}(C)$$

where X_1, X_2, \ldots the particle description of the BD process C (starting at δ_C). By independence of the particles conditionally to their initial condition,

$$\mathbf{P}_{C} \left\{ \prod_{i=1}^{N(t)} \psi_{n}(t, X_{i}) = 1 \right\} = \mathbf{P}_{C} \left\{ \prod_{i=N^{in}+1}^{N(t)} \psi_{n}(t, X_{i}) = 1 \right\} \mathbf{P}_{C} \left\{ \prod_{i=1}^{N^{in}} \psi_{n}(t, X_{i}) = 1 \right\}$$

$$= \mathbf{P}_{0} \left\{ \tau_{n} > t \right\} \prod_{k=1}^{N} g_{n, i_{k}}(t) \quad (6.3)$$

where again, N and $(i_k)_{k=1..N}$ are given by the *labelling* function associated to $C \in \mathcal{E}_n$. Combining relations (6.2) and (6.3) with Lemma 6.2 and summing over all initial conditions ends the proof.

7 Metastability close to z_s

In this section we assume additionally to (H1) and (H2), to fit with [10, 11], that

$$A' < a_i < Ai^{\alpha}, \quad \frac{b_{i+1}}{a_{i+1}} + \frac{\kappa}{i^{\nu}} \le \frac{b_i}{a_i} \quad \text{and} \quad z_s e^{Gi^{-\gamma}} \le \frac{b_i}{a_i} \le z_s e^{G'i^{-\gamma'}}, \tag{H3}$$

for all $i \geq 2$, where $\alpha, \gamma \in (0,1), \gamma', \nu > 0$, κ , A', A, G and G' positives. We also use the terminology of [11] namely a quantity q(z) of z is: exponentially small if $q(z)/(z-z_s)^m$ is bounded for all m > 0 as $z \setminus z_s$ (z converges to z_s and $z > z_s$); and at most algebraically large if $(z-z_s)^{m_0}q(z)$ is bounded for some $m_0 > 0$ as $z \setminus z_s$.

Assumption (H3) ensures the existence of a unique n^* (depending on z) such that $b_{n^*+1}/a_{n^*+1} < z < b_{n^*}/a_{n^*}$. The size n^* is interpreted as the nucleus size: for a cluster $X(t) \le n^*$, X(t) tends to shorten, while for $X(t) > n^*$, it tends to grow. With assumption (H3), $n^* \to \infty$ as $z \setminus z_s$. In [10, 11], n^* is proved to be at most algebraically large. Moreover, the time scale $1/\gamma_{n^*}$ (in Theorem (5.2)) is also at most algebraically large, and J_{n^*} (in Theorem 6.1) is exponentially small. We now choose an initial distribution Π^{in} with support on \mathcal{E}_j , with j independent of z (no generality is claimed here). We have

$$\mathbf{P}_{\Pi^{in}} \left\{ \tau_{n^*} > t \right\} \ge \left(1 - \sum_{i=2}^{j} \mathbf{E}_{\Pi^{in}} \left[C_i \right] \left(1 - \frac{f_i^{n^*}}{Q_i z^i} \right) \right) e^{-J_{n^*} t}, \tag{7.1}$$

which is arbitrary close to one for times $t \ll 1/J_{n^*}$ as $f_i^{n^*}/(Q_iz^i) \to 1$ when $z \setminus z_s$. Moreover, we have

$$\|\mathbf{P}_{\Pi^{\text{i}n}}\left\{\mathbf{C}(t) \in \cdot \mid \tau_{n^*} > t\right\} - \Pi_n^{\text{q}sd}\| \le \left(\frac{H_j^{\text{i}n}}{G^{\text{i}n}} + H_{n^*}^{\text{q}sd}\right) K_{n^*} e^{J_{n^*}t - \gamma_{n^*}t}$$
(7.2)

which is arbitrary small for times $1/\gamma_{n^*} \ll t \ll 1/J_{n^*}$. Indeed, note that $a_iQ_iz^i$ is decreasing up to the size n^* , thus $K_{n^*}^2 \leq \frac{a_1z}{A'}n^*$. Hence, K_{n^*} as well as $H_{n^*}^{qsd}$ are at most algebraically large. Equations (7.1)-(7.2) show the QSD is indeed a metastable state.

References

- [1] W. J. Anderson, Continuous-time Markov chains, Springer Series in Statistics: Probability and its Applications, Springer-Verlag, New York, 1991, An applications-oriented approach. MR 1118840
- [2] R. Becker and W. Döring, *Kinetische behandlung der keimbildung in übersättigten dämpfen*, Annalen der Physik **416** (1935), no. 8, 719–752.
- [3] J. A. Cañizo, A. Einav, B. Lods, *Uniform moment propagation for the Becker–Döring equations*, Proc. R. Soc. Edinb, **149** (2019), 995–1015,
- [4] P. Collet, S. Martínez, and J. San Martín, Quasi-stationary distributions, Probability and its Applications (New York), Springer, Heidelberg, 2013, Markov chains, diffusions and dynamical systems. MR 2986807
- [5] G. R. Grimmett and D. R. Stirzaker, Probability and random processes, third ed., Oxford University Press, New York, 2001. MR 2059709
- [6] E. Hingant and R. Yvinec, Deterministic and stochastic Becker-Döring equations: past and recent mathematical developments, Stochastic processes, multiscale modeling, and numerical methods for computational cellular biology, Springer, Cham, 2017, pp. 175–204. MR 3726370
- [7] _____, The Becker-Döring process: pathwise convergence and phase transition phenomena, J. Stat. Phys. 177 (2019), no. 3, 506–527. MR 4026657
- [8] S. Karlin and J. McGregor, *The classification of birth and death processes*, Trans. Amer. Math. Soc. **86** (1957), 366–400. MR 94854
- [9] J. F. C. Kingman, Markov population processes, J. Appl. Probability 6 (1969), 1–18.MR 254934
- [10] M. Kreer, Classical Becker-Döring cluster equations: rigorous results on metastability and long-time behaviour, Ann. Physik (8) 2 (1993), no. 4, 398–417. MR 1223849
- [11] O. Penrose, Metastable states for the Becker-Döring cluster equations, Comm. Math. Phys. **124** (1989), no. 4, 515–541. MR 1014113
- [12] P. S. Ruzankin, On the rate of Poisson process approximation to a Bernoulli process, J. Appl. Probab. **41** (2004), no. 1, 271–276. MR 2036288
- [13] J. W. P. Schmelzer (ed.), Nucleation theory and applications, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, 2005.

QSD and metastability for the Becker-Döring model

[14] W. Sun, A functional central limit theorem for the Becker-Döring model, J. Stat. Phys. **171** (2018), no. 1, 145–165. MR 3773855

Acknowledgments. E. H. has to thanks the partial support by FONDECYT project n. 11170655.