

Quantization of Chern-Simons topological invariants for H-type and L-type quantum systems

Oscar Randal-Williams,¹ Lokman Tsui,² and Xiao-Gang Wen²

¹*DPMMS, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, UK*

²*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

In 2+1-dimensions (2+1D), a gapped quantum phase with no symmetry (*i.e.* a topological order) can have a thermal Hall conductance $\kappa_{xy} = c \frac{\pi^2 k_B^2}{3h} T$, where the dimensionless c is called chiral central charge. If there is a U_1 symmetry, a gapped quantum phase can also have a Hall conductance $\sigma_{xy} = \nu \frac{e^2}{h}$, where the dimensionless ν is called filling fraction. In this paper, we derive some quantization conditions of c and ν , via a cobordism approach to define Chern-Simons topological invariants which are associated with c and ν . In particular, we obtain quantization conditions that depend on the ground state degeneracies on Riemannian surfaces, and quantization conditions that depend on the type of spacetime manifolds where the topological partition function is non-zero.

CONTENTS

I. Introduction	1	B. Quantization of c and ν for fermionic U_1^f -enriched topological orders with non-zero partition functions	12
A. Notations and conventions	3	VII. Summary	12
II. The vector bundle on the moduli space for H-type bosonic systems with gapped liquid ground states	3	A. Characteristic numbers of 4-manifolds	12
III. Chern-Simons invariants in 2+1D H-type bosonic U_1 -SET and U_1 -SPT orders	3	B. Characteristic numbers of surface bundles over surfaces	13
A. Bosonic gapped liquids with U_1 symmetry in 2-dimensional space	3	1. Definitions and relations	13
B. Examples	5	2. Realising characteristic numbers	14
1. Bosonic K -matrix Abelian topological orders	5	C. Checking (29), (30), (31)	14
2. Bosonic non-Abelian topological orders described by $SU(2)_k$ Chern-Simons Theory	6	1. Spread polynomials $S_n(x)$	14
C. A general point of view	6	2. Symmetric polynomials e_l and p_l	14
IV. Topological invariants in 2+1D H-type fermionic enriched topological orders with U_1^f symmetry	6	3. Checking (29)	15
A. Symmetry twist of fermion system	6	4. Checking (30)	15
B. U_1^f symmetry and spin^c structure	7	5. Checking (31)	15
C. Fermionic gapped liquids with U_1^f symmetry in 2-dimensional space	8	D. Group extension and trivialization	15
D. H-type invertible fermionic U_1^f -enriched topological orders	9	References	16
V. The topological invariants for L-type topological orders	9		
A. Topological partition function for L -type topological order	9		
B. The quantization of c for 2+1D bosonic topological orders with non-zero partition function	10		
C. The quantization of c for 2+1D bosonic topological orders with vanishing partition functions only on non-spin manifolds and for 2+1D fermionic invertible topological orders	10		
D. Framing anomaly	11		
VI. The topological invariants for L-type topological orders with U_1 symmetry	11		
A. Quantization of c and ν for bosonic U_1 -SET orders with non-zero partition functions	11		

I. INTRODUCTION

Different phases of matter are characterized by different orders in them.[1, 2] Topological order[3–5] is a new kind of order beyond Landau symmetry breaking order.[1, 2] It cannot be characterized by the local order parameters associated with the symmetry breaking, but can be characterized by topological quantum field theories.[6] Physically, we need to use new topological quantum numbers to characterize topological orders. In 2+1D, the chiral central charge c [7, 8] (*i.e.* the thermal Hall conductance $\kappa_{xy} = c \frac{\pi^2 k_B^2}{3h} T$ [9]) of the edge excitations is one such topological quantum number. When there is a U_1 symmetry, the Hall conductance $\sigma_{xy} = \nu \frac{e^2}{h}$ is another such topological quantum number. In other words, the chiral central charge c (partially) characterize 2+1D topological orders, and the dimensionless Hall conductance ν , together with c , (partially) characterize 2+1D topological orders with U_1 -symmetry.[10]

In this paper, we like to address the issue of the quantization of those topological quantum numbers. At first sight, it seems that c and ν are not quantized since they do not have to be integers. On the other hand, c and ν must be rational numbers, suggesting that they do satisfy certain quantization conditions which can be complicated. A quantization condition on ν for interacting systems was first obtained by Niu–Thouless–Wu in Ref. [11]. In this paper, we are going to generalize their result and discuss those complicated quantization conditions that apply to all topological orders with U_1 -symmetry.

Before calculating those quantization conditions, we need to distinguish two types of quantum systems: H-type and L-type.[12] The two types of quantum systems have different quantization conditions.

The quantum systems in condensed matter physics are all H-type, *i.e.* are all described by local Hamiltonians defined on smooth spatial manifolds. The quantum systems in high energy theory and quantum field theory, such as the topological quantum field theories, are L-type, *i.e.* are described by local Lagrangian path integrals on smooth spacetime manifolds. Mathematically, the L-type topological orders may correspond to unitary fully extended topological quantum field theories. The H-type topological orders may correspond to topless fully extended topological quantum field theories (or $d + \epsilon$ dimensional fully extended topological quantum field theories, see [13]).

The gapped liquid states[14, 15] with U_1 symmetry include U_1 -symmetry enriched topological (SET) orders[10, 16–18] and U_1 -symmetry protected trivial (SPT) orders.[19–25] For 2+1D H-type gapped states with U_1 symmetry, the quantization of c and ν is determined from the ground state degeneracy on closed genus g surfaces, and the main results are given by eqn. (20) and eqn. (63).

In particular, for H-type invertible gapped states¹ with U_1 symmetry, c and ν satisfy the following quantization conditions:

$$\begin{aligned} \text{bosonic systems: } & c, \nu = 0 \bmod 2. \\ \text{fermionic systems: } & c, \nu, \frac{\nu - c}{2} = 0 \bmod 1. \end{aligned} \quad (1)$$

The even Hall conductance $\nu = 0 \bmod 2$ for bosonic SPT phases was pointed out in Ref. 22–25. We remark that the known H-type invertible topological orders with U_1 symmetry do not saturate the above quantization conditions. For example, we do not know any H-type bosonic invertible topological orders with $c = 2$. Also, we do not know any H-type fermionic U_1^f symmetric invertible topological orders with $\nu - c = 2$.

For 2+1D L-type gapped states with U_1 symmetry, the quantization of c and ν is determined by the non-vanishing topological partition functions on certain types

of spacetime manifolds, such as orientable or spin^c . We find that, for L-type invertible gapped phases with U_1 symmetry, c and ν satisfy the following quantization conditions:

$$\begin{aligned} \text{bosonic systems: } & c = 0 \bmod 8, \nu = 0 \bmod 2. \\ \text{fermionic systems: } & c, \nu, \frac{\nu - c}{8} = 0 \bmod 1. \end{aligned} \quad (2)$$

The above L-type gapped phases with U_1 symmetry have a framing anomaly, *i.e.* the partition functions of the states dependent on the choices of the framing of the spacetime manifold.[6]

If the microscopic Lagrangian path integral is manifestly independent of choices of the framing (for example, only dependent on the diffeomorphism equivalent classes of spacetime metrics), then the resulting partition must also be independent of choices of the framing. The resulting phase is said to be free of framing anomaly. In this case, c must be multiple of 24, and the above quantization conditions reduce to

$$\begin{aligned} \text{bosonic systems: } & c = 0 \bmod 24, \nu = 0 \bmod 2. \\ \text{fermionic systems: } & c = 0 \bmod 24, \nu = 0 \bmod 8. \end{aligned} \quad (3)$$

It is interesting to note that the bosonic E_8 quantum Hall state with $c = 8$ (see eqn. (21) with K given by the E_8 matrix (A5)) is an H-type topological order. But it cannot be realized by a topological quantum field theory with no framing anomaly. In other words, it cannot be realized by a Lagrangian path integral that manifestly only depends on the diffeomorphism equivalent classes of spacetime metrics.

However, the E_8 state can be realized by a Lagrangian path integral with framing anomaly (*i.e.* depends on the additional framing structure of spacetime manifold).[6] One such realization is given by the dynamical Chern–Simons theory eqn. (22) with $K = E_8$. Here we like to remark that it is highly nontrivial to define the dynamical Chern–Simons theory eqn. (22) on spacetime so that we can compute the partition function via a finite calculation. For bosonic Chern–Simons theory eqn. (22) (where K_{II} is even), a non perturbative definition was recently given in Ref. 26, by triangulating the spacetime and giving the triangulation a branching structure. It appears that the branching structure plays a role of framing structure.

On the other hand, the bosonic E_8^3 quantum Hall state (the stacking of three bosonic E_8 quantum Hall state) can be described by a topological quantum field theory with no framing anomaly, *i.e.* it can be realized by a Lagrangian path integral that only depends on the diffeomorphism equivalent classes of spacetime metrics (*i.e.* does not depend on the framing structure of spacetime). An explicit construction of the Lagrangian path integral in terms of SO_∞ non-linear σ -model was given in Ref. 27.

¹ By definition, invertible gapped states have no fractionalized excitations. As a result, they have non-degenerate ground state on closed smooth spatial manifolds.

A. Notations and conventions

We will abbreviate the cup product $a \smile b$ and the wedge product $a \wedge b$ as ab . We will use $\stackrel{n}{=}$ to mean equal up to a multiple of n , and use $\stackrel{d}{=}$ to mean equal up to df (*i.e.* up to a coboundary). We will use $\lfloor x \rfloor$ to denote the greatest integer less than or equal to x , and $\langle l, m \rangle$ for the greatest common divisor of l and m (with $\langle 0, m \rangle \equiv m$).

We introduce a symbol \bowtie to construct fiber bundle $E = F \bowtie B$ from the fiber F and the base space B :

$$\text{pt} \rightarrow F \rightarrow E \rightarrow B \rightarrow \text{pt}. \quad (4)$$

We will also use \bowtie to construct group extension of G by N [28]:

$$1 \rightarrow N \rightarrow N \bowtie_{e_2, \alpha} G \rightarrow G \rightarrow 1. \quad (5)$$

Here $e_2 \in H^2[G; Z(N)]$ and $Z(N)$ is the center of N . Also G may have a non-trivial action on $Z(N)$ via $\alpha : G \rightarrow \text{Aut}(Z(N))$. e_2 and α characterize different group extensions.

Also, we will use $Z_n = \{1, e^{i\frac{2\pi}{n}}, e^{i2\frac{2\pi}{n}}, \dots, e^{i(n-1)\frac{2\pi}{n}}\}$ to denote an Abelian group, where the group multiplication is “ \cdot ”. We use $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$ to denote an integer lifting of Z_n , where “ $+$ ” is done without mod- n . In this sense, \mathbb{Z}_n is not a group under “ $+$ ”. But under a modified equality $\stackrel{n}{=}$, \mathbb{Z}_n is the Z_n group under “ $+$ ”. Similarly, we will use $\mathbb{R}/\mathbb{Z} = [0, 1)$ to denote an \mathbb{R} -lifting of U_1 group. Under a modified equality $\stackrel{1}{=}$, \mathbb{R}/\mathbb{Z} is the U_1 group under “ $+$ ”. In this paper, we have expressions containing the addition “ $+$ ” of \mathbb{Z}_n -valued or \mathbb{R}/\mathbb{Z} -valued, such as $a_1^{\mathbb{Z}_n} + a_2^{\mathbb{Z}_n}$ where $a_1^{\mathbb{Z}_n}$ and $a_2^{\mathbb{Z}_n}$ are \mathbb{Z}_n -valued. Those additions “ $+$ ” are done without mod n or mod 1. In this paper, we also have expressions like $\frac{1}{n}a_1^{\mathbb{Z}_n}$. Such an expression converts a \mathbb{Z}_n -valued $a_1^{\mathbb{Z}_n}$ to a \mathbb{R}/\mathbb{Z} -valued $\frac{1}{n}a_1^{\mathbb{Z}_n}$, by viewing the \mathbb{Z}_n -value as a \mathbb{Z} -value. (In fact, \mathbb{Z}_n is a \mathbb{Z} lifting of Z_n .)

II. THE VECTOR BUNDLE ON THE MODULI SPACE FOR H-TYPE BOSONIC SYSTEMS WITH GAPPED LIQUID GROUND STATES

In this paper, we study topological invariants for gapped liquid [14, 15] ground states without excitations. In order to characterize different classes of gapped liquid ground states, we consider a moduli space – the space of Hamiltonians with on-site symmetry G that have gapped liquid state as the ground state. Here we assume that the Hamiltonians in the moduli space depend on the metrics g_{ij} of the space as well as the G -symmetry twist (*i.e.* the background G -gauge field A_i from gauging the on-site G -symmetry [29]). So for a closed n -dimensional space M^n , with a certain topology as well as a G -principle bundle for the G -symmetry twist on M^n , the moduli space is the space of spatial metrics and G -connections, which

is denoted as $\mathcal{M}_{G \bowtie M^n}$. Here $G \bowtie M^n$ describes the G -principle bundle:

$$\text{pt} \rightarrow G \rightarrow G \bowtie M^n \rightarrow M^n \rightarrow \text{pt}. \quad (6)$$

Also

we assume that the pairs (g_{ij}, A_i) differ by gauge transformations and diffeomorphisms are equivalent and represent the same point of the moduli space $\mathcal{M}_{G \bowtie M^n}$.

Since on every point of the moduli space, the corresponding Hamiltonian is in a gapped liquid phase, the subspace formed by the degenerate ground states can be viewed as the fiber at the point. This way, we obtain a vector bundle over the moduli space $\mathcal{M}_{G \bowtie M^n}$. It was conjectured in Ref. 5 that

the topology of this vector bundle completely characterize the H-type topological orders, *i.e.* gapped liquid phases with symmetry.

In particular, let us consider a closed subspace B in $\mathcal{M}_{G \bowtie M^n}$. The vector bundle on $\mathcal{M}_{G \bowtie M^n}$ reduces to a vector bundle on B . In the next section, we concentrate on this vector bundle on B , and try to relate the vector bundle to topological term in the partition function of the quantum system under consideration. We will follow closely the approach proposed in Ref. 12 and 30.

III. CHERN–SIMONS INVARIANTS IN 2+1D H-TYPE BOSONIC U_1 -SET AND U_1 -SPT ORDERS

A. Bosonic gapped liquids with U_1 symmetry in 2-dimensional space

In this section, we consider gapped liquids with U_1 symmetry in 2-dimensional space for bosonic systems. Let Σ_g be a 2-dimensional closed spatial manifold with a U_1 connection, that describes the twisted global U_1 symmetry. The corresponding U_1 -bundle is given by $U_1 \bowtie \Sigma_g$. In this case, the moduli space $\mathcal{M}_{U_1 \bowtie \Sigma_g}$ of the system is the space of metrics on Σ_g and the U_1 -connections. The degenerate ground states on Σ_g give rise to a vector bundle over the moduli space $\mathcal{M}_{U_1 \bowtie \Sigma_g}$, where the dimension of the vector is given by the ground state degeneracy $D_{U_1 \bowtie \Sigma_g}$.

Let us consider a loop S^1 in the moduli space $\mathcal{M}_{U_1 \bowtie \Sigma_g}$. The holonomy of the vector bundle around the loop is given by a unitary matrix of $D_{U_1 \bowtie \Sigma_g}$ dimension: W_{S^1} . We note that the unitary matrix W_{S^1} is the non-Abelian geometric phase [31] of the degenerate ground states under the adiabatic deformation around the loop S^1 .

From the vector bundle, we can obtain a determinant bundle, whose holonomy around the loop is given by the determinant: $\text{Det}(W_{S^1})$. This phase factor is directly related to the topological term in the effective action

$$S_{eff} = \int_{\Sigma_g \times S^1} d^3x \mathcal{L}_{eff}:$$

$$\text{Det}(W_{S^1}) = \left(e^{i \int_{\Sigma_g \times S^1} d^3x \mathcal{L}_{eff}} \right)^{D_{U_1 \times \Sigma_g}} \quad (7)$$

The effective action may contain a gravitational Chern–Simons term and a U_1 Chern–Simons term:[32–34]

$$\begin{aligned} \mathcal{L}_{eff} d^3x &= -2\pi \frac{c}{24} \omega_3 + \frac{\nu}{4\pi} (A + sA^{SO_2}) d(A + sA^{SO_2}) \\ &= -2\pi \frac{c}{24} \omega_3 + \frac{\nu}{4\pi} A_{eff} dA_{eff} \end{aligned} \quad (8)$$

where A is the U_1 connection 1-form describing the total electromagnetic field, s is the orbital spin carried by the charged bosons,[32, 33] and A^{SO_2} is the time-dependent connection 1-form describing the SO_2 tangent bundle of the curved space. $A_{eff} \equiv A + sA^{SO_2}$ is the effective U_1 connection 1-form. Also ω_3 is the gravitational Chern–Simons term that satisfy $d\omega_3 = p_1$, where p_1 is the first Pontryagin class of the tangent bundle. In the above, c is the chiral central charge of the edge theory, and ν is proportional to the Hall conductance:

$$\sigma_{xy} = \frac{\nu}{2\pi} = \nu \frac{e^2}{h} \quad (9)$$

(in $e = \hbar = 1$ unit).

We note that $\frac{\nu}{2\pi} dA_{eff} = j$ is the 2-cocycle describing the conserved density and current of charged bosons in the ground state. Thus A_{eff} satisfy the quantization condition

$$\int_{\Sigma_g} \frac{\nu}{2\pi} dA_{eff} \in \mathbb{Z} \quad (10)$$

for the closed 2-dimensional space Σ_g . In fact, $\int_{\Sigma_g} \frac{\nu}{2\pi} dA_{eff}$ is the number of bosons in the ground state.

Now assume the S^1 to be the boundary of a 2-dimension subspace B^2 in $\mathcal{M}_{U_1 \times \Sigma_g}$. Eqn. (7) can be rewritten as[35]

$$\text{Det}(W_{S^1}) = e^{D_{U_1 \times \Sigma_g} i 2\pi \int_{\Sigma_g \times B^2} -\frac{c}{24} p_1 + \frac{\nu}{2} c_1^2}, \quad (11)$$

where $a = \frac{A_{eff}}{2\pi}$ and $c_1 = da$ is the first Chern class. If we shrink S^1 to a point, B^2 becomes a closed 2-dimensional subspace. In this case, $\text{Det}(W_{S^1}) = 1$ and

$$D_{U_1 \times \Sigma_g} \int_{\Sigma_g \times B^2} -\frac{c}{24} p_1 + \frac{\nu}{2} c_1^2 \in \mathbb{Z}, \quad (12)$$

for any orientable surface bundle $\Sigma_g \times B^2$ and for any U_1 bundle whose first Chern class c_1 satisfies $\nu \int_{\Sigma_g} c_1 \in \mathbb{Z}$.

This result allows us to obtain quantization condition of chiral central charge c and the filling fraction ν . In particular, every choice of a surface bundle $\Sigma_g \times B^2$ and U_1 bundle will give us a quantization condition of c and

ν . The combination of all those quantization conditions gives us the strictest constraint on the possible values of c and ν . In the following, we will choose some special combination of surface bundles $\Sigma_g \times B^2$ and U_1 bundles to obtain concrete quantization conditions of c and ν .

We note that $\int_{\Sigma_g \times B^2} p_1 = 0 \bmod 12$ for any orientable surface bundles (also called Σ_g -bundles), by (B3). If the genus g of the fiber Σ_g is equal or less than 2, then $\int_{\Sigma_g \times B^2} p_1 = 0$. [36] If $g \geq 3$, then we can always find a base manifold B^2 with a genus equal or less than 111, such that there is a surface bundle $\Sigma_g \times B^2$ with $\int_{\Sigma_g \times B^2} p_1 = \pm 12$. [37] If we choose the U_1 bundle to be trivial $c_1 = 0$, eqn. (12) gives us the quantization condition on c first proposed in Ref. 12 and 30:

$$\frac{c}{2} D_{U_1 \times \Sigma_g} \in \mathbb{Z}, \quad \text{for } g \geq 3, \quad (13)$$

Next, we consider a trivial surface bundle with $S^2 = \Sigma_0^2$ as the fiber: $\Sigma_g \times B^2 = \Sigma_0 \times B^2$. On such a surface bundle, c_1 has a form $c_1 = c_1^B + c_1^\Sigma$ where c_1^Σ lives on Σ_0 . We have

$$\begin{aligned} \frac{1}{2} \nu \int_{\Sigma_0 \times B^2} c_1^2 &= \frac{1}{2} \nu \int_{\Sigma_0 \times B^2} (c_1^B + c_1^\Sigma)^2 \\ &= \nu \int_{\Sigma_0} c_1^\Sigma \times \int_{B^2} c_1^B \end{aligned} \quad (14)$$

Therefore, for U_1 bundles whose Chern class c_1 satisfying $\nu \int_{\Sigma_g} c_1 \in \mathbb{Z}$, $\frac{1}{2} \nu \int_{\Sigma_0 \times B^2} c_1^2$ is always an integer. No constraint on ν is obtained. Also, $\int_{\Sigma_0 \times B^2} p_1 = 0$. Therefore, the trivial surface bundle $\Sigma_g \times B^2 = \Sigma_0 \times B^2$ does not give us any non-trivial quantization conditions for c and ν . So in the following, we will assume $g > 0$ for our space Σ_g .

Next, we consider the surface bundle $\Sigma_g \times B^2 = S_x^1 \times S_y^1 \times S_a^1 \times S_b^1$, where $S_x^1 \times S_y^1 = \Sigma_1$ is the space and $S_a^1 \times S_b^1 = B^2$. We consider the U_1 bundle whose first Chern class has the following form $c_1 = c_1^{xa} + c_1^{yb}$, where c_1^{xa} lives on $S_x^1 \times S_a^1$ and c_1^{yb} lives on $S_y^1 \times S_b^1$. Since $\int_{\Sigma_0 \times B^2} p_1 = 0$ and $\int_{\Sigma_1} c_1^{xa} + c_1^{yb} = \int_{S_x^1 \times S_a^1} c_1^{xa} + \int_{S_y^1 \times S_b^1} c_1^{yb} = 0$, the quantization condition eqn. (12) becomes

$$\begin{aligned} \frac{\nu}{2} D_{U_1 \times \Sigma_1} \int_{S_x^1 \times S_y^1 \times S_a^1 \times S_b^1} (c_1^{xa} + c_1^{yb})^2 \\ = \nu D_{U_1 \times \Sigma_1} \int_{S_x^1 \times S_a^1} c_1^{xa} \int_{S_y^1 \times S_b^1} c_1^{yb}. \end{aligned} \quad (15)$$

We can choose the U_1 bundles such that $\int_{S_x^1 \times S_a^1} c_1^{xa} = 0, \pm 1$ and $\int_{S_y^1 \times S_b^1} c_1^{yb} = 0, \pm 1$. We obtain the following quantization condition

$$\nu D_{U_1 \times \Sigma_1} \in \mathbb{Z}. \quad (16)$$

The above result was first obtained by Ref. 11, using a similar consideration. Because Σ_g is the connected sum

of g Σ_1 's, the above reasoning can be generalized to the case $\Sigma_g \searrow B^2 = \Sigma_g \times S_a^1 \times S_b^1$, which allows us to obtain

$$\nu D_{U_1 \searrow \Sigma_g} \in \mathbb{Z}, \quad g \geq 1. \quad (17)$$

This result generalizes that of Ref. 11.

To obtain an even stronger result, we consider a surface bundle over surface $\Sigma_g \searrow \Sigma_h$. In Appendix B, we show that, for each $g \geq 5$, there are many surface bundles $\Sigma_g \searrow \Sigma_h$ such that each of them has a 2-cocycle $c_1 \in H^2(\Sigma_g \searrow \Sigma_h; \mathbb{Z})$ satisfying $\int_{\Sigma_g} c_1 = 0$. Let $\int_{\Sigma_g \searrow \Sigma_h} c_1^2 = \eta_c$ and $\int_{\Sigma_g \searrow \Sigma_h} p_1 = 3\eta_p$ where η_c and η_p are integers. We find that the set of allowed (η_p, η_c) is given by (see Appendix B)

$$\{(\eta_p, \eta_c) \mid \eta_p \in 4\mathbb{Z}, \eta_c \in \mathbb{Z}\}. \quad (18)$$

We obtain a stronger quantization condition on ν from eqn. (12):

$$\frac{\nu}{2} D_{U_1 \searrow \Sigma_g} \in \mathbb{Z}, \quad g \geq 5. \quad (19)$$

To summarize

for a 2+1D bosonic gapped phase with U_1 symmetry, its chiral central charge c and filling fraction ν satisfy the following quantization conditions

$$\begin{aligned} \frac{c}{2} D_{U_1 \searrow \Sigma_g} &= 0 \bmod 1, & g \geq 3, \\ \nu D_{U_1 \searrow \Sigma_g} &= 0 \bmod 1, & g \geq 1, \\ \frac{\nu}{2} D_{U_1 \searrow \Sigma_g} &= 0 \bmod 1, & g \geq 5. \end{aligned} \quad (20)$$

where $D_{U_1 \searrow \Sigma_g}$ is the ground state degeneracy on closed space Σ_g .

B. Examples

1. Bosonic K -matrix Abelian topological orders

We check $\frac{c}{2} D_{U_1 \searrow \Sigma_g} \in \mathbb{Z}$ for 2+1D bosonic Abelian topological orders described by the following multi-layer wave function characterized by symmetric integral matrix K with even diagonal:

$$\Psi(z_i^I) = \prod_{I, i < j} (z_i^I - z_j^I)^{K_{II}} \prod_{I < J, i, j} (z_i^I - z_j^J)^{K_{IJ}} e^{-\sum_{I, i} \frac{|z_i^I|^2}{4}}, \quad (21)$$

where z_i^I the complex coordinates of the i^{th} boson in I^{th} -layer. The effective field theory of the above state is described by K -matrix Abelian Chern-Simons theory:

$$S_K = \int \frac{K_{IJ}}{4\pi} a_I da_J + \frac{q_I}{2\pi} A da_I. \quad (22)$$

where the charge-vector q_I are integers, describing the U_1 charge of the bosons in I^{th} -layer. It is believed[38] that

the K -matrix Abelian Chern-Simons theory can realize all the 2+1D Abelian topological orders.

After integrating out the matter field a_I , we obtain the effective theory eqn. (8) with

$$c = \text{sgn}(K) = n_+ - n_-, \quad \nu = \mathbf{q}^\top K^{-1} \mathbf{q}.$$

where n_\pm are the number of eigenvalues of K with sign \pm . The ground state degeneracy is given by

$$D_{U_1 \searrow \Sigma_g} = |\text{Det} K|^g.$$

So eqn. (12) states that the product $\text{sgn}(K)|\text{Det} K|^g$ is an even integer for $g > 2$. To check this, note that $\text{sgn}(K) = n \bmod 2$, where n is the dimension of K . So it suffices to check that

$$\text{Det} K \in 2\mathbb{Z} \quad \text{for odd } n. \quad (23)$$

To see this, write

$$\text{Det} K = \sum_{\sigma} (-1)^{\sigma} \prod_{I=1}^n K_{I\sigma(I)} \quad (24)$$

which is summed over the permutations σ of $\{1, \dots, n\}$ and $(-1)^{\sigma}$ is its sign. Note that any σ which involves a diagonal element of K contributes an even term to the sum, so we only need to consider the permutations where $\sigma(I) \neq I$ for all I . Every permutation σ can be expressed as a number of disjoint cycles. The sum of all the lengths of the cycles is n , which is odd by assumption. So there exists a cycle whose length is odd. Such cycle cannot have length 1, since we assumed $\sigma(I) \neq I$. So it has length ≥ 3 . The existence of such a cycle implies $\sigma \neq \sigma^{-1}$, since reversing the cycle gives a different permutation. Moreover σ and σ^{-1} contributes equally to the sum (24), since

$$\prod_{I=1}^n K_{I\sigma^{-1}(I)} = \prod_{I=1}^n K_{\sigma^{-1}(I)I} = \prod_{I=1}^n K_{I\sigma(I)}.$$

So the two terms together is even. So (24) is even.

The conditions on ν , eqn. (17) and eqn. (19), now become

$$\text{Det}(K) \mathbf{q}^\top K^{-1} \mathbf{q} \in \mathbb{Z}, \quad |\text{Det}(K)|^5 \frac{\mathbf{q}^\top K^{-1} \mathbf{q}}{2} \in \mathbb{Z}, \quad (25)$$

for all integral vectors \mathbf{q} . The first expression is satisfied because $\text{Det}(K)K^{-1}$ is an integer matrix. For the second expression, we only need to consider the case when $\text{Det}(K)$ is odd. By (23) this implies the n , the dimension of K , is even. So the submatrix of K by deleting the i^{th} row and i^{th} column has odd dimension for all i . Applying (23) to this submatrix, we see that it has even determinant. Hence the diagonals of the cofactor matrix of K are even, which means $\text{Det}(K)K^{-1}$ has even diagonals.

From this, the second expression reads

$$\begin{aligned} & |\text{Det}(K)|^5 \frac{q_I K_{IJ}^{-1} q_J}{2} \\ &= \frac{1}{2} |\text{Det}(K)|^5 \left(\sum_I q_I^2 K_{II}^{-1} + \sum_{I \neq J} q_I K_{IJ}^{-1} q_J \right) \\ &= |\text{Det}(K)|^5 \left(\frac{\sum_I q_I^2 K_{II}^{-1}}{2} + \sum_{I < J} q_I K_{IJ}^{-1} q_J \right) \in \mathbb{Z}. \end{aligned}$$

From the above result, we find that, for Abelian bosonic topological orders with U_1 symmetry, the filling fraction is quantized as

$$\begin{aligned} \nu &= \frac{2 \times \text{integer}}{\text{gcd}(2D_1, D_1^5)} \\ &= \begin{cases} \frac{1}{D_1} \times \text{integer}, & \text{if } D_1 = \text{even} \\ \frac{2}{D_1} \times \text{integer}, & \text{if } D_1 = \text{odd} \end{cases} \end{aligned} \quad (26)$$

where $D_1 = |\text{Det}(K)|$ is the ground state degeneracy on torus. In particular, for bosonic U_1 -SPT order, $D_1 = |\text{Det}(K)| = 1$ and $\nu = 0 \bmod 2$. We note that the bosonic $\nu = 1/m$ Laughlin state has $D_1 = m = \text{even}$, and saturates the above quantization condition.

2. Bosonic non-Abelian topological orders described by $SU(2)_k$ Chern-Simons Theory

The following wave function

$$\Psi(z_i) = (\chi_k(z_i))^2 \quad (27)$$

describes a bosonic non-Abelian topological orders of charge-1 bosons, where $\chi_k(z_i)$ is the fermion wave function of k -filled Landau levels.[39] The low energy effective theory is the $SU(2)_k$ Chern-Simons theory. The edge is described by a $U_1^{2k}/SU(2)_k$ WZW theory, whose chiral central charge and filling fraction are[39]

$$c = 2k - \frac{3k}{k+2}, \quad \nu = \frac{k}{2}. \quad (28)$$

Using the $SU(2)_k$ Chern-Simons theory, we find that there are $k+1$ anyons in the bulk labeled by $s = m/2$ where $m = 0, 1, \dots, k$. The ground state degeneracy is given by

$$D_{U_1 \times \Sigma_g} = \sum_m S_{0m}^{-2(g-1)} \in \mathbb{Z}$$

where

$$S_{mn} = \sqrt{\frac{2}{k+2}} \sin\left(\frac{(m+1)(n+1)\pi}{k+2}\right)$$

So eqn. (20) reads

$$\frac{c}{2} D_{U_1 \times \Sigma_g} = \frac{-3k}{2(k+2)} \left(\frac{k+2}{2}\right)^{g-1} \sum_{m=1}^{k+1} \csc^{2(g-1)} \frac{m\pi}{k+2} \in \mathbb{Z} \quad (29)$$

which is verified in Appendix C for all integers $k \geq 0$ and $g \geq 3$. We also have

$$\nu D_{U_1 \times \Sigma_g} = \frac{k}{2} \left(\frac{k+2}{2}\right)^{g-1} \sum_{m=1}^{k+1} \csc^{2(g-1)} \frac{m\pi}{k+2} \in \mathbb{Z} \quad (30)$$

for $g \geq 1$, and

$$\frac{\nu}{2} D_{U_1 \times \Sigma_g} = \frac{k}{4} \left(\frac{k+2}{2}\right)^{g-1} \sum_{m=1}^{k+1} \csc^{2(g-1)} \frac{m\pi}{k+2} \in \mathbb{Z} \quad (31)$$

for $g \geq 5$. We also verify the above two relations in Appendix C.

C. A general point of view

In general, we may consider a boson system in d -dimensional space with an internal symmetry described by group G_b . The bosons may carry non-zero integer spin and may transform non-trivially under the spatial rotation SO_d . We embed SO_d into $SO_\infty \equiv SO$. Thus, bosons transform under the full symmetry group $G_{bSO} = G_b \times SO$. On curved spacetime with G_b -symmetry twist, we have a G_{bSO} connection $a^{G_{bSO}}$ on the spacetime. The connection $a^{G_{bSO}}$ is a special one. When we project G_{bSO} to SO : $G_{bSO} \xrightarrow{\pi} SO$, the G_{bSO} connection is projected into an SO connection: $a^{G_{bSO}} \xrightarrow{\pi} a^{SO}$, and such an SO connection a^{SO} must be the connection for the tangent bundle of the spacetime. The low energy effective Lagrangian L_{eff} (8) is a Chern-Simons term for group G_{bSO} . This point of view will help us to understand the effective Lagrangian for fermion system. Since $G_{bSO} = G_b \times SO$, the low energy effective Lagrangian L_{eff} (8) is a sum of the gravitational Chern-Simons term and a Chern-Simons term for group G_b .

IV. TOPOLOGICAL INVARIANTS IN 2+1D H-TYPE FERMIONIC ENRICHED TOPOLOGICAL ORDERS WITH U_1^f SYMMETRY

A. Symmetry twist of fermion system

We have seen that to probe the topological properties of a bosonic gapped liquid phase with symmetry G_b , we can use the symmetry twist described by a G_b -connection a^{SO} , plus the curved spacetime described by the SO -connection a^{SO} of the tangent bundle. In other words, we can use the G_{bSO} -connection $a^{G_{bSO}}$ and effective action $S_{eff}(a^{G_{bSO}})$ to probe the topological properties. Here, we will discuss the symmetry twist for fermion systems, which turns out to be mixed with spacetime curvature.

For a fermion system in d -dimensional space with an internal symmetry, its symmetry group G_f has a central \mathbb{Z}_2^f subgroup:

$$\text{id} \rightarrow \mathbb{Z}_2^f \rightarrow G_f \rightarrow G_b \rightarrow \text{id}. \quad (32)$$

We also write $G_f = \mathbb{Z}_2^f \rtimes_{e_2} G_b$, where e_2 is a group 2-cocycle

$$e_2 \in H^2(G_b; \mathbb{Z}_2^f) \quad (33)$$

describing the group extension (32) (see Appendix D). The fermion carry half-integer spins and transform non-trivially under the spatial rotation SO_d , which is enlarged to SO . Thus, fermions transform under the full symmetry group $G_{fSO} = G_f \rtimes SO$. On curved spacetime with G_f -symmetry twist, we have a G_{fSO} connection $a^{G_{fSO}}$ on the spacetime. The connection $a^{G_{fSO}}$ is a special one. When we project G_{fSO} to SO : $G_{fSO} \xrightarrow{\pi} SO$, the G_{fSO} connection is projected into an SO connection: $a^{G_{fSO}} \xrightarrow{\pi} a^{SO}$, and such an SO connection a^{SO} must be the connection for the tangent bundle of the spacetime. The low energy effective Lagrangian L_{eff} is a Chern–Simons term for group G_{fSO} . Because $G_{fSO} = G_f \rtimes SO \neq G_f \times SO$, the low energy effective Lagrangian L_{eff} (8) in general cannot be written as a sum of the gravitational Chern–Simons term for group SO and a Chern–Simons term for group G_f .

There is another way to describe full fermionic symmetry group $G_{fSO} = G_f \rtimes SO$:

$$G_{fSO} = (G_f \times Spin)/\mathbb{Z}_2^f \quad (34)$$

We note that the spin group $Spin \equiv Spin_\infty = \mathbb{Z}_2^f \rtimes_{w_2} SO$ contains a central subgroup \mathbb{Z}_2^f such that $Spin/\mathbb{Z}_2^f = SO$, where $w_2 \in H^2(SO; \mathbb{Z}_2^f)$ is the group 2-cocycle describing the group extension

$$\text{id} \rightarrow \mathbb{Z}_2^f \rightarrow Spin \rightarrow SO \rightarrow \text{id}. \quad (35)$$

$G_f = \mathbb{Z}_2^f \rtimes_{e_2} G_b$ also contains a central subgroup \mathbb{Z}_2^f such that $G_f/\mathbb{Z}_2^f = G_b$. G_{fSO} is obtained by identify these two \mathbb{Z}_2^f subgroups in $G_f \times Spin$. Let us described the G_{fSO} -connection using this point of view.

We first triangulate the spacetime, where the vertices are labeled by i, j, k, \dots , the links labelled by ij, jk, \dots , etc. A G_f -connection on the triangulation is described by the group elements of G_f on the links: $a_{ij}^{G_f} \in G_f$. We can view $G_f = \mathbb{Z}_2^f \rtimes_{e_2} G_b$ and use the pair $(a_{ij}^{G_b}, a_{ij}^{\mathbb{Z}_2^f})$ to label the group elements of G_f , where $a_{ij}^{G_b} \in G_b$ and $a_{ij}^{\mathbb{Z}_2^f} \in \mathbb{Z}_2^f$. Similarly, a $Spin$ -connection on the triangulation is described by the group elements of $Spin$ on the links: $a_{ij}^{Spin} \in Spin$. We can view $Spin = \mathbb{Z}_2^f \rtimes_{w_2} SO$ and use the pair $(a_{ij}^{SO}, \tilde{a}_{ij}^{\mathbb{Z}_2^f})$ to label the group elements of $Spin$, where $a_{ij}^{SO} \in SO$ and $\tilde{a}_{ij}^{\mathbb{Z}_2^f} \in \mathbb{Z}_2^f$. Now, the G_{fSO} -connection is obtained by requiring $\tilde{a}_{ij}^{\mathbb{Z}_2^f} = a_{ij}^{\mathbb{Z}_2^f}$. Also the SO -connection a_{ij}^{SO} describes the tangent bundle of the spacetime.

Now, let us describe a nearly flat G_{fSO} -connection that describe the curved spacetime and twisted symme-

try. The nearly flat condition is given by

$$a_{ij}^{G_{fSO}} a_{jk}^{G_{fSO}} \approx a_{ik}^{G_{fSO}} \quad (36)$$

for all the triangles ijk . It implies that the G_f -connection is nearly flat

$$(a_{ij}^{G_b}, a_{ij}^{\mathbb{Z}_2^f}) \circ (a_{jk}^{G_b}, a_{jk}^{\mathbb{Z}_2^f}) \approx (a_{ik}^{G_b}, a_{ik}^{\mathbb{Z}_2^f}), \quad (37)$$

and the $Spin$ -connection is nearly flat

$$(a_{ij}^{SO}, a_{ij}^{\mathbb{Z}_2^f}) \circ (a_{jk}^{SO}, a_{jk}^{\mathbb{Z}_2^f}) \approx (a_{ik}^{SO}, a_{ik}^{\mathbb{Z}_2^f}). \quad (38)$$

In other words (see Appendix D)

$$\begin{aligned} a_{ij}^{G_b} a_{jk}^{G_b} &\approx a_{ik}^{G_b}, \\ a_{ij}^{SO} a_{jk}^{SO} &\approx a_{ik}^{SO}, \end{aligned} \quad (39)$$

and

$$\begin{aligned} a_{ij}^{\mathbb{Z}_2^f} + a_{jk}^{\mathbb{Z}_2^f} - a_{ik}^{\mathbb{Z}_2^f} &\stackrel{2}{=} e_2(a_{ij}^{G_b}, a_{jk}^{G_b}) = (e_2)_{ijk}, \\ a_{ij}^{\mathbb{Z}_2^f} + a_{jk}^{\mathbb{Z}_2^f} - a_{ik}^{\mathbb{Z}_2^f} &\stackrel{2}{=} w_2(a_{ij}^{SO}, a_{jk}^{SO}) = (w_2)_{ijk}. \end{aligned} \quad (40)$$

The above can be rewritten as

$$da^{\mathbb{Z}_2^f} \stackrel{2}{=} e_2(a^{G_b}), \quad da^{\mathbb{Z}_2^f} \stackrel{2}{=} w_2(a^{SO}). \quad (41)$$

Since a^{SO} is the connection of the tangent bundle of the spacetime, $w_2(a^{SO}) = w_2$ is the second Stiefel–Whitney class of the tangent bundle of the spacetime. We see that the G_b -connection and the SO -connection on spacetime are not arbitrary. They must satisfy the constraint

$$e_2(a_{ij}^{G_b}, a_{jk}^{G_b}) \stackrel{2}{=} w_2(a_{ij}^{SO}, a_{jk}^{SO}) = w_2. \quad (42)$$

In other words, the nearly flat G_b bundle on the spacetime (describing the symmetry twist) is not arbitrary. Its nearly flat connection a^{G_b} must satisfy

$$e_2(a^{G_b}) \stackrel{2, d}{=} w_2. \quad (43)$$

B. U_1^f symmetry and spin^c structure

Let us consider an example of the above result. A fermion system with fermion number conservation has a U_1^f symmetry. Assume all the odd charges of U_1^f are fermionic, then U_1^f has a \mathbb{Z}_2^f subgroup generated by the π -rotation, which corresponds to the fermion-number-parity. It is more convenient to view U_1^f as a group extension of U_1 by \mathbb{Z}_2^f , i.e. $U_1^f = \mathbb{Z}_2^f \rtimes_{e_2} U_1$ where e_2 is the \mathbb{Z}_2 valued group 2-cocycle that generates $H^2(U_1, \mathbb{Z}_2)$ (see Appendix D).

Using \mathbb{R}/\mathbb{Z} -valued $a^{(\mathbb{R}/\mathbb{Z})^f}$ to label group elements in U_1^f , \mathbb{R}/\mathbb{Z} -valued $a^{\mathbb{R}/\mathbb{Z}}$ to label group elements in U_1 , and \mathbb{Z}_2 -valued $a^{\mathbb{Z}_2}$ to label group elements in \mathbb{Z}_2^f , we find that

$$a^{(\mathbb{R}/\mathbb{Z})^f} = \frac{1}{2} a^{\mathbb{R}/\mathbb{Z}} + \frac{1}{2} a^{\mathbb{Z}_2^f}, \quad (44)$$

Here $\mathbb{R}/\mathbb{Z} = [0, 1)$ and $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$. Now the group multiplication in U_1^f can be rewritten as

$$\begin{aligned} a_1^{(\mathbb{R}/\mathbb{Z})^f} + a_2^{(\mathbb{R}/\mathbb{Z})^f} &= \frac{1}{2}a_1^{\mathbb{R}/\mathbb{Z}} + \frac{1}{2}a_2^{\mathbb{R}/\mathbb{Z}} + \frac{1}{2}a_1^{\mathbb{Z}_2^f} + \frac{1}{2}a_2^{\mathbb{Z}_2^f} \\ &= \frac{1}{2}(a_1^{\mathbb{R}/\mathbb{Z}} + a_2^{\mathbb{R}/\mathbb{Z}} - \lfloor a_1^{\mathbb{R}/\mathbb{Z}} + a_2^{\mathbb{R}/\mathbb{Z}} \rfloor) \\ &\quad + (\frac{1}{2}a_1^{\mathbb{Z}_2^f} + \frac{1}{2}a_2^{\mathbb{Z}_2^f} + \lfloor a_1^{\mathbb{R}/\mathbb{Z}} + a_2^{\mathbb{R}/\mathbb{Z}} \rfloor) \\ &= \frac{1}{2}(a_1^{\mathbb{R}/\mathbb{Z}} + a_2^{\mathbb{R}/\mathbb{Z}} - \lfloor a_1^{\mathbb{R}/\mathbb{Z}} + a_2^{\mathbb{R}/\mathbb{Z}} \rfloor) \\ &\quad + (\frac{1}{2}a_1^{\mathbb{Z}_2^f} + \frac{1}{2}a_2^{\mathbb{Z}_2^f} + e_2(a_1^{\mathbb{R}/\mathbb{Z}}, a_2^{\mathbb{R}/\mathbb{Z}})), \end{aligned} \quad (45)$$

where $\lfloor x \rfloor$ denotes the largest integer smaller than or equal to x , and

$$e_2(a_1^{\mathbb{R}/\mathbb{Z}}, a_2^{\mathbb{R}/\mathbb{Z}}) = \lfloor a_1^{\mathbb{R}/\mathbb{Z}} + a_2^{\mathbb{R}/\mathbb{Z}} \rfloor \quad (46)$$

which is a \mathbb{Z}_2 -valued group 2-cocycle $e_2 \in H^2(U_1; \mathbb{Z}_2)$ that characterizes the group extension of U_1 by \mathbb{Z}_2^f :

$$U_1^f = \mathbb{Z}_2^f \rtimes_{e_2} U_1. \quad (47)$$

Note that $e_2(a_1^{\mathbb{R}/\mathbb{Z}}, a_2^{\mathbb{R}/\mathbb{Z}})$ is a smooth function of $a_1^{\mathbb{R}/\mathbb{Z}}, a_2^{\mathbb{R}/\mathbb{Z}}$ when $0 < a_1^{\mathbb{R}/\mathbb{Z}} < \frac{1}{2}$ and $0 < a_2^{\mathbb{R}/\mathbb{Z}} < \frac{1}{2}$. But it has discontinuities in other places. It turns out that

$$e_2 \stackrel{2, d}{=} c_1^{U_1} \quad (48)$$

where $c_1^{U_1} \in H^2(U_1, \mathbb{Z})$ is the first Chern class of U_1 viewed as group 2-cocycle.

The ‘‘Chern Class’’ for U_1^f and the Chern Class for U_1 are related (see eqn. (44))

$$c_1^{U_1^f} = \frac{1}{2}c_1^{U_1}, \quad (49)$$

where $c_1^{U_1} = da^{\mathbb{R}/\mathbb{Z}}$ and $c_1^{U_1^f} = da^{(\mathbb{R}/\mathbb{Z})^f}$. We like to point out that the U_1^f -connection $a^{(\mathbb{R}/\mathbb{Z})^f}$ corresponds to the electromagnetic gauge potential plus the connection of the curved space:

$$A_{eff} = 2\pi a^{(\mathbb{R}/\mathbb{Z})^f}, \quad (50)$$

and the Chern class $c_1^{U_1^f}$ corresponds to the field strength of the effective U_1^f -connection

$$F_{eff} = dA_{eff} = 2\pi c_1^{U_1^f}. \quad (51)$$

We find that for fermion system with U_1^f symmetry, the Chern number on a closed 2-dimension space C^2 satisfies

$$\int_{C^2} c_1^{U_1^f} = \frac{1}{2\pi} \int_{C^2} F_{eff} \stackrel{1}{=} \frac{1}{2} \int_{C^2} w_2. \quad (52)$$

In other words, on a spin manifold $w_2 \stackrel{2, d}{=} 0$, the Chern number $\frac{1}{2\pi} \int_{C^2} F_{eff}$ is quantized as integers. On non-spin

manifold, the fermion system with U_1^f symmetry can still be defined, provided that the Chern number $\frac{1}{2\pi} \int_{C^2} F_{eff}$ is quantized as half-integers when $\int_{C^2} w_2 \stackrel{2}{=} 1$.

We see that for fermion systems, there is a constraint on the U_1^f connection $a^{U_1^f} = \frac{1}{2\pi} A_{eff}$ and the curved spacetime. When $A_{eff} = 0$, this constraint requires that $w_2 \stackrel{2, d}{=} 0$, *i.e.* the spacetime to be spin. This leads to the general impression that the appearance of fermions requires the spacetime manifold to be spin. Here we see that the appearance of U_1^f fermions requires the spacetime manifold to be spin only when there is no background effective U_1^f gauge field. In the presence of effective background U_1^f gauge field, the spacetime may not be spin. This result can be summarized more precisely by the following statement: the appearance of U_1^f fermions requires the spacetime manifold to be spin^C , and the U_1^f connection $a^{(\mathbb{R}/\mathbb{Z})^f}$ is a spin^C structure.

We know that an orientable manifold is spin if and only if its $w_2 \stackrel{2, d}{=} 0$. Similarly, an orientable manifold is spin^C if and only if $\beta_2 w_2 \stackrel{d}{=} 0$. Here $\beta_2 = \frac{1}{2}d$ is the Bockstein homomorphism. In order words, an orientable manifold M is spin^C if and only if its w_2 is the mod 2 reduction of a \mathbb{Z} -valued 2-cocycle in $H^2(M; \mathbb{Z})$. Also a complex vector bundle of rank n can be viewed as a real vector bundle of rank $2n$. The n^{th} Chern class c_n^U of the complex vector bundle and the $2n^{\text{th}}$ Stiefel–Whitney class are related

$$w_{2n} \stackrel{2, d}{=} c_n^U. \quad (53)$$

This implies that a complex manifold is always spin^C .

C. Fermionic gapped liquids with U_1^f symmetry in 2-dimensional space

Now, we consider a gapped liquids with U_1^f symmetry in 2-dimensional space for fermionic systems. The effective action may contain a gravitational Chern–Simons term and a U_1^f Chern–Simons term:

$$\mathcal{L}_{eff} dx = -2\pi \frac{c}{24} \omega_3 + \frac{\nu}{4\pi} A_{eff} dA_{eff} \quad (54)$$

where A_{eff} is the U_1^f connection 1-form describing the total electromagnetic field, plus the connection from the curved space due to the orbital spin of the fermions. Follow the same discussion for the bosonic case, we find that the quantization conditions of c and ν for fermionic case can be obtained from (see eqn. (12))

$$D_{U_1 \rtimes \Sigma_g} \int_{\Sigma_g \times B^2} -\frac{c}{24} p_1 + \frac{\nu}{8} (c_1^{U_1})^2 \in \mathbb{Z}, \quad (55)$$

for any orientable surface bundle $\Sigma_g \times B^2$ and for any U_1 bundle whose first Chern class $c_1^{U_1}$ satisfies $\frac{1}{2}\nu \int_{\Sigma_g} c_1^{U_1} \in \mathbb{Z}$ and $c_1^{U_1} \stackrel{2, d}{=} w_2$.

Here $c_1^{U_1} = da^{\mathbb{R}/\mathbb{Z}}$, $a^{\mathbb{R}/\mathbb{Z}} = 2\frac{A_{eff}}{2\pi}$ is the U_1 -connection for charge-2 bosons (fermion pairs), and w_2 is the second Stiefel–Whitney class of the tangent bundle of $\Sigma_g \times B^2$.

Next, we choose some special surface bundles $\Sigma_g \times B^2$ and the allowed U_1 bundles to obtain quantization conditions of ν and c . Let us choose the surface bundle to be $\Sigma_g \times B^2 = S_x^1 \times S_y^1 \times S^1 \times S^1$, where $S_x^1 \times S_y^1$ is the space and $S^1 \times S^1 = B^2$. The second Stiefel–Whitney class of such a surface bundle is trivial $w_2 \stackrel{2,d}{=} 0$. Thus the allowed U_1 bundles satisfy $\int_{C^2} c_1^{U_1} = \text{even integers}$. We consider the following form $\text{pf } c_1^{U_1} = c_1^x + c_1^y$, where c_1^x lives on $S_x^1 \times S^1$ and c_1^y lives on $S_y^1 \times S^1$. Since $\int_{\Sigma_0 \times B^2} p_1 = 0$ and $\int_{S_x^1 \times S_y^1} c_1^x + c_1^y = 0$, the quantization condition eqn. (55) becomes

$$\begin{aligned} & \frac{\nu}{8} D_{U_1 \times \Sigma_1} \int_{S_x^1 \times S_y^1 \times S^1 \times S^1} (c_1^x + c_1^y)^2 \\ &= \frac{\nu}{4} D_{U_1 \times \Sigma_1} \int_{S_x^1 \times S^1} c_1^x \int_{S_y^1 \times S^1} c_1^y. \end{aligned} \quad (56)$$

We can choose the U_1 bundle such that $\int_{S_x^1 \times S^1} c_1^x = 0, \pm 2$ and $\int_{S_y^1 \times S^1} c_1^y = 0, \pm 2$. We obtain the following quantization condition

$$\nu D_{U_1 \times \Sigma_1} \in \mathbb{Z}. \quad (57)$$

The above result was first obtained by Ref. 11. Again, because Σ_g , $g \geq 1$, is the connected sum of g Σ_1 's, the above reasoning can be generalized to the case $\Sigma_g \times B^2 = \Sigma_g \times S^1 \times S^1$ whose second Stiefel–Whitney class is still trivial $w_2 \stackrel{2,d}{=} 0$, which allow us to obtain

$$\nu D_{U_1 \times \Sigma_g} \in \mathbb{Z}, \quad g \geq 1. \quad (58)$$

If $\Sigma_g \times B^2$ is spin, then $\int_{\Sigma_g \times B^2} p_1 = 0 \bmod 48$ by (B4). We explain in Appendix B that as long as $g \geq 9$ there is such a spin surface bundle with $\int_{\Sigma_g \times B^2} p_1 = \pm 48$, in which case we may choose the U_1 -bundle to be trivial. This gives us a quantization of c (see Ref. 30):

$$2c D_{U_1 \times \Sigma_g} \in \mathbb{Z}, \quad \text{for } g \geq 9. \quad (59)$$

To obtain additional quantization condition, we consider a more general surface bundle over surface $\Sigma_g \times B^2$. In Appendix B, we show that, for each $g \geq 5$, there are many surface bundles $\Sigma_g \times B^2$ equipped with a 2-cocycle $c_1^{U_1} \in H^2(\Sigma_g \times B^2; \mathbb{Z})$ satisfying (1) $\int_{\Sigma_g} c_1^{U_1} = 0$ and (2) $c_1^{U_1} \stackrel{2,d}{=} w_2$. Let $\int_{\Sigma_g \times B^2} (c_1^{U_1})^2 = \eta_c$ and $\int_{\Sigma_g \times B^2} p_1 = 3\eta_p$ where η_c and η_p are integers. Eqn. (55) becomes

$$D_{U_1 \times \Sigma_g} (-\eta_p \frac{c}{8} + \eta_c \frac{\nu}{8}) \in \mathbb{Z}, \quad (60)$$

In Appendix B, we find that the set of allowed (η_p, η_c) is exactly given by the integer pairs that satisfy

$$\eta_c \stackrel{8}{=} \eta_p \stackrel{4}{=} 0. \quad (61)$$

Now eqn. (60) implies that $\nu D_{U_1 \times \Sigma_g} \in \mathbb{Z}$ and two new quantization conditions

$$\begin{aligned} & c D_{U_1 \times \Sigma_g} \in \mathbb{Z}, \quad g \geq 5, \\ & (-\frac{c}{2} + \frac{\nu}{2}) D_{U_1 \times \Sigma_g} \in \mathbb{Z}, \quad g \geq 5. \end{aligned} \quad (62)$$

To summarize

for a 2+1D fermionic gapped phase with U_1^f symmetry, its chiral central charge c and filling fraction ν satisfy the following quantization conditions

$$\begin{aligned} & c D_{U_1 \times \Sigma_g} = 0 \bmod 1, \quad g \geq 5, \\ & \nu D_{U_1 \times \Sigma_g} = 0 \bmod 1, \quad g \geq 1, \\ & (-c + \nu) D_{U_1 \times \Sigma_g} = 0 \bmod 2, \quad g \geq 5, \end{aligned} \quad (63)$$

where $D_{U_1 \times \Sigma_g}$ is the ground state degeneracy on closed space Σ_g .

D. H-type invertible fermionic U_1^f -enriched topological orders

Let us discuss a simple example. The invertible fermionic U_1^f -enriched topological order are classified by K -matrix (see eqn. (22))

$$K = (1)^{\oplus m} \oplus (-1)^{\oplus n} \quad (64)$$

and charge vector \mathbf{q} with odd-integer components. Its c and ν are given by

$$c = \text{Tr}(K), \quad \nu = \sum_{i=1}^m q_i^2 - \sum_{i=m+1}^{m+n} q_i^2. \quad (65)$$

The ground state degeneracy is always $D_{U_1 \times \Sigma_g} = |\text{Det}(K)| = 1$. We see that $-c + \nu = 0 \bmod 8$, which satisfies (but does not saturate) eqn. (63).

V. THE TOPOLOGICAL INVARIANTS FOR L-TYPE TOPOLOGICAL ORDERS

A. Topological partition function for L-type topological order

Consider a bosonic system in gapped liquid phase described by a path integral on n D spacetime M^n . After integrating out all the dynamical degrees of freedom, we will obtain a partition function that has the following form

$$Z(M^n) = e^{-\int_{M^n} d^n x \varepsilon(x)} Z^{\text{top}}(M^n) \quad (66)$$

where $\varepsilon(x)$ is the energy density of the ground state. The term $Z^{\text{top}}(M^n)$ is called the topological partition function and is a topological invariant of the 2+1D topological order. To be more precise, $Z^{\text{top}}(M^n)$ is a complex

function on \mathcal{M}_{M^n} , which is the moduli space of the spacetime M^n . *i.e.* \mathcal{M}_{M^n} is a space of metrics $g_{\mu\nu}$ of a closed manifold. The $g_{\mu\nu}$'s differ by diffeomorphisms are equivalent and represent the same point of the moduli space \mathcal{M}_{M^n} .

We like to point out that for certain spacetime topologies, the spacetime M^n must contain world-line of point-like excitations, and/or world-sheet of string-like excitations, *etc.* For those certain spacetime topologies $Z^{\text{top}}(M^n) = 0$ on the corresponding moduli space \mathcal{M}_{M^n} . For other spacetime topologies, the spacetime M^n does not contain any excitations. In this case $Z^{\text{top}}(M^n)$ is always non-zero on the corresponding moduli space \mathcal{M}_{M^n} . If $Z^{\text{top}}(M^n)$ was zero at an isolated point (or isolated lower dimensional subspace of \mathcal{M}_{M^n}), then a small perturbation will make $Z^{\text{top}}(M^n)$ non-zero. This causes the diverging change in the effective action $S_{\text{eff}} = \ln(Z^{\text{top}}(M^n))$, which indicates that the system to be gapless. Thus[12]

on the moduli space of gapped systems, $Z^{\text{top}}(M^n)$ is either always zero (when the spacetime must contain excitations), or always non-zero (when the spacetime is filled by gapped ground state).

We also like to conjecture that

on the moduli space of gapped systems, $|Z^{\text{top}}(M^n)|$ is a constant.

Physically, since the system is gapped, the $Z^{\text{top}}(M^n)$ should not depend on the size (*i.e.* the metrics) of the spacetime, we naively expect $Z^{\text{top}}(M^n)$ to be constant. However, the such a naively expectation is not totally correct. We will see that the phase of $Z^{\text{top}}(M^n)$ can depend on the metrics. So we conjecture that $|Z^{\text{top}}(M^n)|$ is a constant.

The metrics dependence of the phase of $Z^{\text{top}}(M^n)$ has a topological reason: they come from the gravitational Chern–Simons terms $\omega_n(a^{SO})$ that depend on the SO connection a^{SO} of the tangent bundle

$$Z^{\text{top}}(M^n) = |Z^{\text{top}}(M^n)| e^{2\pi i \int_{M^n} \kappa \omega_n(a^{SO})}. \quad (67)$$

The gravitational Chern–Simons terms exist only in spacetime dimensions $n \stackrel{4}{=} 3$.

When the tangent bundle of the spacetime is non-trivial, the SO connection a^{SO} of the tangent bundle is not globally defined, and as the result the gravitational Chern–Simons action $\int_{M^n} \kappa \omega_n(a^{SO})$ is not well defined. In this paper, we use a cobordism approach trying to define the difference of two gravitational Chern–Simons actions as

$$\begin{aligned} & \int_{M_1^n} \kappa \omega_n(a^{SO}) - \int_{M_2^n} \kappa \omega_n(a^{SO}) \\ & \equiv \int_{N^{n+1}} \kappa d\omega_n(a^{SO}) = \int_{N^{n+1}} \kappa p(a^{SO}) \end{aligned} \quad (68)$$

where M_1^n and M_2^n are the boundary of N^{n+1} : $\partial N^{n+1} = M_1^n \sqcup -M_2^n$, and $p(a^{SO}) = d\omega_n(a^{SO})$ is a product of Pontryagin classes. Note that a^{SO} in $p(a^{SO})$ is the SO -connection of the tangent bundle for N^{n+1} . Since N^{n+1} satisfying $\partial N^{n+1} = M_1^n \sqcup -M_2^n$ is not unique, the different choices of N^{n+1} may give different $\int_{M_1^n} \kappa \omega_n(a^{SO}) - \int_{M_2^n} \kappa \omega_n(a^{SO})$. However, if the ambiguity is an integer, then $\frac{e^{2\pi i \int_{M_1^n} \kappa \omega_n(a^{SO})}}{e^{2\pi i \int_{M_2^n} \kappa \omega_n(a^{SO})}}$ is well defined. This leads to a quantization of κ :

$$\int_{N^{n+1}} \kappa p(a^{SO}) \in \mathbb{Z}, \quad \partial N^{n+1} = \emptyset. \quad (69)$$

So according to the cobordism approach, the phase of the partition function $e^{2\pi i \int_{M^n} \kappa \omega_n(a^{SO})}$ can be well defined only when κ satisfy a quantization condition eqn. (69).

B. The quantization of c for 2+1D bosonic topological orders with non-zero partition function

Let us apply the above result for 2+1D L-type bosonic topological orders. Here we make a crucial assumption that the partition function is non-zero for all closed *orientable* spacetime manifolds

$$Z^{\text{top}}(M^3) = |Z^{\text{top}}(M^3)| e^{i \frac{2\pi c}{24} \int_{M^3} \omega_3} \neq 0. \quad (70)$$

Not all 2+1D L-type bosonic topological orders satisfy this condition. But at least, the 2+1D L-type bosonic invertible topological orders[12, 40, 41] are believed to satisfy this condition.

For 2+1D L-type bosonic topological orders satisfying this condition, their chiral central charges satisfy

$$\frac{c}{24} \int_{N^4} p_1 \in \mathbb{Z}, \quad \forall \partial N^4 = \emptyset. \quad (71)$$

Since the partition function is non-zero on all closed *orientable* 3-manifold, here we choose N^4 to an arbitrary closed *orientable* 4-manifold. This is a key assumption in our cobordism approach.

We know that for all different closed *orientable* 4-manifolds N^4 , the set $\{\int_{N^4} p_1\}$ is given by $3\mathbb{Z}$. We find that for those L-type bosonic topological orders, their chiral central charges are quantized as

$$\frac{c}{8} \in \mathbb{Z}. \quad (72)$$

We conclude that, at least for the 2+1D L-type bosonic invertible topological orders, their chiral central charges satisfy $c = 0 \bmod 8$.

C. The quantization of c for 2+1D bosonic topological orders with vanishing partition functions only on non-spin manifolds and for 2+1D fermionic invertible topological orders

It was pointed out that a bosonic topological order with emergent fermions must have vanishing topological

partition functions on non-spin manifolds. Here we consider a subclass of 2+1D bosonic topological order with emergent fermions, such that topological partition functions vanishes *only* on non-spin manifolds. The topological partition functions are non-zero on all closed *orientable spin* manifolds. We believe that the partition functions for fermionic invertible topological order satisfy this condition. Also the bosonic topological order obtained by gauging the Z_2^f symmetry in fermionic invertible topological order satisfy this condition. In this case, the quantization condition becomes

$$\frac{c}{24} \int_{N^4} p_1 \in \mathbb{Z}, \quad \forall \partial N^4 = \emptyset, \quad w_2 = 0. \quad (73)$$

i.e. we also require N^4 to be *orientable* and *spin*. As explained in Appendix A, the set of integers which may be realized as $\int_N p_1$ for closed spin 4-manifolds N^4 is $48\mathbb{Z}$. We find that for those simple L-type bosonic topological orders with emergent fermions, their chiral central charges are quantized as

$$2c \in \mathbb{Z}. \quad (74)$$

We conclude that, at least for the 2+1D L-type fermionic invertible topological orders, their chiral central charges satisfy $c = 0 \bmod \frac{1}{2}$.

D. Framing anomaly

We have mentioned that the partition function is a function on the moduli space \mathcal{M}_{M^3} of different closed spacetime manifolds M^3 . In fact, in the presence of gravitational Chern–Simons term, such a statement is incorrect. The partition function is actually a function on the moduli space $\mathcal{M}_{M^3}^F$ of different closed spacetime manifolds M^3 *with framing structure*. In other words, the partition functions for the same manifold but with different framing structures may have different values.

So, what is a framing structure? Give a n -manifold M^n and its tangent bundle TM , its stabilized tangent bundle is given by $TM \oplus \mathbb{R}^\infty$. The manifold M^n is *framable* if the stabilized tangent bundle is trivial. A trivialization of the stabilized tangent bundle (*i.e.* a choice of a global basis of the stabilized tangent bundle) is a framing structure.

It turns out that all orientable 3-manifold M^3 is framable. This gives us a way to define the gravitational Chern–Simons term. If the tangent bundle TM of M^3 is non-trivial, the SO_3 connection a^{SO_3} for the tangent bundle TM cannot be globally defined on M^3 , which makes the gravitational Chern–Simons term $\int_{M^3} \omega_3(a^{SO_3})$ not well defined. But if we embed SO_3 into $SO_\infty \equiv SO$, then the corresponding a^{SO} connection can be globally defined on M^3 (since the stabilized tangent bundle is trivial), which makes the gravitational Chern–Simons term $\int_{M^3} \omega_3(a^{SO})$ well defined. But there are different ways to turn a a^{SO_3} -connection into a globally defined SO -connection a^{SO} , which corresponds to different trivializations (*i.e.* different choices of framing structures). Such

different choices of framing structures can change the gravitational Chern–Simons term $\int_{M^3} \omega_3(a^{SO})$ by an arbitrary integer. We see that if $c \neq 0 \bmod 24$, then the phase of partition function $e^{i \frac{2\pi c}{24} \int_{M^3} \omega_3(a^{SO})}$ will depend on the framing structures and the partition function is not a function on the moduli space \mathcal{M}_{M^3} of 3-manifolds. This phenomenon is called framing anomaly. In this case, the partition function is a function on the moduli space $\mathcal{M}_{M^3}^F$ of 3-manifolds with framing structures.

In general, H-type topological orders have $c \neq 0 \bmod 24$, and thus correspond to L-type topological orders with framing anomaly. The L-type topological orders without framing anomaly must have $c \equiv 0$. In this paper, we will mainly discuss L-type topological orders with framing anomaly. Their partition function is a function on the moduli space $\mathcal{M}_{M^3}^F$ of 3-manifolds with framing structures.

VI. THE TOPOLOGICAL INVARIANTS FOR L-TYPE TOPOLOGICAL ORDERS WITH U_1 SYMMETRY

A. Quantization of c and ν for bosonic U_1 -SET orders with non-zero partition functions

A L-type bosonic U_1 -enriched topological orders may contain two Chern–Simons terms given by eqn. (8). Here we assume that the partition function is non-zero for all closed *orientable* spacetime with any U_1 -bundle on it. Repeating the arguments in the last section, but for spacetime with a U_1 connection $a^{\mathbb{R}/\mathbb{Z}} = \frac{A_{eff}}{2\pi}$, we obtain the following quantization for c and ν , for bosonic U_1 -enriched topological orders whose topological partition functions are non-zero on any closed orientable spacetime:

$$\int_{N^4} -\frac{c}{24} p_1 + \frac{\nu}{2} c_1^2 \in \mathbb{Z}, \quad \forall \partial N^4 = \emptyset \text{ and } \forall U_1 \text{ bundles.} \quad (75)$$

Here N^4 is an arbitrary *orientable* 4-manifold with an arbitrary U_1 -bundle on it. Let $3\eta_p = \int_{N^4} p_1$ and $\eta_c = \int_{N^4} c_1^2$, where η_p and η_c are integers. In Appendix A we show that the pairs (η_p, η_c) which may be realized is given by (\mathbb{Z}, \mathbb{Z}) . This leads to the quantization of c and ν :

$$c = 0 \bmod 8, \quad \nu = 0 \bmod 2. \quad (76)$$

The above result, at least, applies to L-type invertible bosonic U_1 -enriched topological orders and L-type bosonic U_1 -SPT orders.

B. Quantization of c and ν for fermionic U_1^f -enriched topological orders with non-zero partition functions

A L-type fermionic U_1^f -enriched topological orders also contain two Chern–Simons terms given by eqn. (8). For fermionic U_1 -SET orders whose topological partition functions are non-zero on any closed smooth *orientable* spin^C spacetime manifolds, the quantization for c and ν is given by

$$\int_{N^4} -\frac{c}{24}p_1 + \frac{\nu}{8}(c_1^{U_1})^2 \in \mathbb{Z}, \quad \forall \partial N^4 = \emptyset \text{ and } \forall c_1^{U_1} \stackrel{2, d}{=} w_2. \quad (77)$$

Here N^4 is an arbitrary closed smooth *orientable* spin^C 4-manifold, as implied by the condition $c_1^{U_1} \stackrel{2, d}{=} w_2$. Let $3\eta_p = \int_{N^4} p_1$ and $\eta_c = \int_{N^4} (c_1^{U_1})^2$, where η_p and η_c are integers. As explained in Appendix A, we may find such spin^C 4-manifolds realizing any integers η_p and η_c as long as they satisfy

$$\eta_p \stackrel{s}{=} \eta_c. \quad (78)$$

This leads to the quantization of c and ν :

$$\begin{aligned} c &= 0 \bmod 1, & \nu &= 0 \bmod 1, \\ -c + \nu &= 0 \bmod 8. \end{aligned} \quad (79)$$

The above result, at least, applies to L-type invertible fermionic U_1^f -enriched topological orders and L-type fermionic U_1^f -SPT orders.

For L-type fermionic U_1^f -SPT orders, the central charge vanishes $c = 0$ and $\nu = 0 \bmod 8$. This agrees with the result in Ref. 42. We also note that the fermionic invertible U_1^f -SET states discussed in Sec. IV D satisfy the above quantization condition, and actually saturate the quantization condition.

We remark that the combination of invertible fermionic U_1^f -enriched topological orders and fermionic U_1^f -SPT orders (*i.e.* the invertible fermionic gapped liquid phases with U_1^f symmetry) is classified via spin^C cobordism in Ref. 43, which is given by \mathbb{Z}^2 in 2+1D. This agrees with our result that those state are labeled by two integers $\{(\eta_p, \eta_c) \mid \eta_p \stackrel{s}{=} \eta_c\}$. However, the values of c and ν are not discussed in Ref. 43.

VII. SUMMARY

For 2+1D gapped liquid phases with U_1 symmetry, the central charge c and the dimensionless Hall conductance ν are described by the Chern–Simons terms in the partition function. However, for arbitrary smooth spacetime 3-manifolds and for arbitrary U_1 -bundle over spacetime 3-manifolds, it is highly non-trivial to define the Chern–Simons terms. In this paper, we use a cobordism approach to define the Chern–Simons terms. We find that

for H-type quantum systems, the 4-manifolds used in the cobordism approach must be a surface bundle. For L-type quantum systems, the 4-manifolds must have the same type as that of the spacetime 3-manifolds, such that the partition functions are non-zero. This leads to different quantization conditions for c and ν . In particular, for the H-type quantum systems, the quantization of c and ν depends on the ground state degeneracies on Riemannian surfaces. While for the L-type quantum systems, the quantization of c and ν depends on the type of spacetime manifolds where the topological partition function is non-zero.

ORW is supported by the ERC under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 756444), and by a Philip Leverhulme Prize from the Leverhulme Trust. LT is supported by the Croucher Fellowship for Postdoctoral Research. XGW is partially supported by NSF DMS-1664412 and by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651440).

Appendix A: Characteristic numbers of 4-manifolds

The most fundamental invariant of a closed oriented 4-manifold M is its intersection form

$$\begin{aligned} Q_M : H^2(M^4; \mathbb{Z}) \times H^2(M^4; \mathbb{Z}) &\rightarrow \mathbb{Z} \\ (a, b) &\mapsto \langle a \smile b, [M^4] \rangle = \int_M ab. \end{aligned}$$

Under connected sum the intersection form satisfies

$$Q_{M \# N} = Q_M \oplus Q_N.$$

By Poincaré duality it is unimodular, and by commutativity of the cup-product it is symmetric and therefore has a signature, denoted $\sigma(M)$. By a theorem of Thom, M is the boundary of a 5-manifold if and only if its signature is 0.

Hirzebruch’s signature theorem relates the signature to Pontrjagin classes, and for a 4-manifold gives

$$\int_M p_1(TM) = 3 \cdot \sigma(M). \quad (A1)$$

By definition of the Wu class u_2 for any \mathbb{Z}_2 -cohomology class x we have $\int_M x^2 = \int_M x \cdot u_2$. As M is oriented, so its first Stiefel–Whitney class vanishes, and $u_2 \equiv w_2 + w_1^2 = w_2$, so

$$\int_M x^2 = \int_M x \cdot w_2. \quad (A2)$$

Thus if M is spin, *i.e.* $w_2 = 0$, then the form Q_M is even. If M is spin then by Rochlin’s theorem its signature is

divisible by 16, and so

$$\int_M p_1(TM) \stackrel{48}{=} 0. \quad (\text{A3})$$

Finally consider $\text{spin}^{\mathbb{C}}$ 4-manifolds M , with $c_1 \in H^2(M; \mathbb{Z})$ the associated Chern class. Then c_1 reduces modulo 2 to w_2 , so is a characteristic element of Q_M . In this case we may also form the characteristic number $\int_M c_1^2$. By an elementary property of symmetric forms over \mathbb{Z} (see II.5.2 of [44]) we have

$$\int_M c_1^2 \stackrel{s}{=} \sigma(M). \quad (\text{A4})$$

We now describe to what extent these characteristic numbers may be realized.

Oriented 4-manifolds. The manifold $\mathbb{C}P^2$ has $Q_{\mathbb{C}P^2}$ given by the 1-by-1 matrix (1), so has signature 1; similarly $\overline{\mathbb{C}P}^2$ has signature -1 . By taking connected sums we may therefore realize every element of \mathbb{Z} as the signature of an oriented 4-manifold, so may realize every element of $3\mathbb{Z}$ as $\int_M p_1(TM)$.

Oriented 4-manifolds with U_1 -bundle. The manifold $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2$ has signature 0, and cohomology ring $H^*(\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2; \mathbb{Z}) = \mathbb{Z}[x, y]/(x^2 + y^2, x^3, y^3)$. Taking the U_1 -bundle with $c_1 = x$ it has $\int_{\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2} c_1^2 = 1$, and taking instead the U_1 -bundle with $c_1 = y$ it has $\int_{\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2} c_1^2 = -1$. By forming connected sums of $\mathbb{C}P^2$ and $\overline{\mathbb{C}P}^2$ with trivial U_1 -bundle, and $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2$ with the U_1 -bundles just described, we can realize any element of \mathbb{Z}^2 as $(\sigma(N), \int_N c_1^2)$ for an oriented 4-manifold N with a U_1 -bundle over it.

Spin 4-manifolds The 4-manifold $K3$ is spin, and has intersection form

$$Q_{K3} = -E_8^{\oplus 2} \oplus H^{\oplus 3},$$

where

$$H = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad E_8 = \begin{pmatrix} 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 2 \end{pmatrix} \quad (\text{A5})$$

so has signature -16 . By taking connected sums of $K3$ and its orientation-reversed analogue $\overline{K3}$ we may realize every element of $16\mathbb{Z}$ as the signature of a spin 4-manifold, so may realize every element of $48\mathbb{Z}$ as $\int_M p_1(TM)$.

Spin $^{\mathbb{C}}$ 4-manifolds. The manifold $\mathbb{C}P^2$ has cohomology ring $H^*(\mathbb{C}P^2; \mathbb{Z}) = \mathbb{Z}[x]/(x^3)$. Its usual complex structure induces a $\text{spin}^{\mathbb{C}}$ -structure having $c_1 = 3x$, and so

having

$$\begin{aligned} \sigma(\mathbb{C}P^2) &= 1 \\ \frac{1}{8} \left(\sigma(\mathbb{C}P^2) - \int_{\mathbb{C}P^2} c_1^2 \right) &= -1. \end{aligned}$$

The manifold $\mathbb{C}P^1 \times \mathbb{C}P^1$ has cohomology ring $H^*(\mathbb{C}P^1 \times \mathbb{C}P^1; \mathbb{Z}) = \mathbb{Z}[x, y]/(x^2, y^2)$. Its usual complex structure induces a $\text{spin}^{\mathbb{C}}$ -structure having $c_1 = 2x + 2y$, and so having

$$\begin{aligned} \sigma(\mathbb{C}P^1 \times \mathbb{C}P^1) &= 0 \\ \frac{1}{8} \left(\sigma(\mathbb{C}P^1 \times \mathbb{C}P^1) - \int_{\mathbb{C}P^1 \times \mathbb{C}P^1} c_1^2 \right) &= -1. \end{aligned}$$

Therefore by taking connected sums, and changing orientations, there are $\text{spin}^{\mathbb{C}}$ 4-manifolds M realising any values of $\sigma(M)$ and $\int_M c_1^2$ satisfying (A4).

Appendix B: Characteristic numbers of surface bundles over surfaces

Let $\pi : E^4 \rightarrow B^2$ be a smooth fiber bundle, with E and B closed oriented manifolds and fiber the oriented genus g surface Σ_g . Suppose we are given a U_1 -bundle on E , or equivalently a class $c_1 \in H^2(E; \mathbb{Z})$. This appendix describes the various characteristic classes and numbers which arise in this situation, and then explains the extent to which these may be realized.

1. Definitions and relations

Let $T_\pi E = \text{Ker}(D\pi : TE \rightarrow \pi^*TB)$ denote the sub-bundle of the tangent bundle of E consisting of tangent vectors parallel to the fibers of π , giving a decomposition $TE = T_\pi E \oplus \pi^*TB$. The tangent bundle of an orientable surface, such as B , becomes trivial upon the addition of a trivial line bundle, and hence its Stiefel–Whitey and Pontryagin classes vanish. We therefore have

$$\begin{aligned} p_1(TE) &= p_1(T_\pi E) =: p_1 \in H^4(E; \mathbb{Z}) \\ w_2(TE) &= w_2(T_\pi E) =: w_2 \in H^2(E; \mathbb{Z}_2). \end{aligned}$$

In addition, the orientations of TE and TB induce an orientation of the 2-dimensional bundle $T_\pi E$, and we have its Euler class

$$e(T_\pi E) =: e \in H^2(E; \mathbb{Z}).$$

As for any oriented 2-dimensional bundle, we have the identities

$$p_1 = e^2 \in H^4(E; \mathbb{Z}) \quad (\text{B1})$$

$$r_2(e) = w_2 \in H^2(E; \mathbb{Z}_2), \quad (\text{B2})$$

where $r_2 : H^*(-; \mathbb{Z}) \rightarrow H^*(-; \mathbb{Z}_2)$ denotes reduction modulo 2.

Using these classes we may form $\int_E p_1$, which by (B1) agrees with $\int_E e^2$, known as the first Miller–Morita–Mumford class. By (A1) we have $\int_E p_1 = 3 \cdot \sigma(E)$ so $\int_E p_1$ is always divisible by 3, but in fact more is true: as a consequence of the Atiyah–Singer index theorem applied to the fiberwise signature operator we have

$$\int_E p_1 \stackrel{12}{=} 0. \quad (\text{B3})$$

This may be obtained from p. 555 of Ref. 45, using that $B_2 = \frac{1}{6}$. If in addition $w_2 = 0$, then (A3) gives

$$\int_E p_1 \stackrel{48}{=} 0. \quad (\text{B4})$$

Now let us suppose that we are further given a $c_1 \in H^2(E; \mathbb{Z})$. Using this we may form $\int_E c_1^2$ and $\int_E e \cdot c_1$. As e reduces modulo 2 to w_2 by (B2), (A2) gives

$$\int_E c_1^2 \stackrel{2}{=} \int_E e \cdot c_1. \quad (\text{B5})$$

Let us now suppose in addition that $r_2(c_1) = w_2$, i.e. that c_1 provides a fiberwise spin^c -structure, and investigate its consequences. As the Euler class e also reduces to w_2 modulo 2, it follows from the Bockstein sequence that $c_1 = e + 2\bar{c}_1$ for some $\bar{c}_1 \in H^2(E; \mathbb{Z})$. Then we have $\int_E c_1^2 = \int_E e^2 + 4 \int_E (e\bar{c}_1 + \bar{c}_1^2)$ which by the divisibility results above we can write in terms of integers as $\int_E c_1^2 = 12 \cdot \left(\frac{\int_E p_1}{12}\right) + 8 \cdot \left(\frac{\int_E (e\bar{c}_1 + \bar{c}_1^2)}{2}\right)$. This implies

$$\int_E c_1^2 \stackrel{8}{=} \frac{\int_E p_1}{3}, \quad (\text{B6})$$

recovering (A4). Similarly, $\int_E e \cdot c_1 = 12 \cdot \left(\frac{\int_E p_1}{12}\right) + 2 \cdot \int_E e \cdot \bar{c}_1$ so we have

$$\int_E e \cdot c_1 \stackrel{2}{=} 0. \quad (\text{B7})$$

2. Realising characteristic numbers

The Madsen–Weiss theorem [46] and its variants [47, 48] can be used to establish the existence of surface bundles with given geometric structure and characteristic numbers in a highly indirect way, assuming that the genus g is sufficiently large in comparison with the dimension of the base. For the structure considered here, of a surface bundle equipped with a U_1 -bundle (equivalently a second integral cohomology class) on the total space and 2-dimensional base, this has been carried out in [49]. The conclusion is as follows.

Firstly, for any $g \geq 5$ there are oriented Σ_g -bundles $\pi : E^4 \rightarrow B^2$ and classes $c_1 \in H^2(E; \mathbb{Z})$ which realize any values of

$$\int_{\Sigma_g} c_1, \int_E p_1, \int_E c_1^2, \int_E e \cdot c_1$$

as long as they satisfy (B3) and (B5). This is obtained by combining Theorem A (iv) and Theorem C of Ref. 49 and using that every second homology class may be represented by a map from a surface B^2 .

Secondly, for any $g \geq 5$ there are oriented Σ_g -bundles $\pi : E^4 \rightarrow B^2$ and classes $c_1 \in H^2(E; \mathbb{Z})$ satisfying $r_2(c_1) = w_2$ which realize any values of

$$\int_{\Sigma_g} c_1, \int_E p_1, \int_E c_1^2, \int_E e \cdot c_1$$

as long as they satisfy (B3), (B6), and (B7). This is obtained by taking $c_1 = e + 2\bar{c}_1$ and translating the conditions on $\int_{\Sigma_g} \bar{c}_1, \int_E p_1, \int_E \bar{c}_1^2, \int_E e \cdot \bar{c}_1$ from the previous paragraph to conditions on $\int_{\Sigma_g} c_1, \int_E p_1, \int_E c_1^2, \int_E e \cdot c_1$.

Thirdly, the analogous analysis for spin surface bundles though not equipped with a additional line bundle, has been carried out in [50]. It follows from Example 1.11 of loc. cit. that for any $g \geq 9$ there are Spin Σ_g -bundles $\pi : E^4 \rightarrow B^2$ realising any value of $\int_E p_1$ satisfying (B4).

Appendix C: Checking (29), (30), (31)

1. Spread polynomials $S_n(x)$

The spread polynomials $S_n(x)$ are defined by $S_n(\sin^2 \theta) = \sin^2(n\theta)$. They have the explicit form[51]:

$$S_n(x) = x \sum_{m=0}^{n-1} \frac{n}{n-m} \binom{2n-1-m}{m} (-4x)^{n-m-1}$$

which has roots $\{\sin^2(\frac{m\pi}{n}) : m = 0, \dots, n-1\}$. Putting $n = k+2$ and replacing $x \rightarrow 1/x$, the polynomial

$$x^{k+2} S_{k+2}\left(\frac{1}{x}\right) = \sum_{m=0}^{k+1} \alpha_m x^m$$

$$\alpha_m := \frac{k+2}{k+2-m} \binom{2k+3-m}{m} (-4)^{k-m+1}$$

has roots $\{\csc^2 \frac{m\pi}{k+2} : m = 1, \dots, k+1\}$.

2. Symmetric polynomials e_l and p_l

Let $\xi_m := \csc^2 \frac{m\pi}{k+2}$. Their elementary symmetric polynomials are given by

$$e_l := \sum_{1 \leq m_1 < \dots < m_l \leq k+1} \xi_{m_1} \dots \xi_{m_l} = \frac{(-1)^l \alpha_{k-l+1}}{\alpha_{k+1}}$$

$$= \frac{4^l}{(k+2)(l+1)} \binom{k+2+l}{2l+1}$$

for $l \leq k+1$. For $l > k+1$, $e_l := 0$. Define

$$\begin{aligned} E_l &:= \frac{(k+2)^l}{2^l} e_l = 2^l (k+2)^{l-2} \frac{k+2}{l+1} \binom{k+2+l}{2l+1} \\ &= 2^l (k+2)^{l-2} \left[\binom{k+l+3}{2l+2} + \binom{k+l+2}{2l+2} \right]. \end{aligned}$$

Note that $E_1 = \binom{k+3}{3} \in \mathbb{Z}$, which is odd iff $k = 0 \pmod{4}$, so $kE_l \in 4\mathbb{Z}$. For $l \geq 2$, $E_l \in 4\mathbb{Z}$. Moreover $\frac{3E_2}{k+2} \in 2\mathbb{Z}$ and $\frac{E_l}{k+2} \in 2\mathbb{Z}$ for $l \geq 3$.

Define the l -th power sums $p_l := \sum_{m=1}^{k+1} \xi_m^l$. They are related to e_l 's via Newton–Girard Formulas[52]:

$$p_l = (-1)^{l+1} l e_l + \sum_{j=1}^{l-1} (-1)^{j+1} e_j p_{l-j}. \quad (\text{C1})$$

3. Checking (29)

Define $n_l = \frac{3k}{k+2} \frac{(k+2)^l}{2^l} p_l$. The expression eqn. (29) to be checked is

$$n_{g-1} \in 2\mathbb{Z}.$$

We will show this by induction for $g \geq 3$. From (C1) we have

$$n_l = (-1)^{l+1} l k \left(\frac{3E_l}{k+2} \right) + \sum_{j=1}^{l-1} (-1)^{j+1} E_j n_{l-j}. \quad (\text{C2})$$

Note that $n_1 = \frac{(k+3)(k+1)k}{2} \in \mathbb{Z}$, which is odd iff $k = 2 \pmod{4}$. For $l = 2$, we have

$$n_2 = -2k \left(\frac{3E_2}{k+2} \right) + E_1 n_1.$$

where the first term is even since the factor in bracket is even as discussed before. The second term is even since E_1 and n_1 cannot be odd at the same time. For $l \geq 3$, it can be seen that $n_l \in 2\mathbb{Z}$ inductively since both terms of (C2) are even integers.

4. Checking (30)

Define $m_l = k \frac{(k+2)^l}{2^l} p_l$. The expression eqn. (30) to be checked is

$$m_{g-1} \in 2\mathbb{Z}.$$

We will show this by induction for $g \geq 1$. From (C1) we have

$$m_l = (-1)^{l+1} l k E_l + \sum_{j=1}^{l-1} (-1)^{j+1} E_j m_{l-j}. \quad (\text{C3})$$

Note that $m_0 = k(k+1) \in 2\mathbb{Z}$. For $l \geq 1$, the first term in (C3) is even since kE_l is even. Hence $m_l \in 2\mathbb{Z}$ for all $l \geq 0$ by induction.

5. Checking (31)

The expression eqn. (31) to be checked is

$$\frac{m_{g-1}}{2} \in 2\mathbb{Z}.$$

for $g \geq 6$. We will show a stronger result that this holds for all $g \geq 2$ by induction. (C3) can be rewritten as

$$\begin{aligned} \frac{m_l}{2} &= (-1)^{l+1} l \left(\frac{kE_l}{2} \right) + \sum_{j=1}^{l-1} (-1)^{j+1} E_j \left(\frac{m_{l-j}}{2} \right) \\ \frac{m_l}{2} &\stackrel{?}{=} \sum_{j=1}^{l-1} E_j \left(\frac{m_{l-j}}{2} \right) \stackrel{?}{=} \begin{cases} E_1 \left(\frac{m_{l-1}}{2} \right) & \text{for } l \geq 2 \\ 0 & \text{for } l = 1. \end{cases} \quad (\text{C4}) \end{aligned}$$

where we used the fact that $\frac{kE_l}{2} \in 2\mathbb{Z}$ for $l \geq 1$ and $E_j \in 4\mathbb{Z}$ for $j \geq 2$. Note that $\frac{m_0}{2} = \frac{k(k+1)}{2}$ is odd iff $k = 1, 2 \pmod{4}$. Also recall that $E_1 \in \mathbb{Z}$. It is straightforward to read off that $\frac{m_1}{2} \in 2\mathbb{Z}$ and hence $\frac{m_l}{2} \in 2\mathbb{Z}$ for all $l \geq 1$ which follows by induction.

Appendix D: Group extension and trivialization

Consider an extension of a group H

$$A \rightarrow G \rightarrow H \quad (\text{D1})$$

where A is an Abelian group with group multiplication given by $x + y \in A$ for $x, y \in A$. Such a group extension is denoted by $G = A \rtimes H$. It is convenient to label the elements in G as (h, x) , where $h \in H$ and $x \in A$. The group multiplication of G is given by

$$(h_1, x_1)(h_2, x_2) = (h_1 h_2, x_1 + \alpha(h_1) \circ x_2 + e_2(h_1, h_2)). \quad (\text{D2})$$

where e_2 is a function

$$e_2 : H \times H \rightarrow A, \quad (\text{D3})$$

and α is a function

$$\alpha : H \rightarrow \text{Aut}(A). \quad (\text{D4})$$

We see that group extension is defined via e_2 and α . The associativity

$$[(h_1, x_1)(h_2, x_2)](h_3, x_3) = (h_1, x_1)[(h_2, x_2)(h_3, x_3)] \quad (\text{D5})$$

requires that

$$\begin{aligned} x_1 + \alpha(h_1) \circ x_2 + e_2(h_1, h_2) + \alpha(h_1 h_2) \circ x_3 + e_2(h_1 h_2, h_3) \\ = \alpha(h_1) \circ x_2 + \alpha(h_1) \alpha(h_2) \circ x_3 + \alpha(h_1) \circ e_2(h_2, h_3) \\ + x_1 + e_2(h_1, h_2 h_3) \end{aligned} \quad (\text{D6})$$

or

$$\alpha(h_1) \alpha(h_2) = \alpha(h_1 h_2) \quad (\text{D7})$$

and

$$e_2(h_1, h_2) - e_2(h_1, h_2 h_3) + e_2(h_1 h_2, h_3) - \alpha(h_1) \circ e_2(h_2, h_3) = 0. \quad (\text{D8})$$

Such a e_2 is a group 2-cocycle $e_2 \in H^2(\mathcal{B}H; A_\alpha)$, where H has a non-trivial action on the coefficient A as described by α . Also, α is a group homomorphism $\alpha : H \rightarrow \text{Aut}(A)$. We see that the A extension from H to G is described by a group 2-cocycle e_2 and a homomorphism α . Thus we can more precisely denote the group extension by $G = A \rtimes_{e_2, \alpha} H$.

Note that the homomorphism $\alpha : H \rightarrow \text{Aut}(A)$ is in fact the action by conjugation in G ,

$$(h, 0)(\mathbf{1}, x) = (h, \alpha(h) \circ x) = (\mathbf{1}, \alpha(h) \circ x)(h, 0), \\ \Rightarrow (h, 0)(\mathbf{1}, x)(h, 0)^{-1} = (\mathbf{1}, \alpha(h) \circ x). \quad (\text{D9})$$

Thus, α is trivial if and only if A lies in the center of G . This case is called a central extension, where the action α will be omitted.

Our way to label group elements in G :

$$g = (h, x) \in G \quad (\text{D10})$$

defines two projections of G :

$$\pi : G \rightarrow H, \quad \pi(g) = h, \\ \sigma : G \rightarrow A, \quad \sigma(g) = x. \quad (\text{D11})$$

π is a group homomorphism while σ is a generic function.

Using the two projections, $g_1 g_2 = g_3$ can be written as

$$(\pi(g_1), \sigma(g_1))(\pi(g_2), \sigma(g_2)) \\ = [\pi(g_1)\pi(g_2), \sigma(g_1) + \alpha(\pi(g_1)) \circ \sigma(g_2) + e_2(\pi(g_1), \pi(g_2))] \\ = (\pi(g_3), \sigma(g_3)) = (\pi(g_1 g_2), \sigma(g_1 g_2)) \quad (\text{D12})$$

We see that the group cocycle $e_2(h_1, h_2)$ in $H^2(\mathcal{B}H; A)$ can be pullback to give a group cocycle $e_2(\pi(g_1), \pi(g_2))$ in $H^2(\mathcal{B}G; A)$, and such a pullback is a coboundary

$$e_2(\pi(g_1), \pi(g_2)) = -\sigma(g_1) + \sigma(g_1 g_2) - \alpha(\pi(g_1)) \circ \sigma(g_2), \quad (\text{D13})$$

i.e. an element in $B^2(\mathcal{B}G; A_\alpha)$, where G has a non-trivial action on the coefficient A as described by α .

The above result can be put in another form. Consider the homomorphism

$$\varphi : \mathcal{B}G \rightarrow \mathcal{B}H \quad (\text{D14})$$

where $G = A \rtimes_{e_2} H$, and e_2 is a A -valued 2-cocycle on $\mathcal{B}H$. The homomorphism φ sends an link of $\mathcal{B}G$ labeled by $a_{ij}^G \in G$ to an link of $\mathcal{B}H$ labeled by $a_{ij}^H = \pi(a_{ij}^G) \in H$. The pullback of e_2 by φ , $\varphi^* e_2$, is always a coboundary on $\mathcal{B}G$

The above discussion also works for continuous group, if we only consider a neighborhood near the group identity $\mathbf{1}$. In this case, $e_2(h_1, h_2)$ and $\alpha(h)$ are continuous functions on such a neighborhood. But globally, $e_2(h_1, h_2)$ and $\alpha(h)$ may not be continuous functions.

-
- [1] L. D. Landau, Phys. Z. Sowjetunion **11**, 26 (1937).
 - [2] L. D. Landau, Phys. Z. Sowjetunion **11**, 545 (1937).
 - [3] X. G. Wen, Phys. Rev. B **40**, 7387 (1989).
 - [4] X. G. Wen and Q. Niu, Phys. Rev. B **41**, 9377 (1990).
 - [5] X. G. Wen, Int. J. Mod. Phys. B **04**, 239 (1990).
 - [6] E. Witten, Commun.Math. Phys. **121**, 351 (1989).
 - [7] X. G. Wen, Phys. Rev. B **43**, 11025 (1991).
 - [8] X. G. Wen, Phys. Rev. B **41**, 12838 (1990).
 - [9] C. L. Kane and M. P. A. Fisher, Phys. Rev. B **55**, 15832 (1997), cond-mat/9603118.
 - [10] X. Chen, Z.-C. Gu, and X.-G. Wen, Phys. Rev. B **82**, 155138 (2010), arXiv:1004.3835.
 - [11] Q. Niu, D. J. Thouless, and Y.-S. Wu, Phys. Rev. B **31**, 3372 (1985).
 - [12] L. Kong and X.-G. Wen, (2014), arXiv:1405.5858.
 - [13] K. Walker, <https://mathoverflow.net/questions/165830>.
 - [14] B. Zeng and X.-G. Wen, Phys. Rev. B **91**, 125121 (2015), arXiv:1406.5090.
 - [15] B. Swingle and J. McGreevy, Phys. Rev. B **93**, 045127 (2016), arXiv:1407.8203.
 - [16] Y.-M. Lu and A. Vishwanath, Phys. Rev. B **93**, 155121 (2013), arXiv:1302.2634.
 - [17] A. Mesaros and Y. Ran, Phys. Rev. B **87**, 155115 (2013), arXiv:1212.0835.
 - [18] L.-Y. Hung and Y. Wan, Phys. Rev. B **87**, 195103 (2013), arXiv:1302.2951.
 - [19] Z.-C. Gu and X.-G. Wen, Phys. Rev. B **80**, 155131 (2009), arXiv:0903.1069.
 - [20] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Phys. Rev. B **87**, 155114 (2013), arXiv:1106.4772.
 - [21] Z.-C. Gu and X.-G. Wen, Phys. Rev. B **90**, 115141 (2014), arXiv:1201.2648.
 - [22] Y.-M. Lu and A. Vishwanath, Phys. Rev. B **86**, 125119 (2012), arXiv:1205.3156.
 - [23] Z.-X. Liu and X.-G. Wen, Phys. Rev. Lett. **110**, 067205 (2013), arXiv:1205.7024.
 - [24] X. Chen and X.-G. Wen, Phys. Rev. B **86**, 235135 (2012), arXiv:1206.3117.
 - [25] T. Senthil and M. Levin, Phys. Rev. Lett. **110**, 046801 (2013), arXiv:1206.1604.
 - [26] M. DeMarco and X.-G. Wen, (2019), arXiv:1906.08270.
 - [27] X.-G. Wen, Phys. Rev. B **91**, 205101 (2015), arXiv:1410.8477.
 - [28] P. J. Morandi, <https://web.nmsu.edu/~pamorand/notes/> (1997).
 - [29] X.-G. Wen, Phys. Rev. D **88**, 045013 (2013), arXiv:1303.1803.
 - [30] L. Kong and X.-G. Wen, (2020), arXiv:2004.11904.
 - [31] F. Wilczek and A. Zee, Phys. Rev. Lett. **52**, 2111 (1984).
 - [32] X. G. Wen and A. Zee, Phys. Rev. Lett. **69**, 953 (1992), (E) Phys. Rev. Lett. **69**, 3000 (1992).
 - [33] X.-G. Wen and A. Zee, Phys. Rev. Lett. **69**, 3000 (1992).
 - [34] A. Gromov, G. Y. Cho, Y. You, A. G. Abanov,

- and E. Fradkin, *Phys. Rev. Lett.* **114**, 016805 (2015), [arXiv:1410.6812](#).
- [35] B. Zumino, W. Yong-Shi, and A. Zee, *Nucl. Phys. B* **239**, 477 (1984).
- [36] W. Meyer, *Math. Ann.* **201**, 239 (1973).
- [37] H. Endo, *Osaka Journal of Mathematics* **35**, 915 (1998).
- [38] X. G. Wen and A. Zee, *Phys. Rev. B* **46**, 2290 (1992).
- [39] X.-G. Wen, *Phys. Rev. Lett.* **66**, 802 (1991).
- [40] D. S. Freed, (2014), [arXiv:1406.7278](#).
- [41] A. Kapustin, (2014), [arXiv:1404.6659](#).
- [42] X.-G. Wen, *Phys. Rev. B* **89**, 035147 (2014), [arXiv:1301.7675](#).
- [43] D. S. Freed and M. J. Hopkins, (2016), [arXiv:1604.06527](#).
- [44] J. Milnor and D. Husemoller, *Symmetric bilinear forms* (Springer-Verlag, New York-Heidelberg, 1973) pp. viii+147, ergebnisse der Mathematik und ihrer Grenzgebiete, Band 73.
- [45] S. Morita, *Invent. Math.* **90**, 551 (1987).
- [46] I. Madsen and M. Weiss, *Ann. of Math. (2)* **165**, 843 (2007).
- [47] R. L. Cohen and I. Madsen, in *Algebraic geometry—Seattle 2005. Part 1*, Proc. Sympos. Pure Math., Vol. 80 (Amer. Math. Soc., Providence, RI, 2009) pp. 43–76.
- [48] S. Galatius, *Math. Ann.* **334**, 439 (2006).
- [49] J. Ebert and O. Randal-Williams, *Doc. Math.* **17**, 417 (2012).
- [50] O. Randal-Williams, *Adv. Math.* **231**, 482 (2012).
- [51] N. J. Wildberger, *Divine Proportions: Rational trigonometry to universal geometry* (Wild Egg, 2005).
- [52] R. Sroul, in *Programming for Mathematicians* (Springer-Verlag, Berlin, 2000) Chap. 10.12, pp. 278–279.