

Quantum Love

Ram Brustein, Yotam Sherf

Department of Physics,

Ben-Gurion University,

Beer-Sheva 84105, Israel

ramyb@bgu.ac.il, sherfyo@post.bgu.ac.il

Abstract

The response of a gravitating object to an external tidal field is encoded in its Love numbers, which identically vanish for classical black holes (BHs). Here we show, using standard time-independent quantum perturbation theory, that for a quantum BH, generically, the Love numbers are nonvanishing and negative, and that their magnitude depends on the lowest-lying levels of the quantum spectrum of the BH. We calculate the quadrupolar electric quantum Love number of slowly rotating BHs and show that it depends most strongly on the first excited level of the quantum BH. We then compare our results to the same Love number of exotic ultra compact objects and to that of classical compact stars and highlight their different parametric dependence. Finally, we discuss the detectability of the quadrupolar quantum Love number in future precision gravitational-wave observations and show that, under favorable circumstances, its magnitude is large enough to imprint an observable signature on the gravitational waves emitted during the inspiral phase of two moderately spinning BHs.

I. INTRODUCTION

The gravitational-wave (GW) observations by LIGO and the future observations by the planned Laser Interferometer Space Antenna (LISA), offer opportunities for testing strong gravity effects through precision GW measurements during the inspiral phase of a compact binary system [1–4]. As the two companions spiral around each other, they are tidally deformed [5, 6], leaving a specific imprint on the emitted GW waveform [7–12]. The tidal response of each of the companions is quantified in terms of the tidal Love numbers.

The weak external tidal field induces, generically, small nonvanishing mass (electric) and current (magnetic) moments. In the linear response approximation, the moments are proportional to the external tidal field. The largest of these induced moments is typically the mass quadrupole, which is proportional to the quadrupolar tidal field \mathcal{E}_{ab} , $Q_{ab} = -\frac{2}{3}k_2 R^5 \mathcal{E}_{ab}$. Here k_2 is the dimensionless quadrupolar electric tidal Love number and R is the radius of the inspiraling object.

The calculation of k_2 is performed in great detail in [13–15]. Its value is most sensitive to the object compactness¹ $C = M/R$. For the case that C approaches that of a black hole (BH), $C \rightarrow 1/2$, the Love number exhibits a universal decrease, tending precisely to zero in the BH limit. This universal behavior is a consequence of the BH no-hair property [10, 14, 16, 17]. The exact vanishing of k_2 for BHs² and being the largest of the dimensionless Love numbers, makes k_2 a key diagnostic for any deviations from classical general relativity (GR).

In [21], the Love numbers for several exotic ultracompact objects (UCOs) were calculated and were shown not to vanish. The numerical results exhibit a universal, model-independent, logarithmic suppression on the relative deviations from the Schwarzschild radius $R = 2M(1 + \epsilon)$.

We are interested in calculating the Love numbers of large astrophysical BHs. As for any macroscopic object, the Bohr correspondence principle implies that some quantum state corresponds to the classical BH, no matter how large it is. In the following, we use the term “quantum black hole” (QBH) to mean the quantum state that corresponds to a classical BH.

¹ We use relativistic units $G, c = 1$ and consider nonrotating BHs unless stated otherwise.

² In [18, 19] it is claimed that the Love number for spinning BHs in an axisymmetric tidal field ($m \neq 0$) is nonvanishing. The results were challenged in [20].

The QBH is therefore a UCO that possesses a horizon and, in addition, has a discrete spectrum of quantum mechanical energy levels. These energy levels can be viewed as coherent states that correspond to macroscopic, semiclassical excitations of the QBH. In the ground state of the QBH, the exterior geometry is exactly the Schwarzschild geometry. But, when a QBH is in an excited state, it displays deviations from its GR description, and therefore it can be, in principle, distinguished from its classical counterpart.

The classical BH is bald, while the QBH has some “quantum hair” [22–24]. Moreover, the properties of the quantum hair can be entirely explained by an external observer via the Bohr correspondence principle that requires some specific changes to the near horizon geometry, without any need to invoke new physical principles [22, 25]. The amount of information that the quantum hair carries is limited. However, if observed, it could provide unrivaled information on some properties of the spectrum of the QBH [25–29]. Quantum imprints due to tidal heating in the inspiral phase were also studied recently in [30, 31].

We will show that the Love numbers are part of this quantum hair and can, in principle, be observed. In practice, it is k_2 that seems to offer the best opportunity for detection.

Quantum effects for large astrophysical BHs are universally expected to be negligibly small, based on the expectation that the strength of quantum effects is controlled by the extremely small ratio of the Planck length squared to typical curvatures l_P^2/R_S^2 . However, we argue in the following that the strength of quantum effects for QBHs can be much larger.

In GR, the interior of a BH is empty except for a possibly singular core. The firewall argument marked the beginning of a new era in the theory of QBHs [32, 33], indicating that this picture is in need of a substantial revision. Forerunners of the argument and a more recent review can be found in [34–37], respectively.

Putting remnants aside, two main classes of solutions to the firewall problem emerged as possible candidates. In the first class the horizon region is a vacuum, but novel nonlocal physics is introduced to resolve the information paradox: the degrees of freedom very far from the horizon are not distinct from the degrees of freedom inside the horizon [38, 39]. The singularity is often viewed as irrelevant, under the premise that it will be regularized somehow in a way that does not affect the structure of spacetime on horizon scales.

In the second class, BHs are described by nonsingular states that do not collapse under their own gravity. Strong quantum effects “smear” the would-be singularity over horizon-sized length scales. These changes lead to a spectrum of excitations whose characteristic

scale is the horizon rather than the Planck length. The self-consistency of this description of the interior requires a significant departure from semiclassical gravity, as well as some exotic matter which is outside the realm of the standard model [40]. Fuzzballs [41, 42] and the polymer model [43] are in this class. The new physics that resolves the singularity introduces a new scale, and in addition to the Planck scale, the ratio of the two scales can be viewed as a coupling constant. For example, in string theory, this length scale is the string scale l_s , and it is rather the ratio $g_s^2 = l_P^2/l_s^2$ that controls the strength of quantum effects. The magnitude of g_s^2 is expected to be small, but of the order of all other known gauge couplings, $g_s^2 \sim 0.1$.

Here, we present a general, closed expression for both electric (polar) and magnetic (axial) Love numbers (tensor) for QBHs in terms of their spectrum. The calculation is performed in an analogy to the calculation of the polarizability of an atom by using second-order time-independent perturbation theory. We show that the Love numbers are most sensitive to the lowest-lying energy level. From this perspective, the Love numbers do not vanish because the tidal field mixes a small amount of the first excited level with the ground state.

In a follow up paper [44], we describe explicitly the connection between the classical and quantum Love calculations using the ideas presented in [45–48]. We first establish an effective description for the interior fluid modes of ultracompact objects as a collection of driven harmonic oscillators characterized by their frequencies. We then find the appropriate boundary conditions on the perturbed Einstein equations and show that derivation of the quantum Love number of a quantum black hole matches exactly the standard classical calculation of the Love number [13–15], when quantum expectation values are replaced by the corresponding classical quantities, as dictated by the Bohr correspondence principle. The quantum Love number is equal to the classical Love number that is computed in the traditional way. The current paper and [44] have different goals. The goal of the current paper is to study the response of a general quantum system to an external tidal field and demonstrate how it acquires nonvanishing Love numbers. On the other hand, the motivation of [44] is to demonstrate how an object that possesses a horizon can have a nonvanishing Love number. They are similar in that both rely on the interpretation of the nonrelativistic fluid modes as large quantum excitations.

The paper is organized as follows. In Sec. II we review the standard calculation of the atom’s electric polarizability using time-independent perturbation theory. Then, by replac-

ing the external electric field and the dipole moment with the gravitational tidal field and the mass and current moments, respectively, we derive a general expression for the gravitational polarizability of a quantum mechanical object—the Love numbers. Next, in Sec. III, by applying the Bohr correspondence principle we evaluate the Love number and find that it is negative, and its magnitude depends on the lowest-lying levels of the quantum spectrum of the QBH. We demonstrate the ideas by replacing the large excitations spectrum of the QBH with an analogous semiclassical fluidlike description. Then by imposing generic boundary conditions, we provide an explicit expression of the Love number of QBHs. Finally, in Sec. IV we discuss the possible observation of the quantum Love numbers. We show that, under favorable circumstances, future LISA observations could indeed detect them by precise measurement of the spectrum of GWs emitted during the inspiral phase of a binary system of supermassive moderately spinning BHs. In the Appendix, we discuss the promotion of the magnetic Love numbers of a slowly rotating object to tensors and the spin corrections to the tidal Love numbers.

II. QUANTUM LOVE NUMBERS

As a prelude to the calculation of the quantum Love numbers, we briefly recall the analogous calculation of the polarizability of an atom. The atom is placed in a region of an approximately uniform electric field \mathcal{E}_i that is induced by a weak external potential U_{ext} , $\mathcal{E}_i = -\frac{\partial U_{ext}}{\partial x^i}$. The interaction of the atom with the external electric field, is expressed in terms of the dipole moment $D = \int \rho(x')x'dV'$, where the integral is performed over the charge distribution. The interaction is given by $\hat{V}_{int} = -\mathcal{E}_i\hat{D}_i$. The induced dipole moment of the perturbed atom can be calculated in second-order time-independent perturbation theory [49]. In this case, symmetry implies that the atom's linear response to the external electric field is then $\langle\Psi_0|\hat{D}_i|\Psi\rangle = \alpha\mathcal{E}_i$, where $|\Psi_0\rangle = |1,0,0\rangle$ is the ground state of the atom, $|\Psi\rangle$ is the first-order correction to the atom ground state $|\Psi\rangle = |\Psi_0\rangle + \sum_{n,l,m} |n,l,m\rangle \frac{\langle 1,0,0|\hat{V}_{int}|n,l,m\rangle}{\Delta E_{1,n}}$, and α is the electric polarizability,

$$\alpha = \sum_{n,|m|\leq 1} \frac{|\langle\Psi_0|\hat{D}_i|n,l=1,|m|\leq 1\rangle|^2}{\Delta E_{1,n}}, \quad (2.1)$$

and where l and m are the angular quantum numbers, n is the radial quantum number and $\Delta E_{1,n} = E_1 - E_n$.

We derive an expression for the gravitational polarizability—the Love numbers—by replacing the external electric field and the dipole moment by the tidal field and the mass and current moments, respectively.

We consider the inspiral phase of a binary system, where one of the companions is an object of mass M_{ext} on a circular orbit of radius b and the other is a nonrotating QBH of mass M_{BH} and radius R_S . In the early stages of the inspiral, the BH responds to the external slowly varying tidal field that is generated by its companion. For $b \gg R_S$ one can expand the Newtonian potential $U_{ext} = -M_{ext}/|\vec{b} - \vec{x}|$ of the external body in the vicinity of the BH in its local inertial frame, $U(t, x)_{ext} = U_{ext}(0) + \frac{1}{2} \frac{\partial^2 U_{ext}}{\partial x^i \partial x^j} \Big|_0 x^i x^j + \dots$.

The interaction of the QBH with the external field is expressed in terms of the quantum trace-free symmetric mass and current multipole moments, $\hat{Q}^{(l)}$ and $\hat{S}^{(l)}$, these being the quantum counterparts of the classical multipoles [5]. We further assume that the expectation value of the mass and current moments of the BH vanishes in the BH ground state, as dictated by the angular symmetry of the multipole operators and in accordance with the classical no-hair theorems, denoting the ground state of the BH by $|\Psi_0\rangle$, $\langle \Psi_0 | \hat{Q}^{(l)} | \Psi_0 \rangle = 0$, $\langle \Psi_0 | \hat{S}^{(l)} | \Psi_0 \rangle = 0$. Since the external potential is slowly varying, time-independent perturbation theory should be a good approximation.

Let us evaluate explicitly the correction to the ground state energy due to the induced quadrupole, \hat{Q}_{ij} . We follow here the conventions of [13], in analogy to the electric polarizability calculation, $\hat{V}_{int} = -\frac{1}{2} \mathcal{E}_{ij} \hat{Q}_{ij}$, where $\mathcal{E}_{ij} = \frac{\partial^2 U_{ext}}{\partial x^i \partial x^j}$ is the tidal field. The sign of the interaction term is important and leads, generically, to negative quantum Love numbers. For neutron stars, the sign of the interaction term is positive and it leads to positive Love numbers [13, 21, 50]. The physical reason is that for BHs, the mass as a function of the radius $M(R)$ is an *increasing* function, while for neutron stars it is a *decreasing* function (see Fig. 2 of [50]). For UCOs, the sign of the Love number is also, generically, negative.

The leading-order correction to the BH ground state quadrupole is given by

$$\langle \Psi_0 | \hat{Q}_{kl} | \Psi \rangle = -\mathcal{E}_{ij} \sum_{n_r > 1, |m| \leq 2} \frac{\langle \Psi_0 | \hat{Q}_{ij} | n_r, 2, m \rangle \langle n_r, 2, m | \hat{Q}_{kl} | \Psi_0 \rangle}{\Delta E_{1, n_r}}, \quad (2.2)$$

where $\Delta E_{1, n_r} = E_1 - E_{n_r}$. Here the radial number of the ground state Ψ_0 is denoted by $n_r = 1$, so the energy of the ground state is $E_1 = M_{BH}$. Symmetry implies that the BH electric quadrupolar Love number is given by $\frac{1}{2} \langle \Psi_0 | \hat{Q}_{ij} | \Psi \rangle = -\lambda_2 \mathcal{E}_{ij}$. Here λ_2 is the dimensional quadrupolar Love number. The dimensionless Love number k_2 is commonly

defined as $k_2 = \frac{3}{2}R^{-5}\lambda_2$. From Eq. (2.2), it follows that

$$k_2 = -\frac{3}{2R^5} \sum_{n_r > 1, |m| < 2} \frac{1}{2} \frac{|\langle \Psi_0 | \hat{Q}_{ij} | n_r, 2, m \rangle|^2}{|\Delta E_{1, n_r}|}. \quad (2.3)$$

Equation (2.3) is the main result of our paper. It demonstrates that, generically, a quantum mechanical object must have a nonvanishing quadrupolar Love number that depends solely on the quantum state of the object and its energy spectrum. The negative sign of k_2 reflects the fact that the energy of a BH increases when its radius becomes larger, as previously explained.³ As it is previously explained, in [44], we showed explicitly that the quantum Love number is equal to the classical Love number that is computed in the traditional way when quantum expectation values are replaced by the corresponding classical quantities, as dictated by the Bohr correspondence principle.

The general expressions for the higher- l electric and magnetic quantum Love tensors can be obtained by following the steps that led to Eq. (2.2):

$$k_l^E = - \sum_{n_r > 1, l, |m| \leq l} \frac{1}{R^{2l+1}} \frac{(2l-1)!!}{2(l-2)!} \frac{1}{l!} \frac{|\langle \Psi_0 | \hat{Q}^{(l)} | n_r, l, m \rangle|^2}{\Delta E_{1, n_r}}, \quad (2.4)$$

$$k_l^B = - \sum_{n > 1, l, |m| \leq l} \frac{1}{R^{2l+1}} \frac{(l+1)(2l-1)!!}{6(l-2)!} \frac{1}{l!} \frac{|\langle \Psi_0 | \hat{S}^{(l)} | n_r, l, m \rangle|^2}{\Delta E_{1, n_r}}. \quad (2.5)$$

Recently, in [51] it was shown that the magnetic Love numbers of a slowly rotating object should be promoted to tensors. We discuss this in more detail in addition to the spin corrections to the tidal Love number in the Appendix.

Again, the conclusion is that, generically, QBHs must possess nonvanishing Love numbers.

III. ELECTRIC QUADRUPOLEAR QUANTUM LOVE NUMBER

The starting point of our evaluation of k_2 is Eq. (2.3). The external quadrupole tidal field is proportional to the spherical harmonic Y_{20} due to the symmetry of the inspiral trajectory. The induced quadrupole shares this angular dependence. It follows that

$$k_2 = -\frac{3}{4R^5} \sum_{n_r} \frac{|\langle \Psi_0 | \hat{Q} | n_r, 2, 0 \rangle|^2}{|\Delta E_{1, n_r}|}. \quad (3.1)$$

³ This argument is also supported by the *shape Love number* [52, 53].

To calculate k_2 we need to find the discrete quantum spectrum of the QBH. In principle, we should solve the quantum gravity equations and find the spectrum of the BH. Remarkably, this can actually be done for specific models (see, for example, [54]). Here, we rather solve the corresponding classical wave equation and then use the Bohr correspondence principle to find the spectrum in a similar way to the way that the Bohr-Sommerfeld quantization rule was used to find the spectra of atoms. A similar procedure for scalar waves was carried out in [25]. First, we use scaling arguments to estimate k_2 and then support the scaling arguments by a calculation.

On dimensional grounds, the coherent state energy spectrum of macroscopic excitations of the QBH takes the classical form $|\Delta E_{1,n_r}| \sim M_{BH} \omega_{n_r}^2 R^2$, where ω_{n_r} is the frequency of the mode $|n_r, 2, 0\rangle$. The matrix element of the quadrupole operator scales as $|\langle \Psi_0 | \hat{Q} | n_r, 2, 0 \rangle| \sim |\Delta E_{1,n_r}| R^2 \sim M_{BH} \omega_{n_r}^2 R^4$. It follows that each term in the sum in Eq. (3.1) scales as $\frac{1}{R^5} \frac{|\langle \Psi_0 | \hat{Q} | n_r, 2, 0 \rangle|^2}{|\Delta E_{1,n_r}|} \sim \frac{|\Delta E_{1,n_r}|}{R} \sim \frac{|\Delta E_{1,n_r}|}{M_{BH}} \sim \omega_{n_r}^2 R^2$. This semiclassical treatment is supported by observing that the occupation numbers N , in the excited energy levels, scale as $N \hbar \omega_{n_r} \sim M_{BH} \omega_{n_r}^2 R^2$ so $N \sim (\omega_{n_r} R) S_{BH} \gg 1$. We may also use a scaling argument and an explicit calculation to show that the contributions to k_2 in Eq. (3.1) of the excited states above the first excited state are suppressed, so we can approximate the sum over n_r by the contribution from the first excited state. This is a typical situation in most quantum systems. Furthermore, all the terms in the sum are positive, so the approximate value of the magnitude of k_2 is an underestimate. In this case, it is justified to approximate the sum by the contribution of the first excited state. Putting the two scaling arguments together, we get an estimate for k_2 ,

$$k_2 = -\frac{3}{4R^5} \frac{|\langle \Psi_0 | \hat{Q} | 2, 2, 0 \rangle|^2}{|\Delta E_{1,2}|} \sim -\frac{|\Delta E_{1,2}|}{M_{BH}} \sim -\omega_2^2 R^2. \quad (3.2)$$

We now turn to a quantitative evaluation of k_2 , whose aim is to calculate the order unity numerical factor in Eq. (3.2). We emphasize that the estimate in Eq. (3.2) is valid in a model-independent way. The specific model that we discuss will serve to illustrate the procedure in a simple model for which numerical factors can be calculated analytically. Later we parametrize the Love number in terms of the single parameter g^2 and interpret its detectability in terms of the estimate in Eq. (3.2).

Because gravity in the interior of the BH is strongly coupled, one cannot use the semi-classical geometric description in terms of a curved spacetime. It needs to be replaced by

describing gravity as an inertial force in a flat space, a replacement that is allowed by virtue of Einstein's equivalence principle. The specific nature of the excitations in the interior is unimportant and so is the equivalence of the two descriptions of gravity. The only relevant aspect is that excitations are macroscopic, horizon-scale excitations so that applying the Bohr principle is justified.

The idea is that the exotic matter in the interior of the QBH can be effectively viewed as a fluid that supports pulsating modes as for a relativistic star. These fluid modes would exist in addition to the standard spacetime modes of the exterior. The perturbations are divided into two sectors, the fluid modes and spacetime modes. Due to their low speed of sound and the compactness of the QBH, fluid modes are decoupled from the spacetime perturbations as in the Cowling approximation [55–57].

The boundary conditions (BCs) are chosen as follows. Spherical symmetry requires fully reflecting BCs at the center of the QBH. The QBH has an outer surface that behaves just like a classical BH horizon in the classical limit. In this case, the internal fluid modes decouple from the exterior. Then, absence of transmission, or perfect reflection at the outer surface is the correct BC. When quantum effects are small, the outer surface is only partially opaque and so the reflection is not perfect. We found that, quantitatively, both BCs lead to almost identical spectra. Since the analysis is much simpler in the former case, we will impose this BC at the outer surface and find the spectrum of normal modes rather than quasinormal modes.

Thus, the conclusion is that the classical equation that we need to solve is the Laplace equation,

$$\vec{\nabla}^2 \Psi_2(r) = 0, \quad (3.3)$$

with the generic BC $\Psi_{2|r=0} = 0$, and $\Psi'_{2|r=R} = 0$. The solution of Eq. (3.3) is

$$\Psi_2(r) = \mathcal{N}_2 j_2(qr) Y_{20}(\theta, \phi), \quad (3.4)$$

where j_2 is the spherical Bessel function, Y_{20} is the (real) spherical harmonic function with $l = 2$, $m = 0$ and \mathcal{N}_2 is a normalization factor which will be determined later. The BC in this case allows only discrete values on the magnitude of the wave number q ,

$$j'_2(qR) = 0, \quad (3.5)$$

which is very well approximated by

$$q_{n_r} = \left(n_r - \frac{1}{2}\right) \frac{\pi}{R}, \quad n_r = 3, 4, \dots, \quad (3.6)$$

while for $n_r = 2$, the value is somewhat lower,

$$q_2 \simeq 1.06 \frac{\pi}{R}. \quad (3.7)$$

Condition (3.6) can also be viewed as a manifestation of the Bohr quantization condition in the corresponding QBH. Substituting $P = \hbar q$, we find

$$PR = \pi \hbar \left(n_r - \frac{1}{2}\right). \quad (3.8)$$

We need to calculate $|\Delta E_{1,2}|$ and $|\langle \Psi_0 | \hat{Q} | 2, 2, 0 \rangle|$ using the solution $\Psi_{2,2} = \mathcal{N}_{2,2} j_2(q_2 r) Y_{20}$, with the wave number given above. First, because the classical waves are nonrelativistic,

$$|\Delta E_{1,2}| = \frac{1}{2} M_{BH} \omega_2^2 R^2 = \frac{1}{2} M_{BH} g^2 q_2^2 R^2. \quad (3.9)$$

In the last equality, we introduced a parametrized dispersion relation $\omega_2^2 = g^2 q_2^2$, where $g^2 \ll 1$ determines the energy of the first excited level and is the only free parameter of our model. The effective index of refraction in the cavity is $1/g^2$ (see also [22, 25]).

To evaluate the expectation value $|\langle \Psi_0 | \hat{Q} | 2, 2, 0 \rangle|$, Eq. (3.2), we will need a more elaborate calculation. First, we need the general expression for the excitation energies for $n_r \geq 3$,

$$|\Delta E_{1,n_r}| = \frac{1}{2} g^2 M_{BH} \pi^2 \left(n_r - \frac{1}{2}\right)^2, \quad (3.10)$$

where we have absorbed any additional n_r -independent factors into g^2 and assumed that the dispersion relation is the same for all modes. The excitation energy has to be parametrically small compared to the BH mass, $|\Delta E_{1,n_r}| \ll M_{BH}$. This condition restricts the validity of the estimate in Eq. (3.10) and the range of n_r in the sum in Eq. (3.1) (see also the discussion in the subsequent section).

To proceed, the classical quantity that corresponds to the matrix element $|\langle \Psi_0 | \hat{Q} | 2, 2, 0 \rangle|$ is given by

$$|\langle \Psi_0 | \hat{Q} | 2, 2, 0 \rangle| \leftrightarrow \int r^2 dr d\Omega_2 \Delta \rho_{2,2}(r) r^2 Y_{20} \Psi_{2,2}. \quad (3.11)$$

This quantity is evaluated by calculating the effective energy density in the first excited state, $\Delta \rho_{2,2}(r)$, using the following comparison. On one hand,

$$|\Delta E_{1,n_r}| = \int r^2 dr \Delta \rho_{2,n_r}(r). \quad (3.12)$$

On the other hand, to lowest order in g^2 , the energy $|\Delta E_{1,n_r}|$ is proportional to $\omega_{n_r}^2$,

$$\begin{aligned} |\Delta E_{1,n_r}| &= \int r^2 dr d\Omega_2 |\Psi_{2,n_r}|^2 \omega_{n_r}^2 \\ &= |\mathcal{N}_{2,n_r}|^2 \int_0^R r^2 dr j_2^2\left(\frac{\omega_{n_r}}{g} r\right) \omega_{n_r}^2, \end{aligned} \quad (3.13)$$

where we have used Eq. (3.4) and Eq. (3.6) and performed the angular integral. Comparing the two expressions for $|\Delta E_{1,n_r}|$, we find that

$$\Delta\rho(r)_{2,n_r}(r) = \frac{|\Delta E_{1,n_r}|}{I_{2,n_r}} j_2^2\left(\frac{\omega_{n_r}}{g} r\right) \quad (3.14)$$

and

$$|\mathcal{N}_{2,n_r}|^2 = \frac{|\Delta E_{1,n_r}|}{\omega_{n_r}^2 I_{2,n_r}}, \quad (3.15)$$

where $I_{2,n_r} = \frac{R^3}{\pi^3(n_r - \frac{1}{2})^3} \int_0^{\pi(n_r - \frac{1}{2})} y^2 dy j_2^2(y)$. Substituting Eq. (3.14) into expression (3.11) results in the following expression:

$$\begin{aligned} |\langle \Psi_0 | \hat{Q} | n_r, 2, 0 \rangle| &\leftrightarrow \int r^2 dr d\Omega_2 \frac{|\Delta E_{1,n_r}|}{I_{2,n_r}} j_2^3\left(\frac{\omega_{n_r}}{g} r\right) (Y_{20})^2 \\ &= |\Delta E_{1,n_r}| \mathcal{N}_{2,n_r} \frac{I_{4,n_r}}{I_{2,n_r}}, \end{aligned} \quad (3.16)$$

where $I_{4,n_r} = \frac{R^5}{\pi^5(n_r - \frac{1}{2})^5} \int_0^{\pi(n_r - \frac{1}{2})} dy y^4 j_2^3(y)$.

Putting all the pieces together we find that the corresponding expression to the ratio appearing in Eq. (3.1) is the following:

$$\frac{|\langle \Psi_0 | \hat{Q} | n_r, 2, 0 \rangle|^2}{|\Delta E_{1,n_r}|} \leftrightarrow \frac{|\Delta E_{1,n_r}|^2}{\omega_{n_r}^2} \frac{I_{4,n_r}^2}{I_{2,n_r}^3}. \quad (3.17)$$

The sum of terms with $n_r \geq 3$ in Eq. (3.1) is therefore given by

$$\begin{aligned} &\sum_{n_r=3} \frac{|\Delta E_{1,n_r}|^2}{\omega_{n_r}^2} \frac{I_{4,n_r}^2}{I_{2,n_r}^3} \\ &= \frac{1}{4} g^2 M_{BH}^2 R^3 \sum_{n_r=3} \pi(n_r - \tfrac{1}{2}) \left(\tilde{I}_{4,n_r}\right)^2 \left(\tilde{I}_{2,n_r}\right)^{-3}, \end{aligned} \quad (3.18)$$

where we also use the energy spectrum Eq. (3.10) and the integral $\tilde{I}_{2,n_r} = \int_0^{\pi(n_r - \frac{1}{2})} y^2 dy j_2^2(y)$

scales linearly with $\pi(n_r - \frac{1}{2})$, and the integral $\tilde{I}_{4,n_r} = \int_0^{\pi(n_r - \frac{1}{2})} y^4 dy j_2^3(y)$ is approximately a constant. The different scalings arise because of the different scaling of integrals of even

and odd powers of the spherical Bessel function. The final result is that the terms in the sum scale as $1/(\pi(n_r - \frac{1}{2})^2)$, with odd n_r terms being much smaller than even n_r terms. The $n_r = 2$ term is the largest in the sum and next largest term is the $n_r = 4$ term, whose magnitude is about 1/5 of the $n_r = 2$ term.

Once both $|\Delta E_{1,2}|$ and $|\langle \Psi_0 | \hat{Q} | 2, 2, 0 \rangle|$ are known, they can be substituted into Eq. (3.2). The result is given by

$$\begin{aligned} k_2 &= -\frac{3}{16} \frac{1}{q_2 R} \frac{M_{BH}^2}{R^2} \frac{\tilde{J}_4^2}{\tilde{J}_2^3} \omega_2^2 R^2 \\ &= -\frac{3}{16} q_2 R \frac{M_{BH}^2}{R^2} \frac{\tilde{J}_4^2}{\tilde{J}_2^3} g^2, \end{aligned} \quad (3.19)$$

where the integrals $\tilde{J}_2 = \int_0^{q_2 R} y^2 dy j_2^2(y)$ and $\tilde{J}_4 = \int_0^{q_2 R} y^4 dy j_2^3(y)$ can be evaluated analytically. Substituting the numerical values of the integrals and setting $M_{BH}/R = 1/2$, we arrive at our final result,

$$\begin{aligned} k_2 &= -0.09 \omega_2^2 R^2 = -0.18 \frac{|\Delta E_{1,2}|}{M_{BH}} \\ &= -0.99 g^2. \end{aligned} \quad (3.20)$$

As anticipated in Eq. (3.2), k_2 scales as $\omega_2^2 R^2$.

We can compare the value of k_2 in Eq. (3.20) to the values of k_2 for other compact objects. For Neutron stars k_2 is positive and its magnitude is much larger than the value of k_2 in Eq. (3.20). For the exotic UCOs, universal logarithmic dependence was found in [21]. These analyses assumed that some modifications lead to a shift at the UCO outer surface $R = 2M(1 + \epsilon)$ and concluded $k_2 \sim 1/|\ln \epsilon|$ and that it is negative. The real part of the frequency of spacetime modes for these UCOs for the $n = 2$ mode is $\omega_{2,UCO} \sim 1/|\ln \epsilon|$ [58]. For the BH area quantization model [59] (see also [30, 60–62]), $\omega_n = \alpha n / 16\pi R$, with α being a dimensionless coefficient of order unity, so we can apply our semiclassical treatment and from Eq. (3.2) calculate the Love number $k_2 \simeq \frac{3}{16} \left(\frac{\alpha}{8\pi}\right)^2$.

IV. DETECTABILITY

Here, we discuss the possibility of measuring the quantum Love number in future LISA observations of supermassive BH binaries, which LISA can observe from the early stages of the inspiral up to the coalescence. We show that for such binary systems, the sensitivity is

sufficient for possibly detecting the quantum tidal deformation effects for a range of values of g^2 . We include here also the case of moderately spinning BHs whose dimensionless spin parameter is $\chi \lesssim 0.7$. We later show that the main effect of the spin is to modify the radius of the BH for the same mass, $R = M(1 + \sqrt{1 - \chi_i^2})$ and that the direct effect of the spin on the spectrum of the BH can be neglected.

Following [63, 64] (see also [65–67]), we determine for which values of g^2 , the statistical error due to the detector noise is small enough for observing the tidal deformation effects. We also need to include tidal heating effects [68–71] which are present because QBHs possess a horizon. However, we found that these induce small changes to the error estimation.

To estimate the statistical error in measuring the Love number, we use a parameter estimation method based on the Fisher matrix $\Gamma_{ij} = (\frac{\partial h}{\partial \theta^i} | \frac{\partial h}{\partial \theta^j})$, where the inner product $(\cdot | \cdot)$ is defined by $(h_1 | h_2) = 4 \text{Re} \int_{f_{\min}}^{f_{\max}} \frac{\tilde{h}_1(f) \tilde{h}_2^*(f)}{S_n(f)} df$. The LISA noise spectral density is denoted by $S_n(f)$ [1, 72]. The minimal frequency of LISA’s observation band is denoted by f_{\min} , $f_{\min} \approx 10^{-5}$ Hz which corresponds to an observation time of about one year [73]. The maximal frequency f_{\max} is taken to be the frequency at the innermost stable circular orbit (ISCO) [74]. The model signal and the true signal are parametrized by the function $\theta^i = (\ln \mathcal{A}, \ln \mathcal{M}, \ln \eta, \Psi_c, t_c, \chi_1, \chi_2, \Lambda)$, whose arguments are the amplitude \mathcal{A} , the chirp mass $\mathcal{M} = \eta^{3/5} M$, the symmetric mass ratio $\eta = M_1 M_2 / M^2$, the phase Ψ_c , the time at coalescence t_c , the dimensionless spin parameters χ_1, χ_2 and the dimensionless average tidal deformability parameter $\Lambda = \frac{16}{13} \left[\left(1 + 12 \frac{M_2}{M_1}\right) \frac{M_1^5}{M^5} \tilde{\Lambda}_1 + \left(1 + 12 \frac{M_1}{M_2}\right) \frac{M_2^5}{M^5} \tilde{\Lambda}_2 \right]$, where $M = M_1 + M_2$, $\tilde{\Lambda}_i = \lambda_i / M^5$ and λ_i is defined in Sec. II. For this set of parameters, the root-mean-square error in measuring Λ is expressed through the inverse of the Fisher matrix $\sigma_\Lambda = \sqrt{\langle (\Delta \Lambda)^2 \rangle} = \sqrt{(\Gamma^{-1})_{\Lambda\Lambda}}$.

For a binary inspiral, the Fourier transform of the signal is modeled by $\tilde{h}(f, \theta^i) = \mathcal{A} e^{i\Psi}$, where $\Psi = \Psi_{PP} + \Psi_{TD} + \Psi_{TH}$ are the phases of the point-particle, tidal deformability and tidal heating effects, respectively.

The approximation method adopted here is the analytical “TaylorF2 approximant” [75–77]. We include correction terms to the GW phase in the form of spin-orbit, spin-spin and cubic-spin corrections up to 3.5 PN order relative to the leading-order GW term [78, 79], tidal deformability terms to 5 PN and 6 PN order [67, 80, 81], and tidal heating correction term for spinning BHs to the leading 2.5 PN order relative to the leading-order GW term [78, 82]. The amplitude is taken to leading PN order and includes the sky-averaged prefactor

[73].




FIG. 1: The relative statistical error in future LISA observations of $|\sigma_\Lambda/\Lambda|$ for several values of g^2 is shown as a function of the spin parameter χ . We assume equal spin and equal mass $M = 10^6 M_\odot$ companions and that the luminosity distance to the binary system is $D_l = 2$ Gpc. Points below the horizontal dashed line correspond to detections at the 3σ level. The value $g^2 = 0.18$ is a limiting value for which $|\Delta E_{1,2}| = M_{BH}$.

The justification for using the TaylorF2 approximate to estimate the detectability of the Love number of the QBHs is the following. As previously discussed, the only difference between a BH and a QBH is in the response of the QBH to external perturbations. Except for this difference, BHs and QBHs are indistinguishable to a distant external observer as both can be viewed as point masses, being well described by the spherically symmetric vacuum solution. Thus, the TaylorF2 approximation for the QBH and the BH is identical up to the subleading 5th PN order in which tidal deformation effects enter. Consequently, the use of the TaylorF2 approximate for BH-like objects is a standard accepted practice in similar contexts[21, 63, 66].

The results presented in Fig.1 indicate that it is possible to place significant constraints on, or possibly measure the quantum Love number, $|k_2| = 3.96 \times g^2(1 + \sqrt{1 - \chi_i^2})^{-2}$, for

supermassive, moderately spinning binaries ($M \sim 10^6 M_\odot$, $\chi_i \lesssim 0.7$) at luminosity distance $D_l = 2$ Gpc. For example, taking $g^2 = 0.06$ and for spin $\chi = 0.6$ (so $k_2 \approx 0.073$), the relative error $|\sigma_\Lambda/\Lambda| \approx 0.2$ leads to detections at 5σ confidence. Our results suggest that it would be possible to measure $\alpha \simeq 12$ of the area quantization model to better than 1σ confidence.

We wish to emphasize that the effect of degeneracy among the parameters could have been important for determination of the statistical error on Λ . However, as pointed out in [83, 84], even when the degeneracy is maximized, its effect would have increased the relative error on Λ by not more than its square root. We conclude that including the effects of degeneracy is not required at the level of accuracy that we have adopted, as it would not have changed our main conclusion significantly.

V. SUMMARY

In this paper we calculated the Love number of QBHs using standard time-independent quantum perturbation theory. We showed that, unlike classical BHs whose Love numbers vanish, the Love numbers of QBHs are generically nonvanishing and negative and their magnitude depends most strongly on the first excited level of the quantum spectrum. We focused on evaluating the largest Love number k_2 , the electric quadrupolar Love number. Replacing quantum expectation values by the corresponding classical quantities, as dictated by the Bohr correspondence principle, we found that k_2 of nonrotating QBHs takes the universal form

$$k_2 = -\mathcal{N}\omega_2^2 R^2, \quad (5.1)$$

where \mathcal{N} is a positive numerical factor of order unity that is determined by the generic boundary conditions of QBHs, Eqs. (3.3) and (3.4), and the object's excitation spectrum. As shown in Sec. III, the result in Eq. (5.1) is universal and holds for any macroscopic quantum object.

We then proceeded to show that the accumulated dephasing due to the dissipation of tidal deformation in supermassive moderately spinning binaries during ~ 1 year of observation is large enough to induce a significant deviation on the orbital phase. Thus, indicating the detectability of the Love number of QBHs with future precision GW measurements.

Acknowledgments

We thank Zvi Bern, Vitor Cardoso and Julio Parra-Martinez for discussions and Kent Yagi for comments on the manuscript. The research of R. B. and Y. S. was supported by the Israel Science Foundation Grant No. 1294/16. The research of Y. S. was supported by the Negev scholarship. R. B. and Y. S. thank the Theoretical Physics Department, CERN for their hospitality.

Appendix A: Effects of spin

In this appendix, we discuss two effects that depend on the spin of the BH. First, the recent discovery in [51] that magnetic Love numbers of a slowly rotating object should be promoted to tensors and second, we show that the direct effect of the spin on the spectrum of the BH can be neglected, thus justifying the statement in the text that the main effect of the spin is to modify the radius of the BH for the same mass, $R = M(1 + \sqrt{1 - \chi_i^2})$.

Recently, in [51], it was demonstrated that the magnetic Love numbers of a slowly rotating object should be promoted to tensors. For example, following [51], the magnetic Love tensor for $l = 2$ is given by $(k_2)^{ij}_{kl} = (k_2^B)^{ij}_{kl} + (k_2^M)^{ij}_{kl}$. The scalar component $(k_2^B)^{ij}_{kl}$ is related to the magnetic Love number given in Eq. (2.5), $(k_2^B)^{ij}_{kl} = k_2^B \delta_k^i \delta_l^j$. The additional spin induced term is given by

$$(k_2^M)^{ij}_{kl} = - \sum_{n>1, |m|\leq 2} \frac{3}{4R^5} \frac{|\langle \Psi_0 | \hat{S}_{ij}^M | n_r, 2, m \rangle|^2}{\Delta E_{1, n_r}} \times \mathcal{N} (\mathcal{Y}_2^m)^{ij} (\mathcal{Y}_2^m)_{kl}, \quad (\text{A1})$$

where \hat{S}_{ij}^M is the $l = 2$ current moment, $(\mathcal{Y}_2^m)_{kl}$ are azimuthal symmetric-free tensors and \mathcal{N} is a numerical factor that is determined by the orthogonality of the generalized spherical functions (see definitions in [51]). Similarly the magnetic Love tensors for a general l can be obtained.

When the BH is spinning, its spin is coupled to the orbital tidal field. The interaction energy takes the form [85, 86]

$$\begin{aligned} V_{int} &= -Q_{ij} \mathcal{E}^{ij} \\ &= -\lambda_2 (\mathcal{E}_{ij} + 2\alpha \mathcal{B}_{ijk} J^k / M) \mathcal{E}^{ij}, \end{aligned} \quad (\text{A2})$$

where $J^k = M^2 \chi n^k$ is the spin vector (χ is the dimensionless spin parameter), α is a dimensionless coefficient of order unity or less [85] and \mathcal{B}_{ijk} is the $l = 3$ octupolar tidal field.

In this form, since $|\mathcal{B}_{ijk}| \sim M|\mathcal{E}_{ij}|v^3$ it is clear that the spin corrections are 1.5PN order higher than the leading quadrupolar term, and therefore can be neglected.

One can also view this as spin corrections to the quadrupole moment, $\delta Q_{ij} = -\lambda_2 2\alpha \mathcal{B}_{ijk} J^k$, or as spin corrections to the tidal Love number,

$$\lambda_2 \sim \lambda_2^{\chi=0} \left(1 + 2\alpha \left| \frac{\mathcal{B}_{ijk}}{\mathcal{E}_{ij}} \right| \frac{\chi}{M} \right) \sim \lambda_2^{\chi=0} (1 + v^3 \chi). \quad (\text{A3})$$

-
- [1] Pau Amaro-Seoan, H. Audley, S. Babak, J. Baker, E. Barausse, P. Bender, E. Berti, P. Binetruy, M. Born, D. Bortoluzzi, J. Camp, C. Caprini, V. Cardoso, M. Colpi, J. Conklin, N. Cornish, C. Cutler, et al., “Laser Interferometer Space Antenna,” arXiv:1702.00786.
 - [2] B. P. Abbott *et al.* [LIGO Scientific and Virgo], “GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral,” Phys. Rev. Lett. **119**, no.16, 161101 (2017) [arXiv:1710.05832 [gr-qc]].
 - [3] B. P. Abbott *et al.* [LIGO Scientific and Virgo], “GW190425: Observation of a Compact Binary Coalescence with Total Mass $\sim 3.4M_\odot$,” Astrophys. J. Lett. **892**, no.1, L3 (2020) [arXiv:2001.01761 [astro-ph.HE]].
 - [4] P. Amaro-Seoane *et al.* “Low-frequency gravitational-wave science with eLISA/NGO,” Class. Quant. Grav. **29**, 124016 (2012) [arXiv:1202.0839 [gr-qc]].
 - [5] K. S. Thorne, “Multipole Expansions of Gravitational Radiation,” Rev. Mod. Phys. **52**, 299 (1980).
 - [6] L. S. Finn and K. S. Thorne, “Gravitational waves from a compact star in a circular, inspiral orbit, in the equatorial plane of a massive, spinning black hole, as observed by LISA,” Phys. Rev. D **62**, 124021 (2000) [arXiv:gr-qc/0007074 [gr-qc]].
 - [7] E. E. Flanagan and S. A. Hughes, “Measuring gravitational waves from binary black hole coalescences: 1. Signal-to-noise for inspiral, merger, and ringdown,” Phys. Rev. D **57**, 4535-4565 (1998) [arXiv:gr-qc/9701039 [gr-qc]].
 - [8] E. E. Flanagan and T. Hinderer, “Constraining neutron star tidal Love numbers with gravitational wave detectors,” Phys. Rev. D **77**, 021502 (2008) [arXiv:0709.1915 [astro-ph]].
 - [9] T. Hinderer, B. D. Lackey, R. N. Lang and J. S. Read, “Tidal deformability of neutron stars with realistic equations of state and their gravitational wave signatures in binary inspiral,”

- Phys. Rev. D **81**, 123016 (2010) [arXiv:0911.3535 [astro-ph.HE]].
- [10]
- [10] K. Yagi and N. Yunes, “Approximate Universal Relations for Neutron Stars and Quark Stars,” Phys. Rept. **681**, 1-72 (2017) doi:10.1016/j.physrep.2017.03.002 [arXiv:1608.02582 [gr-qc]].
- [11] B. P. Abbott *et al.* [LIGO Scientific and Virgo], “GW170817: Measurements of neutron star radii and equation of state,” Phys. Rev. Lett. **121**, no.16, 161101 (2018) [arXiv:1805.11581 [gr-qc]].
- [12] S. De, D. Finstad, J. M. Lattimer, D. A. Brown, E. Berger and C. M. Biwer, “Tidal Deformabilities and Radii of Neutron Stars from the Observation of GW170817,” Phys. Rev. Lett. **121**, no.9, 091102 (2018) [arXiv:1804.08583 [astro-ph.HE]].
- [13] T. Hinderer, “Tidal Love numbers of neutron stars,” Astrophys. J. **677**, 1216-1220 (2008) [arXiv:0711.2420 [astro-ph]].
- [14] T. Damour and A. Nagar, “Relativistic tidal properties of neutron stars,” Phys. Rev. D **80**, 084035 (2009) [arXiv:0906.0096 [gr-qc]].
- [15] T. Binnington and E. Poisson, “Relativistic theory of tidal Love numbers,” Phys. Rev. D **80**, 084018 (2009) [arXiv:0906.1366 [gr-qc]].
- [16] N. Gürlebeck, “No-hair theorem for Black Holes in Astrophysical Environments,” Phys. Rev. Lett. **114**, no.15, 151102 (2015) [arXiv:1503.03240 [gr-qc]].
- [17] R. A. Porto, “The Tune of Love and the Nature(ness) of Spacetime,” Fortsch. Phys. **64**, no.10, 723-729 (2016) [arXiv:1606.08895 [gr-qc]].
- [18] A. Le Tiec and M. Casals, Phys. Rev. Lett. **126**, no.13, 131102 (2021) doi:10.1103/PhysRevLett.126.131102 [arXiv:2007.00214 [gr-qc]].
- [19] A. Le Tiec, M. Casals and E. Franzin, Phys. Rev. D **103**, no.8, 084021 (2021) doi:10.1103/PhysRevD.103.084021 [arXiv:2010.15795 [gr-qc]].
- [20] H. S. Chia, Phys. Rev. D **104**, no.2, 024013 (2021) doi:10.1103/PhysRevD.104.024013 [arXiv:2010.07300 [gr-qc]].
- [21] V. Cardoso, E. Franzin, A. Maselli, P. Pani and G. Raposo, “Testing strong-field gravity with tidal Love numbers,” Phys. Rev. D **95**, no.8, 084014 (2017) [arXiv:1701.01116 [gr-qc]].
- [22] R. Brustein and A. J. M. Medved, “Quantum hair of black holes out of equilibrium,” Phys. Rev. D **97**, no. 4, 044035 (2018) [arXiv:1709.03566 [hep-th]].
- [23] I. Kourkoulou and J. Maldacena, “Pure states in the SYK model and nearly- AdS_2 gravity,”

- [arXiv:1707.02325 [hep-th]].
- [24] R. Brustein and Y. Zigdon, “Revealing the interior of black holes out of equilibrium in the Sachdev-Ye-Kitaev model,” *Phys. Rev. D* **98**, no. 6, 066013 (2018) [arXiv:1804.09017 [hep-th]].
 - [25] R. Brustein, A. J. M. Medved and K. Yagi, “When black holes collide: Probing the interior composition by the spectrum of ringdown modes and emitted gravitational waves,” *Phys. Rev. D* **96**, no. 6, 064033 (2017) [arXiv:1704.05789 [gr-qc]].
 - [26] S. B. Giddings, “Modulated Hawking radiation and a nonviolent channel for information release,” *Phys. Lett. B* **738**, 92 (2014) [arXiv:1401.5804 [hep-th]].
 - [27] S. B. Giddings, “Nonviolent unitarization: basic postulates to soft quantum structure of black holes,” *JHEP* **1712**, 047 (2017) [arXiv:1701.08765 [hep-th]].
 - [28] R. Brustein and Y. Sherf, “Emission Channels from Perturbed Quantum Black Holes,” *Phys. Rev. D* **100**, no.12, 124005 (2019) [arXiv:1902.08449 [hep-th]].
 - [29] Y. Sherf, “Quantum State of Black-Holes Out of Equilibrium,” arXiv:1911.01236 [hep-th].
 - [30] I. Agullo, V. Cardoso, A. D. Rio, M. Maggiore and J. Pullin, “Potential Gravitational Wave Signatures of Quantum Gravity,” *Phys. Rev. Lett.* **126**, no.4, 041302 (2021) [arXiv:2007.13761 [gr-qc]].
 - [31] S. Datta, “Tidal heating of Quantum Black Holes and their imprints on gravitational waves,” *Phys. Rev. D* **102**, no.6, 064040 (2020) [arXiv:2002.04480 [gr-qc]].
 - [32] A. Almheiri, D. Marolf, J. Polchinski, J. Sully, “Black Holes: Complementarity or Firewalls?,” *JHEP* **1302**, 062 (2013).
 - [33] D. Marolf, J. Polchinski, “Gauge/Gravity Duality and the Black Hole Interior,” *Phys. Rev. Lett.* **111**, 171301 (2013).
 - [34] N. Itzhaki, “Is the black hole complementarity principle really necessary?,” arXiv:hep-th/9607028.
 - [35] S. D. Mathur, “What Exactly is the Information Paradox?,” *Lect. Notes Phys.* **769**, 3 (2009).
 - [36] S. L. Braunstein, S. Pirandola, K. Zyczkowski, “Entangled black holes as ciphers of hidden information,” *Physical Review Letters* **110**, 101301 (2013).
 - [37] S. D. Mathur, “Resolving the black hole causality paradox,” *Gen. Rel. Grav.* **51**, no.2, 24 (2019) [arXiv:1703.03042 [hep-th]].
 - [38] J. Maldacena and L. Susskind, “Cool horizons for entangled black holes,” *Fortsch. Phys.* **61** (2013) 781.

- [39] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian and A. Tajdini, “Replica Wormholes and the Entropy of Hawking Radiation,” JHEP **05**, 013 (2020) [arXiv:1911.12333 [hep-th]].
- [40] R. Brustein, A. J. M. Medved, “Non-Singular Black Holes Interiors Need Physics Beyond the Standard Model,” Fortsch. Phys. **67**, no.10, 1900058 (2019).
- [41] S. D. Mathur, “The Quantum structure of black holes,” Class. Quant. Grav. **23**, R115 (2006).
- [42] K. Skenderis, M. Taylor, “The fuzzball proposal for black holes,” Phys. Rept. **467**, 117 (2008).
- [43] R. Brustein, A. J. M. Medved, “Black holes as collapsed polymers,” Fortsch. Phys. **65**, 0114 (2017) [arXiv:1602.07706].
- [44] R. Brustein and Y. Sherf, “Classical Love for Quantum Blackholes,” Phys. Rev. D **105**, no.2, 024044 (2022) [arXiv:2104.06013 [gr-qc]].
- [45] D. Lai, “Resonant oscillations and tidal heating in coalescing binary neutron stars,” Mon. Not. Roy. Astron. Soc. **270**, 611 (1994) [arXiv:astro-ph/9404062 [astro-ph]].
- [46] W. C. G. Ho and D. Lai, “Resonant tidal excitations of rotating neutron stars in coalescing binaries,” Mon. Not. Roy. Astron. Soc. **308**, 153 (1999) [arXiv:astro-ph/9812116 [astro-ph]].
- [47] S. Chakrabarti, T. Delsate and J. Steinhoff, “Effective action and linear response of compact objects in Newtonian gravity,” Phys. Rev. D **88**, 084038 (2013) [arXiv:1306.5820 [gr-qc]].
- [48] N. Andersson and P. Pnigouras, “Exploring the effective tidal deformability of neutron stars,” Phys. Rev. D **101**, no.8, 083001 (2020) [arXiv:1906.08982 [astro-ph.HE]].
- [49] J. J. Sakurai and J. Napolitano, “Modern Quantum Mechanics,” Boston, USA: Addison-Wesley (2011)
- [50] K. S. Thorne, “Tidal stabilization of rigidly rotating, fully relativistic neutron stars,” Phys. Rev. D **58**, 124031 (1998) [arXiv:gr-qc/9706057 [gr-qc]].
- [51] E. Poisson, “Gravitomagnetic Love tensor of a slowly rotating body: post-Newtonian theory,” Phys. Rev. D **102**, no.6, 064059 (2020) [arXiv:2007.01678 [gr-qc]].
- [52] T. Damour and O. M. Lecian, “On the gravitational polarizability of black holes,” Phys. Rev. D **80**, 044017 (2009) [arXiv:0906.3003 [gr-qc]].
- [53] Poisson, E. and Will, C., Gravity: Newtonian, Post-Newtonian, Relativistic. Cambridge: Cambridge University Press (2014).
- [54] R. Brustein and A. J. M. Medved, “Black holes as collapsed polymers,” Fortsch. Phys. **65**, no.1, 1600114 (2017) [arXiv:1602.07706 [hep-th]].
- [55] K. D. Kokkotas and B. G. Schmidt, “Quasinormal modes of stars and black holes,” Living

- Rev. Rel. **2**, 2 (1999) [arXiv:gr-qc/9909058 [gr-qc]].
- [56] G. Allen, N. Andersson, K. D. Kokkotas and B. F. Schutz, “Gravitational waves from pulsating stars: Evolving the perturbation equations for a relativistic star,” Phys. Rev. D **58**, 124012 (1998) [arXiv:gr-qc/9704023 [gr-qc]].
 - [57] N. Andersson, K. D. Kokkotas and B. F. Schutz, “Space-time modes of relativistic stars,” Mon. Not. Roy. Astron. Soc. **280**, 1230 (1996) [arXiv:gr-qc/9601015 [gr-qc]].
 - [58] E. Maggio, V. Cardoso, S. R. Dolan and P. Pani, “Ergoregion instability of exotic compact objects: electromagnetic and gravitational perturbations and the role of absorption,” Phys. Rev. D **99**, no.6, 064007 (2019) [arXiv:1807.08840 [gr-qc]].
 - [59] J. D. Bekenstein and V. F. Mukhanov, “Spectroscopy of the quantum black hole,” Phys. Lett. B **360**, 7-12 (1995) [arXiv:gr-qc/9505012 [gr-qc]].
 - [60] M. Maggiore, “The Physical interpretation of the spectrum of black hole quasinormal modes,” Phys. Rev. Lett. **100**, 141301 (2008) [arXiv:0711.3145 [gr-qc]].
 - [61] V. F. Foit and M. Kleban, “Testing Quantum Black Holes with Gravitational Waves,” Class. Quant. Grav. **36**, no.3, 035006 (2019) [arXiv:1611.07009 [hep-th]].
 - [62] V. Cardoso, V. F. Foit and M. Kleban, “Gravitational wave echoes from black hole area quantization,” JCAP **08**, 006 (2019) [arXiv:1902.10164 [hep-th]].
 - [63] A. Maselli, P. Pani, V. Cardoso, T. Abdelsalhin, L. Gualtieri and V. Ferrari, “Probing Planckian corrections at the horizon scale with LISA binaries,” Phys. Rev. Lett. **120**, no.8, 081101 (2018) [arXiv:1703.10612 [gr-qc]].
 - [64] A. Maselli, P. Pani, V. Cardoso, T. Abdelsalhin, L. Gualtieri and V. Ferrari, “From micro to macro and back: probing near-horizon quantum structures with gravitational waves,” Class. Quant. Grav. **36**, no.16, 167001 (2019) [arXiv:1811.03689 [gr-qc]].
 - [65] C. Cutler and E. E. Flanagan, “Gravitational waves from merging compact binaries: How accurately can one extract the binary’s parameters from the inspiral wave form?,” Phys. Rev. D **49**, 2658-2697 (1994) [arXiv:gr-qc/9402014 [gr-qc]].
 - [66] K. Yagi and N. Yunes, “Love can be Tough to Measure,” Phys. Rev. D **89**, no.2, 021303 (2014) [arXiv:1310.8358 [gr-qc]].
 - [67] N. Sennett, T. Hinderer, J. Steinhoff, A. Buonanno and S. Ossokine, “Distinguishing Boson Stars from Black Holes and Neutron Stars from Tidal Interactions in Inspiring Binary Systems,” Phys. Rev. D **96**, no.2, 024002 (2017) [arXiv:1704.08651 [gr-qc]].

- [68] E. Poisson and M. Sasaki, “Gravitational radiation from a particle in circular orbit around a black hole. 5: Black hole absorption and tail corrections,” *Phys. Rev. D* **51**, 5753-5767 (1995) [arXiv:gr-qc/9412027 [gr-qc]].
- [69] K. Alvi, “Energy and angular momentum flow into a black hole in a binary,” *Phys. Rev. D* **64**, 104020 (2001) [gr-qc/0107080].
- [70] E. Poisson, “Absorption of mass and angular momentum by a black hole: Time-domain formalisms for gravitational perturbations, and the small-hole / slow-motion approximation,” *Phys. Rev. D* **70**, 084044 (2004) [gr-qc/0407050].
- [71] S. Taylor and E. Poisson, “Nonrotating black hole in a post-Newtonian tidal environment,” *Phys. Rev. D* **78**, 084016 (2008) [arXiv:0806.3052 [gr-qc]].
- [72] T. Robson, N. J. Cornish and C. Liu, “The construction and use of LISA sensitivity curves,” *Class. Quant. Grav.* **36**, no.10, 105011 (2019) [arXiv:1803.01944 [astro-ph.HE]].
- [73] E. Berti, A. Buonanno and C. M. Will, “Estimating spinning binary parameters and testing alternative theories of gravity with LISA,” *Phys. Rev. D* **71**, 084025 (2005) [arXiv:gr-qc/0411129 [gr-qc]].
- [74] M. Favata, “Conservative corrections to the innermost stable circular orbit (ISCO) of a Kerr black hole: A New gauge-invariant post-Newtonian ISCO condition, and the ISCO shift due to test-particle spin and the gravitational self-force,” *Phys. Rev. D* **83**, 024028 (2011) [arXiv:1010.2553 [gr-qc]].
- [75] T. Damour, B. R. Iyer and B. S. Sathyaprakash, “A Comparison of search templates for gravitational waves from binary inspiral,” *Phys. Rev. D* **63**, 044023 (2001) [arXiv:gr-qc/0010009 [gr-qc]].
- [76] K. G. Arun, B. R. Iyer, B. S. Sathyaprakash and P. A. Sundararajan, “Parameter estimation of inspiralling compact binaries using 3.5 post-Newtonian gravitational wave phasing: The Non-spinning case,” *Phys. Rev. D* **71**, 084008 (2005) [arXiv:gr-qc/0411146 [gr-qc]].
- [77] A. Buonanno, B. Iyer, E. Ochsner, Y. Pan and B. S. Sathyaprakash, “Comparison of post-Newtonian templates for compact binary inspiral signals in gravitational-wave detectors,” *Phys. Rev. D* **80**, 084043 (2009) [arXiv:0907.0700 [gr-qc]].
- [78] S. Isoyama and H. Nakano, “Post-Newtonian templates for binary black-hole inspirals: the effect of the horizon fluxes and the secular change in the black-hole masses and spins,” *Class. Quant. Grav.* **35**, no.2, 024001 (2018) [arXiv:1705.03869 [gr-qc]].

- [79] S. Khan, S. Husa, M. Hannam, F. Ohme, M. Pürrer, X. Jiménez Forteza and A. Bohé, “Frequency-domain gravitational waves from nonprecessing black-hole binaries. II. A phenomenological model for the advanced detector era,” *Phys. Rev. D* **93**, no.4, 044007 (2016) [arXiv:1508.07253 [gr-qc]].
- [80] D. Bini, T. Damour and G. Faye, “Effective action approach to higher-order relativistic tidal interactions in binary systems and their effective one body description,” *Phys. Rev. D* **85**, 124034 (2012) [arXiv:1202.3565 [gr-qc]].
- [81] K. Hotokezaka, K. Kyutoku, Y. i. Sekiguchi and M. Shibata, “Measurability of the tidal deformability by gravitational waves from coalescing binary neutron stars,” *Phys. Rev. D* **93**, no.6, 064082 (2016) [arXiv:1603.01286 [gr-qc]].
- [82] V. Cardoso and P. Pani, “Testing the nature of dark compact objects: a status report,” *Living Rev. Rel.* **22**, no.1, 4 (2019) [arXiv:1904.05363 [gr-qc]].
- [83] Marković, Dragoljub, ”Possibility of determining cosmological parameters from measurements of gravitational waves emitted by coalescing, compact binaries”, 1993, *Phys. Rev. D*, 48, 4738
- [84] S. A. Hughes, “Untangling the merger history of massive black holes with LISA,” *Mon. Not. Roy. Astron. Soc.* **331**, 805 (2002) doi:10.1046/j.1365-8711.2002.05247.x [arXiv:astro-ph/0108483 [astro-ph]].
- [85] P. Pani, L. Gualtieri and V. Ferrari, “Tidal Love numbers of a slowly spinning neutron star,” *Phys. Rev. D* **92**, no.12, 124003 (2015) [arXiv:1509.02171 [gr-qc]].
- [86] T. Abdelsalhin, L. Gualtieri and P. Pani, “Post-Newtonian spin-tidal couplings for compact binaries,” *Phys. Rev. D* **98**, no.10, 104046 (2018) [arXiv:1805.01487 [gr-qc]].