Three solutions for a new Kirchhoff-type problem*

Yue Wang[1], Hong-Min Suo[2†]

E-mail: yuewn@sina.cn

1. School of Mathematics and Statistics, Guizhou University, Guiyang 550025*, People's Republic of China*

2. School of Data Science and Information Engineering, Guizhou Minzu University, Guiyang 550025*, People's Republic of China*

arXiv:2008.02838v1 [math.AP] 6 Aug 2020

Abstract: This article concerns on the existence of multiple solutions for a new Kirchhoff-type problem with negative modulus. We prove that there exist three nontrivial solutions when the parameter is enough small via the variational methods and algebraic analysis. Moreover, our fundamental technique is one of the Mountain Pass Lemma, Ekeland variational principle, and Minimax principle.

Keywords: New Kirchhoff-type problem; variational method; algebraic analysis

MSC(2010): 35A15, 35B33, 35J62

1 Introduction and main results

This paper research the following nonlocal problem

$$\begin{cases} -\Big(a - b\displaystyle\int_\Omega |\nabla u|^2 dx\Big)\Delta u = \mu f(x), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \tag{1.1}$$

where $a, b > 0$ are constants, $\mu \in \mathbb{R}$ is a parameter, $f(x)$ is a nonnegative-nonzero function, and Ω is a bounded domain in $\mathbb{R}^N (N \geq 1)$ with smooth boundary $\partial\Omega$. The analysis developed in this paper corresponds to propose an approach based on the idea of considering the nonlocal term with negative modulus which is presents interesting difficulties.

Problem (1.1) is related with the Kirchhoff-type equation as following:

$$-\Big(a + b\int_\Omega |\nabla u|^2 dx\Big)\Delta u = f(x,u), \text{ in } \Omega, \tag{1.2}$$

where $a \geq 0$, $b > 0$, Ω is a bounded domain in $\mathbb{R}^N$ with smooth boundary $\partial\Omega$ or $\Omega = \mathbb{R}^N$, $f : \Omega \times \mathbb{R} \to \mathbb{R}$ is a continual function. Mentioned that Eq.(1.2) is a subproblem of model

$$\varrho h\frac{\partial^2 u}{\partial t^2} - \Big(p_0 + \frac{Eh}{2L}\int_0^L \Big|\frac{\partial u}{\partial x}\Big|^2 dx\Big)\frac{\partial^2 u}{\partial x^2} = f(x,u) \tag{1.3}$$

with $0 < x < L$ and $t \geq 0$, where $u = u(x,t)$ is the lateral displacement, ϱ the mass density, E the Young's modulus, h the cross-section area, L the length, p_0 the initial axial tension. Eq.(1.3) named the Kirchhoff problem as an extension of classical D'Alembert's wave equations for free vibration of elastic strings by Kirchhoff [1] in 1876. When finding the existence of stationary solution, Eq.(1.3) may be express as the Eq.(1.2) and therefore problem (1.2) were named the Kirchhoff-type problem. Eq.(1.2) has been studied by many researchers on whole space $\mathbb{R}^N$ and

*Supported by the NSF of China (11661021,11861021).

†Corresponding author. E-mail:eztf@qq.com.

bounded domain with some boundary conditions, such as [2–12] and their references. Problem (1.2) contains a nonlocal coefficient $(a + b\int_\Omega |\nabla u|^2 dx)$, this leads to that Eq.(1.2) is no longer a pointwise identity and therefore it is often called nonlocal problem. It is worth paying more attentions to Young's modulus, which can also be used in computing tension, when the atoms are pulled apart instead of squeezed together, the strain is negative because the atoms are stretched instead of compressed, this leads to minus Young's modulus. Therefore, a nonlocal coefficient $(a - b\int_\Omega |\nabla u|^2 dx)$ is included in Kirchhoff-type problem, may be an interesting model(see [13]), which is the reason why we call it Kirchhoff-type problem with negative modulus in the abstract.

Indeed, let $a, b > 0$, Yin and Liu in [14] got that there exist at least a nontrivial non-negative solution and a nontrivial non-positive solution for the following problem

$$\begin{cases} -\Big(a - b\displaystyle\int_\Omega |\nabla u|^2 dx\Big)\Delta u = |u|^{p-2}u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded domain in $\mathbb{R}^N$ with smooth boundary, $2 < p < 2^*$, $2^* = \frac{2N}{N-2}$ as $N \geq 3$ and $2^* = +\infty$ as $N = 1, 2$. In addition, infinitely many solutions were got in [15] included the condition above and $p = 2$ via the Ljusternik-Schnirelman type minimax method. Wang et al. [13] obtained that there exist at least two positive solutions when $f(x) \in L^{4/3}(\mathbb{R}^4)$ with $\mu > 0$ small and infinitely many positive solutions with $\mu = 0$ via variational method mainly for the problem

$$-\Big(a - b\int_{\mathbb{R}^4} |\nabla u|^2 dx\Big)\Delta u = |u|^2 u + \mu f(x), \quad \text{in } \mathbb{R}^4.$$

The research interesting in [14] is that the nonlocal coefficient $(a + b\int_\Omega |\nabla u|^2 dx)$ is bounded below but the nonlocal coefficient $(a - b\int_\Omega |\nabla u|^2 dx)$ is not. Different with [14], the research interesting in [13] is that the Kirchhoff-type equation with nonlocal coefficient $(a - b\int_\Omega |\nabla u|^2 dx)$ is a negative modulus problem. For more details about the Kirchhoff problems with negative modulus, we refer readers to [16–30]. Enlightened by the papers above, we consider the problem (1.1), the conclusion state as following theories main via variational method and algebraic analysis.

Theorem 1.1. *Assume that $a, b > 0$ and $f(x) \in L^{\frac{2^*}{2^*-1}}(\Omega)$ is positive a.e., then, there exists $\mu_* > 0$, such that the problem (1.1) has at least three nontrivial solutions for $\mu \in (0, \mu_*)$ and a nontrivial solution for $\mu \in [\mu_*, +\infty)$.*

As the proof as Theorem 1.1, the proof of the existence of solutions for $\mu < 0$ may be miscellaneous by variational method, we shall use new method to overcome the difficulties.

Theorem 1.2. *Assume that $a, b > 0$ and $f(x) \in L^{\frac{2^*}{2^*-1}}(\Omega)$ is positive a.e., then, there exists a constant $\mu_{**} > 0$ such that problem (1.1) has only three solutions for $0 < |\mu| < \mu_{**}$, only two solutions for $\mu = \pm\mu_{**}$ and unique solution for $|\mu| > \mu_{**}$. Moreover, problem (1.1) has infinitely many solutions for $a, b > 0, \mu = 0$.*

Indeed, the condition of $f(x)$ can be replace weakly by $f \in L^2(\Omega)$ and $f(x) > 0$ a.e..

Corollary 1.3. *Assume that $a, b > 0$ and $f(x) \in L^2(\Omega)$ is positive a.e., then, for μ_{**} defined by Theorem 1.2, problem (1.1) has only three solutions for any $0 < |\mu| < \mu_{**}$, only two solutions for $\mu = \pm\mu_{**}$ and unique solution for any $|\mu| > \mu_{**}$.*

The novelty of our results lies in three aspects. Firstly, different with those articles mentioned above, we prove the $(PS)_c$ condition by new method and $c \in [-\frac{a^2}{12b}, +\infty) \setminus \{\frac{a^2}{4b}\}$, what's more, it is not satisfy $(PS)_c$ condition with $c \in (-\infty, -\frac{a^2}{12b}) \cup \{\frac{a^2}{4b}\}$. Secondly, with the help of algebraic analysis, we got that there are at least three nontrivial weak solutions by using one of the Mountain

Pass Lemma, Ekeland variational principle, and Minimax principle. Thirdly, through of a basic fundamental result, we obtain the specific form of μ_{**}.

This article is organized as follows. In section 2, we give some basic knowledge which use to solving the problem. Section 3 contains elementary results and proof of theorem 1.1. In section 4, for the Theorem 1.1, with the help of algebraic analysis, we prove that the existence of three nontrivial weak solutions only by using one of the Mountain Pass Lemma, Ekeland variational principle, and Minimax principle. By using the similar method, we prove main results of the Theorem 1.2 and calculate the μ_{**} exactly.

2 Preliminaries

Throughout this paper we denote by $\to$ (resp. $\rightharpoonup$) the strong (resp. weak) convergence. For any $u, v \in H_0^1(\Omega)$, the inner product is $\langle u, v\rangle = \int_\Omega \nabla u \nabla v dx$ and the norm $\|u\| = \big(\int_\Omega |\nabla u|^2 dx\big)^{\frac{1}{2}}$.

We recall that a function $u \in H_0^1(\Omega)$ is called a weak solution of problem (1.1) if

$$\Big(a - b\int_\Omega |\nabla u|^2 dx\Big)\int_\Omega \nabla u \nabla v dx = \mu \int_\Omega f v dx, \quad \forall\ v \in H_0^1(\Omega).$$

Denote the L^s-norm $\|\cdot\|_s = \big[\int_\Omega |u|^s dx\big]^{\frac{1}{s}}$ for $0 < s < +\infty$ and $\|\cdot\|_\infty = ess\sup\limits_{x\in\Omega} |u|$ (still denoted by $\|\cdot\|_s = \big[\int_\Omega |u|^s dx\big]^{\frac{1}{s}}$) for $s = +\infty$. Set

$$S = \inf_{u\in H_0^1(\Omega)\setminus\{0\}} \frac{\int_\Omega |\nabla u|^2 dx}{\big(\int_\Omega |u|^{2^*} dx\big)^{\frac{2}{2^*}}}, \quad \lambda_1 = \inf_{u\in H_0^1(\Omega)\setminus\{0\}} \frac{\int_\Omega |\nabla u|^2 dx}{\int_\Omega |u|^2 dx}.$$

Let $I(u) : H_0^1(\Omega) \mapsto \mathbb{R}$ be the functional defined by

$$I(u) = \frac{a}{2}\|u\|^2 - \frac{b}{4}\|u\|^4 - \mu\int_\Omega f u dx, \tag{2.1}$$

it can be verify that $I(u) \in C(H_0^1(\Omega), \mathbb{R})$ and the Gâteaux derivative of I given by

$$\langle I'(u), v\rangle = (a - b\|u\|^2)\int_\Omega \nabla u \nabla v dx - \mu\int_\Omega f v dx, \quad \forall\ v \in H_0^1(\Omega). \tag{2.2}$$

If $u \in H_0^1(\Omega)$ such that $I'(u) = 0$, then u is a weak solution of problem (1.1).

3 Proof of the Theorem 1.1

Lemma 3.1. *Assume that $a, b, \mu > 0$, $f \in L^{\frac{2^*}{2^*-1}}(\Omega)$ and $f(x) \geq 0$ a.e., then, I satisfies the $(PS)_c$ condition with $c \in [-\frac{a^2}{12b}, +\infty) \setminus \{\frac{a^2}{4b}\}$, and I does not satisfy the $(PS)_c$ condition at $c = \frac{a^2}{4b}$. That is, for $c \in [-\frac{a^2}{12b}, +\infty) \setminus \{\frac{a^2}{4b}\}$, every (PS) sequence at c has a converge subsequence.*

Proof. Recalling that $\{u_n\} \subset H_0^1(\Omega)$ is a (PS) sequence at c, so that $I(u_n) \to c$ and $I'(u_n) \to 0$ as $n \to \infty$. Let $\{u_n\}$ be a $(PS)_c$ sequence, then, for all $v \in H_0^1(\Omega)$, there holds

$$\begin{cases} I(u_n) = \dfrac{a}{2}\|u_n\|^2 - \dfrac{b}{4}\|u_n\|^4 - \mu\displaystyle\int_\Omega f u_n dx \to c, \\ \langle I'(u_n), v\rangle = \big(a - b\|u_n\|^2\big)\displaystyle\int_\Omega \nabla u_n \nabla v dx - \mu\int_\Omega f v dx \to 0 \end{cases} \tag{3.1}$$

as $n \to \infty$. Especially, taking $v = u_n$ in (3.1), it holds that

$$I(u_n) - \langle I'(u_n), u_n \rangle = \frac{3b}{4}\|u_n\|^4 - \frac{a}{2}\|u_n\|^2 \to c. \tag{3.2}$$

Hence we can obtain that, there is

$$\lim_{n\to\infty} \|u_n\|^2 = \frac{\frac{a}{2} \pm \sqrt{\frac{a^2}{4} + 3bc}}{2 \times \frac{3b}{4}} = \frac{a \pm a\sqrt{1 + \frac{12bc}{a^2}}}{3b} = \frac{a}{3b}\left(1 \pm \sqrt{1 + \frac{12bc}{a^2}}\right).$$

From the above, we have

$$\begin{cases} \lim\limits_{n\to\infty} \|u_n\|^2 < 0, & \text{if} \quad c < -\dfrac{a^2}{12b}; \\ 0 \le \lim\limits_{n\to\infty} \|u_n\|^2 \le \dfrac{2a}{3b}, & \text{if} \quad -\dfrac{a^2}{12b} \le c \le 0; \\ \dfrac{2a}{3b} < \lim\limits_{n\to\infty} \|u_n\|^2 < \dfrac{a}{b}, & \text{if} \quad 0 < c < \dfrac{a^2}{4b}; \\ \lim\limits_{n\to\infty} \|u_n\|^2 = \dfrac{a}{b}, & \text{if} \quad c = \dfrac{a^2}{4b}; \\ \lim\limits_{n\to\infty} \|u_n\|^2 > \dfrac{a}{b}, & \text{if} \quad c > \dfrac{a^2}{4b}. \end{cases} \tag{3.3}$$

Therefore, $c \ge -\frac{a^2}{12b}$ is well defined from (3.3), and there exists no $\lim\limits_{n\to\infty} \|u_n\|^2$ for $c < -\frac{a^2}{12b}$.

Case $\lim\limits_{n\to\infty} \|u_n\|^2 = \frac{a}{b}$, we have $a - b\|u_n\|^2 \to 0$ as $n \to \infty$. By (3,1), we can choose a $v \in H_0^1(\Omega)$ such that $\int_\Omega f v dx = 1$, it holds that $\|v\| < +\infty$ and

$$\left(a - b\|u_n\|^2\right) \int_\Omega \nabla u_n \nabla v dx \to \mu,$$

this leads to that

$$\lim_{n\to\infty} \|u_n\|\|v\| \ge \lim_{n\to\infty} \int_\Omega \nabla u_n \nabla v dx = \mu \lim_{n\to\infty} \left(a - b\|u_n\|^2\right)^{-1} = +\infty,$$

and therefore $\lim\limits_{n\to\infty} \|u_n\| \to +\infty$. Which is a contradiction with $\|u_n\|^2 \to \frac{a}{b}$. So, (3.1) does not hold with all $v \in H_0^1(\Omega)$ and I has no $(PS)_c$ sequence with $c = \frac{a^2}{4b}$.

Case $\lim\limits_{n\to\infty} \|u_n\|^2 \ne \frac{a}{b}$, we have $a - b\|u_n\|^2 \not\to 0$. We declare that $\{u_n\}$ is bounded in $H_0^1(\Omega)$. Otherwise, we can assume that $\|u_n\| \to +\infty$ as $n \to \infty$. In this case, we have

$$0 = \lim_{n\to\infty} \frac{c}{\|u_n\|^4} = \lim_{n\to\infty} I(u_n) = -\frac{b}{4},$$

which is a contradiction by the uniqueness of limit. So $\{u_n\}$ is bounded in $H_0^1(\Omega)$. That is, if necessary, pass to a subsequence (still denoted by $\{u_n\}$) and u_0 in $H_0^1(\Omega)$ such that, for $n \to \infty$,

$$\begin{cases} u_n \rightharpoonup u_0, & \text{weakly in} \quad H_0^1(\Omega), \\ u_n \to u_0, & \text{strongly in} \quad L^q_{\text{loc}}(1 \le q < 2^*), \\ u_n(x) \to u_0(x), & \text{a.e.} \quad x \in \Omega. \end{cases}$$

Hence there is

$$\begin{cases} \langle I'(u_n), u_n \rangle = \left(a - b\|u_n\|^2\right) \int_\Omega \nabla u_n \nabla u_n dx - \mu \int_\Omega f u_n dx \to 0, \\ \langle I'(u_n), u_0 \rangle = \left(a - b\|u_n\|^2\right) \int_\Omega \nabla u_n \nabla u_0 dx - \mu \int_\Omega f u_0 dx \to 0. \end{cases} \tag{3.4}$$

Lebesgue's dominated convergence theorem (see [31, pp.27]) leads to

$$\int_\Omega f u_n dx = \int_\Omega f u_0 dx + o(1). \tag{3.5}$$

From (3.4) and (3.5), we have

$$\big(a-b\|u_n\|^2\big)\int_\Omega \nabla u_n\nabla(u_n-u_0)dx\to 0.$$

Based on $a-b\|u_n\|^2\not\to 0$, one has $u_n\to u_0$ in $H_0^1(\Omega)$. This proof is complete. □

In order to prove main results, it is necessary to make some notes, that is

$$\mathcal{D}^+=\left\{u\in H_0^1(\Omega)\,|\,\int_\Omega fudx>0\right\},\quad \mathcal{D}^-=\left\{u\in H_0^1(\Omega)\,|\,\int_\Omega fudx<0\right\}.$$

Setting $u\in\mathcal{D}^\pm$, then, there is $tu\in\mathcal{D}^\pm$ with $t>0$ and $tu\in\mathcal{D}^\mp$ with $t<0$. For any $u\in\mathcal{D}^\pm$, it is easy to see that $|\int_\Omega fudx|>0$, and therefore, we shall let

$$\Lambda:=\min_{u\in\mathcal{D}^+\bigcup\mathcal{D}^-}\left\{\|u\|^{-1}\Big|\int_\Omega fudx\Big|\right\}=\min_{u\in\mathcal{D}^+}\left\{\|u\|^{-1}\int_\Omega fudx\right\}\le\frac{1}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}.$$

Lemma 3.2. *Assume that $a,b>0$, $f\in L^{\frac{2^*}{2^*-1}}(\Omega)$ and $f(x)\ge 0$ a.e., $u\in H_0^1(\Omega)$, then,*

(i) there exist $r,\rho,\mu_{*1}>0$, such that for any $\mu\in(0,\mu_{*1}]$, it holds

$$\inf_{\|u\|=r}I(u)\ge\rho\quad\text{and}\quad\inf_{\|u\|\le r}I(u):=c_1<0;$$

(ii) There exists $R>0$, such that $\sup_{\|u\|\ge R}I(u)\le 0$ for any $\mu>0$.

Proof. (i) Togethering with the functional (2.1) with Sobolev imbedding inequality, we obtain

$$I(u)\ge\frac{a}{2}\|u\|^2-\frac{b}{4}\|u\|^4-\frac{\mu}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}\|u\|=\|u\|\Big(\frac{a}{2}\|u\|-\frac{b}{4}\|u\|^3-\frac{\mu}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}\Big).$$

We can see that for any $\mu<\frac{a}{9b}(6abS)^{\frac12}\|f\|^{-1}_{\frac{2^*}{2^*-1}}:=\mu_0$ with $r_0=(\frac{2a}{3b})^{\frac12}$, there is

$$\inf_{\|u\|=r_0,\ \mu<\mu_0}I(u)\ge\Big(\frac{2a}{3b}\Big)^{\frac12}\Big[\frac{a}{3}\Big(\frac{2a}{3b}\Big)^{\frac12}-\frac{\mu}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}\Big]:=\rho_0>0.$$

Therefore, in order to calculate its conveniently, it is easy to see that there exist $r=(\frac{2a}{3b})^{\frac12}$, $\rho=\frac{a^2}{9b}$, $\mu_{*1}=\frac{a}{18b}(6abS)^{\frac12}\|f\|^{-1}_{\frac{2^*}{2^*-1}}$, such that, for any $\mu\in(0,\mu_{*1}]$, one has

$$\begin{aligned}\inf_{\|u\|=r,\ \mu\le\mu_*}I(u)&\ge\Big(\frac{2a}{3b}\Big)^{\frac12}\Big[\frac{a}{3}\Big(\frac{2a}{3b}\Big)^{\frac12}-\frac{\mu}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}\Big]\ge\Big(\frac{2a}{3b}\Big)^{\frac12}\Big[\frac{a}{3}\Big(\frac{2a}{3b}\Big)^{\frac12}-\frac{\mu_*}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}\Big]\\&\ge\Big(\frac{2a}{3b}\Big)^{\frac12}\Big[\frac{a}{3}\Big(\frac{2a}{3b}\Big)^{\frac12}-\frac{a}{18b}(6abS)^{\frac12}\|f\|^{-1}_{\frac{2^*}{2^*-1}}\cdot\frac{1}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}\Big]\\&=\Big(\frac{2a}{3b}\Big)^{\frac12}\Big[\frac{a}{3}\Big(\frac{2a}{3b}\Big)^{\frac12}-\frac{a}{6}\Big(\frac{2a}{3b}\Big)^{\frac12}\Big]=\frac{a^2}{9b}=\rho.\end{aligned}$$

Taking $\tilde u\in\mathcal{D}^+\subset H_0^1(\Omega)$ with $\|\tilde u\|=r$, it holds $\|t\tilde u\|\le r$ with $t\to 0^+$, and that

$$\begin{aligned}\lim_{t\to0^+}\frac{I(t\tilde u)}{t}&=\lim_{t\to0^+}\frac1t\Big\{\frac{a}{2}\|tu\|^2-\frac{b}{4}\|tu\|^4-\mu\int_\Omega f\cdot(tu)dx\Big\}\\&=-\mu\int_\Omega f\tilde u dx=-\frac{\mu r}{\|\tilde u\|}\int_\Omega f\tilde u dx\\&\le-\mu r\Lambda=-(\frac{2a}{3b})^{\frac12}\Lambda\mu<0.\end{aligned}$$

Therefore, $c_1 := \inf_{\|u\|\le r} I(u) \le -(\frac{2a}{3b})^{\frac{1}{2}}\Lambda\mu < 0$ is well defined.

(ii) For any $\mu > 0$, by Young's inequality, we have

$$\begin{aligned} I(u) &\le \frac{a}{2}\|u\|^2 - \frac{b}{4}\|u\|^4 + \mu\Big|\int_\Omega f u dx\Big| \\ &\le \frac{a}{2}\|u\|^2 - \frac{b}{4}\|u\|^4 + \frac{a}{2}\|u\|^2 + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}} \\ &= a\|u\|^2 - \frac{b}{4}\|u\|^4 + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}}. \end{aligned} \tag{3.6}$$

So, there exists $R = \frac{2}{b}\Big[a + \big(a^2 + \frac{b\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}}\big)^{\frac{1}{2}}\Big]$, such that $I(u) \le 0$ with $\|u\| \ge R$, thus the conclusion $\sup_{\|u\|\ge R} I(u) \le 0$ can be true. This completes the proof. □

3.1 Proof of the first and second solutions of problem (1.1)

Theorem 3.3. Assume that $a, b > 0$ and $f(x) \in L^{\frac{2^*}{2^*-1}}(\Omega)$ is a positive function a.e., then, for any $\mu \in (0, \mu_*)$(μ_* is defined by lemma 3.2), the problem (1.1) has at least two positive solutions.

Existence of the first solution. Taking $B_r := \{u \in H_0^1(\Omega) \mid \|u\| \le r\}$, where $r = (\frac{2a}{3b})^{\frac{1}{2}}$. By the Lemma 3.2, there exists $\mu_* > 0$ such that $\inf I(B_r) \le -(\frac{2a}{3b})^{\frac{1}{2}}\Lambda\mu < 0$ for any $\mu \in (0, \mu_*)$. By the Ekeland variational principle (see[32, Lemma 1.1]), there is a minimizing sequence $\{u_n\} \subset \overline{B_r}$ such that

$$I(u_n) \le \inf I(B_r) + \frac{1}{n} \quad \text{and} \quad I(u) \ge I(u_n) - \frac{1}{n}\|u - u_n\| \tag{3.7}$$

for all $n \in \mathbb{N}$ and for any $u \in \overline{B_r}$. Therefore, $I(u_n) \to c_1$ and $I'(u_n) \to 0$ in dual space of $H_0^1(\Omega)$ as $n \to \infty$. By Lemma 3.1, there exist a subsequence (still denoted by $\{u_n\}$) and $u_1^* \in B_r$ such that $u_n \to u_1^*$ as $n \to \infty$. Then, $I(u_1^*) = c_1 < 0$ and $I'(u_1^*) = 0$. Moreover, $u_1^* \in \mathcal{D}^+$ and $\|u_1^*\|^2 < \frac{a}{3b}$. This implies that u_1^* is a local minimizer for I. Hence u_1^* is a weak solution of problem (1.1). □

Existence of the second solution. For any $\bar{u} \in \mathcal{D}^+$, there exists $\theta \in (0, 1)$, such that

$$\begin{aligned} \sup I(\bar{u}) &= \sup\Big\{\frac{a}{2}\|\bar{u}\|^2 - \frac{b}{4}\|\bar{u}\|^4 - \mu\int_\Omega f\bar{u}dx\Big\} \\ &\le \sup\Big\{\frac{a}{2}\|\bar{u}\|^2 - \frac{b}{4}\|\bar{u}\|^4 - \mu\Lambda\|\bar{u}\|\Big\} \\ &\le \frac{a^2}{4b} - \theta(\frac{a}{b})^{\frac{1}{2}}\mu\Lambda < \frac{a^2}{4b}. \end{aligned}$$

According to Lemma 3.2, the functional I has mountain pass geometry for any $\mu \in (0, \mu_*)$. For any $e \in H_0^1(\Omega)$ with $\|e\| \ge R\}$(where R defined in the lemma 3.2), we set

$$\Gamma = \Big\{\tau(t) \in C^1\big([0,1], H_0^1(\Omega)\big); \tau(0) = 0, \tau(1) = e\Big\}.$$

By (2.1)–(2.2), $I\big(\tau(t)\big)$ has continuity, $I\big(\tau(0)\big) = 0$, $I\big(\tau(1)\big) \le 0$. So, there is

$$0 < \rho \le c_2 := \inf_{\tau\in\Gamma}\sup_{t\in[0,1]} I\big(\tau(t)\big) \le \sup_{u\in\mathcal{D}^+} I(u) < \frac{a^2}{4b}.$$

By Lemma 3.1 and the mountain pass theorem (see [33, Theorem 2.1–2.4]), there exist u_2^* and a sequence $\{u_k\}$ in $H_0^1(\Omega)$, moreover in $\mathcal{D}^+$, such that $u_k \to u_2^*$ in $H_0^1(\Omega)$, $I(u_k) \to c_2 = I(u_2^*)$ and $I'(u_k) \to 0 = I'(u_2^*)$ in dual space of $H_0^1(\Omega)$. Hence u_2^* is a weak solution of problem (1.1) with $\|u_2^*\| > (\frac{2a}{3b})^{\frac{1}{2}}$. Since $I(u_1^*) < 0 < I(u_2^*)$, we get $u_2^* \ne u_1^*$. □

Proof of that u_1^* and u_2^* are positive. Since $u_i^* \in \mathcal{D}^+ (i = 1, 2)$ are the weak solutions of problem (1.1), we have

$$\Big(a - b\int_\Omega |\nabla u_i^*|^2 dx\Big)\int_\Omega |\nabla u_i^*|^2 dx = \mu \int_\Omega f u_i^* dx > 0.$$

Hence $a - b\|u_i^*\|^2 > 0$. This means that $-\Delta u_i^* = \mu(a - b\|u_i^*\|^2)^{-1} f(x) \geq 0$ and $u_i^* \not\equiv 0$. According to the strong maximum principle, we obtain u_i^* are positive solutions. □

3.2 Proof of the third solution of problem (1.1)

Lemma 3.4. *Assume that $a, b, \mu > 0$, $f \in L^{\frac{2^*}{2^*-1}}(\Omega)$ and $f(x) \geq 0$ a.e., it holds that*

$$\frac{a^2}{4b} < \sup I(u) \leq \frac{a^2}{b} + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}}.$$

Proof. For any $u \in \mathcal{D}^+$, $tu \in \mathcal{D}^-$ with $t < 0$ and we have

$$\begin{aligned}
\sup_{t<0} I(tu) &\geq \sup_{t=-(\frac{a}{b})^{\frac{1}{2}}\|u_0\|^{-1}} \left\{\frac{a}{2}\|tu\|^2 - \frac{b}{4}\|tu\|^4 - \mu\int_\Omega f\cdot(tu)dx\right\} \\
&= \frac{a^2}{4b} + (\frac{a}{b})^{\frac{1}{2}}\frac{\mu}{\|u\|}\left|\int_\Omega f u_0 dx\right| \\
&\geq \frac{a^2}{4b} + \mu\Lambda(\frac{a}{b})^{\frac{1}{2}} > \frac{a^2}{4b}.
\end{aligned} \tag{3.8}$$

Therefore, $\sup I(u) \geq \sup_{u\in\mathcal{D}^-} I(u) > \frac{a^2}{4b}$. Moreover, via the Young's inequality we obtain that

$$\begin{aligned}
\sup I(u) &\leq \frac{a}{2}\|u\|^2 - \frac{b}{4}\|u\|^4 + \mu\Big|\int_\Omega f u dx\Big| \\
&\leq \frac{a}{2}\|u\|^2 - \frac{b}{4}\|u\|^4 + \frac{a}{2}\|u\|^2 + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}} \\
&\leq \max_{t>0}\left\{at^2 - \frac{b}{4}t^4 + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}}\right\} \\
&= \frac{a^2}{b} + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}}.
\end{aligned}$$

Consequently, $\sup I(u) \leq \frac{a^2}{b} + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}}$. This with the (3.8), our proof is complete. □

Theorem 3.5. *Assume that $a, b > 0$ and $f(x) \in L^{\frac{2^*}{2^*-1}}(\Omega)$ is a positive function a.e., then, for any $\mu > 0$, the problem (1.1) has at least a negative solution.*

Existence of the third solution. From Lemma 3.4, the functional I has the supremum. Set

$$\mathcal{F} = \Big\{T_t \in C^1\big(H_0^1(\Omega), H_0^1(\Omega)\big); T_t(u) = tu, t \in \mathbb{R}\Big\},$$

$$\mathcal{A} = \Big\{tu; u \in \mathcal{D}^+ \cup \mathcal{D}^-, t \in \mathbb{R}\Big\},$$

Then, for all $A \in \mathcal{A}$, $T_t(A) \in \mathcal{A}$ are hold for any $T_t \in \mathcal{F}$. Therefore, for any $\mu > 0$, there is

$$\frac{a^2}{4b} < \inf_{A\in\mathcal{A}}\max_{t\in\mathbb{R}}\sup_{\tilde{u}\in\mathcal{A}} I(t\tilde{u}) := c_3 \leq \frac{a^2}{b} + \frac{\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}}.$$

By Lemma 3.1 and applying the Minimax principle (see [34, Theorem 7.3.1] and [35, Theorem 1.5 & Corollary 1.3 in Chapter 3]) for I, there exist u_3^* and a sequence $\{u_m\}$ in $H_0^1(\Omega)$, moreover in $\mathcal{D}^-$, such that $u_m \to u_3^*$ in $H_0^1(\Omega)$, $I(u_m) \to c_3 = I(u_3^*)$ and $I'(u_m) \to 0 = I'(u_3^*)$ in dual space of $H_0^1(\Omega)$. Hence u_3^* is a weak solution of problem (1.1) and $\|u_3^*\|^2 > \frac{a}{b}$. □

Indeed, instead of the Minimax principle, taking $B_R := \{u \in H_0^1(\Omega) \mid \|u\| \leq R\}$ and applying the Ekeland variational principle for $-I$, we can obtain the existence of u_3^* by Lemma 3.1, where $R = \frac{2}{b}\big[a + (a^2 + \frac{b\mu^2}{2aS}\|f\|^2_{\frac{2^*}{2^*-1}})^{\frac{1}{2}}\big]$.

Proof of u_3^* is negative. Since $u_3^* \in \mathcal{D}^-$ is a weak solution of problem (1.1)with $\|u_3^*\|^2 > \frac{a}{b}$, we have $a - b\|u_3^*\|^2 < 0$ and

$$\mu \int_\Omega f u_3^* dx = \Big(a - b\int_\Omega |\nabla u_3^*|^2 dx\Big) \int_\Omega |\nabla u_3^*|^2 dx < 0.$$

Hence $u_3^* \not\equiv 0$ and

$$-\Delta u_3^* = \mu(a - b\|u_3^*\|^2)^{-1} f(x) \leq 0.$$

By the strong minimum principle, we obtain that u_3^* is negative. This proof is complete. □

3.3 Proof of the Theorem 1.1

It is clear that problem (1.1) has at least two positive solutions u_1^* and u_2^* for $\mu \in (0, \mu_{*1}]$ by the Theorem 3.3 and at least a negative solution u_3^* by the Theorem 3.5. Since

$$I(u_1^*) < 0 < I(u_2^*) < \frac{a^2}{4b} < I(u_3^*),$$

we get that u_1^*, u_2^* and u_3^* are different solutions and $0 < \|u_1^*\|^2 < \frac{a}{3b} < \frac{2a}{3b} < \|u_2^*\|^2 < \frac{a}{b} < \|u_3^*\|^2$. For $\mu \in [\mu_{*1}, +\infty)$, the Theorem 3.5 leads to that problem (1.1) owns at least a negative solution. Hence there exists $\mu_* > 0$ such that the problem (1.1) has at least three nontrivial solutions for $\mu \in (0, \mu_{*1})$, and a nontrivial solution for $\mu \in [\mu_{*1}, +\infty)$.

Moreover, for $\|u\|^2 = \frac{4a}{3b}$ and $\mu \leq \frac{a}{72b}(3abS)^{\frac{1}{2}}\|f\|^{-1}_{\frac{2^*}{2^*-1}}$, we can obtain that $I(u) \leq \frac{a^2}{4b}$, $\sup I(u)$ is achieved on $\frac{a}{b} < \|u\|^2 < \frac{4a}{3b}$. Consequently, there exists $\mu_* = \frac{a}{72b}(3abS)^{\frac{1}{2}}\|f\|^{-1}_{\frac{2^*}{2^*-1}} < \mu_{*1}$, such that, problem (1.1) has at least three nontrivial solutions u_1^*, u_2^* and u_3^* for $\mu \in (0, \mu_*)$, and a nontrivial solution u_3^* for $\mu \geq \mu_*$. In addition, it holds that $\frac{a}{b} < \|u_3^*\|^2 < \frac{4a}{3b}$ for $\mu \in (0, \mu_*)$. This proof is completed.

4 Proof of main results via algebraic analysis

Step 1. Let u be a solution, we give the calculated method to other solutions with the help of algebraic analysis. Since u is a solution of problem (1.1), one has

$$\big(a - b\|u\|^2\big) \int_\Omega \nabla u \nabla v dx = \mu \int_\Omega f(x) v dx, \ \forall\ v \in H_0^1(\Omega). \tag{4.1}$$

For given u, we have $\|u\|^2 = \int_\Omega |\nabla u|^2 dx := \alpha > 0$ and it is easy to see that $\alpha \neq \frac{a}{b}$. The existence of three solutions via algebraic analysis, if and only if there exist three values of t such that $tu \in H_0^1(\Omega)$ and

$$\big(a - b\|tu\|^2\big) \int_\Omega \nabla(tu) \nabla v dx = \mu \int_\Omega f(x) v dx, \ \forall\ v \in H_0^1(\Omega). \tag{4.2}$$

It follows from (4.1)–(4.2) that our goal is equivalent to finding all $t \in \mathbb{R}$ such that

$$t\big(a - bt^2\alpha\big) = a - b\alpha. \tag{4.3}$$

It is easy to see that $t_0 = 1$ is a solution of Eq.(4.3), and for extra, Eq.(4.3) has the solutions $t_1 = \frac{1}{2}\Big(-1 + \sqrt{\frac{4a}{b\alpha} - 3}\Big)$ and $t_2 = \frac{1}{2}\Big(-1 - \sqrt{\frac{4a}{b\alpha} - 3}\Big)$ when $\alpha < \frac{4a}{3b}$, $t_1 = t_2 = -\frac{1}{2}$ when $\alpha = \frac{4a}{3b}$ and no real solution when $\alpha > \frac{4a}{3b}$. There is $t_1 = t_0 = 1$ and $t_2 = -2$ when $\alpha = \frac{a}{3b}$ in addition.

We set $u, \bar{u}$ are different solutions of problem (1.1) with $\mu \neq 0$, then,

$$-\big(a - b\|u\|^2\big)\Delta u = \mu f(x) = -\big(a - b\|\bar{u}\|^2\big)\Delta \bar{u}.$$

It holds that $a - b\|u\|^2 \neq 0$ and $a - b\|\bar{u}\|^2 \neq 0$ are constants by (4.1). Consciously,

$$-(a - b\|u\|^2)\Delta u + (a - b\|\bar{u}\|^2)\Delta \bar{u} = 0, \tag{4.4}$$

and $\big[(a - b\|u\|^2)u - (a - b\|\bar{u}\|^2)\bar{u}\big] \in H_0^1(\Omega)$ by $u, \bar{u} \in H_0^1(\Omega)$. Multiplying the equation (4.4) by $\big[(a - b\|u\|^2)u - (a - b\|\bar{u}\|^2)\bar{u}\big]$ and integrating over Ω, we obtain that

$$\Big\|(a - b\|u\|^2)u - (a - b\|\bar{u}\|^2)\bar{u}\Big\|^2 = 0.$$

Hence $u = \frac{a - b\|\bar{u}\|^2}{a - b\|u\|^2}\bar{u}$ and all solutions of problem (1.1) are linear dependence.

Step 2. We give the proof of three solutions with $\mu > 0$ enough small only by using one of the Mountain Pass Lemma, Ekeland variational principle, and Minimax principle.

Three solutions of Theorem 1.1. According to the information above, (1.1) has at least a nontrivial solution (denote by u_1^*, u_2^* or u_3^*) with all $a, b > 0$ and $\mu \in (0, \mu_*)$ by using one of the Mountain Pass Lemma, Ekeland variational principle, and Minimax principle. Moreover, one has

$$0 < \|u_1^*\|^2 < \frac{a}{3b} < \frac{2a}{3b} < \|u_2^*\|^2 < \frac{a}{b} < \|u_3^*\|^2 < \frac{4a}{3b}.$$

Choosing one of u_1^*, u_2^* and u_3^* denoted by u, then, Eq.(4.3) has three different solutions t_0, t_1, t_2 and problem (1.1) has three different solutions u, t_1u, t_2u, them are linear dependence. □

Remark 4.1. From the step 1, we obtain that problem (1.1) has at most three different solutions. Where we proved problem (1.1) has at least 3 solutions with $\mu > 0$ enough small (see Theorem (1.1)), therefore, the problem (1.1) has only three solutions with

$$\Big\{u_1^*, u_2^*, u_3^*\Big\} = \Big\{u_1^*, t_1u_1^*, t_2u_1^*\Big\} = \Big\{u_2^*, t_1u_2^*, t_2u_2^*\Big\} = \Big\{u_3^*, t_1u_3^*, t_2u_3^*\Big\}.$$

Step 3. The specific form of μ_{**} is given exactly. And the problem (1.1) has only three solutions for $0 < |\mu| < \mu_{**}$, two solutions for $|\mu| = \mu_{**}$ and a solution for any $|\mu| > \mu_{**}$.

Proof of Theorem 1.2 with $\mu \neq 0$. Research the following elliptic problem:

$$\begin{cases} -\Delta u = f(x), & \text{in } \Omega, \\ u = 0, & \text{in } \partial\Omega, \end{cases} \tag{4.5}$$

where constant $a > 0$, $f \in L^{\frac{2^*}{2^*-1}}(\Omega)$ and $f(x) \geq 0$ a.e.. Then, problem (4.5) has an unique positive solution $U \in H_0^1(\Omega)$. Moreover, $\int_\Omega \nabla U \nabla v dx = \int_\Omega f v dx$ for all $v \in H_0^1(\Omega)$ and

$$\begin{cases} \|U\|^2 = \displaystyle\int_\Omega fU dx \leq \frac{1}{\sqrt{S}}\|f\|_{\frac{2^*}{2^*-1}}\|U\| \Longrightarrow \|U\|^{-1} \geq \sqrt{S}\|f\|^{-1}_{\frac{2^*}{2^*-1}}, \\ \|U\|^2 = \displaystyle\int_\Omega fU dx \leq \|f\|_2\|U\|_2 \leq \frac{1}{\lambda_1}\|f\|_2\|U\| \Longrightarrow \|U\|^{-1} \geq \lambda_1\|f\|_2^{-1}. \end{cases} \tag{4.6}$$

Consider the function $g(t)$ as following:

$$g(t) = \big(a - b\|tU\|^2\big)t - \mu, \tag{4.7}$$

where $t \in \mathbb{R}$ is variable, $\mu > 0$ is a parameter. If $t = T$ is a zero-point of function $g(t)$, then,

$$\Big(a - b\int_\Omega |\nabla(TU)|^2 dx\Big)\int_\Omega \nabla(TU)\nabla v dx = \mu\int_\Omega f v dx,\ \forall\ v \in H_0^1(\Omega).$$

This means that TU is a solution of problem (1.1). Based on the $g'(t) = a - 3b\|U\|^2 t^2$, we get that the zero-points of $g(t)$ are $t_m = -\sqrt{3ab}(3b\|U\|)^{-1}$ and $t_M = \sqrt{3ab}(3b\|U\|)^{-1}$. Therefore, $g(t)$ is monotonous decreasing along $(-\infty, t_m)$, increasing along $[t_m, t_M]$ and decreasing along $(t_M, +\infty)$,

$$\begin{cases} \min\ g(t) = g(t_m) = \Big(b\|U\|^2 \cdot \dfrac{a}{3b\|U\|^2} - a\Big)\cdot \dfrac{\sqrt{3ab}}{3b\|U\|} - \mu = -\dfrac{2a\sqrt{3ab}}{9b\|U\|} - \mu, \\ \max\ g(t) = g(t_M) = \Big(a - b\|U\|^2 \cdot \dfrac{a}{3b\|U\|^2}\Big)\cdot \dfrac{\sqrt{3ab}}{3b\|U\|} - \mu = \dfrac{2a\sqrt{3ab}}{9b\|U\|} - \mu. \end{cases} \tag{4.8}$$

Consciously, by (4.8), it is easy to get that, there exists $\mu_{**} = 2a\sqrt{3ab}(9b\|U\|)^{-1}$ such that Eq.(4.7) has three solutions T_1, T_2, T_3 for $0 < |\mu| < \mu_{**}$ with

$$T_1 < -\sqrt{3ab}(3b\|U\|)^{-1} < T_2 < \sqrt{3ab}(3b\|U\|)^{-1} < T_3,$$

two solutions T_2 and 'T_1 or T_3' for $|\mu| = \mu_{**}$ with

$$T_1 = -2\sqrt{3ab}(3b\|U\|)^{-1} < T_2 = \sqrt{3ab}(3b\|U\|)^{-1} \text{ or } T_2 = -\sqrt{3ab}(3b\|U\|)^{-1} < T_3 = 2\sqrt{3ab}(3b\|U\|)^{-1},$$

and a solution T_1 or T_3 for $|\mu| > \mu_{**}$ with

$$T_1 < -\sqrt{3ab}(3b\|U\|)^{-1} \text{ or } T_3 > \sqrt{3ab}(3b\|U\|)^{-1}.$$

Besides, we can state that problem (1.1) has three solutions 'T_1U, T_2U, T_3U' for $0 < |\mu| < \mu_{**}$, two solutions '$-2\sqrt{3ab}(3b\|U\|)^{-1}U$, $\sqrt{3ab}(3b\|U\|)^{-1}U$' for $\mu = \mu_{**}$, two solutions '$-\sqrt{3ab}(3b\|U\|)^{-1}U$, $2\sqrt{3ab}(3b\|U\|)^{-1}U$' for $\mu = -\mu_{**}$, a solution 'T_1U' for $\mu > \mu_{**}$ and a solution 'T_3U' for $\mu < -\mu_{**}$, where $U \in H_0^1(\Omega)$ is the unique positive solution of problem (4.5) and $\mu_{**} = 2a\sqrt{3ab}(9b\|U\|)^{-1}$.

Compared with the Step 1, all solutions of problem (1.1) are linear dependence, there are no more solutions than those mentioned above. Hence, there exists $\mu_{**} = 2a\sqrt{3ab}(9b\|U\|)^{-1}$ such that problem (1.1) has only three solutions for $0 < |\mu| < \mu_{**}$, two solutions for $\mu = \pm\mu_{**}$ and a solution for $|\mu| > \mu_{**}$. And by (4.6), $\mu_{**} = 2a\sqrt{3ab}(9b\|U\|)^{-1} \geq 2a\sqrt{3abS}(9b\|f\|_{\frac{2^*}{2^*-1}})^{-1}$. □

At last, we prove the existence of infinitely many solutions of problem (1.1) for $\mu = 0$.

Proof of Theorem 1.2 with $\mu = 0$. For any $u \in H_0^1(\Omega)$, let $V := \sqrt{ab}(b\|u\|)^{-1}u \in H_0^1(\Omega)$, it is clear that $a - b\|V\|^2 = 0$ and $\int_\Omega \nabla V \nabla v dx$ are bounded with all $v \in H_0^1(\Omega)$. So, V is a solution of problem (1.1). According to the arbitrary of $u \in H_0^1(\Omega)$, we get that problem (1.1) has infinitely many solutions when $\mu = 0$. □

We replaced the condition of $f(x)$ before by $f(x) \in L^2(\Omega)$ and $f(x) > 0$ a.e., we can obtain that the conclusion in Corollary 1.3 is clearly due to (4.6) and the progress of step 3, where $U \in H_0^1(\Omega)$ is the unique positive solution of problem (4.5) and $\mu_{**} = 2a\sqrt{3ab}(9b\|U\|)^{-1} \geq 2a\lambda_1\sqrt{3ab}(9b\|f\|_2)^{-1}$.

Acknowledgments

The authors would like to thank the anonymous reviewers and the editors. We would like to thank Professor Wei Wei from Guizhou Education University too, because she has given us many suggestions for the layout of our paper.

This research was supported by the National Natural Science Foundation of China [11861021, 11661021].

Data Availability Statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study, and all data, methods and results have been stated before.

References

[1] Kirchhoff, G.R.: Vorlesungen über Matematische Physik: Mechanik. Druck und von B.G. Teubner, Leipzig (1876). Also 2nd edn (1877); 3rd edn (1883); 4th edn (1897)

[2] Chen, S.T., Zhang, B.L., Tang, X.H.: Existence and non-existence results for Kirchhoff-type problems with convolution nonlinearity. Adv. Nonlinear Anal. **9**(1), 148–167 (2020)

[3] Li, F.Y., Zhang, Y., Zhu, X.L., et al: Ground-state solutions to Kirchhoff-type transmission problems with critical perturbation. J. Math. Anal. Appl. **482**(2), Article ID 123568 (2020)

[4] Faraci, F., Farkas, C.: On a critical Kirchhoff-type problem. Nonlinear Anal. **192**, Article ID 111679 (2020)

[5] Guo, Z.J., Zhang, X.G.: Schrödinger-Kirchhoff equation involving double critical nonlinearities. J. Math. Anal. Appl. **471**(1-2), 358–377 (2019)

[6] Bouizem, Y., Boulaaras, S., Djebbar, B.: Some existence results for an elliptic equation of Kirchhoff-type with changing sign data and a logarithmic nonlinearity. Math. Method. Appl. Sci. **42**(7), 2465–2474 (2019)

[7] Sun Y.J., Tan, Y.X.: Kirchhoff type equations with strong singularities. Commun. Pure Appl. Anal. **18**(1), 181–193 (2019)

[8] Lu, S.S.: Multiple solutions for a Kirchhoff-type equation with general nonlinearity. Adv. Nonlinear Anal. **7**(3), 293–306 (2018)

[9] Li, G.B., Xiang, C.L.: Nondegeneracy of positive solutions to a Kirchhoff problem with critical Sobolev growth. Appl. Math. Lett. **86**, 270–275 (2018)

[10] Chen, S.T., Tang, X.H.: Ground state solutions for asymptotically periodic Kirchhoff-Type equations with asymptotically Cubic or Super-cubic nonlinearities. Mediterr. J. Math. **14**, Article ID 209 (2017).

[11] Lei, C.Y., Liu, G.S., Guo, L.T.: Multiple positive solutions for a Kirchhoff type problem with a critical nonlinearity. Nonlinear Anal. Real World Appl. **31**, 343–355 (2016)

[12] Tang, X.H., Cheng, B.T.: Ground state sign-changing solutions for Kirchhoff type problems in bounded domains. J. Diff. Equ. **261**(4), 2384–2402 (2016)

[13] Wang, Y., Suo, H.M., Lei, C.Y.: Multiple positive solutions for a nonlocal problem involving critical exponent. Electron. J. Diff. Equ. **2017**, Article ID 275 (2017)

[14] Yin, G.S., Liu, J.S.: Existence and multiplicity of nontrivial solutions for a nonlocal problem. Bound. Value Probl. **2015**, Article ID 26 (2015)

[15] Wang, Y., Yang, X.: Infinitely many solutions for a new Kirchhoff type equation with subcritical exponent. Appl. Anal. (2020) https://doi.org/10.1080/00036811.2020.1767288

[16] Wang, Y., Suo, H.M., Lei, C.Y.: Existence and uniqueness of positive solution for a kind of nonlocal problems. Acta Anal. Funct. Appl. **19**(1), 95–103 (2017)

[17] Lei, C.Y., Liao, J.F., Suo, H.M.: Multiple positive solutions for a class of nonlocal problems involving a sign-changing potential. Electron. J. Diff. Equ. **2017**, Article ID 9 (2017)

[18] Lei, C.Y., Chu, C.M., Suo, H.M.: Positive solutions for a nonlocal problem with singularity. Electron. J. Diff. Equ. **2017**, Article ID 85 (2017)

[19] Duan, Y., Sun, X., Li, H.Y.: Existence and multiplicity of positive solutions for a nonlocal problem. J. Nonl. Sci. Appl. **10**, 6056–6061 (2017)

[20] Zhang, J., Zhang Z.Y.: Existence of nontrivial solution for a nonlocal problem with subcritical nonlinearity. Adv. Differ. Equ. **2018**, Article ID 359 (2018)

[21] Liang, J., Suo, H.M., Lei, C.Y.: The Existence of multiple positive solutions for a class of nonlocal problems with critical growth and concave terms. Math. Appl. **32**(1), 39–44 (2019)

[22] Wang, Y., Ye, H.Y., Suo H.M.: Existence of positive solution for a nonlocal problem with Hardy-Sobolev critical exponent. Math. Appl. **32**(2), 452–456 (2019)

[23] Wang, Y., Liang, J.P., Suo, H.M., et al: Classical solutions for nonlocal problems on different unbounded domains. Acta Anal. Funct. Appl. **21**(4), 325–341 (2019)

[24] Hamdani, M.K., Harrabi, A., Mtiri, F., Repovš D.D.: Existence and multiplicity results for a new $p(x)$-Kirchhoff problem. Nonlinear Anal. **190**, ID 111598 (2020)

[25] Wang, Y., Suo, H.M., Wei, W.: Classical Solutions for a kind of new Kirchhoff-type problems without boundary constraint. Acta Math. Sci. **40A**(5): 12 pages, in press, (2020)

[26] Qian, X.T., Chao, W.: Existence of positive solutions for nonlocal problems with indefinite nonlinearity. Bound. Value Probl. **2020**, Article ID 40 (2020)

[27] Wang, Y., Liang, J.P., Suo, H.M.: Existence of multiple solutions for a class of nonlocal near resonance problems. J. Southwest Univ.(Nat. Sci. Ed.), **40**(4):53–58. (2018)

[28] Wang, Y., Ye, H.Y., Lei, J., et al.: Infinitely many classical solutions for Kirchhoff type problem with linear term. J. Guangxi Normal Univ.(Nat. Sci. Ed.), **38**(6): in press, (2020)

[29] Cai, Z.P., Chu, C.M., Lei C.Y.: Existence and multiplicity of positive solutions for a class of nonlocal problems. J. Chongqing Normal Univ.(Nat. Sci.), **35**(01): 84–87. (2018)

[30] Wang, Y., Zhou, Y., Suo, H.M., et al.: The solutions for a kind of critical Kirchhoff problem. J. Chengdu Univ. Tech.(Sci. Tech. Ed.), **47**(2): 120–123. (2020)

[31] Rudin, W.: Real and Complex Analysis, Third Edition. McGraw-Hill, Singapore (1987)

[32] Ekeland, I.: On the variational principle. J. Math. Anal. Appl. **47**, 324–353 (1974)

[33] Ambrosetti, A., Rabinowitz, P.H.: Dual variational methods in critical point theory and applications. J. Funct. Anal. **14**, 349–381 (1973)

[34] Lu, W.D.: *The Variational Method in Differential Equation* (Revised Edtion). Published by the China Science Publishing & Technical Media Ltd., Peking (2003)

[35] Chang, K.C.: Critical Point Theory and its Applications. Published by the Shanghai Scientific & Technical Publishers, Shanghai (1986)