

Further milliarcsecond resolution results on cool giants and binary stars from lunar occultations at Devasthal.

A. Richichi,^{1★} S. Sharma,² T. Sinha,² R. Pandey,² A. Ghosh,² D.K. Ojha,³
A.K. Pandey,² and M.B. Naik³

¹*INAF - Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy*

²*Aryabhatta Research Institute of Observational Sciences, Manora Peak, Naini Tal, 263002 India*

³*Tata Institute of Fundamental Research, Homi Bhabha Road, Colaba, Mumbai - 400005, India*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We report on 26 lunar occultation events observed in the context of a program started at Devasthal in 2017. In addition to the customary observations with the 1.3-m telescope, we report here also the first attempts performed with the TIRCAM2 camera at the 3.6-m DOT telescope in the near-IR. The results consist in the first-time angular diameters for two late-type giants, in a measurement of the well-known AGB pulsating variable SW Vir, and in the measurement of eight small separation binaries one of which is detected for the first time (HR 1860). We also measured the wider binaries SAO 94431 and 55 Tau (twice). The remaining sources were found to be unresolved with upper limits as small as 1 milliarcsecond. We plan to develop further the high-speed capability of the TIRCAM2 instrument, so as to include in our program also more near-infrared, highly extincted sources.

Key words: occultations – stars: binaries: general – stars: atmospheres

1 INTRODUCTION

The scope and methods of the lunar occultation (LO) program at Devasthal have been described in detail in our previous papers (Richichi et al. 2017a, 2018) so that we keep the discussion to a minimum in the present work. We use the LO technique mainly for two purposes: to investigate properties of late-type giants such as angular diameter, effective temperature and in the case of very evolved star the possible presence of circumstellar matter; and to study sub-arcsecond separation binary stars which are either already known but with incomplete data, or discover new systems. In the case of binary stars, adding observations at epochs, position angles or filters not hitherto obtained is crucial to complete the knowledge of the orbit and of the spectral type of the components.

Although obviously limited in the selection of the sources, by the one-dimensional projection of the results and by the fixed-time nature of the events, LO have several advantages that make them still very attractive. Among them, the very efficient use of telescope time, the inexpensive instrumentation, the combination of high sensitivity and high angular resolution, the relatively simple and model-independent data reduction, and being largely not limited

by seeing and telescope diameter while achieving routinely milliarcsecond (mas) resolution.

We report here on 26 LO events, with instrumentation and data analysis briefly summarised in Sect. 2. Eleven of these events include either stars with a resolved angular diameters or binary stars, which we discuss individually in Sect. 3. We also provide a few comments on the unresolved sources.

2 OBSERVATIONS AND DATA ANALYSIS

All events were recorded at Devasthal (India) with two different telescopes and instruments, both operated by the Aryabhatta Research Institute of Observational Sciences (ARIES). A log of the observations is provided in Table 1, which follows the style used in Richichi et al. (2018). The first few columns list the date, time, type of event (disappearance or reappearance), source designation, V magnitude and spectrum. These latter were compiled from the *Simbad* database.

Mainly, we used the 1.3-m telescope equipped with a 512x512 pixels frame transfer ANDOR iXon EMCCD. More details were given in Richichi et al. (2017a). We employed small sub-windows positioned at the center of the detector, with pixel areas as listed under the Sub column in Table 1. Rebinning was also employed as listed in column Bin. The

★ E-mail: andrea4work@posteo.eu

integration time and the effective sampling time in ms are listed under τ and ΔT , respectively. The filters employed were either Cousins R or I.

Also, we employed for the first time for LO work the near-IR camera TIRCAM2 (Naik et al. 2012; Baug et al. 2018) developed by the Tata Institute of Fundamental Research (TIFR), mounted at the Cassegrain main port of the 3.6-m ARIES Devasthal Optical Telescope (DOT; Kumar et al. 2018). TIRCAM2 is a closed-cycle Helium cryo-cooled imaging camera equipped with a Raytheon 512×512 pixels InSb Aladdin III Quadrant focal plane array (FPA). The field of view (FOV) of the TIRCAM2 is $86''.5 \times 86''.5$ with a plate scale of $0''.169$. The full-width at half-maximum of the stellar images was $\sim 0''.7$. TIRCAM2 provides sampling time of ~ 256 ms for the full frame, therefore for LO work we have used a sub-array window of 32×32 pixels to increase the readout rate. The resulting sampling time consists of window readout time (~ 6 ms) as well as other overheads and can vary from 16 ms to 48 ms most of the time. The TIRCAM2 observations, all recorded in a broad-band K filter, were obtained at an early phase of the commissioning of this instrument on the 3.6-m telescope. In fact, three more observations were attempted in October 2017: although the occultations were detected, the data were not useful for a quantitative analysis and are therefore not listed in Table 1.

The last two columns of Table 1 list the signal-to-noise ratio S/N of the best fit, and the type of result. LO predictions are made with our software using a wide range of catalogs, and the data enter our archive with source names which do not follow a particular rule. To help in looking them up, we provide in Table 2 a list of cross-identifications especially aimed at binary star researchers.

In all cases, for our data analysis we convolved the filter transmission with the response of the corresponding detector, and we took into account the finite integration time, the primary diameter and the secondary obstruction.

The data products were FITS cubes, which were converted to photometric light curves using a digital extraction mask hand-tailored in each case to the seeing and image motion of the source. This process was described in Richichi et al. (2018) and references therein. Only about 1–2 s of data were analyzed, corresponding to an apparent lunar motion of $\approx 2''$ on the sky.

The light curves were then analyzed using both a model-dependent (LSM) and a model-independent (CAL) method. These are recalled e.g. in Richichi et al. (2017a). The LSM provides values and errors for free parameters such as angular diameter, or projected separation and fluxes for binary and multiple stars. Scintillation can be accounted for to some extent. CAL is iterative, and provides a brightness profile of the source derived from maximum-likelihood criteria. The errors associated with the parameters (LSM) and the uncertainties on the brightness profiles (CAL) have been discussed in the mentioned references.

3 RESULTS

The list of stars which we found to be resolved or binary is provided in chronological order in Table 3, which also follows the format used in our previous papers. Columns 2 through




Figure 1. Top panel: left, disappearance light curve (dots) for IRC -20478 and best fit by a UD model of 2.57 mas (solid line), including a scintillation correction by Legendre polynomials. Right, reappearance light curve (dots) for ψ Leo and best fit by a UD model of 3.03 mas (solid line). The two curves have been normalized in intensity, and rescaled in x-axis to the time of geometrical occultation. Note that the absolute limb speed is about 50% faster for ψ Leo, see Table 3. Bottom panel: the fit residuals for the two cases, also normalized and offset by arbitrary amounts for clarity.

6 refer to the geometry of the event, listing respectively the measured rate of the event V , its deviation from the predicted rate V_t , the local lunar limb slope ψ , and the effective position and contact angles, PA and CA respectively. Due to limitations in the time sampling, V could not be computed from the data of the TIRCAM2 observations, thus the ψ field is empty and all other quantities are the predicted values. Column 7 lists the best-fitting angular diameter in the uniform disk approximation (UD). For binaries, columns 8 and 9 list the projected separation (along the PA direction from primary to secondary) and the brightness ratio, respectively.

In the following subsections we briefly discuss our results individually for each resolved star, following the same order as Table 3. A brief discussion of the unresolved sources is provided in Sect. 3.13.

3.1 IRC -20478

Only 2 previous LO events were reported for this bright K1/K2 giant, without being resolved (Africano et al. 1976; Eitter & Beavers 1979). We recorded the light curve shown in Fig. 1 under good conditions in R band, allowing us to measure this source for the first time. The data are best fitted by an UD angular diameter of 2.57 ± 0.04 mas, which can be compared to the expected value of 2.6 mas (van Belle 1999). The normalized χ^2 value of this fit is 1.3, while for comparison it is 2.0 for a fit by a point source.

3.2 ψ Leo

No high angular resolution measurements by any technique have been reported for this M giant, apart from a mid-

Table 1. List of observed events

Date	Time (UT)	Type	Source	V	Sp	Filter	Sub	Bin	τ	ΔT	S/N	Notes ¹
2017 Oct 07	20:39	R	μ Cet	4.3	A9IIIp	R	32×32	2×2	2.28	2.75	11.4	Tm
2017 Oct 08	18:21	R	SAO 93529	8.9	F8	R	32×32	2×2	4.00	4.48	5.9	Bin
2017 Oct 08	19:03	R	SAO 93532	6.7	G5	R	32×32	2×2	3.00	3.48	38.4	UR
2017 Oct 09	19:15	R	SAO 94056	8.4	N?	I	32×32	2×2	4.00	4.48	30.5	UR
2017 Oct 25	13:32	D	IRC -20478	4.9	K1/2III	R	16×16	2×2	2.00	2.47	67.4	Dm
2017 Nov 01	12:54	D	IRC +00008	9.7	M5	I	16×16	2×2	3.00	3.47	23.1	UR
2017 Nov 05	21:23	R	55 Tau	6.9	F8V	R	32×32	2×2	4.00	4.48	23.3	WB
2017 Nov 06	00:21	R	DO 11286	10.3		I	32×32	2×2	4.00	4.48	9.3	Bin
2017 Nov 06	17:58	R	SAO 94431	7.2	B3V	R	32×32	2×2	4.00	4.48	6.8	WB
2017 Dec 01	16:35	D	μ Cet	4.3	A9IIIp	I	16×16	2×2	2.50	2.97	52.0	Bin
2017 Dec 30	14:06	D	SAO 93803	7.2	A0	R	16×16	2×2	4.00	4.47	22.9	Bin
2017 Dec 30	17:14	D	SAO 93838	6.6	K1III	I	16×16	2×2	3.00	3.47	41.6	UR
2017 Dec 30	19:14	D	55 Tau	6.9	F8V	I	16×16	2×2	4.00	4.47	22.4	WB
2018 Jan 04	22:20	R	ψ Leo	5.4	M2III	I	16×16	2×2	3.00	3.48	104.7	Dm
2018 Jan 08	20:49	R	SW Vir	6.9	M7III:	I	32×32	2×2	2.28	2.75	26.3	Dm
2018 Apr 04	19:35	R	SAO 159887	8.9	K0III	I	32×32	2×2	4.00	4.48	3.8	UR
2018 May 18 ²	14:32	D	IRC -20156	8.5	C-N5III:	K	32×32	No	6.40	16	30.3	UR
2018 May 21 ²	17:19	D	SAO 98770	9.4	F8	K	32×32	No	6.40	32	18.1	Bin
2018 Oct 21	15:34	D	HD 221925	7.8	F0V	R	32×32	2×2	3.00	3.48	5.4	Bin
2018 Dec 26	19:16	R	SAO 98568	8.0	M2.5III	R	32×32	2×2	4.00	4.48	14.8	UR
2019 Jan 18	16:37	D	HD 36230	8.2	K5	I	16×16	2×2	3.00	3.47	36.3	UR
2019 Jan 18	18:25	D	HR 1860	6.2	B6V	R	16×16	2×2	2.50	2.97	19.2	Bin
2019 Jan 19	16:26	D	SAO 78514	8.0	K2	I	16×16	2×2	3.00	3.48	10.3	UR
2019 Jan 19	18:26	D	AX Gem	9.5	M5	I	16×16	2×2	3.00	3.47	5.1	Tm
2019 Jan 19	19:13	D	SAO 78587	8.6	K0	I	16×16	2×2	3.50	3.97	6.7	UR
2020 Jan 05	15:50	D	HR 797	6.3	A2V	I	21×21	2×2	3.00	3.48	17.3	Bin

¹: Tm = event close to the lunar terminator; UR = unresolved; Dm = resolved diameter ; Bin = binary ; WB = wide binary.²: TIRCAM2 at DOT 3.6 m.**Table 2.** List of source cross-identifications

μ Cet	HD 17094	HIP 12828	SAO 110723	WDS J02449+1007	HR 813
SAO 93529		HIP 16991		WDS J03385+1336AB	
SAO 93532	HD 22682	HIP 17049			
SAO 94056	HD 29496				
IRC -20478	HD 169420	HIP 90289	SAO 186794	WDS J18254-2033AB	21 Sgr, HR6896
IRC +00008			SAO 128708		FG Psc
55 Tau	HD 27383	HIP 20215	SAO 93870	WDS J04199+1631AB	
DO 11286					TYC 1301-780-1
SAO 94431	HD 34251	HIP 24612		WDS J05167+1826AB	
SAO 93803	HD 26380	HIP 19519			
SAO 93838	HD 27029	HIP 19960			
ψ Leo	HD 84194	HIP 47723	SAO 98733	WDS J09437+1401A	16 Leo, HR3866
SW Vir	HD 114961	HIP 64569	SAO 139236		
SAO 159887	HD 147474				
IRC -20156	HD 67507	HIP 39751	SAO 175215		RU Pup
SAO 98770				WDS J09476+1419AB	
HD 221925		HIP 116488	SAO 146801	WDS J23363-0707AB	
SAO 98568	HD 81540	HIP 46311			
HD 36230			SAO 77223		
HR 1860	HD 36589	HIP 26072	SAO 77255		
SAO 78514	HD 46467	HIP 31353			
AX Gem	HD 260525				
SAO 78587	HD 261221				
HR 797	HD 16861	HIP 12647	SAO 93067		85 Cet

IR LO mentioned by [Stecklum et al. \(1999\)](#) who only used it as a test. We could record a LO light curve with high S/N as shown in Fig. 1, leading to an angular diameter of 3.03 ± 0.03 mas. The empirical prediction based on the class and color was 3.6 mas ([van Belle 1999](#)), indicating some pos-

sible discrepancy. The compilation by [Eggleton & Tokovinin \(2008\)](#) does not report suspected multiplicity for ψ Leo. It was also not revealed as a spectroscopic binary in a survey by [Famaey et al. \(2009\)](#).

Table 3. Summary of results: angular sizes (top) and binaries (bottom)

(1) Source	(2) V (m/ms)	(3) (V/V _i)-1	(4) $\psi(^{\circ})$	(5) PA($^{\circ}$)	(6) CA($^{\circ}$)	(7) ϕ_{UD} (mas)	(8) Proj.Sep.(mas)	(9) Br. Ratio
IRC -20478	0.4368	13.7%	6.0	144	60	2.57 \pm 0.04		
ψ Leo	-0.6282	8.1%	7.3	266	152	3.03 \pm 0.03		
SW Vir ¹	-0.588	–	–	160	224	\approx 17		
SAO 93529	-0.7286	2.0%	2.3	41	154		20.3 \pm 2.2	7.9 \pm 1.4
SAO 94431 ¹	-0.746	–	–	51	156		332.6 \pm 1.0	3.45 \pm 0.13
55 Tau	-0.5839	-2.7%	2.4	291	214		137.2 \pm 0.9	3.10 \pm 0.05
DO 11286	-0.4578	7.5%	2.7	34	125		36.0 \pm 1.8	9.33 \pm 0.68
μ Cet	0.5137	61.5%	20.2	146	83		108.6 \pm 0.3	17.9 \pm 0.2
SAO 93803	0.5045	6.9%	3.1	199	-49		63.5 \pm 1.3	18.1 \pm 1.8
55 Tau	0.6569	6.1%	4.7	47	-33		556.3 \pm 3.2	2.64 \pm 0.02
SAO 98770 ¹	0.806	–	–	79	-33		177.0 \pm 0.4	1.29 \pm 0.01
HD 221925 ¹	0.341	–	–	181	-57		165.0 \pm 1.4	2.61 \pm 0.07
HR 1860 ²	0.4553	10.0%	-5.5	215	-51		6.3 \pm 0.3	4.63 \pm 0.16
HR 797	0.3634	10.5%	3.7	123	63		11.2 \pm 1.4	1.51 \pm 0.14

¹: V could not be computed from data, using predicted value. ²: possible triple

3.3 SW Vir

This is a bright AGB star, with a spectral type M7III and a distance of 300 pc which result in a very large angular diameter. As a result, it has been the subject of numerous investigations with a wide range of techniques from LO, to speckle and adaptive optics on large telescopes, to long-baseline interferometry, that we do not attempt to list here. In general, the photospheric angular diameter is found to be around 15–20 mas, with a significant scatter due to the wavelength of observation, to intrinsic pulsation, and to the presence of a circumstellar shell.

Our goal was to measure not only the diameter but to also provide a model-independent brightness profile of both photosphere and circumstellar shell in the style of what was done e.g. for α Tau by Richichi et al. (2017a). Unfortunately, this was not possible due to high-frequency fluctuations in the signal of a not yet understood nature, possibly caused by wind shaking of the telescope. We could only derive a rough estimate of the angular diameter of ≈ 17 mas, in line with expectations.

3.4 SAO 93529

A previous LO of this star was reported by Evans & Edwards (1981), who found it to be a single star. Later on, SAO 93529 was suspected as binary from Hipparcos astrometry and was confirmed in a speckle measurement by Horch et al. (2002) with separation 0''37 along PA= 88°. It has since been reported in several publications, the latest being by Horch et al. (2017) who found a separation of 0''30 along PA= 78° at epoch 2012.1. Given the definite motion exhibited over the available time span, our measurement adds to the database for a possible orbit computation in the future. Horch et al. (2017) also reported $\Delta(880\text{nm}) = 2.7$ mag, to be compared with our $\Delta R = 2.2$ mag.

3.5 SAO 94431 and 55 Tau

These two wide visual and speckle binaries have already been extensively observed and have published orbital elements, so that our measurements are just a mere confirmation. Perhaps of interest are our accurate flux ratio determinations. 55 Tau is a confirmed member of the Hyades open cluster (Perryman et al. 1998).

3.6 DO 11286

This late-type star (IRAS 05255+1832) does not have a spectral classification in the literature, although its $B - V$, $V - K$ colors and the GAIA distance of 1.2 kpc are indicative of a mid-M type giant. It was first detected as binary by Richichi et al. (1997), who could obtain two simultaneous observations in October 1995 from two different sites. The non-detection at one of the sites was used as a constraint to infer a 0''1 to 0''3 actual separation in a southerly direction. Upon revisiting these older data, however, we found that a mistake was made in the sign of the limb slope. The actual PA of the 1995 measurement was in fact 322° and not 106° as reported by Richichi et al. (1997). The ΔK was 2.2 mag. No other high angular resolution measurements are available.

Our present measurement confirms the binary nature of DO 11286, and with ΔI of 2.4 mag, comparable to the previously measured ΔK , it follows that the spectral types are possibly quite similar for the two components. The uncertainty in the estimate of the main spectral type and possible variability prevent us from a more precise classification of the components.

If we assume that the total system mass is about 1 M_{\odot} and that the projected separation of 0''036 in 2017 (or 43 au) is close to the actual one, it would follow that $P \gtrsim 280$ yr. If so, then the 22.1 yr interval between the two LO events would be sufficiently small to warrant combining the projected separations. In this case, we find the actual separation to be 36.3 mas along PA=41°. Additional measurements possibly by AO or speckle at a large telescope would be able to confirm this result.

3.7 μ Cet

This interesting stellar system is a known speckle binary, additionally found to have two more components – detected so far only in the near-IR by LO. The situation has been recently summarized by [Dyachenko et al. \(2018\)](#), who reported two LO events observed from the Russian 6-m telescope as well as simultaneous multi-wavelength speckle observations. The same events were observed also by us at Devashtal, the first one in October 2017 with the 1.3-m and the 3.6-m telescopes, and the second one in December of the same year with the smaller telescope only.

Unfortunately, the first one was with a high lunar phase and close to the terminator. Both telescopes recorded the event, but TIRCAM2 at the 3.6-m was just at the first commissioning of the LO mode and did not allow us to derive a useful light curve beyond the simple detection, while the 1.3-m data are affected by a rapidly varying background due to the sunlit lunar mountains on the terminator and have insufficient S/N to resolve the system.

The December 2017 data are instead of sufficient quality to resolve at least the speckle companion with certainty. The parameters are listed in Table 3. Our $\Delta I = 3.13$ mag is in excellent agreement within errorbars with the LO result by [Dyachenko et al. \(2018\)](#) of 3.16 mag in a narrow continuum filter at 850 nm. The position angle and projected separation could be combined with those of the mentioned paper, but in fact we note that in the presence of a high local limb slope of 20° , which is not too unusual at the given contact angle, the result would have a significant uncertainty. This is in any case less relevant given the extensive speckle coverage of the system on the same date by [Dyachenko et al. \(2018\)](#).

3.8 SAO 93803

Like DO 11286, also SAO 93803 has an interesting record of positive and negative binary detections. They are summarized by [Richichi et al. \(2017b\)](#), who resolved the binary once convincingly and once with borderline S/N in two events in January and October 2016, respectively, both in an SDSS r filter.

Our measurement in a Cousins R filter also detects the companion. [Richichi et al. \(2017b\)](#) estimated that the period could be of order ≈ 50 yr. On this premise, if we neglect the 1.9 yr interval and combine the January 2016 and December 2017 results we obtain a true separation of $0''.123$ along $PA = 78^\circ$ at epoch ≈ 2017.0 . There remains the intriguing aspect of the difference in the brightness ratios, namely $\Delta r = 2.3$ mag and $\Delta R = 3.1$ mag, respectively. This can be at least partly explained by the fact that the transmission of the r and R filters is sufficiently different, with the latter being sensitive to less blue and much more red wavelengths. We note that an AO-aided spectrograph could be used to obtain individual spectra of the components.

3.9 SAO 98770

This F8 star has been known as a binary since several decades, but has a sparse record of observations by visual micrometer ([Heintz 1980](#)), by one previous LO ([Blow et al. 1982](#)), and by two speckle data points ([Mason 1996](#); [Hartkopf et al. 2000](#)). The separation of the components has

been reported between $0''.15$ and $0''.435$, with an upper limit of $< 0''.036$ in one case. Only the LO authors reported a magnitude difference, namely 0.35 mag in a RG610 filter. No orbital elements are known, and there is no available parallax for the SAO 98770 system from either Hipparcos or GAIA. The fact that the separation appears to vary significantly as well as the high proper motion of almost $0''.1$ /yr, however, are suggestive of a rather nearby system with a relatively short period.

Our measurement complements the set of separations, and especially it adds the first near-IR magnitude difference between the components. With $\Delta K = 0.28$ mag, the companion seems to be only slightly cooler than the primary. If the primary is a F8 dwarf, the color difference $R-K=1.18$ mag suggests a spectral type around K0V for the secondary and therefore a combined system mass $\approx 2 M_\odot$. With this as a constraint, we find it difficult to reconcile the wide range of measured separations with periods less than 100 yr, if we have to assume the distance modulus implied by an F8V main component. All these facts make SAO 98770 a very interesting candidate for further high angular resolution studies, e.g. by speckle at a large telescope.

3.10 HD 221925

This relatively wide binary has been reported in about ten publications by both visual and speckle observers starting with [Couteau \(1958\)](#), and we add here the first LO measurement. An orbit determination is still lacking.

3.11 HR 1860

Our LO data resolve HR 1860 as binary by means of both model-dependent and model-independent methods, as shown in Fig. 2. At the GAIA distance of 368 pc, the projected separation of 6.3 mas translates to 2.3 au. The implication is that the orbital period could be as short as very few years. Indeed, as shown by the CAL brightness profile in Fig. 2, there is an indication of a further component even closer in. The LSM analysis indicates a fit convergence for a projected separation of 5 mas and brightness ratio 1:19 against the primary, but as the signal is very close to the noise limit and the improvement in χ^2 is nil, we consider it dubious and mention the possibility of a triple system only for reference to future observers.

No previous high angular resolution measurements are available for this bright B6 dwarf, with the exception of a LO recorded in the visual with a 48-cm telescope by [Trunkovskij \(1987\)](#) who found it to be unresolved. Until now, there had been no indications of binarity in this star from other methods such as spectroscopy ([Abt & Cardona 1984](#); [Chini et al. 2012](#)). This could be explained by a nearly on-sky inclination of the system, and/or by a large difference in mass between the components which would lead to an almost stationary single-lined spectrum.

According to the CHARM2 catalog ([Richichi et al. 2005](#)), a total of 116 LO events of 72 unique B stars have been published until 2004, with 31 or 43% found to be binaries or multiples. This confirms the high binary fraction already inferred for early, massive spectral types by [Chini et al. \(2012\)](#).




Figure 2. Top panel: light curve (dots) for HR 1860, repeated twice with an arbitrary scaling factor and offset. The upper solid line is a fit by a point-like source, the lower solid line is the best fit by a binary star, as described in the text. The times of occultation of the A, B components are marked. The normalized χ^2 values for the two fits are also shown. Middle panel: the residuals for the two fits, offset by arbitrary amounts for clarity. Bottom panel: brightness profile reconstructed by the CAL method, normalized to 1 at the peak of the primary and enlarged for clarity. Note the possible presence of a third component, not included in the fit shown.

3.12 HR 797

This source was first detected as a LO binary by [Blow et al. \(1982\)](#), who reported 10 mas projected separation along $PA=21^\circ$ with $\Delta m = 0.18$ mag in a filter-less blue channel, from an event in November 1981. Subsequent attempts by speckle were either partially successful, being resolved once with 42 mas separation in V band in 1994 but not in two other observations at similar dates by [Mason \(1996\)](#), and not by [Hartkopf et al. \(2012\)](#) in y and I filters.

Our data detection of the binary is shown in Fig. 3. At a GAIA distance of 122 pc, the minimum semi-axis corresponding to the available (projected and non) separations would imply a period shorter than the span between the few available measurements. Further observations are needed to determine orbital elements and dynamical masses. We note that this star is included in a study that uses simulations based on Hipparcos and GAIA proper motions to derive constraints on orbits and masses of companions to nearby stars ([Kervella et al. 2019](#)).




Figure 3. Top panel: light curve (dots) for HR 797, and best fit by binary model with 11.6 mas projected separation. The times of occultation of the A, B components are marked. Bottom panel: the fit residuals.

3.13 Unresolved sources

The remaining unresolved sources, some of which were expected to be binary from previous measurements or resolved in diameter on the basis of their photometry and spectral type, are briefly discussed in the following. When applicable, we compute the upper limit on the angular size of the source using the algorithm described by [Richichi et al. \(1996\)](#), and the expected angular size of late-type giants and evolved stars according to the empirical formula by [van Belle \(1999\)](#). For this group, we do not list here all the references in order to be more concise.

SAO 93532 was found unresolved also in two previous LO events. The same applies to SAO 94056 (two LO events), SAO 93838, SAO 98568, HD 36230 and SAO 78587 (one LO event each). SAO 159887 was a suspected double in one previous LO event but found unresolved by speckle. IRC +00008 and IRC -20156 have no previous high angular resolution observations in the literature and were expected to be resolved with 2.4 and 2.7 mas, respectively. The upper limits from our light curves are 3.8 and 4.5 mas, respectively.

A previous LO event found SAO 78514 as binary with 40 mas projected separation and 1:1.7 brightness ratio in the K band. The event in the present paper shows a ratio < 10 in the I band, which in turn implies a color $I - K > 3$ mag for the secondary. This would indicate a late M spectral type. Finally, AX Gem was resolved in a previous LO with a 3.0 mas UD diameter. Our present data, obtained at a limb position very close to the terminator, do not allow us sufficient S/N to derive a diameter.

We could derive upper limits on the source size for 9 of the 11 unresolved light curves. The average upper limit is 3.2 mas. For comparison, among the detected binary stars listed in Table 3, the one with the smallest upper limit on the angular diameter of the primary is μ Cet, with 0.8 mas.

4 CONCLUSIONS

We have reported on 26 additional LO measurements carried out at Devashtal since our latest paper (Richichi et al. 2018). In addition to the EMCCD on the 1.3-m telescope, we have also used for the first time the near-IR camera TIRCAM2 on the DOT 3.6-m telescope. This instrument was in a commissioning phase at the time of our observations and the time sampling as well as other characteristics were not optimal yet. Of the five events recorded with TIRCAM2, we have reported here on two with a light curve of sufficient quality.

Our results include three resolved angular diameters, two of which are first time measurements of the late-type stars IRC-20478 and ψ Leo, and one of the well known AGB star SW Vir. They also include eight small-separation binaries, of which one is a new detection (HR 1860). The rest are previously known but mostly with very few data points and all without orbital elements, so that our measurements provide valuable additions especially for what concerns the flux ratio and color of the companions. Additionally, we reported on three LO events of two relatively wide binaries.

Of the remaining 11 unresolved sources, a few were expected to be resolved either in diameter or as binaries but we found reasonable justifications for our negative detections. The mean upper limit on the angular size of the unresolved sources was about 3 mas, and around 1 mas in the best cases.

The LO method is confirmed as an ideal technique to achieve high angular resolution comparable to the most sophisticated facilities such as long-baseline interferometry, even with a relatively small telescope and inexpensive commercial instrumentation. The cost to pay are the obvious limitations in the choice and repetition of the sources. We plan to continue our program, and to carry out more near-IR LO observations with an improved fast mode on TIRCAM2 in the near future.

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

ACKNOWLEDGMENTS

This work has made use of the SIMBAD database, operated at CDS, Strasbourg, France. It is a pleasure to thank the technical team in Devashtal, as well as the members of the IR astronomy group at TIFR, for their support during the observing runs. DKO and MBN acknowledge the support of the Department of Atomic Energy, Government of India, under project No. 12-R&D-0200.TFR-5.02-0200.

REFERENCES

Abt, H. A., & Cardona, O. 1984, *ApJ*, 285, 190
 Africano, J. L., Evans, D. S., Fekel, F. C., et al. 1976, *AJ*, 81, 650
 Baug, T., Ojha, D. K., Ghosh, S. K., et al. 2018, *Journal of Astronomical Instrumentation*, 7, 1850003-1881
 Blow, G. L., Chen, P. C., Edwards, D. A., et al. 1982, *AJ*, 87, 1571

Chini, R., Hoffmeister, V. H., Nasseri, A., et al. 2012, *MNRAS*, 424, 1925
 Couteau, P. 1958, *Journal des Observateurs*, 41, 1
 Dyachenko, V., Richichi, A., Balega, Y., et al. 2018, *MNRAS*, 478, 5683
 Eitter, J. J., & Beavers, W. I. 1979, *ApJS*, 40, 475
 Eggleton, P. P., & Tokovinin, A. A. 2008, *MNRAS*, 389, 869
 Evans, D. S., & Edwards, D. A. 1981, *AJ*, 86, 1277
 Famaey, B., Pourbaix, D., Frankowski, A., et al. 2009, *A&A*, 498, 62
 Hartkopf, W. I., Mason, B. D., McAlister, H. A., et al. 2000, *AJ*, 119, 3084
 Hartkopf, W. I., Tokovinin, A., & Mason, B. D. 2012, *AJ*, 143, 42
 Heintz, W. D. 1980, *ApJS*, 44, 111
 Horch, E. P., Robinson, S. E., Meyer, R. D., et al. 2002, *AJ*, 123, 3442
 Horch, E. P., Casetti-Dinescu, D. I., Camarata, M. A., et al. 2017, *AJ*, 153, 212
 Kervella, P., Arenou, F., Mignard, F., et al. 2019, *A&A*, 623, A72
 Kumar, B., Omar, A., Maheswar, G., et al. 2018, *Bulletin de la Societe Royale des Sciences de Liege*, 87, 29
 Mason, B. D. 1996, *AJ*, 112, 2260
 Naik, M. B., Ojha, D. K., Ghosh, S. K., et al. 2012, *Bulletin of the Astronomical Society of India*, 40, 531
 Perryman, M. A. C., Brown, A. G. A., Lebreton, Y., et al. 1998, *A&A*, 331, 81
 Richichi, A., Baffa, C., Calamai, G., et al. 1996, *AJ*, 112, 2786
 Richichi, A., Calamai, G., Leinert, C., et al. 1997, *A&A*, 322, 202
 Richichi, A., Percheron, I., & Khristoforova, M. 2005, *A&A*, 431, 773
 Richichi, A., Dyachenko, V., Pandey, A. K., et al. 2017a, *MNRAS*, 464, 231
 Richichi, A., Tasuya, O., Irawati, P., et al. 2017b, *AJ*, 154, 215
 Richichi, A., Sharma, S., Pandey, A. K., et al. 2018, *New Astron.*, 59, 28
 Stecklum, B., Käufel, H.-U., & Richichi, A. 1999, *The Messenger*, 95, 25
 Trunkovskij, E. M. 1987, *Soviet Astronomy Letters*, 13, 379
 van Belle, G. T. 1999, *PASP*, 111, 1515