

Roots of Chebyshev Polynomials: a purely algebraic approach

Lionel Ponton
lionel.ponton@gmail.com

Abstract

By using purely algebraic tools, we establish well-known properties of roots of Chebyshev polynomials. Especially, we show that these zeros are simple and lie in $(-1, 1)$ and we prove in two ways that they are mostly irrational.

Mathematics Subject Classification (2010). Primary: 12D10, 11J72, 11B83.

Keywords. Chebychev polynomials, location of zeros, irrationality.

1 Introduction

The Chebyshev polynomials form one of the most famous and classical family of polynomials. They are related to many others subjects such as orthogonal polynomials, Jacobi polynomials, sine and cosine functions and approximation theory and may be define in many ways: recurrence relation, differential equation, trigonometric relation, generating function, etc.

In approximation theory, the roots of these polynomials have a key role in connection with the Runge's phenomenon [MH03, p. 143-144]. According to the orthogonality of Chebyshev polynomials [Sny66, p. 16], it is well-know that these zeros are simple and lie in the open interval $(-1, 1)$ [Sny66, p. 7]. Moreover, the trigonometric relations satisfied by Chebyshev polynomials lead to some explicit expressions of these roots of the form $\cos(r\pi)$ where r is a rational number [Sny66, p. 14]. Since the arithmetical properties of this kind of numbers are well-know [NZM91, Th. 6.16, p. 308], it is easy to derive that these roots are mostly irrational [Pon18, Th. 1].

We can note that, even if Chebyshev polynomials are algebraic objects, they are mainly study using calculus and trigonometry. In this paper, we adopt the following point of view: building on an algebraic definition of the Chebyshev polynomials, we derive all the above properties of their roots using only algebraic tools. More precisely, in section 2, we show that all the zeros of Chebyshev polynomials are real and lie in $(-1, 1)$ by means of recursive sequences and very basic computation on complex numbers. Then, in section 3, we establish that these roots are simple using polynomial arithmetic. Finally, in section 4, on the basis of explicit expressions of Chebychev polynomials, we show that their zeros are mostly irrational in two ways: first, by using their expression directly; then, by means of shifted Chebychev polynomials. Note that, even if all the properties of Chebyshev polynomials stated and used in the following are already known, we prove each of them both to provide a self-contained article, and to ensure that it only use algebraic tools.

In that respect, we start with the following definition.

Definition 1. The Chebyshev polynomials of first kind are defined by the recurrence relation:

$$\begin{cases} T_0 = 1, T_1 = X, \\ \forall n \in \mathbb{N}, T_{n+2} = 2XT_{n+1} - T_n \end{cases}$$

and the Chebyshev polynomials of second kind are defined by the recurrence relation:

$$\begin{cases} U_0 = 1, U_1 = 2X, \\ \forall n \in \mathbb{N}, U_{n+2} = 2XU_{n+1} - U_n \end{cases}$$

Throughout this article, we designate \mathbb{N} as the set of nonnegative integers, \mathbb{N}^* as the set of positive integers, \mathbb{Z} as the set of rational integers and \mathbb{C} as the set of complex numbers.

It is easy to check by double induction that, for all $n \in \mathbb{N}$, T_n and U_n are in $\mathbb{Z}[X]$, have degree n and are even or odd according to n is even or odd.

2 The roots of Chebyshev polynomials lie in $(-1, 1)$

Lemma 2. For all $n \in \mathbb{N}$, $T_n(1) = 1$ and $U_n(1) = n + 1$.

Proof. This follows from an obvious double induction on n . □

Lemma 3. Let $w \in \mathbb{C} \setminus \{-1, 1\}$. We define two sequences (t_n) and (u_n) of complex numbers by the recurrence relations:

$$\begin{cases} t_0 = 1, t_1 = w, \\ \forall n \in \mathbb{N}, t_{n+2} = 2wt_{n+1} - t_n \end{cases} \quad \text{and} \quad \begin{cases} u_0 = 1, u_1 = 2w, \\ \forall n \in \mathbb{N}, u_{n+2} = 2wu_{n+1} - u_n \end{cases}$$

Then, there exists a complex number r such that $w = \frac{1}{2}(r + \frac{1}{r})$ and, for all $n \in \mathbb{N}$,

$$t_n = \frac{r^{2n} + 1}{2r^n} \quad \text{and} \quad u_n = \frac{r^{2n+2} - 1}{r^n(r^2 - 1)}.$$

Proof. The characteristic polynomial of (t_n) and (u_n) is $P = X^2 - 2wX + 1$. Let r be a complex root of P . Since the constant term of P is 1, the other complex root of P is $\frac{1}{r}$. Moreover, since $w \notin \{-1, 1\}$, $r \neq \frac{1}{r}$. Thus, there exist complex numbers α, β, γ and δ such that, for all $n \in \mathbb{N}$, $t_n = \alpha r^n + \beta \frac{1}{r^n}$ and $u_n = \gamma r^n + \delta \frac{1}{r^n}$. Since the term of X in P is $-2w$, we have $w = \frac{1}{2}(r + \frac{1}{r})$ and then we deduce from $t_0 = 1$ and $t_1 = w$ that $\alpha = \beta = \frac{1}{2}$ and from $u_0 = 1$ and $u_1 = 2w$ that $\gamma = \frac{r^2}{r^2 - 1}$ and $\delta = \frac{-1}{r^2 - 1}$ which yields the result. □

Theorem 4. Let $n \in \mathbb{N}^*$. All the complex roots of T_n and U_n are real and lie in $(-1, 1)$.

Proof. Let w be a complex root of T_n . Due to Lemma 2 and to the parity of T_n , $w \notin \{-1, 1\}$. Put, for every $k \in \mathbb{N}$, $t_k := T_k(w)$. Then, $t_0 = 1$, $t_1 = w$ and, for all $k \in \mathbb{N}$, $t_{k+2} = 2wt_{k+1} - t_k$. Thus, by Lemma 3, there is a complex number r such that $w = \frac{1}{2}(r + \frac{1}{r})$ and, for all $k \in \mathbb{N}$, $T_k(w) = \frac{r^{2k} + 1}{2r^k}$. Since $T_n(w) = 0$, it follows that $r^{2n} + 1 = 0$ and, in particular, $|r| = 1$. Then, $\bar{r} = \frac{|r|^2}{r} = \frac{1}{r}$ and thus $w = \frac{1}{2}(r + \bar{r}) = \text{Re}(r)$ is a real number. Moreover, $|w| = |\text{Re}(r)| \leq |r| = 1$ and we conclude that $w \in (-1, 1)$. The proof for the roots of U_n is exactly the same. □

3 The roots of Chebyshev polynomials are simple

Lemma 5. For all $n \in \mathbb{N}$, T_n and T_{n+1} are coprime.

Proof. For all $n \in \mathbb{N}$, we put D_n the (monic) gcd of T_n and T_{n+1} . Let $n \in \mathbb{N}$. By definition, D_n divides T_n and T_{n+1} thus, since $T_{n+2} = 2XT_{n+1} - T_n$, D_n divides T_{n+2} too. It follows that D_n divides D_{n+1} . Similarly, since $T_n = 2XT_{n+1} - T_{n+2}$, D_{n+1} divides T_n and thus it divides D_n . Since D_n and D_{n+1} are monic, we conclude that $D_n = D_{n+1}$. Then, (D_n) is constant and, for all $n \in \mathbb{N}$, $D_n = D_0 = \text{gcd}(1, X) = 1$. □

Lemma 6. For all $n \in \mathbb{N}$,

$$T_{n+1} = U_{n+1} - XU_n \quad (1)$$

$$T'_{n+1} = (n+1)U_n \quad (2)$$

$$T_{n+2} = XT_{n+1} - (1 - X^2)U_n \quad (3)$$

$$(1 - X^2)T'_{n+1} + (n+1)(XT_{n+1} - T_n) = 0 \quad (4)$$

$$(n+1)T_{n+1} = XU_n - (1 - X^2)U'_n. \quad (5)$$

Proof. The first three equalities are showed by induction. All these ones are clear if $n = 0$ and $n = 1$ so we only make the induction step. In each case, we assume that the property is true for n and $n+1$.

For (1), we can write

$$\begin{aligned} T_{n+3} &= 2XT_{n+2} - T_{n+1} = 2X(U_{n+2} - XU_{n+1}) - (U_{n+1} - XU_n) \\ &= 2XU_{n+2} - U_{n+1} - X(2XU_{n+1} - U_n) = U_{n+3} - XU_{n+2} \end{aligned}$$

For (2), we use (1):

$$\begin{aligned} T'_{n+3} &= (2XT_{n+2} - T_{n+1})' = 2T_{n+2} + 2XT'_{n+2} - T'_{n+1} \\ &= 2(U_{n+2} - XU_{n+1}) + 2X(n+2)U_{n+1} - (n+1)U_n \\ &= (n+1)(2XU_{n+1} - U_n) + 2U_{n+2} = (n+3)U_{n+2}. \end{aligned}$$

And, finally, for (3),

$$\begin{aligned} T_{n+4} &= 2XT_{n+3} - T_{n+2} = 2X(XT_{n+2} - (1 - X^2)U_{n+1}) - (XT_{n+1} - (1 - X^2)U_n) \\ &= X(2XT_{n+2} - T_{n+1}) - (1 - X^2)(2XU_{n+1} - U_n) \\ &= XT_{n+3} - (1 - X^2)U_{n+2} \end{aligned}$$

Thus, the first three equalities are proved by induction.

For the last two ones, direct calculations succeed, using previous results. For (4), we write:

$$\begin{aligned} (1 - X^2)T'_{n+1} &= (1 - X^2)(n+1)U_n = (n+1)(XT_{n+1} - T_{n+2}) \\ &= (n+1)(XT_{n+1} - (2XT_{n+1} - T_n)) = (n+1)(T_n - XT_{n+1}) \end{aligned}$$

For (5), we differentiate (3). It yields: $T'_{n+2} = T_{n+1} + XT'_{n+1} + 2XU_n - (1 - X^2)U'_n$. But, by (2) and (1), $T'_{n+2} = (n+2)U_{n+1} = (n+2)(T_{n+1} + XU_n)$ and, by (2), $T'_{n+1} = (n+1)U_n$. Then,

$$(n+2)(T_{n+1} + XU_n) = T_{n+1} + X(n+1)U_n + 2XU_n - (1 - X^2)U'_n$$

and thus $(n+1)T_{n+1} = XU_n - (1 - X^2)U'_n$. □

Theorem 7. Let $n \in \mathbb{N}^*$. All the roots of T_n and U_n are simple.

Proof. Let w be a root of T_n . By Lemma 5, T_n and T_{n+1} are coprime thus w is not a root of T_{n+1} . By (3), $T_{n+1}(w) = (1 - w^2)U_{n-1}(w)$ thus $U_{n-1}(w) \neq 0$. But, by (2), $T'_n = nU_{n-1}$ thus $T'_n(w) \neq 0$. Then, w is a simple root of T_n .

Let now w be a root of U_n . By (2), $T'_{n+1}(w) = 0$. Since all the roots of T_{n+1} are simple, $T_{n+1}(w) \neq 0$. But, by (5), $(n+1)T_{n+1}(w) = (w^2 - 1)U'_n(w)$ thus $U'_n(w) \neq 0$ and w is a simple root of U_n . □

4 The roots of Chebyshev polynomials are mostly irrational

Proposition 8. For all $n \in \mathbb{N}$, T_n satisfies the differential equation

$$(1 - X^2)T_n'' - XT_n' + n^2T_n = 0 \quad (6)$$

Proof. We use double induction. The base step is clear. Assume that T_n and T_{n+1} satisfy (6) for a certain nonnegative integer n . Since $T_{n+2} = 2XT_{n+1} - T_n$, we have

$$T_{n+2}' = 2T_{n+1}' + 2XT_{n+1}' - T_n' \quad \text{and} \quad T_{n+2}'' = 4T_{n+1}'' + 2XT_{n+1}'' - T_n''$$

thus

$$\begin{aligned} & (1 - X^2)T_{n+2}'' - XT_{n+2}' + (n+2)^2T_{n+2} \\ &= (1 - X^2)(4T_{n+1}'' + 2XT_{n+1}'' - T_n'') - X(2T_{n+1}' + 2XT_{n+1}' - T_n') + (n+2)^2(2XT_{n+1} - T_n) \\ &= 2X \left[\underbrace{(1 - X^2)T_{n+1}'' - XT_{n+1}' + (n+1)^2T_{n+1}}_{=0} \right] - \left[\underbrace{(1 - X^2)T_n'' - XT_n' + n^2T_n}_{=0} \right] \\ &\quad + 4(1 - X^2)T_{n+1}' - 2XT_{n+1}' + [2(n+1) + 1]2XT_{n+1} - (4n+4)T_n \\ &= 4 \left[(1 - X^2)T_{n+1}' + (n+1)(XT_{n+1} - T_n) \right]. \end{aligned}$$

Then, using (4), we conclude that $(1 - X^2)T_{n+2}'' - XT_{n+2}' + (n+2)^2T_{n+2} = 0$. Hence, the property is true for $n+2$ and Proposition 8 is proved by double induction. \square

Proposition 9. For all $n \in \mathbb{N}^*$,

$$T_n = \frac{n}{2} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k 2^{n-2k}}{n-k} \binom{n-k}{k} X^{n-2k} \quad \text{and} \quad U_n = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k 2^{n-2k} \binom{n-k}{k} X^{n-2k}$$

Proof. Here, we follow [Sny66, p. 24-25]. Let n be a positive integer and put $m = \lfloor \frac{n}{2} \rfloor$. Since T_n is a polynomial in $\mathbb{Z}[X]$ of degree n which has the same parity as n , we can write

$$T_n = \sum_{k=0}^m c_k X^{n-2k}$$

where $c_k \in \mathbb{Z}$ for all $k \in \llbracket 0, m \rrbracket$.

By substituting this expression in (6), one gets

$$(1 - X^2) \sum_{k=0}^m (n-2k)(n-2k-1)c_k X^{n-2k-2} - X \sum_{k=0}^m (n-2k)c_k X^{n-2k-1} + n^2 \sum_{k=0}^m c_k X^{n-2k} = 0$$

i.e., by putting $c_{-1} = 0$,

$$\sum_{k=0}^m ((n-2k+2)(n-2k+1)c_{k-1} - (n-2k)(n-2k-1)c_k - (n-2k)c_k + n^2c_k) X^{n-2k} = 0$$

One deduces that, for all $k \in \llbracket 0, m \rrbracket$,

$$(n-2k+2)(n-2k+1)c_{k-1} + (n^2 - (n-2k)^2)c_k = 0$$

and thus, for all $k \in \llbracket 1, m \rrbracket$

$$c_k = -\frac{(n-2k+1)(n-2k+2)}{4k(n-k)}c_{k-1}.$$

Hence, for all $k \in \llbracket 0, m \rrbracket$,

$$c_k = \frac{(-1)^k}{4^k} c_0 \prod_{j=1}^k \frac{(n-2j+1)(n-2j+2)}{j(n-j)} = \frac{(-1)^k}{4^k} c_0 \frac{n!(n-k-1)!}{k!(n-2k)!(n-1)!}.$$

By an obvious double induction, we get that $c_0 = 2^{n-1}$ thus, for all $k \in \llbracket 0, m \rrbracket$,

$$c_k = (-1)^k 2^{d-1-2k} \frac{d(d-k-1)!}{k!(d-2k)!} = \frac{d}{2} \frac{(-1)^k 2^{d-2k}}{d-k} \binom{d-k}{k}$$

as announced. Moreover, by (2),

$$\begin{aligned} U_n &= \frac{1}{n+1} T'_{n+1} = \frac{1}{n+1} \cdot \frac{n+1}{2} \sum_{k=0}^{\lfloor \frac{n+1}{2} \rfloor} \frac{(-1)^k 2^{n+1-2k}}{n+1-k} \binom{n+1-k}{k} (n+1-2k) X^{n-2k} \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k 2^{n-2k} \frac{(n-k)!}{k!(n-2k)!} X^{n-2k} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k 2^{n-2k} \binom{n-k}{k} X^{n-2k} \end{aligned}$$

because $\lfloor \frac{n+1}{2} \rfloor = \lfloor \frac{n}{2} \rfloor$ if n is even and $n+1-2\lfloor \frac{n+1}{2} \rfloor = 0$ if n is odd. \square

Theorem 10. Let n be a positive integer.

1. The roots of T_n are all irrational if n is even and 0 is the only rational root of T_n if n is odd.
2. The only possible rational roots of U_n are 0, $\frac{1}{2}$ and $-\frac{1}{2}$. Moreover, 0 is a root of U_n if and only if n is odd and $\frac{1}{2}$ and $-\frac{1}{2}$ are roots of U_n if and only if $n \equiv 2 \pmod{3}$.

Proof. Let us put $m = \lfloor \frac{n}{2} \rfloor$.

1. Assume that α is a rational root of T_n . Thus, 2α is a root of $2T_n(\frac{X}{2})$. But, according with Proposition 9,

$$2T_n\left(\frac{X}{2}\right) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \frac{n}{n-k} \binom{n-k}{k} X^{n-2k}.$$

Note that, for all $k \in \llbracket 0, m \rrbracket$, $\frac{n}{n-k} \binom{n-k}{k} = \binom{n-k}{k} + \frac{k}{n-k} \binom{n-k}{k} = \binom{n-k}{k} + \binom{n-k-1}{k-1}$ so $2T_n(\frac{X}{2})$ has integral coefficients. Since its leading term is 1, we deduce from a classic result about algebraic integers (see, for example, [NZM91, Corollary 6.14, p. 308]) that 2α is an integer. Moreover, by Theorem 4, α lies in $(-1, 1)$ thus $\alpha \in \left\{0, -\frac{1}{2}, \frac{1}{2}\right\}$.

Let us recall that by Lemma 3, $T_n(\alpha) = \frac{r^{2d}+1}{2r^d}$ where r is a root of $X^2 - 2\alpha X + 1$. If $\alpha = 0$, we can take $r = i$ and thus $T_n(\alpha) = -\frac{(-1)^{n+1}}{2}$ and we conclude that 0 is a root of T_n if and only if n is odd. If $\alpha = -\frac{1}{2}$ then we can take $r = j := -\frac{1}{2} + i\frac{\sqrt{3}}{2}$. Therefore, $T_n(\alpha) = 0$ if and only if $j^{2n} = -1$. But $j^3 - 1 = (j-1)(j^2 + r + 1) = 0$ thus $j^3 = 1$. It follows that the sequence $(j^d)_{d \in \mathbb{N}}$ is 3-periodic and since $j^0 = 1$, $j = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ and $j^2 = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$ we conclude that $j^{2n} \neq -1$, i.e., $-\frac{1}{2}$ is not a root of T_n . Finally, due to the parity of T_n , $\frac{1}{2}$ is not a root of T_n either.

2. Assume that α is a rational root of U_n . By considering $U_n(\frac{X}{2})$, one proves similarly that $\alpha \in \left\{0, -\frac{1}{2}, \frac{1}{2}\right\}$. The case $\alpha = 0$ leads to the same conclusion. Nonetheless, the case $\alpha = -\frac{1}{2}$ is a bit different. Indeed, one has $U_n(\alpha) = \frac{j^{2n+2}-1}{j^n(j^2-1)}$ and, according to the periodic values of the sequence (j^d) , one sees that $U_n(\alpha) = 0$ if and only if 3 divides $2n+2$ that is to say $n \equiv 2 \pmod{3}$. The same argument of parity concludes the proof. \square

The previous proof depends largely on the result of section 2 and especially on Theorem 4 which ensures that $s \in \{0, -1, 1\}$. In fact, there is a way to derive the irrationality of non-zero roots of T_n by using only the result of the present section. The proof is based on the shifted Chebyshev polynomials.

Definition 11. For all nonnegative integer n , one defines the shifted Chebyshev polynomials (of first kind) T_n^* by:

$$T_n^* = T_n(2X - 1).$$

Proposition 12. For all nonnegative integer n , $T_n^*(X^2) = T_{2n}$.

Proof. One uses double induction. The result is clear if $n = 0$ and $n = 1$. Assume the relation is true for n and $n + 1$. Then,

$$\begin{aligned} T_{n+2}^*(X^2) &= T_{n+2}(2X^2 - 1) = 2(2X^2 - 1)T_{n+1}(2X^2 - 1) - T_n(2X^2 - 1) \\ &= 2(2X^2 - 1)T_{n+1}^*(X^2) - T_n^*(X^2) = 2(2X^2 - 1)T_{2n+2} - T_{2n} \\ &= 4X^2T_{2n+2} - (T_{2n+2} + T_{2n}) - T_{2n+2} = 4X^2T_{2n+2} - 2XT_{2n+1} - T_{2n+2} \\ &= 2X(2XT_{2n+2} - T_{2n+1}) - T_{2n+2} = 2XT_{2n+3} - T_{2n+2} \\ &= T_{2n+4}. \end{aligned}$$

Hence, the relation is true for $n + 2$ and Proposition 12 is proved by induction. \square

Theorem 13. Let n be a positive integer. The only possible rational root of T_n^* is $\frac{1}{2}$.

Proof. It follows from Propositions 9 and 12 that

$$T_n^*(X^2) = n \sum_{k=0}^n \frac{(-1)^k 2^{2n-2k}}{2n-k} \binom{2n-k}{k} X^{2n-2k} = n \sum_{k=0}^n \frac{(-1)^{n-k} 2^{2k}}{n+k} \binom{n+k}{2k} (X^2)^k$$

thus

$$2T_n^*\left(\frac{X}{4}\right) = \sum_{k=0}^n (-1)^{n-k} \frac{2n}{n+k} \binom{n+k}{2k} X^k = \sum_{k=0}^n (-1)^{n-k} \left[2 \binom{n+k}{n-k} - \binom{n+k-1}{n-k} \right] X^k.$$

Hence, $2T_n^*\left(\frac{X}{4}\right)$ has integral coefficients. Assume that α is a rational root of T_n^* . Note that the terms of T_n^* alternate in sign and the constant term of T_n^* is non-zero so $\alpha > 0$. Since $2T_n^*\left(\frac{X}{4}\right)$ is a monic polynomial and has constant term 2, the same argument that for T_n yields that 4α is an integer dividing 2 so $\alpha \in \{\frac{1}{2}, \frac{1}{4}\}$. Moreover, due to the parity of T_n , $1 - \alpha$ is also a rational root of T_n^* so we conclude that the only possible rational root of T_n^* is $\frac{1}{2}$. \square

As an immediate corollary of the previous theorem, we deduce that T_n has no rational root if n is even and 0 is the only rational root of T_n if n is odd.

References

- [MH03] J. C. Mason and D. C. Handscomb, *Chebyshev polynomials*, Chapman & Hall/CRC, Boca Raton, FL, 2003.
- [NZM91] I. Niven, H. S. Zuckerman, and H. L. Montgomery, *An Introduction to the Theory of Numbers*, 5^e ed., John Wiley & Sons, Inc., 1991.
- [Pon18] L. Ponton, *Sur l'irrationalité des racines de certains polynômes usuels (I)*, R.M.S. **129-1** (2018), 26–47.
- [Sny66] M. A. Snyder, *Chebyshev methods in numerical approximation*, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1966.