

Energy corrections due to the noncommutative phase-space of the charged isotropic harmonic oscillator in a uniform magnetic field in 3D

Muhittin Cenk Eser* and Mustafa Riza†

Department of Physics, Eastern Mediterranean University, 99628 Famagusta, North Cyprus

(Dated: January 14, 2022)

In this study, we investigate the effects of noncommutative Quantum Mechanics in three dimensions on the energy levels of a charged isotropic harmonic oscillator in the presence of a uniform magnetic field in the z -direction. The extension of this problem to three dimensions proves to be non-trivial. We obtain the first-order corrections to the energy-levels in closed form in the low energy limit of weak noncommutativity. The most important result we can note is that all energy corrections due to noncommutativity are negative and their magnitude increase with increasing Quantum numbers and magnetic field.

I. INTRODUCTION

With Heisenberg's introduction of the Uncertainty Principle [1], the classical paradigm that position and momentum commute at no cost is crushed. Furthermore, the discussion of a charged particle in an electromagnetic field in the framework of Quantum Mechanics inevitably leads to the introduction of the kinetic momentum operator, which in contrast to the canonical momentum operator, does not commute. The noncommutativity of the kinetic momentum operator indicates that the magnetic field's presence modifies momentum space. These two facts, emerging from Quantum Mechanics' nature, evidently bring up the question, is the assumption of the commutation of the position and momentum operators among themselves an accurate assumption? Or, under which conditions is the assumption of the vanishing commutators of $[x_i, x_j] = 0$ and $[p_i, p_j] = 0$ correct? The field dedicated to the study of non vanishing position and momentum commutators is called Noncommutative Quantum Mechanics. It is evident that in the energy domain of textbook quantum mechanics, the commutators in position and momentum are vanishing. Once the energy is pushed closer towards the Planck Energy, the effects of noncommutativity of the momentum and position operators can be observed [2].

In his pioneering work, Hartland Snyder [3] noticed that in Field Theory Lorenz invariance does not necessarily require noncommutativity of the position and momentum operators. Snyder's work [3] lead to the detailed discussion of the Quantum Field Theory in noncommutative spaces by Szabo [4] and Seiberg and Witten [5]. One of the first formulations of non-relativistic Quantum Mechanics in noncommutative space was presented by Chaturvedi *et al.* in [6]. Based on these ideas, Noncommutative Quantum Mechanics was proposed by Gamboa *et al.* in [7]. Noncommutativity is generally associated with the effect of the geometry of the space [8, 9]. The Klein-Gordon, the Schrödinger, and Pauli-Dirac oscilla-

tors in noncommutative phase-space have been studied by Jian-Hua *et al.* in [10] and Santos and de Melo in [11]. Furthermore, more fundamental problems like the Bohr-van-Leeuwen theorem [12], stating explicitly that magnetization is a purely Quantum Mechanical effect, is discussed in the framework of Noncommutative Quantum Mechanics.

Moreover, the noncommutative geometry in space seems to be a reasonable approach to the limitation of the position uncertainty leading us to the General Uncertainty Principle discussed e.g., by Kempf *et al.* in [13], Das *et al.* in [14], or Bosso *et al.* [15]. Dey *et al.* [16] showed explicitly that noncommutativity of the phase space gives rise to minimum length and minimum momentum uncertainties. Dey *et al.* [2] also stated that the minimum length from noncommutativity is consistent with the General Uncertainty Principle in [13], yielding to $(\Delta x_i)_{min} = l_p \frac{\hbar}{2} \sqrt{\eta \theta}$, where $l_p = \sqrt{\hbar G/c^3} \sim 10^{-35}m$ denotes the Planck-Length. Based on the experimental bound for θ the minimal length is of the order of $10^6 l_p$. Based on the experimental setup proposed in [2] the Planck Length as minimal length can be reached and effects of Quantum Gravity can be observed. Furthermore, the idea of minimum length for resolving the UV singularity motivated noncommutative space-time in Quantum Field Theory. Studies in String Theory [17–19] and in Loop Quantum Gravity [20–22] support this idea. However, only the formation of a black hole provides the necessary conditions for arbitrarily high precision in the position [23, 24]. Consequently, the physical limitation on the shortest distance leads to a UV cutoff [25]. Moreover, the noncommutative phase-space and its space-time symmetry in $2 + 1$ dimensions have been discussed by Kang and Sayipjamal [26].

The relationship between the General Uncertainty Principle as proposed in [13, 15] and noncommutative Quantum Mechanics needs to be analyzed in detail, as both approaches lead to the concept of minimal length uncertainty primarily, and minimal momentum uncertainty in the second instance. A minimal uncertainty in length and momentum leads also to an important conclusion in information theory, namely that the total information in the universe is bounded. This will form a

* muhittin.eser@emu.edu.tr

† mustafa.riza@emu.edu.tr

reasonable extension to Bekenstein's work [27].

This study is dedicated to the noncommutative 3D isotropic harmonic oscillator in a homogeneous magnetic field. The application of the magnetic field to the isotropic harmonic oscillator turns the isotropic harmonic oscillator into an anisotropic harmonic oscillator. The anisotropic harmonic oscillator has a wide range of applications in mathematical physics, Quantum theory, and condensed matter physics commutative as well as non-commutative. In the commutative case, we can find a wide field of applications in the literature. e.g., Petreska has applied the concept of the anisotropic harmonic oscillator to different problems in Quantum Physics [28–32]. On the other hand, it serves also as a perfect model in the discussion of Quantum dots in condensed matter physics [33–43] and atomic physics [44–53]. In the noncommutative case, the discussions are mainly carried out in the noncommutative plane, i.e. noncommutativity is only employed to the xy -plane for both the position and momentum. With respect to this Gao-Feng *et al.* solve the isotropic charged harmonic oscillator in a uniform magnetic field 2D Noncommutative Quantum Mechanics [54]. The isotropic harmonic oscillator in a constant magnetic field is a subset of the anisotropic harmonic oscillator. The anisotropic harmonic oscillator was also discussed under different aspects in the framework of noncommutative Quantum Mechanics [55–57] explicitly. Muhuri *et al.* show in [55] that entanglement induces noncommutativity in space in the example of the anisotropic harmonic oscillator. Furthermore, Ghosh and Nath discuss the impact of noncommutativity on the uncertainty and the Shannon entropy for the 2D anisotropic harmonic oscillator in presence of a magnetic field [56]. The 2D noncommutative anisotropic harmonic oscillator in a homogeneous magnetic field has been discussed by Nath and Roy in [57]. In contrast to the studies cited, this study illuminates the energy corrections due to 3D noncommutativity as a function of the magnetic field in the low energy limit according to [58]. Additionally, various publications are dedicated to the charged Quantum harmonic oscillator in the presence of a constant or time-varying electromagnetic field in noncommutative Quantum Mechanics e.g., [59]. Moreover, the magnetic field's impact on noncommutativity has been discussed in numerous works, especially in the context of the Landau problem [60–72]. There are several discussions on the noncommutative Quantum Hall effect [73–76] as well. The minimally coupled charged harmonic oscillator to the magnetic field in a noncommutative plane has been studied extensively by Jing and Chen [77]. Some more mathematical discussions on the noncommutativity of Quantum Mechanics can be found e.g., in [78–80]. Finally, Hassanabadi *et al.* studied the Dirac oscillator in the presence of the Aharonov-Bohm effect in noncommutative and commutative spaces [81].

The fact that the magnetic field modifies the momentum space leading to the noncommutativity of the kinetic momentum operator on the one hand, and various studies

related to the General Uncertainty Principle, backed also by String theory, suggest that the existence of a minimal length on the other hand, support the approach in Noncommutative Quantum Mechanics including the noncommutativity of the position and the momentum operators. The noncommutativity of the position and momentum operators indicate a minimum length and a minimum momentum. Continuing this train of thought will lead to the conclusion that all physical quantities are quantized and have a minimum size.

Throughout this manuscript we will denote \hat{x}_i as the noncommutative position operator and \hat{p}_i as the noncommutative momentum operator in contrast to the standard position operator x_i and the standard momentum operator p_i . The basic properties of noncommutative phase-space according to e.g., Gamboa *et al.* [7] stating the commutator relationships of the noncommutative position operators and the noncommutative momentum operators as:

$$[\hat{x}_i, \hat{x}_j] = i\theta_{ij}, \quad [\hat{p}_i, \hat{p}_j] = i\eta_{ij} \quad (1)$$

where θ_{ij} and η_{ij} are both antisymmetric tensors. For further reading on antisymmetric tensors, we refer to [82].

Consequently, as one can verify easily, the relationship between noncommutative operators \hat{x}_i and \hat{p}_i with their commutative counterparts can be written as

$$\hat{x}_i = \alpha x_i - \frac{1}{2\alpha\hbar} \theta_{ij} p_j \quad (2)$$

$$\hat{p}_i = \alpha p_i + \frac{1}{2\alpha\hbar} \eta_{ij} x_j, \quad (3)$$

where $\alpha \in (0, 1)$ is the scaling constant related to the noncommutativity of the phase-space and η_{ij} , and θ_{ij} are antisymmetric tensors. So, generally, we can express the tensors η_{ij} and θ_{ij} as following:

$$\eta_{ij} = \eta \lambda_{ij}, \quad (4)$$

$$\theta_{ij} = \theta \lambda_{ij}, \quad (5)$$

where λ_{ij} denotes an antisymmetric tensor [82].

Mathematically, the noncommutativity of the base manifold can be realized by application the Weyl-Moyal star product [83]

$$\begin{aligned} (f \star g)(x, p) &= e^{i\frac{1}{2\alpha^2} \theta_{ij} \partial_i^x \partial_j^x + i\frac{1}{2\alpha^2} \eta_{ij} \partial_i^p \partial_j^p} f(x)g(y) = \\ &= f(x, p)g(x, p) + \frac{i\theta_{ij}}{2\alpha^2} \partial_i^x f \partial_j^x g \Big|_{x_i=x_j} + \frac{i\eta_{ij}}{2\alpha^2} \partial_i^p f \partial_j^p g \Big|_{p_i=p_j} + \\ &\quad + \mathcal{O}(\theta_{ij}^2) + \mathcal{O}(\eta_{ij}^2) + \mathcal{O}(\theta_{ij}\eta_{ij}) \end{aligned} \quad (6)$$

So, the shift from ordinary Quantum Mechanics to Noncommutative Quantum Mechanics is performed by employing the Weyl-Moyal product (6) instead of the ordinary product. Hence, the Noncommutative Time-Independent Schrödinger Equation becomes

$$H(x, p) \star \psi(x) = E\psi(x). \quad (7)$$

By employing the Bopp's shift [84], we can turn the Weyl-Moyal product again to the ordinary product by substituting x and p in the noncommutative equation by \hat{x} and \hat{p} , namely

$$H(x, p) \star \psi(x) = H(\hat{x}, \hat{p})\psi(x). \quad (8)$$

Harko and Liang [58] state that the noncommutativity parameters η and θ can be considered as energy-dependent and that both become sufficiently small in the low energy limit. Employing this fact, gives the justification of the possibility of the application of perturbation theory in the low energy limit.

In light of this, we will discuss the noncommutative charged harmonic oscillator in the presence of a uniform magnetic field employing noncommutativity to all three spacial parameters by including also the z -direction into the noncommutative framework. Therefore, first we will discuss the change to the noncommutative algebra by considering the commutator $[\hat{x}_i, \hat{p}_j]$ in the noncommutative plane and space in section II. In the next section, we will discuss the noncommutative Hamiltonian of the charged particle in a 3D isotropic harmonic oscillator in the presence of a uniform magnetic field where we will expand the Hamiltonian in terms of θ and η . As this Hamiltonian proves to be non-trivial, the corrections to the eigenenergies due to the magnitude of the magnetic field will be calculated in section IV in first-order perturbation theory in η and θ , i.e. in the domain of weak noncommutativity in the low energy limit. Finally, we will carry out a short analysis of the corrections of the eigenenergies in section V on the dependence of the energy corrections on the magnitude of the magnetic field for different values of the Quantum numbers and close this study with some concluding remarks.

II. THE COMMUTATOR $[\hat{x}_i, \hat{p}_j]$ IN THE NONCOMMUTATIVE PLANE AND SPACE

For completeness, let us recall the commutators $[\hat{x}_i, \hat{x}_j]$ and $[\hat{p}_i, \hat{p}_j]$.

$$[\hat{x}_i, \hat{x}_j] = i\theta_{ij}, \quad [\hat{p}_i, \hat{p}_j] = i\eta_{ij} \quad (9)$$

where θ_{ij} and η_{ij} are both antisymmetric tensors.

Yielding to the relationship between noncommutative operators \hat{x}_i and \hat{p}_i with their commutative counterparts

$$\begin{aligned} \hat{x}_i &= \alpha x_i - \frac{1}{2\alpha\hbar}\theta_{ij}p_j \\ \hat{p}_i &= \alpha p_i + \frac{1}{2\alpha\hbar}\eta_{ij}x_j, \end{aligned}$$

where $\alpha \in (0, 1)$ is the scaling constant related to the noncommutativity of the phase-space and η_{ij} , and θ_{ij} are

antisymmetric tensors. So, generally, we can express the tensors η_{ij} and θ_{ij} as following:

$$\begin{aligned} \eta_{ij} &= \eta\lambda_{ij}, \\ \theta_{ij} &= \theta\lambda_{ij}, \end{aligned}$$

where λ_{ij} denotes an antisymmetric tensor.

The difference between the noncommutative plane and space is manifested in the definition of the antisymmetric tensor λ_{ij} . In the noncommutative plane the antisymmetric tensor λ_{ij} is given as:

$$\lambda_{ij} = \begin{cases} 1 & \text{if } ij = 12 \\ -1 & \text{if } ij = 21 \\ 0 & \text{else} \end{cases} \quad (10)$$

By extending the discussion to the 3D noncommutative space, a redefinition of the epsilon tensor is needed $\tilde{\lambda}_{ij}$ is defined as

$$\tilde{\lambda}_{ij} = \begin{cases} 1 & \text{if } ij = 12, 23, 31 \\ -1 & \text{if } ij = 21, 32, 13 \\ 0 & \text{else} \end{cases} \quad (11)$$

Let us first discuss the impact of the extension of the antisymmetric tensor from the noncommutative plane λ_{ij} to the noncommutative space $\tilde{\lambda}_{ij}$ on the commutator $[\hat{x}_i, \hat{p}_j]$. The commutator of the noncommutative position and momentum operators can be calculated straight forward independent of the noncommutativity covering only the plane or the whole space

$$\begin{aligned} [\hat{x}_i, \hat{p}_j] &= \left[\alpha x_i - \frac{1}{2\alpha\hbar}\theta_{ij}p_j, \alpha p_i + \frac{1}{2\alpha\hbar}\eta_{ij}x_j \right] = \\ &= i\hbar\alpha^2\delta_{ij} + i\frac{\theta\eta}{4\alpha^2\hbar}\lambda_{i\mu}\lambda_{j\mu}. \end{aligned} \quad (12)$$

For the 3D noncommutative space $\lambda_{i\mu}\lambda_{j\mu}$ is substituted by $\tilde{\lambda}_{i\mu}\tilde{\lambda}_{j\mu}$.

The difference between the two cases of the noncommutative plane and the noncommutative space is manifested in the product of the antisymmetric tensors $\lambda_{i\mu}\lambda_{j\mu}$ and $\tilde{\lambda}_{i\mu}\tilde{\lambda}_{j\mu}$. Using the properties of the λ tensor (10) for the noncommutative plane, we get for this product

$$\lambda_{i\mu}\lambda_{j\mu} = -\delta_{ij}, \quad (13)$$

where δ_{ij} denotes the Kronecker- δ . Whereas the product of the two λ tensors (11) in the noncommutative space (3D) is

$$\tilde{\lambda}_{i\mu}\tilde{\lambda}_{j\mu} = -3\delta_{ij} + 1. \quad (14)$$

With (13) we get for the commutator (12) in the noncommutative plane

$$[\hat{x}_i, \hat{p}_j] = i\hbar\alpha^2\delta_{ij} - i\frac{\theta\eta}{4\alpha^2\hbar}\delta_{ij}, \quad (15)$$

and with (14) we get for the commutator (12) in the noncommutative space

$$[\hat{x}_i, \hat{p}_j] = i\hbar\alpha^2\delta_{ij} - i\frac{\theta\eta}{4\alpha^2\hbar}(3\delta_{ij} - 1). \quad (16)$$

Ergo, the first effect of the extension from the noncommutative plane (2D) to the noncommutative space (3D) can be seen that the commutator in the plane is non-zero if $i = j$. In contrast, the commutator in the noncommutative space never vanishes.

III. 3D NONCOMMUTATIVE CHARGED HARMONIC OSCILLATOR IN A UNIFORM MAGNETIC FIELD

Our starting point is the commutative Hamiltonian for the charged isotropic harmonic oscillator presence of a uniform magnetic field.

$$H_0(x, p) = \frac{1}{2m} \left(\vec{p} - \frac{q}{c} \vec{A} \right)^2 + \frac{1}{2}m\omega^2 (x^2 + y^2 + z^2) \quad (17)$$

Without loss of generality, we will choose the direction of the uniform magnetic field in the z -direction, i.e., $\vec{B} = B\hat{k}$ yielding to $\vec{A}(\vec{x}, t) = \frac{1}{2}(-yB\hat{i} + xB\hat{j})$ in Coulomb gauge. So, our Hamiltonian $H_0(x, p)$ modifies to

$$H_0(x, p) = \frac{1}{2m} \left(\left(p_x + \frac{qB}{2c}y \right)^2 + \left(p_y - \frac{qB}{2c}x \right)^2 + p_z^2 \right) + \frac{1}{2}m\omega^2 (x^2 + y^2 + z^2). \quad (18)$$

After expanding the Hamiltonian (18) and regrouping the terms we get

$$H_0(x, p) = \frac{1}{2m} (p_x^2 + p_y^2 + p_z^2) - \frac{1}{2}\omega_c L_z + \frac{1}{2}m\tilde{\omega}^2 (x^2 + y^2) + \frac{1}{2}m\omega^2 z^2, \quad (19)$$

where $L_z = xp_y - yp_x$ is the z -component of the angular momentum operator, $\omega_c = \frac{qB}{mc}$ the cyclotron frequency, and $\tilde{\omega}^2 = \omega^2 + \frac{\omega_c^2}{4}$ is the modified frequency of the harmonic oscillator in the xy -plane. Hence, the problem turns into the problem of an anisotropic harmonic oscillator. From equation (8), we know that the Weyl-Moyal product can be turned into a standard product by substituting commutative x and p by the noncommutative operators \hat{x} and \hat{p} , so let us first consider the Hamiltonian $H_0(\hat{x}, \hat{p})$.

$$H_0(\hat{x}, \hat{p}) = \frac{1}{2m} (\hat{p}_x^2 + \hat{p}_y^2 + \hat{p}_z^2) - \frac{1}{2}\omega_c \hat{L}_z + \frac{1}{2}m\tilde{\omega}^2 (\hat{x}^2 + \hat{y}^2) + \frac{1}{2}m\omega^2 \hat{z}^2 \quad (20)$$

All noncommutative operators in the noncommutative phase-space (3D) can be stated explicitly using (2) and (3) together with (4) and (5), respectively.

$$\hat{x} = \alpha x - \frac{\theta}{2\alpha\hbar}p_y + \frac{\theta}{2\alpha\hbar}p_z \quad (21)$$

$$\hat{y} = \alpha y - \frac{\theta}{2\alpha\hbar}p_z + \frac{\theta}{2\alpha\hbar}p_x \quad (22)$$

$$\hat{z} = \alpha z - \frac{\theta}{2\alpha\hbar}p_x + \frac{\theta}{2\alpha\hbar}p_y \quad (23)$$

$$\hat{p}_x = \alpha p_x + \frac{\eta}{2\alpha\hbar}y - \frac{\eta}{2\alpha\hbar}z \quad (24)$$

$$\hat{p}_y = \alpha p_y + \frac{\eta}{2\alpha\hbar}z - \frac{\eta}{2\alpha\hbar}x \quad (25)$$

$$\hat{p}_z = \alpha p_z - \frac{\eta}{2\alpha\hbar}x - \frac{\eta}{2\alpha\hbar}y \quad (26)$$

Based on the position and momentum operators defined in equations (21)-(26), we can construct all other operators needed in this calculation.

As a consequence, the noncommutative angular momentum operator \hat{L}_z can be stated explicitly as following

$$\begin{aligned} \hat{L}_z = \hat{x}\hat{p}_y - \hat{y}\hat{p}_x = & \alpha^2 L_z + \frac{\theta}{2\hbar} (-p_x^2 - p_y^2 + p_x p_z + p_y p_z) + \\ & + \frac{\eta}{2\hbar} (-x^2 - y^2 + xz + yz) + \frac{\theta\eta}{4\alpha^2\hbar^2} (L_x + L_y + L_z). \end{aligned} \quad (27)$$

Furthermore, the sum of the squares of the components of the noncommutative momentum operator $\hat{p}_x^2 + \hat{p}_y^2 + \hat{p}_z^2$ becomes

$$\begin{aligned} \hat{p}_x^2 + \hat{p}_y^2 + \hat{p}_z^2 = & \alpha^2 (p_x^2 + p_y^2 + p_z^2) - \frac{\eta}{\hbar} (L_x + L_y + L_z) + \\ & + \frac{\eta^2}{2\alpha^2\hbar^2} (x^2 - xy + y^2 - xz - yz + z^2), \end{aligned} \quad (28)$$

and the sum of the squares of the x and y components of the noncommutative squared position operator $\hat{x}^2 + \hat{y}^2$ is

$$\begin{aligned} \hat{x}^2 + \hat{y}^2 = & \alpha^2 (x^2 + y^2) + \frac{\theta}{\hbar} (-L_z + (x - y)p_z) + \\ & + \frac{\theta^2}{4\alpha^2\hbar^2} (p_x^2 + p_y^2 + 2p_z^2 - 2p_x p_z - 2p_y p_z), \end{aligned} \quad (29)$$

and finally square of the z component of the noncommutative position operator \hat{z}^2 yields to

$$\hat{z}^2 = \alpha^2 z^2 + \frac{\theta}{\hbar} z (p_y - p_x) + \frac{\theta^2}{4\alpha^2\hbar^2} (p_x - p_y)^2. \quad (30)$$

Substituting (27)-(30) into (20) gives the noncommutative Hamiltonian in the commutative algebra. After regrouping and summarizing all terms, we get the expansion of noncommutative Hamiltonian in the commutative space with respect to the noncommutativity parameters θ and η as

$$H_0(\hat{x}, \hat{p}) = \alpha^2 H_0(x, p) + \frac{\eta}{\hbar} H_\eta(x, p) + \frac{\theta}{\hbar} H_\theta(x, p) + \frac{\eta\theta}{\hbar^2} H_{\eta\theta}(x, p) + \frac{\eta^2}{\hbar^2} H_{\eta^2}(x, p) + \frac{\theta^2}{\hbar^2} H_{\theta^2}(x, p), \quad (31)$$

with

$$H_\eta = -\frac{1}{2m} (L_x + L_y + L_z) - \frac{1}{4}\omega_c (-x^2 - y^2 + xz + yz), \quad (32)$$

$$H_\theta = -\frac{1}{4}\omega_c (-p_x^2 - p_y^2 + p_x p_z + p_y p_z) + \frac{1}{2}m\tilde{\omega}^2 (-L_z + (x - y)p_z) + \frac{1}{2}m\omega^2 z (p_y - p_x), \quad (33)$$

$$H_{\eta\theta} = \frac{\omega_c}{8\alpha^2} (L_x + L_y + L_z), \quad (34)$$

$$H_{\eta^2} = \frac{1}{4m\alpha^2} (x^2 - xy + y^2 - xz - yz + z^2), \quad (35)$$

$$H_{\theta^2} = \frac{1}{4\alpha^2} \left[\frac{1}{2}m\tilde{\omega}^2 (p_x^2 + p_y^2 + 2p_z^2 - 2p_x p_z - 2p_y p_z) + \frac{1}{2}m\omega^2 (p_x - p_y)^2 \right]. \quad (36)$$

Obviously, for $\alpha = 1, \theta = \eta = 0$ we return to the well known commutative case.

IV. PERTURBATIVE APPROACH

According to Harko et al. [58], the contribution of the second-order terms η^2 , θ^2 , and $\eta\theta$ can be considered as small compared to the terms in η and θ in the low energy limit. Consequently, we can determine the effect of the noncommutativity on the binding energy by employing first-order perturbation theory.

To determine the impact of noncommutativity on the energy levels of a charged harmonic oscillator in 3D in the presence of a uniform magnetic field, we first have to revisit the well-known commutative case. The Hamiltonian in the commutative case in cylindrical coordinates is then given as

$$H_0(x, p) = -\frac{\hbar^2}{2m} \left(\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial}{\partial \rho} \right) + \frac{1}{\rho^2} \frac{\partial^2}{\partial \varphi^2} \right) - \frac{1}{2}\omega_c \frac{\hbar}{i} \frac{\partial}{\partial \varphi} + \frac{1}{2}m \left(\omega^2 + \frac{\omega_c^2}{4} \right) \rho^2 - \frac{\hbar^2}{2m} \frac{\partial^2}{\partial z^2} + \frac{1}{2}m\omega^2 z^2, \quad (37)$$

where $x = \rho \cos \varphi$, $y = \rho \sin \varphi$ consequently $\rho^2 = x^2 + y^2$, $L_z = \frac{\hbar}{i} \frac{\partial}{\partial \varphi}$, and $p_\rho^2 = -\hbar^2 \left(\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial}{\partial \rho} \right) \right)$. With $\tilde{\omega}^2 = \omega^2 + \frac{\omega_c^2}{4}$ we get

$$H_0(x, p) = -\frac{\hbar^2}{2m} \left(\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial}{\partial \rho} \right) + \frac{1}{\rho^2} \frac{\partial^2}{\partial \varphi^2} \right) - \frac{1}{2}\omega_c \frac{\hbar}{i} \frac{\partial}{\partial \varphi} + \frac{1}{2}m\tilde{\omega}^2 \rho^2 - \frac{\hbar^2}{2m} \frac{\partial^2}{\partial z^2} + \frac{1}{2}m\omega^2 z^2. \quad (38)$$

In cylindrical coordinates, the time-independent Schrödinger equation for a particle in an isotropic harmonic oscillator in the presence of a uniform magnetic field can be solved by separation of variables as

$$\psi_{n_\rho, \mu, n_z}(x) = \chi(\rho) e^{i\mu\varphi} \zeta(z). \quad (39)$$

After substitution into the time independent Schrödinger equation we get the eigenfunction as:

$$\zeta(z) = \frac{1}{\sqrt{2^n n!}} \left(\frac{m\omega}{\pi \hbar} \right)^{1/4} e^{-\frac{m\omega z}{2\hbar}} H_{n_z} \left(\sqrt{\frac{m\omega}{\hbar}} z \right) \quad (40)$$

$$\chi(\rho) = A_1 \left(\sqrt{2}\rho \right)^{|\mu|} e^{-\frac{m\tilde{\omega}\rho^2}{2\hbar}} U \left(-n_\rho, 1 + |\mu|, \frac{m\tilde{\omega}\rho^2}{2\hbar} \right) + A_2 \left(\sqrt{2}\rho \right)^{|\mu|} e^{-\frac{m\tilde{\omega}\rho^2}{2\hbar}} L_{n_\rho}^{|\mu|} \left(-n_\rho, \frac{m\tilde{\omega}\rho^2}{2\hbar} \right), \quad (41)$$

where $H_n(x)$ denotes the Hermite Polynomials, $U(n, m, x)$ the confluent hypergeometric function of second kind, and $L_n^\alpha(m, x)$ the generalized Laguerre Polynomial [85]. The corresponding eigenvalues are given as:

$$E_{n_\rho, \mu, n_z} = \hbar\tilde{\omega}(2n_\rho + |\mu| + 1) + \frac{1}{2}\hbar\omega_c\mu + \hbar\omega \left(n_z + \frac{1}{2} \right). \quad (42)$$

The corrections to the binding energy for weak non-commutativity in first-order perturbation theory are then according to (31) given as

$$\Delta E_{n_\rho, \mu, n_z}^{(1)} = \frac{\eta}{\hbar} \langle n_\rho, \mu, n_z | H_\eta | n_\rho, \mu, n_z \rangle + \frac{\theta}{\hbar} \langle n_\rho, \mu, n_z | H_\theta | n_\rho, \mu, n_z \rangle \quad (43)$$

Due to the symmetry of the problem, all following matrix elements vanish:

$$\begin{aligned} \langle n_\rho, \mu, n_z | L_x | n_\rho, \mu, n_z \rangle &= \langle n_\rho, \mu, n_z | L_y | n_\rho, \mu, n_z \rangle = \\ &= \langle n_\rho, \mu, n_z | xz | n_\rho, \mu, n_z \rangle = \langle n_\rho, \mu, n_z | yz | n_\rho, \mu, n_z \rangle = \\ &= \langle n_\rho, \mu, n_z | p_x p_z | n_\rho, \mu, n_z \rangle = \langle n_\rho, \mu, n_z | p_y p_z | n_\rho, \mu, n_z \rangle = \\ &= \langle n_\rho, \mu, n_z | y p_z | n_\rho, \mu, n_z \rangle = \langle n_\rho, \mu, n_z | x p_z | n_\rho, \mu, n_z \rangle = \\ &= \langle n_\rho, \mu, n_z | p_x | n_\rho, \mu, n_z \rangle = \langle n_\rho, \mu, n_z | p_y | n_\rho, \mu, n_z \rangle = 0 \end{aligned}$$

So, the only matrix elements that are non-vanishing are

$$\begin{aligned} \Delta E_{n_\rho, \mu, n_z}^{(1)} &= \frac{\eta}{\hbar} \langle n_\rho, \mu, n_z | -\frac{1}{2m} L_z - \frac{\omega_c}{4} \rho^2 | n_\rho, \mu, n_z \rangle + \\ &+ \frac{\theta}{\hbar} \langle n_\rho, \mu, n_z | -\frac{\omega_c}{4} p_\rho^2 - \frac{1}{2}m\tilde{\omega}^2 L_z | n_\rho, \mu, n_z \rangle. \quad (44) \end{aligned}$$

With the help of [86, 87] the lengthy integrals can be solved in closed form, and we get for the first-order corrections in η

$$\Delta E_{\eta}^{(1)} = -\frac{\eta|\mu|}{2m} - \frac{\eta\omega_c}{4m\tilde{\omega}} (2n_{\rho} + |\mu| + 1) \quad (45)$$

and θ

$$\Delta E_{\theta}^{(1)} = -\frac{1}{2}\theta m\tilde{\omega} \left(\tilde{\omega} - \frac{1}{2}\omega_c f(n_{\rho}, |\mu|) \right) \quad (46)$$

with

$$\begin{aligned} f(n_{\rho}, \mu) = & 2 \binom{n_{\rho} + \mu}{\mu} - 4\mu \binom{\mu + n_{\rho} + 2}{n_{\rho} - 1} - \\ & - \mu(1 + \mu) \left[2 \binom{\mu + n_{\rho}}{n_{\rho}} + 4 \binom{\mu + n_{\rho} - 2}{n_{\rho}} + \right. \\ & \left. + \binom{\mu + n_{\rho} + 1}{n_{\rho}} - \binom{\mu + n_{\rho} + 2}{n_{\rho} - 1} \right]. \quad (47) \end{aligned}$$

A short dimensional analysis shows that η has the dimension of $mass^2 \frac{Length^2}{Time^2}$, which corresponds to the momentum squared, and θ has the dimension of $Length^2$. So, the calculated corrections have the correct dimension of energy.

Ergo, we can summarize the results of our calculation in first-order perturbation theory. Recalling the noncommutative Hamiltonian (31), we see that the unperturbed energy is

$$\begin{aligned} E_{n_{\rho}, \mu, n_z}^{(0)} &= \langle n_{\rho}, |\mu|, n_z | \alpha^2 H_0(x, p) | n_{\rho}, |\mu|, n_z \rangle = \\ &= \alpha^2 \left[\hbar\tilde{\omega}(2n_{\rho} + |\mu| + 1) + \frac{1}{2}\hbar\omega_c\mu + \hbar\omega \left(n_z + \frac{1}{2} \right) \right]. \quad (48) \end{aligned}$$

The first-order energy corrections are

$$\begin{aligned} \Delta E_{n_{\rho}, \mu, n_z}^{(1)} &= -\frac{\eta|\mu|}{2m} - \frac{\eta\omega_c}{4m\tilde{\omega}} (2n_{\rho} + |\mu| + 1) - \\ &\quad - \frac{1}{2}\theta m\tilde{\omega} \left(\tilde{\omega} - \frac{1}{2}\omega_c f(n_{\rho}, |\mu|) \right) \quad (49) \end{aligned}$$

with $f(\rho, |\mu|)$ given in (47). These results hold for the situations, where $\eta \ll \hbar m\omega_c$ and $\theta \ll \frac{\hbar}{m\tilde{\omega}}$.

V. DISCUSSION

Recalling one of the motivations for the development of noncommutative Quantum Mechanics was that the kinetic momentum operators do not commute. The cyclotron frequency ω_c is directly proportional to the magnitude of the magnetic field. Therefore, let us examine the effect of the magnetic field on the energy corrections

in the noncommutative phase-space. To see the effect on the corrections clearly, we will consider the energy correction $\Delta E^{(1)}$ normalized by $E^{(0)}$. As the energy corrections are all negative, and there is no change in sign, we will use $|\Delta E^{(1)}/E^{(0)}|$ for plotting the results.

We will employ the atomic unit system and set, therefore \hbar and $m = 1$. We select arbitrarily $\omega = 1$ and vary ω_c between 0.1 and 10. Based on the condition for the validity of the approximation, the values for θ and η have to satisfy

$$\eta \ll \hbar m\omega_c = \omega_c < 0.1 \text{ and} \quad (50)$$

$$\theta \ll \frac{\hbar}{m\tilde{\omega}} = \frac{1}{\sqrt{1 + \frac{\omega_c^2}{4}}} < \frac{1}{\sqrt{1 + \frac{10^2}{4}}} = 0.19. \quad (51)$$

By setting the values $\eta = 0.01$ and $\theta = 0.01$, η and θ satisfy the conditions (50) and (51), respectively.

Very small values for θ need energies close to the Planck energy E_p , that are only available in black holes. Consequently, in this energy scale the values for η would blowing up, and the perturbative approach would not be reasonable anymore. Therefore, as already pointed out, we select the values for θ and η in the low energy limit. In this limit, we may have the chance to observe the effect of the corrections to the energy levels of the anharmonic oscillator due to the changing magnetic field. Therefore, the experiment has to be carried out in an environment where the change of the space-time is still observable. This indicates, that in an experimental setup, where the 3D harmonic oscillator is put in a strong magnetic field, could be method to measure the noncommutativity parameters θ and η .

From (49) it is clear that the function $f(n_{\rho}, |\mu|)$ plays an important role in the corrections. The possible values for $n_{\rho} = 1, 2, 3$ are given in table I.

n_{ρ}	$ \mu $	$f(n_{\rho}, \mu)$
1	0	2
2	0	2
2	1	-28
3	0	2
3	1	-58
3	2	-286

TABLE I. Values for $f(n_{\rho}, |\mu|)$ for $n_{\rho} = 1, 2, 3$ and $|\mu| = 0, \dots, n_{\rho} - 1$

The energy corrections are all negative. In order to show the relation of $E_{n_{\rho}, \mu, n_z}^{(0)}$ and $|\Delta E_{n_{\rho}, \mu, n_z}^{(1)}/E_{n_{\rho}, \mu, n_z}^{(0)}|$ with the magnetic field, we will plot the unperturbed eigenenergy of the 3D isotropic harmonic oscillator in a uniform magnetic field as a function of $\hbar\omega_c$. Exemplarily we select $n_z = 1$ and $\omega = 1$ and $n_{\rho} = 1, 2, 3$ and $|\mu| = 0..n_{\rho} - 1$ for the graphs of these relationships. Any other selection will not change the qualitative behavior of the system.




FIG. 1. For the values, $n_z = 1, n_\rho = 1, \omega = 1$, figure (a) depicts the unperturbed 3D isotropic harmonic oscillator in a uniform magnetic field's eigenenergy as a function of the $\hbar\omega_c$, whereas figure (b) depicts the first-order perturbation normalized by the unperturbed eigenenergy as a function of $\hbar\omega_c$.

The unperturbed eigenenergy $E_{1,0,1}^{(0)}$ from (48) for large ω_c varies asymptotically linearly with ω_c . Whereas the energy correction $\Delta E_{1,0,1}^{(1)}$ varies asymptotically as ω_c^2 . So $|\Delta E_{1,0,1}^{(1)}/E_{1,0,1}^{(0)}|$ will asymptotically vary $\sim \omega_c$, as depicted in figure 1. So, we can conclude that the magnitude of the corrections depends stronger on the magnitude of the magnetic field than the unperturbed energy of the isotropic 3D harmonic oscillator in a uniform magnetic field $E_{n_\rho,\mu,n_z}^{(0)}$. On the other hand, figure 1 shows that for small magnetic fields, the energy corrections decrease until it reaches its local minimum at $\omega_c = 1.79$ before the magnitude of the relative energy corrections starts to increase again towards its asymptotic behavior.




FIG. 2. For the values, $n_z = 1, n_\rho = 2, \omega = 1$, figure (a) depicts the unperturbed 3D isotropic harmonic oscillators in a uniform magnetic field's eigenenergy as a function of the $\hbar\omega_c$ for $\mu = 0$ (blue line) and $\mu = 1$ (orange line), whereas figure (b) depicts the first-order perturbation normalized by the unperturbed energy as a function of $\hbar\omega_c$ for $\mu = 0$ (blue line) and $\mu = 1$ (orange line).

In the case $n_\rho = 2$, the behavior of the eigenenergies of the unperturbed isotropic 3D harmonic oscillator in a uniform magnetic field $E_{2,0,1}^{(0)}$ and $E_{2,1,1}^{(0)}$ and the magnitude of relative energy corrections due to noncommutativity $|\Delta E_{2,0,1}^{(1)}/E_{2,0,1}^{(0)}|$ and $|\Delta E_{2,1,1}^{(1)}/E_{2,1,1}^{(0)}|$ is qualitatively the same as in the case $n_\rho = 1$. The magnitude

of the relative energy correction $|\Delta E_{2,0,1}^{(1)}/E_{2,0,1}^{(0)}|$ first decreases until $\omega_c = 2.58$, where it reaches its absolute minimum before it starts increasing again towards its asymptotic behavior. We can observe the same behavior for $|\Delta E_{2,1,1}^{(1)}/E_{2,1,1}^{(0)}|$ and get a minimum at $\omega_c = 2.71$ for $\mu = 1$. Furthermore, we can identify that for increasing magnetic Quantum number μ , the magnitude of the relative corrections increases.




FIG. 3. For the values, $n_z = 1, n_\rho = 3, \omega = 1$, figure (a) depicts the unperturbed 3D isotropic harmonic oscillators in a uniform magnetic field's eigenenergy as function of the $\hbar\omega_c$ for $\mu = 0$ (blue line), $\mu = 1$ (orange line), and $\mu = 2$ (yellow line), whereas figure (b) depicts the first-order perturbation normalized by the unperturbed energy as a function of $\hbar\omega_c$ for $\mu = 0$ (blue line), $\mu = 1$ (orange line), and $\mu = 2$ (yellow line).

In the case $n_\rho = 3$, the behavior of the energy of the unperturbed isotropic 3D harmonic oscillator in a uniform magnetic field and the magnitude of the relative energy corrections due to noncommutativity is qualitatively the same as in the cases $n_\rho = 1$ and $n_\rho = 2$. As we can see from figure 3, we get increasing corrections $|\Delta E_{3,\mu,1}^{(1)}/E_{3,\mu,1}^{(0)}|$ with increasing magnetic Quantum number μ . The magnitude of the relative energy corrections reach a minimum at $\omega_c = 3.11$ for $\mu = 0$, $\omega_c = 2.81$ for $\mu = 1$, and $\omega_c = 3.20$ for $\mu = 2$ before they start to increase again towards their asymptotic behavior.

Furthermore, we can identify that the increasing magnetic field's impact is increasing for increasing n_ρ and magnetic Quantum number μ . Moreover, the value for ω_c where $|\Delta E_{n_\rho,\mu,1}^{(1)}/E_{n_\rho,\mu,1}^{(0)}|$ becomes minimal increases for the constant μ and increasing n_ρ .

Overall, evidently, the eigenenergies and their first-order corrections strongly depend on the magnitude of the magnetic field. The relative change of the corrections to the magnetic field shows that the corrections increase faster than the eigenenergies with increasing magnetic field for increasing magnetic Quantum numbers.

VI. CONCLUSION

We studied the charged harmonic oscillator in a uniform magnetic field in the extended framework of non-commutative Quantum Mechanics in 3D. In line with

this, without touching the basic definition of the starting point of noncommutative Quantum Mechanics, namely the commutators of $[\hat{x}_i, \hat{x}_j] = i\theta_{ij}$ and $[\hat{p}_i, \hat{p}_j] = i\eta_{ij}$ we extended the antisymmetric tensor λ_{ij} to (11). This extension of the noncommutativity from the noncommutative plane to the noncommutative space gives rise to a change in the algebra of the system. The first result of this is a never-vanishing commutator $[\hat{x}_i, \hat{p}_j]$ for any combination of i and j . Based on this algebra, we investigated the effect of the noncommutativity in 3D to the eigenenergies of the commutative system. The Hamiltonian for the charged isotropic harmonic oscillator in a uniform magnetic field proves to be non-trivial in the noncommutative phase-space (3D). A closed solution could not be obtained in this algebra. Therefore, in the limit of weak noncommutativity, i.e., in the low energy limit, we could obtain the corrections to the eigenenergies in first-order time-independent perturbation theory in closed form. It turns out that the corrections to

the eigenenergies are negative, i.e. the eigenenergies in the noncommutative system are smaller compared to the commutative ones. To analyze the effect of the magnitude of the magnetic field on the energy corrections, we plotted the graphs of the magnitude of the relative energy corrections $|\Delta E_{n_\rho, \mu, n_z}^{(1)} / E_{n_\rho, \mu, n_z}^{(0)}|$ as a function of $\hbar\omega_c$. The analysis showed that the magnitude of the energy corrections $|\Delta E_{n_\rho, \mu, n_z}^{(1)}|$ increases asymptotically for large $\hbar\omega_c$ with $\hbar\omega_c^2$, whereas the unperturbed eigenenergies $E_{n_\rho, \mu, n_z}^{(0)}$ increase with $\hbar\omega_c$ linearly. Ergo, the corrections to the eigenenergies increase faster with respect to $\hbar\omega_c$ than the eigenenergies themselves. This behavior could be also identified in the graphs of the relative corrections of the eigenenergies for the exemplarily selected parameters. This result suggests, that noncommutative Quantum Mechanics can be experimentally studied even in the low energy limit by employing a strong magnetic field to a 3D harmonic oscillator.

-
- [1] W. Heisenberg, Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik, *Z. Phys.* **43**, 172 (1927).
 - [2] S. Dey, A. Bhat, D. Momeni, M. Faizal, A. F. Ali, T. K. Dey, and A. Rehman, Probing noncommutative theories with quantum optical experiments, *Nuclear Physics B* **924**, 578 (2017).
 - [3] H. S. Snyder, Quantized space-time, *Physical Review* **71**, 38 (1947).
 - [4] R. J. Szabo, Quantum field theory on noncommutative spaces, *Physics Reports* **378**, 207 (2003).
 - [5] N. Seiberg and E. Witten, String theory and noncommutative geometry, *Journal of High Energy Physics* **1999**, 032 (1999).
 - [6] S. Chaturvedi, R. Jagannathan, R. Sridhar, and V. Srinivasan, Non-relativistic quantum mechanics in a noncommutative space, *Journal of Physics A: Mathematical and General* **26**, L105 (1993).
 - [7] J. Gamboa, M. Loewe, and J. C. Rojas, Noncommutative quantum mechanics, *Physical Review D* **64** (2001).
 - [8] A. Connes, M. R. Douglas, and A. Schwarz, Noncommutative geometry and matrix theory, *Journal of High Energy Physics* **1998**, 003 (1998).
 - [9] D. Bigatti and L. Susskind, Magnetic fields, branes, and noncommutative geometry, *Phys. Rev. D* **62**, 066004 (2000).
 - [10] W. Jian-Hua, L. Kang, and D. Sayipjamal, Klein-gordon oscillators in noncommutative phase space, *Chinese physics C* **32**, 803 (2008).
 - [11] E. S. Santos and G. R. de Melo, The schroedinger and pauli-dirac oscillators in noncommutative phase space, *International Journal of Theoretical Physics* **50**, 332 (2011).
 - [12] S. Biswas, Bohr-van leeuwen theorem in noncommutative space, *Physics Letters A* **381**, 3723 (2017).
 - [13] A. Kempf, G. Mangano, and R. B. Mann, Hilbert space representation of the minimal length uncertainty relation, *Phys. Rev. D* **52**, 1108 (1995).
 - [14] S. Das, M. P. Robbins, and M. A. Walton, Generalized uncertainty principle corrections to the simple harmonic oscillator in phase space, *Canadian Journal of Physics* **94**, 139 (2016), <https://doi.org/10.1139/cjp-2015-0456>.
 - [15] P. Bosso, S. Das, and R. B. Mann, Planck scale corrections to the harmonic oscillator, coherent, and squeezed states, *Phys. Rev. D* **96**, 066008 (2017).
 - [16] S. Dey, A. Fring, and L. Gouba, -symmetric noncommutative spaces with minimal volume uncertainty relations, *Journal of Physics A: Mathematical and Theoretical* **45**, 385302 (2012).
 - [17] D. J. Gross and P. F. Mende, String theory beyond the planck scale, *Nuclear Physics B* **303**, 407 (1988).
 - [18] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri, and Y. Oz, Large n field theories, string theory and gravity, *Physics Reports* **323**, 183 (2000).
 - [19] J. Magueijo and L. Smolin, String theories with deformed energy-momentum relations, and a possible nontachyonic bosonic string, *Physical Review D* **71**, 026010 (2005).
 - [20] C. Rovelli, Simple model for quantum general relativity from loop quantum gravity, in *Journal of Physics: Conference Series*, Vol. 314 (IOP Publishing, 2011) p. 012006.
 - [21] C. Rovelli, A new look at loop quantum gravity, *Classical and Quantum Gravity* **28**, 114005 (2011).
 - [22] L. Smolin, An invitation to loop quantum gravity, in *Quantum Theory and Symmetries* (World Scientific, 2004) pp. 655–682.
 - [23] S. Doplicher, K. Fredenhagen, and J. E. Roberts, The quantum structure of spacetime at the planck scale and quantum fields, *Communications in Mathematical Physics* **172**, 187 (1995).
 - [24] M. Bronstein, Republication of: Quantum theory of weak gravitational fields, *General Relativity and Gravitation* **44**, 267 (2012).
 - [25] M. A. Kurkov and M. Sakellariadou, Spectral regularisation: induced gravity and the onset of inflation, *Journal of Cosmology and Astroparticle Physics* **2014** (01), 035.
 - [26] L. Kang and D. Sayipjamal, Non-commutative phase space and its space-time symmetry, *Chinese Physics C*

- 34**, 944 (2010).
- [27] J. D. Bekenstein, Universal upper bound on the entropy-to-energy ratio for bounded systems, *Phys. Rev. D* **23**, 287 (1981).
 - [28] Z. Nedelkoski and I. Petreska, Magnetic properties of electrons confined in an anisotropic cylindrical potential, *PHYSICA B-CONDENSED MATTER* **452**, 113 (2014).
 - [29] T. Sandev, I. Petreska, and E. K. Lenzi, Harmonic and anharmonic quantum-mechanical oscillators in noninteger dimensions, *PHYSICS LETTERS A* **378**, 109 (2014).
 - [30] I. Petreska, T. Sandev, Z. Nedelkoski, and L. Pejov, Axially symmetrical molecules in electric and magnetic fields: energy spectrum and selection rules, *Central European Journal of Physics* **11**, 412 (2013).
 - [31] I. Petreska, T. Sandev, G. Ivanovski, and L. Pejov, Splitting of spectra in anharmonic oscillators described by kratzer potential function, *COMMUNICATIONS IN THEORETICAL PHYSICS* **54**, 38 (2010).
 - [32] I. Petreska, G. Ivanovski, and L. Pejov, The perturbation theory model of a spherical oscillator in electric field and the vibrational stark effect in polyatomic molecular species, *SPECTROCHIMICA ACTA PART A-MOLECULAR AND BIOMOLECULAR SPECTROSCOPY* **66**, 985 (2007).
 - [33] P. Stano, C.-H. Hsu, L. C. Camenzind, L. Yu, D. Zumbuehl, and D. Loss, Orbital effects of a strong in-plane magnetic field on a gate-defined quantum dot, *PHYSICAL REVIEW B* **99**, 10.1103/PhysRevB.99.085308 (2019).
 - [34] W. Xie, Third-order nonlinear optical susceptibility of a donor in elliptical quantum dots, *SUPERLATTICES AND MICROSTRUCTURES* **53**, 49 (2013).
 - [35] F. Amiri, H. Shirkani, and M. M. Golshan, Time-evolution of electronic states in a rashba anisotropic two-dimensional quantum dot, *SUPERLATTICES AND MICROSTRUCTURES* **50**, 419 (2011).
 - [36] E. S. Kadantsev and P. Hawrylak, Effective Theory of Electron-Hole Exchange in Semiconductor Quantum Dots, in *INTERNATIONAL CONFERENCE ON THEORETICAL PHYSICS DUBNA-NANO 2010*, Journal of Physics Conference Series, Vol. 248, edited by Osipov, V and Nesterenko, V and Shukrinov, Y (2010) International Conference on Theoretical Physics Dubna-Nano 2010, Dubna, RUSSIA, JUL 05-10, 2010.
 - [37] E. Kadantsev and P. Hawrylak, Theory of exciton fine structure in semiconductor quantum dots: Quantum dot anisotropy and lateral electric field, *PHYSICAL REVIEW B* **81**, 10.1103/PhysRevB.81.045311 (2010).
 - [38] T. Sako and G. H. F. Diercksen, Spectra and correlated wave functions of two electrons confined in a quasi-one-dimensional nanostructure, *PHYSICAL REVIEW B* **75**, 10.1103/PhysRevB.75.115413 (2007).
 - [39] M. Trif, V. N. Golovach, and D. Loss, Spin-spin coupling in electrostatically coupled quantum dots, *PHYSICAL REVIEW B* **75**, 10.1103/PhysRevB.75.085307 (2007).
 - [40] H.-Y. Fan, T.-T. Wang, and Y.-L. Yang, Energy level of electron in an anisotropic quantum dot under a magnetic field by an invariant eigenoperator method, *INTERNATIONAL JOURNAL OF MODERN PHYSICS B* **20**, 5417 (2006).
 - [41] T. Sako, P.-A. Hervieux, and G. H. F. Diercksen, Distribution of oscillator strength in Gaussian quantum dots: An energy flow from center-of-mass mode to internal modes, *PHYSICAL REVIEW B* **74**, 10.1103/PhysRevB.74.045329 (2006).
 - [42] T. Sako and G. Diercksen, Confined quantum systems: spectra of weakly bound electrons in a strongly anisotropic oblate harmonic oscillator potential, *JOURNAL OF PHYSICS-CONDENSED MATTER* **17**, 5159 (2005).
 - [43] T. Sako and G. Diercksen, Confined quantum systems: spectral properties of two-electron quantum dots, *JOURNAL OF PHYSICS-CONDENSED MATTER* **15**, 5487 (2003).
 - [44] T. Honda and T. Sako, Distribution of oscillator strengths and correlated electron dynamics in artificial atoms, *JOURNAL OF PHYSICS B-ATOMIC MOLECULAR AND OPTICAL PHYSICS* **53**, 10.1088/1361-6455/ab9c35 (2020).
 - [45] Y. Zhao, P.-F. Loos, and P. M. W. Gill, Correlation energy of anisotropic quantum dots, *PHYSICAL REVIEW A* **84**, 10.1103/PhysRevA.84.032513 (2011).
 - [46] T. Sako, J. Paldus, and G. H. F. Diercksen, Origin of Hund's multiplicity rule in quasi-two-dimensional two-electron quantum dots, *PHYSICAL REVIEW A* **81**, 10.1103/PhysRevA.81.022501 (2010).
 - [47] F. Prudente, L. Costa, and J. Vianna, A study of two-electron quantum dot spectrum using discrete variable representation method, *JOURNAL OF CHEMICAL PHYSICS* **123**, 10.1063/1.2131068 (2005).
 - [48] W. Zhu and S. Trickey, Analytical solutions for two electrons in an oscillator potential and a magnetic field, *PHYSICAL REVIEW A* **72**, 10.1103/PhysRevA.72.022501 (2005).
 - [49] T. Sako, S. Yamamoto, and G. Diercksen, Confined quantum systems: dipole transition moment of two- and three-electron quantum dots, and of helium and lithium atoms in a harmonic oscillator potential, *JOURNAL OF PHYSICS B-ATOMIC MOLECULAR AND OPTICAL PHYSICS* **37**, 1673 (2004).
 - [50] T. Sako, I. Cernusak, and G. Diercksen, Confined quantum systems: structure of the electronic ground state and of the three lowest excited electronic (1) $\sigma^+(+)(g)$ states of the lithium molecule, *JOURNAL OF PHYSICS B-ATOMIC MOLECULAR AND OPTICAL PHYSICS* **37**, 1091 (2004).
 - [51] T. Sako and G. Diercksen, Confined quantum systems: dipole polarizability of the two-electron quantum dot, the hydrogen negative ion and the helium atom, *JOURNAL OF PHYSICS B-ATOMIC MOLECULAR AND OPTICAL PHYSICS* **36**, 3743 (2003).
 - [52] T. Sako and G. Diercksen, Confined quantum systems: a comparison of the spectral properties of the two-electron quantum dot, the negative hydrogen ion and the helium atom, *JOURNAL OF PHYSICS B-ATOMIC MOLECULAR AND OPTICAL PHYSICS* **36**, 1681 (2003).
 - [53] T. Sako and G. Diercksen, Confined quantum systems: spectral properties of the atoms helium and lithium in a power series potential, *JOURNAL OF PHYSICS B-ATOMIC MOLECULAR AND OPTICAL PHYSICS* **36**, 1433 (2003).
 - [54] W. Gao-Feng, L. Chao-Yun, L. Zheng-Wen, and Q. Shui-Jie, Exact solution to two-dimensional isotropic charged harmonic oscillator in uniform magnetic field in non-commutative phase space, *Chinese Physics C* **32**, 247 (2008).
 - [55] A. Muhuri, D. Sinha, and S. Ghosh, Entanglement induced by noncommutativity: anisotropic harmonic os-

- illator in noncommutative space, EUROPEAN PHYSICAL JOURNAL PLUS **136**, 10.1140/epjp/s13360-020-00972-x (2021).
- [56] P. Ghosh and D. Nath, Information theoretic measures of uncertainty of a noncommutative anisotropic oscillator in a homogeneous magnetic field, PHYSICA A-STATISTICAL MECHANICS AND ITS APPLICATIONS **538**, 10.1016/j.physa.2019.122791 (2020).
 - [57] D. Nath and P. Roy, Noncommutative anisotropic oscillator in a homogeneous magnetic field, Annals of Physics **377**, 115 (2017).
 - [58] T. Harko and S.-D. Liang, Energy-dependent noncommutative quantum mechanics, The European Physical Journal C **79**, 300 (2019).
 - [59] M.-L. Liang and Y. Jiang, Time-dependent harmonic oscillator in a magnetic field and an electric field on the non-commutative plane, Physics Letters A **375**, 1 (2010).
 - [60] J. Mamat, S. Dulat, and H. Mamatabdulla, Landau-like atomic problem on a non-commutative phase space, International Journal of Theoretical Physics **55**, 2913 (2016).
 - [61] P. D. Alvarez, J. L. Cortes, P. A. Horvathy, and M. S. Plyushchay, Super-extended noncommutative landau problem and conformal symmetry, Journal of High Energy Physics (2009).
 - [62] L. R. Ribeiro, E. Passos, C. Furtado, and J. R. Nascimento, Landau analog levels for dipoles in non-commutative space and phase space - landau analog levels for dipoles, European Physical Journal C **56**, 597 (2008).
 - [63] S. Dulat and K. Li, Landau problem in noncommutative quantum mechanics, Chinese Physics C **32**, 92 (2008).
 - [64] P. R. Giri and P. Roy, The non-commutative oscillator, symmetry and the landau problem, European Physical Journal C **57**, 835 (2008).
 - [65] M. Riccardi, Physical observables for noncommutative landau levels, Journal of Physics a-Mathematical and General **39**, 4257 (2006).
 - [66] M. Hatsuda, S. Iso, and H. Umetsu, Noncommutative superspace, supermatrix and lowest landau level, Nuclear Physics B **671**, 217 (2003).
 - [67] P. A. Horvathy, The non-commutative landau problem, Annals of Physics **299**, 128 (2002).
 - [68] O. F. Dayi and L. T. Kelleyane, Wigner functions for the landau problem in noncommutative spaces, Modern Physics Letters A **17**, 1937 (2002).
 - [69] J. Gamboa, F. Mendez, M. Loewe, and J. C. Rojas, The landau problem and noncommutative quantum mechanics, Modern Physics Letters A **16**, 2075 (2001).
 - [70] A. Comtet, On the landau-levels on the hyperbolic plane, Annals of Physics **173**, 185 (1987).
 - [71] R. Iengo and R. Ramachandran, Landau levels in the noncommutative ads2, Journal of High Energy Physics **2002**, 017 (2002).
 - [72] S. Gangopadhyay, A. Saha, and A. Halder, On the landau system in noncommutative phase-space, Physics Letters A **379**, 2956 (2015).
 - [73] B. Harms and O. Micu, Noncommutative quantum hall effect and aharonov-bohm effect, Journal of Physics a-Mathematical and Theoretical **40**, 10337 (2007).
 - [74] F. G. Scholtz, B. Chakraborty, S. Gangopadhyay, and J. Govaerts, Interactions and non-commutativity in quantum hall systems, Journal of Physics a-Mathematical and General **38**, 9849 (2005).
 - [75] B. Basu and S. Ghosh, Quantum hall effect in bilayer systems and the noncommutative plane: A toy model approach, Physics Letters A **346**, 133 (2005).
 - [76] O. F. Dayi and A. Jellal, Hall effect in noncommutative coordinates, Journal of Mathematical Physics **43**, 4592 (2002).
 - [77] J. Jing and J.-F. Chen, Non-commutative harmonic oscillator in magnetic field and continuous limit, The European Physical Journal C **60**, 669 (2009).
 - [78] B. Chakraborty, Z. Kuznetsova, and F. Toppan, Twist deformation of rotationally invariant quantum mechanics, Journal of mathematical physics **51**, 112102 (2010).
 - [79] Z. Kuznetsova and F. Toppan, Effects of twisted noncommutativity in multi-particle hamiltonians, The European Physical Journal C **73**, 2483 (2013).
 - [80] R. Banerjee, A novel approach to noncommutativity in planar quantum mechanics, Modern Physics Letters A **17**, 631 (2002).
 - [81] H. Hassanabadi, S. Hosseini, and S. Zarrinkamar, Dirac oscillator in noncommutative space, Chinese Physics C **38**, 063104 (2014).
 - [82] S. Hess, *Tensors for physics* (Springer, 2015).
 - [83] L. Mezincescu, Star operation in quantum mechanics (2000), arXiv:hep-th/0007046 [hep-th].
 - [84] T. Curtright, D. Fairlie, and C. Zachos, Features of time-independent wigner functions, Physical Review D **58**, 025002 (1998).
 - [85] M. S. Abramowitz and I. Stegun, Ia.(1964), handbook of mathematical functions, Washington: National Bureau of Standards , 923 (1999).
 - [86] H. Srivastava, H. Mavromatis, and R. Alassar, Remarks on some associated laguerre integral results, Applied Mathematics Letters **16**, 1131 (2003).
 - [87] H. A. Mavromatis, An interesting new result involving associated laguerre polynomials, International journal of computer mathematics **36**, 257 (1990).