

On the study of antiprotons yield in hadronic collisions at NICA SPD

Об изучении выхода антипротонов в адронных столкновениях на NICA SPD

V. Alexakhin^{a,1}, A. Guskov^{a,2}, Z. Hayman^{b,3}, R. El-Kholy^{b,4}, A. Tkachenko^{a,5}

^a Joint Institute for Nuclear Research (JINR), Dubna, Russia, 141980

^b Astronomy Department, Faculty of Science, Cairo University, Giza, Egypt, 12613

The study of antiproton yield in p - p and p - d collisions is important for the astrophysical search for dark matter consisting of Weakly Interacting Massive Particles. Refinement of the production cross section, angular and momentum spectra of produced antiprotons in a wide energy range could help to treat the results obtained by the AMS-02 and PAMELA orbit spectrometers. In this paper we present a detailed Monte Carlo study of possible measurements at the planned SPD experiment at the NICA collider that is under construction at JINR (Dubna).

Изучение выхода антипротонов в p - p и p - d столкновениях является важным для астрофизических поисков тёмной материи, состоящей из слабо взаимодействующих массивных частиц (WIMP). Уточнение сечений рождения антипротонов, их угловых и импульсных спектров в широком диапазоне энергий могло бы помочь в интерпретации результатов, полученных на орбитальных спектрометрах AMS-02 и PAMELA. В данной работе мы представляем детальное Монте Карло моделирование возможных измерений в планируемом эксперименте SPD на коллайдере NICA, который строится в ОИЯИ (Дубна).

PACS: 13.85.Ni; 13.85.Tp; 14.80.-j

INTRODUCTION

Dark Matter (DM) is thought to compose more than 24% of the matter-energy content of the Universe [1]. While observations have not yet revealed the origin or nature of DM, there are many theories trying to infer both from available information. One of the most common theories is that the galactic dark halos are built up of thermal relics of the Big Bang called Weakly Interacting Massive Particles (WIMPs) [2]. There is a possibility

¹E-mail: vadim.alexakhin@cern.ch

²E-mail: alexey.guskov@cern.ch

³E-mail: zmhayman@cu.edu.eg

⁴E-mail: relkholy@sci.cu.edu.eg

⁵E-mail: avt@jinr.ru

that, even though DM particles do not interact with Standard Model (SM) particles, they do produce SM particles via pair-annihilation or decay. Powerful experiments, such as AMS-02 and PAMELA, have been trying to detect anomalies in rare components of Cosmic Rays (CRs), especially antiparticles like antiprotons. However, conclusions are hindered by uncertainties.

To be able to detect anomalous components in CRs, it is crucial to measure the production of these components from conventional astrophysical sources with the maximal possible precision. For the secondary antiproton background, the sources are proton-nucleus, nucleus-proton, and nucleus-nucleus collisions of CRs on the Interstellar Medium (ISM). However, almost all secondary antiprotons are a result of collisions of light nuclei. Despite the AMS-02 measuring the antiproton flux with unprecedented accuracy of a few percents [3], no conclusion can be made yet about exotic components due to several sources of uncertainty. The largest source of uncertainties, which can vary from 20 to 50% depending on the energy, is uncertainties on antiproton-production cross sections [4].

In addition to the lack of data of antiproton production in collisions involving helium, data of antiproton production in proton-proton collisions are scarce. Moreover, most of the existing datasets are not corrected for antiproton production via the decay of intermediate hyperons; namely, $\bar{\Lambda}$ and $\bar{\Sigma}^-$; a process that gains significance with increasing energy [5]. In order to keep up with the AMS-02 accuracy and energy range, new accurate measurements of antiproton production are essential. This would enable comparison of data with theoretical models; and, eventually, coming to a conclusion about any exotic signals. These measurements need to cover the proton-beam kinetic-energy range from 1 GeV to 6 TeV, and the pseudorapidity range from 2 to 8 [6].

It is currently planned to measure the antiproton-production cross-sections of p - p and p - He collisions in fixed-target mode by the COMPASS Collaboration [7]. However, the 4π -universality of the Spin Physics Detector (SPD) detector [8] at the Nuclotron-based Ion Collider fAcility (NICA) [9] will enable measurements of antiprotons with high transverse momentum (p_T) with lower beam energy. There is also the possibility of performing proton-deuteron collision measurements, which contribute to the secondary antiproton flux in CRs as well.

NICA is currently under construction at JINR. It is planned to operate the NICA collider with polarized proton and deuteron beams for spin physics study at the SPD detector. In a previous work [10], we performed a preliminary study of antiproton production at the SPD. Monte Carlo simulations generated with Pythia8 have shown that the momenta of the antiprotons produced in p - p collisions, at a center-of-mass (CM) energy $\sqrt{s} = 26$ GeV, would be low enough to identify with the time-of-flight method. They have also shown that with a time resolution of about 100 ps and a flight base of 2 meters, K^-/\bar{p} separation can be achieved for antiprotons with momenta up to 4 GeV, which would include most of the antiproton yield.

In this paper, we study with more detail the antiproton production in hadronic collisions with Monte Carlo simulations specific to the SPD setup using the SPDROOT toolkit.

1. THE SPIN PHYSICS DETECTOR

The Spin Physics Detector is planned as a multipurpose universal 4π detector. It will be able to operate in the polarized p - p collision mode with CM energy up to 27 GeV and luminosity up to about $10^{32} \text{ cm}^{-2} \text{ s}^{-1}$. Another mode would be the polarized d - d collisions at energy up to 13.5 GeV and a luminosity one order lower. Asymmetric p - d collisions are also discussed [8]. The detector consists of the barrel part and two end-caps. The tracking system of the SPD includes the silicon-based vertex detector surrounded by the main tracker, using gas-filled drift straw-tubes as the basic detection element with a spatial resolution of about $100 \mu\text{m}$. The magnetic field (a few configurations are currently under consideration) provides charged-particle momentum measurement with a typical resolution of about 2% at a transverse momentum of 1 GeV/c. The time-of-flight (TOF) system, with a time resolution below 100 ps, provides identification of secondary hadrons in a wide kinematic range. The shashlyk-type electromagnetic calorimeter is responsible for the photon reconstruction and identification of electrons and positrons. The muon system (RS) performs the advanced muon/hadron separation via comprehensive pattern recognition and matching of the track segments to the tracks in the inner part of the detector [8].

This Monte Carlo study simulation was performed in order to estimate the capability of the Spin Physics Detector for the precision measurement of antiproton yield in hadronic collisions at NICA. The study was done using the SPDROOT toolkit, a ROOT- and GEANT4-based software developed for SPD using the FAIRROOT framework [11]. A simplified description of SPD with a quasi-solenoidal magnetic field along the beam axis ($B = 0.4 \text{ T}$) was used. The time resolution of the TOF system used for antiproton identification was supposed to be 70 ps. PYTHIA8 [12] was used as a generator of primary p - p interactions. Since the geometry of the detector is not yet fixed all the results presented in the paper should be treated as indicative.

2. DIRECT PRODUCTION OF ANTIPROTONS IN p - p COLLISIONS

The antiproton-production cross section in p - p collisions multiplied by the average antiproton-multiplicity in the discussed kinematic range is on the level of a few millibarns [10]. That corresponds to an antiproton production rate on a level $> 10^5 \text{ s}^{-1}$. So, the accuracy of the proposed measurements is not limited by the statistical uncertainty. The phase space in terms of the variable x_R and the transverse momentum p_T of produced antiprotons covered by SPD at $\sqrt{s} = 13 \text{ GeV}$ and $\sqrt{s} = 26 \text{ GeV}$ is shown in Fig.1. Here x_R is defined as the ratio of antiproton energy to its maximal possible energy for a particular \sqrt{s} . The geometrical acceptance of the tracking system that affects the track reconstruction efficiency and energy losses of the produced antiprotons, and the magnetic field preventing low-energy particles from reaching the TOF system are taken into account. An antiproton produced at the interaction point was considered detected if it reached the TOF system in the barrel or an end-cap. The radius of the barrel part of the TOF system and the distance from the center of the detector to the end-cap parts are assumed to be 2.0 m each. The detection efficiency as a function of \bar{p} momentum for $\sqrt{s} = 26 \text{ GeV}$ is presented in Fig.2a.




Fig. 1: The kinematic range accessible by p - p collisions at the SPD detector, where the red points represent a CM energy of 13 GeV and the magenta points represent a CM energy of 26 GeV, superimposed on the required measurement range to match the precision level of the AMS-02 measurements [6].




Fig. 2: (a) Detection efficiency as a function of \bar{p} momentum for p - p collisions at $\sqrt{s} = 26$ GeV. (b) t_0 reconstruction accuracy as a function of the number of charged tracks in the event.

Antiproton identification is the most critical issue for the proposed measurements. The situation is worsening with the presence of only one detector plane for the time-of-flight measurement and with the relatively-low multiplicity of secondary charged particles. The algorithm described in Ref. [13] was applied for events with track multiplicities above 3. For each track in the event, three possible hypotheses were assumed: π , K and p . The χ^2 minimization of the time mismatch at the interaction point was performed over all possible combinations of individual-track hypotheses. As a result of such minimization, for each event the interaction time t_0 and the most probable particle IDs for each track could be found. Fig.2b shows the accuracy of t_0 reconstruction as a function of the number of charged tracks in the event. Particle-mass reconstruction using TOF system is illustrated in Fig.3. Safe K^-/\bar{p} separation with a purity of antiproton sample on the level of 1% could be provided for particles with momenta below 3.5 GeV/c.




Fig. 3: Mass reconstruction for π^- , K^- , and \bar{p} using the TOF system. Simulation is performed for $\sqrt{s} = 26$ GeV.

3. ANTIPROTONS FROM ANTI-HYPERONS DECAY

It has been mentioned in [10] that a significant part of antiprotons is produced from the decays of secondary $\bar{\Lambda}$ and $\bar{\Sigma}^-$ anti-hyperons. The ratio Δ_Λ of hyperon-induced to directly produced antiprotons can be expressed as

$$\Delta_\Lambda = \frac{\bar{\Lambda}}{\bar{p}} \times \text{BF}_{\bar{\Lambda} \rightarrow \bar{p}\pi^+} + \frac{\bar{\Sigma}^-}{\bar{p}} \times \text{BF}_{\bar{\Sigma}^- \rightarrow \bar{p}\pi^0}, \quad (1)$$

where $\bar{\Lambda}/\bar{p}$ and $\bar{\Sigma}^-/\bar{p}$ are the hyperon to (promptly produced) antiproton ratios, $\text{BF}_{\bar{\Lambda} \rightarrow \bar{p}\pi^+} = 0.639 \pm 0.005$ and $\text{BF}_{\bar{\Sigma}^- \rightarrow \bar{p}\pi^0} = 0.5157 \pm 0.0003$ are the branching fractions of the corresponding decays. Existing experimental data for the ratio $\bar{\Lambda}/\bar{p}$ are presented in Fig. 4. The relative uncertainty of this quantity at SPD energies is about 12%. As for the $\bar{\Sigma}^-/\bar{p}$ ratio, there is not any trustable experimental data on it; and it was estimated roughly based on the symmetry arguments that $\bar{\Sigma}^-/\bar{\Lambda} = 0.33$ [14]. Thus, $\Delta_\Lambda = (0.81 \pm 0.04) \times \bar{\Lambda}/\bar{p}$.




Fig. 4: $\bar{\Lambda}/\bar{p}$ ratio in proton-proton collisions as measured by several experiments [5]. The range of \sqrt{s} accessible at the NICA SPD is shown in red at the lower left part.

Since hyperons have a macroscopic decay length, their contribution can be accessed at the SPD via reconstruction of the two-body decays $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ and $\bar{\Sigma}^- \rightarrow \bar{p}\pi^0$ in secondary vertices. The fraction of $\bar{\Lambda}$ decaying within the inner part of the SPD setup, averaged over the spectrum, is 93% at $\sqrt{s} = 26 \text{ GeV}$ while almost all $\bar{\Sigma}^-$ decay there. The reconstructed invariant mass of $\bar{p}\pi^+$ from $\bar{\Lambda}$ decay in secondary V-vertices is shown in Fig. 5a. The Gaussian width of the peak is about $2 \text{ GeV}/c^2$. It is fully defined by the uncertainty of the momentum measurement in the SPD tracking system.

For determination of the $\bar{\Sigma}^-$ decay, a secondary vertex should be reconstructed using a track pointing to the primary vertex and associated with $\bar{\Sigma}^-$ and a track associated

with an antiproton. A pair of photons detected in the electromagnetic calorimeter should be assumed to be produced from a π^0 decay in the found secondary vertex. The width of the reconstructed $\bar{\Sigma}^-$ peak is mainly defined by the energy resolution of the calorimeter and equals to 19 MeV as it can be seen from Fig. 5b. Both signals, $\bar{\Lambda}$ and $\bar{\Sigma}^-$, are well-distinguished over the background. That inspires a hope to determine the ratios $\bar{\Lambda}/\bar{p}$ and $\bar{\Sigma}^-/\bar{p}$ and, finally, Δ_Λ with a minimal statistical uncertainty.




Fig. 5: (a) Reconstructed $\bar{p}\pi^+$ invariant mass spectrum with a $\bar{\Lambda}$ signal.
(b) Reconstructed $\bar{p}\pi^0$ invariant mass spectrum with a $\bar{\Sigma}^-$ signal.

4. SUMMARY

Based on the proposal to use the Spin Physics Detector at the NICA collider as an instrument for the precision measurements of the antiproton yield in p - p and p - d collisions required by astrophysical searches for dark matter [10], we performed this Monte Carlo study using a preliminary SPD setup geometry. We refined the kinematic coverage of the detector, studied its possibility to identify directly produced antiprotons with the TOF system, and investigated the conditions of reconstructing the $\bar{\Lambda}$ and $\bar{\Sigma}^-$ hyperon decays. It was shown that the SPD TOF system on the base of 2.0 m can provide purity of K^-/\bar{p} separation better than 99% up to $p = 3.5$ GeV/ c . Accuracy of the interaction time t_0 reconstruction could reach 40-50 ps. The SPD tracking system and electromagnetic calorimeter offer the possibility of effectively reconstructing the decays $\bar{\Lambda} \rightarrow \bar{p}\pi^+$ and $\bar{\Sigma}^- \rightarrow \bar{p}\pi^0$, which are the main sources of secondary antiprotons. The setup resolution of the $\bar{\Lambda}$ and $\bar{\Sigma}^-$ peaks is found to be 2 MeV and 19 MeV, respectively.

We reaffirm our previous conclusions that the supplementary measurements at the Spin Physics Detector could make a sizable contribution to the search of physics beyond the Standard Model.

REFERENCES

1. *Hinshaw G. et al.* [WMAP Collaboration] Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results // [The Astrophysical Journal Supplement Series](#). — 2013. — Sep. — V. 208, no. 2. — P. 19.
2. *Tanabashi M. et al.* [PDG Collaboration] Review of Particle Physics // [Physical Review D](#). — 2018. — Aug. — V. 98, no. 3.
3. *Aguilar M. et al.* [AMS Collaboration] Antiproton Flux, Antiproton-to-Proton Flux Ratio, and Properties of Elementary Particle Fluxes in Primary Cosmic Rays Measured with the Alpha Magnetic Spectrometer on the International Space Station // [Physical Review Letters](#). — 2016. — Aug. — V. 117, no. 9.
4. *Giesen G., Boudaud M., Génolini Y., Poulin V., Cirelli M., Salati P., Serpico P.D.* AMS-02 antiprotons, at last! Secondary astrophysical component and immediate implications for Dark Matter // [Journal of Cosmology and Astroparticle Physics](#). — 2015. — Sep. — V. 2015, no. 09. — P. 023.
5. *Winkler M.W.* Cosmic ray antiprotons at high energies // [Journal of Cosmology and Astroparticle Physics](#). — 2017. — Feb. — V. 2017, no. 02. — P. 048.
6. *Donato F., Korsmeier M., Mauro M.D.* Prescriptions on antiproton cross section data for precise theoretical antiproton flux predictions // [Physical Review D](#). — 2017. — Aug. — V. 96, no. 4.
7. *Adams B. et al.* [COMPASS Collaboration] Letter of Intent: A New QCD facility at the M2 beam line of the CERN SPS (COMPASS++/AMBER). — arXiv:[1808.00848v6](#).
8. *Savin I., Efremov A., Peshekhonov D., Kovalenko A., Teryaev O., Shevchenko O., Nagaichev A., Guskov A., Kukhtin V., Topilin N.* Spin Physics Experiments at NICA-SPD with polarized proton and deuteron beams // [EPJ Web of Conferences](#). — 2015. — V. 85. — P. 02039.
9. *Kekelidze V., Kovalenko A., Lednicky R., Matveev V., Meshkov I., Sorin A., Trubnikov G.* Status of the NICA project at JINR // [EPJ Web of Conferences](#). — 2017. — V. 138. — P. 01027.
10. *Guskov A., El-Kholy R.* On the Possibility to Study Antiproton Production at the SPD Detector at NICA Collider for Dark Matter Search in Astrophysical Experiments // [Physics of Particles and Nuclei Letters](#). — 2019. — May. — V. 16, no. 3. — P. 216–223.
11. *Al-Turany M., Bertini D., Karabowicz R., Kresan D., Malzacher P., Stockmanns T., Uhlig F.* The FairRoot framework // [Journal of Physics: Conference Series](#). — 2012. — Dec. — V. 396, no. 2. — P. 022001.
12. *Sjöstrand T., Ask S., Christiansen J.R., Corke R., Desai N., Ilten P., Mrenna S., Prestel S., Rasmussen C.O., Skands P.Z.* An introduction to PYTHIA 8.2 // [Computer Physics Communications](#). — 2015. — Jun. — V. 191. — P. 159–177.

13. *Adam J. et al.* [ALICE Collaboration] Determination of the event collision time with the ALICE detector at the LHC // [The European Physical Journal Plus](#). — 2017. — Feb. — V. 132, no. 2.
14. *Kappl R., Winkler M.W.* The cosmic ray antiproton background for AMS-02 // [Journal of Cosmology and Astroparticle Physics](#). — 2014. — Sep. — V. 2014, no. 09. — P. 051.