arXiv:2008.04358v3 [math.OC] 11 Jun 2021

# On the Characterization of Generalized Derivatives for the Solution Operator of the Bilateral Obstacle Problem

Anne-Therese Rauls [*], Stefan Ulbrich [†]

December 3, 2021

**Abstract**

We consider optimal control problems for a wide class of bilateral obstacle problems where the control appears in a possibly nonlinear source term. The non-differentiability of the solution operator poses the main challenge for the application of efficient optimization methods and the characterization of Bouligand generalized derivatives of the solution operator is essential for their theoretical foundation and numerical realization. In this paper, we derive specific elements of the Bouligand generalized differential if the control operator satisfies natural monotonicity properties. We construct monotone sequences of controls where the solution operator is Gâteaux differentiable and characterize the corresponding limit element of the Bouligand generalized differential as being the solution operator of a Dirichlet problem on a quasi-open domain. In contrast to a similar recent result for the unilateral obstacle problem [RU19], we have to deal with an opposite monotonic behavior of the active and strictly active sets corresponding to the upper and lower obstacle. Moreover, the residual is no longer a nonnegative functional on $H^{-1}$ and its representation as the difference of two nonnegative Radon measures requires special care. This necessitates new proof techniques that yield two elements of the Bouligand generalized differential. Also for the unilateral case we obtain an additional element to that derived in [RU19].

**Key words**— bilateral obstacle problem, variational inequalities, generalized derivatives, Bouligand generalized differential, optimal control, nonsmooth optimization

**AMS subject classifications**— 47J20, 49J40, 49K40, 49J52, 58C20, 58E35

## 1 Introduction

We consider the optimal control of bilateral obstacle problems

$$\text{Find } y \in K_\psi^\varphi: \quad \langle Ly - f(u), z - y\rangle_{H^{-1}(\Omega),H_0^1(\Omega)} \geq 0 \quad \forall\, z \in K_\psi^\varphi, \tag{1}$$

[*]Departments of Mathematics, TU Darmstadt, Dolivostr. 15, 64293 Darmstadt, Germany, anne-therese.rauls@tu-darmstadt.de

[†]Departments of Mathematics, TU Darmstadt, Dolivostr. 15, 64293 Darmstadt, Germany, stefan.ulbrich@tu-darmstadt.de

with

$$K_\psi^\varphi := \{z \in H_0^1(\Omega) \mid \psi \le z \le \varphi \text{ q.e. in } \Omega\}.$$

Here, $\Omega \subseteq \mathbb{R}^d$ is an open, bounded set, $L \in \mathcal{L}(H_0^1(\Omega), H^{-1}(\Omega))$ is a coercive and strictly T-monotone operator. Here, strict T-monotonicity means that

$$\langle L(y-z), (y-z)_+ \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} > 0$$

for all $y, z \in H_0^1(\Omega)$ with $(y-z)_+ = \sup(0, y-z) \neq 0$, see [Rod87, p. 105]. Furthermore, $f\colon U \to H^{-1}(\Omega)$ is a continuously differentiable and monotone operator on a partially ordered Banach space $U$. The detailed assumptions on $f$ and $U$ will be given below in Assumption 1.2. We assume that the obstacles $\psi, \varphi \in H^1(\Omega)$ are such that $K_\psi^\varphi$ is nonempty. In addition, for some results in this paper we require the following assumption.

**Assumption 1.1**
*We consider lower and upper obstacles $\psi, \varphi \in H^1(\Omega) \cap L^\infty(\Omega)$ and assume there is a constant $c_\psi^\varphi > 0$ such that $\varphi - \psi \ge c_\psi^\varphi$ holds a.e. in $\Omega$.*

It is well known, see for example [Bar84,KS00], that for each $u \in U$, the variational inequality (1) has a unique solution and the solution operator $S_f = S_{\psi,f}^\varphi\colon U \to H_0^1(\Omega)$ is locally Lipschitz continuous.

The optimal control of obstacle problems and elliptic variational inequalities has been studied by many authors, see for example [Bar84,Ber97,BL04,Fri88,IK00,HW18, HK11,KKT03,KW12,MRW15,Mig76,MP84,RU19,RW19,SW13]. By using penalization, relaxation or regularization approaches, optimality conditions have been derived in [Bar84,Ber97,BL04,MP84,IK00,HK11], where [BL04] considers the obstacles for the bilateral case as controls. Numerical solution methods based on these techniques have been developed in [IK00,HK11,KKT03,KW12,MRW15,SW13]. Other approaches consider directly the nonsmooth solution operator. Different optimality systems for the optimal control of the obstacle problem are compared in [HW18]. For the application of nonsmooth optimization methods like bundle methods, the knowledge of at least one element in the generalized differential of the objective function is required. For the finite dimensional obstacle problems, a characterization of the whole Clarke subdifferential of the reduced objective function was obtained in [HR86]. The directional differentiability of solution operators of elliptic variational inequalities and a variational inequality for the directional derivative have been obtained in [Har77, Mig76], see also [CW19]. The resulting structure of Gâteaux derivatives in points of differentiability was used in [RW19] to characterize the full Bouligand generalized differential for the solution operator of the unilateral obstacle problem with distributed control $f(u) = u \in H^{-1}(\Omega)$. Subsequently, for Lipschitz continuous, continuously differentiable, and monotone control operators $f\colon U \to H^{-1}(\Omega)$ on a partially ordered Banach space $U$, an element of the Bouligand generalized differential for the solution operator of the unilateral obstacle problem has been characterized in [RU19] as the solution operator of a variational equation on the inactive set. This forms an analytical foundation to develop error estimators for the numerical computation of subgradients and to apply inexact bundle methods in Hilbert space, see for example [HU19].

In this paper, we derive, based on a characterization of the Gâteaux derivative in points of differentiability, two elements of the Bouligand generalized differential for the solution operator $S_f\colon U \to H_0^1(\Omega)$ of the bilateral obstacle problem (1) in points of nonsmoothness. To this end, we require the following monotonicity property of the control operator $f\colon U \to H^{-1}(\Omega)$.

**Assumption 1.2**
*We assume that the operator $f\colon U \to H^{-1}(\Omega)$ is defined on a partially ordered Banach space $(U, \ge_U)$. Let $f$ be increasing, i.e., $u_1 \ge_U u_2$ implies $f(u_1) \ge f(u_2)$ in $H^{-1}(\Omega)$,*

*i.e.,* $\langle f(u_1) - f(u_2), v\rangle_{H^{-1}(\Omega),H_0^1(\Omega)} \geq 0$ *for all* $v \in H_0^1(\Omega)_+ := \{v \in H_0^1(\Omega) \mid v \geq 0\}$. *Moreover, let* $f$ *be continuously differentiable.*

*In addition, we assume that* $U$ *is separable and that there is a partially ordered Banach space* $(V, \geq_V)$ *such that the positive cone* $\mathcal{P} = \{v \in V \mid v \geq_V 0\}$ *has nonempty interior and* $V$ *is embedded into* $U$. *The order relation* $\geq_V$ *has the property that for all* $v_1, v_2 \in V$ *with* $v_1 \geq_V v_2$ *we have* $v_1 + z \geq_V v_2 + z$ *for all* $z \in V$ *and* $t\, v_1 \geq_V t\, v_2$ *for all scalars* $t \geq 0$. *We assume that the linear embedding* $\iota\colon V \to U$ *is continuous, dense and increasing, i.e., compatible with the order structures in* $V$ *and* $U$. *This means that* $v \in V$ *with* $v \geq_V 0$ *implies* $\iota(v) \geq_U 0$ *in* $U$.

With the help of a generalization of Rademacher's theorem to infinite dimensions, see Theorem 6.1, the assumption allows us to construct for each $u \in U$ a monotonically increasing (or monotonically decreasing) sequence $(u_n)_{n\in\mathbb{N}} \subseteq U$ converging to $u$, such that the locally Lipschitz continuous solution operator $S_f$ is Gâteaux differentiable at $u_n$. In particular, the assumptions on $U$ are fulfilled for $U = L^2(\Omega)$, $U = H^{-1}(\Omega)$ and $U = \mathbb{R}^k$.

Our strategy for computing an element of the Bouligand generalized differential is as follows. Let $u \in U$ and denote by $I(f(u))$ and $A(f(u)) := A_\psi(f(u)) \cup A^\varphi(f(u))$ the inactive and active set of the solution $S_f(u)$, respectively. By representing the residual $LS_f(u) - f(u) \in H^{-1}(\Omega)$ as the difference $\tilde{\xi}_\psi - \tilde{\xi}^\varphi$ of two nonnegative Radon measures (Theorem 3.4), we can define the strictly active set $A_{\mathrm{s}}(f(u)) := A_\psi^{\mathrm{s}}(f(u)) \cup A_{\mathrm{s}}^\varphi(f(u))$ using the measures $\tilde{\xi}_\psi$ and $\tilde{\xi}^\varphi$.

If the solution operator $S_f$ is Gâteaux differentiable at $u$, then $S_f'(u)$ can be obtained as the solution operator of a variational equation on $H_0^1(D)$, where $D$ can be chosen as any quasi-open subset of $\Omega$ satisfying $I(f(u)) \subseteq D \subseteq \Omega \setminus A_{\mathrm{s}}(f(u))$, see Theorem 3.9.

If the solution operator $S_f$ is nonsmooth at $u$, then we can construct a monotonically increasing sequence $(u_n)_{n\in\mathbb{N}} \subseteq U$, where $S_f$ is Gâteaux differentiable, converging to $u$ and $S_f'(u_n)$ can thus be represented as the solution operator of a variational equation on $H_0^1(D_n)$ with $D_n := I(f(u_n)) \cup (A^\varphi(f(u_n)) \setminus A_{\mathrm{s}}^\varphi(f(u_n)))$. By using monotonicity properties of the sequence of sets $(A_{\mathrm{s}}^\varphi(f(u_n)))_{n\in\mathbb{N}}$ and $(A_\psi(f(u_n)))_{n\in\mathbb{N}}$ we show that $(H_0^1(D_n))_{n\in\mathbb{N}}$ converges in the sense of Mosco to $H_0^1(D)$ for $D := I(f(u)) \cup (A^\varphi(f(u)) \setminus A_{\mathrm{s}}^\varphi(f(u)))$, cf. Lemma 5.3, and stability properties of variational equations yield that $(S_f'(u_n))_{n\in\mathbb{N}}$ converges in the strong operator topology to an element in the Bouligand generalized differential of $S_f$ at $u$, which can be characterized as the solution operator of a variational equation on $H_0^1(D)$, see Theorem 6.2. Working with a monotonically decreasing sequence $(u_n)_{n\in\mathbb{N}} \subseteq U$ yields another generalized derivative.

In [RU19], a similar approach has been used for the unilateral obstacle problem. The analysis for the bilateral obstacle problem is more involved and requires new proof techniques for several reasons. First of all, to the best of our knowledge, the decomposition of $LS_f(u) - f(u) \in H^{-1}(\Omega)$ as the difference $\tilde{\xi}_\psi - \tilde{\xi}^\varphi$ of two nonnegative Radon measures has not been established so far and requires care. This representation is, in general, only valid on $H_0^1(\Omega) \cap L^\infty(\Omega)$, see Theorem 3.4, or, alternatively, if the distance of the active sets $A^\varphi(f(u))$ and $A_\psi(f(u))$ is positive (Lemma 3.5), which we demonstrate by giving a counter example in Example 3.6. Next, while for $(u_n)_{n\in\mathbb{N}}$ increasing the choice $D_n := I(f(u_n))$ is possible and monotonically increasing in the unilateral case, the sets $D_n := I(f(u_n)) \cup (A^\varphi(f(u_n)) \setminus A_{\mathrm{s}}^\varphi(f(u_n)))$ do not enjoy monotonicity properties and require the more difficult study of monotonicty properties of the strictly active sets $A_{\mathrm{s}}^\varphi(f(u_n))$, see Lemma 4.5. Also in the unilateral case, the analysis in this paper yields an additional element of the Bouligand generalized differential to that derived in [RU19].

If $L$ is induced by a symmetric coercive bilinear form, then it is well known that

(1) are first order optimality conditions of the problem

$$\min_{y\in H_0^1(\Omega)} \langle \tfrac{1}{2}Ly - f(u), y\rangle_{H^{-1}(\Omega),H_0^1(\Omega)} \quad \text{subject to } y \in K_\psi^\varphi.$$

Then the residual $LS_f(u) - f(u) \in H^{-1}(\Omega)$ corresponds to the Lagrange multiplier and our careful study of its representation as the difference $\tilde{\xi}_\psi - \tilde{\xi}^\varphi$ of two nonnegative Radon measures gives detailed insights into the structure of the Lagrange multiplier. This might be helpful also in other contexts, for example the design and analysis of efficient solution methods for (1).

The paper is organized as follows. Section 2 recalls some basic definitions and results on capacity theory, Sobolev spaces on quasi-open sets and generalized derivatives. In Section 3, monotonicity and differentiability properties of the solution operator are analyzed. The well known variational inequality for the directional derivative is stated and the structure of the critical cone is studied. To this end, a representation of the residual $LS_f(u) - f(u)$ as the difference $\tilde{\xi}_\psi - \tilde{\xi}^\varphi$ of two nonnegative Radon measures is derived and the strictly active sets are defined based on these measures. A counterexample shows that this representation is, in general, only valid on $H_0^1(\Omega) \cap L^\infty(\Omega)$, which requires some care in the sequel. Moreover, a representation of $S_f'(u)$ is derived if $u$ is a point of Gâteaux differentiability. In Section 4 the monotonicity of the strictly active and active sets $A_{\mathrm{s}}^\varphi(f(u_n))$ and $A_\psi(f(u_n))$ is analyzed for monotonically increasing control sequences $(u_n)$ (respectively, of $A_{\mathrm{s}}^\psi(f(u_n))$ and $A^\varphi(f(u_n))$ for monotonically decreasing $(u_n)_{n\in\mathbb{N}}$). This is used in Section 5 to show the Mosco convergence of $(H_0^1(I(f(u_n)) \cup (A^\varphi(f(u_n)) \setminus A_{\mathrm{s}}^\varphi(f(u_n)))))_{n\in\mathbb{N}}$ and $(H_0^1(I(f(u_n)) \cup (A_\psi(f(u_n)) \setminus A_{\mathrm{s}}^\psi(f(u_n)))))_{n\in\mathbb{N}}$, respectively. Section 6 uses now stability properties of variational equalities under Mosco convergence to characterize elements of the Bouligand generalized differential. Finally, an adjoint representation of corresponding Clarke subgradients for an objective functional is derived.

# 2 Fundamental Definitions and Results

Denote by $C_{\mathrm{c}}(\Omega)$ the space of continuous functions on $\Omega$ with compact support contained in $\Omega$ and by $C_{\mathrm{c}}^\infty(\Omega)$ the subspace of infinitely differentiable functions. Furthermore, we denote by $H^1(\Omega)$ the space

$$H^1(\Omega) := \left\{ z \in L^2(\Omega) \mid \tfrac{\partial z}{\partial x_i} \in L^2(\Omega), i = 1, \dots, d \right\},$$

where $\frac{\partial z}{\partial x_i}$ is to be understood in the distributional sense. $H^1(\Omega)$ is equipped with the norm

$$\|z\|_{H^1(\Omega)} = \left( \int_\Omega z^2 + \sum_{i=1}^d \left( \frac{\partial z}{\partial x_i} \right)^2 \mathrm{d}\lambda^d \right)^{1/2}.$$

In this paper, we work with the space $H_0^1(\Omega)$ and we define it as the completion of $C_{\mathrm{c}}^\infty(\Omega)$ in $H^1(\Omega)$. On $H_0^1(\Omega)$ we consider the norm $\|z\|_{H_0^1(\Omega)} := \|\nabla z\|_{L^2(\Omega)}$. The dual space of $H_0^1(\Omega)$ is denoted by $H^{-1}(\Omega)$ and if $\zeta \in H^{-1}(\Omega)$ and $z \in H_0^1(\Omega)$ we use the notation $\langle \zeta, z\rangle$ for the dual pairing.

Note that $z \in H_0^1(\Omega)$ can be extended by zero to an element of $z \in H^1(\mathbb{R}^d)$, since the zero extension of the approximating sequence in $C_{\mathrm{c}}^\infty(\Omega)$ is a Cauchy sequence in $H^1(\mathbb{R}^d)$.

We denote by $H^1(\Omega)_+$, respectively $H_0^1(\Omega)_+$, the respective subsets of nonnegative elements. For $u \in H_0^1(\Omega)$ and $v \in H^1(\Omega)_+$ we often use that $\min(u, v) \in H_0^1(\Omega)$. We use the notation $u_+ := \max(0, u)$, $u_- = -\min(0, u)$, for the positive and negative part of $u \in L^2(\Omega)$ and have $u = u_+ - u_-$. Furthermore, for $n \in \mathbb{N}$ and $u \in H_0^1(\Omega)$, $u_n := \max(-n, \min(u, n))$ is in $H_0^1(\Omega) \cap L^\infty(\Omega)$ and it holds $u_n \to u$ in $H_0^1(\Omega)$, which can be seen by application of Lebesgue's dominated convergence theorem.

## 2.1 Capacity Theory

We quickly recall and clarify the definitions and concepts related to capacity theory that we consider in this paper. For the definitions, see also [ABM14, Sect. 5.82, 5.83], [DZ11, Def. 6.2], [KM92].

**Definition 2.1.** 1. For a set $E \subseteq \Omega$ we define the capacity of $E$ in $\Omega$ by

$$\operatorname{cap}(E) := \inf\{\|z\|^2_{H^1_0(\Omega)} \mid z \in H^1_0(\Omega), z \geq 1 \text{ a.e. in a neighborhood of } E\}. \quad (2)$$

If a property holds on a set $E \subseteq \Omega$ except on a subset of capacity zero, we say that this property holds quasi-everywhere (q.e.) on $E$.

2. We call a set $O \subseteq \Omega$ quasi-open if for all $\varepsilon > 0$ there is an open set $O_\varepsilon \subseteq \Omega$ such that $O \cup O_\varepsilon$ is open and $\operatorname{cap}(O_\varepsilon) < \varepsilon$. A set $A \subseteq \Omega$ is quasi-closed if the complement in $\Omega$ is quasi-open.
3. Let $v\colon \Omega \to \mathbb{R}$ be a function. Then $v$ is quasi-continuous if for all $\varepsilon > 0$ there exists an open set $O_\varepsilon \subseteq \Omega$ such that $v|_{\Omega \setminus O_\varepsilon}$ is continuous and $\operatorname{cap}(O_\varepsilon) < \varepsilon$.
4. Let $E \subseteq \Omega$ be a set. A family $(O_i)_{i\in I}$ of quasi-open subsets of $\Omega$ is called a quasi-covering of $E$ if there is a countable subfamily $(O_{i_n})_{n\in\mathbb{N}}$ satisfying $\operatorname{cap}\left(E \setminus \bigcup_{n\in\mathbb{N}} O_{i_n}\right) = 0$.
5. Let $O \subseteq \Omega$ be quasi-open. Then we define

$$H^1_0(O) := \{z \in H^1_0(\Omega) \mid z = 0 \text{ q.e. on } \Omega \setminus O\}.$$

**Remark 2.2.** If $O \subseteq \Omega$ is open, then the definition of $H^1_0(O)$ in Definition 2.1 coincides with the classical definition of $H^1_0(O)$ that we also use in this paper, see e.g. [AH96, Thm. 9.1.3]. Moreover, the definition of $H^1_0(O)$ for $O \subseteq \Omega$ quasi-open coincides with the definition

$$H^1_0(O) = \bigcap\{H^1_0(G) \mid O \subseteq G \subseteq \Omega, G \text{ open}\}$$

given in [KM92] (up to extension of the elements by 0).

We could also define a capacity Cap for subsets of $\mathbb{R}^d$ by testing with $H^1(\mathbb{R}^d)$-elements in (2) and by considering the infimum over the squared $H^1(\mathbb{R}^d)$-norms. Then $H^1_0(U) = \{v \in H^1(\mathbb{R}^d) \mid v = 0 \text{ Cap-q.e. outside } U\}$ for Cap-quasi-open subsets $U$ of $\mathbb{R}^d$. For $U \subseteq \Omega$, $U$ is Cap-quasi-open if and only if it is quasi-open and the definitions of $H^1_0(U)$ coincide.

**Lemma 2.3**

1. *Let $v \in H^1(\Omega)$. Then $v$ has a quasi-continuous representative $\tilde{v}$. If $\tilde{w}$ is another quasi-continuous representative of $v$, then $\tilde{v} = \tilde{w}$ up to a set of capacity zero.*
2. *Suppose $(v_n)_{n\in\mathbb{N}}, v \subseteq H^1_0(\Omega)$ and $v_n \to v$ in $H^1_0(\Omega)$. Then there is a subsequence $(v_{n_k})_{k\in\mathbb{N}}$ such that $\tilde{v}_{n_k} \to \tilde{v}$ pointwise q.e. for the quasi-continuous representatives.*
3. *Let $O \subseteq \Omega$ be a quasi-open set, let $z \in H^1_0(O)$ and assume that $(O_i)_{i\in I}$ is a quasi-covering of $O$. Then there exists a sequence $(z_n)_{n\in\mathbb{N}} \subseteq H^1_0(O)$ such that $z_n \to z$ and such that each $z_n$ is a finite sum of functions from $\bigcup_{i\in I} H^1_0(O_i)$. If $(O_n)_{n\in\mathbb{N}}$ is a quasi-covering that is increasing in $n$, then we find a sequence $(z_n)_{n\in\mathbb{N}}$ converging to $z$ such that $z_n \in H^1_0(O_n)$. If $z \in H^1_0(\Omega)_+$, then, w.l.o.g., $(z_n)_{n\in\mathbb{N}} \subseteq H^1_0(\Omega)_+$.*
4. *Assume $O \subseteq \Omega$ is quasi-open and suppose $v\colon \Omega \to \mathbb{R}$ is quasi-continuous. Then, $v \geq 0$ a.e. on $O$ if and only if $v \geq 0$ q.e. on $O$.*
5. *Assume $O \subseteq \Omega$ is a quasi-open set. Then there exists $v \in H^1_0(\Omega)_+$ with $\{\tilde{v} > 0\} = O$ up to a set of zero capacity.*

6. *After modification on a subset of capacity zero, each quasi-continuous function $v\colon \Omega \to \mathbb{R}$ is Borel measurable.*

*Proof.* The first statement can be found, e.g., in [DZ11, Chap. 6, Thm. 6.1] or [HKM93, Thm. 4.4], the second in [BS00, Lem. 6.52]. The first part of the third statement can be obtained by combining [KM92, Lem. 2.4 and Lem. 2.10]. For the second part of the third statement use that $(O_n)_{n\in\mathbb{N}}$ is increasing and use the first part. In case $z \geq 0$, the statement in [KM92, Lem. 2.4] and the proof of [KM92, Lem. 2.10] imply that we can choose $(z_n)_{n\in\mathbb{N}} \subseteq H_0^1(\Omega)_+$. We refer to [Wac14, Lem. 2.3] for the fourth statement and to [Vel15, Prop. 2.3.14], [HW18, Lem. 3.6] for the fifth statement. The last statement can be found in [RW19, Sec. 2.1]. □

**Remark 2.4.** Let $v\colon \Omega \to \mathbb{R}$ be a quasi-continuous function. Then the set $\{v > 0\}$ is quasi-open and the set $\{v \geq 0\}$ is quasi-closed. If $v \in H^1(\Omega)$, then by $\{v > 0\}, \{v \geq 0\}$ we always mean the sets $\{\tilde{v} > 0\}, \{\tilde{v} \geq 0\}$, where $\tilde{v}$ is a quasi-continuous and Borel measurable representative. Thus, these sets are Borel measurable, quasi-open, respectively quasi-closed, and determined up to a set of capacity zero.

Throughout the rest of the paper, when considering set equations or inclusions for subsets of $\Omega$, they have to be understood to hold up to an exceptional set of capacity zero.

## 2.2 Generalized Derivatives

We consider the following generalized differential for the solution operator of (1).

**Definition 2.5.** Consider a separable Banach space $X$ and a Hilbert space $Y$. Assume that $T\colon X \to Y$ is a locally Lipschitz continuous operator. The set of Bouligand generalized derivatives of $T$ in $x \in X$ is defined as

$$\begin{aligned} \partial T(x) := \{\Xi \in \mathcal{L}(X,Y) \mid \exists (x_n)_{n\in\mathbb{N}} \subseteq \mathcal{D}_T \text{ with } x_n \to x \\ \text{and } T'(x_n) \to \Xi \text{ in the weak operator topology}\}, \end{aligned} \tag{3}$$

for $\mathcal{D}_T := \{x \in X \mid T \text{ is Gâteaux differentiable at } x \text{ with Gâteaux derivative } T'(x)\}$.

For infinite dimensional spaces, there are several choices of topologies in $X$ and $Y$. Combinations of strong and weak topologies in $X$ and $Y$ lead to four possible definitions that do not coincide, in general. The four versions are defined, e.g. in [CMWC18, Def. 3.1], [RW19, Def. 2.10], and characterized for the solution operators of a nonsmooth semilinear elliptic equation and the unilateral obstacle problem with distributed controls, respectively. The set $\partial T(x)$ as defined in Definition 2.5 is always nonempty as a consequence of Rademacher's theorem for locally Lipschitz continuous mappings. Nevertheless, the generalized derivatives we construct in this paper are also contained in the generalized differential that can be obtained by replacing the weak operator topology in (3) by the strong operator topology. A priori it is not clear that this set is nonempty.

**Remark 2.6.** Let $J\colon H_0^1(\Omega) \times U \to \mathbb{R}$ be a continuously differentiable objective function. Denote by $S_f\colon U \to H_0^1(\Omega)$ the solution operator of (1). We use the notation $\hat{J} := J(S_f(\cdot),\cdot)$ for the reduced objective function and denote by $\partial_C \hat{J}$ Clarke's generalized differential of $\hat{J}\colon U \to \mathbb{R}$. Let $u \in U$ be arbitrary. Then the set inclusion

$$\{\Xi^* J_y(S_f(u),u) + J_u(S_f(u),u) \mid \Xi \in \partial S_f(u)\} \subseteq \partial \hat{J}(u) \subseteq \partial_C \hat{J}(u)$$

holds.

# 3 Properties of the Solution Operator

In this section, we collect properties of the solution operator $S_f$ of (1).

## 3.1 Monotonicity Properties of the Solution Operator

We state two lemmata on monotonicity properties of $S_f$. The next lemma summarizes the monotonicity of $S_f$ with respect to the elements in $U$. The proof is similar to the proof of [Rod87, Sect. 4:5, Thm. 5.1] where the property is shown for the unilateral obstacle problem.

**Lemma 3.1**
*Let $u_1, u_2 \in U$ with $u_1 \geq_U u_2$. Then $S_f(u_1) \geq S_f(u_2)$ a.e. and q.e. in $\Omega$.*

*Proof.* For $i = 1, 2$ set $y_i := S_f(u_i)$. We test the variational inequality characterizing $y_1$ with $z_1 = \max(y_1, y_2) = y_1 + (y_2 - y_1)_+$ and the variational inequality characterizing $y_2$ with $z_2 = \min(y_1, y_2) = y_2 - (y_2 - y_1)_+$, respectively, and obtain

$$0 \leq \langle Ly_1 - f(u_1), z_1 - y_1 \rangle = \langle Ly_1 - f(u_1), (y_2 - y_1)_+ \rangle$$

and

$$0 \leq \langle Ly_2 - f(u_2), z_2 - y_2 \rangle = \langle Ly_2 - f(u_2), -(y_2 - y_1)_+ \rangle.$$

Summing up both inequalities we obtain

$$\langle Ly_1 - Ly_2, (y_2 - y_1)_+ \rangle \geq \langle f(u_1) - f(u_2), (y_2 - y_1)_+ \rangle \geq 0.$$

By strict T-monotonicity, we have $(y_2 - y_1)_+ = 0$, i.e., $y_1 \geq y_2$ a.e. and q.e. in $\Omega$. □

The following lemma establishes monotonicity properties of $S_f$ with respect to one of the obstacles.

**Lemma 3.2**
*Let $\psi_i \in H^1(\Omega)$, $i = 1, 2$, such that $K^{\varphi}_{\psi_i}$ are nonempty, and denote by $y_i$ the corresponding solutions of (1) for fixed $u \in U$. Then $\psi_1 \geq \psi_2$ implies $y_1 \geq y_2$ a.e. and q.e. in $\Omega$.*

*Proof.* We test the variational inequality characterizing $y_1$ with the element $z_1 = \max(y_1, y_2) = y_1 + (y_2 - y_1)_+ \in K^{\varphi}_{\psi_1}$ and the variational inequality characterizing $y_2$ with $z_2 = \min(y_1, y_2) = y_2 - (y_2 - y_1)_+ \in K^{\varphi}_{\psi_2}$, respectively, and obtain

$$0 \leq \langle Ly_1 - f(u), (y_2 - y_1)_+ \rangle, \quad 0 \leq \langle Ly_2 - f(u), -(y_2 - y_1)_+ \rangle.$$

Summing up both inequalities we obtain

$$\langle L(y_1 - y_2), (y_2 - y_1)_+ \rangle \geq 0.$$

By strict T-monotonicity we have $(y_2 - y_1)_+ = 0$, i.e., $y_1 \geq y_2$ a.e. and q.e. in $\Omega$. □

## 3.2 Differentiability Properties of the Solution Operator

We distinguish the following subsets of $\Omega$ for a fixed element $\zeta \in H^{-1}(\Omega)$ that result from the solution $S_{\mathrm{id}}(\zeta)$ of (1) for $f = \mathrm{id}$ being the identity operator on $H^{-1}(\Omega)$. Let $\zeta \in H^{-1}(\Omega)$. By

$$A(\zeta) := \{\omega \in \Omega \mid S_{\mathrm{id}}(\zeta)(\omega) = \psi(\omega) \text{ or } S_{\mathrm{id}}(\zeta)(\omega) = \varphi(\omega)\}$$

we denote the active set. We also distinguish the active sets with respect to $\psi$ and $\varphi$, i.e., we define

$$A_\psi(\zeta) := \{\omega \in \Omega \mid S_{\mathrm{id}}(\zeta)(\omega) = \psi(\omega)\} \quad \text{and} \quad A^\varphi(\zeta) := \{\omega \in \Omega \mid S_{\mathrm{id}}(\zeta)(\omega) = \varphi(\omega)\}.$$

Note that $A(\zeta) = A_\psi(\zeta) \cup A^\varphi(\zeta)$ and that all these sets are quasi-closed sets that are determined up to a set of capacity zero, since we consider quasi-continuous representatives of $S_{\mathrm{id}}(\zeta), \psi, \varphi \in H^1(\Omega)$ in the definition of the active sets. We denote by $I(\zeta)$ the inactive set, i.e. the complement of $A(\zeta)$ in $\Omega$ and by $I_\psi(\zeta) := \Omega \setminus A_\psi(\zeta)$, respectively $I^\varphi(\zeta) := \Omega \setminus A^\varphi(\zeta)$, the inactive sets with respect to the two obstacles.

Let $u, h \in U$. It can be shown that the directional derivative $S_f'(u;h)$ of the solution operator of variational inequality (1) is given by the solution of

$$\begin{aligned} &\text{Find } \eta \in (LS_f(u) - f(u))^\perp \cap T_{K_\psi^\varphi}(S_f(u)) : \\ &\langle L\eta - f'(u;h), z - \eta\rangle \geq 0 \quad \forall\, z \in (LS_f(u) - f(u))^\perp \cap T_{K_\psi^\varphi}(S_f(u)). \end{aligned} \tag{4}$$

Here, $T_{K_\psi^\varphi}(S_f(u))$ is the tangent cone of $K_\psi^\varphi$ at $S_f(u)$, i.e., the closed conic hull of $K_\psi^\varphi - S_f(u)$. When $f$ is the identity operator on $H^{-1}(\Omega)$, the variational inequality (4) follows, e.g., from [Mig76, Thm. 3.3]. Since $S_f$ is locally Lipschitz continuous, $S_f$ is even directionally differentiable in the sense of Hadamard, see [BS00, Prop. 2.49]. When considering a general operator $f\colon U \to H^{-1}(\Omega)$ fulfilling our assumptions, (4) can be obtained using the chain rule for Hadamard directionally differentiable maps, see e.g. [BS00, Prop. 2.47].

By [Mig76, Lem. 3.4], we have

$$T_{K_\psi^\varphi}(S_f(u)) = \{z \in H_0^1(\Omega) \mid z \geq 0 \text{ q.e. on } A_\psi(f(u)), z \leq 0 \text{ q.e. on } A^\varphi(f(u))\}. \tag{5}$$

### 3.2.1 Analysis of the Critical Cone

As in the case with a single obstacle, we want to find a suitable characterization of the critical cone $(LS_{\mathrm{id}}(\zeta) - \zeta)^\perp \cap T_{K_\psi^\varphi}(S_{\mathrm{id}}(\zeta))$ for arbitrary $\zeta \in H^{-1}(\Omega)$. Note that in the case with a single lower obstacle such a characterization is given by $\{z \in H_0^1(\Omega) \mid z \geq 0 \text{ q.e. on } A(\zeta), z = 0 \text{ q.e. on } A_{\mathrm{s}}(\zeta)\}$, see [Wac14, Lem. 3.1]. Here, $A_{\mathrm{s}}(\zeta)$ is the strictly active set, which can also be characterized as in [Wac14, App. A].

A crucial difference to the case with only one obstacle is that $LS_{\mathrm{id}}(\zeta) - \zeta$ is not a nonnegative functional and thus cannot be identified with a positive measure. Instead, we will see that, in some cases, it can be identified with the difference of two nonnegative Radon measures. In general, i.e., when the active sets $A_\psi(\zeta)$ and $A^\varphi(\zeta)$ do not have a positive distance, $LS_{\mathrm{id}}(\zeta) - \zeta$ acts as the difference of two measures on all elements of $H_0^1(\Omega) \cap L^\infty(\Omega)$, but the characterization does not carry over to unbounded elements of $H_0^1(\Omega)$, see Example 3.6.

We define the set of nonnegative Radon measures $\mathcal{M}_+(\Omega)$ on $\Omega$ as

$$\mathcal{M}_+(\Omega) = \{\mu \mid \mu \text{ is a regular, locally finite Borel measure on } \Omega\}.$$

Before we present the theorem on the representation, let us state an auxiliary lemma.

**Lemma 3.3**
*Assume that $v \in L^\infty(\Omega) \cap H_0^1(\Omega)$. Moreover, let $(w_n)_{n\in\mathbb{N}} \subseteq L^\infty(\Omega) \cap H_0^1(\Omega)$ be a sequence with $w_n \to 0$ in $H_0^1(\Omega)$ and $|w_n| \leq C$ a.e. for some $C > 0$ and all $n \in \mathbb{N}$. Then $v\, w_n \to 0$ in $H_0^1(\Omega)$.*

*Proof.* Let us recall that $v\, w_n \in H_0^1(\Omega)$ and $\nabla(v\, w_n) = w_n \nabla v + v \nabla w_n$, see, e.g., [GT01, (7.18)]. Thus,

$$\|v\, w_n\|_{H_0^1(\Omega)} \leq \|w_n \nabla v\|_{L^2(\Omega)} + \|v \nabla w_n\|_{L^2(\Omega)}$$

holds. The second term tends to zero since $v \in L^\infty(\Omega)$ and $\nabla w_n \to 0$ in $L^2(\Omega)$. Moreover, for a subsequence, the term $\|w_n \nabla v\|_{L^2(\Omega)}$ converges to zero aswell. To see this, pick any subsequence and choose a subsubsequence, for simplicity again denoted by $(w_n)_n$, that converges to 0 pointwise q.e., see Lemma 2.3, and thus pointwise $\lambda^d$-a.e. Now, $\|w_n \nabla v\|_{L^2(\Omega)} \to 0$ follows from Lebesgue's dominated convergence theorem and $|w_n \nabla v| \leq C|\nabla v| \in L^2(\Omega)$. The assertion for the whole sequence is obtained by a subsequence-subsequence argument. □

Now, we derive a characterization of the multiplier as the difference of two nonnegative measures. Related results can be found in [Wac18, Lem. 4.2, Thm. 4.3], see also [BB79, GR77]. Therein, the admissible set does not depend on the spatial variable and, thus, the results do not apply to our setting immediately.

**Theorem 3.4**
*Assume that $\psi, \varphi$ fulfill the conditions of Assumption 1.1. Let $\zeta \in H^{-1}(\Omega)$ be arbitrary and set $y := S_{\mathrm{id}}(\zeta)$, $\xi := Ly - \zeta$. Then the following statements hold.*

1. *$\xi \in H^{-1}(\Omega)$ acts as the difference $\tilde{\xi}_\psi - \tilde{\xi}^\varphi$ of nonnegative measures $\tilde{\xi}_\psi, \tilde{\xi}^\varphi \in \mathcal{M}_+(\Omega)$ on all elements of $H_0^1(\Omega) \cap C_{\mathrm{c}}(\Omega)$, i.e.,*

$$\langle \xi, w \rangle = \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi \tag{6}$$

   *holds for all $w \in H_0^1(\Omega) \cap C_{\mathrm{c}}(\Omega)$.*
2. *Let $A \subseteq \Omega$ be an arbitrary Borel set. Then $\mathrm{cap}(A) = 0$ implies $\tilde{\xi}_\psi(A) = \tilde{\xi}^\varphi(A) = 0$.*
3. *The characterization (6) carries over to all $w \in H_0^1(\Omega) \cap L^\infty(\Omega)$. In particular, the quasi-continuous and Borel measurable representatives of $w$ are $\tilde{\xi}_\psi$- and $\tilde{\xi}^\varphi$-integrable.*
4. *Furthermore, it holds $y = \psi$ $\tilde{\xi}_\psi$-a.e. on $\Omega$ and $y = \varphi$ $\tilde{\xi}^\varphi$-a.e. on $\Omega$, i.e., $\tilde{\xi}_\psi(I_\psi(\zeta)) = 0$ and $\tilde{\xi}^\varphi(I^\varphi(\zeta)) = 0$.*
5. *Assume $w \in H_0^1(\Omega) \cap L^1(\tilde{\xi}_\psi)$. Then we have $w \in L^1(\tilde{\xi}^\varphi)$ and (6) holds for $w$. The opposite statement with exchanged roles of $\tilde{\xi}_\psi$ and $\tilde{\xi}^\varphi$ is also true.*

*Proof.* ad 1.: We define

$$v := \frac{y - \psi}{\varphi - \psi}.$$

By the assumptions on $\psi$ and $\varphi$, we have $0 \leq v \leq 1$ and $v \in H^1(\Omega) \cap L^\infty(\Omega)$.

Now, for $w \in H_0^1(\Omega) \cap L^\infty(\Omega)$, we have $v\,w$, $(1-v)\,w \in H_0^1(\Omega)$ and we write

$$\langle \xi, w \rangle = \langle \xi, (1-v)\,w \rangle + \langle \xi, v\,w \rangle.$$

Thus, we deduce

$$\langle \xi, w \rangle = \xi_\psi(w) - \xi^\varphi(w),$$

where $\xi_\psi$, $\xi^\varphi$ are defined by

$$\xi_\psi \colon w \mapsto \langle \xi, (1-v)\,w \rangle, \quad \xi^\varphi \colon w \mapsto \langle \xi, -v\,w \rangle.$$

Note that $\xi_\psi$, $\xi^\varphi$ are nonnegative linear forms on $H_0^1(\Omega) \cap L^\infty(\Omega)$. To see this, assume $w \in H_0^1(\Omega)_+ \cap L^\infty(\Omega)$ and let first $\|w\|_{L^\infty} \leq c_\psi^\varphi$. By definition of $v$, we have $-v\,w + y \in K_\psi^\varphi$ and therefore

$$\xi^\varphi(w) = \langle \xi, -v\,w + y - y \rangle \geq 0. \tag{7}$$

Since $\xi^\varphi$ is linear, (7) holds for all $w \in H_0^1(\Omega)_+ \cap L^\infty(\Omega)$. In a similar fashion, we can show that $\xi_\psi$ is nonnegative on $H_0^1(\Omega)_+ \cap L^\infty(\Omega)$.

In particular, $\xi_\psi$, $\xi^\varphi$ are nonnegative linear forms on $H_0^1(\Omega) \cap C_c(\Omega)$. By [BS00, Lem. 6.53], $\xi_\psi$ and $\xi^\varphi$ have unique nonnegative continuous extensions over $C_c(\Omega)$, also denoted by $\xi_\psi$, respectively $\xi^\varphi$. Moreover, by [BS00, Thm. 6.54], there are unique nonnegative, regular, locally finite Borel measures $\tilde{\xi}_\psi$, $\tilde{\xi}^\varphi$ such that

$$\xi_\psi(w) = \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi \quad \text{and} \quad \xi^\varphi(w) = \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi$$

holds for all $w \in C_c(\Omega)$.

ad 2.: Now, we modify the proof of [BS00, Lem. 6.55] to show that for a Borel set $A \subseteq \Omega$, $\mathrm{cap}(A) = 0$ implies $\tilde{\xi}_\psi(A) = \tilde{\xi}^\varphi(A) = 0$. W.l.o.g. we prove the statement for $\tilde{\xi}^\varphi$.

Let $(\varepsilon_n)_{n\in\mathbb{N}} \subseteq \mathbb{R}_+$ be a sequence with $\varepsilon_n \to 0$ as $n \to \infty$. Fix $n \in \mathbb{N}$. Then we find an open superset $A_n$ of $A$ in $\Omega$ with $\mathrm{cap}(A_n) < \varepsilon_n$. Furthermore, by [HW18, Lem. 3.4], there is $u_n \in H_0^1(\Omega)_+$ satisfying $u_n = 1$ q.e. on $A_n$ as well as $\|u_n\|^2_{H_0^1(\Omega)} = \mathrm{cap}(A_n) < \varepsilon_n$. Moreover, we can assume $u_n \in L^\infty(\Omega)$ and $0 \le u_n \le 1$, since $\min(z, 1) \in H_0^1(\Omega)$ satisfies $\|\min(z,1)\|_{H_0^1(\Omega)} \le \|z\|_{H_0^1(\Omega)}$ for $z \in H_0^1(\Omega)$. By regularity of $\tilde{\xi}^\varphi$, we can find a compact set $K_n \subseteq A_n$ satisfying $\tilde{\xi}^\varphi(A_n) \le \tilde{\xi}^\varphi(K_n) + \varepsilon_n$. Using a smooth version of Urysohn's lemma, there exists a function $g_n \in H_0^1(\Omega) \cap C_c(\Omega)$ with values in $[0,1]$ satisfying $g_n = 1$ on $K_n$ and having compact support in $A_n$. Then we have $1_{K_n} \le g_n \le u_n$ q.e. on $\Omega$.

Now, we conclude

$$\begin{aligned}\tilde{\xi}^\varphi(A_n) \le \tilde{\xi}^\varphi(K_n) + \varepsilon_n \le \int_\Omega g_n \, \mathrm{d}\tilde{\xi}^\varphi + \varepsilon_n = \langle \xi, -v\, g_n\rangle + \varepsilon_n \\ \le \langle \xi, -v\, u_n\rangle + \varepsilon_n \le \|\xi\|_{H^{-1}(\Omega)} \|v\, u_n\|_{H_0^1(\Omega)} + \varepsilon_n.\end{aligned}$$

Using Lemma 3.3, we know that $\|v\, u_n\|_{H_0^1(\Omega)} \to 0$. Now, $\bigcap_{n\in\mathbb{N}} A_n$ is Borel measurable and

$$\tilde{\xi}^\varphi\left(\bigcap_{n\in\mathbb{N}} A_n\right) = 0.$$

Since $A \subseteq \bigcap_{n\in\mathbb{N}} A_n$, we conclude $\tilde{\xi}^\varphi(A) = 0$.

ad 3.: Now, we argue in a similar fashion as in [BS00, Lem. 6.56] to show that each $w \in H_0^1(\Omega) \cap L^\infty(\Omega)$ satisfies

$$\langle \xi, w\rangle = \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi.$$

Let $w \in H_0^1(\Omega) \cap L^\infty(\Omega)$. Then we find $(\overline{w}_n)_{n\in\mathbb{N}} \subseteq C_c^\infty(\Omega)$ with $\overline{w}_n \to w$ in $H_0^1(\Omega)$. Definig $w_n := \max(-\|w\|_{L^\infty(\Omega)}, \min(\overline{w}_n, \|w\|_{L^\infty(\Omega)}))$ we have $w_n \in H_0^1(\Omega) \cap C_c(\Omega)$ and $w_n \to w$ in $H_0^1(\Omega)$ as well as $|w_n| \le \|w\|_{L^\infty(\Omega)}$. Then, Lemma 3.3 yields $v\, w_n \to v\, w$ in $H_0^1(\Omega)$. Therefore, using $\||z|\|_{H_0^1(\Omega)} = \|z\|_{H_0^1(\Omega)}$ for all $z \in H_0^1(\Omega)$, see [ABM14, Cor. 5.8.1], with

$$\| - v\,|w_n - w_m|\|_{H_0^1(\Omega)} = \big\|\,|-v\,|w_n - w_m|\|\big\|_{H_0^1(\Omega)} = \|v\, w_n - v\, w_m\|_{H_0^1(\Omega)}$$

and

$$\|w_n - w_m\|_{L^1(\tilde{\xi}^\varphi)} = \langle \xi, -v\,|w_n - w_m|\rangle \le \|\xi\|_{H^{-1}(\Omega)} \| - v\,|w_n - w_m|\|_{H_0^1(\Omega)}$$

we find that $(w_n)_{n\in\mathbb{N}}$ is a Cauchy sequence in $L^1(\tilde{\xi}^\varphi)$. Similarly, $(w_n)_{n\in\mathbb{N}}$ is a Cauchy sequence in $L^1(\tilde{\xi}_\psi)$. Thus, a subsequence converges pointwise $\tilde{\xi}^\varphi$-a.e. and $\tilde{\xi}_\psi$-a.e. to an element in $L^1(\tilde{\xi}^\varphi) \cap L^1(\tilde{\xi}_\psi)$. Now, pick a subsubsequence that converges pointwise q.e. to $w$. Then, by the second statement of Theorem 3.4 and since the convergence holds except on a Borel set of capacity zero, $w \in L^1(\tilde{\xi}^\varphi) \cap L^1(\tilde{\xi}_\psi)$ and Lebesgue's dominated convergence theorem implies

$$\langle \xi^\varphi, w\rangle = \underbrace{\langle \xi, -v(w-w_n)\rangle}_{\to 0 \text{ as } n\to\infty} + \underbrace{\int_\Omega w_n \,\mathrm{d}\tilde{\xi}^\varphi}_{\to \int_\Omega w \,\mathrm{d}\tilde{\xi}^\varphi \text{ along a subsequence}}.$$

This yields $\langle \xi^\varphi, w\rangle = \int_\Omega w \, d\tilde{\xi}^\varphi$ and, similarly, $\langle \xi_\psi, w\rangle = \int_\Omega w \,\mathrm{d}\tilde{\xi}_\psi$.

ad 4.: We modify the proof of [Wac14, Prop. 2.5]. We consider a smooth cut-off function $\chi \in C_c^\infty(\Omega)$ with $0 \le \chi \le 1$ and $\chi = 1$ on a compact set $K \subseteq \Omega$.

We define $w := \chi\,[(1-v)\,\psi + v\,y] + (1-\chi)\,y$ and obtain $w \in K_\psi^\varphi$. This implies

$$\begin{aligned} 0 \le \langle \xi, w-y\rangle &= \langle \xi, \chi\,(1-v)\,\psi + \chi\,v\,y - \chi\,y\rangle \\ &= \langle \xi, (1-v)\,\chi\,(\psi-y)\rangle = \int_\Omega \chi\,(\psi-y)\,\mathrm{d}\tilde{\xi}_\psi. \end{aligned}$$

Since $\chi\,(\psi-y) \le 0$ q.e. on $\Omega$, and thus, by the second statement of the theorem, $\tilde{\xi}_\psi$-a.e., we conclude $y = \psi$ $\tilde{\xi}_\psi$-a.e. on $K$. Covering $\Omega$ with countably many compact subsets, we infer $y-\psi = 0$ $\tilde{\xi}_\psi$-a.e. on $\Omega$. Similarly, we can show $\varphi - y = 0$ $\tilde{\xi}^\varphi$-a.e. on $\Omega$.

ad 5.: Assume $w \in H_0^1(\Omega) \cap L^1(\tilde{\xi}_\psi)$. We approximate $w$ in $H_0^1(\Omega)$ by $(w_n)_{n\in\mathbb{N}}$ defined via $w_n := \max(-n, \min(n, w))$. Then we have $w_n \to w$ in $H_0^1(\Omega)$ and $w_n \to w$ pointwise $\tilde{\xi}_\psi$-a.e. (after choosing a subsequence). Since $|w_n| \le |w|$ and since $w \in L^1(\tilde{\xi}_\psi)$, we apply Lebesgue's dominated convergence theorem and obtain $w_n \to w$ in $L^1(\tilde{\xi}_\psi)$.

From

$$\begin{aligned} \|w_n - w_m\|_{L^1(\tilde{\xi}^\varphi)} &= \int_\Omega |w_n - w_m| \,\mathrm{d}\tilde{\xi}^\varphi = \int_\Omega |w_n - w_m| \,\mathrm{d}\tilde{\xi}_\psi - \langle \xi, |w_n - w_m|\rangle \\ &\le \|w_n - w_m\|_{L^1(\tilde{\xi}_\psi)} + \|\xi\|_{H^{-1}(\Omega)} \|w_n - w_m\|_{H_0^1(\Omega)} \end{aligned}$$

it follows that $(w_n)_{n\in\mathbb{N}}$ is a Cauchy sequence in $L^1(\tilde{\xi}^\varphi)$ and we can again conclude that $w_n \to w$ in $L^1(\tilde{\xi}^\varphi)$.

From the representation

$$\langle \xi, w_n\rangle = \int_\Omega w_n \,\mathrm{d}\tilde{\xi}_\psi - \int_\Omega w_n \,\mathrm{d}\tilde{\xi}^\varphi$$

for all $n \in \mathbb{N}$, since $w_n \to w$ in $H_0^1(\Omega)$, $L^1(\tilde{\xi}_\psi)$ and $L^1(\tilde{\xi}^\varphi)$, we conclude

$$\langle \xi, w\rangle = \int_\Omega w \,\mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \,\mathrm{d}\tilde{\xi}^\varphi.$$

The opposite statement follows similarly. □

In the subsequent lemma we assume that the active sets $A_\psi(\zeta)$ and $A^\varphi(\zeta)$ have a positive distance. With this condition we mean that there are representative quasi-closed sets Borel measurable $\tilde{A}_\psi(\zeta)$, $\tilde{A}^\varphi(\zeta)$ such that $\operatorname{dist}(\tilde{A}_\psi(\zeta), \tilde{A}^\varphi(\zeta)) > 0$ and the two sets coincide with $A_\psi(\zeta), A^\varphi(\zeta)$ up to a set of capacity zero.

**Lemma 3.5**
*Assume that $\psi, \varphi$ fulfill the conditions of Assumption 1.1. Let $\zeta \in H^{-1}(\Omega)$ be arbitrary and set $y := S_{\mathrm{id}}(\zeta)$ and $\xi := Ly - \zeta$. Suppose $A_\psi(\zeta)$ and $A^\varphi(\zeta)$ have a positive distance. Then, with $\tilde{\xi}_\psi, \tilde{\xi}^\varphi \in \mathcal{M}_+(\Omega)$ as in Theorem 3.4, it holds $H_0^1(\Omega) \subseteq L^1(\tilde{\xi}_\psi) \cap L^1(\tilde{\xi}^\varphi)$ and*

$$\langle \xi, w \rangle = \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi$$

*for all $w \in H_0^1(\Omega)$.*

*Proof.* Since the active sets have a positive distance $C > 0$, we can find an element $v_2$ satisfying $v_2 \in C^\infty(\mathbb{R}^d)$ as well as $v_2 = 1$ on $A_\psi(\zeta)$ and $v_2 = 0$ outside $A_\psi(\zeta) + B_{C/2}$.

Since $v_2$ is smooth, we have $v_2\, w, (v_2 - 1)\, w \in H_0^1(\Omega)$ for all $w \in H_0^1(\Omega)$ and we define the functionals

$$\xi_\psi^2 \colon w \mapsto \langle \xi, v_2\, w \rangle, \quad \xi_2^\varphi \colon w \mapsto \langle \xi, (v_2 - 1)\, w \rangle$$

on $H_0^1(\Omega)$. Since $v_2$ is in $C^\infty(\mathbb{R}^d)$, it is easy to show that $\xi_\psi^2$ and $\xi_2^\varphi$ are bounded linear functionals on $H_0^1(\Omega)$. Moreover, we have

$$\langle \xi, w \rangle = \langle \xi_\psi^2, w \rangle - \langle \xi_2^\varphi, w \rangle$$

for all $w \in H_0^1(\Omega)$.

Assume first $w$ is in $H_0^1(\Omega) \cap L^\infty(\Omega)$. Then we have

$$\int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi = \int_\Omega v_2\, w \, \mathrm{d}\tilde{\xi}_\psi = \int_\Omega v_2\, w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega v_2\, w \, \mathrm{d}\tilde{\xi}^\varphi = \langle \xi, v_2\, w \rangle = \langle \xi_\psi^2, w \rangle.$$

Here, the first equation holds since $\tilde{\xi}_\psi(I_\psi(\zeta)) = 0$ and $w = v_2\, w$ q.e. and thus $\tilde{\xi}_\psi$-a.e. on $A_\psi(\zeta)$, see Theorem 3.4. Similarly, the second equation holds since $v_2\, w = 0$ $\tilde{\xi}^\varphi$-a.e. on $\Omega$.

Let $w \in H_0^1(\Omega)$. Now we have $\max(-n, \min(w, n)) \in H_0^1(\Omega) \cap L^\infty(\Omega)$ and $w_n \to w$ in $H_0^1(\Omega)$. Furthermore,

$$\|w_n - w_m\|_{L^1(\tilde{\xi}_\psi)} = \int_\Omega |w_n - w_m| \, \mathrm{d}\tilde{\xi}_\psi = \langle \xi_\psi^2, |w_n - w_m| \rangle \leq \|\xi_\psi^2\|_{H^{-1}} \| |w_n - w_m| \|_{H_0^1}.$$

Thus, $(w_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $L^1(\tilde{\xi}_\psi)$. Since $w_n \to w$ pointwise q.e. and thus $\tilde{\xi}_\psi$-a.e., see Theorem 3.4, we have $w_n \to w$ in $L^1(\tilde{\xi}_\psi)$. Arguing for $\tilde{\xi}^\varphi$ in a similar fashion, we obtain that

$$\langle \xi, w \rangle = \langle \xi_\psi^2, w \rangle - \langle \xi_2^\varphi, w \rangle = \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi. \qquad \square$$

The following example shows that, in general, i.e., when the active sets do not have a positive distance, the characterization of the functional $LS_{\mathrm{id}}(\zeta) - \zeta$ as the difference of the two measures $\tilde{\xi}_\psi$ and $\tilde{\xi}^\varphi$ does not need to apply for all possible arguments in $H_0^1(\Omega)$. A related example which is independent from the connection to the bilateral obstacle problem can be found in [Wac18, App. 2].

**Example 3.6.** For $d = 2$ and for $0 < \beta < \frac{1}{2}$, consider the function

$$y(x) = \begin{cases} \sin((-\ln(|x|))^\beta), & x \in \Omega := B_\rho(0) \setminus \{0\} \\ 0, & x = 0 \end{cases}, \quad \rho = \exp(-\pi^{1/\beta}) < 1. \tag{8}$$

Then $y|_{\partial B_\rho(0)} = \sin(\pi) = 0$ and $y \in H_0^1(\Omega) \cap L^\infty(\Omega)$, since $|y| \leq 1$ as well as

$$|\nabla y(x)|^2 = \frac{\cos^2((-\ln(|x|))^\beta) \beta^2 (-\ln(|x|))^{2\beta - 2}}{|x|^2} \frac{x_1^2 + x_2^2}{|x|^2} \leq \beta^2 (-\ln(|x|))^{2\beta - 2} |x|^{-2}$$

and thus

$$\|y\|^2_{H^1_0(\Omega)} = \|\nabla y\|^2_{L^2(\Omega)} \le 2\pi\beta^2 \int_0^\rho (-\ln(r))^{2\beta-2} r^{-2} r \, \mathrm{d}r = 2\pi\beta^2 \frac{(-\ln(r))^{2\beta-1}}{1-2\beta}\Big|_0^\rho < \infty,$$

since $\beta < \frac{1}{2}$. Now, we consider the obstacles given by $\psi(x) := \min\left(-\frac{1}{2}, y(x)\right)$, $\varphi(x) := \max\left(\frac{1}{2}, y(x)\right)$.

We have

$$(-\ln(r(t)))^\beta = t \quad \Leftrightarrow \quad r(t) = \exp(-t^{1/\beta})$$

and, for $k \in \mathbb{N}$, we set $r_k^\pm := r(2k\pi \pm \pi/2)$. This choice implies $\rho = r(\pi) > r_1^- > r_1^+ > r_2^- > r_2^+ > \ldots > 0$ and $y(r_k^\pm(\cos t, \sin t)) = \pm 1$ for all $t \in (0, 2\pi)$.

Now, let $\omega_k > 0$ be weights (that will be adjusted below), with $\sum_{k=1}^\infty \omega_k^2 < \infty$ and consider the functional

$$\langle \xi, w\rangle := \sum_{k=1}^\infty \frac{\omega_k}{\sqrt{\ln(r_k^-/r_k^+)}} \int_0^{2\pi} (w(r_k^-(\cos t, \sin t)) - w(r_k^+(\cos t, \sin t))) \, \mathrm{d}t \tag{9}$$

for $w \in H^1_0(\Omega)$. Note that the integral in (9) is well-defined. To see this, observe first that the quasi-continuous Borel measurable representatives of $w$ are unique up to a set of capacity zero. Let now $E \subseteq \Omega$ be a Borel set of capacity zero. Then, for any radius $0 < R < \rho$, by [Hel75, Thm. 7.5], the surface measure $\sigma_R$ on the sphere $\partial B_R$ satisfies $\sigma_R(E \cap \partial B_R) = 0$.

We have $\xi \in H^{-1}(\Omega)$, since

$$\begin{aligned}
|\langle \xi, w\rangle| &\le \sum_{k=1}^\infty \frac{\omega_k}{\sqrt{\ln(r_k^-/r_k^+)}} \int_0^{2\pi} \int_{r_k^+}^{r_k^-} |\nabla w(s(\cos t, \sin t))| \sqrt{s} \frac{1}{\sqrt{s}} \, \mathrm{d}s \, \mathrm{d}t \\
&\le \sum_{k=1}^\infty \frac{\omega_k}{\sqrt{\ln(r_k^-/r_k^+)}} \|\nabla w\|_{L^2(B_{r_k^-}(0)\setminus B_{r_k^+}(0))} \left( \int_0^{2\pi} \int_{r_k^+}^{r_k^-} \frac{1}{s} \, \mathrm{d}s \, \mathrm{d}t \right)^{\frac{1}{2}} \\
&= \sqrt{2\pi} \sum_{k=1}^\infty \omega_k \|\nabla w\|_{L^2(B_{r_k^-}(0)\setminus B_{r_k^+}(0))} \\
&\le \sqrt{2\pi} \left( \sum_{k=1}^\infty \omega_k^2 \right)^{\frac{1}{2}} \left( \sum_{k=1}^\infty \|\nabla w\|^2_{L^2(B_{r_k^-}(0)\setminus B_{r_k^+}(0))} \right)^{\frac{1}{2}} \\
&\le \sqrt{2\pi} \left( \sum_{k=1}^\infty \omega_k^2 \right)^{\frac{1}{2}} \|\nabla w\|_{L^2(\Omega)}.
\end{aligned}$$

Here, we have used the Cauchy Schwarz inequality and the fact that, for all $k \in \mathbb{N}$, the sets $B_{r_k^-}(0) \setminus B_{r_k^+}(0)$ are disjoint subsets of $\Omega$.

For $w \in H^1_0(\Omega) \cap L^\infty(\Omega)$, we have

$$\begin{aligned}
\langle \xi, w\rangle &= \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi, \\
&= \int_\Omega w \, \mathrm{d} \left( \sum_{k=1}^\infty \frac{\omega_k}{\sqrt{\ln(r_k^-/r_k^+) r_k^-}} \sigma_{r_k^-} \right) - \int_\Omega w \, \mathrm{d} \left( \sum_{k=1}^\infty \frac{\omega_k}{\sqrt{\ln(r_k^-/r_k^+) r_k^+}} \sigma_{r_k^+} \right)
\end{aligned}$$

where $\tilde{\xi}_\psi, \tilde{\xi}^\varphi$ are nonnegative finite measures with support in $\{y = \psi\}$ and $\{y = \varphi\}$, respectively. In fact, to show that $\tilde{\xi}^\varphi$ is a finite measure, we observe that

$$\ln(r_k^-/r_k^+) = (2k\pi + \pi/2)^{1/\beta} - (2k\pi - \pi/2)^{1/\beta} \begin{cases} \ge (2k\pi - \pi/2)^{1/\beta - 1} \pi/\beta, \\ \le (2k\pi + \pi/2)^{1/\beta - 1} \pi/\beta. \end{cases} \tag{10}$$

Hence, for any $w \in C(\overline{\Omega})$ with $0 \le w \le 1$

$$0 \le \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi \le \sum_{k=1}^\infty \frac{\omega_k}{\sqrt{\ln(r_k^-/r_k^+)}} \int_0^{2\pi} 1 \, \mathrm{d}t \le 2\pi \left( \sum_{k=1}^\infty \omega_k^2 \right)^{\frac{1}{2}} \left( \sum_{k=1}^\infty \frac{1}{\ln(r_k^-/r_k^+)} \right)^{\frac{1}{2}}$$
$$\le 2\pi \left( \sum_{k=1}^\infty \omega_k^2 \right)^{\frac{1}{2}} \left( \sum_{k=1}^\infty \frac{1}{\pi/\beta(2k\pi - \pi/2)^{1/\beta - 1}} \right)^{\frac{1}{2}} \le C$$

with a constant $C > 0$, since $\beta < 1/2$.

Set $\zeta := Ly - \xi$. We argue that the function $y \in K_\psi^\varphi$ as defined in (8) satisfies the bilateral obstacle problem (1) for $f = \mathrm{id}$, i.e., $y = S_{\mathrm{id}}(\zeta)$. First note that $y \in K_\psi^\varphi \subseteq H_0^1(\Omega) \cap L^\infty(\Omega)$ by the choice of $y, \psi$ and $\varphi$. Now, let $z \in K_\psi^\varphi$ be arbitrary. Then $z - y \ge 0$ q.e. on $\{y = \psi\}$ and $z - y \le 0$ q.e. on $\{y = \varphi\}$. Additionally, we have

$$\mathrm{supp}(\tilde{\xi}_\psi) \setminus \{0\} = \bigcup_{k=1}^\infty \partial B_{r_k^-} \subseteq \{y = \psi\} \tag{11}$$

and

$$\mathrm{supp}(\tilde{\xi}^\varphi) \setminus \{0\} = \bigcup_{k=1}^\infty \partial B_{r_k^+} \subseteq \{y = \varphi\}. \tag{12}$$

This yields

$$\begin{aligned} \langle \xi, z - y \rangle &= \int_\Omega (z - y) \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega (z - y) \, \mathrm{d}\tilde{\xi}^\varphi \\ &= \int_{\{y=\psi\}} (z - y) \, \mathrm{d}\tilde{\xi}_\psi - \int_{\{y=\varphi\}} (z - y) \, \mathrm{d}\tilde{\xi}^\varphi \ge 0 \end{aligned}$$

and we obtain $y = S_{\mathrm{id}}(\zeta)$.

Note that by (11) and (12) we find

$$\mathrm{dist}(A_\psi(\zeta), A^\varphi(\zeta)) = \mathrm{dist}(\{y = \psi\}, \{y = \varphi\}) = 0$$

since $r_k^\pm \searrow 0$. Thus, Lemma 3.5 does not apply.

Now, consider the unbounded function $w(x) = (-\ln(|x|))^\beta - \pi$. Then since

$$|\nabla w(x)|^2 = \frac{\beta^2(-\ln(|x|))^{2\beta-2}}{|x|^2} \frac{x_1^2 + x_2^2}{|x|^2} = \beta^2(-\ln(|x|))^{2\beta-2}|x|^{-2},$$

we have $w \in H_0^1(\Omega)$ as above. With $\omega_k = k^{-1}$, $\beta = \frac{1}{3}$ and by using (10), we obtain the estimate

$$\begin{aligned} \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi &= \sum_{k=1}^\infty \frac{\omega_k}{\sqrt{\ln(r_k^-/r_k^+)}} \int_0^{2\pi} (2k\pi + \tfrac{\pi}{2} - \pi) \, \mathrm{d}t \\ &\ge \sum_{k=1}^\infty \frac{2\pi(2k\pi - \frac{\pi}{2})}{k(2k\pi + \frac{\pi}{2})\sqrt{3\pi}} \ge \sum_{k=1}^\infty \frac{2\pi \, \frac{3}{2}k\pi}{k \, \frac{5}{2}k\pi \, \sqrt{3\pi}} = \infty. \end{aligned}$$

Similarly, we find

$$\int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi \ge \sum_{k=1}^\infty \frac{2\pi \, \frac{1}{2}k\pi}{k \, \frac{5}{2}k\pi \, \sqrt{3\pi}} = \infty.$$

This shows that, in general, the representation (6) does not hold for all $w \in H_0^1(\Omega)$.

In the following lemma, we find a characterization of the critical cone. In parts, the proof is based on the proof of [Wac14, Lem. 3.1].

**Lemma 3.7**
*Assume that $\psi, \varphi$ fulfill the conditions of Assumption 1.1. Let $\zeta \in H^{-1}(\Omega)$ be arbitrary and set $y := S_{\mathrm{id}}(\zeta)$, $\xi := Ly - \zeta$. Then the critical cone*

$$\mathcal{K}_{K_\psi^\varphi}(y, \xi) := T_{K_\psi^\varphi}(y) \cap \xi^\perp$$

*has the following structure. There exist quasi-closed sets $A_\psi^{\mathrm{s}}(\zeta) \subseteq A_\psi(\zeta)$ and $A_{\mathrm{s}}^\varphi(\zeta) \subseteq A^\varphi(\zeta)$ which are unique up to sets of capacity zero such that*

$$\begin{aligned}
\mathcal{K}_{K_\psi^\varphi}(y,\xi) &= \{z \in H_0^1(\Omega) \mid z \geq 0 \text{ q.e. in } A_\psi(\zeta), z \leq 0 \text{ q.e. in } A^\varphi(\zeta) \text{ and } \langle \xi, z\rangle = 0\} \\
&= \left\{z \in H_0^1(\Omega) \mid z \geq 0 \text{ q.e. in } A_\psi(\zeta), z \leq 0 \text{ q.e. in } A^\varphi(\zeta), z = 0 \ \tilde{\xi}_\psi\text{- and } \tilde{\xi}^\varphi\text{-a.e.}\right\} \\
&= \big\{z \in H_0^1(\Omega) \mid z \geq 0 \text{ q.e. in } A_\psi(\zeta), z \leq 0 \text{ q.e. in } A^\varphi(\zeta), \\
&\qquad\qquad z = 0 \text{ q.e. in } A_\psi^{\mathrm{s}}(\zeta) \cup A_{\mathrm{s}}^\varphi(\zeta)\big\},
\end{aligned} \tag{13}$$

*where $\tilde{\xi}_\psi, \tilde{\xi}^\varphi$ denote the decomposition of $\xi$ according to Theorem 3.4.*

*Proof.* Recalling (5), we see that the first equation in (13) holds.

We show the second identity in (13). Assume $z$ is an element of $\mathcal{K}_{K_\psi^\varphi}(y, \xi)$. By polyhedricity, see e.g. [Mig76, Thm. 3.2], we have

$$T_{K_\psi^\varphi}(y) \cap \xi^\perp = \overline{R_{K_\psi^\varphi}(y) \cap \xi^\perp},$$

where $R_{K_\psi^\varphi}(y) := \{z \in H_0^1(\Omega) \mid \exists t > 0, y + tz \in K_\psi^\varphi\}$ denotes the radial cone. Thus, there is a sequence $(z_n)_{n\in\mathbb{N}} \subseteq R_{K_\psi^\varphi}(y) \cap \xi^\perp$ with $z_n \to z$ in $H_0^1(\Omega)$. Since $\psi, \varphi$ are elements of $L^\infty(\Omega)$, and by the structure of $R_{K_\psi^\varphi}(y)$ we conclude $(z_n)_{n\in\mathbb{N}} \subseteq L^\infty(\Omega)$. Using Theorem 3.4, each $z_n$ is integrable with respect to $\tilde{\xi}_\psi$ and $\tilde{\xi}^\varphi$. Let $n \in \mathbb{N}$ be fixed. By $R_{K_\psi^\varphi}(y) \subseteq T_{K_\psi^\varphi}(y)$ we have $z_n \geq 0$ q.e. in $A_\psi(\zeta)$ and $z_n \leq 0$ q.e. in $A^\varphi(\zeta)$. This implies

$$0 = \langle \xi, z_n\rangle = \int_\Omega z_n \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega z_n \, \mathrm{d}\tilde{\xi}^\varphi = \int_{A_\psi(\zeta)} z_n \, \mathrm{d}\tilde{\xi}_\psi - \int_{A^\varphi(\zeta)} z_n \, \mathrm{d}\tilde{\xi}^\varphi, \tag{14}$$

as $\tilde{\xi}_\psi(I_\psi(\zeta))) = 0$ and $\tilde{\xi}^\varphi(I^\varphi(\zeta)) = 0$, see Theorem 3.4. Since $z_n \geq 0$ $\tilde{\xi}_\psi$-a.e. on $A_\psi(\zeta)$ and $z_n \leq 0$ $\tilde{\xi}^\varphi$-a.e. on $A^\varphi(\zeta)$, cf. Theorem 3.4, we conclude that $z_n = 0$ $\tilde{\xi}_\psi$-a.e. on $A_\psi(\zeta)$ and $z_n = 0$ $\tilde{\xi}^\varphi$-a.e. on $A^\varphi(\zeta)$. Using once more that $\tilde{\xi}_\psi(I_\psi(\zeta)) = 0$ and $\tilde{\xi}^\varphi(I^\varphi(\zeta)) = 0$, we can see that this means $z_n = 0$ $\tilde{\xi}_\psi$- and $\tilde{\xi}^\varphi$-a.e. on $\Omega$. Since $z_n \to z$ for a subsequence pointwise q.e. and thus $\tilde{\xi}_\psi$- and $\tilde{\xi}^\varphi$-a.e., see Theorem 3.4, we conclude $z = 0$ $\tilde{\xi}_\psi$- and $\tilde{\xi}^\varphi$-a.e.

Vice versa, assume $z \in T_{K_\psi^\varphi}(y)$ and $z = 0$ $\tilde{\xi}_\psi$- and $\tilde{\xi}^\varphi$-a.e. Using Theorem 3.4, we see that (14) holds (with $z_n$ replaced by $z$) and thus $z \in T_{K_\psi^\varphi}(y) \cap \xi^\perp$ follows.

Thus, we have shown the second equation in (13).

To show the last identity in (13) we note that by [Sto93, Thm. 1], there exist quasi-closed sets $A_\psi^{\mathrm{s}}(\zeta)$ and $A_{\mathrm{s}}^\varphi(\zeta)$ such that

$$\{z \in H_0^1(\Omega) \mid z = 0 \ \tilde{\xi}_\psi\text{-a.e.}\} = \{z \in H_0^1(\Omega) \mid z = 0 \text{ q.e. on } A_\psi^{\mathrm{s}}(\zeta)\} \tag{15}$$

and

$$\{z \in H_0^1(\Omega) \mid z = 0 \ \tilde{\xi}^\varphi\text{-a.e.}\} = \{z \in H_0^1(\Omega) \mid z = 0 \text{ q.e. on } A_{\mathrm{s}}^\varphi(\zeta)\}. \tag{16}$$

We have $y - \psi = 0$ $\tilde{\xi}_\psi$-a.e. and thus $y - \psi = 0$ q.e. on $A_\psi^{\mathrm{s}}(\zeta)$, see (15), which implies $A_\psi^{\mathrm{s}}(\zeta) \subseteq A_\psi(\zeta)$ up to a set of capacity zero. The same arguments apply to show $A_{\mathrm{s}}^\varphi(\zeta) \subseteq A^\varphi(\zeta)$ up to a set of capacity zero. ∎

**Corollary 3.8**
*Assume that $\psi, \varphi$ fulfill the conditions of Assumption 1.1. Let $\zeta \in H^{-1}(\Omega)$ be arbitrary and set $y := S_{\mathrm{id}}(\zeta)$, $\xi := Ly - \zeta$. For $w \in H_0^1(\Omega)$ it holds $w = 0$ q.e. on $A_\psi^{\mathrm{s}}(\zeta)$, respectively $w = 0$ q.e. on $A_{\mathrm{s}}^\varphi(\zeta)$ if and only if $w = 0$ $\tilde{\xi}_\psi$-a.e., respectively $w = 0$ $\tilde{\xi}^\varphi$-a.e. In both cases we have $w \in L^1(\tilde{\xi}_\psi) \cap L^1(\tilde{\xi}^\varphi)$ and $\langle \xi, w \rangle = \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi$.*

*Proof.* The equivalence is implied by the proof of Lemma 3.7, see (15) and (16). The statements $w \in L^1(\tilde{\xi}_\psi) \cap L^1(\tilde{\xi}^\varphi)$ and $\xi, w\rangle = \int_\Omega w \, \mathrm{d}\tilde{\xi}_\psi - \int_\Omega w \, \mathrm{d}\tilde{\xi}^\varphi$ follow from Theorem 3.4. □

In the following sections, we also write $A_{\mathrm{s}}(\zeta) := A_\psi^{\mathrm{s}}(\zeta) \cup A_{\mathrm{s}}^\varphi(\zeta)$ for the strictly active set with respect to both obstacles, we have $A_{\mathrm{s}}(\zeta) \subseteq A(\zeta)$.

Moreover, we will use the notation $A_\psi^{\mathrm{w}}(\zeta) := A_\psi(\zeta) \setminus A_\psi^{\mathrm{s}}(\zeta)$ for the weakly active set with respect to the lower obstacle $\psi$ and $A_{\mathrm{w}}^\varphi(\zeta) := A^\varphi(\zeta) \setminus A_{\mathrm{s}}^\varphi(\zeta)$ for the weakly active set with respect to the upper obstacle $\varphi$. For the sake of completeness, we also introduce the notation $A_{\mathrm{w}}(\zeta) := A_\psi^{\mathrm{w}}(\zeta) \cup A_{\mathrm{w}}^\varphi(\zeta)$ for the weakly active set with respect to upper and lower obstacle.

### 3.2.2 Gâteaux Differentiability of the Solution Operator

As in the case of unilateral obstacle problems, in points $u$ at which $S_f$ is Gâteaux differentiable, we can replace the critical cone in the characterization of the directional derivative by the largest linear subset contained in the critical cone, and by the linear hull of the critical cone, respectively. Both versions yield a characterization of the Gâteaux derivative. The reasoning for these facts in the case of the unilateral obstacle problem can be found in [RU19, Lem. 3.7].

For the bilateral case, characterizations of the Gâteaux derivative are summarized in the following theorem.

**Theorem 3.9**
*Assume the obstacles $\psi, \varphi$ satisfy Assumption 1.1. Suppose $S_f$ is Gâteaux differentiable at $u \in U$. Let $h \in U$. Then $S_f'(u; h)$ is determined by the solution of the variational equation*

$$\text{Find } \eta \in H_0^1(D): \quad \langle L\eta - f'(u; h), z\rangle = 0 \quad \forall\, z \in H_0^1(D) \tag{17}$$

*and $D$ can be chosen as any quasi-open subset of $\Omega$ fulfilling $I(u) \subseteq D \subseteq \Omega \setminus A_{\mathrm{s}}(u)$.*

*Proof.* Assume $S_f$ is Gâteaux differentiable at $u \in U$. Then the map $S_f'(u; \cdot)$ is linear and the image is a linear subspace of $H_0^1(\Omega)$. By the characterization in (4), the image of $S_f'(u; \cdot)$ lies in a linear subspace of the critical cone $(LS_f(u) - f(u))^\perp \cap T_{K_\psi^\varphi}(S_f(u))$. The structure of the critical cone established in Lemma 3.7 implies that $S_f'(u; h) \in H_0^1(I(f(u)))$, since $H_0^1(I(f(u)))$ is the largest linear subset contained in the critical cone. Now $S_f'(u; h)$ solves the variational equation (17) with $D = I(f(u))$, since $H_0^1(I(f(u)))$ is a linear subspace.

Obviously, the image of $S_f'(u; \cdot)$ is also contained in the linear hull of the critical cone, the set $H_0^1(\Omega \setminus A_{\mathrm{s}}(f(u)))$. Assume $z$ is in the critical cone and $h \in U$ is arbitrary. Then

$$\langle LS_f'(u; -h) - f'(u; -h), z - S_f'(u; -h)\rangle \geq 0,$$

which implies

$$\langle LS_f'(u; h) - f'(u; h), -z - S_f'(u; h)\rangle \geq 0.$$

Thus, by linearity arguments we obtain that we can also use test functions from the negative critical cone. Let $z \in H_0^1(\Omega \setminus A_{\mathrm{s}}(f(u)))$ be arbitrary. Since the two

sets $\Omega \setminus (A_{\mathrm{s}}(f(u)) \cup A_\psi(f(u)))$ and $\Omega \setminus (A_{\mathrm{s}}(f(u)) \cup A^\varphi(f(u)))$ are a quasi-covering of $\Omega \setminus A_{\mathrm{s}}(f(u))$, we can find a sequence $(z_\psi^n + z_n^\varphi)_{n\in\mathbb{N}}$ converging to $z$ and fulfilling $z_\psi^n \in H_0^1(\Omega \setminus (A_{\mathrm{s}}(f(u)) \cup A^\varphi(f(u))))$ and $z_n^\varphi \in H_0^1(\Omega \setminus (A_{\mathrm{s}}(f(u)) \cup A_\psi(f(u))))$, see Lemma 2.3. Considering positive and negative parts we write

$$z_n^\varphi = z_n^{\varphi,+} - z_n^{\varphi,-} \quad \text{and } z_\psi^n = z_{\psi,+}^n - z_{\psi,-}^n.$$

The representation in (13) implies that $z_{\psi,+}^n$, $z_{\psi,-}^n$, $-z_n^{\varphi,+}$ and $-z_n^{\varphi,-}$ are elements of the critical cone. This shows

$$\langle LS_f'(u;h) - f'(u;h), z_\psi^n + z_n^\varphi - S_f'(u;h)\rangle \geq 0$$

for all $n \in \mathbb{N}$. Taking the limit and observing that $H_0^1(\Omega\setminus A_{\mathrm{s}}(f(u)))$ is a linear subspace we obtain

$$\langle LS_f'(u;h) - f'(u;h), z\rangle = 0$$

for all $z \in H_0^1(\Omega \setminus A_{\mathrm{s}}(f(u)))$.

Thus, since (17) is a characterization of the Gâteaux derivative for $D = I(f(u))$ and for $D = \Omega\setminus A_{\mathrm{s}}(f(u))$, each set $H_0^1(D)$ with $I(f(u)) \subseteq D \subseteq \Omega\setminus A_{\mathrm{s}}(f(u))$ also yields a characterization for the Gâteaux derivative. □

# 4 Monotonicity of the Active and Strictly Active Sets

In this subsection, the monotonicity of the active and strictly active sets is studied. Within this section, we specify our notation and write $S_{\psi,f}^\varphi$ instead of $S_f$ for the solution operator of (1).

The monotonicity of the active sets is a direct consequence of Lemma 3.1.

**Lemma 4.1**
*Let $u_1, u_2 \in U$ with $u_1 \geq_U u_2$. Then*

1. $A_\psi(f(u_1)) \subseteq A_\psi(f(u_2))$,
2. $A^\varphi(f(u_1)) \supseteq A^\varphi(f(u_2))$.

*Proof.* By Lemma 3.1, we have $S_{\psi,f}^\varphi(u_1) \geq S_{\psi,f}^\varphi(u_2)$ a.e. and by Lemma 2.3 also q.e. in $\Omega$. This implies the statements. □

**Lemma 4.2**
*Suppose the conditions of Assumption 1.1 are satisfied. Let $u \in U$ and let $v \in H_0^1(\Omega)_+$ such that $\{v > 0\} \subseteq \Omega \setminus A_\psi^{\mathrm{s}}(f(u))$. Then $S_{\psi,f}^\varphi(u) = S_{\psi-v,f}^\varphi(u)$.*

*Proof.* Obviously, $S_{\psi,f}^\varphi(u) \geq \psi - v$. Now, let $z \in K_{\psi-v}^\varphi$ be arbitrary. We need to show that

$$\langle LS_{\psi,f}^\varphi(u) - f(u), z - S_{\psi,f}^\varphi(u)\rangle \geq 0.$$

Now, $z = \max(z,\psi) + \min(z-\psi, 0) =: z_1 + z_2$, where $z_1 \in K_\psi^\varphi$ and $z_2 \in H_0^1(\Omega \setminus A_\psi^{\mathrm{s}}(f(u)))_-$. Thus

$$\langle LS_{\psi,f}^\varphi(u) - f(u), z_1 - S_{\psi,f}^\varphi(u)\rangle \geq 0$$

and moreover, since $z_2 = 0$ q.e. on $A_\psi^{\mathrm{s}}(f(u))$ and thus $\tilde\xi_\psi$-a.e., see Theorem 3.4, we have by Corollary 3.8

$$\langle LS_{\psi,f}^\varphi(u) - f(u), z_2\rangle = \int_{A_\psi} z_2 \,\mathrm{d}\tilde\xi_\psi - \int_{A^\varphi} z_2 \,\mathrm{d}\tilde\xi^\varphi = -\int_{A^\varphi} z_2 \,\mathrm{d}\tilde\xi^\varphi \geq 0.$$

Here, $\tilde\xi_\psi$ and $\tilde\xi^\varphi$ are the measures as in Theorem 3.4 that belong to the functional $LS_{\psi,f}^\varphi(u) - f(u) = LS_{\psi,\mathrm{id}}^\varphi(f(u)) - f(u)$. □

**Lemma 4.3**
*Let $\zeta \in H^{-1}(\Omega)$. Then we have $-S^{\varphi}_{\psi,\mathrm{id}}(\zeta) = S^{-\psi}_{-\varphi,\mathrm{id}}(-\zeta)$. Moreover, $A_\psi(\zeta) = \tilde{A}^{-\psi}(-\zeta)$ and $A^\varphi(\zeta) = \tilde{A}_{-\varphi}(-\zeta)$. Here, $\tilde{A}_{-\varphi}(\zeta) := \{\omega \in \Omega \mid S^{-\psi}_{-\varphi,\mathrm{id}}(\zeta)(\omega) = -\varphi(\omega)\}$ and $\tilde{A}^{-\psi}(\zeta) := \{\omega \in \Omega \mid S^{-\psi}_{-\varphi,\mathrm{id}}(\zeta)(\omega) = -\psi(\omega)\}$ denote the respective active sets for $S^{-\psi}_{-\varphi,\mathrm{id}}(\zeta)$.*
*Furthermore, if the conditions of Assumption 1.1 are fulfilled, we have $A^{\mathrm{s}}_\psi(\zeta) = \tilde{A}^{-\psi}_{\mathrm{s}}(-\zeta)$ and $A^{\varphi}_{\mathrm{s}}(\zeta) = \tilde{A}^{\mathrm{s}}_{-\varphi}(-\zeta)$. Here, $\tilde{A}^{-\psi}_{\mathrm{s}}(\zeta), \tilde{A}^{\mathrm{s}}_{-\varphi}(\zeta)$ denote the strictly active sets for $S^{-\psi}_{-\varphi,\mathrm{id}}(\zeta)$.*

*Proof.* First, let us note that for $z \in H^1_0(\Omega)$ it holds $\psi \le z \le \varphi$ if and only if $-\varphi \le -z \le -\psi$ and this implies

$$-K^{\varphi}_{\psi} = K^{-\psi}_{-\varphi}.$$

Thus, $-S^{\varphi}_{\psi,\mathrm{id}}(\zeta) \in K^{-\psi}_{-\varphi}$. Let now $z \in K^{-\psi}_{-\varphi}$ be arbitrary. Then we have

$$\begin{aligned}
&\langle L(-S^{\varphi}_{\psi,\mathrm{id}}(\zeta)) + \zeta, z - (-S^{\varphi}_{\psi,\mathrm{id}}(\zeta))\rangle \\
&= -\langle LS^{\varphi}_{\psi,\mathrm{id}}(\zeta) - \zeta, z - (-S^{\varphi}_{\psi,\mathrm{id}}(\zeta))\rangle \\
&= \langle LS^{\varphi}_{\psi,\mathrm{id}}(\zeta) - \zeta, -z - S^{\varphi}_{\psi,\mathrm{id}}(\zeta)\rangle \\
&\ge 0.
\end{aligned}$$

This yields $-S^{\varphi}_{\psi,\mathrm{id}}(\zeta) = S^{-\psi}_{-\varphi,\mathrm{id}}(-\zeta)$.

Now, $S^{-\psi}_{-\varphi,\mathrm{id}}(-\zeta)(\omega) = -\psi(\omega)$ if and only if $S^{\varphi}_{\psi,\mathrm{id}}(\zeta)(\omega) = \psi(\omega)$ and we have $S^{-\psi}_{-\varphi,\mathrm{id}}(-\zeta)(\omega) = -\varphi(\omega)$ if and only if $S^{\varphi}_{\psi,\mathrm{id}}(\zeta)(\omega) = \varphi(\omega)$, thus $A_\psi(\zeta) = \tilde{A}^{-\psi}(-\zeta)$ and $A^\varphi(\zeta) = \tilde{A}_{-\varphi}(-\zeta)$.

In addition, we have

$$LS^{\varphi}_{\psi,\mathrm{id}}(\zeta) - \zeta = -(LS^{-\psi}_{-\varphi,\mathrm{id}}(-\zeta) - (-\zeta)).$$

This shows the statements for the strictly active sets. □

**Remark 4.4.** If $f\colon U \to H^{-1}(\Omega)$ satisfies $f(-u) = -f(u)$ for all $u \in U$, then $-S^{\varphi}_{\psi,f}(u) = S^{-\varphi}_{-\psi,f}(-u)$.

Now, we check the monotonicity of the strictly active sets.

**Lemma 4.5**
*Let the requirements of Assumption 1.1 be satisfied and let $u_1 \ge_U u_2$. Then it follows*

1. $A^{\mathrm{s}}_\psi(f(u_1)) \subseteq A^{\mathrm{s}}_\psi(f(u_2))$,
2. $A^{\varphi}_{\mathrm{s}}(f(u_1)) \supseteq A^{\varphi}_{\mathrm{s}}(f(u_2))$.

*Proof.* ad 1.: Let

$$\mathcal{U} = \{S^{\varphi}_{\psi,f}(u_1) - \psi < (\varphi - \psi)/2\}. \tag{18}$$

Then $\mathcal{U}$ is quasi-open and $A^{\mathrm{s}}_\psi(f(u_1)) \subseteq A_\psi(f(u_1)) \subseteq \mathcal{U} \subseteq I^\varphi(f(u_1))$ holds.

Assume $\mathcal{U} \setminus A^{\mathrm{s}}_\psi(f(u_2)) \neq \emptyset$ (otherwise the assertion follows directly). Fix $v \in H^1_0(\mathcal{U})_+$ satisfying $\{v > 0\} = \mathcal{U} \setminus A^{\mathrm{s}}_\psi(f(u_2))$, $v < (\varphi - \psi)/2$, see Lemma 2.3.

Let $y_v(t) = S^{\varphi}_{\psi - tv,f}(u_1)$, $t \in [0,1]$, and denote $\bar{y}_v(t) := y_v(t) + tv$. Then

$$\begin{aligned}
&\langle Ly_v(t) - f(u_1), z - y_v(t)\rangle \ge 0 \quad \forall\, z \in K^{\varphi}_{\psi - tv} \\
\iff &\langle L\bar{y}_v(t) - f(u_1) - tLv, \bar{z} - \bar{y}_v(t)\rangle \ge 0 \quad \forall\, \bar{z} \in K^{\varphi + tv}_{\psi} \\
\implies &\langle L\bar{y}_v(t) - f(u_1) - tLv, \bar{z} - \bar{y}_v(t)\rangle \ge 0 \quad \forall\, \bar{z} \in K^{\varphi}_{\psi}.
\end{aligned}$$

Now $\psi \leq \bar{y}_v(t) = y_v(t) + tv \leq y_v(0) + tv \leq \varphi$ by Lemma 3.2 and the definition of $\mathcal{U}$ and $v$.

Hence, $\bar{y}_v(t) \in K_\psi^\varphi$ and the last line shows that $y_v(t) = S^\varphi_{\psi,\mathrm{id}}(T(tv)) - tv$ with $T\colon H_0^1(\Omega) \to H^{-1}(\Omega), v \mapsto f(u_1) + Lv$. Since $S^\varphi_{\psi,\mathrm{id}}$ is directionally differentiable in the Hadamard sense, we can apply the chain rule for the directional derivatives and obtain

$$y_v'(0;1) = (S^\varphi_{\psi,\mathrm{id}})'(T(0);T'(0;v)) - v = (S^\varphi_{\psi,\mathrm{id}})'(f(u_1);Lv) - v.$$

Since $(S^\varphi_{\psi,\mathrm{id}})'(f(u_1);Lv)$ is 0 q.e. on the strictly active set $A_{\mathrm{s}}(f(u_1))$, compare Section 3.2 and, in particular, Lemma 3.7, we have $y_v'(0;1) < 0$ q.e. on $A_{\mathrm{s}}(f(u_1)) \cap \{v > 0\}$.

Thus, by reducing the lower obstacle on a subset of $A^{\mathrm{s}}_\psi(f(u_1))$ the solution with respect to the new obstacle will drop on this set.

Now, we show the statement of the lemma by contradiction. Therefore, assume the set $W \subseteq \Omega$ is a set of positive capacity which is (lower) weakly active for $u_2$ and (lower) strictly active for $u_1$, more precisely, $W \subseteq A^{\mathrm{s}}_\psi(f(u_1)) \subseteq A_\psi(f(u_1)) \subseteq A_\psi(f(u_2))$ and $W \subseteq \Omega \setminus A^{\mathrm{s}}_\psi(f(u_2))$ (up to a set of capacity zero). Then $\mathcal{U}$ as in (18) is a quasi-open neighborhood of $W$ contained in $I^\varphi(f(u_1))$.

Let as above $v \in H_0^1(\mathcal{U})_+$ satisfy $\{v > 0\} = \mathcal{U} \setminus A^{\mathrm{s}}_\psi(f(u_2))$. Then, Lemma 4.2 yields

$$S^\varphi_{\psi-v,f}(u_2) = S^\varphi_{\psi,f}(u_2) \tag{19}$$

and on $W$ we have

$$S^\varphi_{\psi-v,f}(u_1)|_W < S^\varphi_{\psi,f}(u_1)|_W = S^\varphi_{\psi,f}(u_2)|_W \tag{20}$$

by the structure of the directional derivative with respect to the obstacle. Putting Eqs. (19) and (20) together, we see that

$$S^\varphi_{\psi-v,f}(u_2) > S^\varphi_{\psi-v,f}(u_1)$$

on $W$. On the other hand, $S^\varphi_{\psi-v,f}(u_1) \geq S^\varphi_{\psi-v,f}(u_2)$ since $u_1 \geq_U u_2$. Thus, such a set $W$ cannot exist and we conclude $A^{\mathrm{s}}_\psi(u_1) \subseteq A^{\mathrm{s}}_\psi(u_2)$.

ad 2.: By Lemma 4.3, we have $A^\varphi_{\mathrm{s}}(f(u_i)) = \tilde{A}^{\mathrm{s}}_{-\varphi}(-f(u_i))$ for $i = 1, 2$, where we use a similar notation as in Lemma 4.3. Now, the first part of the lemma implies the statement, since

$$A^\varphi_{\mathrm{s}}(f(u_1)) = \tilde{A}^{\mathrm{s}}_{-\varphi}(-f(u_1)) \supseteq \tilde{A}^{\mathrm{s}}_{-\varphi}(-(f(u_2)) = A^\varphi_{\mathrm{s}}(f(u_2)). \qquad \square$$

# 5 Mosco Convergence

For the rest of the paper, we use again the notation $S_f$ for the solution operator of (1).

The following definition goes back to [Mos69]. In this form, the definition can be found, e.g., in [Rod87, Ch. 4:4].

**Definition 5.1.** We say that a sequence $(C_n)_{n\in\mathbb{N}}$ of nonempty, closed, convex subsets of a Banach space $X$ converges to a set $C \subseteq X$ in the sense of Mosco if the following two conditions hold.

1. For each $x \in C$ there exists a sequence $(x_n)_{n\in\mathbb{N}}$ such that $x_n \in C_n$ holds for every $n \in \mathbb{N}$ and such that $x_n \to x$ in $X$.

2. For each subsequence $(x_{n_k})_{k\in\mathbb{N}}$ of a sequence $(x_n)_{n\in\mathbb{N}} \subseteq X$ fulfilling $x_n \in C_n$ for all $n \in \mathbb{N}$ such that for some $x \in X$ we have $x_{n_k} \rightharpoonup x$ in $X$, the weak limit $x$ is in $C$.

Based on this definition, the following result on convergence of solutions of variational inequalities can be established. It is taken from [Rod87, Thm. 4.1], see also [Mos69, Prop. 35.].

**Lemma 5.2**
*Assume $L \in \mathcal{L}(H_0^1(\Omega), H^{-1}(\Omega))$ is coercive and let $C_n$ and $C$ be nonempty closed, convex subsets of $H_0^1(\Omega)$, $n \in \mathbb{N}$, such that $C_n \to C$ in the sense of Mosco. Furthermore, let $(h_n)_{n\in\mathbb{N}}, h \subseteq H^{-1}(\Omega)$ with $h_n \to h$. Then the solutions of*

$$\text{Find } \eta_n \in C_n: \quad \langle L\eta_n - h_n, z - \eta_n\rangle \geq 0 \quad \forall\, z \in C_n$$

*converge strongly in $H_0^1(\Omega)$ to the solution of*

$$\text{Find } \eta \in C: \quad \langle L\eta - h, z - \eta\rangle \geq 0 \quad \forall\, z \in C.$$

Based on this tool, in order to obtain the convergence of the Gâteaux derivatives, which can be characterized as solutions to variational equations, see Theorem 3.9, and in order to characterize the limit, we establish the Mosco convergence of the sets $H_0^1(D_n)$. Depending on either the choice $D_n = I(f(u_n)) \cup A_{\mathrm{w}}^{\varphi}(f(u_n))$ or $D_n = I(f(u_n)) \cup A_{\psi}^{\mathrm{w}}(f(u_n))$, i.e., depending on whether we focus on the inactive set or the complement of the strictly active set with respect to either the upper or the lower obstacle, sequences with different monotone behavior have to be considered.

**Lemma 5.3**
*Suppose Assumption 1.1 is satisfied.*

1. *Let $(u_n)_{n\in\mathbb{N}} \subseteq U$ be an increasing sequence with $u_n \to u$. Then $H_0^1(I(f(u_n)) \cup A_{\mathrm{w}}^{\varphi}(f(u_n))) \to H_0^1(I(f(u)) \cup A_{\mathrm{w}}^{\varphi}(f(u)))$ in the sense of Mosco.*
2. *Let $(u_n)_{n\in\mathbb{N}} \subseteq U$ be a decreasing sequence with $u_n \to u$. Then $H_0^1(I(f(u_n)) \cup A_{\psi}^{\mathrm{w}}(f(u_n))) \to H_0^1(I(f(u)) \cup A_{\psi}^{\mathrm{w}}(f(u)))$ in the sense of Mosco.*

*Proof.* ad 1.: Assume $v \in H_0^1(I(f(u)) \cup A_{\mathrm{w}}^{\varphi}(f(u)))$ and w.l.o.g. $v \geq 0$. We can rewrite the function space as

$$\begin{aligned} &H_0^1(I(f(u)) \cup A_{\mathrm{w}}^{\varphi}(f(u))) \\ &= \{z \in H_0^1(\Omega) \mid z = 0 \text{ q.e. on } A_{\psi}(f(u)) \text{ and } z = 0 \text{ q.e. on } A_{\mathrm{s}}^{\varphi}(f(u))\}. \end{aligned}$$

Since $A_{\mathrm{s}}^{\varphi}(f(u_n)) \subseteq A_{\mathrm{s}}^{\varphi}(f(u))$ for all $n \in \mathbb{N}$, see Lemma 4.5, it holds $v = 0$ q.e. on $A_{\mathrm{s}}^{\varphi}(f(u_n))$ for all $n \in \mathbb{N}$. Since $S_f(u_n) \to S_f(u)$ in $H_0^1(\Omega)$ by continuity of $S_f$, we have $S_f(u_n) \to S_f(u)$ for a subsequence pointwise quasi everywhere, see Lemma 2.3. This means

$$\operatorname{cap}\left(I_{\psi}(f(u)) \setminus \bigcup_{k\in\mathbb{N}} I_{\psi}(f(u_k))\right) = 0,$$

i.e., $(I_{\psi}(f(u_n)))_{n\in\mathbb{N}}$ is a quasi-covering of $I_{\psi}(f(u))$, which is increasing in $n$. We can therefore find nonnegative $v_n \in H_0^1(\Omega)$ with $v_n \to v$ and $v_n = 0$ q.e. on $A_{\psi}(f(u_n))$, see Lemma 2.3. By setting

$$z_n := \min(v_n, v)$$

we have $z_n \in H_0^1(I(f(u_n)) \cup A_{\mathrm{w}}^{\varphi}(f(u_n)))$ for all $n \in \mathbb{N}$ as well as $z_n \to v$.

Let $v_n \in H_0^1(I(f(u_n)) \cup A_{\mathrm{w}}^{\varphi}(f(u_n)))$ for all $n \in \mathbb{N}$. Assume there is a subsequence $(v_{n_k})_{k\in\mathbb{N}}$ with $v_{n_k} \rightharpoonup v$ for some $v \in H_0^1(\Omega)$ as $k \to \infty$. Since $A_{\psi}(f(u)) \subseteq A_{\psi}(f(u_n))$

for all $n \in \mathbb{N}$, we conclude $v \in H_0^1(I_\psi(f(u)))$ by Mazur's lemma. By Corollary 3.8 and Theorem 3.4, from $v_n = 0$ q.e. on $A_{\mathrm{s}}^\varphi(f(u_n))$ and $v_n = 0$ q.e. on $A_\psi(f(u_n))$ it follows

$$\langle LS_f(u_n) - f(u_n), |v_n| \rangle = \int_\Omega |v_n| \, \mathrm{d}\tilde{\xi}_\psi^n - \int_\Omega |v_n| \, \mathrm{d}\tilde{\xi}_n^\varphi = 0$$

for all $n \in \mathbb{N}$. Here, $\tilde{\xi}_\psi^n$ and $\tilde{\xi}_n^\varphi$ are the measures as in Theorem 3.4 that belong to the functional $LS_{\psi,f}^\varphi(u_n) - f(u_n) = LS_{\psi,\mathrm{id}}^\varphi(f(u_n)) - f(u_n)$. From $v_{n_k} \rightharpoonup v$ in $H_0^1(\Omega)$ we conclude $|v_{n_k}| \rightharpoonup |v|$ in $H_0^1(\Omega)$. (To see this, one can use the compact embedding $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$ and $\||z|\|_{H_0^1(\Omega)} = \|z\|_{H_0^1(\Omega)}$ for all $z \in H_0^1(\Omega)$, see [ABM14, Cor. 5.8.1].) Since also $LS_f(u_{n_k}) - f(u_{n_k}) \to LS_f(u) - f(u)$ in $H^{-1}(\Omega)$ we conclude

$$0 = \langle LS_f(u) - f(u), |v| \rangle = -\int_\Omega |v| \, \mathrm{d}\tilde{\xi}^\varphi$$

since $v = 0$ q.e. on $A_\psi(f(u))$. Finally, $v = 0$ $\tilde{\xi}^\varphi$-a.e. on $\Omega$ and thus $v = 0$ q.e. on $A_{\mathrm{s}}^\varphi(f(u))$, see Corollary 3.8.

ad 2.: Again, this part of the lemma follows from the first part of the lemma combined with Lemma 4.3. □

# 6 Generalized Derivatives for the Bilateral Obstacle Problem

In this section, we will find a characterization of two generalized derivatives for the solution operator $S_f$ of (1). To establish this result, we impose the monotonicity assumption Assumption 1.2 on $U$ and $f$ stated in the introduction.

As already indicated in the introduction, the assumptions posed on the positive cone in $V$ will ensure that we can construct monotone convergent sequences in $U$ where the Gâteaux differentiability of the locally Lipschitz continuous solution operator $S_f$ can be guaranteed. The tool that is used is the following generalization of Rademacher's theorem to infinite dimensions, see e.g. [Aro76, Ch. II, Sect.2, Thm. 1], [BL00, Thm. 6.42]. If the space $X$ in Theorem 6.1 is additionally a Hilbert space, a version can be found in [Mig76, Thm. 1.2].

**Theorem 6.1**
*Assume $T\colon X \to Y$ is locally Lipschitz continuous from a separable Banach space $X$ to a Hilbert space $Y$. Then the set $\mathcal{D}_T$ of points at which $T$ is Gâteaux differentiable is a dense subset of $X$.*

In [Aro76], the map $T$ is Lipschitz continuous and defined on an open subset of $X$. By considering neighborhoods of points separately, the formulation as in Theorem 6.1 can be obtained.

Now, we can formulate the main theorem of this paper.

**Theorem 6.2**
*Suppose that Assumption 1.1 and Assumption 1.2 are satisfied and let $u \in U$ be arbitrary. Then a Bouligand generalized derivative for $S_f$ in $u$ is given by the operator $\Xi(u;\cdot) \in \mathcal{L}(U, H_0^1(\Omega))$, where $\Xi(u;h)$ is the unique solution of the variational equation*

$$\text{Find } \eta \in H_0^1(D): \quad \langle L\eta - f'(u;h), z \rangle = 0 \quad \forall\, z \in H_0^1(D). \tag{21}$$

*Here, the sets*

$$D := I(f(u)) \cup A_{\mathrm{w}}^\varphi(f(u)) \quad \text{and} \quad D := I(f(u)) \cup A_\psi^{\mathrm{w}}(f(u))$$

*can be chosen and result in generally different generalized derivatives.*

*Proof.* The proof of [RU19, Prop. 5.5] implies that we can find an increasing, respectively, decreasing, sequence $(u_n)_{n\in\mathbb{N}} \subseteq U$ such that $S_f$ is Gâteaux differentiable at each $u_n$ and such that $u_n$ converges to $u$. Here, Theorem 6.1 is used.

Let us first assume $(u_n)_{n\in\mathbb{N}}$ is an increasing sequence with these properties. Let $h \in U$ be arbitrary. By Theorem 3.9, for each $n \in \mathbb{N}$, $S_f'(u_n; h)$ can be written as the solution of the variational equation

$$\text{Find } \eta_n \in H_0^1(D_n): \quad \langle L\eta_n - f'(u_n; h), z\rangle = 0 \quad \forall\, z \in H_0^1(D_n) \tag{22}$$

with the choice $D_n := I(f(u_n)) \cup A_{\mathrm{w}}^{\varphi}(f(u_n))$.

By Lemma 5.3, we conclude

$$H_0^1(I(f(u_n)) \cup A_{\mathrm{w}}^{\varphi}(f(u_n))) \to H_0^1(I(f(u)) \cup A_{\mathrm{w}}^{\varphi}(f(u)))$$

in the sense of Mosco. Now, Lemma 5.2 implies that $(S_f'(u_n; h))_{n\in\mathbb{N}}$ converges to the solution of (21) with $D := I(f(u)) \cup A_{\mathrm{w}}^{\varphi}(f(u))$. By definition, the resulting operator is a generalized derivative for $S_f$ in $u$.

When considering a decreasing sequence $(u_n)_{n\in\mathbb{N}}$, we use the representation of $S_f'(u_n; h)$ as the solution of the variational equation (22) with the choice $D_n := I(f(u_n)) \cup A_{\psi}^{\mathrm{w}}(f(u_n))$ and obtain the respective Mosco convergence, and thus the convergence of $S_f'(u_n)$ to the solution operator of (21) with $D := I(f(u)) \cup A_{\psi}^{\mathrm{w}}(f(u))$ from the second part of Lemma 5.3. □

## 6.1 Adjoint Representation of Clarke Subgradients

As in the unilateral case, see [RU19, Thm. 5.7], we can find an adjoint representation for the subgradient of a reduced objective function.

Assume $J\colon H_0^1(\Omega) \times U \to \mathbb{R}$ is a continuously differentiable objective function. We consider the optimization problem

$$\begin{aligned} &\min_{y,u} J(y, u) \\ &\text{subject to } y \in K_{\psi}^{\varphi}, \quad \langle Ly - f(u), z - y\rangle \geq 0 \quad \forall\, z \in K_{\psi}^{\varphi}. \end{aligned}$$

We present a formula for two generalized derivatives contained in Clarke's generalized differential that can be obtained for the reduced objective function

$$\hat{J}(u) := J(S_f(u), u)$$

in an arbitrary point $u \in U$.

**Corollary 6.3**
*Suppose that Assumption 1.1 and Assumption 1.2 are satisfied and let $u \in U$ be arbitrary. Denote by $q$ the unique solution of the variational equation*

$$\textit{Find } q \in H_0^1(D), \quad \langle L^* q, v\rangle = \langle J_y(S_f(u), u), v\rangle \quad \forall\, v \in H_0^1(D). \tag{23}$$

*Then the element*

$$f'(u)^* q + J_u(S_f(u), u)$$

*is in Clarke's generalized differential $\partial_C \hat{J}(u)$. In (23), the respective sets*

$$D := I(f(u)) \cup A_{\mathrm{w}}^{\varphi}(f(u)) \quad \text{ or } \quad D := I(f(u)) \cup A_{\psi}^{\mathrm{w}}(f(u))$$

*can be chosen and result in a particular generalized derivative.*

*Proof.* The proof is similar to the proof of [RU19, Thm. 5.4].

Since $L^*$ is coercive, (23) has a unique solution. As stated in Remark 2.6, we have

$$\partial_C \hat{J}(u) \ni \Xi^* J_y(S_f(u), u) + J_u(S_f(u), u) \tag{24}$$

for all $\Xi \in \partial S_f(u)$.

Assume that $q$ solves (23) for $D := I(f(u)) \cup A^{\varphi}_{\mathrm{w}}(f(u))$, respectively $D := I(f(u)) \cup A^{\mathrm{w}}_{\psi}(f(u))$. For $h \in U$, denote by $\Xi(u; h)$ the solution to (21). Now, we have

$$\begin{aligned}
\langle f'(u)^* q, w\rangle_{U^*,U} &= \langle f'(u; w), q\rangle \\
&\overset{(21)}{=} \langle L^* q, \Xi(u; w)\rangle \\
&\overset{(23)}{=} \langle J_y(S_f(u), u), \Xi(u; w)\rangle \\
&= \langle \Xi(u; \cdot)^* J_y(S_f(u), u), w\rangle_{U^*,U}
\end{aligned}$$

for all $w \in U$. Since $\Xi(u; \cdot) \in \partial S_f(u)$, the statement follows from (24). □

# Acknowledgments

This work was supported by the DFG under grant UL158/10-1 within the Priority Program SPP 1962 Non-smooth and Complementarity-based Distributed Parameter Systems: Simulation and Hierarchical Optimization.

# References

[ABM14] Hedy Attouch, Giuseppe Buttazzo, and Gérard Michaille. *Variational analysis in Sobolev and BV spaces*, volume 17 of *MOS-SIAM Series on Optimization*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA; Mathematical Optimization Society, Philadelphia, PA, second edition, 2014. Applications to PDEs and optimization.

[AH96] David R. Adams and Lars Inge Hedberg. *Function spaces and potential theory*, volume 314 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1996.

[Aro76] N. Aronszajn. Differentiability of Lipschitzian mappings between Banach spaces. *Studia Math.*, 57(2):147–190, 1976.

[Bar84] Viorel Barbu. *Optimal control of variational inequalities*, volume 100 of *Research Notes in Mathematics*. Pitman, Boston, MA, 1984.

[BB79] Haïm Brézis and Felix Browder. A property of Sobolev spaces. *Comm. Partial Differential Equations*, 4(9):1077–1083, 1979.

[Ber97] M. Bergounioux. Optimal control of an obstacle problem. *Appl. Math. Optim.*, 36(2):147–172, 1997.

[BL00] Yoav Benyamini and Joram Lindenstrauss. *Geometric nonlinear functional analysis. Vol. 1*, volume 48 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2000.

[BL04] Maïtine Bergounioux and Suzanne Lenhart. Optimal control of bilateral obstacle problems. *SIAM J. Control Optim.*, 43(1):240–255, 2004.

[BS00] J. Frédéric Bonnans and Alexander Shapiro. *Perturbation analysis of optimization problems*. Springer Series in Operations Research. Springer-Verlag, New York, 2000.

[CMWC18] Constantin Christof, Christian Meyer, Stephan Walther, and Christian Clason. Optimal control of a non-smooth semilinear elliptic equation. *Mathematical Control and Related Fields*, 8:247, 2018.

[CW19] Constantin Christof and Gerd Wachsmuth. Differential sensitivity analysis of variational inequalities with locally lipschitz continuous solution operators. *Applied Mathematics & Optimization*, 2019.

[DZ11] Michel C. Delfour and Jean-Paul Zolésio. *Shapes and geometries. Metrics, analysis, differential calculus, and optimization*. Adv. Des. Control. SIAM, Philadelphia, PA, second edition, 2011.

[Fri88] Avner Friedman. *Variational principles and free-boundary problems*. Robert E. Krieger Publishing Co., Inc., Malabar, FL, second edition, 1988.

[GR77] Michel Grun-Rehomme. Caractérisation du sous-différentiel d'intégrandes convexes dans les espaces de Sobolev. *J. Math. Pures Appl. (9)*, 56(2):149–156, 1977.

[GT01] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Volume 224 of Classics in Mathematics. Springer, Berlin, 2001. Reprint of the 1998 edition.

[Har77] Alain Haraux. How to differentiate the projection on a convex set in Hilbert space. Some applications to variational inequalities. *J. Math. Soc. Japan*, 29(4):615–631, 1977.

[Hel75] Lester L. Helms. *Introduction to potential theory*. Robert E. Krieger Publishing Co., Huntington, N.Y., 1975. Reprint of the 1969 edition, Volume 22 of Pure and Applied Mathematics.

[HK11] Michael Hintermüller and Ian Kopacka. A smooth penalty approach and a nonlinear multigrid algorithm for elliptic MPECs. *Comput. Optim. Appl.*, 50(1):111–145, 2011.

[HKM93] Juha Heinonen, Tero Kilpeläinen, and Olli Martio. *Nonlinear potential theory of degenerate elliptic equations*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1993. Oxford Science Publications.

[HR86] Jaroslav Haslinger and Tomáš Roubíček. Optimal control of variational inequalities. Approximation theory and numerical realization. *Appl. Math. Optim.*, 14(1):187–201, 1986.

[HU19] Lukas Hertlein and Michael Ulbrich. An inexact bundle algorithm for nonconvex nonsmooth minimization in Hilbert space. *SIAM J. Control Optim.*, 57(5):3137–3165, 2019.

[HW18] Felix Harder and Gerd Wachsmuth. Comparison of optimality systems for the optimal control of the obstacle problem. *GAMM-Mitt.*, 40(4):312–338, 2018.

[IK00] K. Ito and K. Kunisch. Optimal control of elliptic variational inequalities. *Appl. Math. Optim.*, 41(3):343–364, 2000.

[KKT03] T. Kärkkäinen, K. Kunisch, and P. Tarvainen. Augmented Lagrangian active set methods for obstacle problems. *J. Optim. Theory Appl.*, 119(3):499–533, 2003.

[KM92] Tero Kilpeläinen and Jan Malý. Supersolutions to degenerate elliptic equation on quasi open sets. *Comm. Partial Differential Equations*, 17(3-4):371–405, 1992.

[KS00] David Kinderlehrer and Guido Stampacchia. *An introduction to variational inequalities and their applications*, volume 31 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000. Reprint of the 1980 original.

[KW12] Karl Kunisch and Daniel Wachsmuth. Path-following for optimal control of stationary variational inequalities. *Comput. Optim. Appl.*, 51(3):1345–1373, 2012.

[Mig76] F. Mignot. Contrôle dans les inéquations variationelles elliptiques. *J. Functional Analysis*, 22(2):130–185, 1976.

[Mos69] U. Mosco. Convergence of convex sets and of solutions of variational inequalities. *Adv. Math.*, 3:510–585, 1969.

[MP84] F. Mignot and J.-P. Puel. Optimal control in some variational inequalities. *SIAM J. Control Optim.*, 22(3):466–476, 1984.

[MRW15] Christian Meyer, Andreas Rademacher, and Winnifried Wollner. Adaptive optimal control of the obstacle problem. *SIAM J. Sci. Comput.*, 37(2):A918–A945, 2015.

[Rod87] José-Francisco Rodrigues. *Obstacle problems in mathematical physics*, volume 134 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1987. Notas de Matemática [Mathematical Notes], 114.

[RU19] Anne-Therese Rauls and Stefan Ulbrich. Computation of a Bouligand generalized derivative for the solution operator of the obstacle problem. *SIAM J. Control Optim.*, 57(5):3223–3248, 2019.

[RW19] Anne-Therese Rauls and Gerd Wachsmuth. Generalized derivatives for the solution operator of the obstacle problem. *Set-Valued and Variational Analysis*, Feb 2019.

[Sto93] Peter Stollmann. Closed ideals in Dirichlet spaces. *Potential Anal.*, 2(3):263–268, 1993.

[SW13] Anton Schiela and Daniel Wachsmuth. Convergence analysis of smoothing methods for optimal control of stationary variational inequalities with control constraints. *ESAIM Math. Model. Numer. Anal.*, 47(3):771–787, 2013.

[Vel15] Bozhidar Velichkov. *Existence and regularity results for some shape optimization problems*, volume 19 of *Tesi Sc. Norm. Super. Pisa (N. S.)*. Ed. Norm., Pisa, 2015.

[Wac14] Gerd Wachsmuth. Strong stationarity for optimal control of the obstacle problem with control constraints. *SIAM J. Optim.*, 24(4):1914–1932, 2014.

[Wac18] Gerd Wachsmuth. Pointwise constraints in vector-valued Sobolev spaces: with applications in optimal control. *Appl. Math. Optim.*, 77(3):463–497, 2018.