

Bayesian Analysis on Limiting the Student- t Linear Regression Model

Yoshiko Hayashi *

Advanced Mathematical Institute, Osaka Metropolitan University

February 13, 2024

Abstract

For the outlier problem in linear regression models, the Student- t linear regression model is one of the common methods for robust modeling and is widely adopted in the literature. However, most of them applies it without careful theoretical consideration. This study provides the practically useful and quite simple conditions to ensure that the Student- t linear regression model is robust against an outlier in the y -direction using regular variation theory.

Keywords: Outlier, Linear regression model, Regularly varying distributions, Bayesian modeling

*yoshiko-hayashi@omu.ac.jp

1 Introduction

In regression analysis, outliers in a linear regression model can jeopardize the results by the ordinary least squares (OLS) estimator. The Student- t linear regression model, designed as a linear regression model with the error term having t -distribution, is one of the common methods to solve the outlier problem (Lange et al., 1989). While the Student- t linear regression model has been widely adopted, most studies apply it without careful theoretical consideration.

Bayesian robustness modeling using heavy-tailed distributions, which include t -distribution, provides the theoretical solution for the outlier problem. For a simple Bayesian model, when both the prior distribution and the likelihood of an observation are normal distributions, the posterior distribution is also a normal distribution, and the posterior mean is a weighted average of the mean of prior distribution and the observation. When the prior distribution and the observation are located far from each other and follow a normal distribution, the posterior distribution is far from both pieces of information; this is called *conflict*. For example, when a single observation $x = 15$ follows $N(\mu, 1)$ and the prior of the location parameter follows $N(0, 1)$, then the posterior distribution follows $N(7.5, 0.5)$. In this case, the posterior distribution is not suggested by either the prior distribution or the observation.

For this problem, Dawid (1973) formally provides the theoretical resolution of conflict between the prior distribution and the data, also known as *conflict of information*. He uses the pure location model in which the scale parameter is given, and clarifies how an outlier is automatically ignored in the posterior distribution when the outlier follows a heavy-tailed distribution. This result occurs because we believe the information about the prior distribution more than we believe the observation.

O'Hagan (1990) presents the concept of *credence*, which measures the degree of the tail's information. As Andrade and O'Hagan (2006) mention, credence represents how much we are prepared to believe in one source of information rather than another in the case of conflict; it is represented by the index of a regularly varying function. Andrade and O'Hagan (2011) show that in a univariate model, many observations that are located close enough create a larger credence, which equals the sum of each credence of the observations. When an outlier is far from the group of non-outliers with the same heavy-tailed distribution, the information of the group of non-outliers creates larger credibility, or credence. Thus, the posterior distribution is located closer to the non-outliers, and is robust against the outlier. Andrade and O'Hagan (2011) establish the sufficient conditions for robust modeling against a single outlier in n samples for a univariate model using regular variation theory. The sufficient condition requires the minimum number of non-outliers to be robust against an outlier. O'Hagan and Pericchi (2012) review previous studies on the resolution of the conflict.

O'Hagan (1988) applies heavy-tailed modeling to a Student- t linear regression model without an intercept term under the pure location structure, and demonstrates its robustness. For the model without the intercept term, outlier unconditionally conflicts with non-outliers. Therefore, a univariate model can be directly applied. By contrast, as Peña et al. (2009) mention, we need to be careful about the outlier in the x -direction for the model with an intercept term. Peña et al. (2009) show when the outliers in the x -direction reach infinity, the result of a Student- t linear model does not enable robustness. Peña et al. (2009) examine

the phenomenon using Kullback–Leibler divergence, and propose a down-weighting method that assigns a lower weight to outliers. As Andrade and O’Hagan’s (2011) show the heavy-tailed modeling using t -distribution is a partial robust modeling. Thus the location-scale modeling completely cannot ignores outliers. Gagnon et al. (2020) theoretically develop a robust linear regression model using a super heavy-tailed distribution, which is heavier than t -distribution and provides a wholly robust modeling. Although the Student- t linear regression model provides a partial robustness, the model is widely applied. Thus, it is very important to clarify how the model works as a robust modeling.

Our study investigates the conditions for the Student- t linear model with an intercept term for an outlier in the y -direction by extending Andrade and O’Hagan’s (2011) conditions. For this purpose, first, we investigate the range in which there is conflict between an outlier and non-outliers, which is the necessary condition to apply heavy-tailed modeling. Then, we clarify the condition of the model’s robustness. Heavy-tailed modeling as a resolution




Figure 1: Conflict in linear model: Straight lines show the regression line by OLS and dotted lines show it without the outlier.

of a conflict between an outlier and non-outliers works when the outlier and the mean of the group of non-outliers are located far enough, and the sufficient condition for the number of non-outliers is satisfied. A linear regression model provides the mean of y conditioned on x . Thus, the conflict of information in a linear regression model with an intercept term occurs when an outlier is located far from the regression line, and non-outliers lie close to the regression line created from non-outliers.

The left panel in Figure 1 shows the case in which the outlier conflicts with the group of non-outliers. The figure shows that the outlier is located far from the regression line by OLS. In this case, the Student- t linear regression model is robust against the outlier in the y -direction. This is because the information of the conditional distribution of the outlier is less credible than that for the grouped non-outlier data, under the assumption of the same degrees of freedom of t -distribution for all data, which represent *credence*. Non-outliers in the left panel of Figure 1 are close to each other and create large credence, while the outlier does not belong to the regression line suggested by the grouped data and creates small credence. Meanwhile, as shown in the right panel of Figure 1, when the outlier is in the x -direction, which is called the leverage point, all data, including the outlier, are sufficiently close to the

regression line, and create larger credence than the regression line without the outlier, which is presented by the dotted line in Figure 1. In this case, the straight line in the right panel of Figure 1 has larger credence than the dotted line does.

The rest of the paper is organized as follows. In Section 2, we provide the condition for conflicting information between an outlier and non-outliers in the Student- t linear regression model. Section 3 shows the sufficient conditions for the Student- t linear regression model. Section 4 presents simulation results in a simple linear regression model. Section 5 concludes.

2 Conflicting Information in the Student- t Linear Regression Model

To examine the limitation of the robustness for the Student- t linear model with an intercept term, we consider the following linear regression model. The dependent variable \mathbf{y} is an $n \times 1$ vector, the independent variable \mathbf{X} is an $n \times (k+1)$ full-rank matrix, β is a $(k+1) \times 1$ vector, and \mathbf{u} is an $n \times 1$ vector assumed to be independent and identically distributed:

$$\mathbf{y} = \mathbf{X}\beta + \mathbf{u}, \quad (1)$$

where

$$\mathbf{X} = \begin{bmatrix} 1 & X_{11} & \dots & X_{k1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & X_{1n} & \dots & X_{kn} \end{bmatrix},$$

Consider the residual of the result from OLS for the model in equation (1):

$$\mathbf{y} = \mathbf{X}\hat{\beta}_{\text{ols}} + \mathbf{e}, \quad (2)$$

where

$$\mathbf{e}' = \left[e_{/out}^1, \dots, e_{/out}^{n-1}, e_{out} \right],$$

and subscripts $/out$ and out show a non-outlier and an outlier, respectively.

Assume that the non-outliers are located close enough to the regression line. If the outlier moves away from the regression line and does so faster than the group of non-outliers does, then the residual e_{out} reaches infinity as the outlier reaches infinity in the y -direction. Since non-outliers create combined credence, if one of the non-outliers conflicts with the outlier, then the group of non-outliers conflicts with the outlier. As shown in Figure 1, when an outlier is located close enough to the group of non-outliers in the x -direction, they conflict. Therefore, if the partial derivative of e_{out} with respect to y_{out} is larger than the partial derivative of the

closest non-outlier's residual, the outlier conflicts with the group of non-outliers.

$$\frac{\partial e_{out}}{\partial y_{out}} > \frac{\partial e_{out}^{max}}{\partial y_{out}}, \quad (3)$$

where subscript $/out$ is a non-outlier.

According to Chatterjee and Hadi (1988), the components in inequality (3) are defined as follows:

$$\frac{\partial e_{out}}{\partial y_{out}} = \left(\frac{n-1}{n} \right) - (\mathbf{x}_{out} - \bar{\mathbf{x}})'(\tilde{\mathbf{X}}'\tilde{\mathbf{X}})^{-1}(\mathbf{x}_{out} - \bar{\mathbf{x}}), \quad (4)$$

and

$$\frac{\partial e_{/out}}{\partial y_{out}} = \left(-\frac{1}{n} \right) - (\mathbf{x}_{/out} - \bar{\mathbf{x}})'(\tilde{\mathbf{X}}'\tilde{\mathbf{X}})^{-1}(\mathbf{x}_{out} - \bar{\mathbf{x}}), \quad (5)$$

where $\bar{X}_k = \sum_{i=1}^n X_{ki}/n$ and

$$\tilde{\mathbf{X}} = \begin{bmatrix} X_{11} - \bar{X}_1 & \dots & X_{k1} - \bar{X}_k \\ \vdots & \ddots & \vdots \\ X_{1n} - \bar{X}_1 & \dots & X_{kn} - \bar{X}_k \end{bmatrix}, \quad \mathbf{x}_n = \begin{bmatrix} X_{1n} \\ \vdots \\ X_{kn} \end{bmatrix}, \quad \bar{\mathbf{x}} = \begin{bmatrix} \bar{X}_1 \\ \vdots \\ \bar{X}_k \end{bmatrix}.$$

Substituting equations (4) and (5) in inequality (3), we obtain the following relationship:

$$1 - (\mathbf{x}_{out} - \bar{\mathbf{x}})'(\tilde{\mathbf{X}}'\tilde{\mathbf{X}})^{-1}(\mathbf{x}_{out} - \bar{\mathbf{x}}) > -(\mathbf{x}_{/out} - \bar{\mathbf{x}})'(\tilde{\mathbf{X}}'\tilde{\mathbf{X}})^{-1}(\mathbf{x}_{out} - \bar{\mathbf{x}}). \quad (6)$$

Since $0 \leq (\mathbf{x}_n - \bar{\mathbf{x}})'(\tilde{\mathbf{X}}'\tilde{\mathbf{X}})^{-1}(\mathbf{x}_n - \bar{\mathbf{x}}) \leq 1$ and $(\tilde{\mathbf{X}}'\tilde{\mathbf{X}})$ is a positive definite, the following lemma is obtained.

Lemma 1. (Conflicting information in the Student- t linear regression model)

If the following condition holds, the residual of the outlier e_{out} reaches infinity as y_{out} goes to infinity in the linear regression model.

$$\max[(\mathbf{x}_{out}^j - \bar{\mathbf{x}})'(\mathbf{x}_{out} - \bar{\mathbf{x}})] > 0, \quad j = 1, \dots, n-1.$$

3 Sufficient Conditions for Rejecting an Outlier in the Student- t Linear Regression Model

This section investigates the sufficient conditions for a Student- t linear regression model being robust based on Andrade and O'Hagan's (2011) corollary 4, which shows the conditions for robustness against a single outlier out of n samples in a univariate model. To examine the conditions for the Student- t linear regression model, we adopt the independent Jeffreys priors derived by Fonseca et al. (2008, 2014) under the given degrees of freedom.

As shown in Andrade and O'Hagan (2006), the credence is defined as c for $f(x) \in R_{-c}(c >$

0), where R_{-c} presents $f(x)$ is regularly varying at ∞ with index c . Thus, for t -distribution with d degrees of freedom, the credence is $d+1$ (see Appendix B).

Assume all data, including a single outlier among n observations, is t -distributed with the degree of freedom, d , $t_{(d)}(m, s)$ ¹. Since the t -distribution is a location scale family, the likelihood can be denoted as $p(y_i|\mathbf{X}, \beta, \sigma) = 1/\sigma \times h[(y_i - X'_i\beta)/\sigma]$. X'_i is the i -th row of \mathbf{X} .

For simplicity, assume all data have the same likelihood function and the non-outliers are close enough to the conditional mean $X'_i\beta$. Consider the n -th observation is an outlier.

Model

$$\begin{cases} y_i|\mathbf{X}, \beta, \sigma \stackrel{D}{\sim} p(y_i|X_i, \beta, \sigma) = 1/\sigma \times h[(y_i - X'_i\beta)/\sigma] \text{ independent } (i = 1, \dots, n), \\ \beta_q \stackrel{D}{\sim} p(\beta_q) \propto 1, \quad (q = 0, 1, \dots, k), \\ \sigma \stackrel{D}{\sim} p(\sigma) \propto 1/\sigma, \\ h \in R_{-\rho}, \rho > 1, \quad (i = 1, \dots, n). \end{cases} \quad (7)$$

Theorem 1. (Robustness of an outlier among n observations in the Student- t linear regression model)

Consider n observations in the present model, and Lemma 1 holds, in which the residual $e_n = y_n - x'_n\beta$ reaches infinity as y_n goes to infinity. Then, the following condition holds:

$$\rho < \{n - (k + 1)\}.$$

Then, the posterior distribution partially ignores the outlier:

$$p(\beta, \sigma|\mathbf{X}, \mathbf{y}) \propto \sigma^{\rho-2} p(\mathbf{y}^{(n-1)}|\beta, \sigma, \mathbf{X}^{(n-1)}) \quad \text{as } y_n \rightarrow \infty, \quad (8)$$

where the superscript notation $(n - 1)$ is used to indicate the omission of the n -th observation.

proof: See Appendix B.

Thus, for t -distribution with d degrees of freedom, the sufficient condition in Theorem 1 is $d < \{n - k - 2\}$.

4 Example

We consider the following case of a simple linear regression model with a single outlier:

$$y_i = \beta_0 + \beta_1 x_i + u_i. \quad (i = 1, \dots, n). \quad (9)$$

When the dependent variable of the outlier is larger than the sample mean ($x_{out} > \bar{x}$), the following condition is proposed, in which the residual $e_n = y_n - x'_n\beta$ reaches infinity as

¹The t -distribution, $t_{(d)}(m, s)$, reports mean m and inverse scaling parameter s with d degrees of freedom.

y_n goes to infinity:

$$(x_{/.max} - \bar{x})(x_{out} - \bar{x}) > 0, \quad (10)$$

where $x_{/.max}$ is the maximum value among the non-outliers closest to the outlier. Thus, the following robust range is obtained:

$$\bar{x} < x_{/.max}. \quad (11)$$

Therefore, including the opposite side of outlier $(x_{out} - \bar{x}) < 0$ produces

$$x_{/.min} < \bar{x} < x_{/.max}. \quad (12)$$

Since the sample mean can be arranged as

$$\bar{x} = \frac{\Sigma_{/out} x_{/out}}{n} + \frac{x_{out}}{n}, \quad (13)$$

substituting the equation (13) into equation (12) presents the following range for a simple linear regression model, in which e_{out} reaches infinity as y_{out} goes to infinity:

$$\left(x_{/.min} - \frac{\Sigma_{/out} x_{/out}}{n} \right) < x_{out} < \left(x_{/.max} - \frac{\Sigma_{/out} x_{/out}}{n} \right). \quad (14)$$

These results highlight that the robust range is wider, as the number of non-outliers is larger. In addition, when the independent variable of the outlier is located far from other data, there is no conflict of information irrespective of the value of y .

4.1 Numerical Examples

This subsection investigates the robustness performance in relation to the value of the outlier in the Student- t linear regression model for a simple linear regression. For robustness, the degrees of freedom of the t -distributed errors need to be small enough. Thus, we utilize three degrees of freedom, $u_i \sim t_{(3)}(0, \sigma)$ for the error term. We employ the independent Jeffreys priors; the priors of β_0 and β_1 have uniform distribution, and the prior distribution of σ is $1/\sigma$. Thus, By Theorem 1, the sufficient condition is $3 < n - 4$. The simulated observations are defined as $y_i = 3 + 2x_i + u_i$. We set $x_{/out} = [-2, -1, \dots, 2]$ for the first simulation and $[-2, -1, \dots, 2, -2, -1, \dots, 2]$ for the second one. The error terms are generated from the normal distribution with mean 0 and variance 1. We move the outlier in the x -direction, from -50 to 50, and set $y_{out} = -10$. The left panels of Figure 2 depict the simulated data we use for these simulations. The middle and right panels of Figure 2 show the results of the numerical evaluation of the posterior mean of the parameter β_1 ; the upper right panel illustrates the result of the value for $n = 6$, which satisfies the sufficient condition, and the lower right panel presents it for $n = 11$, which satisfies the condition. The results present that the Student- t linear regression model is robust within the controllable range defined in Lemma 1, which is shown as the vertical dotted lines.




Figure 2: Left panels: Scatterplots of simulated data. Right panels: Posterior mean of the slope β_1 . The upper panel shows the result for $n = 6$, which does not satisfy the sufficient condition, and the lower panel shows the result for $n = 11$, which satisfies the condition. The straight line depicts the result of the Student- t linear regression model, and the dashed line depicts that of the linear regression model with normally distributed error terms. The vertical dotted lines show the range defined in Lemma 1.

5 Concluding remarks

This study extended Andrade and O’Hagan’s (2011) condition for resolving the outlier problem of the Student- t linear regression model. The model treats outliers as a natural outcome of the data, and does not remove them arbitrarily. The condition works when there is conflicting information between outliers and non-outlier. However, in a linear regression model, an outlier does not conflict with non-outliers when the outlier is located far from non-outliers in the x -direction. Thus, we first clarified the range of the presence of conflicting information in a linear regression model. Then, we derived the sufficient condition for robustness of the Student- t linear regression model in the above range. Future research should investigate the conditions for many outliers. Furthermore, it would be interesting to extend this study to a model with unknown degrees of freedom for the t -distribution.

References

- [1] Andrade, J.A.A. and O’Hagan, A. Bayesian robustness modeling using regularly varying distribution. *Bayesian Analysis* **1**, 169–188, 2006.
- [2] Andrade, J.A.A. and O’Hagan, A. Bayesian robustness modelling of location and scale parameters. *Scandinavian Journal of Statistics* **38**(4), 691–711, 2011.
- [3] Bingham, N.H., Goldie, C.M., and Teugels, J.L. *Regular Variation*. Cambridge University Press, Cambridge, 1987.
- [4] Chatterjee, S. and Hadi, A.S. *Sensitivity Analysis in Linear Regression*. Wiley, New York, 1988.
- [5] Dawid, A.P. Posterior expectations for large observations. *Biometrika* **60**(3), 664–667, 1973.
- [6] Fonseca, T.C.O., Ferreira, M.A.R., and Migon, H.S. Objective Bayesian analysis for the Student- t regression model. *Biometrika* **95**(2), 325–333, 2008 (Amendments and Corrections. **101**(1), 251, 2014).
- [7] Gagnon, P., Desgagné, A., and Bédard, M. (2020). A New Bayesian Approach to Robustness Against Outliers in Linear Regression. *Bayesian Analysis* **15**, 389–414, 2020.
- [8] Lange, K.L., Little, R.J.A., and Taylor, J.M.G. Robust statistical modeling using the t distribution. *Journal of the American Statistical Association* **84**(408), 881–892, 1989.
- [9] O’Hagan, A. Modelling with heavy tails. *Bayesian Statistics* **3**, 345–359, 1988.
- [10] O’Hagan, A. (1990), On Outliers and Credence for Location Parameter Inference. *Journal of the American Statistical Association* , **85**, pp.172–176.
- [11] O’Hagan, A. and Pericchi, L. Bayesian heavy-tailed models and conflict resolution: A review. *Brazilian Journal of Probability and Statistics* **26**(4), 372–401, 2012.
- [12] Peña, D., Zamar, R., and Yan, G. Bayesian likelihood robustness in linear models. *Journal of Statistical Planning Inference* **139**(7), 2196–2207, 2009.
- [13] Resnick, S.I. *Heavy-tail Phenomena: Probabilistic and Statistical Modeling*. Springer, New York, 2006.

Appendix A

The tail behavior can be presented by the index of a regularly varying function. The index ρ is defined as follows.

A positive measurable function $f(x)$ is regularly varying at ∞ with index $\rho \in R$ for an arbitrary positive t .

$$\lim_{x \rightarrow \infty} \frac{f(tx)}{f(x)} = t^\rho \quad (\text{A.1})$$

We present it as $f(x) \in R_\rho$ in this study, and $l(x) \in R_0$ is called “slowly varying.” The regularly varying function can be presented as $f(x) = t^\rho l(x)$.

Using the property

$$\lim_{x \rightarrow \infty} \frac{\log(f(x))}{\log x} = \rho, \quad (\text{A.2})$$

we obtain the index for t distribution with the degrees of freedom, d , as

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{\log(p(x; d, \mu, \sigma^2))}{\log x} &= \lim_{x \rightarrow \infty} \frac{(A - \frac{d+1}{2} \log(\{1 + \frac{1}{d}(\frac{x-\mu}{\sigma})^2\}))}{\log x} \\ &= -(d+1), \end{aligned} \quad (\text{A.3})$$

where $A = \log \left(\frac{\Gamma(\frac{d+1}{2})}{\Gamma(\frac{d}{2})\pi^{1/2}d^{1/2}\sigma} \right)$.

Some properties of the regularly varying function used in this study are seen in Bingham et al. (1987) and Resnick (2006).

Appendix B

When scale parameter σ is given, the posterior distribution of β in the model is as follows:

$$\begin{aligned} p(\beta | \sigma, \mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)}) &\propto p(\beta) \cdot \prod_{i=1}^{n-1} h[(y_i - X'_i \beta) / \sigma] \\ &\propto \prod_{i=1}^{n-1} h[(y_i - X'_i \beta) / \sigma] \in R_{-(n-1)\rho}. \end{aligned} \quad (\text{B.1})$$

Applying transformation $\tau = \frac{1}{\sigma}\beta$, which is a $(k+1) \times 1$ vector, we obtain

$$p(\mathbf{y}^{(n-1)} | \sigma, \mathbf{X}^{(n-1)}) = \left(\frac{1}{\sigma} \right)^{n-(k+1)-1} \int \cdots \int_{\tau} \prod_{i=1}^{n-1} h[(y_i / \sigma) - X'_i \tau] d\tau. \quad (\text{B.2})$$

When all elements of \mathbf{X} are given and bounded, $\int \cdots \int_{\tau} \prod_{i=1}^{n-1} h[(y_i / \sigma) - X'_i \tau] d\tau$ in σ is $O(1)$. Thus, as a function of σ , it is slowly varying,

$$\int \cdots \int_{\tau} \prod_{i=1}^{n-1} h[(y_i / \sigma) - x'_i \tau] d\tau \in R_0. \quad (\text{B.3})$$

Thus, the marginal posterior distribution of σ given information $\mathbf{X}^{(n-1)}$ and $\mathbf{y}^{(n-1)}$ be-

comes

$$p(\sigma|\mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)}) \propto p(\sigma) \cdot p(\mathbf{y}^{(n-1)}|\sigma, \mathbf{X}^{(n-1)}) \in R_{-(n-(k+1))}. \quad (\text{B.4})$$

Again, applying transformation $\tau = \frac{1}{\sigma}\beta$ produces the marginal posterior distribution of y_n given information $\mathbf{X}^{(n-1)}$ as

$$p(y_n|\sigma, \mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)}) = \left(\frac{1}{\sigma}\right)^{n-(k+1)} \int \cdots \int_{\tau} h[(y_n/\sigma) - X'_n\tau] \Pi_{i=1}^{n-1} h[(y_i/\sigma) - X'_i\tau] d\tau. \quad (\text{B.5})$$

When non-outliers are located close enough to the regression line, Andrade and O'Hagan's (2011) Proposition 1, which gives the convolution of regularly varying densities being distributed as the sum of them, $f * g(x) \sim f(x) + g(x)$, can be applied as $f(y) = h[(y_n/\sigma) - X'_n\tau]$ and $g(y) = \Pi_{i=1}^{n-1} h[(y_i/\sigma) - X'_i\tau]$. When the residual $e_n = y_n - x'_n\beta$ reaches infinity, as y_n goes to infinity, we obtain

$$\int \cdots \int_{\tau} h[(y_n/\sigma) - X'_n\tau] \Pi_{i=1}^{n-1} h[(y_i/\sigma) - X'_i\tau] d\tau \in R_{-\min(\rho, (n-1)\rho)}. \quad (\text{B.6})$$

Lemma 1 shows the condition for the residual $e_n = y_n - x'_n\beta$ reaching infinity as y_n goes to infinity. Accordingly, the marginal posterior distribution for σ is

$$p(\sigma|\mathbf{y}) = \frac{p(y_n|\sigma, \mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)})p(\sigma|X^{(n-1)}, \mathbf{y}^{(n-1)})}{\int p(y_n|\sigma, \mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)})p(\sigma|\mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)})d\sigma}. \quad (\text{B.7})$$

Next, consider the case in which y_n reaches infinity. As a function of y_n , the posterior distribution of $p(y_n|\sigma, \mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)})$ takes the form of $\frac{1}{\sigma}g\left(\frac{y_n}{\sigma}\right) \in R_{-\rho}$. Thus, by the relationship $g\left(\frac{y_n}{\sigma}\right)/g(y_n) = \sigma^\rho$,

$$\lim_{y_n \rightarrow \infty} p(\sigma|\mathbf{y}) = \frac{\sigma^{\rho-1}p(\sigma|X^{(n-1)}, \mathbf{y}^{(n-1)})}{\lim_{y_n \rightarrow \infty} \int \frac{1}{\sigma}g\left(\frac{y_n}{\sigma}\right)/g(y_n)p(\sigma|\mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)})d\sigma}. \quad (\text{B.8})$$

From (B.4), we obtain

$$p(\sigma|\mathbf{X}^{(n-1)}, \mathbf{y}^{(n-1)}) \propto \sigma^{-(n-(k+1))}l(\sigma). \quad (\text{B.9})$$

Thus, for the dominator of (B.8) to exist, $\rho < n - (k + 1)$ should hold.