

The 4-D Gaussian Random Vector Maximum Conjecture and the 3-D Simplex Mean Width Conjecture

Wei Sun^{a,*}, Ze-Chun Hu^b and Guolie Lan^c

^a Department of Mathematics and Statistics, Concordia University, Canada

^b College of Mathematics, Sichuan University, China

^c School of Economics and Statistics, Guangzhou University, China

Abstract

We prove the four-dimensional Gaussian random vector maximum conjecture. This conjecture asserts that among all centered Gaussian random vectors $X = (X_1, X_2, X_3, X_4)$ with $E[X_i^2] = 1$, $1 \leq i \leq 4$, the expectation $E[\max(X_1, X_2, X_3, X_4)]$ is maximal if and only if all off-diagonal elements of the covariance matrix equal $-\frac{1}{3}$. As a direct consequence, we resolve the three-dimensional simplex mean width conjecture. This latter conjecture is a long-standing open problem in convex geometry, which asserts that among all simplices inscribed into the three-dimensional unit Euclidean ball the regular simplex has the maximal mean width.

Keywords Gaussian random vector, maximum conjecture, tetrahedron, mean width conjecture, derivative formula.

Mathematics Subject Classification (2020) 60E15, 52A40.

*Corresponding author: wei.sun@concordia.ca

1 Introduction and main results

The multivariate Gaussian distribution has played a fundamental role in probability and statistics (see [18] and references therein). Although its study has a long history, there are remarkable new results which have been obtained in recent years. For example, Royen [15] proved the Gaussian correlation inequality (cf. also Latała and Matlak [12]), which has important applications in small ball probabilities (cf. Li [13] and Shao [16]). As another example, based on stochastic calculus, Eldan [6] gave a novel short proof of the celebrated Borell inequality and obtained an almost tight, two-sided, dimension-free robustness estimate for the Gaussian noise stability deficit. However, some unsolved problems still remain. A major one is the Gaussian random vector maximum (GRVM) conjecture. This conjecture asserts that among all centered Gaussian random vectors $X = (X_1, \dots, X_n)$ with $E[X_i^2] = 1$, $1 \leq i \leq n$, the expectation $E[\max(X_1, \dots, X_n)]$ is maximal if and only if all off-diagonal elements of the covariance matrix equal $-\frac{1}{n-1}$, where $n \geq 4$. We would like to point out that the GRVM conjecture is much more difficult than another problem of order statistics of Gaussian random variables. If we add absolute values to X_i 's, then it is known that the expectation $E[\max(|X_1|, \dots, |X_n|)]$ attains maximum if and only if X_1, \dots, X_n are independent (see Šidák [17] and Gluskin [9]).

Besides its own interest in probability and statistics, the GRVM conjecture has a deep connection with convex geometry. It is known that the n -dimensional GRVM conjecture is equivalent to the $(n-1)$ -dimensional simplex mean width (SMW) conjecture (see Litvak [14]). Given a convex body K in \mathbb{R}^n , its support function and mean width are defined by

$$h_K(u) = \max_{x \in K} \langle u, x \rangle, \quad w(K) = 2 \int_{S^{n-1}} h_K(u) dm(u).$$

Hereafter $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ denote the Euclidean inner product and norm, respectively, and m is the normalized Lebesgue measure on the sphere S^{n-1} . The SMW conjecture asserts that among all simplices inscribed into the n -dimensional unit ball the regular simplex has the maximal mean width, where $n \geq 3$.

The SMW conjecture is a long-standing open problem in convex geometry. As pointed out by Gritzmann and Klee [10, Section 9.10.2], several authors ([8, 1, 2, 20]) have assumed the existence of a proof for the conjecture. However, we are not aware of any such proof. As mentioned in recent works by Hug and Schneider [11] and by Börözký and Schneider [3], the n -dimensional SMW conjecture remains open for $n \geq 3$. In the information theory community, it was a general belief that the SMW conjecture is true (cf. [1, 2, 5, 21]) and the SMW conjecture is of interest because of its connection with the famous simplex code conjecture (see [10, Section 9.10.2] or [14] for an introduction). For more history about the GRVM conjecture and the SMW conjecture, we refer the reader to the beautiful survey paper of Litvak [14].

In this paper, we will prove the four-dimensional GRVM conjecture and, equivalently, the three-dimensional SMW conjecture. Let (Ω, \mathcal{F}, P) be a probability space such that, for any nonnegative-definite symmetric real matrix Λ , there exists a Gaussian random vector X on (Ω, \mathcal{F}, P) whose covariance matrix is given by Λ . We denote by \mathcal{G} the set of all four-dimensional centered Gaussian random vectors $X = (X_1, X_2, X_3, X_4)$ with $E[X_1^2] = E[X_2^2] = E[X_3^2] = E[X_4^2] = 1$. Hereafter E

denotes the expectation with respect to P . Define

$$\mathcal{M}(X) = E[\max(X_1, X_2, X_3, X_4)], \quad X \in \mathcal{G}.$$

For $X \in \mathcal{G}$, we denote its covariance matrix by Λ^X or simply by Λ if no confusion arises. We use $\bar{\mathcal{S}}$ and \mathcal{S} to denote respectively the set of all nonnegative-definite and positive-definite 4×4 symmetric real matrices with all diagonal elements equal to 1. For $\Lambda \in \bar{\mathcal{S}}$, take an $X^\Lambda \in \mathcal{G}$ with covariance matrix Λ and define

$$F(\Lambda) = \mathcal{M}(X^\Lambda).$$

Now we state the main results of this paper.

Theorem 1.1 *The function \mathcal{M} defined on \mathcal{G} attains its maximum at X if and only if $\Lambda_{kl}^X = -\frac{1}{3}$ for all $k < l$.*

As a direct consequence of Theorem 1.1, we conclude that the three-dimensional SMW conjecture is true.

Theorem 1.2 *Among all simplices inscribed into the three-dimensional unit Euclidean ball the regular simplex has the maximal mean width.*

The SMW conjecture is a purely geometric problem. We resolve the three-dimensional SMW conjecture by combining probability theory and convex geometry. First, we use a probabilistic method to derive the first order partial derivative formula for the function $F(\Lambda)$. To the best of our knowledge, this formula is unknown before. Based on this novel formula, we analyze the maximum point of the function $\mathcal{M}(X)$. By applying geometric techniques, we discover an intrinsic relation satisfied by the maximum point (see (4.18) below). This important relation is described in terms of the dihedral angles of the tetrahedron $\{X_1, X_2, X_3, X_4\}$. We will show that the regular tetrahedron is the unique tetrahedron that satisfies the relation. To this end, we introduce a special function H (see (4.25) below) and reduce the three-dimensional SMW conjecture to the monotonicity of H . We will use purely analytic techniques to prove that H is a strictly decreasing function. This result has independent interest and some exquisitely delicate inequalities are involved in the proof. We hope the methods developed in this paper can be further developed so as to investigate the higher dimensional GRVM and SMW conjectures.

The remainder of this paper is organized as follows. In Section 2, we derive the first order partial derivative formula for $F(\Lambda)$. By virtue of this formula, we derive in Section 3 the second order partial derivative formula for $F(\Lambda)$ and the explicit formula for $F(\Lambda)$ itself. In Section 4, we prove Theorem 1.1. The proof relies heavily on the monotonicity of the special function H mentioned above. In Section 5, we prove that H is a strictly decreasing function. In Section 6, we make some remarks including non-concavity of the function $F(\Lambda)$ and the lower and upper bounds for the function $\mathcal{M}(X)$. Some auxiliary results are given in the Appendix, Section 7.

2 First order partial derivatives of $F(\Lambda)$

Under the embedding map $I(\Lambda) := (\Lambda_{12}, \Lambda_{13}, \Lambda_{14}, \Lambda_{23}, \Lambda_{24}, \Lambda_{34})$, $\bar{\mathcal{S}}$ and \mathcal{S} become respectively closed and open convex subsets of \mathbb{R}^6 . Let $\Lambda \in \mathcal{S}$. We define

$$f_\Lambda(x) = \frac{1}{\sqrt{(2\pi)^4 \det(\Lambda)}} e^{-\frac{1}{2}x^T \Lambda^{-1} x}, \quad x \in \mathbb{R}^4. \quad (2.1)$$

Hereafter we use T to denote the transpose of a vector or a matrix.

Lemma 2.1 *For $\Lambda \in \mathcal{S}$, we have*

$$\frac{\partial F}{\partial \Lambda_{12}}(\Lambda) = - \int_{-\infty}^{\infty} \int_{-\infty}^t \int_{-\infty}^t f_\Lambda(t, t, x_3, x_4) dx_3 dx_4 dt. \quad (2.2)$$

Similar formulas hold for all $k < l$. Hence

$$\frac{\partial F}{\partial \Lambda_{kl}}(\Lambda) < 0, \quad \forall k < l.$$

Proof. Let $X = (X_1, X_2, X_3, X_4) \in \mathcal{G}$. Note that

$$\lim_{t \uparrow \infty} tP\{\max(X_1, X_2, X_3, X_4) > t\} \leq \sum_{k=1}^4 \lim_{t \uparrow \infty} tP\{X_k > t\} = 0.$$

Then, we have

$$\begin{aligned} & E[\max(X_1, X_2, X_3, X_4)] \\ &= \int_{-\infty}^{\infty} t dP\{\max(X_1, X_2, X_3, X_4) < t\} \\ &= - \int_0^{\infty} t dP\{\max(X_1, X_2, X_3, X_4) > t\} + \int_{-\infty}^0 t dP\{\max(X_1, X_2, X_3, X_4) < t\} \\ &= \int_0^{\infty} P\{\max(X_1, X_2, X_3, X_4) > t\} dt - \int_{-\infty}^0 P\{\max(X_1, X_2, X_3, X_4) < t\} dt \\ &= \int_0^{\infty} [1 - P\{\max(X_1, X_2, X_3, X_4) < t\}] dt - \int_0^{\infty} P\{\max(X_1, X_2, X_3, X_4) < -t\} dt \\ &= \int_0^{\infty} [1 - P\{\max(X_1, X_2, X_3, X_4) < t\} - P\{\min(X_1, X_2, X_3, X_4) > t\}] dt. \end{aligned} \quad (2.3)$$

We have

$$f_\Lambda(x) = \frac{1}{(2\pi)^4} \int_{\mathbb{R}^4} e^{-is^T x - \frac{1}{2}s^T \Lambda s} ds = \frac{1}{(2\pi)^4} \int_{\mathbb{R}^4} e^{-is^T x - \frac{1}{2} \sum_{k,l=1}^4 \Lambda_{kl} s_k s_l} ds, \quad x \in \mathbb{R}^4,$$

and

$$\frac{\partial f_\Lambda}{\partial \Lambda_{kl}} = \frac{\partial^2 f_\Lambda}{\partial x_k \partial x_l}, \quad \forall k < l. \quad (2.4)$$

By (2.3), we get

$$\begin{aligned}
F(\Lambda) &= \int_0^\infty [1 - P\{\max(X_1^\Lambda, \dots, X_4^\Lambda) < t\} - P\{\min(X_1^\Lambda, \dots, X_4^\Lambda) > t\}] dt \\
&= \int_0^\infty \left[1 - \int_{-\infty}^t \cdots \int_{-\infty}^t f_\Lambda(x_1, \dots, x_4) dx_1 \cdots dx_4 - \int_t^\infty \cdots \int_t^\infty f_\Lambda(x_1, \dots, x_4) dx_1 \cdots dx_4 \right] dt.
\end{aligned}$$

Then, we obtain by (2.1) and (2.4) that

$$\begin{aligned}
&\frac{\partial F}{\partial \Lambda_{12}}(\Lambda) \\
&= - \int_0^\infty \left[\int_{-\infty}^t \cdots \int_{-\infty}^t \frac{\partial f_\Lambda}{\partial \Lambda_{12}} dx_1 \cdots dx_4 + \int_t^\infty \cdots \int_t^\infty \frac{\partial f_\Lambda}{\partial \Lambda_{12}} dx_1 \cdots dx_4 \right] dt \\
&= - \int_0^\infty \left[\int_{-\infty}^t \cdots \int_{-\infty}^t \frac{\partial^2 f_\Lambda}{\partial x_1 \partial x_2} dx_1 \cdots dx_4 + \int_t^\infty \cdots \int_t^\infty \frac{\partial^2 f_\Lambda}{\partial x_1 \partial x_2} dx_1 \cdots dx_4 \right] dt \\
&= - \int_0^\infty \left[\int_{-\infty}^t \int_{-\infty}^t f_\Lambda(t, t, x_3, x_4) dx_3 dx_4 + \int_t^\infty \int_t^\infty f_\Lambda(t, t, x_3, x_4) dx_3 dx_4 \right] dt \\
&= - \int_0^\infty \int_{-\infty}^t \int_{-\infty}^t f_\Lambda(t, t, x_3, x_4) dx_3 dx_4 dt + \int_{-\infty}^0 \int_{-\infty}^t \int_{-\infty}^t f_\Lambda(-t, -t, -x_3, -x_4) dx_3 dx_4 dt \\
&= - \int_{-\infty}^\infty \int_{-\infty}^t \int_{-\infty}^t f_\Lambda(t, t, x_3, x_4) dx_3 dx_4 dt \\
&< 0.
\end{aligned}$$

□

Remark 2.2 Lemma 2.1 can apparently be extended to higher dimensional cases.

Suppose $\Lambda \in \bar{\mathcal{S}}$. We choose an $X^\Lambda = (X_1^\Lambda, X_2^\Lambda, X_3^\Lambda, X_4^\Lambda) \in \mathcal{G}$ with covariance matrix Λ . Regard $X_1^\Lambda, X_2^\Lambda, X_3^\Lambda, X_4^\Lambda$ as vectors of the space $L^2(\Omega, \mathcal{F}, P)$ and denote by \mathcal{V} the volume of the tetrahedron with vertexes $\{X_1^\Lambda, X_2^\Lambda, X_3^\Lambda, X_4^\Lambda\}$. Define

$$\tilde{A} = 6\sqrt{2}\mathcal{V}.$$

Note that $(X_1^\Lambda - X_2^\Lambda, X_3^\Lambda - X_2^\Lambda, X_4^\Lambda - X_2^\Lambda)$ is a 3-dimensional centered Gaussian random vector with covariance matrix

$$\Sigma_2^\Lambda := \begin{pmatrix} 2(1 - \Lambda_{12}) & 1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23} & 1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24} \\ 1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23} & 2(1 - \Lambda_{23}) & 1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34} \\ 1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24} & 1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34} & 2(1 - \Lambda_{24}) \end{pmatrix}. \quad (2.5)$$

By the volume formula of a tetrahedron, we get

$$\mathcal{V} = \frac{1}{6} \sqrt{\det(\Sigma_2^\Lambda)}.$$

Hence

$$\tilde{A} = \sqrt{2\det(\Sigma_2^\Lambda)}. \quad (2.6)$$

Let Σ_k^Λ be the covariance matrix of the Gaussian random vector $(X_{l_1}^\Lambda - X_k^\Lambda, X_{l_2}^\Lambda - X_k^\Lambda, X_{l_3}^\Lambda - X_k^\Lambda)$, where $l_1 < l_2 < l_3$ is the increasing arrangement of $\{1, 2, 3, 4\} \setminus \{k\}$. Similar to (2.6), we can show that

$$\tilde{A} = \sqrt{2\det(\Sigma_k^\Lambda)}, \quad k \in \{1, 2, 3, 4\}.$$

Define

$$(\Lambda')_{kl} = 1 - \Lambda_{kl}, \quad k < l,$$

and

$$\begin{aligned} (\tilde{\Lambda})_{12} &= (1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24})(1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23}) - 2(1 - \Lambda_{12})(1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34}) \\ &= (\Lambda')_{12}^2 - (\Lambda')_{12}[(\Lambda')_{13} + (\Lambda')_{14} + (\Lambda')_{23} + (\Lambda')_{24} - 2(\Lambda')_{34}] \\ &\quad + [(\Lambda')_{13} - (\Lambda')_{23}] \cdot [(\Lambda')_{14} - (\Lambda')_{24}]. \end{aligned} \quad (2.7)$$

Similarly, we define $(\tilde{\Lambda})_{kl}$ for $k < l$ by replacing 1 in (2.7) with k , 2 in (2.7) with l , 3 in (2.7) with $\min\{s : 1 \leq s \leq 4, s \neq k, l\}$, and 4 in (2.7) with $\max\{s : 1 \leq s \leq 4, s \neq k, l\}$.

Suppose $\Lambda \in \mathcal{S}$. We have

$$\begin{aligned} &\begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} (\Lambda^{-1})_{11} & (\Lambda^{-1})_{12} & (\Lambda^{-1})_{13} & (\Lambda^{-1})_{14} \\ (\Lambda^{-1})_{21} & (\Lambda^{-1})_{22} & (\Lambda^{-1})_{23} & (\Lambda^{-1})_{24} \\ (\Lambda^{-1})_{31} & (\Lambda^{-1})_{32} & (\Lambda^{-1})_{33} & (\Lambda^{-1})_{34} \\ (\Lambda^{-1})_{41} & (\Lambda^{-1})_{42} & (\Lambda^{-1})_{43} & (\Lambda^{-1})_{44} \end{pmatrix} \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} (\Lambda^{-1})_{11} & \sum_{l=1}^4 (\Lambda^{-1})_{1l} & (\Lambda^{-1})_{13} & (\Lambda^{-1})_{14} \\ \sum_{l=1}^4 (\Lambda^{-1})_{1l} & \sum_{k,l=1}^4 (\Lambda^{-1})_{kl} & \sum_{l=1}^4 (\Lambda^{-1})_{3l} & \sum_{l=1}^4 (\Lambda^{-1})_{4l} \\ (\Lambda^{-1})_{31} & \sum_{l=1}^4 (\Lambda^{-1})_{3l} & (\Lambda^{-1})_{33} & (\Lambda^{-1})_{34} \\ (\Lambda^{-1})_{41} & \sum_{l=1}^4 (\Lambda^{-1})_{4l} & (\Lambda^{-1})_{43} & (\Lambda^{-1})_{44} \end{pmatrix}. \end{aligned} \quad (2.8)$$

Then,

$$\sum_{k,l=1}^4 (\Lambda^{-1})_{kl} > 0.$$

Define

$$A = \sqrt{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}.$$

By (2.8), we get

$$\begin{aligned}
& \left(\begin{array}{cccc} (\Lambda^{-1})_{11} & \sum_{l=1}^4 (\Lambda^{-1})_{1l} & (\Lambda^{-1})_{13} & (\Lambda^{-1})_{14} \\ \sum_{l=1}^4 (\Lambda^{-1})_{1l} & A^2 & \sum_{l=1}^4 (\Lambda^{-1})_{3l} & \sum_{l=1}^4 (\Lambda^{-1})_{4l} \\ (\Lambda^{-1})_{31} & \sum_{l=1}^4 (\Lambda^{-1})_{3l} & (\Lambda^{-1})_{33} & (\Lambda^{-1})_{34} \\ (\Lambda^{-1})_{41} & \sum_{l=1}^4 (\Lambda^{-1})_{4l} & (\Lambda^{-1})_{43} & (\Lambda^{-1})_{44} \end{array} \right)^{-1} \\
&= \left(\begin{array}{cccc} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{array} \right)^{-1} \Lambda \left(\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right)^{-1} \\
&= \left(\begin{array}{cccc} 1 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{array} \right) \Lambda \left(\begin{array}{cccc} 1 & 0 & 0 & 0 \\ -1 & 1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right) \\
&= \left(\begin{array}{cccc} 2(1 - \Lambda_{12}) & \Lambda_{12} - 1 & 1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23} & 1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24} \\ \Lambda_{12} - 1 & 1 & \Lambda_{23} - 1 & \Lambda_{24} - 1 \\ 1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23} & \Lambda_{23} - 1 & 2(1 - \Lambda_{23}) & 1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34} \\ 1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24} & \Lambda_{24} - 1 & 1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34} & 2(1 - \Lambda_{24}) \end{array} \right). \quad (2.9)
\end{aligned}$$

Then, we obtain by (2.6) and (2.9) that

$$A^2 = \frac{\det(\Sigma_2^\Lambda)}{\det(\Lambda)} = \frac{\tilde{A}^2}{2\det(\Lambda)}. \quad (2.10)$$

Theorem 2.3 For $\Lambda \in \mathcal{S}$, we have

$$\frac{\partial F}{\partial \Lambda_{kl}}(\Lambda) = -\frac{1}{4\sqrt{\pi^3(\Lambda')_{kl}}} \arccos \left(\frac{(\tilde{\Lambda})_{kl}}{\sqrt{(\Lambda')_{kl}\tilde{A}^2 + (\tilde{\Lambda})_{kl}^2}} \right), \quad \forall k < l. \quad (2.11)$$

Proof. Without loss of generality, we prove (2.11) for the case that $k = 1, l = 2$. By (2.2), we

get

$$\begin{aligned}
& -\frac{\partial F}{\partial \Lambda_{12}}(\Lambda) \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^t \int_{-\infty}^t f_{\Lambda}(t, t, x_3, x_4) dx_3 dx_4 dt \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^0 \int_{-\infty}^0 f_{\Lambda}(t, t, x_3 + t, x_4 + t) dx_3 dx_4 dt \\
&= \int_{-\infty}^0 \int_{-\infty}^0 \int_{-\infty}^{\infty} f_{\Lambda}(t, t, x_3 + t, x_4 + t) dt dx_3 dx_4 \\
&= \frac{1}{\sqrt{(2\pi)^4 \det(\Lambda)}} \int_{-\infty}^0 \int_{-\infty}^0 \int_{-\infty}^{\infty} e^{-\frac{1}{2} \left\{ \sum_{k,l=1}^4 (\Lambda^{-1})_{kl} t^2 + 2 \left[\sum_{l=1}^4 (\Lambda^{-1})_{3l} x_3 + \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4 \right] t + [(\Lambda^{-1})_{33} x_3^2 + 2(\Lambda^{-1})_{34} x_3 x_4 + (\Lambda^{-1})_{44} x_4^2] \right\}} dt dx_3 dx_4 \\
&= \frac{1}{\sqrt{(2\pi)^4 \det(\Lambda)}} \int_{-\infty}^0 \int_{-\infty}^0 \int_{-\infty}^{\infty} e^{-\frac{1}{2} \left\{ \sqrt{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} t + \frac{\sum_{l=1}^4 (\Lambda^{-1})_{3l} x_3 + \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4}{\sqrt{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \right\}^2} dt \\
&\quad e^{-\frac{1}{2} \left\{ [(\Lambda^{-1})_{33} x_3^2 + 2(\Lambda^{-1})_{34} x_3 x_4 + (\Lambda^{-1})_{44} x_4^2] - \frac{[\sum_{l=1}^4 (\Lambda^{-1})_{3l} x_3 + \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4]^2}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} \right\}} dx_3 dx_4 \\
&= \frac{1}{\sqrt{(2\pi)^3 \det(\Lambda) \sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \int_{-\infty}^0 \int_{-\infty}^0 \\
&\quad e^{-\frac{1}{2} \left\{ [(\Lambda^{-1})_{33} x_3^2 + 2(\Lambda^{-1})_{34} x_3 x_4 + (\Lambda^{-1})_{44} x_4^2] - \frac{[\sum_{l=1}^4 (\Lambda^{-1})_{3l} x_3 + \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4]^2}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} \right\}} dx_3 dx_4 \\
&= \frac{1}{\sqrt{(2\pi)^3 \det(\Lambda) \sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \int_0^{\infty} \int_0^{\infty} \\
&\quad e^{-\frac{1}{2} \left\{ [(\Lambda^{-1})_{33} x_3^2 + 2(\Lambda^{-1})_{34} x_3 x_4 + (\Lambda^{-1})_{44} x_4^2] - \frac{[\sum_{l=1}^4 (\Lambda^{-1})_{3l} x_3 + \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4]^2}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} \right\}} dx_3 dx_4 \\
&:= \frac{1}{\sqrt{(2\pi)^3 \det(\Lambda) \sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \int_0^{\infty} \int_0^{\infty} e^{-\frac{a_1 x_3^2 + b_1 x_4^2 + 2c_1 x_3 x_4}{2A^2}} dx_3 dx_4, \tag{2.12}
\end{aligned}$$

where

$$a_1 := (\Lambda^{-1})_{33} A^2 - \left(\sum_{l=1}^4 (\Lambda^{-1})_{3l} \right)^2, \quad b_1 := (\Lambda^{-1})_{44} A^2 - \left(\sum_{l=1}^4 (\Lambda^{-1})_{4l} \right)^2,$$

and

$$c_1 := (\Lambda^{-1})_{34} A^2 - \left(\sum_{l=1}^4 (\Lambda^{-1})_{3l} \right) \left(\sum_{l=1}^4 (\Lambda^{-1})_{4l} \right). \tag{2.13}$$

Set

$$t^* = -\frac{\sum_{l=1}^4 (\Lambda^{-1})_{3l} x_3 + \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4}{A^2}.$$

Then,

$$\begin{aligned}
& \frac{a_1 x_3^2 + b_1 x_4^2 + 2c_1 x_3 x_4}{A^2} \\
&= [(\Lambda^{-1})_{33} x_3^2 + 2(\Lambda^{-1})_{34} x_3 x_4 + (\Lambda^{-1})_{44} x_4^2] - \frac{[\sum_{l=1}^4 (\Lambda^{-1})_{3l} x_3 + \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4]^2}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} \\
&= (t^*, t^*, x_3 + t^*, x_4 + t^*)(\Lambda^{-1})(t^*, t^*, x_3 + t^*, x_4 + t^*)^T.
\end{aligned}$$

Note that $(x_3, x_4) \neq 0$ implies $(t^*, t^*, x_3 + t^*, x_4 + t^*) \neq 0$. Hence, we have

$$a_1 b_1 - c_1^2 > 0. \quad (2.14)$$

By (2.12) and Lemma 7.1 of the Appendix, we get

$$\int_{-\infty}^{\infty} \int_{-\infty}^t \int_{-\infty}^t f_{\Lambda}(t, t, x_3, x_4) dx_3 dx_4 dt = \frac{A}{\sqrt{(2\pi)^3 \det(\Lambda)(a_1 b_1 - c_1^2)}} \arccos\left(\frac{c_1}{\sqrt{a_1 b_1}}\right). \quad (2.15)$$

Define

$$\Pi = \begin{pmatrix} A^2 & \sum_{l=1}^4 (\Lambda^{-1})_{3l} & \sum_{l=1}^4 (\Lambda^{-1})_{4l} \\ \sum_{l=1}^4 (\Lambda^{-1})_{3l} & (\Lambda^{-1})_{33} & (\Lambda^{-1})_{34} \\ \sum_{l=1}^4 (\Lambda^{-1})_{4l} & (\Lambda^{-1})_{34} & (\Lambda^{-1})_{44} \end{pmatrix}. \quad (2.16)$$

Then, we obtain by (2.8) and (2.9) that

$$\det(\Lambda) \cdot \det(\Pi) = \frac{\det(\Pi)}{\det(\Lambda^{-1})} = 2(1 - \Lambda_{12}). \quad (2.17)$$

Thus,

$$\begin{aligned}
& \det(\Lambda)(a_1 b_1 - c_1^2) \\
&= \det(\Lambda) [((\Lambda^{-1})_{33}(\Lambda^{-1})_{44} - (\Lambda^{-1})_{34}^2)A^4 \\
&\quad - \left((\Lambda^{-1})_{33} \left(\sum_{l=1}^4 (\Lambda^{-1})_{4l} \right)^2 + (\Lambda^{-1})_{44} \left(\sum_{l=1}^4 (\Lambda^{-1})_{3l} \right)^2 - 2(\Lambda^{-1})_{34} \left(\sum_{l=1}^4 (\Lambda^{-1})_{3l} \right) \left(\sum_{l=1}^4 (\Lambda^{-1})_{4l} \right) \right) A^2] \\
&= A^2 \det(\Lambda) \cdot \det(\Pi) \\
&= 2(1 - \Lambda_{12})A^2,
\end{aligned} \quad (2.18)$$

which implies that

$$\frac{A}{\sqrt{(2\pi)^3 \det(\Lambda)(a_1 b_1 - c_1^2)}} = \frac{1}{4\sqrt{\pi^3(1 - \Lambda_{12})}} = \frac{1}{4\sqrt{\pi^3(\Lambda')_{12}}}. \quad (2.19)$$

Recall the block matrix inversion formula

$$\begin{pmatrix} S & T \\ U & V \end{pmatrix}^{-1} = \begin{pmatrix} S^{-1} + S^{-1}T(V - US^{-1}T)^{-1}US^{-1} & -S^{-1}T(V - US^{-1}T)^{-1} \\ -(V - US^{-1}T)^{-1}US^{-1} & (V - US^{-1}T)^{-1} \end{pmatrix}.$$

By (2.9), we get

$$\begin{aligned} \Pi = & \left[\begin{pmatrix} 1 & \Lambda_{23} - 1 & \Lambda_{24} - 1 \\ \Lambda_{23} - 1 & 2(1 - \Lambda_{23}) & 1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34} \\ \Lambda_{24} - 1 & 1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34} & 2(1 - \Lambda_{24}) \end{pmatrix} \right. \\ & \left. - \begin{pmatrix} \Lambda_{12} - 1 \\ 1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23} \\ 1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24} \end{pmatrix} [2(1 - \Lambda_{12})]^{-1} \begin{pmatrix} \Lambda_{12} - 1 \\ 1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23} \\ 1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24} \end{pmatrix}^T \right]^{-1}. \end{aligned} \quad (2.20)$$

Then, we obtain by (2.7), (2.13), (2.16), (2.17) and (2.20) that

$$\begin{aligned} c_1 &= -\det(\Pi) \cdot (\Pi^{-1})_{32} \\ &= -\frac{2(1 - \Lambda_{12})}{\det(\Lambda)} \cdot \left[(1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34}) \right. \\ &\quad \left. - \frac{(1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24})(1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23})}{2(1 - \Lambda_{12})} \right] \\ &= \frac{1}{\det(\Lambda)} [(1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24})(1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23}) \\ &\quad - 2(1 - \Lambda_{12})(1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34})] \\ &= \frac{(\tilde{\Lambda})_{12}}{\det(\Lambda)}. \end{aligned} \quad (2.21)$$

Further, by (2.10), (2.14), (2.18) and (2.21), we get

$$\begin{aligned} 0 &< a_1 b_1 \\ &= (a_1 b_1 - c_1^2) + c_1^2 \\ &= \frac{2\det(\Lambda)(\Lambda')_{12}A^2 + (\tilde{\Lambda})_{12}^2}{[\det(\Lambda)]^2} \\ &= \frac{(\Lambda')_{12}\tilde{A}^2 + (\tilde{\Lambda})_{12}^2}{[\det(\Lambda)]^2}, \end{aligned} \quad (2.22)$$

which together with (2.21) implies that

$$\arccos\left(\frac{c_1}{\sqrt{a_1 b_1}}\right) = \arccos\left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12}\tilde{A}^2 + (\tilde{\Lambda})_{12}^2}}\right). \quad (2.23)$$

Therefore, we obtain by (2.15), (2.19) and (2.23) that

$$\int_{-\infty}^{\infty} \int_{-\infty}^t \int_{-\infty}^t f_{\Lambda}(t, t, x_3, x_4) dx_3 dx_4 dt = \frac{1}{4\sqrt{\pi^3(\Lambda')_{12}}} \arccos\left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12}\tilde{A}^2 + (\tilde{\Lambda})_{12}^2}}\right). \quad (2.24)$$

□

3 Second order partial derivatives and formula of $F(\Lambda)$

Theorem 3.1 For $\Lambda \in \mathcal{S}$, we have

$$\begin{aligned} & \frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{12}}(\Lambda) \\ = & -\frac{1}{8\sqrt{\pi^3}(\Lambda')_{12}^3} \arccos \left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12}\tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} \right) + \frac{1}{4\sqrt{\pi^3}(\Lambda')_{12}\tilde{A}} \\ & \cdot \left[\frac{(\tilde{\Lambda})_{13}[(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]}{4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2} + \frac{(\tilde{\Lambda})_{14}[(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]}{4(\Lambda')_{12}(\Lambda')_{24} - [(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]^2} \right], \end{aligned} \quad (3.1)$$

$$\frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{13}}(\Lambda) = -\frac{(\tilde{\Lambda})_{23}}{2\sqrt{\pi^3}\tilde{A}\{4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2\}},$$

and

$$\frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{34}}(\Lambda) = -\frac{1}{2\sqrt{\pi^3}\tilde{A}}.$$

Similar formulas hold for all $k < l$ if we replace 1 with k , 2 with l , 3 with $\min\{s : 1 \leq s \leq 4, s \neq k, l\}$, and 4 with $\max\{s : 1 \leq s \leq 4, s \neq k, l\}$.

Proof. By (2.2), (2.5)–(2.7), (2.24) and (7.6) of the Appendix, we get

$$\begin{aligned}
& -\frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{12}}(\Lambda) \\
&= \frac{\partial}{\partial \Lambda_{12}} \left\{ \int_{-\infty}^{\infty} \int_{-\infty}^t \int_{-\infty}^t f_{\Lambda}(t, t, x_3, x_4) dx_3 dx_4 dt \right\} \\
&= \frac{1}{8\sqrt{\pi^3}(\Lambda')_{12}^3} \arccos \left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} \right) \\
&\quad - \frac{\sqrt{(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2}}{4\sqrt{\pi^3}(\Lambda')_{12} \tilde{A}} \frac{\partial}{\partial \Lambda_{12}} \left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} \right) \\
&= \frac{1}{8\sqrt{\pi^3}(\Lambda')_{12}^3} \arccos \left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} \right) \\
&\quad - \frac{1}{4\sqrt{\pi^3}(\Lambda')_{12} \tilde{A}} \cdot \left\{ -2(\Lambda')_{12} + (\Lambda')_{13} + (\Lambda')_{14} + (\Lambda')_{23} + (\Lambda')_{24} - 2(\Lambda')_{34} \right. \\
&\quad \left. + \frac{(\tilde{\Lambda})_{12}[(\Lambda')_{13} + (\Lambda')_{23} - (\Lambda')_{12}]}{4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2} + \frac{(\tilde{\Lambda})_{12}[(\Lambda')_{14} + (\Lambda')_{24} - (\Lambda')_{12}]}{4(\Lambda')_{12}(\Lambda')_{24} - [(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]^2} \right\} \\
&= \frac{1}{8\sqrt{\pi^3}(\Lambda')_{12}^3} \arccos \left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} \right) - \frac{1}{4\sqrt{\pi^3}(\Lambda')_{12} \tilde{A}} \\
&\quad \cdot \left[\frac{(\tilde{\Lambda})_{13}[(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]}{4(\Lambda')_{12}(\Lambda')_{23} + [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2} + \frac{(\tilde{\Lambda})_{14}[(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]}{4(\Lambda')_{12}(\Lambda')_{24} - [(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]^2} \right].
\end{aligned}$$

By (2.16), (2.17) and (2.20), we have that

$$\begin{aligned}
& (\Lambda^{-1})_{44} A^2 - \left(\sum_{l=1}^4 (\Lambda^{-1})_{4l} \right)^2 \\
&= \det(\Pi) \cdot (\Pi^{-1})_{22} \\
&= \frac{2(1 - \Lambda_{12})}{\det(\Lambda)} \cdot \left[2(1 - \Lambda_{23}) - \frac{(1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23})^2}{2(1 - \Lambda_{12})} \right] \\
&= \frac{1}{\det(\Lambda)} [4(1 - \Lambda_{12})(1 - \Lambda_{23}) - (1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23})^2] \\
&= \frac{1}{\det(\Lambda)} [4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2]. \tag{3.2}
\end{aligned}$$

By (2.21) and symmetry, we get

$$(\Lambda^{-1})_{14} A^2 - \left(\sum_{l=1}^4 (\Lambda^{-1})_{1l} \right) \left(\sum_{l=1}^4 (\Lambda^{-1})_{4l} \right) = \frac{(\tilde{\Lambda})_{23}}{\det(\Lambda)}. \tag{3.3}$$

Then, we obtain by (2.1), (2.2), (2.4), (2.10), (3.2) and (3.3) that

$$\begin{aligned}
& -\frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{13}}(\Lambda) \\
&= \frac{\partial}{\partial \Lambda_{13}} \left\{ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^t f_{\Lambda}(t, t, x_3, x_4) dx_3 dx_4 dt \right\} \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^t (\partial_1 f_{\Lambda})(t, t, t, x_4) dx_4 dt \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^0 (\partial_1 f_{\Lambda})(t, t, t, x_4 + t) dx_4 dt \\
&= \int_{-\infty}^0 \int_{-\infty}^{\infty} (\partial_1 f_{\Lambda})(t, t, t, x_4 + t) dt dx_4 \\
&= \frac{-1}{\sqrt{(2\pi)^4 \det(\Lambda)}} \int_{-\infty}^0 \int_{-\infty}^{\infty} \left[\sum_{l=1}^4 (\Lambda^{-1})_{1l} t + (\Lambda^{-1})_{14} x_4 \right] \\
&\quad \cdot e^{-\frac{1}{2} \{ \sum_{k,l=1}^4 (\Lambda^{-1})_{kl} t^2 + 2 \sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4 t + (\Lambda^{-1})_{44} x_4^2 \}} dt dx_4 \\
&= \frac{-1}{\sqrt{(2\pi)^4 \det(\Lambda)}} \int_{-\infty}^0 \int_{-\infty}^{\infty} \left[\sum_{l=1}^4 (\Lambda^{-1})_{1l} t + (\Lambda^{-1})_{14} x_4 \right] e^{-\frac{1}{2} \left\{ \sqrt{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} t + \frac{\sum_{l=1}^4 (\Lambda^{-1})_{4l} x_4}{\sqrt{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \right\}^2} dt \\
&\quad \cdot e^{-\frac{1}{2} \left\{ \left[(\Lambda^{-1})_{44} - \frac{(\sum_{l=1}^4 (\Lambda^{-1})_{4l})^2}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} \right] x_4^2 \right\}} dx_4 \\
&= \frac{-1}{\sqrt{(2\pi)^3 \det(\Lambda) \sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \int_{-\infty}^0 \left[\left((\Lambda^{-1})_{14} - \frac{(\sum_{l=1}^4 (\Lambda^{-1})_{1l})(\sum_{l=1}^4 (\Lambda^{-1})_{4l})}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} \right) x_4 \right] \\
&\quad \cdot e^{-\frac{1}{2} \left\{ \left[(\Lambda^{-1})_{44} - \frac{(\sum_{l=1}^4 (\Lambda^{-1})_{4l})^2}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}} \right] x_4^2 \right\}} dx_4 \\
&= \frac{1}{\sqrt{(2\pi)^3 \det(\Lambda) \sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \left[\frac{(\Lambda^{-1})_{14} - \frac{(\sum_{l=1}^4 (\Lambda^{-1})_{1l})(\sum_{l=1}^4 (\Lambda^{-1})_{4l})}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}}{(\Lambda^{-1})_{44} - \frac{(\sum_{l=1}^4 (\Lambda^{-1})_{4l})^2}{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \right] \\
&= \frac{(\tilde{\Lambda})_{23}}{2\sqrt{\pi^3 \tilde{A} \{ 4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2 \}}.
\end{aligned}$$

By (2.1), (2.2), (2.4) and (2.10), we get

$$\begin{aligned}
-\frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{34}}(\Lambda) &= \frac{\partial}{\partial \Lambda_{34}} \left\{ \int_{-\infty}^{\infty} \int_{-\infty}^t \int_{-\infty}^t f_{\Lambda}(t, t, x_3, x_4) dx_3 dx_4 dt \right\} \\
&= \int_{-\infty}^{\infty} f_{\Lambda}(t, t, t, t) dt \\
&= \frac{1}{\sqrt{(2\pi)^4 \det(\Lambda)}} \int_{-\infty}^{\infty} e^{-\frac{\sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}{2} t^2} dt \\
&= \frac{1}{\sqrt{(2\pi)^3 \det(\Lambda) \sum_{k,l=1}^4 (\Lambda^{-1})_{kl}}} \\
&= \frac{1}{2\sqrt{\pi^3 \tilde{A}}}.
\end{aligned}$$

□

Define

$$\mathcal{S}_1 = \{\Lambda \in \bar{\mathcal{S}} : (\Lambda')_{kl} \tilde{A}^2 + (\tilde{\Lambda})_{kl}^2 > 0 \text{ for all } k < l\},$$

and

$$\mathcal{S}_2 = \{\Lambda \in \bar{\mathcal{S}} : \Lambda_{kl} \neq 1 \text{ for all } k < l\}.$$

Lemma 3.2 *We have*

$$\mathcal{S}_1 = \mathcal{S}_2.$$

Proof. Suppose that $\Lambda \notin \mathcal{S}_2$. Then, $\Lambda_{kl} = 1$ for some $k < l$. We assume without loss of generality that $\Lambda_{12} = 1$. By (2.7), we get $(\tilde{\Lambda})_{12} = 0$. Thus, $(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2 = 0$, which implies that $\Lambda \notin \mathcal{S}_1$.

Suppose that $\Lambda \notin \mathcal{S}_1$. Then, $(\Lambda')_{kl} \tilde{A}^2 + (\tilde{\Lambda})_{kl}^2 = 0$ for some $k < l$. We assume without loss of generality that $(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2 = 0$. Take an $X^{\Lambda} \in \mathcal{G}$ with covariance matrix Λ . By (7.3)–(7.5) of the Appendix, we find that either $(X_1^{\Lambda} - X_2^{\Lambda}, X_3^{\Lambda} - X_2^{\Lambda})$ or $(X_1^{\Lambda} - X_2^{\Lambda}, X_4^{\Lambda} - X_2^{\Lambda})$ is a degenerate 2-dimensional Gaussian random vector. Without loss of generality, we assume that $(X_1^{\Lambda} - X_2^{\Lambda}, X_3^{\Lambda} - X_2^{\Lambda})$ is degenerate. Then, there exist constants α and β such that $|\alpha| + |\beta| > 0$ and

$$\alpha(X_1^{\Lambda} - X_2^{\Lambda}) + \beta(X_3^{\Lambda} - X_2^{\Lambda}) = 0.$$

If $\beta = 0$, then $X_1^{\Lambda} = X_2^{\Lambda}$ and thus $\Lambda_{12} = 1$; if $\alpha = 0$, then $X_2^{\Lambda} = X_3^{\Lambda}$ and thus $\Lambda_{23} = 1$. If $\alpha\beta \neq 0$ and $\alpha + \beta = 0$, then $X_1^{\Lambda} = X_2^{\Lambda}$ and thus $\Lambda_{12} = 1$. If $\alpha\beta \neq 0$ and $\alpha + \beta \neq 0$, we get

$$X_2^{\Lambda} = \frac{\alpha}{\alpha + \beta} X_1^{\Lambda} + \frac{\beta}{\alpha + \beta} X_3^{\Lambda}. \quad (3.4)$$

By $E[(X_1^{\Lambda})^2] = E[(X_2^{\Lambda})^2] = E[(X_3^{\Lambda})^2] = 1$ and (3.4), we get

$$1 = \left(\frac{\alpha}{\alpha + \beta} \right)^2 + \left(\frac{\beta}{\alpha + \beta} \right)^2 + \frac{2\alpha\beta}{(\alpha + \beta)^2} \Lambda_{13},$$

which implies that $\Lambda_{13} = 1$. Hence $\Lambda \notin \mathcal{S}_2$.

□

Theorem 3.3 (i) Suppose that $\Lambda \in \mathcal{S}_1$. We have

$$F(\Lambda) = \frac{1}{2\sqrt{\pi^3}} \sum_{k < l} \sqrt{(\Lambda')_{kl}} \arccos \left(\frac{(\tilde{\Lambda})_{kl}}{\sqrt{(\Lambda')_{kl} \tilde{A}^2 + (\tilde{\Lambda})_{kl}^2}} \right). \quad (3.5)$$

(ii) Suppose that $\Lambda_{14} = 1$. We have

$$F(\Lambda) = \frac{1}{2\sqrt{\pi}} \left[\sqrt{(\Lambda')_{12}} + \sqrt{(\Lambda')_{13}} + \sqrt{(\Lambda')_{23}} \right]. \quad (3.6)$$

Proof. (i) We first show that (3.5) holds for $\Lambda \in \mathcal{S}$. Denote the right hand side of (3.5) by $\tilde{F}(\Lambda)$. Let Λ^* be the 4×4 matrix with all elements equal to 1. It is easy to see that $F(\Lambda^*) = \tilde{F}(\Lambda^*) = 0$ and \tilde{F} is continuous at Λ^* . By virtue of characteristic functions, we can show that F is continuous on $\bar{\mathcal{S}}$. Since both F and \tilde{F} are differentiable on \mathcal{S} , to prove that (3.5) holds on \mathcal{S} , we need only show that F and \tilde{F} have the same partial derivatives on \mathcal{S} . Without loss of generality, in the sequel, we show that

$$\frac{\partial F}{\partial \Lambda_{12}} = \frac{\partial \tilde{F}}{\partial \Lambda_{12}}.$$

By (2.11), we get

$$\tilde{F}(\Lambda) = -2 \sum_{k < l} (\Lambda')_{kl} \frac{\partial F}{\partial \Lambda_{kl}}.$$

Hence we need only show that

$$\sum_{k < l} (\Lambda')_{kl} \frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{kl}} = \frac{1}{2} \frac{\partial F}{\partial \Lambda_{12}}, \quad \forall \Lambda \in \mathcal{S}. \quad (3.7)$$

We use a, b, c, d, e, f to represent $\Lambda_{12}, \Lambda_{13}, \Lambda_{14}, \Lambda_{23}, \Lambda_{24}, \Lambda_{34}$, respectively. By (2.22) and (7.3) of the Appendix, we have that

$$g := [4(1-a)(1-d) - (1-a-d+b)^2] \cdot [4(1-a)(1-e) - (1-a-e+c)^2] > 0.$$

By (2.11) and Theorem 3.1, we find that the difference of the two sides of (3.7) multiplied by

$4\sqrt{\pi^3}\tilde{A}g$ is equal to

$$\begin{aligned}
& ((1-b)^2 - (1-b)(1-a+1-c+1-d+1-f-2(1-e)) + (a-d)(c-f))(1-a+(1-d)-(1-b)) \\
& \quad \cdot (4(1-a)(1-e) - (1-a-e+c)^2) \\
& + ((1-c)^2 - (1-c)(1-a+1-b+1-e+1-f-2(1-d)) + (a-e)(b-f))(1-a+(1-e)-(1-c)) \\
& \quad \cdot (4(1-a)(1-d) - (1-a-d+b)^2) \\
& - 2((1-d)^2 - (1-d)(1-a+1-b+1-e+1-f-2(1-c)) + (a-b)(e-f))(1-b) \\
& \quad \cdot (4(1-a)(1-e) - (1-a-e+c)^2) \\
& - 2((1-e)^2 - (1-e)(1-a+1-c+1-d+1-f-2(1-b)) + (a-c)(d-f))(1-c) \\
& \quad \cdot (4(1-a)(1-d) - (1-a-d+b)^2) \\
& - 2((1-b)^2 - (1-b)(1-a+1-c+1-d+1-f-2(1-e)) + (a-d)(c-f))(1-d) \\
& \quad \cdot (4(1-a)(1-e) - (1-a-e+c)^2) \\
& - 2((1-c)^2 - (1-c)(1-a+1-b+1-e+1-f-2(1-d)) + (a-e)(b-f))(1-e) \\
& \quad \cdot (4(1-a)(1-d) - (1-a-d+b)^2) \\
& - 2(1-f)(4(1-a)(1-d) - (1-a-d+b)^2)(4(1-a)(1-e) - (1-a-e+c)^2).
\end{aligned}$$

By direct calculation or more quickly by using Mathematica, we can show that the above multivariate polynomial of (a, b, \dots, f) is equal to 0. We refer the reader to §7.3 for the Mathematica code. Then, (3.7) holds and thus (3.5) holds for $\Lambda \in \mathcal{S}$.

Since F is continuous on $\bar{\mathcal{S}}$ and \tilde{F} is continuous on \mathcal{S}_1 . By the density of \mathcal{S} in $\bar{\mathcal{S}}$, we conclude that (3.5) holds for all $\Lambda \in \mathcal{S}_1$.

(ii) Suppose that $\Lambda_{14} = 1$. We have

$$\begin{aligned}
F(\Lambda) &= E[\max(X_1^\Lambda, X_2^\Lambda, X_3^\Lambda, X_4^\Lambda)] \\
&= E[\max(X_1^\Lambda, X_2^\Lambda, X_3^\Lambda)] \\
&= E[\max(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda, 0)] \\
&= \frac{E[\max(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)] + E[|\max(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)|]}{2}, \tag{3.8}
\end{aligned}$$

and

$$\begin{aligned}
& E[\max(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)] \\
&= \frac{E[(X_1^\Lambda - X_3^\Lambda) + (X_2^\Lambda - X_3^\Lambda)] + E[|(X_1^\Lambda - X_3^\Lambda) - (X_2^\Lambda - X_3^\Lambda)|]}{2} \\
&= \frac{E[|X_1^\Lambda - X_2^\Lambda|]}{2} \\
&= \sqrt{\frac{(\Lambda')_{12}}{\pi}}. \tag{3.9}
\end{aligned}$$

Note that $(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)$ and $(X_3^\Lambda - X_1^\Lambda, X_3^\Lambda - X_2^\Lambda)$ have the same distribution. Then,

$$\begin{aligned}
& E[|\max(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)|] \\
&= \frac{E[|\max(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)|] + E[|\max(X_3^\Lambda - X_1^\Lambda, X_3^\Lambda - X_2^\Lambda)|]}{2} \\
&= \frac{E[|\max(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)|] + E[|\min(X_1^\Lambda - X_3^\Lambda, X_2^\Lambda - X_3^\Lambda)|]}{2} \\
&= \frac{E[|X_1^\Lambda - X_3^\Lambda|] + E[|X_2^\Lambda - X_3^\Lambda|]}{2} \\
&= \sqrt{\frac{(\Lambda')_{13}}{\pi}} + \sqrt{\frac{(\Lambda')_{23}}{\pi}}.
\end{aligned} \tag{3.10}$$

Therefore, (3.6) holds by (3.8)–(3.10). \square

Remark 3.4 Note that if $\Lambda \in \bar{\mathcal{S}}$ satisfying $\Lambda_{kl} = r$ for all $k < l$ and some $r \in \mathbb{R}$, then $-\frac{1}{3} \leq r \leq 1$. If $-\frac{1}{3} \leq r < 1$, by (3.5), (7.1)–(7.3) of the Appendix, and elementary calculation, we get

$$F(\Lambda) = \frac{3\sqrt{1-r} \arccos\left(-\frac{1}{3}\right)}{\sqrt{\pi^3}}. \tag{3.11}$$

Obviously, (3.11) also holds for the case that $r = 1$.

Denote by \mathcal{G}_3 the set of all three-dimensional centered Gaussian random vectors $X = (X_1, X_2, X_3)$ with $E[X_1^2] = E[X_2^2] = E[X_3^2] = 1$. Define

$$\mathcal{M}(X) = E[\max(X_1, X_2, X_3)], \quad X \in \mathcal{G}_3.$$

We use $\bar{\mathcal{S}}_3$ to denote the set of all nonnegative-definite 3×3 symmetric real matrices with all diagonal elements equal to 1. For $\Lambda \in \bar{\mathcal{S}}_3$, take an $X^\Lambda \in \mathcal{G}_3$ with covariance matrix Λ and define

$$F(\Lambda) = \mathcal{M}(X^\Lambda).$$

Corollary 3.5 The function \mathcal{M} defined on \mathcal{G}_3 attains its maximum at X if and only if $\Lambda_{kl}^X = -\frac{1}{2}$ for all $k < l$.

Proof. By (3.6), we know that $F(\Lambda)$ is a strictly concave function defined on $\bar{\mathcal{S}}_3$. Let $\Lambda \in \bar{\mathcal{S}}_3$. Define $\bar{\Lambda} \in \bar{\mathcal{S}}_3$ by

$$\bar{\Lambda}_{kl} = \begin{cases} 1, & \text{if } k = l, \\ \frac{\sum_{k < l} \Lambda_{kl}}{3}, & \text{otherwise.} \end{cases}$$

Then, we have that $F(\Lambda) \leq F(\bar{\Lambda})$ and the equality holds if and only if $\Lambda = \bar{\Lambda}$. Note that if $\Lambda \in \bar{\mathcal{S}}_3$ satisfying $\Lambda_{kl} = r$ for all $k < l$ and some $r \in \mathbb{R}$, then $-\frac{1}{2} \leq r \leq 1$. Therefore, the proof is complete by (3.6). \square

4 Proof of Theorem 1.1

Step 1. First, we show that if $\mathcal{M}(X)$ attains its maximum at $X \in \mathcal{G}$ then X is degenerate, i.e., X_1, X_2, X_3, X_4 are linear dependent. In fact, if X is non-degenerate then $\Lambda^X \in \mathcal{S}$. To simplify notation, in the sequel, we denote Λ^X by Λ . For $\varepsilon > 0$, define Λ^ε by

$$\Lambda_{kl}^\varepsilon = \begin{cases} \Lambda_{12} - \varepsilon, & \text{if } (k, l) = (1, 2) \text{ or } (2, 1), \\ \Lambda_{kl}, & \text{otherwise.} \end{cases}$$

If ε is small enough, then $\Lambda^\varepsilon \in \mathcal{S}$. By Lemma 2.1, we get $F(\Lambda) < F(\Lambda^\varepsilon)$ and hence arrive at a contradiction.

Next, we show that if $\mathcal{M}(X)$ attains its maximum at $X \in \mathcal{G}$, then $\Lambda_{kl} \neq 1$ for any $k < l$. Assume that, for example, $\Lambda_{14} = 1$. By Corollary 3.5, we get

$$F(\Lambda) \leq \frac{3\sqrt{\frac{3}{2}}}{2\sqrt{\pi}} \approx 1.036482 < 1.18862 \approx \frac{3\sqrt{1 + \frac{1}{3} \arccos(-\frac{1}{3})}}{\sqrt{\pi^3}}.$$

Hence X is not a maximum point by Remark 3.4 and we arrive at a contradiction. Thus, $\Lambda \in \mathcal{S}_1$ by Lemma 3.2.

We regard X_1, X_2, X_3, X_4 as vectors of the space $L^2(\Omega, \mathcal{F}, P)$. Denote by $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3, \mathcal{F}_4$ the facets of the tetrahedron $\{X_1, X_2, X_3, X_4\}$ with vertexes $\{X_1, X_2, X_3\}, \{X_1, X_3, X_4\}, \{X_1, X_2, X_4\}, \{X_2, X_3, X_4\}$, respectively. We use α_{ij} , $i < j$, to denote the outer dihedral angle of \mathcal{F}_i and \mathcal{F}_j . By (2.7) and (7.3) of the Appendix, we get

$$\begin{aligned} & \frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12}\tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} \\ = & \frac{(1 + \Lambda_{13} - \Lambda_{12} - \Lambda_{23})(1 + \Lambda_{14} - \Lambda_{12} - \Lambda_{24}) - 2(1 - \Lambda_{12})(1 + \Lambda_{34} - \Lambda_{23} - \Lambda_{24})}{\sqrt{[4(1 - \Lambda_{12})(1 - \Lambda_{23}) - (1 + \Lambda_{13} - \Lambda_{12} - \Lambda_{23})^2][4(1 - \Lambda_{12})(1 - \Lambda_{24}) - (1 + \Lambda_{14} - \Lambda_{12} - \Lambda_{24})^2]}} \\ = & \{\text{Cov}(X_1 - X_2, X_3 - X_2)\text{Cov}(X_1 - X_2, X_4 - X_2) - \text{Var}(X_1 - X_2)\text{Cov}(X_3 - X_2, X_4 - X_2)\} \\ & \cdot \{\text{Var}(X_1 - X_2)\text{Var}(X_3 - X_2) - [\text{Cov}(X_1 - X_2, X_3 - X_2)]^2\}^{-\frac{1}{2}} \\ & \cdot \{\text{Var}(X_1 - X_2)\text{Var}(X_4 - X_2) - [\text{Cov}(X_1 - X_2, X_4 - X_2)]^2\}^{-\frac{1}{2}}. \end{aligned} \quad (4.1)$$

Denote by $\beta_1, \beta_2, \beta_3$ the angles between $\overrightarrow{X_2X_1}$ and $\overrightarrow{X_2X_3}$, $\overrightarrow{X_2X_1}$ and $\overrightarrow{X_2X_4}$, $\overrightarrow{X_2X_3}$ and $\overrightarrow{X_2X_4}$, respectively. Then, we obtain by (4.1) that

$$\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12}\tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} = \frac{\cos \beta_1 \cos \beta_2 - \cos \beta_3}{\sin \beta_1 \sin \beta_2} = \cos \alpha_{13}. \quad (4.2)$$

Similarly, we can show that

$$\begin{aligned}
\frac{(\tilde{\Lambda})_{13}}{\sqrt{(\Lambda')_{13}\tilde{A}^2 + (\tilde{\Lambda})_{13}^2}} &= \cos \alpha_{12}, \\
\frac{(\tilde{\Lambda})_{14}}{\sqrt{(\Lambda')_{14}\tilde{A}^2 + (\tilde{\Lambda})_{14}^2}} &= \cos \alpha_{23}, \\
\frac{(\tilde{\Lambda})_{23}}{\sqrt{(\Lambda')_{23}\tilde{A}^2 + (\tilde{\Lambda})_{23}^2}} &= \cos \alpha_{14}, \\
\frac{(\tilde{\Lambda})_{24}}{\sqrt{(\Lambda')_{24}\tilde{A}^2 + (\tilde{\Lambda})_{24}^2}} &= \cos \alpha_{34}, \\
\frac{(\tilde{\Lambda})_{34}}{\sqrt{(\Lambda')_{34}\tilde{A}^2 + (\tilde{\Lambda})_{34}^2}} &= \cos \alpha_{24}.
\end{aligned} \tag{4.3}$$

Now we show that if $\mathcal{M}(X)$ attains its maximum at $X \in \mathcal{G}$, then X_1, X_2, X_3, X_4 do not lie in the same plane. Assume that X_1, X_2, X_3, X_4 lie in the same plane. Then, two outer dihedral angles of the degenerate tetrahedron $\{X_1, X_2, X_3, X_4\}$ equal 0. Hence, we obtain by Theorem 3.3 that

$$F(\Lambda^X) \leq \frac{4\pi}{2\sqrt{\pi^3}} \approx 1.128379 < 1.18862 \approx \frac{3\sqrt{1 + \frac{1}{3}} \arccos(-\frac{1}{3})}{\sqrt{\pi^3}},$$

which implies that X is not a maximum point by Remark 3.4 and we arrive at a contradiction.

Step 2. Suppose that $\mathcal{M}(X)$ attains its maximum at $X \in \mathcal{G}$. Based on the analysis of Step 1, we know that X_1, X_2, X_3, X_4 are linear dependent and do not lie in the same plane.

First, we focus on the vertex X_1 . We consider two different cases. Case 1: $X_i \neq -X_1$ for any $2 \leq i \leq 4$. Denote by $\mathcal{P}_1, \mathcal{P}_2$ and \mathcal{P}_3 the planes determined by $\{0, X_1, X_2\}, \{0, X_1, X_3\}$ and $\{0, X_1, X_4\}$, respectively. At least one of the three planes $\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3$ is different from the other two planes. We assume without loss of generality that $\mathcal{P}_1 \neq \mathcal{P}_2$ and $\mathcal{P}_1 \neq \mathcal{P}_3$. Case 2: $X_i = -X_1$ for some $2 \leq i \leq 4$, say $X_4 = -X_1$. Denote by \mathcal{P}_1 the plane determined by $\{0, X_1, X_2\}$ and by $\mathcal{P}_2 = \mathcal{P}_3$ the plane determined by $\{0, X_1, X_3, X_4\}$, respectively.

Denote $\theta = (\theta_1, \theta_2, \theta_3, \theta_4, \theta_5)$. Take three independent standard Gaussian random variables U_1, U_2, U_3 as the basis and set (X_1, X_2, X_3, X_4) to be

$$\begin{aligned}
X_1(\theta) &= (0, 0, 1)^T, \\
X_2(\theta) &= (\sin \theta_1, 0, \cos \theta_1)^T, \\
X_3(\theta) &= (\sin \theta_2 \cos \theta_4, \sin \theta_2 \sin \theta_4, \cos \theta_2)^T, \\
X_4(\theta) &= (\sin \theta_3 \cos \theta_5, \sin \theta_3 \sin \theta_5, \cos \theta_3)^T,
\end{aligned}$$

where $\theta_1 \in (0, \pi)$ denotes the angle between $\overrightarrow{0X_1}$ and $\overrightarrow{0X_2}$, $\theta_2, \theta_3 \in (0, 2\pi)$ denote the anti-clockwise angles from $\overrightarrow{0X_1}$ to $\overrightarrow{0X_3}$ and from $\overrightarrow{0X_1}$ to $\overrightarrow{0X_4}$, respectively, and $\theta_4, \theta_5 \in (0, \pi)$ denote

the anti-clockwise angles from \mathcal{P}_1 to \mathcal{P}_2 and from \mathcal{P}_1 to \mathcal{P}_3 , respectively. Then,

$$\begin{aligned}
\Lambda_{12}(\theta) &= \cos \theta_1, \\
\Lambda_{13}(\theta) &= \cos \theta_2, \\
\Lambda_{14}(\theta) &= \cos \theta_3, \\
\Lambda_{23}(\theta) &= \sin \theta_1 \sin \theta_2 \cos \theta_4 + \cos \theta_1 \cos \theta_2, \\
\Lambda_{24}(\theta) &= \sin \theta_1 \sin \theta_3 \cos \theta_5 + \cos \theta_1 \cos \theta_3, \\
\Lambda_{34}(\theta) &= \sin \theta_2 \sin \theta_3 \cos(\theta_4 - \theta_5) + \cos \theta_2 \cos \theta_3.
\end{aligned}$$

For $T \in \mathcal{S}_1$ and $k < l$, define

$$F_{kl}(T) = -\frac{1}{4\sqrt{\pi^3(T')_{kl}}} \arccos \left(\frac{(\tilde{T})_{kl}}{\sqrt{(T')_{kl}\tilde{A}^2 + (\tilde{T})_{kl}^2}} \right). \quad (4.4)$$

Then, F_{kl} is a continuous function on \mathcal{S}_1 . By (2.11), we get

$$F_{kl}(T) = \frac{\partial F}{\partial \Lambda_{kl}}(T), \quad \forall T \in \mathcal{S}.$$

For $T \in \bar{\mathcal{S}}$ and $0 < \delta < 1$, define $T^\delta = (1 - \delta)T + \delta I$, where I is the identity matrix. Then, $T^\delta \in \mathcal{S}$ and T^δ approximates T as $\delta \downarrow 0$. For $1 \leq i \leq 5$, denote by e_i the five-dimensional unit vector with the i -th component equal to 1 and the other four components equal to 0. Define

$$R(\theta) = F(\Lambda(\theta)).$$

By the continuity of the functions F_{kl} , $k < l$, we get

$$\begin{aligned}
& \frac{\partial R}{\partial \theta_i} \\
&= \lim_{\varepsilon \rightarrow 0} \frac{F[\Lambda(\theta + \varepsilon e_i)] - F[\Lambda(\theta)]}{\varepsilon} \\
&= \lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \frac{F[\Lambda^\delta(\theta + \varepsilon e_i)] - F[\Lambda^\delta(\theta)]}{\varepsilon} \\
&= \lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \frac{1}{\varepsilon} \int_0^1 dF[p\Lambda^\delta(\theta + \varepsilon e_i) + (1 - p)\Lambda^\delta(\theta)] \\
&= \sum_{k < l} \left\{ \lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \left(\int_0^1 \frac{\partial F}{\partial \Lambda_{kl}} [p\Lambda^\delta(\theta + \varepsilon e_i) + (1 - p)\Lambda^\delta(\theta)] \cdot \frac{(1 - \delta)[\Lambda_{kl}(\theta + \varepsilon e_i) - \Lambda_{kl}(\theta)]}{\varepsilon} dp \right) \right\} \\
&= \sum_{k < l} \left\{ \lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \int_0^1 \frac{\partial F}{\partial \Lambda_{kl}} [p\Lambda^\delta(\theta + \varepsilon e_i) + (1 - p)\Lambda^\delta(\theta)] dp \cdot \lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} \frac{(1 - \delta)[\Lambda_{kl}(\theta + \varepsilon e_i) - \Lambda_{kl}(\theta)]}{\varepsilon} \right\} \\
&= \sum_{k < l} F_{kl} \frac{\partial \Lambda_{kl}}{\partial \theta_i}.
\end{aligned}$$

Since $\mathcal{M}(X)$ attains its maximum at X , we get

$$\frac{\partial R}{\partial \theta_i} = 0, \quad i = 4, 5,$$

which implies that

$$\begin{aligned} F_{23} \sin \theta_1 \sin \theta_2 \sin \theta_4 + F_{34} \sin \theta_2 \sin \theta_3 \sin(\theta_4 - \theta_5) &= 0, \\ F_{24} \sin \theta_1 \sin \theta_3 \sin \theta_5 - F_{34} \sin \theta_2 \sin \theta_3 \sin(\theta_4 - \theta_5) &= 0. \end{aligned} \quad (4.5)$$

For $1 \leq i \leq 4$, let $0P_i$ be the perpendicular line from 0 to \mathcal{F}_i with $P_i \in \mathcal{F}_i$ and denote its length by \mathcal{L}_i . Note that, for $i < j$, the angle between $0P_i$ and $0P_j$ equals α_{ij} , the outer dihedral angle of \mathcal{F}_i and \mathcal{F}_j . Denote by V_1, V_2, V_3 and V_4 the volumes of the tetrahedrons with vertexes $\{0, X_1, X_2, X_3\}, \{0, X_1, X_3, X_4\}, \{0, X_1, X_2, X_4\}$ and $\{0, X_2, X_3, X_4\}$, respectively. We have

$$\begin{aligned} V_1 &= \frac{1}{6} \sin \theta_1 \cdot |\sin \theta_2| \cdot \sin \theta_4, \\ V_2 &= \frac{1}{6} |\sin \theta_2 \sin \theta_3 \sin(\theta_4 - \theta_5)|, \\ V_3 &= \frac{1}{6} \sin \theta_1 \cdot |\sin \theta_3| \cdot \sin \theta_5. \end{aligned} \quad (4.6)$$

Since each F_{kl} is strictly negative, we obtain by (4.5) and (4.6) that

$$F_{23}V_1 = F_{34}V_2 = F_{24}V_3. \quad (4.7)$$

This implies that $V_1V_2V_3 \neq 0$ since $V_1 > 0$.

Next, by focusing on the vertexes X_2, X_3, X_4 , respectively, and following the above argument, we obtain that

$$\begin{aligned} F_{14}V_3 &= F_{34}V_4 = F_{13}V_1, \\ F_{24}V_4 &= F_{12}V_1 = F_{14}V_2, \\ F_{13}V_2 &= F_{12}V_3 = F_{23}V_4, \end{aligned} \quad (4.8)$$

and $V_1V_2V_3V_4 \neq 0$. Combining (4.7) and (4.8), we get

$$\begin{aligned} V_2 &= V_1 \cdot \frac{F_{12}}{F_{14}} = V_1 \cdot \frac{F_{23}}{F_{34}}, \\ V_3 &= V_1 \cdot \frac{F_{13}}{F_{14}} = V_1 \cdot \frac{F_{23}}{F_{24}}, \\ V_4 &= V_1 \cdot \frac{F_{12}}{F_{24}} = V_1 \cdot \frac{F_{13}}{F_{34}}. \end{aligned} \quad (4.9)$$

Further, we obtain by (4.9) that

$$F_{13} = F_{12} \cdot \frac{V_3}{V_2}, \quad (4.10)$$

$$F_{14} = F_{12} \cdot \frac{V_1}{V_2}, \quad (4.11)$$

$$F_{23} = F_{12} \cdot \frac{V_3}{V_4}, \quad (4.12)$$

$$F_{24} = F_{12} \cdot \frac{V_1}{V_4}, \quad (4.13)$$

$$F_{34} = F_{12} \cdot \frac{V_1 V_3}{V_2 V_4}. \quad (4.14)$$

By (4.10) and (4.11), we get

$$\frac{F_{12}}{V_2} = \frac{F_{13}}{V_3} = \frac{F_{14}}{V_1}. \quad (4.15)$$

By the law of sines for tetrahedrons, we have that

$$\frac{|\overrightarrow{X_1 X_2}| \cdot \text{Area}(\mathcal{F}_2)}{\sin \alpha_{13}} = \frac{|\overrightarrow{X_1 X_3}| \cdot \text{Area}(\mathcal{F}_3)}{\sin \alpha_{12}} = \frac{|\overrightarrow{X_1 X_4}| \cdot \text{Area}(\mathcal{F}_1)}{\sin \alpha_{23}}, \quad (4.16)$$

where $|\cdot|$ denotes the length of a vector. Then, we obtain by (4.2)–(4.4), (4.15) and (4.16) that

$$\frac{\alpha_{13}}{\mathcal{L}_2 \sin \alpha_{13}} = \frac{\alpha_{12}}{\mathcal{L}_3 \sin \alpha_{12}} = \frac{\alpha_{23}}{\mathcal{L}_1 \sin \alpha_{23}}. \quad (4.17)$$

Further, by (4.10)–(4.14) and symmetry, we get

$$\begin{aligned} \frac{\alpha_{13}}{\mathcal{L}_4 \sin \alpha_{13}} &= \frac{\alpha_{14}}{\mathcal{L}_3 \sin \alpha_{14}} = \frac{\alpha_{34}}{\mathcal{L}_1 \sin \alpha_{34}}, \\ \frac{\alpha_{12}}{\mathcal{L}_4 \sin \alpha_{12}} &= \frac{\alpha_{14}}{\mathcal{L}_2 \sin \alpha_{14}} = \frac{\alpha_{24}}{\mathcal{L}_1 \sin \alpha_{24}}, \\ \frac{\alpha_{23}}{\mathcal{L}_4 \sin \alpha_{23}} &= \frac{\alpha_{34}}{\mathcal{L}_2 \sin \alpha_{34}} = \frac{\alpha_{24}}{\mathcal{L}_3 \sin \alpha_{24}}, \end{aligned}$$

which together with (4.17) implies that

$$\frac{\mathcal{L}_1 \mathcal{L}_2 \alpha_{12}}{\sin \alpha_{12}} = \frac{\mathcal{L}_1 \mathcal{L}_3 \alpha_{13}}{\sin \alpha_{13}} = \frac{\mathcal{L}_1 \mathcal{L}_4 \alpha_{14}}{\sin \alpha_{14}} = \frac{\mathcal{L}_2 \mathcal{L}_3 \alpha_{23}}{\sin \alpha_{23}} = \frac{\mathcal{L}_2 \mathcal{L}_4 \alpha_{24}}{\sin \alpha_{24}} = \frac{\mathcal{L}_3 \mathcal{L}_4 \alpha_{34}}{\sin \alpha_{34}}. \quad (4.18)$$

Step 3. Note that X_1 satisfies the following equations:

$$\langle X_1 - P_i, P_i \rangle = 0, \quad 1 \leq i \leq 3.$$

Then, we have that

$$\langle X_1, P_i \rangle = \|P_i\|^2 = \mathcal{L}_i^2, \quad 1 \leq i \leq 3. \quad (4.19)$$

Define the 3×3 matrix Γ by

$$\Gamma = (P_1, P_2, P_3).$$

Given a matrix M , we denote by $\text{diag}(M)$ the row vector consisting of the diagonal elements of M . Then, (4.19) becomes

$$X_1^T \Gamma = \text{diag}(\Gamma^T \Gamma),$$

which implies that

$$X_1^T = \text{diag}(\Gamma^T \Gamma) \cdot \Gamma^{-1}.$$

Define

$$\Sigma = \Gamma^T \Gamma.$$

Then, we obtain by $\|X_1\| = 1$ that

$$\text{diag}(\Sigma) \cdot \Sigma^{-1} \cdot [\text{diag}(\Sigma)]^T = 1. \quad (4.20)$$

Define

$$\tilde{P}_i = \frac{P_i}{\mathcal{L}_i}, \quad 1 \leq i \leq 3, \quad \tilde{\Gamma} = (\tilde{P}_1, \tilde{P}_2, \tilde{P}_3), \quad \tilde{\Sigma} = \tilde{\Gamma}^T \tilde{\Gamma}, \quad \Delta = \begin{pmatrix} \mathcal{L}_1 & 0 & 0 \\ 0 & \mathcal{L}_2 & 0 \\ 0 & 0 & \mathcal{L}_3 \end{pmatrix}.$$

We have

$$\Sigma = \Delta \tilde{\Sigma} \Delta, \quad \Sigma^{-1} = \Delta^{-1} \tilde{\Sigma}^{-1} \Delta^{-1}.$$

Note that

$$\text{diag}(\Delta M) = \text{diag}(M \Delta) = \text{diag}(M) \Delta.$$

Then,

$$\text{diag}(\Sigma) = \text{diag}(\tilde{\Sigma}) \Delta^2 = (1, 1, 1) \Delta^2.$$

Hence, we obtain by (4.20) that

$$(1, 1, 1) \Delta (\tilde{\Sigma})^{-1} \Delta \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 1. \quad (4.21)$$

Note that

$$\tilde{\Sigma} = \begin{pmatrix} 1 & \langle \tilde{P}_1, \tilde{P}_2 \rangle & \langle \tilde{P}_1, \tilde{P}_3 \rangle \\ \langle \tilde{P}_1, \tilde{P}_2 \rangle & 1 & \langle \tilde{P}_2, \tilde{P}_3 \rangle \\ \langle \tilde{P}_1, \tilde{P}_3 \rangle & \langle \tilde{P}_2, \tilde{P}_3 \rangle & 1 \end{pmatrix} = \begin{pmatrix} 1 & \cos \alpha_{12} & \cos \alpha_{13} \\ \cos \alpha_{12} & 1 & \cos \alpha_{23} \\ \cos \alpha_{13} & \cos \alpha_{23} & 1 \end{pmatrix},$$

which is a positive-definite matrix. Thus, we obtain by (4.21) that

$$(\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3) \begin{pmatrix} 1 & \cos \alpha_{12} & \cos \alpha_{13} \\ \cos \alpha_{12} & 1 & \cos \alpha_{23} \\ \cos \alpha_{13} & \cos \alpha_{23} & 1 \end{pmatrix}^{-1} \begin{pmatrix} \mathcal{L}_1 \\ \mathcal{L}_2 \\ \mathcal{L}_3 \end{pmatrix} = 1. \quad (4.22)$$

From now on till the end of this paper, we adopt the convention that $\frac{0}{0} = 1$ and an increasing (decreasing) function means a strictly increasing (decreasing) function. Define

$$f(x) = \frac{\sqrt{1-x^2}}{\arccos x}, \quad -1 \leq x \leq 1. \quad (4.23)$$

Then, f is an increasing function on $[-1, 1]$ with $f(-1) = 0$ and $f(1) = 1$. Let f^{-1} be the inverse function of f . Then, f^{-1} is an increasing function on $[0, 1]$ with $f^{-1}(0) = -1$ and $f^{-1}(1) = 1$.

For $x, y, z \in (0, \infty)$ satisfying $xy < 1, xz < 1, yz < 1$, define

$$\Gamma(x, y, z) = \begin{pmatrix} 1 & f^{-1}(xy) & f^{-1}(xz) \\ f^{-1}(xy) & 1 & f^{-1}(yz) \\ f^{-1}(xz) & f^{-1}(yz) & 1 \end{pmatrix}. \quad (4.24)$$

Further, if $\det(\Gamma(x, y, z)) > 0$, define

$$H(x, y, z) = (x, y, z)[\Gamma(x, y, z)]^{-1} \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \quad (4.25)$$

Denote

$$r_{ij} = \cos \alpha_{ij}, \quad i < j.$$

Then,

$$f(r_{ij}) = \frac{\sin \alpha_{ij}}{\alpha_{ij}}.$$

By (4.18), we may set

$$\gamma := \frac{\mathcal{L}_1 \mathcal{L}_2 \alpha_{12}}{\sin \alpha_{12}} = \frac{\mathcal{L}_1 \mathcal{L}_3 \alpha_{13}}{\sin \alpha_{13}} = \frac{\mathcal{L}_1 \mathcal{L}_4 \alpha_{14}}{\sin \alpha_{14}} = \frac{\mathcal{L}_2 \mathcal{L}_3 \alpha_{23}}{\sin \alpha_{23}} = \frac{\mathcal{L}_2 \mathcal{L}_4 \alpha_{24}}{\sin \alpha_{24}} = \frac{\mathcal{L}_3 \mathcal{L}_4 \alpha_{34}}{\sin \alpha_{34}}. \quad (4.26)$$

Then, we have that

$$\mathcal{L}_i \mathcal{L}_j = \gamma f(r_{ij}), \quad i < j. \quad (4.27)$$

Define

$$u_i = \frac{\mathcal{L}_i}{\sqrt{\gamma}}, \quad 1 \leq i \leq 4. \quad (4.28)$$

By (4.27), we get

$$u_i u_j = f(r_{ij}), \quad r_{ij} = f^{-1}(u_i u_j), \quad i < j. \quad (4.29)$$

Then, we obtain by (4.22) and (4.29) that

$$H(u_1, u_2, u_3) = \frac{1}{\gamma}.$$

Further, by symmetry, we can show that

$$H(u_1, u_2, u_3) = H(u_1, u_2, u_4) = H(u_1, u_3, u_4) = H(u_2, u_3, u_4) = \frac{1}{\gamma}. \quad (4.30)$$

Step 4. By (4.30) and Theorem 5.1 of the next section, we conclude that

$$u_1 = u_2 = u_3 = u_4,$$

which together with (4.26)–(4.28) implies that

$$\alpha_{12} = \alpha_{13} = \cdots = \alpha_{34}. \quad (4.31)$$

It is well-known that if the six dihedral angles of a given tetrahedron are congruent, then the tetrahedron is regular (cf. e.g. [19]). Then, we obtain by (4.31) that

$$(\Lambda')_{12} = (\Lambda')_{13} = \cdots = (\Lambda')_{34}.$$

Therefore, $\Lambda_{kl} = -\frac{1}{3}$ for all $k < l$ by Remark 3.4. □

5 Monotonicity of H

Let the function H be defined as in (4.25). For $x, y \in (0, \infty)$ satisfying $xy < 1$, define

$$U_{xy} = \left\{ z \in \left(0, \frac{1}{\max\{x, y\}} \right) : \det(\Gamma(x, y, z)) > 0 \right\}.$$

In this section, we will establish the following crucial result, which has been used in the above section to prove Theorem 1.1.

Theorem 5.1 *Let $w_1, w_2 \in (0, \infty)$ satisfying $w_1 w_2 < 1$. Suppose that $w_3, w_4 \in U_{w_1 w_2}$ satisfying $w_3 < w_4$. Then, $H(w_1, w_2, w_3) > H(w_1, w_2, w_4)$.*

Proof. Let $(x, y, z) \in (0, \infty)^3$ satisfying $xy, xz, yz < 1$. Define

$$\varsigma = f^{-1}(xy), \quad \eta = f^{-1}(xz), \quad \xi = f^{-1}(yz), \quad (5.1)$$

and

$$\theta = \arccos \varsigma, \quad \nu = \arccos \eta, \quad \mu = \arccos \xi. \quad (5.2)$$

Then, $(\theta, \nu, \mu) \in (0, \pi)^3$ and

$$x = \sqrt{\frac{f(\cos \theta) f(\cos \nu)}{f(\cos \mu)}}, \quad y = \sqrt{\frac{f(\cos \theta) f(\cos \mu)}{f(\cos \nu)}}, \quad z = \sqrt{\frac{f(\cos \mu) f(\cos \nu)}{f(\cos \theta)}}. \quad (5.3)$$

For $(\theta, \nu, \mu) \in (0, \pi)^3$, we define

$$J(\theta, \nu, \mu) = H(x, y, z),$$

where (x, y, z) is given by (5.3).

Step 1. By (4.24), (5.1) and (5.2), we have

$$\begin{aligned} & \det(\Gamma(x, y, z)) \\ &= 1 - [f^{-1}(xy)]^2 - [f^{-1}(xz)]^2 - [f^{-1}(yz)]^2 + 2f^{-1}(xy)f^{-1}(xz)f^{-1}(yz) \\ &= 1 - \varsigma^2 - \eta^2 - \xi^2 + 2\varsigma\eta\xi \\ &= 1 - \cos^2 \theta - \cos^2 \nu - \cos^2 \mu + 2 \cos \theta \cos \nu \cos \mu \\ &= \sin^2 \mu \sin^2 \nu - (\cos \theta - \cos \mu \cos \nu)^2. \end{aligned} \quad (5.4)$$

Then,

$$\det(\Gamma(x, y, z)) > 0 \Leftrightarrow \begin{cases} \theta \in (|\mu - \nu|, \mu + \nu), & \text{if } \mu + \nu \leq \pi, \\ \theta \in (|\mu - \nu|, 2\pi - (\mu + \nu)), & \text{if } \mu + \nu > \pi. \end{cases} \quad (5.5)$$

Let $\mu, \nu \in (0, \pi)$. Define

$$V_{\mu\nu} = \{\theta \in (0, \pi) : \text{condition (5.5) holds}\}.$$

We will show that $J(\cdot, \nu, \mu)$ is an increasing function on $V_{\mu\nu}$.

By (4.24), (4.25) and (5.1), we get

$$\begin{aligned} & H(x, y, z) \\ &= \frac{1}{1 - [f^{-1}(xy)]^2 - [f^{-1}(xz)]^2 - [f^{-1}(yz)]^2 + 2f^{-1}(xy)f^{-1}(xz)f^{-1}(yz)} \\ & \quad \cdot \{x^2(1 - [f^{-1}(yz)]^2) + y^2(1 - [f^{-1}(xz)]^2) + z^2(1 - [f^{-1}(xy)]^2) \\ & \quad + 2xy[f^{-1}(xz)f^{-1}(yz) - f^{-1}(xy)] + 2xz[f^{-1}(xy)f^{-1}(yz) - f^{-1}(xz)] \\ & \quad + 2yz[f^{-1}(xy)f^{-1}(xz) - f^{-1}(yz)]\} \\ &= \frac{1}{1 - \varsigma^2 - \eta^2 - \xi^2 + 2\varsigma\eta\xi} \left\{ \frac{f(\varsigma)f(\eta)(1 - \xi^2)}{f(\xi)} + \frac{f(\varsigma)f(\xi)(1 - \eta^2)}{f(\eta)} + \frac{f(\xi)f(\eta)(1 - \varsigma^2)}{f(\varsigma)} \right. \\ & \quad \left. + 2f(\varsigma)(\eta\xi - \varsigma) + 2f(\eta)(\varsigma\xi - \eta) + 2f(\xi)(\varsigma\eta - \xi) \right\}. \end{aligned} \quad (5.6)$$

For $\tau \in (-1, 1)$, define

$$\varsigma = \xi\eta + \tau\sqrt{1 - \xi^2}\sqrt{1 - \eta^2}.$$

Then,

$$\cos \theta = \cos \mu \cos \nu + \tau \sin \mu \sin \nu, \quad (5.7)$$

and hence

$$\begin{aligned} & 1 - \varsigma^2 - \eta^2 - \xi^2 + 2\varsigma\eta\xi \\ &= 1 - \cos^2 \theta - \cos^2 \nu - \cos^2 \mu + 2 \cos \theta \cos \nu \cos \mu \\ &= \sin^2 \mu \sin^2 \nu - (\cos \theta - \cos \mu \cos \nu)^2 \\ &= (1 - \tau^2) \sin^2 \mu \sin^2 \nu. \end{aligned} \quad (5.8)$$

In this step, we assume without loss of generality that $\mu \geq \nu$. Then,

$$\begin{cases} \theta \in (\mu - \nu, \mu + \nu), & \text{if } \mu + \nu \leq \pi, \\ \theta \in (\mu - \nu, 2\pi - (\mu + \nu)), & \text{if } \mu + \nu > \pi. \end{cases}$$

Denote

$$a = \mu + \nu, \quad b = \mu - \nu. \quad (5.9)$$

By (4.23), (5.2) and (5.6)–(5.9), we get

$$\begin{aligned}
& J(\theta, \nu, \mu) \\
&= H(x, y, z) \\
&= \frac{1}{(1-\tau^2)\sin^2\mu\sin^2\nu} \left\{ f(\cos\theta) \left(\frac{\mu\sin\mu\sin\nu}{\nu} + \frac{\nu\sin\nu\sin\mu}{\mu} - 2\tau\sin\mu\sin\nu \right) \right. \\
&\quad + \frac{\sin\mu\sin\nu(1-\cos^2\theta)}{\mu\nu f(\cos\theta)} \\
&\quad \left. + \frac{2\sin\nu\sin\mu(\tau\cos\mu\sin\nu - \sin\mu\cos\nu)}{\nu} + \frac{2\sin\mu\sin\nu(\tau\sin\mu\cos\nu - \cos\mu\sin\nu)}{\mu} \right\} \\
&= \frac{1}{(1-\tau^2)\mu\nu\sin\mu\sin\nu} \left\{ f(\cos\theta) (\mu^2 + \nu^2 - 2\tau\mu\nu) + \frac{1-\cos^2\theta}{f(\cos\theta)} \right. \\
&\quad \left. + 2\mu(\tau\cos\mu\sin\nu - \sin\mu\cos\nu) + 2\nu(\tau\sin\mu\cos\nu - \cos\mu\sin\nu) \right\} \\
&= \frac{2[\cos(\mu-\nu) - \cos(\mu+\nu)]}{[(\mu+\nu)^2 - (\mu-\nu)^2] \cdot [\cos\theta - \cos(\mu+\nu)] \cdot [\cos(\mu-\nu) - \cos\theta]} \\
&\quad \cdot \left\{ \frac{\sin\theta}{\theta} \left[(\mu-\nu)^2 + \frac{[(\mu+\nu)^2 - (\mu-\nu)^2] \cdot [\cos(\mu-\nu) - \cos\theta]}{\cos(\mu-\nu) - \cos(\mu+\nu)} \right] + \theta\sin\theta \right. \\
&\quad + \frac{2[\cos\theta - \cos(\mu-\nu)]}{\cos(\mu-\nu) - \cos(\mu+\nu)} [(\mu+\nu)\sin(\mu+\nu) - (\mu-\nu)\sin(\mu-\nu)] \\
&\quad \left. - 2(\mu-\nu)\sin(\mu-\nu) \right\} \\
&= \frac{2(\cos b - \cos a)}{(a^2 - b^2)(\cos\theta - \cos a)(\cos b - \cos\theta)} \left\{ \frac{\sin\theta}{\theta} \left[b^2 + \frac{(a^2 - b^2)(\cos b - \cos\theta)}{\cos b - \cos a} \right] + \theta\sin\theta \right. \\
&\quad \left. + \frac{2(\cos\theta - \cos b)}{\cos b - \cos a} (a\sin a - b\sin b) - 2b\sin b \right\} \\
&= \frac{2(\cos b - \cos a)}{(a^2 - b^2)(\cos\theta - \cos a)(\cos b - \cos\theta)} \left\{ \frac{\sin\theta}{\theta} \cdot \frac{a^2(\cos b - \cos\theta) + b^2(\cos\theta - \cos a)}{\cos b - \cos a} \right. \\
&\quad + \frac{\theta\sin\theta(\cos b - \cos\theta)}{\cos b - \cos a} + \frac{\theta\sin\theta(\cos\theta - \cos a)}{\cos b - \cos a} \\
&\quad \left. - \frac{2a\sin a(\cos b - \cos\theta)}{\cos b - \cos a} - \frac{2b\sin b(\cos\theta - \cos a)}{\cos b - \cos a} \right\} \\
&= \frac{2}{a^2 - b^2} \left\{ \frac{a^2\sin\theta + \theta^2\sin\theta - 2a\theta\sin a}{\theta(\cos\theta - \cos a)} + \frac{b^2\sin\theta + \theta^2\sin\theta - 2b\theta\sin b}{\theta(\cos b - \cos\theta)} \right\} \\
&= \frac{2}{a^2 - b^2} \left\{ \frac{(a-\theta)^2\sin\theta + 2a\theta(\sin\theta - \sin a)}{\theta(\cos\theta - \cos a)} - \frac{(b-\theta)^2\sin\theta + 2b\theta(\sin\theta - \sin b)}{\theta(\cos\theta - \cos b)} \right\} \\
&:= \frac{2}{a^2 - b^2} G(a, b, \theta).
\end{aligned}$$

We will show that $\partial_\theta G(a, b, \theta) > 0$. For $u \in [0, \theta) \cup (\theta, 2\pi - \theta)$, define

$$K(u, \theta) = \frac{(u - \theta)^2 \sin\theta + 2u\theta(\sin\theta - \sin u)}{\theta(\cos\theta - \cos u)}. \quad (5.10)$$

Then,

$$G(a, b, \theta) = K(a, \theta) - K(b, \theta),$$

and

$$\partial_\theta G(\theta, a, b) = \partial_\theta K(a, \theta) - \partial_\theta K(b, \theta). \quad (5.11)$$

By (5.10), we get

$$\begin{aligned} & \partial_\theta K(u, \theta) \\ = & \frac{(u^2 + \theta^2)\theta(1 - \cos \theta \cos u) - (u^2 - \theta^2) \sin \theta (\cos \theta - \cos u) - 2u\theta^2 \sin u \sin \theta}{\theta^2 (\cos \theta - \cos u)^2} \\ := & P(u, \theta). \end{aligned} \quad (5.12)$$

By virtue of L'Hôpital's rule, we obtain that

$$\begin{aligned} & \lim_{u \rightarrow \theta} P(u, \theta) \\ = & \frac{1}{\theta^2} \lim_{u \rightarrow \theta} \frac{2u\theta(1 - \cos \theta \cos u) - 2u \sin \theta (\cos \theta - \cos u) - \sin u (u^2 + \theta^2)(\sin \theta - \theta \cos \theta) - 2u\theta^2 \cos u \sin \theta}{2 \sin u (\cos \theta - \cos u)} \\ = & \frac{1}{\theta^2} \left\{ -\theta + \lim_{u \rightarrow \theta} \frac{2u\theta(1 - \cos \theta \cos u) - \sin u (u^2 + \theta^2)(\sin \theta - \theta \cos \theta) - 2u\theta^2 \cos u \sin \theta}{2 \sin u (\cos \theta - \cos u)} \right\} \\ = & \frac{1}{\theta^2} \left\{ -\theta + \lim_{u \rightarrow \theta} \frac{2u\theta(1 - \cos \theta \cos u) - \sin u \sin \theta (u^2 + \theta^2)}{2 \sin \theta (\cos \theta - \cos u)} \right. \\ & \left. + \lim_{u \rightarrow \theta} \frac{\theta \cos \theta \sin u (u^2 + \theta^2) - 2u\theta^2 \cos u \sin \theta}{2 \sin \theta (\cos \theta - \cos u)} \right\} \\ = & \frac{1}{\theta^2} \left\{ -\theta + \frac{\theta^3}{\sin^2 \theta} \right\} \\ = & \frac{\theta^2 - \sin^2 \theta}{\theta \sin^2 \theta} \\ := & P(\theta). \end{aligned}$$

We have

$$\begin{aligned}
& P(u, \theta) > P(\theta) \\
\Leftrightarrow & [(u^2 + \theta^2)\theta(1 - \cos \theta \cos u) - (u^2 - \theta^2) \sin \theta (\cos \theta - \cos u) - 2u\theta^2 \sin u \sin \theta] \sin^2 \theta \\
& > \theta(\theta^2 - \sin^2 \theta)(\cos \theta - \cos u)^2 \\
\Leftrightarrow & \{(u^2 + \theta^2)\theta[1 - \cos(u - \theta)] - (u^2 - \theta^2) \sin \theta (\cos \theta - \cos u) + (u - \theta)^2 \theta \sin u \sin \theta\} \sin^2 \theta \\
& > \theta(\theta^2 - \sin^2 \theta)(\cos \theta - \cos u)^2 \\
\Leftrightarrow & \left[2(u^2 + \theta^2)\theta \sin^2 \left(\frac{u - \theta}{2} \right) - 2(u^2 - \theta^2) \sin \theta \sin \left(\frac{u - \theta}{2} \right) \sin \left(\frac{u + \theta}{2} \right) \right. \\
& \quad \left. + (u - \theta)^2 \theta \left(\sin^2 \left(\frac{u + \theta}{2} \right) - \sin^2 \left(\frac{u - \theta}{2} \right) \right) \right] \sin^2 \theta \\
& > 4\theta(\theta^2 - \sin^2 \theta) \sin^2 \left(\frac{u - \theta}{2} \right) \sin^2 \left(\frac{u + \theta}{2} \right) \\
\Leftrightarrow & \left[(u + \theta)^2 \theta \sin^2 \left(\frac{u - \theta}{2} \right) - 2(u^2 - \theta^2) \sin \theta \sin \left(\frac{u - \theta}{2} \right) \sin \left(\frac{u + \theta}{2} \right) \right. \\
& \quad \left. + (u - \theta)^2 \theta \sin^2 \left(\frac{u + \theta}{2} \right) \right] \sin^2 \theta \\
& > 4\theta(\theta^2 - \sin^2 \theta) \sin^2 \left(\frac{u - \theta}{2} \right) \sin^2 \left(\frac{u + \theta}{2} \right) \\
\Leftrightarrow & \left[\frac{(u + \theta)^2 \theta}{\sin^2 \left(\frac{u - \theta}{2} \right)} - \frac{2(u^2 - \theta^2) \sin \theta}{\sin \left(\frac{u - \theta}{2} \right) \sin \left(\frac{u + \theta}{2} \right)} + \frac{(u - \theta)^2 \theta}{\sin^2 \left(\frac{u - \theta}{2} \right)} \right] \sin^2 \theta > 4\theta(\theta^2 - \sin^2 \theta) \\
\Leftrightarrow & \left[\frac{u + \theta}{\sin \left(\frac{u + \theta}{2} \right)} - \frac{u - \theta}{\sin \left(\frac{u - \theta}{2} \right)} \right]^2 \theta \sin^2 \theta + \frac{2(u^2 - \theta^2)(\theta - \sin \theta) \sin^2 \theta}{\sin \left(\frac{u - \theta}{2} \right) \sin \left(\frac{u + \theta}{2} \right)} > 4\theta(\theta^2 - \sin^2 \theta). \tag{5.13}
\end{aligned}$$

Case 1: Suppose that $u \in (\theta, 2\pi - \theta)$.

Define

$$g(w) = \frac{w}{\sin w}, \quad w \in (0, \pi).$$

Then,

$$g'(w) = \frac{\sin w - w \cos w}{\sin^2 w} > 0,$$

which implies that $g(w)$ is an increasing function for $w \in (0, \pi)$. Hence

$$\frac{u^2 - \theta^2}{\sin \left(\frac{u - \theta}{2} \right) \sin \left(\frac{u + \theta}{2} \right)} = \frac{u + \theta}{\sin \left(\frac{u + \theta}{2} \right)} \cdot \frac{u - \theta}{\sin \left(\frac{u - \theta}{2} \right)} = 4g \left(\frac{u + \theta}{2} \right) g \left(\frac{u - \theta}{2} \right)$$

is an increasing function for $u \in (\theta, 2\pi - \theta)$.

We have

$$g''(w) = \frac{w(1 + \cos^2 w) - 2 \sin w \cos w}{\sin^3 w} > 0,$$

which implies that $g'(w)$ is an increasing function for $w \in (0, \pi)$. Hence

$$\partial_u \left(\frac{\frac{u + \theta}{2}}{\sin \left(\frac{u + \theta}{2} \right)} \right) - \partial_u \left(\frac{\frac{u - \theta}{2}}{\sin \left(\frac{u - \theta}{2} \right)} \right) = \frac{1}{2} \left[g' \left(\frac{u + \theta}{2} \right) - g' \left(\frac{u - \theta}{2} \right) \right] > 0,$$

which implies that

$$\frac{\frac{u+\theta}{2}}{\sin\left(\frac{u+\theta}{2}\right)} - \frac{\frac{u-\theta}{2}}{\sin\left(\frac{u-\theta}{2}\right)}$$

is an increasing function for $u \in (\theta, 2\pi - \theta)$. Thus,

$$\left[\frac{u+\theta}{\sin\left(\frac{u+\theta}{2}\right)} - \frac{u-\theta}{\sin\left(\frac{u-\theta}{2}\right)} \right]^2 \theta \sin^2 \theta + \frac{2(u^2 - \theta^2)(\theta - \sin \theta) \sin^2 \theta}{\sin\left(\frac{u-\theta}{2}\right) \sin\left(\frac{u+\theta}{2}\right)}$$

is an increasing function for $u \in (\theta, 2\pi - \theta)$.

Case 2: Suppose that $u \in [0, \theta)$.

We have

$$\frac{\frac{u+\theta}{2}}{\sin\left(\frac{u+\theta}{2}\right)} - \frac{\frac{u-\theta}{2}}{\sin\left(\frac{u-\theta}{2}\right)} = \frac{\frac{\theta+u}{2}}{\sin\left(\frac{\theta+u}{2}\right)} - \frac{\frac{\theta-u}{2}}{\sin\left(\frac{\theta-u}{2}\right)} = g\left(\frac{u+\theta}{2}\right) - g\left(\frac{\theta-u}{2}\right),$$

which implies that

$$\frac{\frac{u+\theta}{2}}{\sin\left(\frac{u+\theta}{2}\right)} - \frac{\frac{u-\theta}{2}}{\sin\left(\frac{u-\theta}{2}\right)}$$

is an increasing function for $u \in [0, \theta)$.

We have

$$\begin{aligned} (\ln g)''(w) &= \frac{g''(w)g(w) - (g'(w))^2}{g^2(w)} \\ &= \frac{w^2 - \sin^2 w}{g^2(w) \sin^4(w)} \\ &> 0, \end{aligned}$$

and

$$\begin{aligned} &\partial_u \left(\frac{u^2 - \theta^2}{\sin\left(\frac{u-\theta}{2}\right) \sin\left(\frac{u+\theta}{2}\right)} \right) > 0 \\ \Leftrightarrow &\partial_u \left(\frac{\frac{\theta+u}{2}}{\sin\left(\frac{\theta+u}{2}\right)} \cdot \frac{\frac{\theta-u}{2}}{\sin\left(\frac{\theta-u}{2}\right)} \right) > 0 \\ \Leftrightarrow &g'\left(\frac{\theta+u}{2}\right) g\left(\frac{\theta-u}{2}\right) - g'\left(\frac{\theta-u}{2}\right) g\left(\frac{\theta+u}{2}\right) > 0 \\ \Leftrightarrow &\frac{g'\left(\frac{\theta+u}{2}\right)}{g\left(\frac{\theta+u}{2}\right)} > \frac{g'\left(\frac{\theta-u}{2}\right)}{g\left(\frac{\theta-u}{2}\right)} \\ \Leftrightarrow &(\ln g)'\left(\frac{\theta+u}{2}\right) > (\ln g)'\left(\frac{\theta-u}{2}\right). \end{aligned}$$

Then,

$$\frac{u^2 - \theta^2}{\sin\left(\frac{u-\theta}{2}\right) \sin\left(\frac{u+\theta}{2}\right)}$$

is an increasing function for $u \in [0, \theta)$. Thus,

$$\left[\frac{u + \theta}{\sin\left(\frac{u+\theta}{2}\right)} - \frac{u - \theta}{\sin\left(\frac{u-\theta}{2}\right)} \right]^2 \theta \sin^2 \theta + \frac{2(u^2 - \theta^2)(\theta - \sin \theta) \sin^2 \theta}{\sin\left(\frac{u-\theta}{2}\right) \sin\left(\frac{u+\theta}{2}\right)}$$

is an increasing function for $u \in [0, \theta)$.

By virtue of L'Hôpital's rule, we get

$$\begin{aligned} & \lim_{u \rightarrow \theta} \left\{ \left[\frac{u + \theta}{\sin\left(\frac{u+\theta}{2}\right)} - \frac{u - \theta}{\sin\left(\frac{u-\theta}{2}\right)} \right]^2 \theta \sin^2 \theta + \frac{2(u^2 - \theta^2)(\theta - \sin \theta) \sin^2 \theta}{\sin\left(\frac{u-\theta}{2}\right) \sin\left(\frac{u+\theta}{2}\right)} \right\} \\ &= \left(\frac{2\theta}{\sin \theta} - 2 \right)^2 \theta \sin^2 \theta + 8\theta \sin \theta (\theta - \sin \theta) \\ &= 4\theta(\theta^2 - \sin^2 \theta). \end{aligned} \tag{5.14}$$

Then, we obtain by (5.13) and (5.14) that $P(u, \theta) > P(\theta)$ if $u \in (\theta, 2\pi - \theta)$ and $P(u, \theta) < P(\theta)$ if $u \in [0, \theta)$. Thus, by (5.11) and (5.12), we get

$$\partial_\theta G(\theta, a, b) = P(a, \theta) - P(b, \theta) = [P(a, \theta) - P(\theta)] + [P(\theta) - P(b, \theta)] > 0. \tag{5.15}$$

Step 2. Let $x, y \in (0, \infty)$ satisfying $xy < 1$. We will show that either $U_{xy} = \emptyset$ or there exist $z_1 < z_2$ such that $U_{xy} = (z_1, z_2)$. By symmetry, we assume without loss of generality that $x \leq y$ in this step. Then, $\mu \leq \nu$.

By (5.4), we get

$$\det(\Gamma(x, y, z)) = 0 \Leftrightarrow \mu = 2\pi - (\nu + \theta), \theta - \nu, \text{ or } \nu - \theta.$$

By (5.1) and (5.2), we have

$$\frac{\sin \mu}{\mu} = \frac{y \sin \nu}{x \nu}. \tag{5.16}$$

If $\mu = 2\pi - (\nu + \theta)$, by (5.16), we get

$$\frac{\sin(2\pi - (\nu + \theta))}{2\pi - (\nu + \theta)} = \frac{y \sin \nu}{x \nu}. \tag{5.17}$$

Since $\frac{\sin w}{w}$ is a decreasing function for $w \in (0, \pi)$, there exists at most one ν satisfying (5.17).

If $\mu = \theta - \nu$, then $\nu < \theta$. By (5.16), we get

$$\frac{\sin(\theta - \nu)}{\theta - \nu} = \frac{y \sin \nu}{x \nu}. \tag{5.18}$$

Since $\frac{\sin w}{w}$ is a decreasing function for $w \in (0, \pi)$, there exists at most one ν satisfying (5.18). Moreover, there exists a ν satisfying (5.18) only if

$$\frac{y \sin \theta}{x \theta} < 1. \tag{5.19}$$

If $\mu = \nu - \theta$, then $\nu > \theta$. By (5.16), we get

$$\frac{\sin(\nu - \theta)}{\nu - \theta} = \frac{y \sin \nu}{x \nu}. \quad (5.20)$$

Define

$$W(w) = \frac{w \sin(w - \theta)}{(w - \theta) \sin w}, \quad w \in [\theta, \pi).$$

We have

$$W'(w) = \frac{(\sin \theta)w(w - \theta) - \theta \sin w \sin(w - \theta)}{[(w - \theta) \sin w]^2}.$$

Then,

$$W'(w) > 0 \Leftrightarrow \frac{\sin(w - \theta)}{(w - \theta)} \cdot \frac{\sin w}{w} < \frac{\sin \theta}{\theta}.$$

Since $\frac{\sin w}{w}$ is a decreasing function for $w \in (0, \pi)$, we get $W'(w) > 0$ for $w \in (\theta, \pi)$. Thus, there exists at most one ν satisfying (5.20). Moreover, since W is an increasing function on $[\theta, \pi)$ and

$$W(\theta) = \frac{\theta}{\sin \theta},$$

there exists a ν satisfying (5.20) only if

$$\frac{y \sin \theta}{x \theta} > 1. \quad (5.21)$$

By comparing conditions (5.19) and (5.21), we conclude that it is impossible that both $\mu = \theta - \nu$ and $\mu = \nu - \theta$ have solutions. Hence, $\det(\Gamma(x, y, z)) = 0$ has at most two solutions. Note that

$$\begin{aligned} \det(\Gamma(x, y, 0)) &:= -1 - \varsigma^2 + 2\varsigma < 0, \\ \det\left(\Gamma\left(x, y, \frac{1}{y}\right)\right) &:= -\varsigma^2 - \left[f^{-1}\left(\frac{x}{y}\right)\right]^2 - 2\varsigma f^{-1}\left(\frac{x}{y}\right) < 0. \end{aligned}$$

If $\det(\Gamma(x, y, z)) = 0$ has at most one solution, then $U_{xy} = \emptyset$. If $\det(\Gamma(x, y, z)) = 0$ has two solutions $z_1 < z_2$, then we have that

$$\det(\Gamma(x, y, z)) < 0 \quad \text{for } z \in (0, z_1) \cup \left(z_2, \frac{1}{y}\right),$$

and

$$\det(\Gamma(x, y, z)) > 0, \quad \forall z \in (z_1, z_2) \quad \text{or} \quad \det(\Gamma(x, y, z)) < 0, \quad \forall z \in (z_1, z_2).$$

Hence $\det(\Gamma(x, y, z)) > 0$ has no solution or $\det(\Gamma(x, y, z)) > 0$ for any $z \in (z_1, z_2)$. Thus, $U_{xy} = \emptyset$ or $U_{xy} = (z_1, z_2)$.

Step 3. Define

$$V = \{(\theta, \nu, \mu) \in (0, \pi)^3 : \text{condition (5.5) holds}\}.$$

Then, V is an open subset of $(0, \pi)^3$. Let $(\theta_0, \nu_0, \mu_0) \in V$. Then, there exists $\varepsilon > 0$ such that

$$V_0 := [\theta_0 - \varepsilon, \theta_0 + \varepsilon] \times [\nu_0 - \varepsilon, \nu_0 + \varepsilon] \times [\mu_0 - \varepsilon, \mu_0 + \varepsilon] \subset V.$$

By Step 1 and symmetry of the function J , we conclude that for any $(\theta, \nu, \mu), (\theta', \nu', \mu') \in V_0$ satisfying $\theta \leq \theta', \nu \leq \nu'$ and $\mu \leq \mu'$, we have

$$J(\theta, \nu, \mu) \leq J(\theta', \nu', \mu'),$$

and the equality holds if and only if $(\theta, \nu, \mu) = (\theta', \nu', \mu')$.

Define

$$U = \{(x, y, z) \in (0, \infty)^3 : xy, xz, yz < 1, \det(\Gamma(x, y, z)) > 0\}.$$

Let $(x_0, y_0, z_0) \in U$. Since f is a homeomorphic map between U and V , there exists $\epsilon > 0$ such that for any $(x, y, z), (x', y', z') \in [x_0 - \epsilon, x_0 + \epsilon] \times [y_0 - \epsilon, y_0 + \epsilon] \times [z_0 - \epsilon, z_0 + \epsilon]$ with $x \leq x', y \leq y'$ and $z \leq z'$, we have

$$H(x, y, z) \geq H(x', y', z'),$$

and the equality holds if and only if $(x, y, z) = (x', y', z')$.

Let $w_1, w_2 \in (0, \infty)$ satisfying $w_1 w_2 < 1$. Suppose that $w_3, w_4 \in U_{w_1 w_2}$ satisfying $w_3 < w_4$. By Step 2, we know that there exist $z_1 < z_2$ such that $U_{w_1 w_2} = (z_1, z_2)$. By the above analysis, we know that for any $z \in (z_1, z_2)$, there exists $\epsilon_z > 0$ such that for any $p, q \in [z - \epsilon_z, z + \epsilon_z]$ with $p < q$, we have

$$H(w_1, w_2, p) > H(w_1, w_2, q).$$

Therefore, the proof is complete by virtue of the Heine-Borel theorem. \square

Remark 5.2 *To obtain the monotonicity of the function H , we prove in Step 1 the monotonicity of the function $G(\cdot, a, b)$ by establishing (5.15). We believe that the following stronger result holds.*

$$\partial_u P(u, \theta) > 0, \quad \text{for any } u \in [0, \theta) \bigcup (\theta, 2\pi - \theta) \text{ and } \theta \in (0, \pi). \quad (5.22)$$

In the Appendix, we will show that (5.22) is equivalent to the following new inequality.

$$\begin{aligned} & (\cos \theta - \cos u)^2 [2u\theta(1 - \cos \theta \cos u) - 2u \sin \theta (\cos \theta - \cos u) + \sin u \sin \theta (u^2 - 3\theta^2)] \\ & + (\cos \theta - \cos u) [2u\theta^2 \sin \theta (\sin^2 u + 1 - \cos \theta \cos u) - (u^2 + \theta^2) \sin u (\sin^2 \theta + 1 - \cos \theta \cos u)] \\ & > 0, \quad \text{for any } u \in [0, \theta) \bigcup (\theta, 2\pi - \theta) \text{ and } \theta \in (0, \pi). \end{aligned} \quad (5.23)$$

So far we have not been able to prove (5.23), however, it has been verified by computer.

6 Some remarks

In this section, we make some remarks. We hope they are helpful for considering the higher dimensional GRVM and SMW conjectures.

6.1 Non-concavity of $F(\Lambda)$

If F is a concave function on $\bar{\mathcal{S}}$, then the sufficiency of Theorem 1.1 can be established directly by Lemma 2.1 and Theorem 3.3. Unfortunately, we find that F is not a concave function. Consider

$$\Lambda = \begin{pmatrix} 1 & 0.93 & 0.91 & 0.90 \\ 0.93 & 1 & 0.75 & 0.77 \\ 0.91 & 0.75 & 1 & 0.75 \\ 0.90 & 0.77 & 0.75 & 1 \end{pmatrix},$$

and $\bar{\Lambda} \in \mathcal{S}$ such that

$$(\bar{\Lambda})_{kl} = \frac{\sum_{s < t} \Lambda_{st}}{6}, \quad k < l.$$

By (3.5) and the software R, we get

$$F(\Lambda) - F(\bar{\Lambda}) = 0.0003994782,$$

which implies that F is non-concave.

We would like to take this opportunity to point out that numerical calculation via computer has played an important role in discovering some results of this paper. Besides using Mathematica to quickly verify (3.7) and using the software R to check non-concavity of F , we have used computer to check the monotonicity of H . It had greatly improved our confidence before we were able to give the proof of Theorem 5.1 at last.

We now give an affirmative result on the concavity of the function $F(\cdot)$ around some particular Λ . Let $X \in \mathcal{G}$. We denote by S the tetrahedron with vertexes $\{X_1, X_2, X_3, X_4\}$. If none of its dihedral angles is obtuse, then S is said to be nonobtuse. For $\Lambda \in \mathcal{S}$, we denote the Hessian matrix of $-F(\Lambda)$ by

$$\mathcal{H}(\Lambda) = - \left(\frac{\partial^2 F}{\partial \Lambda_{pq} \partial \Lambda_{kl}} \right), \quad p < q, k < l.$$

Proposition 6.1 *Suppose that S is nonobtuse. Then, all diagonal elements of $\mathcal{H}(\Lambda^X)$ are positive and $\det(\mathcal{H}(\Lambda^X)) > 0$.*

Proof. Let $X \in \mathcal{G}$. To simplify notation, we denote Λ^X by Λ . First, we show that

$$- \frac{\partial^2 F}{\partial \Lambda_{kl} \partial \Lambda_{kl}} > 0, \quad \forall k < l. \quad (6.1)$$

Denote by Υ_2^X the vertex Gramian of S associated with vertex X_2 , which is equal to the covariance matrix of the Gaussian random vector $\left(\frac{X_1 - X_2}{\sqrt{2}}, \frac{X_3 - X_2}{\sqrt{2}}, \frac{X_4 - X_2}{\sqrt{2}} \right)$ as defined in (7.1). Since S is nonobtuse, each facet of S is nonobtuse (cf. [7] and [4, Proposition 2.8]). Hence the three facets meeting at vertex X_2 are all nonobtuse, which implies that (cf. (7.1) of the Appendix)

$$r_1 \geq 0, \quad r_2 \geq 0, \quad r_3 \geq 0. \quad (6.2)$$

By [4, Theorem 2.4], Υ_2^X is a weakly diagonally dominant Stieltjes matrix. Together with (7.2) of the Appendix, this implies that

$$\tilde{\Lambda}_{kl} \leq 0, \quad \forall k < l. \quad (6.3)$$

Then, we obtain by (3.1), (6.2) and (6.3) that

$$-\frac{\partial^2 F}{\partial \Lambda_{12} \partial \Lambda_{12}} > 0.$$

Similarly, we can show that (6.1) holds for all $k < l$.

We now show that $\det(\mathcal{H}(\Lambda)) > 0$. Denote by $\Phi(\Lambda)$ the 6×6 diagonal matrix with

$$\begin{aligned} \Phi(\Lambda)_{11} &= \frac{1}{8\sqrt{\pi^3(\Lambda')_{12}^3}} \arccos \left(\frac{(\tilde{\Lambda})_{12}}{\sqrt{(\Lambda')_{12}\tilde{A}^2 + (\tilde{\Lambda})_{12}^2}} \right), \\ \Phi(\Lambda)_{22} &= \frac{1}{8\sqrt{\pi^3(\Lambda')_{13}^3}} \arccos \left(\frac{(\tilde{\Lambda})_{13}}{\sqrt{(\Lambda')_{13}\tilde{A}^2 + (\tilde{\Lambda})_{13}^2}} \right), \\ &\dots\dots\dots \\ \Phi(\Lambda)_{66} &= \frac{1}{8\sqrt{\pi^3(\Lambda')_{34}^3}} \arccos \left(\frac{(\tilde{\Lambda})_{34}}{\sqrt{(\Lambda')_{34}\tilde{A}^2 + (\tilde{\Lambda})_{34}^2}} \right). \end{aligned}$$

Define

$$\Psi(\Lambda) = \mathcal{H}(\Lambda) - \Phi(\Lambda). \quad (6.4)$$

Similar to (3.7), we can show that

$$\sum_{k < l} (\Lambda')_{kl} \frac{\partial^2 F}{\partial \Lambda_{pq} \partial \Lambda_{kl}} = \frac{1}{2} \frac{\partial F}{\partial \Lambda_{pq}}, \quad \forall p < q. \quad (6.5)$$

Define

$$v = ((\Lambda')_{12}, (\Lambda')_{13}, \dots, (\Lambda')_{34}),$$

and

$$u = \left(\frac{\partial F}{\partial \Lambda_{12}}, \frac{\partial F}{\partial \Lambda_{13}}, \dots, \frac{\partial F}{\partial \Lambda_{34}} \right).$$

Then, (6.5) can be written as

$$-\mathcal{H}(\Lambda)v^T = \frac{1}{2}u^T. \quad (6.6)$$

By Theorem 2.3, Theorem 3.1, (6.4) and (6.6), we get

$$\Psi(\Lambda)v^T = 0,$$

which implies that $\det(\Psi(\Lambda)) = 0$. Therefore, we obtain by (6.1) and (6.4) that

$$\begin{aligned}
\det(\mathcal{H}(\Lambda)) &= \det(\Phi(\Lambda)) + \det(\Psi(\Lambda)) + \det(\Phi(\Lambda)) \cdot \text{Trace}[\Phi(\Lambda)^{-1}\Psi(\Lambda)] \\
&= \det(\Phi(\Lambda)) \cdot \{1 + \text{Trace}[\Phi^{-1}(\Lambda)\Psi(\Lambda)]\} \\
&= \det(\Phi(\Lambda)) \cdot \text{Trace}[\Phi^{-1}(\Lambda)\mathcal{H}(\Lambda)] \\
&= \left(\prod_{i=1}^6 \Phi_{ii}(\Lambda) \right) \left(\sum_{i=1}^6 \frac{\mathcal{H}_{ii}(\Lambda)}{\Phi_{ii}(\Lambda)} \right) \\
&> 0.
\end{aligned}$$

□

6.2 Bounds for $\mathcal{M}(X)$

In this subsection, we derive the lower and upper bounds for the function $\mathcal{M}(X)$.

Proposition 6.2 *Let $X \in \mathcal{G}$ with covariance matrix Λ . Then,*

$$\frac{1}{4\sqrt{\pi}} \sum_{k < l} \sqrt{(\Lambda')_{kl}} \leq E[\max(X_1, X_2, X_3, X_4)] \leq \frac{1}{3\sqrt{\pi}} \sum_{k < l} \sqrt{(\Lambda')_{kl}}. \quad (6.7)$$

Proof. We define $X^I = \max(X_1, X_2, X_3, X_4)$, $X^{IV} = \min(X_1, X_2, X_3, X_4)$, and let X^{II} and X^{III} be the second and third largest values of X_1, X_2, X_3, X_4 , respectively. Note that

$$E[X^{IV}] = -E[X^I], \quad E[X^{III}] = -E[X^{II}].$$

We have

$$\begin{aligned}
&E[X^{IV}] \\
&= E[X_1 \wedge X_2 \wedge X_3 \wedge X_4] \\
&= E[(X_1 - X_4) \wedge (X_2 - X_4) \wedge (X_3 - X_4) \wedge 0] \\
&= E[(X_1 - X_4) \wedge (X_2 - X_4) \wedge (X_3 - X_4)] \\
&\quad - E[\{(X_1 - X_4) \wedge (X_2 - X_4) \wedge (X_3 - X_4)\} \vee 0] \\
&= E[X_1 \wedge X_2 \wedge X_3] - E[(X_1 - X_4)^+ \wedge (X_2 - X_4)^+ \wedge (X_3 - X_4)^+] \\
&= E[X_1 \wedge X_2 \wedge X_3] - E[\{(X_1 - X_4) \wedge (X_2 - X_4) \wedge (X_3 - X_4)\} \cdot 1_{\{X_4 = \min_{1 \leq i \leq 4} X_i\}}] \\
&= E[X_1 \wedge X_2 \wedge X_3] - E[\{X_1 \wedge X_2 \wedge X_3\} \cdot 1_{\{X_4 = \min_{1 \leq i \leq 4} X_i\}}] + E[X_4 \cdot 1_{\{X_4 = \min_{1 \leq i \leq 4} X_i\}}] \\
&= E[X_1 \wedge X_2 \wedge X_3] - E[X^{III} \cdot 1_{\{X_4 = \min_{1 \leq i \leq 4} X_i\}}] + E[X^{IV} \cdot 1_{\{X_4 = \min_{1 \leq i \leq 4} X_i\}}]. \quad (6.8)
\end{aligned}$$

Similarly, we have

$$E(X^{IV}) = E[X_2 \wedge X_3 \wedge X_4] - E[X^{III} \cdot 1_{\{X_1 = \min_{1 \leq i \leq 4} X_i\}}] + E[X^{IV} \cdot 1_{\{X_1 = \min_{1 \leq i \leq 4} X_i\}}], \quad (6.9)$$

$$E(X^{IV}) = E[X_3 \wedge X_4 \wedge X_1] - E[X^{III} \cdot 1_{\{X_2 = \min_{1 \leq i \leq 4} X_i\}}] + E[X^{IV} \cdot 1_{\{X_2 = \min_{1 \leq i \leq 4} X_i\}}], \quad (6.10)$$

$$E(X^{IV}) = E[X_4 \wedge X_1 \wedge X_2] - E[X^{III} \cdot 1_{\{X_3 = \min_{1 \leq i \leq 4} X_i\}}] + E[X^{IV} \cdot 1_{\{X_3 = \min_{1 \leq i \leq 4} X_i\}}]. \quad (6.11)$$

By (3.6), we get

$$E[X_1 \wedge X_2 \wedge X_3] = -\frac{1}{2\sqrt{\pi}} \left[\sqrt{(\Lambda')_{12}} + \sqrt{(\Lambda')_{13}} + \sqrt{(\Lambda')_{23}} \right], \quad (6.12)$$

$$E[X_2 \wedge X_3 \wedge X_4] = -\frac{1}{2\sqrt{\pi}} \left[\sqrt{(\Lambda')_{23}} + \sqrt{(\Lambda')_{24}} + \sqrt{(\Lambda')_{34}} \right], \quad (6.13)$$

$$E[X_3 \wedge X_4 \wedge X_1] = -\frac{1}{2\sqrt{\pi}} \left[\sqrt{(\Lambda')_{34}} + \sqrt{(\Lambda')_{13}} + \sqrt{(\Lambda')_{14}} \right], \quad (6.14)$$

$$E[X_4 \wedge X_1 \wedge X_2] = -\frac{1}{2\sqrt{\pi}} \left[\sqrt{(\Lambda')_{14}} + \sqrt{(\Lambda')_{24}} + \sqrt{(\Lambda')_{12}} \right]. \quad (6.15)$$

By summing up (6.8)–(6.11) and using (6.12)–(6.15), we get

$$\begin{aligned} 4E[X^{IV}] &= E[X_1 \wedge X_2 \wedge X_3] + E[X_2 \wedge X_3 \wedge X_4] + E[X_3 \wedge X_4 \wedge X_1] + E[X_4 \wedge X_1 \wedge X_2] \\ &\quad - E[X^{III}] + E[X^{IV}] \\ &= -\frac{1}{\sqrt{\pi}} \sum_{k < l} \sqrt{(\Lambda')_{kl}} - E[X^{III}] + E[X^{IV}]. \end{aligned}$$

Then, we have that

$$E[X^{III}] = 3E[X^I] - \frac{1}{\sqrt{\pi}} \sum_{k < l} \sqrt{(\Lambda')_{kl}}, \quad (6.16)$$

and

$$E[X^{II}] = -3E[X^I] + \frac{1}{\sqrt{\pi}} \sum_{k < l} \sqrt{(\Lambda')_{kl}}. \quad (6.17)$$

Therefore, (6.7) holds by (6.16), (6.17) and the fact that $E[X^{III}] \leq E[X^{II}] \leq E[X^I]$. \square

7 Appendix

7.1 Some auxiliary results of Λ

Suppose that $X \in \mathcal{G}$ with covariance matrix $\Lambda \in \mathcal{S}$. Then, $\left(\frac{X_1-X_2}{\sqrt{2}}, \frac{X_3-X_2}{\sqrt{2}}, \frac{X_4-X_2}{\sqrt{2}}\right)$ is a 3-dimensional centered Gaussian random vector with covariance matrix

$$\begin{aligned}\Upsilon_2^X &:= \begin{pmatrix} 1 - \Lambda_{12} & \frac{1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23}}{2} & \frac{1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24}}{2} \\ \frac{1 - \Lambda_{12} + \Lambda_{13} - \Lambda_{23}}{2} & 1 - \Lambda_{23} & \frac{1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34}}{2} \\ \frac{1 - \Lambda_{12} + \Lambda_{14} - \Lambda_{24}}{2} & \frac{1 - \Lambda_{23} - \Lambda_{24} + \Lambda_{34}}{2} & 1 - \Lambda_{24} \end{pmatrix} \\ &:= \begin{pmatrix} a_0 & r_1 & r_2 \\ r_1 & b_0 & r_3 \\ r_2 & r_3 & c_0 \end{pmatrix}.\end{aligned}\tag{7.1}$$

We have

$$\begin{aligned}(\Upsilon_2^X)^{-1} &= \frac{1}{\det(\Upsilon_2^X)} \begin{pmatrix} b_0 c_0 - r_3^2 & r_2 r_3 - c_0 r_1 & r_1 r_3 - b_0 r_2 \\ r_2 r_3 - c_0 r_1 & a_0 c_0 - r_2^2 & r_1 r_2 - a_0 r_3 \\ r_1 r_3 - b_0 r_2 & r_1 r_2 - a_0 r_3 & a_0 b_0 - r_1^2 \end{pmatrix} \\ &:= \frac{1}{\det(\Upsilon_2^X)} \begin{pmatrix} a & \rho_1 & \rho_2 \\ \rho_1 & b & \rho_3 \\ \rho_2 & \rho_3 & c \end{pmatrix}.\end{aligned}$$

By elementary calculation, we get

$$\begin{aligned}\rho_1 &= \frac{1}{4} \tilde{\Lambda}_{23}, & \rho_2 &= \frac{1}{4} \tilde{\Lambda}_{24}, & \rho_3 &= \frac{1}{4} \tilde{\Lambda}_{12}, \\ a + \rho_1 + \rho_2 &= -\frac{1}{4} \tilde{\Lambda}_{34}, \\ b + \rho_1 + \rho_3 &= -\frac{1}{4} \tilde{\Lambda}_{13}, \\ c + \rho_2 + \rho_3 &= -\frac{1}{4} \tilde{\Lambda}_{14}.\end{aligned}\tag{7.2}$$

Then,

$$\begin{aligned}(\Lambda')_{12} \tilde{A}^2 + (\tilde{\Lambda})_{12}^2 &= 16 [a_0 \cdot \det(\Upsilon_2^X) + \rho_3^2] \\ &= 16(a_0 b_0 - r_1^2)(a_0 c_0 - r_2^2) \\ &= \{4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2\} \\ &\quad \cdot \{4(\Lambda')_{12}(\Lambda')_{24} - [(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]^2\}.\end{aligned}\tag{7.3}$$

By

$$\begin{aligned}(\tilde{\Lambda})_{12} &= 4(r_1 r_2 - a_0 r_3), \\ 4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2 &= 4(a_0 c_0 - r_2^2),\end{aligned}\tag{7.4}$$

and

$$4(\Lambda')_{12}(\Lambda')_{24} - [(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]^2 = 4(a_0b_0 - r_1^2), \quad (7.5)$$

we get

$$\begin{aligned} & -2(\Lambda')_{12} + (\Lambda')_{13} + (\Lambda')_{14} + (\Lambda')_{23} + (\Lambda')_{24} - 2(\Lambda')_{34} \\ & + \frac{(\tilde{\Lambda})_{12}[(\Lambda')_{13} + (\Lambda')_{23} - (\Lambda')_{12}]}{4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2} + \frac{(\tilde{\Lambda})_{12}[(\Lambda')_{14} + (\Lambda')_{24} - (\Lambda')_{12}]}{4(\Lambda')_{12}(\Lambda')_{24} - [(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]^2} \\ = & 4r_3 - 2r_1 - 2r_2 + \frac{2(r_1r_2 - a_0r_3)(c_0 - r_2)}{a_0c_0 - r_2^2} + \frac{2(r_1r_2 - a_0r_3)(b_0 - r_1)}{a_0b_0 - r_1^2} \\ = & \frac{2r_2(c_0r_1 - r_2r_3 + a_0r_3 - r_1r_2 - a_0c_0 + r_2^2)}{a_0c_0 - r_2^2} + \frac{2r_1(b_0r_2 - r_1r_3 + a_0r_3 - r_1r_2 - a_0b_0 + r_1^2)}{a_0b_0 - r_1^2} \\ = & \frac{-2r_2(b + \rho_1 + \rho_3)}{a_0c_0 - r_2^2} + \frac{-2r_1(c + \rho_2 + \rho_3)}{a_0b_0 - r_1^2} \\ = & \frac{(\tilde{\Lambda})_{13}[(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]}{4(\Lambda')_{12}(\Lambda')_{23} - [(\Lambda')_{12} + (\Lambda')_{23} - (\Lambda')_{13}]^2} + \frac{(\tilde{\Lambda})_{14}[(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]}{4(\Lambda')_{12}(\Lambda')_{24} - [(\Lambda')_{12} + (\Lambda')_{24} - (\Lambda')_{14}]^2}. \quad (7.6) \end{aligned}$$

7.2 Calculation of a double integral

In this subsection, we let $A > 0$ and

$$\Psi = \begin{pmatrix} a_1 & c_1 \\ c_1 & b_1 \end{pmatrix}$$

be a positive-definite matrix. Define

$$\mathcal{I} = \int_0^\infty \int_0^\infty e^{-\frac{a_1y^2 + b_1z^2 + 2c_1yz}{2A^2}} dydz.$$

Lemma 7.1 *We have*

$$\mathcal{I} = \frac{A^2}{\sqrt{a_1b_1 - c_1^2}} \arccos \left(\frac{c_1}{\sqrt{a_1b_1}} \right). \quad (7.7)$$

Proof. The two eigenvalues of Ψ are

$$\lambda_1 = \frac{a_1 + b_1 + \sqrt{(a_1 - b_1)^2 + 4c_1^2}}{2}, \quad \lambda_2 = \frac{a_1 + b_1 - \sqrt{(a_1 - b_1)^2 + 4c_1^2}}{2}.$$

Then, there exists an orthogonal matrix O such that

$$O \begin{pmatrix} a_1 & c_1 \\ c_1 & b_1 \end{pmatrix} O^T = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}. \quad (7.8)$$

Denote

$$O = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix},$$

where $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. One can check that if $c_1 = 0$, then $\theta = 0$; if $c_1 \neq 0$, then

$$\tan \theta = \frac{\lambda_1 - a_1}{c_1} = \frac{\sqrt{(a_1 - b_1)^2 + 4c_1^2} - (a_1 - b_1)}{2c_1},$$

which implies that if $c_1 > 0$, then $\theta \in (0, \frac{\pi}{2})$; if $c_1 < 0$, then $\theta \in (-\frac{\pi}{2}, 0)$.

Define

$$Q = \begin{pmatrix} \sqrt{\lambda_2} & 0 \\ 0 & \sqrt{\lambda_1} \end{pmatrix} O = \begin{pmatrix} \sqrt{\lambda_2} \cos \theta & \sqrt{\lambda_2} \sin \theta \\ -\sqrt{\lambda_1} \sin \theta & \sqrt{\lambda_1} \cos \theta \end{pmatrix}.$$

By (7.8), we get

$$\begin{aligned} Q \begin{pmatrix} a_1 & c_1 \\ c_1 & b_1 \end{pmatrix} Q^T &= \begin{pmatrix} \sqrt{\lambda_2} & 0 \\ 0 & \sqrt{\lambda_1} \end{pmatrix} O \begin{pmatrix} a_1 & c_1 \\ c_1 & b_1 \end{pmatrix} O^T \begin{pmatrix} \sqrt{\lambda_2} & 0 \\ 0 & \sqrt{\lambda_1} \end{pmatrix} \\ &= \begin{pmatrix} \sqrt{\lambda_2} & 0 \\ 0 & \sqrt{\lambda_1} \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} \sqrt{\lambda_2} & 0 \\ 0 & \sqrt{\lambda_1} \end{pmatrix} \\ &= \begin{pmatrix} \lambda_1 \lambda_2 & 0 \\ 0 & \lambda_1 \lambda_2 \end{pmatrix}. \end{aligned}$$

Define $\begin{pmatrix} y \\ z \end{pmatrix} = Q^T \begin{pmatrix} u \\ v \end{pmatrix}$. Then

$$\mathcal{I} = \sqrt{\lambda_1 \lambda_2} \iint_{D_Q} e^{-\frac{\lambda_1 \lambda_2 (u^2 + v^2)}{2A^2}} du dv,$$

where

$$D_Q = \left\{ (u, v) \in \mathbb{R}^2 : 0 \leq \sqrt{\lambda_2}(\cos \theta)u - \sqrt{\lambda_1}(\sin \theta)v, \ 0 \leq \sqrt{\lambda_2}(\sin \theta)u + \sqrt{\lambda_1}(\cos \theta)v \right\}.$$

If $c_1 = 0$, then $\theta = 0$ and hence $D_Q = \{(u, v) \in \mathbb{R}^2 : u \geq 0, v \geq 0\}$. We have

$$\begin{aligned} \mathcal{I} &= \sqrt{\lambda_1 \lambda_2} \int_0^{\frac{\pi}{2}} \int_0^\infty r e^{-\frac{\lambda_1 \lambda_2 r^2}{2A^2}} dr d\phi \\ &= \frac{\pi A^2}{2\sqrt{\lambda_1 \lambda_2}}. \end{aligned}$$

If $c_1 > 0$, then $\theta \in (0, \frac{\pi}{2})$. We have

$$D_Q = \left\{ (u, v) \in \mathbb{R}^2 : -\sqrt{\frac{\lambda_2}{\lambda_1}} \cdot (\tan \theta)u \leq v \leq \sqrt{\frac{\lambda_2}{\lambda_1}} \cdot \frac{u}{\tan \theta} \right\}.$$

Then,

$$\begin{aligned}
\mathcal{I} &= \sqrt{\lambda_1 \lambda_2} \int_{\arctan(-\sqrt{\frac{\lambda_2}{\lambda_1}} \tan \theta)}^{\arctan(\sqrt{\frac{\lambda_2}{\lambda_1}} \frac{1}{\tan \theta})} \int_0^\infty r e^{-\frac{\lambda_1 \lambda_2 r^2}{2A^2}} dr d\phi \\
&= \frac{A^2}{\sqrt{\lambda_1 \lambda_2}} \arctan \left(\frac{\sqrt{\lambda_1 \lambda_2}}{(\lambda_1 - \lambda_2) \cos \theta \sin \theta} \right) \\
&= \frac{A^2}{\sqrt{a_1 b_1 - c_1^2}} \arctan \left(\frac{\sqrt{a_1 b_1 - c_1^2}}{c_1} \right) \\
&= \frac{A^2}{\sqrt{a_1 b_1 - c_1^2}} \arccos \left(\frac{c_1}{\sqrt{a_1 b_1}} \right).
\end{aligned}$$

If $c_1 < 0$, then $\theta \in (-\frac{\pi}{2}, 0)$. We have

$$D_Q = \left\{ (u, v) \in \mathbb{R}^2 : -\sqrt{\frac{\lambda_2}{\lambda_1}} \cdot (\tan \theta) u \leq v, \sqrt{\frac{\lambda_2}{\lambda_1}} \cdot \frac{u}{\tan \theta} \leq v \right\}.$$

Then,

$$\begin{aligned}
\mathcal{I} &= \sqrt{\lambda_1 \lambda_2} \int_{\arctan(-\sqrt{\frac{\lambda_2}{\lambda_1}} \tan \theta)}^{\pi + \arctan(\sqrt{\frac{\lambda_2}{\lambda_1}} \frac{1}{\tan \theta})} \int_0^\infty r e^{-\frac{\lambda_1 \lambda_2 r^2}{2A^2}} dr d\phi \\
&= \frac{A^2}{\sqrt{\lambda_1 \lambda_2}} \left\{ \pi - \arctan \left(\frac{-\sqrt{\lambda_1 \lambda_2}}{(\lambda_1 - \lambda_2) \cos \theta \sin \theta} \right) \right\} \\
&= \frac{A^2}{\sqrt{a_1 b_1 - c_1^2}} \left\{ \pi - \arctan \left(\frac{-\sqrt{a_1 b_1 - c_1^2}}{c_1} \right) \right\} \\
&= \frac{A^2}{\sqrt{a_1 b_1 - c_1^2}} \arccos \left(\frac{c_1}{\sqrt{a_1 b_1}} \right).
\end{aligned}$$

Therefore, (7.7) holds for any case. □

7.3 Mathematica code for verifying (3.7)

```

In[1]:= Expand[((1 - b)^2 - (1 - b)*(1 - a + 1 - c + 1 - d + 1 - f -
2*(1 - e)) + (a - d)*(c - f))*(1 -
a + (1 - d) - (1 -
b))*(4*(1 - a)*(1 - e) - (1 - a - e + c)^2) + ((1 - c)^2 - (1 -
c)*(1 - a + 1 - b + 1 - e + 1 - f - 2*(1 - d)) + (a - e)*(b -
f))*(1 -
a + (1 - e) - (1 - c))*(4*(1 - a)*(1 - d) - (1 - a - d + b)^2) -
2*((1 - d)^2 - (1 - d)*(1 - a + 1 - b + 1 - e + 1 - f -
2*(1 - c)) + (a - b)*(e - f))*(1 -
b)*(4*(1 - a)*(1 - e) - (1 - a - e + c)^2) -
2*((1 - e)^2 - (1 - e)*(1 - a + 1 - c + 1 - d + 1 - f -

```

```

2*(1 - b)) + (a - c)*(d - f))*(1 -
c)*(4*(1 - a)*(1 - d) - (1 - a - d + b)^2) -
2*((1 - b)^2 - (1 - b)*(1 - a + 1 - c + 1 - d + 1 - f -
2*(1 - e)) + (a - d)*(c - f))*(1 -
d)*(4*(1 - a)*(1 - e) - (1 - a - e + c)^2) -
2*((1 - c)^2 - (1 - c)*(1 - a + 1 - b + 1 - e + 1 - f -
2*(1 - d)) + (a - e)*(b - f))*(1 -
e)*(4*(1 - a)*(1 - d) - (1 - a - d + b)^2) -
2*(1 - f)*(4*(1 - a)*(1 - d) - (1 - a - d + b)^2)*(4*(1 - a)*(1 -
e) - (1 - a - e + c)^2)]
Out[1]= 0

```

7.4 Derivation of inequality (5.23)

Let $K(u, \theta)$ and $P(u, \theta)$ be defined as in (5.10) and (5.12), respectively. We have

$$\begin{aligned}
& \partial_u P(u, \theta) \\
&= \frac{1}{\theta^2(\cos \theta - \cos u)^4} \left\{ (\cos \theta - \cos u)^2 [2u\theta(1 - \cos \theta \cos u) - 2u \sin \theta (\cos \theta - \cos u) \right. \\
&\quad \left. - \sin u(u^2 + \theta^2)(\sin \theta - \theta \cos \theta) - 2u\theta^2 \cos u \sin \theta] \right. \\
&\quad \left. + 2 \sin u (\cos \theta - \cos u) [(u^2 - \theta^2) \sin \theta (\cos \theta - \cos u) + 2u\theta^2 \sin u \sin \theta - (u^2 + \theta^2)\theta(1 - \cos \theta \cos u)] \right\} \\
&= \frac{1}{\theta^2(\cos \theta - \cos u)^4} \left\{ (\cos \theta - \cos u)^2 [(2u\theta(1 - \cos \theta \cos u) + 2 \sin u \sin \theta(u^2 - \theta^2)) \right. \\
&\quad \left. - (2u \sin \theta (\cos \theta - \cos u) + \sin u(u^2 + \theta^2)(\sin \theta - \theta \cos \theta) + 2u\theta^2 \cos u \sin \theta)] \right. \\
&\quad \left. + (\cos \theta - \cos u) [4u\theta^2 \sin^2 u \sin \theta - 2(u^2 + \theta^2)\theta \sin u(1 - \cos \theta \cos u)] \right\}. \tag{7.9}
\end{aligned}$$

On the other hand, by elementary calculation, we get

$$\partial_u K(u, \theta) = \frac{2u\theta(1 - \cos \theta \cos u) - (u^2 + \theta^2) \sin \theta \sin u + 2(\cos \theta - \cos u)(u \sin \theta - \theta \sin u)}{\theta(\cos \theta - \cos u)^2},$$

and

$$\begin{aligned}
& \partial_\theta \partial_u K(u, \theta) \\
&= \frac{1}{\theta^2(\cos \theta - \cos u)^4} \left\{ (\cos \theta - \cos u)^2 [\sin u(u^2 + \theta^2)(\sin \theta - \theta \cos \theta) - 2u \sin \theta (\cos \theta - \cos u) \right. \\
&\quad \left. - 4\theta^2 \sin \theta \sin u + 2u\theta(\theta \cos u \sin \theta + 1 - \cos \theta \cos u)] \right. \\
&\quad \left. + (\cos \theta - \cos u) [4u\theta^2 \sin \theta (1 - \cos \theta \cos u) - 2\theta \sin^2 \theta \sin u(u^2 + \theta^2)] \right\}. \tag{7.10}
\end{aligned}$$

Then, we obtain by (5.12), (7.9) and (7.10) that

$$\begin{aligned}
& \partial_u P(u, \theta) \\
&= \partial_u \partial_\theta K(u, \theta) \\
&= \frac{\partial_u P(u, \theta) + \partial_\theta \partial_u K(u, \theta)}{2} \\
&= \frac{1}{\theta^2 (\cos \theta - \cos u)^4} \left\{ (\cos \theta - \cos u)^2 [2u\theta(1 - \cos \theta \cos u) - 2u \sin \theta (\cos \theta - \cos u) \right. \\
&\quad \left. + \sin u \sin \theta (u^2 - 3\theta^2)] \right. \\
&\quad \left. + (\cos \theta - \cos u) [2u\theta^2 \sin \theta (\sin^2 u + 1 - \cos \theta \cos u) - (u^2 + \theta^2) \theta \sin u (\sin^2 \theta + 1 - \cos \theta \cos u)] \right\}.
\end{aligned}$$

Therefore, $\partial_u P(u, \theta) > 0$ for any $u \in [0, \theta) \cup (\theta, 2\pi - \theta)$ and $\theta \in (0, \pi)$ is equivalent to (5.23) holds.

Acknowledgments This work was supported by the Natural Sciences and Engineering Research Council of Canada, and the National Natural Science Foundation of China (Grant No. 11771309, No. 11871184 and No. 61973096).

References

- [1] A.V. Balakrishnan. Research problem No. 9: geometry. *Bull. Amer. Math. Soc.* 69 737-738 (1963).
- [2] A.V. Balakrishnan. Signal selection for space communication channels. In: *Advances in Communication Systems*, (ed. by A.V. Balakrishnan), Academic Press, 1-31 (1965).
- [3] K. Böröczky Jr. and R. Schneider. Circumscribed simplices of minimal mean width. *Beiträge Algebra Geom.* 48 217-224 (2007).
- [4] J. Brandts and A. Cihangir. Geometric aspects of the symmetric inverse M-matrix problem. *Linear Algebra Appl.* 506 33-81 (2016).
- [5] B. Dunbridge. Asymmetric signal design for the coherent Gaussian channel. *IEEE Trans. Inform. Theory.* IT-13 422-431 (1967).
- [6] R. Eldan. A two-sided estimate for the Gaussian noise stability deficit. *Invent. Math.* 201 561-624 (2015).
- [7] M. Fiedler. Über qualitative Winkeleigenschaften der Simplexe. *Czechoslovak Math. J.* 7(82) 463-478 (1957).
- [8] E.N. Gilbert. A comparison of signaling alphabets. *Bell System Tech. J.* 31 504-522 (1952).
- [9] E.D. Gluskin. Extremal properties of orthogonal parallelepipeds and their applications to the geometry of Banach spaces. *Math. USSR Sbornik.* 64 85-96 (1989).

- [10] P. Gritzmann and V. Klee. On the complexity of some basic problems in computational convexity: II. Volume and mixed volumes. In: T. Bisztriczky et al. (Eds.), *Polytopes: Abstract, Convex and Computational* (Scarborough 1993), pp. 373-466, NATO ASI Series C, vol. 440, Kluwer (1994).
- [11] D. Hug and R. Schneider. Large typical cells in Poisson-Delaunay mosaics. *Rev. Roumaine Math. Pures Appl.* 50 657-670 (2005).
- [12] R. Latała and D. Matlak. Royen's proof of the Gaussian correlation inequality. *Geometric Aspects of Functional Analysis*, 265-275. Springer (2017).
- [13] W.V. Li. A Gaussian correlation inequality and its applications to small ball probabilities. *Electr. Comm. Probab.* 4 111-118 (1999).
- [14] A.E. Litvak. Around the simplex mean width conjecture. *Analytic Aspects of Convexity*. Springer INdAM Series, 25 73-84 (2018).
- [15] T. Royen. A simple proof of the Gaussian correlation conjecture extended to multivariate gamma distributions. *Far East J. Theoret. Stat.* 48 139-145 (2014).
- [16] Q.M. Shao. A Gaussian correlation inequality and its applications to the existence of small ball constant. *Stoch. Proc. Appl.* 107 269-287 (2003)
- [17] Z. Šidák. Rectangular confidence regions for the means of multivariate normal distributions. *J. Am. Stat. Assoc.* 62 626-633 (1967).
- [18] Y.L. Tong. *The Multivariate Normal Distribution*. Springer (1990).
- [19] 1978 USAMO Problems/Problem 4. https://artofproblemsolving.com/wiki/index.php/1978_USAMO_Problems/Problem_4.
- [20] D.W. Walkup. A simplex with a large cross-section. *Amer. Math. Monthly* 75 34-36 (1968).
- [21] C.L. Weber. *Elements of Detection and Signal Design*. McGraw-Hill (1968).