

ARTICLE TYPE

Accuracy of magnitudes in pre-telescopic star catalogues

Philipp Protte | Susanne M Hoffmann

¹Physikalisch-Astronomische Fakultät,
Friedrich-Schiller-Universität
Jena, Germany

Correspondence

Susanne M Hoffmann, Email:
susanne.hoffmann@uni-jena.de

Historical star magnitudes from catalogues by Ptolemy (137 AD), al-Šūfī (964) and Tycho Brahe (1602/27) are converted to the Johnson V-mag scale and compared to modern day values from the HIPPARCOS catalogue. The deviations (or “errors”) are tested for dependencies on three different observational influences. The relation between historical and modern magnitudes is found to be linear in all three catalogues as it had previously been shown for the Almagest data by Hearnshaw (1999). A slight dependency on the colour index (B-V) is shown throughout the data sets and al-Šūfī’s as well as Brahe’s data also give fainter values for stars of lower culmination height (indicating extinction). In all three catalogues, a star’s estimated magnitude is influenced by the brightness of its immediate surroundings. After correction for the three effects, the remaining variance within the magnitude errors can be considered as approximate accuracy of the pre-telescopic magnitude estimates. The final converted and corrected magnitudes are available via the Vizier catalogue access tool (Ochsenbein, Bauer, & Marcout, 2000).

KEYWORDS:

magnitudes, history of astronomy, almagest, extinction, naked eye observation

1 | INTRODUCTION

Many processes in astronomy have long timescales, especially questions on the evolution of stars (just recently, the timescale of Betelgeuse’s supernova was publicly discussed). Further examples include close binary systems such as cataclysmic variables (CVs) and their nova eruption behaviour (S. M. Hoffmann, Vogt, & Protte, 2019; Shara et al., 2017; Vogt, Hoffmann, & Tappert, 2019), or even supernovae in close binary systems which have the potential to eject runaway stars (Neuhäuser, Gießler, & Hambaryan, 2019). All these questions on the evolution of astronomical objects require long-term observations but our telescopic surveys only reach back for a few decades (in cases of CVs) or roughly two centuries (in cases of sunspot observations (Neuhäuser, Arlt, & Richter, 2018; Neuhäuser & Neuhäuser, 2016)).¹ Aiming for conclusions on long-term evolution it is, thus, desirable to

include data from non-telescopic observations which could possibly provide a much longer baseline: Far Eastern tradition, for instance, recorded transient phenomena (such as novae, supernovae, and comets) more systematically than the Western one. However, one of the biggest questions in these terms is the transformation of any ancient (or old) description of the *brightness* of the phenomenon. Only in very few cases (e.g. 437, SN 1572) the historical records mention daylight visibility. In a handful of cases (e.g. 1175, 1203, 1596, and 1603 according to Ho (1962), and (Xu, Pankenier, & Jiang, 2000, 129–146)), the description refers to the brightness giving Mars, Saturn, or bright stars like Capella (α Aur) and Antares (α Sco) as comparison.

Although we can look up the brightnesses of the planets and fixed stars in a modern star catalogue or model their brightness at a certain date with our knowledge on their variability, it appears worthwhile to study the accuracy of such historical estimates.

As commonly known, Argelander in the 19th century defined a clear method to estimate the magnitude of a given object by

¹Sporadic telescopic observations have, of course, existed for a bit longer but *systematic surveys* have not been common practice from the early beginning on.

comparing it to a couple of stars in the vicinity. The method(s) of earlier astronomers to derive the magnitude of a star or transient object are yet unknown. As, therefore, the numbers in historical star catalogues are hardly reproducible, we try to derive a better understanding of their scattering, error bars, and their dependencies.

1.1 | Dependencies and open questions

In particular, the visual appearance of a celestial point source depends on many influences, e. g. the brightness of the background, the local and temporal conditions of the atmosphere and the constitution of the observer's eye. Currently, we cannot consider the observer's eye and we are not even sure that the author of a text book really observed every data point on his own and that not students or assistants were helping or even taking over the measurement. The interplay of the human eye's lens and the atmospheric conditions cause optical effects such as (i) reddening of stars close to the horizon (extinction), (ii) blurring of bright stars due to humidity or sandstorm, and (iii) the impression of rays, horns, or fuzziness of bright objects (see APPENDIX A:) which might affect the estimate of the magnitude. The background brightness of the sky depends on the density of stars in a particular area, the zodiacal light as well as geophysical influences such as the omnipresent airglow (discovered by Ångström (1869)) and the presence or absence of meteor showers (Siedentopf, 1959). As this work aims to make historical star catalogues usable for modern research, our goal is to find an algorithm for how to deal with historical magnitudes. We assume that magnitudes for a star catalogue have not been observed only one time by one person but cross checked by the assistants of the historical book author. As even Ptolemy mentions the difficulties (and errors) of observations close to the horizon, we assume that they estimated magnitudes at highest possible altitude for a given star. Therefore, the remaining influences to be considered in Section 3 are the dependency of the human brightness estimation on the following questions:

- Redder stars appear fainter, so how does the colour of the star influence the estimate? Can the historical numbers be improved by application of a colour correction?
- The atmosphere influences the appearance by extinction: Does a correction improve the conversion of historical magnitudes into modern ones?
- Observational bias of the environment: How does the presence of bright stars and a background of many faint stars (e.g. in the Milky Way) influence the historical brightness estimation?

We analyze these questions in Section 3 by using the data of three historical star catalogues introduced in Section 2 which,

of course, had already been analyzed by other scholars before us. Their results are, therefore, summarized in the following subsection.

1.2 | Placement among previous works

The idea to represent different brightnesses by numbers goes back to Antiquity. Pliny the Elder's (+1st century) words suggest that Hipparchus (−2nd century) might have measured these magnitudes but this cannot be verified or falsified (S. Hoffmann, 2017, p. 92 and 194). The first surviving appearance of the magnitude scale is in Ptolemy's *Almagest* from the 2nd century (Hearnshaw, 1996, p. 4) and from there it was copied to the Arabic and Latin science culture but the numbers always remained estimations. With the dawning of electric photometry, astrophotography and the necessity of exact values from telescopic observations, the 19th century took some efforts to develop a mathematically exact scale. Pogson (1856)'s system finally prevailed but it differs from the ancient scale because mathematics in the meantime had introduced negative numbers and the zero. Additionally, any type of logarithmic law can only approximate the human sense and it neglects personal influences of the observer (guessing errors). There is no easy conversion from historical magnitudes because the new scale was used for the new star catalogues.

Although the *Almagest*'s star catalogue has thoroughly been analysed and discussed with respect to its record of star positions, there are only a handful (e.g. Hearnshaw, 1999; Schaefer, 2013) of recent investigations into the star's magnitudes. Even fewer authors consider other pre-telescopic magnitude estimations, which are given most notably by al-Šūfī and Tycho Brahe. Recently, Verbunt & van Gent (2010a, 2012a) released online versions of the three catalogues by Ptolemy, Ulugh Bēg and Tycho Brahe with Ulugh Bēg's catalogue containing magnitude estimations that were adopted from al-Šūfī's *Book of the fixed stars*.

The computer-readable data makes it easier than ever to evaluate and statistically compare the three catalogues. If the data optimally converted and corrected for systematic deviations, it might be possible to utilise the ancient magnitudes for investigations of stellar evolution and variability (as examples for such endeavours see Mayer (1984) and Hertzog (1984) although the former's results were later refuted by Hearnshaw (1999)). All the effects on magnitude estimates, mentioned at the end of Section 1.1 have already been analysed during the second half of the 19th and the first half of the 20th century (for an elaborate example see Zinner, 1926) but there is no systematic query of all three catalogues based on the conversion method, introduced by Hearnshaw (1999) while using modern computer-aided statistical procedures.

2 | ON THE ORIGINAL CATALOGUES

We evaluated three star catalogues with measurements from different epochs and cultural backgrounds, beginning with the one, featured in Ptolemy’s *Μαθηματικὴ Σύνταξις* (engl.: *Mathematical treatise*, commonly known by its Arabic name *Almagest*, 137 AD). This work contains the oldest extensive data set of stellar brightness and Ptolemy also was the first astronomer to verifiably make use of a numeric scale: the magnitudes. He assigned the brightest stars to the magnitude 1 and the remaining ones into five gradually fainter classes, labelled 2–6. For some stars he added qualifiers, saying a star was either slightly brighter or fainter than the given magnitude.

The second catalogue is the one by Ulugh Bēg from around 1437 AD which contained the first independent, comprehensive position measurements in 1300 years, yet adopted (see Knobel, 1917) its magnitude data from Abd al-Rahman al-Šūfī’s ‘Book of fixed Stars’ (for a modern english translation see Hafez, 2010), which he most likely composed around 964 AD. in the city of Shiraz (Hafez, 2010, p.64). His list of stars is explicitly based on Ptolemy’s catalogue, containing almost the same set of stars with positions, only corrected for precession. However, al-Šūfī was only the second astronomer to systematically assign magnitudes to all the entries in his catalogue, using the same numerical scale, as Ptolemy. Al-Šūfī’s catalogue served as an important source for many subsequent Islamic-Arabic astronomers who used his data or cited his texts (see Hafez, 2010, p.66 ff). One of those was Ulugh Bēg, who, when he compiled his own star catalogue in 1437, adopted the magnitudes (and in 27 cases also the positions (Verbunt & van Gent, 2012b)) from the *Book of the fixed Stars*.

Lastly, we included Tycho Brahe’s star catalogue from 1602/1627 which again consists of newly gathered data for positions and magnitudes. Brahe was the first modern European scholar to compile a comprehensive original star catalogue. The results were first published as a 777-star catalogue in 1602 shortly after his death but a handwritten copy of a more extensive catalogue had already been sent to several astronomers during the 1590’s. Finally, in 1627 Johannes Kepler published an edition of the list containing 1004 stars which was very similar to the manuscript version (Verbunt & van Gent, 2010a). While Tycho set new standards of precision for position measurements, he adopted Ptolemy’s magnitude scale with no finer graduation than those of his predecessors. As pointed out by most analyses of the different versions of his catalogue (e.g. Baily, 1843), the 1627 version even omits the brighter-/fainter-qualifiers that were still included in the previous release.

Figure 1 shows the chronology and data transfer of the four catalogues.




FIGURE 1 Transfer of position and magnitude data between the four ancient catalogues. Dashed arrows indicate an uncertain amount of influence, bold arrows show that data was copied almost “word-for-word”.

3 | ANALYSIS OF THE MAGNITUDES

For the historical magnitudes we used the data, given by Verbunt & van Gent (2010b, 2012b). Additional modern data (e.g. exact position, V-mag and colour index) have been taken from the VizieR release of the HIPPARCOS catalogue (ESA, 1997). Verbunt and van Gent used the translation of Ulugh Bēg’s catalogue by Knobel (1917) for their digitalisation and the latter, knowing that Ulugh Bēg adapted al-Šūfī’s magnitudes, entered them directly from al-Šūfī’s *Book of the fixed Stars* to minimise translation errors. Therefore the catalogue designated as Ulugh Bēg’s is actually a hybrid catalogue and as only the magnitudes are analysed, they are referred to as ‘al-Šūfī’s magnitudes’ and abbreviated as m_S and the catalogue as S .

Brahe’s catalogue was released in different editions, where only his 777-star list from 1602 includes qualifiers to specify magnitudes beyond 1-mag steps. The data given by Verbunt and van Gent includes a total of 1007 stars, merged from all available editions and the qualifiers were included in our analysis whenever they were given.

For the analysis we excluded double entries as well as stars without safe modern identification or magnitude (e.g. for double stars). Furthermore, stars designated as “faint” or “nebulous” were left out and the two brightest stars (Sirius and Canopus) were excluded from most analyses as significant “outliers”. The used data sets are summarised in Table 1, where the last set (PSB) only includes stars that were existent in all three catalogues and had concordant modern identifications. This set was used to examine the covariance between the historical catalogues. Different magnitude designations are used throughout the analysis: m_{HIP} are the modern V-mag values taken from the HIPPARCOS catalogue, while m_{old} and m




FIGURE 2 Relation between historical and modern magnitude scales for each star of the data subsets P, S and B. Although the scatter plot does not look particularly linear, the regression can be justified (see text) and is used to convert the historical values to the modern magnitude scale.

designate the historical magnitudes before and after conversion, respectively. Additional indices (P,S,B) might be added when talking about data from a certain author. Finally, we define $\delta m = m - m_{\text{HIP}}$.

TABLE 1 The analysed catalogues. Identifications taken from Verbunt & van Gent (2010b, 2012b). PSB is the intersection of P,S,B, containing only stars that are concordantly identified in all three catalogues.

Abbr.	Included authors	# of stars
P	Ptolemy	990
S	Ulugh Bēg	988
B	Brahe	937
PSB	Ptolemy+Ulugh Bēg +Brahe	695

3.1 | Conversion of the magnitude scales

The historical magnitude values are not identical with modern, photometric V-band magnitudes. While today's magnitudes of the stars in our reduced catalogue cover a continuous range from ~ 0 to ~ 6 , in the pre-telescopic era they were based on estimated assignments into 6 discrete groups. Nevertheless intermediate steps between these groups were used by all three authors as differently formulated 'qualifiers' which indicate if the star is slightly brighter or fainter than the denoted magnitude.

The disparate definitions result in two problems: on the one hand, the data shows a different range for both magnitudes (roughly $1 \dots 6 \leftrightarrow 0 \dots 6$). Therefore, a direct comparison of both values in form of a difference $m_{\text{old}} - m_{\text{HIP}}$ will be biased towards showing large positive values for brighter stars. To minimise any dependency of the difference on a star's brightness, an adequate conversion formula for m_{old} is needed.

The other problem, however, has to be tackled first: The intermediate qualifiers, given by the historical observers, do not imply an exact value. Trying to convert the qualifiers into numerical divergence, most previous authors added $+0.33$ mag for the 'fainter-qualifier' and -0.33 mag for the 'brighter-qualifier', but also ± 0.3 mag and ± 0.5 mag have been applied. In an attempt to find the best approximation, we compared the 'two-step-system' (± 0.33 mag) with the 'one-step-system' (± 0.5 mag).

Looking at the average modern magnitude of each group of stars with qualifiers (e.g. 2(f) – stars a little fainter than 2nd mag, or 3(b) – stars a little brighter than 3rd mag), the 'two-step-system' is found to show several inconsistencies. For example, the Almagest's 2(f)-stars which would be identified with $m_{\text{old}}^P = 2.33$ mag are fainter on average than the 3(b)-stars, identified with $m_{\text{old}}^P = 2.67$ mag. The 'one-step-system' on the other hand, is consistent in almost all cases and is therefore adopted for our analysis. The applied values of m_{old} and m_{HIP} are shown for each star in the scatter plots of Figure 2. It might be added at this point that most of the following analysis was done with 0.33 mag-steps before the 'one-step-system' was chosen. The differences in the results were negligible in almost all cases.

The modern magnitude scale is logarithmic in regard to the

light flux, a fact that corresponds to (and historically derives from) the logarithmic perception of brightness in the human eye (Weber-Fechner-Law). Therefore the relation between m_{old} and m_{HIP} should be approximately linear which is not evident in any of the three sub-figures of Figure 2, due to the large scattering. But even averaging the m_{HIP} for each step of m_{old} , as it has been done in many previous works (most recently Schaefer, 2013), does not yield a linear correlation, but rather implies a curved function. Thus, instead of trying to find a consistent conversion *formula* from m_{old} to m , most authors applied an empirical method: The modern averages are immediately used as m for every star within the respective step.

Contrary to all previous (and some subsequent) studies, Hearnshaw (1999) showed that the linear relation can indeed be found in Ptolemy's data when switching the dependent with the independent variable. He argues that taking the mean modern values for each historical magnitude is statistically invalid because the variable with larger uncertainties should be averaged. A quick look at the brightest² known variable stars shows that their variability (see Figure 3) is usually much smaller than the resulting uncertainties of the historic magnitudes (see Figure 9). Applying the same procedure as




FIGURE 3 Histogram showing the distribution of variability amplitudes among the 1,137 known variable stars of $m_{max}, m_{min} < 6$ mag. The possible magnitude error, induced by variability is small, compared to the overall uncertainties of historical magnitudes in most cases.

Hearnshaw (1999), we found linear relations for m_{old}^P and for m_{old}^S and m_{old}^B , as well which can be seen in Figure 4 where m_{old} is averaged for bins of m_{HIP} . It is therefore justified to convert the m_{old} to a new variable called m by applying a

linear conversion formula:

$$m = a \cdot m_{old} + t \quad \text{with} \quad a = \frac{1}{b} \quad t = -\frac{t'}{b} \quad (1)$$

Where b and t' are the regression coefficients from Figure 2 :

$$\begin{aligned} b_P &= 0.75 \pm 0.02 & t'_P &= 0.92 \pm 0.08 \\ b_S &= 0.86 \pm 0.02 & t'_S &= 0.69 \pm 0.07 \\ b_B &= 0.93 \pm 0.02 & t'_B &= 0.43 \pm 0.10 \end{aligned} \quad (2)$$

From b_P we can calculate $a_P = 1.33^{+0.04}_{-0.03}$, meaning that one Ptolemaic magnitude corresponds to 1.33 modern magnitudes (concordant with $a_P = 1.36$, as found by Hearnshaw (1999)). In the same way, the values for the other two authors are: $a_S = 1.16^{+0.03}_{-0.02}$ and $a_B = 1.08^{+0.02}_{-0.03}$. After converting the historical magnitudes to the modern scale, we can now define the error variable $\delta m = m - m_{HIP}$ for each star.




FIGURE 4 Average historical magnitudes for modern-magnitude-bins of 0.1 mag in the data sets P, S, B. Error bars are the SEM where bin-size $N > 2$ and else the averaged SD of all other bins. The histograms in the background show the relative number of stars within each bin. The regression here is only for visualisation of the linearity while the actual coefficients are taken from Figure 2.

² $m_{max}, m_{min} < 6$ mag

3.2 | Analysis of the magnitude errors

3.2.1 | Variance and covariance

In a first step, the distribution of the magnitude errors δm is analysed for each of the three star catalogues P, S, B, as well as the common catalogue PSB. Table 2 gives mean values and standard deviations for both cases with the squared correlation coefficients between the δm added for the shared data set. The

TABLE 2 Mean values μ and std.dev. σ of the δm in the single-author catalogues (P, S, B) and for the shared catalogue (PSB), including squared correlation coefficients r^2 .

[mag]	P, S, B	PSB		
		δm_P	δm_S	δm_B
δm_P	$\mu : 0.00$ $\sigma : 0.79$	$\mu : -0.01$ $\sigma : 0.73$		
δm_S	$\mu : -0.01$ $\sigma : 0.64$	$r^2 : 0.51$	$\mu : -0.05$ $\sigma : 0.58$	
δm_B	$\mu : 0.02$ $\sigma : 0.76$	$r^2 : 0.18$	$r^2 : 0.16$	$\mu : 0.00$ $\sigma : 0.72$

mean values of δm for the three single catalogues come close to zero but al-Šūfi's data in the common list shows a slightly larger offset. This might be due to a selection effect in the shared catalogue where certain stars were omitted. The standard deviation shows similar values for Ptolemy's and Brahe's catalogues but a significantly slimmer scattering for al-Šūfi's magnitude errors. Finally, the r^2 -values indicate a strong correlation between Ptolemy's and al-Šūfi's magnitude errors while Brahe's data seems to be largely independent. The correlation is visualised in the scatter plots in Figure 5, including covariance-ellipses that contain $\sim 95\%$ of the data points. The




FIGURE 5 Correlation between the δm of the three catalogues within data set PSB. 2σ -covariance ellipses are given for each scatter plot.

surprise because al-Šūfi takes Ptolemy's magnitude estimation as a basis for his own (see also Figure 1). He even gives literal references like: "The fourth [star] [...] is much greater then [sic] 4th magnitude, but it was mentioned by Ptolemy as 4th magnitude exactly." (Hafez, 2010, p. 154). From these kind of comments it can be assumed that al-Šūfi only changed a star's magnitude if he deemed it distinctly erroneous and therefore left a large fraction of them unchanged, causing this dependency.

The distinctly weaker correlation between Ptolemy's and Brahe's data might as well be due to some of the latter's magnitudes being influenced by the *Almagest*.

3.2.2 | Dependence on the colour index

The transmission spectrum of the modern V-band filter is quite similar but not identical with the spectral sensitivity function of the human eye. In fact there are at least two such sensitivity functions (one for daylight- or *photopic* vision and one for night- or *scotopic* vision) and the V-band curve lies in between of them both. Generally, the photopic vision is employed under brighter ambient light and means that the eyes' cones are active and we can perceive colour. In contrast, the scotopic vision is the extreme case where our vision depends solely on the eyes' rods which can only differentiate between bright and dark but not detect colour (e.g. Clauss & Clauss, 2018, p. 178). As the rods are more sensitive to shorter wavelengths, the sensitivity function for the scotopic vision is shifted towards blue colours, making blue stars a little brighter than red stars of the same V-band magnitude. The upcoming analysis is restricted to stars of $m_{\text{HIP}} > 2.3$, excluding those stars which are bright enough to be seen with photopic vision (indicated by them having a colour to the naked eye)³. In this case we would expect the calculated δm to be something like a colour index between the V-band filter and the human eye-“filter” for $m > 2.3$ mag. A plot of two colour indices (e.g. $(I - J)$ against $(K - L)$) shows an almost linear relation with a slope ∂_{BV} that can be calculated from the effective wavelengths of the four filters (Ballesteros, 2012). We performed a linear regression analysis for plots of δm against the $(B - V)$ -values for the three catalogues.

To better visualise the tendency among the data, averages of δm were calculated for $(B - V)$ -bins of 0.1 mag. The result can be seen in the left column of Figure 6, together with a scatter plot of the entire catalogues. A linear model was fitted to the scatter plots and the parameters are shown in Table 3. From the slopes of the linear regression, it is possible to calculate the effective wavelength of the human eye (under the given premises). The three catalogues yield values from 527 nm to 532 nm with error bars of ± 6 nm. Both, the increase of δm for

correlation between Ptolemy and al-Šūfi should come as no

³The exact threshold of colour vision in terms of star magnitudes is not clearly defined but from observational experience, 2.3 mag could be an approximate value.

reddish stars as seen in Figure 6 , as well as the calculated effective wavelength falling short of the V-band filter ($\lambda_{eff} = 548$ nm), agree with the above assumption of predominantly scotopic vision. Nevertheless, the calculated wavelength is longer than what would be expected for exclusively scotopic vision (~ 507 nm, see CVRL, 1995). The calculated regression coefficients make it possible to systematically adjust m with regard to the colour index of each star (see section 4.1). Lastly, a similar analysis was attempted for the brighter stars, but given the small number of stars with $m < 2.3$ mag and the large standard deviation of δm , the statistical analysis did not yield significant results for any of the three catalogues.

TABLE 3 Parameters of the fitted models in Figure 6 . For the two linear models ∂ is the slope and θ the intercept. n is the number of stars included in each model and R^2 the fraction of variance, explained by the model. For statistical testing of the models, see 4.2.

	Colour		Extinction		Background	
	∂_{BV}	θ_{BV}	k_{fit}	∂_β	θ_β	
P	0.11	-0.06	-0.01	-0.37	-0.69	
S	0.14	-0.06	0.09	-0.41	-0.75	
B	0.14	-0.06	0.05	-0.30	-0.53	
	n	R^2	n	R^2	n	R^2
P	922	0.006	992	0.0003	990	0.024
S	925	0.017	990	0.022	989	0.045
B	886	0.011	938	0.016	936	0.016

3.2.3 | Extinction features

Even though the effect of atmospheric extinction is obvious in its existence for everyone who has watched the (night-) sky with some attention, no discussion of the effect can be found in any of the works containing the three catalogues. If the observers had completely ignored the extinction, it would have to be expected that stars with low culmination altitudes were estimated too faint. Schaefer (2013) analyses the dependency and comes to the conclusion that all three catalogues are in some way “corrected” for extinction⁴.

With the new conversion, a similar analysis is shown in the middle column of Figure 6 , as a scatter plot of all stars, and as averaged δm -values for 3° -bins of culmination altitude.

⁴That does not necessarily mean they were explicitly corrected by some formula or observational procedure but could also mean to just estimate slightly brighter magnitudes for low standing stars. Trying to observe stars at their highest position could also be considered such a correction and is presupposed for the following analysis.




FIGURE 6 The dependency of δm on colour index $B-V$ (**left**), maximum culmination altitude alt (**middle**) and background brightness β (**right**) within catalogues P, S and B. Grey scatter plots are the single stars and bold coloured dots are mean values of bins of the independent variable with SEM-error bars. The bold lines show models, fitted to the scatter plots for each dependency. Stars with $m_{HIP} < 2.3$ mag were excluded from the colour-analysis. For the middle column, the dashed lines show expected extinction coefficients of $k = 0.15$ and 0.25 mag/X. The β -bins correspond to the levels of grey in Figs. 7 and 8 . All model parameters can be found in Table 3 .

Schaefer gives the following extinction function:

$$\delta m = k \cdot \left[(\sin(alt) + 0.025 \cdot e^{-11 \cdot \sin(alt)})^{-1} - 1 \right] \quad (3)$$

With the extinction coefficient k , given in magnitudes per air-mass X and the horizontal altitude alt of the respective star. In application to the historical catalogues, the altitude (or rather the culmination point of a star) can be calculated from the geographic latitude ϕ of the observer and the declination δ of the star at the time of observation.

$$alt = 90\lambda - |\delta - \phi| \quad (4)$$

The extinction curves are plotted within each sub-figure for $k = 0.25 \text{ mag}/X$ (as suggested by Schaefer, 2013), as well as $k = 0.15 \text{ mag}/X$. Pickering (2002) assumes such a value for a pre-industrial atmosphere. Additionally, models according to equation 3 were fitted to the data in Figure 6 with the resulting parameters k_{fit} listed in Table 3. Schaefer's general result is reproduced with close to no (in fact even a slight but insignificant negative) extinction effect showing in the *Almagest*'s data. In contrast, an effect is clearly visible in al-Šūfī's magnitudes but it still falls short of the expected intensity for Schaefer's extinction coefficient. However, the data could almost agree with the lower extinction coefficient of $k = 0.15 \text{ mag}/X$. In Brahe's magnitude estimations, a weak extinction effect can be found but again, it falls way short of the plotted models. Although the effect is weaker than expected, it can be corrected for at least in al-Šūfī's and Brahe's data, as a clear systematic deviation can be found there.

It should be noted that the models, fitted in Figure 6 are very sensitive to single extreme δm at low altitudes, which might occur due to falsely identified stars or other sporadic errors. Furthermore, it also seems to be highly controversial what extinction coefficient would have to be expected for a pre-industrial atmosphere (see Hearnshaw, 1999; Pickering, 2002; Schaefer, 2013)⁵ and lastly, the actual effect found in each catalogue also depends on the exact method by which the magnitudes were estimated which can only be speculated about.

3.2.4 | Star maps and background brightness

The dependency of δm on the colour index, as well as the culmination altitude, are both effects that can be understood and modelled in a (bio-)physical sense. However, there seem to be further trends within the data, which can be found looking at the spatial distribution of the δm . Figures 7 and 8 show maps of all stars within catalogues P,S,B. The δm -values are rounded into 5 bins and the stars coloured accordingly. The

exact colour-scales can be found within both figures. Additionally, the maps show a kind of "background brightness" or rather a summed brightness of stars per area, which is depicted by a graduated grey-scale. The actual area from which the flux is summed are not the grey rectangular fields, but circles – or rather cones in actual 3D-space – around the centre of each field. These cones have a uniform radius of 4° each, so the summed flux can easily be converted to mag/sr . The grey fields are not of perfectly uniform solid angle but were necessary for the visualisation, as they cover the whole projection without gaps or overlaps. The fields cover 5° of latitude each and a longitude segment that corresponds best to 5° of a great circle while still guaranteeing an integer amount of segments within the 360° circle of latitude. For the summation, all stars within the HIPPARCOS catalogue (ESA, 1997), between 6 mag and 10 mag were used (a total of 112,914 stars). Only faint stars were chosen for several reasons:

- Assure that a star's background is not primarily defined by the star itself.
- Keep the differences in surface brightness small, even for a high spatial resolution (i.e. small grey fields).
- The density of those dimmer stars corresponds well to the perceived brightness of the actual night sky (e.g. the Milky Way (MW) is clearly visible)

Looking at the maps, we can find areas within each catalogue where stars are predominantly estimated too faint ("red areas"). More specifically, we can make the following observations for catalogues P and S:

1. The similarity of catalogues P and S can be found once more in the maps.
2. Nevertheless, Al-Šūfī seems to have reworked many of Ptolemy's most southern stars to fainter values.
3. Both show "red areas" throughout the MW and especially around the centre of the galaxy ($240 < \lambda < 300^\circ / 0 < \beta < 30^\circ$).
4. Like Ptolemy, Al-Šūfī still estimates many stars in the area $300 < \lambda < 360^\circ / -30 < \beta < 30^\circ$ as too bright. However the visibility limit shifts away from those stars and towards the galaxy centre for al-Šūfī's time.
5. Al-Šūfī generally has fewer extreme mistakes, depicting his lower standard deviation in δm .

Points 2 and 4 mostly explain the stronger extinction effect in al-Šūfī's data (see Figure 6) but can only partly (2.) be considered to be really caused by extinction. Apart from that, it seems obvious that both authors show a tendency to estimate stars in bright areas too faint.

⁵Possibly, the worldwide 2020 Corona-lockdown might bring new insights into this question.




FIGURE 7 Plots of star catalogues P (top figure) and S (bottom figure) in ecliptical coordinates (equinox J2000). The stars' size is according to their m_{HIP} and colour according to their δm . Green stars were estimated too bright and red ones too faint (see scale). The pink lines mark the Milky Way ($\pm 10^\circ$ of gal. lat.), the orange lines are the southern visibility limits of the respective time and place. The background depicts the summed brightness of stars in the area as grey-scale. More precisely, the flux of all stars between $6 \text{ mag} < m_{\text{HIP}} < 10 \text{ mag}$ from the HIPPARCOS catalogue is summed and given as surface brightness of the respective area. For details on background-colouring, see text. There is a clear tendency of many too brightly estimated stars in darker areas and vice versa. .




FIGURE 8 Plots of star catalogue B in ecliptical coordinates (equinox J2000). For detailed description, see Figure 7

Brahe's map differs considerably from P and S, showing the following notable features:

1. The brightest parts of the MW are missing, due to Brahe's northern geographic latitude.
2. "Red areas" can be found from the galaxy centre (only partly visible) along the visibility limit (VL) within $270 < \lambda < 330^\circ$.
3. However, other areas along the VL are not particularly red.
4. Another large "red area" can be found at $30 < \lambda < 120^\circ$, roughly along the MW.
5. Again, other parts of the MW do not show any clear tendency.

It becomes obvious, where the extinction feature (see middle column of Figure 6) stems from in Brahe's case (2.) but it is remarkable that the effect is obvious within this area and completely vanishes for other longitudes. A dependency on the background brightness can also be found in some parts of the map. For all three catalogues it is very notable that effects of background brightness and extinction are partly visible but can not be the sole reason of every "red" or "green area".

Concerning the dependency of δm on the background brightness, one might want to explain it by the varying degree of

dark-adaptions for differently bright areas. However, single bright stars would have the strongest effect here and those are excluded from the background brightness, shown in the maps⁶. So the whole phenomenon seems to be of rather psychological nature which makes it harder to quantify theoretically. Nevertheless, several authors (Hearnshaw, 1999; Zinner, 1926) have described and analysed the dependency but mostly restricted themselves to a comparison between stars within and outside of the Milky Way. Going a step further, we used the value of background brightness from the maps (Figs. 7, 8) to plot average δm for 10 bins of surface brightness. The mean values of each bin are shown in the right column of Figure 6 together with a scatter plot of all stars. The decline of δm for darker backgrounds becomes clearly visible. As there is no available mathematical model to describe the expected dependency of δm on the background brightness, a linear regression is the simplest approximation. The regression parameters (see Table 3) can again be used to correct the values of δm for the described effect. Of course, the correction can only be made if the exact same background brightness values are calculated for every star which might be laborious for anyone trying to make use of the correction formula (5). As an alternative, the average δm for stars within and without the Milky Way ($\pm 10^\circ$

⁶looking at the surroundings of the brightest stars, there is no clear trend in any of the maps, either way.

TABLE 4 Average δm of stars within (δm_{MW}) and without (δm_{notMW}) the Milky Way.

Cat.	δm_{MW}	δm_{notMW}
P	0.23 ± 0.06	-0.06 ± 0.03
S	0.24 ± 0.05	-0.07 ± 0.02
B	0.08 ± 0.06	0.01 ± 0.03

of galactic lat.) is given in Table 4 and can be used for the correction formula (5) instead.

4 | CONCLUSIONS

The above analysis investigated different systematical dependencies within the magnitude data of the three historical catalogues. It was shown that stars are often estimated slightly fainter, if ...

- they are red
- they stand close to the southern visibility limit
- they are seen in bright areas of the night sky.

4.1 | Correction formula

The effects were described quantitatively and can therefore be corrected. As the single corrective terms are small in comparison to the overall variance in δm , they can be assumed to be independent from one another and the correction takes the form of a simple additive parameter Δ_i for each of the three models which is subtracted from the initial converted magnitude. This yields a new magnitude m^* for each star which can be considered the best approximation to the modern V-mag scale.

$$m^* = \frac{m_{old} - t'}{b} - \Delta_{B-V} - \Delta_{Ext} - \Delta_{\beta} \quad (5)$$

Where t' and b are given in equation (2). The single Δ_i are then the respective models which were fitted to the data, or for the background brightness it can also be the alternative model based on the position within or without the Milky Way.

$$\begin{aligned} \Delta_{B-V} &= \partial_{B-V} \cdot (B - V) + \theta_{B-V} \\ \Delta_{Ext} &= k \cdot [(\sin(alt) + 0.025 \cdot e^{-11 \cdot \sin(alt)})^{-1} - 1] \\ \Delta_{\beta} &= \partial_{\beta} \cdot \beta + \theta_{\beta} \quad \text{or} \quad = \delta m_{MW/notMW} \end{aligned}$$

The empirical parameters ∂_{B-V} , θ_{B-V} , k , ∂_{β} , θ_{β} can be taken from Figure 6 and $\delta m_{MW/notMW}$ is given in Table 4.

4.2 | Statistical significance

Looking at the three single correction models, it is conspicuous that the variance, explained by the models is much lower than the residual variance induced by the wide scattering of the δm . In fact the R^2 -values (see Table 3) which express the fraction of the variance, explained by the models, remain at a few percent for all nine models. However, when testing the models for significance against a null hypothesis H_0 which predicts zero correlation, H_0 can be dismissed (meaning $F_n > F_{1,n-2,0.99}$) in almost all cases on a 95 % or higher confidence level. This is due to the high number of δm values and suggests that the corrections – however little variance they might explain – are in fact significant. The only model, not showing statistical significance is the extinction correction for the *Almagest* data, which showed a negative (and therefore nonsensical) value of k_{fit} . Nevertheless, when applying the corrections, the low values

TABLE 5 The models in Figure 6 were tested for statistical significance against the hypothesis H_0 of no dependency. The resulting parameters F_n were calculated from R^2 and n (see Table 3) and are given together with the confidence level (CL) on which H_0 can be rejected.

	Color		Extinction		Background	
	F_n	CL	F_n	CL	F_n	CL
P	5.6	95%	0.3	–	23.9	99.9%
S	16.0	99.9%	22.2	99.9%	46.5	99.9%
B	9.8	99%	15.4	99.9%	15.0	99.9%

$$F_{1,n-2,0.95/0.99/0.999} = 3.9 / 6.7 / 10.9 \text{ for } 886 < n < 992$$

of R^2 lead to an almost negligible “improvement” of the δm ’s standard deviations. Figure 9 shows the distribution of the δm before and after the correction formula was applied.

4.3 | Final Remarks on the accuracy of pre-telescopic magnitudes

So after all, what is the accuracy of pre-telescopic magnitude estimations and how can they best be converted to their corresponding V-mag values? From the three major catalogues which contain original magnitudes, Ptolemy’s *Almagest* and Tycho Brahe’s data can be considered largely independent and both show uncertainties of very similar size. Al-Šūfi’s estimations, on the other hand can be seen as an – in most instances – improved version of Ptolemy’s data which show a significantly higher accuracy. The distribution of magnitude errors δm in all three catalogues follows almost Gaussian curves, so the doubled standard deviations 2σ can be considered error bars on a




FIGURE 9 Distribution of the δm before and after the corrections, applied in section 3.2. The histograms of bin size 0.25 mag show approximate normal distributions for all three catalogues. The corrections result in only minor changes of the distributions. Overall, Al-Šūfī’s magnitude errors show the lowest variance.

95% confidence level.

Any magnitude, taken directly from one of the catalogues to be used for studies of transient observations and longterm variabilities or even processes of stellar evolution, should first be converted before comparing them to modern V-mag values. We recommend, adding the brighter- / fainter-qualifiers as 0.5 mag-steps to the original magnitudes and then employing formula (1) to attain the Johnson V-magnitude. The resulting values should be sufficient for most applications and come with error bars of:

$$2\sigma_P = 1.59 \text{ mag}, \quad 2\sigma_S = 1.26 \text{ mag}, \quad 2\sigma_B = 1.52 \text{ mag}$$

In comparison, Hearnshaw (1999) calculates standard deviations between 0.41 and 0.72 mag for most groups of Ptolemaic magnitudes while Zinner (1926) gives values between 0.44 and 0.60 mag as “mean errors”⁷ to adopt the corrected values, even though the error bars are not distinctly reduced by the correction. Those magnitudes can either be attained by equation (5) or taken from the online catalogue, provided by the authors (see APPENDIX B:) Using the corrected values seems especially necessary when analysing only certain groups of stars which might otherwise be systematically biased. This could for example be red giants which all show high values of (B-V) or stars within a certain constellation which might all be in an especially bright or dark part of the night sky.

⁷probably mean *absolute* errors which are always smaller than (or equal to) the standard deviation. Both values are given for the historical magnitudes m_P where our own standard deviation results in $\sigma'_P = 0.79 \text{ mag}/1.33 = 0.59 \text{ mag}$ and therefore in good concordance with the previous studies. In some cases it might be sensible

Other than that, the analysis of dependencies on colour, extinction and background brightness might also be used to investigate otherwise unrelated questions like the extinction coefficient of the pre-industrial atmosphere or even the effective absorption wavelength of the human eye (under naked eye observation conditions).

As we conclude that the error bars of the magnitudes in historical catalogues are ~ 1.3 to ~ 1.6 mag, almost all ($> 93\%$) variabilities of the naked eye stars (as displayed in Fig. 3) are covered by the error bars which makes it virtually impossible to detect longterm variabilities

It should, however, be kept in mind that statistics mean little for a single data point. As we know, for particular cases, the ancient observers must have recognised changes in brightness of less than 1.3 mag. After all, it was possible to naked eye observe the brightness drop of Betelgeuse (α Ori) in winter 2019/ 2020 for many laymen⁸ and there are hypotheses that the variability of Algol (β Per) had been known in ancient Egypt (Jetsu et al., 2013). Cases like these are possible for individual stars which are in a region with appropriate naked eye comparison stars. That is why, our statistical error bars should be considered the general first step but for some handpicked individual stars careful case studies appear worthwhile.

⁸Also, Aboriginal Australians seem to have discovered Betelgeuses’ variability (Schaefer, 2018)

ACKNOWLEDGMENTS

P.P. thanks Prof. Bradley Schaefer, Dr. Rob van Gent and Prof F. Richard Stephenson for answering my requests and helping me get a first grip of the topic during my Master Thesis.

We furthermore thank our referee, Prof. John Hearnshaw, for his recommendations and reassuring comments.

S.H. thanks the Free State of Thuringia for the employment at the Friedrich Schiller University of Jena, Germany.

We thank Ralph Neuhäuser (AIU, Friedrich-Schiller-Universität Jena) for the initiative of investigating historical data and to create room for transdisciplinary projects.

This research has thankfully made use of Vizier catalogue access tool, CDS, Strasbourg, France (DOI:10.26093/cds/vizier, (Ochsenbein et al., 2000)), and the VSX variable star catalogue of the American Association of Variable star Observers (AAVSO) (Watson, Henden, & Price, 2006).

Author contributions

This analysis of pre-telescopic magnitudes was mostly written by PP in the context of his master thesis. SH developed the idea from the thesis towards this contribution and offered extensive support, advice and revision during the completion of the thesis as well as the article. Additionally, SH contributed large parts of the introduction and the final remarks.

How cite this article: Protte, P., Hoffmann, S.M., (2020), Accuracy of magnitudes in pre-telescopic star catalogues, *AN*,

APPENDIX A: OPTICAL INFLUENCES OF HUMAN VISION ON MAGNITUDE ESTIMATES

The pictures in figure A1 show that a bright object (in this case, Venus) can be described as ‘having rays’, ‘horned’, ‘hairy’, or ‘fuzzy’. The photos were taken in central Europe (April 4th to 6th 2020) under normal *clear* weather conditions. The rays and horns of bright objects are not only an effect of the weather but are produced by the interplay of a lens (of the eye as well as of camera optics) and its entrance pupil with an entering wavefront. Passing through a lens with a limited entrance pupil, the wavefronts are described by the Zernike polynomials Z producing the known effects like astigmatism Z_2 , coma Z_3 , the trefoil effect Z_3 (three rays), spherical aberration Z_4 , and higher orders of aberration in the perfectly spherical lens. The effect is caused by the limited size of the pupil (López-Gil et al., 2007) and unevenness of the border increases the effect, as

well as astigmatism of the lens itself. The irises of both, camera and eye, are limitations of the pupil and the polygonal shape of the mechanical iris of a camera lens as well as the muscles at the edge of the eye both increase such effects: The photos of these rays do in fact show roughly the same as what the eye sees.

With atmospheric conditions of the desert or in tropical climate (with sandstorm or humidity) the atmospheric effects become stronger and the bright point source is blurred; the beam of light from the star does not enter the pupil parallelly and the constraints for applying the Zernike polynomials directly are not fulfilled perfectly anymore. The atmospheric influence can even lead to less rays, simply showing the blurred Airy disks around the bright object instead (rightmost picture, with cirrus clouds on April 10th).

APPENDIX B: ONLINE-ONLY MATERIAL

We prepared the catalogues according to our suggestions in the above work. The data files will be uploaded in CDS as soon as the paper is published. It will be available at CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/AN>

REFERENCES

- Baily, F. 1843, *memras*, 13, 1.
 Ballesteros, F. J. 2012, Feb, *EPL (Europhysics Letters)*, 97, 34008.
 Clauss, W., & Clauss, C. 2018, *Humanbiologie kompakt* (2. ed.), Springer Berlin Heidelberg.
 CVRL. 1995, Data for luminous efficiency functions., Website of the Colour and Vision Research Laboratory. Retrieved from <http://www.cvrl.org/> (Last accessed: June 10th, 2019)
 ESA. 1997, Feb, *VizieR Online Data Catalog*, I/239. Retrieved from http://vizier.u-strasbg.fr/viz-bin/VizieR-3?-source=I/239/hip_main
 Fujiwara, T., & Yamaoka, H. 2005, June, *Journal of Astronomical History and Heritage*, 8(1), 39-47.
 Fujiwara, T., Yamaoka, H., & Miyoshi, S. J. 2004, March, *A&A*, 416, 641-646.
 Granzer, T. 2014, February, Data for Johnson-Cousins filter curves., Website of the Leibnitz-Institute fÄijr Astrophysik Potsdam. Retrieved from <https://www.aip.de/en/research/facilities/stella/instruments/data> (Last accessed: June 10th, 2019)
 Hafez, I. 2010. Abd al-Rahman al-Sufi and his book of the fixed stars: a journey of re-discovery. (Doctoral dissertation, James Cook University). Retrieved from <http://eprints.jcu.edu.au/28854/>
 Hearnshaw, J. B. 1996, *The Measurement of Starlight, Two Centuries of Astronomical Photometry*.
 Hearnshaw, J. B. 1999, *New A Rev.*, 43, 403-410.
 Hertzog, K. P. 1984, August, *MNRAS*, 209, 533-541.
 Ho, P. Y. 1962, *Vistas in Astronomy*, 5.
 Hoffmann, S. 2017, *Hipparchus Himmelsglobus: Ein Bindeglied in*




FIGURE A1 Naked eye appearance of a bright object in contrast to fainter stars (here: Venus, -4.6 mag, next to the central part of the cluster of the Pleiades, stars of 2.9 to 4.3 mag). Photos are shown as taken with Canon 600D and 300 mm tele objective, without any astronomical preparation, without any filter or any processing in order to display the real impression for the human eye.

- der babylonisch-griechischen Astrometrie? Springer Fachmedien Wiesbaden.
- Hoffmann, S. M., Vogt, N., & Protte, P. 2019, *AN*.
- Jetsu, L., Porceddu, S., Lyytinen, J., Kajatkari, P., Lehtinen, J., Markkanen, T., & Toivari-Viitala, J. 2013, August, *ApJ*, 773(1), 1. doi:
- Knobel, E. B. 1917, Ulugh Beg's catalogue of stars: revised from all Persian manuscripts existing in Great Britain, with a vocabulary of Persian and Arabic words.
- López-Gil, N., Rucker, F. J., Stark, L. R., Badar, M., Borgovan, T., Burke, S., & Kruger, P. B. 2007, *Vision Research*, 47(6), 755 - 765. Retrieved from <http://www.sciencedirect.com/science/article/pii/S0042698906003488> doi:
- Mayer, P. 1984, April, *The Observatory*, 104, 77-80.
- Neuhäuser, R., Arlt, R., & Richter, S. 2018, May, *Astronomische Nachrichten*, 339(4), 219-267.
- Neuhäuser, R., Gießler, F., & Hambaryan, V. V. 2019, September, *MNRAS*, 2261.
- Neuhäuser, R., & Neuhäuser, D. L. 2016, June, *Astronomische Nachrichten*, 337(6), 581.
- Ochsenbein, F., Bauer, P., & Marcout, J. 2000, April, *A&AS*, 143, 23-32. doi:
- Pickering, K. A. 2002, Sep, *DIO*, 12, 51-58.
- Pogson, N. 1856, nov, *MNRAS*, 17, 12-15. doi:
- Ångström, J. A. 1869, jan, *Annalen der Physik*, 213(5), 161-163. doi:
- Schaefer, B. E. 1993, *Vistas in Astronomy*, 36, 311-361.
- Schaefer, B. E. 2013, February, *Journal for the History of Astronomy*, 44, 47-74.
- Schaefer, B. E. 2018, April, *Journal of Astronomical History and Heritage*, 21(1), 7-12.
- Shara, M. M., Ilkiewicz, K., Mikolajewska, J. et al. 2017, *Nature*, 548, 558-560.
- Siedentopf, H. 1959, jan, *Naturwissenschaften*, 46(9), 309-314. doi:
- Toomer, G. J. 1984, Ptolemy's Almagest. Princeton: Princeton University Press.
- Verbunt, F., & van Gent, R. H. 2010a, June, *A&A*, 516, A28.
- Verbunt, F., & van Gent, R. H. 2010b, March, *VizieR Online Data Catalog*, 351. Retrieved from <http://vizier.u-strasbg.fr/viz-bin/VizieR-3?-source=J/A%2bA/516/A28/Tycho-Brahe>
- Verbunt, F., & van Gent, R. H. 2012a, August, *A&A*, 544, A31.
- Verbunt, F., & van Gent, R. H. 2012b, June, *VizieR Online Data Catalog*, 354. Retrieved from <http://vizier.cfa.harvard.edu/viz-bin/VizieR?-source=J/A+A/544/A31>
- Vogt, N., Hoffmann, S. M., & Tappert, C. 2019, *Astron. Nachr.*, 752, 340.
- Watson, C. L., Henden, A. A., & Price, A. 2006, May, *Society for Astronomical Sciences Annual Symposium*, 25, 47.
- Xu, Z., Pankenier, D. W., & Jiang, Y. 2000, East Asian Archaeoastronomy. Gordon and Breach Science Publishers.
- Zinner, E. 1926, *Veroeffentlichungen der Remeis-Sternwarte zu Bamberg*, 2.

How cite this article: Protte, P., Hoffmann, S.M., (2020), Accuracy of magnitudes in pre-telescopic star catalogues, *AN*, .