Relative Reshetikhin-Turaev invariants, hyperbolic cone metrics and discrete Fourier transforms I

Ka Ho Wong and Tian Yang

Abstract

We propose the Volume Conjecture for the relative Reshetikhin-Turaev invariants of a closed oriented 3-manifold with a colored framed link inside it whose asymptotic behavior is related to the volume and the Chern-Simons invariant of the hyperbolic cone metric on the manifold with singular locus the link and cone angles determined by the coloring. We prove the conjecture in the case that the ambient 3-manifold is obtained by doing an integral surgery along some components of a fundamental shadow link and the complement of the link in the ambient manifold is homeomorphic to the fundamental shadow link complement, for sufficiently small cone angles. Together with Costantino and Thurston's result that all compact oriented 3-manifolds with toroidal or empty boundary can be obtained by doing an integral surgery along some components of a suitable fundamental shadow link, this provides a possible approach of solving Chen-Yang's Volume Conjecture for the Reshetikhin-Turaev invariants of closed oriented hyperbolic 3-manifolds. We also introduce a family of topological operations (the change of pair operations) that connect all pairs of a closed oriented 3-manifold and a framed link inside it that have homeomorphic complements, which correspond to doing the partial discrete Fourier transforms to the corresponding relative Reshetikhin-Turaev invariants. As an application, we find a Poisson Summation Formula for the discrete Fourier transforms.

1 Introduction

We propose the Volume Conjecture for the relative Reshetikhin-Turaev invariants of a closed oriented 3-manifold with a colored framed link inside it whose asymptotic behavior is related to the volume and the Chern-Simons invariant of the hyperbolic cone metric on the manifold with singular locus the link and cone angles determined by the coloring. See Conjecture 1.1, and Sections 2.1 and 2.2 for a review of the relative Reshetikhin-Turaev invariants and the hyperbolic cone manifolds.

We prove the conjecture in the case that the ambient 3-manifold is obtained by doing an integral surgery along some components of a fundamental shadow link and the complement of the link in the ambient manifold is homeomorphic to the fundamental shadow link complement, for sufficiently small cone angles. See Theorem 1.2, and Section 2.4 for a review of the fundamental shadow links. A result of Costantino and Thurston [10] shows that all compact oriented 3-manifolds with toroidal or empty boundary can be obtained by doing an integral surgery along some components of a suitable fundamental shadow link. On the other hand, it is expected that hyperbolic cone metrics interpolate the complete cusped hyperbolic metric on the 3-manifold with toroidal boundary and the smooth hyperbolic metric on the Dehn-filled 3-manifold, corresponding to the colors running from $\frac{r-1}{2}$ to 0 or r-2. Therefore, if one can push the cone angles in Theorem 1.2 from sufficiently small all the way up to 2π , then one proves the Volume Conjecture for the Reshetikhin-Turaev invariants of closed oriented hyperbolic 3-manifolds proposed by Chen and the second author [6].

This thus suggests a possible approach of solving Chen-Yang's Volume Conjecture. In [45], we prove Conjecture 1.1 for all pairs (M, K) such that $M \setminus K$ is homeomorphic to the figure-8 knot complement in S^3 with all possible cone angles, showing the plausibility of this new approach.

We also introduce a family of the change of pair operations (see Section 1.2) that connect all pairs of a closed oriented 3-manifold and a framed link inside it that have homeomorphic complements, which correspond to doing the partial discrete Fourier transforms (see Section 1.3) to the corresponding relative Reshetikhin-Turaev invariants. As an application, we find a Poisson Summation formula for the discrete Fourier transforms (see Formula (1.1)).

1.1 Volume Conjecture for the relative Reshetikhin-Turaev invariants

Conjecture 1.1. Let M be a closed oriented 3-manifold and let L be a framed hyperbolic link in M with n components. For an odd integer $r \ge 3$, let $\mathbf{m} = (m_1, \ldots, m_n)$ and let $\operatorname{RT}_r(M, L, \mathbf{m})$ be the r-th relative Reshetikhin-Turaev invariant of M with L colored by \mathbf{m} and evaluated at the root of unity $q = e^{\frac{2\pi\sqrt{-1}}{r}}$. For a sequence $\mathbf{m}^{(r)} = (m_1^{(r)}, \ldots, m_n^{(r)})$, let

$$\theta_k = \left| 2\pi - \lim_{r \to \infty} \frac{4\pi m_k^{(r)}}{r} \right|$$

and let $\theta = (\theta_1, \dots, \theta_n)$. If $M_{L_{\theta}}$ is a hyperbolic cone manifold consisting of M and a hyperbolic cone metric on M with singular locus L and cone angles θ , then

$$\lim_{r \to \infty} \frac{4\pi}{r} \log \operatorname{RT}_r(M, L, \mathbf{m}^{(r)}) = \operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(M_{L_{\theta}}) \qquad \text{mod } \sqrt{-1} \pi^2 \mathbb{Z},$$

where r varies over all positive odd integers.

We note that if $M = S^3$, then Conjecture 1.1 covers Kashaev's Volume Conjecture for the colored Jones polynomials of hyperbolic links [20, 25, 26, 12] and its generalization [27], at the root of unity $q = e^{\frac{2\pi\sqrt{-1}}{r}}$. See also [16] and [8, Section 4.2] for a discussion of the values at the root $q = e^{\frac{\pi\sqrt{-1}}{r}}$. If the framed link $L = \emptyset$ or the coloring $\mathbf{m} = \mathbf{0}$ or $\mathbf{r} - \mathbf{2}$, then Conjecture 1.1 covers Chen-Yang's Volume Conjecture for the Reshetikhin-Turaev invariants of closed oriented hyperbolic 3-manifolds.

The main result of this paper is the following Theorem 1.2 (see also Theorem 3.1 for a more precise statement), where the change of pair operation is described in the next section.

Theorem 1.2. Conjecture 1.1 is true for all pairs (M, L) obtained by doing a change of pair operation from the pair (M_c, L_{FSL}) with sufficiently small cone angles, where $M_c = \#^{c+1}(S^2 \times S^1)$ and L_{FSL} is a fundamental shadow link in M_c .

As a consequence of Theorem 1.2, we prove the Generalized Volume Conjecture [27, 16] for the colored Jones polynomials of the universal families of links respectively considered by Purcell [31], van der Veen [41] and Kumar [22]. See Proposition 6.2 and Theorems 6.3 and 6.5 for more details.

1.2 The change of pair operations

Let M be a closed oriented 3-manifold and let L be a framed link in M. In this section, we introduce a topological operation that changes the pair (M, L) with out changing the complement $M \setminus L$, and show that these operations connect all such pairs that have homeomorphic complements.

Suppose $L = L_1 \cup \cdots \cup L_n$. For each component $L_i : [0, 1] \times S^1 \to M$ of L, we call $L_i(\{0\} \times S^1) \subset M$ the core curve of L_i and $L_i(\{1\} \times S^1) \subset M$ the parallel copy. Let (I, J) be a partition of $\{1, \ldots, n\}$, $L_I = \bigcup_{i \in I} L_i$ and $L_J = \bigcup_{j \in J} L_j$. For each $i \in I$, let L_i^* be the framed knot in $M \setminus L$ whose core curve is isotopic to the meridian of the tubular neighborhood of L_i , and let $L_I^* = \bigcup_{i \in I} L_i^*$. Let M^* be the closed 3-manifold obtained from M by doing the surgery along L_I and let L^* be the framed link obtained from



Figure 1: M is obtained from S^3 by the surgery along L', and $L = L_1 \cup L_2$. $(I, J) = (\{1\}, \{2\})$. M^* is obtained from M by doing the surgery along L_1 , hence is obtained from S^3 by doing the surgery along $L' \cup L_1$; and $L^* = L_1^* \cup L_2$.

L by replacing L_I by L_I^* , i.e., $M^* = M_{L_I}$ and $L^* = L_I^* \cup L_J$. The change of pair operation $T_{(L_I;L_I^*)}$ is defined by sending (M, L) to (M^*, L^*) . See Figure 1.

By the way it is chosen, the core curve of each L_i^* intersects the parallel copy of L_i once, hence in M^* the core curve of each L_i^* is isotopic to the core of the filled in solid torus of the surgery along L_i . As a consequence, we have

Proposition 1.3. If (M^*, L^*) is obtained form (M, L) be doing a change of pair operation, then $M^* \ L^*$ is homeomorphic to $M \ L$.

Conversely, if (M^*, L^*) is a pair such that $M^* \ L^*$ is homeomorphic to $M \ L$, then M^* is obtained from M by doing a rational Dehn-surgery along some components L_I of L. By e.g. [35, p. 273], M^* can be obtained from M by doing an integral surgery along a framed link L' obtained from L_I by iteratively linking in framed unknots, corresponding to doing a sequence of the change of pair operations. As a consequence, we have

Proposition 1.4. Every two pairs (M, L) and (M^*, L^*) such that $M \setminus L$ is homeomorphic to $M^* \setminus L^*$ are related by a sequence of the change of pairs operations.

1.3 Relationship with discrete Fourier transform

In the computation of the relative Reshetikhin-Turaev invariants, the change of pair operation corresponds to replacing the coloring \mathbf{m}_I on L_I by the Kirby colorings Ω_r and cabling L_I^* by the Chebyshev polynomials corresponding to the new coloring \mathbf{n}_I , sending $\operatorname{RT}_r(M, L, \mathbf{m})$ to $\operatorname{RT}_r(M^*, L^*, (\mathbf{n}_I, \mathbf{m}_J))$, where \mathbf{m}_J is the coloring on L_J . See Section 2.1 for a review of the relative Reshetikhin-Turaev invariants and Figure 2 for an example.

These operations pictorially represent the discrete Fourier transforms. See [1] for the original definition and [2] for an exposition in the language of skein theory. To be more precise, let $I_r = \{1, 2, \ldots, r - 2\}$ and $\mu_r = \frac{\sqrt{2} \sin \frac{\pi}{r}}{\sqrt{r}}$ in the SU(2) theory and at $q = e^{\frac{\pi\sqrt{-1}}{r}}$, and let $I_r = \{0, 2, \ldots, r - 3\}$ and $\mu_r = \frac{2 \sin \frac{2\pi}{r}}{\sqrt{r}}$ in the SO(3) theory and at $q = e^{\frac{2\pi\sqrt{-1}}{r}}$. Let

 $f: \mathbf{I}_r^n \to \mathbb{C}$



Figure 2: For the computation of $\operatorname{RT}_r(M, L, (m_1, m_2))$ on the left, L' is cabled by the Kirby coloring Ω_r and L_1 and L_2 are respectively cabled by the Chebyshev polynomials e_{m_1} and e_{m_2} . For $\operatorname{RT}_r(M^*, L^*, (n_1, m_2))$ on the right, L' and L_1 are cabled by Ω_r , L_1^* is cabled by e_{n_1} and L_2 is cabled by e_{m_2} . The framings are omitted in the Figure.

be a complex valued function on I_r^n for some positive integer n. Let (I, J) be a partition of $\{1, \ldots, n\}$ and let $\mathbf{n}_I = (n_i)_{i \in I}$ be a |I|-tuple of elements of I_r . Then the \mathbf{n}_I -th partial discrete Fourier coefficient of f is the function

$$\widehat{f}(\mathbf{n}_I): \mathbf{I}_r^J \to \mathbb{C}$$

defined for all \mathbf{m}_J in \mathbf{I}_r^J by

$$\widehat{f}(\mathbf{n}_I)(\mathbf{m}_J) = \mu_r^{|I|} \sum_{\mathbf{m}_I} \prod_{i \in I} \mathrm{H}(m_i, n_i) f(\mathbf{m}_I, \mathbf{m}_J),$$

where the sum is over all |I|-tuples $\mathbf{m}_I = (m_i)_{i \in I}$ of elements of \mathbf{I}_r , and

$$\mathbf{H}(m,n) = (-1)^{m+n} \frac{q^{(m+1)(n+1)} - q^{-(m+1)(n+1)}}{q - q^{-1}}.$$

Since the coefficients $H(m_i, n_i)$ above are exactly the coefficients of the following skein-theoretical computation

$$\left\langle \underbrace{\stackrel{}{\longleftarrow} \Omega_{\mathbf{r}}}_{\mathbf{I}} \mathbf{n} \right\rangle = \mu_{r} \sum_{m \in \mathbf{I}_{r}} \mathbf{H}(m, n) \left\langle \mathbf{I} \mathbf{m} \right\rangle,$$

the relative Reshetikhin-Turaev invariants $\operatorname{RT}_r(M^*, L^*, (\mathbf{n}_I, \mathbf{m}_J))$ of the pair (M^*, L^*) obtained from (M, L) by a change of pair operation $T_{(L_I; L_I^*)}$ is up to scalar the value at \mathbf{m}_J of the \mathbf{n}_I -th partial discrete Fourier coefficient of the function $\operatorname{RT}_r(M, L, _-) : \operatorname{I}_r^n \to \mathbb{C}$. The scalar is a power of q depending on the framings of L_I^* .

It is proved in [12, 4] that Turaev-Viro invariant of the complement $M \setminus L$ can be computed by the relative Reshetikhin-Turaev invariants of the pair (M, L).

Proposition 1.5 ([12, 4]).

$$\operatorname{TV}_r(M \smallsetminus L) = c \sum_{\mathbf{m}} |\operatorname{RT}_r(M, L, \mathbf{m})|^2,$$

where the sum is over all multi-elements \mathbf{m} of \mathbf{I}_r , and the constant c equals 1 in the SU(2) theory and equals $2^{\operatorname{rank}H_2(M \setminus L,\mathbb{Z}_2)}$ in the SO(3) theory.

By Proposition 1.3, if (M^*, L^*) and (M, L) differ by a change of pair operation, then $M \setminus L$ and $M^* \setminus L^*$ are homeomorphic to each other. As a consequence, we have

Proposition 1.6.

$$\sum_{\mathbf{m}} \left| \mathrm{RT}_r(M, L, \mathbf{m}) \right|^2 = \sum_{\mathbf{n}} \left| \mathrm{RT}_r(M^*, L^*, \mathbf{n}) \right|^2,$$

where the sums are over all multi-elements \mathbf{m} and \mathbf{n} of I_r .

Propositions 1.3, 1.4, 1.5 and 1.6 together provide infinitely many different ways to compute the Turaev-Viro invariants of $M \setminus L$, all of which are up to scalar related by a sequence of partial discrete Fourier transforms. It is hopeful that among these different expressions, some are suitable for the purpose of solving the Volume Conjecture for the Turaev-Viro invariants.

Finally, as a special case, let $f = \operatorname{RT}_r(M, L, _) : \operatorname{I}_r^n \to \mathbb{C}$ for a pair (M, L), and suppose that $I = \{1, \ldots, n\}$ and all the framings of L_I^* are zero and (M^*, L^*) is obtained from (M, L) by $T_{(L_I, L_I^*)}$. Then $\operatorname{RT}_r(M^*, L^*, \mathbf{n}) = \widehat{f}(\mathbf{n})$ is the **n**-th (full) discrete Fourier coefficient of f. As a consequence of Proposition 1.6, we have

$$\sum_{\mathbf{m}} |f(\mathbf{m})|^2 = \sum_{\mathbf{n}} |\widehat{f}(\mathbf{n})|^2, \qquad (1.1)$$

where m and n are over all multi-elements of I_r . This could be considered as a Poisson Summation Formula for the discrete Fourier transforms. (See also [2] for an asymptotic version of the Poisson Summation Formula in the setting of Yokota invariants for colored planar graphs.)

1.4 Outline of the proof

We follow the guideline of Ohtsuki's method. In Proposition 3.5, we compute the relative Reshetikhin-Turaev invariants of (M, L) writing them as a sum of values of a holomorphic function f_r at integer points. The function f_r comes from Faddeev's quantum dilogarithm function. Using Poisson Summation Formula, we in Proposition 3.7 write the invariants as a sum of the Fourier coefficients of f_r computed in Propositions 3.6. In Proposition 5.2 we show that the critical value of the functions in the leading Fourier coefficients has real part the volume and imaginary part the Chern-Simons invariant. The key observation there is a relationship between the asymptotics of quantum 6j-symbols and the Neumann-Zagier potential function (Proposition 4.1), which is of interest in its own right. Then we estimate the leading Fourier coefficients in Sections 5.3 using the Saddle Point Method (Proposition 5.1). Finally, we estimate the non-leading Fourier coefficients and the error term respectively in Sections 5.4 and 5.5 showing that they are neglectable, and prove Theorem 3.1 in Section 5.6.

Acknowledgments. The authors would like to thank Giulio Belletti, Effie Kalfagianni, Sanjay Kumar, Zhengwei Liu, Feng Luo, Hongbin Sun and Roland van der Veen for helpful discussions. The first author would like to thank the organizers of the "Nearly Carbon Neutral Geometric Topology Conference 2020" for the invitation and for providing a friendly environment for inspiring discussions. The second author is partially supported by NSF Grant DMS-1812008.

2 Preliminaries

2.1 Relative Reshetikhin-Turaev invariants

In this article we will follow the skein theoretical approach of the Reshetikhin-Turaev invariants [5, 23] and focus on the SO(3)-theory and the values at the root of unity $q = e^{\frac{2\pi\sqrt{-1}}{r}}$ for odd integers $r \ge 3$.

A framed link in an oriented 3-manifold M is a smooth embedding L of a disjoint union of finitely many thickened circles $S^1 \times [0, \epsilon]$, for some $\epsilon > 0$, into M. The Kauffman bracket skein module $K_r(M)$ of M is the \mathbb{C} -module generated by the isotopic classes of framed links in M modulo the follow two relations:

(1) Kauffman Bracket Skein Relation:
$$A = e^{\frac{\pi\sqrt{-1}}{r}} + e^{-\frac{\pi\sqrt{-1}}{r}}$$

(2) Framing Relation: $L \cup = \left(-e^{\frac{2\pi\sqrt{-1}}{r}} - e^{-\frac{2\pi\sqrt{-1}}{r}}\right) L.$

There is a canonical isomorphism

$$\rangle : \mathrm{K}_r(\mathrm{S}^3) \to \mathbb{C}$$

defined by sending the empty link to 1. The image $\langle L \rangle$ of the framed link L is called the Kauffman bracket of L.

Let $K_r(A \times [0, 1])$ be the Kauffman bracket skein module of the product of an annulus A with a closed interval. For any link diagram D in \mathbb{R}^2 with k ordered components and $b_1, \ldots, b_k \in K_r(A \times [0, 1])$, let

$$\langle b_1, \ldots, b_k \rangle_D$$

be the complex number obtained by cabling b_1, \ldots, b_k along the components of D considered as a element of $K_r(S^3)$ then taking the Kauffman bracket $\langle \rangle$.

On $K_r(A \times [0, 1])$ there is a commutative multiplication induced by the juxtaposition of annuli, making it a \mathbb{C} -algebra; and as a \mathbb{C} -algebra $K_r(A \times [0, 1]) \cong \mathbb{C}[z]$, where z is the core curve of A. For an integer $n \ge 0$, let $e_n(z)$ be the n-th Chebyshev polynomial defined recursively by $e_0(z) = 1$, $e_1(z) = z$ and $e_n(z) = ze_{n-1}(z) - e_{n-2}(z)$. Let $I_r = \{0, 2, \dots, r-3\}$ be the set of even integers in between 0 and r - 2. Then the Kirby coloring $\Omega_r \in K_r(A \times [0, 1])$ is defined by

$$\Omega_r = \mu_r \sum_{n \in \mathbf{I}_r} [n+1]e_n,$$

where

$$\mu_r = \frac{2\sin\frac{2\pi}{r}}{\sqrt{r}}$$

and [n] is the quantum integer defined by

$$[n] = \frac{e^{\frac{2n\pi\sqrt{-1}}{r}} - e^{-\frac{2n\pi\sqrt{-1}}{r}}}{e^{\frac{2\pi\sqrt{-1}}{r}} - e^{-\frac{2\pi\sqrt{-1}}{r}}}.$$

Let M be a closed oriented 3-manifold and let L be a framed link in M with n components. Suppose M is obtained from S^3 by doing a surgery along a framed link L', D(L') is a standard diagram of L' (ie, the blackboard framing of D(L') coincides with the framing of L'). Then L adds extra components to D(L') forming a linking diagram $D(L \cup L')$ with D(L) and D(L') linking in possibly a complicated way. Let U_+ be the diagram of the unknot with framing $1, \sigma(L')$ be the signature of the linking matrix of L' and $\mathbf{m} = (m_1, \ldots, m_n)$ be a multi-elements of I_r . Then the r-th relative Reshetikhin-Turaev invariant of M with L colored by \mathbf{m} is defined as

$$\operatorname{RT}_{r}(M,L,\mathbf{m}) = \mu_{r} \langle e_{m_{1}}, \dots, e_{m_{n}}, \Omega_{r}, \dots, \Omega_{r} \rangle_{D(L \cup L')} \langle \Omega_{r} \rangle_{U_{+}}^{-\sigma(L')}.$$
(2.1)

(+1)

Note that if $L = \emptyset$ or $m_1 = \cdots = m_n = 0$, then $\operatorname{RT}_r(M, L, \mathbf{m}) = \operatorname{RT}_r(M)$, the *r*-th Reshetikhin-Turaev invariant of M; and if $M = S^3$, then $\operatorname{RT}_r(M, L, \mathbf{m}) = \mu_r \operatorname{J}_{\mathbf{m}, L}(q^2)$, the value of the m-th unnormalized colored Jones polynomial of L at $t = q^2$.

2.2 Hyperbolic cone manifolds

According to [7], a 3-dimensional hyperbolic cone-manifold is a 3-manifold M, which can be triangulated so that the link of each simplex is piecewise linear homeomorphic to a standard sphere and M is equipped with a complete path metric such that the restriction of the metric to each simplex is isometric to a hyperbolic geodesic simplex. The singular locus L of a cone-manifold M consists of the points with no neighborhood isometric to a ball in a Riemannian manifold. It follows that

- (1) L is a link in M such that each component is a closed geodesic.
- (2) At each point of L there is a *cone angle* θ which is the sum of dihedral angles of 3-simplices containing the point.
- (3) The restriction of the metric on $M \setminus L$ is a smooth hyperbolic metric, but is incomplete if $L \neq \emptyset$.

Hodgson-Kerckhoff [18] proved that hyperbolic cone metrics on M with singular locus L are locally parametrized by the cone angles provided all the cone angles are less than or equal to 2π , and Kojima [21] proved that hyperbolic cone manifolds (M, L) are globally rigid provided all the cone angles are less than or equal to π . It is expected to be globally rigid if all the cone angles are less than or equal to 2π .

Given a 3-manifold N with boundary a union of tori T_1, \ldots, T_n , a choice of generators (u_i, v_i) for each $\pi_1(T_i)$ and pairs of relatively prime integers (p_i, q_i) , one can do the $(\frac{p_1}{q_1}, \ldots, \frac{p_k}{q_k})$ -Dehn filling on N by attaching a solid torus to each T_i so that $p_i u_i + q_i v_i$ bounds a disk. If $H(u_i)$ and $H(v_i)$ are respectively the logarithmic holonomy for u_i and v_i , then a solution to

$$p_i \mathbf{H}(u_i) + q_i \mathbf{H}(v_i) = \sqrt{-1}\theta_i \tag{2.2}$$

near the complete structure gives a cone-manifold structure on the resulting manifold M with the cone angle θ_i along the core curve L_i of the solid torus attached to T_i ; it is a smooth structure if $\theta_1 = \cdots = \theta_n = 2\pi$.

In this setting, the Chern-Simons invariant for a hyperbolic cone manifold (M, L) can be defined by using the Neumann-Zagier potential function [28]. To do this, we need a framing on each component, namely, a choice of a curve γ_i on T_i that is isotopic to the core curve L_i of the solid torus attached to T_i . We choose the orientation of γ_i so that $(p_i u_i + q_i v_i) \cdot \gamma_i = 1$. Then we consider the following function

$$\frac{\Phi(\mathrm{H}(u_1),\ldots,\mathrm{H}(u_n))}{\sqrt{-1}} - \sum_{i=1}^n \frac{\mathrm{H}(u_i)\mathrm{H}(v_i)}{4\sqrt{-1}} + \sum_{i=1}^n \frac{\theta_i\mathrm{H}(\gamma_i)}{4},$$

where Φ is the Neumann-Zagier potential function (see [28]) defined on the deformation space of hyperbolic structures on $M \setminus L$ parametrized by the holonomy of the meridians $\{H(u_i)\}$, characterized by

$$\begin{cases}
\frac{\partial \Phi(\mathrm{H}(u_1),\ldots,\mathrm{H}(u_n))}{\partial \mathrm{H}(u_i)} = \frac{\mathrm{H}(v_i)}{2}, \\
\Phi(0,\ldots,0) = \sqrt{-1} \left(\mathrm{Vol}(M \smallsetminus L) + \sqrt{-1} \mathrm{CS}(M \smallsetminus L) \right) \mod \pi^2 \mathbb{Z},
\end{cases}$$
(2.3)

where $M \setminus L$ is with the complete hyperbolic metric. Another important feature of Φ is that it is even in each of its variables $H(u_i)$.

Following the argument in [28, Sections 4 & 5], one can prove that if the cone angles of components of L are $\theta_1, \ldots, \theta_n$, then

$$\operatorname{Vol}(M_{L_{\theta}}) = \operatorname{Re}\left(\frac{\Phi(\operatorname{H}(u_{1}), \dots, \operatorname{H}(u_{n}))}{\sqrt{-1}} - \sum_{i=1}^{n} \frac{\operatorname{H}(u_{i})\operatorname{H}(v_{i})}{4\sqrt{-1}} + \sum_{i=1}^{n} \frac{\theta_{i}\operatorname{H}(\gamma_{i})}{4}\right).$$
(2.4)

Indeed, in this case, one can replace the 2π in Equations (33) (34) and (35) of [28] by θ_i , and as a consequence can replace the $\frac{\pi}{2}$ in Equations (45), (46) and (48) by $\frac{\theta_i}{4}$, proving the result.

In [46], Yoshida proved that when $\theta_1 = \cdots = \theta_n = 2\pi$,

$$\operatorname{Vol}(M) + \sqrt{-1} \operatorname{CS}(M) = \frac{\Phi(\operatorname{H}(u_1), \dots, \operatorname{H}(u_n))}{\sqrt{-1}} - \sum_{i=1}^n \frac{\operatorname{H}(u_i) \operatorname{H}(v_i)}{4\sqrt{-1}} + \sum_{i=1}^n \frac{\theta_i \operatorname{H}(\gamma_i)}{4} \mod \sqrt{-1}\pi^2 \mathbb{Z}$$

Therefore, we can make the following

Definition 2.1. The Chern-Simons invariant of a hyperbolic cone manifold $M_{L_{\theta}}$ with a choice of the framing $(\gamma_1, \ldots, \gamma_n)$ is defined as

$$\operatorname{CS}(M_{L_{\theta}}) = \operatorname{Im}\left(\frac{\Phi(\operatorname{H}(u_1), \dots, \operatorname{H}(u_n))}{\sqrt{-1}} - \sum_{i=1}^n \frac{\operatorname{H}(u_i)\operatorname{H}(v_i)}{4\sqrt{-1}} + \sum_{i=1}^n \frac{\theta_i\operatorname{H}(\gamma_i)}{4}\right) \mod \pi^2 \mathbb{Z}.$$

Then together with (2.4), we have

$$\operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(M_{L_{\theta}}) = \frac{\Phi(\operatorname{H}(u_{1}), \dots, \operatorname{H}(u_{n}))}{\sqrt{-1}} - \sum_{i=1}^{n} \frac{\operatorname{H}(u_{i}) \operatorname{H}(v_{i})}{4\sqrt{-1}} + \sum_{i=1}^{n} \frac{\theta_{i} \operatorname{H}(\gamma_{i})}{4} \mod \sqrt{-1} \pi^{2} \mathbb{Z}$$
(2.5)

Remark 2.2. It is an interesting question to find a direct geometric definition of the Chern-Simons invariants for hyperbolic cone manifolds.

2.3 Quantum 6*j*-symbols

A triple (m_1, m_2, m_3) of even integers in $\{0, 2, \dots, r-3\}$ is *r*-admissible if

(1)
$$m_i + m_j - m_k \ge 0$$
 for $\{i, j, k\} = \{1, 2, 3\},\$

(2) $m_1 + m_2 + m_3 \leq 2(r-2).$

For an *r*-admissible triple (m_1, m_2, m_3) , define

$$\Delta(m_1, m_2, m_3) = \sqrt{\frac{\left[\frac{m_1 + m_2 - m_3}{2}\right]! \left[\frac{m_2 + m_3 - m_1}{2}\right]! \left[\frac{m_3 + m_1 - m_2}{2}\right]!}{\left[\frac{m_1 + m_2 + m_3}{2} + 1\right]!}}$$

with the convention that $\sqrt{x} = \sqrt{|x|}\sqrt{-1}$ when the real number x is negative.

A 6-tuple $(m_1, ..., m_6)$ is *r*-admissible if the triples (m_1, m_2, m_3) , (m_1, m_5, m_6) , (m_2, m_4, m_6) and (m_3, m_4, m_5) are *r*-admissible

Definition 2.3. The quantum 6j-symbol of an r-admissible 6-tuple (m_1, \ldots, m_6) is

$$\begin{vmatrix} m_1 & m_2 & m_3 \\ m_4 & m_5 & m_6 \end{vmatrix} = \sqrt{-1}^{-\sum_{i=1}^6 m_i} \Delta(m_1, m_2, m_3) \Delta(m_1, m_5, m_6) \Delta(m_2, m_4, m_6) \Delta(m_3, m_4, m_5) \\ \sum_{k=\max\{T_1, T_2, T_3, T_4\}}^{\min\{Q_1, Q_2, Q_3\}} \frac{(-1)^k [k+1]!}{[k-T_1]! [k-T_2]! [k-T_3]! [k-T_4]! [Q_1-k]! [Q_2-k]! [Q_3-k]!},$$

where $T_1 = \frac{m_1 + m_2 + m_3}{2}$, $T_2 = \frac{m_1 + m_5 + m_6}{2}$, $T_3 = \frac{m_2 + m_4 + m_6}{2}$ and $T_4 = \frac{m_3 + m_4 + m_5}{2}$, $Q_1 = \frac{m_1 + m_2 + m_4 + m_5}{2}$, $Q_2 = \frac{m_1 + m_3 + m_4 + m_6}{2}$ and $Q_3 = \frac{m_2 + m_3 + m_5 + m_6}{2}$.

Here we recall a classical result of Costantino [8] which was originally stated at the root of unity $q = e^{\frac{\pi\sqrt{-1}}{r}}$. At the root of unity $q = e^{\frac{2\pi\sqrt{-1}}{r}}$, see [4, Appendix] for a detailed proof.

Theorem 2.4 ([8]). Let $\{(m_1^{(r)}, \ldots, m_6^{(r)})\}$ be a sequence of r-admissible 6-tuples, and let

$$\theta_i = \left| \pi - \lim_{r \to \infty} \frac{2\pi m_i^{(r)}}{r} \right|.$$

If $\theta_1, \ldots, \theta_6$ are the dihedral angles of a truncated hyperideal tetrahedron Δ , then as r varies over all the odd integers

$$\lim_{r \to \infty} \frac{2\pi}{r} \log \left| \begin{array}{cc} m_1^{(r)} & m_2^{(r)} & m_3^{(r)} \\ m_4^{(r)} & m_5^{(r)} & m_6^{(r)} \end{array} \right|_{q=e^{\frac{2\pi\sqrt{-1}}{r}}} = Vol(\Delta).$$

Closely related, a triple $(\alpha_1, \alpha_2, \alpha_3) \in [0, 2\pi]^3$ is *admissible* if

- (1) $\alpha_i + \alpha_j \alpha_k \ge 0$ for $\{i, j, k\} = \{1, 2, 3\},\$
- (2) $\alpha_i + \alpha_j + \alpha_k \leq 4\pi$.

A 6-tuple $(\alpha_1, \ldots, \alpha_6) \in [0, 2\pi]^6$ is *admissible* if the triples $\{1, 2, 3\}, \{1, 5, 6\}, \{2, 4, 6\}$ and $\{3, 4, 5\}$ are admissible.

2.4 Fundamental shadow links

In this section we recall the construction and basic properties of the fundamental shadow links. The building blocks for the fundamental shadow links are truncated tetrahedra as in the left of Figure 3. If we take c building blocks $\Delta_1, \ldots, \Delta_c$ and glue them together along the triangles of truncation, we obtain a (possibly non-orientable) handlebody of genus c + 1 with a link in its boundary consisting of the edges of the building blocks, such as in the right of Figure 3. By taking the orientable double (the orientable double covering with the boundary quotient out by the deck involution) of this handlebody, we obtain a link L_{FSL} inside $M_c = \#^{c+1}(S^2 \times S^1)$. We call a link obtained this way a *fundamental shadow link*, and its complement in M_c a *fundamental shadow link complement*.



Figure 3: The handlebody on the right is obtained from the truncated tetrahedron on the left by identifying the triangles on the top and the bottom by a horizontal reflection and the triangles on the left and the right by a vertical reflection.

The fundamental importance of the family of the fundamental shadow links is the following.

Theorem 2.5 ([10]). Any compact oriented 3-manifold with toroidal or empty boundary can be obtained from a suitable fundamental shadow link complement by doing an integral Dehn-filling to some of the boundary components.

A hyperbolic cone metric on M_c with singular locus L_{FSL} and with sufficiently small cone angles $\theta_1, \ldots, \theta_n$ can be constructed as follows. For each $s \in \{1, \ldots, c\}$, let e_{s_1}, \ldots, e_{s_6} be the edges of the building block Δ_s , and θ_{s_j} be the cone angle of the component of L containing e_{s_j} . If θ_i 's are sufficiently small, then $\{\frac{\theta_{s_1}}{2}, \ldots, \frac{\theta_{s_6}}{2}\}$ for the set of dihedral angles of a truncated hyperideal tetrahedron, by abuse of notation still denoted by Δ_s . Then the hyperbolic cone manifold M_c with singular locus L_{FSL} and cone angles $\theta_1, \ldots, \theta_n$ is obtained by glueing Δ_s 's together along isometries of the triangles of truncation, and taking the double. In this metric, the logarithmic holonomy of the meridian u_i of the tubular neighborhood $N(L_i)$ of L_i satisfies

$$\mathbf{H}(u_i) = \sqrt{-1}\theta_i. \tag{2.6}$$

Let l_i be the length of the closed geodesic L_i . Since the manifold comes from taking double, we can choose a longitude v_i on the boundary of $N(L_i)$ with $u_i \cdot v_i = 1$ and with the logarithmic holomony satisfying

$$\mathbf{H}(v_i) = -l_i. \tag{2.7}$$

In this way, a framing on L_i gives an integer p_i in the way that the parallel copy of L_i on $N(L_i)$ is isotopic to the curve representing $p_i u_i + v_i$.

Proposition 2.6 ([8, 9]). If $L_{FSL} = L_1 \cup \cdots \cup L_n \subset M_c$ is a framed fundamental shadow link with framing p_i on L_i , and $\mathbf{m} = (m_1, \ldots, m_n)$ is a coloring of its components with even integers in $\{0, 2, \ldots, r-3\}$, then

$$\operatorname{RT}_{\mathbf{r}}(M_{c}, L_{FSL}, \mathbf{m}) = \left(\frac{2\sin\frac{2\pi}{r}}{\sqrt{r}}\right)^{-c} \prod_{i=1}^{n} q^{\frac{p_{i}m_{i}(m_{i}+2)}{2}} \prod_{s=1}^{c} \begin{vmatrix} m_{s_{1}} & m_{s_{2}} & m_{s_{3}} \\ m_{s_{4}} & m_{s_{5}} & m_{s_{6}} \end{vmatrix},$$

where m_{s_1}, \ldots, m_{s_6} are the colors of the edges of the building block Δ_s inherited from the color **m** on L_{FSL} .

Together with Theorem 2.4 and the construction of the hyperbolic cone structure, we see that Conjecture 1.1 is true for (M_c, L_{FSL}) with sufficiently small cone angles. This was first proved by Costantino in [8] at the root of unity $q = e^{\frac{\pi\sqrt{-1}}{r}}$.

2.5 Dilogarithm and quantum dilogarithm functions

Let $\log : \mathbb{C} \setminus (-\infty, 0] \to \mathbb{C}$ be the standard logarithm function defined by

$$\log z = \log |z| + \sqrt{-1} \arg z$$

with $-\pi < \arg z < \pi$.

The dilogarithm function $Li_2 : \mathbb{C} \setminus (1, \infty) \to \mathbb{C}$ is defined by

$$\operatorname{Li}_2(z) = -\int_0^z \frac{\log(1-u)}{u} du$$

where the integral is along any path in $\mathbb{C} \setminus (1, \infty)$ connecting 0 and z, which is holomorphic in $\mathbb{C} \setminus [1, \infty)$ and continuous in $\mathbb{C} \setminus (1, \infty)$.

The dilogarithm function satisfies the follow properties (see eg. Zagier [47]).

(1)

$$\operatorname{Li}_{2}\left(\frac{1}{z}\right) = -\operatorname{Li}_{2}(z) - \frac{\pi^{2}}{6} - \frac{1}{2}\left(\log(-z)\right)^{2}.$$
(2.8)

(2) In the unit disk $\{z \in \mathbb{C} \mid |z| < 1\}$,

$$\text{Li}_{2}(z) = \sum_{n=1}^{\infty} \frac{z^{n}}{n^{2}}.$$
(2.9)

(3) On the unit circle $\{z = e^{2\sqrt{-1}\theta} \mid 0 \leq \theta \leq \pi\},\$

$$\text{Li}_{2}(e^{2\sqrt{-1}\theta}) = \frac{\pi^{2}}{6} + \theta(\theta - \pi) + 2\sqrt{-1}\Lambda(\theta).$$
 (2.10)

Here $\Lambda : \mathbb{R} \to \mathbb{R}$ is the Lobachevsky function defined by

$$\Lambda(\theta) = -\int_0^\theta \log|2\sin t| dt,$$

which is an odd function of period π . See eg. Thurston's notes [37, Chapter 7].

The following variant of Faddeev's quantum dilogarithm functions [13, 14] will play a key role in the proof of the main result. Let $r \ge 3$ be an odd integer. Then the following contour integral

$$\varphi_r(z) = \frac{4\pi\sqrt{-1}}{r} \int_{\Omega} \frac{e^{(2z-\pi)x}}{4x\sinh(\pi x)\sinh(\frac{2\pi x}{r})} dx$$
(2.11)

defines a holomorphic function on the domain

$$\left\{ z \in \mathbb{C} \mid -\frac{\pi}{r} < \operatorname{Re} z < \pi + \frac{\pi}{r} \right\},$$

where the contour is

$$\Omega = (-\infty, -\epsilon] \cup \{z \in \mathbb{C} \mid |z| = \epsilon, \operatorname{Im} z > 0\} \cup [\epsilon, \infty),$$

for some $\epsilon \in (0, 1)$. Note that the integrand has poles at $n\sqrt{-1}$, $n \in \mathbb{Z}$, and the choice of Ω is to avoid the pole at 0.

The function $\varphi_r(z)$ satisfies the following fundamental properties, whose proof can be found in [44, Section 2.3].

Lemma 2.7. (1) For $z \in \mathbb{C}$ with $0 < \operatorname{Re} z < \pi$,

$$1 - e^{2\sqrt{-1}z} = e^{\frac{r}{4\pi\sqrt{-1}}\left(\varphi_r\left(z - \frac{\pi}{r}\right) - \varphi_r\left(z + \frac{\pi}{r}\right)\right)}.$$
(2.12)

(2) For $z \in \mathbb{C}$ with $-\frac{\pi}{r} < \operatorname{Re} z < \frac{\pi}{r}$,

$$1 + e^{r\sqrt{-1}z} = e^{\frac{r}{4\pi\sqrt{-1}}\left(\varphi_r(z) - \varphi_r\left(z + \pi\right)\right)}.$$
(2.13)

Using (2.12) and (2.13), for $z \in \mathbb{C}$ with $\pi + \frac{2(n-1)\pi}{r} < \text{Re}z < \pi + \frac{2n\pi}{r}$, we can define $\varphi_r(z)$ inductively by the relation

$$\prod_{k=1}^{n} \left(1 - e^{2\sqrt{-1}\left(z - \frac{(2k-1)\pi}{r}\right)} \right) = e^{\frac{r}{4\pi\sqrt{-1}}\left(\varphi_r\left(z - \frac{2n\pi}{r}\right) - \varphi_r(z)\right)},\tag{2.14}$$

extending $\varphi_r(z)$ to a meromorphic function on \mathbb{C} . The poles of $\varphi_r(z)$ have the form $(a+1)\pi + \frac{b\pi}{r}$ or $-a\pi - \frac{b\pi}{r}$ for all nonnegative integer a and positive odd integer b.

Let $q = e^{\frac{2\pi\sqrt{-1}}{r}}$, and let

$$(q)_n = \prod_{k=1}^n (1 - q^{2k}).$$

Lemma 2.8. (1) For $0 \le n \le r - 2$,

$$(q)_n = e^{\frac{r}{4\pi\sqrt{-1}} \left(\varphi_r\left(\frac{\pi}{r}\right) - \varphi_r\left(\frac{2\pi n}{r} + \frac{\pi}{r}\right)\right)}.$$
(2.15)

(2) For $\frac{r-1}{2} \leq n \leq r-2$,

$$(q)_n = 2e^{\frac{r}{4\pi\sqrt{-1}}\left(\varphi_r\left(\frac{\pi}{r}\right) - \varphi_r\left(\frac{2\pi n}{r} + \frac{\pi}{r} - \pi\right)\right)}.$$
(2.16)

We consider (2.16) because there are poles in $(\pi, 2\pi)$, and to avoid the poles we move the variables to $(0, \pi)$ by subtracting π .

Since

$$\{n\}! = (-1)^n q^{-\frac{n(n+1)}{2}}(q)_n,$$

as a consequence of Lemma 2.8, we have

(

Lemma 2.9. (1) For $0 \le n \le r - 2$,

$$\{n\}! = e^{\frac{r}{4\pi\sqrt{-1}} \left(-2\pi \left(\frac{2\pi n}{r}\right) + \left(\frac{2\pi}{r}\right)^2 (n^2 + n) + \varphi_r \left(\frac{\pi}{r}\right) - \varphi_r \left(\frac{2\pi n}{r} + \frac{\pi}{r}\right) \right)}.$$
(2.17)

(2) For $\frac{r-1}{2} \leq n \leq r-2$,

$$\{n\}! = 2e^{\frac{r}{4\pi\sqrt{-1}}\left(-2\pi\left(\frac{2\pi n}{r}\right) + \left(\frac{2\pi}{r}\right)^2(n^2+n) + \varphi_r\left(\frac{\pi}{r}\right) - \varphi_r\left(\frac{2\pi n}{r} + \frac{\pi}{r} - \pi\right)\right)}.$$
(2.18)

The function $\varphi_r(z)$ and the dilogarithm function are closely related as follows.

Lemma 2.10. (1) For every z with $0 < \text{Re}z < \pi$,

$$\varphi_r(z) = \operatorname{Li}_2(e^{2\sqrt{-1}z}) + \frac{2\pi^2 e^{2\sqrt{-1}z}}{3(1 - e^{2\sqrt{-1}z})} \frac{1}{r^2} + O\left(\frac{1}{r^4}\right).$$
(2.19)

(2) For every z with $0 < \text{Re}z < \pi$,

$$\varphi_r'(z) = -2\sqrt{-1}\log(1 - e^{2\sqrt{-1}z}) + O\left(\frac{1}{r^2}\right).$$
(2.20)

(3) [30, Formula (8)(9)]

$$\varphi_r\left(\frac{\pi}{r}\right) = \operatorname{Li}_2(1) + \frac{2\pi\sqrt{-1}}{r}\log\left(\frac{r}{2}\right) - \frac{\pi^2}{r} + O\left(\frac{1}{r^2}\right).$$

3 Computation of the relative Reshetikhin-Turaev invariants

The goal of this section is to compute the relative Reshetikhin-Turaev invariants of (M, L). In Proposition 3.5, we write the invariants as a sum of values of a fixed holomorphic function at the integer points. The holomorphic function comes from Faddeev's quantum dilogarithm function. Using the Poisson Summation Formula, we in Proposition 3.7 write the invariants as a sum of the Fourier coefficients of the holomorphic function, which is computed in Propositions 3.6.

Let $L_{\text{FSL}} = L_1 \cup \cdots \cup L_n$ be a fundamental shadow link in $M_c = \#^{c+1}(S^2 \times S^1)$, and let $L' \subset S^3$ be the disjoint union of c + 1 unknots with the 0-framings by doing surgery along which we get M_c . Let (I, J) be a partition of $\{1, \ldots, n\}$, and let (M, L) be the pair obtained from (M_c, L_{FSL}) by doing a change of pair operation $T_{(L_I;L_I^*)}$ as introduced in Section 1.2, i.e., $M = (M_c)_{L_I}$ and $L = L_I^* \cup L_J$, where $L_I^* = \bigcup_{i \in I} L_i^*$ and L_i^* is the framed unknot in $M_c \setminus L_{\text{FSL}}$ with the core curve isotopic to the meridian of the tubular neighborhood of L_i . Let \mathbf{n}_I be a coloring of L_I^* and let \mathbf{m}_J be a coloring of L_J . Then Theorem 1.2 can be rephrased as follows. **Theorem 3.1.** For $i \in I$, let

$$\theta_i = \Big| 2\pi - \lim_{r \to \infty} \frac{4\pi n_i}{r} \Big|;$$

and for $j \in J$, let

$$\theta_j = \left| 2\pi - \lim_{r \to \infty} \frac{4\pi m_j}{r} \right|.$$

Let $\theta = (\theta_1, \dots, \theta_n)$ and let $M_{L_{\theta}}$ be the hyperbolic cone manifold consisting of M and a hyperbolic cone metric on M with singular locus L and cone angles θ . Then there exists an $\epsilon > 0$ such that if all the cone angles θ_k are less than ϵ , then as r varies over all the odd integers

$$\lim_{r \to \infty} \frac{4\pi}{r} \log \operatorname{RT}_r(M, L, (\mathbf{n}_I, \mathbf{m}_J)) = \operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(M_{L_{\theta}}) \qquad \text{mod } \sqrt{-1} \pi^2 \mathbb{Z}$$

The goal of the rest of this paper is to prove Theorem 3.1.

Suppose L_i^* has the framing q_i and L_i has the framing p_i for each $i \in I$, and L_j has the framing p_j for each $j \in J$. Then the *r*-th relative Reshetikin-Turaev invariant of M with L colored by $(\mathbf{n}_I, \mathbf{m}_J)$ can be computed as

$$\operatorname{RT}_{r}(M,L,(\mathbf{n}_{I},\mathbf{m}_{J})) = \left(\frac{2\sin\frac{2\pi}{r}}{\sqrt{r}}\right)^{|I|-c} e^{-\sigma(L'\cup L_{I})\left(-\frac{3}{r}-\frac{r+1}{4}\right)\sqrt{-1}\pi} \prod_{i\in I} q^{\frac{q_{i}n_{i}(n_{i}+2)}{2}} \prod_{j\in J} q^{\frac{p_{j}m_{j}(m_{j}+2)}{2}} \sum_{\mathbf{m}_{I}} \prod_{i\in I} q^{\frac{p_{i}m_{i}(m_{i}+2)}{2}} [(m_{i}+1)(n_{i}+1)] \prod_{s=1}^{c} \begin{vmatrix} m_{s_{1}} & m_{s_{2}} & m_{s_{3}} \\ m_{s_{4}} & m_{s_{5}} & m_{s_{6}} \end{vmatrix} ,$$

$$(3.1)$$

where the sum is over all multi-even integers $\mathbf{m}_I = (m_i)_{i \in I}$ in $\{0, 2, \dots, r-3\}$, and m_{s_1}, \dots, m_{s_6} are the colors of the edges of the building block Δ_s inherited from the colors on L_{FSL} .

In the rest of this section, we aim to write $\operatorname{RT}_r(M, L, (\mathbf{n}_I; \mathbf{m}_J))$ into a sum of integrals using the Poisson Summation Formula. This requires writing the invariant into the sum of the values of a fixed holomorphic function. To this end, we look at the a single quantum 6*j*-symbol first.

Definition 3.2. An *r*-admissible 6-tuple (m_1, \ldots, m_6) is of the hyperideal type if for $\{i, j, k\} = \{1, 2, 3\}, \{1, 5, 6\}, \{2, 4, 6\}$ and $\{3, 4, 5\}, \{2, 4, 6\}$ and $\{3, 4, 5\}, \{2, 4, 6\}$ and $\{3, 4, 5\}, \{3, 4, 5\},$

- (1) $0 \leq m_i + m_j m_k < r 2$,
- (2) $r-2 \leq m_i + m_j + m_k \leq 2(r-2),$

Definition 3.3. A 6-tuple $(\alpha_1, \ldots, \alpha_6) \in [0, 2\pi]^6$ is of the hyperideal type if for $\{i, j, k\} = \{1, 2, 3\}, \{1, 5, 6\}, \{2, 4, 6\}$ and $\{3, 4, 5\}, \{1, 5, 6\}, \{2, 4, 6\}$ and $\{3, 4, 5\}, \{1, 5, 6\}, \{2, 4, 6\}$ and $\{3, 4, 5\}, \{1, 5, 6\}, \{2, 4, 6\}$ and $\{3, 4, 5\}, \{1, 5, 6\}, \{2, 4, 6\}, \{2, 4, 6\}, \{2, 4, 6\}, \{2, 4, 6\}, \{2, 4, 6\}, \{3, 4, 5\}, \{3, 4, 5\}, \{3, 4, 5\}, \{3, 4, 5\}, \{3, 4, 5\}, \{3, 4, 5\}, \{3, 4, 5\}, \{3, 4, 5\}, \{4$

- (1) $0 \leq \alpha_i + \alpha_j \alpha_k \leq 2\pi$,
- (2) $2\pi \leq \alpha_i + \alpha_j + \alpha_k \leq 4\pi$.

We notice that the six numbers $|\alpha_1 - \pi|, \ldots, |\alpha_6 - \pi|$ are the dihedral angles of an ideal or a hyperideal tetrahedron if and only if $(\alpha_1, \ldots, \alpha_6)$ is of the hyperideal type.

As a consequence of Lemma 2.9 we have

Proposition 3.4. The quantum 6*j*-symbol at the root of unity $q = e^{\frac{2\pi\sqrt{-1}}{r}}$ can be computed as

$$\begin{vmatrix} m_1 & m_2 & m_3 \\ m_4 & m_5 & m_6 \end{vmatrix} = \frac{1}{2\{1\}} \sum_{k=\max\{T_1, T_2, T_3, T_4\}}^{\min\{Q_1, Q_2, Q_3, r-2\}} e^{\frac{r}{4\pi\sqrt{-1}}U_r\left(\frac{2\pi m_1}{r}, \dots, \frac{2\pi m_6}{r}, \frac{2\pi k}{r}\right)},$$

where U_r is defined as follows. If (m_1, \ldots, m_6) is of hyperideal type, then

$$U_{r}(\alpha_{1},...,\alpha_{6},\xi) = \pi^{2} - \left(\frac{2\pi}{r}\right)^{2} + \frac{1}{2}\sum_{i=1}^{4}\sum_{j=1}^{3}(\eta_{j} - \tau_{i})^{2} - \frac{1}{2}\sum_{i=1}^{4}\left(\tau_{i} + \frac{2\pi}{r} - \pi\right)^{2} + \left(\xi + \frac{2\pi}{r} - \pi\right)^{2} - \sum_{i=1}^{4}(\xi - \tau_{i})^{2} - \sum_{j=1}^{3}(\eta_{j} - \xi)^{2} - 2\varphi_{r}\left(\frac{\pi}{r}\right) - \frac{1}{2}\sum_{i=1}^{4}\sum_{j=1}^{3}\varphi_{r}\left(\eta_{j} - \tau_{i} + \frac{\pi}{r}\right) + \frac{1}{2}\sum_{i=1}^{4}\varphi_{r}\left(\tau_{i} - \pi + \frac{3\pi}{r}\right) - \varphi_{r}\left(\xi - \pi + \frac{3\pi}{r}\right) + \sum_{i=1}^{4}\varphi_{r}\left(\xi - \tau_{i} + \frac{\pi}{r}\right) + \sum_{j=1}^{3}\varphi_{r}\left(\eta_{j} - \xi + \frac{\pi}{r}\right),$$
(3.2)

where $\tau_1 = \frac{\alpha_1 + \alpha_2 + \alpha_3}{2}$, $\tau_2 = \frac{\alpha_1 + \alpha_5 + \alpha_6}{2}$, $\tau_3 = \frac{\alpha_2 + \alpha_4 + \alpha_6}{2}$ and $\tau_4 = \frac{\alpha_3 + \alpha_4 + \alpha_5}{2}$, $\eta_1 = \frac{\alpha_1 + \alpha_2 + \alpha_4 + \alpha_5}{2}$, $\eta_2 = \frac{\alpha_1 + \alpha_3 + \alpha_4 + \alpha_6}{2}$ and $\eta_3 = \frac{\alpha_2 + \alpha_3 + \alpha_5 + \alpha_6}{2}$. If (m_1, \ldots, m_6) is not of the hyperideal type, then U_r will be changed according to Lemma 2.9.

As a consequence, we have

Proposition 3.5.

$$\operatorname{RT}_{r}(M, L, (\mathbf{n}_{I}, \mathbf{m}_{J})) = \kappa_{r} \sum_{\mathbf{m}_{I}, \mathbf{k}} \Big(\sum_{\epsilon_{I}} g_{r}^{\epsilon_{I}}(\mathbf{m}_{I}, \mathbf{k}) \Big),$$

where

$$\kappa_r = \frac{2^{|I|-2c}}{\{1\}^c} \left(\frac{\sin\frac{2\pi}{r}}{\sqrt{r}}\right)^{|I|-c} e^{\left(-\sigma(L'\cup L_I)(-\frac{3}{r}-\frac{r+1}{4})-\frac{r}{4}(\sum_{i\in I}q_i+\sum_{i=1}^np_i+2|I|)\right)\sqrt{-1}\pi},$$

 $\epsilon_I = (\epsilon_i)_{i \in I} \in \{1, -1\}^I$ runs over all multi-signs, $\mathbf{m}_I = (m_i)_{i \in I}$ runs over all multi-even integers in $\{0, 2, \ldots, r-3\}$ so that for each $s \in \{1, \ldots, c\}$ the triples $(m_{s_1}, m_{s_2}, m_{s_3}), (m_{s_1}, m_{s_5}, m_{s_6}), (m_{s_2}, m_{s_4}, m_{s_6})$ and $(m_{s_3}, m_{s_4}, m_{s_5})$ are r-adminssible, and $\mathbf{k} = (k_1, \ldots, k_c)$ runs over all multiintegers with each k_s lying in between $\max\{T_{s_i}\}$ and $\min\{Q_{s_i}, r-2\}$, with

$$g_r^{\epsilon_I}(\mathbf{m}_I, \mathbf{k}) = e^{\frac{2\pi\sqrt{-1}}{r} \left(\sum_{i \in I} q_i n_i + \sum_{i=1}^n p_i m_i + \sum_{i \in I} \epsilon_i (m_i + n_i + 1)\right) + \frac{r}{4\pi\sqrt{-1}} \mathcal{W}_r^{\epsilon_I}(\frac{2\pi\mathbf{m}_I}{r}, \frac{2\pi\mathbf{k}}{r})}$$

where $\frac{2\pi\mathbf{m}_I}{r} = \left(\frac{2\pi m_i}{r}\right)_{i\in I}, \frac{2\pi\mathbf{k}}{r} = \left(\frac{2\pi k_1}{r}, \dots, \frac{2\pi k_c}{r}\right), and$

$$\mathcal{W}_{r}^{\epsilon_{I}}(\alpha_{I},\xi) = -\sum_{i\in I} q_{i}(\beta_{i}-\pi)^{2} - \sum_{j\in J} p_{j}(\alpha_{j}-\pi)^{2}$$
$$-\sum_{i\in I} p_{i}(\alpha_{i}-\pi)^{2} - \sum_{i\in I} 2\epsilon_{i}(\alpha_{i}-\pi)(\beta_{i}-\pi) + \sum_{s=1}^{c} U_{r}(\alpha_{s_{1}},\dots,\alpha_{s_{6}},\xi_{s})$$

with $\beta_i = \frac{2\pi n_i}{r}$ for $i \in I$ and $\alpha_j = \frac{2\pi m_j}{r}$ for $j \in J$.

Proof. We have

$$q^{\frac{n(n+2)}{2}} = \left(e^{\frac{\pi\sqrt{-1}}{4}}\right)^{-r} q^{\frac{1}{2}\left(n-\frac{r}{2}\right)^2 + n},$$
$$q^{(m+1)(n+1)} = \left(e^{\frac{\pi\sqrt{-1}}{2}}\right)^{-r} q^{\left(m-\frac{r}{2}\right)\left(n-\frac{r}{2}\right) + m + n + 1},$$

and

$$q^{-(m+1)(n+1)} = -\left(e^{\frac{\pi\sqrt{-1}}{2}}\right)^{-r} q^{-\left(m-\frac{r}{2}\right)\left(n-\frac{r}{2}\right)-m-n-1}.$$

Then the result follows from (3.1) and Proposition 3.4.

We notice that the summation in Proposition 3.5 is finite, and to use the Poisson Summation Formula, we need an infinite sum over integral points. To this end, we consider the following regions and a bump function over them.

Let
$$\beta_i = \frac{2\pi n_i}{r}$$
 for $i \in I$, $\alpha_i = \frac{2\pi m_i}{r}$ for $i = 1, \dots, n$, $\xi_s = \frac{2\pi k_s}{r}$ for $s = 1, \dots, c$, $\tau_{s_i} = \frac{2\pi T_{s_i}}{r}$ for $i = 1, \dots, 4$, and $\eta_{s_j} = \frac{2\pi Q_{s_j}}{r}$ for $j = 1, 2, 3$. For a fixed $(\alpha_j)_{j \in J}$, let

$$\mathbf{D}_{\mathbf{A}} = \left\{ (\alpha_I, \xi) \in \mathbb{R}^{|I|+c} \, \middle| \, (\alpha_{s_1}, \dots, \alpha_{s_6}) \text{ is admissible, } \max\{\tau_{s_i}\} \leqslant \xi_s \leqslant \min\{\eta_{s_j}, 2\pi\}, s = 1, \dots, c \right\}.$$

and let

$$D_{H} = \Big\{ (\alpha_{I}, \xi) \in D_{A} \ \Big| \ (\alpha_{s_{1}}, \dots, \alpha_{s_{6}}) \text{ is of the hyperideal type, } s = 1, \dots, c \Big\}.$$

For a sufficiently small $\delta > 0$, let

$$\mathbf{D}_{\mathbf{H}}^{\delta} = \Big\{ (\alpha_I, \xi) \in \mathbf{D}_{\mathbf{H}} \ \Big| \ d((\alpha_I, \xi), \partial \mathbf{D}_{\mathbf{H}}) < \delta \Big\},\$$

where d is the Euclidean distance on \mathbb{R}^n . Let $\psi : \mathbb{R}^{|I|+c} \to \mathbb{R}$ be the C^{∞} -smooth bump function supported on (D_H, D_H^{δ}) , ie,

$$\begin{cases} \psi(\alpha_I, \xi) = 1, & (\alpha_I, \xi) \in \overline{\mathcal{D}_{\mathcal{H}}^{\delta}} \\ 0 < \psi(\alpha_I, \xi) < 1, & (\alpha_I, \xi) \in \mathcal{D}_{\mathcal{H}} \smallsetminus \overline{\mathcal{D}_{\mathcal{H}}^{\delta}} \\ \psi(\alpha_I, \xi) = 0, & (\alpha_I, \xi) \notin \mathcal{D}_{\mathcal{H}}, \end{cases}$$

and let

$$f_r^{\epsilon_I}(\mathbf{m}_I, \mathbf{k}) = \psi\Big(\frac{2\pi\mathbf{m}_I}{r}, \frac{2\pi\mathbf{k}}{r}\Big)g_r^{\epsilon_I}(\mathbf{m}_I, \mathbf{k}).$$

In Proposition 3.5, \mathbf{m}_I runs over multi-even integers. On the other hand, to use the Poisson Summation Formula, we need a sum over all multi-integers. For this purpose, we for each $i \in I$ let $m_i = 2m'_i$ and let $\mathbf{m}'_I = (m'_i)_{i \in I}$. Then by Proposition 3.5,

$$\operatorname{RT}_r(M, L, (\mathbf{n}_I, \mathbf{m}_J)) = \kappa_r \sum_{(\mathbf{m}_I', \mathbf{k}) \in \mathbb{Z}^{|I|+c}} \left(\sum_{\epsilon_I \in \{1, -1\}^I} f_r^{\epsilon_I}(2\mathbf{m}_I', \mathbf{k}) \right) + \text{error term.}$$

Let

$$f_r = \sum_{\epsilon_I \in \{1, -1\}^I} f_r^{\epsilon_I}$$

Then

$$\operatorname{RT}_r(M, L, (\mathbf{n}_I, \mathbf{m}_J)) = \kappa_r \sum_{(\mathbf{m}'_I, \mathbf{k}) \in \mathbb{Z}^{|I|+c}} f_r(2\mathbf{m}'_I, \mathbf{k}) + \text{error term.}$$

Since f_r is C^{∞} -smooth and equals zero out of D_H , it is in the Schwartz space on $\mathbb{R}^{|I|+c}$. Then by the Poisson Summation Formula (see e.g. [36, Theorem 3.1]),

$$\sum_{(\mathbf{m}_I',\mathbf{k})\in\mathbb{Z}^{|I|+c}}f_r\big(2\mathbf{m}_I',\mathbf{k}\big)=\sum_{(\mathbf{a}_I,\mathbf{b})\in\mathbb{Z}^{|I|+c}}\widehat{f}_r(\mathbf{a}_I,\mathbf{b}),$$

where $\mathbf{a}_I = (a_i)_{i \in I} \in \mathbb{Z}^I$, $\mathbf{b} = (b_1, \dots, b_c) \in \mathbb{Z}^c$ and $\widehat{f}_r(\mathbf{a}_I, \mathbf{b})$ is the $(\mathbf{a}_I, \mathbf{b})$ -th Fourier coefficient of f_r defined by

$$\widehat{f}_r(\mathbf{a}_I, \mathbf{b}) = \int_{\mathbb{R}^{|I|+c}} f_r(\mathbf{m}_I', \mathbf{k}) e^{\sum_{i \in I} 2\pi \sqrt{-1} a_i m_i' + \sum_{s=1}^c 2\pi \sqrt{-1} b_s k_s} d\mathbf{m}_I' d\mathbf{k}$$

where $d\mathbf{m}'_I d\mathbf{k} = \prod_{i \in I} dm'_i \prod_{s=1}^c dk_s$.

By a change of variable, and by changing $2m'_i$ back to m_i , the Fourier coefficients can be computed as

Proposition 3.6.

$$\widehat{f}_r(\mathbf{a}_I, \mathbf{b}) = \sum_{\epsilon_I \in \{1, -1\}^I} \widehat{f_r^{\epsilon_I}}(\mathbf{a}_I, \mathbf{b})$$

with

$$\widehat{f_r^{\epsilon_I}}(\mathbf{a}_I, \mathbf{b}) = \frac{r^{|I|+c}}{2^{2|I|+c} \cdot \pi^{|I|+c}} \int_{\mathrm{D}_{\mathrm{H}}} \psi(\alpha_I, \xi) e^{\left(\sum_{i \in I} q_i \beta_i + \sum_{i=1}^n p_i \alpha_i + \sum_{i \in I} \epsilon_i (\alpha_i + \beta_i + \frac{2\pi}{r})\right)\sqrt{-1}} \cdot e^{\frac{r}{4\pi\sqrt{-1}} \left(\mathcal{W}_r^{\epsilon_I}(\alpha_I, \xi) - \sum_{i \in I} 2\pi a_i \alpha_i - \sum_{s=1}^c 4\pi b_s \xi_s\right)} d\alpha_I d\xi,$$

where $d\alpha_I d\xi = \prod_{i \in I} d\alpha_i \prod_{s=1}^c d\xi_s$, and

$$\mathcal{W}_{r}^{\epsilon_{I}}(\alpha_{I},\xi) = -\sum_{i\in I} q_{i}(\beta_{i}-\pi)^{2} - \sum_{j\in J} p_{j}(\alpha_{j}-\pi)^{2}$$
$$-\sum_{i\in I} p_{i}(\alpha_{i}-\pi)^{2} - \sum_{i\in I} 2\epsilon_{i}(\alpha_{i}-\pi)(\beta_{i}-\pi) + \sum_{s=1}^{c} U_{r}(\alpha_{s_{1}},\dots,\alpha_{s_{6}},\xi_{s}).$$

Proposition 3.7.

$$\operatorname{RT}_r(M, L, (\mathbf{n}_I, \mathbf{m}_J)) = \kappa_r \sum_{(\mathbf{a}_I, \mathbf{b}) \in \mathbb{Z}^{|I|+c}} \widehat{f}_r(\mathbf{a}_I, \mathbf{b}) + \operatorname{error term.}$$

We will estimate the leading Fourier coefficients, the non-leading Fourier coefficients and the error term respectively in Sections 5.3, 5.4 and 5.5, and prove Theorem 3.1 in Section 5.6.

4 Relationship with the Neumann-Zagier potential function

The goal of this section is to show the relationship between $W_r^{\epsilon_I}$ and the Neumann-Zagier potential function [28] of the fundamental shadow link complement $M_c \setminus L_{FSL}$. To this end, we need to first look at the function U coming from a single 6*j*-symbol, and to recall its relationship with the volume of a truncated hyperideal tetrahedron.

By Lemma 2.10, $W_r^{\epsilon_I}$ is approximated by the holomorphic function W^{ϵ_I} defined below, which will play an important role later. (The approximation will be specified in the proof of Proposition 5.5.) The function W^{ϵ_I} is defined by

$$\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi) = -\sum_{i \in I} q_{i}(\beta_{i} - \pi)^{2} - \sum_{j \in J} p_{j}(\alpha_{j} - \pi)^{2}$$
$$-\sum_{i \in I} p_{i}(\alpha_{i} - \pi)^{2} - \sum_{i \in I} 2\epsilon_{i}(\alpha_{i} - \pi)(\beta_{i} - \pi) + \sum_{s=1}^{c} U(\alpha_{s_{1}}, \dots, \alpha_{s_{6}}, \xi_{s})$$

with \boldsymbol{U} defined by

$$U(\alpha_{1},...,\alpha_{6},\xi) = \pi^{2} + \frac{1}{2} \sum_{i=1}^{4} \sum_{j=1}^{3} (\eta_{j} - \tau_{i})^{2} - \frac{1}{2} \sum_{i=1}^{4} (\tau_{i} - \pi)^{2} + (\xi - \pi)^{2} - \sum_{i=1}^{4} (\xi - \tau_{i})^{2} - \sum_{j=1}^{3} (\eta_{j} - \xi)^{2} - 2Li_{2}(1) - \frac{1}{2} \sum_{i=1}^{4} \sum_{j=1}^{3} Li_{2} (e^{2i(\eta_{j} - \tau_{i})}) + \frac{1}{2} \sum_{i=1}^{4} Li_{2} (e^{2i(\tau_{i} - \pi)}) - Li_{2} (e^{2i(\xi - \pi)}) + \sum_{i=1}^{4} Li_{2} (e^{2i(\xi - \tau_{i})}) + \sum_{j=1}^{3} Li_{2} (e^{2i(\eta_{j} - \xi)}).$$

$$(4.1)$$

We note that U defines a holomorphic function on the region

 $B_{\mathrm{H},\mathbb{C}} = \left\{ (\alpha,\xi) \in \mathbb{C}^7 \mid \mathrm{Re}(\alpha) \text{ is of the hyperideal type, } \max\{\mathrm{Re}(\tau_i)\} \leqslant \mathrm{Re}(\xi) \leqslant \min\{\mathrm{Re}(\eta_j), 2\pi\} \right\}$ where $\alpha = (\alpha_1, \dots, \alpha_6)$ and $\mathrm{Re}(\alpha) = (\mathrm{Re}(\alpha_1), \dots, \mathrm{Re}(\alpha_6))$; and \mathcal{W}^{ϵ_I} is continuous on

$$D_{\mathrm{H},\mathbb{C}} = \left\{ (\alpha_I, \xi_1) \in \mathbb{C}^{|I|+c} \mid (\mathrm{Re}(\alpha_I), \mathrm{Re}(\xi)) \in \mathrm{D}_{\mathrm{H}} \right\}$$

and for any $\delta > 0$ is analytic on

$$\mathbf{D}_{\mathbf{H},\mathbb{C}}^{\delta} = \left\{ (\alpha_I, \xi_1) \in \mathbb{C}^{|I|+c} \mid (\operatorname{Re}(\alpha_I), \operatorname{Re}(\xi)) \in \mathbf{D}_{\mathbf{H}}^{\delta} \right\},\$$

where $\operatorname{Re}(\alpha_I) = (\operatorname{Re}(\alpha_i))_{i \in I}$ and $\operatorname{Re}(\xi) = (\operatorname{Re}(\xi_1), \dots, \operatorname{Re}(\xi_c))$. Let

$$\mathbf{B}_{\mathrm{H}} = \mathbf{B}_{\mathrm{H},\mathbb{C}} \cap \mathbb{R}^7$$

Then by (2.10), for $(\alpha_1, \ldots, \alpha_6, \xi) \in B_H$,

$$U(\alpha_1, \dots, \alpha_6, \xi) = 2\pi^2 + 2\sqrt{-1}V(\alpha_1, \dots, \alpha_6, \xi)$$
(4.2)

for $V : B_H \to \mathbb{R}$ defined by

$$V(\alpha_{1},...,\alpha_{6},\xi) = \delta(\alpha_{1},\alpha_{2},\alpha_{3}) + \delta(\alpha_{1},\alpha_{5},\alpha_{6}) + \delta(\alpha_{2},\alpha_{4},\alpha_{6}) + \delta(\alpha_{3},\alpha_{4},\alpha_{5}) - \Lambda(\xi) + \sum_{i=1}^{4} \Lambda(\xi - \tau_{i}) + \sum_{j=1}^{3} \Lambda(\eta_{j} - \xi),$$
(4.3)

where δ is defined by

$$\delta(x, y, z) = -\frac{1}{2}\Lambda\Big(\frac{x+y-z}{2}\Big) - \frac{1}{2}\Lambda\Big(\frac{y+z-x}{2}\Big) - \frac{1}{2}\Lambda\Big(\frac{z+x-y}{2}\Big) + \frac{1}{2}\Lambda\Big(\frac{x+y+z}{2}\Big).$$

A result of Costantino [8] shows that for each $\alpha = (\alpha_1, \dots, \alpha_6)$ of the hyperideal type, there exists a unique $\xi(\alpha)$ so that $(\alpha, \xi(\alpha)) \in B_H$ and

$$\frac{\partial V(\alpha,\xi)}{\partial \xi}\Big|_{\xi=\xi(\alpha)} = 0.$$

Indeed, he proves that for each α , V is strictly concave down in ξ with derivatives $\pm \infty$ at the boundary points of the interval of ξ , hence there is a unique critical point $\xi(\alpha)$ at which V achieves the absolute maximum. Moreover, by using the Murakami-Yano formula [27, 40] he shows that

$$V(\alpha,\xi(\alpha)) = \operatorname{Vol}(\Delta_{|\alpha-\pi|}), \tag{4.4}$$

the volume of the ideal or the truncated hyperideal tetrahedron with dihedral angles $|\alpha_1 - \pi|, \ldots, |\alpha_6 - \pi|$. Let l_1, \ldots, l_6 be the lengths of the edges of $\Delta_{|\alpha - \pi|}$, and let $u_i = 2\sqrt{-1}|\alpha_i - \pi|$ for $i \in \{1, \ldots, 6\}$. Then by the Schläfi formula, we have

$$\frac{\partial U(\alpha,\xi(\alpha))}{\partial u_i} = -\frac{l_i}{2}.$$
(4.5)

For $\alpha = (\alpha_1, \ldots, \alpha_6) \in \mathbb{C}^6$ such that $(\operatorname{Re}(\alpha_1), \ldots, \operatorname{Re}(\alpha_6))$ is of the hyperideal type, we let $\xi(\alpha)$ be such that

$$\frac{\partial U(\alpha,\xi)}{\partial \xi}\Big|_{\xi=\xi(\alpha)} = 0.$$
(4.6)

Following the idea of [27], see also [3], it is proved that $e^{-2\sqrt{-1}\xi(\alpha)}$ satisfies a concrete quadratic equation. Therefore, for each such α , there is at most one $\xi(\alpha)$ such that $(\alpha, \xi(\alpha)) \in B_{H,\mathbb{C}}$. At this point, we do not know whether $(\alpha, \xi(\alpha)) \in B_{H,\mathbb{C}}$ for all such α , but in the next section we will show that it is the case if all $\operatorname{Re}(\alpha_1), \ldots, \operatorname{Re}(\alpha_6)$ are sufficiently close to π .

For $s \in \{1, \ldots, c\}$, let $\alpha_s = (\alpha_{s_1}, \ldots, \alpha_{s_6})$. For a fixed $\alpha_J = (\alpha_j)_{j \in J}$, define the following function

$$\mathcal{U}(\alpha_I) = \sum_{s=1}^{c} U(\alpha_s, \xi(\alpha_s))$$

for all α_I such that $(\alpha_s, \xi(\alpha_s)) \in B_{H,\mathbb{C}}$ for all $s \in \{1, \ldots, c\}$.

The next proposition shows that with an appropriate choice of the meridians and longitudes, the value of \mathcal{U} coincides with the value of the Neumann-Zagier potential function [28] defined on a neighborhood of the complete structure in the deformation space of $M_c \smallsetminus L_{FSL}$.

Proposition 4.1. For each component T_i of the boundary of $M_c \setminus L_{FSL}$, choose the basis (u_i, v_i) of $\pi_1(T_i)$ as in (2.6) and (2.7), and let Φ be the Neumann-Zagier potential function characterized by

$$\begin{pmatrix}
\frac{\partial \Phi(\mathrm{H}(u_1),\ldots,\mathrm{H}(u_n))}{\partial \mathrm{H}(u_i)} = \frac{\mathrm{H}(v_i)}{2}, \\
\Phi(0,\ldots,0) = \sqrt{-1} \left(\mathrm{Vol}(M_c \smallsetminus L_{FSL}) + \sqrt{-1} \mathrm{CS}(M_c \smallsetminus L_{FSL}) \right) \mod \pi^2 \mathbb{Z},
\end{cases}$$
(4.7)

where $M_c \setminus L_{FSL}$ is with the complete hyperbolic metric. If $H(u_i) = \pm 2\sqrt{-1}(\alpha_i - \pi)$ for each $i \in \{1, \ldots, n\}$, then

$$\mathcal{U}(\alpha_I) = 2c\pi^2 + \Phi(\mathbf{H}(u_1), \dots, \mathbf{H}(u_n)).$$

Proof. We assume that $H(u_i) = 2\sqrt{-1}(\alpha_i - \pi)$, and the case $H(u_i) = -2\sqrt{-1}(\alpha_i - \pi)$ follows from the fact that Φ is even in its variables.

From the construction of L_{FSL} in Section 2.4, the component $L_i = \bigcup e_{s_j}$, where e_{s_j} is the *j*-th edge of the building block Δ_s for *s* coming from a subset *S* of $\{1, \ldots, c\}$ and *j* coming from a subset of $\{1, \ldots, 6\}$ depending on *s*. Let l_i be the lengths of L_i and let l_{s_j} be the length of e_{s_j} for each s_j . Then by (2.6), (2.7) and (4.5), we have

$$\frac{\partial \mathcal{U}(\alpha_I)}{\partial \mathcal{H}(u_i)} = \sum_{s \in S} \frac{\partial \mathcal{U}(\alpha_s, \xi(\alpha_s))}{\partial \mathcal{H}(u_i)} = \sum_{s_j} -\frac{l_{s_j}}{2} = -\frac{l_i}{2} = \frac{\mathcal{H}(v_i)}{2}.$$
(4.8)

This means $\frac{\partial \mathcal{U}}{\partial \mathrm{H}(u_i)}$ and $\frac{\partial \Phi}{\partial \mathrm{H}(u_i)}$ coincide for purely imaginary variables for each $i \in I$. Then by Lemma 4.2 below, $\frac{\partial \mathcal{U}}{\partial \mathrm{H}(u_i)}$ and $\frac{\partial \Phi}{\partial \mathrm{H}(u_i)}$ coincide, verifying the first equality of (4.7).

For the second equality of (4.7), we have $\mathcal{U}(\pi, \ldots, \pi) = 2c\pi^2 + 2cv_8\sqrt{-1}$, where v_8 is the volume of the regular ideal octahedron. Since $\operatorname{Vol}(M_c \setminus L_{FSL}) = 2cv_8$ and $\operatorname{CS}(M_c \setminus L_{FSL}) = 0$, we have

$$\mathcal{U}(\pi,\ldots,\pi) = 2c\pi^2 + \Phi(0,\ldots,0).$$

Lemma 4.2. Suppose D is a domain of \mathbb{C}^n and F_1 and F_2 are two holomorphic functions on D. If F_1 and F_2 coincide on $D \cap (\sqrt{-1}\mathbb{R})^n$, then F_1 and F_2 coincide on D.

Proof. We use induction on n. If n = 1, then the result follows from the Identity Theorem of a single variable analytic function. Now suppose the result is true for $n \leq k$. For each fixed $(z_2, \ldots, z_k) \in (\sqrt{-1}\mathbb{R})^{k-1}$, by the assumption of the lemma, we have $F_1(z_1, z_2, \ldots, z_k) = F_2(z_1, z_2, \ldots, z_k)$ for any purely imaginary z_1 . Then by the single variable case $F_1(z_1, z_2, \ldots, z_k) = F_2(z_1, z_2, \ldots, z_k)$ for any complex z_1 . This equality can also be understood as for any fixed complex z_1 , $F_1(z_1, z_2, \ldots, z_k) = F_2(z_1, z_2, \ldots, z_k)$ for all purely imaginary (z_2, \ldots, z_k) . Then by the reduction hypothesis, we have $F_1(z_1, z_2, \ldots, z_k) = F_2(z_1, z_2, \ldots, z_k) = F_2(z_1, z_2, \ldots, z_k)$ for all (z_2, \ldots, z_k) .

5 Asymptotics

The goal of this section is to prove Theorem 3.1. The main tool we use is Proposition 5.1, which is a generalization of the standard Saddle Point Approximation [29]. For the readers' convenience, we include a proof of Proposition 5.1 in Appendix A.

Proposition 5.1. Let $D_{\mathbf{z}}$ be a region in \mathbb{C}^n and let $D_{\mathbf{a}}$ be a region in \mathbb{R}^k . Let $f(\mathbf{z}, \mathbf{a})$ and $g(\mathbf{z}, \mathbf{a})$ be complex valued functions on $D_{\mathbf{z}} \times D_{\mathbf{a}}$ which are holomorphic in \mathbf{z} and smooth in \mathbf{a} . For each positive integer r, let $f_r(\mathbf{z}, \mathbf{a})$ be a complex valued function on $D_{\mathbf{z}} \times D_{\mathbf{a}}$ holomorphic in \mathbf{z} and smooth in \mathbf{a} . For a fixed $\mathbf{a} \in D_{\mathbf{a}}$, let $f^{\mathbf{a}}$, $g^{\mathbf{a}}$ and $f^{\mathbf{a}}_r$ be the holomorphic functions on $D_{\mathbf{z}}$ defined by $f^{\mathbf{a}}(\mathbf{z}) = f(\mathbf{z}, \mathbf{a})$, $g^{\mathbf{a}}(\mathbf{z}) = g(\mathbf{z}, \mathbf{a})$ and $f^{\mathbf{a}}_r(\mathbf{z}) = f_r(\mathbf{z}, \mathbf{a})$. Suppose $\{\mathbf{a}_r\}$ is a convergent sequence in $D_{\mathbf{a}}$ with $\lim_r \mathbf{a}_r = \mathbf{a}_0$, $f^{\mathbf{a}_r}_r$ is of the form

$$f_r^{\mathbf{a}_r}(\mathbf{z}) = f^{\mathbf{a}_r}(\mathbf{z}) + \frac{\upsilon_r(\mathbf{z}, \mathbf{a}_r)}{r^2},$$

 $\{S_r\}$ is a sequence of embedded real n-dimensional closed disks in D_z sharing the same boundary, and \mathbf{c}_r is a point on S_r . Suppose $\{\mathbf{c}_r\}$ is convergent in D_z with $\lim_r \mathbf{c}_r = \mathbf{c}_0$. If and for each r

- (1) \mathbf{c}_r is a critical point of $f^{\mathbf{a}_r}$ in $D_{\mathbf{z}}$,
- (2) $\operatorname{Re} f^{\mathbf{a}_r}(\mathbf{c}_r) > \operatorname{Re} f^{\mathbf{a}_r}(\mathbf{z})$ for all $\mathbf{z} \in S \setminus \{\mathbf{c}_r\}$,

- (3) the Hessian matrix $\operatorname{Hess}(f^{\mathbf{a}_r})$ of $f^{\mathbf{a}_r}$ at \mathbf{c}_r is non-singular
- (4) $|g^{\mathbf{a}_r}(\mathbf{c}_r)|$ is bounded from below by a positive constant independent of r,
- (5) $|v_r(\mathbf{z}, \mathbf{a}_r)|$ is bounded from above by a constant independent of r on $D_{\mathbf{z}}$, and
- (6) the Hessian matrix $\operatorname{Hess}(f^{\mathbf{a}_0})$ of $f^{\mathbf{a}_0}$ at \mathbf{c}_0 is non-singular,

then

$$\int_{S_r} g^{\mathbf{a}_r}(\mathbf{z}) e^{rf_r^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z} = \left(\frac{2\pi}{r}\right)^{\frac{n}{2}} \frac{g^{\mathbf{a}_r}(\mathbf{c}_r)}{\sqrt{-\det\operatorname{Hess}(f^{\mathbf{a}_r})(\mathbf{c}_r)}} e^{rf^{\mathbf{a}_r}(\mathbf{c}_r)} \left(1 + O\left(\frac{1}{r}\right)\right).$$

In the rest of this paper, we assume that $\theta_1, \ldots, \theta_n$ are sufficiently close to 0, or equivalently, $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are sufficiently close to π . In the special case that $\beta_i = \alpha_j = \pi$ for all $i \in I$ and $j \in J$, a direct computation shows that $\xi(\pi, \ldots, \pi) = \frac{7\pi}{4}$. For $\delta > 0$, we denote by $D_{\delta,\mathbb{C}}$ the L^1 δ -neighborhood of $(\pi, \ldots, \pi, \frac{7\pi}{4}, \ldots, \frac{7\pi}{4})$ in $\mathbb{C}^{|I|+c}$, that is

$$\mathbf{D}_{\delta,\mathbb{C}} = \Big\{ (\alpha_I, \xi) \in \mathbb{C}^{|I|+c} \ \Big| \ d_{L^1}\Big((\alpha_I, \xi_1), \Big(\pi, \dots, \pi, \frac{7\pi}{4}, \dots, \frac{7\pi}{4}\Big) \Big) < \delta \Big\},\$$

where d_{L^1} is the real L^1 -norm on \mathbb{C}^n defined by

$$d_{L^1}(\mathbf{x}, \mathbf{y}) = \max_{i \in \{1, \dots, n\}} \{ |\operatorname{Re}(x_i) - \operatorname{Re}(y_i)|, |\operatorname{Im}(x_i) - \operatorname{Im}(y_i)| \}$$

where $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$. We will also consider the region

$$\mathbf{D}_{\delta} = \mathbf{D}_{\delta,\mathbb{C}} \cap \mathbb{R}^{|I|+\epsilon}$$

5.1 Critical points and critical values of W^{ϵ_I}

Suppose $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are sufficiently close to π . For $i \in I$, let $\theta_i = 2|\beta_i - \pi|$, and let $\mu_i = 1$ if $\beta_i \ge \pi$ and let $\mu_i = -1$ if $\beta_i \le \pi$ so that $\mu_i \theta_i = 2(\beta_i - \pi)$.

Proposition 5.2. For each $i \in I$, let $H(u_i)$ be the logarithmic holonomy of u_i of the hyperbolic cone manifold $M_{L_{\theta}}$ and let

$$\alpha_i = \pi + \frac{\epsilon_i \mu_i \sqrt{-1}}{2} \mathbf{H}(u_i).$$
(5.1)

For $s \in \{1, ..., c\}$, let $\alpha_s = (\alpha_{s_1}, ..., \alpha_{s_6})$ and let $\xi(\alpha_s)$ be as defined in (4.6). Then W^{ϵ_I} has a critical point

$$z^{\epsilon_I} = \left(\left(\alpha_i \right)_{i \in I}, \left(\xi(\alpha_s) \right)_{s=1}^c \right)$$

in $D_{\delta,\mathbb{C}}$ with critical value

$$2c\pi^2 + \sqrt{-1} \Big(\operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(M_{L_{\theta}}) \Big).$$

Proof. For $i \in I$, if θ_i is sufficiently close to 0, then the come metric is sufficiently close to the complete metric. As a consequence, $H(u_i)$ is sufficiently close to 0, and hence α_i is sufficiently close to π . Then by the continuity of $\xi(\alpha_s), z^{\epsilon_I} \in D_{\delta,\mathbb{C}}$ for sufficiently small $\theta_1, \ldots, \theta_n$.

For any $(\alpha_i)_{i \in I}$ such that $(\alpha_I, (\xi(\alpha_s)_{s=1}^c) \in D_{H,C})$, and for $s \in \{1, \ldots, c\}$,

$$\frac{\partial \mathcal{W}^{\epsilon_I}}{\partial \xi_s}\Big|_{\left((\alpha_i)_{i\in I}, (\xi(\alpha_s))_{s=1}^c\right)} = \frac{\partial U(\alpha_s)}{\partial \xi_s}\Big|_{\xi(\alpha_s)} = 0.$$
(5.2)

In particular, for each $s \in \{1, \ldots, c\}$,

$$\frac{\partial \mathcal{W}^{\epsilon_I}}{\partial \xi_s}\Big|_{z^{\epsilon_I}} = 0.$$
(5.3)

By the Chain Rule, for each $s \in \{1, \ldots, c\}$ and $i \in I$,

$$\frac{\partial U(\alpha_s,\xi(\alpha_s))}{\partial \alpha_i}\Big|_{\alpha_s} = \frac{\partial U}{\partial \alpha_i}\Big|_{(\alpha_s,\xi(\alpha_s))} + \frac{\partial U}{\partial \xi}\Big|_{(\alpha_s,\xi(\alpha_s))} \cdot \frac{\partial \xi(\alpha_s)}{\partial \alpha_i}\Big|_{\alpha_s} = \frac{\partial U}{\partial \alpha_i}\Big|_{(\alpha_s,\xi(\alpha_s))},$$

hence

$$\frac{\partial \left(\sum_{s=1}^{c} U(\alpha_s, \xi_s)\right)}{\partial \alpha_i}\Big|_{z^{\epsilon_I}} = \frac{\partial \mathcal{U}}{\partial \alpha_i}\Big|_{\alpha_I} = -\epsilon_i \mu_i \sqrt{-1} \mathbf{H}(v_i),$$

where the last equation comes from (4.8) and (5.1). As a consequence, for each $i \in I$, we have

$$\frac{\partial \mathcal{W}^{\epsilon_I}}{\partial \alpha_i}\Big|_{z^{\epsilon_I}} = -2p_i(\alpha_i - \pi) - 2\epsilon_i(\beta_i - \pi) + \frac{\partial \mathcal{U}}{\partial \alpha_i}\Big|_{\alpha_I}
= -\epsilon_i\mu_i\sqrt{-1}\Big(p_i\mathrm{H}(u_i) - \sqrt{-1}\theta_i + \mathrm{H}(v_i)\Big) = 0,$$
(5.4)

where the last equality comes from the $(p_i, 1)$ -Dehn filling equation (2.2) with the cone angle θ_i . Equations (5.3) and (5.4) show that z^{ϵ_I} is a critical point of W^{ϵ_I} .

To compute the critical value, by Proposition 4.1, we first have

$$\mathcal{U}(\alpha_I) = 2c\pi^2 + \Phi(\mathcal{H}(u_1), \dots, \mathcal{H}(u_n)),$$
(5.5)

where $\alpha_I = (\alpha_i)_{i \in I} = \left(\pi + \frac{\epsilon_i \mu_i \sqrt{-1}}{2} \mathbf{H}(u_i)\right)_{i \in I}$.

For each $i \in I$, let $\gamma_i = -u_i + q_i(p_iu_i + v_i)$ so that it is the curve on the boundary of a tubular neighborhood of L_i^* that is isotopic to L_i^* given by the framing q_i of L_i^* and with the orientation so that $(p_iu_i + v_i) \cdot \gamma_i = 1$. Then we have $\theta_i = 2\mu_i(\beta_i - \pi)$, $H(u_i) = -2\epsilon_i\mu_i\sqrt{-1}(\alpha_i - \pi)$,

$$\mathbf{H}(v_i) = \sqrt{-1}\theta_i - p_i \mathbf{H}(u_i) = 2\mu_i \sqrt{-1}(\beta_i - \pi) + 2p_i \epsilon_i \mu_i \sqrt{-1}(\alpha_i - \pi)$$

and

$$H(\gamma) = -H(u_i) + q_i(p_i H(u_i) + H(v_i)) = 2\epsilon_i \mu_i \sqrt{-1}(\alpha_i - \pi) + 2q_i \mu_i \sqrt{-1}(\beta_i - \pi).$$

As a consequence, we have

=

$$-\sum_{i\in I} \frac{\mathrm{H}(u_i)\mathrm{H}(v_i)}{4} + \sum_{i\in I} \frac{\sqrt{-1}\theta_i\mathrm{H}(\gamma_i)}{4}$$

= $-\sum_{i\in I} q_i(\beta_i - \pi)^2 - \sum_{i\in I} p_i(\alpha_i - \pi)^2 - \sum_{i\in I} 2\epsilon_i(\alpha_i - \pi)(\beta_i - \pi).$ (5.6)

For each $j \in J$, let $\gamma_j = p_j u_j + v_j$ so that it is the curve on the boundary of a tubular neighborhood of L_j that is isotopic to L_j given by the framing p_j of L_j and with the orientation so that $u_j \cdot \gamma_j = 1$. Then we have $\theta_j = 2|\alpha_j - \pi|$, $H(u_j) = 2\sqrt{-1}|\alpha_j - \pi|$, $H(v_j) = -l_j$ and

$$\mathbf{H}(\gamma_j) = p_j \mathbf{H}(u_j) + \mathbf{H}(v_j) = 2p_j \sqrt{-1} |\alpha_j - \pi| - l_j.$$

As a consequence, we have

$$-\sum_{j\in J} \frac{\mathrm{H}(u_j)\mathrm{H}(v_j)}{4} + \sum_{j\in J} \frac{\sqrt{-1}\theta_j\mathrm{H}(\gamma_j)}{4} = \sum_{j\in J} -p_j(\alpha_j - \pi)^2.$$
 (5.7)

Putting (5.5), (5.6), (5.7) and (2.5) together, we have

$$\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}}) = \mathcal{U}(\alpha_{I}) - \sum_{i \in I} q_{i}(\beta_{i} - \pi)^{2} - \sum_{i=1}^{n} p_{i}(\alpha_{i} - \pi)^{2} - \sum_{i \in I} 2\epsilon_{i}(\alpha_{i} - \pi)(\beta_{i} - \pi)$$
$$= 2c\pi^{2} + \Phi(\mathcal{H}(u_{1}), \dots, \mathcal{H}(u_{n})) - \sum_{i=1}^{n} \frac{\mathcal{H}(u_{i})\mathcal{H}(v_{i})}{4} + \sum_{i=1}^{n} \frac{\sqrt{-1}\theta_{i}\mathcal{H}(\gamma_{i})}{4}$$
$$= 2c\pi^{2} + \sqrt{-1} \Big(\operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1}\operatorname{CS}(M_{L_{\theta}}) \Big).$$

5.2 Convexity of \mathcal{W}^{ϵ_I}

Proposition 5.3. There exists a $\delta_0 > 0$ such that if all $\{\alpha_j\}_{j \in J}$ are in $(\pi - \delta_0, \pi + \delta_0)$, then for any ϵ_I , Im $\mathcal{W}^{\epsilon_I}(\alpha_I, \xi)$ is strictly concave down in $\{\operatorname{Re}(\alpha_i)\}_{i \in I}$ and $\{\operatorname{Re}(\xi_s)\}_{s=1}^c$, and is strictly concave up in $\{\operatorname{Im}(\alpha_i)\}_{i \in I}$ and $\{\operatorname{Im}(\xi_s)\}_{s=1}^c$ on $\operatorname{D}_{\delta_0,\mathbb{C}}$.

Proof. We first consider the special case $\{\alpha_i\}_{i \in I}$ and $\{\xi_s\}_{s=1}^c$ are real. In this case,

Im
$$\mathcal{W}^{\epsilon_I}(\alpha_I,\xi) = \sum_{s=1}^c 2V(\alpha_{s_1},\ldots,\alpha_{s_6},\xi_s)$$

with V defined in (4.3).

At $(\pi, \ldots, \pi, \frac{7\pi}{4})$, we have $\frac{\partial^2 \text{Im}V}{\partial \alpha_{s_i}^2} = -2$ for $s_i \in I \cap \{s_1, \ldots, s_6\}$, $\frac{\partial^2 V}{\partial \alpha_{s_i} \alpha_{s_j}} = -1$ for $s_i \neq s_j$ in $I \cap \{s_1, \ldots, s_6\}$, $\frac{\partial^2 \text{Im}V}{\partial \alpha_{s_i} \xi_s} = 2$ for $s_i \in I \cap \{s_1, \ldots, s_6\}$ and $\frac{\partial^2 \text{Im}V}{\partial \xi_s^2} = -8$. Then a direct computation shows that, at $(\pi, \ldots, \pi, \frac{7\pi}{4})$, the Hessian matrix of ImV in $\{\text{Re}(\alpha_i)\}_{i \in I \cap \{s_1, \ldots, s_6\}}$ and $\text{Re}(\xi_s)$ is negative definite. As a consequence, the Hessian matrix of Im \mathcal{W}^{ϵ_I} in $\{\text{Re}(\alpha_i)\}_{i \in I}$ and $\{\text{Re}(\xi_s)\}_{s=1}^c$ is negative definite at $(\pi, \ldots, \pi, \frac{7\pi}{4}, \ldots, \frac{7\pi}{4})$.

Then by the continuity, there exists a sufficiently small $\delta_0 > 0$ such that for all $\{\alpha_j\}_{j\in J}$ in $(\pi - \delta_0, \pi + \delta_0)$ and $(\alpha_I, \xi) \in D_{\delta_0,\mathbb{C}}$, the Hessian matrix of $\operatorname{Im} \mathcal{W}^{\epsilon_I}$ with respect to $\{\operatorname{Re}(\alpha_i)\}_{i\in I}$ and $\{\operatorname{Re}(\xi_s)\}_{s=1}^c$ is still negative definite, implying that $\operatorname{Im} \mathcal{W}^{\epsilon_I}$ is strictly concave down in $\{\operatorname{Re}(\alpha_i)\}_{i\in I}$ and $\{\operatorname{Re}(\xi_s)\}_{s=1}^c$ on $D_{\delta_0,\mathbb{C}}$. Since \mathcal{W}^{ϵ_I} is holomorphic, $\operatorname{Im} \mathcal{W}^{\epsilon_I}$ is strictly concave up in $\{\operatorname{Im}(\alpha_i)\}_{i\in I}$ and $\{\operatorname{Im}(\xi_s)\}_{s=1}^c$ on $D_{\delta_0,\mathbb{C}}$.

Proposition 5.4. If all $\{\alpha_j\}_{j\in J}$ are in $(\pi - \delta_0, \pi + \delta_0)$, then the Hessian matrix Hess \mathcal{W}^{ϵ_I} of \mathcal{W}^{ϵ_I} with respect to $\{\alpha_i\}_{i\in I}$ and $\{\xi_s\}_{s=1}^c$ is non-singular on $D_{\delta_0,\mathbb{C}}$.

Proof. By Proposition 5.3, the real part of the Hess W^{ϵ_I} is negative definite. Then by [24, Lemma], it is nonsingular.

5.3 Asymptotics of the leading Fourier coefficients

Proposition 5.5. Suppose $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are in $\{\pi - \epsilon, \pi + \epsilon\}$ for a sufficiently small $\epsilon > 0$. For $\epsilon_I \in \{1, -1\}^I$, let z^{ϵ_I} be the critical point of W^{ϵ_I} described in Proposition 5.2. Then

$$\widehat{f_r^{\epsilon_I}}(0,\ldots,0) = \frac{C^{\epsilon_I}(z^{\epsilon_I})}{\sqrt{-\det\operatorname{Hess}\mathcal{W}^{\epsilon_I}(z^{\epsilon_I})}} e^{\frac{r}{4\pi} \left(\operatorname{Vol}(M_{L_\theta}) + \sqrt{-1}\operatorname{CS}(M_{L_\theta})\right)} \left(1 + O\left(\frac{1}{r}\right)\right)$$

where each $C^{\epsilon_I}(z^{\epsilon_I})$ depends continuously on $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$; and when $\beta_i = \alpha_j = \pi$,

$$C^{\epsilon_I}(z^{\epsilon_I}) = (-1)^{\sum_{i \in I} q_i + \sum_{i=1}^n p_i + c} \frac{r^{\frac{|I| - c}{2}}}{2^{\frac{3|I| + c}{2}} \pi^{\frac{|I| + c}{2}}}$$

For the proof of Proposition 5.5, we need the following lemma.

Lemma 5.6. For each $\epsilon_I \in \{1, -1\}^I$ and any fixed $\{\alpha_j\}_{j \in J}$,

$$\max_{\mathrm{D}_{\mathrm{H}}} \mathrm{Im} \mathcal{W}^{\epsilon_{I}} \leqslant \mathrm{Im} \mathcal{W}^{\epsilon_{I}} \left(\pi, \dots, \pi, \frac{7\pi}{4}, \dots, \frac{7\pi}{4} \right) = 2cv_{8}$$

where v_8 is the volume of the regular ideal octahedron, and the equality holds if and only if $\alpha_1 = \cdots = \alpha_n = \pi$ and $\xi_1 = \cdots = \xi_c = \frac{7\pi}{4}$.

Proof. On D_H , we have

Im
$$\mathcal{W}^{\epsilon_I}(\alpha_I,\xi) = \sum_{s=1}^c 2V(\alpha_{s_1},\ldots,\alpha_{s_6},\xi_s)$$

for V defined in (4.3). Then the result is a consequence of the result of Costantino [8] and the Murakami-Yano formula [27] (see Ushijima [40] for the case of hyperideal tetrahedra). Indeed, by [8], for a fixed $\alpha = (\alpha_1, \ldots, \alpha_6)$ of the hyperideal type, the function $f(\xi)$ defined by $f(\xi) = V(\alpha, \xi)$ is strictly concave down and the unique maximum point $\xi(\alpha)$ exists and lies in $(\max\{\tau_i\}, \min\{\eta_j, 2\pi\})$, ie, $(\alpha, \xi(\alpha)) \in B_H$. Then by [40], $V(\alpha, \xi(\alpha)) = \operatorname{Vol}(\Delta_{|\alpha-\pi|})$, the volume of the truncated hyperideal tetrahedron $\Delta_{|\alpha-\pi|}$ with dihedral angles $|\alpha_1 - \pi|, \ldots, |\alpha_6 - \pi|$. Since $\xi(\pi, \ldots, \pi) = \frac{7\pi}{4}$ and the regular ideal cetahedron $\Delta_{(0,\ldots,0)}$ has the maximum volume among all the truncated hyperideal tetrahedra, $V(\pi, \ldots, \pi, \frac{7\pi}{4}) = v_8 = \operatorname{Vol}(\Delta_{(0,\ldots,0)}) \ge \operatorname{Vol}(\Delta_{|\alpha-\pi|}) = V(\alpha, \xi(\alpha)) \ge V(\alpha, \xi)$ for any $(\alpha, \xi) \in B_H$.

For the equality part, suppose $(\alpha_I, \xi) \neq (\pi, \dots, \pi, \frac{7\pi}{4}, \dots, \frac{7\pi}{4})$. If $(\alpha_{s_1}, \dots, \alpha_{s_6}) \neq (\pi, \dots, \pi)$ for some $s \in \{1, \dots, c\}$, then $\operatorname{Im}\mathcal{W}^{\epsilon_I}(\alpha_I, \xi) \leq 2\operatorname{Vol}(\Delta_{|\alpha-\pi|}) + 2(c-1)v_8 < 2cv_8$. If $(\alpha_I) = (\pi, \dots, \pi)$ but $\xi_s \neq \frac{7\pi}{4}$ for some $s \in \{1, \dots, c\}$, then the strict concavity of $f(\xi)$ implies that $\operatorname{Im}\mathcal{W}^{\epsilon_I}(\pi, \dots, \pi, \xi) < \operatorname{Im}\mathcal{W}^{\epsilon_I}(\pi, \dots, \pi, \frac{7\pi}{4}, \dots, \frac{7\pi}{4}) = 2cv_8$.

Proof of Proposition 5.5. Let $\delta_0 > 0$ be as in Proposition 5.3. By Lemma 5.3, Proposition 5.6 and the compactness of $D_H \setminus D_{\delta_0}$,

$$2cv_8 > \max_{\mathrm{D}_{\mathrm{H}} \smallsetminus \mathrm{D}_{\delta_0}} \mathrm{Im} \mathcal{W}^{\epsilon_I}.$$

By Proposition 5.2 and continuity, if $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are sufficiently close to π , then the critical point z^{ϵ_I} of \mathcal{W}^{ϵ_I} as in Proposition 5.2 lies in $D_{\delta_0,\mathbb{C}}$, and $\operatorname{Im} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I}) = \operatorname{Vol}(M_{L_{\theta}})$ is sufficiently close to $2cv_8$ so that

$$\mathrm{Im}\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}}) > \max_{\mathrm{D}_{\mathrm{H}}\smallsetminus\mathrm{D}_{\delta_{0}}}\mathrm{Im}\mathcal{W}^{\epsilon_{I}}.$$

Therefore, we only need to estimate the integral on D_{δ_0} . To do this, we consider as drawn in Figure 4 the surface $S^{\epsilon_I} = S_{\text{top}}^{\epsilon_I} \cup S_{\text{side}}^{\epsilon_I}$ in $\overline{D_{\delta_0,\mathbb{C}}}$, where

$$S_{\mathsf{top}}^{\epsilon_I} = \{ (\alpha_I, \xi) \in \mathcal{D}_{\delta_0, \mathbb{C}} \mid ((\operatorname{Im}(\alpha_I)), \operatorname{Im}(\xi)) = \operatorname{Im}(z^{\epsilon_I}) \}$$

and

$$S_{\text{side}}^{\epsilon_I} = \{ (\alpha_I, \xi) + t\sqrt{-1} \text{Im}(z^{\epsilon_I}) \mid (\alpha_I, \xi) \in \partial \mathcal{D}_{\delta_0}, t \in [0, 1] \}.$$

By analyticity, the integral remains the same if we deform the domain from D_{δ_0} to S^{ϵ_I} .

By Proposition 5.3, $\text{Im}\mathcal{W}^{\epsilon_I}$ is concave down on $S_{\text{top}}^{\epsilon_I}$. Since z^{ϵ_I} is the critical points of $\text{Im}\mathcal{W}^{\epsilon_I}$, it is the only absolute maximum on $S_{\text{top}}^{\epsilon_I}$.

On the side $S_{\text{side}}^{\epsilon_I}$, for each $(\alpha_I, \xi) \in \partial D_{\delta_0}$, we consider the function

$$g_{(\alpha_I,\xi)}^{\epsilon_I}(t) = \operatorname{Im} \mathcal{W}^{\epsilon_I} \left((\alpha_I,\xi) + t\sqrt{-1} \operatorname{Im}(z^{\epsilon_I}) \right)$$



Figure 4: The deformed surface S^{ϵ_I}

on [0, 1]. By Lemma 5.3, $g_{(\alpha_I,\xi)}^{\epsilon_I}(t)$ is concave up for any $(\alpha_I,\xi) \in \partial D_{\delta_0}$. As a consequence, $g_{(\alpha_I,\xi)}^{\epsilon_I}(t) \leq \max\{g_{(\alpha_I,\xi)}^{\epsilon_I}(0), g_{(\alpha_I,\xi)}^{\epsilon_I}(1)\}$. Now by the previous two steps, since $(\alpha_I,\xi) \in \partial D_{\delta_0}$,

$$g_{(\alpha_I,\xi)}^{\epsilon_I}(0) = \operatorname{Im} \mathcal{W}^{\epsilon_I}(\alpha_I,\xi) < \operatorname{Im} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I});$$

and since $(\alpha_I, \xi) + \sqrt{-1} \text{Im}(z^{\epsilon_I}) \in S_{\text{top}}^{\epsilon_I}$,

$$g_{(\alpha_I,\xi)}^{\epsilon_I}(0) = \operatorname{Im} \mathcal{W}^{\epsilon_I}((\alpha_I,\xi) + \sqrt{-1} \operatorname{Im}(z^{\epsilon_I})) < \operatorname{Im} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I}).$$

As a consequence,

$$\operatorname{Im} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I}) > \max_{\substack{S_{\operatorname{side}}^{\epsilon_I}}} \operatorname{Im} \mathcal{W}^{\epsilon_I}.$$

Therefore, we proved that z^{ϵ_I} is the unique maximum point of $\operatorname{Im} \mathcal{W}^{\epsilon_I}$ on $S^{\epsilon_I} \cup (D_H \setminus D_{\delta_0})$, and \mathcal{W}^{ϵ_I} has critical value $2c\pi^2 + \sqrt{-1} (\operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1} \operatorname{Vol}(M_{L_{\theta}}))$ at z^{ϵ_I} .

By Proposition 5.4, det Hess $\mathcal{W}^{\epsilon_I}(z^{\epsilon_I}) \neq 0$.

Finally, we estimate the difference between $W_r^{\epsilon_I}$ and W^{ϵ_I} . By Lemma 2.10, (3), we have

$$\varphi_r\left(\frac{\pi}{r}\right) = \operatorname{Li}_2(1) + \frac{2\pi\sqrt{-1}}{r}\log\left(\frac{r}{2}\right) - \frac{\pi^2}{r} + O\left(\frac{1}{r^2}\right);$$

and for z with $0 < \text{Rez} < \pi$ we have

$$\varphi_r\left(z+\frac{k\pi}{r}\right) = \varphi_r(z) + \varphi_r'(z)\frac{k\pi}{r} + O\left(\frac{1}{r^2}\right).$$

Then by Lemma 2.10, in $\left\{ (\alpha_I, \xi) \in \overline{\mathbb{D}_{\mathrm{H},\mathbb{C}}^{\delta}} \mid |\mathrm{Im}(\alpha_i)| < L \text{ for } i \in I, |\mathrm{Im}(\xi_s)| < L \text{ for } s \in \{1, \ldots, c\} \right\}$ for some L > 0,

$$\mathcal{W}_r^{\epsilon_I}(\alpha_I,\xi) = \mathcal{W}^{\epsilon_I}(\alpha_I,\xi) - \frac{4c\pi\sqrt{-1}}{r}\log\left(\frac{r}{2}\right) + \frac{4\pi\sqrt{-1}\kappa(\alpha_I,\xi)}{r} + \frac{\nu_r(\alpha_I,\xi)}{r^2},$$

with

$$\begin{split} &\kappa(\alpha_{I},\xi) \\ = \sum_{s=1}^{c} \left(\frac{1}{2}\sum_{i=1}^{4}\sqrt{-1}\tau_{s_{i}} - \sqrt{-1}\xi_{s} - \sqrt{-1}\pi - \frac{\sqrt{-1}\pi}{2} \right. \\ &+ \frac{1}{4}\sum_{i=1}^{4}\sum_{j=1}^{3}\log\left(1 - e^{2\sqrt{-1}(\eta_{s_{j}} - \tau_{s_{i}})}\right) - \frac{3}{4}\sum_{i=1}^{4}\log\left(1 - e^{2\sqrt{-1}(\tau_{s_{i}} - \pi)}\right) \\ &+ \frac{3}{2}\log\left(1 - e^{2\sqrt{-1}(\xi_{s} - \pi)}\right) - \frac{1}{2}\sum_{i=1}^{4}\log\left(1 - e^{2\sqrt{-1}(\xi_{s} - \tau_{s_{i}})}\right) - \frac{1}{2}\sum_{j=1}^{3}\log\left(1 - e^{2\sqrt{-1}(\eta_{s_{j}} - \xi_{s})}\right) \Big) \end{split}$$

and $|\nu_r(\alpha_I, \xi)|$ bounded from above by a constant independent of r. Then

$$e^{\left(\sum_{i\in I}q_{i}\beta_{i}+\sum_{i=1}^{n}p_{i}\alpha_{i}+\sum_{i\in I}\epsilon_{i}(\alpha_{i}+\beta_{i}+\frac{2\pi}{r})\right)\sqrt{-1}+\frac{r}{4\pi\sqrt{-1}}}\mathcal{W}_{r}^{\epsilon_{I}}(\alpha_{I},\xi)}$$

$$=\left(\frac{r}{2}\right)^{-c}e^{\left(\sum_{i\in I}q_{i}\beta_{i}+\sum_{i=1}^{n}p_{i}\alpha_{i}+\sum_{i\in I}\epsilon_{i}(\alpha_{i}+\beta_{i})\right)\sqrt{-1}+\kappa(\alpha_{I},\xi)}\cdot e^{\frac{r}{4\pi\sqrt{-1}}\left(\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)+\frac{\nu_{r}(\alpha_{I},\xi)-\sum_{i\in I}\epsilon_{i}8\pi^{2}}{r^{2}}\right)}.$$

Now let $D_{\mathbf{z}} = \left\{ (\alpha_I, \xi) \in \overline{\mathrm{D}_{\mathrm{H},\mathbb{C}}^{\delta}} \mid |\mathrm{Im}(\alpha_i)| < L \text{ for } i \in I, |\mathrm{Im}(\xi_s)| < L \text{ for } s \in \{1, \ldots, c\} \right\}$ for some L > 0. Let $\mathbf{a}_r = ((\beta_i)_{i \in I}, (\alpha_j)_{j \in J})$ (recall that $\beta_i = \frac{2\pi n_i}{r}$ and $\alpha_j = \frac{2\pi m_j}{r}$ depends on r), $f^{\mathbf{a}_r}(\alpha_I, \xi) = \mathcal{W}^{\epsilon_I}(\alpha_I, \xi), g^{\mathbf{a}_r}(\alpha_I, \xi) = \psi(\alpha_I, \xi) e^{(\sum_{i \in I} q_i \beta_i + \sum_{i=1}^n p_i \alpha_i + \sum_{i \in I} \epsilon_i (\alpha_i + \beta_i))\sqrt{-1} + \kappa(\alpha_I, \xi)}, f^{\mathbf{a}_r}(\alpha_I, \xi) = \mathcal{W}^{\epsilon_I}_r(\alpha_I, \xi) + \frac{4c\pi\sqrt{-1}}{r} \log\left(\frac{r}{2}\right), v_r(\alpha_I, \xi) = \nu_r(\alpha_I, \xi) - \sum_{i \in I} \epsilon_i 8\pi^2, S_r = S^{\epsilon_I} \cup (\mathrm{D}_{\mathrm{H}} \setminus \mathrm{D}_{\delta_0})$ and z^{ϵ_I} is the critical point of f in D. Then all the conditions of Proposition 5.1 are satisfied with and the result follows.

When $\beta_i = \alpha_j = \pi$, a direct computation shows that

$$C^{\epsilon_{I}}(z^{\epsilon_{I}}) = \frac{r^{|I|+c}}{2^{2|I|+c}\pi^{|I|+c}} \left(\frac{2\pi}{r}\right)^{\frac{|I|+c}{2}} \left(\frac{r}{2}\right)^{-c} g\left(\pi, \dots, \pi, \frac{7\pi}{4}, \dots, \frac{7\pi}{4}\right)$$
$$= (-1)^{\sum_{i \in I} q_{i} + \sum_{i=1}^{n} p_{i} + c} \frac{r^{\frac{|I|-c}{2}}}{2^{\frac{3|I|+c}{2}}\pi^{\frac{|I|+c}{2}}}.$$

Corollary 5.7. If $\epsilon > 0$ is sufficiently small and all $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are in $\{\pi - \epsilon, \pi + \epsilon\}$, then

$$\sum_{\epsilon_I \in \{1,-1\}^I} \frac{C^{\epsilon_I}(z^{\epsilon_I})}{\sqrt{-\det \operatorname{Hess} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I})}} \neq 0.$$

Proof. If $\beta_i = \alpha_j = \pi$ for all $i \in I$ and $j \in J$, then all $z^{\epsilon_I} = (\pi, \dots, \pi, \frac{7\pi}{4}, \frac{7\pi}{4})$ and all \mathcal{W}^{ϵ_I} are the same functions. As a consequence, all the $C^{\epsilon_I}(z^{\epsilon_I})$'s and all Hessian determinants det $\text{Hess}\mathcal{W}^{\epsilon_I}(z^{\epsilon_I})$'s are the same at this point, imply that the sum is not equal to zero. Then by continuity, if ϵ is small enough, then the sum remains none zero.

Remark 5.8. We suspect that all $C^{\epsilon_I}(z^{\epsilon_I})$'s and all det $\text{Hess}\mathcal{W}^{\epsilon_I}(z^{\epsilon_I})$'s are always the same for any given $\{\beta_i\}_{i\in I}$ and $\{\alpha_j\}_{j\in J}$.

5.4 Estimate of the other Fourier coefficients

Proposition 5.9. Suppose $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are in $\{\pi - \epsilon, \pi + \epsilon\}$ for a sufficiently small $\epsilon > 0$. If $(\mathbf{a}_I, \mathbf{b}) \neq (0, \dots, 0)$, then

$$\left|\widehat{f_r^{\epsilon_I}}(\mathbf{a}_I, \mathbf{b})\right| < O\left(e^{\frac{r}{4\pi}\left(\operatorname{Vol}(M_{L_{\theta}}) - \epsilon'\right)}\right)$$

for some $\epsilon' > 0$.

Proof. Recall that if $\beta_i = \alpha_j = \pi$ for all $i \in I$ and $j \in J$, then the total derivative

$$D\mathcal{W}^{\epsilon_I}\left(\pi,\ldots,\pi,\frac{7\pi}{4},\ldots,\frac{7\pi}{4}\right) = (0,\ldots,0).$$

Hence there exists a $\delta_1 > 0$ and an $\epsilon > 0$ such that if $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are in $\{\pi - \epsilon, \pi + \epsilon\}$, then for all $(\alpha_I, \xi) \in D_{\delta_1, \mathbb{C}}$ and for any unit vector $\mathbf{u} = ((u_i)_{i \in I}, (w_s)_{s=1}^c) \in \mathbb{R}^{|I|+c}$, the directional derivatives

$$|D_{\mathbf{u}}\mathrm{Im}\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)| = \Big|\sum_{i\in I} u_{i}\frac{\partial\mathrm{Im}\mathcal{W}^{\epsilon_{I}}}{\partial\mathrm{Im}(\alpha_{i})} + \sum_{s=1}^{c} w_{s}\frac{\partial\mathrm{Im}\mathcal{W}^{\epsilon_{I}}}{\partial\mathrm{Im}(\xi_{s})}\Big| < \frac{2\pi - \epsilon''}{2\sqrt{2|I| + 2c}}$$

for some $\epsilon'' > 0$.

On $D_{\rm H}$, we have

$$\operatorname{Im}\left(\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)-\sum_{i\in I}2\pi a_{i}\alpha_{i}-\sum_{s=1}^{c}4\pi b_{s}\xi_{s}\right)=\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi).$$

Then by Lemma 5.3, Proposition 5.6 and the compactness of $D_H \ D_{\delta_1}$,

$$2cv_8 > \max_{\mathbf{D}_{\mathbf{H}} \smallsetminus \mathbf{D}_{\delta_1}} \operatorname{Im} \left(\mathcal{W}^{\epsilon_I}(\alpha_I, \xi) - \sum_{i \in I} 2\pi a_i \alpha_i - \sum_{s=1}^c 4\pi b_s \xi_s \right) + \epsilon'''$$

for some $\epsilon''' > 0$. By Proposition 5.2 and continuity, if $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are sufficiently close to π , then the critical point z^{ϵ_I} of \mathcal{W}^{ϵ_I} as in Proposition 5.2 lies in $D_{\delta_1,\mathbb{C}}$, and $\operatorname{Im} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I}) = \operatorname{Vol}(M_{L_{\theta}})$ is sufficiently close to $2cv_8$ so that

$$\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}}) > \max_{D_{H} \smallsetminus D_{\delta_{1}}} \operatorname{Im}\left(\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi) - \sum_{i \in I} 2\pi a_{i}\alpha_{i} - \sum_{s=1}^{c} 4\pi b_{s}\xi_{s}\right) + \epsilon^{\prime\prime\prime}.$$
(5.8)

Therefore, we only need to estimate the integral on D_{δ_1} .

If $(\mathbf{a}_I, \mathbf{b}) \neq (0, \dots, 0)$, then there is at least one of $\{a_i\}_{i \in I}$ or $\{b_s\}_{s=1}^c$ that is nonzero. Without loss of generality, assume that $a_1 \neq 0$.

If $a_1 > 0$, then consider the surface $S^+ = S^+_{\text{top}} \cup S^+_{\text{side}}$ in $\overline{D_{\delta_1,\mathbb{C}}}$ where

$$S_{top}^{+} = \{ (\alpha_I, \xi) \in \mathcal{D}_{\delta_1, \mathbb{C}} \mid (\operatorname{Im}(\alpha_I), \operatorname{Im}(\xi)) = (\delta_1, 0, \dots, 0) \}$$

and

$$S_{\text{side}}^{+} = \{ (\alpha_{I}, \xi) + (t\sqrt{-1}\delta_{1}, 0, \dots, 0) \mid (\alpha_{I}, \xi) \in \partial \mathbf{D}_{\delta_{1}}, t \in [0, 1] \}.$$

On the top, for any $(\alpha_I, \xi) \in S_{top}^+$, by the Mean Value Theorem,

$$\begin{aligned} \left| \operatorname{Im} \mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}}) - \operatorname{Im} \mathcal{W}^{\epsilon_{I}}(\alpha_{I}, \xi) \right| &= \left| D_{\mathbf{u}} \operatorname{Im} \mathcal{W}^{\epsilon_{I}}(z) \right| \cdot \left\| z^{\epsilon_{I}} - (\alpha_{I}, \xi) \right\| \\ &< \frac{2\pi - \epsilon''}{2\sqrt{2|I| + 2c}} \cdot 2\sqrt{2|I| + 2c} \delta_{1} \\ &= 2\pi \delta_{1} - \epsilon'' \delta_{1}, \end{aligned}$$

where z is some point on the line segment connecting z^{ϵ_I} and (α_I, ξ) , $\mathbf{u} = \frac{z^{\epsilon_I} - (\alpha_I, \xi)}{\|z^{\epsilon_I} - (\alpha_I, \xi)\|}$ and $2\sqrt{2|I| + 2c}\delta_1$ is the diameter of $D_{\delta_1,\mathbb{C}}$. Then

$$\operatorname{Im}\left(\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)-\sum_{i\in I}2\pi a_{i}\alpha_{i}-\sum_{s=1}^{c}4\pi b_{s}\xi_{s}\right)=\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)-2\pi a_{1}\delta_{1}$$
$$<\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}})+2\pi \delta_{1}-\epsilon''\delta_{1}-2\pi \delta_{1}$$
$$=\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}})-\epsilon''\delta_{1}.$$

On the side, for any point $(\alpha_I, \xi) + (t\sqrt{-1}\delta_1, 0, \dots, 0) \in S^+_{\text{side}}$, by the Mean Value Theorem again, we have

$$\left|\operatorname{Im}\mathcal{W}^{\epsilon_{I}}((\alpha_{I},\xi)+(t\sqrt{-1}\delta_{1},0,\ldots,0))-\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)\right|<\frac{2\pi-\epsilon''}{2\sqrt{2|I|+2c}}t\delta_{1}.$$

Then

$$\operatorname{Im} \mathcal{W}^{\epsilon_{I}} \left((\alpha_{I}, \xi) + (t\sqrt{-1}\delta_{1}, 0, \dots, 0) \right) - 2\pi a_{1}t\delta_{1} < \operatorname{Im} \mathcal{W}^{\epsilon_{I}} (\alpha_{I}, \xi) + \frac{2\pi - \epsilon''}{2\sqrt{2|I| + 2c}} t\delta_{1} - 2\pi t\delta_{1} < \operatorname{Im} \mathcal{W}^{\epsilon_{I}} (\alpha_{I}, \xi) < \operatorname{Im} \mathcal{W}^{\epsilon_{I}} (z^{\epsilon_{I}}) - \epsilon''',$$

where the last inequality comes from that $(\alpha_I, \xi) \in \partial D_{\delta_1} \subset D_H \setminus D_{\delta_1}$ and (5.8). Now let $\epsilon' = \min\{\epsilon''\delta_1, \epsilon'''\}$, then on $S^+ \cup (D_H \setminus D_{\delta_1})$,

$$\operatorname{Im}\left(\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)-\sum_{i\in I}2\pi a_{i}\alpha_{i}-\sum_{s=1}^{c}4\pi b_{s}\xi_{s}\right)<\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}})-\epsilon',$$

and the result follows.

If $a_1 < 0$, then we consider the surface $S^- = S^-_{top} \cup S^-_{side}$ in $\overline{D_{\delta_1,\mathbb{C}}}$ where

$$S_{top}^{-} = \{ (\alpha_I, \xi) \in \mathcal{D}_{\delta_1, \mathbb{C}} \mid (\operatorname{Im}(\alpha_I), \operatorname{Im}(\xi)) = (-\delta_1, 0, \dots, 0) \}$$

and

$$S_{\text{side}}^{-} = \{ (\alpha_I, \xi) - (t\sqrt{-1}\delta_1, 0, \dots, 0) \mid (\alpha_I, \xi) \in \partial \mathcal{D}_{\delta_1}, t \in [0, 1] \}.$$

Then the same estimate as in the previous case proves that on $S^- \cup (D_H \setminus D_{\delta_1})$,

$$\operatorname{Im}\left(\mathcal{W}^{\epsilon_{I}}(\alpha_{I},\xi)-\sum_{i\in I}2\pi a_{i}\alpha_{i}-\sum_{s=1}^{c}4\pi b_{s}\xi_{s}\right)<\operatorname{Im}\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}})-\epsilon',$$

from which the result follows.

5.5 Estimate of the error term

The goal of this section is to estimate the error term in Proposition 3.7.

Proposition 5.10. Suppose $\{\alpha_j\}_{j\in J}$ are in $\{\pi - \epsilon, \pi + \epsilon\}$ for a sufficiently small $\epsilon > 0$. Then the error term in Proposition 3.7 is less than $O(e^{\frac{r}{4\pi}(\operatorname{Vol}(M_{L_{\theta}})-\epsilon')})$ for some $\epsilon' > 0$.

For the proof we need the following estimate, which first appeared in [15, Proposition 8.2] for $q = e^{\frac{\pi\sqrt{-1}}{r}}$, and for the root $q = e^{\frac{2\pi\sqrt{-1}}{r}}$ in [11, Proposition 4.1].

Lemma 5.11. For any integer 0 < n < r and at $q = e^{\frac{2\pi\sqrt{-1}}{r}}$,

$$\log |\{n\}!| = -\frac{r}{2\pi} \Lambda\left(\frac{2n\pi}{r}\right) + O\left(\log(r)\right).$$

Proof of Proposition 5.10. For a fixed $\alpha_J = (\alpha_j)_{j \in J}$, let

$$M_{\alpha_J} = \max\left\{\sum_{s=1}^{c} 2V(\alpha_{s_1}, \dots, \alpha_{s_6}, \xi_s) \mid (\alpha_I, \xi) \in \partial \mathbf{D}_{\mathbf{H}} \cup (\mathbf{D}_{\mathbf{A}} \setminus \mathbf{D}_{\mathbf{H}})\right\}$$

where V is as defined in (4.3). We note that if $\beta_i = \alpha_j = \pi$ for all $i \in I$ and $j \in J$, the $D_H = D_A$ and $\partial D_H \cup (D_A \setminus D_H) = \partial D_H$. Then By Lemma 5.6,

$$M_{\pi,\ldots,\pi} < 2cv_8;$$

and by continuity, if ϵ is sufficiently small, then

$$M_{\alpha_J} < \operatorname{Vol}(M_{L_{\theta}})$$

for all $\{\alpha_j\}_{j\in J}$ in $\{\pi - \epsilon, \pi + \epsilon\}$.

Now by Lemma 5.11 and the continuity, for $\epsilon' = \frac{\operatorname{Vol}(M_{L_{\theta}}) - M_{\alpha_J}}{2}$, we can choose a sufficiently small $\delta > 0$ so that if $\left(\frac{2\pi \mathbf{m}_I}{r}, \frac{2\pi \mathbf{k}}{r}\right) \notin \mathbf{D}_{\mathrm{H}}^{\delta}$, then

$$\left|g_r^{\epsilon_I}(\mathbf{m}_I, \mathbf{k})\right| < O\left(e^{\frac{r}{4\pi}(M_{\alpha_J} + \epsilon')}\right) = O\left(e^{\frac{r}{4\pi}(\operatorname{Vol}(M_{L_\theta}) - \epsilon')}\right).$$

Let ψ be the bump function supported on (D_H, D_H^{δ}) . Then the error term in Proposition 3.7 is less than $O(e^{\frac{r}{4\pi}(\text{Vol}(M_{L_{\theta}})-\epsilon')})$.

5.6 Proof of Theorem 3.1

Proof of Theorem 3.1. Let $\epsilon > 0$ be sufficiently small so that the conditions of Propositions 5.5, 5.9 and 5.10 and of Corollary 5.7 are satisfied, and suppose $\{\beta_i\}_{i \in I}$ and $\{\alpha_j\}_{j \in J}$ are all in $(\pi - \epsilon, \pi + \epsilon)$. By Propositions 3.6, 3.7, 5.5, 5.9 and 5.10, we have

$$\operatorname{RT}_{r}(M, L, (\mathbf{n}_{I}, \mathbf{m}_{J})) = \kappa_{r} \Big(\sum_{\epsilon_{I} \in \{1, -1\}^{I}} \widehat{f_{r}^{\epsilon_{I}}}(0, \dots, 0) \Big) \Big(1 + O\left(e^{\frac{r}{2\pi}(-\epsilon')}\right) \Big)$$
$$= \kappa_{r} \Big(\sum_{\epsilon_{I} \in \{1, -1\}^{I}} \frac{C^{\epsilon_{I}}(z^{\epsilon_{I}})}{\sqrt{-\det\operatorname{Hess}}\mathcal{W}^{\epsilon_{I}}(z^{\epsilon_{I}})} \Big) e^{\frac{r}{4\pi\sqrt{-1}} \Big(2c\pi^{2} + \sqrt{-1}\left(\operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1}\operatorname{CS}(M_{L_{\theta}})\right)\Big)} \Big(1 + O\left(\frac{1}{r}\right) \Big).$$

By Proposition 3.5, we have

$$\lim_{r \to \infty} \frac{4\pi}{r} \log \kappa_r = \left(\sigma(L' \cap L_I) - \sum_{i \in I} q_i - \sum_{i=1}^n p_i - 2|I|\right) \sqrt{-1}\pi^2;$$

and by Corollary 5.7, we have

$$\sum_{\epsilon_I \in \{1,-1\}^I} \frac{C^{\epsilon_I}(z^{\epsilon_I})}{\sqrt{-\det \operatorname{Hess} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I})}} \neq 0,$$

and hence

$$\lim_{r \to \infty} \frac{4\pi}{r} \log \Big(\sum_{\epsilon_I \in \{1, -1\}^I} \frac{C^{\epsilon_I}(z^{\epsilon_I})}{\sqrt{-\det \operatorname{Hess} \mathcal{W}^{\epsilon_I}(z^{\epsilon_I})}} \Big) = 0.$$

Therefore,

$$\lim_{r \to \infty} \frac{4\pi}{r} \log \operatorname{RT}_r(M, L, (\mathbf{n}_I, \mathbf{m}_J))$$

$$= \left(\sigma(L' \cap L_I) - \sum_{i \in I} q_i - \sum_{i=1}^n p_i - 2|I| \right) \sqrt{-1} \pi^2 - 2c\sqrt{-1}\pi^2 + \operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(M_{L_{\theta}})$$

$$= \operatorname{Vol}(M_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(M_{L_{\theta}}) \quad \text{mod } \sqrt{-1}\pi^2 \mathbb{Z},$$

which completes the proof.

6 Some concrete examples

The goal of this section is to show that several important families of links have complements homeomorphic to fundamental shadow link complements $M_c \ L_{FSL}$, and are obtained from (M_c, L_{FSL}) by doing a change of pair operation, including the twisted octahedral fully augmented links considered by Purcell [31] and van der Veen [41], and the family \mathcal{U} considered by Kumar [22, Theorem 4.1]. As a consequence, Conjecture 1.1, the Volume Conjecture for the Turaev-Viro invariants [6] and the Generalized Volume Conjecture [27, 16] hold, and the answer to [12, Question 1.7] is positive, for these families of examples. See Theorems 6.1, 6.3 and 6.5. It is worth mentioning that both the family of the twisted octahedral fully augmented links and the family \mathcal{U} are universal families in the sense that every link in S^3 is a sublink of a member of these families.

We first look at the twisted octahedral fully augmented links. Following the construction in [41], we start with a trivalent graph T that is homomorphic to the 1-skeleton of a Euclidean tetrahedron as shown on the left of Figure 5, and apply a sequence of the triangle move as show on the right of Figure 5 to get a trivalent graph G.



Figure 5

Then we color some edges of G by red in a way that each vertex is adjacent to exactly one red edge. Such a coloring of the red edges always exists because for the initial graph T we can color any pair of the opposite edges by red, and then for each triangle move, we color the new edge opposite to the red edge in the old graph by red. See Figure 6.



Figure 6

For each choice of the red edges of G, we can construct a link in S^3 as follows. As shown in Figure 7, we first circulate each red edge by a trivial loop (the belt). Then we replace each red edge by a pair of



Figure 7

arcs parallel to it and connect the ends of the two arcs respectively to the edges of G that are originally adjacent to the red edge. Finally, we do a certain number of (possibly half) twists to each pair of parallel arcs. In this way, we get a twisted octahedral fully augmented link in S^3 . See Figure 8 for a concrete example starting from the trivalent graph T.



Figure 8

The twisted octahedral fully augmented links were also described in [31] using the dual nerve of the graph. Namely, we start with the graph T and consider its dual nerve, which is a graph T^* homeomorphic to T with triangular faces. After doing a sequence of the central subdivisions of the faces, we get a graph G^* with triangular faces. Then we choose a collection of red edges such that each triangular face contains exactly one red edge. Finally we consider the dual graph G of G^* and color the dual edge of the red edges of G^* by red, and change the red as shown in Figure 7. In this way, we obtain a link in S^3 . Since the triangle move is dual to the central subdivision (see Figure 9), the links constructed in this way are exactly the twisted octahedral fully augmented links.



Figure 9

In [41], van der Veen prove that the complement of a twisted octahedral fully augmented link is homeomorphic to some fundamental shadow link complement. Together with the result of [4], we have

Theorem 6.1 ([41, 4]). Suppose L is a framed twisted octahedral fully augmented link in S^3 . Then as r varies over all odd integers,

$$\lim_{r \to \infty} \frac{2\pi}{r} \log \mathrm{TV}_r(S^3 \backslash L) = \mathrm{Vol}(S^3 \backslash L).$$

The main observation of this section is the following Proposition 6.2, which is a refinement of the result of [41].

Proposition 6.2. Let L be a framed twisted octahedral fully augmented link. Then (S^3, L) is obtained from (M_c, L_{FSL}) by doing a change of pair operation, where $M_c = \#^{c+1}S^2 \times S^1$ for some positive integer c and $L_{FSL} \subset M_c$ is a fundamental shadow link. As a consequence of Proposition 1.3, $S^3 \setminus L$ is homeomorphic to $M_c \setminus L_{FSL}$.

Proof. Suppose $L = L_1 \cup \cdots \cup L_n$. Let I be the subset of $\{1, \ldots, n\}$ such that $\{L_i\}_{i \in I}$ are the belt components of L, and let $L_I = \bigcup_{i \in I} L_i$. Let $J = \{1, \ldots, n\} \setminus I$, and let $L_J = \bigcup_{j \in J} L_j$. Recall that each L_i corresponds to an red edge of the graph G which splits into two parallel arcs with a number of twits. Later we will prove the result in two steps. In Step 1 we consider the case that there is no twist or only a half-twists for each $i \in I$. In Step 2 we consider the general case.

We first observe that L_J lies in a tubular neighborhood of the 1-skeleton of a truncated polyhedron P_G in S^3 obtained by gluing truncated tetrahedra together along the triangles of truncation. Indeed, we can regard the initial trivalent graph T as the set of edges of a truncated tetrahedron (see Figure 10), and regard a triangular move as attaching another truncated tetrahedron along the triangle of truncation (see Figure 11).



Figure 10: In a truncated polyhedron, we only consider the intersection of two faces as an edge, and do not consider the intersection of a face and a triangle of truncation as an edge.



Figure 11

In this way, we obtain a truncated polyhedron P_G in $\mathbb{R}^3 \subset S^3$, and the edges of the graph G correspond to the edges of P_G . As shown in Figure 12, the two parallel arcs from splitting the red edge can be drawn in a tubular neighborhood of the red arc. See Figure 13 for a concrete example.



Figure 12



Figure 13

Step 1. Suppose in the construction of L there is no twist or only a half twist for each pair of arcs from splitting the red edges. For each $i \in I$, we let L'_i be L_i with the 0-framing which is possibly different from the original framing of L_i , and let L^*_i be the framed trivial loop around L_i with the 0-framing. Let $L'_I = \bigcup_{i \in I} L'_i$. We first claim that by doing the change of pair operation $T(L'_I; L^*_I)$ to the pair (S^3, L) we get the pair $(M_{|I|}, L_{FSL})$, where $M_{|I|} = \#^{|I|}(S^2 \times S^1)$ and L_{FSL} is a fundamental shadow link in $M_{|I|}$. See Figure 14.



Figure 14

Indeed, let $N_I = \bigcup_{i \in I} N(L_i)$ be the union of the tubular neighborhoods of the belts $\{L_i\}_{i \in I}$. Then $S^3 \setminus N_I$ is a handlebody $H_{|I|}$ of genus $|I|, L_I^* \cup L_J$ is a link in $H_{|I|}$, and doing 0-surgeries along $\{L_i\}_{i \in I}$ is the same as taking the double of $H_{|I|}$, which is homeomorphic to $M_{|I|}$. On the other hand, the handlebody $H_{|I|}$ can also be considered as obtained from the polyhedron P_G by gluing the two triangles of truncation at the end of each red edge via the orientation reversing affine homeomorphism identifying the two end points of the red edge.

If there is no twist for all pairs of arcs from splitting the red edges, then the link $L_I^* \cup L_J$ corresponds to the link L_{FSL} consisting of the union of the edges of P_G . From the construction in Section 2.4, L_{FSL} in the double of $H_{|I|}$ is a fundamental shadow link in $M_{|I|}$. See Figure 15 for a concrete example.



Figure 15

If there is a half twist on the pair of arcs from splitting red edge e_i circulated by L_i , for some $i \in I$, then we glue the two triangles of truncation at the end of e_i via the orientation preserving affine home-

omorphism identifying the two end points e_i . In this way, we obtain a non-orientable handlebody $H'_{|I|}$ whose orientable double is homeomorphic to $M_{|I|}$, and the link $L_I^* \cup L_J$ corresponds to the link L_{FSL} consisting of the union of the edges of P_G , which is a fundamental shadow link in $M_{|I|}$.

For each $i \in I$, we let L_i^{**} be the framed trivial loop around L_i^* with the same framing as that of L_i . Then we claim that (S^3, L) can be obtained from $(M_{|I|}, L_{FSL})$ by doing the change of pair operation $T(L_I^*, L_I^{**})$, proving the result in Case 1. See Figure 16.



Figure 16

This could be seen by the following the second Kirby Moves (handle slides) which do not change the pair. For each $i \in I$, we first slide L_i^{**} over L_i' as in Figure 17 to get a framed trivial loop isotopic to L_i . Suppose L_j and $L_{j'}$ (with j possibly the same as j') are the components of L circulated by L_i . Then we



Figure 17

slide L_j and $L_{j'}$ over L_i^* as in Figure 18 so that $L'_i \cup L_i^*$ is a Hopf link with 0-framings unlinked with the rest of L. Doing these operations for each $i \in I$, we get the original link L in the 3-manifold obtained from S^3 by doing a surgery along the disjoint union of |I| Hopf links with 0-framings, which is still S^3 .



Figure 18

Step 2. Suppose in the construction of L, the pair of arcs from splitting the red edge circulated by L_i is twisted p_i times or p_i and a half times. Let $(M_{|I|}, L_{FSL})$ be the fundamental shadow link constructed in Step 1. For each $i \in I$, let $L_i^{*'}$ be L_i^* with the $(-p_i)$ -framing, and let $L_I^{*'} = \bigcup_{i \in I} L_i^{*'}$. Still let L_i^{**} be the

framed trivial loop around L_i^* with the same framing as that of L_i . We claim that (S^3, L) can be obtained from $(M_{|I|}, L_{\text{FSL}})$ by doing the change of pair operation $T(L_I^{*\prime}, L_I^{**})$, proving the result in Case 2.

Similar to Case 1, we first slide L_i^{**} over L_i' as in Figure 19 to get a framed trivial loop isotopic to L_i . By e.g. [35, p. 273], since $L_i^{*'}$ is a trivial loop around L_i' and



Figure 19

$$\frac{1}{p_i} = 0 - \frac{1}{-p_i}$$

doing the surgery along the union of L'_i and L''_i respectively with framings 0 and $(-p_i)$ is equivalent to doing a $\frac{1}{p_i}$ -surgery along a loop isotopic to L_i , which is the same as doing p_i full twists along the strip circulated by L_i . See Figure 20. This completes the proof.



Figure 20

By Theorem 3.1, Propositions 6.2 and the relationship between the Resherikhin-Turaev invarints and the colored Jones polynomials that

$$\mathrm{RT}_r(S^3, L, \mathbf{m}) = \mu_r \mathrm{J}_{\mathbf{m}, L}(q^2),$$

we have

Theorem 6.3. Suppose L is a framed twisted octahedral fully augmented with n components. For a sequence $\mathbf{m}^{(r)} = (m_1^{(r)}, \dots, m_n^{(r)})$ of colorings of L, let

$$\theta_k = \left| 2\pi - \lim_{r \to \infty} \frac{4\pi m_k^{(r)}}{r} \right|$$

and let $\theta = (\theta_1, \dots, \theta_n)$. If all the θ_i 's are sufficiently small, then

$$\lim_{r \to \infty} \frac{4\pi}{r} \log \mathcal{J}_{L,\mathbf{m}^{(r)}}\left(e^{\frac{4\pi i}{r}}\right) = \operatorname{Vol}(S^3_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(S^3_{L_{\theta}}) \qquad \text{mod } \sqrt{-1}\pi^2 \mathbb{Z},$$

where r varies over all odd integers.

Next, we consider the universal family \mathcal{U} of links in S^3 considered by Kumar [22] whose complements are homeomorphic to some fundamental shadow link complements. Together with the result of [4], he proved that Chen-Yang's Volume Conjecture for the Turaev-Viro invariants is true for the complements of these families of links. The way he found the fundamental shadow links is essentially by doing a change of pair operation along the belt components of the links. Then by the same argument in Step 1 of the proof of Proposition 6.2, we have

Proposition 6.4. Let L be a framed link in U of [22]. Then (S^3, L) is obtained from (M_c, L_{FSL}) by doing a change of pair operation.

As a consequence of Theorem 3.1 and Proposition 6.4, we have

Theorem 6.5. Suppose L is a framed link of n components which is a member of U of [22]. For a sequence $\mathbf{m}^{(r)} = (m_1^{(r)}, \ldots, m_n^{(r)})$ of colorings of L, let

$$\theta_k = \left| 2\pi - \lim_{r \to \infty} \frac{4\pi m_k^{(r)}}{r} \right|$$

and let $\theta = (\theta_1, \dots, \theta_n)$. If all the θ_i 's are sufficiently small, then

$$\lim_{r \to \infty} \frac{4\pi}{r} \log \mathcal{J}_{L,\mathbf{m}^{(r)}}\left(e^{\frac{4\pi i}{r}}\right) = \operatorname{Vol}(S^3_{L_{\theta}}) + \sqrt{-1} \operatorname{CS}(S^3_{L_{\theta}}) \qquad \text{mod } \sqrt{-1}\pi^2 \mathbb{Z},$$

where r varies over all odd integers.

Remark 6.6. The proof of Proposition 6.2 essentially provides an algorithm of constructing a fundamental shadow link from a given twisted octahedral fully augmented link or a given element of \mathcal{U} . In [41], van der Veen for each link L in S^3 provided an algorithm of constructing a twisted octahedral fully augmented link L' such that $S^3 \ L$ is obtained from $S^3 \ L'$ by $\frac{1}{0}$ -filling suitable boundary components. In [22], Kumar for each link L in S^3 considered as the closure of a braid provided an algorithm of constructing a link L' in \mathcal{U} such that $S^3 \ L$ is obtained from $S^3 \ L'$ by $\frac{1}{0}$ -filling suitable boundary components. Therefore, together with Propositions 6.2 and 6.4, we for each L in S^3 have two algorithms of constructing a fundamental shadow link L_{FSL} in M_c such that $S^3 \ L$ is obtained from $M_c \ L_{FSL}$ by filling suitable boundary components. It is an interesting question to know whether the universal families of the twisted octahedral fully augmented links and \mathcal{U} are actually the same family.

A A proof of Proposition 5.1

The goal of this appendix is to prove Proposition 5.1. We need the following two Lemmas whose proof are included at the end of the appendix.

Lemma A.1. For any $\epsilon > 0$, there exists a $\delta > 0$ such that (1)

$$\int_{-\epsilon}^{\epsilon} e^{-rz^2} dz = \sqrt{\frac{\pi}{r}} + O(e^{-\delta r}),$$

and

(2)

$$\int_{-\epsilon}^{\epsilon} z^2 e^{-rz^2} dz = \frac{1}{2} \sqrt{\frac{\pi}{r^3}} + O(e^{-\delta r}).$$

Lemma A.2. (Complex Morse Lemma) Let $D_{\mathbf{z}}$ be a region in \mathbb{C}^n , let $D_{\mathbf{a}}$ be a region in \mathbb{R}^k , and let $f: D_{\mathbf{z}} \times D_{\mathbf{a}} \to \mathbb{C}$ be a complex valued function that is holomorphic in $\mathbf{z} \in D_{\mathbf{z}}$ and smooth in $\mathbf{a} \in D_{\mathbf{a}}$. For $\mathbf{a} \in D_{\mathbf{a}}$, let $f^{\mathbf{a}}: D_{\mathbf{z}} \to \mathbb{C}$ be the function defined by $f^{\mathbf{a}}(\mathbf{z}) = f(\mathbf{z}, \mathbf{a})$. Suppose for each $\mathbf{a} \in D_{\mathbf{a}}$, $f^{\mathbf{a}}$ has a non-degenerate critical point $c_{\mathbf{a}}$ which smoothly depends on \mathbf{a} . Then for each $\mathbf{a}_0 \in D_{\mathbf{a}}$, there exists an open set $V \subset \mathbb{C}^n$ containing $\mathbf{0}$, an open set $A \subset D_{\mathbf{a}}$ containing \mathbf{a}_0 , and a smooth function $\psi: V \times A \to D_{\mathbf{z}}$ such that, if we denote $\psi^{\mathbf{a}}(\mathbf{Z}) = \psi(\mathbf{Z}, \mathbf{a})$, then for each $\mathbf{a} \in D_{\mathbf{a}}, \mathbf{z} = \psi^{\mathbf{a}}(\mathbf{Z})$ is a holomorphic change of variable on V such that

$$\psi^{\mathbf{a}}(\mathbf{0}) = \mathbf{c}_{\mathbf{a}},$$
$$f^{\mathbf{a}}(\psi^{\mathbf{a}}(\mathbf{Z})) = f^{\mathbf{a}}(\mathbf{c}_{\mathbf{a}}) - Z_1^2 - \dots - Z_n^2,$$

and

$$\det\left(D(\psi^{\mathbf{a}})(\mathbf{0})\right) = \frac{2^{\frac{n}{2}}}{\sqrt{-\det\operatorname{Hess}(f^{\mathbf{a}})(\mathbf{c}_{\mathbf{a}})}}$$

Proof of Proposition 5.1. We write $\mathbf{z} = (z_1, \ldots, z_n) \in \mathbb{C}^n$, $\mathbf{Z} = (Z_1, \ldots, Z_n) \in \mathbb{C}^n$, $\mathbf{W} = (W_1, \ldots, W_n) \in \mathbb{C}^n$, $d\mathbf{z} = dz_1 \ldots dz_n$ and $\mathbf{0} = (0, \ldots, 0) \in \mathbb{C}^n$.

We first consider the special case $\mathbf{c}_r = \mathbf{0}, S_r = [-\epsilon, \epsilon]^n \subset \mathbb{R}^n \subset \mathbb{C}^n$, and

$$f^{\mathbf{a}_r}(\mathbf{z}) = -\sum_{i=1}^n z_i^2$$

for each r. In this case, let

$$\sigma_r^{\mathbf{a}_r}(\mathbf{z}) = \upsilon_r(\mathbf{z}, \mathbf{a}_r) \int_0^1 e^{\frac{\upsilon_r(\mathbf{z}, \mathbf{a}_r)}{r}s} ds.$$

Then we can write

$$e^{\frac{\upsilon_r(\mathbf{z},\mathbf{a}_r)}{r}} = 1 + \frac{\sigma_r^{\mathbf{a}_r}(\mathbf{z})}{r},$$

and

$$g^{\mathbf{a}_r}(\mathbf{z})e^{rf_r^{\mathbf{a}_r}(\mathbf{z})} = g^{\mathbf{a}_r}(\mathbf{z})e^{rf^{\mathbf{a}_r}(\mathbf{z})} + \frac{1}{r}g^{\mathbf{a}_r}(\mathbf{z})\sigma_r^{\mathbf{a}_r}(\mathbf{z})e^{rf^{\mathbf{a}_r}(\mathbf{z})}.$$
(A.1)

Since $|v_r(\mathbf{z}, \mathbf{a}_r)| < M$ for some M > 0 independent of r,

$$|\sigma_r^{\mathbf{a}_r}(\mathbf{z})| < M \int_0^1 e^{\frac{M}{r}s} ds = M\left(\frac{e^{\frac{M}{r}} - 1}{\frac{M}{r}}\right) < 2M.$$

Since $\{\alpha_r\}$ is convergent and g is smooth in a, if M is big enough, then $|g^{\mathbf{a}_r}(\mathbf{z})| < M$ for all $z \in S_r = [-\epsilon, \epsilon]^n$ for r large enough. By Lemma A.1 (1), we have

$$\left| \int_{S_r} \frac{1}{r} g^{\mathbf{a}_r}(\mathbf{z}) \sigma_r(\mathbf{z}) e^{rf^{\mathbf{a}_r}(\mathbf{z})} \right| < \frac{2M^2}{r} \int_{S_r} e^{rf^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z}$$

$$= \frac{2M^2}{r} \left(\frac{\pi}{r}\right)^{\frac{n}{2}} + O(e^{-\delta r}) = O\left(\frac{1}{\sqrt{r^{n+2}}}\right).$$
(A.2)

By the Taylor Theorem, we have

$$g^{\mathbf{a}_r}(\mathbf{z}) = g^{\mathbf{a}_r}(\mathbf{0}) + \sum_{i=1}^n \frac{\partial}{\partial z_i} g^{\mathbf{a}_r}(\mathbf{0}) z_i + \sum_{i \neq j} \frac{\partial^2}{\partial z_i z_j} g^{\mathbf{a}_r}(\mathbf{0}) z_i z_j + \sum_{i=1}^n h_i^{\mathbf{a}_r}(\mathbf{z}) z_i^2$$

for some holomorphic functions $h_i^{\mathbf{a}_r}(\mathbf{z}), i = 1, \dots, n$, on D. Then by Lemma A.1 (1), we have

$$\int_{S_r} g^{\mathbf{a}_r}(\mathbf{0}) e^{rf^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z} = g^{\mathbf{a}_r}(\mathbf{0}) \left(\frac{\pi}{r}\right)^{\frac{n}{2}} + O\left(\frac{1}{r}\right)$$
(A.3)

Since each $z_i e^{-rz_i^2}$ is odd, we have

$$\int_{-\epsilon}^{\epsilon} z_i e^{-rz_i^2} dz_i = 0.$$

As a consequence, we have

$$\int_{-\epsilon}^{\epsilon} \frac{\partial}{\partial z_i} g^{\mathbf{a}_r}(\mathbf{0}) z_i e^{rf^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z} = 0, \qquad (A.4)$$

for each *i*, and

$$\int_{-\epsilon}^{\epsilon} \frac{\partial^2}{\partial z_i z_j} g^{\mathbf{a}_r}(\mathbf{0}) z_i z_j e^{rf^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z} = 0$$
(A.5)

for each $i \neq j$. Since $\{\mathbf{a}_r\}$ is convergent and h_i^{α} is smooth in \mathbf{a} , if M is big enough, then $|h_{\mathbf{a}_r,i}(\mathbf{z})| < M$ for all $\mathbf{z} \in S_r$, $i \in \{1, \ldots, n\}$ for r large enough. By Lemma A.1 we have for each $i \in \{1, \ldots, n\}$

$$\left|\int_{S_r} h_i^{\mathbf{a}_r}(\mathbf{z}) z_i^2 e^{rf^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z}\right| < M \left(\int_{-\epsilon}^{\epsilon} z_i^2 e^{-rz_i^2} dz_i\right) \prod_{j \neq i} \left(\int_{-\epsilon}^{\epsilon} e^{-rz_j^2} dz_j\right) = O\left(\frac{1}{\sqrt{r^{n+2}}}\right).$$
(A.6)

Putting (A.3), (A.4), (A.5) and (A.6) together, we have the result for this special case.

For the general case, by assumption (6) of Proposition 5.1, for a sufficiently close to \mathbf{a}_0 , $f^{\mathbf{a}}$ has a unique non-degenerate critical point $\mathbf{c}_{\mathbf{a}}$ in a sufficiently small neighborhood of \mathbf{c}_0 . Then we can apply Lemma A.2 to the function f and \mathbf{a}_0 . Let V, A and ψ respectively be the two open sets and the change of variable function as described in Lemma A.2. For r sufficiently large, let

$$U_r = \psi^{\mathbf{a}_r} \Big(\prod_{i=1}^n \left\{ Z_i \in \mathbb{C} \mid -\epsilon < \operatorname{Re}(Z_i) < \epsilon, -\epsilon < \operatorname{Im}(Z_i) < \epsilon \right\} \Big)$$

for some sufficiently small $\epsilon > 0$. Let $\operatorname{Vol}(S_r \setminus U_r)$ be the Euclidean volume of $S_r \setminus U_r$. By the compactness of $S_r \setminus U_r$ and by assumptions (2), (4) and (5) of Proposition 5.1, there exist constants M > 0 and $\delta > 0$ independent of r such that $|g^{\mathbf{a}_r}(\mathbf{z})| < M$, $\operatorname{Vol}(S_r \setminus U_r) < M$ and

$$\operatorname{Re} f_r^{\mathbf{a}_r}(\mathbf{z}) < \operatorname{Re} f^{\mathbf{a}_r}(\mathbf{c}_r) - \delta$$
 (A.7)

on $S_r \smallsetminus U$ for r large enough. Then

$$\left|\int_{S_r \smallsetminus U} g^{\mathbf{a}_r}(\mathbf{z}) e^{r f_r^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z}\right| < M^2 \left(e^{r(\operatorname{Re}f^{\mathbf{a}_r}(\mathbf{z})(\mathbf{c}_r) - \delta)} \right) = O\left(e^{r(\operatorname{Re}f^{\mathbf{a}_r}(\mathbf{z})(\mathbf{c}_r) - \delta)} \right).$$
(A.8)

In Figure 21 below, the shaded region is where $\operatorname{Re}(-\sum_{i=1}^{n} Z_{i}^{2}) < 0$. For each \mathbf{a}_{r} , in $\overline{(\psi^{\mathbf{a}_{r}})^{-1}(U_{r})}$ there is a homotopy H_{r} from $\overline{(\psi^{\mathbf{a}_{r}})^{-1}(S_{r} \cap U_{r})}$ to $[-\epsilon, \epsilon]^{n} \subset \mathbb{R}^{n}$ defined by "pushing everything down" to the real part. Let $S'_{r} = H_{r}(\partial(\psi^{\mathbf{a}_{r}})^{-1}(S_{r} \cap U_{r}) \times [0, 1])$. Then $\overline{(\psi^{\mathbf{a}_{r}})^{-1}(S_{r} \cap U_{r})}$ is homotopic to $S'_{r} \cup [-\epsilon, \epsilon]^{n}$. Then by analyticity,

$$\int_{S_{r}\cap U} g^{\mathbf{a}_{r}}(\mathbf{z}) e^{rf_{r}^{\mathbf{a}_{r}}(\mathbf{z})} d\mathbf{z}$$

$$= \int_{(\psi^{\mathbf{a}_{r}})^{-1}(S_{r}\cap U)} g^{\mathbf{a}_{r}}(\psi^{\mathbf{a}_{r}}(\mathbf{Z})) \det \mathcal{D}(\psi^{\mathbf{a}_{r}}(\mathbf{Z})) e^{rf_{r}^{\mathbf{a}_{r}}(\psi^{\mathbf{a}_{r}}(\mathbf{Z}))} d\mathbf{Z}$$

$$= \int_{S_{r}'} g^{\mathbf{a}_{r}}(\psi^{\mathbf{a}_{r}}(\mathbf{Z})) \det \mathcal{D}(\psi^{\mathbf{a}_{r}}(\mathbf{Z})) e^{rf_{r}^{\mathbf{a}_{r}}(\psi^{\mathbf{a}_{r}}(\mathbf{Z}))} d\mathbf{Z} + \int_{[-\epsilon,\epsilon]^{n}} g^{\mathbf{a}_{r}}(\psi(\mathbf{Z})) \det \mathcal{D}(\psi^{\mathbf{a}_{r}}(\mathbf{Z})) e^{rf_{r}^{\mathbf{a}_{r}}(\psi^{\mathbf{a}_{r}}(\mathbf{Z}))} d\mathbf{Z}$$
(A.9)





Since $\psi^{\mathbf{a}_r}(S') \subset S_r \smallsetminus U_r$, by (A.7)

$$\int_{S'_r} g^{\mathbf{a}_r}(\psi^{\mathbf{a}_r}(\mathbf{Z})) \det \mathcal{D}(\psi^{\mathbf{a}_r}(\mathbf{Z})) e^{rf_r^{\mathbf{a}_r}(\psi^{\mathbf{a}_r}(\mathbf{Z}))} d\mathbf{Z} = \int_{\psi^{\mathbf{a}_r}(S'_r)} g^{\mathbf{a}_r}(\mathbf{z}) e^{rf_r^{\mathbf{a}_r}(\mathbf{z})} d\mathbf{z} = O\left(e^{r(\operatorname{Re}f^{\mathbf{a}_r}(\mathbf{c}_r) - \delta)}\right);$$
(A.10)

and by the special case

$$\begin{split} &\int_{[-\epsilon,\epsilon]^n} g^{\mathbf{a}_r}(\psi^{\mathbf{a}_r}(\mathbf{Z})) \det \mathcal{D}(\psi^{\mathbf{a}_r}(\mathbf{Z})) e^{rf_r^{\mathbf{a}_r}(\psi^{\mathbf{a}_r}(\mathbf{Z}))} d\mathbf{Z} \\ &= e^{rf^{\mathbf{a}_r}(\mathbf{c}_r)} \int_{[-\epsilon,\epsilon]^n} g^{\mathbf{a}_r}(\psi^{\mathbf{a}_r}(\mathbf{Z})) \det \mathcal{D}(\psi^{\mathbf{a}_r}(\mathbf{Z})) e^{r\left(-\sum_{i=1}^n Z_i^2 + \frac{v_r(\mathbf{z},\mathbf{a}_r)}{r^2}\right)} d\mathbf{Z} \\ &= e^{rf^{\mathbf{a}_r}(\mathbf{c}_r)} g^{\mathbf{a}_r}(\psi^{\mathbf{a}_r}(\mathbf{0})) \det \mathcal{D}(\psi^{\mathbf{a}_r}(\mathbf{0})) \left(\frac{\pi}{r}\right)^{\frac{n}{2}} \left(1 + O\left(\frac{1}{r}\right)\right) \\ &= \left(\frac{2\pi}{r}\right)^{\frac{n}{2}} \frac{g^{\mathbf{a}_r}(\mathbf{c}_r)}{\sqrt{-\det \operatorname{Hess}(f^{\mathbf{a}_r})(\mathbf{c}_r)}} e^{rf^{\mathbf{a}_r}(\mathbf{c}_r)} \left(1 + O\left(\frac{1}{r}\right)\right). \end{split}$$

Together with (A.8), (A.9) and (A.10), we have the result.

Proof of Lemma A.1. For (1), we have

$$\int_{-\epsilon}^{\epsilon} e^{-rz^2} dz = \int_{-\infty}^{\infty} e^{-rz^2} dz - \int_{-\infty}^{-\epsilon} e^{-rz^2} dz - \int_{\epsilon}^{\infty} e^{-rz^2} dz,$$

where the first term

$$\int_{-\infty}^{\infty} e^{-rz^2} dz = \sqrt{\frac{\pi}{r}}$$

is a Gaussian integral, and the other two terms

$$\int_{-\infty}^{-\epsilon} e^{-rz^2} dz = \int_{\epsilon}^{\infty} e^{-rz^2} dz \leqslant \int_{\epsilon}^{\infty} e^{-r\epsilon z} dz = \frac{e^{-r\epsilon^2}}{r\epsilon} = O(e^{-\delta r}).$$

For (2), by integration by parts, we have

$$\int_{-\epsilon}^{\epsilon} e^{-rz^2} dz = z e^{-rz^2} \Big|_{-\epsilon}^{\epsilon} + 2r \int_{-\epsilon}^{\epsilon} z^2 e^{-rz^2} dz,$$

hence by (1)

$$\int_{-\epsilon}^{\epsilon} z^2 e^{-rz^2} dz = \frac{1}{2r} \left(\int_{-\epsilon}^{\epsilon} e^{-rz^2} dz - 2\epsilon e^{-r\epsilon^2} \right) = \frac{1}{2} \sqrt{\frac{\pi}{r^3}} + O(e^{-\delta r}).$$

Proof of Lemma A.2. By doing the linear transformation $(\mathbf{z}, \mathbf{a}) \mapsto (\mathbf{z} + \mathbf{c}_{\mathbf{a}}, \mathbf{a})$, we may assume that that $\mathbf{c}_{\mathbf{a}} = \mathbf{0}$ for all $\mathbf{a} \in D_{\mathbf{a}}$. Then by the Taylor Theorem, for each $\mathbf{a} \in D_{\mathbf{a}}$ and $\mathbf{z} \in D_{\mathbf{z}}$, we can write

$$f^{\mathbf{a}}(\mathbf{z}) = f^{\mathbf{a}}(\mathbf{0}) + \sum_{i=1}^{n} z_i b_i^{\mathbf{a}}(\mathbf{z})$$

for some holomorphic functions $b_i^{\mathbf{a}}$, i = 1, ..., n. Since **0** is a critical point of $f^{\mathbf{a}}$, we have

$$b_i^{\mathbf{a}}(\mathbf{0}) = \frac{\partial}{\partial z_i} f^{\mathbf{a}}(\mathbf{0}) = 0.$$

As a result, by Taylor theorem again, we can write

$$f^{\mathbf{a}}(\mathbf{z}) = f^{\mathbf{a}}(\mathbf{0}) + \sum_{i=1}^{n} z_i b_i^{\mathbf{a}}(\mathbf{z}) = f^{\mathbf{a}}(\mathbf{0}) + \sum_{i,j=1}^{n} z_i z_j h_{ij}^{\mathbf{a}}(\mathbf{z})$$

for some holomorphic functions $h_{ij}^{\mathbf{a}}, i, j = 1, \dots, n$. Since

$$\sum_{i,j=1}^{n} z_i z_j h_{ij}^{\mathbf{a}}(\mathbf{z}) = \sum_{i,j=1}^{n} z_i z_j \Big(\frac{h_{ij}^{\mathbf{a}}(\mathbf{z}) + h_{ji}^{\mathbf{a}}(\mathbf{z})}{2}\Big),$$

we may assume that $h_{ij}^{\mathbf{a}}$ is symmetric in *i* and *j*. Since **0** is a non-degenerate critical point of $f^{\mathbf{a}}$, and

$$\frac{\partial^2}{\partial z_i z_j} f^{\mathbf{a}}(\mathbf{0}) = 2h_{ij}^{\mathbf{a}}(\mathbf{0}),$$

we have $det(h_{ij}^{\mathbf{a}}(\mathbf{0})) \neq 0$.

Next, suppose for some m with $0 \leq m \leq n$, there exist an open set $V_m \subset \mathbb{C}^n$ containing $\mathbf{0}$, an open set $A_m \subset D_{\mathbf{a}}$ containing \mathbf{a}_0 , and a smooth function $\psi_m : V_m \times A_m \to \mathbb{C}^n$ such that, if we denote $\psi_m^{\mathbf{a}}(\mathbf{Z}) = \psi_m(\mathbf{Z}, \mathbf{a})$, then $\psi_m^{\mathbf{a}}$ gives a holomorphic change of variable with

$$f^{\mathbf{a}}(\psi_{m}^{\mathbf{a}}(\mathbf{Z})) = f^{\mathbf{a}}(\mathbf{0}) - Z_{1}^{2} - \dots - Z_{m-1}^{2} + \sum_{i,j=m}^{n} Z_{i}Z_{j}H_{m,ij}^{\mathbf{a}}(\mathbf{Z}),$$

where $H_{m,ij}^{\mathbf{a}}(\mathbf{Z})$ is holomorphic in \mathbf{Z} and symmetric in *i* and *j*. Based on this, we are going to find an open set V_{m+1} of \mathbb{C}^n containing $\mathbf{0}$, an open set $A_{m+1} \subset A_m$ containing \mathbf{a}_0 , and a smooth function $\psi_{m+1} : V_{m+1} \times A_{m+1} \to \mathbb{C}^n$ such that, if we denote $\psi_{m+1}^{\mathbf{a}}(\mathbf{Z}) = \psi_{m+1}(\mathbf{Z}, \mathbf{a})$, then $\psi_{m+1}^{\mathbf{a}}$ gives a holomorphic change of variable with

$$f^{\mathbf{a}}(\psi_{m+1}^{\mathbf{a}}(\mathbf{Z})) = f^{\mathbf{a}}(\mathbf{0}) - Z_1^2 - \dots - Z_m^2 + \sum_{i,j=m+1}^n Z_i Z_j H_{m+1,ij}^{\mathbf{a}}(\mathbf{Z})$$

for some holomorphic functions $H_{m+1,ij}^{\mathbf{a}}(\mathbf{Z})$ that are symmetric in *i* and *j*.

To do so, we by the Chain Rule have

$$\frac{\partial^2 f^{\mathbf{a}}}{\partial Z_i Z_j}(\psi_m^{\mathbf{a}}(\mathbf{0})) = (D\psi_m^{\mathbf{a}}(\mathbf{0}))^T \Big(\frac{\partial^2 f^{\mathbf{a}}}{\partial z_i z_j}(\mathbf{0})\Big) (D\psi_m^{\mathbf{a}}(\mathbf{0})),$$

where $D\psi_m^{\mathbf{a}}(\mathbf{0})$ is the Jacobian matrix of $\psi_m^{\mathbf{a}}$ at $\mathbf{0}$. Thus, we have

$$2^{m-1}\det(2H_{m,ij}^{\mathbf{a}}(\mathbf{0})) = \det\left(\frac{\partial^2 f^{\mathbf{a}}}{\partial Z_i Z_j}(\psi_m^{\mathbf{a}}(\mathbf{0}))\right) \neq 0,$$

implying that $(H_{m,ij}^{\mathbf{a}}(\mathbf{0}))$ is a $(n-m+1)\times(n-m+1)$ non-singular matrix. Therefore, there exists $k \ge m$ such that $H_{m,km}^{\mathbf{a}}(\mathbf{0}) \ne 0$. Reordering the variables if necessary, we may assume that $H_{m,mm}^{\mathbf{a}}(\mathbf{0}) \ne 0$. By continuity of $H_{m,mm}^{\mathbf{a}}(\mathbf{Z})$ in \mathbf{Z} and \mathbf{a} , there exists an open set $V'_m \subset V_m$ containing $\mathbf{0}$ and an open set $A'_m \subset A_m$ containing \mathbf{a}_0 such that $H^{\mathbf{a}}_{m,mm}(\mathbf{0}) \neq 0$ for all $(\mathbf{Z}, \mathbf{a}) \in V'_m \times A'_m$. Then we can let

$$\widetilde{H}_{m,ij}^{\mathbf{a}}(\mathbf{Z}) = \frac{H_{m,ij}^{\mathbf{a}}(\mathbf{Z})}{H_{m,mm}^{\mathbf{a}}(\mathbf{Z})}$$

and have

$$f^{\mathbf{a}}(\psi_{m}^{\mathbf{a}}(\mathbf{Z})) = f^{\mathbf{a}}(\mathbf{0}) - Z_{1}^{2} - \dots - Z_{m-1}^{2} + \sum_{i,j=m}^{n} Z_{i}Z_{j}H_{m,ij}^{\mathbf{a}}(\mathbf{Z})$$

$$= f^{\mathbf{a}}(\mathbf{0}) - Z_{1}^{2} - \dots - Z_{m-1}^{2} + H_{m,mm}^{\mathbf{a}}(\mathbf{Z}) \sum_{i,j=m}^{n} Z_{i}Z_{j}\widetilde{H}_{m,ij}^{\mathbf{a}}(\mathbf{Z})$$

$$= f^{\mathbf{a}}(\mathbf{0}) - Z_{1}^{2} - \dots - Z_{m-1}^{2} + H_{m,mm}^{\mathbf{a}}(\mathbf{Z}) \Big(Z_{m} + \sum_{j=m+1}^{n} Z_{j}\widetilde{H}_{m,mj}^{\mathbf{a}}(\mathbf{Z}) \Big)^{2}$$

$$- H_{m,mm}^{\mathbf{a}}(\mathbf{Z}) \Big[\Big(\sum_{j=m+1}^{n} Z_{j}\widetilde{H}_{m,mj}^{\mathbf{a}}(\mathbf{Z}) \Big)^{2} + \sum_{i,j=m+1}^{n} Z_{i}Z_{j}\widetilde{H}_{m,ij}^{\mathbf{a}}(\mathbf{Z}) \Big].$$

Define $\mathbf{W} = \mathbf{W}(\mathbf{Z})$ by

$$W_l = Z_l$$

for $l \neq m$, and

$$W_m = \sqrt{-H_{m,mm}^{\mathbf{a}}(\mathbf{Z})} \left(Z_m + \sum_{j=m+1}^n Z_j \widetilde{H}_{m,mj}^{\mathbf{a}}(\mathbf{Z}) \right)$$

We note that

$$\frac{\partial W_l}{\partial Z_k}(\mathbf{0}) = \delta_{l,k}$$

for $l \neq m$, and

$$\frac{\partial W_m}{\partial Z_k}(\mathbf{0}) = \sqrt{-H_{mm}^{\mathbf{a}}(\mathbf{0})} \Big(\delta_{m,k} + \sum_{j=m+1}^n \delta_{j,k} \widetilde{H}_{m,mj}^{\mathbf{a}}(\mathbf{0})\Big).$$

Then the Jacobian matrix $DW(\mathbf{0})$ is an upper triangular matrix with the (m, m)-th entry $\sqrt{-H_{m,mm}^{\mathbf{a}}(\mathbf{0})} \neq 0$ 0 and all the other diagonal entries 1, hence $\det DW(\mathbf{0}) \neq 0$. Now consider the map $G: V'_m \times A'_m \to \mathbb{C}^n \times \mathbb{R}^k$ defined by

$$G(\mathbf{Z}, \mathbf{a}) = (\mathbf{W}(\mathbf{Z}), \mathbf{a})$$

Then the Jacobian matrix $DG(\mathbf{0}, \mathbf{a}_0)$ is of the form

$$DG(\mathbf{0}, \mathbf{a}_0) = \begin{pmatrix} DW(\mathbf{0}) & * \\ 0 & I_k \end{pmatrix},$$

where I_k is the $k \times k$ identity matrix. Moreover, $\det(DG(\mathbf{0}, \mathbf{a}_0)) = \det(DW(\mathbf{0})) \neq 0$. Thus, by the Inverse Function Theorem, there exists an open set $V''_m \subset V'_m$ and an open subset with compact closure $A_{m+1} \subset A'_m$ containing \mathbf{a}_0 such that $G: V''_m \times A_{m+1} \to \mathbb{C}^n \times A_m$ is a diffeomorphism to its image. By slightly shrinking A_{m+1} if necessary, $G(V''_m \times A_{m+1})$ contains an open subset of the form $V_{m+1} \times A_{m+1}$. For each $\mathbf{a} \in A_{m+1}$, let $\psi^{\mathbf{a}}_{m+1} = \psi^{\mathbf{a}}_m \circ \mathbf{W}^{-1}: V_{m+1} \to D_{\mathbf{z}}$. Then we have

$$f^{\mathbf{a}}(\psi_{m+1}^{\mathbf{a}}(\mathbf{W})) = f^{\mathbf{a}}(\mathbf{0}) - W_1^2 - \dots - W_m^2 + \sum_{i,j=m+1}^n W_i W_j H_{m+1,ij}^{\mathbf{a}}(\mathbf{W})$$

for some holomorphic functions $H_{m+1,ij}^{\mathbf{a}}(\mathbf{Z})$ that are symmetric in *i* and *j*.

Inductively doing the above procedure on m, and letting $V = V_n$, $A = A_n$ and $\psi^{\mathbf{a}} = \psi_n^{\mathbf{a}}$, we prove the result. Moreover, by the Chain Rule, we have

$$\operatorname{Hess}(f^{\mathbf{a}} \circ \psi^{\mathbf{a}})(\mathbf{0})) = (D\psi^{\mathbf{a}}(\mathbf{0}))^{T} \Big(\operatorname{Hess}(f^{\mathbf{a}})(\mathbf{0}) \Big) (D\psi^{\mathbf{a}}(\mathbf{0})).$$

Since $\text{Hess}(f^{\mathbf{a}} \circ \psi^{\mathbf{a}})(\mathbf{0})$ is equal to the negative of the $n \times n$ identity matrix, by taking the determinant on both sides, we get

$$\det\left(D(\psi^{\mathbf{a}})(\mathbf{0})\right) = \frac{2^{\frac{\mu}{2}}}{\sqrt{-\det\operatorname{Hess}(f^{\mathbf{a}})(\mathbf{0})}}.$$

References

- J. Barrett. Geometrical measurements in three-dimensional quantum gravity, International Journal of Modern Physics A, 18(supp02):97–113, 2003.
- [2] G. Belletti, A maximum volume conjecture for hyperbolic polyhedra, Preprint: arXiv:2002.01904.
- [3] G. Belletti and T. Yang, *Discrete Fourier transform, quantum 6j-symbols and deeply truncated tetrahedra*, in preparation.
- [4] G. Belletti, R. Detcherry, E. Kalfagianni, and T. Yang, *Growth of quantum 6j-symbols and applications to the Volume Conjecture*, to appear in J. Differential Geom.
- [5] C. Blanchet, N. Habegger, G. Masbaum and P. Vogel, *Three-manifold invariants derived from the Kauffman bracket*, Topology 31 (1992), no. 4, 685–699.
- [6] Q. Chen and T. Yang, Volume Conjectures for the Reshetikhin-Turaev and the Turaev-Viro Invariants, Quantum Topol. 9 (2018), no. 3, 419–460.
- [7] D. Cooper, C. Hodgson and S. Kerckhoff, *Three-dimensional orbifolds and cone-manifolds. With a postface by Sadayoshi Kojima*, MSJ Memoirs, 5. Mathematical Society of Japan, Tokyo, 2000. x+170 pp. ISBN: 4-931469-05-1.
- [8] F. Costantino, 6*j*-symbols, hyperbolic structures and the volume conjecture, Geom. Topol. 11 (2007), 1831–1854.

- [9] F. Costantino, *Colored Jones invariants of links in* $S^3 \#^k S^2 \times S^1$ *and the volume conjecture,* J. London Math. Soc. 76 (2007), no. 2, 1–15.
- [10] F. Costantino and D. Thurston, 3-manifolds efficiently bound 4-manifolds, J. Topol. 1 (2008), no. 3, 703–745.
- [11] R. Detcherry and E. Kalfagianni, *Gromov norm and Turaev-Viro invariants of 3-manifolds*, to appear in Ann. Sci. de l'Ecole Normale Sup..
- [12] R. Detcherry, E. Kalfagianni and T. Yang, *Turaev-Viro invariants, colored Jones polynomials and volume*, Quantum Topol. 9 (2018), no. 4, 775–813.
- [13] L. Faddeev, Discrete Heisenberg-Weyl group and modular group, Lett. Math. Phys. 34 (1995), no. 3, 249–254.
- [14] L. Faddeev, R. Kashaev and A. Volkov, *Strongly coupled quantum discrete Liouville theory*, I. Algebraic approach and duality. Comm. Math. Phys. 219 (2001), no. 1, 199–219.
- [15] S. Garoufalidis and T. Le, Asymptotics of the colored Jones function of a knot, Geom. Topol. 15 (2011), no. 4, 2135–2180.
- [16] S. Gukov, *Three-dimensional quantum gravity, Chern-Simons theory, and the A-polynomial*, Comm. Math. Phys. 255 (2005) 577–627.
- [17] C. Hodgson, Degeneration and regeneration of geometric structures on three-manifolds (foliations, Dehn surgery), Thesis (Ph.D.) Princeton University. 1986. 175 pp.
- [18] C. Hodgson and S. Kerckhoff, *Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery*, J. Differential Geom. 48 (1998), no. 1, 1–59.
- [19] R. Kashaev, A link invariant from quantum dilogarithm, Modern Phys. Lett. A 10 (1995), no. 19, 1409–1418.
- [20] R. Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. 39 (1997), no. 3, 269–275.
- [21] S. Kojima, *Deformations of hyperbolic 3-cone-manifolds*, J. Differential Geom. 49 (1998), no. 3, 469–516.
- [22] S. Kumar, Fundamental shadow links realized as links in S³, https://arxiv.org/abs/2005.11447
- [23] W. Lickorish, *The skein method for three-manifold invariants*, J. Knot Theory Ramifications 2 (1993), no. 2, 171–194.
- [24] D. London, A note on matrices with positive definite real part, Proc. Amer. Math. Soc. 82 (1981), no. 3, 322–324.
- [25] H. Murakami and J. Murakami, *The colored Jones polynomials and the simplicial volume of a knot*, Acta Math. 186 (2001), no. 1, 85–104.
- [26] H. Murakami, J. Murakami, M. Okamoto, T. Takata and Y. Yokota, *Kashaev's conjecture and the Chern-Simons invariants of knots and links*, Experiment. Math. 11 (2002) 427–435.
- [27] H. Murakami and Y. Yokota, *The colored Jones polynomial of the figure eight knot and its Dehn surgery spaces*, J. Reine Angew. Math. 607 (2007), 47–68.

- [28] W. Neumann and D. Zagier, *Volumes of hyperbolic three-manifolds*, Topology 24 (1985), no. 3, 307–332.
- [29] T. Ohtsuki, On the asymptotic expansion of the Kashaev invariant of the 5₂ knot, Quantum Topol. 7 (2016), no. 4, 669–735.
- [30] T. Ohtsuki, On the asymptotic expansion of the quantum SU(2) invariant at $q = \exp(4\pi\sqrt{-1/N})$ for closed hyperbolic 3-manifolds obtained by integral surgery along the figure-eight knot, Algebr. Geom. Topol. 18 (2018), no. 7, 4187–4274.
- [31] J.Purcell, An introduction to fully augmented links, http://users.monash.edu/~jpurcell/ papers/expos-auglinks.pdf
- [32] N. Reshetikhin and V. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, Comm. Math. Phys. 127 (1990), no. 1, 1–26.
- [33] N. Reshetikhin and V. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. 103 (1991), no. 3, 547–597.
- [34] J. Roberts, Skein theory and Turaev-Viro invariants, Topology 34 (1995), no. 4, 771–787.
- [35] D. Rolfsen, *Knots and links*, 2nd printing with corrections, Mathematics Lecture Series 7, Publish or Perish, Inc. (1990).
- [36] E. Stein and R. Shakarchi, *Fourier analysis*, An introduction. Princeton Lectures in Analysis, 1. Princeton University Press, Princeton, NJ, 2003. xvi+311 pp. ISBN: 0-691-11384-X.
- [37] W. Thurston, *The geometry and topology of 3-manifolds*, Princeton Univ. Math. Dept. (1978). Available from http://msri.org/publications/books/gt3m/.
- [38] V. Turaev, Quantum invariants of knots and 3-manifolds, Second revised edition. de Gruyter Studies in Mathematics, 18. Walter de Gruyter & Co., Berlin, 2010.
- [39] V. Turaev and O. Viro, *State sum invariants of 3-manifolds and quantum 6j-symbols*, Topology 31 (1992), no. 4, 865–902.
- [40] Ushijima, A volume formula for generalised hyperbolic tetrahedra, Non-Euclidean geometries, 249–265, Math. Appl. (N. Y.), 581, Springer, New York, 2006.
- [41] R. van der Veen, *The volume conjecture for augmented knotted trivalent graph*, Algebr. Geom. Topol., Volume 9, Number 2 (2009), 691–722.
- [42] K. Walker, On Wittens 3-manifold invariants, preprint.
- [43] E. Witten, Quantum field theory and the Jones polynomial, Comm. Math. Phys. 121 (3): 351–399.
- [44] K. H. Wong and T. Yang, On the Volume Conjecture for hyperbolic Dehn-filled 3-manifolds along the figure-eight knot, Preprint, arXiv:2003.10053.
- [45] K. H. Wong and T. Yang, *Relative Reshetikhin-Turaev invariants, hyperbolic cone metrics and discrete Fourier transforms II*, in preparation.
- [46] T. Yoshida, The η -invariant of hyperbolic 3-manifolds, Invent. Math. 81 (1985), no. 3, 473–514.

[47] D. Zagier, *The dilogarithm function*, Frontiers in number theory, physics, and geometry. II, 365, Springer, Berlin, 2007.

Ka Ho Wong Department of Mathematics Texas A&M University College Station, TX 77843, USA (daydreamkaho@math.tamu.edu)

Tian Yang Department of Mathematics Texas A&M University College Station, TX 77843, USA (tianyang@math.tamu.edu)