On the winding number for particle trajectories in a disk-like vortex patch of the Euler equations

Kyudong Choi^{*} In-Jee Jeong[†]

March 11, 2021

Abstract

We consider vortex patch solutions of the incompressible Euler equations in the plane. It is shown that the winding number around the origin for most particles in the patch grows linearly in time when the initial patch is close to a disk enough.

1 Introduction

We consider the incompressible Euler equation in vorticity form in \mathbb{R}^2 :

$$\partial_t \omega + u \cdot \nabla \omega = 0 \quad \text{for } x \in \mathbb{R}^2 \text{ and for } t > 0,$$

$$\omega|_{t=0} = \omega_0 \quad \text{for } x \in \mathbb{R}^2.$$
(1.1)

where the Biot-Savart law is given by $u = (K * \omega)$ with

$$K(x) := \frac{1}{2\pi} \frac{x^{\perp}}{|x|^2} = \frac{1}{2\pi} \left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2} \right)$$

When ω_0 lies on $L^1 \cap L^\infty$, the existence and uniqueness of a global-in-time weak solution is due to Yudovich [12]. We are concerned with vortex patch solutions; when $\omega_0 = \mathbf{1}_{\Omega_0}$ for some bounded measurable set $\Omega_0 \subset \mathbb{R}^2$, the corresponding Yudovich solution ω has the form of $\omega(t) = \mathbf{1}_{\Omega_t}$ where $\Omega_t = \{\phi_x(t) \in \mathbb{R}^2 \mid x \in \Omega_0\}$. Here $\phi_x(\cdot)$ is the unique solution to the system

$$\frac{d}{dt}\phi_x(t) = u(t,\phi_x(t)) \quad \text{for } t > 0 \quad \text{and} \quad \phi_x(0) = x,$$

^{*}Department of Mathematical Sciences, Ulsan National Institute of Science and Technology. Email: kchoi@unist.ac.kr

[†]Department of Mathematical Sciences, Seoul National University. E-mail: injee_j@snu.ac.kr

Key words: 2D Euler, vortex patch, stability, winding number, large time behavior, particle trajectory. 2010 AMS Mathematics Subject Classification: 76B47, 35Q35

which is well-defined thanks to the estimate

$$\|u(t)\|_{L^{\infty}} \le C \|\omega_0\|_{L^1 \cap L^{\infty}}, \quad t \ge 0$$
(1.2)

and the log-Lipschitz estimate

$$|u(t,x) - u(t,x')| \le C \|\omega_0\|_{L^1 \cap L^\infty} |x - x'| \ln(\frac{1}{|x - x'|}), \quad |x - x'| \le \frac{1}{2}, \quad t \ge 0.$$
(1.3)

The above estimates in turn guarantee that for each T > 0, there exists constants c = c(T), C = C(T) > 0 satisfying

$$c|x - x'|^{\exp(C\|\omega_0\|_{L^1 \cap L^{\infty}} t)} \le |\phi_x(t) - \phi_{x'}(t)| \le C|x - x'|^{\exp(-C\|\omega_0\|_{L^1 \cap L^{\infty}} t)}, \quad x, x' \in \mathbb{R}^2, \quad t \in [0, T]$$

$$(1.4)$$

(e.g. see [13, Sec. 8.2.3] or [14, Ch. 2]).

In this paper, we produce an estimate on the winding number around the origin for fluid particle trajectories when the initial patch is disk-like. Here we say that a patch on Ω is disk-like if the measure of the symmetric difference

$$\Omega \bigtriangleup B_r := (\Omega \backslash B_r) \cup (B_r \backslash \Omega)$$

is small enough for some r > 0 where we denote $B_r := \{x \in \mathbb{R}^2 \mid |x| < r\}$. Under the assumption that $x \in \mathbb{R}^2 \setminus \{0\}$ is a point whose trajectory never hits the origin (i.e. $\phi_x(t) \neq 0$ for all $t \geq 0$)¹, we define the notion of winding number of x up to time t > 0 is defined by the integral

$$N_x(t) := \frac{1}{2\pi} \int_0^t \frac{u_{tan}(s, \phi_x(s))}{|\phi_x(s)|} ds,$$

where u_{tan} is the speed in the angular direction with respect to the origin (see Definition (1.6)). It is not difficult to check that this definition coincides with the usual notion of winding number (for curves in \mathbb{R}^2 not intersecting the origin) applied to the curve $\gamma_x : [0,t] \mapsto \mathbb{R} \setminus \{0\}$ defined by $\gamma_x(s) = \phi_x(s)$. For example, we observe that for the circular vortex patch supported on the unit disc $D := B_1$, we have, for any t > 0,

$$\frac{N_x(t)}{t} = \frac{1}{4\pi}, \quad \forall x \in D \setminus \{0\},$$
(1.5)

simply because $u_{tan}(t, x) = \frac{1}{2}|x|$ for $t \ge 0$ and $x \in D \setminus \{0\}$.

There are several difficulties in treating the winding number. First, this quantity may decrease in time (particles can "unwind") and second, there is no uniform in space bound for the ratio u(t,x)/|x|. Rather, even assuming that the velocity vanishes at the origin (u(t,0) = 0), taking x' = 0 in (1.3) shows that the quantity may diverge like $C \log(1/|x|)$ as $|x| \to 0$, which is indeed

¹In Lemma 2.4 we shall prove that the set of points in \mathbb{R}^2 whose trajectory hits the origin after some time is of zero measure.

sharp for vortex patch solutions (e.g. see [2]).

1.1 Main result

We consider the initial vorticity given by the patch on some bounded measurable set Ω_0 of unit strength, whose solution is identified with moving domains Ω_t . We shall assume that Ω_0 is a small perturbation of the unit disc D. We are now ready to state our main result on winding number for disk-like patches.

Theorem 1.1. For any R > 0, there exist constants $\delta_0 \in (0,1)$ and C > 0 such that if

 $\Omega_0 \subset B_R$ and $|D \triangle \Omega_0| < \delta_0$,

then, by denoting $\delta := |D \triangle \Omega_0|$, for any T > 0, there exists a set $H_T \subset \Omega_0$ such that

$$|H_T| \ge |\Omega_0| - C\delta^{1/2}$$

and

$$\left|\frac{N_x(T)}{T} - \frac{1}{4\pi}\right| \le C\delta^{1/12}, \quad x \in H_T.$$

In addition, there exists a set $H \subset \Omega_0$ such that

$$|H| \ge |\Omega_0| - C\delta^{1/12}$$

and

$$\liminf_{t \to \infty} \left| \frac{N_x(t)}{t} - \frac{1}{4\pi} \right| \le C \delta^{1/12}, \quad x \in H.$$

Recalling (1.5), the above result states that most particles wind around the origin roughly the same number of times as particles in the case of the disc. This can be interpreted as a more refined, dynamical in nature, notion of stability; the classical stability results (Wan and Pulvirenti [19] and Sideris and Vega [17]) simply say that for initially disc-like patches, the shape of the patch remains close to the disc for all times.

Moreover, control on the winding number could be useful to prove certain *instability* results for vortex patches and more generally for solutions to the 2D Euler equations. The basic idea is as follows: if two fluid particles rotate around the origin with different angular velocities while roughly keeping their respective distance from the origin, any curve connecting the particles gets stretched linearly in time. This observation was used in [9] to obtain spiral formation and [7] for perimeter growth.

Notations

We collect the notations here that are used throughout the paper.

- Given r > 0 and $x \in \mathbb{R}^2$, we define $B_r(x) := \{y \in \mathbb{R}^2 \mid |y x| < r\}$ and $B_r := B_r(0)$. We denote the unit disc by $D := B_1$. The complement is denoted by D^C .
- Given $\epsilon > 0$ and $i \in \mathbb{N} \cup \{0\}$, we set $B_i^{\epsilon} := B_{2^{-i}\epsilon}$ and $A_i^{\epsilon} := B_i^{\epsilon} \setminus B_{i+1}^{\epsilon}$.

• The radial and tangential part of the velocity around the origin are defined by

$$u = u_{rad} \frac{x}{|x|} + u_{tan} \frac{x^{\perp}}{|x|}.$$
 (1.6)

Here we denote $x^{\perp} = (-x_2, x_1)$ for $x = (x_1, x_2)$.

2 Proof

Proof of Theorem 1.1. Fix R > 0 and let $\omega(t) = \mathbf{1}_{\Omega_t}$ be the solution of (1.1) with initial data $\omega_0 = \mathbf{1}_{\Omega_0}$ where $\Omega_0 \subset B_R$. Take any $\delta_0 > 0$ satisfying $\delta_0^{1/6} \leq 1/2$ and assume

$$\delta := |\Omega_0 \triangle D| < \delta_0.$$

If $\delta = 0$, then we can take H_T , $H = D \setminus \{0\}$ by (1.5). From now on, we assume $\delta > 0$.

Throughout the paper, C denotes a positive constant which may depend on R > 0, may change from line to line, but it is independent of $\delta_0, \delta > 0$.

2.1 L¹-stability of a disk patch

We recall the L^1 -stability result for the circular vortex patch from [17]:

Lemma 2.1 ([17, Theorem 3]). For any bounded open set $\Omega_0 \subset \mathbb{R}^2$ and for any r > 0, we have

$$\|\mathbf{1}_{\Omega_t} - \mathbf{1}_{B_r}\|_{L^1}^2 \le 4\pi \cdot \sup_{x \in \Omega_0 \triangle B_r} \|x\|^2 - r^2\| \cdot \|\mathbf{1}_{\Omega_0} - \mathbf{1}_{B_r}\|_{L^1} \quad for \ any \ t > 0$$

Note that $\|\mathbf{1}_A - \mathbf{1}_B\|_{L^1} = \int_{A \triangle B} 1 dx = |A \triangle B|$. Applying the above lemma with our $\Omega_0 \subset B_R$ and $\mathbf{1}_D$, we obtain that

$$|\Omega_t \triangle D| \le C\sqrt{\delta}, \quad t > 0. \tag{2.1}$$

From this L^1 bound, we obtain that (e.g. see [6, 9]).

Lemma 2.2. Under the above assumptions on Ω_0 , we have with $u(t) = K * \mathbf{1}_{\Omega(t)}$ and $u_D = K * \mathbf{1}_D$ that

$$\|u(t) - u_D\|_{L^{\infty}} \le C\delta^{1/4}$$

for all $t \geq 0$. In particular, we have

$$\left|\frac{u_{tan}(t,x)}{|x|} - \frac{1}{2}\right| \le C \frac{\delta^{1/4}}{|x|}, \quad x \in D \setminus \{0\}.$$
(2.2)

Proof. From the estimate $|K(x)| \leq C/|x|$, we have

$$||K * f||_{L^{\infty}} \le C ||f||_{L^{1}}^{1/2} ||f||_{L^{\infty}}^{1/2}$$

for any $f \in (L^1 \cap L^\infty)(\mathbb{R}^2)$ (e.g. see Lemma 2.1. in [11]). Thus, we can estimate, for any $x \in \mathbb{R}^2$,

$$u(t,x) - u_D(x)| \le \|K * (\mathbf{1}_{\Omega_t} - \mathbf{1}_D)\|_{L^{\infty}} \le C \|\mathbf{1}_{\Omega_t} - \mathbf{1}_D\|_{L^1}^{1/2} \|\mathbf{1}_{\Omega_t} - \mathbf{1}_D\|_{L^{\infty}}^{1/2} \le C |\Omega_t \triangle D|^{1/2}.$$

Thus, by (2.1), we get the first estimate. Taking the tangential components and using that $u_{D,tan} = |x|/2$ for $x \in D \setminus \{0\}$, we obtain (2.2).

Next, we estimate time spent near the origin for particles. A similar estimate can be found in [6, Lemma 2.2].

Lemma 2.3. Under the above assumptions on Ω_0 , for any T > 0, we have

$$\int_{\Omega_0} \left(\frac{1}{T} \int_0^T \mathbf{1}_{B_r}(\phi_x(t)) dt \right) dx \le Cr^2, \quad r > 0$$

and

$$\int_{\Omega_0} \left(\frac{1}{T} \int_0^T \mathbf{1}_{D^C}(\phi_x(t)) dt \right) dx \le C\sqrt{\delta}.$$

Proof. We compute, using that the flow map $x \mapsto \phi_x(t)$ is area-preserving, for r > 0 and t > 0,

$$\int_{\Omega_0} \mathbf{1}_{B_r}(\phi_x(t)) dx = \int_{\Omega_t \cap B_r} 1 \, dx \le |B_r| = \pi r^2,$$

and, by (2.1),

$$\int_{\Omega_0} \mathbf{1}_{D^C}(\phi_x(t)) dx = \int_{\Omega_t \cap D^C} 1 \, dx \le \int_{\Omega_t \triangle D} 1 \, dx = |\Omega_t \triangle D| \le C \sqrt{\delta}.$$

Now the desired bounds follow from integrating in time and applying Fubini's theorem.

2.2 Set of particles touching the origin

We prove the following result:

Lemma 2.4. Let $\phi_x(t)$ be the flow map associated with a Yudovich solution. Then, we have

$$|\{x \in \mathbb{R}^2 | \phi_x(t) = 0 \text{ for some } t \ge 0\}| = 0.$$

This result is not trivial since for Yudovich solutions (in particular vortex patches) since the flow map is in general only Hölder continuous with Hölder exponent decaying to zero as $t \to \infty$. On the other hand, there are space-filling curves in the plane which is actually Hölder continuous with exponent 1/2 (Hilbert's curve).

Proof of Lemma 2.4. First step is to prove that the inverse of the flow map is Hölder continuous with respect to time. For any given $t \ge 0$, we consider $\phi(t, x) = \phi_x(t)$ as a bijection $\mathbb{R}^2 \to \mathbb{R}^2$, and denote the inverse map by $A(t, \cdot)$; that is, $A(t, \phi(t, x)) = x$. For this purpose, take some $t, t' \ge 0$ and observe that

$$\phi(t,x) - \phi(t',x) = \int_{t'}^t u(\tau,\phi(\tau,x)) \, d\tau$$

gives

$$\frac{|\phi(t,x) - \phi(t',x)|}{|t - t'|} \le C \|u\|_{L^{\infty}_{t,x}} \le C \|\omega_0\|_{L^1 \cap L^{\infty}}.$$

Then, for $0 \le t' \le t \le 1$ and for $x \in \mathbb{R}^2$, we first take a, a' such that $\phi(t', a') = \phi(t, a) = x$, so that

$$A(t,x) - A(t',x) = a - a'.$$

We have

$$\phi(t', a') - \phi(t, a') = \phi(t, a) - \phi(t, a'),$$

and the left hand side is bounded in absolute value by $C \|\omega_0\|_{L^1 \cap L^{\infty}} |t - t'|$. Using this together with the lower bound given in (1.4), we deduce that there exist absolute constants c, C > 0 such that

$$c|a - a'|^{\exp(Ct\|\omega_0\|_{L^1 \cap L^\infty})} \le C\|\omega_0\|_{L^1 \cap L^\infty} |t - t'|.$$

Hence we obtain

$$|A(t,x) - A(t',x)| \le C(\|\omega_0\|_{L^1 \cap L^\infty} |t - t'|)^{\exp(-Ct\|\omega_0\|_{L^1 \cap L^\infty})}, \quad x \in \mathbb{R}^2, \quad 0 \le t' \le t \le 1.$$
(2.3)

Now we fix any $\alpha \in (1/2, 1)$. Given the estimate (2.3), we may find constants $T_0 \in (0, 1)$ and $C_0 > 0$ depending only on $\|\omega_0\|_{L^1 \cap L^\infty}$ such that

$$|A(t,x) - A(t',x)| \le C_0 |t - t'|^{\alpha}, \quad x \in \mathbb{R}^2, \quad 0 \le t' \le t \le T_0.$$
(2.4)

We can now prove that

$$|\{x \in \mathbb{R}^2 | \phi_x(t) = 0 \text{ for some } t \in [0, T_0)\}| = 0.$$

To see this, observe that the set of points x satisfying $\phi_x(t) = 0$ for some $t \in [0, T_0)$ is simply given by $\{A(t, 0) | t \in [0, T_0)\} =: \mathcal{A}([0, T_0))$. Fix any integer $n \ge 1$ and consider the points $x_k := A(\frac{kT_0}{n}, 0)$ for $k = 0, 1, \ldots, (n-1)$. From (2.4), we see that

$$\mathcal{A}([0,T_0)) \subset \bigcup_{k=0}^{n-1} B_{C_0(\frac{T_0}{n})^{\alpha}}(x_k),$$

which implies that

$$|\mathcal{A}([0,T_0))| \le C n^{1-2\alpha} T_0^{2\alpha}.$$

In the last estimate, C, T_0 are independent of $n \ge 1$, and therefore we conclude that $|\mathcal{A}([0, T_0))| = 0$ thanks to $\alpha > 1/2$.

We now write $\phi(t; s, a)$ for the location of the particle at time t which was a at time s. For each integer $m \ge 0$, we consider the set

$$U_m = \{\phi(mT_0; s, 0) \mid s \in [mT_0, (m+1)T_0)\}.$$

Note that $\mathcal{A}([0,T_0)) = U_0$. Applying the same argument (with initial data ω_0 replaced with $\omega(t = mT_0)$), we deduce that $|U_m| = 0$. This is possible only because T_0 depends only on $||\omega||_{L^1 \cap L^\infty}$, which is independent of time. Next, since $A(t, \cdot)$ is measure preserving for any $t \ge 0$, we see that $|A(mT_0, U_m)| = 0$. The set $A(mT_0, U_m)$ is precisely the collection of points which touches the origin at some time $[mT_0, (m+1)T_0)$. This finishes the proof.

2.3 Estimate of winding number

Denote

$$\epsilon := \delta^{1/6} > 0.$$

Observe $\epsilon = \delta^{1/6} < \delta_0^{1/6} \le 1/2$. Fix T > 0 and define

$$G_i^T(x) := \frac{1}{T} \int_0^T \mathbf{1}_{B_i^{\epsilon}}(\phi_x(t)) dt, \quad i \ge 0$$

(recall the notation $B_i^{\epsilon} := B_{2^{-i}\epsilon}$) and

$$G_{-1}^{T}(x) := \frac{1}{T} \int_{0}^{T} \mathbf{1}_{D^{C}}(\phi_{x}(t)) dt$$

Lemma 2.3 states that

$$\int_{\Omega_0} G_i^T(x) dx \le C (2^{-i} \epsilon)^2, \quad i \ge 0,$$

and, from Chebyshev's inequality, we obtain

$$\left| \left\{ x \in \Omega_0 \, | \, G_i^T(x) \ge (2^{-i}\epsilon)^{\frac{3}{2}} \right\} \right| \le C(2^{-i}\epsilon)^{\frac{1}{2}} = C2^{-i/2}\delta^{1/12}, \quad i \ge 0.$$

Similarly, we have

$$\left| \left\{ x \in \Omega_0 \, | \, G_{-1}^T(x) \ge 2\epsilon \right\} \right| \le C \frac{\sqrt{\delta}}{\epsilon} = C \delta^{1/3} \le C \delta^{1/12}.$$

Now we define

$$H_T := \{ x \in \Omega_0 \, | \, G_i^T(x) < (2^{-i}\epsilon)^{\frac{3}{2}}, \, \forall i \ge 0 \text{ and } G_{-1}^T(x) < 2\epsilon \text{ and } \phi_x(t) \ne 0, \, \forall t \in [0,T] \}.$$

Recalling Lemma 2.4, we have

$$|H_T| \ge |\Omega_0| - C\delta^{1/12}.$$

For any fixed $x \in H_T$, we split the time integral as follows:

$$\int_{0}^{T} \frac{u_{tan}(t,\phi_{x}(t))}{|\phi_{x}(t)|} dt = \int_{[0,T] \setminus (\bigcup_{i \ge -1} I_{i})} \frac{u_{tan}(t,\phi_{x}(t))}{|\phi_{x}(t)|} dt + \sum_{i \ge -1} \int_{I_{i}} \frac{u_{tan}(t,\phi_{x}(t))}{|\phi_{x}(t)|} dt,$$

where

$$I_i := \{ t \in [0, T] \, | \, \phi_x(t) \in A_i^{\epsilon} \}, \quad i \ge 0$$

(recall the notation $A_i^\epsilon := B_i^\epsilon ackslash B_{i+1}^\epsilon$) and

$$I_{-1} := \{ t \in [0, T] \, | \, \phi_x(t) \in D^c \}.$$

For $i \ge 0$, we have that $|I_i| \le TG_i^T(x) \le T(2^{-i}\epsilon)^{\frac{3}{2}}$ and $|I_{-1}| \le CT\epsilon$. Thus we get

 $|(\cup_{i\geq -1}I_i)| \leq CT\epsilon.$

For $i \ge 0$, we also estimate, using the uniform bound $||u(t)||_{L^{\infty}} \le C(1 + |\Omega_0|) \le C$ which follows from (1.2),

$$\left| \int_{I_i} \frac{u_{tan}(t, \phi_x(t))}{|\phi_x(t)|} dt \right| \le C \int_{I_i} \frac{1}{|\phi_x(t)|} dt \le C |I_i| 2^i \epsilon^{-1} \le CT \epsilon^{1/2} 2^{-i/2}.$$

In the case i = -1, we simply use $|\phi_x(t)| \ge 1$ together with $||u(t)||_{L^{\infty}} \le C$ to obtain

$$\left| \int_{I_{-1}} \frac{u_{tan}(t, \phi_x(t))}{|\phi_x(t)|} dt \right| \le C |I_{-1}| \le C T \epsilon.$$

Hence we obtain

$$\left|\sum_{i\geq -1} \int_{I_i} \frac{u_{tan}(t, \phi_x(t))}{|\phi_x(t)|} dt\right| \leq CT\sqrt{\epsilon}.$$

Note that the above constant C > 0 is independent of $x \in H_T$.

Lastly, for $t \in [0,T] \setminus (\cup_{i \geq -1} I_i)$ we have that

$$\epsilon \le |\phi_x(t)| < 1,$$

from which it follows by (2.2) that

$$\left|\frac{u_{tan}(t,\phi_x(t))}{|\phi_x(t)|} - \frac{1}{2}\right| \le C \frac{\delta^{1/4}}{|\phi_x(t)|} \le C \frac{\delta^{1/4}}{\epsilon} \le C \delta^{1/12}.$$

Therefore, we obtain the lower bound

$$\int_{[0,T]\setminus(\cup_{i\geq -1}I_i)}\frac{u_{tan}(t,\phi_x(t))}{|\phi_x(t)|}dt \ge T(1-C\epsilon)\left(\frac{1}{2}-C\delta^{1/12}\right).$$

Collecting all the estimates, we obtain

$$\frac{1}{T} \int_0^T \frac{u_{tan}(t, \phi_x(t))}{|\phi_x(t)|} dt \ge (1 - C\epsilon) \left(\frac{1}{2} - C\delta^{1/12}\right) - C\sqrt{\epsilon} \ge \frac{1}{2} - C\delta^{1/12}.$$

This gives the lower bound

$$\frac{N_x(T)}{T} \ge \frac{1}{4\pi} - C\delta^{1/12},$$

while the upper bound can be obtained similarly. The proof for the first statement is now complete. To see the second statement, we just observe that with $T_n := n$

$$H := \limsup_{n} H_{T_n} = \bigcap_{m \ge 1} \bigcup_{n \ge m} H_{T_n}$$

has measure greater than $|\Omega_0| - C\delta^{1/12}$.

Remark 2.5. Since Yudovich's log-Lipschitz estimate in principle allows for arbitrarily fast rotation, it seems like an interesting problem to ask whether there is an initial data $\omega_0 \in L^1 \cap L^\infty$ which has a trajectory winding around the origin arbitrarily many times in a fixed time interval. This is related to the (difficult) question of whether instantaneous spiraling is possible for vortex patches. We refer the interested readers to discussions in [8, Sec. 5–6], [10, 16].

Remark 2.6. In a recent related work [6], the travel distance for fluid particles

$$d_x(t) := \int_0^t |u(s,\phi_x(s))| ds$$

was considered, and it was established that for most particles on the initial vortex patch, this quantity grows linearly in time when the initial patch is disk-like.

Remark 2.7. It is also interesting to ask a similar question about winding number (or travel distance) on well-known solutions other than a disk, whose certain stability has been known. It might include Kirchhoff's ellipse [18], Lamb dipoles [1], Hill's vortex [5], shear flows [3] and rectangles in a strip [4], and so on.

Acknowledgement

KC has been supported by the National Research Foundation of Korea (NRF-2018R1D1A1B07043065) and by the Research Fund (1.200085.01) of UNIST(Ulsan National Institute of Science & Technology). IJ has been supported by the Science Fellowship of POSCO TJ Park Foundation and the National Research Foundation of Korea grant 2019R1F1A1058486. IJ thanks Sun-Chul Kim for telling about the article [15].

References

- [1] Ken Abe and Kyudong Choi. Stability of Lamb dipoles. preprint, arXiv:1911.01795.
- [2] H. Bahouri and J.-Y. Chemin. Équations de transport relatives á des champs de vecteurs non-lipschitziens et mécanique des fluides. Arch. Rational Mech. Anal., 127(2):159–181, 1994.
- [3] Jacob Bedrossian and Nader Masmoudi. Inviscid damping and the asymptotic stability of planar shear flows in the 2D Euler equations. *Publ. Math. Inst. Hautes Études Sci.*, 122:195– 300, 2015.
- [4] Jennifer Beichman and Sergey Denisov. 2D Euler equation on the strip: stability of a rectangular patch. Comm. Partial Differential Equations, 42(1):100–120, 2017.
- [5] Kyudong Choi. Stability of Hill's spherical vortex. preprint, arXiv:2011.06808.
- [6] Kyudong Choi. On the estimate of distance traveled by a particle in a disk-like vortex patch. Appl. Math. Lett., 97:67–72, 2019.
- [7] Kyudong Choi and In-Jee Jeong. Growth of perimeter for vortex patches in a bulk. Appl. Math. Lett., 113:106857, 9, 2021.
- [8] Tarek M. Elgindi and In-Jee Jeong. On singular vortex patches, I: Well-posedness issues. Memoirs of the AMS, to appear, arXiv:1903.00833.
- [9] Tarek M. Elgindi and In-Jee Jeong. On singular vortex patches, II: long-time dynamics. Trans. Amer. Math. Soc., 373(9):6757–6775, 2020.
- [10] V. Elling and M. V. Gnann. Variety of unsymmetric multibranched logarithmic vortex spirals. European J. Appl. Math., 30(1):23–38, 2019.
- [11] Dragoş Iftimie, Thomas C. Sideris, and Pascal Gamblin. On the evolution of compactly supported planar vorticity. Comm. Partial Differential Equations, 24(9-10):1709–1730, 1999.
- [12] V. I. Judovič. Non-stationary flows of an ideal incompressible fluid. Ž. Vyčisl. Mat. i Mat. Fiz., 3:1032–1066, 1963.
- [13] Andrew J. Majda and Andrea L. Bertozzi. Vorticity and incompressible flow, volume 27. Cambridge University Press, Cambridge, 2002.
- [14] Carlo Marchioro and Mario Pulvirenti. Mathematical theory of incompressible nonviscous fluids, volume 96 of Applied Mathematical Sciences. Springer-Verlag, New York, 1994.
- [15] T. B. Mitchell and L. F. Rossi. The evolution of Kirchhoff elliptic vortices. *Physics of Fluids*, 20(5):054103, 2008.
- [16] D. I. Pullin. On similarity flows containing two-branched vortex sheets. In Mathematical aspects of vortex dynamics (Leesburg, VA, 1988), pages 97–106. SIAM, Philadelphia, PA, 1989.
- [17] Thomas C. Sideris and Luis Vega. Stability in L¹ of circular vortex patches. Proc. Amer. Math. Soc., 137(12):4199–4202, 2009.

- [18] Yun Tang. Nonlinear stability of vortex patches. Trans. Amer. Math. Soc., 304(2):617–638, 1987.
- [19] Y. H. Wan and M. Pulvirenti. Nonlinear stability of circular vortex patches. Comm. Math. Phys., 99(3):435–450, 1985.