

Noncritical Weighted Hardy's Inequalities with compact perturbations

HIROSHI ANDO and TOSHIO HORIUCHI

Abstract

Let Ω be a bounded domain of $\mathbb{R}^N$ ($N \geq 1$) whose boundary $\partial\Omega$ is a C^2 compact manifolds. In the present paper we shall study a variational problem relating the weighted Hardy inequalities with sharp missing terms established in [4]. As weights we adopted powers of the distance function $\delta(x)$ to the boundary $\partial\Omega$.¹

1 Introduction

Let Ω be a bounded domain of $\mathbb{R}^N$ ($N \geq 1$) whose boundary $\partial\Omega$ is a C^2 compact manifolds. In [4] we have established N dimensional weighted Hardy's inequalities with weight function being powers of the distance function $\delta(x) = \text{dist}(x, \partial\Omega)$ to the boundary $\partial\Omega$. In this paper we shall study a variational problem relating to these new inequalities.

We prepare more notations to describe our results. Let $1 < p < \infty$ and $\alpha < 1 - 1/p$. By $L^p(\Omega, \delta^{\alpha p})$ we denote the space of Lebesgue measurable functions with weight $\delta^{\alpha p}$, for which

$$\|u\|_{L^p(\Omega, \delta^{\alpha p})} = \left(\int_{\Omega} |u|^p \delta^{\alpha p} dx \right)^{1/p} < \infty. \quad (1.1)$$

$W_{\alpha,0}^{1,p}(\Omega)$ is given by the completion of $C_c^\infty(\Omega)$ with respect to the norm defined by

$$\|u\|_{W_{\alpha,0}^{1,p}(\Omega)} = \| |\nabla u| \|_{L^p(\Omega, \delta^{\alpha p})}. \quad (1.2)$$

Then $W_{\alpha,0}^{1,p}(\Omega)$ becomes a Banach space with the norm $\|\cdot\|_{W_{\alpha,0}^{1,p}(\Omega)}$. Under these preparation we recall the noncritical weighted Hardy inequality in [4]. In particular, we have the simplest one:

$$\int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx \geq \mu \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx \quad \text{for } u \in W_{\alpha,0}^{1,p}(\Omega), \quad (1.3)$$

where μ is a positive constant independent of u . If $\alpha = 0$ and $p = 2$, then (1.3) is a well-known Hardy's inequality and valid for a bounded domain Ω of $\mathbb{R}^N$ with Lipschitz boundary (c.f. [6], [8], [11], [13]). If Ω is convex and $\alpha = 0$, then (1.3) with $\mu = (1 - 1/p)^p$ holds for arbitrary $1 < p < \infty$ (see [13], [14]).

¹Keywords: Weighted Hardy's inequalities, nonlinear variational problem, Weak Hardy property, p -Laplace operator with weights,
2000 mathematics Subject Classification: Primary 35J70, Secondary 35J60, 34L30, 26D10.
This research was partially supported by Grant-in-Aid for Scientific Research (No.16K05189) and (No.15H03621).

The best possible μ in (1.3) is given by the quantity

$$\inf_{u \in W_{\alpha,0}^{1,p}(\Omega), u \neq 0} \frac{\int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx}{\int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx}, \quad (1.4)$$

which depends on p, α and Ω .

In this paper we consider the following variational problem

$$J_{\lambda}^{\alpha} = \inf_{u \in W_{\alpha,0}^{1,p}(\Omega), u \neq 0} \chi_{\lambda}^{\alpha}(u) \quad (1.5)$$

where $\lambda \in \mathbb{R}$ and

$$\chi_{\lambda}^{\alpha}(u) = \frac{\int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx - \lambda \int_{\Omega} |u|^p \delta^{\alpha p} dx}{\int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx}. \quad (1.6)$$

Note that J_0^{α} gives the best constant in (1.3). Clearly, the function $\lambda \mapsto J_{\lambda}^{\alpha}$ is non-increasing on $\mathbb{R}$ and $J_{\lambda}^{\alpha} \rightarrow -\infty$ as $\lambda \rightarrow \infty$.

Remark 1.1. *It is worthy to remark that (1.3) is never valid in the critical case that $\alpha \geq 1 - 1/p$. Nevertheless, we have established in this case a variant of weighted Hardy's inequalities in [4] (cf. [10]). In a coming paper [3], we shall treat general weighted Hardy's inequalities with compact perturbations and study relating variational problems including the critical case that $\alpha \geq 1 - 1/p$.*

This paper is organized in the following way: The main result is described in Section 2. Section 3 is devoted to the proof of main result.

2 Main results

Our main result is the following.

Theorem 2.1. *Assume that Ω is a bounded domain of class C^2 in $\mathbb{R}^N$. Assume that $1 < p < \infty$ and $\alpha < 1 - 1/p$. Then there exists a constant $\lambda^* \in \mathbb{R}$ such that:*

1. *If $\lambda \leq \lambda^*$, then $J_{\lambda}^{\alpha} = \Lambda_{\alpha,p}$. If $\lambda > \lambda^*$, then $J_{\lambda}^{\alpha} < \Lambda_{\alpha,p}$.*

Here

$$\Lambda_{\alpha,p} = \left(1 - \alpha - \frac{1}{p}\right)^p. \quad (2.1)$$

Moreover, it holds that:

2. *If $\lambda < \lambda^*$, then the infimum J_{λ}^{α} in (1.5) is not attained.*
3. *If $\lambda > \lambda^*$, then the infimum J_{λ}^{α} in (1.5) is attained.*

Remark 2.1. 1. *In Theorem 2.1, it remains for $\lambda = \lambda^*$ of the open problem whether the infimum $J_{\lambda^*}^{\alpha}$ in (1.5) is attained or not.*

2. *For the case of $\alpha = 0$ and $p = 2$, it is shown that the infimum J_{λ}^0 in (1.5) is attained if and only if $\lambda > \lambda^*$. See [6].*
3. *For the case of $\alpha = 0$ and $\lambda = 0$, the value of the infimum J_0^0 in (1.5) and its attainability are studied in [13].*

4. In the assertion 3 of Theorem 2.1, if $\lambda > \lambda^*$ then the minimizer u for the variational problem (1.5) is a non-trivial weak solution of the following Euler-Lagrange equation:

$$-\operatorname{div}(\delta^{\alpha p} |\nabla u|^{p-2} \nabla u) - \lambda \delta^{\alpha p} |u|^{p-2} u = J_{\lambda}^{\alpha} \delta^{(\alpha-1)p} |u|^{p-2} u \quad \text{in } \mathcal{D}'(\Omega).$$

Corollary 2.1. *Under the same assumptions as in Theorem 2.1, there exists a constant $\lambda \in \mathbb{R}$ such that for $u \in W_{\alpha,0}^{1,p}(\Omega)$*

$$\int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx \geq \Lambda_{\alpha,p} \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx + \lambda \int_{\Omega} |u|^p \delta^{\alpha p} dx. \quad (2.2)$$

For each small $\eta > 0$, by Ω_{η} we denote a tubular neighborhood of $\partial\Omega$;

$$\Omega_{\eta} = \{x \in \Omega : \delta(x) = \operatorname{dist}(x, \partial\Omega) < \eta\}. \quad (2.3)$$

Then we have the following inequality of Hardy type which is crucial in the proof of Theorem 2.1.

Theorem 2.2. *Assume that Ω is a bounded domain of class C^2 in $\mathbb{R}^N$. Assume that $1 < p < \infty$ and $\alpha < 1 - 1/p$. Assume that η is a sufficiently small positive number. Then we have that for $u \in W_{\alpha,0}^{1,p}(\Omega)$*

$$\int_{\Omega_{\eta}} |\nabla u|^p \delta^{\alpha p} dx \geq \Lambda_{\alpha,p} \int_{\Omega_{\eta}} |u|^p \delta^{(\alpha-1)p} dx, \quad (2.4)$$

where $\Lambda_{\alpha,p}$ is defined by (2.1).

In [4] we have more precise estimate than (2.4).

Corollary 2.2. *Under the same assumptions as in Theorem 2.2, there exists a positive constant γ such that for $u \in W_{\alpha,0}^{1,p}(\Omega)$*

$$\int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx \geq \gamma \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx. \quad (2.5)$$

For any bounded domain $\Omega \subset \mathbb{R}^N$ we can prove the following:

Theorem 2.3. *Assume that Ω is a bounded domain of $\mathbb{R}^N$. Assume that $1 < p < \infty$ and $\alpha < 1 - 1/p$. Then the followings are equivalent with each other.*

1. *There exists a positive number γ such that the inequality (2.5) is valid for every $u \in W_{\alpha,0}^{1,p}(\Omega)$.*
2. *For a sufficiently small positive number η , there exists a positive number κ such that the inequality (2.4) with $\Lambda_{\alpha,p}$ replaced by κ is valid for every $u \in W_{\alpha,0}^{1,p}(\Omega)$.*

For the proofs of Theorem 2.2, Corollary 2.2 and Theorem 2.3, see in [4].

3 Proof of Theorem 2.1

In this section, we give the proof of Theorem 2.1.

3.1 Upper bound of J_λ^α

First, we prove the assertion 1 of Theorem 2.1.

Lemma 3.1. *Let $1 < p < \infty$ and $\alpha < 1 - 1/p$. For any $\varepsilon > 0$ and any $\eta > 0$ there exists a function $h \in W_{\alpha,0}^{1,p}((0,\eta))$ such that*

$$\int_0^\eta |h'(t)|^p t^{\alpha p} dt \leq (\Lambda_{\alpha,p} + \varepsilon) \int_0^\eta |h(t)|^p t^{(\alpha-1)p} dt, \quad (3.1)$$

where $\Lambda_{\alpha,p}$ is defined by (2.1).

Proof. Since the inequality (3.1) is invariant with respect to scaling, we may assume that $\eta = 2$. Put

$$h(t) = \begin{cases} t^\beta & \text{if } t \in (0, 1), \\ 2 - t & \text{if } t \in [1, 2] \end{cases}$$

with $\beta > 1 - \alpha - 1/p$. Then we see that $h \in W_{\alpha,0}^{1,p}((0,2))$,

$$\int_0^2 |h'(t)|^p t^{\alpha p} dt = \frac{\beta^p}{p(\beta - 1 + \alpha + 1/p)} + C_{\alpha,p} \quad (3.2)$$

and

$$\int_0^2 |h(t)|^p t^{(\alpha-1)p} dt = \frac{1}{p(\beta - 1 + \alpha + 1/p)} + D_{\alpha,p}, \quad (3.3)$$

where

$$C_{\alpha,p} = \int_1^2 t^{\alpha p} dt \quad \text{and} \quad D_{\alpha,p} = \int_1^2 (2 - t)^p t^{(\alpha-1)p} dt$$

are constants independent of β . It follows from (3.2) and (3.3) that

$$\frac{\int_0^2 |h'(t)|^p t^{\alpha p} dt}{\int_0^2 |h(t)|^p t^{(\alpha-1)p} dt} \longrightarrow \Lambda_{\alpha,p} \quad \text{as } \beta \rightarrow 1 - \alpha - \frac{1}{p} + 0,$$

which implies (3.1) with $\eta = 2$. Therefore we obtain the desired conclusion. $\square$

Lemma 3.2. *Let Ω be a bounded domain of class C^2 in $\mathbb{R}^N$. Let $1 < p < \infty$ and $\alpha < 1 - 1/p$. Then it holds that*

$$J_\lambda^\alpha \leq \Lambda_{\alpha,p} \quad (3.4)$$

for all $\lambda \in \mathbb{R}$.

Proof. Since the boundary $\partial\Omega$ is of class C^2 , there exists an $\eta_0 > 0$ such that for any $\eta \in (0, \eta_0)$ and every $x \in \Omega_\eta$ we have a unique point $\sigma(x) \in \partial\Omega$ satisfying $\delta(x) = |x - \sigma(x)|$. The mapping

$$\Omega_\eta \ni x \mapsto (\delta(x), \sigma(x)) = (t, \sigma) \in (0, \eta) \times \partial\Omega$$

is a C^2 diffeomorphism, and its inverse is given by

$$(0, \eta) \times \partial\Omega \ni (t, \sigma) \mapsto x(t, \sigma) = \sigma + t \cdot n(\sigma) \in \Omega_\eta,$$

where $n(\sigma)$ is the inward unit normal to $\partial\Omega$ at $\sigma \in \partial\Omega$. For each $t \in (0, \eta)$, the mapping

$$\partial\Omega \ni \sigma \mapsto \sigma_t(\sigma) = x(t, \sigma) \in \Sigma_t = \{x \in \Omega : \delta(x) = t\}$$

is also a C^2 diffeomorphism of $\partial\Omega$ onto Σ_t , and its Jacobian satisfies

$$|\text{Jac } \sigma_t(\sigma) - 1| \leq ct \quad \text{for any } \sigma \in \partial\Omega, \quad (3.5)$$

where c is a positive constant depending only on η_0 , $\partial\Omega$ and the choice of local coordinates. Since $n(\sigma)$ is orthogonal to Σ_t at $\sigma_t(\sigma) = \sigma + t \cdot n(\sigma) \in \Sigma_t$, it follows that for every integrable function v in Ω_η

$$\begin{aligned} \int_{\Omega_\eta} v(x) dx &= \int_0^\eta dt \int_{\Sigma_t} v(\sigma_t) d\sigma_t \\ &= \int_0^\eta dt \int_{\partial\Omega} v(x(t, \sigma)) |\text{Jac } \sigma_t(\sigma)| d\sigma, \end{aligned} \quad (3.6)$$

where $d\sigma$ and $d\sigma_t$ denote surface elements on $\partial\Omega$ and Σ_t , respectively. Hence (3.6) together with (3.5) implies that for every integrable function v in Ω_η

$$\int_0^\eta (1 - ct) dt \int_{\partial\Omega} |v(x(t, \sigma))| d\sigma \leq \int_{\Omega_\eta} |v(x)| dx \quad (3.7)$$

$$\leq \int_0^\eta (1 + ct) dt \int_{\partial\Omega} |v(x(t, \sigma))| d\sigma. \quad (3.8)$$

Let $\varepsilon > 0$, and let $\eta \in (0, \eta_0)$. Take $h \in W_{\alpha,0}^{1,p}((0, \eta))$ be a function satisfying (3.1). Put

$$u(x) = \begin{cases} h(\delta(x)) & \text{if } x \in \Omega_\eta, \\ 0 & \text{if } x \in \Omega \setminus \Omega_\eta. \end{cases} \quad (3.9)$$

Since $|\nabla u(x)| = |h'(\delta(x))|$ for $x \in \Omega_\eta$ by $|\nabla \delta(x)| = 1$, it follows from (3.8) that

$$\int_{\Omega_\eta} |\nabla u|^p \delta^{\alpha p} dx \leq (1 + c\eta) |\partial\Omega| \int_0^\eta |h'(t)|^p t^{\alpha p} dt, \quad (3.10)$$

which implies $u \in W_{\alpha,0}^{1,p}(\Omega)$ by $\text{supp } u \subset \Omega_\eta$. On the other hand, by (3.7) and (3.9) we have that

$$\int_{\Omega_\eta} |u|^p \delta^{(\alpha-1)p} dx \geq (1 - c\eta) |\partial\Omega| \int_0^\eta |h(t)|^p t^{(\alpha-1)p} dt. \quad (3.11)$$

Since $\text{supp } u \subset \Omega_\eta$, by combining (3.10), (3.11) and the estimate

$$\int_{\Omega_\eta} |u|^p \delta^{\alpha p} dx \leq \eta^p \int_{\Omega_\eta} |u|^p \delta^{(\alpha-1)p} dx,$$

we obtain that

$$\chi_\lambda^\alpha(u) \leq \frac{1 + c\eta}{1 - c\eta} \frac{\int_0^\eta |h'(t)|^p t^{\alpha p} dt}{\int_0^\eta |h(t)|^p t^{(\alpha-1)p} dt} + |\lambda| \eta^p.$$

This together with (3.1) implies that

$$J_\lambda^\alpha \leq \frac{1 + c\eta}{1 - c\eta} (\Lambda_{\alpha,p} + \varepsilon) + |\lambda| \eta^p. \quad (3.12)$$

Letting $\eta \rightarrow +0$ in (3.12), (3.4) follows. Therefore it concludes the proof. $\square$

Lemma 3.3. *Let Ω be a bounded domain of class C^2 in $\mathbb{R}^N$. Let $1 < p < \infty$ and $\alpha < 1 - 1/p$. Then there exists a $\lambda \in \mathbb{R}$ such that $J_\lambda^\alpha = \Lambda_{\alpha,p}$.*

Proof. Let $\eta > 0$ be a sufficiently small number as in Theorem 2.2. For any $u \in W_{\alpha,0}^{1,p}(\Omega) \setminus \{0\}$, by using Hardy's inequality (2.4) and the estimate

$$\int_{\Omega \setminus \Omega_\eta} |u|^p \delta^{(\alpha-1)p} dx \leq \eta^{-p} \int_{\Omega} |u|^p \delta^{\alpha p} dx,$$

we have that

$$\begin{aligned} \Lambda_{\alpha,p} \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx &= \Lambda_{\alpha,p} \int_{\Omega_\eta} |u|^p \delta^{(\alpha-1)p} dx + \Lambda_{\alpha,p} \int_{\Omega \setminus \Omega_\eta} |u|^p \delta^{(\alpha-1)p} dx \\ &\leq \int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx + \Lambda_{\alpha,p} \eta^{-p} \int_{\Omega} |u|^p \delta^{\alpha p} dx, \end{aligned}$$

which implies that

$$\chi_\lambda^\alpha(u) \geq \Lambda_{\alpha,p}$$

for $\lambda \leq -\Lambda_{\alpha,p} \eta^{-p}$. Consequently, it holds that $J_\lambda^\alpha \geq \Lambda_{\alpha,p}$ for $\lambda \leq -\Lambda_{\alpha,p} \eta^{-p}$. This together with (3.4) implies the desired conclusion. $\square$

Lemma 3.4. *Let Ω be a bounded domain of class C^2 in $\mathbb{R}^N$. Let $1 < p < \infty$ and $\alpha < 1 - 1/p$. Then the function $\lambda \mapsto J_\lambda^\alpha$ is Lipschitz continuous on $\mathbb{R}$.*

Proof. Let $\lambda, \bar{\lambda} \in \mathbb{R}$. Then it holds that for any $u \in W_{\alpha,0}^{1,p}(\Omega) \setminus \{0\}$

$$|\chi_\lambda^\alpha(u) - \chi_{\bar{\lambda}}^\alpha(u)| = |\lambda - \bar{\lambda}| \frac{\int_{\Omega} |u|^p \delta^{\alpha p} dx}{\int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx} \leq M^p |\lambda - \bar{\lambda}|,$$

where $M = \sup_{x \in \Omega} \delta(x)$ is a positive constant depending only on Ω . Hence we see that

$$|J_\lambda^\alpha - J_{\bar{\lambda}}^\alpha| \leq M^p |\lambda - \bar{\lambda}|$$

for $\lambda, \bar{\lambda} \in \mathbb{R}$. It completes the proof. $\square$

Proof of the assertion 1 of Theorem 2.1. By Lemma 3.3 and $\lim_{\lambda \rightarrow \infty} J_\lambda^\alpha = -\infty$, the set $\{\lambda \in \mathbb{R} : J_\lambda^\alpha = \Lambda_{\alpha,p}\}$ is non-empty and upper bounded. Hence the $\sup\{\lambda \in \mathbb{R} : J_\lambda^\alpha = \Lambda_{\alpha,p}\}$ exists finitely. Put

$$\lambda^* = \sup\{\lambda \in \mathbb{R} : J_\lambda^\alpha = \Lambda_{\alpha,p}\}. \quad (3.13)$$

Since the function $\lambda \mapsto J_\lambda^\alpha$ is non-increasing on $\mathbb{R}$, it follows from Lemma 3.2 and Lemma 3.3 that $J_\lambda^\alpha = \Lambda_{\alpha,p}$ for $\lambda < \lambda^*$ and $J_\lambda^\alpha < \Lambda_{\alpha,p}$ for $\lambda > \lambda^*$. Further, by Lemma 3.4 we have the equality $J_{\lambda^*}^\alpha = \Lambda_{\alpha,p}$. Therefore the assertion 1 of Theorem 2.1 is valid. $\square$

3.2 J_λ^α is not attained when $\lambda < \lambda^*$

Next, we prove the assertion 2 of Theorem 2.1.

Proof of the assertion 2 of Theorem 2.1. Suppose that for some $\lambda < \lambda^*$ the infimum J_λ^α in (1.5) is attained at an element $u \in W_{\alpha,0}^{1,p}(\Omega) \setminus \{0\}$. Then, by the assertion 1 of Theorem 2.1, we have that

$$\chi_\lambda^\alpha(u) = J_\lambda^\alpha = \Lambda_{\alpha,p} \quad (3.14)$$

and for $\lambda < \bar{\lambda} < \lambda^*$

$$\chi_{\bar{\lambda}}^{\alpha}(u) \geq J_{\bar{\lambda}}^{\alpha} = \Lambda_{\alpha,p}. \quad (3.15)$$

From (3.14) and (3.15) it follows that

$$(\bar{\lambda} - \lambda) \int_{\Omega} |u|^p \delta^{\alpha p} dx \leq 0.$$

Since $\bar{\lambda} - \lambda > 0$, we conclude that

$$\int_{\Omega} |u|^p \delta^{\alpha p} dx = 0,$$

which contradicts $u \neq 0$ in $W_{\alpha,0}^{1,p}(\Omega)$. Therefore it completes the proof. $\square$

3.3 Attainability of J_{λ}^{α} when $\lambda > \lambda^*$

At last, we prove the assertion 3 of Theorem 2.1.

Let $\{u_k\}$ be a minimizing sequence for the variational problem (1.5) normalized so that

$$\int_{\Omega} |u_k|^p \delta^{(\alpha-1)p} dx = 1 \quad \text{for all } k. \quad (3.16)$$

Since $\{u_k\}$ is bounded in $W_{\alpha,0}^{1,p}(\Omega)$, by taking a suitable subsequence, we may assume that there exists a $u \in W_{\alpha,0}^{1,p}(\Omega)$ such that

$$\nabla u_k \xrightarrow{\text{weak}} \nabla u \quad \text{in } (L^p(\Omega, \delta^{\alpha p}))^N, \quad (3.17)$$

$$u_k \xrightarrow{\text{weak}} u \quad \text{in } L^p(\Omega, \delta^{(\alpha-1)p}) \quad (3.18)$$

and

$$u_k \longrightarrow u \quad \text{in } L^p(\Omega, \delta^{\alpha p}) \quad (3.19)$$

by Hardy's inequality (2.5) and the compact embedding $W_{\alpha,0}^{1,p}(\Omega) \hookrightarrow L^p(\Omega, \delta^{\alpha p})$.

Under these preparation we establish the properties of concentration and compactness for the minimizing sequence, respectively.

Proposition 3.1. *Let Ω be a bounded domain of class C^2 in $\mathbb{R}^N$. Let $1 < p < \infty$ and $\alpha < 1 - 1/p$. Let $\lambda \in \mathbb{R}$. Let $\{u_k\}$ be a minimizing sequence for (1.5) satisfying (3.16), (3.17), (3.18) and (3.19) with $u = 0$. Then it holds that*

$$\nabla u_k \longrightarrow 0 \quad \text{in } (L_{\text{loc}}^p(\Omega))^N \quad (3.20)$$

and

$$J_{\lambda}^{\alpha} = \Lambda_{\alpha,p}. \quad (3.21)$$

Proof. Let $\eta > 0$ be a sufficiently small number as in Theorem 2.2. By Hardy's inequality (2.4) and (3.16) we have that

$$\begin{aligned} \int_{\Omega_{\eta}} |\nabla u_k|^p \delta^{\alpha p} dx &\geq \Lambda_{\alpha,p} \int_{\Omega_{\eta}} |u_k|^p \delta^{(\alpha-1)p} dx \\ &= \Lambda_{\alpha,p} \left(1 - \int_{\Omega \setminus \Omega_{\eta}} |u_k|^p \delta^{(\alpha-1)p} dx \right), \end{aligned}$$

and so

$$\begin{aligned} \chi_\lambda^\alpha(u_k) &\geq \Lambda_{\alpha,p} \left(1 - \int_{\Omega \setminus \Omega_\eta} |u_k|^p \delta^{(\alpha-1)p} dx \right) \\ &\quad + \int_{\Omega \setminus \Omega_\eta} |\nabla u_k|^p \delta^{\alpha p} dx - \lambda \int_{\Omega} |u_k|^p \delta^{\alpha p} dx. \end{aligned} \quad (3.22)$$

Since

$$\int_{\Omega \setminus \Omega_\eta} |u_k|^p \delta^{(\alpha-1)p} dx \leq \eta^{-p} \int_{\Omega} |u_k|^p \delta^{\alpha p} dx,$$

it follows from (3.19) with $u = 0$ that

$$\lim_{k \rightarrow \infty} \int_{\Omega \setminus \Omega_\eta} |u_k|^p \delta^{(\alpha-1)p} dx = 0. \quad (3.23)$$

Hence, by (3.22), (3.23) and (3.19) with $u = 0$, we obtain that

$$\limsup_{k \rightarrow \infty} \int_{\Omega \setminus \Omega_\eta} |\nabla u_k|^p \delta^{\alpha p} dx \leq J_\lambda^\alpha - \Lambda_{\alpha,p}.$$

Since $J_\lambda^\alpha - \Lambda_{\alpha,p} \leq 0$ by Lemma 3.2, we conclude that

$$\lim_{k \rightarrow \infty} \int_{\Omega \setminus \Omega_\eta} |\nabla u_k|^p \delta^{\alpha p} dx = 0, \quad (3.24)$$

and so

$$\lim_{k \rightarrow \infty} \int_{\Omega \setminus \Omega_\eta} |\nabla u_k|^p dx = 0.$$

This shows (3.20). Moreover, letting $k \rightarrow \infty$ in (3.22), it follows from (3.23), (3.24) and (3.19) with $u = 0$ that

$$J_\lambda^\alpha \geq \Lambda_{\alpha,p}.$$

This together with Lemma 3.2 implies (3.21). Consequently it completes the proof. $\square$

Proposition 3.2. *Let Ω be a bounded domain of class C^2 in $\mathbb{R}^N$. Let $1 < p < \infty$ and $\alpha < 1 - 1/p$. Let $\lambda \in \mathbb{R}$. Let $\{u_k\}$ be a minimizing sequence for (1.5) satisfying (3.16), (3.17), (3.18) and (3.19) with $u \neq 0$. Then it holds that*

$$J_\lambda^\alpha = \min(\Lambda_{\alpha,p}, \chi_\lambda^\alpha(u)). \quad (3.25)$$

In addition, if $J_\lambda^\alpha < \Lambda_{\alpha,p}$, then it holds that

$$J_\lambda^\alpha = \chi_\lambda^\alpha(u), \quad (3.26)$$

namely u is a minimizer for (1.5), and

$$u_k \longrightarrow u \quad \text{in } W_{\alpha,0}^{1,p}(\Omega). \quad (3.27)$$

Proof. Let $\eta > 0$ be a sufficiently small number as in Theorem 2.2. Then we have (3.22) by the same arguments as in the proof of Proposition 3.1. By the estimate

$$\int_{\Omega \setminus \Omega_\eta} |u_k - u|^p \delta^{(\alpha-1)p} dx \leq \eta^{-p} \int_{\Omega} |u_k - u|^p \delta^{\alpha p} dx,$$

(3.19) implies that

$$\lim_{k \rightarrow \infty} \int_{\Omega \setminus \Omega_\eta} |u_k|^p \delta^{(\alpha-1)p} dx = \int_{\Omega \setminus \Omega_\eta} |u|^p \delta^{(\alpha-1)p} dx. \quad (3.28)$$

Since it follows from (3.17) that $\nabla u_k \rightharpoonup \nabla u$ weakly in $(L^p(\Omega \setminus \Omega_\eta, \delta^{\alpha p}))^N$, by weakly lower semi-continuity of the L^p -norm, we see that

$$\begin{aligned} \liminf_{k \rightarrow \infty} \int_{\Omega \setminus \Omega_\eta} |\nabla u_k|^p \delta^{\alpha p} dx &\geq \left(\liminf_{k \rightarrow \infty} \|\nabla u_k\|_{L^p(\Omega \setminus \Omega_\eta, \delta^{\alpha p})} \right)^p \\ &\geq \|\nabla u\|_{L^p(\Omega \setminus \Omega_\eta, \delta^{\alpha p})}^p \\ &= \int_{\Omega \setminus \Omega_\eta} |\nabla u|^p \delta^{\alpha p} dx. \end{aligned} \quad (3.29)$$

Hence, by letting $k \rightarrow \infty$ in (3.22), from (3.19), (3.28) and (3.29) it follows that

$$\begin{aligned} J_\lambda^\alpha &\geq \Lambda_{\alpha,p} \left(1 - \int_{\Omega \setminus \Omega_\eta} |u|^p \delta^{(\alpha-1)p} dx \right) \\ &\quad + \int_{\Omega \setminus \Omega_\eta} |\nabla u|^p \delta^{\alpha p} dx - \lambda \int_{\Omega} |u|^p \delta^{\alpha p} dx. \end{aligned} \quad (3.30)$$

Letting $\eta \rightarrow +0$ in (3.30), we obtain that

$$\begin{aligned} J_\lambda^\alpha &\geq \Lambda_{\alpha,p} \left(1 - \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx \right) \\ &\quad + \int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx - \lambda \int_{\Omega} |u|^p \delta^{\alpha p} dx. \end{aligned} \quad (3.31)$$

Since it holds that

$$0 < \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx \leq \liminf_{k \rightarrow \infty} \int_{\Omega} |u_k|^p \delta^{(\alpha-1)p} dx = 1 \quad (3.32)$$

by $u \neq 0$, (3.16), (3.18) and weakly lower semi-continuity of the L^p -norm, we have from (3.31) and (3.32) that

$$\begin{aligned} J_\lambda^\alpha &\geq \Lambda_{\alpha,p} \left(1 - \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx \right) + \chi_\lambda^\alpha(u) \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx \\ &\geq \min(\Lambda_{\alpha,p}, \chi_\lambda^\alpha(u)). \end{aligned} \quad (3.33)$$

This together with Lemma 3.2 implies (3.25). Moreover, by (3.25) and (3.33), we conclude that

$$J_\lambda^\alpha = \Lambda_{\alpha,p} \left(1 - \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx \right) + \chi_\lambda^\alpha(u) \int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx. \quad (3.34)$$

In addition, if $J_\lambda^\alpha < \Lambda_{\alpha,p}$, then $J_\lambda^\alpha = \chi_\lambda^\alpha(u)$ by (3.25), and so, it follows from (3.34) and (3.16) that

$$\int_{\Omega} |u|^p \delta^{(\alpha-1)p} dx = 1 = \lim_{k \rightarrow \infty} \int_{\Omega} |u_k|^p \delta^{(\alpha-1)p} dx. \quad (3.35)$$

(3.18) and (3.35) imply that

$$u_k \longrightarrow u \quad \text{in } L^p(\Omega, \delta^{(\alpha-1)p}). \quad (3.36)$$

Further, by (3.16), (3.19), (3.26) and (3.35), we obtain that

$$\begin{aligned}\int_{\Omega} |\nabla u_k|^p \delta^{\alpha p} dx &= \chi_{\lambda}^{\alpha}(u_k) + \lambda \int_{\Omega} |u_k|^p \delta^{\alpha p} dx \\ &\longrightarrow \chi_{\lambda}^{\alpha}(u) + \lambda \int_{\Omega} |u|^p \delta^{\alpha p} dx = \int_{\Omega} |\nabla u|^p \delta^{\alpha p} dx.\end{aligned}$$

This together with (3.17) implies that

$$\nabla u_k \longrightarrow \nabla u \quad \text{in } (L^p(\Omega, \delta^{\alpha p}))^N, \quad (3.37)$$

which shows (3.27). Consequently it completes the proof. $\square$

Proof of the assertion 3 of Theorem 2.1. Let $\lambda > \lambda^*$. Then $J_{\lambda}^{\alpha} < \Lambda_{\alpha,p}$ by the assertion 1 of Theorem 2.1. Let $\{u_k\}$ be a minimizing sequence for (1.5) satisfying (3.16), (3.17), (3.18) and (3.19). Then we see that $u \neq 0$ by Proposition 3.1. Therefore, by applying Proposition 3.2, we conclude that $\chi_{\lambda}^{\alpha}(u) = J_{\lambda}^{\alpha}$, namely u is a minimizer for (1.5). It finishes the proof. $\square$

References

- [1] Adimurthi, N. Nirmalendu, M. Chaudhuri, M. Ramaswamy, An improved Hardy-Sobolev inequality and its application, *Proceedings of the American Mathematical Society*, **Vol.130**, No.2 (2001), 489–505.
- [2] H.Ando, T.Horiuchi, Missing terms in the weighted Hardy-Sobolev inequalities and its application, *Kyoto J. Math.* **Vol.52**, No.4 (2012), 759–796.
- [3] H. Ando, T. Horiuchi, General Weighted Hardy’s Inequalities with compact perturbations, in preparation.
- [4] H. Ando, T. Horiuchi, X. Liu, One dimensional Weighted Hardy’s Inequalities and application, *Journal Mathematical Inequality*, to appear.
- [5] H. Ando, T. Horiuchi, E. Nakai, Weighted Hardy inequalities with infinitely many sharp missing terms, *Mathematical Journal of Ibaraki University*, **Vol.46** (2014), 9–30.
- [6] H.Brezis, M.Marcus, Hardy’s inequalities revisited, *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* 4^e série, tome **25**, No.1-2 (1997), 217–237.
- [7] Z. Chen, Y. Shen, Sharp Hardy-Sobolev inequalities with general weights and remainder terms, *Journal of inequalities and applications*, (2009), Article ID 419845, 24pages doi:10.1155/2009/419845.
- [8] E. B. Davis, The Hardy constant, *Quart. J. Math. Oxford*, (2) **Vol.46** (1995), 417–431.
- [9] A. Detalla, T. Horiuchi, H. Ando, Missing terms in Hardy-Sobolev inequalities and its application, *Far East Journal of Mathematical Sciences*, **Vol.14**, No.3 (2004), 333–359.
- [10] T. Horiuchi, Hardy’s inequalities with non-doubling weights and sharp remainders, in preparation.

- [11] A. Kufner, B. Opic, Hardy-type Inequalities, *Pitman Research Notes in Mathematics series*, **Vol.219**, London, Longman Group UK Limited, (1990).
- [12] E. H. Lieb and M. Loss, Analysis, *American Mathematical Society*, (2001).
- [13] M.Marcus, V.J.Mizel, Y.Pinchover, On the best constant for Hardy's inequality in $\mathbb{R}^n$, *Trans. Amer. Math. Soc.*, **Vol.350**, No.8 (1998), 3237–3255.
- [14] T. Matskewich, P. E. Sobolevskii, The best possible constant in a generalized Hardy's inequality for convex domains in $\mathbb{R}^n$, *Nonlinear Analysis TMA*, **Vol.28** (1997), 1601–1610.
- [15] V. G. Maz'ja, Sobolev spaces (2nd edition), Springer, (2011).

Hiroshi Ando:

Department of Mathematics, Faculty of Science, Ibaraki University,
Mito, Ibaraki, 310-8512, Japan;
hiroshi.ando.math@vc.ibaraki.ac.jp

Toshio Horiuchi:

Department of Mathematics, Faculty of Science, Ibaraki University,
Mito, Ibaraki, 310-8512, Japan;
toshio.horiuchi.math@vc.ibaraki.ac.jp