

A Note on the Gaussian Minimum Conjecture

Yang-Fan Zhong, Ting Ma, Ze-Chun Hu*

College of Mathematics, Sichuan University, China

Abstract

Let $n \geq 2$ and $(X_i, 1 \leq i \leq n)$ be a centered Gaussian random vector. The Gaussian minimum conjecture says that $E(\min_{1 \leq i \leq n} |X_i|) \geq E(\min_{1 \leq i \leq n} |Y_i|)$, where $Y_1, \dots, Y_n$ are independent centered Gaussian random variables with $E(X_i^2) = E(Y_i^2)$ for any $i = 1, \dots, n$. In this note, we will show that this conjecture holds if and only if $n = 2$.

MSC: Primary 60E15; Secondary 62H12

Keywords: Gaussian Minimum Conjecture; Gaussian random vector

1 Introduction and main result

Let $n \geq 2$ and $(X_i, 1 \leq i \leq n)$ be a centered Gaussian random vector. The well-known Šidák's inequality ([7], [1]) says that

$$\mathbb{E} \left(\max_{1 \leq i \leq n} |X_i| \right) \leq \mathbb{E} \left(\max_{1 \leq i \leq n} |Y_i| \right), \quad (1.1)$$

where $Y_1, \dots, Y_n$ are independent centered Gaussian random variables with $E(X_i^2) = E(Y_i^2)$ for any $i = 1, \dots, n$.

If we replace 'max' by 'min' in Šidák's inequality, Gordon et al. proved among other things that ([2], [3])

$$\mathbb{E} \left(\min_{1 \leq i \leq n} |X_i| \right) \geq \frac{1}{2} \mathbb{E} \left(\min_{1 \leq i \leq n} |Y_i| \right). \quad (1.2)$$

Note that in [2] and [3], the authors proved the inequality (1.2) without any condition on the joint distribution of $(X_1, \dots, X_n)$.

Wenbo V. Li and Qiman Shao conjectured that when $(X_i, 1 \leq i \leq n)$ is a centered Gaussian random vector, $\frac{1}{2}$ in the inequality (1.2) can be removed, i.e.

$$\mathbb{E} \left(\min_{1 \leq i \leq n} |X_i| \right) \geq \mathbb{E} \left(\min_{1 \leq i \leq n} |Y_i| \right), \quad (1.3)$$

which is called the *Gaussian Minimum Conjecture* (see [4], [8]).

Now we state the main result of this note.

Theorem 1.1 *The Gaussian Minimum Conjecture holds if and only if $n = 2$.*

*Corresponding author: zchu@scu.edu.cn

Remark 1.2 Professor Qiman Shao told the third author that the Gaussian Minimum Conjecture for $n = 2$ can be easily proved based on the following fact:

$$\begin{aligned}
P(\min(|X_1|, |X_2|) > x) &= 1 - P(|X_1| \leq x) - P(|X_2| \leq x) + P(|X_1| \leq x, |X_2| \leq x) \\
&\geq 1 - P(|X_1| \leq x) - P(|X_2| \leq x) + P(|X_1| \leq x)P(|X_2| \leq x) \\
&= 1 - P(|Y_1| \leq x) - P(|Y_2| \leq x) + P(|Y_1| \leq x)P(|Y_2| \leq x) \\
&= 1 - P(|Y_1| \leq x) - P(|Y_2| \leq x) + P(|Y_1| \leq x, |Y_2| \leq x) \\
&= P(\min(|Y_1|, |Y_2|) > x),
\end{aligned}$$

where the Gaussian correlation inequality was used. In next section, we will give a different proof for $n = 2$ based on an explicit formula for $E(\min(|X_1|, |X_2|))$ (see (2.14) below).

As to the minimum value point of $\mathbb{E}(\min_{1 \leq i \leq n} |X_i|)$ with $E(X_i^2) = 1, \forall i = 1, \dots, n$, for $n \geq 3$, we refer to [5, Conjecture 5.1]. In addition, we would like to mention a related conjecture, the Gaussian random vector maximum (GRVM) conjecture, which asserts that among all centered Gaussian random vectors $X = (X_1, \dots, X_n)$ with $E[X_i^2] = 1, 1 \leq i \leq n$, the expectation $E[\max(X_1, \dots, X_n)]$ is maximal if and only if all off-diagonal elements of the covariance matrix equal $-\frac{1}{n-1}$, where $n \geq 3$. The paper [9] proved the GRVM conjecture for $n = 3, 4$.

2 Proof of Theorem 1.1

2.1 Necessity

In this part, we will show that if $n \geq 3$, the inequality (1.3) does not hold for some n -dimensional centered Gaussian random vector $(X_1, \dots, X_n)$ and independent centered Gaussian random variables $Y_1, \dots, Y_n$ with $E(X_i^2) = E(Y_i^2), \forall i = 1, \dots, n$.

(i) $n = 3$. R. van Handel gave the following counterexample for $n = 3$ (see [6]). Let Y_1, Y_2, Y_3 be three independent standard Gaussian random variables. Define $X_1 = (Y_1 - Y_2)/\sqrt{2}, X_2 = (Y_2 - Y_3)/\sqrt{2}, X_3 = (Y_3 - Y_1)/\sqrt{2}$. R. van Handel checked numerically that $0.32 \sim \mathbb{E}(\min_{1 \leq i \leq 3} |X_i|) < \mathbb{E}(\min_{1 \leq i \leq 3} |Y_i|) \sim 0.34$, which tell us that (1.3) does not hold for $n = 3$. In fact, we can give the exact values of $\mathbb{E}(\min_{1 \leq i \leq 3} |X_i|)$ and $\mathbb{E}(\min_{1 \leq i \leq 3} |Y_i|)$.

At first, we calculate $\mathbb{E}(\min_{1 \leq i \leq 3} |Y_i|)$. The density function $p_1(x, y, z)$ of (Y_1, Y_2, Y_3) can be expressed by

$$p_1(x, y, z) = \frac{1}{(2\pi)^{3/2}} e^{-\frac{x^2+y^2+z^2}{2}}.$$

By the symmetry, we have

$$\begin{aligned}
\mathbb{E}\left(\min_{1 \leq i \leq 3} |Y_i|\right) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (|x| \wedge |y| \wedge |z|) \frac{1}{(2\pi)^{3/2}} e^{-\frac{x^2+y^2+z^2}{2}} dx dy dz \\
&= \frac{8}{(2\pi)^{3/2}} \int_0^{\infty} \int_0^{\infty} \int_0^{\infty} (x \wedge y \wedge z) e^{-\frac{x^2+y^2+z^2}{2}} dx dy dz \\
&= \frac{8 \cdot 3!}{(2\pi)^{3/2}} \int_0^{\infty} \left(\int_0^z e^{-\frac{y^2+z^2}{2}} \left(\int_0^y x e^{-\frac{x^2}{2}} dx \right) dy \right) dz \\
&= \frac{48}{(2\pi)^{3/2}} \int_0^{\infty} \left(\int_0^z e^{-\frac{y^2+z^2}{2}} \left(1 - e^{-\frac{y^2}{2}} \right) dy \right) dz \\
&= \frac{12\sqrt{2}}{\pi^{3/2}} \left[\int_0^{\infty} \left(\int_0^z e^{-\frac{y^2+z^2}{2}} dy \right) dz - \int_0^{\infty} \left(\int_0^z e^{-\frac{2y^2+z^2}{2}} dy \right) dz \right]. \tag{2.1}
\end{aligned}$$

Define a function

$$F(a) := \int_0^{\infty} \left(\int_0^z e^{-\frac{ay^2+z^2}{2}} dy \right) dz, \quad a > 0,$$

and a set

$$D := \{(y, z) \in \mathbf{R}^2 : 0 \leq y \leq z\}.$$

Define a transformation

$$(y, z) := T(u, v) = (u/\sqrt{a}, v).$$

Denote by D_T the original image of D under T . Then we have

$$D_T = \{(u, v) \in \mathbf{R}^2 : T(u, v) \in D\} = \{(u, v) \in \mathbf{R}^2 : 0 \leq u/\sqrt{a} \leq v\}.$$

Now, we have

$$\begin{aligned} F(a) &= \iint_D e^{-\frac{ay^2+z^2}{2}} dydz = \frac{1}{\sqrt{a}} \iint_{D_T} e^{-\frac{u^2+v^2}{2}} dudv \\ &= \frac{1}{\sqrt{a}} \cdot \left(\frac{\pi}{2} - \arctan \frac{1}{\sqrt{a}} \right) \cdot \int_0^\infty e^{-\frac{r^2}{2}} r dr \\ &= \frac{1}{\sqrt{a}} \arctan \sqrt{a}. \end{aligned} \tag{2.2}$$

By (2.1) and (2.2), we get

$$\begin{aligned} \mathbb{E} \left(\min_{1 \leq i \leq 3} |Y_i| \right) &= \frac{12\sqrt{2}}{\pi^{3/2}} (F(1) - F(2)) \\ &= \frac{12\sqrt{2}}{\pi^{3/2}} \left(\arctan 1 - \frac{1}{\sqrt{2}} \arctan \sqrt{2} \right) \\ &= \frac{12}{\pi^{3/2}} \left(\frac{\sqrt{2}\pi}{4} - \arctan \sqrt{2} \right). \end{aligned} \tag{2.3}$$

Next, we come to calculate $\mathbb{E}(\min_{1 \leq i \leq 3} |X_i|)$. Note that $X_3 = -(X_1 + X_2)$, and the covariance matrix of (X_1, X_2) is $\Sigma = \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$. It follows that $\det \Sigma = \frac{3}{4}$, and $\Sigma^{-1} = \begin{pmatrix} 4/3 & 2/3 \\ 2/3 & 4/3 \end{pmatrix}$. Then the density function $p_2(x, y)$ of (X_1, X_2) can be expressed by

$$p_2(x, y) = \frac{1}{2\pi\sqrt{3/4}} e^{-\frac{\frac{4}{3}(x^2+y^2+xy)}{2}} = \frac{1}{\sqrt{3}\pi} e^{-\frac{2(x^2+y^2+xy)}{3}}.$$

By the symmetry, we have

$$\begin{aligned}
\mathbb{E} \left(\min_{1 \leq i \leq 3} |X_i| \right) &= \frac{1}{\sqrt{3}\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (|x| \wedge |y| \wedge |x+y|) e^{-\frac{2(x^2+y^2+xy)}{3}} dx dy \\
&= \frac{1}{\sqrt{3}\pi} \left[\int_0^{\infty} \int_0^{\infty} (x \wedge y) e^{-\frac{2(x^2+y^2+xy)}{3}} dx dy + \int_{-\infty}^0 \int_{-\infty}^0 ((-x) \wedge (-y)) e^{-\frac{2(x^2+y^2+xy)}{3}} dx dy \right. \\
&\quad + \int_{-\infty}^0 \int_0^{\infty} ((-x) \wedge y \wedge |x+y|) e^{-\frac{2(x^2+y^2+xy)}{3}} dx dy \\
&\quad \left. + \int_0^{\infty} \int_{-\infty}^0 (x \wedge (-y) \wedge |x+y|) e^{-\frac{2(x^2+y^2+xy)}{3}} dx dy \right] \\
&= \frac{2}{\sqrt{3}\pi} \left[\int_0^{\infty} \int_0^{\infty} (x \wedge y) e^{-\frac{2(x^2+y^2+xy)}{3}} dx dy + \int_0^{\infty} \int_0^{\infty} (x \wedge y \wedge |x-y|) e^{-\frac{2(x^2+y^2-xy)}{3}} dx dy \right] \\
&= \frac{4}{\sqrt{3}\pi} \left[\int_0^{\infty} \left(\int_0^y x e^{-\frac{2(x^2+y^2+xy)}{3}} dx \right) dy + \int_0^{\infty} \left(\int_0^y (x \wedge (y-x)) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \right] \\
&= \frac{4}{\sqrt{3}\pi} \left[\int_0^{\infty} \left(\int_0^y x e^{-\frac{2(x^2+y^2+xy)}{3}} dx \right) dy + \int_0^{\infty} \left(\int_0^{y/2} x e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \right. \\
&\quad \left. + \int_0^{\infty} \left(\int_{y/2}^y (y-x) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \right] \\
&= \frac{4}{\sqrt{3}\pi} \left[\int_0^{\infty} \left(\int_0^y x e^{-\frac{2(x^2+y^2+xy)}{3}} dx \right) dy + \int_0^{\infty} \left(\int_0^{y/2} x e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \right. \\
&\quad \left. + \int_0^{\infty} \left(\int_{y/2}^y y e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy - \int_0^{\infty} \left(\int_{y/2}^y x e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \right] \\
&=: \frac{4}{\sqrt{3}\pi} [G_1 + G_2 + G_3 - G_4], \tag{2.4}
\end{aligned}$$

where

$$\begin{aligned}
G_1 &:= \int_0^{\infty} \left(\int_0^y x e^{-\frac{2(x^2+y^2+xy)}{3}} dx \right) dy, & G_2 &:= \int_0^{\infty} \left(\int_0^{y/2} x e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy, \\
G_3 &:= \int_0^{\infty} \left(\int_{y/2}^y y e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy, & G_4 &:= \int_0^{\infty} \left(\int_{y/2}^y x e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy.
\end{aligned}$$

We have

$$\begin{aligned}
G_1 &= \int_0^{\infty} \left(\int_0^y \left(x + \frac{y}{2} \right) e^{-\frac{2(x^2+y^2+xy)}{3}} dx \right) dy - \frac{1}{2} \int_0^{\infty} \left(\int_0^y \left(y + \frac{x}{2} \right) e^{-\frac{2(x^2+y^2+xy)}{3}} dx \right) dy \\
&\quad + \frac{1}{4} \int_0^{\infty} \left(\int_0^y x e^{-\frac{2(x^2+y^2+xy)}{3}} dx \right) dy \\
&= \frac{G_1}{4} + \int_0^{\infty} \left(\int_0^y \left(x + \frac{y}{2} \right) e^{-\frac{2\left(x+\frac{y}{2}\right)^2 + \frac{3}{2}y^2}} dx \right) dy - \frac{1}{2} \int_0^{\infty} \left(\int_x^{\infty} \left(y + \frac{x}{2} \right) e^{-\frac{2\left(y+\frac{x}{2}\right)^2 + \frac{3}{2}x^2}} dy \right) dx \\
&= \frac{G_1}{4} + \int_0^{\infty} e^{-\frac{y^2}{2}} \left(\int_{y/2}^{3y/2} u e^{-\frac{2u^2}{3}} du \right) dy - \frac{1}{2} \int_0^{\infty} e^{-\frac{x^2}{2}} \left(\int_{3x/2}^{\infty} v e^{-\frac{2v^2}{3}} dv \right) dx \\
&= \frac{G_1}{4} + \frac{3}{4} \left[\int_0^{\infty} e^{-\frac{y^2}{2}} \left(e^{-\frac{y^2}{6}} - e^{-\frac{3y^2}{2}} \right) dy - \frac{1}{2} \int_0^{\infty} e^{-\frac{x^2}{2}} \cdot e^{-\frac{3x^2}{2}} dx \right] \\
&= \frac{G_1}{4} + \frac{3}{8} \left(2 \int_0^{\infty} e^{-\frac{2y^2}{3}} dy - 3 \int_0^{\infty} e^{-2y^2} dy \right).
\end{aligned}$$

It follows that

$$G_1 = \frac{1}{2} \left(2 \int_0^\infty e^{-\frac{2y^2}{3}} dy - 3 \int_0^\infty e^{-2y^2} dy \right). \quad (2.5)$$

We have

$$\begin{aligned} G_2 &= \int_0^\infty \left(\int_0^{y/2} \left(x - \frac{y}{2} \right) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy + \frac{1}{2} \int_0^\infty \left(\int_0^{y/2} \left(y - \frac{x}{2} \right) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \\ &\quad + \frac{1}{4} \int_0^\infty \left(\int_0^{y/2} x e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \\ &= \frac{G_2}{4} + \int_0^\infty \left(\int_0^{y/2} \left(x - \frac{y}{2} \right) e^{-\frac{2\left(x-\frac{y}{2}\right)^2 + \frac{3y^2}{2}}{3}} dx \right) dy + \frac{1}{2} \int_0^\infty \left(\int_{2x}^\infty \left(y - \frac{x}{2} \right) e^{-\frac{2\left(y-\frac{x}{2}\right)^2 + \frac{3x^2}{2}}{3}} dy \right) dx \\ &= \frac{G_2}{4} + \int_0^\infty e^{-\frac{y^2}{2}} \left(\int_{-y/2}^0 u e^{-\frac{2u^2}{3}} du \right) dy + \frac{1}{2} \int_0^\infty e^{-\frac{x^2}{2}} \left(\int_{3x/2}^\infty v e^{-\frac{2v^2}{3}} dv \right) dx \\ &= \frac{G_2}{4} + \frac{3}{4} \left[\int_0^\infty e^{-\frac{y^2}{2}} \left(e^{-\frac{y^2}{6}} - 1 \right) dy + \frac{1}{2} \int_0^\infty e^{-\frac{x^2}{2}} \cdot e^{-\frac{3x^2}{2}} dx \right] \\ &= \frac{G_2}{4} + \frac{3}{8} \left(2 \int_0^\infty e^{-\frac{2y^2}{3}} dy - 2 \int_0^\infty e^{-\frac{y^2}{2}} dy + \int_0^\infty e^{-2x^2} dx \right). \end{aligned}$$

It follows that

$$G_2 = \frac{1}{2} \left(2 \int_0^\infty e^{-\frac{2y^2}{3}} dy - 2 \int_0^\infty e^{-\frac{y^2}{2}} dy + \int_0^\infty e^{-2x^2} dx \right). \quad (2.6)$$

We have

$$\begin{aligned} G_3 &= \int_0^\infty \left(\int_{y/2}^y \left(y - \frac{x}{2} \right) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy + \frac{1}{2} \int_0^\infty \left(\int_{y/2}^y \left(x - \frac{y}{2} \right) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \\ &\quad + \frac{1}{4} \int_0^\infty \left(\int_{y/2}^y y e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \\ &= \frac{G_3}{4} + \int_0^\infty \left(\int_x^{2x} \left(y - \frac{x}{2} \right) e^{-\frac{2\left(y-\frac{x}{2}\right)^2 + \frac{3x^2}{2}}{3}} dy \right) dx + \frac{1}{2} \int_0^\infty \left(\int_{y/2}^y \left(x - \frac{y}{2} \right) e^{-\frac{2\left(x-\frac{y}{2}\right)^2 + \frac{3y^2}{2}}{3}} dx \right) dy \\ &= \frac{G_3}{4} + \int_0^\infty e^{-\frac{x^2}{2}} \left(\int_{x/2}^{3x/2} u e^{-\frac{2u^2}{3}} du \right) dx + \frac{1}{2} \int_0^\infty e^{-\frac{y^2}{2}} \left(\int_0^{y/2} v e^{-\frac{2v^2}{3}} dv \right) dy \\ &= \frac{G_3}{4} + \frac{3}{4} \left[\int_0^\infty e^{-\frac{x^2}{2}} \left(e^{-\frac{x^2}{6}} - e^{-\frac{3x^2}{2}} \right) dx + \frac{1}{2} \int_0^\infty e^{-\frac{y^2}{2}} \left(1 - e^{-\frac{y^2}{6}} \right) dy \right] \\ &= \frac{G_3}{4} + \frac{3}{8} \left(\int_0^\infty e^{-\frac{y^2}{2}} dy + \int_0^\infty e^{-\frac{2x^2}{3}} dx - 2 \int_0^\infty e^{-2x^2} dx \right). \end{aligned}$$

It follows that

$$G_3 = \frac{1}{2} \left(\int_0^\infty e^{-\frac{y^2}{2}} dy + \int_0^\infty e^{-\frac{2x^2}{3}} dx - 2 \int_0^\infty e^{-2x^2} dx \right). \quad (2.7)$$

We have

$$\begin{aligned}
G_4 &= \int_0^\infty \left(\int_{y/2}^y \left(x - \frac{y}{2} \right) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy + \frac{1}{2} \int_0^\infty \left(\int_{y/2}^y \left(y - \frac{x}{2} \right) e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \\
&\quad + \frac{1}{4} \int_0^\infty \left(\int_{y/2}^y x e^{-\frac{2(x^2+y^2-xy)}{3}} dx \right) dy \\
&= \frac{G_4}{4} + \int_0^\infty \left(\int_{y/2}^y \left(x - \frac{y}{2} \right) e^{-\frac{2(x-\frac{y}{2})^2 + \frac{3y^2}{2}}{3}} dx \right) dy + \frac{1}{2} \int_0^\infty \left(\int_x^{2x} \left(y - \frac{x}{2} \right) e^{-\frac{2(y-\frac{x}{2})^2 + \frac{3x^2}{2}}{3}} dy \right) dx \\
&= \frac{G_4}{4} + \int_0^\infty e^{-\frac{y^2}{2}} \left(\int_0^{y/2} u e^{-\frac{2u^2}{3}} du \right) dy + \frac{1}{2} \int_0^\infty e^{-\frac{x^2}{2}} \left(\int_{x/2}^{3x/2} v e^{-\frac{2v^2}{3}} dv \right) dx \\
&= \frac{G_4}{4} + \frac{3}{4} \left[\int_0^\infty e^{-\frac{y^2}{2}} \left(1 - e^{-\frac{y^2}{6}} \right) dy + \frac{1}{2} \int_0^\infty e^{-\frac{x^2}{2}} \left(e^{-\frac{x^2}{6}} - e^{-\frac{3x^2}{2}} \right) dx \right] \\
&= \frac{G_4}{4} + \frac{3}{8} \left(2 \int_0^\infty e^{-\frac{y^2}{2}} dy - \int_0^\infty e^{-\frac{2y^2}{3}} dy - \int_0^\infty e^{-2y^2} dy \right).
\end{aligned}$$

It follows that

$$G_4 = \frac{1}{2} \left(2 \int_0^\infty e^{-\frac{y^2}{2}} dy - \int_0^\infty e^{-\frac{2y^2}{3}} dy - \int_0^\infty e^{-2y^2} dy \right). \quad (2.8)$$

By (2.4)-(2.8), we obtain

$$\begin{aligned}
\mathbb{E} \left(\min_{1 \leq i \leq 3} |X_i| \right) &= \frac{6}{\sqrt{3}\pi} \left(2 \int_0^\infty e^{-\frac{2y^2}{3}} dy - \int_0^\infty e^{-2y^2} dy - \int_0^\infty e^{-\frac{y^2}{2}} dy \right) \\
&= \frac{6}{\sqrt{3}\pi} \cdot \frac{\sqrt{2\pi}}{2} \left(2\sqrt{\frac{3}{4}} - \sqrt{\frac{1}{4}} - 1 \right) \\
&= \frac{3(2-\sqrt{3})}{\sqrt{2\pi}}.
\end{aligned} \quad (2.9)$$

Hence, we get

$$\begin{aligned}
\mathbb{E} \left(\min_{1 \leq i \leq 3} |Y_i| \right) - \mathbb{E} \left(\min_{1 \leq i \leq 3} |X_i| \right) &= \frac{12}{\pi^{3/2}} \left(\frac{\sqrt{2\pi}}{4} - \arctan \sqrt{2} \right) - \frac{3(2-\sqrt{3})}{\sqrt{2\pi}} \\
&= \frac{12}{\pi^{3/2}} \left(\frac{\sqrt{6\pi}}{8} - \arctan \sqrt{2} \right) > 0,
\end{aligned}$$

since

$$\tan \left(\arctan \sqrt{2} \right) = \sqrt{2} \sim 1.414, \text{ and } \tan \frac{\sqrt{6\pi}}{8} \sim 1.434.$$

(ii) $n \geq 4$. Without loss of generality, we only consider the case that $n = 4$. We use proof by contradiction. Suppose that (1.3) holds for $n = 4$. Let $Y_i, X_i, i = 1, 2, 3$ be the same as in the above example. Let Y_4 be a standard Gaussian random variable independent of (Y_1, Y_2, Y_3) . Then by the assumption, for any $a > 0$, we have

$$\mathbb{E} \left(\left(\min_{1 \leq i \leq 3} |X_i| \right) \wedge |aY_4| \right) \geq \mathbb{E} \left(\left(\min_{1 \leq i \leq 3} |Y_i| \right) \wedge |aY_4| \right).$$

Letting $a \rightarrow \infty$, by the monotone convergence theorem, we obtain that

$$\mathbb{E} \left(\min_{1 \leq i \leq 3} |X_i| \right) \geq \mathbb{E} \left(\min_{1 \leq i \leq 3} |Y_i| \right).$$

It is a contradiction. Hence for any $M > 0$, there exists $a_0 > M$ such that

$$\mathbb{E} \left(\left(\min_{1 \leq i \leq 3} |X_i| \right) \wedge |a_0 Y_4| \right) < \mathbb{E} \left(\left(\min_{1 \leq i \leq 3} |Y_i| \right) \wedge |a_0 Y_4| \right).$$

2.2 Sufficiency

In this part, we will show that the inequality (1.3) holds if $n = 2$.

Write $X_1 = x_1 f_1, X_2 = x_2 f_2$, where both f_1 and f_2 have the standard normal distribution $N(0, 1)$. Without loss of generality, we can assume that $x_1, x_2 > 0$. Further we can assume that $x_1 = 1, x_2 = a \in (0, 1]$.

Denote by Σ_1 the covariance matrix of (f_1, f_2) . By the symmetry of the distribution $N(0, 1)$, we can assume that $\Sigma_1 = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$, where $0 \leq \rho \leq 1$. Then the covariance matrix Σ_2 of $(f_1, a f_2)$ can be expressed by $\Sigma_2 = \begin{pmatrix} 1 & a\rho \\ a\rho & a^2 \end{pmatrix}$. It follows that if $\rho \in [0, 1)$, then

$$\Sigma_2^{-1} = \frac{1}{a^2(1-\rho^2)} \begin{pmatrix} a^2 & -a\rho \\ -a\rho & 1 \end{pmatrix},$$

and thus the density function of $(f_1, a f_2)$ is

$$p(x, y) = \frac{1}{2\pi\sqrt{a^2(1-\rho^2)}} \exp \left(-\frac{a^2 x^2 + y^2 - 2a\rho xy}{2a^2(1-\rho^2)} \right).$$

At first, we assume that $\rho \in [0, 1)$. By the symmetry, we have

$$\begin{aligned} & \mathbb{E}(|f_1| \wedge |a f_2|) \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (|x| \wedge |y|) \frac{1}{2\pi\sqrt{a^2(1-\rho^2)}} e^{-\frac{a^2 x^2 + y^2 - 2a\rho xy}{2a^2(1-\rho^2)}} dx dy \\ &= \frac{1}{2\pi\sqrt{a^2(1-\rho^2)}} \left[\int_0^{\infty} \int_0^{\infty} (x \wedge y) e^{-\frac{a^2 x^2 + y^2 - 2a\rho xy}{2a^2(1-\rho^2)}} dx dy \right. \\ & \quad + \int_{-\infty}^0 \int_{-\infty}^0 ((-x) \wedge (-y)) e^{-\frac{a^2 x^2 + y^2 - 2a\rho xy}{2a^2(1-\rho^2)}} dx dy + \int_0^{\infty} \int_{-\infty}^0 (x \wedge (-y)) e^{-\frac{a^2 x^2 + y^2 - 2a\rho xy}{2a^2(1-\rho^2)}} dx dy \\ & \quad \left. + \int_{-\infty}^0 \int_0^{\infty} ((-x) \wedge y) e^{-\frac{a^2 x^2 + y^2 - 2a\rho xy}{2a^2(1-\rho^2)}} dx dy \right] \\ &= \frac{1}{\pi\sqrt{a^2(1-\rho^2)}} \left[\int_0^{\infty} \int_0^{\infty} (x \wedge y) e^{-\frac{a^2 x^2 + y^2 - 2a\rho xy}{2a^2(1-\rho^2)}} dx dy + \int_0^{\infty} \int_0^{\infty} (x \wedge y) e^{-\frac{a^2 x^2 + y^2 + 2a\rho xy}{2a^2(1-\rho^2)}} dx dy \right]. \end{aligned}$$

Define

$$I(\vartheta) := \int_0^{\infty} \int_0^{\infty} (x \wedge y) e^{-\frac{a^2 x^2 + y^2 + 2a\vartheta xy}{2a^2(1-\rho^2)}} dx dy.$$

Then

$$\mathbb{E}(|f_1| \wedge |a f_2|) = \frac{1}{\pi\sqrt{a^2(1-\rho^2)}} [I(\rho) + I(-\rho)].$$

We have

$$\begin{aligned} I(\vartheta) &= \int_0^{\infty} \int_0^{\infty} (x \wedge y) e^{-\frac{a^2 x^2 + y^2 + 2a\vartheta xy}{2a^2(1-\rho^2)}} dx dy \\ &= \int_0^{\infty} \left(\int_0^y x e^{-\frac{a^2 x^2 + y^2 + 2a\vartheta xy}{2a^2(1-\rho^2)}} dx \right) dy + \int_0^{\infty} \left(\int_0^x y e^{-\frac{a^2 x^2 + y^2 + 2a\vartheta xy}{2a^2(1-\rho^2)}} dy \right) dx. \end{aligned}$$

Define

$$J(\alpha, \beta, \gamma) := \int_0^\infty \left(\int_0^y x e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy,$$

where $\alpha > 0, \beta > 0, \alpha\beta - \gamma^2 > 0$. Then we have

$$I(\vartheta) = J(a^2, 1, a\vartheta) + J(1, a^2, a\vartheta),$$

and thus

$$\mathbb{E}(|f_1| \wedge |af_2|) = \frac{1}{\pi\sqrt{a^2(1-\rho^2)}} [J(a^2, 1, a\rho) + J(1, a^2, a\rho) + J(a^2, 1, -a\rho) + J(1, a^2, -a\rho)]. \quad (2.10)$$

In the following, we come to calculate the function $J(\alpha, \beta, \gamma)$. We have

$$\begin{aligned} J(\alpha, \beta, \gamma) &= \int_0^\infty \left(\int_0^y x e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy \\ &= \int_0^\infty \left(\int_0^y \left(x + \frac{\gamma}{\alpha} y \right) e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy - \frac{\gamma}{\alpha} \int_0^\infty \left(\int_0^y y e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy \\ &=: J_1(\alpha, \beta, \gamma) - J_2(\alpha, \beta, \gamma), \end{aligned} \quad (2.11)$$

where

$$\begin{aligned} J_1(\alpha, \beta, \gamma) &:= \int_0^\infty \left(\int_0^y \left(x + \frac{\gamma}{\alpha} y \right) e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy, \\ J_2(\alpha, \beta, \gamma) &:= \frac{\gamma}{\alpha} \int_0^\infty \left(\int_0^y y e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy. \end{aligned}$$

We have

$$\begin{aligned} J_1(\alpha, \beta, \gamma) &= \int_0^\infty \left(\int_0^y \left(x + \frac{\gamma}{\alpha} y \right) e^{-\frac{\alpha \left(x + \frac{\gamma}{\alpha} y \right)^2 + \frac{\alpha\beta - \gamma^2}{\alpha} y^2}{2a^2(1-\rho^2)}} dx \right) dy \\ &= \int_0^\infty e^{-\frac{(\alpha\beta - \gamma^2)y^2}{2a^2(1-\rho^2)\alpha}} \left(\int_{\frac{\gamma}{\alpha}y}^{(1+\frac{\gamma}{\alpha})y} u e^{-\frac{\alpha u^2}{2a^2(1-\rho^2)}} du \right) dy \\ &= \frac{a^2(1-\rho^2)}{\alpha} \int_0^\infty e^{-\frac{(\alpha\beta - \gamma^2)y^2}{2a^2(1-\rho^2)\alpha}} \left(e^{-\frac{\gamma^2 y^2}{2a^2(1-\rho^2)\alpha}} - e^{-\frac{(\alpha+\gamma)^2 y^2}{2a^2(1-\rho^2)\alpha}} \right) dy \\ &= \frac{a^2(1-\rho^2)}{\alpha} \int_0^\infty \left(e^{-\frac{\beta y^2}{2a^2(1-\rho^2)}} - e^{-\frac{(\alpha+\beta+2\gamma)y^2}{2a^2(1-\rho^2)}} \right) dy \\ &= \frac{a^2(1-\rho^2)}{\alpha} \cdot \frac{\sqrt{2\pi}}{2} \left(\frac{a\sqrt{1-\rho^2}}{\sqrt{\beta}} - \frac{a\sqrt{1-\rho^2}}{\sqrt{\alpha+\beta+2\gamma}} \right) \\ &= \frac{\sqrt{2\pi}a^3(1-\rho^2)^{\frac{3}{2}}}{2\alpha} \left(\frac{1}{\sqrt{\beta}} - \frac{1}{\sqrt{\alpha+\beta+2\gamma}} \right), \end{aligned} \quad (2.12)$$

and

$$\begin{aligned}
J_2(\alpha, \beta, \gamma) &= \frac{\gamma}{\alpha} \int_0^\infty \left(\int_0^y \left(y + \frac{\gamma}{\beta} x \right) e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy \\
&\quad - \frac{\gamma}{\alpha} \int_0^\infty \left(\int_0^y \left(\frac{\gamma}{\beta} x \right) e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy \\
&= -\frac{\gamma^2}{\alpha\beta} \int_0^\infty \left(\int_0^y x e^{-\frac{\alpha x^2 + \beta y^2 + 2\gamma xy}{2a^2(1-\rho^2)}} dx \right) dy \\
&\quad + \frac{\gamma}{\alpha} \int_0^\infty \left(\int_x^\infty \left(y + \frac{\gamma}{\beta} x \right) e^{-\frac{\beta(y+\frac{\gamma}{\beta}x)^2 + \frac{\alpha\beta-\gamma^2}{\beta}x^2}{2a^2(1-\rho^2)}} dy \right) dx \\
&= -\frac{\gamma^2}{\alpha\beta} J(\alpha, \beta, \gamma) + \frac{\gamma}{\alpha} \int_0^\infty e^{-\frac{(\alpha\beta-\gamma^2)x^2}{2a^2(1-\rho^2)\beta}} \left(\int_{\frac{(\beta+\gamma)x}{\beta}}^\infty u e^{-\frac{\beta u^2}{2a^2(1-\rho^2)}} du \right) dx \\
&= -\frac{\gamma^2}{\alpha\beta} J(\alpha, \beta, \gamma) + \frac{\gamma}{\alpha} \int_0^\infty e^{-\frac{(\alpha\beta-\gamma^2)x^2}{2a^2(1-\rho^2)\beta}} \cdot \frac{a^2(1-\rho^2)}{\beta} e^{-\frac{(\beta+\gamma)^2 x^2}{2a^2(1-\rho^2)\beta}} dx \\
&= -\frac{\gamma^2}{\alpha\beta} J(\alpha, \beta, \gamma) + \frac{a^2(1-\rho^2)\gamma}{\alpha\beta} \int_0^\infty e^{-\frac{(\alpha+\beta+2\gamma)x^2}{2a^2(1-\rho^2)}} dx \\
&= -\frac{\gamma^2}{\alpha\beta} J(\alpha, \beta, \gamma) + \frac{\sqrt{2\pi}a^3(1-\rho^2)^{\frac{3}{2}}\gamma}{2\alpha\beta\sqrt{\alpha+\beta+2\gamma}}. \tag{2.13}
\end{aligned}$$

By (2.11) - (2.13), we get

$$\begin{aligned}
J(\alpha, \beta, \gamma) &= \frac{1}{1 - \frac{\gamma^2}{\alpha\beta}} \left[\frac{\sqrt{2\pi}a^3(1-\rho^2)^{\frac{3}{2}}}{2\alpha} \left(\frac{1}{\sqrt{\beta}} - \frac{1}{\sqrt{\alpha+\beta+2\gamma}} \right) - \frac{\sqrt{2\pi}a^3(1-\rho^2)^{\frac{3}{2}}\gamma}{2\alpha\beta\sqrt{\alpha+\beta+2\gamma}} \right] \\
&= \frac{\sqrt{2\pi}a^3(1-\rho^2)^{\frac{3}{2}}}{2} \cdot \frac{\beta}{\alpha\beta-\gamma^2} \left(\frac{1}{\sqrt{\beta}} - \frac{\beta+\gamma}{\beta\sqrt{\alpha+\beta+2\gamma}} \right),
\end{aligned}$$

which together with (2.10) implies that

$$\begin{aligned}
&\mathbb{E}(|f_1| \wedge |af_2|) \\
&= \frac{1}{\pi\sqrt{a^2(1-\rho^2)}} [J(a^2, 1, a\rho) + J(1, a^2, a\rho) + J(a^2, 1, -a\rho) + J(1, a^2, -a\rho)] \\
&= \frac{1}{\pi\sqrt{a^2(1-\rho^2)}} \cdot \frac{\sqrt{2\pi}a^3(1-\rho^2)^{\frac{3}{2}}}{2} \left[\frac{1}{a^2(1-\rho^2)} \left(1 - \frac{1+a\rho}{\sqrt{a^2+1+2a\rho}} \right) \right. \\
&\quad \left. + \frac{a^2}{a^2(1-\rho^2)} \left(\frac{1}{a} - \frac{a^2+a\rho}{a^2\sqrt{1+a^2+2a\rho}} \right) + \frac{1}{a^2(1-\rho^2)} \left(1 - \frac{1-a\rho}{\sqrt{a^2+1-2a\rho}} \right) \right. \\
&\quad \left. + \frac{a^2}{a^2(1-\rho^2)} \left(\frac{1}{a} - \frac{a^2-a\rho}{a^2\sqrt{a^2+1-2a\rho}} \right) \right] \\
&= \frac{a^2(1-\rho^2)}{\sqrt{2\pi}} \cdot \frac{2(1+a) - \sqrt{1+a^2+2a\rho} - \sqrt{1+a^2-2a\rho}}{a^2(1-\rho^2)} \\
&= \frac{2(1+a) - \sqrt{1+a^2+2a\rho} - \sqrt{1+a^2-2a\rho}}{\sqrt{2\pi}}. \tag{2.14}
\end{aligned}$$

If $\rho = 1$, then $f_2 = f_1$ a.s. Note that $a \in (0, 1]$. Then we have

$$\begin{aligned}\mathbb{E}(|f_1| \wedge |af_2|) &= a\mathbb{E}(|f_1|) \\ &= a \int_{-\infty}^{\infty} |x| \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx \\ &= \frac{2a}{\sqrt{2\pi}} \int_0^{\infty} x e^{-\frac{x^2}{2}} dx \\ &= \frac{2a}{\sqrt{2\pi}}.\end{aligned}$$

In addition, if $\rho = 1$, we have

$$\frac{2(1+a) - \sqrt{1+a^2+2a\rho} - \sqrt{1+a^2-2a\rho}}{\sqrt{2\pi}} = \frac{2a}{\sqrt{2\pi}}.$$

Hence (2.14) holds for any $\rho \in [0, 1]$.

For any $a \in (0, 1]$, and any $\rho \in (0, 1)$, we have

$$\frac{d \left[\frac{2(1+a) - \sqrt{1+a^2+2a\rho} - \sqrt{1+a^2-2a\rho}}{\sqrt{2\pi}} \right]}{d\rho} = \frac{a}{\sqrt{2\pi}} \left(\frac{1}{\sqrt{1+a^2-2a\rho}} - \frac{1}{\sqrt{1+a^2+2a\rho}} \right) > 0.$$

Hence for any $a \in (0, 1]$, $\frac{2(1+a) - \sqrt{1+a^2+2a\rho} - \sqrt{1+a^2-2a\rho}}{\sqrt{2\pi}}$ is a strictly increasing function in $\rho \in [0, 1]$. Hence it reaches its minimum value at $\rho = 0$, i.e. the inequality (1.3) holds. The proof is complete.

Acknowledgments We thank Guolie Lan, Qiman Shao and Wei Sun for helpful discussions. This work was supported by the National Natural Science Foundation of China (No. 11771309).

References

- [1] Gluskin E.D: Extremal properties of orthogonal parallelepipeds and their applications to the geometry of Banach spaces, Math. USSR Sbornik, 64, 85-96 (1989).
- [2] Gordon Y., Litvak A., Schütt C., Werner E.: Minima of sequences of Gaussian random variables, C. R. Acad. Paris, Ser. 1 340, 445-448 (2005).
- [3] Gordon Y., Litvak A., Schütt C., Werner E.: On the minimum of several random variables, Proc. Amer. Math. Soc. 134(12), 3665-3675 (2006).
- [4] Li W. B.: L1: Introduction, Overview and Applications. CBMS Lectures at UAH, June 4-8, 2012. http://jamesyli.com/wenboli_backup/papers/CBMS-10Lectures.pdf
- [5] Litvak A. E. Around the simplex mean width conjecture. Analytic Aspects of Convexity. Springer INdAM Series, 25, 73-84 (2018).
- [6] Mallat S., Zeitouni O.: A conjecture concerning optimality of the Karhunen-Loève basis in nonlinear reconstruction, arXiv:1109.0489v2, 2011.
- [7] Šidák, Z.: Rectangular confidence regions for the means of multivariate normal distributions, J. Am. Stat. Assoc. 62, 626-633 (1967).
- [8] Shao Q. M.: In memory of Wenbo V. Li's contribution, High dimensional probability VII, 281-291, Progr. Probab., 71, Springer, 2016.
- [9] Sun W., Hu Z.-C., Lan G. L. The 4-D Gaussian random vector maximum conjecture and the 3-D simplex mean width conjecture, arXiv: 2008.04827v1, 2020.