RANKIN-SELBERG INTEGRALS FOR LOCAL SYMMETRIC SQUARE FACTORS ON GL(2)

YEONGSEONG JO

ABSTRACT. Let π be an irreducible admissible (complex) representation of GL(2) over a nonarchimedean characteristic zero local field with odd residual characteristic. In this paper we prove the equality between the local symmetric square *L*-function associated to π arising from integral representations and the corresponding Artin *L*-function for its Langlands parameter through the local Langlands correspondence. With this in hand, we show the stability of local symmetric γ factors attached to π under highly ramified twists.

1. INTRODUCTION

Let F be a p-adic field with $p \neq 2$. We study the Rankin-Selberg integral of the local symmetric square L-functions for an irreducible admissible (complex) representation π of $GL_2(F)$, introduced by Yamana [44]. The ultimate goal of this paper is twofold. The first aim is to show that the automorphic symmetric square L-function attached to π by the theory of integral representations is equal to the corresponding Artin L-function for its Langlands parameter via the local Langlands correspondence.

To elaborate our result more rigorously, let q be the cardinality of the residue field of F. The local symmetric square L-function $L(s, \pi, \text{Sym}^2)$ is defined as the unique normalized generator of a $\mathbb{C}[q^{\pm s/2}]$ -fractional ideal spanned by Rankin-Selberg integrals for the space of good sections. The generator is oftentimes referred as the greatest common divisor (gcd) [23]. Let $\text{Sym}^2 : GL_2(\mathbb{C}) \to$ $GL_3(\mathbb{C})$ be the symmetric square representation. We can relate to the local symmetric square L-function $L(s, \text{Sym}^2(\rho(\pi)))$ of Artin type, where ρ stands for the local Langlands correspondence.

Theorem 1 (The equality). Let π be an irreducible admissible representation of $GL_2(F)$ and $\rho(\pi)$ the associated Langlands parameter. Then we have

$$L(s, \pi, \operatorname{Sym}^2) = L(s, \operatorname{Sym}^2(\rho(\pi))).$$

The discrete series case for $GL_n(F)$ has already been proven by the work of Yamana [44]. It was accomplished in [14] that local symmetric square *L*-functions for $GL_n(F)$ from the Langlands-Shahidi method [34] coincide with counterpart Artin *L*-functions. One immediate corollary of Theorem 1 is the factorization

(1.1)
$$L(s,\pi\times\pi) = L(s,\omega_{\pi})L(s,\pi,\operatorname{Sym}^2) = L(s,\pi,\wedge^2)L(s,\pi,\operatorname{Sym}^2)$$

where ω_{π} is the central character of π , and $L(s, \pi \times \pi)$, $L(s, \omega_{\pi})$ and $L(s, \pi, \wedge^2)$ denote, respectively, the Rankin-Selberg *L*-function for $GL_2(F) \times GL_2(F)$ [17], the Tate *L*-function [39], and the Jacquet-Shalika exterior square *L*-function for $GL_2(F)$ (See [19]). As opposed to contending that (1.1) is the definition in Gelbart and Jacquet [11, §3], we provide a natural way to define *L*-functions and express them directly in terms of inducing data.

²⁰²⁰ Mathematics Subject Classification. Primary; 11F70, Secondary; 11F66, 11F85, 22E50.

Key words and phrases. Exceptional poles and Bernstein-Zelevinsky derivatives, Howe vectors, Local symmetric square L-functions, Stability of symmetric square γ -factors.

The second purpose of this note is to prove the stability of symmetric square local factors. Let ψ be a fixed additive character of the field F. The Rankin-Selberg integrals satisfy a functional equation to define γ -factors $\gamma(s, \pi, \operatorname{Sym}^2, \psi)$. In contrast with the γ -factor being a rational function in $\mathbb{C}(q^{-s/2})$, an ε -factor $\varepsilon(s, \pi, \operatorname{Sym}^2, \psi)$ appearing in the functional equation is exponential, namely, a unit in $\mathbb{C}[q^{\pm s/2}]$.

Theorem 2 (The analytic stability). Let π and σ be irreducible admissible representations of $GL_2(F)$ sharing the same central character. For every sufficiently highly ramified characters χ of F^{\times} , identified as a character of $GL_2(F)$ through the determinant, we obtain

$$\gamma(s,\pi\otimes\chi,\mathrm{Sym}^2,\psi)=\gamma(s,\sigma\otimes\chi,\mathrm{Sym}^2,\psi)$$

In this situation, the L and epsilon factors stabilize as well:

$$L(s,\pi\otimes\chi,\mathrm{Sym}^2)=L(s,\sigma\otimes\chi,\mathrm{Sym}^2)=1\quad and\quad \varepsilon(s,\pi\otimes\chi,\mathrm{Sym}^2,\psi)=\varepsilon(s,\sigma\otimes\chi,\mathrm{Sym}^2,\psi).$$

The analogous result has been settled for GL_n over a non-archimedean field F with characteristic zero [8] and with positive characteristic [9] in the formulation of Langlands-Shahidi local coefficients.

The Rankin-Selberg method plays a profound role in analyzing Langlands automorphic L-functions in company with the Langlands-Shahidi method [8–10, 34] and the doubling method [13,43]. The main subject of the Rankin-Selberg convolution is to determine the integral representation which admits a factorization into an Euler product. More importantly the global integral ought to produce the symmetric square L-function with unramified data. This method originated from the construction of Shimura [35] for n = 2. Later this integral was reformulated in adelic language by Gelbart and Jacquet [11] for n = 2 and by Patterson and Piatetski-Shapiro [30] for n = 3. In the early 1990's Bump and Ginzburg [3] extended the method to arbitrary n. Afterward Takeda [37,38] carried out the twisted case for general n. However the integrals of Gelbart-Jacquet and Takeda are not taken over $GL_2(F)$ so the family of integrals does not afford a $GL_2(F)$ -trilinear form which is an essential ingredient to characterize the exceptional poles of L-function. For this reason, we mainly treat a modified local Zeta integral introduced by Yamana [44].

One significant difference from the setting of Jacquet, Piatetski-Shapiro, and Shalika [17], is that in comparison to Fourier transforms, the functional equation involves the intertwining operator. In the literature one might be temped to try the space of holomorphic sections to generate a $\mathbb{C}[q^{\pm s/2}]$ fractional ideal but the poles of this fractional ideal only contribute the regular *L*-function. The notion of good sections appeared in the work of Piatetski-Shapiro and Rallis [32]. Taking a lead of their methodology, the set of "good sections" consists of holomorphic sections and the image of holomorphic sections under the normalized operator. In turn the additional pole coming from the normalized operator is attributed to the exceptional *L*-function. The noteworthy discovery is that adapting good sections compensates the lack of symmetry of the functional equation caused by just using holomorphic sections. For the sake of explaining it, the operator is a bijection on the space of good sections although holomorphic sections are not necessarily mapped to themselves.

The terminologies of "exceptional" and "regular" parts of L-functions have been extensively exploited in the construction of L-functions for $GSp_4(F)$. We refer to [33] and the references therein, which are all based on the work of Piatetski-Shapiro [31]. The method of good sections was reshaped and rapidly developed in the perspective of the doubling method [13, 43]. Nevertheless it takes several years for this approach to emerge in the study of non-archimedean local L-functions through integral representations. Kaplan [23] applied good sections to L-functions for $SO_{2m}(F) \times GL_n(F)$ and recently Chen [6] implemented the study of Asai cube L-functions for $GL_2(F)$, which is precisely what is used by Piatetski-Shapiro and Rallis [32] in the contexture of Rankin triple product for $GL_2(F)$. Thankfully the main result of [20] asserts that L-functions supplied by auxiliary variables of Schwartz-Bruhat functions and good sections are in fact the same in the framework of numerous $GL_n(F)$ -type cases such as local Rankin-Selberg, Asai, and exterior square L-functions.

A major shortcoming of utilizing the Rankin-Selberg method is that one needs to prove multiplicativity of γ -factors. Cogdell and Piatetski-Shapiro [7] devised a systematic machinery to compute local L-functions without relying on multiplicative properties. The crux of their observation is to interpret the occurrence of poles of exceptional L-functions for Bernstein-Zelevinsky derivatives of representations [2] as the appearance of various distinguished representations. In order to control the location of poles, they suggest suitably deforming representations for which L-functions are tractable. This technique is adapted to tackle the problem of computing Asai [28] and Bump-Friedberg L-functions [29] by Matringe and exterior square L-functions by the author [18, 19]. See [4] also for the derivatives and exceptional poles on archimedean places. Along the line of this prototype, Kable [22] initiated the project to understand the structure of the symmetric square L-function in the late 1990's. To gain an intimate knowledge of that L-function, Kable was led to examine all the derivatives of exceptional representations. After that, the remaining task was the dyadic case and this computation was concluded by Kaplan [24] and Yamana [44] independently. As a continuation of their direction, we complete the particular case of n = 2. It might be possible to reduce the local and global functorial lifts of Gelbart and Jacquet [11] to our main results. However it is our belief that the context of the present paper will work out the higher ranked case of $GL_n(F)$ and we plan to do so in near future.

The stability of symmetric square L-functions is known for $GL_2(F)$ [11, (6.4)]. Nonetheless the stability of L-functions does not directly imply the stability of the corresponding gamma and epsilon factors. In principle, our result should follow from the equality between Rankin-Selberg and Langlands-Shahidi local symmetric square γ -factors. To the author's understanding, the matching of two types of γ -factors is unfortunately not recorded anywhere. In this regard, a proof of stability of γ -factors within the context of integral representations has its own merit. Our approach in the proof is rooted on asymptotic analysis of partial Bessel functions associated with Howe vectors which can be viewed as an extension of the work [5,45] to the framework of the metaplectic group.

The interested reader will notice that many of our calculation apply over any non-archimedean local field. The restriction to the characteristic on the field F throughout this article comes from the intrinsic nature of the global integral built in number fields [37, 38], which is sufficient for its global application. Nevertheless the Bump and Ginzburg global integral [3] is actually constructed over (global) function fields. We pursue an in-depth investigation on the comparison of L-functions and stability of γ -factors in the positive characteristic case.

The rest of the paper is organized as follows. Section 2 contains the preliminary, including a brief review of the metaplectic group, exceptional representations and Rankin-Selberg integrals. Section 3 is concerned with the exceptional and regular *L*-function. At the end of Section 3 we deduce the factorization formula. The local symmetric square *L*-functions are computed in Section 4 and Section 5 is devoted to the stability of local factors.

2. The Rankin-Selberg Integrals

2.1. The metaplectic group. Let F be a non-archimedean local field of characteristic zero with odd residual characteristic. We let \mathcal{O} be the ring of integers of F, \mathfrak{p} the unique prime ideal of \mathcal{O} , and ϖ a uniformizer, so $\mathfrak{p} = (\varpi)$. We normalize the absolute value by $|\varpi|^{-1} = |\mathcal{O}/\mathfrak{p}| = q$. For each subgroup $H(F) \subset GL_2(F)$, we often write H for H(F) when the base field is clear from the context. Let B = TN denote the Borel subgroup of upper triangular matrices, where

$$T = \left\{ t(a,b) := \begin{pmatrix} a \\ b \end{pmatrix} \middle| a, b \in F^{\times} \right\}$$

is the maximal torus made of diagonal matrices and

$$N = \left\{ n(x) := \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \middle| x \in F \right\}$$

is the unipotent radical of B. Let Z denote the center of GL_2 and let A denote the subtorus

$$A = \{t(a,1) \mid a \in F^{\times}\}.$$

Let

$$\overline{N} = \left\{ \overline{n}(x) := \begin{pmatrix} 1 \\ x & 1 \end{pmatrix} \middle| x \in F \right\}$$

be the unipotent subgroup opposed to N and $\overline{B} = T\overline{N}$ the lower triangular Borel subgroup. We write

$$w_2 = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}$$

to denote the long Weyl element in GL_2 . Let W denote the Weyl group defined by $W = N_{GL_2}(T)/T = \{I_2, w_2\}$. We recall the Bruhat decomposition

$$GL_2 = B \cup Bw_2N,$$

with uniqueness of expression, that is, every $g \notin B$ has the unique expression of the form $g = bw_2 n$, $b \in B, n \in N$. We denote by P the mirabolic subgroup given by

$$P = \left\{ \begin{pmatrix} a & x \\ & 1 \end{pmatrix} \middle| a \in F^{\times}, x \in F \right\} \cong A \ltimes N.$$

Put $K = GL_2(\mathcal{O})$, the standard maximal compact subgroup of $GL_2(F)$.

The content of § 2.1 and § 2.2 is basically a summary of necessary definitions and essential results in [37,38,44]. For a, a' and $b \in F^{\times}$, the Hilbert symbol is a map $(\cdot, \cdot)_F : F^{\times} \times F^{\times} \to \{\pm 1\}$ satisfying (See [42, Chapter VIII. §5])

- (1) $(a,b)_F(a',b)_F = (aa',b)_F$
- (2) $(a,b)_F(b,a)_F = 1$
- (3) $(a, -a)_F(a, 1-a)_F = 1$

(4) $\{a \mid (a, y)_F = 1 \text{ for all } y \in F^{\times}\} = (F^{\times})^2$, where $(F^{\times})^2 = \{a^2 \mid a \in F^{\times}\}$.

It is noteworthy that $(a,b)_F = 1$ for all $a, b \in \mathcal{O}^{\times}$ if and only if |2| = 1 (cf. [25, Section 1.1.3]). Hence the Hilbert symbol is unramified. We set

$$oldsymbol{X} egin{pmatrix} a & b \ c & d \end{pmatrix} = egin{pmatrix} c & ext{if } c
eq 0, \ d & ext{if } c = 0. \end{bmatrix}$$

The Kubota 2-cocycle $\sigma_2: GL_2 \times GL_2 \to \{\pm 1\}$ is defined by

$$\sigma_2(g_1,g_2) = \left(\det(g_1), \frac{\boldsymbol{X}(g_1g_2)}{\boldsymbol{X}(g_1)}\right)_F \left(\frac{\boldsymbol{X}(g_1g_2)}{\boldsymbol{X}(g_1)}, \frac{\boldsymbol{X}(g_1g_2)}{\boldsymbol{X}(g_2)}\right)_F.$$

The metaplectic double cover \widetilde{GL}_2 is a non-trivial central extension of GL_2 by $\{\pm 1\}$:

$$1 \longrightarrow \{\pm 1\} \longrightarrow \widetilde{GL}_2 \xrightarrow{pr} GL_2 \longrightarrow 1,$$

where pr is a canonical projection given by $pr(g,\xi) = g$ for $g \in GL_2$ and $\xi \in \{\pm 1\}$. As a set, GL_2 is realized to be

$$\widetilde{GL}_2 = GL_2(F) \times \{\pm 1\} = \{(g,\xi) \mid g \in GL_2(F), \xi \in \{\pm 1\}\}$$

and the group law is defined by

$$(g_1,\xi_1) \cdot (g_2,\xi_2) = (g_1g_2,\sigma_2(g_1,g_2)\xi_1\xi_2).$$

It is known that there exists a compact subgroup \mathcal{K} of GL_2 which splits in GL_2 (cf. [37, §1.1; 44, 1B]), that is to say, there is a continuous map $s_2 : GL_2 \to \{\pm 1\}$ such that $\sigma_2(k_1, k_2) = s_2(k_1)s_2(k_2)s_2(k_1k_2)$ for all $k_1, k_2 \in \mathcal{K}$. If the residue characteristic of F is odd (that being said, |2| = 1) then we can take $\mathcal{K} = K$ (See [37, §1.1]) and $s_2(k_1)s_2(k_2)s_2(k_1k_2) = 1$ for all $k_1, k_2 \in \mathcal{K}$.

With this choice of s_2 , the section $\mathcal{K} \to \widetilde{GL}_2$ defined by $k \mapsto (k, s_2(k))$ is what is called the canonical lift of Kazhdan and Patterson [26]. We define another 2-cocycle τ_2 by

 $\tau_2(g_1, g_2) = \sigma_2(g_1, g_2) s_2(g_1) s_2(g_2) s_2(g_1 g_2)$ for $g_1, g_2 \in GL_2$.

The choice of s_2 , and hence τ_2 , is not unique. However as explained in [37, P.181], we assume that s_2 is chosen to be trivial so that τ_2 coincides with σ_2 . We define a set theoretic section \mathfrak{s} by $\mathfrak{s}(g) = (g, 1)$ (cf. [12, (1.1.4)]). We remark that the multiplication in the image $\mathfrak{s}(GL_2)$ is given via σ_2 , by means of

$$(g_1, 1)(g_2, 1) = (g_1g_2, \sigma_2(g_1, g_2))$$
 for $g_1, g_2 \in GL_2$.

For every m, let K_m be the m-th congruence subgroup, that is, $K_m = \{k \in K \mid k \equiv I_2 \pmod{\mathfrak{p}^m}\}$. Then the collection $\{\mathfrak{s}(K_m) \mid m \geq 0\}$ is a basis of compact open neighborhoods of the identity in \widetilde{GL}_2 and the topology of \widetilde{GL}_2 as a locally compact group is determined by the embedding $\mathfrak{s}: K \to \widetilde{GL}_2$.

We introduce the basic property of Hilbert symbol. For $g \in GL_2$, $g = n_1 t w n_2$ for some $n_1, n_2 \in N$, $t \in T$, and $w \in W$. We define t to be the map $GL_2 \to T$, given by $\mathfrak{t}(g) = \mathfrak{t}(n_1 t w n_2) = t$.

Lemma 2.1. [1, Theorem 7] Suppose that $n \in N$, $g \in GL_2$ and $w \in W$. Let t = t(a,b) and t' = t(a',b') be elements in T. Then we have

- (1) $\sigma_2(n,g) = \sigma_2(g,n) = 1 \text{ for all } n \in N, g \in GL_2$
- (2) $\sigma_2(t,t') = (a,b')_F \text{ for all } t, t' \in T$
- (3) $\sigma_2(t,g) = \sigma_2(t,\mathfrak{t}(g))$ for all $t \in T, g \in GL_2$
- (4) $\sigma_2(w,g) = \sigma_2(\mathfrak{t}(wg)\mathfrak{t}(g)^{-1}, -\mathfrak{t}(g))$ for all $w \in W, g \in GL_2$.

(3) and (4) reduces the calculation to (2) and then we compute the cocycle explicitly by Hilbert symbols. Then we deduce the following general formula.

Lemma 2.2. [1, Theorem 7] Let $g, g' \in GL_2$ and suppose $g = n_1 twn_2$. Then

 $\sigma_2(g,g') = \sigma_2(t,wn_2g')\sigma_2(w,n_2g').$

The right-hand side can be computed by (3) and (4) of Lemma 2.1. For each subgroup $H \subset GL_2$, we let $\tilde{H} = pr^{-1}(H)$ the metaplectic preimage of H under pr. \mathfrak{s} splits H whenever the cocycle σ_2 is trivial on $H \times H$. In this case we simply denote by H^* the image $\mathfrak{s}(H)$. Then \tilde{H} is the direct product of $\{\pm 1\}$ with H^* . We see from Lemma 2.1 that \mathfrak{s} splits the following subgroups N, A, and

$$Z^{2} = \left\{ \begin{pmatrix} a \\ & a \end{pmatrix} \middle| a \in (F^{\times})^{2} \right\}.$$

Accordingly we denote $\mathfrak{s}(N)$, $\mathfrak{s}(A)$, and $\mathfrak{s}(Z^2)$ by N^* , A^* , and Z^{2^*} . In particular \mathfrak{s} splits W if and only if $(-1, -1)_F = 1$ (See [1, Section 5]). We note that $\widetilde{GL}_1 = GL_1 \times \{\pm 1\}$, where the product is given by the direct product. Also we define $\widetilde{F^{\times}}$ to be $\widetilde{F^{\times}} = F^{\times} \times \{\pm 1\}$ as a set but the product is $(a_1, \xi_1) \cdot (a_2, \xi_2) = (a_1a_2, (a_1, a_2)_F\xi_1\xi_2).$

We know from Lemma 2.1 that $\sigma_2(a_1I_2, a_2I_2) = (a_1, a_2)_F$. Hence \widetilde{Z} is not the center of \widetilde{GL}_2 . As a matter of the fact, \widetilde{Z}^2 is the center of \widetilde{GL}_2 and \widetilde{Z} is isomorphic to $\widetilde{F^{\times}}$. We also note that \widetilde{Z} is the center of $\widetilde{GL}_2^{(2)}$, where $\widetilde{GL}_2^{(2)}$ is the metaplectic preimage of

$$GL_2^{(2)} = \{ g \in GL_2 \mid \det(g) \in (F^{\times})^2 \}.$$

We fix a non-trivial continuous additive character ψ of F. We define $\mathfrak{f}(\psi)$, the conductor of ψ , to be the smallest positive integer m such that ψ is trivial on \mathfrak{p}^m . For each $a \in F^{\times}$, we denote by ψ_a the additive character defined by $\psi_a(x) = \psi(ax)$. The map $F \to \mathbb{C}^{\times}$ defined by $x \mapsto \psi(x^2)$ is what Weil called a *character of second degree*. The Weil index $\gamma(\psi)$ of ψ [42] is an eighth root of unity attached to any character of second degree ψ . Likewise we can define $\gamma(\psi_a)$ for each $a \in F$. We put

$$\mu_{\psi}(a) = \frac{\gamma(\psi_a)}{\gamma(\psi)}$$

What is particularly important is that

$$\mu_{\psi}(ab) = \mu_{\psi}(a)\mu_{\psi}(b)(a,b)_F \quad \text{and} \quad \mu_{\psi}(ab^2) = \mu_{\psi}(a)$$

for $a, b \in F^{\times}$. Hence it extends to a group homomorphism $\widetilde{F^{\times}} \to \mathbb{C}$ defined by $(a, \xi) \mapsto \xi \mu_{\psi}(a)$.

2.2. Exceptional representations. In §2.2, we review some technical results on the exceptional representations. Let H be a subgroup of GL_2 . We let $\mathbf{1}_H$ denote the trivial character on H. Let π denote an irreducible admissible representation of \widetilde{H} . π is said to be genuine if $\pi((1,\xi)h) = \xi\pi(h)$ for all $\xi \in \{\pm 1\}$ and $h \in H$, that is, each element in $(1,\xi) \in \widetilde{H}$ acts as a multiplication by ξ . Any representation π of H can be pulled back to a non-genuine representation of \widetilde{H} by composing it with the canonical projection $pr : \widetilde{GL}_2 \to GL_2$. In particular, for a Borel subgroup B, we view the modular character of δ_B as a character on \widetilde{B} in this way.

As we discussed before, each element in N can be written in the form $(1,\xi)n^*$ for $n^* \in N^*$ and $\xi \in \{\pm 1\}$ and $(1,\xi) \in \widetilde{T}$. We can check by exploiting Lemma 2.1 that $\mathfrak{s}(t)\mathfrak{s}(n)\mathfrak{s}(t)^{-1} = \mathfrak{s}(tnt^{-1})$ for all $t \in T$ and $n \in N$. Thus $\widetilde{B} = \widetilde{T}N^*$ and N^* is normalized by \widetilde{T} . We also have $\widetilde{T} \cap N^* = \{(1,1)\}$. For the maximal torus $T \subset B$, we let

$$T^e = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \middle| ab^{-1} \text{ is square } \right\}.$$

The metaplectic preimage \tilde{T}^e is a maximal abelian subgroup of \tilde{T} . We define a character ω^{ψ} on \tilde{T}^e by $\omega^{\psi}((1,\xi)\mathfrak{s}(t(a,b))) = \xi\mu_{\psi}(b)^{-1}$. For t(a,b) and $t(a',b') \in \tilde{T}^e$, Lemme 2.1 implies that $\sigma_2(t(a,b),t(a',b')) = (a,a')_F$. We can conclude that ω^{ψ} is indeed a genuine character of \tilde{T}^e . The exceptional representation θ^{ψ} [26] is the unique irreducible quotient of the normalized induced representation $\mathrm{Ind}_{\tilde{T}^e N^*}^{\widetilde{GL}_2}(\omega^{\psi} \otimes \delta_B^{1/4})$, isomorphic to the unique irreducible subrepresentation of $\mathrm{Ind}_{\tilde{T}^e N^*}^{\widetilde{GL}_2}(\omega^{\psi} \otimes \delta_B^{-1/4})$. Here normalized induction means that $\mathrm{Ind}_{\tilde{T}^e N^*}^{\widetilde{GL}_2}(\omega^{\psi} \otimes \delta_B^{1/4})$ is unitarizable whenever $\omega^{\psi} \otimes \delta_B^{1/4}$ is unitarizable. Let \tilde{T}^2 be the inverse image of T^2 , where $T^2 = \{t^2 \mid t \in T\}$ is the set of square elements in the torus. $\tilde{T}^2 \tilde{Z}^2$ is the center of \tilde{T} . In general a character ω of $\tilde{T}^2 \tilde{Z}^2$ is called exceptional if $\omega(\mathfrak{s}(x^2, x^{-2})) = |x|$ for all $x \in F^{\times}$. The restriction of $\omega^{\psi} \otimes \delta_B^{1/4}$ to $\tilde{T}^2 \tilde{Z}^2$ is an exceptional character.

Let η be a character of F^{\times} . For a ramified character η , let $\mathfrak{f}(\eta)$ be the conductor of η defined to be the smallest integer m such that η is trivial on $1 + \mathfrak{p}^m$. A character η of F^{\times} is said to be quadratic if $\eta^2 = 1$. We define a genuine character $\tilde{\eta}$ of \tilde{Z}^2 by

$$\widetilde{\eta}((1,\xi)\mathfrak{s}(z)) = \xi\eta(a), \quad z = \begin{pmatrix} a \\ & a \end{pmatrix} \in Z^2.$$

We embed GL_1 into GL_2 via the map $a \mapsto \begin{pmatrix} a \\ 1 \end{pmatrix}$. Then we extend $\mathbf{1}_{GL_1}$ to the representation $\widetilde{\mathbf{1}}_{GL_1} \boxtimes \widetilde{\eta}$ of the semidirect product $(\widetilde{GL}_1 \times \widetilde{Z}^2) \ltimes N^*$ by letting \widetilde{Z}^2 act by $\widetilde{\eta}$ and N^* act trivially, where $\widetilde{\mathbf{1}}_{GL_1}$ is the non-genuine character on \widetilde{GL}_1 (trivial extension) given by $(a, \xi) \mapsto \xi$. For $s \in \mathbb{C}$, we define a normalized induced representation (cf. [38, p.132])

$$I(s,\eta) = \operatorname{Ind}_{\widetilde{Z}^{2}\widetilde{P}}^{\widetilde{GL}_{2}}((\widetilde{\eta} \boxtimes \widetilde{\mathbf{1}}_{GL_{1}}) \otimes \delta_{B}^{s/4}).$$

equipped with the natural action of GL_2 on $I(s,\eta)$ by a right translation R. Let us look at the transformation under \widetilde{Z}^2 . We write $(1,\xi)\mathfrak{s}(z) \in \widetilde{Z}^2$ with $z = aI_2$ and $a \in (F^{\times})^2$. For $f_s \in I(s,\eta)$ we have

(2.1)
$$f_s((1,\xi)\mathfrak{s}(z)) = \xi \eta(a) f_s(I_2).$$

The two sides of the functional equation in (2.4) involve slightly different induced representations. For $w \in W$, we denote $\mathfrak{s}(w)$ simply by the same symbol w when there is no danger of confusion. We construct the induced representation occurring the left hand side of the functional equation. In our situation, it means that

$$J(-s,\eta) = \operatorname{Ind}_{\widetilde{Z}^{2} \ w_{2} \widetilde{A} N^{*}}^{\widetilde{GL}_{2}}((\widetilde{\eta} \boxtimes ^{w_{2}} \widetilde{\mathbf{1}}_{GL_{1}}) \otimes \delta_{B}^{-s/4}),$$

where the twisted representation ${}^{w_2}\widetilde{\mathbf{1}}_{GL_2}$ of $\widetilde{\mathbf{1}}_{GL_2}$, to be the representation of ${}^{w_2}\widetilde{A} = w_2\widetilde{A}w_2^{-1}$, is given by ${}^{w_2}\widetilde{\mathbf{1}}_{GL_1}(\widetilde{a}) = \widetilde{\mathbf{1}}_{GL_1}(w_2^{-1}\widetilde{a}w_2)$ for $\widetilde{a} \in {}^{w_2}\widetilde{A}$.

2.3. Rankin-Selberg integrals and good sections. We investigate the basic definition of L-functions and basic existence theorems. Everything stated in § 2.3 without any specific reference is found in [44, §3]. Let π be an irreducible admissible generic representation of GL_2 , ω_{π} its central character, and $\mathcal{W}(\pi, \psi)$ its Whittaker model. Let θ^{ψ} denote an exceptional representation of \widetilde{GL}_2 . Then there exists a unique non-zero Whittaker functional λ on $V_{\theta\psi}$ such that

(2.2)
$$\lambda \left[\theta^{\psi} \left((1,\xi) \mathfrak{s} \begin{pmatrix} a \\ a \end{pmatrix} \mathfrak{s} \begin{pmatrix} 1 & x \\ 1 \end{pmatrix} \right) v \right] = \xi \mu_{\psi}(a)^{-1} \psi^{-1}(x) \lambda(v)$$

for $a \in F^{\times}$, $x \in F$ and $v \in V_{\theta^{\psi}}$ [12, §2.1]. We let $\mathcal{W}(\theta^{\psi}, \psi^{-1})$ denote the Whittaker model of θ^{ψ} . We obtain the Whittaker model $\mathcal{W}(\theta^{\psi}, \psi^{-1})$ by setting $(W_{\theta^{\psi}})_v(\tilde{g}) = \lambda(\theta^{\psi}(\tilde{g})v)$ for $v \in V_{\theta^{\psi}}$ and $\tilde{g} \in \widetilde{GL}_2$. We put $W_{\theta^{\psi}} = (W_{\theta^{\psi}})_v$.

For $W \in \mathcal{W}(\pi, \psi)$, $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, and $f_{2s-1} \in I(2s-1, \omega_{\pi}^{-1})$, we associate the Zeta integral [37,44]

(2.3)
$$I(W, W_{\theta^{\psi}}, f_{2s-1}) = \int_{Z^2 N \setminus GL_2} W(g) W_{\theta^{\psi}}(\mathfrak{s}(g)) f_{2s-1}(\mathfrak{s}(g)) dg$$

This integral is absolute convergent for $\operatorname{Re}(s)$ sufficiently large. By means of Properties (2.1) and (2.2), the group Z^2 acts on the product $W_{\theta\psi}(-)f_{2s-1}(-)$ as ω_{π}^{-1} . Therefore the integral $I(W, W_{\theta\psi}, f_{2s-1})$ is well-defined in the sense that Z^2 acts trivially for the integrand.

There is an intertwining operator

$$M(s,\eta): I(s,\eta) \to J(-s,\eta)$$

given by the formula

$$M(s,\eta)f_s(\widetilde{g}) = \int_F f_s\left(\mathfrak{s}\begin{pmatrix}1\\1\end{pmatrix}\mathfrak{s}\begin{pmatrix}1&x\\1\end{pmatrix}\widetilde{g}\right) dx$$

This integral converges absolutely for $\operatorname{Re}(s)$ large and is defined by meromorphic continuation otherwise. The intertwining operator is mainly utilized later in §5.2. Now it is proven in [44, Proposition 3.14] that there exists a rational function in $\mathbb{C}(q^{-s/2})$ enjoying the following functional equation

(2.4)
$$I(W, W_{\theta^{\psi}}, M(2s-1, \omega_{\pi}^{-1})f_{2s-1}) = \Gamma(s, \pi, \operatorname{Sym}^2, \psi)I(W, W_{\theta^{\psi}}, f_{2s-1}).$$

As we have seen in [44], we normalize the intertwining operator by

$$\hat{N}(s,\eta,\psi) = \gamma(s,\eta^{-2},\psi)M(s,\eta)$$

We want to understand an involution of group $g \mapsto {}^{\iota}g$ of GL_2 defined by ${}^{\iota}g := w_2 {}^{t}g^{-1}w_2$. An automorphism ${}^{\iota}: \widetilde{g} \mapsto {}^{\iota}\widetilde{g}$ is called a lift of the involution if ${}^{\iota}\xi = \xi$ and $p({}^{\iota}\widetilde{g}) = {}^{\iota}p(\widetilde{g})$ for all $\xi \in \{\pm 1\}$ and $\widetilde{g} \in \widetilde{GL}_2$. Kable [21; 44, Proposition 1.3] constructed a lift $\widetilde{g} \mapsto {}^{\iota}\widetilde{g}$ of $g \mapsto {}^{\iota}g$ to \widetilde{GL}_2 satisfying

$$\mathfrak{s}(t(a,b)) = \mathfrak{s}({}^{\iota}t(a,b))(b,a)_F, \quad {}^{\iota}\widetilde{z} = \widetilde{z}^{-1}, \quad {}^{\iota}({}^{\iota}\widetilde{g}) = \widetilde{g}, \quad {}^{\iota}\mathfrak{s}(n) = \mathfrak{s}({}^{\iota}n)$$

for all $t(a,b) \in T$, $\tilde{z} \in Z^2$, $\tilde{g} \in GL_2$ and $n \in N$. Furthermore if the residual characteristic of F is odd and $f: K \to \widetilde{GL}_2$ is a homomorphism, then $f({}^{\iota}k) = {}^{\iota}f(k)$ for all $k \in K$. Following Yamana [44], we then define a \mathbb{C} -linear map

$$N(s,\eta,\psi):I(s,\eta)\to I(-s,\eta^{-1})$$

by setting $N(s, \eta, \psi) f_s(\tilde{g}) = {}^{\iota} [\hat{N}(s, \eta, \psi) f_s](\tilde{g}) = [\hat{N}(s, \eta, \psi) f_s]({}^{\iota}\tilde{g})$. The normalized operator satisfies the functional equation.

Proposition 2.3. Let $N(s, \eta, \psi)$ be a normalized operator as above. Then we have

$$N(-s, \eta^{-1}, \psi^{-1}) \circ N(s, \eta, \psi) = \operatorname{Id}$$

Proof. The induced representation $I(s, \eta)$ can be viewed as a subrepresentation of a genuine principal series representation $\operatorname{Ind}_{\widetilde{T}^e N^*}^{\widetilde{GL}_2}(\mu_s)$, where μ_s is an extension to \widetilde{T}^e of the genuine character of $\widetilde{T}^2 \widetilde{Z}^2$ defined by

$$\mu_s \left(\mathfrak{s} \begin{pmatrix} a \\ b \end{pmatrix} \right) = \eta(b)\mu_{\psi}(b)|a|^{\frac{s}{4}}|b|^{-\frac{s}{4}} = \eta(b)\mu_{\psi}(b)\delta_B^{\frac{s}{4}} \begin{pmatrix} a \\ b \end{pmatrix}$$

for $t(a, b) \in T^2 Z^2$. Shahidi [9, 10, 34] defines the Plancherel measure $\mu(s, \eta)$ associated with η by (2.5) $M(-s, \eta) \circ M(s, \eta) = \mu(s, \eta)^{-1} \cdot \text{Id.}$

It is a priori a rational function in $\mathbb{C}(q^{-s/4})$. As described in [10, (4.9),(4.11),(9.22)], the formula we seek for $\mu(s,\eta)^{-1}$ is therefore

$$\begin{aligned} q^{\mathfrak{f}(\psi)-\mathfrak{f}(\eta^2)} \frac{L(s,\eta^{-2})L(-s,\eta^2)}{L(1-s,\eta^2)L(1+s,\eta^{-2})} &= \varepsilon(s,\eta^{-2},\psi)^{-1}\varepsilon(-s,\eta^2,\psi^{-1})^{-1}\frac{L(s,\eta^{-2})L(-s,\eta^2)}{L(1-s,\eta^2)L(1+s,\eta^{-2})} \\ &= \gamma(s,\eta^{-2},\psi)^{-1}\gamma(-s,\eta^2,\psi^{-1})^{-1} \end{aligned}$$

and the rational function $\alpha_{\psi}(s,\eta)$ in [44, Lemma 3.5] is given by

$$\varepsilon(s,\eta^{-2},\psi)^{-1}\varepsilon(-s,\eta^2,\psi^{-1})^{-1} = q^{\mathfrak{f}(\psi)-\mathfrak{f}(\eta^2)}.$$

Indeed the formula is originally stated for \widetilde{SL}_2 [10, (9.22)]. Nevertheless the very recent result [10, Corollary 10.2] allows us to relate the Plancherel measure associated to a genuine representation of \widetilde{GL}_2 with that of \widetilde{SL}_2 . If we incorporate this into normalized operators, (2.5) can be rewritten as $N(-s, \eta^{-1}, \psi^{-1}) \circ N(s, \eta, \psi) = \text{Id.}$

In what follows we refer to Waldspurger [40, §4] for a through treatment of the notion of sections. We observe that $\widetilde{K} \simeq K^* \times \{\pm 1\}$ is a compact open subgroup [21, §5]. A \widetilde{K} -finite function $f: \mathbb{C} \times \widetilde{GL}_2 \to \mathbb{C}$ such that the mapping $\widetilde{g} \mapsto f(s, \widetilde{g})$ belonging to $I(s, \eta)$ for all s is called a section. A section $f_s \in I(s, \eta)$ is called a standard section if its restriction to \widetilde{K} is independent of s. Let $V_{std}(s, \eta)$ denote the space of standard sections. The space of holomorphic sections is defined by $V_{hol}(s, \eta) = \mathbb{C}[q^{-s/4}, q^{s/4}] \otimes_{\mathbb{C}} V_{std}(s, \eta)$. The elements of $V_{rat}(s, \eta) = \mathbb{C}(q^{-s/4}) \otimes_{\mathbb{C}} V_{std}(s, \eta)$ are called rational sections. In order to incorporate sections in normalized operator, we need to allow sections to vary arithmetically in s. To this end, Piatetski-Shapiro and Rallis introduce the family of good sections [32].

Definition 2.4. We define the space $V_{good}(2s-1,\eta)$ of good sections to comprise the following:

(i) $V_{hol}(2s-1,\eta)$ (ii) $N(1-2s,\eta^{-1},\psi^{-1}) [V_{hol}(1-2s,\eta^{-1})].$

According to Proposition 3.11 (4) of [44], Definition 2.4 agrees with that of Yamana when η is unitary. The good section is closed under the normalized intertwining operator in the sense that $N(2s - 1, \eta, \psi) [V_{good}(2s - 1, \eta)] \subset V_{good}(1 - 2s, \eta^{-1})$. If $f_{2s-1} \in V_{good}(2s - 1, \omega_{\pi}^{-1})$, then $I(W, W_{\theta^{\psi}}, f_{2s-1})$ is a rational function of $q^{-s/2}$. Let $\mathcal{I}(\pi)$ be a subspace of $\mathbb{C}(q^{-s/2})$ generated by integrals $I(W, W_{\theta^{\psi}}, f_{2s-1})$ for $W \in \mathcal{W}(\pi, \psi), W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, and $f_{2s-1} \in V_{good}(2s - 1, \omega_{\pi}^{-1})$. Each such integral can be written with a common denominator (cf. [44, Proposition 3.8-(3)]) and $\mathcal{I}(\pi)$ is closed under multiplication by $q^{s/2}$. Then $\mathcal{I}(\pi)$ is a $\mathbb{C}[q^{-s/2}, q^{s/2}]$ -fractional ideal and in particular it contains 1. Hence we can always find a normalized generator of the form $P(q^{-s/2})^{-1}$ where the polynomial $P(X) \in \mathbb{C}[X]$ satisfies P(0) = 1.

Definition 2.5. We define the symmetric square L-function by $L(s, \pi, \text{Sym}^2) = P(q^{-s/2})^{-1}$, the normalized generator of the fractional ideal $\mathcal{I}(\pi)$ formed by the family of integrals $I(W, W_{\theta^{\psi}}, f_{2s-1})$ for $W \in \mathcal{W}(\pi, \psi), W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, and $f_{2s-1} \in V_{good}(2s-1, \omega_{\pi}^{-1})$.

Let π^{ι} denote the representation of GL_2 on the same space V_{π} but with the action $\pi^{\iota}(g) = \pi({}^{\iota}g)$. If π is irreducible, then $\pi^{\iota} = \tilde{\pi}$, the contragredient representation (See [7, §2.1]). Define $\widetilde{W} : GL_2 \to \mathbb{C}$ as $\widetilde{W}(g) = W({}^{\iota}g) = W(w_2{}^{t}g^{-1}w_2)$. Then we have $\widetilde{W} \in \mathcal{W}(\tilde{\pi}, \psi^{-1})$. For $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, we define a ψ -Whittaker function $\widetilde{W}_{\theta^{\psi}}$ by $\widetilde{W}_{\theta^{\psi}}(\tilde{g}) = W_{\theta^{\psi}}({}^{\iota}\tilde{g})$ for $\tilde{g} \in \widetilde{GL}_2$. We note that $\widetilde{W}_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi^{-1}}, \psi)$ (See [44, Lemma 1.9 (4)]). With the normalized operator, there is a rational function $\gamma(s, \pi, \operatorname{Sym}^2, \psi)$ in $\mathbb{C}(q^{-s/2})$ such that we have

$$I(\widetilde{W}, \widetilde{W}_{\theta^{\psi}}, N(2s-1, \omega_{\pi}^{-1}, \psi) f_{2s-1}) = \gamma(s, \pi, \operatorname{Sym}^2, \psi) I(W, W_{\theta^{\psi}}, f_{2s-1})$$

The ε -factor is defined as the ratio

(2.6)
$$\varepsilon(s,\pi,\operatorname{Sym}^2,\psi) = \frac{\gamma(s,\pi,\operatorname{Sym}^2,\psi)L(s,\pi,\operatorname{Sym}^2)}{L(1-s,\tilde{\pi},\operatorname{Sym}^2)}$$

With the ε -factor in hand, the functional equation can be written in the form

(2.7)
$$\frac{I(\tilde{W}, \tilde{W}_{\theta^{\psi}}, N(2s-1, \omega_{\pi}^{-1}, \psi)f_{2s-1})}{L(1-s, \tilde{\pi}, \operatorname{Sym}^2)} = \varepsilon(s, \pi, \operatorname{Sym}^2, \psi) \frac{I(W, W_{\theta^{\psi}}, f_{2s-1})}{L(s, \pi, \operatorname{Sym}^2)}$$

Converting $I(\widetilde{W}, \widetilde{W}_{\theta^{\psi}}, N(2s - 1, \omega_{\pi}^{-1}, \psi) f_{2s-1})$ on the left hand side of (2.7) as in the proof of Proposition 3.14 of [44], the function equation (2.7) becomes

(2.8)
$$\frac{I(W, W_{\theta^{\psi}}, \hat{N}(2s-1, \omega_{\pi}^{-1}, \psi) f_{2s-1})}{L(1-s, \tilde{\pi}, \operatorname{Sym}^2)} = \varepsilon(s, \pi, \operatorname{Sym}^2, \psi) \frac{I(W, W_{\theta^{\psi}}, f_{2s-1})}{L(s, \pi, \operatorname{Sym}^2)}$$

Proposition 2.6. The epsilon factor $\varepsilon(s, \pi, \text{Sym}^2, \psi)$ satisfies the following functional equation;

$$\varepsilon(1-s, \widetilde{\pi}, \operatorname{Sym}^2, \psi^{-1})\varepsilon(s, \pi, \operatorname{Sym}^2, \psi) = 1.$$

Furthermore $\varepsilon(s, \pi, \operatorname{Sym}^2, \psi)$ is a unit in $\mathbb{C}[q^{\pm s/2}]$.

Proof. The proof is akin to that of [20, Theorem 3.11]. We omit the complete details. \Box

Comparing (2.4) with (2.8), the two factors $\gamma(s, \pi, \text{Sym}^2, \psi)$ and $\Gamma(s, \pi, \text{Sym}^2, \psi)$ are related by the following way.

Proposition 2.7. As functions in $\mathbb{C}(q^{-s/2})$, we have

$$\Gamma(s, \pi, \operatorname{Sym}^2, \psi) = \frac{\gamma(s, \pi, \operatorname{Sym}^2, \psi)}{\gamma(2s - 1, \omega_{\pi}^2, \psi)}.$$

We return to the integral $I(W, W_{\theta^{\psi}}, f_{2s-1})$ in (2.3). Since $W_{\theta^{\psi}}$ and f_{2s-1} are genuine, their product $\tilde{g} \mapsto W_{\theta^{\psi}}(\tilde{g})f_{2s-1}(\tilde{g})$ must factor through the natural map $\widetilde{GL}_2 \to GL_2$. We may therefore view the function $W(g)W_{\theta^{\psi}}(\mathfrak{s}(g))f_{2s-1}(\mathfrak{s}(g))$ as a function on GL_2 . For convenience, we will omit the section \mathfrak{s} by letting g denote $\mathfrak{s}(g)$ for $g \in GL_2$ throughout Section 3 and Section 4.

3. Exceptional Poles and Derivatives

3.1. Derivatives and Whittaker models. We establish Whittaker models for derivatives of exceptional representations. A topological group is called an ℓ -group if it is a Hausdorff space and has a neighborhood base at the identity consisting of compact open subgroups. If G is any ℓ -group, we denote by Rep(G) the category of smooth complex G-modules. Let P_1 denote the identity matrix I_2 . Kable [22] manufactures four functors

$$\Phi^{-}: \operatorname{Rep}(\widetilde{P}) \to \operatorname{Rep}(\widetilde{P}_{1}) \qquad \Psi^{-}: \operatorname{Rep}(\widetilde{P}) \to \operatorname{Rep}(\widetilde{GL}_{1}) \\ \Phi^{+}: \operatorname{Rep}(\widetilde{P}_{1}) \to \operatorname{Rep}(\widetilde{P}) \qquad \Psi^{+}: \operatorname{Rep}(\widetilde{GL}_{1}) \to \operatorname{Rep}(\widetilde{P})$$

which he attributes to Bernstein and Zelevinsky [2]. For $\tau \in \operatorname{Rep}(\widetilde{P})$, $\Phi^{-}(\tau) = \tau/\tau(N^*,\psi)$ is the twisted Jacquet functor or the twisted localization functor, where $\tau(N^*,\psi) = \langle \tau(\mathfrak{s}(n))v - \psi(n)v | n \in N, v \in \tau \rangle$ and $\Psi^{-}(\tau) = \tau/\tau(N^*, \mathbf{1})$ is the Jacquet functor, where $\tau(N^*, \mathbf{1}) = \langle \tau(\mathfrak{s}(n))v - v | n \in N, v \in \tau \rangle$. It is crucial to note that actions of the groups \widetilde{P}_1 and \widetilde{GL}_1 on $\Phi^{-}(\tau)$ and $\Psi^{-}(\tau)$ are normalized by a suitable modulus character $|\det|^{-\frac{1}{2}}$. For $\mathbf{1} \in \operatorname{Rep}(\widetilde{P}_1)$, we put $\Phi^{+}(\mathbf{1}) = \operatorname{c-ind}_{\widetilde{P}_1N^*}^{\widetilde{P}_2}((\mathbf{1} \boxtimes \psi) \otimes |\det|^{\frac{1}{2}})$, where the induction is an unnormalized compactly induced induction. For $\sigma \in \operatorname{Rep}(\widetilde{GL}_1)$, the induction functor Ψ^+ is given by $\Psi^+(\sigma) = \operatorname{ind}_{\widetilde{GL}_1N^*}^{\widetilde{P}}((\sigma \boxtimes \mathbf{1}) \otimes |\det|^{\frac{1}{2}})$. We emphasize that all four functors take genuine representations into genuine representations.

For our purpose, we reconstruct Whittaker models for the first derivative of exceptional representations [12, §3] in the context of Cogdell and Piatetski-Shapiro [7, §1]. Let us mention that what Gelbart and Piatetski-Shapiro denoted by r_{χ} with $\chi = \mathbf{1}_{F^{\times}}$ a trivial character in [12, Proposition 2.3.3] corresponds to what we mean by θ^{ψ} in this paper. Even if $(\theta^{\psi}, V_{\theta^{\psi}})$ is irreducible, $\theta^{\psi}|_{\tilde{P}}$ will not be irreducible and so we shall be forced to build a natural filtration by \tilde{P} -submodules

$$\{0\} \subset \tau_2 \subset \tau_1 = \theta^{\psi}|_{\widetilde{P}}$$

such that $\tau_2 \simeq \operatorname{c-ind}_{\widetilde{P}_1N^*}^{\widetilde{P}_2}(\mathbf{1} \boxtimes \psi) \simeq \Phi^+(\tau_{(1)})$ and $\tau_1/\tau_2 \simeq \Psi^+(\theta^{\psi^{(1)}})$, where the representations $\tau_{(1)} \in \operatorname{Rep}(\widetilde{P}_1)$ and $\theta^{\psi^{(1)}} \in \operatorname{Rep}(\widetilde{GL}_1)$ are defined by $\tau_{(1)} = \Phi^-(\theta^{\psi}|_{\widetilde{P}})$ and $\theta^{\psi^{(1)}} = \Psi^-(\theta^{\psi}|_{\widetilde{P}})$. The proof of these statements can be found in the work of Bernstein and Zelevinsky [2; 41, (2.3)].

We consider the Kirillov map

(3.1)
$$v \mapsto (W_{\theta^{\psi}})_v \left(\mathfrak{s} \begin{pmatrix} a \\ & 1 \end{pmatrix}\right)$$

which takes $V_{\theta^{\psi}}$ to a space of complex valued functions on F^{\times} . Furthermore each map $a \mapsto (W_{\theta^{\psi}})_v \left(\mathfrak{s} \begin{pmatrix} a \\ 1 \end{pmatrix}\right)$ is locally constant on F^{\times} and compactly supported in F. According to [12, §3.1], the Kirillov map (3.1) is injective, that is, $(W_{\theta^{\psi}})_v \left(\mathfrak{s} \begin{pmatrix} a \\ 1 \end{pmatrix}\right) = 0$ implies that v = 0. Therefore the representation of \tilde{P} has a realization on the space of functions on F^{\times} , which is called the *Kirillov model*, $K(\theta^{\psi}, \psi^{-1})$.

Proposition 3.1. [12, §3.1 (Gelbart and Piatetski-Shapiro)] Let $\theta_{(0)}^{\psi} = \theta^{\psi}|_{\tilde{P}}$. Then in terms of Whittaker model $\theta_{(0)}^{\psi}$, we have

$$K(\theta^{\psi},\psi^{-1}) := \mathcal{W}(\theta^{\psi}_{(0)},\psi^{-1}) = \left\{ W : \mathfrak{s} \begin{pmatrix} a \\ & 1 \end{pmatrix} \mapsto W \left(\mathfrak{s} \begin{pmatrix} a \\ & 1 \end{pmatrix} \right) \ \middle| \ W \in \mathcal{W}(\theta^{\psi},\psi^{-1}), a \in F^{\times} \right\}$$

endowed with the natural action of \widetilde{P} by right translation.

We analyze the behavior of the Kirillov function $W_{\theta^{\psi}}\left(\mathfrak{s}\begin{pmatrix}a\\1\end{pmatrix}\right)$ near 0. Let $\mathcal{S}(F^{\times})$ be the space of locally constant functions $\varphi: F^{\times} \to \mathbb{C}$ with compact supports in F^{\times} . The representation $R_{\theta^{\psi}}$ of \widetilde{B} on the space $\mathcal{S}(F^{\times})$ is given by

$$R_{\theta^{\psi}}\begin{pmatrix}a & x\\ & 1\end{pmatrix}\varphi(y) = \psi(xy)\varphi(ya) \quad \text{and} \quad R_{\theta^{\psi}}\begin{pmatrix}z\\ & z\end{pmatrix}\varphi(y) = (y,z)_{F}\mu_{\psi}(z)^{-1}\varphi(y).$$

Then $\mathcal{S}(F^{\times})$ is nothing but the model of $\mathcal{W}(\Phi^+(\tau_{(1)}), \psi^{-1})$ under the map $\varphi \in \mathcal{S}(F^{\times}) \mapsto W \in \mathcal{W}(\Phi^+(\tau_{(1)}), \psi^{-1})$, where $\varphi(y) = W\left(\mathfrak{s}\begin{pmatrix} y\\ & 1 \end{pmatrix}\right)$.

Proposition 3.2. [12, §3.2 (Cogdell, Gelbart, and Piatetski-Shapiro)] Let $\theta_{(0)}^{\psi} = \theta^{\psi}|_{\widetilde{P}}$ and $\theta_{(1)}^{\psi} = \Phi^{-}(\theta^{\psi}|_{\widetilde{P}})$. As a \widetilde{P} -modules, $\Phi^{+}(\theta_{(1)}^{\psi})$ has a model of space of functions

$$\mathcal{W}(\Phi^+(\theta^{\psi}_{(1)}),\psi^{-1}) = \left\{ W: \mathfrak{s} \begin{pmatrix} a \\ 1 \end{pmatrix} \mapsto W\left(\mathfrak{s} \begin{pmatrix} a \\ 1 \end{pmatrix} \right) \mid W \in \mathcal{W}(\theta^{\psi},\psi^{-1}), a \in F^{\times} \text{ and there} \\ \text{exists } N > 0 \text{ such that } W\left(\mathfrak{s} \begin{pmatrix} a \\ 1 \end{pmatrix} \right) = 0 \text{ whenever } |a| < q^{-N} \right\} \simeq \mathcal{S}(F^{\times})$$

that is isomorphic to $\mathcal{W}(\theta_{(0)}^{\psi}, \psi^{-1})(N^*, \mathbf{1}) = \langle \theta^{\psi}(\mathfrak{s}(n))W - W \mid W \in \mathcal{W}(\theta^{\psi}, \psi^{-1}), n \in N \rangle.$

As in [12, §3.2], $\mathcal{W}(\Phi^+(\theta_{(1)}^{\psi}), \psi^{-1})$ is the Whittaker model which Gelbart and Piatetski-Shapiro denote by $K_0(\pi)$ and $K(\theta^{\psi}, \psi^{-1})/\mathcal{S}(F^{\times}) \simeq \mathcal{W}(\theta_{(0)}^{\psi}, \psi^{-1})/\mathcal{W}(\theta_{(0)}^{\psi}, \psi^{-1})(N^*, \mathbf{1})$ is the Jacquet module which they denote by $J(\pi)$. Before we turn to the Rankin-Selberg integrals, we illustrate the connection between the Whittaker model for θ^{ψ} and the first derivative $\theta^{\psi^{(1)}}$. For every $b \in F^{\times}$, we define a quadratic character χ_b of F^{\times} by $\chi_b(a) = (b, a)_F$. Let $\nu(g) = |\det(g)|$ denote the unramified determinant character of GL_1 or GL_2 . In virtue of Kable [22, Theorem 5.2], let us write

(3.2)
$$\theta^{\psi^{(1)}} \otimes \nu^{\frac{1}{4}} \simeq \bigoplus_{b \in (F^{\times})^2 \setminus F^{\times}} \chi_b.$$

The character χ_b is an irreducible subrepresentation of the representation $\theta^{\psi^{(1)}}$ with the normalized quotient model $\mathcal{W}(\theta_{(0)}^{\psi}, \psi^{-1})/\mathcal{W}(\theta_{(0)}^{\psi}, \psi^{-1})(N^*, \mathbf{1})$. We denote $\theta_{(0),b}^{\psi}$ by the inverse image of χ_b in $\theta_{(0)}^{\psi}$ for the canonical normalized projection from $\theta_{(0)}^{\psi}$ onto $\theta^{\psi^{(1)}}$ and by $\mathcal{W}(\theta_{(0),b}^{\psi}, \psi^{-1})$ the corresponding subspace of $\mathcal{W}(\theta_{(0)}^{\psi}, \psi^{-1})$. Let $\mathcal{S}(F)$ denote the space of complex-valued locally constant functions with compact supports in F. We reformulate [12, Proposition 3.4] and [12, Remark 3.3.7] in the framework of the Cogdell and Piatetski-Shapiro interpretation of derivatives [7, Corollary to Proposition 1.7]. **Theorem 3.3** (Cogdell, Gelbart, and Piatetski-Shapiro). With notations as above, we have the following: for every $W_{\theta_{(0),b}^{\psi}} \in \mathcal{W}(\theta_{(0),b}^{\psi}, \psi^{-1})$ and for all φ_{\circ} in $\mathcal{S}(F)$ locally constant and supported in a sufficient small neighborhood of 0, there exists a character χ_b and a constant $c \in \mathbb{C}$ such that

$$W_{\theta_{(0),b}^{\psi}}\left(\mathfrak{s}\begin{pmatrix}a\\&1\end{pmatrix}\right)\varphi_{\circ}(a) = \begin{cases} c|a|^{\frac{1}{4}}\chi_{b}(a)\varphi_{\circ}(a), & a \in (F^{\times})^{2} \cdot b\\ 0, & a \notin (F^{\times})^{2} \cdot b. \end{cases}$$

Conversely, for any character χ_b there exist $W_{\theta_{(0),b}^{\psi}} \in \mathcal{W}(\theta_{(0),b}^{\psi}, \psi^{-1})$ and $\varphi_{\circ} \in \mathcal{S}(F)$ non-vanishing at zero such that

$$\begin{array}{ll} \chi_b(a)\varphi_\circ(a), & a \in (F^\times)^2 \cdot b \\ 0, & a \notin (F^\times)^2 \cdot b \end{array} = |a|^{-\frac{1}{4}} W_{\theta_{(0),b}^{\psi}} \left(\mathfrak{s} \begin{pmatrix} a \\ & 1 \end{pmatrix} \right) \varphi_\circ(a).$$

The basic properties of Section 3.1 remain unchanged in the usual GL_2 -setting. The best reference for the theory of derivatives and Whittaker functions on GL_n is [7].

3.2. Exceptional poles and distinctions. In §3.2 we investigate the exceptional and regular Zeta integral. For each $W \in \mathcal{W}(\pi, \psi)$, $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, and $f_{2s-1} \in V_{hol}(2s-1, \omega_{\pi}^{-1})$, we define the regular integral

$$I_{reg}(W, W_{\theta^{\psi}}, f_{2s-1}) = \int_{Z^2 N \setminus GL_2} W(g) W_{\theta^{\psi}}(g) f_{2s-1}(g) dg$$

which converges for $\operatorname{Re}(s)$ large and extends to \mathbb{C} as a function of $\mathbb{C}(q^{-s/2})$. We will use the following property that holomorphic sections are GL_2 -stable to define a $\mathbb{C}[q^{\pm s/2}]$ -fractional ideal.

Lemma 3.4. For $f_{2s-1} \in V_{hol}(2s-1,\eta)$ and $g \in GL_2$, $R(\mathfrak{s}(g))f_{2s-1}$ is a holomorphic section.

Proof. The holomorphic sections are slightly different from those treated in [20, Lemma 4.1]. Observing that $(F^{\times})^2 \setminus F^{\times}$ is finite, the argument [20, Lemma 4.1; 23, Claim 2.1] can easily be modified to deal with our corresponding holomorphic sections.

The \mathbb{C} -vector space

$$\mathcal{I}_{reg}(\pi) = \langle I_{reg}(W, W_{\theta^{\psi}}, f_{2s-1}) \mid W \in \mathcal{W}(\pi, \psi), W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}), f_{2s-1} \in V_{hol}(2s-1, \omega_{\pi}^{-1}) \rangle$$

spanned by the integrals $I_{reg}(W, W_{\theta^{\psi}}, f_{2s-1})$ is a $\mathbb{C}[q^{\pm s/2}]$ -fractional ideal of $\mathbb{C}(q^{-s/2})$. For $W \in \mathcal{W}(\pi, \psi), W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ and $s \in \mathbb{C}$, we define the simple zeta integral

$$I_{(0)}(s; W, W_{\theta^{\psi}}) = \int_{F^{\times}} W \begin{pmatrix} a \\ & 1 \end{pmatrix} W_{\theta^{\psi}} \begin{pmatrix} a \\ & 1 \end{pmatrix} |a|^{\frac{s}{2} - \frac{3}{4}} d^{\times} a$$

 $I_{(0)}(s, W, W_{\theta^{\psi}})$ converges to an element of $\mathbb{C}(q^{-s/2})$ for $\operatorname{Re}(s)$ large enough. Let $\mathcal{I}_{(0)}(\pi)$ denote a \mathbb{C} -vector space of $\mathbb{C}(q^{-s/2})$ spanned by $I_{(0)}(s, W, W_{\theta^{\psi}})$ as W is taken over $\mathcal{W}(\pi, \psi)$ and $W_{\theta^{\psi}}$ runs through $\mathcal{W}(\theta^{\psi}, \psi^{-1})$. Then $\mathcal{I}_{(0)}(\pi)$ is a $\mathbb{C}[q^{\pm s/2}]$ -fractional ideal which is related to $\mathcal{I}_{reg}(\pi)$ by Proposition below. For any open compact set K° of K and any element k of K, we first define the test function $\mathbb{1}_{kK^{\circ},2s-1} \in I(2s-1,\eta)$ by

$$(3.3) \quad \mathbb{1}_{kK^{\circ},2s-1}(\widetilde{g}) = \begin{cases} \xi\eta(z)\delta_B(p)^{(2s+1)/4} & \text{if } \widetilde{g} = (1,\xi)\mathfrak{s}(z)\mathfrak{s}(p)\mathfrak{s}(k)\mathfrak{s}(k^{\circ}), z \in Z^2, p \in P, k^{\circ} \in K^{\circ} \\ 0 & \text{if } \widetilde{g} \notin \widetilde{Z}^2 \widetilde{P}\mathfrak{s}(k)\mathfrak{s}(K^{\circ}). \end{cases}$$

Proposition 3.5. As $\mathbb{C}[q^{s/2}, q^{-s/2}]$ -fractional ideals, we have $\mathcal{I}_{(0)}(\pi) = \mathcal{I}_{reg}(\pi)$. Furthermore we obtain the inclusion $\mathcal{I}_{(0)}(\pi) \subset \mathcal{I}(\pi)$ of $\mathbb{C}[q^{s/2}, q^{-s/2}]$ -fractional ideals.

Proof. Let $f_{2s-1} \in V_{hol}(2s-1,\omega_{\pi}^{-1})$ be given. We write $f_{2s-1} = \sum_{i=1}^{m} P_i(q^{s/2},q^{-s/2})f_{2s-1}^{(i)}$ with $P_i(X) \in \mathbb{C}[X]$ and $f_{2s-1}^{(i)} \in V_{std}(2s-1,\omega_{\pi}^{-1})$. We let $K_{\circ} \subset K$ be a compact open subgroup which stabilizes each $f_{2s-1}^{(i)}$ as well as W and $W_{\theta\psi}$. Write $K = \bigcup_j k_j K_{\circ}$. Since the integrand is left N-invariant and right K_{\circ} -invariant, we reach

$$I_{reg}(W, W_{\theta^{\psi}}, f_{2s-1}) = c_1 \sum_{i,j} P_i(q^{\pm s/2}) \int_{Z^2 \setminus T} [\pi(k_j)W](t) [\theta^{\psi}(k_j)W_{\theta^{\psi}}](t) [R(k_j)f_{2s-1}^{(i)}](t)\delta_B(t)^{-1} dt$$

by the Iwasawa decomposition. Here c_1 is a volume of K_{\circ} . Z^2 has a finite index in Z. Taking the product T = ZA into account, the integral $I_{reg}(W, W_{\theta^{\psi}}, f_{2s-1})$ can be decomposed as a finite sum of the form

$$\begin{split} I_{reg}(W, W_{\theta^{\psi}}, f_{2s-1}) &= c_1 \sum_{i,j} \sum_{(F^{\times})^2 \setminus F^{\times}} P_i(q^{s/2}, q^{-s/2}) \omega_{\pi}(b) \mu_{\psi}(b)^{-1} f_{2s-1}^{(i)}(t(b, b) k_j) \\ & \times \int_{F^{\times}} [\pi(k_j) W] \begin{pmatrix} a \\ 1 \end{pmatrix} [\theta^{\psi}(k_j) W_{\theta^{\psi}}] \begin{pmatrix} a \\ 1 \end{pmatrix} \chi_b^{-1}(a) |a|^{\frac{s}{2} - \frac{3}{4}} d^{\times} a \\ &= \sum_{i,j} \sum_{(F^{\times})^2 \setminus F^{\times}} Q_i(q^{s/2}, q^{-s/2}; b, k_j) I_{(0)}(s, \pi(k_j) W, \theta^{\psi}(k_j) W_{\theta^{\psi}}) \end{split}$$

with $Q_i(X; b, k_j) \in \mathbb{C}[X]$. The second equality follows from [44, Lemma 1.9 (3)]) that $\theta^{\psi} \otimes \chi_b^{-1} \simeq \theta^{\psi}$ for a quadratic character χ_b^{-1} . This confirms that $\mathcal{I}_{reg}(\pi) \subset \mathcal{I}_{(0)}(\pi)$.

For the reverse inclusion, we choose K' for $W \in \mathcal{W}(\pi, \psi)$ and $W_{\theta\psi} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ to be right invariant under K'. We then take f_{2s-1} a characteristic function of $\mathbb{1}_{K',2s-1}$ in (3.3). The integral $I_{reg}(W, W_{\theta\psi}, f_{2s-1})$ reduces to

$$I_{reg}(W, W_{\theta^{\psi}}, f_{2s-1}) = c_2 \int_{F^{\times}} W \begin{pmatrix} a \\ & 1 \end{pmatrix} W_{\theta^{\psi}} \begin{pmatrix} a \\ & 1 \end{pmatrix} |a|^{\frac{s}{2} - \frac{3}{4}} d^{\times} a = c_2 I_{(0)}(s, W, W_{\theta^{\psi}}),$$

where c_2 is a volume of K'. Therefore we obtain the desired inclusion $\mathcal{I}_{reg}(\pi) \supset \mathcal{I}_{(0)}(\pi)$.

According to Theorem 7.1 of [3] or Proposition 3.8 (4) of [44], we have the following non-vanishing result.

Proposition 3.6. There exist $W \in \mathcal{W}(\pi, \psi)$ and $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ such that $I_{(0)}(s, W, W_{\theta^{\psi}}) = 1$.

As a consequence, $\mathcal{I}_{(0)}(\pi)$ contains 1. We can find a polynomial $P(X) \in \mathbb{C}[X]$ such that P(0) = 1and $1/P(q^{-s/2})$ generates the fractional ideal $\mathcal{I}_{reg}(\pi) = \mathcal{I}_{(0)}(\pi)$. We denote by the regular Lfunction

$$L_{reg}(s, \pi, \text{Sym}^2) = L_{(0)}(s, \pi, \text{Sym}^2) = \frac{1}{P(q^{-s/2})}$$

We assume that $\frac{I(W, W_{\theta^{\psi}}, f_{2s-1})}{L_{reg}(s, \pi, \operatorname{Sym}^2)}$ has a pole for some $W \in \mathcal{W}(\pi, \psi)$, $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, and $f_{2s-1} \in V_{good}(2s-1, \omega_{\pi})$. Such poles are called *exceptional poles*. Then the poles of the ratio $L(s, \pi, \operatorname{Sym}^2)/L_{reg}(s, \pi, \operatorname{Sym}^2)$ are exactly exceptional poles of $L(s, \pi, \operatorname{Sym}^2)$. We now show that poles of this fraction are all simple.

Proposition 3.7. The ratio $\frac{L(s, \pi, \text{Sym}^2)}{L_{reg}(s, \pi, \text{Sym}^2)}$ has simple poles and in particular $s = s_0$ is the pole of $L(2s, \omega_{\pi}^2)$ if $s = s_0$ is an exceptional pole of $L(s, \pi, \text{Sym}^2)$.

Before proceeding the proof, we would like to mention the analytic property of the intertwining operator $M(2s-1, \omega_{\pi}^{-1})$ in [38, Lemma 4.3; 44, Lemma 3.2].

Lemma 3.8 (Takeda). The operator

$$\frac{1}{L(2s-1,\omega_{\pi}^2)}M(2s-1,\omega_{\pi}^{-1}):I(2s-1,\omega_{\pi}^{-1})\to J(1-2s,\omega_{\pi}^{-1})$$

is holomorphic for all $s \in \mathbb{C}$.

Proof of Proposition 3.7. For any $f_{2s-1} \in V_{good}(2s-1,\omega_{\pi}^{-1})$, Lemma 3.8 is exploited to take $P_i(q^{\pm s/2}) \in \mathbb{C}[q^{\pm s/2}]$ and $f_{2s-1}^{(i)} \in V_{std}(2s-1,\omega_{\pi}^{-1})$ such that $I(W, W_{\theta^{\psi}}, f_{2s-1})$ is a finite sum of the form

$$I(W, W_{\theta^{\psi}}, f_{2s-1}) = L(2s, \omega_{\pi}^2) \sum_{i=1}^m P_i(q^{\pm s/2}) I(W, W_{\theta^{\psi}}, f_{2s-1}^{(i)})$$

Indeed the fraction $\frac{I(W, W_{\theta^{\psi}}, f_{2s-1}^{(i)})}{L_{reg}(s, \pi, \text{Sym}^2)}$ is entire. This implies that $L(s, \pi, \text{Sym}^2)^{-1}$ divides a product

 $L(2s, \omega_{\pi}^2)^{-1}L_{reg}(s, \pi, \operatorname{Sym}^2)^{-1}$ in $\mathbb{C}[q^{\pm s/2}]$. Thus the poles of the ratio $\frac{L(s, \pi, \operatorname{Sym}^2)}{L_{reg}(s, \pi, \operatorname{Sym}^2)}$ are found among the poles of $L(2s, \omega_{\pi}^2)$.

In the spirit of [33], we define *exceptional* L-functions in such a way that exceptional poles can be regular but $L(s, \pi, \text{Sym}^2) \neq L_{reg}(s, \pi, \text{Sym}^2)$ precisely if there exist exceptional poles.

Definition 3.9. We let

$$L_{ex}(s, \pi, \text{Sym}^2) = \prod_{s_0} (1 - q^{s_0/2} q^{-s/2})^{-1}$$

where s_0 is taken over all exceptional poles of $L_{ex}(s, \pi, \text{Sym}^2)$.

With this said, we can factor $L(s, \pi, \text{Sym}^2)$ as

(3.4)
$$L(s,\pi,\operatorname{Sym}^2) = L_{ex}(s,\pi,\operatorname{Sym}^2)L_{reg}(s,\pi,\operatorname{Sym}^2).$$

When it is clear what ω_{π} we are working with, we will abuse the notation by letting $\omega_{\pi} = \omega_{\pi} \circ \det$. Now we would like to characterize the occurrence of exceptional poles. To this end, we need the following lemma [44, Lemma 1.15].

Lemma 3.10 (Yamana). Let ω_{π} be a quadratic character of F^{\times} . The representation $I(1, \omega_{\pi}^{-1})$ has the unique irreducible quotient, which is isomorphic to $\theta^{\psi^{-1}} \otimes \omega_{\pi}$. Furthermore the quotient map $I(1, \omega_{\pi}^{-1}) \to \theta^{\psi^{-1}} \otimes \omega_{\pi}$ is realized as the intertwining operator $M(1, \omega_{\pi}^{-1})$.

It is worthwhile noting from [12, §1.3; 44, Remark 1.7] that $\theta^{\psi^{-1}}$ is independent of ψ hence we may suppress the superscript ψ . We say that an irreducible admissible representation (π, V_{π}) of GL_2 is θ -distinguished if $\pi \otimes \theta^{\psi} \otimes \theta^{\psi^{-1}}$ admits a nonzero GL_2 -invariant trilinear form. A θ distinguished representation is characterized in terms of poles of symmetric square L-functions for self-dual representations at s = 0.

Proposition 3.11. [44, Corollary 3.9] We assume that $\dim_{\mathbb{C}} \operatorname{Hom}_{GL_2}(\pi \otimes \theta^{\psi} \otimes I(1, \omega_{\pi}^{-1}), \mathbb{C}) \leq 1$ and ω_{π} is a quadratic character. Then the following conditions are equivalent:

- (i) $\pi \otimes \omega_{\pi}$ is θ -distinguished.
- (i) $\operatorname{Hom}_{GL_2}(\pi \otimes \theta^{\psi} \otimes \theta^{\psi^{-1}} \otimes \omega_{\pi}, \mathbb{C}) \simeq \operatorname{Hom}_{GL_2}(\pi \otimes \theta^{\psi} \otimes I(1, \omega_{\pi}^{-1}), \mathbb{C}) \neq 0$ (iii) There is a non-zero functional $\Lambda : \pi \otimes \theta^{\psi} \otimes I(1, \omega_{\pi}^{-1}) \to \mathbb{C}$ which factors through the quotient $map \ \pi \otimes \theta^{\psi} \otimes I(1, \omega_{\pi}^{-1}) \to \pi \otimes \theta^{\psi} \otimes \theta^{\psi^{-1}} \otimes \omega_{\pi}.$

According to [44, Theorem 2.14], one dimensionality of the space $\operatorname{Hom}_{GL_2}(\pi \otimes \theta^{\psi} \otimes I(1, \omega_{\pi}^{-1}), \mathbb{C})$ is available for irreducible unitary representations of GL_2 . Though this might be hold for all irreducible admissible representations of GL_2 , the author does not know if the same technique in the proof of [44, Theorem 2.14] applies to these representations at this moment. To overcome this, we exploit the deformation of representations in §4. We finally come to give a characterization of exceptional poles.

Theorem 3.12. Let π be an irreducible admissible generic representation of GL_2 . Suppose that $\dim_{\mathbb{C}} \operatorname{Hom}_{GL_2}(\pi \otimes \theta^{\psi} \otimes I(1, \omega_{\pi}^{-1}), \mathbb{C}) \leq 1$. If $L_{ex}(s, \pi, \operatorname{Sym}^2)$ has a pole at s = 0, then π is θ -distinguished.

Proof. The proof is inspired by that of [20, Theorem 4.7]. We provide the complete detail to make thing concrete. The fraction

$$\frac{I(W, W_{\theta^{\psi}}, f_{2s-1})}{L(s, \pi, \text{Sym}^2)} = \frac{I(W, W_{\theta^{\psi}}, f_{2s-1})}{L_{ex}(s, \pi, \text{Sym}^2)L_{reg}(s, \pi, \text{Sym}^2)}$$

is entire in s, so we can evaluate at s = 0. For s = 0, the Zeta integral defines an intertwining map $T_0: \mathcal{W}(\pi, \psi) \times \mathcal{W}(\theta^{\psi}, \psi^{-1}) \times V_{good}(2s - 1, \omega_{\pi}^{-1}) \to \mathbb{C}$,

$$(W, W_{\theta^{\psi}}, f_{2s-1}) \mapsto \left. \frac{I(W, W_{\theta^{\psi}}, f_{2s-1})}{L(s, \pi, \operatorname{Sym}^2)} \right|_{s=0}$$

Suppose that $L_{ex}(s, \pi, \text{Sym}^2)$ has a pole at s = 0. Then the family $I(W, W_{\theta^{\psi}}, f_{2s-1})$ for good sections $f_{2s-1} \in V_{good}(2s-1, \omega_{\pi}^{-1})$ has a higher order pole at s = 0 than the family $I(W, W_{\theta^{\psi}}, f_{2s-1})$ for holomorphic sections $f_{2s-1} \in V_{hol}(2s-1, \omega_{\pi}^{-1})$ so $T_0(W, W_{\theta^{\psi}}, f_{2s-1})$ is identically zero for $f_{2s-1} \in V_{hol}(2s-1, \omega_{\pi}^{-1})$. It follows from Definition 2.5 of L-functions that T_0 is a non-zero element. Hence we choose an element $h_{2s-1} \in V_{good}(2s-1, \omega_{\pi}^{-1})$ which is not in holomorphic sections and that

$$\frac{I(W, W_{\theta^{\psi}}, h_{2s-1})}{L(s, \pi, \operatorname{Sym}^2)}\Big|_{s=0} \quad \text{is non zero.}$$

Then $h_{2s-1} = N(1-2s, \omega_{\pi}, \psi^{-1}) f_{1-2s}$ for some $f_{1-2s} \in V_{hol}(1-2s, \omega_{\pi})$. Since the exceptional *L*-function $L_{ex}(s, \pi, \text{Sym}^2)^{-1}$ divides $L(2s, \omega_{\pi}^2)^{-1}$ in $\mathbb{C}[q^{\pm s/2}]$, ω_{π} is the non-trivial quadratic chacter of F^{\times} . Thus we obtain

$$0 \neq \left. \frac{I(W, W_{\theta^{\psi}}, h_{2s-1})}{L(s, \pi, \operatorname{Sym}^2)} \right|_{s=0} = \left. \frac{I(W, W_{\theta^{\psi}}, N(1-2s, \omega_{\pi}, \psi^{-1})f_{1-2s})}{L(s, \pi, \operatorname{Sym}^2)} \right|_{s=0}$$
$$= c_3 \left. \frac{\varepsilon(1-2s, \mathbf{1}_{F^{\times}}, \psi^{-1})}{L(1-2s, \mathbf{1}_{F^{\times}})} \cdot \frac{I(W, W_{\theta^{\psi}}, {}^{\iota}[\hat{N}(1-2s, \omega_{\pi}, \psi^{-1})f_{1-2s}])}{L_{reg}(s, \pi, \operatorname{Sym}^2)} \right|_{s=0}$$

with a non-zero constant $c_3 = \frac{\operatorname{Res}_{s=0}L(2s, \mathbf{1}_{F^{\times}})}{\operatorname{Res}_{s=0}L_{ex}(s, \pi, \operatorname{Sym}^2)} \neq 0$. Consequently, Lemma 3.10 implies that the above non-trivial functional

$$(W, W_{\theta^{\psi}}, f_1) \mapsto c_3 \frac{\varepsilon(1 - 2s, \mathbf{1}_{F^{\times}}, \psi^{-1})}{L(1 - 2s, \mathbf{1}_{F^{\times}})} \cdot \frac{I(W, W_{\theta^{\psi}}, {}^{\iota}[\hat{N}(1 - 2s, \omega_{\pi}, \psi^{-1})f_{1 - 2s}])}{L_{reg}(s, \pi, \text{Sym}^2)} \bigg|_{s=0}$$

factors through the quotient $\pi \otimes \theta^{\psi} \otimes I(1, \omega_{\pi}^{-1}) \to \pi \otimes \theta^{\psi} \otimes \theta^{\psi^{-1}} \otimes \omega_{\pi}$. Appealing to [44, Lemma 1.9 (3)], one gets $\theta^{\psi^{-1}} \otimes \omega_{\pi} \simeq \theta^{\psi^{-1}}$. We conclude from Proposition 3.11 that π is θ -distinguished. \Box

For an exceptional pole at $s = s_0$, Theorem 3.12 becomes the following.

Corollary 3.13. Let π be an irreducible admissible generic representation of GL_2 . If $s = s_0$ is a pole of $L_{ex}(s, \pi, \text{Sym}^2)$ and $\dim_{\mathbb{C}} \text{Hom}_{GL_2}(\pi \nu^{\frac{s_0}{2}} \otimes \theta^{\psi} \otimes I(1, \omega_{\pi\nu^{s_0/2}}^{-1}), \mathbb{C}) \leq 1$, then $\pi \nu^{\frac{s_0}{2}}$ is θ -distinguished.

Proof. We know from [44, Remark 3.13] that $V_{good}(2s-1, \omega_{\pi\nu^{s_0/2}}^{-1}) \otimes \nu^{\frac{s_0}{2}} \simeq V_{good}(2(s+s_0)-1, \omega_{\pi}^{-1})$ and $V_{hol}(2s-1, \omega_{\pi\nu^{s_0/2}}^{-1}) \otimes \nu^{\frac{s_0}{2}} \simeq V_{hol}(2(s+s_0)-1, \omega_{\pi}^{-1})$. Therefore $L_{ex}(s, \pi, \text{Sym}^2)$ has a pole at $s = s_0$ if and only if $L_{ex}(s, \pi\nu^{\frac{s_0}{2}}, \text{Sym}^2) = L_{ex}(s+s_0, \pi, \text{Sym}^2)$ has a pole at s = 0. The result right away follows from Theorem 3.12.

We shall see later (Proposition 4.5) a kind of the converse of Corollary 3.13.

3.3. Regular *L*-functions and factorizations. We examine the regular pole at $s = s_0$ of the family $\mathcal{I}_{(0)}(\pi)$. Let d_{s_0} be its maximal order in this family. The integral $I_{(0)}(s, W, W_{\theta^{\psi}})$ has the Laurent expansion, which in case we write

$$I_{(0)}(s; W, W_{\theta^{\psi}}) = \frac{B_{(0), s_0}(W, W_{\theta^{\psi}})}{(q^{s/2} - q^{s_0/2})^{d_{s_0}}} + \text{higher order terms}$$

where W and $W_{\theta^{\psi}}$ belong to the space $\mathcal{W}(\pi_{(0)}, \psi)$ and $\mathcal{W}(\theta_{(0)}^{\psi}, \psi^{-1})$ respectively (cf. Proposition 3.1). The function W of $\mathcal{W}(\Phi^+(\pi_{(1)}), \psi)$ realized as a function $W\begin{pmatrix} a \\ 1 \end{pmatrix}$ on F^{\times} has a multiplicative support in a (cf. Proposition 3.2). Then the non-trivial bilinear form $B_{(0),s_0}$ vanishes on $\mathcal{W}(\Phi^+(\pi_{(0)}), \psi)$. As a representation of P, $\pi_{(0)}/\Phi^+(\pi_{(1)})$ is isomorphic to $\Psi^+(\pi^{(1)})$. Hence $B_{(0),s_0}$ defines a non-trivial bilinear form on $\mathcal{W}(\Psi^+(\pi^{(1)}), \psi) \times \mathcal{W}(\Phi^+(\theta_{(0)}^{\psi}), \psi^{-1})$ which is quasi-invariant with respect to the action of \tilde{P} . But by [21, §4, Proposition 4.3] (cf. [2, Proposition 3.7]), there is no non-trivial quasi-invariant pairing between $\Psi^+(\pi^{(1)})$ and $\Phi^+(\theta_{(0)}^{\psi})$. Therefore we may view $B_{(0),s_0}$ as a non-zero bilinear form on the space $\mathcal{W}(\Psi^+(\pi^{(1)}), \psi) \times \mathcal{W}(\Psi^+(\theta^{\psi^{(1)}}), \psi^{-1})$. Having the description of the bilinear from $B_{(0),s_0}$ in hand, we are in a position to show the main factorization.

Proposition 3.14. Let π be an irreducible admissible generic representation of GL_2 such that all of its derivatives are completely reducible. Then

$$L_{reg}(s, \pi, \text{Sym}^2)^{-1} = l.c.m.\{L(s, \pi_i^{(1)} \times \pi_i^{(1)})^{-1}\}$$

where the lease common multiple is with respect to divisibility in $\mathbb{C}[q^{s/2}, q^{-s/2}]$ and is taken over all irreducible constituents $\pi_i^{(1)}$ of $\pi^{(1)}$.

Proof. The proof proceeds as in those of [7, Proposition 2.3; 29, Proposition 4.14]. We take a pole $s = s_0$ of the regular *L*-function $L_{reg}(s, \pi, \text{Sym}^2)$ and let d_{s_0} be its order in $L_{reg}(s, \pi, \text{Sym}^2)$. Then it occurs as a pole of order d_{s_0} of the simple Zeta integral

$$I_{(0)}(s; W, W_{\theta^{\psi}}) = \int_{F^{\times}} W \begin{pmatrix} a \\ & 1 \end{pmatrix} W_{\theta^{\psi}} \begin{pmatrix} a \\ & 1 \end{pmatrix} |a|^{\frac{s}{2} - \frac{3}{4}} d^{\times} a$$

for some W in $\mathcal{W}(\pi, \psi)$ and $W_{\theta^{\psi}}$ in $\mathcal{W}(\theta^{\psi}, \psi^{-1})$. Moreover for any $\varphi \in \mathcal{S}(F)$ with $\varphi(0) = 1$ and $W_{\theta^{\psi}}$ in $\mathcal{W}(\theta^{\psi}, \psi^{-1})$, the integral

$$I^{1}_{(0)}(s;W,W_{\theta^{\psi}}) := \int_{F^{\times}} W\begin{pmatrix}a\\&1\end{pmatrix} W_{\theta^{\psi}}\begin{pmatrix}a\\&1\end{pmatrix} (1-\varphi(a))|a|^{\frac{s}{2}-\frac{3}{4}} d^{\times}a.$$

is always entire. Therefore $s = s_0$ is a pole of order d_{s_0} of the integral

$$I^0_{(0)}(s;W,W_{\theta^{\psi}}) := \int_{F^{\times}} W \begin{pmatrix} a \\ & 1 \end{pmatrix} W_{\theta^{\psi}} \begin{pmatrix} a \\ & 1 \end{pmatrix} \varphi(a) |a|^{\frac{s}{2} - \frac{3}{4}} d^{\times}a.$$

after writing $I_{(0)}(s; W, W_{\theta^{\psi}}) = I^0_{(0)}(s; W, W_{\theta^{\psi}}) + I^1_{(0)}(s; W, W_{\theta^{\psi}})$. Let b_1, b_2, \dots, b_s be the distinct representatives of the left coset $(F^{\times})^2 \setminus F^{\times}$. As we have indicated in (3.2), we decompose $\pi^{(1)}$ and

 $\theta^{\psi^{(1)}}$ into direct sums of simple factors $\pi^{(1)} \simeq \bigoplus_i \pi_i^{(1)}$ and $\theta^{\psi^{(1)}} \simeq \bigoplus_j (\chi_{b_j} \otimes \nu^{-\frac{1}{4}})$. Then W and $W_{\theta^{\psi}}$ can be expressed as $W = W_1 \oplus \cdots \oplus W_t$ and $W_{\theta^{\psi}} = W_{b_1} \oplus \cdots \oplus W_{b_s}$ with each W_i and W_{b_j} projecting on some characters $\pi_i^{(1)}$ and $\chi_{b_j} \otimes \nu^{-\frac{1}{4}}$ respectively. There exist *i* and *j* such that $s = s_0$ is a pole of order d_{s_0} of the integral

$$I^{0}_{(0)}(s;W_{i},W_{b_{j}}) = \int_{F^{\times}} W_{i} \begin{pmatrix} a \\ & 1 \end{pmatrix} W_{b_{j}} \begin{pmatrix} a \\ & 1 \end{pmatrix} \varphi(a)|a|^{\frac{s}{2}-\frac{3}{4}} d^{\times}a.$$

We take φ to be a characteristic function of a small neighborhood of 0. Upon applying [7, Corollary to Proposition 1.7] accompanied with Theorem 3.3, we see that

$$I_{(0)}^{0}(s;W_{i},W_{b_{j}}) = |b_{j}|^{\frac{s}{2}} \int_{F^{\times}} \pi_{i}^{(1)}(z^{2}b_{j})\chi_{b_{j}}(z^{2}b_{j})\varphi(z^{2}b_{j})|z|^{s} d^{\times}z = \alpha q^{-\beta s/2} \int_{F^{\times}} \pi_{i}^{(1)}(z)^{2}\varphi(z^{2}b_{j})|z|^{s} d^{\times}z,$$

where $\alpha q^{-\beta s/2}$ is a unit in $\mathbb{C}[q^{-s/2}]$ with $\alpha \in \mathbb{C}$ and $\beta \in \mathbb{Z}$. The last equality follows from $\chi^2_{b_i} = \mathbf{1}_{F^{\times}}$. However as we further replace $\varphi(z^2b_i)$ by $\varphi(z)$, the integral simplifies to a multiple of a standard Tate integral $I^{0}_{(0)}(s, W_{i}, W_{b_{j}}) = \alpha q^{-\beta s/2} I(s; \pi_{i}^{(1)} \times \pi_{i}^{(1)}, \varphi)$, where

$$I(s;\pi_i^{(1)} \times \pi_i^{(1)},\varphi) := \int_{F^{\times}} \pi_i^{(1)}(z)\pi_i^{(1)}(z)\varphi(z)|z|^s \ d^{\times}z.$$

Hence $I(s; \pi_i^{(1)} \times \pi_i^{(1)}, \varphi)$ has a pole of order d_{s_0} at $s = s_0$ and $L(s, \pi_i^{(1)} \times \pi_i^{(1)})$ as well. Now let $s = s_0$ be a pole of an order d_{s_0} of $L(s, \pi_i^{(1)} \times \pi_i^{(1)})$ for some *i*. It is evident that there exist a character $\pi^{(1)}$ and ϕ in $\mathcal{S}(F)$ which does not vanish at zero such that the Tate integral $I(s; \pi_i^{(1)} \times \pi_i^{(1)}, \phi)$ contributes to the pole of the order d_{s_0} at $s = s_0$. We observe that φ may in fact be taken to be a characteristic function of a neighborhood of 0 small enough because $I(s; \pi_i^{(1)} \times \pi_i^{(1)}, \phi - \varphi)$ is entire. The choice of φ allows us to choose $\varphi(z^2)$ in place of $\varphi(z)$. We pick the identity $b_j = 1$ as a coset representative of $(F^{\times})^2 \cdot 1$. Thanks to [7, Corollary to Proposition 1.7] along with Theorem 3.3, we choose W_i and W_{b_j} mapping to characters $\pi_i^{(1)}$ and χ_{b_j} via the natural normalized projection map such that

$$\begin{split} I(s;\pi_i^{(1)} \times \pi_i^{(1)},\varphi) &= \int_{F^{\times}} \pi_i^{(1)}(z^2)\chi_{b_j}(z^2)\varphi(z^2)|z|^s \ d^{\times}z \\ &= \int_{(F^{\times})^2} W_i \begin{pmatrix} z \\ & 1 \end{pmatrix} W_{b_j} \begin{pmatrix} z \\ & 1 \end{pmatrix} \varphi(z)|z|^{\frac{s}{2}-\frac{3}{4}} \ d^{\times}z = \int_{F^{\times}} W_i \begin{pmatrix} a \\ & 1 \end{pmatrix} W_{b_j} \begin{pmatrix} a \\ & 1 \end{pmatrix} \varphi(a)|a|^{\frac{s}{2}-\frac{3}{4}} \ d^{\times}a. \end{split}$$

With help of [19, Lemma 2.5], this integral $I(s; \pi_i^{(1)} \times \pi_i^{(1)}, \varphi)$ equals to $I_{reg}(s; W'_i, W_{b_j})$ for some $W'_i \in \mathcal{W}(\pi, \psi)$ and hence $s = s_0$ is a pole of order d_{s_0} of $L_{reg}(s, \pi, \text{Sym}^2)$.

We know from [7; 19, §4] that $L(s, \chi, \wedge^2) = 1$ and $L(s, \chi \times \chi) = L_{ex}(s, \chi \times \chi)$ for any character χ . The main reference for the unexplained notation $L_{ex}(s, \chi \times \chi)$ is [7, §2.2]. In order to obtain visibly consistent expressions of [7, Theorem 2.1] or [19], it is preferable to write either $L_{ex}(s, \chi, \text{Sym}^2)$ or $L(s, \chi, \text{Sym}^2)$ for $L(s, \chi \times \chi)$ when it is clear from the context. Combining Proposition 3.14 with (3.4) yields the following theorem.

Theorem 3.15. Let π be an irreducible admissible generic representation of GL_2 such that all of its derivatives are completely reducible and $\pi^{(0)} = \pi$. Then

$$L(s, \pi, \text{Sym}^2)^{-1} = l.c.m.\{L(s, \pi_i^{(j)}, \text{Sym}^2)^{-1}\}$$

where the least common multiple is with respect to divisibility in $\mathbb{C}[q^{s/2}, q^{-s/2}]$ and is taken over all j with $0 \le j \le 1$ and for all constituents $\pi_i^{(1)}$ of $\pi^{(1)}$.

4. Deformation and Specialization

4.1. Principal series representations and L-functions. In $\S4.1$ we employ a deformation method proposed by Cogdell and Piatetski-Shapiro [7, \S]. The advantage of this approach is to obtain a certain class of induced representations for which we can explicitly determine the poles of L-functions by means of shifting the location of those poles. For an admissible representation τ (which is not necessarily irreducible), we say that τ is of Whittaker type if $\operatorname{Hom}_{GL_2}(\tau, \operatorname{Ind}_N^{GL_2}(\psi))$ is of dimension 1 [7, 17, 29]. Let $\pi = \operatorname{Ind}_B^{GL_2}(\chi_1 \boxtimes \chi_2)$ be a normalized induced representation, where each χ_i is a character of F^{\times} . Let \mathcal{D}_{π} denote the complex manifold $(\mathbb{C}/\frac{2\pi i}{\log(q)}\mathbb{Z})^2$. We denote by \mathcal{D}_s the rescaled complex manifold $(\mathbb{C}/\frac{4\pi i}{\log(q)}\mathbb{Z})$. The isomorphism $\mathcal{D}_{\pi} \to (\mathbb{C}^{\times})^2$ is defined by $u = (u_1, u_2) \mapsto q^u := (q^{u_1}, q^{u_2}).$ For each $u \in \mathcal{D}_{\pi}$ we set $\pi_u = \operatorname{Ind}_B^{GL_2}(\chi_1 \nu^{u_1} \boxtimes \chi_2 \nu^{u_2}).$ Then π_u is nothing but a representation of Whittaker type for every $u \in \mathcal{D}_{\pi}$.

Definition 4.1. We say that $u = (u_1, u_2) \in \mathcal{D}_{\pi}$ is in general position if it satisfies the following conditions:

- (1) A representation $\pi_u = \operatorname{Ind}_B^{GL_2}(\chi_1 \nu^{u_1} \boxtimes \chi_2 \nu^{u_2})$ is irreducible. (2) The two characters $\pi_u^{(1,0)} = \chi_2 \nu^{u_2}$ and $\pi_u^{(0,1)} = \chi_1 \nu^{u_1}$ are distinct.
- (3) $L(s, \chi_1 \nu^{u_1}, \text{Sym}^2)$ and $L(s, \chi_2 \nu^{u_2}, \text{Sym}^2)$ do not have any common poles.
- (4) If $i \in \{1, 2\}$, then $L(s, \chi_i \nu^{u_i}, \text{Sym}^2)$ and $L(s, \chi_1 \nu^{u_1} \times \chi_2 \nu^{u_2})$ do not have any common poles. (5) If s = e is a pole of $L(2s, \omega_{\pi}^2)$, then the dimension of the space
 - $\operatorname{Hom}_{GL_2}(\sigma \otimes \theta^{\psi} \otimes I(1, \omega_{\sigma}^{-1}), \mathbb{C}) \text{ with } \sigma = \operatorname{Ind}_{B}^{GL_2}(\chi_1 \nu^{(u_1 u_2 + e)/2} \boxtimes \chi_2 \nu^{(-u_1 + u_2 + e)/2})$ is at most 1.

The condition (1) and (2) assert that outside the hyperplane defining general position, all of the derivatives are completely reducible such that each constitute is irreducible and generic. As a consequence Theorem 3.15 is applicable to the deformed representation π_u in general position. The purpose of (5) is to make it feasible to exploit the following result about distinguished representations which play an important role in our study of exceptional poles.

Theorem 4.2 ([24], Kaplan). Let π be an irreducible admissible generic representation of GL_2 and χ a character of F^{\times} .

- (i) [Theorem 4.4] The induced representation $\operatorname{Ind}_B^{GL_2}(\chi \boxtimes \chi^{-1})$ is θ -distinguished. (ii) [Corollary 4.19] If $\operatorname{Hom}_{GL_2}(\pi \otimes \theta^{\psi} \otimes \theta^{\psi^{-1}}, \mathbb{C}) \neq 0$, then $\pi \simeq \widetilde{\pi}$.

We now confirm that off a finite number of the hyperplanes in u, the deformed representation π_u is in general position.

Proposition 4.3. Let π be as above, the element u in \mathcal{D}_{π} that is not in general position belongs to a finite number of affine hyperplanes.

Proof. The conditions (1), (2), (3), and (4) are explained in [29, Proposition 5.1]. As described in [29, Proposition 5.1], (5) can be checked along the line of [44, Theorem 2.14].

The removed affine hyperplanes defining general position do not depend on $s \in \mathbb{C}$. Before going into the computation, let us recall what is known as Hartogs theorem in the view of the (complex) algebraic geometry.

Theorem 4.4 ([27, Chapter IV, §4, Theorem 4.6.7], Hartogs theorem). Let \mathcal{M} be an n-dimensional complex manifold with $n \geq 2$. If $\mathcal{N} \subseteq \mathcal{M}$ is an analytic subset of codimension 2 or more, then every holomorphic function on $\mathcal{M} - \mathcal{N}$ extends to a holomorphic function on \mathcal{M} .

The following proposition is the starting point of computing L-functions for principal series representations and is needed for demonstrating the agreement of arithmetic and analytic L-functions which we shall proceed to provide.

Proposition 4.5. Let $\pi = \operatorname{Ind}_{B}^{GL_{2}}(\chi_{1} \boxtimes \chi_{2})$ be a principal series representation of GL_{2} . Let $u = (u_{1}, u_{2}) \in \mathcal{D}_{\pi}$ be in general position and $\pi_{u} = \operatorname{Ind}_{B}^{GL_{2}}(\chi_{1}\nu^{u_{1}} \boxtimes \chi_{2}\nu^{u_{2}})$ the deformed representation. Then we have

$$L_{ex}(s, \text{Ind}_B^{GL_2}(\chi_1 \nu^{u_1} \boxtimes \chi_2 \nu^{u_2}), \text{Sym}^2) = L_{ex}(s, \chi_1 \nu^{u_1} \times \chi_2 \nu^{u_2}).$$

Proof. First we suppose that $L_{ex}(s, \operatorname{Ind}_{B}^{GL_{2}}(\chi_{1}\nu^{u_{1}}\boxtimes\chi_{2}\nu^{u_{2}}), \operatorname{Sym}^{2})$ has a pole at $s = s_{0}$. As explained in the proof of Corollary 3.13, this is amount to saying that $L_{ex}(s, \nu^{\frac{s_0}{2}} \operatorname{Ind}_B^{GL_2}(\chi_1 \nu^{u_1} \boxtimes \chi_2 \nu^{u_2}), \operatorname{Sym}^2)$ has a pole at s = 0. Proposition 3.7 says that the pole s = 0 is simple and that it appears amongst that of $L(2s, \omega_{\pi_{\mu}\nu^{s_0/2}}^2)$. Unraveling the unramified twist, $L(2s, \omega_{\pi}^2)$ has a poles at s = e with $e = u_1 + u_2 + s_0$ and σ in Definition 4.1-(5) becomes $\pi_u \nu^{s_0/2}$. We apply Theorem 3.12 to $\pi_u \nu^{s_0/2}$ and deduce from Theorem 4.2 that $\pi_u \nu^{s_0/2}$ is self-contragredient. In other words

$$\operatorname{Ind}_{B}^{GL_{2}}(\chi_{1}\nu^{u_{1}}\boxtimes\chi_{2}\nu^{u_{2}})^{\sim}\simeq\operatorname{Ind}_{B}^{GL_{2}}(\chi_{1}\nu^{u_{1}}\boxtimes\chi_{2}\nu^{u_{2}})\nu^{s_{0}}.$$

The only way this is possible is that

- (i) $(\chi_1 \nu^{u_1})^{\sim} \simeq \chi_1 \nu^{u_1 + s_0}$ and $(\chi_2 \nu^{u_2})^{\sim} \simeq \chi_2 \nu^{u_2 + s_0}$ (ii) $(\chi_1 \nu^{u_1})^{\sim} \simeq \chi_2 \nu^{u_2 + s_0}$.

In (i) the locus is defined by two independent equations $q^{-(2u_1+s)}\chi_1^2(\varpi) = 1$ and $q^{-(2u_2+s)}\chi_2^2(\varpi) =$ 1, after evaluating both sides at ϖ , and hence will be of codimension 2. Due to Bernstein's Theorem [7, §3], viewing $I(W_u, W_{\theta^{\psi}}, f_{2s-1})$ as rational functions in $\mathbb{C}(q^{-s/2}, q^{-u})$, every singularities of the integral $I(W_u, W_{\theta^{\psi}}, f_{2s-1})$ must be accounted for the form (*ii*) $(\chi_1 \nu^{u_1})^{\sim} \simeq \chi_2 \nu^{u_2+s_0}$ by Hartogs theorem, Theorem 4.4. As a result $L_{ex}(s, \chi_1 \nu^{u_1} \times \chi_2 \nu^{u_2})$ has a pole at $s = s_0$.

Next we assume that $L_{ex}(s, \chi_1 \nu^{u_1} \times \chi_2 \nu^{u_2})$ has a pole at $s = s_0$. Equivalently $L_{ex}(s, \chi_1 \nu^{u_1 + \frac{s_0}{2}} \times$ $\chi_2 \nu^{u_2 + \frac{s_0}{2}}$ possesses a pole at s = 0. Then we have $(\chi_1 \nu^{u_1 + \frac{s_0}{2}})^{\sim} \simeq \chi_2 \nu^{u_2 + \frac{s_0}{2}}$ and the principal series representation $\pi_u \nu^{\frac{s_0}{2}} = \operatorname{Ind}_B^{GL_2}(\chi_1 \nu^{u_1 + \frac{s_0}{2}} \boxtimes \chi_2 \nu^{u_2 + \frac{s_0}{2}})$ is self-contragredient. In terms of central characters, this says that $\omega^2_{\pi_u \nu^{s_0/2}}$ is trivial. Now Theorem 4.2 assures that $\pi_u \nu^{s_0/2}$ is θ -distinguished. At this point, we essentially repeat the argument of [44, Theorem 3.7-(1)] for completeness. We further assume that $L_{ex}(s, \pi_u \nu^{\frac{s_0}{2}}, \text{Sym}^2)$ is holomorphic at s = 0. Since Lfunctions $L(s, \chi_1 \nu^{u_1 + \frac{s_0}{2}} \times \chi_2 \nu^{u_2 + \frac{s_0}{2}})^{-1}$, $L(s, \chi_1 \nu^{u_1 + \frac{s_0}{2}}, \operatorname{Sym}^2)^{-1}$ and $L(s, \chi_2 \nu^{u_2 + \frac{s_0}{2}}, \operatorname{Sym}^2)^{-1}$ are relatively prime, Theorem 3.15 implies that $L(s, \pi_u \nu^{\frac{s_0}{2}}, \text{Sym}^2)$ is holomorphic at s = 0. Remembering a quadratic character $\omega_{\pi_u\nu^{s_0/2}}$, Lemma 3.16 in [44] enables us to take $W \in \mathcal{W}(\pi_u\nu^{\frac{s_0}{2}},\psi)$, $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ and $f_{2s-1} \in V_{good}(2s-1, \omega_{\pi, \nu^{s_0/2}}^{-1})$ such that

(4.1)
$$M(1, \omega_{\pi_u\nu^{s_0/2}}^{-1})f_1 = 0 \quad \text{and} \quad \lim_{s \to 1} I(W, W_{\theta^{\psi}}, \hat{N}(2s - 1, \omega_{\pi_u\nu^{s_0/2}}^{-1})f_{2s-1}) \neq 0.$$

Evaluating the both side of the functional equation (2.8) at s = 1, we arrive at

$$0 \neq \frac{I(W, W_{\theta^{\psi}}, \hat{N}(1, \omega_{\pi_{u}\nu^{s_{0}/2}}^{-1})f_{1})}{L(0, \pi_{u}\nu^{s_{0}/2}, \operatorname{Sym}^{2})} = \varepsilon(1, \pi, \operatorname{Sym}^{2}, \psi) \frac{I(W, W_{\theta^{\psi}}, f_{1})}{L(1, \pi_{u}\nu^{s_{0}/2}, \operatorname{Sym}^{2})}$$

But the first condition of (4.1) enforces that the linear functional

$$(W, W_{\theta^{\psi}}, f_1) \mapsto \varepsilon(1, \pi, \operatorname{Sym}^2, \psi) \frac{I(W, W_{\theta^{\psi}}, f_1)}{L(1, \pi_u \nu^{s_0/2}, \operatorname{Sym}^2)}$$

cannot factor through the quotient map $\pi_u \nu^{s_0/2} \otimes \theta^{\psi} \otimes I(1, \omega_{\pi_u \nu^{s_0/2}}^{-1}) \to \pi_u \nu^{s_0/2} \otimes \theta^{\psi} \otimes \theta^{\psi^{-1}} \otimes \omega_{\pi_u \nu^{s_0/2}}$ and hence $\pi_u \nu^{s_0/2}$ is not θ -distinguished making the use of Proposition 3.11. This is a contradiction and we conclude that $L_{ex}(s, \pi_u \nu^{\frac{s_0}{2}}, \operatorname{Sym}^2)$ has a pole at s = 0, which is equivalent to saying that $L_{ex}(s, \pi_u, \operatorname{Sym}^2)$ has a pole at $s = s_0$.

We shift our gear to the principal series representation $\pi = \text{Ind}_B^{GL_2}(\chi_1 \boxtimes \chi_2)$ of GL_2 . The representation is a possibly reducible representation. Nonetheless all the theories in Section 3.2 still go through and all the constructions of $I(W, W_{\theta^{\psi}}, f_{2s-1}), L(s, \pi, \text{Sym}^2), L_{ex}(s, \pi, \text{Sym}^2)$ and $L_{reg}(s, \pi, \text{Sym}^2)$ are completely carried over without any adjustments.

Theorem 4.6. Let $\pi = \operatorname{Ind}_B^{GL_2}(\chi_1 \boxtimes \chi_2)$ be a principal series representation of GL_2 . Let $u = (u_1, u_2) \in \mathcal{D}_{\pi}$ be in general position and $\pi_u = \operatorname{Ind}_B^{GL_2}(\chi_1 \nu^{u_1} \boxtimes \chi_2 \nu^{u_2})$ the deformed representation. Then we have the following:

 $\begin{aligned} \text{(i)} \ \ L(s, \operatorname{Ind}_{B}^{GL_{2}}(\chi_{1}\nu^{u_{1}} \boxtimes \chi_{2}\nu^{u_{2}}), \operatorname{Sym}^{2}) &= L(s + u_{1} + u_{2}, \chi_{1} \times \chi_{2}) \prod_{1 \leq i \leq 2} L(s + 2u_{i}, \chi_{i}, \operatorname{Sym}^{2}). \\ \text{(ii)} \ \ L(s, \operatorname{Ind}_{B}^{GL_{2}}(\chi_{1} \boxtimes \chi_{2}), \operatorname{Sym}^{2})^{-1} \ \ divides \ \ L(s, \chi_{1} \times \chi_{2})^{-1} \prod_{1 \leq i \leq 2} L(s, \chi_{i}, \operatorname{Sym}^{2})^{-1} \ \ in \ \mathbb{C}[q^{\pm s/2}], \\ \text{that is, there is a } Q(X) \in \mathbb{C}[X] \ such \ that \\ \ \ L(s, \operatorname{Ind}_{B}^{GL_{2}}(\chi_{1} \boxtimes \chi_{2}), \operatorname{Sym}^{2}) = Q(q^{-s/2})L(s, \chi_{1} \times \chi_{2}) \prod_{1 \leq i \leq 2} L(s, \chi_{i}, \operatorname{Sym}^{2}). \end{aligned}$

Proof. The first item is a direct consequence of Proposition 4.5 together with Theorem 3.15. Regarding the second item, it follows from the standard Bernstein's argument of the continuation principle [7, §3] that $I(W_u, W_{\theta^{\psi}}, f_{2s-1})$ defines a rational function in $\mathbb{C}(q^{\pm s/2}, q^{\pm u})$ as we vary in u. According to (i), the fraction

(4.2)
$$\frac{I(W_u, W_{\theta^{\psi}}, f_{2s-1})}{L(s+u_1+u_2, \chi_1 \times \chi_2) \prod_{1 \le i \le 2} L(s+2u_i, \chi_i, \text{Sym}^2)}$$

has no poles on Zariski open set of u in general position. Our remaining task is to the ration (4.2) in fact lies in $\mathbb{C}[q^{\pm s/2}, q^{\pm u}]$. But the proof of holomorphy continues as in [7, Proposition 4.1], (See [18, Proposition 5.3; 29, Lemma 5.1] for further refinements of basic ideas). If we now specialize at u = 0 we conclude that

$$\frac{I(W, W_{\theta^{\psi}}, f_{2s-1})}{L(s, \chi_1 \times \chi_2) \prod_{1 \le i \le 2} L(s, \chi_i, \operatorname{Sym}^2)}$$

has no poles for all $W \in \mathcal{W}(\pi, \psi), W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ and $f_{2s-1} \in V_{acod}(2s-1, \omega_{\pi}^{-1}).$

As elucidated in the Langlands-Shahidi method [8, 9, 34], the unit appearing in Proposition 4.7-

(*ii*) will be presumably 1. This is so-called the *multiplicativity* of γ -factors. However manifesting the multiplicativity property requires manipulating integrals in a delicate manner. Apparently the best we can do at this point is to obtain the weak multiplicativity which is enough for the applications therein.

Proposition 4.7. Let $\pi = \operatorname{Ind}_{B}^{GL_2}(\chi_1 \boxtimes \chi_2)$ be a principal series representation of GL_2 . Let $u = (u_1, u_2) \in \mathcal{D}_{\pi}$ be in general position and $\pi_u = \operatorname{Ind}_{B}^{GL_2}(\chi_1 \nu^{u_1} \boxtimes \chi_2 \nu^{u_2})$ the deformed representation. Then we have the following:

(i) $\gamma(s, \pi_u, \operatorname{Sym}^2, \psi)$ and $\gamma(s + u_1 + u_2, \chi_1 \times \chi_2, \psi) \prod_{1 \le i \le 2} \gamma(s + 2u_i, \chi_i, \operatorname{Sym}^2, \psi)$ are equal up to a unit in $\mathbb{C}[q^{\pm s/2}, q^{\pm u}]$. (ii) $\gamma(s, \pi, \operatorname{Sym}^2, \psi)$ and $\gamma(s, \chi_1 \times \chi_2, \psi) \prod_{1 \le i \le 2} \gamma(s, \chi_1, \operatorname{Sym}^2, \psi)$ are equal up to a unit in $\mathbb{C}[q^{\pm s/2}]$.

(*ii*) $\gamma(s, \pi, \operatorname{Sym}^2, \psi)$ and $\gamma(s, \chi_1 \times \chi_2, \psi) \prod_{1 \le i \le 2} \gamma(s, \chi_i, \operatorname{Sym}^2, \psi)$ are equal up to a unit in $\mathbb{C}[q^{\pm s/2}]$. *Proof.* The proof is standard [18, Proposition 5.4; 29, Proposition 5.5] and is due to Cogdell and

Proof. The proof is standard [18, Proposition 5.4; 29, Proposition 5.5] and is due to Cogdell and Piatetski-Shapiro [7, Proposition 4.3] by applying Proposition 2.6, Theorem 4.6, and Proposition 5.7 to our setting. \Box

To proceed further we adopt the terminology from [7,29]. We say that an admissible representation of GL_2 is of Langlands type if Π is of the form $\operatorname{Ind}_B^{GL_2}(\pi_1\nu^{u_1} \boxtimes \pi_2\nu^{u_2})$, where each π_i is a square integrable representation of GL_{n_i} , $n_1 + n_2 = 2$, each u_i is real and they are ordered so that $u_1 \geq u_2$. Let π be an irreducible admissible representation of GL_2 . A representation of the form $\chi \circ$ det with χ a character of F^{\times} is an example of an irreducible and admissible but not generic representation of GL_2 . Regardless of being generic, π can be realized as the unique Langlands quotient of Langlands type $\Pi = \operatorname{Ind}_B^{GL_2}(\pi_1\nu^{u_1}\boxtimes \pi_2\nu^{u_2})$ which is of Whittaker type. The L-function $L(s, \pi, \operatorname{Sym}^2)$ is defined to be

$$L(s, \pi, \operatorname{Sym}^2) := L(s, \Pi, \operatorname{Sym}^2).$$

Lemma 4.8. Let $\pi = \operatorname{Ind}_B^{GL_2}(\pi_1 \boxtimes \pi_2)$ be a principal series representation of GL_2 with each π_i a character of F^{\times} (possibly a representation of Langlands type). Then $L(s, \pi_2, \operatorname{Sym}^2)^{-1}$ divides $L(s, \pi, \operatorname{Sym}^2)^{-1}$ in $\mathbb{C}[q^{\pm s/2}]$, that is, $L(s, \pi_2, \operatorname{Sym}^2) = Q(q^{-s/2})L(s, \pi, \operatorname{Sym}^2)$ for some $Q(X) \in \mathbb{C}[X]$.

Proof. Jacquet, Piatetski-Shapiro, and Shalika established in [17, Proposition 9.1] that for a character π_2 and a Schwartz-Bruhat function $\varphi \in \mathcal{S}(F)$, there exists $W \in \mathcal{W}(\pi, \psi)$ such that

(4.3)
$$W\begin{pmatrix}a\\&1\end{pmatrix} = \pi_2(a)\varphi(a)|a|^{\frac{1}{2}}.$$

The divisibility of *L*-functions follows exactly in the same way as we argued in the proof of Proposition 3.14, appealing to the property above (4.3) instead of [7, Corollary to Proposition 1.7].

We denote by $P \sim Q$ that the ratio is a unit in $\mathbb{C}[q^{s/2}, q^{-s/2}]$ for two rational functions $P(q^{-s/2})$ and $Q(q^{-s/2})$ in $\mathbb{C}(q^{-s/2})$. Unlike the Langlands-Shahidi method [8,9] which takes the following formalism (4.4) to be the definition, proving the inductive formula (4.4) is the crucial subject in the theory of Rankin-Selberg integrals.

Theorem 4.9. Let $\pi = \text{Ind}_B^{GL_2}(\pi_1 \nu^{u_1} \boxtimes \pi_2 \nu^{u_2})$ be a representation of Langlands type of GL_2 . Then we have

(4.4)
$$L(s,\pi,\mathrm{Sym}^2) = L(s+u_1+u_2,\pi_1\times\pi_2)\prod_{1\le i\le 2}L(s+2u_i,\pi_i,\mathrm{Sym}^2).$$

Proof. The proof is identical with those employed in [7, Theorem 4.1; 29, Theorem 5.1]. \Box

As a result, Theorem 4.9 in the case of irreducible unramified representations yields:

Corollary 4.10. Let $\pi = \text{Ind}_B^{GL_2}(\mu_1 \boxtimes \mu_2)$ be an irreducible unramified representation of GL_2 . Then we have

$$L(s, \pi, \operatorname{Sym}^2) = \prod_{1 \le i \le j \le 2} \frac{1}{1 - \mu_i(\varpi)\mu_j(\varpi)q^{-s}}$$

Corollary 4.10 provides an affirmative answer to Yamana's question [44, §3H] for the unramified GL_2 -case at least.

4.2. The Langlands correspondence and the equality of *L*-functions. We denote by W'_F the Weil-Deligne group of *F* (see for example [39]). We then denote by $L(s, \text{Sym}^2(\rho))$ the Artin *L*-function attached to the 2-dimensional (complex) Frobenius semi-simple representation ρ of W'_F , where $\text{Sym}^2(\rho)$ is the symmetric square of ρ .

Theorem 4.11. Let $\pi \mapsto \rho(\pi)$ be the local Langlands correspondence from the set of isomorphism classes of 2-dimensional complex representations of W'_F to that of the isomorphism classes of irreducible admissible representations of $GL_2(F)$. Then we have the following equality of L-functions:

$$L(s, \pi, \operatorname{Sym}^2) = L(s, \operatorname{Sym}^2(\rho(\pi)))$$

Proof. The case of π a discrete series representation (also called a special or Steinberg representation) was established in [44], so it remains to consider the case when π is the induced representation of Langlands type. The proof in this case is a consequence of Theorem 4.9 accompanied with the local Langlands correspondence. We refer the reader to the beginning of [29, Section 5.3] for further details.

We turn attention to the factorization

$$L(s,\rho(\pi)\otimes\rho(\pi))=L(s,\wedge^2(\rho(\pi)))L(s,\mathrm{Sym}^2(\rho(\pi)))$$

according to Langlands formalism. We derive from [18, Theorem 5.14] and [19, Proposition 4.1] pertaining $L(s, \pi, \wedge^2) = L(s, \omega_{\pi})$ that the analogous factorization is available in the realm of integral representations.

Corollary 4.12. Let π be an irreducible admissible representation of $GL_2(F)$. Then we obtain

$$L(s, \pi \times \pi) = L(s, \pi, \wedge^2)L(s, \pi, \operatorname{Sym}^2) = L(s, \omega_{\pi})L(s, \pi, \operatorname{Sym}^2),$$

where $L(s, \pi, \wedge^2)$ is the Jacquet-Shalika L-function defined in [19].

5. Stability of γ -factors

5.1. Howe vectors and Bessel functions. In §5.1, we extend the basic theory of Howe vectors [5,45] to metaplectic groups \widetilde{GL}_2 . Each of the constructions in turn had its origin in the work of R. Howe [15]. We assume that ψ is unramified, that is, ψ is trivial on \mathcal{O} while $\psi(\varpi^{-1}\mathcal{O}) \neq 1$. We put $d_m = t(\varpi^{-m}, \varpi^m)$ and let $J_m = d_m K_m d_m^{-1}$. Then J_m is given by

$$J_m = \begin{pmatrix} 1 + \mathfrak{p}^m & \mathfrak{p}^{-m} \\ \mathfrak{p}^{3m} & 1 + \mathfrak{p}^m \end{pmatrix}.$$

For $k = (k_{ij}) \in K_m$, we define a character τ_m of K_m by

$$\tau_m(k) = \psi(\varpi^{-2m}k_{12})$$

As ψ is unramified, we can check that τ_m is indeed a character on K_m . We define a character ψ_m on J_m by

$$\psi_m(j) = \tau_m(d_m^{-1}jd_m) \text{ for } j \in J_m.$$

We can also see that ψ and ψ_m agree on N_m where $N_m = N \cap J_m$. For the rest of Section 5 we let π be an irreducible admissible representation of GL_2 . If π is generic $\pi \simeq \mathcal{W}(\pi, \psi)$ while if π is not generic then we set $\mathcal{W}(\Pi, \psi)$ to be a Whittaker model for π where Π is the induced representation of Langlands type having π as its unique irreducible quotient. We fix a Whittaker function $W \in \mathcal{W}(\pi, \psi)$ such that $W(I_2) = 1$. For $m \ge 1$, we define a function W_m on GL_2 by

$$W_m(g) = \frac{1}{\operatorname{vol}(N_m)} \int_{N_m} W(gn) \psi^{-1}(n) \ dn.$$

The construction of Howe Whittaker functions can be adapted to the metaplectic setting with GL_2 replaced by GL_2 . We take $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ to be $W_{\theta^{\psi}}(I_2) = 1$. In like manner we define a function $W_{\theta^{\psi},m}$ on GL_2 for $m \ge 1$ by

$$W_{\theta^{\psi},m}(\widetilde{g}) = \frac{1}{\operatorname{vol}(N_m)} \int_{N_m} W_{\theta^{\psi}}(\widetilde{g}\mathfrak{s}(n))\psi(n) \ dn.$$

 $W_{\theta^{\psi},m}$ is simply W_1 in the notation of [12, Lemma 4.1.1]. Many of results in §5.1 resemble those in the setting of the general linear group GL_2 [36, Lemma 4.1; 45, §5.1], henceforth we only remark on the nature of differences or omit the proof most of the time.

Lemma 5.1. We choose $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ so that $W_{\theta^{\psi}}(I_2) = 1$. Let L be a positive integer such that $\theta^{\psi}(\mathfrak{s}(K_L))W_{\theta^{\psi}} = W_{\theta^{\psi}}$. Then we have

- (1) $W_{\theta^{\psi},m}(I_2) = 1.$
- (2) If $m \ge L$ then $W_{\theta^{\psi},m}(\widetilde{g}\mathfrak{s}(j)) = \psi_m^{-1}(j)W_{\theta^{\psi},m}(\widetilde{g})$ for all $j \in J_m$.
- (3) If $m \geq k$, then

$$W_{\theta^{\psi}.m}(\tilde{g}) = \frac{1}{\operatorname{vol}(N_m)} \int_{N_m} W_{\theta^{\psi},k}(\tilde{g}\mathfrak{s}(n))\psi(n)dn$$

We now fix $W \in \mathcal{W}(\pi, \psi)$ and $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ satisfying $W(I_2) = W_{\theta^{\psi}}(I_2) = 1$. Let L be an integer such that W and $W_{\theta\psi}$ are right invariant under K_L . The vector $W_{\theta\psi,m}$ is called Howe vectors of θ^{ψ} if $m \geq L$. A parallel notation W_m is applied to W.

Lemma 5.2. Let $m \ge L$ and $W_{\theta^{\psi},m}$ be as in Lemma 5.1. Then we have

- (1) $W_{\theta^{\psi} m}(\mathfrak{s}(t(a,1))) \neq 0$ if and only if $a \in 1 + \mathfrak{p}^{m}$.
- (2) If $W_{\theta^{\psi},m}(\mathfrak{s}(t(a,1))\mathfrak{s}(w_2)) \neq 0$, then $a \in \mathfrak{p}^{-3m}$.

In virtue of Lemma 5.1-(2) the Whittaker function $W_{\theta^{\psi},m}$ satisfies

$$W_{\theta^{\psi},m}(\mathfrak{s}(n_1)\widetilde{g}\mathfrak{s}(n_2)) = \psi^{-1}(n_1)\psi^{-1}(n_2)W_{\theta^{\psi},m}(\widetilde{g}) \text{ for all } n_1 \in N, n_2 \in N_m \text{ and } \widetilde{g} \in \widetilde{GL}_2.$$

The transformation implies that $W_{\theta^{\psi},m}$ really behaves partially like a Bessel function and $W_{\theta^{\psi},m}$ is said to be a partial Bessel function. In particular we are interested in the full Bessel function attached to the representation θ^{ψ} for the further analysis. For $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ the integral

$$\ell(W,x) = \int_{N_m} W_{\theta^{\psi}} \left(\mathfrak{s} \begin{pmatrix} x \\ & 1 \end{pmatrix} \mathfrak{s}(w_2) \mathfrak{s}(n) \right) \psi(n) dn$$

converges in the sense of stabilizing for large m depending on x. See [12, Lemma 4.1]. It defines a Whittaker functional on $V_{\theta^{\psi}}$ for fixed $x \in F$, as does the functional $v \mapsto W_{\theta^{\psi},v}(I_2)$. It follows from the uniqueness of Whittaker functional $[12, \S2.1]$ that there is the constant proportionality as a function of x when v varies, that is,

$$j_{\theta^{\psi}}(x)W_{\theta^{\psi},v}(I_2) = \int_{N_m} W_{\theta^{\psi},v} \left(\mathfrak{s} \begin{pmatrix} x \\ & 1 \end{pmatrix} \mathfrak{s}(w_2)\mathfrak{s}(n) \right) \psi(n) dn.$$

The function $j_{\theta^{\psi}}(x)$ is called the *Bessel function*. The basic behavior of Bessel functions is carried out as a part of Ph.D. Thesis by Wang [41, p. 37] supervised by Rallis. The proof is in turn based on the work of Gelbart and Piatetski-Shapiro [12, Proposition 4.4.2] and Soudry [36, Lemma 4.1].

Proposition 5.3. Let $L \ge 1$ and $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$ be as in Lemma 5.1.

(1) (Asymptotics for Bessel function on a certain range) For every $m \ge L$, there is a positive constant $C_m = q^{9m}$ such that if $|x| > C_m$, then

$$j_{\theta^{\psi}}(x) = \int_{xu^{-2} \in 1+\mathfrak{p}^{3m}} (-x, u)_F \mu_{\psi}(u)^{-1} \psi\left(xu^{-1} + u\right) du.$$

- (2) (Bessel functions) Let M be a positive integer such that $q^{6M} \ge C_L$. If $i \ge M$, then $\operatorname{vol}(N_{3i})W_{\theta^{\psi},3i}(\mathfrak{s}(t(x,1))\mathfrak{s}(w_2)) = j_{\theta^{\psi}}(x) \quad \text{for all} \quad |x| \le q^{6i}.$
- (3) (Local uniform smoothness) Let $i \ge M$. For $x \in \mathfrak{p}^{-6i}$ and $a \in 1 + \mathfrak{p}^{3i}$,

$$j_{\theta\psi}(xa) = j_{\theta\psi}(x).$$

5.2. Intertwining operators. In §5.2, we are concerned with computing the image of the test function (3.3) under the intertwining operator. We denote by χ a character of F^{\times} . Let X be an open compact subgroup of N. For $x \in X$ and i > 0, we set

$$A(x,i) = \{ \overline{n} \in \overline{N} \mid \overline{n}x \in Z^2 P \overline{N}_i \}.$$

Lemma 5.4. For any positive integer c there exists an integer $i_1 = i_1(X, c)$ such that for all $i \ge i_1$, $x \in X$ and $\overline{n} \in A(x, i)$ we have

(5.1)
$$\overline{n}x = t(\alpha, \alpha)t(\beta, 1)n\overline{n}_0$$

with $\alpha, \beta \in 1 + \mathfrak{p}^c$, $n \in N$ and $\overline{n}_0 \in \overline{N}_i$. Furthermore there is an integer $i_2 = i_2(X)$ such that for all $i \geq i_2$ we have $A(x, i) = \overline{N}_i$.

Proof. We mimic a train of the standard argument from [5, Lemma 3.7]. Since $\overline{n}(y)n(x) \in Z^2 P \overline{N}_i$, $\overline{n}(y)n(x)$ takes a form

$$\overline{n}(y)n(x) = \begin{pmatrix} \alpha\beta & v \\ & \alpha \end{pmatrix} \overline{n}(\overline{y})$$

for $\alpha \in (F^{\times})^2$, |x| < M and $\overline{y} \in \mathfrak{p}^{3i}$. For convenience, we rewrite it as

(5.2)
$$\begin{pmatrix} 1 - x\overline{y} & x \\ -\overline{y} & 1 \end{pmatrix} = n(x)\overline{n}(-\overline{y}) = \overline{n}(-y) \begin{pmatrix} \alpha\beta & v \\ \alpha \end{pmatrix} = \begin{pmatrix} \alpha\beta & v \\ -\alpha\beta y & -vy+\alpha \end{pmatrix}.$$

Looking at the left entries in (5.2), we have $1 - x\overline{y} = \alpha\beta$ and $\overline{y} = \alpha\beta y$. It is quickly seen from the condition |x| < M that for any positive integer c we can pick $i_1(X, c)$ such that $\alpha\beta = 1 - x\overline{y} \in 1 + \mathfrak{p}^c$. Taking the determinant on the both sides of (5.2), we obtain $\alpha^2\beta = 1$, which tells us that $\alpha \in 1 + \mathfrak{p}^c$. Hence β must be in $1 + \mathfrak{p}^c$. Now we put $i_2(X) := i_1(X, 1)$ and get $y \in \mathfrak{p}^{3i}$ from $y = (\alpha\beta)^{-1}\overline{y}$. \Box

Let us rewrite (3.3) in the current setting for later use. Given a positive integer *i* and a complex number $s \in \mathbb{C}$, we define the following function $f_{2s-1}^i \in I(2s-1, \omega_{\pi\otimes\chi}^{-1})$ on \widetilde{GL}_2 by

$$f_{2s-1}^{i}(\widetilde{g}) = \begin{cases} \xi \omega_{\pi \otimes \chi}^{-1}(a) \delta_{B}(p)^{(2s+1)/4} & \text{if } \widetilde{g} = (1,\xi) \mathfrak{s}(t(a,a)) \mathfrak{s}(p) \mathfrak{s}(\overline{n}) & \text{with } a \in (F^{\times})^{2}, p \in P, \overline{n} \in \overline{N}_{i}, \\ 0 & \text{otherwise.} \end{cases}$$

The section is the \widetilde{GL}_2 -analogue to that of [5, Lemma 3.8]. We let \widetilde{f}_{1-2s}^i denote $M(2s-1, \omega_{\pi\otimes\chi}^{-1})f_{2s-1}^i$ the image of f_{2s-1}^i under the intertwining operator $M(2s-1, \omega_{\pi\otimes\chi}^{-1})$. We now evaluate f_{2s-1}^i .

Proposition 5.5. Let X be a compact open subset of N. Then there exists an integer $I(X, \omega_{\pi \otimes \chi})$ such that for all $i \ge I(X, \omega_{\pi \otimes \chi})$ and $x \in X$, we have

$$\widetilde{f}_{1-2s}^{i}(\mathfrak{s}(w_2)\mathfrak{s}(n(x))) = \operatorname{vol}(\overline{N}_i)$$

Proof. We choose a positive integer c such that $\omega_{\pi\otimes\chi}$ is trivial on $1+\mathfrak{p}^c$. Then we choose $I(X, \omega_{\pi\otimes\chi})$ to be the maximum of $i_1(X, c)$ and $i_2(X)$. Since

$$\widetilde{f}_{1-2s}^{i}(\mathfrak{s}(w_2)\mathfrak{s}(n(x))) = \int_F f_{2s-1}(\mathfrak{s}(w_2)\mathfrak{s}(n(y)\mathfrak{s}(w_2)\mathfrak{s}(n(x)))dx,$$

the value of \widetilde{f}_{1-2s}^i is completely determined by the support of the function $f_{2s-1}(\mathfrak{s}(\overline{n}(y)n(x)))$. In fact $\overline{N}^*N^* = \mathfrak{s}(\overline{N}N)$ because $\sigma_2(\overline{n}(y), n(x)) = 1$ for all $x, y \in F$. Then elements $\overline{n}(y)n(x)$ in $Z^2P\overline{N}$

contribute to the support. By writing $\overline{n}(y)n(x) = t(\alpha, \alpha)t(\beta, 1)n\overline{n}_0$ as in (5.1), Lemma 5.4 implies that

$$f_{2s-1}(\mathfrak{s}(w_2)\mathfrak{s}(n(y)\mathfrak{s}(w_2)\mathfrak{s}(n(x))) = \omega_{\pi\otimes\chi}^{-1}(\alpha)\chi^{-2}(\alpha)\delta_B^{\frac{2s+1}{4}}(t(\beta,1))f(\mathfrak{s}(\overline{n}_0)) = 1.$$

But again by Lemma 5.4, this only happens when $n(x) \in A(x,i) = \overline{N}_i$. Therefore we have $\widetilde{f}_{1-2s}^i(\mathfrak{s}(w_2)\mathfrak{s}(n(x))) = \operatorname{vol}(\overline{N}_i)$ as claimed.

5.3. Dependence on ψ . In order to move to twisted Rankin-Selberg integrals, we need some preparation. For $W \in \mathcal{W}(\pi, \psi)$, $W_{\theta^{\psi}} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, $f_{2s-1} \in I(2s-1, \omega_{\pi\otimes\chi}^{-1})$ and a complex number s, we define the integral

$$I(W, W_{\theta^{\psi}}, f_{2s-1}, \chi) = \int_{Z^2 N \setminus GL_2} W(g) W_{\theta^{\psi}}(\mathfrak{s}(g)) f_{2s-1}(\mathfrak{s}(g)) \chi(\det(g)) dg.$$

By the usage of (2.6), we reframe the functional equation of the γ -factor (2.8) by

$$I(W, W_{\theta^{\psi}}, \hat{N}(2s-1, \omega_{\pi}^{-1})f_{2s-1}, \chi) = \gamma(s, \pi \otimes \chi, \text{Sym}^{2}, \psi)I(W, W_{\theta^{\psi}}, f_{2s-1}, \chi).$$

The main theme of §5.3 is to form so-called the dependence on ψ , analogous to [12, Lemma 5.1.3] and [45, Lemma 2.8], in the setting of $\gamma(s, \pi \otimes \chi, \text{Sym}^2, \psi)$.

Proposition 5.6. For any $a \in F^{\times}$, let $\mu_{\psi_a}(a) = \mu_{\psi}(a)(a, a)_F$. Then

$$\gamma(s,\pi\otimes\chi,\mathrm{Sym}^2,\psi_a)=\mu_{\psi_a}(a)^{-1}\omega_{\pi}(a)^{3}\chi(a)^{6}|a|^{3\left(s-\frac{1}{2}\right)}\gamma(s,\pi\otimes\chi,\mathrm{Sym}^2,\psi).$$

Proof. Let $W^a(g) := W(t(a,1)g)$ and $W^a_{\theta\psi}(\mathfrak{s}(g)) := W_{\theta\psi}(\mathfrak{s}(t(a,1))\mathfrak{s}(g))$. We can check that W^a is in $\mathcal{W}(\pi,\psi_a)$ and $W^a_{\theta\psi}$ belongs to $\mathcal{W}(\theta\psi_a,\psi_a^{-1})$ (cf. [12, Lemma 5.1.3]). For $W^a \in \mathcal{W}(\pi,\psi_a)$ and $W^a_{\theta\psi} \in \mathcal{W}(\theta^{\psi_a},\psi_a^{-1})$ and $f_{2s-1} \in I(2s-1,\omega_{\pi\otimes\chi}^{-1})$, we have

$$\begin{split} I(W^{a}, W^{a}_{\theta^{\psi}}, f_{2s-1}, \chi) \\ &= \int_{Z^{2}N \setminus GL_{2}} W(t(a, 1)g) W_{\theta^{\psi}}(\mathfrak{s}(t(a, 1))\mathfrak{s}(g)) f_{2s-1}(\mathfrak{s}(t(a^{-1}, 1))\mathfrak{s}(t(a, 1))\mathfrak{s}(g)) \chi(\det(g)) dg. \end{split}$$

By changing variables $t(a, 1)g \mapsto g$, we get

$$I(W^{a}, W^{a}_{\theta^{\psi}}, f_{2s-1}, \chi) = \chi(a)^{-1} |a|^{-\frac{2s+1}{4}} \int_{Z^{2}N \setminus GL_{2}} W(g) W_{\theta^{\psi}}(\mathfrak{s}(g)) f_{2s-1}(\mathfrak{s}(g)) \chi(\det(g)) dg.$$

The dual side needs to be treated more carefully. For $W^a \in \mathcal{W}(\pi, \psi_a)$, $W^a_{\theta\psi} \in \mathcal{W}(\theta^{\psi_a}, \psi_a^{-1})$ and $f_{2s-1} \in I(2s-1, \omega_{\pi\otimes\chi}^{-1})$, we break t(a, 1) into $t(a, a)t(1, a^{-1})$ and I_2 into $t(1, a)t(1, a^{-1})$. Then

$$\begin{split} &I(W^{a}, W^{a}_{\theta^{\psi}}, \hat{N}(s, \omega^{-1}_{\pi\otimes\chi}, \psi_{a})f_{2s-1}, \chi) \\ &= \omega_{\pi}(a)\mu_{\psi}(a)^{-1}(a, a^{-1})_{F}\int_{Z^{2}N\setminus GL_{2}}W(t(1, a^{-1})g)W_{\theta^{\psi}}(\mathfrak{s}(t(1, a^{-1}))\mathfrak{s}(g)) \\ &\times \hat{N}(s, \omega^{-1}_{\pi\otimes\chi}, \psi_{a})f_{2s-1}(\mathfrak{s}(t(1, a))\mathfrak{s}(t(1, a^{-1}))\mathfrak{s}(g))\chi(a)\chi(\det(t(1, a^{-1})g))dg. \end{split}$$

Changing an additive character implies a change to measures as the measure are chosen relative to the additive character. We define the fixed measure dx with respect to ψ as in §5.1 and we denote by $d_a x$ the measure adapted to ψ_a . Then $d_a x = |a|^{\frac{1}{2}} dx$. Let $M_a(2s - 1, \omega_{\pi \otimes \chi}^{-1}) f_{2s-1}$ denote the application of the intertwining operator to f_{2s-1} given by

$$M_a(2s-1,\omega_{\pi\otimes\chi}^{-1})f_{2s-1}(\mathfrak{s}(g)) = \int_F f_{2s-1}(\mathfrak{s}(w_2)\mathfrak{s}(n(x))\mathfrak{s}(g))d_ax.$$

Using $\gamma(2s-1, \omega_{\pi}^2 \chi^4, \psi_a) = |a|^{2s-\frac{3}{2}} \omega_{\pi}^2(a) \chi^4(a) \gamma(2s-1, \omega_{\pi}^2 \chi^4, \psi)$ (cf. [39, (3.2.3)]), $\hat{N}(2s-1, \omega_{\pi\otimes\chi}^{-1}, \psi_a)$ can be written in terms of the one for ψ as

$$\begin{split} \hat{N}(2s-1,\omega_{\pi\otimes\chi}^{-1},\psi_a)f_{2s-1}(\mathfrak{s}(t(1,a))) \\ &= |a|^{2s-\frac{3}{2}}\omega_{\pi}^2(a)\chi^4(a)\gamma(2s-1,\omega_{\pi}^2\chi^4,\psi)M_a(2s-1,\omega_{\pi\otimes\chi}^{-1})f_{2s-1}(\mathfrak{s}(t(1,a))) \\ &= |a|^{2s-\frac{3}{2}}\omega_{\pi}^2(a)\chi^4(a)\gamma(2s-1,\omega_{\pi}^2\chi^4,\psi)|a|^{\frac{2s-3}{4}+\frac{1}{2}}M(2s-1,\omega_{\pi\otimes\chi}^{-1})f_{2s-1}. \end{split}$$

By making the change of variables $t(1, a^{-1})g \mapsto g$, we achieve that

$$\begin{split} I(W^{a}, W^{a}_{\theta\psi}, \dot{N}(s, \omega_{\pi\otimes\chi}^{-1}, \psi_{a})f_{2s-1}, \chi) \\ &= \mu_{\psi_{a}}(a)^{-1}|a|^{\frac{5}{2}s-\frac{7}{4}}\omega_{\pi}^{3}(a)\chi^{5}(a)\int_{Z^{2}N\setminus GL_{2}}W(g)W_{\theta\psi}(\mathfrak{s}(g))\hat{N}(2s-1, \omega_{\pi\otimes\chi}^{-1}, \psi)f_{2s-1}(\mathfrak{s}(g))\chi(\det(g))dg \end{split}$$

as desired.

Henceforth it suffices to prove the stability of γ -factor with ψ of conductor \mathcal{O} . In contrast to γ -factors in the Langlands-Shahidi method [9, Theorem 5.1-(iv)], the auxiliary factor $\mu_{\psi_a}(a)^{-1}$ appears due to the presence of the exceptional representation θ^{ψ} .

5.4. **Proof of stability.** Before we launch our computation, let us fix our measures. Because $N(F), \overline{N}(F) \simeq F$ and $A(F) \simeq F^{\times}$, we identify the measure on groups N(F) and $\overline{N}(F)$ with an additive measure dx on F and the measure on A(F) with the multiplicative measure $d^{\times}a$ on F^{\times} . We normalize our additive measure so that $vol(\mathcal{O}, dx) = 1$. We take the multiplicative measure $d^{\times}a = da/|a|$. Under this normalization, $d^{\times}a$ assigns \mathcal{O}^{\times} to (q-1)/q and \mathfrak{p}^m to q^{-m} .

Our immediate goal is to construct explicit Howe vectors $W_m \in \mathcal{W}(\pi, \psi), W_{\theta^{\psi},m} \in \mathcal{W}(\theta^{\psi}, \psi^{-1})$, a section $f_{2s-1}^i \in I(2s-1, \omega_{\pi\otimes\chi}^{-1})$, and a highly ramified character χ such that $I(W_m, W_{\theta^{\psi},m}, f_{2s-1}^i, \chi)$ is a constant.

Proposition 5.7. Let π be an irreducible admissible representation of GL_2 . For every $i \ge m$ and $m \ge \max\{L, \mathfrak{f}(\chi)\}$, we have

(5.3)
$$I(W_m, W_{\theta^{\psi}, m}, f^i_{2s-1}, \chi) = q^{-3i-m}$$

Proof. We compute $I(W_m, W_{\theta^{\psi}, m}, f_{2s-1}^i, \chi)$ on the open dense subset $NZ^2 \setminus NT\overline{N}$ of $NZ^2 \setminus GL_2$. The decomposition of the Haar measure $dg = \delta_B(a)^{-1} dn d^{\times} z d^{\times} a d\overline{n}$ leads us to get

$$\begin{split} I(W_m, W_{\theta^{\psi}, m}, f_{2s-1}^i, \chi) &= \int_{Z^2 \setminus Z} \int_{F^{\times}} \int_F W_m(t(z, z)t(a, 1)\overline{n}(x)) W_{\theta^{\psi}, m}(\mathfrak{s}[t(z, z)t(a, 1)\overline{n}(x)]) \\ &\times f_{2s-1}^i(\mathfrak{s}[t(z, z)t(a, 1)\overline{n}(x)]) \chi(z^2a) |a|^{-1} dx d^{\times} a d^{\times} z. \end{split}$$

Now we would like to write $\mathfrak{s}[t(z,z)t(a,1)\overline{n}(x)] = \mathfrak{s}(t(z,z))\mathfrak{s}(t(a,1))\mathfrak{s}(\overline{n}(x))$. This equality does not hold in general but the fact that both $W_{\theta^{\psi},m}$ and f_{2s-1}^i are genuine make it possible to do this maneuver. This leads us to exploit the property of f_{2s-1}^i and we obtain

$$I(W_m, W_{\theta^{\psi}, m}, f_{2s-1}^i, \chi) = \int_{F^{\times}} \int_{\mathfrak{p}^{3i}} W_m(t(a, 1)\overline{n}(x)) W_{\theta^{\psi}, m}(\mathfrak{s}(t(a, 1))\mathfrak{s}(\overline{n}(x)))\chi(a) |a|^{\frac{2s+1}{4}-1} dx d^{\times} a.$$

Since $i \ge m$, we have $\mathfrak{p}^{3i} \subseteq \mathfrak{p}^{3m}$. In the light of Lemma 5.1 and [45, Lemma 5.1] together, we obtain $W_m(t(a,1)\overline{n}(x)) = W_m(t(a,1))$ and $W_{\theta^{\psi},m}(\mathfrak{s}(t(a,1))\mathfrak{s}(\overline{n}(x))) = W_{\theta^{\psi},m}(\mathfrak{s}(t(a,1)))$. It follows

from Lemma 5.2 joined with [45, Lemma 5.2] that

$$\begin{split} I(W_m, W_{\theta^{\psi}, m}, f_{2s-1}^i, \chi) &= q^{-3i} \int_{1 + \mathfrak{p}^m} W_m(t(a, 1)) W_{\theta^{\psi}, m}(t(a, 1)) \chi(a) |a|^{\frac{2s+1}{4} - 1} dx d^{\times} a \\ &= q^{-3i} \int_{1 + \mathfrak{p}^m} \chi(a) d^{\times} a. \end{split}$$

Then the assumption $m \geq \max\{L, \mathfrak{f}(\chi)\}$ gives rise to $I(W_m, W_{\theta^{\psi}, m}, f_{2s-1}^i, \chi) = q^{-3i-m}$, which concludes the proof.

We express the difference of Rankin-Selberg integrals in terms of the Mellin transform of a product of certain Bessel functions. In the case of $GL_2(F) \times GL_2(F)$, (5.4) is basically the content of a crucial lemma of Soudry [36, Lemma 4.5].

Proposition 5.8 (The Mellin transform). Let π and σ be irreducible admissible representations of GL_2 having the same central character $\omega = \omega_{\pi} = \omega_{\sigma}$. We fix $W^1 \in \mathcal{W}(\pi, \psi)$ and $W^2 \in \mathcal{W}(\sigma, \psi)$ as above, and form Howe Whittaker functions W_m^1 and W_m^2 , respectively. For $m \ge 6L$ and $i \ge 6L$ $\max\{m+1, I(N_m, \omega_{\pi\otimes\chi})\}, we have$

$$\begin{split} &I(W_m^1, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\pi\otimes\chi}^{-1}) f_{2s-1}^i, \chi) - I(W_m^2, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\sigma\otimes\chi}^{-1}) f_{2s-1}^i, \chi) \\ &= q^{-3i-m+3L} \int_{F^{\times}} [W_{3L}^1(t(a, 1)w_2) - W_{3L}^2(t(a, 1)w_2)] j_{\theta^{\psi}}(a) \omega(a)^{-1} \mu_{\psi}(a^{-1})^{-1} \chi^{-1}(a) |a|^{-\frac{2s+1}{4}} d^{\times} a. \end{split}$$

In particular the integral can be assumed to range over the compact set \mathfrak{p}^{-9L} .

Proof. We take the difference and compute the dual side of the functional equation on the dense open subset $NZ^2 \setminus NTw_2N$. Then we have

$$\begin{split} &I(W_m^1, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\pi\otimes\chi}^{-1}) f_{2s-1}^i, \chi) - I(W_m^2, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\sigma\otimes\chi}^{-1}) f_{2s-1}^i, \chi) \\ &= \int_{Z^2 \backslash Z} \int_{F^{\times}} \int_F [W_m^1(t(z, z)t(1, a)w_2n(x)) - W_m^2(t(z, z)t(1, a)w_2n(x))] \\ &\quad \times W_{\theta^{\psi}, m}(\mathfrak{s}[t(z, z)t(1, a)w_2n(x)]) \widetilde{f}_{1-2s}^i(\mathfrak{s}[t(z, z)t(1, a)w_2n(x)]) \chi(z^2a) |a| dx d^{\times} a d^{\times} z. \end{split}$$

As we have seen before, we may write $\mathfrak{s}[t(z,z)t(1,a)w_2n(x)] = \mathfrak{s}(t(z,z))\mathfrak{s}(t(1,a))\mathfrak{s}(w_2)\mathfrak{s}(n(x)),$ namely, the "genuineness" of $W_{\theta^{\psi},m}$ and \tilde{f}_{1-2s}^i eliminates the discrepancy of those expressions. By [45, Proposition 5.3], we obtain

$$W_m^1(t(z,z)t(1,a)w_2n(x)) = W_m^2(t(z,z)t(1,a)w_2n(x))$$

for all $n(x) \in N - N_m$. Therefore

$$\begin{split} &I(W_m^1, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\pi\otimes\chi}^{-1}) f_{2s-1}^i, \chi) - I(W_m^2, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\sigma\otimes\chi}^{-1}) f_{2s-1}^i, \chi) \\ &= \int_{Z^2 \backslash Z} \int_{F^{\times}} \int_{\mathfrak{p}^{-m}} [W_m^1(t(z, z)t(1, a)w_2n(x)) - W_m^2(t(z, z)t(1, a)w_2n(x))] \\ &\times W_{\theta^{\psi}, m}(\mathfrak{s}(t(z, z))\mathfrak{s}(t(1, a))\mathfrak{s}(w_2)\mathfrak{s}(n(x))) \\ &\times \widetilde{f}_{1-2s}^i(\mathfrak{s}(t(z, z))\mathfrak{s}(t(1, a))\mathfrak{s}(w_2)\mathfrak{s}(n(x)))\chi(z^2a)|a|dxd^{\times}ad^{\times}z. \end{split}$$

We concentrate on the support of t(z, z). For the moment we take Lemma 5.9 below for granted. **Lemma 5.9.** If $i \ge m+1$ and $n(x) \in \mathfrak{p}^{-m}$, then

$$\widetilde{f}_{1-2s}^{i}(\mathfrak{s}(t(z,z))\mathfrak{s}(t(1,a))\mathfrak{s}(w_{2})\mathfrak{s}(n(x))) = \begin{cases} \omega_{\pi\otimes\chi}^{-1}(z)|a|^{\frac{2s-3}{4}}\widetilde{f}_{1-2s}^{i}(\mathfrak{s}(w_{2})\mathfrak{s}(n(x))) & \text{if } t(z,z)\in Z^{2}\\ 0 & \text{otherwise.} \end{cases}$$

Let us resume our argument. Lemma 5.9 aligned with Proposition 5.5 implies that

$$\begin{split} &I(W_m^1, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\pi\otimes\chi}^{-1}) f_{2s-1}^i, \chi) - I(W_m^2, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\sigma\otimes\chi}^{-1}) f_{2s-1}^i, \chi) \\ &= q^{-3i} \int_{F^{\times}} \int_{\mathfrak{p}^{-m}} [W_m^1(t(1, a) w_2 n(x)) - W_m^2(t(1, a) w_2 n(x))] W_{\theta^{\psi}, m}(\mathfrak{s}(t(1, a)) \mathfrak{s}(w_2) \mathfrak{s}(n(x)))) \\ & \quad \times \chi(a) |a|^{\frac{2s-3}{4}+1} dx d^{\times} a. \end{split}$$

We see from Lemma 5.1 and [45, Lemma 5.1] that $W_m^i(t(1,a)w_2n(x)) = \psi(x)W_m^i(t(1,a)w_2)$ and $W_{\theta^{\psi},m}(\mathfrak{s}(t(1,a))\mathfrak{s}(w_2)\mathfrak{s}(n(x))) = \psi(x)^{-1}W_{\theta^{\psi},m}(\mathfrak{s}(t(1,a))\mathfrak{s}(w_2))$ for i = 1, 2 and all $x \in \mathfrak{p}^{-m}$. The integral becomes

$$\begin{split} &I(W_m^1, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\pi \otimes \chi}^{-1}) f_{2s-1}^i, \chi) - I(W_m^2, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\sigma \otimes \chi}^{-1}) f_{2s-1}^i, \chi) \\ &= q^{-3i+m} \int_{F^{\times}} [W_m^1(t(1, a)w_2) - W_m^2(t(1, a)w_2)] W_{\theta^{\psi}, m}(\mathfrak{s}(t(1, a))\mathfrak{s}(w_2))\chi(a) |a|^{\frac{2s+1}{4}} d^{\times}a. \end{split}$$

We concern with removing the dependence of m. To this end, we know from [45, Lemma 5.1, Proposition 5.3] that

(5.5)

$$W_m^1(t(1,a)w_2) - W_m^2(t(1,a)w_2) = \frac{1}{\operatorname{vol}(N_m)} \int_{N_{3L}} [W_{3L}^1(t(1,a)w_2n) - W_{3L}^2(t(1,a)w_2n)]\psi^{-1}(n)dn = \frac{\operatorname{vol}(N_{3L})}{\operatorname{vol}(N_m)} [W_{3L}^1(t(1,a)w_2) - W_{3L}^2(t(1,a)w_2)].$$

We take m to be $m \ge 6L$. Appealing to Proposition 5.3-(2), we relate the partial Bessel function with the full Bessel function by

(5.6)
$$\operatorname{vol}(N_m)W_{\theta^{\psi},m}(\mathfrak{s}(t(a,1))\mathfrak{s}(w_2)) = j_{\theta^{\psi}}(a) \quad \text{for all} \quad |a| \le q^{2m}$$

Applying the change of variables $a \mapsto a^{-1}$ and then putting (5.5) and (5.6) together, we have

$$\begin{split} &I(W_m^1, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\pi\otimes\chi}^{-1}) f_{2s-1}^i, \chi) - I(W_m^2, W_{\theta^{\psi}, m}, M(2s-1, \omega_{\sigma\otimes\chi}^{-1}) f_{2s-1}^i, \chi) \\ &= q^{-3i-m+3L} \int_{F^{\times}} [W_{3L}^1(t(a, 1)w_2) - W_{3L}^2(t(a, 1)w_2)] j_{\theta^{\psi}}(a) \omega(a)^{-1} \mu_{\psi}(a^{-1})^{-1} \chi^{-1}(a) |a|^{-\frac{2s+1}{4}} d^{\times} a. \end{split}$$

Here a is in fact taken over the compact set \mathfrak{p}^{-9L} utilizing [45, Lemma 5.2].

Proof of Lemma 5.9. We compute the intertwining operator \tilde{f}_{1-2s}^i by brutal force. Since

$$\widetilde{f}_{1-2s}^i(\mathfrak{s}(t(z,z))\mathfrak{s}(t(1,a))\mathfrak{s}(w_2)\mathfrak{s}(n(x))) = \int_F f_{2s-1}^i(\mathfrak{s}(w_2)\mathfrak{s}(n(y))\mathfrak{s}(t(z,z))\mathfrak{s}(t(1,a))\mathfrak{s}(w_2)\mathfrak{s}(n(x))dy,$$

the support of $\widetilde{f}_{1-2s}^i(\mathfrak{s}(t(z,z))\mathfrak{s}(t(1,a))\mathfrak{s}(w_2)\mathfrak{s}(n(x)))$ boils down to the support of $f_{2s-1}^i(\mathfrak{s}(t(z,z))\mathfrak{s}(t(a,1))\mathfrak{s}(w_2)\mathfrak{s}(n(y))\mathfrak{s}(w_2)\mathfrak{s}(n(x))).$

We assume
$$t(z,z)t(a,1)\overline{n}(y)n(x) = \begin{pmatrix} \alpha\beta & v \\ & \alpha \end{pmatrix}\overline{n}(\overline{y})$$
 for $\alpha \in (F^{\times})^2$, $x \in \mathfrak{p}^{-m}$ and $\overline{y} \in \mathfrak{p}^{3i}$. Then
 $\begin{pmatrix} za & zax \\ zy & zxy + z \end{pmatrix} = \begin{pmatrix} \alpha\beta + \delta\overline{y} & v \\ \alpha\overline{y} & \alpha \end{pmatrix}$.

Focusing on the bottom entries $zy = \alpha \overline{y}$ and $zxy + z = \alpha$, we conclude $z = \alpha(1 - x\overline{y})$. According to the surjective map [5, Lemma 3.3], $1 - x\overline{y}$, being an element of $1 + \mathfrak{p}^{2m+3}$, is in fact a square element. Therefore z belongs to $(F^{\times})^2$.

We are now in the position to present our main result.

Theorem 5.10. Let π and σ be irreducible admissible representations of GL_2 having the same central character. Then for every sufficiently highly ramified character χ of F^{\times} we have

$$\gamma(s, \pi \otimes \chi, \operatorname{Sym}^2, \psi) = \gamma(s, \sigma \otimes \chi, \operatorname{Sym}^2, \psi)$$

Proof. We may assume that $m \ge \max\{\mathfrak{f}(\chi), 6L\}$. We enlarge i so that $i \ge \max\{m+1, I(N_m, \omega_{\pi\otimes\chi})\}$. Substituting (5.3) and (5.4) in the functional equation (2.4), then simply implies:

(5.7)
$$\Gamma(s, \pi \otimes \chi, \operatorname{Sym}^2, \psi) - \Gamma(s, \sigma \otimes \chi, \operatorname{Sym}^2, \psi) = q^{3L} \int_{\mathfrak{p}^{-9L}} [W_{3L}^1(t(a, 1)w_2) - W_{3L}^2(t(a, 1)w_2)] j_{\theta^{\psi}}(a)\omega(a)^{-1} \mu_{\psi}(a^{-1})^{-1} \chi^{-1}(a) |a|^{-\frac{2s+1}{4}} d^{\times} a.$$

To complete the proof, we must now observe from Proposition 5.3-(3) that the integrand

$$a \mapsto [W_{3L}^1(t(a,1)w_2) - W_{3L}^2(t(a,1)w_2)]j_{\theta^{\psi}}(a)\omega(a)^{-1}\mu_{\psi}(a^{-1})^{-1}|a|^{-\frac{2s+4}{4}}$$

is right invariant under $1 + \mathfrak{p}^{6L}$ for $a \in \mathfrak{p}^{-9L}$. Taking χ a sufficiently highly ramified character, the integral in (5.7) vanishes. Since Tate's gamma factor $\gamma(2s-1, \omega_{\pi\otimes\chi}^2, \psi)$ stabilizes under highly ramified twist, we conclude from Proposition 2.7 that

$$\gamma(s, \pi \otimes \chi, \operatorname{Sym}^2, \psi) = \gamma(s, \sigma \otimes \chi, \operatorname{Sym}^2, \psi)$$

as expected.

We illustrate an important consequence of this that we highlight below. We recover the result of Gelbart and Jacquet [11, (6.4)].

Theorem 5.11. Let π and σ be irreducible admissible representations of GL_2 having the same central character. Then for every sufficiently highly ramified character χ of F^{\times} we have

$$L(s, \pi \otimes \chi, \operatorname{Sym}^2) = L(s, \sigma \otimes \chi, \operatorname{Sym}^2) = 1.$$

Moreover $\varepsilon(s, \pi \otimes \chi, \operatorname{Sym}^2, \psi) = \varepsilon(s, \sigma \otimes \chi, \operatorname{Sym}^2, \psi).$

Proof. In virtue of Yamana [44, Lemma 3.15] concerning the pole of the symmetric square L-function, this can be proved exactly as in Jacquet and Shalika [16, Proposition 5.1].

Acknowledgments. The author wishes to thank J. Cogdell for pointing out the unfinished project in A. Kable [22] and for encouraging us to write this article. The second part "stability of γ -factors" grows out of several discussions with M. Krishnamurthy. We wish to express our gratitude to M. Krishnamurthy for keen insights and for fruitful mathematical communications. We also thank every members of the number theory and the representation theory group at the University of Iowa for vibrating environments when this paper was written. Finally, we would like to convey our sincere appreciation to the anonymous referee for invaluable suggestions which improve exposition and organization of our paper.

References

- William D. Banks, Jason Levy, and Mark R. Sepanski, *Block-compatible metaplectic cocycles*, J. Reine Angew. Math. 507 (1999), 131–163.
- [2] I. N. Bernstein and A. V. Zelevinsky, Induced representations of reductive p-adic groups. I, Ann. Sci. École Norm. Sup. (4) 10 (1977), no. 4, 441–472.
- [3] Daniel Bump and David Ginzburg, Symmetric square L-functions on GL(r), Ann. of Math. (2) 136 (1992), no. 1, 137–205.
- [4] Jingsong Chai, Some results on archimedean Rankin-Selberg integrals, Pacific J. Math. 273 (2015), no. 2, 277–305.
- [5] Jingsong Chai and Qing Zhang, A strong multiplicity one theorem for SL₂, Pacific J. Math. 285 (2016), no. 2, 345–374.

- Shih-Yu Chen, Gamma factors for the Asai cube representation, to appear in Mathematische Zeitschrift. https://arxiv.org/abs/arXiv:1904.07844.
- [7] J. W. Cogdell and I. I. Piatetski-Shapiro, *Derivatives and L-functions for GL_n*, Representation theory, number theory, and invariant theory, Progr. Math., vol. 323, Birkhäuser/Springer, Cham, 2017, pp. 115–173.
- [8] J. W. Cogdell, F. Shahidi, and T.-L. Tsai, Local Langlands correspondence for GL_n and the exterior and symmetric square ε -factors, Duke Math. J. **166** (2017), no. 11, 2053–2132.
- [9] Radhika Ganapathy and Luis Lomelí, On twisted exterior and symmetric square γ-factors, Ann. Inst. Fourier (Grenoble) 65 (2015), no. 3, 1105–1132 (English, with English and French summaries).
- [10] Fan Gao, Freydoon Shahidi, and Dani Szpruch, Local coefficients and gamma factors for principal series of covering groups, to appear in Mem. Amer. Math. Soc. https://arxiv.org/abs/arXiv:1902.02686v3.
- [11] Stephen Gelbart and Hervé Jacquet, A relation between automorphic representations of GL(2) and GL(3), Ann. Sci. École Norm. Sup. (4) 11 (1978), no. 4.
- [12] Stephen Gelbart and I. I. Piatetski-Shapiro, Distinguished representations and modular forms of half-integral weight, Invent. Math. 59 (1980), no. 2, 145–188.
- [13] Michael Harris, Stephen S. Kudla, and William J. Sweet, Theta dichotomy for unitary groups, J. Amer. Math. Soc. 9 (1996), no. 4, 941–1004.
- [14] Guy Henniart, Correspondance de Langlands et fonctions L des carrés extérieur et symétrique, Int. Math. Res. Not. IMRN 4 (2010), 633–673 (French).
- [15] Roger Howe, Classification of Irreducible Representations of $GL_2(F)$, preprint, I.H.E.S., Bures-sur-Yvette, France (1978).
- [16] Hervé Jacquet and Joseph Shalika, A lemma on highly ramified ϵ -factors, Math. Ann. 271 (1985), no. 3, 319–332.
- [17] H. Jacquet, I. I. Piatetskii-Shapiro, and J. A. Shalika, *Rankin-Selberg convolutions*, Amer. J. Math. 105 (1983), no. 2, 367–464.
- [18] Yeongseong Jo, Derivatives and exceptional poles of the local exterior square L-function for GL_m, Math. Z. 294 (2020), no. 3-4, 1687–1725.
- [19] _____, Factorization of the local exterior square L-function of GL_m , Manuscripta Math. 162 (2020), no. 3-4, 493–536.
- [20] _____, Rankin-Selberg L-functions via good sections, Forum Math. 32 (2020), no. 4, 1039–1074.
- [21] Anthony C. Kable, The main involutions of the metaplectic group, Proc. Amer. Math. Soc. 127 (1999), no. 4, 955–962.
- [22] _____, The tensor product of exceptional representations on the general linear group, Ann. Sci. Ecole Norm. Sup. (4) **34** (2001), no. 5, 741–769 (English, with English and French summaries).
- [23] Eyal Kaplan, On the gcd of local Rankin-Selberg integrals for even orthogonal groups, Compos. Math. 149 (2013), no. 4, 587–636.
- [24] _____, The characterization of theta-distinguished representations of GL(n), Israel J. Math. **222** (2017), no. 2, 551–598.
- [25] _____, The double cover of odd general spin groups, small representations, and applications, J. Inst. Math. Jussieu 16 (2017), no. 3, 609–671.
- [26] D. A. Kazhdan and S. J. Patterson, Metaplectic forms, Inst. Hautes Études Sci. Publ. Math. 59 (1984), 35–142.
- [27] Stanisław Łojasiewicz, Introduction to complex analytic geometry, Birkhäuser Verlag, Basel, 1991. Translated from the Polish by Maciej Klimek.
- [28] Nadir Matringe, Conjectures about distinction and local Asai L-functions, Int. Math. Res. Not. IMRN 9 (2009), 1699–1741.
- [29] _____, On the local Bump-Friedberg L-function, J. Reine Angew. Math. 709 (2015), 119–170.
- [30] S. J. Patterson and I. I. Piatetski-Shapiro, The symmetric-square L-function attached to a cuspidal automorphic representation of GL₃, Math. Ann. 283 (1989), no. 4, 551–572.
- [31] I. I. Piatetski-Shapiro, L-functions for GSp₄, Pacific J. Math. Special Issue (1997), 259–275. Olga Taussky-Todd: in memoriam.
- [32] I. Piatetski-Shapiro and Stephen Rallis, Rankin triple L functions, Compositio Math. 64 (1987), no. 1, 31–115.
- [33] Ralf Schmidt and Long Tran, Zeta integrals for GSp(4) via Bessel models, Pacific J. Math. 296 (2018), no. 2, 437–480.
- [34] Freydoon Shahidi, Twisted endoscopy and reducibility of induced representations for p-adic groups, Duke Math.
 J. 66 (1992), no. 1, 1–41.
- [35] Goro Shimura, On the holomorphy of certain Dirichlet series, Proc. London Math. Soc. (3) 31 (1975), no. 1, 79–98.
- [36] David Soudry, The L and γ factors for generic representations of $GSp(4, k) \times GL(2, k)$ over a local non-Archimedean field k, Duke Math. J. 51 (1984), no. 2, 355–394.

- [37] Shuichiro Takeda, The twisted symmetric square L-function of GL(r), Duke Math. J. 163 (2014), no. 1, 175–266.
- [38] _____, On a certain metaplectic Eisenstein series and the twisted symmetric square L-function, Math. Z. 281 (2015), no. 1-2, 103–157.
- [39] J. Tate, Number theoretic background, Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 3–26.
- [40] J.-L. Waldspurger, La formule de Plancherel pour les groupes p-adiques (d'après Harish-Chandra), J. Inst. Math. Jussieu 2 (2003), no. 2, 235–333 (French, with French summary).
- [41] Chian-Jen Wang, On the existence of cuspidal distinguished representations of metaplectic groups, ProQuest LLC, Ann Arbor, MI, 2003. Thesis (Ph.D.)-The Ohio State University.
- [42] André Weil, *Basic number theory*, Classics in Mathematics, Springer-Verlag, Berlin, 1995. Reprint of the second (1973) edition.
- [43] Shunsuke Yamana, L-functions and theta correspondence for classical groups, Invent. Math. 196 (2014), no. 3, 651–732.
- [44] _____, Local symmetric square L-factors of representations of general linear groups, Pacific J. Math. 286 (2017), no. 1, 215–256.
- [45] Qing Zhang, A local converse theorem for U(1,1), Int. J. Number Theory 13 (2017), no. 8, 1931–1981.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY, IA 52242, USA *E-mail address*: jo.59@buckeyemail.osu.edu