

VARENTROPY OF PAST LIFETIMESFRANCESCO BUONO,* *Universit degli Studi di Napoli Federico II*MARIA LONGOBARDI,** *Universit degli Studi di Napoli Federico II***Abstract**

In Reliability Theory, uncertainty is measured by the Shannon entropy. Recently, in order to analyze the variability of such measure, varentropy has been introduced and studied. In this paper we define a new concept of varentropy for past lifetime distributions as the variability of uncertainty in past lifetime. We analyze the relation of this varentropy with the corresponding past entropy and other concepts of interest in reliability and we study the effect of linear and monotonic transformations on it. Finally, we evaluate past varentropy for random variables following a proportional reversed hazard rate model.

Keywords: Past and inactivity lifetimes; Reversed hazard rate function; Varentropy; Proportional reversed hazards model

2010 Mathematics Subject Classification: Primary 62N05

Secondary 60E15

1. Introduction

Let X be an absolutely continuous random variable with support $D \subseteq (0, +\infty)$, describing the lifetime of a living being or of an item. Let F be its cumulative distribution function (cdf), \bar{F} its survival function (sf) and f its probability density function (pdf).

* Postal address: Dipartimento di Matematica e Applicazioni "Renato Caccioppoli", Universit degli Studi di Napoli Federico II, Via Cintia, 80126, Naples, Italy

** Postal address: Dipartimento di Biologia, Universit degli Studi di Napoli Federico II, Via Cintia, 80126, Naples, Italy

In reliability, in information theory, in finance, in theoretical neurobiology and in several other fields, a well-known measure of uncertainty for X is the differential entropy or Shannon information measure, defined as

$$H(X) = \mathbb{E}[-\log f(X)] = - \int_0^{+\infty} f(x) \log f(x) dx,$$

where \log is the natural logarithm (see Shannon [27]il).

Ebrahimi [16] and Muliere et al. [26] studied the residual entropy as measure of information for the residual lifetime $X_t = (X - t | X \geq t)$. The cdf of the residual lifetime is expressed as

$$f_{X_t}(x) = \frac{f(x+t)}{\bar{F}(t)}, \quad x > 0,$$

whereas the residual entropy can be written as

$$H(X_t) = - \int_t^{+\infty} \frac{f(x)}{\bar{F}(t)} \log \frac{f(x)}{\bar{F}(t)} dx.$$

For all $t \in D$, the past lifetime of X is ${}_tX = (X | X \leq t)$, with cdf and pdf expressed as

$$F_{{}_tX}(x) = \frac{F(x)}{F(t)}, \quad f_{{}_tX}(x) = \frac{f(x)}{F(t)} \quad x \in (0, t)$$

and the inactivity time of X is $X_{(t)} = (t - X | X \leq t)$, where its sf and pdf are

$$\bar{F}_{(t)}(x) = \frac{F(t-x)}{F(t)}, \quad f_{(t)}(x) = \frac{f(t-x)}{F(t)} \quad x \in (0, t)$$

(see Li and Zuo [23] for details).

Di Crescenzo and Longobardi [9] studied the past entropy to measure the uncertainty about the past lifetime. The past entropy is defined as

$$H({}_tX) = \mathbb{E}[-\log f_{{}_tX}({}_tX)] = - \int_0^t \frac{f(x)}{F(t)} \log \frac{f(x)}{F(t)} dx. \quad (1.1)$$

The mean inactivity time is a function of a great importance in the context of reliability theory. It is the mean value of $X_{(t)}$ and it is expressed as

$$m^*(t) = \mathbb{E}(X_{(t)}) = \frac{1}{F(t)} \int_0^t F(t-x) dx = \frac{1}{F(t)} \int_0^t F(x) dx, \quad (1.2)$$

(see Izadkhan [20] for further details).

Notice that the past entropy is also the entropy of the inactivity time, that is, $H({}_tX) = H(X_{(t)})$.

We define, for $t \in D$, the reversed hazard rate function of X as

$$q(t) = \lim_{\Delta t \rightarrow 0^+} \frac{1}{\Delta t} \mathbb{P}(X \geq t - \Delta t | X \leq t) = \frac{f(t)}{F(t)}. \quad (1.3)$$

This function is the instantaneous failure rate occurring immediately before the time point t , i.e., the failure occurs just before the time point t , given that the unit has not survived longer than t (see Block and Savits [2] and Finkelstein [17] for more details about the reversed hazard rate function). Recall also that the cumulative reversed hazard rate function is defined as

$$\Lambda^*(t) = \int_t^{+\infty} q(x) dx = -\log F(t). \quad (1.4)$$

The past entropy can also be expressed in different ways (see [9]):

$$H({}_tX) = -\Lambda^*(t) - \frac{1}{F(t)} \int_0^t f(x) \log f(x) dx \quad (1.5)$$

$$= 1 - \frac{1}{F(t)} \int_0^t f(x) \log q(x) dx. \quad (1.6)$$

By differentiating both sides of (1.5), we get the following expression for the derivative of the past entropy:

$$H'({}_tX) = q(t)[1 - H({}_tX) - \log q(t)]. \quad (1.7)$$

In recent literature several new definitions of entropy of a continuous distribution have been proposed: weighted version and its residual and past ones (see [10]) and many different version of cumulative entropies and their applications (see for instance [6], [7], [8], [11], [12], [13], [14] and [24]).

We recall, also, that the distribution function, the reversed hazard rate function and the mean inactivity lifetimes are linked by the following relations:

$$F(t) = \exp \left[- \int_t^{+\infty} q(x) dx \right] \quad (1.8)$$

$$q(t) = \frac{1 - (m^*(t))'}{m^*(t)}. \quad (1.9)$$

In Section 2, the definitions of varentropy and residual varentropy are recalled and a new varentropy of past lifetime distributions is defined. Moreover, some examples of past varentropy are exhibited and some relations with other important concepts of reliability theory are given. The behaviour of such a measure under certain transformations is also described.

In Section 3, we give some bounds for past varentropy. Finally, we consider the proportional reversed hazard rate model and evaluate the past varentropy of the family of random variables following this model.

2. Past varentropy

The differential entropy can be seen as expectation of the information contained in X . In order to measure the concentration of this information content around the entropy, recently a new definition called varentropy was given (see also Bobkov and Madiman [3]). The varentropy is defined as

$$V(X) = \text{Var}(-\log f(X)) = \text{Var}(\log f(X)) = \mathbb{E}[(\log f(X))^2] - [H(X)]^2 \quad (2.1)$$

$$= \int_0^{+\infty} f(x)[\log f(x)]^2 dx - \left[\int_0^{+\infty} f(x) \log f(x) dx \right]^2. \quad (2.2)$$

In the literature there are several papers about the study of the varentropy and its properties, we may refer to Arikan [1] and Madiman and Wang [25]. In the discrete case, we can write the entropy and the varentropy of a random variable X with support $\{x_i, i \in I\}$ and discrete density function p_X , as

$$H(X) = - \sum_{i \in I} p_X(x_i) \log p_X(x_i)$$

and

$$V(X) = \sum_{i \in I} p_X(x_i) [\log p_X(x_i)]^2 - [H(X)]^2,$$

respectively.

Di Crescenzo and Paolillo [15] extended the definition of the varentropy (2.1) to the residual lifetimes introducing the residual varentropy that is

$$\begin{aligned} V(X_t) &= \text{Var}(-\log f_{X_t}(X_t)) = \text{Var}(\log f_{X_t}(X_t)) = \mathbb{E}[(\log f_{X_t}(X_t))^2] - [H(X_t)]^2 \\ &= \int_t^{+\infty} \frac{f(x)}{\overline{F}(t)} \left[\log \frac{f(x)}{\overline{F}(t)} \right]^2 dx - \left[\int_t^{+\infty} \frac{f(x)}{\overline{F}(t)} \log \frac{f(x)}{\overline{F}(t)} dx \right]^2 \\ &= \frac{1}{\overline{F}(t)} \int_t^{+\infty} f(x)(\log f(x))^2 dx - (\Lambda(t) + H(X_t))^2, \end{aligned}$$

where, in analogy with (1.4), $\Lambda(t) = -\log \overline{F}(t)$.

Recently, a great importance is given to the study of past lifetime distributions, because, in many realistic situation, it is reasonable that uncertainty is related to the

past. Let us suppose to have a system observed at a certain time t and then it is found failed. So the uncertainty is referred to the past and the concentration around it can be investigated. For this reason, we introduce and study the past varentropy, i.e., the varentropy of the past lifetime. We define, for all $t \in D$, the past varentropy as

$$\begin{aligned} V({}_tX) &= \text{Var}(-\log f_{tX}({}_tX)) = \text{Var}(\log f_{tX}({}_tX)) = \mathbb{E}[(\log f_{tX}({}_tX))^2] - [H({}_tX)]^2 \\ &= \int_0^t \frac{f(x)}{F(t)} \left[\log \frac{f(x)}{F(t)} \right]^2 dx - \left[\int_0^t \frac{f(x)}{F(t)} \log \frac{f(x)}{F(t)} dx \right]^2 \end{aligned} \quad (2.3)$$

$$= \frac{1}{F(t)} \int_0^t f(x)(\log f(x))^2 dx - (\Lambda^*(t) + H({}_tX))^2, \quad (2.4)$$

where $\Lambda^*(t)$ and $H({}_tX)$ are given in (1.4) and (1.1), respectively. We note that the past varentropy is also the varentropy of the inactivity time, that is, $V({}_tX) = V(X_{(t)})$.

In the following example we evaluate the past entropy and the past varentropy for some distributions of interest in reliability theory.

Example 2.1. • Let X be a random variable with uniform distribution over $(0, b)$, i.e., $X \sim U(0, b)$, $b > 0$. Hence, for $t \in (0, b)$ we have

$$H({}_tX) = \log t$$

$$V({}_tX) = 0.$$

- Let X be a random variable with exponential distribution, i.e., $X \sim \text{Exp}(\lambda)$, $\lambda > 0$. Then, for $t > 0$ we have

$$H({}_tX) = 1 + \log \left(\frac{1 - e^{-\lambda t}}{\lambda} \right) - \frac{\lambda t e^{-\lambda t}}{1 - e^{-\lambda t}}$$

$$\begin{aligned} V({}_tX) &= 1 - 2\lambda^2 \log \lambda + 2 \log \lambda \\ &\quad + \frac{e^{-\lambda t}}{1 - e^{-\lambda t}} \left[-\lambda^2 t^2 + 2\lambda^3 t \log \lambda - 2\lambda t \log \lambda - \frac{\lambda^2 t^2 e^{-\lambda t}}{1 - e^{-\lambda t}} \right]. \end{aligned}$$

- Let X be a random variable such that $f(x) = 2x$ and $F(x) = x^2$, $x \in (0, 1)$. Hence, for $t \in (0, 1)$ we have

$$\begin{aligned} H({}_tX) &= \frac{1}{2} + \log \frac{t}{2} \\ V({}_tX) &= \frac{1}{4} + 4 \log 2 \log t + \log^2 t^2 - 2 \log 2t \log t^2. \end{aligned}$$

In the next proposition, we obtain an expression for the derivative of the past varentropy.

Proposition 2.1. *For all $t \in D$, the derivative of the past varentropy is*

$$V'({}_tX) = -q(t) [V({}_tX) - (H({}_tX) + \log q(t))^2].$$

Proof. By differentiating both sides of (2.4), we get

$$\begin{aligned} V'({}_tX) &= \frac{q(t)}{F(t)} \int_0^t f(x)(\log f(x))^2 dx + q(t)(\log f(t))^2 \\ &= -2(\Lambda^*(t) + H({}_tX))(-q(t) + H'({}_tX)), \end{aligned} \quad (2.5)$$

where $q(t)$ is defined in (1.3). Hence, recalling (1.7) and (2.4), from (2.5) we get

$$\begin{aligned} V'({}_tX) &= -q(t) [V({}_tX) + (\Lambda^*(t) + H({}_tX))^2 - (\log f(t))^2 \\ &\quad - 2(\Lambda^*(t) + H({}_tX))(H({}_tX) + \log q(t))], \end{aligned}$$

and, after some calculations, we get the thesis. \square

It is useful to study the monotonicity of this function and to establish if, under some conditions, the past varentropy can be constant.

Theorem 2.1. (i) *Let the past varentropy $V({}_tX)$ be constant, i.e., $V({}_tX) = v \geq 0$, for all $t \in D$. Then*

$$|H({}_tX) + \log q(t)| = \sqrt{v}, \quad \forall t \in D.$$

(ii) *Let $c \in \mathbb{R}$. If*

$$H({}_tX) + \log q(t) = c \quad \forall t \in D \quad (2.6)$$

then

$$V({}_tX) = c^2 + \frac{V(X) - c^2}{F(t)} \quad \forall t \in D.$$

Proof. (i) From Proposition 2.1, we get that if the past varentropy is constant, then

$$(H({}_tX) + \log q(t))^2 = v \geq 0,$$

and so

$$|H({}_tX) + \log q(t)| = \sqrt{v}, \quad \forall t \in D.$$

(ii) From Proposition 2.1 and from the assumption (2.6), we get

$$V'({}_tX) = -q(t)(V({}_tX) - c^2)$$

with the boundary condition

$$\lim_{t \rightarrow \sup D} V({}_tX) = V(X).$$

Solving the above linear ordinary differential equation we get the thesis. \square

Remark 2.1. We refer to Corollary 2.2 of Kunda et al. [22] for distributions which satisfy the relation expressed in (2.6).

Recall that the generalized reversed hazard rate is defined, for $\alpha \in \mathbb{R}$, as

$$q_\alpha(t) = \frac{f(t)}{[F(t)]^{1-\alpha}}, \quad t \in D \quad (2.7)$$

(see Buono et al. [4]). Notice that, by choosing $\alpha = 0$ in (2.7), we get $q_0(t) = q(t)$, i.e., q_0 is the usual reversed hazard rate function. In the following theorem, we study a necessary and sufficient condition in terms of past entropy, under which the generalized reversed hazard rate function is constant.

Theorem 2.2. *Let X be an absolutely continuous random variable with support D and let $c \in \mathbb{R}$. The generalized reversed hazard rate function of X with parameter $1 - c$, that is, $q_{1-c}(t)$, is constant, such that*

$$q_{1-c}(t) = \frac{f(t)}{[F(t)]^c} = e^{c-H(X)}, \quad \forall t \in D$$

if, and only if, (2.6) holds.

Proof. Let us suppose that $q_{1-c}(t) = e^{c-H(X)}, \forall t \in D$. From (1.3) and (1.5) we get

$$\begin{aligned} H({}_tX) + \log q(t) &= \log f(t) - \frac{1}{F(t)} \int_0^t f(x) \log f(x) dx \\ &= \log f(t) + \frac{1}{F(t)} \left[H(X) + \int_t^{+\infty} f(x) \log f(x) dx \right]. \end{aligned}$$

From the hypothesis, we get

$$\int_t^{+\infty} f(x) \log f(x) dx = -H(X)\overline{F}(t) - cF(t) \log F(t),$$

and so

$$H({}_tX) + \log q(t) = H(X) + \log \frac{f(t)}{[F(t)]^c} = c,$$

i.e., the thesis.

Conversely, from (1.7) we obtain

$$c = H({}_tX) + \log q(t) = 1 - \frac{H'({}_tX)}{q(t)},$$

and so

$$H'({}_tX) = (1 - c)q(t).$$

By integrating both sides from t and $+\infty$ and recalling (1.8), we get

$$H(X) - H({}_tX) = (1 - c)\Lambda^*(t).$$

Hence,

$$\begin{aligned} c = H({}_tX) + \log q(t) &= H(X) + \log F(t) - \log(F(t))^c + \log q(t) \\ &= H(X) + \log q_{1-c}(t), \end{aligned}$$

and we get the thesis. \square

In the following proposition, we study the past varentropy under linear transformations. We recall that if

$$Y = aX + b, \quad a > 0, \quad b \geq 0,$$

then the past entropies of X and Y are connected by (see [9])

$$H({}_tY) = H\left(\frac{t-b}{a}X\right) + \log a \quad \forall t. \quad (2.8)$$

Proposition 2.2. *Let $Y = aX + b$, with $a > 0$ and $b \geq 0$. Then, for their past varentropies, we have*

$$V({}_tY) = V\left(\frac{t-b}{a}X\right), \quad \forall t. \quad (2.9)$$

Proof. From $Y = aX + b$ we know that $F_Y(x) = F_X\left(\frac{x-b}{a}\right)$ and $f_Y(x) = \frac{1}{a}f_X\left(\frac{x-b}{a}\right)$. Hence, from (2.3) and (2.8), we get

$$V({}_tY) = \int_0^{\frac{t-b}{a}} \frac{f_X(x)}{F_X\left(\frac{t-b}{a}\right)} \left(\log \frac{\frac{1}{a}f_X(x)}{F_X\left(\frac{t-b}{a}\right)} \right)^2 dx - \left(H\left(\frac{t-b}{a}X\right) + \log a \right)^2. \quad (2.10)$$

By writing

$$\log \frac{\frac{1}{a}f_X(x)}{F_X\left(\frac{t-b}{a}\right)} = \log \frac{f_X(x)}{F_X\left(\frac{t-b}{a}\right)} - \log a,$$

and developing the two squares in (2.10), after some calculations we get the thesis. \square

If ϕ is any continuous, differentiable and strictly monotonic function and

$$Y = \phi(X),$$

then the past entropies of X and Y are related, for all $t > 0$, by

$$H(tY) = \begin{cases} H(\phi^{-1}(t)X) + \mathbb{E}[\log \phi'(X)|X < \phi^{-1}(t)], & \text{if } \phi \text{ is strictly increasing,} \\ H(X_{\phi^{-1}(t)}) + \mathbb{E}[\log(-\phi'(X))|X > \phi^{-1}(t)], & \text{if } \phi \text{ is strictly decreasing} \end{cases} \quad (2.11)$$

(see [9]). In the following result, the past varentropy under monotonic transformations is studied.

Proposition 2.3. *Let $Y = \phi(X)$, where ϕ is a continuous, differentiable and strictly monotonic function.*

(i) *If ϕ is strictly increasing, then*

$$\begin{aligned} V(tY) &= V(\phi^{-1}(t)X) - 2\mathbb{E}\left[\log \frac{f_X(X)}{F_X(\phi^{-1}(t))} \log \phi'(X) \middle| X < \phi^{-1}(t)\right] \\ &\quad + \text{Var}[\log \phi'(X)|X < \phi^{-1}(t)] - 2H(\phi^{-1}(t)X) [\log \phi'(X)|X < \phi^{-1}(t)]. \end{aligned}$$

(ii) *If ϕ is strictly decreasing, then*

$$\begin{aligned} V(tY) &= V(X_{\phi^{-1}(t)}) - 2\mathbb{E}\left[\log \frac{f_X(X)}{F_X(\phi^{-1}(t))} \log(-\phi'(X)) \middle| X > \phi^{-1}(t)\right] \\ &\quad + \text{Var}[\log(-\phi'(X))|X > \phi^{-1}(t)] - 2H(X_{\phi^{-1}(t)}) [\log(-\phi'(X))|X > \phi^{-1}(t)]. \end{aligned}$$

Proof. Let us suppose ϕ strictly increasing. From $Y = \phi(X)$ we know that $F_Y(x) = F_X(\phi^{-1}(x))$ and $f_Y(x) = \frac{f_X(\phi^{-1}(x))}{\phi'(\phi^{-1}(x))}$. Hence, from (2.3) and (2.11), we get

$$\begin{aligned} V(tY) &= \int_0^{\phi^{-1}(t)} \frac{f_X(x)}{F_X(\phi^{-1}(t))} \left(\log \frac{f_X(x)}{F_X(\phi^{-1}(t))} - \log \phi'(x) \right)^2 dx \\ &\quad - [H(\phi^{-1}(t)X) + \mathbb{E}[\log \phi'(X)|X < \phi^{-1}(t)]]^2. \end{aligned}$$

Hence, by developing the two squares in the previous equality, and observing that

$$\begin{aligned} \int_0^{\phi^{-1}(t)} \frac{f_X(x)}{F_X(\phi^{-1}(t))} (\log \phi'(x))^2 dx &= \mathbb{E}^2[\log \phi'(X)|X < \phi^{-1}(t)] \\ &= \text{Var}[\log \phi'(X)|X < \phi^{-1}(t)], \end{aligned}$$

we obtain the result.

The proof of (ii) is similar. □

3. Bounds for past varentropy

We recall the definition of variance inactivity time function, that is the variance of the inactivity time $X_{(t)}$ and can be expressed as

$$(\sigma^2(t))^* = \text{Var}(X_{(t)}) = \text{Var}(t - X | X \leq t) = \frac{2}{F(t)} \int_0^t dy \int_0^y F(x) dx - (m^*(t))^2,$$

where $m^*(t)$ is defined in (1.2). (For more properties of the variance inactivity time function, see Kandil et al. [21]).

Theorem 3.1. *Let $X_{(t)}$ be the inactivity time at time t of X , and let the mean inactivity time $m^*(t)$ and the variance inactivity time $(\sigma^2(t))^*$ be finite. Then, for all $t \in D$,*

$$V({}_tX) \geq (\sigma^2(t))^* \left[\mathbb{E}(\omega'_{(t)}(X_{(t)})) \right]^2,$$

where the function $\omega_{(t)}(x)$ is defined by

$$(\sigma^2(t))^* \omega_{(t)}(x) f_{X_{(t)}}(x) = \int_0^x (m^*(z) - z) f_{X_{(t)}}(z) dz, \quad x > 0. \quad (3.1)$$

Proof. First of all, notice that $V({}_tX) = V(X_{(t)})$. If X is a random variable with pdf f , mean μ and variance σ^2 , we have

$$\text{Var}[g(X)] \geq \sigma^2 [\mathbb{E}(\omega(X)g'(X))]^2, \quad (3.2)$$

where $\omega(x)$ is defined by $\sigma^2 \omega(x) f(x) = \int_0^x (\mu - z) f(z) dz$ (see Cacoullos and Papathanasiou [5]). Hence, by choosing $X_{(t)}$ as the random variable in (3.2), and $g(x) = -\log f_{X_{(t)}}(x)$, we obtain

$$\text{Var}(-\log f_{X_{(t)}}(X_{(t)})) = V({}_tX) \geq (\sigma^2(t))^* \left[\mathbb{E} \left(\omega_{(t)}(X_{(t)}) \frac{f'_{X_{(t)}}(X_{(t)})}{f_{X_{(t)}}(X_{(t)})} \right) \right]^2. \quad (3.3)$$

By differentiating now both sides of (3.1), we get

$$\omega_{(t)}(x) \frac{f'_{X_{(t)}}(x)}{f_{X_{(t)}}(x)} = \frac{m^*(t) - x}{(\sigma^2(t))^*} - \omega'_{(t)}(x),$$

and then by (3.3)

$$\begin{aligned} V({}_tX) &\geq (\sigma^2(t))^* \left[\mathbb{E} \left(\frac{m^*(t) - X_{(t)}}{(\sigma^2(t))^*} - \omega'_{(t)}(X_{(t)}) \right) \right]^2 \\ &= (\sigma^2(t))^* \left[\mathbb{E}(\omega'_{(t)}(X_{(t)})) \right]^2. \end{aligned}$$

□

Theorem 3.2. *Let X be a non-negative random variable with support D and log-concave pdf f . Then*

$$V({}_tX) \leq 1 \quad \text{for all } t \in D.$$

Proof. Observe that if $f(x)$ is log-concave, then also $f_t(x) = \frac{f(x)}{F(t)}$ is log-concave. It follows from Theorem 2.3 of Fradelizi et al. [18] that if X has a log-concave pdf, then $V(X) \leq 1$ and so the proof follows from this remark. \square

4. Proportional reversed hazard rate model

In this section we consider the reversed proportional hazard rate model, see Gupta and Gupta [19] for details.

The family of random variables $\{X^{(a)} : a > 0\}$ follows a proportional reversed hazard rate model if there exists a non-negative random variable X with cdf F and pdf f such that

$$F^{(a)}(t) = \mathbb{P}(X^{(a)} \leq t) = [F(t)]^a, \quad f^{(a)}(t) = a[F(t)]^{(a-1)}f(t), \quad t > 0. \quad (4.1)$$

We remark that the model takes the name from the fact that the reversed hazard rate functions of the random variables in the family are proportional to the reversed hazard rate function of X :

$$q^{(a)}(t) = \frac{f^{(a)}(t)}{F^{(a)}(t)} = a \frac{f(t)}{F(t)} = a q(t).$$

Moreover, the cumulative reversed hazard rate function can be expressed as

$$\Lambda^{(a)*}(t) = -\log F^{(a)}(t) = a\Lambda^*(t).$$

The proportional reversed hazard rate model finds applications with parallel systems. In fact, if we have a system composed by n units in parallel and characterized by i.i.d. lifetimes X_1, \dots, X_n with cdf $F(t)$, the lifetime of the system is given by $X^{(n)} = \max\{X_1, \dots, X_n\}$. Then, we have $F_{X^{(n)}}(t) = [F(t)]^n$, i.e., the system satisfies the proportional reversed hazard rate model (4.1) with $a = n$.

In order to evaluate the past varentropy of $X^{(a)}$, we determine the past entropy of

$X^{(a)}$. This can be expressed as

$$\begin{aligned} H\left({}_tX^{(a)}\right) &= -\Lambda^{(a)*}(t) - \frac{1}{[F(t)]^a} \int_0^t f^{(a)}(x) \log f^{(a)}(x) dx \\ &= -a\Lambda^*(t) - \frac{1}{[F(t)]^a} \int_0^{[F(t)]^a} \gamma(y; a) dy, \end{aligned}$$

with the change of variable $y = [F(x)]^a$, and where $\gamma(y; a) = \log [ay^{1-1/a} f(F^{-1}(y^{1/a}))]$.

Hence, we obtain the past varentropy of $X^{(a)}$ as

$$\begin{aligned} V\left({}_tX^{(a)}\right) &= \frac{1}{[F(t)]^a} \int_0^t f^{(a)}(x) (\log f^{(a)}(x))^2 dx - \left[\frac{1}{[F(t)]^a} \int_0^t f^{(a)}(x) \log f^{(a)}(x) dx \right]^2 \\ &= \frac{1}{[F(t)]^a} \int_0^{[F(t)]^a} [\gamma(y; a)]^2 dy - \left[\frac{1}{[F(t)]^a} \int_0^{[F(t)]^a} \gamma(y; a) dy \right]^2. \end{aligned}$$

Conclusions

In this paper, we have introduced and studied the past varentropy. It is related to the past entropy, which is a measure of information about the past lifetime distribution. In particular, the past varentropy provides the variability of the information given by the past entropy. We have showed several properties of the past varentropy, such as its behaviour under linear or monotonic transformations and what happens if it is constant. In addition, we have given bounds for the past varentropy and an application to the proportional reversed hazard rate model.

Acknowledgements

Francesco Buono and Maria Longobardi are partially supported by the GNAMPA research group of INdAM (Istituto Nazionale di Alta Matematica) and MIUR-PRIN 2017, Project "Stochastic Models for Complex Systems" (No. 2017 JFFHSH).

References

- [1] Arikan, E. (2016). Varentropy decreases under polar transform. *IEEE Transactions on Information Theory*, **62**, 3390–3400.
- [2] Block, H. W., Savits, T. H. (1998). The Reversed Hazard Rate Function. *Probability in the Engineering and Informational Sciences* **12**: 69–90.

- [3] Bobkov, S., Madiman, M. (2011). Concentration of the information in data with log-concave distributions.. *Annals of Probability*, **39**, 1528–1543.
- [4] Buono, F., Longobardi, M., Szymkowiak, M. (2020). On Generalized Reversed Aging Intensity Functions. *Preprint under review*.
- [5] Cacoullos, T., Papathanasiou, V. (1989). Characterizations of distributions by variance bounds. *Statistics & Probability Letters*, **7**(5), 351–356.
- [6] Cal, C., Longobardi, M., Ahmadi, J. (2017). Some properties of cumulative Tsallis entropy. *Physica A: Statistical Mechanics and its Applications*, **486**, 1012–1021.
- [7] Cal, C., Longobardi, M., Navarro, J. (2020). Properties for generalized cumulative past measures of information. *Probability in the Engineering and Informational Sciences*, **34** (1), 92–111.
- [8] Cal, C., Longobardi, M., Psarrakos, G. (2019). A family of weighted distributions based on the mean inactivity time and cumulative past entropies . *Ricerche di Matematica* (in press). DOI: 10.1007/s11587-019-00475-7
- [9] Di Crescenzo, A., Longobardi, M. (2002). Entropy-based measure of uncertainty in past lifetime distributions. *Journal of Applied Probability*, **39**, 434–440.
- [10] Di Crescenzo, A., Longobardi, M. (2006). On weighted residual and past entropies. *Scientiae Math. Japon.*, **64**, 255–266.
- [11] Di Crescenzo, A., Longobardi, M. (2009). On cumulative entropies. *J. Stat. Plann. Infer.*, **139**, 4072–4087.
- [12] Di Crescenzo, A., Longobardi, M. (2009). On cumulative entropies and lifetime estimations. In: Methods and Models in Artificial and Natural Computation, IWINAC 2009, *Lecture Notes in Computer Science* , **5601**, 132–141. Springer-Verlag, Berlin.
- [13] Di Crescenzo, A., Longobardi, M. (2012). Neuronal data analysis based on the empirical cumulative entropy. In: Computer Aided Systems Theory, EUROCAST 2011, Part I, *Lecture Notes in Computer Science* , **6927**, 72–79. Springer-Verlag, Berlin.
- [14] Di Crescenzo, A., Longobardi, M. (2013). Stochastic comparisons of cumulative entropies. In: “Stochastic Orders in Reliability and Risk, In Honor of Professor Moshe Shaked” (H. Li and X. Li, eds.), *Lecture Notes in Statistics*, **208**, 167–182. Springer, New York .
- [15] Di Crescenzo, A., Paolillo, L. (2020). Analysis and applications of the residual varentropy of random lifetimes. *arXiv*: 2003.04932.
- [16] Ebrahimi, N. (1996). How to measure uncertainty in the residual life time distribution. *Sankhya: Series A*, **58**, 48–56.

- [17] Finkelstein, M. S. (2002). On the Reversed Hazard Rate. *Reliability Engineering & System Safety* **78**: 71–75.
- [18] Fradelizi, M., Madiman, M., Wang, L. (2016). Optimal Concentration of Information Content for Log-Concave Densities. In: Houdr C., Mason D., Reynaud-Bouret P., Rosiski J. (eds) *High Dimensional Probability VII. Progress in Probability*, **71**, 45–60.
- [19] Gupta, R. C., Gupta, R. D. (2007). Proportional reversed hazard rate model and its applications. *Journal of Statistical Planning and Inference* **137**: 3525–3536.
- [20] Izadkhan, S. (2014). Mean Inactivity Time Function, Associated Orderings, and Classes of Life Distributions. *IEEE Transactions on Reliability*, **63**, 593–602.
- [21] Kandil, A. M., Kayid, M., Mahdy, M. (2011). Variance Inactivity Time Function and its Reliability Properties. *The 45th Annual Conference on Statistics, Computer Science and operations Research ISRR*, Cairo-Egypt, 94–113.
- [22] Kunda, C., Nanda, A., Maiti, S. (2010). Some distributional results through past entropy. *Journal of Statistical Planning and Inference* **140**: 1280–1291.
- [23] Li, X., Zuo, M. J. (2004). Stochastic Comparison of Residual Life and Inactivity Time at a Random Time. *Stochastic Models*, **20** (2), 229–235.
- [24] Longobardi, M. (2014). Cumulative measures of information and stochastic orders. *Ricerche Mat.*, **63**, 209–223.
- [25] Madiman, M., Wang, L. (2014). An optimal varentropy bound for log-concave distributions. *International Conference on Signal Processing and Communications (SPCOM)*, Bangalore, doi: 10.1109/SPCOM.2014.6983953.
- [26] Muliere, P., Parmigiani, G., Polson, N. G. (1993). A Note on the Residual Entropy Function. *Probability in the Engineering and Informational Sciences*, **7**, 413–420.
- [27] Shannon, C. E. (1948). A mathematical theory of communication. *Bell System Technical Journal*, **27**, 379–423.
- [28] Tahmasebi, S., Longobardi, M., Foroghi, F., Lak, F. (2019). An extension of weighted generalized cumulative past measure of information *Ricerche di Matematica* (in press). DOI: 10.1007/s11587-019-00448-w