

Gauging away the Stoner model: Engineering unconventional metallic ferromagnetism with artificial gauge fields

Joshuah T. Heath* ¹, Kevin S. Bedell¹,

[1] Physics Department, Boston College, Chestnut Hill, Massachusetts 02467, USA
heathjo@bc.edu

January 10, 2022

Abstract

For nearly a century, the Stoner model has dominated research in itinerant ferromagnetism, yet recent work on ultracold, optically trapped Fermi gases suggests that phase separation of spins occurs independent of long-range magnetic order. In this paper, we consider the breakdown of the Stoner criterion in a Landau-Fermi liquid with a weak non-zero gauge field. Due to a process analogous to Kohn's theory of a metal-insulator transition, we find the stability of a Fermi liquid phase in the absence of non-quasiparticle contributions to the response is strongly dependent on Landau parameters of mixed partial waves. Our work paves the way for the description of novel spintronic hardware in the language of Landau-Fermi liquid transport.

The study of artificial gauge fields has recently gained traction as a means of exploring physics beyond conventional electromagnetism [1–4]. The creation of a spin-dependent gauge field through synthetic spin-orbit (SO) coupling is of particular interest in the field of spintronics as a means to create a finite spin current [1,5–11], facilitating the development of novel solid state devices which exploit the electron's spin degrees of freedom. As we approach the inevitable termination of Moore's law, such machinery (and, subsequently, the gauge fields which support them) will prove to be indispensable for the development and implementation of new information storage and logic devices.

In addition to a spin-dependent gauge field, the generation of spin polarization necessary for spintronic hardware usually requires the presence of a non-equilibrium spin bath, by which spin-polarized transport can be studied [12] and relaxation back to equilibrium measured and ultimately controlled for the efficient implementation of fast switching [13,14]. The itinerant ferromagnetism required for such non-equilibrium behavior was originally described in terms of the Stoner paradigm [15,16], in which a repulsive exchange interaction in the spin-1/2 Fermi gas results in a segregation of delocalized spins into distinct domains [17–19]. Nevertheless, a significant number of theoretical [20–23] and experimental [24,25] works have suggested that quantum correlational effects neglected in Stoner's original mean-field argument competes with ferromagnetism. Similar work on ultracold gases of ⁶Li atoms suggests that the phenomenon of spin segregation occurs independently of ferromagnetic ordering [26–28], once

again in violation of Stoner's model. As the former point contradicts the theory of ferromagnetic Fermi liquids [29–35] and the latter contradicts the standard Pomeranchuk argument, the applicability of Landau-Fermi liquid theory to materials of spintronic interest is called into question if Stoner's argument remains problematic, threatening to backtrack significant progress made in the marriage of the two fields [12, 36–41].

In this paper, we show that a stable metallic phase can coexist with ferromagnetic order in the absence of spin segregation as long as the system supports some finite, weak gauge field. A non-zero vector potential results in a universal translation of the entire Fermi sea, and subsequently induces a novel interdependence between Landau parameters of unequal partial waves if we are to ensure adiabatic continuity with the non-interacting system remains valid. By taking into account dynamically-generated instabilities of zero sound, we find that the metallic phase for non-zero gauge field is protected by phase separation irrespective of long-range ferromagnetic ordering. This differs greatly from the traditional Stoner model, where phase separation and magnetic order occur synchronously. In this way, we show that something as simple as a gauge field can induce non-trivial momentum dependence in the near-equilibrium degrees of freedom which stabilize the Fermi liquid ansatz against time-reversal symmetry breaking.

Fermi surface instabilities of a standard Landau-Fermi liquid occur when the free energy functional

$$\delta\mathcal{F} \approx \sum_{p\sigma} (\epsilon_{p\sigma} - \mu) \delta n_{p\sigma} + \frac{1}{2} \sum_{pp'\sigma\sigma'} f_{pp'}(\sigma, \sigma') \delta n_{p\sigma} \delta n_{p'\sigma'} \quad (1)$$

becomes negative, where $\epsilon_{p\sigma}$ is the quasiparticle energy, $\delta n_{p\sigma} = n_{p\sigma} - n_{p\sigma}^0$ is the displacement of the distribution function, and the Landau parameter $f_{pp'}(\sigma, \sigma')$ is the second functional derivative of the total energy with respect to $\delta n_{p\sigma}$. Expanding the Landau parameters in terms of spherical harmonics, we can recast the above as

$$\delta\mathcal{F} = \frac{N(0)}{8} \sum_{\ell} \frac{1}{2\ell+1} \left[\nu_{\ell s}^2 \left(1 + \frac{F_{\ell}^s}{2\ell+1} \right) + \nu_{\ell a}^2 \left(1 + \frac{F_{\ell}^a}{2\ell+1} \right) \right] \quad (2)$$

where $f_{pp'}(\sigma, \sigma') = \frac{1}{N(0)} \sum_{\ell} (F_{\ell}^s + \sigma \cdot \sigma' F_{\ell}^a) P_{\ell}(\cos \theta)$, $N(0)$ is the density of states, and the Fermi surface distortion $\nu_{\ell s/a} = \sum_{\ell} (\nu_{\ell \uparrow} \pm \nu_{\ell \downarrow}) P_{\ell}(\cos \theta)$ is defined by $\delta n_p = -\frac{\partial n_k^0}{\partial \epsilon_k} \nu_p$.

In the s-wave channel, the stability condition $\delta\mathcal{F} > 0$ rests solely on the values of the symmetric (F_0^s) and anti-symmetric (F_0^a) $\ell = 0$ Landau parameters. Because the condition $1 + F_0^s < 0$ ($1 + F_0^a < 0$) leads to a negative compressibility (spin susceptibility), violations of the Pomeranchuk condition in the (anti)-symmetric Landau parameter mark the onset of a structural phase transition mediated by long-wavelength charge density (spin) fluctuations. However, note that a violation of the stability condition in *one* of the Landau parameters does not necessarily imply a complete breakdown of the Fermi liquid ansatz. In the limit of weak (i.e., local) ferromagnetism, the symmetric and anti-symmetric Landau parameters are related by $F_0^s = -F_0^a/(1 + 2F_0^a)$ [42, 43], and only then (for general non-zero ν_{0s} and ν_{0a}) does the resultant ferromagnetic instability occur concurrently with the segregation of spin populations at the Fermi surface. It therefore becomes apparent that a Landau-Fermi description of the itinerant ferromagnetism demonstrated in ultracold gases of ^6Li atoms [27, 28] must be realized as an "unconventional" variant of ferromagnetism characterized by higher non-vanishing orbital partial waves [37].

Higher-order $\ell > 0$ Pomeranchuk instabilities have been the subject of recent interest, with previous studies suggesting that such instabilities in the spin channel are accompanied by a dynamically-generated, non-relativistic spin-orbit (SO) coupling [36,37]. The existence of an "emergent" SO coupling from low-energy collective phenomenon was brought into question by [44], where the divergence of the quasiparticle susceptibility $\chi(\mathbf{q}, \omega)$ in the $\ell = 1$ channel is removed by virtue of a vanishing vertex coupling between Landau quasiparticles and bare electrons [45]; i.e., the non-quasiparticle contribution to the susceptibility exactly cancels the divergence in the quasiparticle contribution. As a result, the full disappearance of an $\ell = 1$ instability can only be realized for a specific $\mathbf{k}$ -dependence of the form factor $\lambda_{\ell=1}^{s/a}(\mathbf{k})$ [46]. A generic order parameter with different functional dependence of $\lambda_{\ell=1}^{s/a}(\mathbf{k})$ will result in the onset of Pomeranchuk instabilities for $\ell > 0$, while Pomeranchuk instabilities towards phases with spontaneously generated charge/spin currents prove to be impossible [47,48]. To remain consistent with the recent literature, we limit our study to those systems where the order parameter of the resultant phase has the same symmetry as the charge/spin current. This corresponds to a nearly isotropic system where the high-energy and low-energy contributions to the susceptibility do not enjoy a direct correlation [46].

The dynamic SO coupling described by [36,37] can be reformulated in terms of an SU(2) gauge field [1,49–53]. The effects of a gauge potential $\mathbf{A}_\sigma$ on the Fermi surface has been noted previously in connection with Kohn's theorem [54,55] and the anomalous Hall effect [56], yet a detailed study of the resultant Pomeranchuk instability has not been explored. To address this issue, we consider a general vector potential $\mathbf{A}_\sigma$, where $\mathbf{A}_\uparrow = \mathbf{A}_\downarrow \equiv \mathbf{A}$ for a symmetric (i.e., electromagnetic-like or spin-independent) gauge field and $\mathbf{A}_\uparrow = -\mathbf{A}_\downarrow \equiv \mathbf{A}$ for an anti-symmetric (i.e., Rashba/Dresselhaus-like [57–62] or spin-dependent) gauge. Note that in the presence of the many-body SO coupling which generates a spin-dependent gauge field, the Landau parameter takes on a more complicated form as a generalized tensor under rotations of spatial momenta [63]. In addition to terms proportional to $\delta_{\alpha\bar{\alpha}}\delta_{\alpha'\bar{\alpha}'}$ and $(\sigma_i)_{\alpha\bar{\alpha}}(\sigma_j)_{\alpha'\bar{\alpha}'}$ (where the subscripts denote matrix elements and σ_i is the Pauli matrix), the presence of a net spin polarization and finite spin-orbit coupling implies terms proportional to $(\sigma_i)_{\alpha\bar{\alpha}}\delta_{\alpha'\bar{\alpha}'}$ and $\delta_{\alpha\bar{\alpha}}(\sigma_i)_{\alpha'\bar{\alpha}'}$ in the Landau interaction parameter, with generic coefficients proportional to the magnitude of the momenta $\mathbf{p}$ and $\mathbf{p}'$. This is a direct consequence of a net spin polarization of all electrons in the Fermi sea and, hence, a net displacement of the Fermi seas of up- and down-spin particles in opposite directions, resulting in the violation of spherical symmetry of the Fermi surface and intrinsic dependence of $f_{pp'}(\sigma, \sigma')$ on the azimuthal quantum number m . For this reason, we restrict ourselves to the limit $\frac{e}{c}\mathbf{A}_\sigma \ll \mathbf{p}_F$, allowing us to take the previously mentioned scalar decomposition of the Landau parameter in terms of F_ℓ^s and F_ℓ^a . Such an approximation has similarly been taken in the chiral Fermi liquid description of electron systems with weak Rashba [40] and Dresselhaus [64] SO coupling, 2D electron gases with zero external magnetic field [65], and weakly polarized Fermi liquids in a quasi-equilibrium state [66].

For some weak non-zero gauge field, the momentum of every particle in the Fermi sea is shifted by the same amount: $\mathbf{p}_F \rightarrow \mathbf{p}_F - \mathbf{A}_\sigma$ [55,56], taking units where $e = c = 1$. The resultant Fermi surface distortion can be found by expanding the Heaviside step function

about $p_F(\theta, \phi) = p_F - A_\sigma$:

$$\begin{aligned}\delta n_{p-A_\sigma, \sigma} &= \Theta(p_F(\theta, \phi) - p + A_\sigma) - \Theta(p_F - p) \\ &= \delta n_{p\sigma} - A_\sigma \delta(p_F - p) - A_\sigma \delta p_F \frac{\partial}{\partial p} \delta(p_F - p) \\ &\quad - \frac{1}{2} A_\sigma^2 \frac{\partial}{\partial p} \delta(p_F - p)\end{aligned}\quad (3)$$

where $\delta p_F \equiv p_F(\theta, \phi) - p_F$ defines the anisotropic shift of the Fermi momentum at finite interaction. The first term $\delta n_{p\sigma} = -\delta p_F \delta(p_F - p) - \frac{1}{2} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p)$ is identical to what is seen in a traditional Fermi liquid, while the last three terms are specific to Fermi liquids with some finite gauge. After a tedious but straightforward calculation, we can expand the first expression in Eqn. (1) for $\mathbf{A}_\sigma \neq 0$ in terms of partial waves, yielding

$$\sum_{p\sigma} (\epsilon_{p-A_\sigma, \sigma} - \mu) \delta n_{p-A_\sigma, \sigma} \approx \frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left\{ (\nu_{\ell s}^2 + \nu_{\ell a}^2) + \frac{16A}{p_F} \epsilon_F \nu_{\ell, s/a} \right\} + \frac{48A^2}{p_F^2} \epsilon_F^2 \right] \quad (4)$$

where $\nu_{\ell, s/a} = \nu_{\uparrow} \pm \nu_{\downarrow}$ for the symmetric and anti-symmetric gauges, respectively. In the above derivation, we assume that $\epsilon_F \gg \nu_{s/a}$ to keep the expression tractable, and have set $|\mathbf{A}_\sigma| \equiv A$.

A similar but somewhat more complex calculation can be performed on the second term in Eqn. (1) for finite gauge field (see Supplemental Material). When the dust settles, we are left with

$$\begin{aligned}\sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta n_{p-A_\sigma, \sigma} \delta n_{p'-A_{\sigma'}, \sigma'} \\ \approx \frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left(\frac{1}{2\ell+1} \{ F_\ell^s \nu_{\ell s}^2 + F_\ell^a \nu_{\ell a}^2 \} + \frac{8A}{p_F} \frac{F_\ell^s}{2\ell+1} \epsilon_F \nu_{\ell, s/a} \delta_{\ell 0} \right) + \frac{64A^2}{9p_F^2} F_1^{s/a} \epsilon_F^2 \right] \quad (5)\end{aligned}$$

where $F_\ell^{s/a}$ is F_ℓ^s for the symmetric gauge field and F_ℓ^a for the anti-symmetric gauge. Note that the second term is proportional to F_ℓ^s independent of what variety of gauge field we consider, and only remains finite for the $\ell = 0$ channel.

We can see from the above that the presence of a non-zero gauge field refashions the original $\ell = 0$ Pomeranchuk instability into a generalized condition between $F_1^{s/a}$ and $F_0^{s/a}$. This is a direct consequence of the Landau-Fermi free energy transforming into a second-order polynomial with respect to the Fermi surface distortion ν_ℓ , as opposed to simply being dependent on ν_ℓ^2 as seen in a conventional Fermi liquid. The presence of a linear ν_ℓ and constant term from the gauge field puts new constraints on an equilibrium value of ν_ℓ to ensure that such distortions are real. This yields the following stability conditions for the spin symmetric and spin anti-symmetric gauge fields, assuming $F_0^{s/a} < -1$:

$$\mathbf{A}_\uparrow = \mathbf{A}_\downarrow, \quad F_0^s < -1 \rightarrow F_1^s \geq \frac{9}{4} \left(\frac{1}{1+F_0^s} + F_0^s \right) \quad (6a)$$

$$\mathbf{A}_\uparrow = -\mathbf{A}_\downarrow, \quad F_0^a < -1 \rightarrow F_1^a \geq \frac{9}{4} \left(\frac{1-3F_0^a+F_0^s(4+F_0^s)}{1+F_0^a} \right) \quad (6b)$$

where we assume that $\nu_{0a} = 0$ for the symmetric gauge field and $\nu_{0s} = 0$ for the anti-symmetric gauge. If we instead take $\nu_{0s} = 0$ ($\nu_{0a} = 0$) for the (anti-)symmetric gauge, Eqns. (6a) and (6b) are simplified to $F_1^{s/a} \geq -27/4$. Whereas the $\ell = 0$ stability condition for $\mathbf{A}_\sigma = 0$ is solely determined by $F_0^{s/a}$, the presence of some weak finite gauge field implies that the stability of the Fermi surface has strong dependence on higher-order Landau parameters. As the spin susceptibility is calculated from the change in the *local* energy of a quasiparticle as the result of spin displacement [67], the value of the spin susceptibility is not effected by a uniform translation of the Fermi sea and is thus a gauge-invariant quantity. This suggests that ferromagnetic ordering still occurs when $F_0^a < -1$ for $\mathbf{A}_\sigma \neq 0$, albeit "disentangled" from the phase separation that plagues the local model.

In Fig. 1, we can see under what conditions the Fermi liquid is stable for the symmetric gauge. The left-hatched region corresponds to the only combination of Landau parameters F_0^s and F_1^s where equilibrium Fermi liquid theory completely breaks down in the presence of a spin-symmetric gauge. The right-hatched region corresponds to a dynamical instability; i.e., where there exists some critical $\nu_{0s} \neq 0$ that pushes $\delta\mathcal{F} < 0$. Such instabilities have previously been noted in ferromagnetic Fermi liquids with spin-orbit coupling, where the collective mode can drive the system to a Lifshitz transition [68]. Only in the white region above the top blue line of Fig. 1 is the Fermi liquid phase completely free from instabilities for all values of $\nu_{\ell s}$. In Fig. 2, we see where the Fermi liquid is stable in the anti-symmetric gauge, with the regime of phase separation for $\nu_{0s} \sim 0$ reduced to a small sliver near $F_0^s \sim -2$ as $F_0^a \rightarrow -1$ from below. Here, the right-hatched region corresponds to those F_1^a values where a dynamical instability is possible for $F_0^a \in [-2, 0]$, while the left-hatched corresponds to where $\delta\mathcal{F} < 0$ indefinitely in this range. While both the symmetric and anti-symmetric gauge fields increase the stability of the Fermi surface, the latter nearly eliminates the threat of phase separation at near-equilibrium if we are in close proximity to a ferromagnetic instability.

Physically, we associate the newfound stability of the itinerant metallic phase with the condensation of gauge bosons. This conclusion is hinted at by repeating the calculation of Eqns. (4)+(5) for the $\ell = 1$ channel, in which we find the constraint $\frac{3}{8}(-13 - \sqrt{89}) < F_1^{s/a} < \frac{3}{8}(-13 + \sqrt{89})$ if we want to simultaneously have $1 + F_1^s/3 < 0$ and $\delta\mathcal{F} \geq 0$. This implies that, in the presence of a gauge, the fundamental carriers of the gauge field Bose condense and subsequently lead to the breakdown of Galilean invariance [69]. Otherwise, the effective mass of the Landau quasiparticle $m^*/m = 1 + F_1^s/3$ could become negative, resulting in a Fermi surface instability and hence a contradiction [70]. The breakdown of Galilean invariance is of fundamental importance to our study, otherwise the addition of a spin-independent gauge field wouldn't affect the stability of the system. At a fundamental level, the introduction of either a natural or synthetic electromagnetic interaction through a gauge principle is well-known to be incompatible with Galilean field theories [71]. The argument from condensation is a direct consequence of a nonintegrable phase attached to the electron field operators (as previously noted in the context of Goldstone boson condensation [72]), and can be viewed as a manifestation of the Aharonov-Bohm effect [73]. As in ^3He - ^4He mixtures, a finite superfluid velocity defines a preferred frame and Galilean invariance of the fermionic system is subsequently lost. The fermionic system we consider can therefore be viewed as a non-Galilean-invariant Fermi liquid. Such a variation of the traditional Landau-Fermi liquid has already been noted to exist in the presence of a lattice geometry when the liquid phase exhibits low carrier concentrations [74].




Figure 1: Visual representation of the Pomeranchuk instability for the symmetric gauge field (Eqn. (6a)) for $\nu_{0a} = 0$. In a stable Landau-Fermi liquid, F_1^s is forbidden to enter the dark-red left-hatched region, while the light-red right-hatched region is characterized by a dynamical instability. For $\nu_{0s} = 0$, the value of F_1^s is constrained to be above $-27/4$ (green dashed line).

A stronger argument for this condensation can be made by mapping the gauge bosons in our system to the Goldstone excitations in a macroscopic ring topology. In the metal-insulator transition of a topologically-trivial metal, condensation of the phonon mode in the dual geometry causes a distortion which breaks ring periodicity, producing a finite current in the simply connected space when we take the thermodynamic limit [75]. In this way, the condensation of the gauge field induces the formation of a metallic phase via symmetry breaking in the ordered state of the dual ring. The presence of finite- $\mathbf{A}$ dependence in the system's energy would then signify a metallic phase, while the absence of any gauge field-dependence would signify an insulating state [54,75]. Our calculations of $\ell = 0$ and $\ell = 1$ Fermi surface stabilities with a finite gauge field can be seen as a manifestation of this phenomenon in a Landau-Fermi liquid.

Finally, we would like to point out a subtlety in the above calculations concerning the appearance of apparent dynamical *stabilities* brought about by a finite gauge. In the limit of small Fermi surface distortion, the stability condition determined by Eqns. (4)+(5) reduces to $1 + \frac{4}{27}F_1^{s/a} \geq 0$. When this bound is broken, the Fermi liquid ground state is stable only for some non-zero values of $\nu_{0,s/a} \neq 0$. A calculation of the Drude weight $D_{s/a}$ yields a similar expression, with the Kohn relation [54,55,76] yielding

$$\begin{aligned} D_{s/a} &\equiv \left. \frac{\partial^2 E(\mathbf{A}_\sigma)}{\partial \mathbf{A}_\sigma^2} \right|_{\mathbf{A}_\sigma=0} \\ &= \frac{9\pi n}{m^*} \left(1 + \frac{4}{27}F_1^{s/a} \right) \end{aligned} \quad (7)$$

where $E(\mathbf{A}_\sigma)$ is the energy density. While a positive Drude weight signals a stable metallic phase, a negative Drude weight for the symmetric gauge field implies an unstable maximum and an unusual paramagnetic response [77]. For the anti-symmetric gauge, the negative Drude weight implies a negative spin stiffness and hence an instability away from ferromagnetic ordering [78]. These results suggest that systems with $F_1^{s/a} \leq -\frac{27}{4}$ are marked by instabilities not captured by the above analysis.

We interpret the dynamic fragility of the $\mathbf{A}_\sigma \neq 0$ Fermi liquid phase for $F_1^{s/a} \leq -\frac{27}{4}$ as a violation of Mermin's theorem on the sufficient conditions for the propagation of zero sound [79]. In a nutshell, Mermin's seminal work relies upon four ingredients: (1) the stability condition Eqn. (2); (2) the Landau kinetic equation; (3) the relation between the forward scattering amplitude $a_{pp'}^{s/a}$ and the Landau parameter:

$$a_{pp'}^{s/a} = f_{pp'} + \sum_{p''} f_{pp''}^{s/a} \frac{\partial n_{p''}^0}{\partial \epsilon_{p''}} a_{p''p'}^{s/a} \quad (8)$$

and (4) the exclusion principle, which requires that $a_{pp}^s + a_{pp}^a = 0$. With these four conditions, it can be shown that the Landau kinetic equation has at least one eigenvalue λ of unity, and hence forward scattering is sufficient for the existence of regular or spin zero sound in a Fermi liquid [79]. In the presence of a gauge, the forward scattering amplitude and the exclusion principle are identical as in the $\mathbf{A}_\sigma = 0$ case. The stability for $\mathbf{A}_\sigma \neq 0$ has been discussed previously, and the previously homogenous kinetic equation is modified to include a term proportional to $\frac{\cos \theta}{\eta - \cos \theta} (\mathbf{A}_\sigma \cdot \frac{\mathbf{p}}{m})$. Combining this with the forward scattering sum rule, the kinetic equation becomes




Figure 2: Visual representation of the Pomeranchuk instability for the anti-symmetric gauge field (Eqn. (6b)) for $\nu_{0s} = 0$. For $F_0^a > -1$, the Fermi liquid experiences a dynamical instability when F_1^a is below the solid green line. For $F_0^a < -1$, phase separation occurs when F_1^a is below the solid blue line. For $\nu_{0s} = 0$, the value of F_1^s is constrained to be above $-27/4$ (green dashed line).

$$\nu = \left(\frac{\cos \theta}{\eta Y} + a^{s/a} \right) \nu + \cos \theta \left(\frac{(Y-1)\nu + X}{Y\eta} \right) \left\{ \frac{1 - f^{s/a}}{1 + f^{s/a}} \right\} \quad (9)$$

where η is the dimensionless phase velocity of zero sound, $Y \equiv a_{p,p-A}^{\sigma\sigma} \frac{\partial n_p^0}{\partial \epsilon_p}$, $X \approx \frac{A_\sigma p_F}{m} \cos \theta$, and the notation $\sum_{p'} B_{pp'} \frac{\partial n_{p'}^0}{\partial \epsilon_{p'}} \nu_{p'} \equiv -B\nu$ is taken for some observable $B = a_{pp'}^{s/a}$ or $f_{pp'}^{s/a}$. In the limit $A_\sigma \ll p_F$ and $\eta \rightarrow \infty$, Eqn. (9) reduces to $\nu = a^{s/a} \nu$ as in the original formulation. In the gauge-free theory $a_{pp'}^{s/a}$ is bounded to be below one from Eqn. (2). By calculating the terms in the kinetic equation, one can show [79] that there exists an η' such that $\lambda_{\eta'} > 1$, implying that there exists some η'' between η' and ∞ where $\lambda_{\eta''} = 1$. In the presence of a finite gauge, however, the modification of the Pomeranchuk condition results in the forward scattering amplitude taking values greater than one for $1 + F_0^{s/a} < 0$. This implies that forward scattering is not a sufficient condition for the propagation of zero sound if a weak gauge field is present. The onset of this instability can be seen explicitly by calculating the terms of interest in Eqn. (9):

$$\cos \theta \nu \equiv \eta^{-1} \int \frac{d\hat{n}}{4\pi} |\chi(\hat{n})|^2 \cos \theta = \frac{1}{2} - \frac{\eta^{-1} \mathbb{A}^2}{2} \int_{\cos \theta_0}^1 \frac{dx}{\eta - x} \quad (10a)$$

$$\begin{aligned} B\nu &\equiv \int \frac{d\hat{n}}{4\pi} \int \frac{d\hat{n}'}{4\pi} \chi^*(\hat{n}) B \chi(\hat{n}') \\ &= \frac{B \mathbb{A}^2}{4} \iint_{\cos \theta_0}^1 \frac{dx}{\eta - x} \frac{dx'}{\eta - x'} \end{aligned} \quad (10b)$$

where we have used the ansatz $\chi(\hat{n}) = \mathbb{A}/(\eta - \cos \theta)$, with $\mathbb{A}$ taken as a normalization constant and $0 < \theta < \theta_0$. As we approach $\eta = 1$ from above, the coefficient of B diverges. Rearranging Eqn. (9), we therefore find a constraint on the value of ν assuming some finite gauge:

$$\nu < \frac{X}{\eta} \cos \theta \left(\frac{\cos \theta}{\eta} + a^{s/a} \right)^{-1} \quad (11)$$

The existence of zero sound for $\mathbf{A}_\sigma \neq 0$ then requires that (1) the right-hand side of Eqn. (9) yields a solution < 1 for some η and (2) ν doesn't break the bound set by Eqn. (11). As the Fermi surface is only stable for larger, finite ν when $F_1^{s/a} < -27/4$, the calculation above suggests any metallic phase in this regime will be fragile to the disappearance of zero sound and subsequently a breakdown of conventional Landau-Fermi liquid theory in agreement with the Drude weight result.

In summary, we have investigated Fermi surface and zero sound instabilities in a Landau-Fermi liquid with weak, non-zero spin symmetric and spin anti-symmetric gauge fields $\mathbf{A}_\sigma$. Regardless of the magnitude of this gauge, we find that all Fermi liquids with $1 + F_0^{s/a} < 0$ are stable assuming that $1 + \frac{4}{27} F_1^{s/a} > 0$ and $\nu_{\ell, s/a}$ does not differ appreciably from zero. As a consequence, Landau-Fermi liquid theory is consistent with an itinerant form of ferromagnetism independent of phase separation as long as $\mathbf{A}_\sigma \neq 0$. Such a result complements the work done in [44, 46] on Fermi surface instabilities for partial waves $\ell > 0$; whereas these

studies suggest an $\ell = 1$ Pomeranchuk instability is forbidden under certain conditions of the form factor where high- and low-energy contributions to the susceptibility are directly correlated, our work shows $\ell = 0$ instabilities are eliminated in the presence of weak gauge fields and where non-quasiparticle contributions are absent from the response. The robust metallic phase we find is a clear manifestation of gauge boson condensation previously introduced in the context of a metal-insulator transition [75]. Experimentally, our work provides a physical explanation for the violation of the Stoner model in ultracold gases of ^6Li [26–28], as the degenerate mixture is confined to the lowest hyperfine states as the result of Zeeman splitting in a weak magnetic field and, hence, a spin-symmetric gauge potential. Our work similarly compliments existing studies of 2D spin-orbit coupled Fermi gases in a harmonic potential, where a Rashba-type gauge has been predicted to stabilize itinerant ferromagnetism via modification of the single-particle dispersion [80, 81].

Overall, the main message of this paper is a positive one: that a major ingredient for spintronic devices (i.e., a spin-dependent gauge field) serves to stabilize the itinerant ferromagnetism required for coherent spin manipulation. Our results strongly suggest that optically-trapped gases of neutral atoms such as ^6Li [82] and ^{40}K [83–85] are ideal candidates for spintronic hardware, as the resultant non-Abelian gauge fields [10, 52, 62, 86–92] in dimensions $D \geq 2$ legitimize a description of spin transport in the well-established language of Landau-Fermi liquid theory while simultaneously permitting maximal control over the gauge field itself.

Acknowledgements.—One of the authors (J.T.H.) thanks Matthew Gochan for useful feedback. We would also like to acknowledge Shou-Cheng Zhang for his work on unconventional ferromagnetism, which served as one of the catalysts for this project. This research was partially supported by the John H. Rourke endowment fund at Boston College.

References

- [1] M. S. Shikakhwa, S. Turgut and N. K. Pak, *Gauge covariance and spin-current conservation in the gauge-field formulation of systems with spin-orbit coupling*, J. Phys. A: Math. Theor. **45**(10), 105305 (2012).
- [2] J. Dalibard, F. Gerbier, G. Juzeliūnas and P. Öhberg, *Colloquium: Artificial gauge potentials for neutral atoms*, Rev. Mod. Phys. **83**, 1523 (2011), doi:10.1103/RevModPhys.83.1523.
- [3] V. Galitski and I. B. Spielman, *Spin-orbit coupling in quantum gases*, Nature **494**(7435), 49 (2013).
- [4] M. Aidelsburger, S. Nascimbene and N. Goldman, *Artificial gauge fields in materials and engineered systems*, C. R. Phys. **19**(6), 394 (2018).
- [5] S. Datta and B. Das, *Electronic analog of the electro-optic modulator*, Appl. Phys. Lett. **56**(7), 665 (1990).
- [6] S.-Q. Shen, *Spin transverse force on spin current in an electric field*, Phys. Rev. Lett. **95**, 187203 (2005), doi:10.1103/PhysRevLett.95.187203.

- [7] H. Marrows C. and J. Hickey B., *New directions in spintronics*, Philos. Trans. Royal Soc. A **369**(1948), 3027 (2011).
- [8] A. Fert, *Nobel lecture: Origin, development, and future of spintronics*, Rev. Mod. Phys. **80**, 1517 (2008), doi:10.1103/RevModPhys.80.1517.
- [9] P. A. Grünberg, *Nobel lecture: From spin waves to giant magnetoresistance and beyond*, Rev. Mod. Phys. **80**, 1531 (2008), doi:10.1103/RevModPhys.80.1531.
- [10] K. Premasiri and X. P. A. Gao, *Tuning spin-orbit coupling in 2D materials for spintronics: a topical review*, J. Phys.: Condens. Matter **31**(19), 193001 (2019).
- [11] Y. Liu, C. Zeng, J. Zhong, J. Ding, Z. M. Wang and Z. Liu, *Spintronics in Two-Dimensional Materials*, Nano-Micro Lett. **12**(1), 93 (2020).
- [12] H. Li and K. S. Bedell, *Quantum spin transport and dynamics through a ferromagnetic/normal metal junction*, Phys. Rev. B **92**, 224437 (2015), doi:10.1103/PhysRevB.92.224437.
- [13] Y. Nishikawa, A. Tackeuchi, S. Nakamura, S. Muto and N. Yokoyama, *All-optical picosecond switching of a quantum well etalon using spin-polarization relaxation*, Appl. Phys. Lett. **66**(7), 839 (1995).
- [14] I. Žutić, J. Fabian and S. Das Sarma, *Spintronics: Fundamentals and applications*, Rev. Mod. Phys. **76**, 323 (2004), doi:10.1103/RevModPhys.76.323.
- [15] E. C. Stoner, *LXXX. Atomic moments in ferromagnetic metals and alloys with non-ferromagnetic elements*, London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science **15**(101), 1018 (1933).
- [16] E. C. Stoner, *Collective electron ferromagnetism in metals and alloys*, J. Phys. Radium **12**(3), 372 (1951).
- [17] P. T. Grochowski, T. Karpiuk, M. Brewczyk and K. Rzazewski, *Unified description of dynamics of a repulsive two-component fermi gas*, Phys. Rev. Lett. **119**, 215303 (2017), doi:10.1103/PhysRevLett.119.215303.
- [18] V. Penna and L. Salasnich, *Itinerant ferromagnetism of two-dimensional repulsive fermions with Rabi coupling*, New J. Phys. **19**(4), 043018 (2017).
- [19] V. Penna and L. Salasnich, *Polarization in a three-dimensional Fermi gas with Rabi coupling*, J. Phys. B: At. Mol. Opt. Phys. **52**(3), 035301 (2019).
- [20] M. C. Gutzwiller, *Effect of correlation on the ferromagnetism of transition metals*, Phys. Rev. Lett. **10**, 159 (1963), doi:10.1103/PhysRevLett.10.159.
- [21] M. C. Gutzwiller, *Correlation of electrons in a narrow s band*, Phys. Rev. **137**, A1726 (1965), doi:10.1103/PhysRev.137.A1726.
- [22] H. Zhai, *Correlated versus ferromagnetic state in repulsively interacting two-component fermi gases*, Phys. Rev. A **80**, 051605 (2009), doi:10.1103/PhysRevA.80.051605.

- [23] X. Cui and T.-L. Ho, *Phase separation in mixtures of repulsive fermi gases driven by mass difference*, Phys. Rev. Lett. **110**, 165302 (2013), doi:10.1103/PhysRevLett.110.165302.
- [24] G.-B. Jo, Y.-R. Lee, J.-H. Choi, C. A. Christensen, T. H. Kim, J. H. Thywissen, D. E. Pritchard and W. Ketterle, *Itinerant Ferromagnetism in a Fermi Gas of Ultracold Atoms*, Science **325**(5947), 1521 (2009).
- [25] C. Sanner, E. J. Su, W. Huang, A. Keshet, J. Gillen and W. Ketterle, *Correlations and pair formation in a repulsively interacting fermi gas*, Phys. Rev. Lett. **108**, 240404 (2012), doi:10.1103/PhysRevLett.108.240404.
- [26] G. Valtolina, F. Scazza, A. Amico, A. Burchianti, A. Recati, T. Enss, M. Inguscio, M. Zaccanti and G. Roati, *Exploring the ferromagnetic behaviour of a repulsive Fermi gas through spin dynamics*, Nat. Phys. **13**(7), 704 (2017).
- [27] A. Amico, F. Scazza, G. Valtolina, P. E. S. Tavares, W. Ketterle, M. Inguscio, G. Roati and M. Zaccanti, *Time-resolved observation of competing attractive and repulsive short-range correlations in strongly interacting fermi gases*, Phys. Rev. Lett. **121**, 253602 (2018), doi:10.1103/PhysRevLett.121.253602.
- [28] G. M. Koutentakis, S. I. Mistakidis and P. Schmelcher, *Interplay of phase separation and itinerant magnetism for correlated few fermions in a double-well*, New J. Phys. **22**(6), 063058 (2020).
- [29] A. Abrikosov and I. Dzialoshinskii, Journal of Experimental and Theoretical Physics **8**, 535 (1958).
- [30] I. Dzyaloshinskii and P. Kondratenko, Journal of Experimental and Theoretical Physics **43**, 1036 (1975).
- [31] K. S. Bedell and K. B. Blagoev, *Quantum spin hydrodynamics and a new spin-current mode in ferromagnetic metals*, Philos. Mag. Lett. **81**(7), 511 (2001).
- [32] Y. Zhang, P. Farinas and K. Bedell, *The "Higgs" Amplitude Mode in Weak Ferromagnetic Metals*, Acta Phys. Pol. A **127**(2), 153 (2015).
- [33] V. P. Mineev, *Transverse spin dynamics in a spin-polarized fermi liquid*, Phys. Rev. B **69**, 144429 (2004), doi:10.1103/PhysRevB.69.144429.
- [34] V. P. Mineev, *Theory of transverse spin dynamics in a polarized fermi liquid and an itinerant ferromagnet*, Phys. Rev. B **72**, 144418 (2005), doi:10.1103/PhysRevB.72.144418.
- [35] V. P. Mineev, *Fermi liquid theory and ferromagnetism*, arXiv (2011), 1111.3208.
- [36] C. Wu and S.-C. Zhang, *Dynamic generation of spin-orbit coupling*, Phys. Rev. Lett. **93**, 036403 (2004), doi:10.1103/PhysRevLett.93.036403.
- [37] C. Wu, K. Sun, E. Fradkin and S.-C. Zhang, *Fermi liquid instabilities in the spin channel*, Phys. Rev. B **75**, 115103 (2007), doi:10.1103/PhysRevB.75.115103.
- [38] N. Manyala, J. F. DiTusa, G. Aeppli and A. P. Ramirez, *Doping a semiconductor to create an unconventional metal*, Nature **454**(7207), 976 (2008).

- [39] S.-S. Zhang, X.-L. Yu, J. Ye and W.-M. Liu, *Collective modes of spin-orbit-coupled fermi gases in the repulsive regime*, Phys. Rev. A **87**, 063623 (2013), doi:10.1103/PhysRevA.87.063623.
- [40] A. Ashrafi, E. I. Rashba and D. L. Maslov, *Theory of a chiral fermi liquid: General formalism*, Phys. Rev. B **88**, 075115 (2013), doi:10.1103/PhysRevB.88.075115.
- [41] X. Montiel and M. Eschrig, *Generation of pure superconducting spin current in magnetic heterostructures via nonlocally induced magnetism due to landau fermi liquid effects*, Phys. Rev. B **98**, 104513 (2018), doi:10.1103/PhysRevB.98.104513.
- [42] J. R. Engelbrecht and K. S. Bedell, *Robustness of a local fermi liquid against ferromagnetism and phase separation*, Phys. Rev. Lett. **74**, 4265 (1995), doi:10.1103/PhysRevLett.74.4265.
- [43] K. B. Blagoev, J. R. Engelbrecht and K. S. Bedell, *S -wave superconductivity in weak ferromagnetic metals*, Philos. Mag. Lett. **78**(2), 169 (1998).
- [44] E. I. Kiselev, M. S. Scheurer, P. Wölfle and J. Schmalian, *Limits on dynamically generated spin-orbit coupling: Absence of $l = 1$ pomeranchuk instabilities in metals*, Phys. Rev. B **95**, 125122 (2017), doi:10.1103/PhysRevB.95.125122.
- [45] P. Wölfle, *Quasiparticles in condensed matter systems*, Rep. Prog. Phys. **81**(3), 032501 (2018).
- [46] Y.-M. Wu, A. Klein and A. V. Chubukov, *Conditions for $l = 1$ pomeranchuk instability in a fermi liquid*, Phys. Rev. B **97**, 165101 (2018), doi:10.1103/PhysRevB.97.165101.
- [47] A. J. Leggett, *Theory of a superfluid fermi liquid. i. general formalism and static properties*, Phys. Rev. **140**, A1869 (1965), doi:10.1103/PhysRev.140.A1869.
- [48] A. V. Chubukov, A. Klein and D. L. Maslov, *Fermi-Liquid Theory and Pomeranchuk Instabilities: Fundamentals and New Developments*, J. Exp. Theor. Phys. **127**(5), 826 (2018).
- [49] J. Fröhlich and U. M. Studer, *Gauge invariance and current algebra in nonrelativistic many-body theory*, Rev. Mod. Phys. **65**(3), 733 (1993).
- [50] P.-Q. Jin, Y.-Q. Li and F.-C. Zhang, *$SU(2) \times U(1)$ unified theory for charge, orbit and spin currents*, J. Phys. A: Math. Gen. **39**(22), 7115 (2006).
- [51] N. Hatano, R. Shirasaki and H. Nakamura, *Non-abelian gauge field theory of the spin-orbit interaction and a perfect spin filter*, Phys. Rev. A **75**, 032107 (2007), doi:10.1103/PhysRevA.75.032107.
- [52] B. Berche, N. Bolivar, A. Lopez and E. Medina, *Gauge field theory approach to spin transport in a 2D electron gas*, Cond. Matt. Phys. **12**, 707 (2009), Invited contribution to Statphys conference, June 2009, Lviv (Ukraine).
- [53] B. Berche, N. Bolívar, A. López and E. Medina, *Gauge transformations of spin-orbit interactions in graphene*, Eur. Phys. J. B **88**(8), 198 (2015).

- [54] W. Kohn, *Theory of the insulating state*, Phys. Rev. **133**, A171 (1964), doi:10.1103/PhysRev.133.A171.
- [55] K. S. Bedell and P. F. Farinas, *Kohn's theorem and correlation functions for a fermi liquid*, Phys. Rev. Lett. **74**, 4285 (1995), doi:10.1103/PhysRevLett.74.4285.
- [56] F. D. M. Haldane, *Berry curvature on the fermi surface: Anomalous hall effect as a topological fermi-liquid property*, Phys. Rev. Lett. **93**, 206602 (2004), doi:10.1103/PhysRevLett.93.206602.
- [57] G. Dresselhaus, *Spin-orbit coupling effects in zinc blende structures*, Phys. Rev. **100**, 580 (1955), doi:10.1103/PhysRev.100.580.
- [58] Y. A. Bychkov and E. I. Rashba, *Oscillatory effects and the magnetic susceptibility of carriers in inversion layers*, J. Phys. C: Solid State Phys. **17**(33), 6039 (1984).
- [59] M. C. Beeler, R. A. Williams, K. Jiménez-García, L. J. LeBlanc, A. R. Perry and I. B. Spielman, *The spin Hall effect in a quantum gas*, Nature **498**(7453), 201 (2013).
- [60] C. J. Kennedy, G. A. Siviloglou, H. Miyake, W. C. Burton and W. Ketterle, *Spin-orbit coupling and quantum spin hall effect for neutral atoms without spin flips*, Phys. Rev. Lett. **111**, 225301 (2013), doi:10.1103/PhysRevLett.111.225301.
- [61] A. Dyrdał, V. K. Dugaev and J. Barnaś, *Spin-resolved orbital magnetization in rashba two-dimensional electron gas*, Phys. Rev. B **94**, 205302 (2016), doi:10.1103/PhysRevB.94.205302.
- [62] O. Fialko, J. Brand and U. Zülicke, *Fragility of the fractional quantum spin Hall effect in quantum gases*, New J. Phys. **16**(2), 025006 (2014).
- [63] T. Fujita and K. F. Quader, *Spin-orbit coupling in fermi-liquid theory*, Phys. Rev. B **36**, 5152 (1987), doi:10.1103/PhysRevB.36.5152.
- [64] A. Kumar and D. L. Maslov, *Effective lattice model for the collective modes in a fermi liquid with spin-orbit coupling*, Phys. Rev. B **95**, 165140 (2017), doi:10.1103/PhysRevB.95.165140.
- [65] A. Shekhter, M. Khodas and A. M. Finkel'stein, *Chiral spin resonance and spin-hall conductivity in the presence of the electron-electron interactions*, Phys. Rev. B **71**, 165329 (2005), doi:10.1103/PhysRevB.71.165329.
- [66] K. S. Bedell and H. P. Dahal, *Spin waves in quasiequilibrium spin systems*, Phys. Rev. Lett. **97**, 047204 (2006), doi:10.1103/PhysRevLett.97.047204.
- [67] D. Pines and P. Nozières, *The Theory of Quantum Liquids*, Westview Press, Boulder, CO, ISBN 0-201-09429-0 (1989).
- [68] Y. Zhang and K. S. Bedell, *Spin orbit magnetism and unconventional superconductivity*, Phys. Rev. B **87**, 115134 (2013), doi:10.1103/PhysRevB.87.115134.
- [69] I. Khalatnikov, *Journal of Experimental and Theoretical Physics* **28**, 1014 (1968).

- [70] Y. Zhang, V. M. Yakovenko and S. Das Sarma, *Dispersion instability in strongly interacting electron liquids*, Phys. Rev. B **71**, 115105 (2005), doi:10.1103/PhysRevB.71.115105.
- [71] S. K. Wong, *Gauge invariance and Galilean invariance*, Nuovo Cim. B **4**(2), 300 (1971).
- [72] M. Wadati, H. Matsumoto and H. Umezawa, *Extended objects created by goldstone bosons*, Phys. Rev. D **18**, 520 (1978), doi:10.1103/PhysRevD.18.520.
- [73] Y. Hosotani, *Dynamics of non-integrable phases and gauge symmetry breaking*, Ann. Phys. **190**(2), 233 (1989).
- [74] H. K. Pal, V. I. Yudson and D. L. Maslov, *Resistivity of non-Galilean-invariant Fermi and non-Fermi liquids*, Lith. J. Phys. **52**(2) (2012).
- [75] K. Mouloupoulos and N. W. Ashcroft, *General theory of the metal-insulator transition*, Phys. Rev. B **45**, 11518 (1992), doi:10.1103/PhysRevB.45.11518.
- [76] B. S. Shastry and B. Sutherland, *Twisted boundary conditions and effective mass in heisenberg-ising and hubbard rings*, Phys. Rev. Lett. **65**, 243 (1990), doi:10.1103/PhysRevLett.65.243.
- [77] D. Poilblanc, *Twisted boundary conditions in cluster calculations of the optical conductivity in two-dimensional lattice models*, Phys. Rev. B **44**, 9562 (1991), doi:10.1103/PhysRevB.44.9562.
- [78] Y. Alavirad and J. D. Sau, *Ferromagnetism and its stability from the one-magnon spectrum in twisted bilayer graphene*, arXiv (2019), 1907.13633.
- [79] N. D. Mermin, *Existence of zero sound in a fermi liquid*, Phys. Rev. **159**, 161 (1967), doi:10.1103/PhysRev.159.161.
- [80] S.-S. Zhang, W.-M. Liu and H. Pu, *Itinerant chiral ferromagnetism in a trapped rashba spin-orbit-coupled fermi gas*, Phys. Rev. A **93**, 043602 (2016), doi:10.1103/PhysRevA.93.043602.
- [81] W. E. Liu, S. Chesi, D. Webb, U. Zülicke, R. Winkler, R. Joynt and D. Culcer, *Generalized stoner criterion and versatile spin ordering in two-dimensional spin-orbit coupled electron systems*, Phys. Rev. B **96**, 235425 (2017), doi:10.1103/PhysRevB.96.235425.
- [82] L. W. Cheuk, A. T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr and M. W. Zwierlein, *Spin-injection spectroscopy of a spin-orbit coupled fermi gas*, Phys. Rev. Lett. **109**, 095302 (2012), doi:10.1103/PhysRevLett.109.095302.
- [83] P. Wang, Z.-Q. Yu, Z. Fu, J. Miao, L. Huang, S. Chai, H. Zhai and J. Zhang, *Spin-orbit coupled degenerate fermi gases*, Phys. Rev. Lett. **109**, 095301 (2012), doi:10.1103/PhysRevLett.109.095301.
- [84] Z. Fu, L. Huang, Z. Meng, P. Wang, L. Zhang, S. Zhang, H. Zhai, P. Zhang and J. Zhang, *Production of Feshbach molecules induced by spin-orbit coupling in Fermi gases*, Nat. Phys. **10**(2), 110 (2013).

- [85] L. Huang, Z. Meng, P. Wang, P. Peng, S.-L. Zhang, L. Chen, D. Li, Q. Zhou and J. Zhang, *Experimental realization of two-dimensional synthetic spin-orbit coupling in ultracold Fermi gases*, Nat. Phys. **12**(6), 540 (2016).
- [86] K. Osterloh, M. Baig, L. Santos, P. Zoller and M. Lewenstein, *Cold atoms in non-abelian gauge potentials: From the hofstadter "moth" to lattice gauge theory*, Phys. Rev. Lett. **95**, 010403 (2005), doi:10.1103/PhysRevLett.95.010403.
- [87] A. M. Dudarev, R. B. Diener, I. Carusotto and Q. Niu, *Spin-orbit coupling and berry phase with ultracold atoms in 2d optical lattices*, Phys. Rev. Lett. **92**, 153005 (2004), doi:10.1103/PhysRevLett.92.153005.
- [88] N. Goldman, A. Kubasiak, P. Gaspard and M. Lewenstein, *Ultracold atomic gases in non-abelian gauge potentials: The case of constant wilson loop*, Phys. Rev. A **79**, 023624 (2009), doi:10.1103/PhysRevA.79.023624.
- [89] D. L. Campbell, G. Juzeliūnas and I. B. Spielman, *Realistic rashba and dres-selhaus spin-orbit coupling for neutral atoms*, Phys. Rev. A **84**, 025602 (2011), doi:10.1103/PhysRevA.84.025602.
- [90] M. Aidelsburger, M. Atala, M. Lohse, J. T. Barreiro, B. Paredes and I. Bloch, *Realization of the hofstadter hamiltonian with ultracold atoms in optical lattices*, Phys. Rev. Lett. **111**, 185301 (2013), doi:10.1103/PhysRevLett.111.185301.
- [91] N. Goldman, A. Kubasiak, A. Bermudez, P. Gaspard, M. Lewenstein and M. A. Martin-Delgado, *Non-abelian optical lattices: Anomalous quantum hall effect and dirac fermions*, Phys. Rev. Lett. **103**, 035301 (2009), doi:10.1103/PhysRevLett.103.035301.
- [92] I. I. Satija, D. C. Dakin, J. Y. Vaishnav and C. W. Clark, *Physics of a two-dimensional electron gas with cold atoms in non-abelian gauge potentials*, Phys. Rev. A **77**, 043410 (2008), doi:10.1103/PhysRevA.77.043410.

Supplemental Material:

Explicit calculation of the Pomeranchuk instability condition at finite gauge

In this section, we will briefly go over the calculation of Eqns. (4) and (5). We begin with Eqn. (4), where we explicitly write down the first term in the Landau free energy:

$$\begin{aligned}
& \sum_{p\sigma} (\epsilon_{p-A_\sigma, \sigma} - \mu) \delta n_{p-A_\sigma, \sigma} \\
& \approx v_F \sum_{p\sigma} (p - p_F - A_\sigma) \left\{ -\delta p_F \delta(p_F - p) - A_\sigma \delta(p_F - p) - \frac{1}{2} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p) - A_\sigma \delta p_F \frac{\partial}{\partial p} \delta(p_F - p) \right. \\
& \quad \left. - \frac{1}{2} A_\sigma^2 \frac{\partial}{\partial p} \delta(p_F - p) \right\} \\
& \approx -v_F \sum_{p\sigma} \left\{ (p - p_F) \delta p_F \delta(p_F - p) + \frac{1}{2} \delta p_F^2 (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \right. \\
& \quad + A_\sigma \left[-\delta p_F \delta(p_F - p) + (p - p_F) \delta(p_F - p) - \frac{1}{2} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p) + \delta p_F (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \right] \\
& \quad \left. + A_\sigma^2 \left[-\delta(p_F - p) - \delta p_F \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{2} (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \right] \right\} \quad (12)
\end{aligned}$$

where we ignore terms cubic in A_σ and higher. The first terms are just the gauge-independent terms:

(1)

$$-v_F \sum_{p\sigma} (p - p_F) \delta p_F \delta(p_F - p) \rightarrow -\frac{v_F}{(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \delta p_F \int dp p^2 (p - p_F) \delta(p - p_F) = \boxed{0} \quad (13)$$

(2)

$$\begin{aligned}
& -\frac{v_F}{2} \sum_{p\sigma} \delta p_F^2 (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \rightarrow \\
& -\frac{v_F}{2(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \delta p_F^2 \int dp p^2 (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \\
& = -\frac{v_F}{2(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \delta p_F^2 \left\{ p^2 (p - p_F) \delta(p_F - p) \Big|_0^{\infty} - \int dp (3p^2 - 2pp_F) \delta(p_F - p) \right\} \\
& = \frac{v_F p_F^2}{2(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \delta p_F^2 \\
& = \frac{N(0)}{4} \sum_{\sigma} \sum_{\ell_1, \ell_2} \nu_{\ell_1\sigma} \nu_{\ell_2\sigma} \int_{-1}^1 \frac{d(\cos\theta)}{2} P_{\ell_1}(\cos\theta) P_{\ell_2}(\cos\theta) \\
& = \boxed{\frac{N(0)}{4} \sum_{\ell\sigma} \frac{\nu_{\ell\sigma}^2}{2\ell+1}} \tag{14}
\end{aligned}$$

These are the terms that remain when we take $A_{\sigma} = 0$. Now we will consider those that are linear in A_{σ} :

$$-v_F \sum_{p\sigma} A_{\sigma} \left[-\delta p_F \delta(p_F - p) + (p - p_F) \delta(p_F - p) - \frac{1}{2} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p) + \delta p_F (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \right] \tag{15}$$

(1)

$$v_F \sum_{p\sigma} \delta p_F \delta(p_F - p) \rightarrow \frac{v_F}{(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \delta p_F \int dp p^2 \delta(p_F - p) = \boxed{\frac{N(0)v_F}{2} \sum_{\sigma} \frac{\nu_{\ell\sigma}}{2\ell+1} \delta_{\ell 0}} \tag{16}$$

(2)

$$-v_F \sum_{p\sigma} (p - p_F) \delta(p_F - p) \rightarrow -\frac{v_F}{(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \int dp p^2 (p - p_F) \delta(p_F - p) = \boxed{0} \tag{17}$$

(3)

$$\frac{v_F}{2} \sum_{p\sigma} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p) \rightarrow \frac{v_F}{2(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \delta p_F^2 \int dp p^2 \frac{\partial}{\partial p} \delta(p_F - p) = \boxed{-\frac{N(0)}{2p_F} \sum_{\ell\sigma} \frac{\nu_{\ell\sigma}^2}{2\ell+1}} \tag{18}$$

(4)

$$-v_F \sum_{p\sigma} \delta p_F (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) = \boxed{\frac{N(0)v_F}{2} \sum_{\ell\sigma} \frac{\nu_{\ell\sigma}}{2\ell+1} \delta_{\ell 0}} \tag{19}$$

The term that is linear in A_σ then becomes

$$\begin{aligned}
& \frac{N(0)}{4} \sum_{\ell\sigma} \frac{A_\sigma}{2\ell+1} \{4v_F\nu_{\ell\sigma}\delta_{\ell 0} - 2\nu_{\ell\sigma}^2/p_F\} \\
&= \frac{N(0)}{4} \sum_{\ell} \left[\frac{A_\uparrow}{2\ell+1} (4v_F\nu_{\ell\uparrow} - 2\nu_{\ell\uparrow}^2/p_F) + \frac{A_\downarrow}{2\ell+1} (4v_F\nu_{\ell\downarrow} - 2\nu_{\ell\downarrow}^2/p_F) \right] \\
&= \begin{cases} \left[\frac{N(0)}{4} \sum_{\ell} \left[\frac{A}{2\ell+1} \left(4v_F\nu_{\ell s} - \frac{1}{p_F} (\nu_{\ell s}^2 + \nu_{\ell a}^2) \right) \right] \right], & A_\uparrow = A_\downarrow \\ \left[\frac{N(0)}{4} \sum_{\ell} \left[\frac{A}{2\ell+1} \left(4v_F\nu_{\ell a} - \frac{2}{p_F} \nu_{\ell s}\nu_{\ell a} \right) \right] \right], & A_\uparrow = -A_\downarrow \end{cases} \quad (20)
\end{aligned}$$

Where we have used the fact that

$$\nu_{\ell\uparrow}^2 + \nu_{\ell\downarrow}^2 = \frac{1}{2} (\nu_{\ell s}^2 + \nu_{\ell a}^2) \quad (21)$$

$$\nu_{\ell\uparrow}^2 - \nu_{\ell\downarrow}^2 = (\nu_{\ell\uparrow} + \nu_{\ell\downarrow})(\nu_{\ell\uparrow} - \nu_{\ell\downarrow}) = \nu_{\ell s}\nu_{\ell a} \quad (22)$$

We will now deal with quadratic terms:

$$-v_F \sum_{p\sigma} A_\sigma^2 \left[-\delta(p_F - p) - \delta p_F \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{2}(p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \right] \quad (23)$$

(1)

$$v_F \sum_{p\sigma} \delta(p_F - p) \rightarrow \frac{v_F}{(2\pi\hbar)^3} \sum_{p\sigma} \int d\Omega \int dp p^2 \delta(p_F - p) = \boxed{\sum_{\sigma} \frac{N(0)v_F^2}{2}} \quad (24)$$

(2)

$$v_F \sum_{p\sigma} \delta p_F \frac{\partial}{\partial p} \delta(p_F - p) \rightarrow \frac{v_F}{(2\pi\hbar)^3} \sum_{p\sigma} \int d\Omega \delta p_F \int dp p^2 \frac{\partial}{\partial p} \delta(p_F - p) = \boxed{-\frac{N(0)v_F}{p_F} \sum_{\ell\sigma} \frac{\nu_{\ell\sigma}}{2\ell+1} \delta_{\ell 0}} \quad (25)$$

(3)

$$-\frac{v_F}{2} \sum_{p\sigma} (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) \rightarrow -\frac{v_F}{2(2\pi\hbar)^3} \sum_{\sigma} \int d\Omega \int dp p^2 (p - p_F) \frac{\partial}{\partial p} \delta(p_F - p) = \boxed{\sum_{\sigma} \frac{N(0)v_F^2}{4}} \quad (26)$$

The quadratic terms are therefore given below:

$$\frac{N(0)}{4} \sum_{\sigma} A_{\sigma}^2 \left\{ 3v_F^2 - \sum_{\ell} \frac{4v_F \nu_{\ell\sigma}/p_F}{2\ell+1} \delta_{\ell 0} \right\} \quad (27)$$

Note that the above can be simplified as follows:

$$\frac{N(0)}{4} \sum_{\sigma} A_{\sigma}^2 \left\{ 3v_F^2 - \sum_{\ell} \frac{4v_F \nu_{\ell\sigma}/p_F}{2\ell+1} \delta_{\ell 0} \right\} = \frac{N(0)}{4} A^2 \left[6v_F^2 - \sum_{\ell} \frac{4v_F}{p_F(2\ell+1)} \nu_{\ell s} \delta_{\ell 0} \right] \quad (28)$$

Because we only care about terms up to quadratic in A , the above becomes

$$\begin{aligned} & \sum_{p\sigma} (\epsilon_{p-A\sigma, \sigma} - \mu) \delta n_{p-A\sigma, \sigma} \\ & \rightarrow \begin{cases} \frac{N(0)}{4} \left[\sum_{\ell} \frac{1}{2\ell+1} \left\{ \frac{1}{2}(\nu_{\ell s}^2 + \nu_{\ell a}^2) + A \left\{ 4v_F \nu_{\ell s} - \frac{1}{p_F} (\nu_{\ell s}^2 + \nu_{\ell a}^2) \right\} - \frac{4A^2}{m^*} \nu_{\ell s} \delta_{\ell 0} \right\} + 6v_F^2 A^2 \right], & A_{\uparrow} = A_{\downarrow} \\ \frac{N(0)}{4} \left[\sum_{\ell} \frac{1}{2\ell+1} \left\{ \frac{1}{2}(\nu_{\ell s}^2 + \nu_{\ell a}^2) + 2A \nu_{\ell a} \left\{ 2v_F - \frac{\nu_{\ell s}}{p_F} \right\} - \frac{4A^2}{m^*} \nu_{\ell s} \delta_{\ell 0} \right\} + 6v_F^2 A^2 \right], & A_{\uparrow} = -A_{\downarrow} \end{cases} \\ & = \begin{cases} \frac{N(0)}{4} \left[\sum_{\ell} \frac{1}{2\ell+1} \left\{ \frac{1}{2}(\nu_{\ell s}^2 + \nu_{\ell a}^2) + \frac{A}{p_F} \{ 8\epsilon_F \nu_{\ell s} - (\nu_{\ell s}^2 + \nu_{\ell a}^2) \} - 8 \frac{A^2 \epsilon_F}{p_F^2} \nu_{\ell s} \delta_{\ell 0} \right\} + \frac{24\epsilon_F^2 A^2}{p_F^2} \right], & A_{\uparrow} = A_{\downarrow} \\ \frac{N(0)}{4} \left[\sum_{\ell} \frac{1}{2\ell+1} \left\{ \frac{1}{2}(\nu_{\ell s}^2 + \nu_{\ell a}^2) + 2 \frac{A}{p_F} \nu_{\ell a} \{ 4\epsilon_F - \nu_{\ell s} \} - 8 \frac{A^2 \epsilon_F}{p_F^2} \nu_{\ell s} \delta_{\ell 0} \right\} + \frac{24\epsilon_F^2 A^2}{p_F^2} \right], & A_{\uparrow} = -A_{\downarrow} \end{cases} \\ & \approx \boxed{\begin{cases} \frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left\{ (\nu_{\ell s}^2 + \nu_{\ell a}^2) + \frac{16A}{p_F} \epsilon_F \nu_{\ell s} \right\} + \frac{48A^2}{p_F^2} \epsilon_F^2 \right], & A_{\uparrow} = A_{\downarrow} \\ \frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left\{ (\nu_{\ell s}^2 + \nu_{\ell a}^2) + \frac{16A}{p_F} \epsilon_F \nu_{\ell a} \right\} + \frac{48A^2}{p_F^2} \epsilon_F^2 \right], & A_{\uparrow} = -A_{\downarrow} \end{cases}} \end{aligned}$$

where we have used the fact that $\epsilon_F \gg \nu_s$ and $\epsilon_F \gg \nu_a$. This completes the derivation of Eqn. (4).

We will now consider the gauge-dependence of the quadratic term in the Landau-Fermi free energy; i.e., the derivation of Eqn. (5):

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta n_{p-A_\sigma, \sigma} \delta n_{p'-A_{\sigma'}, \sigma'} \\
&= \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \\
&\times \left\{ -\delta p_F \delta(p_F - p) - A_\sigma \delta(p_F - p) - \frac{1}{2} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p) - A_\sigma \delta p_F \frac{\partial}{\partial p} \delta(p_F - p) - \frac{1}{2} A_\sigma^2 \frac{\partial}{\partial p} \delta(p_F - p) \right\} \\
&\times \left\{ -\delta p'_F \delta(p_F - p') - A_{\sigma'} \delta(p_F - p') - \frac{1}{2} \delta p_F'^2 \frac{\partial}{\partial p'} \delta(p_F - p') - A_{\sigma'} \delta p'_F \frac{\partial}{\partial p'} \delta(p_F - p') - \frac{1}{2} A_{\sigma'}^2 \frac{\partial}{\partial p'} \delta(p_F - p') \right\} \\
&\approx \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \left\{ \delta p_F \delta p'_F \delta(p_F - p) \delta(p_F - p') \right. \\
&\quad + A_\sigma \left[\delta p'_F \delta(p_F - p') \delta(p_F - p) + \delta p_F \delta p'_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{2} \delta p_F'^2 \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right] \\
&\quad + A_{\sigma'} \left[\delta p_F \delta(p_F - p) \delta(p_F - p') + \delta p'_F \delta p_F \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') + \frac{1}{2} \delta p_F^2 \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \right] \\
&\quad + A_\sigma^2 \left[\frac{1}{2} \delta p'_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{4} \delta p_F'^2 \frac{\partial}{\partial p'} \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \right] \\
&\quad + A_{\sigma'}^2 \left[\frac{1}{2} \delta p_F \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') + \frac{1}{4} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right] \\
&\quad + A_\sigma \cdot A_{\sigma'} \left[\delta(p_F - p) \delta(p_F - p') + \delta p'_F \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right. \\
&\quad \left. + \delta p_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \delta p_F \delta p'_F \frac{\partial}{\partial p} \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right] \tag{29}
\end{aligned}$$

where we only keep terms up to quadratic dependence in A_σ . To begin, we look at the gauge-independent term:

(1)

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta p_F \delta p'_F \delta(p_F - p) \delta(p_F - p') \\
& \rightarrow \sum_{\sigma\sigma'} \frac{1}{(2\pi\hbar)^6} \int d\Omega \int d\Omega' \int dp p^2 \int dp' p'^2 f_{pp'\sigma\sigma'} \delta(p_F - p) \delta(p_F - p') \delta p_F \delta p'_F \\
& = \frac{p_F^4}{(2\pi\hbar)^6} \sum_{\sigma\sigma'} \int d\Omega \delta p_F \int d\Omega' \delta p'_F f_{pp'\sigma\sigma'} \\
& = \frac{16\pi^2 p_F^4}{(2\pi\hbar)^6 v_F^2} \sum_{\ell\ell'\ell''\sigma\sigma'} \nu_{\ell\sigma} \nu_{\ell'\sigma'} \int_{-1}^1 \frac{d(\cos\theta)}{2} \int_{-1}^1 \frac{d(\cos\theta')}{2} f_{\ell''\sigma'\sigma''} P_\ell(\cos\theta) P_{\ell'}(\cos\theta') P_{\ell''}(\cos\theta') P_{\ell''}(\cos\theta) \\
& = \frac{N(0)}{4} \sum_{\ell\sigma\sigma'} \nu_{\ell\sigma} \nu_{\ell\sigma'} \frac{1}{2\ell+1} \frac{F_{\ell\sigma\sigma'}}{2\ell+1} \\
& = \frac{N(0)}{4} \sum_{\ell} \frac{1}{2\ell+1} \frac{1}{2\ell+1} \{ \nu_{\ell\uparrow}^2 F_{\ell\uparrow\uparrow} + 2\nu_{\ell\uparrow}\nu_{\ell\downarrow} F_{\ell\uparrow\downarrow} + \nu_{\ell\downarrow}^2 F_{\ell\downarrow\downarrow} \} \\
& = \boxed{\frac{N(0)}{4} \sum_{\ell} \frac{1}{2\ell+1} \frac{1}{2\ell+1} \{ F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2 \}} \tag{30}
\end{aligned}$$

We will now calculate the terms linear in A_σ :

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \\
& \times \left\{ A_\sigma \left[\delta p'_F \delta(p_F - p') \delta(p_F - p) + \delta p_F \delta p'_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{2} \delta p_F'^2 \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right] \right. \\
& \quad \left. + A_{\sigma'} \left[\delta p_F \delta(p_F - p) \delta(p_F - p') + \delta p'_F \delta p_F \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') + \frac{1}{2} \delta p_F^2 \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \right] \right\} \\
& = 2 \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \\
& \times \left\{ A_\sigma \left[\delta p'_F \delta(p_F - p') \delta(p_F - p) + \delta p_F \delta p'_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{2} \delta p_F'^2 \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right] \right\} \tag{31}
\end{aligned}$$

(1)

$$\sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta p'_F \delta(p_F - p') \delta(p_F - p) = \boxed{\frac{N(0)v_F}{4} \sum_{\ell'\sigma\sigma'} \nu_{\ell'\sigma'} \frac{1}{2\ell'+1} \frac{F_{\ell'\sigma\sigma'}}{2\ell'+1} \delta_{\ell'0}} \tag{32}$$

(2)

$$\sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta p_F \delta p'_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) = \boxed{-\frac{N(0)}{2p_F} \sum_{\ell\sigma\sigma'} \nu_{\ell\sigma} \nu_{\ell\sigma'} \frac{1}{2\ell+1} \frac{F_{\ell\sigma\sigma'}}{2\ell+1}} \tag{33}$$

(3)

$$\begin{aligned}
& \frac{1}{2} \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta p_F'^2 \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \\
& \rightarrow \frac{1}{2(2\pi\hbar)^6} \sum_{\sigma\sigma'} \int d\Omega \int d\Omega' \int dp p^2 \int dp' p'^2 f_{pp'\sigma\sigma'} \delta p_F'^2 \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \\
& = -\frac{16\pi^2 p_F^3}{(2\pi\hbar)^6 v_F^2} \sum_{\ell\ell'\ell''\sigma\sigma'} \nu_{\ell\sigma'} \nu_{\ell'\sigma'} f_{\ell''\sigma\sigma'} \int_{-1}^1 \frac{d(\cos\theta)}{2} P_{\ell''}(\cos\theta) \int_{-1}^1 \frac{d(\cos\theta')}{2} P_{\ell}(\cos\theta') P_{\ell'}(\cos\theta') P_{\ell''}(\cos\theta') \\
& = -\frac{p_F^3}{4\pi^2 \hbar^6 v_F^2} \sum_{\ell\ell'\ell''\sigma\sigma'} \nu_{\ell\sigma'} \nu_{\ell'\sigma'} f_{\ell''\sigma\sigma'} \frac{1}{2\ell''+1} \delta_{\ell''0} \begin{pmatrix} \ell & \ell' & \ell'' \\ 0 & 0 & 0 \end{pmatrix}^2 \\
& = -\frac{N(0)}{4p_F} \sum_{\ell\sigma\sigma'} \nu_{\ell\sigma}^2 \frac{F_{0\sigma\sigma'}}{2\ell+1}
\end{aligned} \tag{34}$$

Where we have used the fact that

$$\begin{pmatrix} \ell & \ell' & 0 \\ 0 & 0 & 0 \end{pmatrix} = (-1)^{\frac{\ell+\ell'}{2}} \sqrt{\frac{(\ell'-\ell)!(\ell-\ell')!}{\ell+\ell'+1}} \frac{1}{\left(\frac{\ell'-\ell}{2}\right)! \left(\frac{\ell-\ell'}{2}\right)!} = -\frac{1}{\sqrt{2\ell+1}} \delta_{\ell\ell'} \tag{35}$$

We can now write the linear term in A_σ :

$$\begin{aligned}
& 2\frac{N(0)}{4} \sum_{\ell\sigma\sigma'} A_\sigma \left[\nu_{\ell\sigma} \frac{v_F}{2\ell+1} \frac{F_{\ell\sigma\sigma'}}{2\ell+1} \delta_{\ell 0} - \frac{2}{p_F} \nu_{\ell\sigma} \nu_{\ell'\sigma'} \frac{1}{2\ell+1} \frac{F_{\ell\sigma\sigma'}}{2\ell+1} - \frac{1}{p_F} \nu_{\ell\sigma}^2 \frac{F_{0\sigma\sigma'}}{2\ell+1} \right] \\
& = \frac{N(0)}{2} \sum_{\ell\sigma\sigma'} A_\sigma \frac{1}{2\ell+1} \left[\nu_{\ell\sigma} v_F \frac{F_{\ell\sigma\sigma'}}{2\ell+1} \delta_{\ell 0} - \frac{1}{p_F} \left\{ 2\nu_{\ell\sigma} \nu_{\ell'\sigma'} \frac{F_{\ell\sigma\sigma'}}{2\ell+1} + \nu_{\ell\sigma}^2 F_{0\sigma\sigma'} \right\} \right] \\
& = \frac{N(0)}{4} \sum_{\ell} \frac{1}{2\ell+1} \frac{1}{2\ell+1} \left[\sum_{\sigma\sigma'} 2A_\sigma \nu_{\ell\sigma} v_F F_{\ell\sigma\sigma'} \delta_{\ell 0} - \sum_{\sigma\sigma'} A_\sigma \frac{2}{p_F} \left\{ 2\nu_{\ell\sigma} \nu_{\ell'\sigma'} F_{\ell\sigma\sigma'} + \nu_{\ell\sigma}^2 F_{0\sigma\sigma'} (2\ell+1) \right\} \right]
\end{aligned} \tag{36}$$

Simplifying the above cases piece by piece, we find the following:

(1)

$$\begin{aligned}
\sum_{\sigma\sigma'} A_\sigma 2\nu_{\ell\sigma} v_F F_{\ell\sigma\sigma'} \delta_{\ell 0} & = 2A_\uparrow \nu_{\ell\uparrow} v_F \delta_{\ell 0} (F_{\ell\uparrow\uparrow} + F_{\ell\uparrow\downarrow}) + 2A_\downarrow \nu_{\ell\downarrow} v_F \delta_{\ell 0} (F_{\ell\downarrow\uparrow} + F_{\ell\downarrow\downarrow}) \\
& = \begin{cases} 4v_F A F_\ell^s \nu_{\ell s} \delta_{\ell 0}, & A_\uparrow = A_\downarrow \\ 4v_F A F_\ell^s \nu_{\ell a} \delta_{\ell 0}, & A_\uparrow = -A_\downarrow \end{cases}
\end{aligned}$$

(2)

$$\begin{aligned}
& \sum_{\sigma\sigma'} A_\sigma \frac{2}{p_F} \left\{ 2\nu_{\ell\sigma}\nu_{\ell\sigma'} F_{\ell\sigma\sigma'} + \nu_{\ell\sigma}^2 F_{0\sigma\sigma'}(2\ell+1) \right\} \\
&= \begin{cases} \frac{4A}{p_F} \left\{ \nu_{\ell s}^2 [F_\ell^s + \frac{1}{2}(2\ell+1)F_0^s] + \nu_{\ell a}^2 [F_\ell^a + \frac{1}{2}(2\ell+1)F_0^s] \right\}, & A_\uparrow = A_\downarrow \\ \frac{4A}{p_F} \nu_{\ell s}\nu_{\ell a} \left\{ F_\ell^s + F_\ell^a + (2\ell+1)F_0^s \right\}, & A_\uparrow = -A_\downarrow \end{cases}
\end{aligned}$$

The term linear in A_σ can then be written as the following:

$$\begin{aligned}
& \frac{N(0)}{4} \sum_\ell \frac{1}{2\ell+1} \frac{1}{2\ell+1} \left[\sum_{\sigma\sigma'} 2A_\sigma \nu_{\ell\sigma} \nu_F F_{\ell\sigma\sigma'} \delta_{\ell 0} - \sum_{\sigma\sigma'} A_\sigma \frac{2}{p_F} \left\{ 2\nu_{\ell\sigma}\nu_{\ell\sigma'} F_{\ell\sigma\sigma'} + \nu_{\ell\sigma}^2 F_{0\sigma\sigma'}(2\ell+1) \right\} \right] \\
&= \frac{N(0)}{4} \sum_\ell \frac{1}{2\ell+1} \frac{1}{2\ell+1} \\
& \quad \times 4A \begin{cases} \nu_F F_\ell^s \nu_{\ell s} \delta_{\ell 0} - \frac{1}{p_F} \left\{ \nu_{\ell s}^2 [F_\ell^s + \frac{1}{2}(2\ell+1)F_0^s] + \nu_{\ell a}^2 [F_\ell^a + \frac{1}{2}(2\ell+1)F_0^s] \right\}, & A_\uparrow = A_\downarrow \\ \nu_F F_\ell^s \nu_{\ell a} \delta_{\ell 0} - \frac{1}{p_F} \nu_{\ell s} \nu_{\ell a} \left\{ F_\ell^s + F_\ell^a + (2\ell+1)F_0^s \right\}, & A_\uparrow = -A_\downarrow \end{cases}
\end{aligned} \tag{37}$$

We will now tackle the terms that go as A_σ^2 :

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \left\{ A_\sigma^2 \left[\frac{1}{2} \delta p_F' \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{4} \delta p_F'^2 \frac{\partial}{\partial p'} \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \right] \right. \\
& \quad \left. + A_{\sigma'}^2 \left[\frac{1}{2} \delta p_F \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') + \frac{1}{4} \delta p_F^2 \frac{\partial}{\partial p} \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right] \right\} \\
&= \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \left\{ A_\sigma^2 \left[\delta p_F' \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \frac{1}{2} \delta p_F'^2 \frac{\partial}{\partial p'} \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \right] \right\} \tag{38}
\end{aligned}$$

Each term is calculated one-by-one:

(1)

$$\sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta p_F' \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) = \boxed{-\frac{N(0)v_F}{2p_F} \sum_{\ell\ell'\sigma\sigma'} \nu_{\ell\sigma} \frac{1}{2\ell+1} \frac{F_{\ell\sigma\sigma'}}{2\ell+1} \delta_{\ell 0}} \tag{39}$$

(2)

$$\begin{aligned}
& \frac{1}{2} \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta p_F'^2 \frac{\partial}{\partial p'} \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \\
& \rightarrow \frac{1}{2(2\pi\hbar)^6} \int d\Omega \int d\Omega' \int dp p^2 \int dp' p'^2 f_{pp'\sigma\sigma'} \delta p_F'^2 \frac{\partial}{\partial p'} \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \\
& = \frac{4p_F^2}{2(2\pi\hbar)^6} \int d\Omega \int d\Omega' \delta p_F'^2 f_{pp'\sigma\sigma'} \\
& = \frac{64\pi^2 p_F^2}{2(2\pi\hbar)^6 v_F^2} \sum_{\ell\ell'\ell''\sigma\sigma'} \nu_{\ell'\sigma'} \nu_{\ell'\sigma'} f_{\ell''\sigma\sigma'} \int_{-1}^1 \frac{d(\cos\theta)}{2} \int_{-1}^1 \frac{d(\cos\theta')}{2} P_\ell(\cos\theta') P_{\ell'}(\cos\theta') P_{\ell''}(\cos\theta) P_{\ell''}(\cos\theta') \\
& = \frac{p_F^2}{2\pi^4 \hbar^6 v_F^2} \sum_{\ell\ell'\ell''\sigma\sigma'} \nu_{\ell'\sigma'} \nu_{\ell'\sigma'} f_{\ell''\sigma\sigma'} \frac{1}{2\ell''+1} \delta_{\ell''0} \begin{pmatrix} \ell & \ell' & \ell'' \\ 0 & 0 & 0 \end{pmatrix}^2 \\
& = \boxed{\frac{N(0)}{2p_F^2} \sum_{\ell\sigma\sigma'} \nu_{\ell\sigma}^2 \frac{F_{0\sigma\sigma'}}{2\ell+1}} \tag{40}
\end{aligned}$$

Thus, the above result in the following term in A_σ^2 :

$$\begin{aligned}
& 2 \sum_{\ell\sigma\sigma'} A_\sigma^2 \left\{ -\frac{N(0)}{2p_F} \nu_{\ell\sigma} \frac{1}{2\ell+1} \frac{F_{\ell\sigma\sigma'}}{2\ell+1} \delta_{\ell 0} + \frac{N(0)}{2p_F^2} \nu_{\ell\sigma}^2 \frac{F_{0\sigma\sigma'}}{2\ell+1} \right\} \\
& = \frac{N(0)}{4} \sum_{\ell} \frac{1}{2\ell+1} \frac{1}{2\ell+1} 4 \sum_{\sigma\sigma'} A_\sigma^2 \left\{ -\frac{\nu_{\ell\sigma}}{m^*} F_{\ell\sigma\sigma'} \delta_{\ell 0} + \frac{\nu_{\ell\sigma}^2}{p_F^2} F_{0\sigma\sigma'} (2\ell+1) \right\} \tag{41}
\end{aligned}$$

We look at each term separately:

$$\begin{aligned}
-\sum_{\sigma\sigma'} A_\sigma^2 \frac{\nu_{\ell\sigma}}{m^*} F_{\ell\sigma\sigma'} \delta_{\ell 0} & = -\frac{\delta_{\ell 0}}{m^*} \{ A_{\uparrow\uparrow}^2 \nu_{\uparrow} F_{\ell\uparrow\uparrow} + A_{\uparrow\downarrow}^2 \nu_{\uparrow} F_{\ell\uparrow\downarrow} + A_{\downarrow\downarrow}^2 \nu_{\downarrow} F_{\ell\downarrow\uparrow} + A_{\downarrow\uparrow}^2 \nu_{\downarrow} F_{\ell\downarrow\downarrow} \} \\
& = -\frac{2}{m^*} A^2 \nu_{\ell s} F_\ell^s \delta_{\ell 0} \tag{42}
\end{aligned}$$

$$\begin{aligned}
\frac{2\ell+1}{p_F^2} \sum_{\sigma\sigma'} A_\sigma^2 \nu_{\ell\sigma}^2 F_{0\sigma\sigma'} & = \frac{2\ell+1}{p_F^2} \{ A_{\uparrow\uparrow}^2 \nu_{\uparrow}^2 (F_{0\uparrow\uparrow} + F_{0\uparrow\downarrow}) + A_{\downarrow\downarrow}^2 \nu_{\downarrow}^2 (F_{0\downarrow\uparrow} + F_{0\downarrow\downarrow}) \} \\
& = (2\ell+1) \frac{A^2}{p_F^2} F_0^s (\nu_{\ell s}^2 + \nu_{\ell a}^2) \tag{43}
\end{aligned}$$

Hence, we find that the term that goes as A_σ^2 goes as

$$\begin{aligned}
& \frac{N(0)}{4} \sum_{\ell} \frac{1}{2\ell+1} \frac{1}{2\ell+1} 4 \sum_{\sigma\sigma'} A_\sigma^2 \left\{ -\frac{\nu_{\ell\sigma}}{m^*} F_{\ell\sigma\sigma'} \delta_{\ell 0} + \frac{\nu_{\ell\sigma}^2}{p_F^2} F_{0\sigma\sigma'} (2\ell+1) \right\} \\
& = \boxed{\frac{N(0)}{4} \sum_{\ell} \frac{1}{2\ell+1} \frac{1}{2\ell+1} \left[4A^2 \left\{ -\frac{2}{m^*} \nu_{\ell s} F_\ell^s \delta_{\ell 0} + (2\ell+1) \frac{1}{p_F^2} F_0^s (\nu_{\ell s}^2 + \nu_{\ell a}^2) \right\} \right]} \tag{44}
\end{aligned}$$

We will now deal with the terms that go as $A_\sigma \cdot A_{\sigma'}$:

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} A_\sigma \cdot A_{\sigma'} \left[\delta(p_F - p)\delta(p_F - p') + \delta p'_F \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right. \\
& \quad \left. + \delta p_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \delta p_F \delta p'_F \frac{\partial}{\partial p} \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right] \\
&= \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} A_\sigma \cdot A_{\sigma'} \\
& \quad \times \left[\delta(p_F - p)\delta(p_F - p') + 2\delta p_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) + \delta p_F \delta p'_F \frac{\partial}{\partial p} \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \right]
\end{aligned} \tag{45}$$

We go through each term separately, being careful to include a $\cos \chi$ term due to the dot product: (1)

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \cos \theta \cos \theta' \delta(p_F - p)\delta(p_F - p') \\
&= \frac{p_F^4}{4\pi^4 \hbar^6} \sum_{\ell\sigma\sigma'} f_{\ell\sigma\sigma'} \int_{-1}^1 \frac{d(\cos \theta)}{2} \cos \theta P_\ell(\cos \theta) \int_{-1}^1 \frac{d(\cos \theta')}{2} \cos \theta' P_\ell(\cos \theta') \\
&= \boxed{\frac{N(0)}{4} \sum_{\sigma\sigma'} \frac{F_{1\sigma\sigma'}}{9} v_F^2}
\end{aligned} \tag{46}$$

where we have used the fact that $\cos \chi = \cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi')$ and the integral of ϕ and ϕ' from 0 to 2π of $\cos(\phi - \phi')$ is zero.

(2)

$$\begin{aligned}
& 2 \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \cos \theta \cos \theta' \delta p_F \delta(p_F - p') \frac{\partial}{\partial p} \delta(p_F - p) \\
&= -\frac{64\pi^2 p_F^3}{(2\pi \hbar)^6 v_F} \sum_{\ell\ell'\sigma\sigma'} \nu_{\ell\sigma} f_{\ell'\sigma\sigma'} \int_{-1}^1 \frac{d(\cos \theta)}{2} P_\ell(\cos \theta) P_{\ell'}(\cos \theta) \cos \theta \int_{-1}^1 \frac{d(\cos \theta')}{2} P_{\ell'}(\cos \theta') \cos \theta' \\
&= -\frac{N(0)v_F}{p_F} \sum_{\ell\sigma\sigma'} \nu_{\ell\sigma} \frac{F_{1\sigma\sigma'}}{3} \begin{pmatrix} \ell & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}^2 \\
&= -\frac{N(0)v_F}{p_F} \sum_{\ell\sigma\sigma'} \nu_{\ell\sigma} \frac{F_{1\sigma\sigma'}}{3} \left\{ \frac{1}{3} \delta_{\ell 0} + \frac{2}{15} \delta_{\ell 2} \right\} \\
&\approx \boxed{-\frac{N(0)}{4} \sum_{\ell\sigma\sigma'} \frac{4}{9} F_{1\sigma\sigma'} \frac{\nu_{\ell\sigma}}{m^*} \delta_{\ell 0}}
\end{aligned} \tag{47}$$

where we have used the fact that

$$\begin{pmatrix} \ell & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = (-1)^{\frac{\ell}{2}+1} \ell! \sqrt{\frac{(2-\ell)!}{(3+\ell)!}} \frac{\frac{\ell}{2}+1}{(1-\frac{\ell}{2})! (\frac{\ell}{2})!} = -\frac{1}{\sqrt{3}} \delta_{\ell 0} + \sqrt{\frac{2}{15}} \delta_{\ell 2} \tag{48}$$

and we have assumed that we can safely ignore all terms dependent on $\ell > 1$.

(3)

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \cos \theta \cos \theta' \delta p_F \delta p'_F \frac{\partial}{\partial p} \delta(p_F - p) \frac{\partial}{\partial p'} \delta(p_F - p') \\
& \rightarrow \frac{1}{(2\pi\hbar)^6} \int d\Omega \delta p_F \cos \theta \int d\Omega' f_{pp'\sigma\sigma'} \delta p'_F \cos \theta' \int dp p^2 \frac{\partial}{\partial p} \delta(p_F - p) \int dp' p'^2 \frac{\partial}{\partial p'} \delta(p_F - p') \\
& = \frac{N(0)}{p_F^2} \sum_{\ell\ell'\ell''\sigma\sigma'} \nu_{\ell\sigma} \nu_{\ell'\sigma'} F_{\ell''\sigma\sigma'} \begin{pmatrix} \ell'' & \ell & 1 \\ 0 & 0 & 0 \end{pmatrix}^2 \begin{pmatrix} \ell'' & \ell' & 1 \\ 0 & 0 & 0 \end{pmatrix}^2 \\
& \approx \frac{N(0)}{p_F^2} \sum_{\ell\ell'\ell''\sigma\sigma'} \nu_{\ell\sigma} \nu_{\ell'\sigma'} F_{\ell''\sigma\sigma'} \left\{ \frac{\delta_{\ell 0} \delta_{\ell'' 1}}{3} + \frac{\delta_{\ell 1} \delta_{\ell'' 0}}{3} \right\} \left\{ \frac{\delta_{\ell' 0} \delta_{\ell'' 1}}{3} + \frac{\delta_{\ell' 1} \delta_{\ell'' 0}}{3} \right\} \\
& = \boxed{\frac{N(0)}{4} \sum_{\ell\sigma\sigma'} \nu_{\ell\sigma} \nu_{\ell\sigma'} \left\{ \frac{4F_{1\sigma\sigma'}}{3p_F^2} \delta_{\ell 0} + \frac{4F_{0\sigma\sigma'}}{3p_F^2} \delta_{\ell 1} \right\}} \quad (49)
\end{aligned}$$

where we have used the fact that

$$\begin{pmatrix} \ell'' & \ell & 1 \\ 0 & 0 & 0 \end{pmatrix}^2 = \frac{(\ell - \ell'')^2 (1 + \ell + \ell'')}{(\ell + \ell'')(2 + \ell + \ell'')\Gamma(2 + \ell - \ell'')\Gamma(2 - \ell + \ell'')} \quad (50)$$

for $\ell'' + \ell$ odd and $\ell - 1 \leq \ell'' \leq 1 + \ell$. For $\ell = 0$, $\ell'' = 1$ while for $\ell = 1$, $\ell'' = 0$ or $\ell'' = 2$. Ignoring all terms higher than $\ell = 1$, we can then say that

$$\begin{pmatrix} \ell'' & \ell & 1 \\ 0 & 0 & 0 \end{pmatrix}^2 \approx \frac{1}{3} \delta_{\ell 0} \delta_{\ell'' 1} + \frac{1}{3} \delta_{\ell 1} \delta_{\ell'' 0} \quad (51)$$

Bringing all terms that go as $A_\sigma \cdot A_{\sigma'}$ together, we find that it yields the following:

$$\begin{aligned}
& \frac{N(0)}{4} \sum_{\sigma\sigma'} A_\sigma A_{\sigma'} \frac{F_{1\sigma\sigma'}}{9} v_F^2 - \frac{N(0)}{4} \sum_{\ell\sigma\sigma'} A_\sigma A_{\sigma'} \frac{4}{9} F_{1\sigma\sigma'} \frac{\nu_{\ell\sigma}}{m^*} \delta_{\ell 0} \\
& + \frac{N(0)}{4} \sum_{\ell\sigma\sigma'} A_\sigma A_{\sigma'} \nu_{\ell\sigma} \nu_{\ell\sigma'} \left\{ \frac{4F_{1\sigma\sigma'}}{3p_F^2} \delta_{\ell 0} + \frac{4F_{0\sigma\sigma'}}{3p_F^2} \delta_{\ell 1} \right\} \quad (52)
\end{aligned}$$

$$= \frac{N(0)}{4} \frac{1}{3} \sum_{\sigma\sigma'} A_\sigma A_{\sigma'} \left[\frac{F_{1\sigma\sigma'}}{3} \left\{ v_F^2 - 4 \sum_{\ell} \frac{\nu_{\ell\sigma}}{m^*} \delta_{\ell 0} \right\} + \frac{4}{p_F^2} \sum_{\ell} \nu_{\ell\sigma} \nu_{\ell\sigma'} \{ F_{1\sigma\sigma'} \delta_{\ell 0} + F_{0\sigma\sigma'} \delta_{\ell 1} \} \right] \quad (53)$$

We can then bring all the terms that go as $A_\sigma \cdot A_{\sigma'}$ together, and we simplify piece-by-piece:

$$\begin{aligned}
& \sum_{\sigma\sigma'} A_\sigma A_{\sigma'} \frac{F_{1\sigma\sigma'}}{3} \left\{ v_F^2 - 4 \sum_\ell \frac{\nu_{\ell\sigma}}{m^*} \delta_{\ell 0} \right\} \\
&= \frac{v_F^2}{3} (A_\uparrow^2 (F_1^s + F_1^a) + 2A_\uparrow A_\downarrow (F_1^s - F_1^a) + A_\downarrow^2 (F_1^s + F_1^a)) \\
&- \frac{4}{m^*} \sum_\ell \delta_{\ell 0} (\nu_{\ell\uparrow} A_\uparrow^2 (F_1^s + F_1^a) + \nu_{\ell\uparrow} A_\uparrow A_\downarrow (F_1^s - F_1^a) + \nu_{\ell\downarrow} A_\downarrow A_\uparrow (F_1^s - F_1^a) + \nu_{\ell\downarrow} A_\downarrow^2 (F_1^s + F_1^a)) \\
&= \begin{cases} \frac{4F_1^s}{3} A^2 \left\{ v_F^2 - \sum_\ell \frac{4\nu_{\ell s}}{m^*} \delta_{\ell 0} \right\}, & A_\uparrow = A_\downarrow \\ \frac{4F_1^a}{3} A^2 \left\{ v_F^2 - \sum_\ell \frac{4\nu_{\ell s}}{m^*} \delta_{\ell 0} \right\}, & A_\uparrow = -A_\downarrow \end{cases} \quad (54)
\end{aligned}$$

For the second term, we have

$$\begin{aligned}
\sum_{\ell\sigma\sigma'} A_\sigma A_{\sigma'} \frac{4}{p_F^2} \nu_{\ell\sigma} \nu_{\ell\sigma'} \{F_{1\sigma\sigma'} \delta_{\ell 0} + F_{0\sigma\sigma'} \delta_{\ell 1}\} &\equiv \frac{4}{p_F^2} \sum_{\ell\sigma\sigma'} A_\sigma A_{\sigma'} \nu_{\ell\sigma} \nu_{\ell\sigma'} \tilde{F}_{\ell\sigma\sigma'} \\
&= \begin{cases} \frac{4A^2}{p_F^2} \sum_\ell \left(\tilde{F}_\ell^s \nu_{\ell s}^2 + \tilde{F}_\ell^a \nu_{\ell a}^2 \right), & A_\uparrow = A_\downarrow \\ \frac{4A^2}{p_F^2} \sum_\ell \left(\tilde{F}_\ell^s \nu_{\ell a}^2 + \tilde{F}_\ell^a \nu_{\ell s}^2 \right), & A_\uparrow = -A_\downarrow \end{cases} \quad (55)
\end{aligned}$$

We can then simplify the $A_\sigma \cdot A_{\sigma'}$ to the following:

$$\begin{aligned}
& \frac{N(0)}{4} \frac{1}{3} \sum_{\sigma\sigma'} A_\sigma A_{\sigma'} \left[\frac{F_{1\sigma\sigma'}}{3} \left\{ v_F^2 - 4 \frac{\nu_{\ell\sigma}}{m^*} \delta_{\ell 0} \right\} + \frac{4}{p_F^2} \nu_{\ell\sigma} \nu_{\ell\sigma'} \{F_{1\sigma\sigma'} \delta_{\ell 0} + F_{0\sigma\sigma'} \delta_{\ell 1}\} \right] \\
&= \frac{N(0)}{4} \left[\frac{4F_1^s}{9} v_F^2 + \sum_\ell A^2 \left\{ \begin{aligned} & -\frac{16F_1^s}{9m^*} \nu_{\ell s} \delta_{\ell 0} + \frac{4}{3p_F^2} \left(\tilde{F}_\ell^s \nu_{\ell s}^2 + \tilde{F}_\ell^a \nu_{\ell a}^2 \right), & A_\uparrow = A_\downarrow \\ & -\frac{16F_1^a}{9m^*} \nu_{\ell s} \delta_{\ell 0} + \frac{4}{3p_F^2} \left(\tilde{F}_\ell^s \nu_{\ell a}^2 + \tilde{F}_\ell^a \nu_{\ell s}^2 \right), & A_\uparrow = -A_\downarrow \end{aligned} \right\} \right] \quad (56)
\end{aligned}$$

Let's try to simplify the total quadratic contribution to the free energy. We'll start with the case of $A_\uparrow = A_\downarrow$:

$$\begin{aligned}
& \frac{N(0)}{8} \left[\sum_{\ell} \left(\frac{1}{2\ell+1} \frac{1}{2\ell+1} \right. \right. \\
& \quad \times \left\{ F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2 + 4A \left(v_F F_{\ell}^s \nu_{\ell s} \delta_{\ell 0} - \frac{1}{p_F} \left\{ \nu_{\ell s}^2 \left[F_{\ell}^s + \frac{1}{2}(2\ell+1)F_0^s \right] + \nu_{\ell a}^2 \left[F_{\ell}^a + \frac{1}{2}(2\ell+1)F_0^s \right] \right\} \right) \right\} \\
& \quad + 4A^2 \left\{ -\frac{2F_{\ell}^s}{m^*(2\ell+1)^2} \nu_{\ell s} \delta_{\ell 0} + \frac{F_0^s}{p_F^2(2\ell+1)^2} (\nu_{\ell s}^2 + \nu_{\ell a}^2) - \frac{16F_1^s}{9m^*} \nu_{\ell s} \delta_{\ell 0} + \frac{4}{3p_F^2} \left(\tilde{F}_{\ell}^s \nu_{\ell s}^2 + \tilde{F}_{\ell}^a \nu_{\ell a}^2 \right) \right\} \left. \right) + \frac{4F_1^s}{9} v_F^2 \Big] \\
& = \frac{N(0)}{8} \left[\sum_{\ell} \left(\frac{1}{2\ell+1} \frac{1}{2\ell+1} \right. \right. \\
& \quad \times \left\{ F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2 + \frac{4A}{p_F} \left(2\epsilon_F F_{\ell}^s \nu_{\ell s} \delta_{\ell 0} - \left\{ \nu_{\ell s}^2 \left[F_{\ell}^s + \frac{1}{2}(2\ell+1)F_0^s \right] + \nu_{\ell a}^2 \left[F_{\ell}^a + \frac{1}{2}(2\ell+1)F_0^s \right] \right\} \right) \right\} \Big] \\
& \quad + \frac{4A^2}{p_F^2} \left\{ -4\epsilon_F \delta_{\ell 0} \left[8F_1^s + \frac{F_{\ell}^s}{(2\ell+1)^2} \right] \nu_{\ell s} + \left[\frac{F_0^s}{(2\ell+1)^2} + \frac{4\tilde{F}_{\ell}^s}{3} \right] \nu_{\ell s}^2 + \left[\frac{F_0^s}{(2\ell+1)^2} + \frac{4\tilde{F}_{\ell}^a}{3} \right] \nu_{\ell a}^2 \right\} \left. \right) + \frac{64A^2}{9p_F^2} F_1^s \epsilon_F^2 \Big] \\
& \approx \left[\frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left(\frac{1}{2\ell+1} \{ F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2 \} + \frac{8A}{p_F} \frac{F_{\ell}^s}{2\ell+1} \epsilon_F \nu_{\ell s} \delta_{\ell 0} \right) + \frac{64A^2}{9p_F^2} F_1^s \epsilon_F^2 \right] \right] \quad (57)
\end{aligned}$$

We now do the same simplification for the spin gauge $A_{\uparrow} = -A_{\downarrow}$:

$$\begin{aligned}
& \frac{N(0)}{8} v \sum_{\ell} \left(\frac{1}{2\ell+1} \frac{1}{2\ell+1} \left\{ F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2 + 4A \left(v_F F_{\ell}^s \nu_{\ell a} \delta_{\ell 0} - \frac{1}{p_F} \nu_{\ell s} \nu_{\ell a} \left\{ F_{\ell}^s + F_{\ell}^a + (2\ell+1)F_0^s \right\} \right) \right\} \right. \\
& \quad + 4A^2 \left\{ -\frac{2F_{\ell}^s}{m^*(2\ell+1)^2} \nu_{\ell s} \delta_{\ell 0} + \frac{F_0^s}{p_F^2(2\ell+1)^2} (\nu_{\ell s}^2 + \nu_{\ell a}^2) - \frac{16F_1^a}{m^*} \nu_{\ell s} \delta_{\ell 0} + \frac{4}{3p_F^2} \left(\tilde{F}_{\ell}^s \nu_{\ell a}^2 + \tilde{F}_{\ell}^a \nu_{\ell s}^2 \right) \right\} \left. \right) + \frac{4F_1^a}{9} v_F^2 \Big] \\
& = \frac{N(0)}{8} \left[\sum_{\ell} \left(\frac{1}{2\ell+1} \frac{1}{2\ell+1} \left\{ F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2 + \frac{4A}{p_F} \nu_{\ell a} \left(2\epsilon_F F_{\ell}^s \delta_{\ell 0} - \nu_{\ell s} \left\{ F_{\ell}^s + F_{\ell}^a + (2\ell+1)F_0^s \right\} \right) \right\} \right) \right. \\
& \quad + \frac{4A^2}{p_F^2} \left\{ -4\epsilon_F \frac{F_{\ell}^s}{(2\ell+1)^2} \nu_{\ell s} \delta_{\ell 0} + \frac{F_0^s}{(2\ell+1)^2} (\nu_{\ell s}^2 + \nu_{\ell a}^2) - \frac{32}{9} \epsilon_F^2 F_1^a \nu_{\ell s} \delta_{\ell 0} + \frac{4}{3} \left(\tilde{F}_{\ell}^s \nu_{\ell a}^2 + \tilde{F}_{\ell}^a \nu_{\ell s}^2 \right) \right\} \left. \right) + \frac{64A^2}{9p_F^2} F_1^a \epsilon_F^2 \Big] \\
& \approx \left[\frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left(\frac{1}{2\ell+1} \{ F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2 \} + \frac{8A}{p_F} \frac{F_{\ell}^s}{2\ell+1} \epsilon_F \nu_{\ell a} \delta_{\ell 0} \right) + \frac{64A^2}{9p_F^2} F_1^a \epsilon_F^2 \right] \right] \quad (58)
\end{aligned}$$

Putting this altogether, we find that

$$\begin{aligned}
& \sum_{pp'\sigma\sigma'} f_{pp'\sigma\sigma'} \delta n_{p-A_{\sigma},\sigma} \delta n_{p'-A_{\sigma'},\sigma'} \\
&= \begin{cases} \left[\frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left(\frac{1}{2\ell+1} \{F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2\} + \frac{8A}{p_F} \frac{F_{\ell}^s}{2\ell+1} \epsilon_F \nu_{\ell s} \delta_{\ell 0} \right) + \frac{64A^2}{9p_F^2} F_1^s \epsilon_F^2 \right], & A_{\uparrow} = A_{\downarrow} \right. \\ \left. \left[\frac{N(0)}{8} \left[\sum_{\ell} \frac{1}{2\ell+1} \left(\frac{1}{2\ell+1} \{F_{\ell}^s \nu_{\ell s}^2 + F_{\ell}^a \nu_{\ell a}^2\} + \frac{8A}{p_F} \frac{F_{\ell}^s}{2\ell+1} \epsilon_F \nu_{\ell a} \delta_{\ell 0} \right) + \frac{64A^2}{9p_F^2} F_1^a \epsilon_F^2 \right] \right] & A_{\uparrow} = -A_{\downarrow} \end{cases}
\end{aligned}$$

This completes the calculation of Eqn. (5).