

INVARIANT MEASURE FOR 2D STOCHASTIC CAHN-HILLIARD-NAVIER-STOKES EQUATIONS

ZHAOYANG QIU

ABSTRACT. Using the Maslowski and Seidler method, the existence of invariant measure for 2-dimensional stochastic Cahn-Hilliard-Navier-Stokes equations with multiplicative noise is proved in state space $L_x^2 \times H^1$, working with the weak topology. Also, the existence of global pathwise solution is investigated using the stochastic compactness argument.

1. INTRODUCTION

The study of dynamical behaviour of solutions to SPDEs is a major subject, especially in geophysical flow, climate dynamics, gene regulation system, chemical reaction systems. Here we consider the existence of invariant measure to the 2-dimensional stochastic Cahn-Hilliard-Navier-Stokes(CH-NS) equations with multiplicative noise in a smooth bounded domain \mathcal{D} ,

$$\begin{cases} du - (\nu_0 \Delta u - (u \cdot \nabla)u - \nabla p + \mu \nabla \phi)dt = h(u, \nabla \phi)d\mathcal{W}, \\ d\phi + (u \cdot \nabla)\phi dt - \nu_2 \Delta \mu dt = 0, \\ \mu = -\nu_1 \Delta \phi + \kappa f(\phi), \\ \nabla \cdot u = 0, \end{cases} \quad (1.1)$$

which describe the motion of two immiscible mixture fluid, consisting of the Navier-Stokes equations governing the fluid velocity and a convective Cahn-Hilliard equation which is responsible for the order parameter standing for the relative concentration of one of the fluids. For further physical background, see [6, 16, 19–21].

In equations (1.1), u, p and ϕ denote the velocity, pressure and the phase parameter, respectively. ν_0 is the viscosity of the fluid, here we assume that $\nu_0 \equiv 1$. Here μ stands for the chemical potential of binary mixtures which is the variational derivative of the quantity (free energy functional):

$$E(\phi) = \int_{\mathcal{D}} \frac{\nu_1}{2} |\nabla \phi|^2 + \kappa F(\phi) dx,$$

where $F(\phi) = \int_0^\phi f(r)dr$, the two physical parameters ν_1, κ describe the interaction between two phases. Especially, the constant $\nu_1 > 0$ is related to the thickness of the interface between the two fluids. $\nu_2 > 0$ is the mobility constant. In physical background,

Date: January 28, 2022.

2010 Mathematics Subject Classification. 35Q35, 76D05, 35R60, 60F10.

Key words and phrases. Stochastic Cahn-Hilliard-Navier-Stokes equations, pathwise solution, invariant measure, bounded weakly continuous.

a representative example of F is logarithmic type, that is,

$$F(s) = c_0[(1+s)\ln(1+s) + (1-s)\ln(1-s)] - c_1 s^2, c_1 > c_0 > 0,$$

where $s \in (-1, 1)$. Usually, we use a polynomial approximation of the type $F(\phi) = C_1\phi^4 - C_2\phi^2$ taking place of the type of logarithmic, where C_1, C_2 are two positive constants.

The system (1.1) is equipped with the following natural no-flux boundary conditions

$$\left. \frac{\partial \phi}{\partial \mathbf{n}} \right|_{\partial \mathcal{D}} = \left. \frac{\partial \Delta \phi}{\partial \mathbf{n}} \right|_{\partial \mathcal{D}} = 0, \quad (1.2)$$

where \mathbf{n} is the outward normal on the boundary $\partial \mathcal{D}$. It is easily to check that above conditions imply

$$\left. \frac{\partial \mu}{\partial \mathbf{n}} \right|_{\partial \mathcal{D}} = 0, \quad (1.3)$$

together with equation (1.1)₂ yields the conservation form of phase parameter ϕ , that is

$$\langle \phi(t) \rangle := \frac{1}{|\mathcal{D}|} \int_{\mathcal{D}} \phi(t, x) dx = \langle \phi(0) \rangle,$$

for all $t > 0$, where $|\mathcal{D}|$ is the Lebesgue measure of \mathcal{D} . For the boundary condition of u , we impose the following Dirichlet boundary condition

$$u|_{\partial \mathcal{D}} = 0. \quad (1.4)$$

Moreover, the initial data is

$$u(x, 0) = u_0(x), \quad \phi(x, 0) = \phi_0(x). \quad (1.5)$$

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space. \mathcal{W} is a cylindrical Wiener process defined on the Hilbert space \mathcal{H} , which is adapted to the complete, right continuous filtration $\{\mathcal{F}_t\}_{t \geq 0}$. Namely, $\mathcal{W} = \sum_{k \geq 1} e_k \beta_k$ with $\{e_k\}_{k \geq 1}$ being the complete orthonormal basis of \mathcal{H} and $\{\beta_k\}_{k \geq 1}$ being a sequence of independent standard one-dimensional Brownian motions. Considering an auxiliary space $\mathcal{H}_0 \supset \mathcal{H}$, define by

$$\mathcal{H}_0 = \left\{ h = \sum_{k \geq 1} \alpha_k e_k : \sum_{k \geq 1} \alpha_k^2 k^{-2} < \infty \right\},$$

with the norm $\|h\|_{\mathcal{H}_0}^2 = \sum_{k \geq 1} \alpha_k^2 k^{-2}$. Observe that the mapping $\Phi : \mathcal{H} \rightarrow \mathcal{H}_0$ is Hilbert-Schmidt. We also have that $\mathcal{W} \in C([0, \infty), \mathcal{H}_0)$ almost surely, see [7].

Although the complicated phenomenon of mixture fluid is still far from being well-understood, since its importance in physics and the mathematical challenge, CH-NS equations already received widely attention both in deterministic and stochastic areas, see [10, 13–17, 19, 25, 27].

The study of invariant measure could be traced back to the 1930's, see [26], the well-known method of proving the existence of invariant measure was introduced by Krylov-Bogoliubov which was limited to the state space of Markov process lay in the locally compact spaces. Then, the relative theory was extended to case of the infinite dimensional Banach spaces, see [3, 11]. Both of the cases highly require, first the semigroup \mathbf{P}_t is Feller,

second, find a Borel probability measure ξ on the state space X and T_0 such that the set of measures

$$\mathcal{M} = \left\{ \frac{1}{T} \int_0^T \mathbf{P}_t^* \xi dt; T > T_0 \right\} \quad (1.6)$$

is tight where \mathbf{P}_t^* is the dual semigroup, see [8], which requires us to find a auxiliary set compactly embedding into the state space.

First, in the background of SPDEs, the weak solution generally possesses the property of continuous in state space under the weak topology, second, in our case, even if for the phase parameter ϕ , we have the auxiliary set $H^2 \hookrightarrow H^1$ compactly, the high-order nonlinear term $f(\phi)$ and the special constitution cause the difficulty to get the bound for $\int_0^t \mathbb{E} \|\Delta \phi\|_{L_x^2}^2 ds \sim O(t)$ for t large. These two reasons push us to make use of the method developed by Maslowski and Seidler [23] extending the Krylov-Bogoliubov theory to work with the weak topology which simplifies the proof of (1.6). The method is widely occupied recently, see [5] for beam equation and nonlinear wave equation, [4] for the Navier-Stokes equation in unbounded domain, [1] for the damped 2D Euler equation.

In order to construct the transition semigroup to 2-dimensional stochastic CH-NS system (1.1)-(1.5), the existence of global pathwise solution as a cornerstone will be shown by a stochastic compactness approach instead of the Galerkin-type approximation convergence method applied by [2, 25]. The specific strategy is, following the Yamada-Watanabe theorem, to verify the existence of martingale solution (weak in probability sense), then the known uniqueness result as well as the Gyöngy-Krylov characterization allows us to recover the compactness. Here, we impose the following two conditions on noise intensity operator h : there exist two constants K_0, K_1 such that

$$\mathbf{C.1} \quad \|h(u, \nabla \phi)\|_{L_2(\mathcal{H}; X)}^2 \leq K_0(1 + \|(u, \nabla \phi)\|_X^2),$$

$$\mathbf{C.2} \quad \|h(u_1, \nabla \phi_1)\|_{L_2(\mathcal{H}; X)}^2 \leq K_1(1 + \|(u_1 - u_2, \nabla(\phi_1 - \phi_2))\|_X^2),$$

where $L_2(\mathcal{H}, X)$ denotes the collection of Hilbert-Schmidt operators, the set of all linear operators G from \mathcal{H} to the Banach space X , with the norm $\|G\|_{L_2(\mathcal{H}, X)}^2 = \sum_{k \geq 1} \|G e_k\|_X^2$.

Required by the tightness of (1.6) on space $(L_x^2 \times H^1)_w$, the quantity

$$\int_0^t \mathbb{E} \|\nabla \phi\|_{L_x^2}^2 ds$$

at most has a linear growth with respect to time t . Unlike the coupled SPDEs with the "symmetric" construction, here it seems impossible to acquire this quantity directly by taking inner product in equation (1.1)₂. In addition, the high-order nonlinear term $f(\phi)$ also makes the estimate challenging which forces us to consider the most physically specific case $f(\phi) = C_1 \phi^3 - C_2 \phi$, see Remark 3.2. To overcome these difficulties, two extra conditions are imposed on constant K_0 and the mobility constant ν_2 , that is,

$$\mathbf{C.3} \quad \text{the fixed constant } K_0 \text{ in } \mathbf{C.1} \text{ satisfies } CK_0 \leq 2, \text{ where } C = C(\mathcal{D}) \text{ is constant,}$$

$$\mathbf{C.4} \quad \text{the mobility constant } \nu_2 \text{ is large.}$$

For further technique details, see the Step 2 of Lemma 5.2.

The rest of paper goes as follows. We present the weak formula of the system in Section 2. In Section 3 we mainly focus on the global existence and uniqueness of the pathwise solution, also the necessary estimates are given. In section 4, we provide the auxiliary result which is crucial in both establishing the existence of pathwise solution and the sequentially weakly continuous of transition semigroup. The existence of invariant measure is proved in

Section 5. An Appendix is also included afterwards to state the results that are frequently used in the paper.

2. WEAK FORMULATION

At the beginning, we provide some preliminaries. Throughout the paper, we use the notation

$$\begin{aligned} L_t^p X &:= L^p(0, T; X), \quad L_\omega^p X := L^p(\Omega; X), \\ C_t^\alpha X &:= C^\alpha([0, T]; X), \quad W_t^{\alpha, p} X := W^{\alpha, p}(0, T; X) \end{aligned}$$

for $\alpha \in (0, 1], p \geq 1$, where X is the Banach space and Ω is the sample space. Here, we use the fractional order Sobolev space with respect to time t , since the evolution equations forced by noise only have α -order Hölder continuous in t for $\alpha \in (0, \frac{1}{2})$, see [9, 12]. Denote X' as the dual of Banach space X , \mathbb{E} as the mathematical expectation. The notation $\langle \cdot, \cdot \rangle$ defines the dual between the space X and its dual space X' .

Let X_w be the Hilbert space X with the weak topology. Introduce

$$C([0, T]; X_w) := \text{the space of } X \text{ valued weakly continuous function,}$$

endowed with the weak topology such that the mapping

$$u \mapsto \langle u, h \rangle$$

is continuous for any $h \in X$, which is a quasi-Polish space, for more details see [5], and

$$(L_t^p X)_w := \text{the space } L^p(0, T; X) \text{ with the weak topology.}$$

We say a functional $f : X \rightarrow \mathbb{R}$ is sequentially weakly continuous if $x_n \rightarrow x$ weakly star in X , then $f(x_n) \rightarrow f(x)$, where X is the Banach space, and we use $C_b(X_w)$ to stand for the bounded weakly continuous function.

Denote $L_x^p, p \geq 1$ the usual space of p -order Lebesgue integrable function on space domain \mathcal{D} and

$$H^{s, p} := \left\{ u \in L_x^p; \frac{\partial^i u}{\partial x} \in L_x^p, i = 1, \dots, s \right\},$$

endowed with the norm

$$\|u\|_{H^{s, p}}^p = \|u\|_{L_x^p}^p + \sum_{i=1}^s \|\partial^i u\|_{L_x^p}^p.$$

Next, introduce the spaces of the fluid velocity u , denote

$$\mathbf{V} = \{u \in C_c^\infty(\mathcal{D}); \operatorname{div} u = 0 \text{ in } \mathcal{D}\}.$$

Define by L_x^p the closure of \mathbf{V} with respect to the norm of L_x^p , and H^s the closure of \mathbf{V} with respect to the norm of H^s .

The norm $\|(u, v)\|_{L_x^2} := \|u\|_{L_x^2} + \|v\|_{L_x^2}$ for $u \in L_x^2, v \in L_x^2$.

Define the Stokes operator by

$$A_0 u = -P \Delta u, \text{ for all } u \in D(A_0) := \{u \in H^2 \cap L_x^2\}$$

where P is the Helmholtz-Leray projection. By the definition, we could have the operator A_0 is positive and self-adjoint on L_x^2 , and A_0^{-1} is compact. Therefore, there exists a sequence $\{e_k\}_{k \geq 1}$ being an orthonormal basis for L_x^2 of the Stokes operator A_0 .

Also introduce the operator $A_1 \phi = -\Delta \phi$ for any

$$\phi \in D(A_1) = \{\phi \in H^2(\mathcal{D}) : \partial_n \phi|_{\partial \mathcal{D}} = 0\}.$$

The Poincaré inequality implies the H^2 norm is equivalent to the norm $\|A_1(\cdot)\|_{L_x^2} + |\langle \cdot \rangle|$.

To define the variational setting, we next define $(b_0, b_1, b_2) : D(A_0) \times D(A_0) \rightarrow L_x^2$, $L_x^2 \times D(A_1) \rightarrow L_x^2$, $D(A_0) \times D(A_1) \rightarrow L_x^2$ as the bilinear operators such that

$$\begin{aligned}\langle b_0(u, v), w \rangle &= \int_{\mathcal{D}} (u \cdot \nabla v) \cdot w dx = B_0(u, v, w), \\ \langle b_1(\mu, \phi), w \rangle &= \int_{\mathcal{D}} \mu (\nabla \phi \cdot w) dx = B_1(\mu, \phi, w), \\ \langle b_2(u, \phi), \rho \rangle &= \int_{\mathcal{D}} (u \cdot \nabla \phi) \cdot \rho dx = B_2(u, \phi, \rho).\end{aligned}$$

Note that,

$$b_0(u, u) = Pu \cdot \nabla u, \quad b_1(\mu, \phi) = P\mu \nabla \phi$$

and for $u \in L_x^2, v \in H^1, \phi \in H^1$

$$B_0(u, v, v) = 0, \quad B_2(u, \phi, \phi) = 0. \quad (2.1)$$

With all these in hand, we could rewrite system (1.1) into the following form

$$\begin{cases} du + A_0 u dt + b_0(u, u) dt - b_1(\nu_1 A_1 \phi, \phi) dt = h(u, \nabla \phi) dW, \\ d\phi + \nu_2 A_1 \mu dt + b_2(u, \phi) dt = 0, \\ \mu = \nu_1 A_1 \phi + \kappa f(\phi), \end{cases} \quad (2.2)$$

understanding on space $(H^1)' \times (H^1)'$ (the weak sense).

Remark 2.1. Since $\nabla F(\phi) = f(\phi) \nabla \phi$, so $\mu \nabla \phi = -\nu_1 \Delta \phi \cdot \nabla \phi + \kappa \nabla F(\phi)$. The term $\nabla F(\phi)$ could be absorbed into the pressure. Hence, we could replace $b_1(\mu, \phi)$ by $b_1(\nu_1 A_1 \phi, \phi)$.

In order to handle the stochastic term, the following Burkholder-Davis-Gundy inequality is essential, for an X -valued predictable process $\varpi \in L^2(\Omega; L_{loc}^2([0, \infty), L_2(\mathcal{H}, X)))$ by taking $\varpi_k = \varpi e_k$, it holds

$$\mathbb{E} \left[\sup_{t \in [0, T]} \left\| \int_0^t \varpi dW \right\|_X^p \right] \leq c_p \mathbb{E} \left(\int_0^T \|\varpi\|_{L_2(\mathcal{H}, X)}^2 dt \right)^{\frac{p}{2}} = c_p \mathbb{E} \left(\int_0^T \sum_{k \geq 1} \|\varpi_k\|_X^2 dt \right)^{\frac{p}{2}},$$

for any $p \geq 1$.

3. EXISTENCE, UNIQUENESS AND THE NECESSARY ESTIMATES

In this section, we give the existence, uniqueness and the necessary estimates of the solution to system (1.1)-(1.5). To simplify the notation, we denote

$$\mathbb{H} := L_x^2 \times H^1, \quad \tilde{\mathbb{H}} := H^1 \times H^2.$$

Definition 3.1. (Martingale solution) Let Λ be a Borel probability measure on \mathbb{H} with $\int_{\mathbb{H}} |x|^p d\Lambda \leq C$ for a constant C . $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}, \mathcal{W}, u, \phi)$ is a martingale solution to the system (1.1)-(1.4) with the initial data Λ if the following conditions hold:

1. $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ is a stochastic basis and $\{\mathcal{F}_t\}_{t \geq 0}$ is complete right-continuous filtration, \mathcal{W} is a cylindrical Wiener process,
2. (u, ϕ) is the \mathcal{F}_t progressive measurable processes with

$$u \in L_{\omega}^p(L_t^{\infty} L_x^2 \cap L_t^2 H^1), \quad \phi \in L_{\omega}^p(L_t^{\infty} H^1 \cap L_t^2 H^2),$$

3. $\Lambda = \mathbb{P} \circ (u_0, \phi_0)^{-1}$,
4. for any $v \in H^1, \bar{v} \in H^1, t \in [0, T]$, (u, ϕ) satisfies \mathbb{P} -a.s.

$$\begin{cases} (u(t), v) + \int_0^t \langle A_0 u, v \rangle + B_0(u, u, v) - B_1(\nu_1 A_1 \phi, \phi, v) ds \\ \quad = (u_0, v) + \int_0^t (h(u, \nabla \phi), v) dW, \\ (\phi(t), \bar{v}) + \int_0^t \nu_2 \langle A_1 \mu, \bar{v} \rangle + B_2(u, \phi, \bar{v}) ds = (\phi_0, \bar{v}), \end{cases}$$

with

$$\mu = \nu_1 A_1 \phi + \kappa f(\phi), \quad \mathbb{P}\text{-a.s. in } \mathcal{D} \times (0, T).$$

Definition 3.2. (Pathwise solution) Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the fixed probability space. (u, ϕ) is a global pathwise solution to the system (1.1)-(1.5) if the following conditions hold:

1. (u, ϕ) is \mathcal{F}_t progressive measurable processes such that

$$u \in L_\omega^p(L_t^\infty L_x^2 \cap L_t^2 H^1), \quad \phi \in L_\omega^p(L_t^\infty H^1 \cap L_t^2 H^2),$$

2. for any $v \in H^1, \bar{v} \in H^1, t \in [0, T]$, (u, ϕ) satisfies \mathbb{P} -a.s.

$$\begin{cases} (u(t), v) + \int_0^t \langle A_0 u, v \rangle + B_0(u, u, v) - B_1(\nu_1 A_1 \phi, \phi, v) ds \\ \quad = (u_0, v) + \int_0^t (h(u, \nabla \phi), v) dW, \\ (\phi(t), \bar{v}) + \int_0^t \nu_2 \langle A_1 \mu, \bar{v} \rangle + B_2(u, \phi, \bar{v}) ds = (\phi_0, \bar{v}), \end{cases}$$

with

$$\mu = \nu_1 A_1 \phi + \kappa f(\phi), \quad \mathbb{P}\text{-a.s. in } \mathcal{D} \times (0, T).$$

Definition 3.3. (Uniqueness) If $(u_1, \phi_1), (u_2, \phi_2)$ are two solutions of system (1.1)-(1.5). Suppose that the initial data $(u_1(0), \phi_1(0))$ and $(u_2(0), \phi_2(0))$ satisfy

$$(u_1(0), \phi_1(0)) = (u_2(0), \phi_2(0)),$$

then the pathwise uniqueness holds if

$$\mathbb{P}\left\{(u_1(t, x), \phi_1(t, x)) = (u_2(t, x), \phi_2(t, x)); \quad \forall t \in [0, T]\right\} = 1.$$

Theorem 3.1. Assume that the initial data (u_0, ϕ_0) satisfies $\|(u_0, \phi_0)\|_{\mathbb{H}} \leq C_3$ for constant $C_3 > 0$, the operator h satisfies the Conditions **C.1** and **C.2**. Then, there exists a unique global pathwise solution to system (1.1)-(1.5) in the sense of Definitions 3.2, 3.3.

Remark 3.1. The proof of Theorem 3.1 was given in pioneering work [25], for deterministic case see [16]. To identify the limit, the author implemented the Galerkin-type approximation convergence in mean square introduced by [2]. Here, unlike [25], we use the stochastic compactness argument to establish the existence of pathwise solution.

Proof. For the proof of existence, the crucial technique details is the same with the result built in Section 4, therefore, here we only give a simplify proof .

Existence. Step 1. For the existence of finite-dimensional Galerkin approximate solutions, we refer the readers to [25], and also the sequence of approximate solutions (u_n, ϕ_n) satisfies the following uniform a priori estimates in n for all $p \geq 2$

$$u_n \in L_\omega^p(L_t^\infty L_x^2 \cap L_t^2 H^1), \quad \phi_n \in L_\omega^p(L_t^\infty H^1 \cap L_t^2 H^2), \quad \mu_n \in L_\omega^p(L_t^2 H^1). \quad (3.1)$$

Remark 3.2. The proof of the a priori estimates (3.1) is actually simpler than (5.17). Here, we do not have to handle the term $(F(\phi), 1)$, since it has the bound from below. Moreover, we could use the stochastic Gronwall lemma to absorb the term $\mathbb{E} \int_0^t \|\nabla \phi\|_{L_x^2}^2 ds$ arising from noise term, getting the upper bound with the exponential growth of t which is not suitable for proving the tightness of (1.6) on the space \mathbb{H}_w .

To order to obtain the tightness of the set of probability measures induced by the law of (u_n, ϕ_n) , we need the following necessary estimate:

there exists constant C such that for any $\alpha \in (0, \frac{1}{2})$, $\tilde{\alpha} \in (0, 1]$

$$\mathbb{E} \|u_n\|_{C_t^\alpha(H^{-1, \frac{3}{2}})}^2 + \mathbb{E} \|\phi_n\|_{C_t^{\tilde{\alpha}}(H^{-1})}^2 \leq C, \quad (3.2)$$

where C is independent of n .

To simplify the notation, we use (u, ϕ) replacing (u_n, ϕ_n) . Note that, for a.s. ω , and for any $\epsilon > 0$, there exists $t_1, t_2 \in [0, T]$ such that

$$\sup_{t_0 \neq t'_0} \frac{\left\| \int_{t_0}^{t'_0} Pu \cdot \nabla u ds \right\|_{H^{-1, \frac{3}{2}}}}{|t'_0 - t_0|^\alpha} \leq \frac{\left\| \int_{t_1}^{t_2} Pu \cdot \nabla u ds \right\|_{H^{-1, \frac{3}{2}}}}{|t_2 - t_1|^\alpha} + \epsilon.$$

Taking $t = \frac{3}{2}$, $s = 2$, $r = 2$ in Lemma 6.1 and using Hölder's inequality, for $\alpha \leq 1$,

$$\begin{aligned} \mathbb{E} \sup_{t, t' \in [0, T]} \frac{\left\| \int_{t'}^t Pu \cdot \nabla u ds \right\|_{H^{-1, \frac{3}{2}}}}{|t - t'|^\alpha} &\leq \mathbb{E} \frac{\left\| \int_{t_1}^{t_2} Pu \cdot \nabla u ds \right\|_{H^{-1, \frac{3}{2}}}}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq \mathbb{E} \frac{\int_{t_1}^{t_2} \|Pu \cdot \nabla u\|_{H^{-1, \frac{3}{2}}} ds}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq \mathbb{E} \frac{\int_{t_1}^{t_2} \|u\|_{H^1} \|\nabla u\|_{H^{-1}} ds}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq \frac{\left(\mathbb{E} \sup_{t \in [0, T]} \|u\|_{L_x^2}^2 \right)^{\frac{1}{2}} \left(\mathbb{E} \left[\int_{t_1}^{t_2} \|\nabla u\|_{H^{-1}} ds \right]^2 \right)^{\frac{1}{2}}}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq |t_2 - t_1|^{1-\alpha} \left(\mathbb{E} \sup_{t \in [0, T]} \|u\|_{L_x^2}^2 \right)^{\frac{1}{2}} \left(\mathbb{E} \int_{t_1}^{t_2} \|u\|_{H^1}^2 ds \right)^{\frac{1}{2}} + \epsilon \\ &\leq C. \end{aligned} \quad (3.3)$$

Similarly, one can estimate

$$\begin{aligned} \mathbb{E} \sup_{t, t' \in [0, T]} \frac{\left\| \int_{t'}^t P(A_1 \phi \cdot \nabla \phi) ds \right\|_{H^{-1, \frac{3}{2}}}}{|t - t'|^\alpha} &\leq \mathbb{E} \frac{\left\| \int_{t_1}^{t_2} A_1 \phi \cdot \nabla \phi ds \right\|_{H^{-1, \frac{3}{2}}}}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq \mathbb{E} \frac{\int_{t_1}^{t_2} \|A_1 \phi\|_{H^{-1}} \|\nabla \phi\|_{H^1} ds}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq |t_2 - t_1|^{1-\alpha} \left(\mathbb{E} \sup_{t \in [0, T]} \|\nabla \phi\|_{L_x^2}^2 \right)^{\frac{1}{2}} \left(\mathbb{E} \int_{t_1}^{t_2} \|\nabla \phi\|_{H^1}^2 ds \right)^{\frac{1}{2}} + \epsilon \end{aligned}$$

$$\leq C, \quad (3.4)$$

also, taking $t = 1, s = 3, r = 2$ in Lemma 6.1 and using the embedding $L_x^2 \hookrightarrow H^{-1,3}$,

$$\begin{aligned} \mathbb{E} \sup_{t, t' \in [0, T]} \frac{\left\| \int_{t'}^t u \cdot \nabla \phi ds \right\|_{H^{-1}}}{|t - t'|^\alpha} &\leq \mathbb{E} \frac{\left\| \int_{t_1}^{t_2} u \cdot \nabla \phi ds \right\|_{H^{-1}}}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq \mathbb{E} \frac{\int_{t_1}^{t_2} \|u\|_{H^{-1,3}} \|\nabla \phi\|_{H^1} ds}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq \mathbb{E} \frac{\int_{t_1}^{t_2} \|u\|_{L_x^2} \|\nabla \phi\|_{H^1} ds}{|t_2 - t_1|^\alpha} + \epsilon \\ &\leq |t_2 - t_1|^{1-\alpha} \left(\mathbb{E} \sup_{t \in [0, T]} \|u\|_{L_x^2}^2 \right)^{\frac{1}{2}} \left(\mathbb{E} \int_{t_1}^{t_2} \|\nabla \phi\|_{H^1}^2 ds \right)^{\frac{1}{2}} + \epsilon \\ &\leq C. \end{aligned} \quad (3.5)$$

By the Burkholder-Davis-Gundy inequality, the Condition **C.1** and (3.1), for $v \in H^1$

$$\begin{aligned} \mathbb{E} \left(\sup_{t_0 \neq t'_0} \frac{\left| \int_{t'_0}^{t'_0} (h(u, \nabla \phi), v) d\mathcal{W} \right|}{|t'_0 - t_0|^\alpha} \right)^p &\leq \mathbb{E} \frac{\left(\int_{t_1}^{t_2} \|h(u, \nabla \phi)\|_{L_2(\mathcal{H}; L_x^2)}^2 \|v\|_{L_x^2}^2 dt \right)^{\frac{p}{2}}}{|t_2 - t_1|^{\alpha p}} + \epsilon^p \\ &\leq \frac{CK_0^{\frac{p}{2}} (t_2 - t_1)^{\frac{p}{2}} \mathbb{E}(1 + \|(u, \nabla \phi)\|_{L_t^\infty L_x^2})^p}{|t_1 - t_2|^{\alpha p}} + \epsilon^p \\ &\leq CK_0^{\frac{p}{2}} |t_2 - t_1|^{(\frac{1}{2}-\alpha)p} + \epsilon^p \leq C, \end{aligned} \quad (3.6)$$

for any $\alpha < \frac{1}{2}$. For the linear term, by (3.1) we easily get

$$\mathbb{E} \left\| \int_0^t A_0 u ds \right\|_{C_t^1(H^{-1})}^2 \leq C, \quad \mathbb{E} \left\| \int_0^t A_1 \mu ds \right\|_{C_t^1(H^{-1})}^2 \leq C. \quad (3.7)$$

Taking into account of estimates (3.3)-(3.7), we obtain the desired bound (3.2).

Step 2. Next, we could use the bounds (3.1)(3.2) to establish the tightness of the probability measure set \mathcal{L}^n induced by the law of (u_n, ϕ_n) , the further details of the proof is same with the argument as Lemma 4.1 which is implemented for the solutions (u_l, ϕ_l) corresponding to initial data $(u_{0,l}, \phi_{0,l})$, then, we could identify the limit using the technique details we present in Step 2 of the proof of Proposition 4.1, establishing the existence of global martingale solution which is weak in both probability sense and PDE sense in the sense of Definition 3.1.

Step 3. To obtain the existence of global pathwise solution which is strong in probability sense, we shall recover the compactness of approximate sequences (u_n, ϕ_n) on the original space $(\Omega, \mathcal{F}, \mathbb{P})$ applying the Gyöngy-Krylov characterization, for the proof see Appendix. Then, passing the limit, the global pathwise solution follows. \square

4. CONTINUOUS DEPENDENCE OF INITIAL DATA

Let $(u_{0,l}, \phi_{0,l})$ be a sequence of initial data with

$$\|(u_{0,l}, \phi_{0,l})\|_{\mathbb{H}} \leq C_3 \text{ for all } l \in \mathbb{N}^+, \quad (4.1)$$

according to the Theorem 3.1, there exists a sequence global pathwise solutions (u_l, ϕ_l) to system (1.1)-(1.4) corresponding to initial data $(u_{0,l}, \phi_{0,l})$ and the fixed stochastic basis $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}, \mathcal{W})$, meanwhile, these solutions still enjoy the following uniform bounds

$$\mathbb{E} \left(\|u_l\|_{C_t^\alpha(H^{-1, \frac{3}{2}})}^2 + \|\phi_l\|_{C_t^{\tilde{\alpha}}(H^{-1})}^2 \right) \leq C \quad (4.2)$$

and

$$\mathbb{E} \left(\|(u_l, \phi_l)\|_{L_t^\infty \mathbb{H}}^p + \|(u_l, \phi_l)\|_{L_t^2 \widetilde{\mathbb{H}}}^p + \|\mu_l\|_{L_t^2 H^1}^p \right) \leq C, \quad (4.3)$$

for any $p \geq 2, \alpha \in (0, \frac{1}{2}), \tilde{\alpha} \in (0, 1]$, where constant C is independent of l .

This section is mainly to show the following continuous dependence result:

Proposition 4.1. *Assume that initial data $(u_{0,l}, \phi_{0,l})$ is an \mathbb{H} -valued sequence which converges weakly to $(u_0, \phi_0) \in \mathbb{H}$, and (4.1) holds for certain constant C_3 . Then, there exist a new probability space $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}})$, a new subsequence $(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mu}_{l_k}, \widetilde{\mathcal{W}}_{l_k})$ and the processes $(\tilde{u}, \tilde{\phi}, \tilde{\mu}, \widetilde{\mathcal{W}})$, such that*

$$(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mu}_{l_k}, \widetilde{\mathcal{W}}_{l_k}) \text{ and } (u_{l_k}, \phi_{l_k}, \mu_{l_k}, \mathcal{W}) \text{ have the same joint distribution in } \mathcal{X}, \quad (4.4)$$

$$(\tilde{u}, \tilde{\phi}) \text{ and } (u, \phi) \text{ have the same joint distribution in } \mathcal{X} \quad (4.5)$$

and

$$(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mu}_{l_k}, \widetilde{\mathcal{W}}_{l_k}) \rightarrow (\tilde{u}, \tilde{\phi}, \tilde{\mu}, \widetilde{\mathcal{W}}), \text{ in the topology of } \mathcal{X}, \text{ } \widetilde{\mathbb{P}}\text{-a.s.} \quad (4.6)$$

where the (u, ϕ) is the pathwise solution with the initial data (u_0, ϕ_0) . Moreover, we have for all $p \geq 1$

$$\widetilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|(\tilde{u}_{l_k}, \tilde{\phi}_{l_k})\|_{\mathbb{H}}^{2p} \right) + \widetilde{\mathbb{E}} \left(\int_0^T \|(\tilde{u}_{l_k}, \tilde{\phi}_{l_k})\|_{\mathbb{H}}^2 + \|\tilde{\mu}_{l_k}\|_{H^1}^2 dt \right)^p \leq C. \quad (4.7)$$

In addition, $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}}, \tilde{u}, \tilde{\phi}, \widetilde{\mathcal{W}})$ is a martingale solution of the system (1.1) in the sense of Definition 3.1.

Remark 4.1. For the Galerkin approximate solutions (u_n, ϕ_n) relative to the finite dimensional initial data $(u_{0,n}, \phi_{0,n})$ and the fixed stochastic basis $(\Omega, \mathcal{F}, \mathbb{P}, \mathcal{W})$, we also have the same results as (4.4), (4.6): there exist a new probability space $(\widehat{\Omega}, \widehat{\mathcal{F}}, \widehat{\mathbb{P}})$, a new subsequence $(\hat{u}_{n_k}, \hat{\phi}_{n_k}, \hat{\mu}_{n_k}, \widehat{\mathcal{W}}_{n_k})$ and the processes $(\hat{u}, \hat{\phi}, \hat{\mu}, \widehat{\mathcal{W}})$, such that

$$(\hat{u}_{n_k}, \hat{\phi}_{n_k}, \hat{\mu}_{n_k}, \widehat{\mathcal{W}}_{n_k}) \text{ and } (u_{n_k}, \phi_{n_k}, \mu_{n_k}, \mathcal{W}) \text{ have the same joint distribution in } \mathcal{X},$$

and

$$(\hat{u}_{n_k}, \hat{\phi}_{n_k}, \hat{\mu}_{n_k}, \widehat{\mathcal{W}}_{n_k}) \rightarrow (\hat{u}, \hat{\phi}, \hat{\mu}, \widehat{\mathcal{W}}), \text{ in the topology of } \mathcal{X}, \text{ } \widehat{\mathbb{P}}\text{-a.s..}$$

The key point of showing above continuous dependence result is to acquire the tightness of the probability measures induced by the law of (u_l, ϕ_l) . Define the set

$$\mathcal{X} = \mathcal{X}_{u,\phi} \times \mathcal{X}_\mu \times \mathcal{X}_\mathcal{W},$$

where

$$\mathcal{X}_{u,\phi} := C([0, T]; \mathbb{H}_w) \cap L_t^2 \mathbb{H} \cap (L_t^2 \widetilde{\mathbb{H}})_w, \quad \mathcal{X}_\mu := L_t^2 L_x^2 \cap (L_t^2 H^1)_w, \quad \mathcal{X}_\mathcal{W} := C_t(\mathcal{H}_0)$$

and \mathbb{H}_w denotes the spaces L_x^2, H^1 with the weak topology of L_x^2, H^1 respectively. Let Γ be the smallest topology on \mathcal{X} such that four natural embedding from \mathcal{X} are continuous.

Define the measure set

$$\mathcal{L}^l(B) = \mathbb{P}\{(u_l, \phi_l, \mu_l, \mathcal{W}) \in B\},$$

for $B \in \mathcal{B}(\mathcal{X})$ which is the smallest σ -algebra containing the family Γ .

Lemma 4.1. *The measure set \mathcal{L}^l is tight on space (\mathcal{X}, Γ) .*

Proof. It is enough to show that every measure set

$$\begin{aligned} \mathcal{L}_u^l(B_1) &:= \mathbb{P}\{u_l \in B_1\}, \quad \mathcal{L}_\phi^l(B_2) := \mathbb{P}\{\phi_l \in B_2\}, \\ \mathcal{L}_\mathcal{W}^l(B_3) &:= \mathbb{P}\{\mathcal{W} \in B_3\}, \quad \mathcal{L}_\mu^l(B_4) := \mathbb{P}\{\mu_l \in B_4\}, \end{aligned}$$

is tight on $\mathcal{X}_u, \mathcal{X}_\phi, \mathcal{X}_\mu, \mathcal{X}_\mathcal{W}$ respectively, for $B_1 \in \mathcal{B}(\mathcal{X}_u), B_2 \in \mathcal{B}(\mathcal{X}_\phi), B_3 \in \mathcal{B}(\mathcal{X}_\mathcal{W}), B_4 \in \mathcal{B}(\mathcal{X}_\mu)$, where $\mathcal{X}_u := C([0, T]; (L_x^2)_w) \cap L_t^2 L_x^2 \cap (L_t^2 H^1)_w$, $\mathcal{X}_\phi := C([0, T]; H_w^1) \cap L_t^2 H^1 \cap (L_t^2 H^2)_w$.

Claim 1. The probability measure set \mathcal{L}_u^l is tight on \mathcal{X}_u .

Define the set

$$\begin{aligned} B_{1,K}^u &= \left\{ u_l \in L_t^2 H^1 \cap C_t^\alpha(H^{-1, \frac{3}{2}}) : \|u_l\|_{L_t^2 H^1} + \|u_l\|_{C_t^\alpha(H^{-1, \frac{3}{2}})} \leq K \right\}, \\ B_{2,K}^u &= \left\{ u_l \in L_t^\infty L_x^2 \cap C_t^\alpha(H^{-1, \frac{3}{2}}) : \|u_l\|_{L_t^\infty L_x^2} + \|u_l\|_{C_t^\alpha(H^{-1, \frac{3}{2}})} \leq K \right\}. \end{aligned}$$

The Aubin-Lions compact embedding, see Lemma 6.2

$$L_t^2 H^1 \cap C_t^\alpha(H^{-1, \frac{3}{2}}) \hookrightarrow L_t^2 L_x^2, \quad L_t^\infty L_x^2 \cap C_t^\alpha(H^{-1, \frac{3}{2}}) \hookrightarrow C([0, T]; (L_x^2)_w)$$

implies that the set $B_{1,K}^u, B_{2,K}^u$ is relatively compact in $L_t^2 L_x^2, C([0, T]; (L_x^2)_w)$. Considering (4.2), (4.3), by the Chebyshev inequality we have

$$\begin{aligned} \mathcal{L}_u^l((B_{1,K}^u \cap B_{2,K}^u)^c) &\leq \mathbb{P}\left(\|u_l\|_{L_t^2 H^1} > \frac{K}{2}\right) + \mathbb{P}\left(\|u_l\|_{C_t^\alpha(H^{-1, \frac{3}{2}})} > \frac{K}{2}\right) \\ &\quad + \mathbb{P}\left(\|u_l\|_{L_t^\infty L_x^2} > \frac{K}{2}\right) + \mathbb{P}\left(\|u_l\|_{C_t^\alpha(H^{-1, \frac{3}{2}})} > \frac{K}{2}\right) \\ &\leq \frac{C}{K} \mathbb{E}\left(\|u_l\|_{L_t^2 H^1} + \|u_l\|_{L_t^\infty L_x^2} + \|u_l\|_{C_t^\alpha(H^{-1, \frac{3}{2}})}\right) \leq \frac{C}{K}. \end{aligned}$$

Note that, by the Alaoglu-Banach theorem, the set

$$B_{3,K}^u := \left\{ u_l \in L_t^2 H^1 : \|u_l\|_{L_t^2 H^1} \leq K \right\}$$

is relatively compact on path space $(L_t^2 H^1)_w$. The bound (4.3) and the Chebyshev inequality again, yield

$$\mathcal{L}_u^l((B_{3,K}^u)^c) \leq \frac{C}{K}.$$

We obtain the tightness of set \mathcal{L}_u^l .

Claim 2. The probability measure set \mathcal{L}_ϕ^l is tight on \mathcal{X}_ϕ .

In a similar way to the **Claim 1**, define the set

$$\begin{aligned} B_{1,K}^\phi &= \left\{ \phi_l \in L_t^2 H^2 \cap C_t^\alpha(H^{-1}) : \|\phi_l\|_{L_t^2 H^1} + \|\phi_l\|_{C_t^\alpha(H^{-1})} \leq K \right\}, \\ B_{2,K}^\phi &= \left\{ \phi_l \in L_t^\infty H^1 \cap C_t^\alpha(H^{-1}) : \|\phi_l\|_{L_t^\infty H^1} + \|\phi_l\|_{C_t^\alpha(H^{-1})} \leq K \right\}, \\ B_{3,K}^\phi &= \left\{ \phi_l \in L_t^2 H^2 : \|\phi_l\|_{L_t^2 H^2} \leq K \right\}. \end{aligned}$$

Applying the Aubin-Lions compact embedding

$$L_t^2 H^2 \cap C_t^\alpha H^{-1} \hookrightarrow L_t^2 H^1, L_t^\infty H^1 \cap C_t^\alpha(H^{-1}) \hookrightarrow C([0, T]; H_w^1),$$

we infer that $B_{1,K}^\phi, B_{2,K}^\phi$ is relatively compact in $L_t^2 H^1, C([0, T]; H_w^1)$ respectively. Also, by the Alaoglu-Banach theorem, the set $B_{3,K}^\phi$ is relatively compact in $(L_t^2 H^2)_w$.

By bounds (4.2), (4.3), we deduce that the probability measure set \mathcal{L}_ϕ^l is tight.

Claim 3. The probability measure set \mathcal{L}_μ^l is tight on \mathcal{X}_μ .

Since

$$\begin{aligned} & \mathbb{E} \left(\int_0^T \int_0^T \frac{\|f(\phi(t_1)) - f(\phi(t_2))\|_{H^{-1, \frac{3}{2}}}^p}{|t_1 - t_2|^{1+\alpha p}} dt_1 dt_2 \right)^{\frac{1}{p}} \\ & \leq C \mathbb{E} \left(\int_0^T \int_0^T \frac{\|\phi(t_1) - \phi(t_2)\|_{H^{-1}}^p \|\phi^2(t_1) + \phi^2(t_2)\|_{H^1}^p}{|t_1 - t_2|^{1+\alpha p}} dt_1 dt_2 \right)^{\frac{1}{p}} \\ & \leq C \mathbb{E} \left(\sup_{t_1, t_2 \in [0, T]} \frac{\|\phi(t_1) - \phi(t_2)\|_{H^{-1}}}{|t_1 - t_2|^{\frac{1}{p} + \alpha}} \right. \\ & \quad \times \left. \int_0^T \int_0^T \|(\phi(t_1), \phi(t_2))\|_{L_x^6}^p \|\nabla \phi(t_1), \nabla \phi(t_2)\|_{L_x^3}^p dt_1 dt_2^{\frac{1}{p}} \right) \\ & \leq C \left[\mathbb{E} \left(\sup_{t_1, t_2 \in [0, T]} \frac{\|\phi(t_1) - \phi(t_2)\|_{H^{-1}}}{|t_1 - t_2|^{\frac{1}{p} + \alpha}} \right)^2 \right]^{\frac{1}{2}} \\ & \quad \times \left[\mathbb{E} \left(\int_0^T \int_0^T \|(\phi(t_1), \phi(t_2))\|_{H^1}^p \|(\nabla \phi(t_1), \nabla \phi(t_2))\|_{H^2}^p dt_1 dt_2 \right)^{\frac{2}{p}} \right]^{\frac{1}{2}} \end{aligned}$$

taking $p = 2, \alpha = \frac{1}{2}$, by bounds (4.2), (4.3), the Hölder inequality, we have

$$\mathbb{E} \|\mu_l\|_{W^{\frac{1}{2}, 2}(0, T; H^{-3})} \leq C,$$

together with bound (4.3), using the same argument as above, we could infer that \mathcal{L}_μ^l is tight on \mathcal{X}_μ .

Since the sequence \mathcal{W} is only one element and thus, the set $\mathcal{L}_\mathcal{W}$ is weakly compact. Lemma 4.1 follows. \square

Proof of Proposition 4.1. Step 1. With the tightness of probability measure set \mathcal{L}^l in hand, from the Skorokhod-Jakubowski representation theorem (Theorem 6.2), we obtain (4.4) directly, and

$$(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mu}_{l_k}, \tilde{\mathcal{W}}_{l_k}) \rightarrow (\tilde{u}, \tilde{\phi}, \tilde{\mu}, \tilde{\mathcal{W}}), \text{ in the topology of } \mathcal{X}, \tilde{\mathbb{P}}\text{-a.s.}$$

In particular, the chemical potential μ is a nonlinear term, we need to verify that $\tilde{\mu} = -\nu_1 \Delta \tilde{\phi} + \kappa f(\tilde{\phi})$, a.e. on $(0, T) \times \mathcal{D}$. Note that, by the mean-value theorem, there exists $\theta \in (0, 1)$ such that

$$\begin{aligned} \|f(\tilde{\phi}_{l_k}) - f(\tilde{\phi})\|_{L_t^2 L_x^2} &\leq C \left\| |\theta \tilde{\phi} + (1 - \theta) \tilde{\phi}_{l_k}|^2 (\tilde{\phi}_{l_k} - \tilde{\phi}) \right\|_{L_t^2 L_x^2} \\ &\leq C \left\| |\theta \tilde{\phi} + (1 - \theta) \tilde{\phi}_{l_k}|^2 \right\|_{L_x^3} \|\tilde{\phi}_{l_k} - \tilde{\phi}\|_{L_x^6} \Big\|_{L_t^2} \\ &\leq C \|(\tilde{\phi}_{l_k}, \tilde{\phi})\|_{L_t^\infty H^1}^2 \|\tilde{\phi}_{l_k} - \tilde{\phi}\|_{L_t^2 H^1}. \end{aligned}$$

The result follows from the fact that $\|\tilde{\phi}_{l_k} - \tilde{\phi}\|_{L_t^2 H^1} \rightarrow 0$, $\tilde{\mathbb{P}}$ -a.s. as $k \rightarrow \infty$. Since $(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mu}_{l_k})$ and $(u_{l_k}, \phi_{l_k}, \mu_{l_k})$ have the same joint distribution, we obtain (4.7) from (4.3).

Step 2. We show that $(\tilde{u}, \tilde{\phi}, \tilde{\mu}, \tilde{\mathcal{W}})$ also satisfies the system (2.2). Again, since the sequences $(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mu}_{l_k}, \tilde{\mathcal{W}}_{l_k})$ and $(u_{l_k}, \phi_{l_k}, \mu_{l_k}, \mathcal{W})$ have the same joint distribution, we have $(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mu}_{l_k}, \tilde{\mathcal{W}}_{l_k})$ satisfies the system (2.2), the proof is standard, see [30]. Next, we identify the limit by passing $k \rightarrow \infty$.

For any $v \in H^2$, decomposing

$$B_0(\tilde{u}_{l_k}, \tilde{u}_{l_k}, v) - B_0(\tilde{u}, \tilde{u}, v) = B_0(\tilde{u}, \tilde{u}_{l_k} - \tilde{u}, v) + B_0(\tilde{u}_{l_k} - \tilde{u}, \tilde{u}_{l_k}, v).$$

By the convergence $\tilde{u}_{l_k} \rightarrow \tilde{u}$ in $L_t^2 L_x^2$, $\tilde{\mathbb{P}}$ -a.s. and Lemma 6.1, we have $\tilde{\mathbb{P}}$ -a.s.

$$\int_0^t |B_0(\tilde{u}_{l_k} - \tilde{u}, \tilde{u}_{l_k}, v)| ds \leq \|v\|_{H^2} \|\tilde{u}_{l_k} - \tilde{u}\|_{L_t^2 L_x^2} \|\tilde{u}_{l_k}\|_{L_t^2 H^1} \rightarrow 0.$$

Furthermore, using the bound $\tilde{u} \in L_t^2 H^1$ and the convergence $\tilde{u}_{l_k} \rightarrow \tilde{u}$ in $(L_t^2 H^1)_w$, $\tilde{\mathbb{P}}$ -a.s. we infer that

$$\int_0^t B_0(\tilde{u}, \tilde{u}_{l_k} - \tilde{u}, v) ds \rightarrow 0.$$

By Lemma 6.1, the Hölder inequality and the bound (4.7),

$$\begin{aligned} \tilde{\mathbb{E}} \left(\int_0^T B_0(\tilde{u}_{l_k}, \tilde{u}_{l_k}, v) dt \right)^p &\leq \tilde{\mathbb{E}} \left(\int_0^T \|\tilde{u}_{l_k}\|_{L_x^2} \|\tilde{u}_{l_k}\|_{H^1} \|v\|_{H^{1,3}} dt \right)^p \\ &\leq \|v\|_{H^2}^p \tilde{\mathbb{E}} \left[\sup_{t \in [0, T]} \|\tilde{u}_{l_k}\|_{L_x^2}^p \left(\int_0^T \|\tilde{u}_{l_k}\|_{H^1} dt \right)^p \right] \end{aligned}$$

$$\begin{aligned}
&\leq CT^{\frac{p}{2}} \|v\|_{H^2}^p \tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\tilde{u}_{l_k}\|_{L_x^2}^{2p} \right) \tilde{\mathbb{E}} \left(\int_0^T \|\tilde{u}_{l_k}\|_{H^1}^2 dt \right)^p \\
&\leq C.
\end{aligned} \tag{4.8}$$

Since

$$A_1 \phi \cdot \nabla \phi = \nabla \left(\frac{|\nabla \phi|^2}{2} \right) - \operatorname{div}(\nabla \phi \otimes \nabla \phi),$$

then, we have

$$\begin{aligned}
&B_1(\nu_1 A_1 \tilde{\phi}_{l_k}, \tilde{\phi}_{l_k}, v) - B_1(\nu_1 A_1 \tilde{\phi}, \tilde{\phi}, v) \\
&= \int_{\mathcal{D}} \nu_1 (\nabla \tilde{\phi}_{l_k} \otimes \nabla \tilde{\phi}_{l_k} - \nabla \tilde{\phi} \otimes \nabla \tilde{\phi}) : \nabla v dx \\
&= \int_{\mathcal{D}} \nu_1 ((\nabla \tilde{\phi}_{l_k} - \nabla \tilde{\phi}) \otimes \nabla \tilde{\phi}_{l_k} + \nabla \tilde{\phi} \otimes (\nabla \tilde{\phi}_{l_k} - \nabla \tilde{\phi})) : \nabla v dx.
\end{aligned}$$

By the convergence $\tilde{\phi}_{l_k} \rightarrow \tilde{\phi}$ in $L_t^2 H^1$, $\tilde{\mathbb{P}}$ -a.s. and the bound (4.7), we have $\tilde{\mathbb{P}}$ -a.s.

$$\begin{aligned}
\int_0^t \int_{\mathcal{D}} |((\nabla \tilde{\phi}_{l_k} - \nabla \tilde{\phi}) \otimes \nabla \tilde{\phi}_{l_k}) : \nabla v| dx ds &\leq \|\nabla \tilde{\phi}_{l_k} - \nabla \tilde{\phi}\|_{L_t^2 L_x^2} \|\nabla \tilde{\phi}_{l_k}\|_{L_t^2 L_x^6} \|\nabla v\|_{L_x^3} \\
&\leq \|\nabla \tilde{\phi}_{l_k} - \nabla \tilde{\phi}\|_{L_t^2 L_x^2} \|\nabla \tilde{\phi}_{l_k}\|_{L_t^2 H^1} \|\nabla v\|_{L_x^3} \rightarrow 0.
\end{aligned}$$

Also by the convergence $\tilde{\phi}_{l_k} \rightarrow \tilde{\phi}$ in $(L_t^2 H^2)_w$ and the bound $\tilde{\phi} \in L_t^2 H^2$, $\tilde{\mathbb{P}}$ -a.s. we have

$$\int_0^t \int_{\mathcal{D}} \nabla \tilde{\phi} \otimes (\nabla \tilde{\phi}_{l_k} - \nabla \tilde{\phi}) : \nabla v dx ds \rightarrow 0, \quad \tilde{\mathbb{P}}\text{-a.s.}$$

By Lemma 6.1, the Hölder inequality and the bound (4.7), we have

$$\begin{aligned}
\tilde{\mathbb{E}} \left(\int_0^T B_1(\nu_1 A_1 \tilde{\phi}_{l_k}, \tilde{\phi}_{l_k}, v) dt \right)^p &\leq C \|v\|_{H^{1,3}}^p \tilde{\mathbb{E}} \left(\int_0^T \|A_1 \tilde{\phi}_{l_k}\|_{H^{-1}} \|\nabla \tilde{\phi}_{l_k}\|_{H^1} dt \right)^p \\
&\leq CT^{\frac{p}{2}} \|v\|_{H^{1,3}}^p \tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\nabla \tilde{\phi}_{l_k}\|_{L^2}^{2p} \right) \tilde{\mathbb{E}} \left(\int_0^T \|\nabla \tilde{\phi}_{l_k}\|_{H^1}^2 dt \right)^p \\
&\leq C.
\end{aligned} \tag{4.9}$$

Applying the same argument, we could infer that for any $\bar{v} \in H^1$

$$\int_0^t B_2(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \bar{v}) - B_2(\tilde{u}, \tilde{\phi}, \bar{v}) ds \rightarrow 0, \quad \tilde{\mathbb{P}}\text{-a.s.}$$

and

$$\tilde{\mathbb{E}} \left(\int_0^T B_2(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \bar{v}) dt \right)^p \leq C. \tag{4.10}$$

Finally, we pass the limit in stochastic term. Using the Condition **C.2**

$$\|h(\tilde{u}_{l_k}, \nabla \tilde{\phi}_{l_k}) - h(\tilde{u}, \nabla \tilde{\phi})\|_{L_2(\mathcal{H}; L_x^2)} \leq K_1 \|(\tilde{u}_{l_k} - \tilde{u}, \nabla(\tilde{\phi}_{l_k} - \tilde{\phi}))\|_{L_x^2},$$

together with the Proposition 4.1 (4.6) yields

$$\|h(\tilde{u}_{l_k}, \nabla \tilde{\phi}_{l_k}) - h(\tilde{u}, \nabla \tilde{\phi})\|_{L_2(\mathcal{H}; L_x^2)} \rightarrow 0, \quad \text{a.e. } (t, \omega) \in [0, T] \times \tilde{\Omega}.$$

Combining the convergence and the bound (4.7), we have from the Vitali convergence theorem (see Theorem 6.1),

$$h(\tilde{u}_{l_k}, \nabla \tilde{\phi}_{l_k}) \rightarrow h(\tilde{u}, \nabla \tilde{\phi}), \text{ in } L_\omega^p L_t^2 L_2(\mathcal{H}; L_x^2),$$

which implies the convergence in probability in $L_t^2 L_2(\mathcal{H}; L_x^2)$. Moreover, from Proposition 4.1 (4.6), we have

$$\widetilde{\mathcal{W}}_{l_k} \rightarrow \widetilde{\mathcal{W}}, \text{ in probability in } C_t(\mathcal{H}_0).$$

Then, applying the Lemma 6.3, we infer that

$$\int_0^t h(\tilde{u}_{l_k}, \nabla \tilde{\phi}_{l_k}) d\widetilde{\mathcal{W}}_{l_k} \rightarrow \int_0^t h(\tilde{u}, \nabla \tilde{\phi}) d\widetilde{\mathcal{W}},$$

in probability in $L_t^2 L_2(\mathcal{H}; L_x^2)$. Using the Itô isometry and the Condition **C.1**, we have

$$\begin{aligned} \widetilde{\mathbb{E}} \left| \int_0^T h(\tilde{u}_{l_k}, \nabla \tilde{\phi}_{l_k}) d\widetilde{\mathcal{W}}_{l_k} \right|^2 &= C \widetilde{\mathbb{E}} \int_0^T \|h(\tilde{u}_{l_k}, \nabla \tilde{\phi}_{l_k})\|_{L_2(\mathcal{H}; L_x^2)}^2 dt \\ &\leq CK_0 \left(1 + \widetilde{\mathbb{E}} \int_0^T \|\tilde{u}_{l_k}\|_{L_x^2}^2 + \|\nabla \tilde{\phi}_{l_k}\|_{L_x^2}^2 dt \right) \\ &\leq C. \end{aligned} \quad (4.11)$$

Proposition 4.1 (4.6) gives $(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}) \rightarrow (\tilde{u}, \tilde{\phi})$, $\widetilde{\mathbb{P}}$ -a.s. in $C([0, T]; \mathbb{H}_w)$, using the dominated convergence theorem we get for all $t \in [0, T]$, $\widetilde{\mathbb{P}}$ -a.s.

$$\lim_{k \rightarrow \infty} \int_0^t |(\tilde{u}_{l_k} - \tilde{u}, v)|^2 ds \rightarrow 0, \quad \lim_{k \rightarrow \infty} \int_0^t |(\tilde{\phi}_{l_k} - \tilde{\phi}, \bar{v})|^2 ds \rightarrow 0,$$

which combines with the bound (4.7), we obtain

$$\widetilde{\mathbb{E}} \int_0^T |(\tilde{u}_{l_k} - \tilde{u}, v)|^2 dt \rightarrow 0, \quad \widetilde{\mathbb{E}} \int_0^T |(\tilde{\phi}_{l_k} - \tilde{\phi}, \bar{v})|^2 dt \rightarrow 0.$$

By the same way, we have

$$\widetilde{\mathbb{E}} |(\tilde{u}_{l_k}(0) - \tilde{u}(0), v)|^2 \rightarrow 0, \quad \widetilde{\mathbb{E}} |(\tilde{\phi}_{l_k}(0) - \tilde{\phi}(0), \bar{v})|^2 \rightarrow 0.$$

For the linear term, since $(\tilde{u}_{l_k}, \tilde{\mu}_{l_k}) \rightarrow (\tilde{u}, \tilde{\mu})$, $\widetilde{\mathbb{P}}$ -a.s. in $(L_t^2(\mathbb{H}^1 \times H^1))_w$, we infer that

$$\int_0^t \langle A_0 \tilde{u}_{l_k}, v \rangle ds \rightarrow \int_0^t \langle A_0 \tilde{u}, v \rangle ds, \quad \int_0^t \langle A_1 \tilde{\mu}_{l_k}, \bar{v} \rangle ds \rightarrow \int_0^t \langle A_1 \tilde{\mu}, \bar{v} \rangle ds.$$

In addition, we have

$$\widetilde{\mathbb{E}} \left| \int_0^T \langle A_0 \tilde{u}_{l_k}, v \rangle dt \right|^2 \leq C \|v\|_{\mathbb{H}^1}^2 \widetilde{\mathbb{E}} \int_0^T \|\nabla \tilde{u}_{l_k}\|_{L_x^2}^2 dt \leq C, \quad (4.12)$$

$$\widetilde{\mathbb{E}} \left| \int_0^T \nu_2 \langle A_1 \tilde{\mu}_{l_k}, \bar{v} \rangle dt \right|^2 \leq C \|\bar{v}\|_{H^1}^2 \widetilde{\mathbb{E}} \int_0^T \|\nabla \tilde{\mu}_{l_k}\|_{L_x^2}^2 dt \leq C. \quad (4.13)$$

All the constants C in above are independent of l_k .

Next, define the functional

$$\mathbb{F}_1(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \widetilde{\mathcal{W}}_{l_k}, v) = (\tilde{u}_{l_k}(0), v) - \int_0^t \langle A_0 \tilde{u}_{l_k}, v \rangle ds - \int_0^t B_0(\tilde{u}_{l_k}, \tilde{u}_{l_k}, v) ds$$

$$\begin{aligned}
& + \int_0^t B_1(\nu_1 A_1 \tilde{\phi}_{l_k}, \tilde{\phi}_{l_k}, v) ds + \left\langle \int_0^t h(\tilde{u}_{l_k}, \nabla \tilde{\phi}_{l_k}) d\tilde{\mathcal{W}}_{l_k}, v \right\rangle, \\
\mathbb{F}_2(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \bar{v}) &= (\tilde{\phi}_{l_k}(0), \bar{v}) - \int_0^t \nu_2 \langle A_1 \tilde{\mu}_{l_k}, \bar{v} \rangle ds - \int_0^t B_2(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \bar{v}) ds
\end{aligned}$$

and

$$\begin{aligned}
\bar{\mathbb{F}}_1(\tilde{u}, \tilde{\phi}, \tilde{\mathcal{W}}, v) &= (\tilde{u}(0), v) - \int_0^t \langle A_0 \tilde{u}, v \rangle ds - \int_0^t B_0(\tilde{u}, \tilde{u}, v) ds \\
&+ \int_0^t B_1(\nu_1 A_1 \tilde{\phi}, \tilde{\phi}, v) ds + \left\langle \int_0^t h(\tilde{u}, \nabla \tilde{\phi}) d\tilde{\mathcal{W}}, v \right\rangle, \\
\bar{\mathbb{F}}_2(\tilde{u}, \tilde{\phi}, \bar{v}) &= (\tilde{\phi}(0), \bar{v}) - \int_0^t \nu_2 \langle A_1 \tilde{\mu}, \bar{v} \rangle ds - \int_0^t B_2(\tilde{u}, \tilde{\phi}, \bar{v}) ds.
\end{aligned}$$

Combining the bounds (4.8)-(4.13) and the convergence properties, using the Vitali convergence theorem, we conclude that for any set $B \in [0, T]$

$$\mathbb{E} \int_0^T 1_B(\mathbb{F}_1(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mathcal{W}}_{l_k}, v) - \bar{\mathbb{F}}_1(\tilde{u}, \tilde{\phi}, \tilde{\mathcal{W}}, v)) dt \rightarrow 0, \quad (4.14)$$

$$\mathbb{E} \int_0^T 1_B(\mathbb{F}_2(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mathcal{W}}_{l_k}, \bar{v}) - \bar{\mathbb{F}}_2(\tilde{u}, \tilde{\phi}, \tilde{\mathcal{W}}, \bar{v})) dt \rightarrow 0, \quad (4.15)$$

where $1_{\{\cdot\}}$ stands for the indicator function.

Since $(\tilde{u}_{l_k}, \tilde{\phi}_{l_k})$ satisfies the system (2.2) in the weak sense, that is,

$$((\tilde{u}_{l_k}, v), (\tilde{\phi}_{l_k}, \bar{v})) = (\mathbb{F}_1(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mathcal{W}}_{l_k}, v), \mathbb{F}_2(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}, \tilde{\mathcal{W}}_{l_k}, \bar{v})), \quad \tilde{\mathbb{P}}\text{-a.s.}$$

From (4.14)-(4.15), we conclude that for any $t \in [0, T]$, $v \in H^2$, $\bar{v} \in H^1$ and $\tilde{\mathbb{P}}\text{-a.s.}$

$$(\tilde{u}, v) = \bar{\mathbb{F}}_1(\tilde{u}, \tilde{\phi}, \tilde{\mathcal{W}}, v), \quad (\tilde{\phi}, \bar{v}) = \bar{\mathbb{F}}_2(\tilde{u}, \tilde{\phi}, \bar{v}).$$

Finally, by the density argument, the above equality holds for any $(v, \bar{v}) \in H^1 \times H^1$. We obtain that the $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}, \tilde{u}, \tilde{\phi}, \tilde{\mathcal{W}})$ is a martingale solution of the system (1.1)-(1.4).

Step 3. Finally, we show (4.5) holds. Since (u, ϕ) is a global pathwise solution with the initial data (u_0, ϕ_0) and the fixed stochastic basis $(\Omega, \mathcal{F}, \mathbb{P}, \mathcal{W})$, also $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}, \tilde{u}, \tilde{\phi}, \tilde{\mathcal{W}})$ is a martingale solution of the system (1.1)-(1.4) with the initial data Λ , where $\Lambda = \mathbb{P} \circ (u_0, \phi_0)^{-1}$, and the solution of system (1.1)-(1.4) is pathwise unique, so unique in law, we deduce that (4.5) holds.

This completes the proof. \square

5. EXISTENCE OF INVARIANT MEASURE

In this section, we show the main result: existence of invariant measure. We say a probability measure ξ on $\mathcal{B}(X_w)$ is an invariant measure if

$$\int_X \mathbf{P}_t \varphi d\xi = \int_X \varphi d\xi, \quad \text{for all } t \geq 0, \quad \varphi \in C_b(X_w).$$

Note that, if the space X is separable, then $\mathcal{B}(X_w) = \mathcal{B}(X)$.

For any bounded Borel function $\varphi \in \mathcal{B}_b(\mathbb{H})$, we define

$$(\mathbf{P}_t \varphi)(U_0) = \mathbb{E}[\varphi(U(t, U_0))], \quad U_0 \in \mathbb{H}, \quad (5.1)$$

where $U(t, U_0) = (u(t, u_0), \phi(t, \phi_0))$ is the pathwise solution of system (1.1)-(1.5) starting from the initial data $U_0 = (u_0, \phi_0)$, which is a Markov process. The proof of Markov property is standard, see [1, 4]. We also have $\mathbf{P}_{t+s}\varphi(\cdot) = \mathbf{P}_t\mathbf{P}_s\varphi(\cdot)$ for $t, s \geq 0$.

Lemma 5.1. *Let $\varphi : \mathbb{H} \rightarrow \mathbb{R}$ be the bounded sequentially weakly continuous function. Then, we have*

$$\mathbf{P}_t : C_b(\mathbb{H}_w) \rightarrow C_b(\mathbb{H}_w),$$

particularly, if $U_{0,l} \rightarrow U_0$ weakly in \mathbb{H} , then

$$(\mathbf{P}_t\varphi)(U_{0,l}) \rightarrow (\mathbf{P}_t\varphi)(U_0).$$

Proof. The proof is inspired by [4]. From the Theorem 3.1, corresponding to the initial data $U_{0,l}$, there exists a sequence \mathbb{H} -valued \mathcal{F}_t progressive measurable processes $U_l = (u_l, \phi_l)$ which is the solution of system (1.1)-(1.5). Using the Proposition 4.1, there exist a new probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, a new subsequence $(\tilde{u}_{l_k}, \tilde{\phi}_{l_k})$ and the processes $(\tilde{u}, \tilde{\phi})$, such that

$$(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}) \text{ and } (u_{l_k}, \phi_{l_k}), (\tilde{u}, \tilde{\phi}) \text{ and } (u, \phi) \text{ have the same joint distribution in } \mathcal{X}_{u,\phi},$$

and

$$(\tilde{u}_{l_k}, \tilde{\phi}_{l_k}) \rightarrow (\tilde{u}, \tilde{\phi}), \text{ in the topology of } \mathcal{X}_{u,\phi}, \tilde{\mathbb{P}}\text{-a.s.}$$

where $\tilde{U} = (\tilde{u}, \tilde{\phi})$ satisfies the system (2.2).

Since $\tilde{U}_{l_k} := (\tilde{u}_{l_k}, \tilde{\phi}_{l_k}) \rightarrow \tilde{U}$ weakly in $C([0, T]; \mathbb{H}_w)$, this together with the fact that φ is a bounded sequentially weakly continuous function, yields

$$\varphi(\tilde{U}_{l_k}) \rightarrow \varphi(\tilde{U}), \text{ in } \mathbb{R}, \tilde{\mathbb{P}}\text{-a.s.}$$

Since the function φ is bounded, we infer from the dominated convergence theorem

$$\tilde{\mathbb{E}} \left[\varphi(\tilde{U}_{l_k}(t; \tilde{U}_{0,l_k})) \right] \rightarrow \tilde{\mathbb{E}} \left[\varphi(\tilde{U}(t, \tilde{U}_0)) \right], \text{ as } k \rightarrow \infty.$$

Since the processes \tilde{U}_{l_k} and U_{l_k} , \tilde{U} and U have the same distribution which lead to

$$\tilde{\mathbb{E}} \left[\varphi(\tilde{U}_{l_k}(t; \tilde{U}_{0,l_k})) \right] = \mathbb{E} [\varphi(U_{l_k}(t; U_{0,l_k}))] = (\mathbf{P}_t\varphi)(U_{0,l_k}), \quad (5.2)$$

$$\tilde{\mathbb{E}} \left[\varphi(\tilde{U}(t; \tilde{U}_0)) \right] = \mathbb{E} [\varphi(U(t; U_0))] = (\mathbf{P}_t\varphi)(U_0). \quad (5.3)$$

Therefore, by (5.2), (5.3), we have

$$\lim_{k \rightarrow \infty} (\mathbf{P}_t\varphi)(U_{0,l_k}) = (\mathbf{P}_t\varphi)(U_0).$$

Using the sub-subsequence argument, we have the original sequence

$$\lim_{k \rightarrow \infty} (\mathbf{P}_t\varphi)(U_{0,l}) = (\mathbf{P}_t\varphi)(U_0).$$

Obviously, $\mathbf{P}_t\varphi$ from \mathbb{H} into \mathbb{R} is bounded. This completes the proof. \square

Lemma 5.2. *Let $U(t)$ be the solution of system (2.2) with the initial data U_0 . Then, there exists T_0 such that for every $\varepsilon > 0$, there exists $R > 0$ such that*

$$\sup_{T > T_0} \frac{1}{T} \int_0^T (\mathbf{P}_t^* \delta_{U_0})(\mathbb{H} \setminus B_R) dt \leq \varepsilon,$$

where $B_R := \{U(t) \in \mathbb{H} : \|U(t)\|_{\mathbb{H}} \leq R\}$.

Proof. Step 1. We first show that there exists constant C independent of t such that

$$\int_0^t \mathbb{E} \|U(s)\|_{\mathbb{H}}^2 ds \leq C(1+t).$$

Using the Itô formula to function $\|u\|_{L_x^2}^2$, by (2.1), we have

$$\begin{aligned} \|u\|_{L_x^2}^2 + 2 \int_0^t \|\nabla u\|_{L_x^2}^2 ds &= \|u_0\|_{L_x^2}^2 + 2 \int_0^t B_1(\nu_1 A_1 \phi, \phi, u) ds + 2 \int_0^t (h(u, \nabla \phi) d\mathcal{W}, u) \\ &\quad + \int_0^t \|h(u, \nabla \phi)\|_{L_2(\mathcal{H}; L_x^2)}^2 ds. \end{aligned} \quad (5.4)$$

Taking inner product with μ on both sides of equation (2.2)₂, we have

$$\begin{aligned} \nu_1 \|\nabla \phi\|_{L_x^2}^2 + 2\kappa(F(\phi), 1) + 2\nu_2 \int_0^t \|\nabla \mu\|_{L_x^2}^2 ds + 2 \int_0^t B_2(u, \phi, \mu) ds \\ = \nu_1 \|\nabla \phi_0\|_{L_x^2}^2 + 2\kappa(F(\phi_0), 1). \end{aligned} \quad (5.5)$$

Taking the sum of (5.4)-(5.5), using the relation

$$B_2(u, \phi, \mu) = B_1(\nu_1 A_1 \phi, \phi, u),$$

we have

$$\begin{aligned} \|u\|_{L_x^2}^2 + \nu_1 \|\nabla \phi\|_{L_x^2}^2 + 2\kappa(F(\phi), 1) + 2 \int_0^t \|\nabla u\|_{L_x^2}^2 + \nu_2 \|\nabla \mu\|_{L_x^2}^2 ds \\ = \|u_0\|_{L_x^2}^2 + \nu_1 \|\nabla \phi_0\|_{L_x^2}^2 + 2\kappa(F(\phi_0), 1) + 2 \int_0^t (h(u, \nabla \phi) d\mathcal{W}, u) \\ + \int_0^t \|h(u, \nabla \phi)\|_{L_2(\mathcal{H}; L_x^2)}^2 ds. \end{aligned} \quad (5.6)$$

Taking expectation on both sides of (5.6), we arrive

$$\begin{aligned} \mathbb{E} \left(\|u\|_{L_x^2}^2 + \nu_1 \|\nabla \phi\|_{L_x^2}^2 + 2\kappa(F(\phi), 1) + 2 \int_0^t \|\nabla u\|_{L_x^2}^2 + \nu_2 \|\nabla \mu\|_{L_x^2}^2 ds \right) \\ = \|u_0\|_{L_x^2}^2 + \nu_1 \|\nabla \phi_0\|_{L_x^2}^2 + 2\kappa(F(\phi_0), 1) + \mathbb{E} \int_0^t \|h(u, \nabla \phi)\|_{L_2(\mathcal{H}; L_x^2)}^2 ds. \end{aligned} \quad (5.7)$$

The stochastic term is a square integral martingale which its expectation vanished. Indeed,

$$\mathbb{E} \int_0^t \|h(u, \nabla \phi)\|_{L_2(\mathcal{H}; L_x^2)}^2 ds \leq K_0 \mathbb{E} \int_0^t 1 + (\|u\|_{L_x^2}^2 + \|\nabla \phi\|_{L_x^2}^2) ds \leq C. \quad (5.8)$$

By (5.8) again, we get from (5.7)

$$\begin{aligned} \mathbb{E} \left(\|u\|_{L_x^2}^2 + \nu_1 \|\nabla \phi\|_{L_x^2}^2 + 2\kappa C_1 \|\phi\|_{L_x^4}^4 + 2 \int_0^t \|\nabla u\|_{L_x^2}^2 + \nu_2 \|\nabla \mu\|_{L_x^2}^2 ds \right) \\ \leq \|u_0\|_{L_x^2}^2 + \nu_1 \|\nabla \phi_0\|_{L_x^2}^2 + 2\kappa(F(\phi_0), 1) \\ + K_0 \mathbb{E} \int_0^t 1 + (\|u\|_{L_x^2}^2 + \|\nabla \phi\|_{L_x^2}^2) ds + 2\kappa C_2 \mathbb{E} \|\phi\|_{L_x^2}^2 \\ \leq \|u_0\|_{L_x^2}^2 + \nu_1 \|\nabla \phi_0\|_{L_x^2}^2 + 2\kappa(F(\phi_0), 1) \end{aligned}$$

$$\begin{aligned}
& + K_0 \mathbb{E} \int_0^t 1 + (\|u\|_{L_x^2}^2 + \|\nabla \phi\|_{L_x^2}^2) ds + 2\kappa C_2 |\mathcal{D}|^{\frac{1}{2}} \mathbb{E} \|\phi\|_{L_x^4}^2 \\
& \leq \|u_0\|_{L_x^2}^2 + \nu_1 \|\nabla \phi_0\|_{L_x^2}^2 + 2\kappa(F(\phi_0), 1) + 2\delta_0 \kappa C_2 |\mathcal{D}|^{\frac{1}{2}} \mathbb{E} \|\phi\|_{L_x^4}^4 + C(\delta_0, \kappa, C_2, \mathcal{D}) + K_0 t \\
& + K_0 \mathbb{E} \int_0^t \|u\|_{L_x^2}^2 + \|\nabla \phi\|_{L_x^2}^2 ds,
\end{aligned} \tag{5.9}$$

where we choose δ_0 small such that $2\delta_0 \kappa C_2 |\mathcal{D}|^{\frac{1}{2}} \leq 2\kappa C_1$.

The key point is to control the last term $\mathbb{E} \int_0^t \|\nabla \phi\|_{L_x^2}^2 ds$. Taking inner product with ϕ with (2.2)₃, then integral in time t and taking expectation, we get

$$\nu_1 \mathbb{E} \int_0^t \|\nabla \phi\|_{L_x^2}^2 ds + \kappa \mathbb{E} \int_0^t (F(\phi), 1) ds = \mathbb{E} \int_0^t (\mu, \phi) ds. \tag{5.10}$$

Note that, by (2.2)₃ and the boundary condition (1.2), we get

$$\langle \mu \rangle = \kappa \langle f(\phi) \rangle.$$

Using the Hölder inequality and the Schwartz inequality, we get

$$\begin{aligned}
\langle f(\phi) \rangle &= \frac{1}{|\mathcal{D}|} \int_{\mathcal{D}} f(\phi) dx = \frac{1}{|\mathcal{D}|} \int_{\mathcal{D}} C_1 \phi^3 - C_2 \phi dx \\
&\leq \frac{1}{|\mathcal{D}|} \left(C_1 |\mathcal{D}|^{\frac{1}{4}} \int_{\mathcal{D}} \phi^4 dx^{\frac{3}{4}} + C_2 |\mathcal{D}|^{\frac{1}{2}} \int_{\mathcal{D}} \phi^2 dx^{\frac{1}{2}} \right) \\
&\leq \frac{C_1 \delta}{|\mathcal{D}|} \int_{\mathcal{D}} \phi^4 dx + \frac{C_2 \delta}{|\mathcal{D}|} \int_{\mathcal{D}} \phi^2 dx + \frac{C}{\delta},
\end{aligned} \tag{5.11}$$

where δ is constant selected later.

Using the Hölder inequality, the Schwartz inequality and boundary condition (1.2), estimate (5.11), the right hand term of (5.10) could be controlled as follows

$$\begin{aligned}
\mathbb{E} \int_0^t (\mu, \phi) ds &\leq \mathbb{E} \int_0^t \|\mu\|_{L_x^2} \|\phi\|_{L_x^2} ds \leq \frac{1}{2\delta} \mathbb{E} \int_0^t \|\mu\|_{L_x^2}^2 ds + \frac{\delta}{2} \mathbb{E} \int_0^t \|\phi\|_{L_x^2}^2 ds \\
&\leq \frac{C}{2\delta} \mathbb{E} \int_0^t \|\nabla \mu\|_{L_x^2}^2 + \|\langle \mu \rangle\|_{L_x^2}^2 ds + \frac{C\delta}{2} \mathbb{E} \int_0^t \|\nabla \phi\|_{L_x^2}^2 + \|\langle \phi \rangle\|_{L_x^2}^2 ds \\
&\leq \frac{C}{2\delta} \mathbb{E} \int_0^t \|\nabla \mu\|_{L_x^2}^2 + \frac{C_1 \delta}{\sqrt{|\mathcal{D}|}} \int_{\mathcal{D}} \phi^4 dx + \frac{C(\mathcal{D})}{\delta} ds \\
&\quad + \frac{C\delta}{2} \mathbb{E} \int_0^t \|\nabla \phi\|_{L_x^2}^2 + \|\langle \phi_0 \rangle\|_{L_x^2}^2 ds + \frac{C_2 \delta}{|\mathcal{D}|} \mathbb{E} \int_0^t \|\nabla \phi\|_{L_x^2}^2 + \|\langle \phi_0 \rangle\|_{L_x^2}^2 ds.
\end{aligned} \tag{5.12}$$

Here, choosing δ, K_0 small such that

$$\max \left\{ \frac{C\delta}{2} + \frac{C_2 \delta^2}{2|\mathcal{D}|} + K_0, \frac{C_1 \delta}{\sqrt{|\mathcal{D}|}} \right\} \leq \min\{\nu_1, \kappa\},$$

then choosing $2\nu_2 > \frac{C}{2\delta}$. The Dirichlet boundary condition gives

$$\|u\|_{L_x^2} \leq C \|\nabla u\|_{L_x^2}, \tag{5.13}$$

where constant C depends on \mathcal{D} . Combining (5.9), (5.10) and (5.12), (5.13), we get

$$\begin{aligned} & \mathbb{E} \left(\|u\|_{L_x^2}^2 + C \int_0^t \|\phi\|_{L_x^4}^4 ds + C \int_0^t \|\nabla \phi\|_{L_x^2}^2 ds \right. \\ & \quad \left. + \int_0^t (2 - K_0 C) \|\nabla u\|_{L_x^2}^2 + (2\nu_2 - \frac{C}{2\delta}) \|\nabla \mu\|_{L_x^2}^2 ds \right) \\ & \leq C(C_1, C_2, \kappa, \nu_1, \delta, \mathcal{D}) + C(C_1, C_2, \kappa, \nu_1, \delta, \mathcal{D})t + C_2 \kappa \mathbb{E} \int_0^t \|\phi\|_{L_x^2}^2 ds \\ & \quad + \|u_0\|_{L_x^2}^2 + \nu_1 \|\nabla \phi_0\|_{L_x^2}^2 + 2\kappa(F(\phi_0), 1). \end{aligned} \quad (5.14)$$

Using the Sobolev embedding $H^1 \hookrightarrow L_x^4 \hookrightarrow L_x^2$, we have

$$|2\kappa(F(\phi_0), 1)| \leq C(\|\phi_0\|_{L_x^4}^4 + \|\phi_0\|_{L_x^2}^2) \leq C(1 + \|\phi_0\|_{H^1}^4).$$

According to the Condition **C.4**, by (5.14) and above estimate, we have

$$\begin{aligned} & \mathbb{E} \left(C \int_0^t \|\phi\|_{L_x^4}^4 ds + C \int_0^t \|\phi\|_{H^1}^2 ds + \int_0^t (2 - K_0 C) \|\nabla u\|_{L_x^2}^2 ds \right) \\ & \leq C(C_1, C_2, C_3, \kappa, \nu_1, \delta, \mathcal{D}) + C(C_1, C_2, \kappa, \nu_1, \delta, \mathcal{D})t + (C + C_2 \kappa) \mathbb{E} \int_0^t \|\phi\|_{L_x^2}^2 ds. \end{aligned} \quad (5.15)$$

Since

$$\begin{aligned} (C + C_2 \kappa) \mathbb{E} \int_0^t \|\phi\|_{L_x^2}^2 ds & \leq C(C_2, \kappa, \mathcal{D}) \mathbb{E} \int_0^t \|\phi\|_{L_x^4}^2 ds \\ & \leq \delta_1 C(C_2, \kappa, \mathcal{D}) \mathbb{E} \int_0^t \|\phi\|_{L_x^4}^4 ds + C(\delta_1, C_2, \kappa, \mathcal{D})t. \end{aligned} \quad (5.16)$$

Choosing δ_1 small, combining (5.15), (5.16) and the Condition **C.3**, we have

$$\int_0^t \mathbb{E} \left(\|\nabla u\|_{L_x^2}^2 + \|\phi\|_{H^1}^2 \right) ds \leq C + Ct,$$

where constant C depends on $(C_1, C_2, C_3, \kappa, \nu_1, \nu_2, \mathcal{D}, K_0)$ but independent of t .

By (5.13) again, we conclude

$$\int_0^t \mathbb{E} \left(\|u\|_{L_x^2}^2 + \|\phi\|_{H^1}^2 \right) ds \leq C + Ct, \quad (5.17)$$

where constant C depends on $(C_1, C_2, C_3, \kappa, \nu_1, \nu_2, \mathcal{D}, K_0)$ but independent of t .

Step 2. By the Chebyshev inequality and (5.17),

$$\begin{aligned} \frac{1}{T} \int_0^T (\mathbf{P}_t^* \delta_{U_0})(\mathbb{H} \setminus B_R) dt &= \frac{1}{T} \int_0^T \mathbb{P}\{\|U(t)\|_{\mathbb{H}} > R\} dt \\ &\leq \frac{1}{R^2 T} \int_0^T \mathbb{E} \|U(t)\|_{\mathbb{H}}^2 dt \\ &\leq \frac{C + CT}{R^2 T}. \end{aligned}$$

This completes the proof. \square

Theorem 5.1. *Suppose that the Conditions C.1-C.4 hold, then the transition semigroup defined by (5.1) has an invariant measure.*

Proof. Combining the Lemmas 5.1, 5.2, Theorem 5.1 follows from the Maslowski and Seidler theory, Proposition 3.1 in [23]. \square

Remark 5.1. We believe that the Theorem 5.1 could be extended to the case of equation (1.1)₂ also being forced by a multiplicative noise. For such a system, we given the well-posedness of solution in paper [27] with extra boundary condition corresponding to the phenomenon of vesicle membranes. Here, to use the stochastic compactness argument, only minor modification is required, that is, $\phi \in C_t^{\tilde{\alpha}}(H^{-1})$, \mathbb{P} a.s. for $\tilde{\alpha}$ less than $\frac{1}{2}$ strictly.

6. APPENDIX

At the beginning, we introduce lemma used frequently for controlling the nonlinear term.

Lemma 6.1. [29, Lemma 3] *Let $r^* = \begin{cases} \frac{dr}{d-r}, & \text{if } r < d, \\ \text{any finite nonnegative real number,} & \text{if } r = d, \\ \infty, & \text{if } r > d, \end{cases}$*

where $d = 2, 3$ is the dimension. For $1 \leq r \leq \infty, 1 \leq s \leq \infty$, if $\frac{1}{r} + \frac{1}{s} \leq 1$, and $\frac{1}{r^} + \frac{1}{s} = \frac{1}{t}$, $f \in H^{1,r}$ and $g \in H^{-1,s}$, then $fg \in H^{-1,t}$, that is,*

$$\|fg\|_{H^{-1,t}} \leq \|f\|_{H^{1,r}} \|g\|_{H^{-1,s}}.$$

To establish the compactness property of a family of solutions, the following Aubin-Lions lemma and Skorokhod-Jakubowsk representation theorem are commonly used. For more background we recommend Corollary 5 of [28] and Theorem 2.4 of [7]. The Vitali convergence theorem is applied to identify the limit (see for example, Chapter 3 of [22]).

Lemma 6.2. (The Aubin-Lions Lemma) *Suppose that $X_1 \subset X_0 \subset X_2$ are Banach spaces and X_1 and X_2 are reflexive and the embedding of X_1 into X_0 is compact. Then for any $1 \leq p < \infty, 0 < \alpha < 1$, the embedding*

$$L_t^p X_1 \cap C_t^\alpha X_2 \hookrightarrow L_t^p X_0, \quad L_t^\infty X_1 \cap C_t^\alpha X_2 \hookrightarrow C_t((X_1)_w)$$

and

$$L_t^p X_1 \cap W_t^{\alpha,r} X_2 \hookrightarrow L_t^p X_0,$$

where $\alpha > 0$ if $r \geq p$, is compact, respectively.

Theorem 6.1. (The Vitali convergence theorem) *Let $p \geq 1, \{X_n\}_{n \geq 1} \in L^p$ and $X_n \rightarrow X$ in probability. Then, the following are equivalent*

- (1). $X_n \rightarrow X$ in L^p ;
- (2). the sequence $|X_n|^p$ is uniformly integrable;
- (3). $\mathbb{E}|X_n|^p \rightarrow \mathbb{E}|X|^p$.

Theorem 6.2. (The Skorokhod-Jakubowsk representative theorem) *Let X be a topological space. If the set of probability measures $\{\nu_n\}_{n \geq 1}$ on $\mathcal{B}(X)$ is tight, then there exists a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a sequence of random variables u_n, u such that theirs laws are ν_n, ν and $u_n \rightarrow u, \mathbb{P}$ -a.s. as $n \rightarrow \infty$.*

To pass the limit in stochastic integral of solution sequence, we take the following result from [9, Lemma 2.1].

Lemma 6.3. *Assume that G_ε is a sequence of X -valued $\mathcal{F}_t^\varepsilon$ predictable processes such that*

$$G_\varepsilon \rightarrow G, \text{ in probability in } L_t^2 L_2(\mathcal{H}; X),$$

and the cylindrical Wiener process sequence \mathcal{W}_ε satisfies

$$\mathcal{W}_\varepsilon \rightarrow \mathcal{W}, \text{ in probability in } C_t \mathcal{H}_0,$$

then,

$$\int_0^t G_\varepsilon d\mathcal{W}_\varepsilon \rightarrow \int_0^t G d\mathcal{W}, \text{ in probability in } L_t^2 X.$$

Lemma 6.4. *[18, Lemma 1.1] (The Gyöngy-Krylov Lemma) Let X be a complete separable metric space and suppose that $\{Y_n\}_{n \geq 0}$ is a sequence of X -valued random variables on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $\{\mathcal{L}_{m,n}\}_{m,n \geq 1}$ be the set of joint laws of $\{Y_n\}_{n \geq 1}$, that is*

$$\mathcal{L}_{m,n}(E) := \mathbb{P}\{(Y_n, Y_m) \in E\}, \quad E \in \mathcal{B}(X \times X).$$

Then $\{Y_n\}_{n \geq 1}$ converges in probability if and only if for every subsequence of the joint probability laws $\{\mu_{m_k, n_k}\}_{k \geq 1}$, there exists a further subsequence that converges weakly to a probability measure μ such that

$$\mathcal{L}\{(Y_1, Y_2) \in X \times X : Y_1 = Y_2\} = 1.$$

Proof of Step 3 of Theorem 3.1. Invoked by [9, 18], define by $\mathcal{L}_{n,m}$ the joint law

$(u_n, \phi_n, \mu_n; u_m, \phi_m, \mu_m; \mathcal{W})$ on the path space $\mathcal{X} = \mathcal{X}_u \times \mathcal{X}_\phi \times \mathcal{X}_\mu \times \mathcal{X}_u \times \mathcal{X}_\phi \times \mathcal{X}_\mu \times \mathcal{X}_\mathcal{W}$, where

$$\mathcal{X}_u := C_t(H^{-1}) \cap L_t^2 L_x^2, \quad \mathcal{X}_\phi := C_t(L_x^2) \cap L_t^2 H^1, \quad \mathcal{X}_\mu := L_t^2 L_x^2, \quad \mathcal{X}_\mathcal{W} := C_t(\mathcal{H}_0),$$

and $\{u_n, \phi_n; u_m, \phi_m\}_{n,m \geq 1}$ are two sequences of approximate solutions to system (2.2). Using the same argument as Lemma 4.1, we infer that,

the collection of joint laws $\{\mathcal{L}_{m,n}\}_{n,m \geq 1}$ is tight on \mathcal{X} .

For any subsequence $\{\mathcal{L}_{n_k, m_k}\}_{k \geq 1}$, there exists a measure \mathcal{L} such that $\{\mathcal{L}_{n_k, m_k}\}_{k \geq 1}$ converges weakly to \mathcal{L} . Applying the Skorokhod-Jakubowski representation theorem, we have a new probability space $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$ and \mathcal{X} -valued random variables

$$(\hat{u}_{n_k}, \hat{\phi}_{n_k}, \hat{\mu}_{n_k}; \hat{u}_{m_k}, \hat{\phi}_{m_k}, \hat{\mu}_{m_k}; \widehat{\mathcal{W}}_k) \text{ and } (\hat{u}_1, \hat{\phi}_1, \hat{\mu}_1; \hat{u}_2, \hat{\phi}_2, \hat{\mu}_2; \widehat{\mathcal{W}})$$

such that

$$\begin{aligned} \hat{\mathbb{P}}\{(\hat{u}_{n_k}, \hat{\phi}_{n_k}, \hat{\mu}_{n_k}; \hat{u}_{m_k}, \hat{\phi}_{m_k}, \hat{\mu}_{m_k}; \widehat{\mathcal{W}}_k) \in \cdot\} &= \mathcal{L}_{n_k, m_k}(\cdot), \\ \hat{\mathbb{P}}\{(\hat{u}_1, \hat{\phi}_1, \hat{\mu}_1; \hat{u}_2, \hat{\phi}_2, \hat{\mu}_2; \widehat{\mathcal{W}}) \in \cdot\} &= \mathcal{L}(\cdot) \end{aligned}$$

and

$$(\hat{u}_{n_k}, \hat{\phi}_{n_k}, \hat{\mu}_{n_k}; \hat{u}_{m_k}, \hat{\phi}_{m_k}, \hat{\mu}_{m_k}; \widehat{\mathcal{W}}_k) \rightarrow (\hat{u}_1, \hat{\phi}_1, \hat{\mu}_1; \hat{u}_2, \hat{\phi}_2, \hat{\mu}_2; \widehat{\mathcal{W}}), \quad \hat{\mathbb{P}}\text{-a.s.}$$

in the topology of \mathcal{X} . Analogously, this argument can be applied to both

$$(\hat{u}_{n_k}, \hat{\phi}_{n_k}, \widehat{\mathcal{W}}_k), (\hat{u}_1, \hat{\phi}_1, \widehat{\mathcal{W}}) \quad \text{and} \quad (\hat{u}_{m_k}, \hat{\phi}_{m_k}, \widehat{\mathcal{W}}_k), (\hat{u}_2, \hat{\phi}_2, \widehat{\mathcal{W}})$$

to show that $(\hat{u}_1, \hat{\phi}_1, \widehat{\mathcal{W}})$ and $(\hat{u}_2, \hat{\phi}_2, \widehat{\mathcal{W}})$ are two martingale solutions relative to the same stochastic basis $\hat{\mathcal{S}} := (\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}}, \{\hat{\mathcal{F}}_t\}_{t \geq 0}, \widehat{\mathcal{W}})$.

In addition, we have $\mathcal{L}_{n,m} \rightharpoonup \mathcal{L}$ where \mathcal{L} is defined by

$$\mathcal{L}(\cdot) = \widehat{\mathbb{P}}\{(\hat{u}_1, \hat{\phi}_1, \hat{\mu}_1; \hat{u}_2, \hat{\phi}_2, \hat{\mu}_2) \in \cdot\}.$$

The uniqueness implies that $\mathcal{L}\{(u_1, \phi_1, \mu_1; u_2, \phi_2, \mu_2) \in \mathcal{X} : (u_1, \phi_1, \mu_1) = (u_2, \phi_2, \mu_2)\} = 1$.

The both conditions of Lemma 6.4 are verified, we deduce that the sequence (u_n, ϕ_n, μ_n) defined on the original probability space $(\Omega, \mathcal{F}, \mathbb{P})$ converges \mathbb{P} -a.s. in the topology of $\mathcal{X}_u \times \mathcal{X}_\phi \times \mathcal{X}_\mu$ to processes (u, ϕ, μ) . \square

ACKNOWLEDGMENTS

Z. Qiu's research is supported by the CSC under grant No.201806160015.

REFERENCES

- [1] H. Bessaih, B. Ferrario. Invariant measures for stochastic damped 2D Euler equations. *Commun. Math. Phys.*, 377, (2020), 531-549.
- [2] H. Breckner. Galerkin approximation and the strong solution of the Navier-Stokes equation. *J. Appl. Math. Stoch. Anal.*, 13(3), (2000), 239-259.
- [3] Z. Brzeźniak, D. Gatarek. Martingale solutions and invariant measures for stochastic evolution equations in Banach spaces. *Stochastic Process. Appl.*, 84(2), (1999), 187-225.
- [4] Z. Brzeźniak, E. Motyl, M. Ondrejat. Invariant measure for the stochastic Navier-Stokes equations in unbounded 2D domains. *Ann. Probab.*, 45(5), (2017), 3145-3201.
- [5] Z. Brzeźniak, M. Ondrejot, J. Seidler. Invariant measures for stochastic nonlinear beam and wave equations. *J. Differential Equations*, 260, (2016), 4157-4179.
- [6] T. Blesgen. A generalization of the Navier-Stokes equations to two-phase flows. *J. Phys. D: Appl. Phys.*, 32(10), (1999), 1119.
- [7] G. Da Prato, J. Zabczyk. *Stochastic equations in infinite dimensions*, Cambridge university press, 1992.
- [8] G. Da Prato, J. Zabczyk. *Ergodicity for infinite dimensional systems*. Cambridge University Press, 1996.
- [9] A. Debussche, N. Glatt-Holtz, R. Temam. Local martingale and pathwise solutions for an abstract fluids model. *Phys. D: Nonlinear Phenomena*, 240(14-15), (2011), 1123-1144.
- [10] G. Deugoué, T.T. Medjo. Convergence of the solution of the stochastic 3D globally modified Cahn-Hilliard-Navier-Stokes equations, *J. Differential Equations*, 265(2), (2018), 545-592.
- [11] F. Flandoli. Dissipativity and invariant measures for stochastic Navier-Stokes equations. *NoDEA Nonlinear Differential Equations Appl.*, 1(4), (1994), 403-423.
- [12] F. Flandoli, D. Gatarek. Martingale and stationary solutions for stochastic Navier-Stokes equations. *Probab. Theory Relat. Fields*, 102(3), (1995), 367-391.
- [13] S. Frigeri, C.G. Gal, M. Grasselli. On nonlocal Cahn-Hilliard-Navier-Stokes systems in two dimensions, *J. Nonlinear Sci.*, 26(4), (2016), 847-893.
- [14] S. Frigeri, M. Grasselli, P. Krejci. Strong solutions for two-dimensional nonlocal Cahn-Hilliard-Navier-Stokes systems in two dimensions, *J. Differential Equations*, 255(9), (2013), 2587-2614.
- [15] S. Frigeri, E. Rocca, J. Sprekels. Optimal distributed control of a nonlocal Cahn-Hilliard-Navier-Stokes systems in two dimensions, *SIAM J. Control Optim.*, 54(1), (2015), 221-250.
- [16] C.G. Gal, M. Grasselli. Asymptotic behavior of a Cahn-Hilliard-Navier-Stokes system in 2D, *Ann. Ins. H. Poincaré Anal. Non Linéaire*, 27(1), (2010), 401-436.
- [17] C.G. Gal, M. Grasselli, A. Miranville. Cahn-Hilliard-Navier-Stokes systems with moving contact lines. *Calc. Var. Partial Differential Equations*, 55(3), (2016), 50.
- [18] I. Gyöngy, N. Krylov. Existence of strong solutions for Itô's stochastic equations via approximations. *Probab. Theory Relat. Fields*, 105(2), (1996), 143-158.
- [19] A. Giorgini, A. Miranville, R. Temam. Uniqueness and Regularity for the Navier-Stokes-Cahn-Hilliard System. *SIAM J. Math. Anal.*, 51(3), (2019), 2535-2574.

- [20] M.E. Gurtin, D. Polignone, J. Vinals. Two-phase binary fluids and immiscible fluids described by an order parameter. *Math. Models Meth. Appl. Sci.*, 6(06), (1996), 815-831.
- [21] M. Heida, J. Málek, K.R. Rajagopal. On the development and generalizations of Cahn-Hilliard equations within a thermodynamic framework. *Z. Angew. Math. Phys.*, 63(1), (2012), 145-169.
- [22] O. Kallenberg. *Foundations of Modern Probability: Second Edition*. New York, NY: Springer: Probability and Its Applications, 2002.
- [23] B. Maslowski, J. Seidler. On sequentially weakly Feller solutions to SPDE's. *Atti della Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei Mat. Appl.*, 10(2), (1999), 69-78.
- [24] J.C. Maxwell. On stresses in rarified gases arising from inequalities of temperature. *Philos. Trans. R. Soc.*, 170, (1879), 231-256.
- [25] T.T. Medjo. On the existence and uniqueness of solution to a stochastic 2D Cahn-Hilliard-Navier-Stokes model, *J. Differential Equations*, 263(2), (2017), 1028-1054.
- [26] J.C. Oxtoby, S.M. Ulam. On the existence of a measure invariant under a transformation. *Ann. Math.*, 560-566, (1939).
- [27] Z. Qiu, H. Wang. Large deviation principle for the 2D stochastic CahnHilliardNavierStokes equations. *Z. Angew. Math. Phys.*, 79, (2020), 1-29.
- [28] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Ann. Math. Pura. Appl.*, 146(1), (1986), 65-96.
- [29] J. Simon. Nonhomogeneous viscous incompressible fluids: existence of velocity, density, and pressure. *SIAM J. Math. Anal.*, 21(5), (1990), 1093-1117.
- [30] D. Wang, H. Wang. Global existence of martingale solutions to the three-dimensional stochastic compressible Navier-Stokes equations. *Differential and Integral Equations*, 28(11/12), (2015), 1105-1154.

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY,
WUHAN, 430074, CHINA.

E-mail address: zhqmath@163.com