

HIDDEN POSITIVITY AND A NEW APPROACH TO NUMERICAL COMPUTATION OF HAUSDORFF DIMENSION: HIGHER ORDER METHODS

RICHARD S. FALK AND ROGER D. NUSSBAUM

ABSTRACT. In [14], the authors developed a new approach to the computation of the Hausdorff dimension of the invariant set of an iterated function system or IFS. In this paper, we extend this approach to incorporate high order approximation methods. We again rely on the fact that we can associate to the IFS a parametrized family of positive, linear, Perron-Frobenius operators L_s , an idea known in varying degrees of generality for many years. Although L_s is not compact in the setting we consider, it possesses a strictly positive C^m eigenfunction v_s with eigenvalue $R(L_s)$ for arbitrary m and all other points z in the spectrum of L_s satisfy $|z| \leq b$ for some constant $b < R(L_s)$. Under appropriate assumptions on the IFS, the Hausdorff dimension of the invariant set of the IFS is the value $s = s_*$ for which $R(L_s) = 1$. This eigenvalue problem is then approximated by a collocation method at the extended Chebyshev points of each subinterval using continuous piecewise polynomials of arbitrary degree r . Using an extension of the Perron theory of positive matrices to matrices that map a cone K to its interior and explicit a priori bounds on the derivatives of the strictly positive eigenfunction v_s , we give rigorous upper and lower bounds for the Hausdorff dimension s_* , and these bounds converge rapidly to s_* as the mesh size decreases and/or the polynomial degree increases.

1. INTRODUCTION

In this paper, we continue previous work in finding rigorous estimates for the Hausdorff dimension of invariant sets for iterated function systems or IFS's. To describe the framework of the problem we are considering, we let $S \subset \mathbb{R}$ be a nonempty compact set, and for some positive integer m , let $\theta_p : S \rightarrow S$ and $g_p : S \rightarrow [0, \infty] \in C^m(S)$ for $1 \leq p \leq n < \infty$. If θ_p are contraction mappings, it is known that there exists a unique, compact, nonempty set $C \subset S$ such that $C = \cup_{p=1}^n \theta_p(C)$. The set C is called the invariant set for the IFS $\{\theta_p : 1 \leq p \leq n\}$.

For $s > 0$, define a bounded linear map $L_s : C(S) \rightarrow C(S)$, (often called a *Perron-Frobenius operator* or *linear transfer operator*) by

$$(1.1) \quad (L_s f)(t) = \sum_{p=1}^n [g_p(t)]^s f(\theta_p(t)), \quad t \in S.$$

Date: original submitted August 24, 2020; revised version February 26, 2021.

2000 Mathematics Subject Classification. Primary 11K55, 37C30; Secondary: 65J10.

Key words and phrases. Hausdorff dimension, positive transfer operators, continued fractions.

Under additional appropriate hypotheses (stated in the next section), L_s , considered as a map from $C^m(S) \mapsto C^m(S)$, has a strictly positive eigenfunction $v_s \in C^m(S)$ with algebraically simple eigenvalue $\lambda_s = R(L_s)$, the spectral radius of L_s . In addition, all other points z in the spectrum of L_s satisfy $|z| \leq b$ for some constant $b < R(L_s)$. A more precise statement of this result, along with other conclusions, is given in Theorem 2.1 in the next section. Note that in the C^m setting, L_s is, in general, not compact, has positive essential spectral radius and cannot be the limit in operator norm of a sequence of finite dimensional linear operators. These difficulties do not usually arise if L_s can be studied in a Banach space of complex analytic functions; and there is an extensive literature concerning the spectral theory of Perron-Frobenius operators which map a Banach space of analytic functions into itself. We prefer to work in the more general C^m setting so as to provide tools which also can be applied to some non-analytic examples, e.g., as in Section 5 of [14].

The aim of this paper is to derive an approximation scheme that allows us to estimate $R(L_s)$ by the spectral radius of an associated matrix \mathbf{L}_s which approximates the operator L_s in a weak sense and then to obtain rigorous bounds on the error $|R(L_s) - R(\mathbf{L}_s)|$. We then use this approximation scheme to estimate s_* , the unique number $s \geq 0$ such that $R(L_s) = 1$. Under appropriate assumptions, s_* equals the Hausdorff dimension of the invariant set associated to the IFS. This observation about Hausdorff dimension has been made, in varying degrees of generality by many authors. See, for example, [6], [7], [4], [9], [10], [13], [15], [18], [20], [19], [22], [23], [24], [25], [31], [30], [35], [37], [38], [39], [41], and [8]. There is also a large literature on the approximation of linear transform operators, not necessarily related to the computation of Hausdorff dimension, and often assuming the maps are analytic. We do not attempt to survey that literature, other than to cite one recent paper, [1], which has some connections to our work here, and contains many references to that literature.

In previous work, [14], the authors presented a new approach to the problem described in the preceding paragraph. We obtained rigorous upper and lower bounds for the Hausdorff dimension s_* , and these bounds exhibited second order convergence to s_* as the mesh size decreases. The approximate matrix was obtained by a collocation method using continuous piecewise linear functions, motivated by the fact that if such functions are nonnegative at the mesh points, they are nonnegative at all points of the interval in which they are defined. This property leads to nonnegative matrix approximations of the operator L_s . One would like these matrices to mimic the properties of the continuous operator L_s , which means they should satisfy the conclusions of the Perron theorem for positive matrices (matrices with strictly positive entries), i.e., they should have an eigenvalue of multiplicity one equal to the spectral radius of the matrix with corresponding positive eigenvector and all other eigenvalues of the matrix should have modulus less than the spectral radius. This is not true for nonnegative matrices, however, unless they have an additional property. One such property that would guarantee this is that the matrix \mathbf{L}_s be *primitive*, i.e., there exists a positive integer p such that \mathbf{L}_s^p is a positive matrix. Note that if \mathbf{L}_s is *irreducible*, then the first two properties hold, but there can be other eigenvalues of the same modulus as the spectral radius. Unfortunately,

the approximation scheme used led to matrices which are neither primitive nor irreducible. The remedy to obtain the desired properties was to note that the cone K of nonnegative vectors is not the natural cone in which such matrices should be studied. Using a more general notion of positivity of an operator L in which L maps a cone K into itself, one can still obtain the conclusions of the Perron theorem. This is important since we use the spectral radius of the approximate matrix \mathbf{L}_s to approximate the spectral radius of L_s and the fact that there is a single dominant eigenvalue enables us to calculate it efficiently using some variant of the power method.

In this paper, we analyze a similar method obtained by approximation using higher order piecewise polynomials. As we shall see, the matrices resulting from the approximation scheme appear to be even more problematic, since they are not even nonnegative. Despite this fact, the use of an alternative cone, in place of the standard cone of nonnegative vectors, allows us to show that the conclusions of the classical Perron theorem also hold for the matrices of this paper. There is a substantial abstract theory which has been developed for finite dimensional linear operators which are positive in the sense that they map a cone into itself. The survey paper [43], references in [43], and appendices A and B in [27] provide a good starting point. However, the difficulty lies in finding such a cone that fits the application under study. We use the term *hidden positivity* to call attention to the fact that we are able to find such a cone for the approximate operators developed in this paper.

The cone we use is easiest to describe in the case of continuous, piecewise linear functions, and is defined as follows. On the interval $[0, 1]$, for fixed integer N , let $h = 1/N$ and $x_i = ih$, $i = 0, 1, \dots, N$. The space of continuous, piecewise linear functions is just the finite dimensional space of continuous functions that restricted to each subinterval $[x_i, x_{i+1}]$ are linear functions. Since a function w in this space is completely determined by its values $w_i = w(x_i)$, $i = 0, 1, \dots, N$ (the *degrees of freedom* of w), we can also view w as the vector $[w_0, \dots, w_N]$. For any integer $M > 0$, we then define the cone K_M by

$$K_M = \{w : w_i \leq \exp(M|x_i - x_j|)w_j, \quad i, j = 0, 1, \dots, N\}.$$

The cone for higher order piecewise polynomials is similar, but its description is more involved because of the more complicated nature of the degrees of freedom of such functions. The details are provided in Section 3.

One technical difference between piecewise linear functions and higher order piecewise polynomials is that in order to obtain the results described above, we must consider approximations $\mathbf{L}_{s,\nu}$ of the matrix L_s^ν , where ν depends on the degree r of the piecewise polynomial approximation. As we observe in the next section, the operator L_s^ν has the same form as L_s , i.e.,

$$(L_s^\nu f)(t) = \sum_{\omega \in \Omega_\nu} [g_\omega(t)]^s f(\theta_\omega(t)),$$

where for $\nu \geq 1$, $\Omega_\nu = \{\omega = (p_1, p_2, \dots, p_\nu) : 1 \leq p_j \leq n \text{ for } 1 \leq j \leq \nu\}$; and for $\omega = (p_1, p_2, \dots, p_\nu) \in \Omega_\nu$,

$$\theta_\omega(t) = (\theta_{p_1} \circ \theta_{p_2} \circ \dots \circ \theta_{p_\nu})(t),$$

and $g_\omega(t)$ is defined in the next section. We note that under the weaker assumption that θ_ω is a contraction mapping for all $\omega \in \Omega_\nu$, there exists a unique compact set C such that $C = \cup_{\omega \in \Omega_\nu} \theta_\omega(C)$ and that necessarily $C = \cup_{p=1}^n \theta_p(C)$. By using the matrix $\mathbf{L}_{s,\nu}$, one reduces the domain of the operator to a finite set of subintervals whose total length is much less than the original length of the domain, resulting in many fewer mesh points. The downside, however, is that this advantage is completely offset by the increase in the number of terms in the operator L_s for each time the map is iterated, i.e., from n to n^ν . We note that since we have not found any case where the method fails if we do not iterate the matrix, we conjecture that this extra condition is an artifact of the method of proof, and not the method itself.

To obtain the conclusions of the Perron theorem, the key result is to show that for some $0 < M' < M$,

$$(1.2) \quad \mathbf{L}_{s,\nu}(K_M \setminus \{0\}) \subset K_{M'} \setminus \{0\}.$$

This enables us to apply results from the literature on mappings of a cone to itself to obtain the desired conclusions. Details of this connection, along with references to the relevant literature, are described in Section 4.

A main goal of our approach, in addition to proposing a new approximation scheme, is to provide rigorous upper and lower bounds for the Hausdorff dimension of the underlying IFS. This will follow directly if we are able to derive rigorous error bounds for $|[R(L_s)]^\nu - R(\mathbf{L}_{s,\nu})|$. In the case of piecewise linear functions, we obtained the bounds by using a simple and well-known result (c.f. Lemma 2.2 of [14]) that if A is a nonnegative matrix and w a vector with strictly positive components, then if for all components k , (i) if $(Aw)_k \geq \lambda w_k$, then $R(A) \geq \lambda$ and (ii) if $(Aw)_k \leq \lambda w_k$, then $R(A) \leq \lambda$. Here, we use an analogous result for a matrix mapping a cone K to itself, in which we replace \leq by \leq_K , i.e. $u \leq_K v$ if and only if $v - u \in K$.

Another key tool for obtaining rigorous upper and lower bounds for the Hausdorff dimension s_* , is to obtain and use explicit a priori bounds on the quantity $D^q v_s(x)/v_s(x)$ of the strictly positive eigenfunction v_s of L_s , where $D^q v_s$ denotes the q -th derivative of v_s . Such estimates are derived in Section 7.

In order to improve the efficiency of our computation, we consider in Section 7 the possibility of replacing the original interval $S = [a, b]$ by a smaller interval $S_0 \subset S$ such that $\theta_\beta(S_0) \subset S_0$ for $\beta \in \mathcal{B}$. In particular, for the maps $\theta_\beta = 1/(x + \beta)$, setting $\gamma = \min\{\beta : \beta \in \mathcal{B}\}$ and $\Gamma = \max\{\beta : \beta \in \mathcal{B}\}$, we can reduce the interval S to $[a_\infty, b_\infty]$, where

$$a_\infty = -\frac{\gamma}{2} + \sqrt{(\gamma/2)^2 + (\gamma/\Gamma)} \quad \text{and} \quad b_\infty = -\frac{\Gamma}{2} + \sqrt{(\Gamma/2)^2 + (\Gamma/\gamma)} = \frac{\Gamma}{\gamma} a_\infty.$$

For example, for the set $\{1, 2\}$, we reduce the interval $[0, 1]$ to $[(\sqrt{3}-1)/2, \sqrt{3}-1]$ of length 0.366, while for the set $\{10, 11\}$, we reduce the interval $[0, 1]$ to $[0.0901, .0991]$ of length .009.

A main result of the paper (Theorem 8.6) says that under appropriate hypotheses, there is a computable constant H , such that

$$[R([1 + Hh^r]^{-1} \mathbf{L}_{s,\nu})]^{1/\nu} \leq \lambda_s \leq [R([1 - Hh^r]^{-1} \mathbf{L}_{s,\nu})]^{1/\nu},$$

where h denotes the maximum mesh size and r the degree of the piecewise polynomial approximation. Using these inequalities, we can obtain rigorous upper and lower bounds on the Hausdorff dimension of the invariant set associated with the transfer operator L_s as follows. Let s_l and s_u denote values of s satisfying

$$[1 - Hh^r]^{-1}R(\mathbf{L}_{s_u, \nu}) < 1, \quad [1 + Hh^r]^{-1}R(\mathbf{L}_{s_l, \nu}) > 1.$$

It follows immediately from Theorem 8.6 that $\lambda_{s_u}^\nu < 1$ and $\lambda_{s_l}^\nu > 1$. Since the spectral radius λ_s of L_s is a strictly decreasing function of s , there will be a value s_* satisfying $s_l < s_* < s_u$ for which $\lambda_{s_*}^\nu = 1$, or equivalently $\lambda_{s_*} = 1$. The value s_* gives the Hausdorff dimension s_* of the invariant set associated with the transfer operator L_s . Since $s_u - s_l$ is of order h^r , by choosing h to be sufficiently small and/or r to be sufficiently large, we obtain a highly accurate estimate for s_* . As noted above, for a given s , $R([1 \pm Hh^r]^{-1}\mathbf{L}_{s, \nu})$ is easily computed by variants of the power method for eigenvalues, since the largest eigenvalue has multiplicity one and is the only eigenvalue of its modulus. Our theoretical results imply that $\mathbf{L}_{s, \nu}$ has an eigenvector w in $K := K_M$ with eigenvalue $R(\mathbf{L}_{s, \nu})$ and that this eigenvector can be computed to high accuracy. Still, one might be concerned about possible errors in the computation of $R(\mathbf{L}_{s, \nu})$ and w . However, independently of how a purported eigenvector $w \in K$ for $\mathbf{L}_{s, \nu}$ is found, if $\alpha w \leq_K \mathbf{L}_{s, \nu} w \leq_K \beta w$, Lemma 4.1 in Section 4 implies that $\alpha \leq R(\mathbf{L}_{s, \nu}) \leq \beta$. This provides a means of giving rigorous bounds for $R(\mathbf{L}_{s, \nu})$.

In Section 9, we present results of computations of the Hausdorff dimension s of invariant sets in $[0, 1]$ arising from continued fraction expansions. In this much studied case, one defines $\theta_p = 1/(x + p)$, for p a positive integer and $x \in [0, 1]$; and for a subset $\mathcal{B} \subset \mathbb{N}$, one considers the IFS $\{\theta_p : p \in \mathcal{B}\}$ and seeks estimates on the Hausdorff dimension of the invariant set $C = C(\mathcal{B})$ for this IFS. This problem has previously been considered by many authors. See [3], [6], [7], [15], [18], [20], [19], [23], [24], and [17]. In this case, (1.1) becomes

$$(L_s f)(x) = \sum_{p \in \mathcal{B}} \left(\frac{1}{x + p} \right)^{2s} f\left(\frac{1}{x + p} \right), \quad 0 \leq x \leq 1,$$

and one seeks a value $s \geq 0$ for which $\lambda_s := R(L_s) = 1$. Several of the papers listed above contain a large number of computations to various degrees of accuracy of the Hausdorff dimension of the IFS $\{\theta_p : p \in \mathcal{B}\}$, for various choices of the set \mathcal{B} . An early paper, [20], gives results for over 30 choices of \mathcal{B} , containing between two and five terms in the set \mathcal{B} , with results reported to an accuracy between 10^{-6} and 10^{-19} , depending on the problem studied. A *Mathematica* code implementing the algorithm is also provided. In [23], computations to four decimal places are given for over 35 choices of the set \mathcal{B} , ranging from two terms, to as many as 34 (this computation is to three decimal places), and also includes a computation of $E[1, 2]$, ($\mathcal{B} = \{1, 2\}$), accurate to 54 decimal places. In [24], eight examples of \mathcal{B} , consisting of two terms, are computed with accuracies ranging from 10^{-13} to 10^{-52} , depending on the choice of \mathcal{B} , although the authors note that for the sets [10, 11], and [100, 10, 000], they were able to compute to accuracies of 10^{-61} and 10^{-122} , respectively. This depends on the fact that the speed of convergence of their methods depends on the size of the smallest value of $p \in \mathcal{B}$. In [26], the Hausdorff dimension of $E[1, 2]$ is rigorously computed to 100 decimal places, although more digits are computed. It is less clear how well some of the approximation schemes

employed in these papers work when $|\mathcal{B}|$ is moderately large or when different real analytic functions $\hat{\theta}_j : [0, 1] \rightarrow [0, 1]$ are used. Here and in [14], in the one dimensional case, we present an alternative approach with much wider applicability that only requires the maps in the IFS to be C^m , for some finite value of m . As an illustration, we considered in [14], perturbations of the IFS for the middle thirds Cantor set for which the corresponding contraction maps are C^3 , but not C^4 .

The computations in Section 9 include choices of various sets of continued fractions, maximum mesh size h , piecewise polynomial degree r , and number of iterations ν , (where $\nu = 1$ corresponds to the original map), including choices of ν for which the hypotheses of our theorems are satisfied, but also computations which obtain the same results when the mappings are not iterated. These results support our conjecture that our method also works in the non-iterated situation. To facilitate computation of further examples, a *Matlab* code is provided in <https://sites.math.rutgers.edu/~falk/hausdorff/codes.html>.

An outline of the paper is as follows. In the next section, we introduce further notation and state some preliminary results we will use in our analysis. Section 3 contains a description of the approximate problem and the cone we use to analyze it. Section 4 contains the theoretical results we will need to show that the matrices arising from the approximation scheme satisfy the conclusions of the Perron theorem. In Section 5, the main result is to determine conditions under which the matrix $\mathbf{L}_{s,\nu}$ satisfies (1.2). These conditions involve a number of constants, which we then estimate in Section 6, ultimately deriving bounds for $R(L_s)$ in terms of $[R(\mathbf{L}_{s,\nu})]^{1/\nu}$. In Section 7, we consider a method for reducing the size of the interval S on which the problem is defined, with the aim of reducing the number of mesh points that will be needed in the approximation scheme. In so doing, we are also able to improve the bound on two constants which are used in the error estimate for $R(L_s)$. Recall that condition (1.2) requires determining for each constant M , a constant $0 < M' < M$ such that (1.2) is satisfied. In Section 8, we provide a procedure for determining this constant. Finally, the numerical computations described above are given in Section 9.

It would be of considerable interest to extend the methods of this paper to the two dimensional case, e.g., to the problem of obtaining rigorous estimates for the Hausdorff dimension of sets of complex continued fractions. We conjecture that such an extension can be done, but we leave it as an open problem for possible future work.

2. NOTATION AND PRELIMINARIES

Let $C(S)$ denote the Banach space of continuous functions $f : S \rightarrow \mathbb{R}$, where S is a compact subset of \mathbb{R} . Assume

(H0): For $1 \leq p \leq n < \infty$, $\theta_p : S \rightarrow S$ is a Lipschitz map.

(H1): For $1 \leq p \leq n < \infty$, $g_p : S \rightarrow [0, \infty)$ is a nonnegative continuous function which is not identically zero. In addition, there exists a constant $M_0 > 0$ such that

$$g_p(t_1) \leq g_p(t_2) \exp(M_0|t_1 - t_2|), \quad \forall t_1, t_2 \in S, \quad 1 \leq p \leq n.$$

We note that it is easy to show that (H1) is equivalent to assuming that $g_p(t) > 0$ for all $t \in S$, and

$$|\ln(g_p(t_1)) - \ln(g_p(t_2))| \leq M_0|t_1 - t_2|, \quad \forall t_1, t_2 \in S, \quad 1 \leq p \leq n.$$

For $s > 0$, define a bounded linear map $L_s : C(S) \rightarrow C(S)$ (often called a *Perron-Frobenius operator*) by (1.1), i.e.

$$(L_s f)(t) = \sum_{p=1}^n [g_p(t)]^s f(\theta_p(t)), \quad t \in S.$$

We shall need to consider the ν th iterate of L_s , L_s^ν . For $\nu \geq 1$, let $\Omega_\nu = \{\omega = (p_1, p_2, \dots, p_\nu) : 1 \leq p_j \leq n \text{ for } 1 \leq j \leq \nu\}$; and for $\omega = (p_1, p_2, \dots, p_\nu) \in \Omega_\nu$, define

$$\theta_\omega(t) = (\theta_{p_1} \circ \theta_{p_2} \circ \dots \circ \theta_{p_\nu})(t)$$

and

$$g_\omega(t) = g_{p_1}(\theta_{p_2} \circ \dots \circ \theta_{p_\nu}(t)) g_{p_2}(\theta_{p_3} \circ \dots \circ \theta_{p_\nu}(t)) \dots g_{p_{\nu-1}}(\theta_{p_\nu}(t)) g_{p_\nu}(t).$$

The reader can verify (e.g., see [35]) that for all $f \in C(S)$,

$$(L_s^\nu f)(t) = \sum_{\omega \in \Omega_\nu} [g_\omega(t)]^s f(\theta_\omega(t)).$$

Note that L_s^ν has the same form as L_s , except with index set Ω_ν . To analyze the operator L_s^ν , we shall need stronger assumptions than (H0). We will thus assume

(H2): (H0) is satisfied and there exist constants $C_0 \geq 1$ and κ , $0 < \kappa < 1$, such that for all integers $\nu \geq 1$, all $\omega \in \Omega_\nu$, and all $t_1, t_2 \in S$,

$$|\theta_\omega(t_1) - \theta_\omega(t_2)| \leq C_0 \kappa^\nu |t_1 - t_2|.$$

Assuming (H1) and (H2), one can prove that for all $\omega \in \Omega_\nu$ and all $t_1, t_2 \in S$,

$$g_\omega(t_1) \leq \exp(M'_0 |t_1 - t_2|) g_\omega(t_2),$$

where $M'_0 = M_0 C_0 [(1 - \kappa^\nu)/(1 - \kappa)]$. The proof is left to the reader. The reader will notice that the above framework carries over to the more general case that S is a compact metric space with metric ρ . (H1) and (H2) take the same form except that $|t_1 - t_2|$ is replaced by $\rho(t_1, t_2)$.

The following result provides some theoretical background which will be essential for our later work concerning the operator L_s . This theorem is a special case of Corollary 6.6 in [34]. We refer to Section 3 of [33] for a brief discussion of the essential spectrum, which is mentioned in Theorem 2.1 below.

Theorem 2.1. *Assume hypotheses (H1) and (H2) are satisfied, that S is a finite union of compact intervals, and that L_s is given by (1.1), where $s > 0$. Assume also that $\theta_i \in C^m(S)$ and $g_i \in C^m(S)$ for some positive integer m . Let $\Lambda_s : Y := C^m(S) \rightarrow Y$ be the bounded linear operator given by (1.1), but considered as a map of $Y \mapsto Y$, so $L_s(f) = \Lambda_s(f)$ for $f \in Y$. If $R(L_s)$ (respectively, $R(\Lambda_s)$) denotes the spectral radius of L_s (respectively, of Λ_s) and $\rho(\Lambda_s)$ denotes the essential spectral radius of Λ_s and κ is as in (H2), then*

$$\rho(\Lambda_s) \leq \kappa^m R(\Lambda_s), \quad R(\Lambda_s) = R(L_s) := \lambda_s > 0.$$

Let $\hat{\Lambda}_s$ denote the complexification of Λ_s . If $\sigma(\hat{\Lambda}_s)$ denotes the spectrum of $\hat{\Lambda}_s$, and we define $\sigma(\Lambda_s) := \sigma(\hat{\Lambda}_s)$, then if $z \in \sigma(\Lambda_s)$ and $\rho(\Lambda_s) < |z|$, z is an isolated point of $\sigma(\Lambda_s)$ and is an eigenvalue of Λ_s of finite algebraic multiplicity. Also, there exists $b_s < \lambda_s$ such that

$$\sigma(\Lambda_s) \setminus \{\lambda_s\} \subset \{z \in \mathbb{C} : |z| \leq |b_s|\}.$$

There exists a strictly positive eigenfunction $v_s \in C^m(S)$ with eigenvalue $\lambda_s > 0$, and λ_s is an algebraically simple eigenvalue of Λ_s . If $u \in Y$ and $u(t) > 0$ for all $t \in S$, there exists a positive real number α (dependent on u) such that

$$(2.1) \quad \lim_{k \rightarrow \infty} \left(\frac{1}{\lambda_s} \right)^k \Lambda_s^k(u) = \alpha v_s,$$

where the convergence is in the norm topology on Y .

Remark 2.1. In our work here, it will be important to have estimates on

$$\sup \left\{ \frac{|(d^j v_s / dt^j)(t)|}{v_s(t)} : t \in S \right\},$$

where $1 \leq j \leq m$. Note that if we take $u := 1$ in (2.1), we find that for $t \in S$ and $1 \leq j \leq m$,

$$(2.2) \quad \frac{|(d^j v_s / dt^j)(t)|}{v_s(t)} = \lim_{k \rightarrow \infty} \frac{|\sum_{\omega \in \Omega_k} (d^j g_\omega^s / dt^j)(t)|}{\sum_{\omega \in \Omega_k} g_\omega(t)^s},$$

and the convergence in (2.2) is uniform in $t \in S$. If u is as in (2.1), we also obtain from (2.1) that

$$\lim_{k \rightarrow \infty} \frac{\Lambda_s^{k+1} u}{\Lambda_s^k u} = \lambda_s,$$

where the convergence to the constant function λ_s is in the norm topology on Y .

3. APPROXIMATION OF THE SPECTRAL RADIUS OF L_s

Returning to the notation of (1.1), we want to approximate $R(L_s)$ by the spectral radius of an appropriate finite dimensional linear map \mathbf{L}_s . To do so, we assume that $S = [a, b]$ in (H1) and (H2), with $a < b$ and let \hat{S} denote a union of disjoint subintervals $[a_i, b_i] \subset [a, b]$, $i = 1, \dots, I$. We also assume throughout this section that $\theta_p(\hat{S}) \subset \hat{S}$ for $1 \leq p \leq n$. Further subdivide each interval $[a_i, b_i]$ into N_i equally spaced subintervals $[t_{j-1}^i, t_j^i]$, $j = 1, \dots, N_i$ of width

$$(3.1) \quad h_i = (b_i - a_i) / N_i, \quad 1 \leq i \leq I.$$

Set $h = \max_{1 \leq i \leq I} h_i$.

Next let $\{c_{j,k}^i\}_{k=0}^r \in [t_{j-1}^i, t_j^i]$, with $c_{j,0}^i = t_{j-1}^i$, $c_{j,r}^i = t_j^i$, and $c_{j,k}^i < c_{j,k+1}^i$ for $0 \leq k < r$. Given values $\{F_{j,k}^i\} = F(c_{j,k}^i)$, we then define on \hat{S} , a piecewise polynomial \mathcal{F} as follows: For $t_{j-1}^i \leq x \leq t_j^i$, $1 \leq j \leq N_i$, and $1 \leq i \leq I$,

$$(3.2) \quad \mathcal{F}|_{[t_{j-1}^i, t_j^i]}(x) = \mathcal{F}_j^i(x) = \sum_{k=0}^r l_{j,k}^i(x) F_{j,k}^i,$$

where

$$l_{j,k}^i(x) = \frac{\prod_{\substack{l=0 \\ l \neq k}}^r (x - c_{j,l}^i)}{\prod_{\substack{l=0 \\ l \neq k}}^r (c_{j,k}^i - c_{j,l}^i)}.$$

Since $c_{j,r}^i = c_{j+1,0}^i$, $\mathcal{F} \in \mathbf{V}_h^r$, the space of continuous piecewise polynomials of degree $\leq r$, whose degrees of freedom are the $Nr + I := Q$ values $F_{j,k}^i$, where $N = \sum_{i=1}^I N_i$.

We note that we can simplify our expressions by choosing points $\{\hat{c}_k\}_{k=0}^r \in [-1, 1]$, with $\hat{c}_k < \hat{c}_{k+1}$ for $0 \leq k < r$, $\hat{c}_0 = -1$ and $\hat{c}_r = 1$. If we then define

$$c_{j,k}^i = t_{j-1}^i + h_i(1 + \hat{c}_k)/2,$$

and write $x \in [t_{j-1}^i, t_j^i] \subset [a_i, b_i]$ in the form $x = t_{j-1}^i + h_i(1 + \hat{x})/2$, where $\hat{x} \in [-1, 1]$, we obtain

$$(3.3) \quad l_{j,k}^i(x) = \hat{l}_k(\hat{x}) = \frac{\prod_{\substack{l=0 \\ l \neq k}}^r (\hat{x} - \hat{c}_l)}{\prod_{\substack{l=0 \\ l \neq k}}^r (\hat{c}_k - \hat{c}_l)}.$$

Because we seek to make use of high order piecewise polynomials, it is important to choose the points \hat{c}_k to avoid the large errors that can occur in polynomial interpolation due to Runge's phenomenon (e.g., when equally spaced interpolation points are used). Since, for our analysis, we shall need the function $\mathcal{F}(x)$ in (3.2) to be continuous, we choose the points \hat{c}_k to be the extended Chebyshev points in $[-1, 1]$ given by

$$(3.4) \quad \hat{c}_k = -\cos\left(\frac{2k+1}{2r+2}\pi\right) / \cos\left(\frac{\pi}{2r+2}\right), \quad k = 0, \dots, r,$$

obtained by rescaling the usual Chebyshev nodes. Then

$$(3.5) \quad c_{j,k}^i = t_{j-1}^i + \frac{h_i}{2} \left(1 - \left[\cos\left(\frac{2k+1}{2r+2}\pi\right) / \cos\left(\frac{\pi}{2r+2}\right)\right]\right), \quad k = 0, \dots, r.$$

We note that another possible choice is to use the augmented Chebyshev points, consisting of the roots of the Chebyshev polynomial of degree $r-1$ shifted to the interval $[t_{j-1}^i, t_j^i]$ plus the endpoints t_{j-1}^i and t_j^i .

With this notation, we can now define, for $s > 0$, the linear map $\mathbf{L}_s : \mathbb{R}^Q \rightarrow \mathbb{R}^Q$. If $\mathbf{F} = \{F_{j,k}^i\} \in \mathbb{R}^Q$, we define

$$\mathbf{L}_s(\mathbf{F})(c_{j,k}^i) = \sum_{p=1}^n g_p(c_{j,k}^i)^s \mathcal{F}(\theta_p(c_{j,k}^i)),$$

where \mathcal{F} is defined above. Equivalently, we can also think of the operator \mathbf{L}_s as a map from the space $\mathbf{V}_h^r \rightarrow \mathbf{V}_h^r$, if we replace \mathbf{F} by \mathcal{F} , and given the values $\{G_{j,k}^i\} = \sum_{p=1}^n g_p(c_{j,k}^i)^s \mathcal{F}(\theta_p(c_{j,k}^i))$, we define $\mathcal{G}(x)$ as follows: For $t_{j-1}^i \leq x \leq t_j^i$, $1 \leq j \leq N_i$, and $1 \leq i \leq I$,

$$\mathcal{G}|_{[t_{j-1}^i, t_j^i]}(x) = \mathcal{G}_j^i(x) = \sum_{k=0}^r l_{j,k}^i(x) G_{j,k}^i.$$

Given a positive real number M , we next define $K_M \subset \mathbb{R}^Q$ as $\{\mathbf{F} \in \mathbb{R}^Q\}$ such that for all $\xi = c_{j,k}^i$ and $\eta = c_{j',k'}^{i'}$, with $1 \leq i, i' \leq I$, $1 \leq j \leq N_i$, $1 \leq j' \leq N_{i'}$, and $0 \leq k, k' \leq r$,

$$(3.6) \quad F(\xi) \leq \exp(M|\xi - \eta|)F(\eta).$$

Note that to verify (3.6), it suffices to verify it whenever ξ and η are two consecutive points in the linear ordering inherited from \mathbb{R} of the points $\{c_{j,k}^i\}$.

One can easily verify that if $\mathbf{F} \in K_M$, then either (a) $F_{j,k}^i = 0$ for all $1 \leq i \leq I$, $1 \leq j \leq N_i$, and $0 \leq k \leq r$, or (b) $F_{j,k}^i > 0$ for all i, j, k in this range. In case (b), one has for all $1 \leq i, i' \leq I$, $1 \leq j \leq N_i$, $1 \leq j' \leq N_{i'}$, $0 \leq k, k' \leq r$,

$$(3.7) \quad |\ln(F_{j,k}^i) - \ln(F_{j',k'}^{i'})| \leq M|c_{j,k}^i - c_{j',k'}^{i'}|.$$

and (3.7) implies that (3.6) holds.

One might hope to prove that the spectral radius $R(\mathbf{L}_s)$ of \mathbf{L}_s closely approximates the spectral radius $R(L_s)$, and we shall see that this is true if the Lipschitz constant $C_0\kappa$ in (H2) (corresponding to the case $\nu = 1$ and the operator $R(L_s)$) and the constant h in (3.1) are sufficiently small. However, if $C_0\kappa$ is not sufficiently small, we can instead work with the operator L_s^ν , where ν is a positive integer, and the corresponding Lipschitz constant in (H2) is then $C_0\kappa^\nu$. This in turn means that we will have to replace \mathbf{L}_s by $\mathbf{L}_{s,\nu} : \mathbb{R}^Q \rightarrow \mathbb{R}^Q$, where, ν is a positive integer and in our earlier notation,

$$(3.8) \quad (\mathbf{L}_{s,\nu}(\mathbf{F}))(c_{j,k}^i) = \sum_{\omega \in \Omega_\nu} g_\omega(c_{j,k}^i)^s \mathcal{F}(\theta_\omega(c_{j,k}^i)).$$

4. CONES, POSITIVE EIGENVECTORS, AND BIRKHOFF'S CONTRACTION CONSTANT

As noted in Section 1, we would like to have the approximating matrices defined in the previous section mimic the properties of the infinite dimensional, bounded linear operator L_s , which means they should satisfy the conclusions of the Perron theorem for positive matrices, i.e., they should have an eigenvalue of multiplicity one equal to the spectral radius of the matrix with corresponding positive eigenvector and that all other eigenvalues of the matrix should have modulus less than the spectral radius. However, the matrix \mathbf{L}_s defined in the previous section is not even a nonnegative matrix once the degree r of the piecewise polynomial is > 1 . The reason for this, seen by constructing the matrix, is that the Lagrange basis functions for a polynomial of degree > 1 are not always positive.

The remedy, also used in the case $r = 1$, when the resulting matrix was nonnegative, but not *primitive* or *irreducible*, is to base the analysis on a cone different from the usual cone of nonnegative functions. More precisely, by using the cone K_M defined in the previous section, we shall show that the conclusions of the classical Perron theorem also hold for the matrices of this paper.

To outline our method of proof, it is convenient to describe, at least in the finite dimensional case, some basic definitions and classical theorems concerning linear

maps $L : \mathbb{R}^N \rightarrow \mathbb{R}^N$ which leave a cone $K \subset \mathbb{R}^N$ invariant. In doing so, we shall closely follow the analogous description in [14]. Recall that a closed subset K of \mathbb{R}^N is called a closed cone if (i) $ax+by \in K$ whenever $a \geq 0, b \geq 0, x \in K$ and $y \in K$ and (ii) if $x \in K \setminus \{0\}$, then $-x \notin K$. If K is a closed cone, K induces a partial ordering on \mathbb{R}^N denoted by \leq_K (or simply \leq , if K is obvious) by $u \leq_K v$ if and only if $v - u \in K$. If $u, v \in K$, we shall say that u and v are *comparable* (with respect to K) and we shall write $u \sim_K v$ if there exist positive scalars a and b such that $v \leq_K au$ and $u \leq_K bv$. *Comparable with respect to K* partitions K into equivalence classes of comparable elements. We shall henceforth assume that $\text{int}(K)$, the interior of K , is nonempty. Then an easy argument shows that all elements of $\text{int}(K)$ are comparable. Generally, if $x_0 \in K$ and $K_{x_0} := \{x \in K : x \sim_K x_0\}$, all elements of K_{x_0} are comparable.

Following standard notation, if $u, v \in K$ are comparable elements, we define

$$M(u/v; K) = \inf\{\beta > 0 : u \leq \beta v\},$$

$$m(u/v; K) = M(v/u; K)^{-1} = \sup\{\alpha > 0 : \alpha v \leq u\}.$$

If u and v are comparable elements of $K \setminus \{0\}$, we define Hilbert's projective metric $d(u, v; K)$ by

$$d(u, v; K) = \ln(M(u/v; K)) + \ln M(v/u; K).$$

We make the convention that $d(0, 0; K) = 0$. If $x_0 \in K \setminus \{0\}$, then for all $u, v, w \in K_{x_0}$, one can prove that (i) $d(u, v; K) \geq 0$, (ii) $d(u, v; K) = d(v, u; K)$, and (iii) $d(u, v; K) + d(v, w; K) \geq d(u, w; K)$. Thus d restricted to K_{x_0} is almost a metric, but $d(u, v; K) = 0$ if and only if $v = tu$ for some $t > 0$ and generally, $d(su, tv; K) = d(u, v; K)$ for all $u, v \in K_{x_0}$ and all $s > 0$ and $t > 0$. If $\|\cdot\|$ is any norm on \mathbb{R}^N and $S := \{u \in \text{int}(K) : \|u\| = 1\}$ (or, more generally, if $x_0 \in K \setminus \{0\}$ and $S = \{x \in K_{x_0} : \|x\| = 1\}$), then $d(\cdot, \cdot; K)$, restricted to $S \times S$, gives a metric on S ; and it is known that S is a complete metric space with this metric.

With these preliminaries, we can describe a special case of the Birkhoff-Hopf theorem. We refer to [2], [21], and [40] for the original papers and to [12] and [11] for an exposition of a general version of this theorem and further references to the literature. We remark that P. P. Zabreiko, M. A. Krasnosel'skij, Y. V. Pokornyi, and A. V. Sobolev independently obtained closely related theorems; and we refer to [28] for details. If K is a closed cone as above, $S = \{x \in \text{int}(K) : \|x\| = 1\}$, and $L : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a linear map such that $L(\text{int}(K)) \subset \text{int}(K)$, we define $\Delta(L; K)$, the *projective diameter* of L by

$$\Delta(L; K) = \sup\{d(Lx, Ly; K) : x, y \in K \text{ and } Lx \sim_K Ly\}$$

$$= \sup\{d(Lx, Ly; K) : x, y \in \text{int}(K)\}.$$

The Birkhoff-Hopf theorem implies that if $\Delta := \Delta(L; K) < \infty$, then L is a contraction mapping with respect to Hilbert's projective metric. More precisely, if we define $\lambda = \tanh(\frac{1}{4}\Delta) < 1$, then for all $x, y \in K \setminus \{0\}$ such that $x \sim_K y$, we have

$$d(Lx, Ly; K) \leq \lambda d(x, y; K),$$

and the constant λ is optimal.

If we define $\Phi : S \rightarrow S$ by $\Phi(x) = L(x)/\|L(x)\|$, it follows that Φ is a contraction mapping with a unique fixed point $v \in S$, and v is necessarily an eigenvector of

L with eigenvalue $r(L) := r =$ the spectral radius of L . Furthermore, given any $x \in \text{int}(K)$, there are explicitly computable constants M and $c < 1$ (see Theorem 2.1 in [12]) such that for all $k \geq 1$,

$$\|L^k(x)/\|L^k(x)\| - v\| \leq Mc^k;$$

and the latter inequality is exactly the sort of result we need. Furthermore, it is proved in Theorem 2.3 of [12] that $r = r(L)$ is an algebraically simple eigenvalue of L and that if $\sigma(L)$ denotes the spectrum of L and $q(L)$ denotes the *spectral clearance* of L ,

$$q(L) := \sup\{|z|/r(L) : z \in \sigma(L), z \neq r(L)\},$$

then $q(L) < 1$ and $q(L)$ can be explicitly estimated.

The main issue, then, is to find a suitable cone satisfying the hypotheses outlined above. We shall show in the sections that follow that the cone K_M , defined in the previous section, is such a cone. To do so, we shall show that there exists M' , $0 < M' < M$, such that $L(K_M \setminus \{0\}) \subset K_{M'} \setminus \{0\}$. After correcting the typo in the formula for $d_2(f, g)$ on page 286 of [29], it follows from Lemma 2.12 on page 284 of [29] that

$$\sup\{d(f, g; K_M) : f, g \in K_{M'} \setminus \{0\}\} \leq 2 \ln \left(\frac{M + M'}{M - M'} \right) + 2 \exp(M'(b - a)) < \infty,$$

where now $S := [a, b]$ in (H1) and (H2) (c.f. Section 3). This implies that $\Delta(L; K_M) < \infty$, which in turn implies that L has a normalized eigenvector $v \in K_{M'}$ with positive eigenvalue $r = r(L) =$ the spectral radius of L . Furthermore, r has algebraic multiplicity 1, $q(L) < 1$, and $\lim_{k \rightarrow \infty} \|L^k(x)/\|L^k(x)\| - v\| = 0$ for all $x \in K_M \setminus \{0\}$. Thus it suffices to prove, for an appropriate map L , that for some $M' < M$,

$$(4.1) \quad L(K_M \setminus \{0\}) \subset K_{M'} \setminus \{0\}.$$

We note that if the map L satisfies (4.1), then it is not difficult to show that $L(K_M \setminus \{0\}) \subset \text{int}(K_M)$. An alternative approach is then to apply Theorem 4.4 of [44], which concludes that $R(L)$ is a simple eigenvalue, greater than the magnitude of any other eigenvalue, and that an eigenvector corresponding to $R(L)$ lies in $\text{int}(K)$. In any case, the key step is to show for an appropriate matrix L and cone K_M , that (4.1) is satisfied.

A key part of the paper is to obtain upper and lower bounds on $R(L_s)$ using the approximations developed in this paper. To do so, we will use an extension to cones of a well known result for positive matrices.

Lemma 4.1. *Suppose $L(K_M \setminus \{0\}) \subset K_{M'}$ and $\mathcal{V}_s \in K_M \setminus \{0\}$. Then if there exists positive constants α and β such that*

$$\alpha \mathcal{V}_s \leq_{K_M} L \mathcal{V}_s \leq_{K_M} \beta \mathcal{V}_s,$$

then $\alpha \leq R(L) \leq \beta$.

5. THEORY FOR THE DISCRETE APPROXIMATION

The main result of this section, Theorem 5.6, is to show that under appropriate hypotheses, the matrix operator $\mathbf{L}_{s,\nu}$ defined in Section 3, satisfies

$$\mathbf{L}_{s,\nu}(K_M(T) \setminus \{0\}) \subset K_{M'}(T) \setminus \{0\},$$

where T will be as in (5.1) below.

Throughout this section, we shall assume that (H1) and (H2) are satisfied and that $S = [a, b]$ with $a < b$, where S is as in (H1) and (H2). We shall also assume that (H3), given below, is satisfied, and we shall use the notation of (H1), (H2), and (H3).

(H3): For a given positive integer ν , there exist pairwise disjoint, nonempty compact intervals $[a_i, b_i] \subset S$, $1 \leq i \leq I$, (where $S := [a, b]$ is as in (H0) - (H2)) with the following property: For every $\omega \in \Omega_\nu$, there exists $i = i(\omega)$, $1 \leq i \leq I$, such that $\theta_\omega(S) \subset [a_i, b_i]$.

Remark 5.1. Assume that (H0)-(H2) are satisfied and that for some positive integer ν' , $\theta_{\omega_1}(S)$ and $\theta_{\omega_2}(S)$ are disjoint whenever ω_1 and ω_2 are unequal elements of $\Omega_{\nu'}$. Label the elements of $\Omega_{\nu'}$ as ω_i , $1 \leq i \leq I$, and define $[a_i, b_i] = \theta_{\omega_i}(S)$. Then for all positive integers $\nu \geq \nu'$, (H3) is satisfied. More generally, one could, for $1 \leq i \leq I$, take $[a_i, b_i]$ to be any interval contained in $[a, b]$ such that $\theta_{\omega_i}(S) \subset [a_i, b_i]$ as long as $[a_i, b_i] \cap [a_j, b_j] = \emptyset$ for $1 \leq i, j \leq I$. Note that (H3) is also trivially satisfied if we take $I = 1$ and $[a_1, b_1] = [a, b]$.

Remark 5.2. In the context of (H3), it is possible by relabeling to assume that $b_i < a_{i+1}$ for $1 \leq i < I$, so the intervals are linearly ordered as subsets of \mathbb{R} . Thus we shall make this assumption if convenient.

We now follow the notation of Section 3. If we define

$$(5.1) \quad T := \{c_{j,k}^i : 1 \leq i \leq I, 1 \leq j \leq N_i, 0 \leq k \leq r\},$$

then T is a finite subset of \mathbb{R} and a compact metric space. Recall that we consider the finite dimensional vector space $V = V(T)$ of dimension $Q = Nr + I$ of all maps $F : T \rightarrow \mathbb{R}$ and $K_M(T)$ is then defined as in Section 4 or (3.6), i.e., $F \in K_M(T) \setminus \{0\}$ if and only if

$$|\ln(F(\xi)) - \ln(F(\eta))| \leq M|\xi - \eta|$$

for all $\xi, \eta \in T$. Note that V is linearly isomorphic to \mathbb{R}^Q .

A central question for our approach is to determine under what conditions on $\mathbf{L}_{s,\nu}$ (see (3.8)) $\mathbf{L}_{s,\nu}(K_M(T)) \subset K_{M'}(T)$ for some M, M' with $0 < M' < M$. To do so, we begin by recalling some classical results.

Lemma 5.1. (See [32], [36], pages 121-123, or [42]). Let $p(t)$ be a real-valued polynomial of degree $\leq r$. Then

$$\max_{-1 \leq t \leq 1} |p'(t)| \leq r^2 \max_{-1 \leq t \leq 1} |p(t)|.$$

A proof of the following estimate is given in [5] and refinements for $r \geq 5$ can be found in [16].

Lemma 5.2. *If $\hat{l}_k(\hat{x})$, $0 \leq k \leq r$, is defined by (3.3), then*

$$\max_{-1 \leq \hat{x} \leq 1} \sum_{k=0}^r |\hat{l}_k(\hat{x})| \leq \frac{2}{\pi} \ln(r+1) + 3/4 := \psi(r),$$

where \ln denotes natural logarithm.

It will also be convenient to have some elementary estimates concerning the numbers $c_{j,k}^i$, $1 \leq j \leq N_i$, $0 \leq k \leq r$, in (3.5). If x is a real number, $[x]$ denotes the greatest integer $m \leq x$. If r is an integer, it follows that $[r/2] = r/2$ if r is even and $[r/2] = (r-1)/2$ if r is odd. The next lemma is a straightforward exercise and is left to the reader.

Lemma 5.3. *Let the numbers $c_{j,k}^i$ be defined by (3.5). Then for $0 \leq k \leq r$ and $1 \leq i \leq I$,*

$$\begin{aligned} |c_{j,k}^i - c_{j,[r/2]}^i| &\leq h_i/2, \quad \text{if } r \text{ is even,} \\ |c_{j,k}^i - c_{j,[r/2]}^i| &\leq h_i/2[1 + \tan(\pi/[2r+2])], \quad \text{if } r \text{ is odd.} \end{aligned}$$

For $1 \leq j \leq N_i$ and $0 \leq k < r$,

$$\begin{aligned} \min_{1 \leq j \leq N_i, 0 \leq k < r} |c_{j,k}^i - c_{j,k+1}^i| &= 2h_i[\sin(\pi/[2r+2])]^2, \\ \max_{1 \leq j \leq N_i, 0 \leq k < r} |c_{j,k}^i - c_{j,k+1}^i| &= h_i \tan(\pi/[2r+2]) \quad \text{if } r \text{ is odd,} \\ \max_{1 \leq j \leq N_i, 0 \leq k < r} |c_{j,k}^i - c_{j,k+1}^i| &= h_i \sin(\pi/[2r+2]) \quad \text{if } r \text{ is even.} \end{aligned}$$

Since the first result in Lemma 5.3 will be used later, we define the constant $\eta(r)$ for r a positive integer by:

$$(5.2) \quad \eta(r) = \begin{cases} \frac{1}{2}, & \text{if } r \text{ even} \\ \frac{1}{2}[1 + \tan(\pi/[2r+2])], & \text{if } r \text{ odd} \end{cases}.$$

In addition, for ν a positive integer and $\omega \in \Omega_\nu$, we define constants $M_0(\nu)$ and $c(\nu)$ such that for all $\omega \in \Omega_\nu$,

$$(5.3) \quad g_\omega \in K_{M_0(\nu)}(S) \quad \text{and} \quad \text{lip}(\theta_\omega|_S) \leq c(\nu).$$

We already know (see Section 2) that $M_0(\nu) \leq M_0 C_0(1 - \kappa^\nu)/(1 - \kappa)$, where M_0 is as in (H1) and C_0 and κ are as in (H2); and (H2) implies that $c(\nu) \leq C_0 \kappa^\nu$. However, in specific examples which we shall study later, these estimates can be significantly improved.

Lemma 5.4. *Suppose that $\tau \in \mathbb{R}$, $\epsilon > 0$, and \hat{c}_k , $0 \leq k \leq r$ is a normalized extended Chebyshev point as in (3.4), and $c_k = \tau + \frac{\epsilon}{2}(1 + \hat{c}_k)$. If $x \in [\tau, \tau + \epsilon]$, let $\hat{x} \in [-1, 1]$ denote the unique point such that $x = \tau + \frac{\epsilon}{2}[1 + \hat{x}]$. Assume that $M > 0$, $\Gamma = \{c_k : 0 \leq k \leq r\}$ and $F \in K_M(\Gamma) \setminus \{0\}$, so $|\ln(F(\xi)) - \ln(F(\eta))| \leq M|\xi - \eta|$ for all $\xi, \eta \in \Gamma$. Let $\mathcal{F} : [\tau, \tau + \epsilon] \rightarrow \mathbb{R}$ denote the unique polynomial of degree $\leq r$ such that $\mathcal{F}(c_k) = F(c_k)$ for $0 \leq k \leq r$. Let $\eta(r)$ be as in (5.2) and $\psi(r)$ as in Lemma 5.2 and define $u = M\epsilon\eta(r)$. If*

$$\psi(r)u \exp(u) < 1,$$

then $\mathcal{F}(x) > 0$ for all $x \in [\tau, \tau + \epsilon]$. If $\psi(r)u \exp(u) < 1$ and

$$C := \frac{[2\eta(r)r^2\psi(r)] \exp(u)M}{1 - \psi(r)u \exp(u)},$$

then for all $x, y \in [\tau, \tau + \epsilon]$,

$$|\ln(\mathcal{F}(x)) - \ln(\mathcal{F}(y))| \leq C|x - y|.$$

Proof. Recall that for $0 \leq k \leq r$,

$$l_k(x) = \frac{\prod_{\substack{l=0 \\ l \neq k}}^r (x - c_l)}{\prod_{\substack{l=0 \\ l \neq k}}^r (c_k - c_l)}, \quad \hat{l}_k(\hat{x}) = \frac{\prod_{\substack{l=0 \\ l \neq k}}^r (\hat{x} - \hat{c}_l)}{\prod_{\substack{l=0 \\ l \neq k}}^r (\hat{c}_k - \hat{c}_l)},$$

and $l_k(x) = \hat{l}_k(\hat{x})$ for $x = \tau + \frac{\epsilon}{2}[1 + \hat{x}]$ and, writing $F_k = F(c_k)$,

$$\mathcal{F}(x) = \sum_{k=0}^r l_k(x)F(c_k) = \sum_{k=0}^r l_k(x)F_k.$$

Recalling that $\sum_{k=0}^r l_k(x) = 1$ for all $x \in [\tau, \tau + \epsilon]$, we obtain

$$\begin{aligned} \mathcal{F}(x) &= \sum_{k=0}^r l_k(x)F_k = F_{[r/2]} \left(1 + \sum_{\substack{k=0 \\ k \neq [r/2]}}^r l_k(x) \left[\frac{F_k}{F_{[r/2]}} - 1 \right] \right) = F_{[r/2]}[1 + \phi(x)] \\ &= F_{[r/2]} \left(1 + \sum_{\substack{k=0 \\ k \neq [r/2]}}^r \hat{l}_k(\hat{x}) \left[\frac{F_k}{F_{[r/2]}} - 1 \right] \right) := F_{[r/2]}[1 + \hat{\phi}(\hat{x})], \end{aligned}$$

where as usual, $x = \tau + \frac{\epsilon}{2}[1 + \hat{x}]$, $\hat{x} \in [-1, 1]$.

Since $F \in K_M(\Gamma) \setminus \{0\}$, we have for $0 \leq k \leq r$, $k \neq [r/2]$,

$$\exp(-M|c_k - c_{[r/2]}|) \leq \frac{F_k}{F_{[r/2]}} \leq \exp(M|c_k - c_{[r/2]}|).$$

Because Lemma 5.3 (with $h_i = \epsilon$) implies that $|c_k - c_{[r/2]}| \leq \eta(r)\epsilon$,

$$\exp(-M\eta(r)\epsilon) - 1 \leq \frac{F_k}{F_{[r/2]}} - 1 \leq \exp(M\eta(r)\epsilon) - 1.$$

Using the mean value theorem and writing $u = M\eta(r)\epsilon$, it follows that

$$-u \leq \frac{F_k}{F_{[r/2]}} - 1 \leq u \exp(u),$$

so

$$\left| \frac{F_k}{F_{[r/2]}} - 1 \right| \leq u \exp(u).$$

Using Lemma 5.2, it follows that

$$(5.4) \quad |\hat{\phi}(\hat{x})| \leq \sum_{\substack{k=0 \\ k \neq [r/2]}}^r |\hat{l}_k(\hat{x})| \left| \frac{F_k}{F_{[r/2]}} - 1 \right| \leq \psi(r)u \exp(u),$$

so if $\psi(r)u \exp(u) < 1$, $1 + \hat{\phi}(\hat{x}) > 0$, and $\mathcal{F}(x) > 0$ for all $x \in [\tau, \tau + \epsilon]$. For the remainder of the proof, we assume that $\psi(r)u \exp(u) < 1$.

If $x, y \in [\tau, \tau + \epsilon]$, our previous calculations show that

$$\begin{aligned} |\ln \mathcal{F}(x) - \ln \mathcal{F}(y)| &= \left| \ln[1 + \phi(x)] - \ln[1 + \phi(y)] \right| \\ &= \left| \int_{1+\phi(y)}^{1+\phi(x)} \frac{1}{s} ds \right| \leq \left| \frac{\phi(x) - \phi(y)}{2} \right| \left[\frac{1}{1 + \phi(x)} + \frac{1}{1 + \phi(y)} \right], \end{aligned}$$

where we have used the fact that $(1/s)$ is a convex function and hence the integral is bounded by the trapezoidal rule approximation.

Now, by the mean value theorem, for some $\hat{\xi}$ lying between \hat{x} and \hat{y} and hence $\in [-1, 1]$,

$$\begin{aligned} |\phi(x) - \phi(y)| &= |\hat{\phi}(\hat{x}) - \hat{\phi}(\hat{y})| = |\hat{\phi}'(\hat{\xi})| |\hat{x} - \hat{y}| \\ &\leq \frac{2}{\epsilon} |x - y| \max_{-1 \leq \hat{\xi} \leq 1} |\hat{\phi}'(\hat{\xi})| \leq \frac{2}{\epsilon} |x - y| r^2 \max_{-1 \leq \hat{\xi} \leq 1} |\hat{\phi}(\hat{\xi})|, \end{aligned}$$

where in the last step we have used Markov's polynomial inequality (Lemma 5.2).

Recalling our earlier estimate for $|\hat{\phi}(\hat{\xi})|$ in (5.4), we obtain

$$|\phi(x) - \phi(y)| \leq 2r^2 \psi(r) \eta(r) M \exp(u) |x - y|$$

and

$$\frac{1}{2} \left[\frac{1}{1 + \phi(x)} + \frac{1}{1 + \phi(y)} \right] \leq \frac{1}{1 - \psi(r) u \exp(u)},$$

which implies that

$$|\ln(\mathcal{F}(x)) - \ln(\mathcal{F}(y))| \leq C |x - y|,$$

with the constant C defined in the statement of the lemma. \square

Lemma 5.5. *Let notation be as in Section 3 and T be as defined by (5.1). Suppose that $F : T \rightarrow \mathbb{R}$ is an element of $K_M(T) \setminus \{0\}$ and let $\mathcal{F} : \cup_{i=1}^I [a_i, b_i] \rightarrow \mathbb{R}$ be defined by (3.2). For $1 \leq i \leq I$, define $u_i = M h_i \eta(r)$ and assume that*

$$\psi(r) u_i \exp(u_i) < 1.$$

Then $\mathcal{F}(x) > 0$ for all $x \in [a_i, b_i]$. If we define C_i by

$$(5.5) \quad C_i := \frac{[2\eta(r)r^2\psi(r)] \exp(u_i) M}{1 - \psi(r) u_i \exp(u_i)},$$

then for all $x, y \in [a_i, b_i]$,

$$(5.6) \quad |\ln(\mathcal{F}(x)) - \ln(\mathcal{F}(y))| \leq C_i |x - y|.$$

Proof. Recall that $[a_i, b_i] = \cup_{j=1}^{N_i} [t_{j-1}^i, t_j^i]$, where $t_j^i - t_{j-1}^i = h_i = (b_i - a_i)/N_i$. If we write $h_i = \epsilon$, note that whenever $x, y \in [t_{j-1}^i, t_j^i]$ for some j , Lemma 5.5 implies that (5.6) is satisfied. Thus we can assume that $x, y \in [a_i, b_i]$ and that there does not exist j , $1 \leq j \leq N_i$, such that x and y are both elements of $[t_{j-1}^i, t_j^i]$. We can also assume that $x < y$ and select j_1 , $1 \leq j_1 \leq N_i$, such that $x \in [t_{j_1-1}^i, t_{j_1}^i]$ and j_2 , $1 \leq j_2 \leq N_i$, such that $y \in [t_{j_2-1}^i, t_{j_2}^i]$. By our assumptions, it must be true that $j_1 < j_2$. If we apply Lemma 5.5 to $\mathcal{F}(x)$ and $\mathcal{F}(t_{j_1}^i)$, we obtain

$$|\ln(\mathcal{F}(t_{j_1}^i)) - \ln(\mathcal{F}(x))| \leq C_i (t_{j_1}^i - x).$$

Similarly, if we apply Lemma 5.5 to $\mathcal{F}(t_{j_2-1}^i)$ and $\mathcal{F}(y)$, we obtain

$$|\ln(\mathcal{F}(y)) - \ln(\mathcal{F}(t_{j_2-1}^i))| \leq C_i(y - t_{j_2-1}^i).$$

Since $\mathcal{F}(t_{j_1}^i) = F(t_{j_1}^i)$ and $\mathcal{F}(t_{j_2-1}^i) = F(t_{j_2-1}^i)$ and $F \in K_M(T)$, we obtain

$$|\ln(\mathcal{F}(t_{j_2-1}^i)) - \ln(\mathcal{F}(t_{j_1}^i))| \leq M(t_{j_2-1}^i - t_{j_1}^i) \leq C_i(t_{j_2-1}^i - t_{j_1}^i),$$

where we have used the fact that $C_i > M$. Combining these inequalities, we find that

$$\begin{aligned} |\ln(\mathcal{F}(y)) - \ln(\mathcal{F}(x))| &\leq |\ln(\mathcal{F}(y)) - \ln(\mathcal{F}(t_{j_2-1}^i))| \\ &\quad + |\ln(\mathcal{F}(t_{j_2-1}^i)) - \ln(\mathcal{F}(t_{j_1}^i))| + |\ln(\mathcal{F}(t_{j_1}^i)) - \ln(\mathcal{F}(x))| \leq C_i(y - x), \end{aligned}$$

which proves Lemma 5.5. \square

Up to this point, we have only used the fact that $F \in K_M(T)$, where T is defined in (5.1) and notation is as in Section 3. We now exploit the fact that $\text{lip}(\theta_\omega|_S) \leq c(\nu)$ for all $\omega \in \Omega_\nu$.

Theorem 5.6. *Let notation be as in Section 3 and for positive reals $M' < M$, let $K_M(T)$ and $K_{M'}(T)$ be as defined earlier. Recall that $h_i = (b_i - a_i)/N_i$, $1 \leq i \leq I$, and $h = \max\{h_i : 1 \leq i \leq I\}$. Assume that hypotheses (H1), (H2), and (H3) are satisfied, and that*

$$(5.7) \quad \psi(r)u \exp u < 1,$$

where we now set

$$u = Mh\eta(r).$$

If $F \in K_M(T) \setminus \{0\}$ and \mathcal{F} is the piecewise polynomial approximation of F of degree $\leq r$ on $\hat{S} = \cup_{i=1}^I [a_i, b_i]$, then $\mathcal{F}(x) > 0$ for all $x \in \hat{S}$.

Define $C := \max\{C_i : 1 \leq i \leq I\}$, where C_i is defined by (5.5) and let $\mathbf{L}_{s,\nu} : V(T) := \mathbb{R}^Q \rightarrow V(T)$ be defined by (3.8). Assume the above hypotheses are satisfied and also assume that

$$(5.8) \quad c(\nu)C := \frac{c(\nu)[2\eta(r)r^2\psi(r)] \exp(u)M}{1 - \psi(r)u \exp(u)} < M' - sM_0(\nu).$$

Then it follows that $\mathbf{L}_{s,\nu}(K_M(T) \setminus \{0\}) \subset K_{M'}(T) \setminus \{0\}$.

Proof. Suppose that $F \in K_M(T) \setminus \{0\}$, which implies that $F(\xi) > 0$ for all $\xi \in T$. Since $u \geq u_i$ for $1 \leq i \leq I$, (5.7) implies that $\psi(r)u_i \exp u_i < 1$ for $1 \leq i \leq I$. It follows from Lemma 5.5 that $\mathcal{F}(x) > 0$ for all $x \in [a_i, b_i]$, $1 \leq i \leq I$, so $\mathcal{F}(x) > 0$ for all $x \in \hat{S}$. Since (H1) implies that $g_\omega(x)^s > 0$ for all $x \in S = [a, b]$ and for all $\omega \in \Omega_\nu$

$$\mathbf{L}_{s,\nu}(F)(\xi) = \sum_{\omega \in \Omega_\nu} [g_\omega(\xi)]^s \mathcal{F}(\theta_\omega(\xi)),$$

and $g_\omega(x)^s \mathcal{F}(\theta_\omega(x)) > 0$ for all $x \in \hat{S}$ and certainly for all $\xi \in T$, it suffices to prove that the map $\xi \mapsto g_\omega(\xi)^s \mathcal{F}(\theta_\omega(\xi)) \in K_{M'}(T) \setminus \{0\}$ for every $\omega \in \Omega_\nu$. We know that for all $x, y \in S$,

$$|\ln(g_\omega(x)^s) - \ln(g_\omega(y)^s)| \leq sM_0(\nu)|x - y|,$$

so it suffices to prove that for all $x, y \in \hat{S}$,

$$|\ln(\mathcal{F}(\theta_\omega(x))) - \ln(\mathcal{F}(\theta_\omega(y)))| \leq [M' - sM_0(\nu)] |x - y|.$$

For each fixed $\omega \in \Omega_\nu$, (H3) implies that there exists $i = i(\omega)$ such that $\theta_\omega(\hat{S}) \subset [a_i, b_i]$. Writing $x' = \theta_\omega(x) \in [a_i, b_i]$ and $y' = \theta_\omega(y) \in [a_i, b_i]$, Lemma 5.5 implies that

$$|\ln(\mathcal{F}(x')) - \ln(\mathcal{F}(y'))| \leq C|x' - y'| \leq c(\nu)C|x - y|,$$

so (5.8) completes the proof. \square

Remark 5.3. Assume that $\psi(r)u \exp(u) < 1$. Notice that for a given positive integer r , a necessary condition that (5.8) be satisfied is that

$$(5.9) \quad c(\nu)2r^2\psi(r)\eta(r) < \frac{M' - sM_0(\nu)}{M}.$$

For a given $M' < M$, if (5.9) is satisfied, then (5.8) will be satisfied if h is sufficiently small.

Remark 5.4. The reader will note that in our definition of $\mathbf{L}_{s,\nu}(F)$ for $F \in V(T)$, we arrange that $\mathcal{F}|_{[a_i, b_i]}$ is a piecewise polynomial map of degree $\leq r$. In some applications, it is desirable to make $\mathcal{F}|_{[a_i, b_i]}$ a piecewise polynomial map of degree $\leq r_i$, which leads to a generalization of the definition of $\mathbf{L}_{s,\nu}$. An analogue of Theorem 5.6 which handles this more general case can be proved by an argument similar to the proof of Theorem 5.6. Because of considerations of length, we omit the proof.

6. ESTIMATING $R(L_s)$ BY THE SPECTRAL RADIUS OF $\mathbf{L}_{s,\nu}$

In the previous section (c.f. Theorem 5.6), we determined conditions under which

$$\mathbf{L}_{s,\nu}(K_M(T) \setminus \{0\}) \subset K_{M'}(T) \setminus \{0\}$$

for some $M' < M$. The main result of this section is to show that under this condition, $R(\mathbf{L}_{s,\nu})$, the spectral radius of $\mathbf{L}_{s,\nu}$, satisfies

$$\lambda_s^\nu(1 - Hh^r) \leq R(\mathbf{L}_{s,\nu}) \leq \lambda_s^\nu(1 + Hh^r)$$

for some constant H to be specified below. Using this result, we obtain the following explicit bounds on the spectral radius λ_s of L_s .

$$[R([1 + Hh^r]^{-1}\mathbf{L}_{s,\nu})]^{1/\nu} \leq \lambda_s \leq [R([1 - Hh^r]^{-1}\mathbf{L}_{s,\nu})]^{1/\nu},$$

where the entries of the matrices $[1 + Hh^r]^{-1}\mathbf{L}_{s,\nu}$ and $[1 - Hh^r]^{-1}\mathbf{L}_{s,\nu}$ differ by $O(h^r)$.

Throughout this section, we shall assume the hypotheses and notation of (H1), (H2), and (H3); and $v_s(\cdot)$ will always denote the positive eigenvector v_s of L_s guaranteed by Theorem 2.1. In particular, S and $[a_i, b_i]$, $1 \leq i \leq I$ will be as in (H3) and (as can be guaranteed by relabeling), we shall assume that $b_i < a_{i+1}$ for $1 \leq i < I$.

We shall further denote by E and χ , constants for which the following two inequalities are satisfied.

$$(6.1) \quad \sup_{a \leq x \leq b} \frac{|d^p v_s(x)/dx^p|}{v_s(x)} \leq E(s, p) := E,$$

where p is a positive integer, and

$$(6.2) \quad v_s(x_1) \leq v_s(x_2) \exp(2s|x_1 - x_2|/\chi),$$

and for all $x_1, x_2 \in S$, where $\chi := \chi(s, \{\theta_i, g_i : 1 \leq i \leq n\})$. Using Theorem 2.1 and Remark 2.1, we shall see, in the next section, that for some interesting examples, it is possible to obtain sharp estimates on the constants E and χ such that (6.1) and (6.2) below are satisfied. These estimates will refine earlier results in [14].

Using the notation of Section 3, if $\mathcal{V}_s(x)$ is the piecewise polynomial interpolant of $v_s(x)$ of degree $\leq r$ at the extended Chebyshev points in $[a_i, b_i]$, $1 \leq i \leq I$, then on each subinterval $[t_{j-1}^i, t_j^i]$, $j = 1, \dots, N_i$, we have, using standard estimates for polynomial interpolation,

$$v_s(x) - \mathcal{V}_s(x) = \frac{v_s^{(r+1)}(\xi_x)}{(r+1)!} \prod_{k=0}^r (x - c_{j,k}^i),$$

for some $\xi_x \in [t_{j-1}^i, t_j^i]$. If we write, as done previously,

$$c_{j,k}^i = t_{j-1}^i + h_i(1 + \hat{c}_k)/2, \quad x = t_{j-1}^i + h_i(1 + \hat{x})/2, \quad \hat{x} \in [-1, 1],$$

then for $x \in [a_i, b_i]$,

$$|v_s(x) - \mathcal{V}_s(x)| \leq |v_s^{(r+1)}(\xi_x)| h_i^{r+1} m_{r+1},$$

where

$$(6.3) \quad m_{r+1} = \frac{1}{2^{r+1}(r+1)!} \max_{\hat{x} \in [-1, 1]} \left| \prod_{k=0}^r (\hat{x} - \hat{c}_k) \right|.$$

Using (6.1) and (6.2), we see that

$$|v_s^{(r+1)}(\xi_x)| \leq E \exp(2sh_i/\chi) v_s(x),$$

so

$$|v_s(x) - \mathcal{V}_s(x)| \leq E h_i^{r+1} m_{r+1} v_s(x) \exp(2sh_i/\chi).$$

Defining, for $1 \leq i \leq I$,

$$(6.4) \quad G_{r,i} := E m_r \exp(2sh_i/\chi),$$

(6.3) implies that for $1 \leq i \leq I$ and $x \in [a_i, b_i]$

$$(6.5) \quad (1 - G_{r+1,i} h_i^{r+1}) v_s(x) \leq \mathcal{V}_s(x) \leq (1 + G_{r+1,i} h_i^{r+1}) v_s(x).$$

In order to make (6.3) explicit, we need a formula for $\max_{\hat{x} \in [-1, 1]} \left| \prod_{k=0}^r (\hat{x} - \hat{c}_k) \right|$. The result and proof, which we provide below, are slight modifications of the well-known corresponding bound and proof when \hat{c}_k are taken to be the zeros of the standard Chebyshev polynomial.

Lemma 6.1. *If $r \geq 2$ is a positive integer and \hat{c}_k is defined by (3.4), then*

$$\max_{\hat{x} \in [-1, 1]} \left| \prod_{k=0}^r (\hat{x} - \hat{c}_k) \right| = \frac{1}{2^r} \left[\frac{1}{\cos(\pi/[2r+2])} \right]^{r+1}.$$

Proof. If we define $w = \hat{x} \cos(\pi/[2r+2])$, where $|\hat{x}| \leq 1$, we have $|w| \leq \cos(\pi/[2r+2])$. For notational convenience, we write $\alpha = 1/\cos(\pi/[2r+2])$, and we obtain

$$\begin{aligned} & \max_{\hat{x} \in [-1,1]} \left| \prod_{k=0}^r (\hat{x} - \hat{c}_k) \right| \\ &= \alpha^{r+1} \max \left\{ \left| \prod_{k=0}^r (w + \cos([2k+1]\pi/[2r+2])) \right| : |w| \leq \cos(\pi/[2r+2]) \right\} \\ &= \alpha^{r+1} \max \left\{ \left| \prod_{k=0}^r (w - \cos([2k+1]\pi/[2r+2])) \right| : |w| \leq \cos(\pi/[2r+2]) \right\}. \end{aligned}$$

If we define $q_{r+1}(w) = \prod_{k=0}^r (w - \cos([2k+1]\pi/[2r+2]))$, $q_{r+1}(w)$ is a polynomial of degree $r+1$ which vanishes at the points $\cos([2k+1]\pi/[2r+2])$, $0 \leq k \leq r$, and has leading term w^{r+1} ; and these properties uniquely determine $q_{r+1}(w)$.

Recall that for integers $r \geq 0$, $\cos([r+1]\theta) = p_{r+1}(\cos \theta)$ for $0 \leq \theta \leq \pi$, where $p_{r+1}(w)$ is the *Chebyshev* polynomial of degree $r+1$. These polynomials satisfy $p_1(w) = w$, $p_2(w) = 2w^2 - 1$, and for $r \geq 2$, the recurrence relation $p_{r+1}(w) = 2wp_r(w) - p_{r-1}(w)$. Using the recursion relation for $p_{r+1}(w)$ and induction, it also follows that the coefficient of w^{r+1} in $p_{r+1}(w)$ is 2^r . Since $\cos([r+1]\theta) = 0$ when $\theta = [2k+1]\pi/[2r+2]$ for $0 \leq k \leq r$, we see that $p_{r+1}(w) = 0$ when $w = \cos([2k+1]\pi/[2r+2])$, for $0 \leq k \leq r$. It follows that for $w = \cos(\theta)$ and $0 \leq \theta \leq \pi$,

$$q_{r+1}(w) = \frac{1}{2^r} p_{r+1}(w) = \frac{1}{2^r} \cos([r+1]\theta),$$

so

$$\max_{|w| \leq 1} |q_{r+1}(w)| = 1/2^r.$$

However,

$$\begin{aligned} & \max\{|q_{r+1}(w)| : |w| \leq \cos(\pi/[2r+2])\} \\ &= \frac{1}{2^r} \max\{|\cos([r+1]\theta)| : \pi/[2(r+1)^2] \leq \theta \leq (2r+1)\pi/(2r+2)\}. \end{aligned}$$

Since $\pi/[2(r+1)^2] < \pi/(r+1) < (2r+1)\pi/(2r+2)$ and $|\cos([r+1]\pi/(r+1))| = 1$,

$$\frac{1}{2^r} \max\{|q_{r+1}(w)| : |w| \leq \cos(\pi/[2r+2])\} = \frac{1}{2^r},$$

which completes the proof. \square

If E and χ are defined by (6.1) and (6.2), we can use Lemma 6.1 to estimate the constant $G_{r+1,i}$ in (6.4) more precisely:

$$(6.6) \quad G_{r+1,i} = E \exp(2sh_i/\chi) \left[\frac{1}{(r+1)!} \right] \left[\frac{1}{2 \cos(\pi/[2r+2])} \right]^{r+1} \frac{1}{2^r},$$

and with this estimate of $G_{r+1,i}$, (6.5) is satisfied for all $x \in [a_i, b_i]$, $1 \leq i \leq I$. For notational convenience, we define h and G_{r+1} by $h = \max_{1 \leq i \leq I} h_i$ and

$$(6.7) \quad G_{r+1} = \max_{1 \leq i \leq I} G_{r+1,i} = E \exp(2sh/\chi) \left[\frac{1}{(r+1)!} \right] \left[\frac{1}{2 \cos(\pi/[2r+2])} \right]^{r+1} \frac{1}{2^r}.$$

Lemma 6.2. Define $\lambda_s = R(\Lambda_s) = R(L_s)$, where Λ_s and L_s are as in Theorem 2.1. Let $[a_i, b_i]$, $1 \leq i \leq I$, be as in (H3). For $1 \leq i \leq I$, let N_i , h_i and \mathcal{V}_s be as defined in the fourth paragraph of this section. Assume that, for $1 \leq i \leq I$,

$$[\sin(\pi/(2r+2))]^2 h_i \leq a_{i+1} - b_i.$$

Define $h_{\min} = \min_{1 \leq i \leq I} h_i$ and $\mu = h/h_{\min}$. Then we have for all $x \in [a_i, b_i]$, $1 \leq i \leq I$,

$$(6.8) \quad (1 - G_{r+1,i} h_i^{r+1}) v_s(x) \leq \mathcal{V}_s(x) \leq (1 + G_{r+1,i} h_i^{r+1}) v_s(x),$$

and

$$(6.9) \quad (1 - G_{r+1} h^{r+1}) \lambda_s^\nu v_s(x) \leq (L_s^\nu \mathcal{V}_s)(x) \leq (1 + G_{r+1} h^{r+1}) \lambda_s^\nu v_s(x).$$

If we define M_1 by

$$(6.10) \quad M_1 = \left[\mu \frac{G_{r+1} h^r}{1 - G_{r+1}^2 h^{2r+2}} \right] \left[\frac{1}{\sin(\pi/[2r+2])} \right]^2 + \frac{2s}{\chi}$$

and

$$T = \{c_{j,k}^i : 1 \leq i \leq I, 1 \leq j \leq N_i, 0 \leq k \leq r\} \subset S,$$

then $\mathcal{V}_s|_T \in K(2s/\chi; T)$ and $L_s^\nu \mathcal{V}_s|_T = \mathbf{L}_{s,\nu}(\mathcal{V}_s|_T) \in K(M_1; T)$.

Proof. To simplify the exposition, we shall denote G_{r+1} as G . Equation (6.5) gives (6.8) and (6.2) implies that $v_s \in K(2s/\chi; S)$. Since $\mathcal{V}_s|_T = v_s|_T$, it follows that $\mathcal{V}_s|_T \in K(2s/\chi; S)$. If we observe that $1 - Gh^{r+1} \leq 1 - G_{r+1,i} h_i^{r+1}$ and $1 + G_{r+1,i} h_i^{r+1} \leq 1 + Gh^{r+1}$ for $1 \leq i \leq I$, we derive from (6.5) that for $1 \leq i \leq I$ and $x \in [a_i, b_i]$,

$$(1 - Gh^{r+1}) v_s(x) \leq \mathcal{V}_s(x) \leq (1 + Gh^{r+1}) v_s(x).$$

Applying L_s^ν to this inequality, we obtain (6.9) and in particular, (6.9) holds for all $x \in T$. A little thought shows that for all $x \in T$,

$$[L_s^\nu \mathcal{V}_s](x) = (\mathbf{L}_{s,\nu}(\mathcal{V}_s|_T))(x).$$

If $x, y \in T \cap [a_i, a_{i+1}]$, $1 \leq i \leq I$, and $x \neq y$, we obtain from (6.9) that

$$\begin{aligned} (L_s^\nu \mathcal{V}_s)(x) &\leq (1 + Gh^{r+1}) \lambda_s^\nu v_s(x) \leq (1 + Gh^{r+1}) \exp(2s|x-y|/\chi) v_s(y) \\ &\leq \frac{1 + Gh^{r+1}}{1 - Gh^{r+1}} \exp(2s|x-y|/\chi) (L_s^\nu \mathcal{V}_s)(y). \end{aligned}$$

Taking logarithms on both sides of the above inequality, and noting that x and y are interchangeable in the inequality, we find that

$$|\ln(L_s^\nu \mathcal{V}_s)(x) - \ln(L_s^\nu \mathcal{V}_s)(y)| \leq \frac{2s}{\chi} |x-y| + [\ln(1 + Gh^{r+1}) - \ln(1 - Gh^{r+1})].$$

Using the trapezoidal rule and the convexity of $u \mapsto 1/u$,

$$\begin{aligned} \ln(1 + Gh^{r+1}) - \ln(1 - Gh^{r+1}) &= \int_{1-Gh^{r+1}}^{1+Gh^{r+1}} \frac{1}{u} du \\ &\leq \frac{1}{2} \left[\frac{1}{1 - Gh^{r+1}} + \frac{1}{1 + Gh^{r+1}} \right] [2Gh^{r+1}] = \frac{2Gh^{r+1}}{1 - G^2 h^{2r+2}}. \end{aligned}$$

To prove that $L_s^\nu \mathcal{V}_s|_T \in K(M_1; T)$, it will suffice to prove that

$$(6.11) \quad \frac{2s}{\chi} |x - y| + \frac{2Gh^{r+1}}{1 - G^2h^{2r+2}} \leq \frac{2s}{\chi} |x - y| + \left[\mu \frac{Gh^r}{1 - G^2h^{2r+2}} \right] \left[\frac{1}{\sin(\pi/[2r+2])} \right]^2 |x - y|,$$

whenever $x, y \in ([a_i, b_i] \cap T) \cup \{a_{i+1}\}$ for $1 \leq i \leq I$ and whenever $x, y \in ([a_I, b_I] \cap T)$. (Of course, we assume, as we can, that $x \neq y$). A calculation shows that this will be true if

$$2h \leq \mu \left[\frac{1}{\sin(\pi/[2r+2])} \right]^2 |x - y|.$$

If $x, y \in [a_i, b_i]$, we know that $|x - y| \geq 2h_i [\sin(\pi/(2r+2))]^2$, so it suffices to prove that $h \leq \mu h_i$, which follows from the definition of μ . We can assume that $x < y$, so if $x, y \in [a_i, b_i]$, the same argument applies. If $y = a_{i+1}$, $|x - y| \geq |a_{i+1} - b_i|$, and we assume that $|a_{i+1} - b_i| \geq 2h_i [\sin(\pi/(2r+2))]^2$, so again the same argument applies and gives (6.11). \square

Remark 6.1. If $I = 1$, the condition on $a_{i+1} - b_i$ is vacuous and $\mu = 1$.

Our next lemma will play a crucial role in relating $R(\mathbf{L}_{s,\nu})$ to $R(L_s^\nu)$.

Lemma 6.3. Let notation and assumptions be as in Lemma 6.2. Let $G := G_{r+1}$ be as in (6.7) and M_1 as in (6.10). Assume that $H := H_{r+1}$ is a constant with $H > G$ and assume that $h < 1$. Define M_2 by

$$M_2 = M_1 + \frac{G}{H} \left[\frac{\mu}{1 - [(G/H)h]^2} \right] \frac{1}{[\sin(\pi/[2r+2])]^2}.$$

If $K = K(M_2; T)$, we have

$$(6.12) \quad \lambda_s^\nu \mathcal{V}_s(1 - Hh^r) \leq_K \mathbf{L}_{s,\nu} \mathcal{V}_s|_T \leq_K \lambda_s^\nu \mathcal{V}_s(1 + Hh^r).$$

Proof. Our previous results show that $(L_s^\nu \mathcal{V}_s)(x) = (\mathbf{L}_{s,\nu} \mathcal{V}_s)(x)$ for all $x \in T$, and, for all $x \in S = [a, b]$,

$$\lambda_s^\nu \mathcal{V}_s(x) \frac{1 - Gh^{r+1}}{1 + Gh^{r+1}} \leq (L_s^\nu \mathcal{V}_s)(x) \leq \lambda_s^\nu \mathcal{V}_s(x) \frac{1 + Gh^{r+1}}{1 - Gh^{r+1}}.$$

Recalling that $\mathcal{V}_s(x) = v_s(x)$ for $x \in T$, we have for $x \in T$,

$$\begin{aligned} \lambda_s^\nu \mathcal{V}_s(x)(1 + Hh^r) - (L_s^\nu \mathcal{V}_s)(x) &\leq \lambda_s^\nu \mathcal{V}_s(x)(1 + Hh^r) - \lambda_s^\nu(1 - Gh^{r+1})\mathcal{V}_s(x) \\ &= \lambda_s^\nu(Hh^r + Gh^{r+1})\mathcal{V}_s(x) = \lambda_s^\nu h^r \mathcal{V}_s(x)(1 + [G/H]h)H. \end{aligned}$$

If $y \in T$, a similar argument shows that

$$\begin{aligned} \lambda_s^\nu \mathcal{V}_s(y)(1 + Hh^r) - (L_s^\nu \mathcal{V}_s)(y) &\geq \lambda_s^\nu \mathcal{V}_s(y)(1 + Hh^r) - \lambda_s^\nu(1 + Gh^{r+1})\mathcal{V}_s(y) \\ &= \lambda_s^\nu h^r \mathcal{V}_s(y)(1 - [G/H]h)H. \end{aligned}$$

Using Lemma 6.2 and the above estimates, we find that

$$(6.13) \quad \frac{\lambda_s^\nu \mathcal{V}_s(x)(1 + Hh^r) - (L_s^\nu \mathcal{V}_s)(x)}{\lambda_s^\nu \mathcal{V}_s(y)(1 + Hh^r) - (L_s^\nu \mathcal{V}_s)(y)} \leq \frac{\mathcal{V}_s(x)(1 + [G/H]h)}{\mathcal{V}_s(y)(1 - [G/H]h)} \leq \exp(M_1|x - y|) \frac{1 + [G/H]h}{1 - [G/H]h}.$$

The right half of (6.12) will follow from (6.13) if we prove that, for all $x, y \in T$ with $x \neq y$,

$$(6.14) \quad \exp(M_1|x-y|) \frac{1 + [G/H]h}{1 - [G/H]h} \leq \exp(M_2|x-y|).$$

As in Lemma 6.2, it suffices to verify (6.14) for all points $x \neq y$, $x, y \in [a_i, a_{i+1}] \cap T$, $1 \leq i \leq I$, where $a_{I+1} = b_I$.

Arguing as in Lemma 6.2, we see that

$$\ln(1 + [G/H]h) - \ln(1 - [G/H]h) \leq \frac{G}{H} \frac{2h}{1 - ([G/H]h)^2}.$$

If we take the log of both sides of (6.14), it suffices to prove that

$$\begin{aligned} M_1|x-y| + \frac{G}{H} \frac{2h}{1 - ([G/H]h)^2} \\ \leq M_1|x-y| + \frac{G}{H} \left[\frac{\mu}{1 - ([G/H]h)^2} \right] \frac{1}{[\sin(\pi/[2r+2])]^2} |x-y|. \end{aligned}$$

As was proved in Lemma 6.2, all $x, y \in [a_i, a_{i+1}] \cap T$ with $x \neq y$ satisfy $|x-y| \geq 2h_i[\sin(\pi/(2r+2))]^2$. Since $2|x-y| \geq 2h_i/[\sin(\pi/[2r+2])]^2$, we see after simplification, the above inequality will be satisfied if $h \leq \mu h_i$, which holds by the definition of μ . This proves the right hand side of (6.12). The proof of the left hand side of inequality (6.12) follows by an exactly analogous argument and is left to the reader. \square

Our next theorem connects $R(\mathbf{L}_{s,\nu})$ and $R(L_s^\nu)$. To use the theorem, we shall need to estimate various constants, and we shall carry this out in the next section for an important class of examples.

Theorem 6.4. *Let notation and assumptions be as in Lemma 6.2 and let H and M_2 be as in Lemma 6.3. Assume that ν, h, s and r have been selected so that $(\mathbf{L}_{s,\nu}(K(M;T))) \subset K(M';T)$ where $0 < M' < M$ and $M \geq M_2$ (see Theorem 5.6). Then we have that $R(\mathbf{L}_{s,\nu})$, the spectral radius of $\mathbf{L}_{s,\nu}$, satisfies*

$$\lambda_s^\nu(1 - Hh^r) \leq R(\mathbf{L}_{s,\nu}) \leq \lambda_s^\nu(1 + Hh^r).$$

Proof. Our previous results show that $\mathbf{L}_{s,\nu}$ has a unique, strictly positive eigenvector $w_{s,\nu} \in K(M;T)$ with $\|w_{s,\nu}\| = 1$. The eigenvalue corresponding to $w_{s,\nu}$ is $R(\mathbf{L}_{s,\nu})$. Furthermore, for every $u \in K(M;T) \setminus \{0\}$, $\lim_{m \rightarrow \infty} (\mathbf{L}_{s,\nu}^m u / \|\mathbf{L}_{s,\nu}^m u\|) = w_{s,\nu}$, with convergence in the sup norm topology on \mathbb{R}^Q .

If we use (6.12), but define $K = K(M;T)$, then because $M \geq M_2$ and $K(M;T) \supset K(M_2;T)$, we obtain

$$\lambda_s^\nu \mathcal{V}_s(1 - Hh^r) \leq_K \mathbf{L}_{s,\nu} \mathcal{V}_s \leq_K \lambda_s^\nu \mathcal{V}_s(1 + Hh^r).$$

The theorem now follows directly from Lemma 4.1. \square

Our ultimate goal has been to provide rigorous upper and lower bounds on $\lambda_s = R(L_s)$ in terms of the eigenvalues of computable matrices, as was done in [14]. This follows immediately from Theorem 6.4.

Theorem 6.5. *Under the hypotheses of Theorem 6.4, we have*

$$[(1 + Hh^r)^{-1}R(\mathbf{L}_{s,\nu})]^{1/\nu} \leq \lambda_s \leq [(1 - Hh^r)^{-1}R(\mathbf{L}_{s,\nu})]^{1/\nu},$$

where the entries of the matrices $[1 + Hh^r]^{-1}\mathbf{L}_{s,\nu}$ and $[1 - Hh^r]^{-1}\mathbf{L}_{s,\nu}$ differ by $O(h^r)$.

7. CALCULATING THE OPTIMAL INTERVAL $[a, b]$ AND ESTIMATING E AND χ .

Throughout this section, we shall assume at least the hypotheses of Theorem 2.1, so L_s has a strictly positive C^m eigenfunction v_s . We shall take $S = [a, b]$, $a < b$ in (H2). If S_0 is a closed, nonempty subset of S and $\theta_i(S_0) \subset S_0$ for $1 \leq i \leq n$, then $L_s : C(S) \rightarrow C(S)$ induces a bounded linear operator $L_{s,S_0} : C(S_0) \rightarrow C(S_0)$. It is often desirable to replace the original interval $[a, b]$ by a smaller interval (or union of intervals) $S_0 \subset [a, b]$ such that $\theta_i(S_0) \subset S_0$ for $1 \leq i \leq n$, and we first describe a class of examples for which this can be easily done. Note that $v_s|_{S_0}$ is strictly positive; and since $L_{s,S_0}(v_s|_{S_0}) = R(L_s)(v_s|_{S_0})$, the following lemma implies that $R(L_{s,S_0}) = R(L_s)$. Although this is a special case of another well-known result, a proof is provided for the readers' convenience.

Lemma 7.1. *Let S_0 be a compact metric space, $W := C(S_0)$ and $P = \{f \in W : f(t) \geq 0, \forall t \in S_0\}$. Assume that $L : W \rightarrow W$ is a bounded linear operator such that $L(P) \subset P$. If $w \in W$ and $w(t) > 0$ for all $t \in S_0$, then*

$$R(L) = \lim_{k \rightarrow \infty} \|L^k\|^{1/k} = \lim_{k \rightarrow \infty} \|L^k w\|^{1/k}.$$

If, in addition, $Lw = \lambda w$, then $\lambda = R(L)$; and there exists a constant $C \geq 1$ such that $\|L^k\| \leq C\lambda^k$ for all positive integers k .

Proof. Since S_0 is compact, there exists $\alpha > 0$ such that $w(t) \geq \alpha$ for all $t \in S_0$. If $f \in W$ and $\|f\| \leq 1$, it follows that for all $t \in S_0$,

$$-(1/\alpha)w(t) \leq f(t) \leq (1/\alpha)w(t).$$

Because L is order-preserving in the partial ordering from P ,

$$-(1/\alpha)(L^k w)(t) \leq (L^k f)(t) \leq (1/\alpha)(L^k w)(t)$$

for all positive integers k , which implies that

$$\|L^k\|^{1/k} = \left(\sup\{\|L^k f\| : f \in W, \|f\| \leq 1\} \right)^{1/k} \leq (1/\alpha)^{1/k} \|L^k w\|^{1/k}.$$

We also have that

$$(1/\alpha)^{1/k} \|L^k w\|^{1/k} \leq (1/\alpha)^{1/k} \|w\|^{1/k} \|L^k\|^{1/k}.$$

Since $\lim_{k \rightarrow \infty} \|L^k\|^{1/k} = R(L)$ and $\lim_{k \rightarrow \infty} (1/\alpha)^{1/k} = \lim_{k \rightarrow \infty} \|w\|^{1/k} = 1$, we conclude that $\lim_{k \rightarrow \infty} \|L^k w\|^{1/k} = R(L)$.

If $Lw = \lambda w$, the above argument shows that

$$(1/\alpha)\lambda^k w(t) \leq (L^k f)(t) \leq (1/\alpha)\lambda^k w(t),$$

which implies $\|L^k\| \leq \frac{1}{\alpha} \|w\| \lambda^k$, so the lemma is satisfied with $C = \frac{1}{\alpha} \|w\| \geq 1$. \square

Let $S_0 = [\mathbf{a}_0, \mathbf{b}_0]$ be a compact interval of reals, $\mathbf{a}_0 < \mathbf{b}_0$, and let \mathcal{B} be a finite set of real numbers. For each $\beta \in \mathcal{B}$, $\theta_\beta : S_0 \rightarrow \mathbb{R}$. We make the following hypothesis:

- (H4) (i) For each $\beta \in \mathcal{B}$, $\theta_\beta : S_0 \rightarrow S_0$ and θ_β is a continuous map.
- (ii) There exists $\gamma \in \mathcal{B}$ and $\Gamma \in \mathcal{B}$ such that for all $x \in S_0$ and all $\beta \in \mathcal{B}$, $\theta_\Gamma(x) \leq \theta_\beta(x) \leq \theta_\gamma(x)$.
- (iii) $x \mapsto \theta_\gamma(x)$ and $x \mapsto \theta_\Gamma(x)$ are strictly decreasing functions on S_0 .

The example we have in mind is that \mathcal{B} is a finite set of distinct real numbers with $\beta \geq \gamma > 0$ for all $\beta \in \mathcal{B}$ and $\theta_\beta(x) = 1/(x + \beta)$ and $S_0 = [0, 1/\gamma]$, but there seems no gain in specializing at this point.

Lemma 7.2. *Assume (H4). Define $\mathbf{a}_1 = \theta_\Gamma(\mathbf{b}_0)$ and $\mathbf{b}_1 = \theta_\gamma(\mathbf{a}_0)$. Then $\mathbf{a}_0 \leq \mathbf{a}_1 \leq \mathbf{b}_1 \leq \mathbf{b}_0$, $\mathbf{a}_1 < \mathbf{b}_1$, and $\theta_\beta(x) \in [\mathbf{a}_1, \mathbf{b}_1]$ for all $x \in S_0$ and all $\beta \in \mathcal{B}$.*

Proof. Property (i) in (H4) implies that $\mathbf{a}_0 \leq \mathbf{a}_1 \leq \mathbf{b}_0$ and $\mathbf{a}_0 \leq \mathbf{b}_1 \leq \mathbf{b}_0$. Property (ii) implies that $\theta_\Gamma(\mathbf{b}_0) = \mathbf{a}_1 \leq \theta_\gamma(\mathbf{b}_0)$ and Property (iii) implies that $\theta_\gamma(\mathbf{b}_0) < \theta_\gamma(\mathbf{a}_0) = \mathbf{b}_1$, so $\mathbf{a}_1 < \mathbf{b}_1$. For all $x \in [\mathbf{a}_0, \mathbf{b}_0]$ and all $\beta \in \mathcal{B}$, $\theta_\Gamma(x) \leq \theta_\beta(x)$ (Property (ii)) and $\theta_\Gamma(\mathbf{b}_0) \leq \theta_\Gamma(x)$ (Property (iii)), so $\mathbf{a}_1 \leq \theta_\beta(x)$. Similarly, $\theta_\beta(x) \leq \theta_\gamma(x)$ and $\theta_\gamma(x) \leq \theta_\gamma(\mathbf{a}_0) = \mathbf{b}_1$, so $\theta_\beta(x) \leq \mathbf{b}_1$. \square

Lemma 7.3. *Assume (H4). Also assume that for $1 \leq j \leq k$, we have found an increasing sequence of reals $\mathbf{a}_0 \leq \mathbf{a}_1 \leq \dots \leq \mathbf{a}_k$ and a decreasing sequence of reals $\mathbf{b}_0 \geq \mathbf{b}_1 \geq \dots \geq \mathbf{b}_k$ such that $\mathbf{b}_j - \mathbf{a}_j > 0$ for $1 \leq j \leq k$ and $\theta_\beta([\mathbf{a}_j, \mathbf{b}_j]) \subset [\mathbf{a}_{j+1}, \mathbf{b}_{j+1}]$ for $0 \leq j \leq k-1$ and all $\beta \in \mathcal{B}$. Define $\mathbf{a}_{k+1} = \theta_\Gamma(\mathbf{b}_k)$ and $\mathbf{b}_{k+1} = \theta_\gamma(\mathbf{a}_k)$. Then we have $\mathbf{a}_k \leq \mathbf{a}_{k+1}$, $\mathbf{b}_{k+1} \leq \mathbf{b}_k$, $\mathbf{b}_{k+1} - \mathbf{a}_{k+1} > 0$ and $\theta_\beta([\mathbf{a}_k, \mathbf{b}_k]) \subset [\mathbf{a}_{k+1}, \mathbf{b}_{k+1}]$ for all $\beta \in \mathcal{B}$.*

Proof. Apply Lemma 7.2 with $[\mathbf{a}_k, \mathbf{b}_k]$ taking the place of $[\mathbf{a}_0, \mathbf{b}_0]$ and $\theta_\Gamma(\mathbf{b}_k) = \mathbf{a}_{k+1}$ taking the place of \mathbf{a}_1 and $\theta_\gamma(\mathbf{a}_k) = \mathbf{b}_{k+1}$ taking the place of \mathbf{b}_1 . \square

It follows from Lemma 7.3 that if (H4) holds and if we inductively define sequences \mathbf{a}_k and \mathbf{b}_k by $\mathbf{a}_{k+1} = \theta_\Gamma(\mathbf{b}_k)$ and $\mathbf{b}_{k+1} = \theta_\gamma(\mathbf{a}_k)$ for $k \geq 0$, then for all integers $k \geq 0$, $\mathbf{a}_k < \mathbf{b}_k$, $\mathbf{a}_{k+1} \geq \mathbf{a}_k$, $\mathbf{b}_{k+1} \leq \mathbf{b}_k$, and $\theta_\beta([\mathbf{a}_k, \mathbf{b}_k]) \subset [\mathbf{a}_{k+1}, \mathbf{b}_{k+1}]$ for all $\beta \in \mathcal{B}$. It follows that $\lim_{k \rightarrow \infty} \mathbf{a}_k := \mathbf{a}_\infty$ and $\lim_{k \rightarrow \infty} \mathbf{b}_k := \mathbf{b}_\infty$ both exist.

Lemma 7.4. *Assume (H4) and let notation be as above. Then $\theta_\beta([\mathbf{a}_\infty, \mathbf{b}_\infty]) \subset [\mathbf{a}_\infty, \mathbf{b}_\infty]$ for all $\beta \in \mathcal{B}$ and $\theta_\gamma(\mathbf{a}_\infty) = \mathbf{b}_\infty$ and $\theta_\Gamma(\mathbf{b}_\infty) = \mathbf{a}_\infty$, so $\theta_\Gamma \circ \theta_\gamma(\mathbf{a}_\infty) = \mathbf{a}_\infty$ and $\theta_\gamma \circ \theta_\Gamma(\mathbf{b}_\infty) = \mathbf{b}_\infty$. If $\beta_1, \beta_2, \dots, \beta_k$ are elements of \mathcal{B} and $x \in [\mathbf{a}_0, \mathbf{b}_0]$, then $(\theta_{\beta_1} \circ \theta_{\beta_2} \circ \dots \circ \theta_{\beta_k})(x) \in [\mathbf{a}_k, \mathbf{b}_k]$. If $(\theta_{\beta_1} \circ \theta_{\beta_2} \circ \dots \circ \theta_{\beta_k})(x) = x$ for some $x \in [\mathbf{a}_0, \mathbf{b}_0]$ and some elements $\beta_1, \beta_2, \dots, \beta_k$ of \mathcal{B} , then $x \in [\mathbf{a}_\infty, \mathbf{b}_\infty]$.*

Proof. Because $\lim_{k \rightarrow \infty} \mathbf{a}_k := \mathbf{a}_\infty$, $\lim_{k \rightarrow \infty} \mathbf{b}_k := \mathbf{b}_\infty$, $\theta_\gamma(\mathbf{a}_k) = \mathbf{b}_{k+1}$, and $\theta_\Gamma(\mathbf{b}_k) = \mathbf{a}_{k+1}$, it follows from the continuity of θ_γ and θ_Γ that $\theta_\gamma(\mathbf{a}_\infty) = \mathbf{b}_\infty$ and $\theta_\Gamma(\mathbf{b}_\infty) = \mathbf{a}_\infty$.

If $x \in [\mathbf{a}_0, \mathbf{b}_0]$ and $\beta_1, \beta_2, \dots, \beta_k$ are elements of \mathcal{B} , repeated applications of Lemma 7.2 show that $\theta_{\beta_k}(x) \in [\mathbf{a}_1, \mathbf{b}_1]$, $(\theta_{\beta_{k-1}} \circ \theta_{\beta_k})(x) \in [\mathbf{a}_2, \mathbf{b}_2]$, and generally

that $(\theta_{\beta_1} \circ \theta_{\beta_2} \circ \cdots \circ \theta_{\beta_k})(x) \in [\mathbf{a}_k, \mathbf{b}_k]$. If $x \in [\mathbf{a}_0, \mathbf{b}_0]$ and $\beta_1, \beta_2, \dots, \beta_k$ are elements of \mathcal{B} are such that $(\theta_{\beta_1} \circ \theta_{\beta_2} \circ \cdots \circ \theta_{\beta_k})(x) = x$, it follows that $x \in [\mathbf{a}_k, \mathbf{b}_k]$. Now the same argument can be repeated to show that $x \in [\mathbf{a}_{2k}, \mathbf{b}_{2k}]$ and generally that $x \in [\mathbf{a}_{mk}, \mathbf{b}_{mk}]$ for every positive integer m . Since $\bigcap_{m \geq 1} [\mathbf{a}_{mk}, \mathbf{b}_{mk}] = [\mathbf{a}_\infty, \mathbf{b}_\infty]$, we conclude that $x \in [\mathbf{a}_\infty, \mathbf{b}_\infty]$. \square

Remark 7.1. *Under the hypotheses of Lemma 7.4, if $(\theta_\Gamma \circ \theta_\gamma)(x) = x$ or $(\theta_\gamma \circ \theta_\Gamma)(x) = x$ for some $x \in [\mathbf{a}_0, \mathbf{b}_0]$, then $\mathbf{a}_\infty \leq x \leq \mathbf{b}_\infty$, so \mathbf{a}_∞ is the least fixed point of $\theta_\Gamma \circ \theta_\gamma$ in $[\mathbf{a}_0, \mathbf{b}_0]$ and \mathbf{b}_∞ is the greatest fixed point of $\theta_\gamma \circ \theta_\Gamma$ in $[\mathbf{a}_0, \mathbf{b}_0]$.*

Lemma 7.4 provides a way of obtaining invariant intervals J , such that $\theta_\beta(J) \subset J$ for all $\beta \in \mathcal{B}$. However, it is frequently the case that we have more information than given in (H4), and then one can give more flexible methods to find invariant intervals. The following lemma, whose proof we omit, describes a commonly occurring class of examples where such methods are available.

Lemma 7.5. *Let hypotheses and notation be as in Lemma 7.4. Suppose also that there exist intervals $J_1 = [x_1, \mathbf{a}_\infty]$ and $J_2 = [\mathbf{b}_\infty, x_2]$ with $\mathbf{a}_0 \leq x_1 < \mathbf{a}_\infty$ and $\mathbf{b}_\infty < x_2 < \mathbf{b}_0$, such that $\theta_\gamma(J_2) \subset [\mathbf{a}_\infty, \mathbf{b}_\infty]$, $\theta_\Gamma(J_1) \subset [\mathbf{a}_\infty, \mathbf{b}_\infty]$, $\text{lip}(\theta_\gamma|_{J_1}) \leq c_1$, $\text{lip}(\theta_\Gamma|_{J_2}) \leq c_2$, and $c_1 c_2 < 1$. If $\xi_1 \in J_1$ is chosen so that $\xi_2 = \theta_\gamma(\xi_1) \in J_2$, then $\theta_\beta([\xi_1, \xi_2]) \subset [\xi_1, \xi_2]$ for all $\beta \in \mathcal{B}$. Similarly, if $\eta_2 \in J_2$ is chosen so that $\eta_1 = \theta_\Gamma(\eta_2) \in J_1$, then $\theta_\beta([\eta_1, \eta_2]) \subset [\eta_1, \eta_2]$ for all $\beta \in \mathcal{B}$.*

We shall use \mathcal{B} as an index set, so the operator L_s can be written

$$(L_s f)(x) = \sum_{\beta \in \mathcal{B}} g_\beta(x)^s f(\theta_\beta(x)),$$

where $\theta_\beta(S_0) \subset S_0$ for all $\beta \in \mathcal{B}$ and $S_0 = [\mathbf{a}_0, \mathbf{b}_0]$. If the conditions of Theorem 2.1 are satisfied, L_s has a strictly positive, C^m eigenfunction. Assuming (H4) and the hypotheses of Theorem 2.1, the observation in the first paragraph of this section implies that to compute $R(L_s)$, we can, in the notation of Lemma 7.4, replace $[\mathbf{a}_0, \mathbf{b}_0]$ by $[\mathbf{a}_\infty, \mathbf{b}_\infty]$ or by $[\mathbf{a}_k, \mathbf{b}_k]$ for any integer $k \geq 1$. In fact, we could use any interval $J \subset [\mathbf{a}_0, \mathbf{b}_0]$ with $\theta_\beta(J) \subset J$ for all $\beta \in \mathcal{B}$ (compare Lemma 7.5).

For the remainder of this section, we shall assume

(H5) \mathcal{B} is a finite set of distinct real numbers and $\gamma = \min\{\beta : \beta \in \mathcal{B}\} \geq 1$. For every $\beta \in \mathcal{B}$, we define $\theta_\beta : [0, 1/\gamma] \rightarrow [0, 1/\gamma]$ by $\theta_\beta(x) = (x + \beta)^{-1}$.

We shall write $\Gamma = \max\{\beta : \beta \in \mathcal{B}\}$ and $\gamma = \min\{\beta : \beta \in \mathcal{B}\}$ and always assume that $\gamma < \Gamma$. The reader can check that $\{\theta_\beta : \beta \in \mathcal{B}\}$ satisfies the conditions of (H4) with $\theta_\gamma(x) = (x + \gamma)^{-1}$ and $\theta_\Gamma(x) = (x + \Gamma)^{-1}$. Using the calculations in the following paragraph, the reader can check that the conditions of Lemma 7.5 are also satisfied.

We assume that the sequences $\{\mathbf{a}_k : k \geq 1\}$ and $\{\mathbf{b}_k : k \geq 1\}$ are defined as in Lemmas 7.3 and 7.4, with $\mathbf{a}_0 = 0$ and $\mathbf{b}_0 = 1/\gamma$, and \mathbf{a}_∞ and \mathbf{b}_∞ defined as in Lemma 7.4. Since \mathbf{a}_∞ is a fixed point of $\theta_\Gamma \circ \theta_\gamma$ in $[0, 1/\gamma]$ and \mathbf{b}_∞ is a fixed point of $\theta_\gamma \circ \theta_\Gamma$ in $[0, 1/\gamma]$, one can easily solve the equations

$$x = (\theta_\Gamma \circ \theta_\gamma)(x) = \frac{x + \gamma}{\Gamma x + (1 + \Gamma \gamma)} \quad \text{and} \quad x = (\theta_\gamma \circ \theta_\Gamma)(x) = \frac{x + \Gamma}{\gamma x + (1 + \Gamma \gamma)}$$

to obtain

$$(7.1) \quad \mathfrak{a}_\infty = -\frac{\gamma}{2} + \sqrt{(\gamma/2)^2 + (\gamma/\Gamma)} \quad \text{and} \quad \mathfrak{b}_\infty = -\frac{\Gamma}{2} + \sqrt{(\Gamma/2)^2 + (\Gamma/\gamma)}.$$

One can verify that $\mathfrak{b}_\infty = (\Gamma/\gamma)\mathfrak{a}_\infty$, so $0 < \mathfrak{a}_\infty < \mathfrak{b}_\infty < 1/\gamma$.

Since our index set is \mathcal{B} , we slightly abuse previous notation and, for a positive integer ν , we define the set of ordered ν -tuples of elements of \mathcal{B} by

$$\Omega_\nu = \{(\beta_1, \beta_2, \dots, \beta_\nu) : \beta_j \in \mathcal{B} \text{ for } 1 \leq j \leq \nu\}.$$

For each $\omega = (\beta_1, \beta_2, \dots, \beta_\nu) \in \Omega_\nu$, we define $\theta_\omega = \theta_{\beta_1} \circ \theta_{\beta_2} \circ \dots \circ \theta_{\beta_\nu}$. Our first task is to estimate $c(\nu)$ (see (5.3)), which gives an upper bound for $\text{lip}(\theta_\omega)$, $\omega \in \Omega_\nu$.

If $\omega = (\beta_1, \beta_2, \dots, \beta_\nu) \in \Omega_\nu$ and $\beta \in \mathcal{B}$, define a matrix

$$M_\beta = \begin{pmatrix} 0 & 1 \\ 1 & \beta \end{pmatrix}.$$

It is proved in Section 6 of [14] that

$$M = M_{\beta_1} M_{\beta_2} \dots M_{\beta_\nu} = \begin{pmatrix} A_{\nu-1} & A_\nu \\ B_{\nu-1} & B_\nu \end{pmatrix},$$

where A_j and B_j are defined inductively by $A_0 = 0$, $A_1 = 1$, $B_0 = 1$, $B_1 = \beta_1$, and generally, for $1 \leq j \leq \nu$, by

$$(7.2) \quad A_{j+1} = A_{j-1} + \beta_{j+1}A_j, \quad B_{j+1} = B_{j-1} + \beta_{j+1}B_j.$$

Note that $\det(M_\beta) = -1$ so $\det(M) = (-1)^\nu$. Standard results for Möbius transforms now imply that for $x \in [\mathfrak{a}_k, \mathfrak{b}_k]$, $0 \leq k \leq \infty$,

$$(7.3) \quad \begin{aligned} (\theta_{\beta_1} \circ \theta_{\beta_2} \circ \dots \circ \theta_{\beta_\nu})(x) &= \frac{A_{\nu-1}x + A_\nu}{B_{\nu-1}x + B_\nu}, \\ \frac{d}{dx}(\theta_{\beta_1} \circ \theta_{\beta_2} \circ \dots \circ \theta_{\beta_\nu})(x) &= \frac{(-1)^\nu}{(B_{\nu-1}x + B_\nu)^2}. \end{aligned}$$

If we define $\tilde{B}_0 = 1$, $\tilde{B}_1 = \gamma$, and $\tilde{B}_{j+1} = \tilde{B}_{j-1} + \gamma\tilde{B}_j$ for $j \geq 1$, then because $\gamma \leq \beta$ for all $\beta \in \mathcal{B}$, it is straightforward to prove that $\tilde{B}_j \leq B_j$ for $0 \leq j \leq \nu$, where B_j is defined by (7.2). It follows that for all $\omega \in \Omega_\nu$ and $x \in [\mathfrak{a}_k, \mathfrak{b}_k]$,

$$|\theta'_\omega(x)| \leq [\tilde{B}_{\nu-1}\mathfrak{a}_k + \tilde{B}_\nu]^{-2},$$

which implies that, for $\theta_\omega : [\mathfrak{a}_k, \mathfrak{b}_k] \rightarrow \mathbb{R}$,

$$(7.4) \quad \max\{\text{lip}(\theta_\omega) : \omega \in \Omega_\nu\} = [\tilde{B}_{\nu-1}\mathfrak{a}_k + \tilde{B}_\nu]^{-2} := c(\nu).$$

It remains to give an exact formula for the right hand side of (7.4). The linear difference equation $\tilde{B}_{j+1} = \tilde{B}_{j-1} + \gamma\tilde{B}_j$ has solutions of the form λ^j for $j \geq 0$, which leads to the formula $\lambda^{n+1} = \lambda^{n-1} + \gamma\lambda^n$, or for $\lambda \neq 0$, $\lambda^2 = 1 + \lambda\gamma$. Hence

$$(7.5) \quad \lambda = \lambda_+ = \frac{\gamma}{2} + \frac{1}{2}\sqrt{\gamma^2 + 4}, \quad \lambda = \lambda_- = \frac{\gamma}{2} - \frac{1}{2}\sqrt{\gamma^2 + 4}.$$

The general solution of the difference equation is then

$$(7.6) \quad c_1\lambda_+^j + c_2\lambda_-^j = \tilde{B}_j, \quad j \geq 0,$$

where c_1 and c_2 must be chosen so that $\tilde{B}_0 = 1$ and $\tilde{B}_1 = \gamma$. A calculation gives

$$(7.7) \quad c_1 = \frac{\sqrt{\gamma^2 + 4} + \gamma}{2\sqrt{\gamma^2 + 4}}, \quad c_2 = \frac{\sqrt{\gamma^2 + 4} - \gamma}{2\sqrt{\gamma^2 + 4}}.$$

Summarizing the above discussion, we obtain

Lemma 7.6. *Assume (H4) and consider θ_β , $\beta \in \mathcal{B}$, as a map of $[\mathbf{a}_k, \mathbf{b}_k]$ to itself, for $0 \leq k \leq \infty$, where $\mathbf{a}_0 = 0$, $\mathbf{b}_0 = 1$, $\mathbf{a}_k = \theta_\Gamma(\mathbf{b}_{k-1})$ and $\mathbf{b}_k = \theta_\gamma(\mathbf{a}_{k-1})$ for $k \geq 1$ and \mathbf{a}_∞ and \mathbf{b}_∞ are given by (7.1). Then for $j \geq 1$, \tilde{B}_j is given by (7.6), where λ_+ and λ_- are given by (7.5) and c_1 and c_2 by (7.7).*

Remark 7.2. *Because $\lambda_+ > 1$ and $-1 < \lambda_- = -1/\lambda_+ < 0$ for all $\gamma > 0$, $c_1 \lambda_+^j$ is the dominant term in (7.6) as j increases; and one can check that $|c_2 \lambda_-^j| < 1/2$ for all $j \geq 0$. Of course, for moderate values of j , one can easily compute \tilde{B}_j from its recurrence formula. It is clear that the constant $c(\nu)$ in (7.4) is minimized by working on the interval $[\mathbf{a}_\infty, \mathbf{b}_\infty]$.*

Assuming (H5), we now define for $s > 0$, $L_s : C([0, 1]) \rightarrow C([0, 1])$ by

$$(L_s f)(x) = \sum_{\beta \in \mathcal{B}} |\theta'_\beta(x)|^s f(\theta_\beta(x)) := \sum_{\beta \in \mathcal{B}} g_\beta(x)^s f(\theta_\beta(x)).$$

It is well-known that for ν a positive integer,

$$(L_s^\nu f)(x) = \sum_{\omega \in \Omega_\nu} |\theta'_\omega(x)|^s f(\theta_\omega(x)) := \sum_{\omega \in \Omega_\nu} g_\omega(x)^s f(\theta_\omega(x)).$$

Because $\theta_\omega([\mathbf{a}_k, \mathbf{b}_k]) \subset [\mathbf{a}_k, \mathbf{b}_k]$ for $0 \leq k \leq \infty$ and $\omega \in \Omega_\nu$, we can also consider L_s^ν as a map of $C([\mathbf{a}_k, \mathbf{b}_k]) \mapsto C([\mathbf{a}_k, \mathbf{b}_k])$ and as noted earlier, this does not change the spectral radius of L_s^ν . Thus, we shall consider L_s^ν as a map from $C([\mathbf{a}_k, \mathbf{b}_k])$ into itself, with optimal results obtained by taking $k = \infty$.

We need to find a constant $M_0(\nu)$ (compare (5.3)) such that for all $\omega \in \Omega_\nu$, $g_\omega(x) := |\theta'_\omega(x)| \in K(M_0(\nu); [\mathbf{a}_k, \mathbf{b}_k])$. In this case, this is equivalent to proving that for all $\omega \in \Omega_\nu$, $x \mapsto \ln(|\theta'_\omega(x)|)$ is a Lipschitz map on $[\mathbf{a}_k, \mathbf{b}_k]$ with Lipschitz constant $\leq M_0(\nu)$. If $\omega = (\beta_1, \beta_2, \dots, \beta_\nu)$, we have by (7.3), that

$$|\theta'_\omega(x)| = \frac{1}{(B_{\nu-1}x + B_\nu)^2},$$

so

$$\ln(|\theta'_\omega(x)|) = -2 \ln(B_{\nu-1}x + B_\nu).$$

Thus it suffices to choose $M_0(\nu)$ so that for all $x \in [\mathbf{a}_k, \mathbf{b}_k]$ and all $\omega \in \Omega_\nu$,

$$\begin{aligned} \left| \frac{d}{dx} \ln(|\theta'_\omega(x)|) \right| &= 2 \frac{B_{\nu-1}}{B_{\nu-1}x + B_\nu} \\ &= 2 \left[\frac{1}{x + (B_\nu/B_{\nu-1})} \right] \leq 2 \left[\frac{1}{\mathbf{a}_k + (B_\nu/B_{\nu-1})} \right] \leq M_0(\nu). \end{aligned}$$

If we define $x_j = B_{j-1}/B_j$ for $1 \leq j < \nu$, then since $B_{j+1} = B_{j-1} + \beta_{j+1}B_j$ for $1 \leq j \leq \nu$, we get $B_{j+1}/B_j = B_{j-1}/B_j + \beta_{j+1}$ or $x_{j+1} = 1/(x_j + \beta_{j+1}) = \theta_{\beta_{j+1}}(x_j)$ for $1 \leq j \leq \nu$. Since $x_1 = 1/\beta_1 \in [\mathbf{a}_1, \mathbf{b}_1] = [1/(\Gamma + 1/\gamma), 1/\gamma]$, it follows from

Lemma 7.2 that $x_{j+1} \in [\mathbf{a}_{j+1}, \mathbf{b}_{j+1}]$ for $1 \leq j < \nu$, so $1/x_{j+1} = B_{j+1}/B_j \in [\mathbf{b}_{j+1}^{-1}, \mathbf{a}_{j+1}^{-1}]$ and $\mathcal{B}_\nu/B_{\nu-1} \geq \mathbf{b}_\nu^{-1}$. It follows that for $\omega \in \Omega_\nu$ and $x \in [\mathbf{a}_k, \mathbf{b}_k]$, we can take

$$(7.8) \quad M_0(\nu) = 2/(\mathbf{a}_k + \mathbf{b}_\nu^{-1}).$$

By adding the exponent s , one easily derives that for all $\omega \in \Omega_\nu$, $\nu \geq 1$, and $0 \leq k \leq \infty$,

$$(7.9) \quad g_\omega(\cdot)^s = |\theta'_\omega(\cdot)|^s \in K(2s/(\mathbf{a}_k + \mathbf{b}_\nu^{-1}); [\mathbf{a}_k, \mathbf{b}_k]).$$

Note that we could replace \mathbf{b}_ν^{-1} by $\mathbf{a}_{\nu-1} + \gamma$.

We summarize the above discussion in the following lemma.

Lemma 7.7. *Assume (H5) and let \mathbf{a}_k and \mathbf{b}_k , $0 \leq k \leq \infty$, be as described in Lemma 7.6. If $\omega \in \Omega_\nu$, $\nu \geq 1$, the map $x \mapsto \ln(|\theta'_\omega(x)|^s)$, $x \in [\mathbf{a}_k, \mathbf{b}_k]$ is Lipschitz with Lipschitz constant $\leq 2s/(\mathbf{a}_k + \mathbf{b}_\nu^{-1})$, so (7.9) is satisfied.*

It remains to estimate, for $0 \leq k \leq \infty$, $\max_{x \in [\mathbf{a}_k, \mathbf{b}_k]} |(d^j v_s/dx^j)(x)|/v_s(x)$, where v_s denotes the unique (to within normalization) strictly positive eigenfunction of L_s . The basic idea is to exploit (2.2), as was done in Section 6 of [14], but our results will refine those in [14].

Our previous calculations show that for $x \in [\mathbf{a}_k, \mathbf{b}_k]$, $0 \leq k \leq \infty$, we have

$$g_\omega(x)^s = |\theta'_\omega(x)|^s = \frac{1}{B_{\nu-1}^{2s}(x + B_\nu/B_{\nu-1})^{2s}}.$$

It follows that for $j \geq 1$, and letting D denote d/dx , we have

$$(7.10) \quad \frac{(-1)^j (D^j[(g_\omega)^s])(x)}{g_\omega(x)} = \frac{(2s)(2s+1) \cdots (2s+j-1)}{(x + B_\nu/B_{\nu-1})^j}.$$

The same argument used in Lemma 7.7 shows that

$$(7.11) \quad \mathbf{b}_\nu^{-1} \leq B_\nu/B_{\nu-1} \leq \mathbf{a}_\nu^{-1},$$

so if $x \in [\mathbf{a}_k, \mathbf{b}_k]$, we derive from (7.10) and (7.11) that

$$(7.12) \quad \frac{(2s)(2s+1) \cdots (2s+j-1)}{[\mathbf{b}_k + \mathbf{a}_\nu^{-1}]^j} \leq \frac{(-1)^j (D^j[(g_\omega)^s])(x)}{g_\omega(x)^s} \leq \frac{(2s)(2s+1) \cdots (2s+j-1)}{(\mathbf{a}_k + \mathbf{b}_\nu^{-1})^j}.$$

It follows from (7.12) that for $x \in [\mathbf{a}_k, \mathbf{b}_k]$,

$$(7.13) \quad \frac{(2s)(2s+1) \cdots (2s+j-1)}{[\mathbf{b}_k + \mathbf{a}_\nu^{-1}]^j} \leq \frac{(-1)^j \sum_{\omega \in \Omega_\nu} (D^j[(g_\omega)^s])(x)}{\sum_{\omega \in \Omega_\nu} g_\omega(x)^s} \leq \frac{(2s)(2s+1) \cdots (2s+j-1)}{(\mathbf{a}_k + \mathbf{b}_\nu^{-1})^j}.$$

Taking limits as $\nu \rightarrow \infty$ in (7.13) and using (2.2), we find that for $x \in [\mathbf{a}_k, \mathbf{b}_k]$,

$$(7.14) \quad \frac{(2s)(2s+1) \cdots (2s+j-1)}{[\mathbf{b}_k + \mathbf{a}_\infty^{-1}]^j} \leq \frac{(-1)^j (d^j v_s/dx^j)(x)}{v_s(x)}$$

$$\leq \frac{(2s)(2s+1) \cdots (2s+j-1)}{(\mathfrak{a}_k + \mathfrak{b}_\infty^{-1})^j}.$$

Notice that we can replace \mathfrak{a}_∞^{-1} by $\mathfrak{b}_\infty + \Gamma$ and \mathfrak{b}_∞^{-1} by $\mathfrak{a}_\infty + \gamma$ in (7.14).

As one can easily see, the lower bound in (7.14) increases as k increases and the upper bound decreases as k increases, so the optimal bounds are obtained when $k = \infty$ and apply to the interval $[\mathfrak{a}_\infty, \mathfrak{b}_\infty]$.

We summarize the above results in the following lemma.

Lemma 7.8. *Let v_s denote the unique (to with normalization) strictly positive eigenfunction of L_s . Assume (H4) and let \mathfrak{a}_k and \mathfrak{b}_k , $k \geq 0$ be as in Lemma 7.4. Then v_s satisfies (7.14).*

Remark 7.3. *Since, in Lemma 7.8, we have specified the coefficient g_β and the maps θ_β for $\beta \in \mathcal{B}$, Lemma 7.8 gives us a simple formula for the constant $E(s, p) = E$ in (6.1).*

$$\max_{x \in [\mathfrak{a}_k, \mathfrak{b}_k]} \frac{|(d^p v_s / dx^p)(x)|}{v_s(x)} \leq \frac{(2s)(2s+1) \cdots (2s+p-1)}{(\mathfrak{a}_k + \mathfrak{b}_\infty^{-1})^p} := E,$$

where p and k are positive integers and $1 \leq k \leq \infty$. Here we have allowed the interval to vary with k , but we may eventually restrict to $k = \infty$.

It remains to find a constant χ (compare (6.2)) such that for all $x_1, x_2 \in [\mathfrak{a}_k, \mathfrak{b}_k]$,

$$v_s(x_1) \leq \exp(2s|x_1 - x_2|/\chi) v_s(x_2).$$

It follows from (7.14) that if $\mathfrak{a}_k \leq x_1 \leq x_2 \leq \mathfrak{b}_k$, then

$$-\int_{x_1}^{x_2} \frac{v'_s(x)}{v_s(x)} dx = \ln(v_s(x_1)) - \ln(v_s(x_2)) \leq \frac{2s}{\mathfrak{a}_k + \mathfrak{b}_\infty^{-1}} |x_2 - x_1|,$$

which implies that

$$(7.15) \quad v_s(x_1) \leq \exp(2s|x_1 - x_2|/[\mathfrak{a}_k + \mathfrak{b}_\infty^{-1}]) v_s(x_2).$$

If $x_2 \leq x_1$, we know that $v_s(x_2) \geq v_s(x_1)$, so (7.15) is satisfied for all $x_1, x_2 \in [\mathfrak{a}_k, \mathfrak{b}_k]$. In particular, (7.15) is satisfied if the roles of x_1 and x_2 are reversed, which implies that $x \mapsto \ln(v_s(x))$ is Lipschitz on $[\mathfrak{a}_k, \mathfrak{b}_k]$ with Lipschitz constant $2s/(\mathfrak{a}_k + \mathfrak{b}_\infty^{-1})$. Summarizing, we have

Lemma 7.9. *If $\chi = \mathfrak{a}_k + \mathfrak{b}_\infty^{-1}$, (6.2) is satisfied on $[\mathfrak{a}_k, \mathfrak{b}_k]$.*

8. COMPUTATION OF $R(L_s)$

In this section we shall describe how to use the results of Sections 5-7 to obtain rigorous, high order estimates for $R(L_s) = \lambda_s$. As a subcase, we shall obtain rigorous estimates for the Hausdorff dimension of certain fractal objects described by iterated function systems.

For simplicity, we shall restrict attention to the class of maps $\theta_\beta : [0, 1] \rightarrow [0, 1]$, where $\theta_\beta(x) = 1/(x + \beta)$ for $\beta \in \mathcal{B}$ and \mathcal{B} as in (H5) of Section 7. For $s \geq 0$, we define $L_s : C[0, 1] \rightarrow C[0, 1]$ by

$$(L_s f)(x) = \sum_{\beta \in \mathcal{B}} |\theta'_\beta(x)|^s f(\theta_\beta(x)).$$

Recall (see Lemmas 7.3 and 7.4) that we define $\mathbf{a}_0 = 0$, $\mathbf{b}_0 = 1/\gamma$, $\mathbf{a}_{k+1} = \theta_\Gamma(\mathbf{b}_k)$, $\mathbf{b}_{k+1} = \theta_\gamma(\mathbf{a}_k)$, $\mathbf{a}_\infty = \lim_{k \rightarrow \infty} \mathbf{a}_k$, and $\mathbf{b}_\infty = \lim_{k \rightarrow \infty} \mathbf{b}_k$. Since $\theta_\beta([a_k, b_k]) \subset [a_k, b_k]$ for $0 \leq k \leq \infty$ and for all $\beta \in \mathcal{B}$, and since L_s has a strictly positive eigenvector on $[0, 1/\gamma]$, L_s induces a bounded linear operator $L_{s, [\mathbf{a}_k, \mathbf{b}_k]} : C([\mathbf{a}_k, \mathbf{b}_k]) \rightarrow C([\mathbf{a}_k, \mathbf{b}_k])$ and $R(L_s) = R(L_{s, [\mathbf{a}_k, \mathbf{b}_k]})$. Various constants are optimized by working on $[\mathbf{a}_\infty, \mathbf{b}_\infty]$, so we shall abuse notation and also use L_s to denote L_s as an operator on $C([\mathbf{a}_\infty, \mathbf{b}_\infty])$ (or, sometimes, $C([\mathbf{a}_k, \mathbf{b}_k])$). For a given positive integer r , we assume that (H3) is satisfied, but with $S := [\mathbf{a}_\infty, \mathbf{b}_\infty]$.

Thus $[a_i, b_i] \subset S$, $1 \leq i \leq I$, denote pairwise disjoint intervals that satisfy the conditions of (H3). Given positive integers N_i , $1 \leq i \leq I$, we write $h_i = (b_i - a_i)/N_i$. As in Section 3 (see (3.5)), we define mesh points $c_{j,k}^i \in [a_i, b_i]$, $1 \leq i \leq I$, $1 \leq j \leq N_i$, $0 \leq k \leq r$ and $T := \{c_{j,k}^i\}$ for i, j, k in the ranges given above. As in Section 3, if v_s is the positive eigenvector of L_s on S , $\mathcal{V}_s : \hat{S} := \cup_{i=1}^I [a_i, b_i] \rightarrow \mathbb{R}$ is the polynomial interpolant of v_s of degree $\leq r$ on $[t_{j-1}^i, t_j^i]$ for $1 \leq i \leq I$, $1 \leq j \leq N_i$, so $\mathcal{V}_s(x) = v_s(x)$ for all $x \in T$.

Our general approach will be as follows: Given $s > 0$, we must find r, ν, M and h such that the conditions of Theorem 5.6 are satisfied. First, we choose a positive integer $r \geq 2$, where r is the piecewise polynomial degree in (3.6). Once r has been chosen, we select a positive integer ν such that (compare Remark 5.3)

$$(8.1) \quad c(\nu)[2\eta(r)r^2\psi(r)] := \kappa_1 < 1.$$

Here $c(\nu)$ is as in (5.3); and for our case an exact formula for $c(\nu)$ is provided by (7.4); where we shall take $\mathbf{a}_k = \mathbf{a}_\infty$ in (7.4). Also, $\psi(r)$ is as in Lemma 5.2 and $\eta(r)$ as in (5.2). As a practical matter, we demand that κ_1 not be too close to 1, say $\kappa_1 \leq 4/5$. Note that for fixed r , this means that ν must be sufficiently large and hence $c(\nu)$ sufficiently small, so that (8.1) is satisfied. We next choose κ_2 with $\kappa_1 < \kappa_2 < 1$ and κ_2 not too close to 1. A simple choice is $\kappa_2 = (1 + \kappa_1)/2$. We define (see Theorem 5.6), $M' = \kappa_2 M$. If we write $u := M\eta(r)h$, the conditions of Theorem 5.6 take the form

$$(8.2) \quad \psi(r)u \exp(u) < 1$$

$$(8.3) \quad \frac{\kappa_1 \exp(u)}{1 - \psi(r)u \exp(u)} < \kappa_2 - \frac{sM_0(\nu)}{M}.$$

Here $M_0(\nu)$ is as in (5.3); and in our case Lemma 7.7 insures that $M_0(\nu) \leq 2/(\mathbf{a}_\infty + \mathbf{a}_{\nu-1} + \gamma)$.

Since $\exp(u)/(1 - \psi(r)u \exp(u)) > 1$, (8.3) implies that

$$\kappa_1 < \kappa_2 - sM_0(\nu)/M.$$

We choose $M > 0$ such that

$$(8.4) \quad M = \frac{4s}{\mathfrak{a}_\infty + \mathfrak{a}_{\nu-1} + \gamma} \frac{1}{\kappa_2 - \kappa_1} \geq 2sM_0(\nu)/(\kappa_2 - \kappa_1),$$

which implies that

$$\kappa_2 - sM_0(\nu)/M \geq \kappa_2 - (\kappa_2 - \kappa_1)/2 > \kappa_1.$$

Also note that since $\mathfrak{a}_\infty + \mathfrak{a}_{\nu-1} + \gamma < \chi$, we have that

$$(8.5) \quad M \geq \frac{4s}{\chi} \frac{1}{\kappa_2 - \kappa_1}.$$

Given an M that satisfies (8.4), we can choose $h = \max_i h_i$ sufficiently small, i.e., $h \leq \mathfrak{h}_0$, that (8.2) and (8.3) are satisfied. Recall, however, that we also have to insure that the constant M , defined by (8.4) also satisfies $M \geq M_2$, where M_2 is as in Lemma 6.3 and M_1 is given by (6.10). As we shall see, this may require a further restriction on the size of h .

The constants M_1 and M_2 are defined in terms of $G_{r+1} := G$ (compare (6.7)), $\chi := 2\mathfrak{a}_\infty + \gamma$, s , and H (compare Lemma 6.3), and it is desirable to choose H to be small. The constant E in G_{r+1} is given by Remark 7.3 with $p = r + 1$. By using (6.6) in Section 7 and the estimates for E and χ in Lemmas 7.8 and 7.9, we find that we can write

$$G = E \exp\left(\frac{2sh}{\mathfrak{a}_\infty + \mathfrak{b}_\infty^{-1}}\right) \frac{1}{(r+1)!} \left[\frac{1}{2 \cos(\pi/(2r+2))} \right]^{r+1} \frac{1}{2^r},$$

where

$$(8.6) \quad E := \frac{(2s)(2s+1) \cdots (2s+r)}{(\mathfrak{a}_\infty + \mathfrak{b}_\infty^{-1})^{r+1}} = \frac{(2s)(2s+1) \cdots (2s+r)}{(2\mathfrak{a}_\infty + \gamma)^{r+1}},$$

and we have used the fact that $\mathfrak{b}_\infty = 1/(\mathfrak{a}_\infty + \gamma)$. Note that in the application of (7.14) for E , one must take $j = r + 1$. A calculation gives

$$(8.7) \quad G := G_{r+1} = 2 \exp\left(\frac{2sh}{2\mathfrak{a}_\infty + \gamma}\right) \left[\frac{(2s)(2s+1) \cdots (2s+r)}{(2)(4)(6) \cdots (2r+2)} \right] \\ \cdot \left[\frac{1}{2\mathfrak{a}_\infty + \gamma} \right]^{r+1} \left[\frac{1}{2 \cos(\pi/(2r+2))} \right]^{r+1}.$$

Finally, we define

$$(8.8) \quad D := D_{r+1} = \left[\frac{1}{\sin(\pi/(2r+2))} \right]^2 G_{r+1}$$

and

$$(8.9) \quad H_{r+1} = \mu D_{r+1} \frac{\chi}{2s} = \mu \left[\frac{1}{\sin(\pi/(2r+2))} \right]^2 G_{r+1} \frac{(2\mathfrak{a}_\infty + \gamma)}{2s}.$$

To estimate M_1 and M_2 , we shall need estimates on these quantities.

Lemma 8.1. *Assume that $r \geq 2$, $0 < s < 2$, and $2\mathfrak{a}_\infty + \gamma \geq 1$. Then G_r is a decreasing function of r .*

Proof. For $r \geq 2$ and $0 < s \leq 2$, since $2j + 2 \geq 2s + j$ for $j \geq 2$, it follows that

$$\prod_{j=0}^r \frac{2s+j}{2j+2} \leq \left(\frac{2s}{2}\right) \left(\frac{2s+1}{4}\right) = s \frac{2s+1}{4}.$$

By using the Taylor series for $\cos(\theta)$, we see that

$$\frac{1}{2 \cos(\pi/(2r+2))} \leq \frac{1}{2 - [\pi/(2r+2)]^2}.$$

Using these estimates in (8.7) gives

$$G_{r+1} \leq 2 \exp\left(\frac{2sh}{2a_\infty + \gamma}\right) \left[s \frac{2s+1}{4}\right] \left[\frac{1}{2a_\infty + \gamma}\right]^{r+1} \left[\frac{1}{2 - [\pi/(2r+2)]^2}\right]^{r+1},$$

which implies that $\lim_{r \rightarrow \infty} G_{r+1} = 0$. Furthermore, for $r \geq 2$ and $2a_\infty + \gamma \geq 1$, another calculation shows that

$$\frac{G_{r+1}}{G_r} = \left[\frac{2s+r}{2r+2}\right] \left[\frac{1}{2a_\infty + \gamma}\right] \left[\frac{1}{2 \cos(\pi/[2r+2])}\right] \left[\frac{\cos(\pi/(2r))}{\cos(\pi/(2r+2))}\right]^r < 1,$$

so G_r is a decreasing function for integer $r \geq 2$. \square

If we set

$$u_{r+1} = \frac{1}{\chi} \left[\frac{1}{2 \cos(\pi/(2r+2))}\right],$$

a calculation gives

$$u_3 = \frac{1}{\chi\sqrt{3}}, \quad u_4 = \frac{1}{\chi\sqrt{2+\sqrt{2}}},$$

which gives

$$(8.10) \quad \begin{aligned} G_3 &= 2s \left[\frac{1}{\chi\sqrt{3}}\right]^3 \left[\frac{2s+1}{4}\right] \left[\frac{s+1}{3}\right] \exp(2sh/\chi), \\ G_4 &= 2s \left[\frac{1}{\chi\sqrt{2+\sqrt{2}}}\right]^2 \left[\frac{1}{\chi}\right]^4 \left[\frac{2s+1}{4}\right] \left[\frac{s+1}{3}\right] \left[\frac{2s+3}{8}\right] \exp(2sh/\chi), \end{aligned}$$

$D_3 = 4G_3$, and $D_4 = [4/(2 - \sqrt{2})]G_4$. Then

Lemma 8.2. *If $0 \leq s \leq 3/2$, $r \geq 2$, and $\chi := 2a_\infty + \gamma \geq 1$,*

$$\frac{G_{r+2}}{G_{r+1}} \leq \left[\frac{3}{4\sqrt{2+\sqrt{2}}}\right] \frac{1}{\chi}, \quad \text{so} \quad G_{r+2} \leq \left[\frac{3}{4\sqrt{2+\sqrt{2}}}\left(\frac{1}{\chi}\right)\right]^{r-1} G_3.$$

Proof. A calculation gives

$$\frac{G_{r+2}}{G_{r+1}} = \left[\frac{2s+r+1}{2r+4}\right] u_{r+2} \left[\frac{u_{r+2}}{u_{r+1}}\right]^{r+1}.$$

Since $0 < u_{r+2} < u_{r+1}$, it follows that

$$\frac{G_{r+2}}{G_{r+1}} \leq \left[\frac{2s+r+1}{2r+4}\right] \frac{1}{\chi} \left[\frac{1}{2 \cos(\pi/(2r+4))}\right],$$

and if $r \geq 2$ and $s \leq 3/2$,

$$\frac{G_{r+2}}{G_{r+1}} \leq \left[\frac{3}{4\chi}\right] \left[\frac{1}{2 \cos(\pi/8)}\right] \leq \left[\frac{3}{4[\sqrt{2+\sqrt{2}}]}\right] \frac{1}{\chi}$$

and so

$$G_{r+2} \leq \left[\frac{3}{4[\sqrt{2+\sqrt{2}}]} \left(\frac{1}{\chi} \right) \right]^{r-1} G_3.$$

Furthermore, since we assume that $\chi \geq 1$,

$$(8.11) \quad G_3 = \left[\frac{2s}{\chi} \right] \left[\frac{1}{\chi^2 3\sqrt{3}} \right] \left[\frac{2s+1}{4} \right] \left[\frac{s+1}{3} \right] \exp(2sh/\chi) \\ \leq \left[\frac{2s}{\chi^3} \right] \left[\frac{5}{18\sqrt{3}} \right] \exp(3h) \leq \sqrt{3}/2,$$

for $h \leq \ln(9/5)/3 \leq 0.2$. \square

Lemma 8.3. Assume $0 \leq s \leq 3/2$, $r \geq 2$, $\chi \geq 1$, and D_{r+1} is as in (8.8). Then

$$D_{r+2} \leq \left[\frac{3}{4\chi} \right]^{r-1} D_3.$$

Proof. A calculation gives

$$\frac{D_{r+2}}{D_{r+1}} = \left[\frac{u_{r+2}}{u_{r+1}} \right]^{r+1} u_{r+2} \left[\frac{\sin(\pi/[2r+2])}{\sin(\pi/[2r+4])} \right]^2 \left[\frac{2s+r+1}{2r+4} \right] \\ \leq \frac{1}{\chi} \left[\frac{1}{2 \cos(\pi/(2r+4))} \right] \left[\frac{\sin(\pi/[2r+2])}{\sin(\pi/[2r+4])} \right]^2 \left[\frac{r+4}{2r+4} \right].$$

Using Taylor series expansions for $\sin(u)$, $u \geq 0$, we have $\sin(u) \leq u$ and $\sin(u) \geq u - u^3/6$. Hence,

$$\frac{\sin(\pi/[2r+2])}{\sin(\pi/[2r+4])} \leq \left[\frac{2r+4}{2r+2} \right] \left[1 - \frac{1}{6} \left(\frac{\pi}{2r+4} \right)^2 \right]^{-1}.$$

Noting that for $r \geq 2$, the expression on the right hand side of the above is a decreasing function of r , as are the other two functions of r in the bound for D_{r+2}/D_{r+1} , we see that an upper bound for D_{r+2}/D_{r+1} is obtained by setting $r = 2$ in each of the expressions above. This gives

$$\frac{D_{r+2}}{D_{r+1}} \leq \left[\frac{1}{2\chi \cos(\pi/8)} \right] \left[\frac{4}{3} \right]^2 \left[1 - \frac{1}{6} \left(\frac{\pi}{8} \right)^2 \right]^{-2} \left[\frac{3}{4} \right] \leq \frac{3}{4\chi},$$

and so

$$D_{r+2} \leq \left[\frac{3}{4\chi} \right] D_{r+1} \leq \left[\frac{3}{4\chi} \right]^{r-1} D_3. \quad \square$$

The following bound on H_r is a direct consequence of the above estimates.

Lemma 8.4. Assume $0 \leq s \leq 3/2$, $r \geq 2$. Then

$$H_{r+2} \leq \mu \left[\frac{\chi}{2s} \right] \left[\frac{3}{4\chi} \right]^{r-1} 4G_3.$$

Lemma 8.5. Suppose $0 \leq s \leq 3/2$, $r \geq 2$, M_1 is as in (6.10), M_2 is as in Lemma 6.3, and $H_{r+1} = \mu D_{r+1}(\chi/2s)$. Then

$$M_1 = \frac{\mu D_{r+1} h^r}{1 - G_{r+1}^2 h^{2r+2}} + \frac{2s}{\chi},$$

and

$$\begin{aligned}
(8.12) \quad M_2 &= M_1 + \left[\frac{2s}{\chi} \right] \frac{1}{1 - [(G_{r+1}/H_{r+1})h]^2} \\
&= \frac{2s}{\chi} \left[1 + \frac{H_{r+1}h^r}{1 - G_{r+1}^2 h^{2r+2}} + \left(1 - h^2 \frac{2s}{\mu\chi} \sin^2(\pi/[2r+2]) \right)^{-1} \right].
\end{aligned}$$

Proof. Applying the definitions of M_1 , M_2 , D_{r+1} , G_{r+1} , and H_{r+1} , we get

$$\begin{aligned}
M_2 &= M_1 + \left[\frac{\mu G_{r+1}}{H_{r+1}} \right] \left[\frac{1}{1 - [(G_{r+1}/H_{r+1})h]^2} \right] \left[\frac{1}{[\sin(\pi/[2r+2])]^2} \right] \\
&= M_1 + \left[\frac{2s}{\chi} \right] \frac{1}{1 - [(G_{r+1}/H_{r+1})h]^2} \\
&= \frac{2s}{\chi} + \frac{\mu D_{r+1}h^r}{1 - G_{r+1}^2 h^{2r+2}} + \left[\frac{2s}{\chi} \right] \frac{1}{1 - [(G_{r+1}/H_{r+1})h]^2} \\
&= \frac{2s}{\chi} + \left[\frac{2s}{\chi} \right] \frac{H_{r+1}h^r}{1 - G_{r+1}^2 h^{2r+2}} + \left[\frac{2s}{\chi} \right] \frac{1}{1 - (2s/[\mu\chi])[\sin(\pi/[2r+2])h]^2} \\
&= \frac{2s}{\chi} \left[1 + \frac{H_{r+1}h^r}{1 - G_{r+1}^2 h^{2r+2}} + \left(1 - h^2 \frac{2s}{\mu\chi} \sin^2(\pi/[2r+2]) \right)^{-1} \right].
\end{aligned}$$

□

Remark 8.1. Note that D_{r+1} has the factor $2s/\chi$, so the identity (8.12) for M_2 does not blow up as $s \rightarrow 0$.

Remark 8.2. If we replace in G_3 and D_3 , the quantity $\exp(2sh/\chi)$ by $\exp(2s\mathfrak{h}_0/\chi)$, where \mathfrak{h}_0 is chosen so that (8.2) and (8.3) are satisfied for $0 < h \leq \mathfrak{h}_0$, then we easily obtain a bound for M_2 in the form

$$(8.13) \quad M_2 \leq \frac{2s}{\chi} \left[2 + H_{r+1}h^r(1 + \tilde{c}h^{2r+2}) + ch^2 \right].$$

where \tilde{c} and c are easily computable from (8.12). Note that if $0 \leq s \leq 3/2$, $\chi = 2\mathfrak{a}_\infty + \gamma \geq 1$, and $r \geq 2$, then $\chi \geq 1$ and $\sin^2(\pi/[2r+2]) \leq 1/4$. So, for $h \leq 1/\sqrt{3}$,

$$\left(1 - h^2 \frac{2s}{\mu\chi} \sin^2(\pi/[2r+2]) \right)^{-1} \leq (1 - 3h^2/4)^{-1} \leq 1 + h^2,$$

i.e., we can take $c = 1$. Similarly, for $h \leq 0.2$, $G_{r+1}^2 \leq 3/4$, so we can also take $\tilde{c} = 1$.

Recall that we have to insure that $M > M_2$. From (8.5) we have that

$$M \geq \left[\frac{4s}{\chi} \right] \frac{1}{\kappa_2 - \kappa_1}.$$

Comparing this expression to the bound for M_2 given in (8.13) and noting that $\kappa_2 - \kappa_1 < 1$, M will be $> M_2$ if we choose $h \leq \mathfrak{h}_1$ sufficiently small so that it also satisfies

$$(8.14) \quad 2 + H_{r+1}h^r(1 + h^{2r+2}) + h^2 \leq 2/(\kappa_2 - \kappa_1).$$

We can now state versions of Theorem 6.4 and Theorem 6.5 in the context of this section.

Theorem 8.6. *Assume that $r \geq 2$, $\chi \geq 1$, and $0 \leq s \leq 3/2$, and let ν, κ_1 , and κ_2 be as described at the beginning of this section. Let M_2 and H_{r+1} be as described in Lemma 8.5 and select M such that (8.4) is satisfied. Finally, assume that $h = \max_{i \in I} h_i \leq \min(\mathfrak{h}_0, \mathfrak{h}_1, 0.2)$. Then, with $u = M\eta(r)h$, (8.2) and (8.3) are satisfied and $M > M_2$. Furthermore, (compare Theorem 6.4) $\mathbf{L}_{s,\nu}(K(M;T)) \subset K(M';T)$, where $M' = \kappa_2 M < M$. In addition, we have that for $H = H_{r+1}$,*

$$\lambda_s^\nu(1 - Hh^r) \leq R(\mathbf{L}_{s,\nu}) \leq \lambda_s^\nu(1 + Hh^r).$$

Proof. The fact that $M_2 > M$ follows directly from the computations above. Our selection of r, ν, h, M and $M' = \kappa_2 M$ shows that the inequality in Theorem 5.2 is satisfied, so $\mathbf{L}_{s,\nu}(K(M;T)) \subset K(M';T)$. The inequality for $R(\mathbf{L}_{s,\nu})$ in Theorem 8.6 follows directly from Theorem 6.4. \square

Theorem 8.7. *Under the hypotheses of Theorem 8.6, we have*

$$[(1 + Hh^r)^{-1}R(\mathbf{L}_{s,\nu})]^{1/\nu} \leq \lambda_s \leq [(1 - Hh^r)^{-1}R(\mathbf{L}_{s,\nu})]^{1/\nu},$$

where the entries of the matrices $[1 + Hh^r]^{-1}\mathbf{L}_{s,\nu}$ and $[1 - Hh^r]^{-1}\mathbf{L}_{s,\nu}$ differ by $O(h^r)$.

Using the inequalities of Theorem 8.6, we can obtain rigorous upper and lower bounds on the Hausdorff dimension s_* of the invariant set associated with the transfer operator L_s as follows. Let s_l and s_u denote values of s satisfying

$$(1 - Hh^r)^{-1}R(\mathbf{L}_{s_u,\nu}) < 1, \quad (1 + Hh^r)^{-1}R(\mathbf{L}_{s_l,\nu}) > 1.$$

It follows immediately from Theorem 8.6 that $\lambda_{s_u}^\nu < 1$ and $\lambda_{s_l}^\nu > 1$. Since the spectral radius λ_s of L_s is a decreasing function of s , there will be a value s_* satisfying $s_l < s_* < s_u$ for which $\lambda_{s_*}^\nu = 1$, or equivalently $\lambda_{s_*} = 1$. The value s_* gives the Hausdorff dimension s_* of the invariant set associated with the transfer operator L_s .

9. NUMERICAL COMPUTATIONS

In this final section, we present results of computations of the Hausdorff dimension s for various choices of sets of continued fractions, maximum mesh size h , piecewise polynomial degree r , and number of iterations ν of the map, where $\nu = 1$ corresponds to the original map. These computations include choices of the above parameters (especially the number of iterations ν), for which the hypotheses of our theorems are satisfied, but also computations which obtain the same results when the mappings are not iterated (denoted by $\nu = 1^*$ in Table 9.1 below). We note the obvious fact that as the number of iterations increase, the complexity of the operator increases, so the time to compute the maximum eigenvalue of the resulting matrix operator increases as well. To keep the time involved in the various computations to be reasonable, we have elected to compute to fewer digits those sets with a large value of ν , especially if the set E contains many digits. We have also done additional computations, not reported in Table 9.1, which indicate that the method is robust with respect to the choices of h , r , and ν .

TABLE 9.1. Computation of Hausdorff dimension s for various choices of sets of continued fractions, maximum mesh size h , piecewise polynomial degree r , and number of iterations ν .

Set E	r	h	ν	
				$s =$
E[1,2]	14	0.0002	7	
				$s = 0.531\ 280\ 506\ 277\ 205\ 141\ 624\ 468\ 647\ 368\ 471\ 785\ 493\ 059\ 109\ 018\ 398$
E[1,3]	8	5.0e-05	6	
				$s = 0.454\ 489\ 077\ 661\ 828\ 743\ 845\ 777\ 611\ 651$
E[1,4]	8	5.0e-05	6	
				$s = 0.411\ 182\ 724\ 774\ 791\ 776\ 844\ 805\ 904\ 696$
E[2,3]	8	5.0e-05	3	
				$s = 0.337\ 436\ 780\ 806\ 063\ 636\ 304\ 494\ 910\ 387$
E[2,4]	8	5.0e-05	3	
				$s = 0.306\ 312\ 768\ 052\ 784\ 030\ 277\ 908\ 307\ 445$
E[3,4]	8	5.0e-05	3	
				$s = 0.263\ 737\ 482\ 897\ 426\ 558\ 759\ 863\ 384\ 275$
E[3,7]	18	.01	3	
				$s = 0.224\ 923\ 947\ 191\ 778\ 989\ 184\ 480\ 593\ 490$
E[10,11]	20	.002	2	
				$s = 0.146\ 921\ 235\ 390\ 783\ 463\ 311\ 108\ 628\ 515\ 904\ 073\ 067\ 083\ 129\ 676\ 755$
E[100, 10,000]	20	.002	1	
				$s = 0.052\ 246\ 592\ 638\ 658\ 878\ 652\ 588\ 416\ 300\ 508\ 181\ 012\ 676\ 284\ 431\ 681$
E[1,2,3]	5	.0001	5	
				$s = 0.705\ 660\ 908\ 028\ 738$
E[1,3,4]	5	.0001	5	
				$s = 0.604\ 242\ 257\ 756\ 515$
E[1,3,5]	8	.001	6	
				$s = 0.581\ 366\ 821\ 182\ 975$
E[1,4,7]	6	.001	6	
				$s = 0.517\ 883\ 757\ 006\ 911$
E[2,3,4]	16	.005	4	
				$s = 0.480\ 696\ 222\ 317\ 573\ 041\ 322\ 515\ 564\ 711$
E[1,2,3,4]	8	.005	6	
				$s = 0.788\ 945\ 557\ 483\ 153$
E[2,3,4,5]	16	.005	4	
				$s = 0.559\ 636\ 450\ 164\ 776\ 713\ 312\ 144\ 913\ 530$
E[1,2,3,4,5]	5	.0005	5	
				$s = 0.836\ 829\ 443\ 681\ 209$
E[2,4,6,8,10]	7	.005	3	
				$s = 0.517\ 357\ 030\ 937\ 017$
E[1,...,10]	10	.01	1*	
				$s = 0.925\ 737\ 591\ 146\ 765$
E[1,...,34]	10	.01	1*	
				$s = 0.980\ 419\ 625\ 226\ 980$
E[1,3,5,...,33]	10	.01	1*	
				$s = 0.770\ 516\ 008\ 717\ 163$
E[2,4,6,...,34]	10	.01	1*	
				$s = 0.633\ 471\ 970\ 241\ 089$

In addition to the results presented in Table 9.1, we have also used our method to compute the Hausdorff dimension of the set $E[1, 2]$ with degree $r = 36$, $h = .001$, and $\nu = 1$ using multiple precision with 108 digits. Although this choice does not satisfy the hypotheses of our theorem, the result agrees to 100 decimal places with the result in [26]. While we do not have a proof that our method also works in the non-iterated situation, we do not have any examples where it fails. We conjecture that the limitation is the method of proof and not the underlying method.

We next discuss how to control the size of some of the constants that appear in the error estimates. We begin with the constant $\mu = \max_i h_i / \min_i h_i$. Recall that to satisfy hypothesis (H3), for a given positive integer ν , we need to determine pairwise disjoint, nonempty compact intervals $[a_i, b_i] \subset S$, $1 \leq i \leq I$, such that for every $\omega \in \Omega_\nu$, there exists $i = i(\omega)$, $1 \leq i \leq I$, such that $\theta_\omega(S) \subset [a_i, b_i]$. By the results in the previous section, we can take $S = [\mathbf{a}_\infty, \mathbf{b}_\infty]$ and we then order the $\omega \in \Omega_\nu$, such that the sets $\theta_\omega(S)$ are ordered with $a_i < a_{i+1}$. Although we could use the domain consisting of the union of the sets $\theta_\omega(S)$, this can lead to very small subinterval sizes. Instead, we determine a new domain by iterating only ν' times, while still using the mappings obtained by ν iterations to calculate the mapping L . The constant ν' is determined so that the length of the smallest interval will be $\geq h_{\max}/\mu$.

Although we have not done the computations using interval arithmetic, we have only included the number of digits in each computation that we expect to be correct, which is always less than the number of digits provided by *Matlab* for the precision we have specified. For computations that use more digits than provided by standard *Matlab* computations, we have used the *Advanpix* multiprecision toolbox.

Because the theory developed in the previous sections involves the computation and estimates for many constants and parameters, we next provide, for the benefit of the reader, details for the specific example of $E[1, 4, 7]$, corresponding to the computation for $r = 6$, $h = 0.001$, $\nu = 6$, and $s = 0.518$, shown in Table 9.1. The computational domain is determined by only iterating $\nu' = 2$ times and the total number of subintervals used is 191. With these choices, $\gamma = 1$ and $\Gamma = 7$. Then by (7.1),

$$\mathbf{a}_\infty = -\frac{\gamma}{2} + \sqrt{(\gamma/2)^2 + (\gamma/\Gamma)} \approx 0.127 \quad \text{and} \quad \mathbf{b}_\infty = (\Gamma/\gamma)\mathbf{a}_\infty \approx 0.8875.$$

Working on the interval $[\mathbf{a}_\infty, \mathbf{b}_\infty]$, we have $\chi = \mathbf{a}_\infty + \mathbf{b}_\infty^{-1} = 2\mathbf{a}_\infty + \gamma \approx 1.25$. From (7.5)–(7.7), we have for $\nu = 6$, that

$$c(\nu) = [\tilde{B}_{\nu-1}\mathbf{a}_\infty + \tilde{B}_\nu]^{-2} = 0.0051$$

From (5.2), we have that $\eta(r) = 1/2$ and from Lemma 5.2 that

$$\psi(r) = (2/\pi) \ln(r+1) + 3/4 \approx 2.0$$

From (7.8), since we are working on the interval $[\mathbf{a}_\infty, \mathbf{b}_\infty]$, we take

$$M_0(\nu) = 2/(\mathbf{a}_\infty + \mathbf{b}_\nu^{-1}) = 2/(\mathbf{a}_\infty + \mathbf{a}_{\nu-1} + \gamma) = 1.5954$$

Setting

$$\kappa_1 = c(\nu)2\eta(r)r^2\psi(r) = 0.364 < 1,$$

we choose $\kappa_2 = (1 + \kappa_1)/2 = 0.6823$, so that $\kappa_1 < \kappa_2 < 1$. Next, we choose

$$M = \frac{4s}{\mathfrak{a}_\infty + \mathfrak{a}_{\nu-1} + \gamma} \frac{1}{\kappa_2 - \kappa_1} = 5.2022$$

and $M' = \kappa_2 M$. One of the conditions on the mesh size $h = \max_{i \in I} h_i$ is that it be sufficiently small so that (8.2) and (8.3) are satisfied. For our example, setting $u = M\eta(r)h$, we have

$$\begin{aligned} \psi(r)u \exp(u) &= .0052 < 1, \\ \frac{\kappa_1 \exp(u)}{1 - \psi(r)u \exp(u)} &= 0.3674 < 0.5234 = \kappa_2 - \frac{sM_0(\nu)}{M}. \end{aligned}$$

The second condition, coming from (8.14) is that

$$(9.1) \quad 2 + H_{r+1}(h^r + h^{2r+2}) + h^2 \leq 2/(\kappa_2 - \kappa_1),$$

where, combining (8.8), (8.9), and Lemma 8.3, we have

$$H_{r+1} \leq \mu \frac{\chi}{2s} D_{r+1} \leq \mu \frac{\chi}{2s} \left[\frac{3}{4\chi} \right]^{r-2} D_3 \leq \mu \frac{\chi}{2s} \left[\frac{3}{4\chi} \right]^{r-2} 4G_3,$$

where

$$G_3 = 2s \left[\frac{1}{\chi\sqrt{3}} \right]^3 \left[\frac{2s+1}{4} \right] \left[\frac{s+1}{3} \right] \exp(2sh/\chi).$$

Combining these results and setting μ , the ratio of the maximum subinterval size to the minimum subinterval size, $= 3.76$, the value calculated by the computer code, we get

$$H_{r+1} \leq 0.0608.$$

Since $2/(\kappa_2 - \kappa_1) \geq 6$, it is clear that (9.1) is satisfied.

We thus have satisfied the conditions of Theorems 8.6 and 8.7, which guarantee the rigorous bounds we use to obtain rigorous upper and lower bounds on the Hausdorff dimension of the set $E[1, 4, 7]$.

REFERENCES

1. O. F. Bandtlow and J. Slipantschuk, *Lagrange approximation of transfer operators associated with homomorphic data*, available from [arXiv.org:2004.03534](https://arxiv.org/abs/2004.03534)
2. G. Birkhoff, *Extensions of Jentzsch's theorem*, Trans. Amer. Math. Soc. 85 (1957), 219-227. MR 0087058
3. Jean Bourgain and Alex Kontorovich, *On Zaremba's conjecture*, Ann. of Math. (2) **180** (2014), no. 1, 137-196. MR 3194813
4. Rufus Bowen, *Hausdorff dimension of quasicircles*, Inst. Hautes Études Sci. Publ. Math. (1979), no. 50, 11-25. MR 556580 (81g:57023)
5. L. Brutman, *On the Lebesgue function for polynomial interpolation*, SIAM J. Numer. Anal. 15 (1978), no. 4, 694-704. MR 0510554
6. R. T. Bumby, *Hausdorff dimension of Cantor sets*, J. Reine Angew. Math. 331 (1982), 192-206. MR 0647383
7. ———, *Hausdorff dimension of sets arising in number theory*, Number Theory (New York, 1983-84) (Lecture Notes in Mathematics, 1135). Springer, 1985, 1-8. MR 0803348
8. V. Chousionis, D. Leykekhman, and M. Urbanski, *On the dimension spectrum of infinite sub-systems of continued fractions*, Trans. Amer. Math. Soc., 373 (2020), 1009-1042. MR 4068257
9. T. W. Cusick, *Continuants with bounded digits*, Mathematika **24** (1977), no. 2, 166-172. MR 0472721 (57 #12413)

10. ———, *Continuants with bounded digits. II*, *Mathematika* **25** (1978), no. 1, 107–109. MR 0498413 (58 #16539)
11. S. P. Eveson and R. D. Nussbaum, *An elementary proof of the Birkhoff-Hopf theorem*, *Math. Proc. Cambridge Philos. Soc.* 117, Issue 1, (1995), 31–55. MR 1297895
12. ———, *Applications of the Birkhoff-Hopf theorem to the spectral theory of positive linear operators*, *Math. Proc. Cambridge Philos. Soc.* 117 (1995), no. 3, 491–512. MR 1317492
13. Kenneth Falconer, *Techniques in fractal geometry*, John Wiley & Sons, Ltd., Chichester, 1997. MR 1449135 (99f:28013)
14. R. S. Falk and R. D. Nussbaum, *C^m eigenfunctions of Perron-Frobenius operators and a new approach to numerical computation of Hausdorff dimension: applications in \mathbb{R}^1* , *Fractal Geom.* 5 (2018), no. 3, 279–337. MR 3827801
15. I. J. Good, *The fractional dimensional theory of continued fractions*, *Proc. Cambridge Philos. Soc.* **37** (1941), 199–228. MR 0004878 (3,75b)
16. R. Günttner, *Evaluation of Lebesgue constants*, *SIAM J. Numer. Anal.* 17 (1980), no. 4, 512–520. MR 0584726
17. Stefan-M. Heinemann and Mariusz Urbański, *Hausdorff dimension estimates for infinite conformal IFSs*, *Nonlinearity* **15** (2002), no. 3, 727–734. MR 1901102 (2003c:37029)
18. D. Hensley, *The Hausdorff dimensions of some continued fraction Cantor sets*, *J. Number Theory* 33 (1989), 182–198. MR 1034198
19. ———, *Continued fraction Cantor sets, Hausdorff dimension, and functional analysis*, *J. Number Theory* **40** (1992), no. 3, 336–358. MR 1154044 (93c:11058)
20. ———, *A polynomial time algorithm for the Hausdorff dimension of continued fraction Cantor sets*, *J. Number Theory* 58 (1996), 9–45. MR 1387719
21. Eberhard Hopf, *An inequality for positive linear integral operators*, *J. Math. Mech.* **12** (1963), 683–692. MR 0165325 (29 #2614)
22. John E. Hutchinson, *Fractals and self-similarity*, *Indiana Univ. Math. J.* **30** (1981), no. 5, 713–747. MR 625600 (82h:49026)
23. Oliver Jenkinson, *On the density of Hausdorff dimensions of bounded type continued fraction sets: the Texan conjecture*, *Stoch. Dyn.* **4** (2004), no. 1, 63–76. MR 2069367 (2005m:28021)
24. O. Jenkinson and M. Pollicott, *Computing the dimension of dynamically defined sets: E_2 and bounded continued fractions*, *Ergodic Theory Dynam. Systems* 21 (2001), no. 5, 1429–1445. MR 1855840
25. ———, *Calculating Hausdorff dimensions of Julia sets and Kleinian limit sets*, *Amer. J. Math.* **124** (2002), no. 3, 495–545. MR 1902887 (2003c:37064)
26. ———, *Rigorous effective bounds on the Hausdorff dimension of continued fraction Cantor sets: a hundred decimal digits for the dimension of E_2* , *Adv. Math.* 325 (2018), 87–115. MR 3742587
27. B. Lemmens and R. D. Nussbaum, *Nonlinear Perron-Frobenius Theory*, *Cambridge Tracts in Mathematics*, 189 (2012), Cambridge University Press, Cambridge.
28. M. A. Krasnosel'skij, Je. A. Lifshits, and A. V. Sobolev, *Positive linear systems*, *Sigma Series in Applied Mathematics*, vol. 5, Heldermann Verlag, Berlin, 1989, The method of positive operators, Translated from the Russian by Jürgen Appell. MR 1038527 (91f:47051)
29. Bas Lemmens and Roger Nussbaum, *Birkhoff's version of Hilbert's metric and its applications in analysis*, *Handbook of Hilbert geometry*, IRMA Lect. Math. Theor. Phys., vol. 22, Eur. Math. Soc., Zürich, 2014, pp. 275–303. MR 3329884
30. R. Daniel Mauldin and Mariusz Urbański, *Dimensions and measures in infinite iterated function systems*, *Proc. London Math. Soc.* (3) **73** (1996), no. 1, 105–154. MR 1387085 (97c:28020)
31. ———, *Graph directed Markov systems*, *Cambridge Tracts in Mathematics*, vol. 148, Cambridge University Press, Cambridge, 2003, Geometry and dynamics of limit sets. MR 2003772 (2006e:37036)
32. A. A. Markov, *On a question by D. I. Mendeleev*, *Zap. Imp. Akad. Nauk SPb.* (1890) 62: 1–24.
33. R. D. Nussbaum, *The radius of the essential spectrum*, *Duke Math. J.* 37 (1970), 473–478. MR 0264434
34. ———, *C^m positive eigenvectors for linear operators arising in the computation of Hausdorff dimension*, *Integral Equations Operator Theory* 84 (2016), no. 3, 357–393.
35. R. D. Nussbaum, A. Priyadarshi, and S. Verduyn Lunel, *Positive operators and Hausdorff dimension of invariant sets*, *Trans. Amer. Math. Soc.*, 364 (2012), no. 2, 1029–1066. MR 2846362

36. T. Rivlin, *The Chebyshev polynomials: from Approximation Theory to Algebra and Number Theory*, Dover Publications, Inc., Mineola, New York, 2020 (second edition).
37. David Ruelle, *Thermodynamic formalism*, Encyclopedia of Mathematics and its Applications, vol. 5, Addison-Wesley Publishing Co., Reading, Mass., 1978, The mathematical structures of classical equilibrium statistical mechanics, With a foreword by Giovanni Gallavotti and Gian-Carlo Rota. MR 511655 (80g:82017)
38. ———, *Bowen's formula for the Hausdorff dimension of self-similar sets*, Scaling and self-similarity in physics (Bures-sur-Yvette, 1981/1982), Progr. Phys., vol. 7, Birkhäuser Boston, Boston, MA, 1983, pp. 351–358. MR 733478 (85d:58051)
39. Hans Henrik Rugh, *On the dimensions of conformal repellers. Randomness and parameter dependency*, Ann. of Math. (2) **168** (2008), no. 3, 695–748. MR 2456882 (2010b:37131)
40. H. Samelson, *On the Perron-Frobenius theorem*, Michigan Math. J. 4 (1957), 57–59. MR 0086041 (19,114e)
41. Andreas Schief, *Self-similar sets in complete metric spaces*, Proc. Amer. Math. Soc. **124** (1996), no. 2, 481–490. MR 1301047 (96h:28015)
42. A. Shadrin, *Twelve proofs of the Markov inequality*, D.K. Dimitrov, et al. (Eds.), Approximation Theory: A Volume Dedicated to Borislav Bojanov, Marin Drinov Acad. Publ. House, Sofia (2004), 233–299. MR 2117091
43. B.-S. Tam, *A cone- theoretic approach to the spectral theory of positive linear operators: The finite dimensional case*, Taiwanese J. Math. 5(2), (2001), 207–277. MR 1832167
44. J. S. Vandergraft, *Spectral Pproperties of matrices which have invariant cones*, SIAM J. Appl. Math., 16(6), (1968), 1208–1222. MR 244284

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

Email address: falk@math.rutgers.edu

URL: <http://www.math.rutgers.edu/~falk/>

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

Email address: nussbaum@math.rutgers.edu

URL: <http://www.math.rutgers.edu/~nussbaum/>