

Monochromatization of e^+e^- colliders with a large crossing angle

V. I. Telnov^{1,*}

¹*Budker Institute of Nuclear Physics SB RAS, Novosibirsk, 630090, Russia
Novosibirsk State University, Novosibirsk, 630090, Russia*

(Dated: 31 August 2020)

The relative center-of-mass energy spread σ_W/W at e^+e^- colliders is $\mathcal{O}(10^{-3})$, which is much larger than the widths of narrow resonances produced in the s -channel in e^+e^- collisions. This fact greatly lowers the resonance production rates of the J/ψ , $\psi(2S)$, $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ mesons and makes it extremely difficult to observe resonance production of the Higgs boson. Thus, a significant reduction of the center-of-mass energy spread would open up great opportunities in the search for new physics in rare decays of narrow resonances, the search for new narrow states with small $\Gamma_{e^+e^-}$, the study of true muonium and tauonium, etc. The existing monochromatization scheme is only suitable for head-on collisions, while e^+e^- colliders with crossing angles (the so-called Crab Waist collision scheme) can provide significantly higher luminosity due to reduced collision effects. In this paper, we propose a new monochromatization method for colliders with a large crossing angle. The contribution of the beam energy spread to the spread of the center-of-mass energy is canceled by introducing an appropriate energy-angle correlation at the interaction point; $\sigma_W/W \sim (3-5) \times 10^{-6}$ appears possible. Limitations of the proposed method are also considered.

PACS numbers: 29.20

The point-like nature of the electron and a narrow energy spread are important advantages of e^+e^- colliders. The energy spread occurs due to synchrotron radiation (SR) in the rings as well as beamstrahlung at the IP (important for Z and H factories). The energy spread due to SR depends mainly on the beam energy E_0 and magnetic radius of the ring R , and only weakly on the specific design of the collider. For uniform rings without damping wigglers $\sigma_E/E \approx 0.86 \times 10^{-3} E[\text{GeV}]/\sqrt{R[\text{m}]}$. The energy spreads for some of the existing and planned e^+e^- rings are given in Table I.

TABLE I. Beam energy at circular e^+e^- colliders

	VEPP-2000	BEPC-II	SuperKEKB	FCC-ee
E_0 , GeV	1	~ 2	4-7	62.5
$2\pi R$, km	0.024	0.24	3	100
σ_E/E , 10^{-3}	~ 0.6	~ 0.5	0.7	0.6 (w/o BS)

One can see that the invariant mass spread $\sigma_W/W = (1/\sqrt{2})\sigma_E/E \sim (0.35-0.5) \times 10^{-3}$. This spread is much greater than the widths of the narrow e^+e^- resonances J/ψ , $\psi(2S)$, $\Upsilon(1S)$, $\Upsilon(2S)$, $\Upsilon(3S)$ and the Higgs boson, see Table II. The resonance width Γ is the full width at half maximum, so one should compare Γ/m and $2.36 \sigma_W/W \approx (0.8-1.2) \times 10^{-3}$.

TABLE II. Width of some narrow e^+e^- resonances

	J/ψ	$\psi(2S)$	$\Upsilon(1S)$	$\Upsilon(2S)$	$\Upsilon(3S)$	$H(125)$
m , GeV/ c^2	3.097	3.686	9.460	10.023	10.355	125
Γ , keV	93	300	54	32	20.3	4200
Γ/m , 10^{-5}	3	8	0.57	0.32	0.2	3.4
$2.36 \sigma_W/W$	~ 35	~ 13	~ 180	~ 310	~ 500	~ 30

One of the promising directions for particle physics is the study of rare and forbidden processes sensitive

to new physics. Therefore, J/ψ and Υ factories with a narrow invariant-mass spread would be good candidates for future experimental facilities. In the case of a large continuum background, the signal-to-noise ratio $S/\sqrt{B} \propto (\mathcal{L}_{\text{int}}/\sigma_W)/\sqrt{\mathcal{L}_{\text{int}}} = \sqrt{\mathcal{L}_{\text{int}}/\sigma_W}$; therefore, the integrated luminosity required to observe a rare decay of a known resonance (or to observe a narrow resonance with a very small $\Gamma_{e^+e^-}$) $\mathcal{L}_{\text{int}} \propto (1/\sigma_W)^2$. A 100-fold improvement in monochromaticity for Υ -mesons would be equivalent to a luminosity increase by a factor of $100^2 = 10000$! In the absence of a background, monochromatization lowers the observable branching limit proportionally to σ_W .

The first consideration of energy monochromatization in e^+e^- collisions dates back to mid-1970s [1]. In the proposed scheme (Fig. 1), beams collide head-on and have a horizontal or vertical energy dispersion at the interaction point (IP), opposite in sign for the e^+ and e^- beams. As a result, the particles collide with opposite energy deviations, $E_0 + \Delta E$ and $E_0 - \Delta E$, and their invariant mass $W \approx 2\sqrt{E_1 E_2} \approx 2E_0 - (\Delta E)^2/E_0$ is very close to $2E_0$. This monochromatization scheme was con-




FIG. 1. Existing monochromatization scheme for head-on collisions [1].

sidered by many authors in 1980s–1990s [2–8] for use in c - τ and B -factory projects (i.e., in the energy range of the ψ and Υ resonances); however, none of the proposals were implemented as they all resulted in an unaccept-

able loss of luminosity. A new wave of interest in this topic is connected with the FCC-ee and CEPC projects, where one of the processes of great interest is $e^+e^- \rightarrow H$, which has a tiny cross section and is observable only if $\sigma_W \lesssim \Gamma_H$ [9, 10].

The new generation of circular e^+e^- colliders (DAΦNE[11], SuperKEKB[13], $c\tau$ [14], FCC-ee[15], CEPC[16]) rely on the so-called “crab-waist” collision scheme [11, 12], where the beams collide at an angle $\theta_c \gg \sigma_x/\sigma_z$. When the luminosity is limited by the tune shift, characterized by the beam-beam strength parameter ξ_y , the crab-waist collision scheme allows much higher luminosities to be reached compared to head-on collisions. For the same beam currents, the maximum luminosity for head-on collisions $\mathcal{L} \propto 1/\sigma_z$, where σ_z is the bunch length, while for collisions at an angle $\mathcal{L} \propto 1/\beta_y$, where the vertical beta function $\beta_y \sim \sigma_x/\theta_c$ can be ~ 20 – 30 times smaller than σ_z ; as a result, the luminosity can be higher by the same factor. In existing designs, the crossing angle θ_c varies between 30 mrad (FCC-ee) and 83 mrad (SuperKEKB). In what follows, we propose and explore significant modifications to this collision scheme aimed at achieving monochromatization.

First, let us consider the mass resolution in the unmodified collision scheme with a crossing angle, Fig 2. The invariant mass of the produced system




FIG. 2. Collisions with crossing angles.

$$\begin{aligned} W^2 &= (P_1 + P_2)^2 = 2m^2 + 2(E_1 E_2 - \vec{p}_1 \vec{p}_2) \approx \\ &\approx 2E_1 E_2 (1 + \cos(\theta_1 + \theta_2)). \end{aligned} \quad (1)$$

Here, we neglect the terms of the relative order $(m/E)^2$ and choose units so that $c = 1$. The contribution of the vertical angular spread is negligible in all practical cases.

By differentiating this formula while assuming that the energies and the angles are independent and setting $\theta_1 = \theta_2 = \theta_c/2$ and $E_1 = E_2 = E$, we find the relative mass spread

$$\left(\frac{\sigma_W}{W}\right)^2 = \frac{1}{2} \left(\frac{\sigma_E}{E}\right)^2 + \frac{1}{2} \frac{\sin^2 \theta_c}{(1 + \cos \theta_c)^2} \sigma_\theta^2, \quad (2)$$

where θ_c is the beam crossing angle, σ_E is the beam energy spread, and σ_θ is the beam angular spread at the IP, which is determined by the horizontal beam emittance ε_x and beta function β_x^* at the IP: $\sigma_\theta = \sqrt{\varepsilon_x/\beta_x^*}$. For head-on collisions, the second term vanishes, and the mass resolution is determined solely by the beam energy




FIG. 3. Collisions with the energy-angle correlation.

spread. In the aforementioned colliders with the crab-waist scheme, the contribution of beam energy spread is also dominant.

The presently proposed monochromatization method is based on the fact that the invariant mass W^2 depends on both the beam energies and their crossing angle. The second term in Eq. 2 reflects the natural stochastic beam spread due to the horizontal beam emittance and cannot be avoided; however, the first term can be suppressed very significantly, as we shall demonstrate. For simplicity of further consideration, hereinafter we omit the second term (i.e., assume $\sigma_\theta = 0$) as it is independent of the first and can be added back later. In the proposed method, we provide the beams with an angular dispersion such that a beam particle arrives to the IP with a horizontal angle that depends on its energy: the higher the energy, the larger the angle, Fig. 3. We can choose such a dispersion that when a particle coming from the left, with energy $E_0 + dE$ and angle $\theta = \theta_c/2 + d\theta$, collides with a particle coming from the right, with the nominal (average) energy E_0 and angle $\theta_c/2$, they produce the same invariant mass as two colliding particles that both have the nominal energies and angles, E_0 and $\theta_c/2$. From Eq. 1, we obtain the required condition:

$$(E_0 + dE)E_0(1 + \cos(\theta_c + d\theta)) = E_0^2(1 + \cos \theta_c). \quad (3)$$

In the linear approximation, this gives the required angular dispersion (the same for both particles)

$$d\theta_i = \frac{1 + \cos \theta_c}{\sin \theta_c} \frac{dE_i}{E_0}. \quad (4)$$

In what follows, the exact dispersion relation (Eq. 3) will be called “nonlinear”, while Eq. 4 will be referred to as “linear” dispersion. Taking the first derivative of W^2 (Eq. 1) and substituting the linear dispersion from Eq. 4, one can confirm that the resulting variation of the invariant mass is indeed zero:

$$\begin{aligned} dW^2 &= 2E_2((1 + \cos \theta_c) dE_1 - E_1 \sin \theta_c d\theta_1) + \\ &+ 2E_1((1 + \cos \theta_c) dE_2 - E_2 \sin \theta_c d\theta_2) = 0. \end{aligned} \quad (5)$$

Note that the proposed monochromatization method works even for unequal beam energies. For simplicity, in the rest of the paper we will usually assume $E_1 = E_2$.

If W were a product of two functions that depend, respectively, only on the parameters of the left or the right colliding particles, then the contribution of the beam energy spreads σ_{E_i} to σ_W could have been completely zeroed using the nonlinear dispersion (Eq. 3). However, the contributions of the two beams are not factorized due to the $\cos(\theta_1 + \theta_2)$ term in Eq. 1. Therefore, a second-order contribution from the energy spread remains; below, we will find it for both the linear and nonlinear dispersions.

Since the first derivative of W is zero (Eq. 5), we must use the quadratic term of the Taylor series:

$$dW^2 = (1/2!) d^2W^2(E_1, E_2, \theta_1, \theta_2), \quad (6)$$

where W^2 is given by Eq. 1 and

$$d^2W^2 = \left(dE_1 \frac{\partial}{\partial E_1} + dE_2 \frac{\partial}{\partial E_2} + d\theta_1 \frac{\partial}{\partial \theta_1} + d\theta_2 \frac{\partial}{\partial \theta_2} \right)^2 W^2. \quad (7)$$

Then, in the resulting expression we replace $d\theta_1$ and $d\theta_2$ by dE_1 and dE_2 using Eq. 4. As a result, we get an expression in the form $dW/W = a((dE_1)^2 + (dE_2)^2) + b(dE_1 dE_2)$. In the case of ideal nonlinear dispersion (Eq. 3), only the unfactorized term $\cos(\theta_1 + \theta_2)$ contributes to dW ; therefore, $a = 0$, and dW is described by the term $b(dE_1 dE_2)$, which depends on both beams. In the case of linear dispersion (Eq. 4), both terms contribute and, in addition to fluctuations, there is also a small shift of the mean invariant mass, ΔW .

In the case of Gaussian beam energy distributions with r.m.s. spread σ_E , we have $\overline{(dE)^2} = \sigma_E^2$, $\sigma(dE)^2 = \sqrt{2}\sigma_E^2$, $\overline{dE_1 dE_2} = 0$, $\sigma(dE_1 dE_2) = \sigma_E^2$, the mass spreads from the two beams must be summed quadratically. In addition, the fluctuations of the first and the second terms are independent and must be summed quadratically.

Finally, for ideal nonlinear dispersion (Eq. 3), the mass spread due to the beam energy spread

$$\left(\frac{\sigma_W}{W} \right)_E = \frac{\sigma_E^2}{2E^2} \frac{1 + \cos \theta_c}{\sin^2 \theta_c}, \quad \frac{\Delta W}{W} = 0. \quad (8)$$

For linear dispersion (Eq. 4)

$$\left(\frac{\sigma_W}{W} \right)_E = \frac{\sigma_E^2}{2E^2} \left[\left(1 + \frac{1 + \cos \theta_c}{\sin^2 \theta_c} \right)^2 + \left(\frac{1 + \cos \theta_c}{\sin^2 \theta_c} \right)^2 \right]^{1/2}, \quad (9)$$

$$\frac{\Delta W}{W} = \frac{\sigma_E^2}{2E^2} \left(1 + \frac{1 + \cos \theta_c}{\sin^2 \theta_c} \right). \quad (10)$$

The total invariant mass spread is the sum of the residual contribution of the energy spread (Eq. 8 or Eq. 9) and the second term of Eq. 2, which is due to the angular spread:

$$\left(\frac{\sigma_W}{W} \right)^2 = \left(\frac{\sigma_W}{W} \right)_E^2 + \frac{1}{2} \frac{\sin^2 \theta_c}{(1 + \cos \theta_c)^2} \sigma_\theta^2 \quad (11)$$




FIG. 4. Monochromaticity of collisions vs collision angle.

These formulas have been verified by direct simulation. The dependence of the invariant mass spread on the collision angle θ_c is shown in Fig. 4. It can be seen that linear dispersion works almost as well as the best-case nonlinear dispersion. Now, let us discuss some of the limitations of this monochromatization method.

First, the required dispersion (Eq. 4) is unacceptably large at small collision angles. The large horizontal angular spread at the IP requires very strong final quadrupole magnets that are located in the places with high dispersion, which leads to strong synchrotron radiation and, as a consequence, to an additional energy spread and deterioration of the horizontal emittance ε_x . For the energy spread $\sigma_E/E = 0.5 \times 10^{-3}$, the r.m.s. angular spread due to the dispersion is $\sigma_{\theta,d} = 1.87 \times 10^{-3}$ for $\sin \theta_c = 0.5$ and 0.5×10^{-3} at $\sin \theta_c = 1$. There are fewer problems at large collision angles; however, W decreases as $W(\theta_c)/W(0) = \cos(\theta_c/2)$, which is 0.966, 0.949, 0.926, 0.707 for $\sin \theta_c = 0.5, 0.6, 0.7, 1$, respectively. Therefore, we take $\sin \theta_c = 0.5$ as a reasonable benchmark for further estimates.

The horizontal angular spread at the IP $\sigma_\theta = \sqrt{\varepsilon_x/\beta_x^*}$, where β_x^* is the value of the horizontal beta function at the IP. From Fig. 4, we see that in order for the contributions of the residual energy spread and the horizontal angular spread (only due to emittance) to be equal, the latter should be $\sigma_\theta \lesssim 10^{-5}$ for $\sin \theta_c = 0.5$. The horizontal emittance at the 7 (4) GeV KEK Super B factory is $\varepsilon_x = 4.8 (3.3) \times 10^{-9}$ m. Specialized synchrotron-radiation sources have (and are planning) $\varepsilon_x < 10^{-10}$ m at energies $E = 3-6$ GeV. The maximum value of β_x^* is limited by the distance between the IP and the final quadrupole, which is about 1 m. So, $\sigma_\theta = \sqrt{10^{-10}/1} \sim 10^{-5}$ can be thought of as a feasible ultimate target that achieves $\sigma_W/W \approx 2.5 \times 10^{-6}$, which is about 150 to 200 times better than that at current and past e^+e^- storage rings.

Let us consider further limitations. Below, we consider the following effects:

- the increase of ε_x and σ_E/E due to radiation in the final focus quadrupoles;
- the increase of ε_x and σ_E/E in the magnetic field of the detector;
- the increase of ε_x and σ_E/E in the part of the ring where dispersion is created;
- the beam-beam attraction, which influences the collision angle.

The difference in the horizontal positions of the particles with energies $E_0 + dE$ and E_0 is $dx = D_x dE/E$, where D_x is the horizontal dispersion function. Synchrotron radiation in regions with $D_x \neq 0$ leads to an increase of the emittance [17]:

$$\Delta\varepsilon_x = A \int \frac{\mathcal{H}(s)}{|\rho|^3} ds, \quad A = \frac{55}{48\sqrt{3}} \frac{r_e^2 \gamma^5}{\alpha}. \quad (12)$$

where $\alpha = e^2/\hbar c \approx 1/137$, $r_e = e^2/mc^2$ is the classical electron radius, $\gamma = E/mc^2$. The dispersion invariant $\mathcal{H}(s) = \beta_x D_x'^2 + 2\alpha_x D_x D_x' + \gamma_x D_x^2$, where $\beta_x, \alpha_x, \gamma_x$ are optical functions (Twist parameters), $\alpha_x = -\beta_x/2$, $\gamma_x = (1 + \alpha_x^2)/\beta_x$.

Synchrotron radiation in quadrupoles and the dispersion sections (in addition to the ring magnets) also leads to an increase in the energy spread:

$$\Delta\left(\frac{\sigma_E}{E}\right)^2 = A \int \frac{ds}{|\rho|^3}. \quad (13)$$

To understand the importance of these effects, let us estimate the equilibrium energy spread and horizontal emittance when they are caused only by these effects but their damping is provided by synchrotron radiation in the rings with damping times $\tau_E = 3R^2/2\gamma^3 cr_e$, $\tau_x = 2\tau_E$. For one interaction point, they are

$$\Delta\left(\frac{\sigma_E}{E}\right)^2 = \frac{\tau_E A}{2T_{\text{rev}}} \int \frac{ds}{|\rho|^3} \approx 10.8 r_e \gamma^2 R \int \frac{ds}{|\rho|^3} \quad (14)$$

$$\Delta\varepsilon_x = \frac{\tau_x A}{2T_{\text{rev}}} \int \frac{\mathcal{H}(s) ds}{|\rho|^3} \approx 21.6 r_e \gamma^2 R \int \frac{\mathcal{H}(s) ds}{|\rho|^3} \quad (15)$$

Let us estimate the radiation effects in the final quadrupole of length l_q located at a distance F from the IP. The particles enter the quadrupole from the IP at angles given by Eq. 4 and exit parallel to its axis, therefore

$$\frac{1}{\rho} \sim \frac{1 + \cos \theta_c}{l_q \sin \theta_c} \frac{dE}{E}; \quad \left\langle \frac{ds}{|\rho|^3} \right\rangle \approx \frac{1.7}{l_q^2} \left(\frac{1 + \cos \theta_c}{\sin \theta_c} \right)^3 \left(\frac{\sigma_E}{E} \right)^3, \quad (16)$$

where the coefficient 1.7 is obtained numerically for $F = l_q$ after averaging over the energies (although these calculations are estimates, I keep the numerical coefficients where possible).

Similarly, for $F = l_q$ we obtain

$$\frac{\mathcal{H}(s) ds}{|\rho|^3} \approx \frac{4}{\beta_x} \left(\frac{1 + \cos \theta_c}{\sin \theta_c} \right)^5 \left(\frac{\sigma_E}{E} \right)^3, \quad (17)$$

where β_x is the horizontal beta function on the outside of the quadrupole. Substituting to Eqs. 16, 17 and multiplying by 2 to account for the quadrupoles from both sides of the IP, we obtain a reasonable estimate of the additional energy spread and emittance due to radiation in the final quadrupoles:

$$\Delta\left(\frac{\sigma_E}{E}\right)^2 \approx \frac{40 r_e \gamma^2 R}{F^2} \left(\frac{1 + \cos \theta_c}{\sin \theta_c} \right)^3 \left(\frac{\sigma_E}{E} \right)^3 \propto \frac{E^5}{\sqrt{R}}; \quad (18)$$

$$\Delta\varepsilon_x = 170 \frac{r_e \gamma^2 R}{\beta_x} \left(\frac{1 + \cos \theta_c}{\sin \theta_c} \right)^5 \left(\frac{\sigma_E}{E} \right)^3 \propto \frac{E^5}{\sqrt{R}}. \quad (19)$$

For example: for $E = 5$ GeV, $R = 500$ m, $\sin \theta_c = 0.5$, $\sigma_E/E = 0.5 \times 10^{-3}$, $F = \beta_x = 1$ m, we get $\Delta(\sigma_E/E)^2 = 3.6 \times 10^{-11}$, $\Delta\varepsilon_x = 2.1 \times 10^{-9}$ m. Thus, the effect on the energy spread is negligible, and the increase of the horizontal emittance is two times smaller than the SuperKEKB emittance.

For FCC-ee with $R \approx 15$ km and $E = 50$ GeV, the energy spread will still be acceptable but the emittance will be 4 orders of magnitude greater than its design value, 3×10^{-10} m. Unfortunately, this means that this method of monochromatization does not work at high energies ($E \gtrsim 5$ GeV).

Let us now consider the influence of the magnetic field of the detector solenoid. The dispersion in the solenoid (created outside the detector for this method of monochromatization) is proportional to the distance from the IP and at $z = F$ is the same as in the last quadrupole. So, to evaluate the effects in the detector we can compare the field in the quadrupole and the effective transverse field of the solenoid $B_{s,\perp} = B_s \sin(\theta_c/2)$. The field in the quadrupoles for particles with $E = E_0$ can be found from $l_q/\rho \approx (\sigma_E/E)(1 + \cos \theta_c)/\sin \theta_c$, which gives $B_q \sim 0.06 E[\text{GeV}]/l_q[\text{m}]$ T. For $E = 5$ GeV and $l_q = 1$ m, $B_q \sim 0.03$ T. The effective detector field for $B_s = 1$ T and $\sin \theta_c = 0.5$ is $B_{s,\perp} \sim 0.25$ T, which is 8 times greater than that in the quadrupoles. This is a very serious problem because, as shown above, $\Delta\varepsilon_x$ due to radiation in the quadrupoles is already close to the acceptable limit for $E = 5$ GeV. Possible solutions: a) compensate the field of the detector solenoid on the axis with an antisolenoid (they are used in any case in storage rings for the removal of x - y coupling), leaving only a short free space ($\sim \pm 10$ cm) for detecting the produced particles, and at the same time reduce B_s . In this approach, the solenoid and quadrupole contributions would be comparable at $B_s \sim 0.5$ T, which is acceptable, in principle, for this collider energy; b) to use a toroidal magnetic field, as in the ATLAS muon detector, in which case the magnetic field at the axis is zero. So, this is a serious problem, but unlike the same problem in the quadrupoles, it has solutions.

Let us estimate what distance is needed to create the required dispersion $D_x = 1.5 F(1 + \cos \theta_c)/\sin \theta_c$ at the

entrance to the final focus quadrupoles (the factor 1.5 is due to the difference in dispersions at the quadrupole entrance and exit for $F = l_q$). In order for β_x^* at the IP to be as large as possible ($\sim F$), the β_0 function at the entrance to the quadrupole should also be $\beta_0 \sim F$ (because $F^2 \approx \beta_x^* \beta_0$). The length of the dispersion system is determined by the increase of emittance due to radiation. We consider the scheme shown in Fig. 5 with explanatory notes.




FIG. 5. A scheme of the chromatic section used for estimates.

For this scheme, $\int \mathcal{H}(s) ds / |\rho|^3 \approx 20 (D_x)^5 / L^5 \beta_0$, where $L = 1.5l$ is the total length. Substituting into Eqs. 16, 17 and multiplying by 2 (both sides of the IP), we obtain the additional emittance due to radiation in the dispersion sections (for $\beta_0 = F$)

$$\Delta \varepsilon_x \sim 6500 \frac{r_e \gamma^2 R F^4}{L^5} \left(\frac{1 + \cos \theta_c}{\sin \theta_c} \right)^5. \quad (20)$$

For $E = 5$ GeV, $R = 500$ m, $\sin \theta_c = 0.5$ and $F = 1$ m, we get $\Delta \varepsilon_x \sim 10^{-9}$ m at $L = 230$ m. It would be 2 times shorter for $F = 0.7$ m and $\sin \theta_c = 0.7$.

Our consideration of radiation effects in the final quadrupoles and chromatic sections have shown that this monochromatization method works well at and below the energies of the Υ mesons. For the above example ($2E = 10$ GeV, $R = 500$ m, $\sin \theta_c = 0.5$) with the horizontal emittance limited by Eq. 19 one can have an angular spread $\sigma_{\theta_x} \sim \sqrt{\varepsilon_x / \beta_x^*} \sim \sqrt{2 \times 10^{-9} / 1} \sim 4.5 \times 10^{-5}$, which corresponds to the mass width $\sigma_W / W \sim 8.5 \times 10^{-6}$. In this case the monochromaticity is determined by the angular spread, see Fig. 4. Similar considerations for $\sin \theta_c = 0.6(0.7)$ give $\sigma_W / W \sim 6.1(4.5) \times 10^{-6}$. These values of the monochromaticity (for $\sin \theta_c = 0.5-0.7$) are 70–110 times better than at the existing e^+e^- storage rings.

According to Eq. 19 the horizontal emittance limit due to radiation in final quadrupoles varies as E^5 / \sqrt{R} and it becomes insignificant for low energies. As noted, the existing and planned synchrotron radiation sources with $E = 3-6$ GeV have emittances of $\varepsilon_x < 10^{-10}$ m, then the angular spread $\sigma_{\theta_x} \sim \sqrt{10^{-10} / 1} = 10^{-5}$ and $\sigma_W / W \sim 2.5 \times 10^{-6}$ is in reach, that corresponds to a 150-fold improvement in monochromaticity. In the energy region of J/Ψ and below, the intra-beam scattering is important, and realistic numbers can be obtained only after careful consideration. With decreasing energy, the emittance ε_x and σ_{θ_x} increase, therefore the optimum crossing angle will decrease, see Fig. 4.

At high-luminosity e^+e^- factories, the number of particles per bunch is large, $N \approx (5-10) \times 10^{10}$. The question arises: how does the collision angle change due to the attraction of the beams? Simple estimates indicate that this effect can be problematic. However, a detailed examination unexpectedly shows that beam attraction does not affect the invariant mass of the colliding particles. Indeed, let us consider relativistic particles with the energy E that are attracted by an opposing relativistic beam that creates an electric field \mathcal{E} and a magnetic field $B \approx \mathcal{E}$. At distance ds , the particle receives energy $dE = e\mathcal{E} \sin \theta_c ds \approx eB \sin \theta_c ds$ and an additional angle $d\theta \approx (e\mathcal{E} \cos \theta_c + eB) ds / E \approx eB(1 + \cos \theta_c) ds / E$. Substituting dE and $d\theta$ for this and similar opposite particle in Eq. 1, we show that $dW = 0$!

A few words about the possible loss of luminosity due to monochromatization. The only difference compared to the SuperKEKB design would be a larger crossing angle, 500 mrad instead of 90 mrad. The luminosity $\mathcal{L} \propto N(Nf) / (\sigma_z \sigma_y \theta_c)$. For the same beams, an increase of the crossing angle by a factor of 6 would mean a loss of luminosity by the same factor. However, the collision effects would become weaker, and one can partially compensate for the loss by increasing N and decreasing σ_z . Other limitations are present, and so the resulting luminosity would be lower, perhaps by a factor of two or three. Such a decrease is acceptable because monochromatization would significantly increase the effective luminosity, $\propto 1/\sigma_W^2$ when studying rare decays in presence of a large continuum background.

In conclusion, a new method of monochromatization of e^+e^- collisions is being proposed, which works at large crossing angles ($\theta_c \gtrsim 0.3-0.5$ rad) and allows high luminosities due to reduced collision effects (as in the crab-waist collision scheme). The contribution of the beam energy spread to the invariant mass is compensated by introducing an appropriate energy-angle correlation at the IP. This method requires an energy dispersion in the quadrupoles, at the IP it is equal to zero. The main problem of this method is the increase of the horizontal emittance due to synchrotron radiation in quadrupole magnets, which limits its applicability to $2E_0 \lesssim 10$ GeV. There are other problems, such as the influence of the

detector field, which can be overcome but require non-standard solutions. The achievable invariant-mass spread is $\sigma_W/W \sim (3-5) \times 10^{-6}$, which is about 100 times better than at the past and existing e^+e^- storage rings. There are two main directions in particle physics: higher energies—or very high luminosities at relatively low energies. Monochromatization is a very natural next step in the development of the next generation of luminosity-frontier colliders. It can increase by several orders of magnitude the effective luminosity in the study of rare decays or looking for narrow states with a small $\Gamma_{e^+e^-}$. The full potential of this method can be realized at the very narrow $\Upsilon(nS)$ resonances as well as at lower energies, where a lot of interesting physics is also present. This paper is just the beginning; the next step toward realistic projects requires the efforts of accelerator designers.

This work was supported by Russian Ministry of Education and Science and RFBR-DFG Grant No 20-52-12056.

* telnov@inp.nsk.su

- [1] A. Renieri, “Possibility of Achieving Very High-Energy Resolution in electron-Positron Storage Rings,” Laboratori Nazionali di Frascati Report No. LNF-75/6(R).
- [2] I. Y. Protopopov, A. N. Skrinsky and A. A. Zholents, “Energy monochromatization of particle interaction in storage rings,” IYF-79-06.
- [3] A. A. Avdienko *et al.*, “The Project of Modernization of the VEPP-4 Storage Ring for Monochromatic Experiments in the Energy Range of J/ψ and Υ Mesons,” Proc. 12th Intern. Conf. High Energy Accelerators, Fermilab, 1983, p. 186.
- [4] K. Wille and A. W. Chao, “Investigation of a Monochromator Scheme for SPEAR,” SLAC/AP-32 (1984).
- [5] M. Jowett, “Feasibility of a Monochromator Scheme in LEP,” CERN LEP Note 544, September (1985).
- [6] Yu. I. Alexahin, A. Dubrovin, A. A. Zholents, “Proposal on a Tau-Charm Factory with Monochromatization,” Proc. 2nd European Particle Accelerator Conference, Nice, France, 12–16 June 1990, p. 398.
- [7] A. Zholents, “Polarized J/ψ Mesons at a Tau-Charm Factory with a Monochromator Scheme,” CERN SL/97-27, June 1992.
- [8] A. Faus-Golfe and J. Le Duff, “Versatile DBA and TBA Lattices for a Tau-Charm Factory with and without Beam Monochromatization,” Nucl. Instr. Methods A 372 (1996).
- [9] D. d’Enterria, “Higgs physics at the Future Circular Collider,” PoS **ICHEP2016**, 434 (2017).
- [10] M. A. Valdivia Garcia, A. Faus-Golfe and F. Zimmermann, “Towards a Mono-chromatization Scheme for Direct Higgs Production at FCC-ee,” CERN-ACC-2016-0077.
- [11] M. Zobov, D. Alesini *et al.*, “Test of crab-waist collisions at DAFNE Phi factory,” Phys. Rev. Lett. **104**, 174801 (2010).
- [12] M. Bona *et al.* [M. Bona *et al.* “SuperB: Conceptual Design Report,” INFN-AE-07-02, arXiv:0709.0451 [hep-ex].
- [13] Y. Ohnishi *et al.*, “Accelerator design at SuperKEKB,” PTEP **2013**, 03A011 (2013)
- [14] A. E. Bondar *et al.* “Project of a Super Charm-Tau factory at the Budker Institute of Nuclear Physics in Novosibirsk,” Yad. Fiz. **76**, no. 9, 1132–1145 (2013).
- [15] A. Abada *et al.* “FCC-ee: The Lepton Collider,” Eur. Phys. J. ST **228**, no. 2, 261–623 (2019).
- [16] “CEPC Conceptual Design Report: Volume 1 - Accelerator,” arXiv:1809.00285.
- [17] H. Wiedemann, “Particle Accelerator Physics” (Springer-Verlag, 2007).