

TOWERS OF BUBBLES FOR YAMABE-TYPE EQUATIONS AND FOR THE BRÉZIS-NIRENBERG PROBLEM IN DIMENSIONS $n \geq 7$

BRUNO PREMOSELLI

ABSTRACT. Let (M, g) be a closed locally conformally flat Riemannian manifold of dimension $n \geq 7$ and of positive Yamabe type. If ξ_0 denotes a non-degenerate critical point of the mass function we prove the existence, for any $k \geq 1$ and $\varepsilon > 0$, of a *positive* blowing-up solution u_ε of

$$\Delta_g u_\varepsilon + (c_n S_g + \varepsilon h) u_\varepsilon = u_\varepsilon^{2^*-1},$$

that blows up like the superposition of k positive bubbles concentrating at different speeds at ξ_0 . The method of proof combines a finite-dimensional reduction with the sharp pointwise analysis of solutions of a linear problem. As another application of this method of proof we construct *sign-changing* blowing-up solutions u_ε for the Brézis-Nirenberg problem

$$\Delta_\xi u_\varepsilon - \varepsilon u_\varepsilon = |u_\varepsilon|^{\frac{4}{n-2}} u_\varepsilon \text{ in } \Omega, \quad u_\varepsilon = 0 \text{ on } \partial\Omega$$

on a smooth bounded open set $\Omega \subset \mathbb{R}^n$, $n \geq 7$, that look like the superposition of k positive bubbles of alternating sign.

1. INTRODUCTION AND STATEMENT OF THE RESULTS

1.1. Positive towers of bubbles in the Riemannian case. Let (M, g) be a closed smooth connected Riemannian manifold, that is compact without boundary, of dimension $n \geq 3$. Assume (M, g) is of positive Yamabe type. The main goal of this work is to construct blowing-up families of positive solutions $(u_\varepsilon)_\varepsilon$ of critical nonlinear elliptic equations of the following form:

$$\Delta_g u_\varepsilon + h_\varepsilon u_\varepsilon = u_\varepsilon^{2^*-1}, \tag{1.1}$$

where $\Delta_g = -\operatorname{div}_g(\nabla \cdot)$, $2^* = \frac{2n}{n-2}$ and where h_ε converges in $C^1(M)$ as $\varepsilon \rightarrow 0$. We construct sequences that display *towering* bubbling phenomena when $h_\varepsilon \rightarrow c_n S_g$, where $c_n = \frac{n-2}{4(n-1)}$ and where S_g is the scalar curvature of (M, g) .

When $h_\varepsilon \equiv c_n S_g$ equation (1.1) is the celebrated Yamabe equation. The existence of a positive solution in this case was proven through the work of Yamabe [42], Aubin [1], Trudinger [41] and Schoen [37]. The question of the compactness of the set of positive solutions of the Yamabe equation on manifolds of positive Yamabe type has generated a vast amount of work in the last decades. On the standard sphere (\mathbb{S}^n, g_0) , $n \geq 3$, the set of positive

solutions is not compact, but if (M, g) is not conformally diffeomorphic to the standard sphere, the set of positive solutions of the Yamabe equation is compact when the dimension of the manifold satisfies $3 \leq n \leq 24$, as was shown by Khuri, Marques and Schoen [18] (previous results were obtained Li and Zhu [23], Druet [9], Marques [24] and Li and Zhang [21]). In dimensions $n \geq 25$ Brendle [4] and Brendle and Marques [5] constructed examples of manifolds for which the set of positive solutions is not compact. From the PDE point of view, understanding the compactness of solutions to the Yamabe equation (and, more generally, of larger classes of nonlinear critical elliptic equations) relies on establishing sharp blow-up estimates near blow-up points. A crucial step in the proof of the compactness of the Yamabe equation consists in showing that, in dimensions $3 \leq n \leq 24$, all the possible blow-up points of a sequence of positive solutions must be *isolated* and *simple* (following the terminology of Schoen [38]) i.e. that around each blow-up point ξ_0 the solution can be approximated by a so called *standard bubble*

$$W_k(x) \sim \frac{\mu_k^{\frac{n-2}{2}}}{\left(\mu_k^2 + \frac{d_g(x, \xi_k)^2}{n(n-2)}\right)^{\frac{n-2}{2}}} \quad \text{for some } \xi_k \in M \text{ and } \mu_k \rightarrow 0. \quad (1.2)$$

For more general equations like (1.1) one may wonder if compactness properties – or, more generally, *stability* properties according to the terminology in Hebey [13] – remain true for positive solutions when h_ε in (1.1) is not everywhere equal to $c_n S_g$. There the picture becomes more complicated. If $n = 3$, no *positive* blowing-up positive solutions exist as long as the mass of the limiting operator is positive as proved by Li-Zhu [23]. If we denote by h_∞ the C^1 limit of h_ε and if (M, g) is not conformally diffeomorphic to the standard sphere, the mass of $\Delta_g + h_\infty$ remains positive even if $h_\infty > c_3 S_g$ everywhere, provided h_∞ is not too large. When $n \geq 4$, Druet [9] proved that (1.1) does not have any positive blowing-up solution when $h_\infty < c_n S_g$ everywhere, or when $n = 4, 5$ and $h_\varepsilon \leq c_n S_g$ for all ε and the manifold is not conformally equivalent to the round sphere.

If $h_\varepsilon > c_n S_g$ somewhere, the situation is subtler and blow-up is likely to occur when $n \geq 4$. We focus in this paper in the following special case of (1.1):

$$\Delta_g u + (c_n S_g + \varepsilon h) u = u^{2^*-1}, \quad (1.3)$$

where $h \in C^0(M)$ and $\varepsilon > 0$. If $n \geq 4$, positive single-bubble blowing-up solutions or multi-bubble solutions with isolated concentration points do exist as shown by Esposito, Pistoia and Vetois in [12]. Pistoia and Vaira proved in [29], when $h \equiv 1$ and $n \geq 7$, the existence of blowing-up solutions of (1.3) possessing a so-called *clustering* blow-up point at a non-degenerate non-zero

minimum point of the Weyl's tensor. In these so-called *clustering* configurations the solutions blow-up as a finite number of peaks, of comparable heights, whose centers converge to the same point. Clustering configurations in dimensions 4 and 5 have also been constructed by Thizy and Vétois [40] and, for families of equations like (1.1), in dimensions $n \geq 6$ by Robert and Vétois [35].

We address here the question of existence of so-called *towering bubbling configurations* for positive solutions of (1.3). These are finite-energy blowing-up positive solutions that blow-up as a superposition of positive bubbles concentrating at different speed, and whose centers converge to the same point – these configurations are also sometimes referred to as *towers of bubbles*. We prove that *towering* blow-up points for (1.3) can be constructed in dimension $n \geq 7$ on locally conformally flat manifolds. Our main result is as follows:

Theorem 1.1. *Let (M, g) be a closed locally conformally flat Riemannian manifold of dimension $n \geq 7$ of positive Yamabe type. Let ξ_0 denote a non-degenerate critical point of the mass function $\xi \mapsto m(\xi)$. Let $k \geq 1$. Then there exists a family $(u_\varepsilon)_\varepsilon$ of positive solutions of*

$$\Delta_g u_\varepsilon + (c_n S_g + \varepsilon) u_\varepsilon = u_\varepsilon^{2^*-1}$$

that blows-up with k bubbles at ξ_0 , and for which ξ_0 is a towering blow-up point.

For the definition of the Riemannian mass we refer to (2.1) below. Morabito, Pistoia and Vaira in [26] first proved the existence of positive towering bubbling configurations on non-locally conformally flat manifolds, at a point ξ_0 where the Weyl's tensor does not vanish, under the stringent assumption that the manifold is symmetric with respect to ξ_0 . In the present work we treat the locally conformally flat case that was left open, and we perform the construction without any symmetry assumption on M : this considerably increases the complexity of the proof and requires us to develop a new analytical approach. Towering bubble configurations for positive solutions similar to the ones constructed in this paper do not exist in dimensions $n \leq 6$, as can be seen by an adaptation of the arguments in Druet [8] (see also Hebey [13], Chapter 8). In low dimensions $n = 4, 5$, the closest analogue to a positive towers of bubbles is made of isolated peaks concentrating at different speeds and has been built in Premoselli-Thizy [32]. Theorem 1.1 thus completes the constructive picture of the possible blow-up scenarios for positive solutions of (1.3).

We stated Theorem 1.1 when $h \equiv 1$ for simplicity. If one allows a more general function $h \in C^1(M)$ (compare with (1.3)) it is still possible to prove the existence of towering blow-up points. The nondegeneracy assumption on ξ_0 , in particular, can be lifted if one chooses h suitably (see e.g. the end of the proof of Theorem 1.1 at the very end of Section 5 where this topic is discussed).

1.2. Sign-changing towers of bubbles for the Brézis-Nirenberg problem. The method we use to prove Theorem 1.1 also applies to the construction of *sign-changing* solutions to the classical Brézis-Nirenberg problem [6]

$$\begin{cases} \Delta_\xi u - \varepsilon u = |u|^{\frac{4}{n-2}} u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1.4)$$

where ξ is the Euclidean metric, $\Delta_\xi = -\sum_{i=1}^n \partial_i^2$, $\Omega \subset \mathbb{R}^n$, $n \geq 3$ is an open and bounded smooth domain. It is well known that (1.4) possesses positive blowing-up solutions, when $\varepsilon > 0$ is small enough and $n \geq 4$, blowing-up at distinct isolated points (see Rey [33] and Musso and Pistoia [28]). In the sign-changing case and when $n \geq 7$, Iacopetti and Vaira [16] built *sign-changing* solutions to problem (1.4) whose shape looks like the superposition of bubbles with alternating sign centered at the origin, provided the domain Ω is symmetric with respect to the origin. This is in sharp contrast with the *positive* case where positive towers of bubbles, at least on symmetric domains, cannot occur for $\varepsilon > 0$ small enough as was proven by Cerqueti [7] using ideas of Li [20, 22]. We also mention Iacopetti and Vaira [17] where sign-changing solutions with asymptotically vanishing negative part have been constructed.

The analysis that we develop in this paper to prove Theorem 1.1 adapts to the Euclidean case of the Brézis-Nirenberg problem. Without assuming any symmetry in the domain we are able to prove the existence of sign-changing towering bubbling configurations:

Theorem 1.2. *Let Ω be a smooth bounded subset of \mathbb{R}^n , $n \geq 7$, and let ξ_0 denote a non-degenerate critical point of the Robin function $\xi \mapsto H(\xi, \xi)$. Let $k \geq 1$. There exists a family $(u_\varepsilon)_\varepsilon$ of sign-changing solutions of*

$$\Delta_\xi u_\varepsilon - \varepsilon u_\varepsilon = |u_\varepsilon|^{\frac{4}{n-2}} u_\varepsilon \quad (1.5)$$

that blows-up as the superposition of k bubbles with alternating sign at ξ_0 .

The existence of radial towers of bubbles centered at the origin on the ball had been previously obtained by Esposito-Ghoussoub-Pistoia-Vaira [11] in the more general context of Hardy-Schrödinger equations. Theorem 1.2 is again a generalisation of this construction to arbitrary non-symmetric domains. For the precise definition of the Robin function we refer to (6.4). As for Theorem 1.1 the assumptions on ξ_0 can be relaxed if a general h is taken into account in the operator $\Delta_\xi - \varepsilon h$, see the arguments at the end of Section 6. The existence of sign-changing towers of bubbles for problem (1.4) is likely to be true only in dimensions $n \geq 7$, and a proof that sign-changing towers of bubbles made of two bubbles cannot appear in dimensions 3 to 6 can be found in Ben Ayed-El Medhi-Pacella [2, 3] and Iacopetti and Pacella [15]. Let us also mention

an asymptotic description of low-energy sign-changing solutions in dimensions $n \geq 7$ in Iacopetti [14].

1.3. Strategy of proof and outline of the paper. The proofs of Theorems 1.1 and 1.2 follow the strategy that was introduced in Premoselli [30, 31] and applied in Premoselli-Thizy [32]. The blowing-up family $(u_\varepsilon)_\varepsilon$ that we construct in Theorem 1.1 has the following form:

$$u_\varepsilon = \sum_{i=1}^k W_{i,\varepsilon} + \varphi_\varepsilon,$$

where $W_{i,\varepsilon}$ are, at first-order, Riemannian standard bubbles of respective parameters $\mu_{k,\varepsilon} \ll \mu_{k-1,\varepsilon} \ll \dots \ll \mu_{1,\varepsilon}$ and centered at points $\xi_{i,\varepsilon}$ and φ_ε is a remainder. The points ξ_ℓ satisfy $d_g(\xi_{\ell-1}, \xi_\ell) = O(\mu_{\ell-1})$ for $2 \leq \ell \leq k$, they all converge to the same point $\xi_0 \in M$, and the parameters are related according to (2.8) below. The main idea of the present paper is to construct a remainder φ_ε satisfying

$$\left\| \frac{\varphi_\varepsilon}{\sum_{i=1}^k W_{i,\varepsilon}} \right\|_{L^\infty(M)} = o(1) \quad (1.6)$$

(more precise estimates are actually required, see (4.5)). This contrasts with the approach followed in Morabito-Pistoia-Vaira [26]: in the highly symmetric setting of [26] all the bubbles were centered in the center of symmetry of the manifold and a suitable φ_ε was constructed in H^1 by splitting it in k distinct remainders with a triangular dependence on the bubble parameters. Since we work here on a general non-symmetric background, where the centers of the bubbles also have to be chosen, the energy estimates of [26] fail to capture precisely enough the interactions between the bubbles $W_{i,\varepsilon}$. We thus have to construct a remainder φ_ε which comes with sharp global pointwise estimates such as (1.6).

Constructing φ_ε that satisfies (1.6) is the main challenge of this paper. It requires to prove sharp pointwise estimates for solutions of a linear equation (see Proposition 3.2 below). These estimates are obtained by means of a bubble-tree-like analysis, which is carried on at a significant precision. Once this linear theory is available, φ_ε is constructed *via* a standard fixed-point procedure, which now takes place in strong weighted spaces. This construction of φ_ε applies both to positive and sign-changing solutions. The sharp pointwise estimates that we obtain on φ_ε , combined with the finite-dimensional reduction, also provide another way to obtain *a priori* pointwise blow-up estimates for finite-energy solutions of (1.3), similar to the ones in Druet-Hebey-Robert [10] and to the symmetry estimates of Li-Zhang [21] and Khuri-Marques-Schoen [18] (see Section 4).

The outline of the paper is as follows. In Section 2 we introduce the notations and the bubbling profiles used in the construction. In Section 3 we prove sharp pointwise estimates for solutions of the linearization of (1.3). This is achieved in Proposition 3.2 and is the core of the analysis of the paper. In section 4 we perform the nonlinear construction to solve (1.1) up to kernel elements. In section 5 we perform an asymptotic expansion of the coefficients of the kernel elements and conclude the proof of Theorem 1.1. Theorem 1.2 is then proven in 6. Finally, the Appendix gathers a few technical results used throughout the paper. Although Theorem 1.1 is proven in the locally conformally flat case, we prove the results in sections 3 and 4 in the general case, since the analysis carries on and might be applied in future works. We also take into account a general h as in (1.3) since it does not create additional difficulties.

Acknowledgments: The author warmly thanks A. Pistoia for many fruitful and motivating discussions during the preparation of this work, and in particular for pointing out the possible application to the Brézis-Nirenberg problem, which led to Theorem 1.2. The author was supported by a FNRS CdR grant J.0135.19 and by the Fonds Thélam.

2. NOTATIONS AND APPROXIMATE SOLUTIONS

2.1. The ansatz. Let (M, g) be a compact Riemannian manifold without boundary of dimension $n \geq 3$. The celebrated conformal normal coordinates result of Lee-Parker [19] states the existence of a smooth function $\Lambda : M \times M \rightarrow (0, +\infty)$ such that for any $\xi \in M$, the conformal metric $g_\xi = \Lambda_\xi^{\frac{4}{n-2}} g$ satisfies:

$$S_{g_\xi}(\xi) = 0, \quad \nabla S_{g_\xi}(\xi) = 0 \quad \text{and} \quad \Delta_{g_\xi} S_{g_\xi}(\xi) = \frac{1}{6} |W_g(\xi)|_g^2,$$

where we have let $\Lambda_\xi = \Lambda(\xi, \cdot)$ and where S_h and W_h denote respectively the scalar curvature and the Weyl tensor of a metric h . If (M, g) is locally conformally flat in a neighbourhood of some $\xi \in M$, g_ξ is flat in a geodesic ball centered at ξ . If h is any metric of positive Yamabe type on M we denote by G_h the Green function of the conformal laplacian $\Delta_h + c_n S_h$. Let $\xi \in M$ and consider the metric $g_\xi = \Lambda_\xi^{\frac{4}{n-2}} g$. If (M, g) is locally conformally flat in a neighbourhood of ξ the Green function G_{g_ξ} expands as

$$G_{g_\xi}(\xi, \exp_\xi^{g_\xi}(y)) = \frac{1}{(n-2)\omega_{n-1}} |y|^{2-n} + m(\xi) + O(|y|), \quad (2.1)$$

for $|y|$ small, where we denoted by $\exp_\xi^{g_\xi}$ the exponential map at ξ for g_ξ . The quantity $m(\xi)$, which is a smooth function of ξ , is called the *mass* at ξ . If (M, g) is not conformally flat in a neighbourhood of $\xi \in M$ we have instead,

when $n \geq 7$:

$$G_{g_\xi}(\xi, \exp_\xi^{g_\xi}(y)) = \frac{1}{(n-2)\omega_{n-1}}|y|^{2-n} + O(|y|^{6-n}). \quad (2.2)$$

We refer to Lee-Parker [19] for a proof of these facts.

The bubbling profiles we will use in this work are inspired from the profiles in Esposito-Pistoia-Vétois [12], with a suitable modification to smoothen them. Let $0 < r_0 < \frac{1}{2}i_g(M)$ be such that

$$2r_0 < \frac{1}{2}i_{g_\xi}(M) \quad \text{for any } \xi \in M \quad (2.3)$$

and let $\chi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a smooth compactly supported function satisfying $\chi(r) = r$ for $r \leq r_0$ and $\chi(r) = 2r_0$ for $r \geq 2r_0$. If $\xi \in M$ and $\mu > 0$, we define the bubble $W_{\mu,\xi}$ as:

$$W_{\mu,\xi}(x) = \Phi(\xi, x) \frac{\mu^{\frac{n-2}{2}}}{\left(\mu^2 + \frac{\chi(d_{g_\xi}(\xi, x)^2)}{n(n-2)}\right)^{\frac{n-2}{2}}}, \quad (2.4)$$

where d_{g_ξ} is the geodesic distance for the metric g_ξ and where we have let, for all $\xi, x \in M$:

$$\Phi(\xi, x) = (n-2)\omega_{n-1}G_{g_\xi}(\xi, x)\Lambda_\xi(x)\chi(d_{g_\xi}(\xi, x))^{n-2}. \quad (2.5)$$

This ansatz satisfies in particular:

$$W_{\mu,\xi}(x) = \Phi(\xi, x)\mu^{\frac{n-2}{2}}\left(\mu^2 + \frac{d_{g_\xi}(\xi, x)^2}{n(n-2)}\right)^{1-\frac{n}{2}} \quad \text{if } d_{g_\xi}(\xi, x) \leq r_0.$$

As shown in [12], $W_{\mu,\xi}$ satisfies:

$$\Delta_g W_{\mu,\xi} + c_n S_g W_{\mu,\xi} - W_{\mu,\xi}^{2^*-1} = \begin{cases} \frac{O(\mu^{\frac{n+2}{2}})}{\left(\mu^2 + \frac{d_{g_\xi}(\xi, x)^2}{n(n-2)}\right)^2} & (M, g) \text{ is l.c.f} \\ \frac{O(\mu^{\frac{n+2}{2}})}{\left(\mu^2 + \frac{d_{g_\xi}(\xi, x)^2}{n(n-2)}\right)^{\frac{n-2}{2}}} & \text{otherwise,} \end{cases} \quad (2.6)$$

where “l.c.f” means here *locally conformally flat in a neighbourhood of ξ* . We will use the same abbreviation throughout the paper, and there should be no confusion since our analysis will take place in a neighbourhood of a point $\xi_1 \in M$. In (2.6) we also used the following notation:

$$A = O(B) \text{ or, equivalently, } A \lesssim B$$

if there exists a constant $C > 0$ independent of ε such that $|A| \leq C|B|$. Similarly, we will write $A = o(B)$ if $|A||B|^{-1}$ goes to 0 as $\varepsilon \rightarrow 0$.

2.2. The parameters. We introduce the parameters of our construction. Let $A > 1$ be a large positive constant, $k \geq 1$ be an integer. Let $(t_i)_{1 \leq i \leq k} \in [A^{-1}, A]^k$ and $\varepsilon > 0$. For any $1 \leq \ell \leq k$ define:

$$\mu_\ell = \mu_{\ell, \varepsilon}(t_\ell) = \varepsilon^{\gamma_\ell} t_\ell,$$

where

$$\gamma_\ell = \begin{cases} \frac{n-2}{2(n-4)} \left(\frac{n-2}{n-6} \right)^{\ell-1} - \frac{1}{2} & (M, g) \text{ is l.c.f} \\ \left(\frac{n-2}{n-6} \right)^{\ell-1} - \frac{1}{2} & \text{otherwise} \end{cases} \quad (2.7)$$

With this definition it is easily seen that we have

$$\varepsilon^{1+2\gamma_\ell} = \varepsilon^{\frac{n-2}{2}(\gamma_\ell - \gamma_{\ell-1})}.$$

In particular, for fixed values of $(t_i)_{1 \leq i \leq k} \in [A^{-1}, A]^k$, the μ_ℓ satisfy:

$$\mu_1 \sim \begin{cases} \varepsilon^{\frac{1}{n-4}} & (M, g) \text{ l.c.f} \\ \varepsilon^{\frac{1}{2}} & \text{otherwise} \end{cases} \quad \text{and} \quad \varepsilon \mu_{\ell+1}^2 \sim \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} \quad (2.8)$$

for any $1 \leq \ell \leq k-1$, where \sim denotes the order of growth in ε as $\varepsilon \rightarrow 0$.

Denote by $B(0, 1)$ the open unit ball in \mathbb{R}^n and define

$$S_k = [A^{-1}, A] \times M \times \left([A^{-1}, A] \times B(0, 1) \right)^{k-1}. \quad (2.9)$$

Let $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$. For any $\varepsilon > 0$, we define the points (ξ_2, \dots, ξ_k) by:

$$\text{for any } 2 \leq \ell \leq k, \quad \xi_\ell = \exp_{\xi_{\ell-1}}^{g_{\xi_{\ell-1}}}(\mu_{\ell-1} z_\ell). \quad (2.10)$$

Recall that for $\xi \in M$, $\exp_\xi^{g_\xi}$ is the exponential map at ξ for the metric $g_\xi = \Lambda_\xi^{\frac{4}{n-2}}$. In (2.10) we omitted the dependence on the parameters for the sake of clarity but keep in mind that we have:

$$\xi_2 = \xi_{2, \varepsilon}(t_1, z_2) \quad , \quad \dots \quad , \quad \xi_\ell = \xi_{\ell, \varepsilon}(t_{\ell-1}, z_\ell).$$

For $1 \leq \ell \leq k$ we will let

$$W_\ell(x) = W_{\mu_\ell, \xi_\ell}(x) \quad (2.11)$$

where W_{μ_ℓ, ξ_ℓ} is as in (2.4). By definition W_ℓ only depends on the parameters $(t_{\ell-1}, t_\ell, z_\ell)$ when $\ell \geq 2$, and only on (t_1, ξ_1) when $\ell = 1$. In this setting W_k denotes the highest bubble (the one that concentrates faster) while W_1 is the lowest one. In the course of the proof we will need to analyse the mutual pointwise interactions between these bubbles. Following the notion of radius of influence introduced in Druet [8] (see also Hebey [13], Chapter 8) We define for this the following sets:

$$B_1 = M, B_{k+1} = \emptyset \quad \text{and} \quad B_\ell = B_{g_{\xi_\ell}}(\xi_\ell, \sqrt{\mu_\ell \mu_{\ell-1}}) \text{ for } 2 \leq \ell \leq k, \quad (2.12)$$

where $B_{g_{\xi_\ell}}$ denotes the geodesic ball for the metric g_{ξ_ℓ} . For any family $(t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{i,\varepsilon}, z_{i,\varepsilon})_{2 \leq i \leq k})$ of elements of S_k it is easily checked by (2.10) that, for ε small enough, the balls B_ℓ are nested:

$$B_k \subset B_{k-1} \subset \cdots \subset B_1.$$

It is also easily checked that, for any $1 \leq \ell \leq k$,

$$W_\ell \lesssim W_{\ell-1} \text{ in } B_{\ell-1} \setminus B_\ell \text{ and } W_{\ell-1} \lesssim W_\ell \text{ in } B_\ell. \quad (2.13)$$

In particular, for any $1 \leq \ell \leq k$ there holds

$$\sum_{j=1}^k W_j \lesssim W_\ell \quad \text{on } B_\ell \setminus B_{\ell+1}.$$

2.3. Kernel elements. Let $h \in C^0(M)$ and $\varepsilon > 0$. We will work with the following H^1 scalar product in M :

$$\langle u, v \rangle = \int_M \left(\langle \nabla u, \nabla v \rangle + (c_n S_g + \varepsilon h) uv \right) dv_g. \quad (2.14)$$

Let $1 \leq \ell \leq k$, $0 \leq j \leq n$ and let $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$. The approximate kernel elements associated to the ℓ -th bubble on the manifold are defined by:

$$\begin{aligned} Z_{\ell,0}(x) &= Z_{\ell,0,\mu_\ell,\xi_\ell}(x) \\ &= \Lambda_{\xi_\ell}(x) \chi(d_{g_{\xi_\ell}}(\xi_\ell, x)) \\ &\quad \times \mu_\ell^{\frac{n-2}{2}} \left(\frac{d_{g_{\xi_\ell}}(\xi_\ell, x)^2}{n(n-2)} - \mu_\ell^2 \right) \left(\mu_\ell^2 + \frac{d_{g_{\xi_\ell}}(\xi_\ell, x)^2}{n(n-2)} \right)^{-\frac{n}{2}}, \\ Z_{\ell,j}(x) &= Z_{\ell,j,\mu_\ell,\xi_\ell}(x) \\ &= \Lambda_{\xi_\ell}(x) \chi(d_{g_{\xi_\ell}}(\xi_\ell, x)) \\ &\quad \times \mu_\ell^{\frac{n}{2}} \left\langle \left(\exp_{\xi_\ell}^{g_{\xi_\ell}} \right)^{-1}(x), e_j(\xi_\ell) \right\rangle_{g_{\xi_\ell}(\xi_\ell)} \left(\mu_\ell^2 + \frac{d_{g_{\xi_\ell}}(\xi_\ell, x)^2}{n(n-2)} \right)^{-\frac{n}{2}}. \end{aligned} \quad (2.15)$$

In the previous expression (e_j) is a local orthonormal basis defined in $B_g(\xi_1, 2r_0)$. It is always well-defined and depends smoothly on the choice of the point since we can always choose a covering of parallel-type on M , see [36]. Let also

$$U_0(x) = \left(1 + \frac{|x|^2}{n(n-2)} \right)^{1-\frac{n}{2}} \quad \text{for any } x \in \mathbb{R}^n \quad (2.16)$$

and, for $1 \leq j \leq n$ and $x \in \mathbb{R}^n$,

$$V_0(x) = \frac{\frac{|x|^2}{n(n-2)} - 1}{\left(1 + \frac{|x|^2}{n(n-2)} \right)^{\frac{n}{2}}} \quad \text{and} \quad V_j(x) = x_j \left(1 + \frac{|x|^2}{n(n-2)} \right)^{-\frac{n}{2}}. \quad (2.17)$$

For $1 \leq \ell \leq k$ we will let, for $x \in M$:

$$\theta_\ell(x) = \mu_\ell + d_{g_{\xi_\ell}}(\xi_\ell, x). \quad (2.18)$$

For $1 \leq \ell \leq k$ the functions $Z_{\ell,j}$ in (2.15) satisfy the following equations in M :

$$\Delta_g Z_{\ell,0} + c_n S_g Z_{\ell,0} = (2^* - 1) W_\ell^{2^*-2} Z_{\ell,0} + \begin{cases} O(\mu_\ell^{\frac{n-2}{2}}) & (M, g) \text{ is l.c.f} \\ O(\mu_\ell^2 W_\ell) & \text{otherwise,} \end{cases} \quad (2.19)$$

and, for $1 \leq j \leq n$,

$$\Delta_g Z_{\ell,j} + c_n S_g Z_{\ell,j} = (2^* - 1) W_\ell^{2^*-2} Z_{\ell,j} + \begin{cases} O(\mu_\ell^{\frac{n}{2}}) & \text{if } (M, g) \text{ is l.c.f} \\ O(\mu_\ell W_\ell) & \text{otherwise,} \end{cases} \quad (2.20)$$

Explicit computations yield, for $1 \leq i < j \leq k$ and $0 \leq p, q \leq n$:

$$\langle Z_{i,p}, Z_{i,q} \rangle = \|\nabla V_p\|_{L^2(\mathbb{R}^n)}^2 \delta_{pq} + O(\mu_i^2) \quad (2.21)$$

and

$$\langle Z_{i,p}, Z_{j,q} \rangle = O\left(\left(\frac{\mu_j}{\mu_i}\right)^{\frac{n-2}{2}}\right). \quad (2.22)$$

The proofs of (2.19) and (2.20) can be found in [12] (Lemma 6.3) while we refer to the Appendix of this paper for the proof of (2.21) and (2.22). We finally let, for $1 \leq \ell \leq k$:

$$K_\ell = \text{Span}\{Z_{i,j}, 1 \leq i \leq \ell, 0 \leq j \leq n\}, \quad (2.23)$$

and we denote by K_ℓ^\perp its orthogonal with respect to the scalar product (2.14).

3. SHARP POINTWISE ASYMPTOTICS FOR THE LINEAR PROBLEM

3.1. Statement of the result. Let $k \geq 1$ be an integer and, for any $0 < \varepsilon \leq 1$, let $(t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{i,\varepsilon}, z_{i,\varepsilon})_{2 \leq i \leq k}) \in S_k$ be a family of parameters, where S_k is as in (2.9). For the sake of clarity we will drop the dependence in ε and, until the end of the paper, we will always denote by W_1, \dots, W_k the bubbles obtained as in (2.11) with μ_i and ξ_i given by (2.7) and (2.10) for this choice of the family $(t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{i,\varepsilon}, z_{i,\varepsilon})_{2 \leq i \leq k})$. In what follows the notation $f(u)$ will either denote $f(u) = |u|^{2^*-2}u$ or $f(u) = (u_+)^{2^*-1}$, where $u_+ = \max(u, 0)$. We recall, as a starting point, the well-known linear theory in $H^1(M)$ for (1.3):

Proposition 3.1. *Let $k \geq 1$ be an integer. We fix $(\kappa_i)_{1 \leq i \leq n} \in \{-1, 1\}^k$. There exists $\varepsilon_0 > 0$ such that the following holds: for any $\varepsilon \in (0, \varepsilon_0)$ and $(t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{i,\varepsilon}, z_{i,\varepsilon})_{2 \leq i \leq k}) \in S_k$, $h \in C^0(M)$ and for any $k_\varepsilon \in L^{\frac{2n}{n+2}}(M)$, there exists a unique $\varphi_\varepsilon \in K_k^\perp$ satisfying:*

$$\Pi_{K_k^\perp} \left(\varphi_\varepsilon - (\Delta_g + c_n S_g + \varepsilon h)^{-1} \left[(2^* - 1) f' \left(\sum_{i=1}^k \kappa_i W_i \right) \varphi_\varepsilon + k_\varepsilon \right] \right) = 0. \quad (3.1)$$

Here $\Pi_{K_k^\perp}$ denotes the orthogonal projection on K_k^\perp for (2.14), and K_k is as in (2.23). This function satisfies in addition

$$\|\varphi_\varepsilon\|_{H^1(M)} \lesssim \|k_\varepsilon\|_{L^{\frac{2n}{n+2}}}. \quad (3.2)$$

Proof. This is a by now very classical result. We refer for instance to Robert-Vétois [36], Esposito-Pistoia-Vétois [12] or Musso-Pistoia [27] for a proof. \square

Equation (3.1) implies the existence of real numbers $\lambda_{ij} = \lambda_{ij,\varepsilon}$, with $1 \leq i \leq k$ and $0 \leq j \leq n$, such that

$$\begin{aligned} \Delta_g \varphi_\varepsilon + (c_n S_g + \varepsilon h) \varphi_\varepsilon - f' \left(\sum_{i=1}^k \kappa_i W_i \right) \varphi_\varepsilon &= k_\varepsilon \\ &+ \sum_{\substack{i=1..k \\ j=0..n}} \lambda_{ij} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j}, \end{aligned} \quad (3.3)$$

where $Z_{i,j}$ is as in (2.15). By standard elliptic theory, if $k_\varepsilon \in C^0(M)$, $\varphi_\varepsilon \in C^1(M)$. Let $h \in C^0(M)$ be fixed and assume that $\varepsilon > 0$ is small enough so that $\Delta_g + c_n S_g + \varepsilon h$ is coercive. For any fixed ε and for any $k_\varepsilon \in C^0(M)$ we denote the unique function φ_ε given by Proposition 3.1 by

$$\varphi_\varepsilon := \mathcal{L}_\varepsilon(k_\varepsilon). \quad (3.4)$$

We investigate in this section equation (3.3) in more detail. Assuming that a suitable pointwise control holds on k_ε we prove refined pointwise estimates on φ_ε . The pointwise control we will assume on k_ε is the following:

$$\begin{aligned} |k_\varepsilon| \lesssim & \sum_{i=1}^k \left\{ \begin{array}{ll} \mu_i^{\frac{n+2}{2}} \theta_i^{-4} & \text{If } (M, g) \text{ is l.c.f. around } \xi_1 \\ \varepsilon W_i & \text{If } (M, g) \text{ is not l.c.f.} \end{array} \right\} \\ & + \left\{ \begin{array}{ll} W_k^{2^*-2} W_{k-1} & \text{in } B_k \\ \vdots & \\ W_i^{2^*-2} (W_{i+1} + W_{i-1}) & \text{in } B_i \setminus B_{i+1}, \\ \vdots & \\ W_1^{2^*-2} W_2 & \text{in } B_1 \setminus B_2 = M \setminus B_2 \end{array} \right\}, \end{aligned} \quad (3.5)$$

where in (3.5) the index i in the right-hand side runs from 2 to $k-1$ and where θ_i is defined in (2.18). It is easily seen, using (2.6) and (2.13), that (3.5) is satisfied for the following particular choice of k_ε :

$$k_\varepsilon = (\Delta_g + c_n S_g + \varepsilon h) \left(\sum_{i=1}^k \kappa_i W_i \right) - f \left(\sum_{i=1}^k \kappa_i W_i \right), \quad (3.6)$$

where we choose $f(u) = (u_+)^{2^*-1}$ if all the κ_i are equal to 1 and $f(u) = |u|^{2^*-2}u$ otherwise.

The following proposition is the main result of this section and the core of the analysis of this paper.

Proposition 3.2. *Let $k \geq 1$ be an integer, $(\kappa_i)_{1 \leq i \leq n} \in \{-1, 1\}^k$ and let $\varepsilon_0 > 0$ be as in Proposition 3.1. For any $\varepsilon \in (0, \varepsilon_0)$ let $(t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{i,\varepsilon}, z_{i,\varepsilon})_{2 \leq i \leq k}) \in S_k$, $k_\varepsilon \in C^0(M)$ and let $\varphi_\varepsilon \in K_k^\perp$ be the unique solution of*

$$\begin{aligned} \Delta_g \varphi_\varepsilon + (c_n S_g + \varepsilon h) \varphi_\varepsilon - f' \left(\sum_{i=1}^k \kappa_i W_i \right) \varphi_\varepsilon \\ = k_\varepsilon + \sum_{\substack{i=1..k \\ j=0..n}} \lambda_{ij} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j}, \end{aligned} \quad (3.7)$$

where the λ_{ij} are uniquely defined by (3.1) and $f(u) = |u|^{2^*-2}u$ or $f(u) = (u_+)^{2^*-1}$. Assume that k_ε satisfies (3.5). Then:

- For any $1 \leq \ell \leq k$,

$$\sum_{j=0..n} |\lambda_{\ell j}| \lesssim \varepsilon \mu_\ell^2. \quad (3.8)$$

- For any $2 \leq \ell \leq k$, and any $x \in B_\ell \setminus B_{\ell+1}$:

$$|\varphi_\varepsilon(x)| \lesssim \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbb{1}_{\{\theta_{\ell+1}(x) \leq \mu_\ell\}}, \quad (3.9)$$

where the notation $\mathbb{1}_{\{\theta_{\ell+1}(x) \leq \mu_\ell\}}$ indicates that this term vanishes when $\theta_{\ell+1}(x) > \mu_\ell$, and with the convention that $B_{k+1} = \emptyset$ and $W_{k+1} = 0$.

- For any $x \in B_1 \setminus B_2 = M \setminus B_2$:

$$\begin{aligned} |\varphi_\varepsilon(x)| \lesssim & \left\{ \begin{array}{ll} \frac{\mu_1^{\frac{n+2}{2}}}{\theta_1(x)^2} & \text{if } (M, g) \text{ is l.c.f. around } \xi_1, \\ \frac{\mu_1^{\frac{n+2}{2}}}{\theta_1(x)^{n-4}} & \text{if } (M, g) \text{ is not l.c.f.} \end{array} \right\} \\ & + \frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbb{1}_{\{\theta_2(x) \leq \mu_1\}}. \end{aligned} \quad (3.10)$$

The constants appearing in (3.8), (3.9), (3.10) are independent of ε and of the choices of $(t_1, \xi_1, (t_{i,\varepsilon}, z_{i,\varepsilon})_{2 \leq i \leq k}) \in S_k$, U_1, \dots, U_k and k_ε .

In Proposition 3.2, B_ℓ is as in (2.12) and θ_ℓ as in (2.18).

Let us comment on this result. Proposition 3.2 yields a pointwise control on φ_ε and on the λ_{ij} assuming only (3.5): no previous control on φ_ε or on the λ_{ij} was assumed. The estimates (3.9) and (3.10) have a deeper meaning. Let

indeed G_ε be the Green's function of $\Delta_g + c_n S_g + \varepsilon h$ in M . Assuming that k_ε satisfies (3.5), straightforward computations using (A.1) and (A.2) give:

$$\int_M G_\varepsilon(x, \cdot) |k_\varepsilon| dv_g \lesssim \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbb{1}_{\{\theta_{\ell+1}(x) \leq \mu_\ell\}}$$

for $x \in B_\ell \setminus B_{\ell+1}$, $2 \leq \ell \leq k$, and

$$\int_M G_\varepsilon(x, \cdot) |k_\varepsilon| dv_g \lesssim \begin{cases} \frac{\mu_1^{\frac{n+2}{2}}}{\theta_1(x)^2} & \text{if } (M, g) \text{ is l.c.f. around } \xi_1, \\ \frac{\mu_1^{\frac{n+2}{2}}}{\theta_1(x)^{n-4}} & \text{if } (M, g) \text{ is not l.c.f.} \end{cases} + \frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbb{1}_{\{\theta_2(x) \leq \mu_1\}}.$$

if $x \in B_1 \setminus B_2 = M \setminus B_2$. Compare the last two estimates with (3.9) and (3.10): Proposition 3.2 therefore shows that a solution $\varphi_\varepsilon \in K_k^\perp$ of (3.7) behaves *as if it satisfied the much simpler equation*

$$\Delta_g \varphi_\varepsilon + (c_n S_g + \varepsilon h) \varphi_\varepsilon = k_\varepsilon + \sum_{\substack{i=1 \dots k \\ j=0 \dots n}} \lambda_{ij} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j}, \quad (3.11)$$

with the λ_{ij} satisfying (3.8). Proposition 3.2 thus shows that the term $f'(\sum_{i=1}^k \kappa_i U_i) \varphi_\varepsilon$ has no incidence on the behavior of φ_ε , provided $\varphi_\varepsilon \in K_k^\perp$: it therefore has to be seen as a considerably more involved analogue of Proposition 3.1, one which yields sharp pointwise estimates on φ_ε . The usefulness of Proposition 3.2 in the proof of Theorems 1.1 and 1.2 becomes clear when anticipating on Section 4 below. In Section 4 we will perform a nonlinear fixed-point argument to solve (1.3) up to kernel elements. Thanks to Proposition 3.2 we will be able to perform the point-fixing argument in suitably weighted spaces for the C^0 norm. An accurate choice of the weight will reduce the point-fixing to an application of Proposition 3.2 with an error term given by (3.6) and by a negligible (and controlled) nonlinear correction.

The proof of Proposition 3.2 goes through an a priori bubble-tree analysis. We iteratively obtain refined pointwise estimates on $|\varphi_\varepsilon|$ on each annular region $B_i \setminus B_{i+1}$ starting from B_k (which corresponds to the highest bubble). These estimates depend on the pointwise behaviour of φ_ε on all of M and on the λ_{ij} and are heavily intertwined. We then propagate these estimates until $M \setminus B_1$, we perform a second-order one-bubble analysis on the lowest bubble in $M \setminus B_1$ to prove (3.10), and we propagate it back to the highest bubble to obtain (3.9). The λ_{ij} themselves are iteratively estimated in the course of the proof.

3.2. A representation formula. We start by proving a short lemma that will be crucial in the proof of Proposition 3.2. It is a representation formula for the linearised operator at the standard bubble in \mathbb{R}^n :

Lemma 3.3. *Let $\varphi \in C_c^\infty(\mathbb{R}^n)$ and let, for any $x \in \mathbb{R}^n$,*

$$\Pi(\varphi)(x) = \sum_{j=0}^n \left(\frac{\int_{\mathbb{R}^n} \varphi V_j U_0^{2^*-2} dy}{\int_{\mathbb{R}^n} V_j^2 U_0^{2^*-2} dy} \right) V_j(x)$$

be the orthogonal projection of φ on $\text{Span}\{V_j, 0 \leq j \leq n\}$ for the $D^{1,2}$ scalar product in \mathbb{R}^n . Then, for any $x \in \mathbb{R}^n$:

$$|\varphi(x) - \Pi(\varphi)(x)| \lesssim \int_{\mathbb{R}^n} |x - y|^{2-n} |\Delta_\xi \varphi - (2^* - 1)U_0^{2^*-2}\varphi|(y) dy \quad (3.12)$$

where V_j and U_0 are as in (2.16) and (2.17).

Proof. Let g_0 be the round metric on the sphere. Let $N \in \mathbb{S}^n$ and let $\pi_N : \mathbb{S}^n \setminus \{-N\} \rightarrow \mathbb{R}^n$ the stereographic projection sending N to 0. There holds

$$(\pi_N^{-1})^* g_0 = U^{\frac{4}{n-2}} \xi,$$

where ξ is the Euclidean metric and $U(x)^{\frac{4}{n-2}} = \frac{4}{(1+|x|^2)^2}$. As a consequence, $dv_{g_0} = U^{\frac{2n}{n-2}} dx$. For $\varphi \in C_c^\infty(\mathbb{R}^n)$ and $x \in \mathbb{S}^n$ we let:

$$\tilde{\varphi}(x) = \frac{\varphi(\pi_N(x))}{U(\pi_N(x))}.$$

This defines a smooth function $\tilde{\varphi}$ on \mathbb{S}^n with compact support in $\mathbb{S}^n \setminus \{-N\}$, and there holds for any $\varphi, \psi \in C_c^\infty(\mathbb{R}^n)$,

$$\int_{\mathbb{S}^n} \tilde{\varphi} \tilde{\psi} dv_{g_0} = \int_{\mathbb{R}^n} \varphi \psi U^{2^*-2} dx. \quad (3.13)$$

By the conformal invariance of the conformal laplacian we have furthermore, for $x \in \mathbb{S}^n$:

$$\left(\Delta_{g_0} + \frac{n(n-2)}{4} \right) \tilde{\varphi}(x) = U(\pi_N(x))^{1-2^*} \Delta_\xi \varphi(\pi_N(x)),$$

so that for any $\varphi, \psi \in C_c^\infty(\mathbb{R}^n)$;

$$\int_{\mathbb{S}^n} \left(\langle \nabla \tilde{\varphi}, \nabla \tilde{\psi} \rangle + \frac{n(n-2)}{4} \tilde{\varphi} \tilde{\psi} \right) dv_{g_0} = \int_{\mathbb{R}^n} \langle \nabla \varphi, \nabla \psi \rangle dx$$

and

$$(\Delta_{g_0} - n) \tilde{\varphi}(x) = U(\pi_N(x))^{1-2^*} \left(\Delta_\xi \varphi(\pi_N(x)) - \frac{n(n+2)}{(1+|x|^2)^2} \varphi(\pi_N(x)) \right) \quad (3.14)$$

for any $x \in \mathbb{S}^n$. Let

$$K = \{f \in H^1(\mathbb{S}^n) \text{ such that } \Delta_{g_0} f - n f = 0\}.$$

An orthonormal basis of K , for the L^2 scalar product (3.13) on (\mathbb{S}^n, g_0) , is given by the family (K_0, \dots, K_n) where

$$K_i = c_i^{-1} x_i, \quad c_i^2 = \int_{\mathbb{S}^n} x_i^2 dv_{g_0}, \quad 0 \leq i \leq n,$$

and where the x_i are the coordinate functions restricted to \mathbb{S}^n . Let $G(x, y)$ denote the Green's function (see e.g. [34]) for the operator $\Delta_{g_0} - n$ on \mathbb{S}^n . Standard properties of the Green's function ensure that

$$|G(x, y)| \leq C d_{g_0}(x, y)^{2-n} \text{ for } x \neq y$$

and that, for any smooth function ψ in \mathbb{S}^n and any $z \in \mathbb{S}^n$,

$$\psi(z) - \Pi_K(\psi)(z) = \int_{\mathbb{S}^n} G(z, y) (\Delta_{g_0} \psi(y) - n\psi(y)) dv_{g_0}(y),$$

where $\Pi_K(\psi) = \sum_{j=0}^n (\int_{\mathbb{S}^n} \psi K_j dv_{g_0}) K_j$ is the L^2 projection of ψ over K . Let now $x \in \mathbb{R}^n$, and apply the latter to $\psi = \tilde{\varphi}$ and $z = \pi_N^{-1}(x)$. Letting $\tilde{K}_i(x) = U(x) K_i(\pi_N^{-1}(x))$ we thus get, by (3.13) and (3.14):

$$\varphi(x) - \Pi(\varphi)(x) = \int_{\mathbb{R}^n} \tilde{G}(x, y) \left(\Delta_\xi \varphi(y) - \frac{n(n+2)}{(1+|y|^2)^2} \varphi(y) \right) dy, \quad (3.15)$$

where we have let

$$\Pi(\varphi)(x) = \sum_{j=0}^n \left(\int_{\mathbb{R}^n} \varphi \tilde{K}_j U^{2^*-2} dx \right) \tilde{K}_j(x)$$

and, for any $x \neq y$ in \mathbb{R}^n ,

$$\tilde{G}(x, y) = U(x) G(\pi_N^{-1}(x), \pi_N^{-1}(y)) U(y).$$

Standard properties of the stereographic projection yield

$$d_{g_0}(\pi_N^{-1}(x), \pi_N^{-1}(y)) \gtrsim \frac{|x - y|}{\sqrt{(1+|x|^2)(1+|y|^2)}}$$

so that

$$|\tilde{G}(x, y)| \lesssim |x - y|^{2-n} \quad \text{for any } x \neq y \text{ in } \mathbb{R}^n.$$

Finally, if $\varphi \in C_c^\infty(\mathbb{R}^n)$, letting $z = \frac{x}{\sqrt{n(n-2)}}$ and $\hat{\varphi}(x) = \varphi(z)$ gives

$$\Delta_\xi \hat{\varphi}(x) - (2^* - 1) U_0^{2^*-2}(x) \hat{\varphi}(x) = \frac{1}{n(n-2)} \left(\Delta_\xi \varphi(z) - \frac{n(n+2)}{(1+|z|^2)^2} \varphi(z) \right).$$

It remains to apply (3.15) at $\frac{x}{\sqrt{n(n-2)}}$ and to perform the scaling $y \mapsto \frac{y}{\sqrt{n(n-2)}}$ in the integral. The proof is concluded by remarking that in the chart π_N the coordinate functions (x_0, \dots, x_n) on \mathbb{S}^n write respectively as

$$\frac{1 - |x|^2}{1 + |x|^2}, \frac{2x_1}{1 + |x|^2}, \dots, \frac{2x_n}{1 + |x|^2},$$

which gives in turn

$$\tilde{K}_0\left(\frac{x}{\sqrt{n(n-2)}}\right) = c_0^{-1} 2^{\frac{n-2}{2}} V_0(x) \text{ and } \tilde{K}_i\left(\frac{x}{\sqrt{n(n-2)}}\right) = c_i^{-1} \frac{2^{\frac{n}{2}}}{n(n-2)} V_i(x)$$

for all $1 \leq i \leq n$, where the V_i are given by (2.17). \square

3.3. Proof of Proposition 3.2, Part 1: a recursive estimate. Let $k \geq 1$ be an integer, $\{\kappa_i\}_{1 \leq i \leq k} \in \{\pm 1\}^k$ and $\varepsilon_0 > 0$ be as in Proposition 3.1. For any $\varepsilon \in (0, \varepsilon_0)$ let $(t_1, \xi_1, (t_{i,\varepsilon}, z_{i,\varepsilon})_{2 \leq i \leq k}) \in S_k$, $h \in C^0(M)$, $k_\varepsilon \in C^0(M)$ satisfying (3.5), and let $\varphi_\varepsilon \in K_k^\perp$ be the unique solution of (3.7), where the λ_{ij} are uniquely defined by (3.1). By (3.2) and (3.5) we easily get

$$\|\varphi_\varepsilon\|_{H^1} = o(1)$$

as $\varepsilon \rightarrow 0$, and hence integrating (3.7) against $Z_{i,j}$ for all $1 \leq i \leq k$ and $0 \leq j \leq n$ yields

$$|\lambda_{ij}| = o(1) \quad \text{for } 1 \leq i \leq k, \quad 0 \leq j \leq n. \quad (3.16)$$

For any $2 \leq i \leq k$, we define

$$C_i = B_{g_{\xi_i}}(\xi_i, 4\sqrt{\mu_i \mu_{i-1}}) \setminus B_{g_{\xi_{i+1}}}(\xi_{i+1}, \frac{1}{4}\sqrt{\mu_{i+1} \mu_i}), \quad (3.17)$$

and

$$C_1 = B_{g_{\xi_1}}(\xi_1, 2r_0) \setminus B_{g_{\xi_2}}(\xi_2, \frac{1}{4}\sqrt{\mu_2 \mu_1}), \quad (3.18)$$

where r_0 is as in (2.3). We also define

$$\begin{aligned} a_i &= a_{i,\varepsilon} := \left\| \frac{\phi_\varepsilon}{W_i} \right\|_{L^\infty(C_i)} \quad \text{for } 2 \leq i \leq k, \\ a_1 &= a_{1,\varepsilon} := \left\| \frac{\phi_\varepsilon}{W_1} \right\|_{L^\infty(M \setminus B_{g_{\xi_2}}(\xi_2, \frac{1}{4}\sqrt{\mu_2 \mu_1}))}, \end{aligned} \quad (3.19)$$

and, for a given $1 \leq i \leq k$,

$$\Lambda_i := \sum_{j=0..n} |\lambda_{ij}|. \quad (3.20)$$

By definition of a_i , φ_ε satisfies $|\varphi_\varepsilon| \leq a_i W_i$ in C_i . The proof of Proposition 3.2 consists in an explicit estimation of the a_i .

For a fixed $1 \leq \ell \leq k$, we will say that the property (H_ℓ) is satisfied if the following two properties $(H1_\ell)$ and $(H2_\ell)$ are both satisfied:

Property $(H1_\ell)$: for any $p \in \{\ell, \dots, k\}$ and for any $x \in C_p$,

$$\begin{aligned}
|\varphi_\varepsilon(x)| &\lesssim \varepsilon \mu_p^{3-\frac{n}{2}} \sum_{j \geq p} a_j + \sum_{j \leq p-1} \mu_j^{1-\frac{n}{2}} \Lambda_j + \mu_{p-1}^{1-\frac{n}{2}} \left(\frac{\mu_p}{\theta_p(x)} \right)^2 \\
&+ \sum_{j \geq p+1} \mu_{j-1}^2 a_j W_j(x) + \varepsilon \left(1 + \sum_{j \leq p} a_j \right) \theta_p^2(x) W_p(x) \\
&+ (1 + a_p) \frac{\theta_{p+1}(x)^2}{\mu_p^2} W_{p+1}(x) \mathbb{1}_{\{\theta_{p+1}(x) \leq \mu_p\}}
\end{aligned} \tag{H1_\ell}$$

Property $(H2_\ell)$: for any $p \in \{\ell, \dots, k\}$,

$$\Lambda_p \lesssim \varepsilon \mu_p^2 \left(1 + \sum_{i=1}^k a_i \right). \tag{H2_\ell}$$

A word on the notations is in order here. In $(H1_\ell)$ and $(H2_\ell)$ we take the following conventions: $W_{k+1} = 0$ and $\mu_0 = \frac{1}{\mu_1}$. In the rest of the proof of Proposition 3.2 both the expressions $\varepsilon \mu_\ell^2$ and $\left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}}$ shall appear throughout the computations. When $\ell \geq 2$, of course, they are of the same order in ε by (2.8). They might differ when $\ell = 1$, however, according to whether (M, g) is locally conformally flat around ξ_1 or not. It will be understood, by the choice of our convention $\mu_0 = \frac{1}{\mu_1}$, that the term $\left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}}$ becomes μ_1^{n-2} when $\ell = 1$.

The core of the proof of Proposition 3.2 consists in proving recursively that (H_ℓ) is true for any $1 \leq \ell \leq k$.

Lemma 3.4. *For any $1 \leq \ell \leq k$, (H_ℓ) is true.*

Proof. We prove Lemma 3.4 by induction. Since the proof of the initialisation and the induction part are very close we prove them together. Let $1 \leq \ell \leq k$ be an integer. **If $\ell = k$, we do not assume anything. If $\ell \leq k-1$, we assume that $(H_{\ell+1})$ holds true.** The proof of (H_ℓ) goes through a series of steps.

Step 1: a local rescaling. We let $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a smooth nonnegative function with $\eta \equiv 1$ in $[0, \frac{3}{2}]$ and $\eta \equiv 0$ in $\mathbb{R} \setminus [0, 2]$. We let in the following:

$$\hat{B}_\ell = B_0 \left(2 \sqrt{\frac{\mu_{\ell-1}}{\mu_\ell}} \right) \text{ if } \ell \geq 2, \quad \hat{B}_1 = B_0 \left(\frac{2r_0}{\mu_1} \right)$$

where r_0 is as in (2.3). We also let

$$\hat{B}_{\ell+1} = B \left(z_{\ell+1}, \sqrt{\frac{\mu_{\ell+1}}{\mu_\ell}} \right).$$

For any $x \in \hat{B}_\ell$, we let

$$\hat{\varphi}_\varepsilon(x) = \mu_\ell^{\frac{n-2}{2}} (\Lambda_{\xi_\ell}^{-1} \varphi_\varepsilon) (\exp_{\xi_\ell}^{g_{\xi_\ell}}(\mu_\ell x)) \cdot \eta\left(\sqrt{\frac{\mu_\ell}{\mu_{\ell-1}}} x\right). \quad (3.21)$$

In the following we will let $x_\ell = \exp_{\xi_\ell}^{g_{\xi_\ell}}(\mu_\ell x)$ if $x \in \hat{B}_\ell$ and we will define

$$\hat{g}_\ell(x) = (\exp_{\xi_\ell}^{g_{\xi_\ell}})^* g(\mu_\ell x).$$

Let also, for $x \in \hat{B}_\ell$ and $1 \leq i \leq k$, $0 \leq j \leq n$:

$$\hat{W}_i(x) = \mu_\ell^{\frac{n-2}{2}} (\Lambda_{\xi_\ell}^{-1} W_i)(x_\ell) \quad \text{and} \quad \hat{Z}_{i,j}(x) = \mu_\ell^{\frac{n-2}{2}} (\Lambda_{\xi_\ell}^{-1} Z_{i,j})(x_\ell).$$

By the conformal invariance property of the conformal laplacian and (3.7), the function $\hat{\varphi}_\varepsilon$ satisfies the following equation: if $x \in \hat{B}_\ell$,

$$\begin{aligned} & \left(\Delta_{\hat{g}_\ell} + \mu_\ell^2 c_n S_{g_{\xi_\ell}}(x_\ell) + \mu_\ell^2 \varepsilon h(x_\ell) \Lambda_{\xi_\ell}(x_\ell)^{2-2^*} \right) \left(\hat{\varphi}_\varepsilon - \sum_{\substack{i=1..k \\ j=0..n}} \lambda_{ij} \hat{Z}_{i,j} \right) \\ &= \mu_\ell^{\frac{n+2}{2}} \Lambda_{\xi_\ell}(x_\ell)^{1-2^*} k_\varepsilon(x_\ell) + \mu_\ell^2 \Lambda_{\xi_\ell}(x_\ell)^{2-2^*} f' \left(\sum_{i=1}^k \kappa_i W_i \right)(x_\ell) \hat{\varphi}_\varepsilon \\ &+ O \left(\frac{\mu_\ell}{\mu_{\ell-1}} |\hat{\varphi}_\varepsilon| + \sqrt{\frac{\mu_\ell}{\mu_{\ell-1}}} |\nabla \hat{\varphi}_\varepsilon| \right) \cdot \mathbb{1}_{\frac{3}{2} \sqrt{\frac{\mu_{\ell-1}}{\mu_\ell}} \leq |x| \leq 2 \sqrt{\frac{\mu_{\ell-1}}{\mu_\ell}}} \end{aligned} \quad (3.22)$$

If (M, g) is locally conformally flat around ξ_1 the metric $\Lambda_{\xi_\ell}^{\frac{4}{n-2}} g$ is flat, and then

$$(\Delta_{\hat{g}_\ell} + \mu_\ell^2 c_n S_{g_{\xi_\ell}}(x_\ell)) \hat{\varphi}_\varepsilon(x) = \Delta_\xi \hat{\varphi}_\varepsilon(x)$$

in \tilde{B}_ℓ , where ξ is the Euclidean metric. Otherwise, fine properties of the conformal normal coordinates (see e.g. [12], Lemma 6.3) give

$$\begin{aligned} (\Delta_{\hat{g}_\ell} + \mu_\ell^2 c_n S_{g_{\xi_\ell}}(x_\ell)) \hat{\varphi}_\varepsilon(x) &= \Delta_\xi \hat{\varphi}_\varepsilon(x) + O \left(\mu_\ell^2 |x|^2 |\nabla^2 \hat{\varphi}_\varepsilon(x)| \right) + O \left(\mu_\ell^3 |x|^2 |\nabla \hat{\varphi}_\varepsilon(x)| \right) \\ &+ O \left(\mu_\ell^4 |x|^2 |\hat{\varphi}_\varepsilon| \right). \end{aligned}$$

By (2.13) we have

$$\left| f' \left(\sum_{i=1}^k \kappa_i W_i \right)(x_\ell) - (2^* - 1) W_\ell(x_\ell)^{2^*-2} \right| \lesssim W_\ell^{2^*-3} (W_{\ell+1} + W_{\ell-1}) \quad (3.23)$$

in $B_\ell \setminus B_{\ell+1}$, where the term $W_{\ell-1}$ is assumed to vanish when $\ell = 1$, which yields after scaling and by (2.8):

$$\mu_\ell^2 \Lambda_{\xi_\ell}(x_\ell)^{2-2^*} \left| f' \left(\sum_{i=1}^k \kappa_i W_i \right) (x_\ell) - (2^* - 1) W_\ell(x_\ell)^{2^*-2} \right|$$

$$\lesssim \begin{cases} \varepsilon \mu_\ell^2 (1 + |x|)^{n-6} + \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} \frac{(1 + |x|)^{n-6}}{\left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^{n-2}} & \text{in } \hat{B}_\ell \setminus \hat{B}_{\ell+1} \\ \sum_{i=\ell+1}^k \hat{W}_i^{2^*-2}(x) & \text{in } \hat{B}_{\ell+1} \end{cases}.$$

By (2.4) and the choice of the conformal correction Φ we also have, with (2.1) and (2.2):

$$\left| \mu_\ell^2 \Lambda_{\xi_\ell}(x_\ell)^{2-2^*} W_\ell^{2^*-2}(x_\ell) - U_0^{2^*-2}(x) \right| \lesssim \begin{cases} \mu_\ell^{n-2} (1 + |x|)^{n-6} & \text{if } (M, g) \text{ is l.c.f.} \\ \mu_\ell^4 & \text{otherwise.} \end{cases}$$

Finally, the definition of the functions $Z_{i,j}$ in (2.15) and straightforward computations using (2.19) and (2.20) yield that, for any $1 \leq i \leq k$,

$$\left| \left(\Delta_{\hat{g}_\ell} + \mu_\ell^2 c_n S_{g_{\xi_\ell}}(x_\ell) + \mu_\ell^2 \varepsilon h(x_\ell) \Lambda_{\xi_\ell}(x_\ell)^{2-2^*} \right) \sum_{j=0..n} \lambda_{ij} \hat{Z}_{i,j} \right|$$

$$\lesssim \begin{cases} \left(\frac{\mu_\ell}{\mu_i} \right)^{\frac{n+2}{2}} \Lambda_i & \text{if } i \leq \ell - 1 \\ \Lambda_\ell (1 + |x|)^{-n-2} & \text{if } i = \ell \\ \Lambda_i \hat{W}_i(x)^{2^*-1} & \text{if } i \geq \ell + 1 \end{cases}$$

Note also that we have $\hat{Z}_{\ell,j}(x) = V_j(x)$ for any $x \in \hat{B}_\ell$, where V_j is as in (2.17). Gathering the previous estimates in (3.22) shows that $\hat{\varphi}_\varepsilon$ satisfies:

$$\begin{aligned}
& \left| \left(\Delta_\varepsilon - (2^* - 1)U_0^{2^*-2} \right) \hat{\varphi}_\varepsilon \right| \lesssim \varepsilon \mu_\ell^2 |\hat{\varphi}_\varepsilon| + \mu_\ell^{\frac{n+2}{2}} |k_\varepsilon(x_\ell)| \\
& + \frac{\Lambda_\ell}{(1+|x|)^{n+2}} + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n+2}{2}} \Lambda_j + \sum_{i \geq \ell+1} \Lambda_i \hat{W}_i(x)^{2^*-1} \\
& + \left\{ \begin{array}{l} \varepsilon \mu_\ell^2 (1+|x|)^{n-6} \\ + \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} \frac{(1+|x|)^{n-6}}{\left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^{n-2}} \quad \text{in } \hat{B}_\ell \setminus \hat{B}_{\ell+1} \\ \sum_{i=\ell+1}^k \hat{W}_i(x)^{2^*-2} \quad \text{in } \hat{B}_{\ell+1} \end{array} \right\} \cdot |\hat{\varphi}_\varepsilon| \\
& + \left\{ \begin{array}{ll} \mu_\ell^{n-2} (1+|x|)^{n-6} |\hat{\varphi}_\varepsilon| & (M, g) \text{ l.c.f.} \\ \mu_\ell^4 (1+|x|^2) |\hat{\varphi}_\varepsilon| + \mu_\ell^3 |x|^2 |\nabla \hat{\varphi}_\varepsilon(x)| + \mu_\ell^2 |x|^2 |\nabla^2 \hat{\varphi}_\varepsilon(x)| & \text{otherwise} \end{array} \right. \\
& + \left(\frac{\mu_\ell}{\mu_{\ell-1}} |\hat{\varphi}_\varepsilon| + \sqrt{\frac{\mu_\ell}{\mu_{\ell-1}}} |\nabla \hat{\varphi}_\varepsilon| \right) \cdot \mathbf{1}_{\frac{3}{2} \sqrt{\frac{\mu_{\ell-1}}{\mu_\ell}} \leq |x| \leq 2 \sqrt{\frac{\mu_{\ell-1}}{\mu_\ell}}}.
\end{aligned} \tag{3.24}$$

When $\ell \geq 2$ or when $\ell = 1$ and (M, g) is locally conformally flat the term $\mu_\ell^{n-2} (1+|x|)^{n-6} |\hat{\varphi}_\varepsilon|$ is absorbed into the term $\varepsilon \mu_\ell^2 (1+|x|)^{n-6} |\hat{\varphi}_\varepsilon|$. Recall also that when $\ell = 1$ the term $\varepsilon \mu_\ell^2 (1+|x|)^{n-6} |\hat{\varphi}_\varepsilon|$, which is due to the contribution of $W_\ell^{2^*-3} W_{\ell-1}$, vanishes.

Step 2: a first pointwise estimate. In this step we prove the following rough estimate:

Lemma 3.5. *Let $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$. Then*

$$\begin{aligned}
& |\hat{\varphi}_\varepsilon(x) - \Pi(\hat{\varphi}_\varepsilon)(x)| \\
& \lesssim \varepsilon \mu_\ell^2 \sum_{j \geq \ell} a_j + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} \Lambda_j + \frac{\varepsilon \mu_\ell^2 (1 + \sum_{j \leq \ell} a_j)}{(1+|x|)^{n-4}} \\
& + \frac{\Lambda_\ell}{(1+|x|)^{n-2}} + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1+|x|)^2} \\
& \sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \hat{W}_j + \frac{\mu_\ell^2 a_\ell}{(1+|x|)^{n-4}} \mathbf{1}_{n.l.c.f} \\
& + (1 + a_\ell) \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^2 \hat{W}_{\ell+1}(x) \mathbf{1}_{\{|x - z_{\ell+1}| \leq 2\}},
\end{aligned} \tag{3.25}$$

where $\Pi(\hat{\varphi}_\varepsilon)$ is defined in Lemma 3.3. The notation $\mathbb{1}_{n.l.c.f}$ indicates that this term vanishes if (M, g) is locally conformally flat.

Proof. We apply (3.12) to $\hat{\varphi}_\varepsilon$. As already mentioned, if $\ell = k$ we do not assume anything, but if $\ell \leq k-1$ we assume that $(H_{\ell+1})$ holds. Let $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$ (which only implies $x \in \hat{B}_k$ if $\ell = k$).

• First, straightforward computations using (3.19) and Giraud's lemma show that

$$\int_{\hat{B}_\ell \setminus \hat{B}_{\ell+1}} \varepsilon \mu_\ell^2 |\hat{\varphi}_\varepsilon(y)| |x - y|^{2-n} dy \lesssim \frac{\varepsilon \mu_\ell^2 a_\ell}{(1 + |x|)^{n-4}}.$$

Since $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$ we also obtain, if $\ell \leq k-1$,

$$\begin{aligned} & \int_{\hat{B}_{\ell+1}} \varepsilon \mu_\ell^2 |\hat{\varphi}_\varepsilon(y)| |x - y|^{2-n} dy \\ & \lesssim \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^{2-n} \varepsilon \mu_\ell^2 \int_{\hat{B}_{\ell+1}} |\hat{\varphi}_\varepsilon(y)| dy \\ & \lesssim \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^{2-n} \varepsilon \mu_\ell^{1-\frac{n}{2}} \int_{B_{\ell+1}} |\varphi_\varepsilon(y)| dv_g(y) \\ & \lesssim \left(\sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \right) \hat{W}_{\ell+1}(x) + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1 + |x|)^2}, \end{aligned}$$

where we used the induction property $(H1_{\ell+1})$ to obtain the last line. Gathering these two estimates yields

$$\begin{aligned} & \int_{\hat{B}_\ell} \varepsilon \mu_\ell^2 |\hat{\varphi}_\varepsilon(y)| |x - y|^{2-n} dy \lesssim \frac{\varepsilon \mu_\ell^2 a_\ell}{(1 + |x|)^{n-4}} \\ & + \left(\sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \right) \hat{W}_{\ell+1}(x) + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1 + |x|)^2}. \end{aligned} \tag{3.26}$$

• Using now (3.5), (A.1) and (A.2) we obtain

$$\begin{aligned} & \int_{\hat{B}_\ell} \mu_\ell^{\frac{n+2}{2}} |k_\varepsilon(x_\ell)| |x - y|^{2-n} dy \lesssim \frac{\varepsilon \mu_\ell^2}{(1 + |x|)^{n-4}} + \frac{\mu_\ell^n}{(1 + |x|)^2} \mathbb{1}_{l.c.f.} \\ & + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1 + |x|)^2} + \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^2 \hat{W}_{\ell+1} \mathbb{1}_{\{|x - z_{\ell+1}| \leq 2\}}. \end{aligned} \tag{3.27}$$

• We now compute the terms involving the λ_{ij} . On one side, Giraud's lemma together with straightforward computations yields

$$\begin{aligned} \int_{\hat{B}_\ell} \left(\frac{\Lambda_\ell}{(1+|x|)^{n+2}} + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n+2}{2}} \Lambda_j \right) |x-y|^{2-n} dy \\ \lesssim \frac{\Lambda_\ell}{(1+|x|)^{n-2}} + \frac{\mu_{\ell-1}}{\mu_\ell} \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n+2}{2}} \Lambda_j \\ \lesssim \frac{\Lambda_\ell}{(1+|x|)^{n-2}} + \varepsilon \mu_\ell^2 \sum_{j \leq \ell-1} \Lambda_j, \end{aligned} \quad (3.28)$$

where the sum is empty if $\ell = 1$. On the other side, straightforward computations yield

$$\int_{\hat{B}_\ell} \sum_{i \geq \ell+1} \Lambda_i \hat{W}_i(y)^{2^*-1} |x-y|^{2-n} dy \lesssim \sum_{i \geq \ell+1} \Lambda_i \hat{W}_i(x).$$

This sum is empty if $\ell = k$. If $\ell \leq k-1$, we use in addition the induction property ($H2_{\ell+1}$): by (2.8) and by (2.13) we get

$$\int_{\hat{B}_\ell} \sum_{i \geq \ell+1} \Lambda_i \hat{W}_i(y)^{2^*-1} |x-y|^{2-n} dy \lesssim \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} \frac{\sum_{j=1}^k a_j}{(1+|x|)^{n-2}}. \quad (3.29)$$

• We now turn to the third line in the right-hand side of (3.24). If $\ell \leq k-1$, using the induction property ($H1_{\ell+1}$) we get that

$$\begin{aligned} \int_{\hat{B}_{\ell+1}} \sum_{i \geq \ell+1} \hat{W}_i(y)^{2^*-2} |\hat{\varphi}_\varepsilon(y)| |x-y|^{2-n} dy \lesssim \left(\sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \right) \hat{W}_{\ell+1}(x) \\ + \varepsilon \mu_\ell^2 \frac{(\sum_{j \leq \ell} a_j)}{(1+|x|)^{n-2}}. \end{aligned} \quad (3.30)$$

With the definition of a_ℓ in (3.19) we obtain with Giraud's lemma

$$\int_{\hat{B}_\ell \setminus \hat{B}_{\ell+1}} \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} (1+|y|)^{n-6} |\hat{\varphi}_\varepsilon(y)| |x-y|^{2-n} dy \lesssim \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{a_\ell}{(1+|x|)^2}, \quad (3.31)$$

while using (A.2) below yields

$$\begin{aligned} \int_{\hat{B}_\ell \setminus \hat{B}_{\ell+1}} \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} \frac{(1+|y|)^{n-6}}{\left(\frac{\mu_{\ell+1}}{\mu_\ell} + |y-z_{\ell+1}| \right)^{n-2}} |\hat{\varphi}_\varepsilon(y)| |x-y|^{2-n} dy \\ \lesssim a_\ell \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x-z_{\ell+1}| \right)^2 \hat{W}_{\ell+1}(x) \mathbf{1}_{\{|x-z_{\ell+1}| \leq 2\}}. \end{aligned} \quad (3.32)$$

• Mimicking the previous computations we also obtain, using $(H1_{\ell+1})$ when $\ell \leq k-1$, that when (M, g) is locally conformally flat

$$\begin{aligned} \int_{\hat{B}_\ell} \mu_\ell^{n-2} (1+|y|)^{n-6} |\hat{\varphi}_\varepsilon(y)| |x-y|^{2-n} dy &\lesssim \left(\sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \right) \hat{W}_{\ell+1}(x) \\ &\quad + \frac{\mu_\ell^{n-2} (1+a_\ell)}{(1+|x|)^2} \end{aligned} \quad (3.33)$$

holds, while when (M, g) is not locally conformally flat around ξ_1 we obtain:

$$\begin{aligned} \int_{\hat{B}_\ell} \mu_\ell^4 (1+|y|^2) |\hat{\varphi}_\varepsilon(y)| |x-y|^{2-n} dy &\lesssim \frac{\mu_\ell^4 (1+a_\ell)}{(1+|x|)^{n-6}} + \frac{\mu_\ell^4 \sum_{j \geq \ell} a_j}{(1+|x|)^{n-2}} \\ &\quad + \sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \hat{W}_j(x). \end{aligned} \quad (3.34)$$

• The only terms in (3.24) that remain to be estimated involve derivatives of $\hat{\varphi}_\varepsilon$. We control them with the following Claim:

Claim 3.6. *Let $1 \leq i \leq k$. For any $x \in C_i$, where C_i is as in (3.17), there holds*

$$\begin{aligned} &|\varphi_\varepsilon(x)| + \tilde{\theta}_i(x) |\nabla \varphi_\varepsilon(x)| + \tilde{\theta}_i(x)^2 |\nabla^2 \varphi_\varepsilon(x)| \\ &\lesssim (a_i + \sum_{p \geq i} \Lambda_p) W_i(x) + \mu_{i-1}^{1-\frac{n}{2}} \left(\frac{\mu_i}{\theta_i(x)} \right)^2 + \frac{\theta_{i+1}(x)^2}{\mu_i^2} W_{i+1}(x) \mathbb{1}_{\{\theta_{i+1}(x) \leq \mu_i\}} \\ &\quad + \sum_{p \leq i-1} \Lambda_p \mu_p^{1-\frac{n}{2}}, \end{aligned}$$

where we have let

$$\tilde{\theta}_i(x) = \min(\theta_i(x), \theta_{i+1}(x)).$$

Proof. The proof is a simple application of elliptic regularity results. We keep the notations of (3.21). By (3.24) it is easily seen that for any $y \in \hat{B}_i \setminus \hat{B}_{i+1}$

$$\begin{aligned} &\left| \left(\Delta_{\hat{g}_i} + O(\mu_i^2) \right) \left(\hat{\varphi}_\varepsilon - \sum_{\substack{p=1..k \\ q=0..n}} \lambda_{p,q} \hat{Z}_{p,q} \right)(y) \right| \lesssim \frac{a_i}{(1+|y|)^{n+2}} \\ &\quad + \frac{\varepsilon \mu_i^2}{(1+|y|)^{n-2}} + \left(\frac{\mu_{i+1}}{\mu_i} \right)^{\frac{n-2}{2}} \frac{1}{(1+|y|)^4} \frac{1}{\left(\frac{\mu_{i+1}}{\mu_i} + |z_{i+1} - y| \right)^{n-2}}. \end{aligned}$$

Let $(y_\varepsilon)_\varepsilon$ be a sequence of points in $\hat{B}_i \setminus \hat{B}_{i+1}$.

We first assume that $|y_\varepsilon - z_{i+1}| \leq 2$. For any $z \in B_0(3/2) \setminus B_0(2/3)$ we let

$$\tilde{\varphi}_\varepsilon(z) = \left(\hat{\varphi}_\varepsilon - \sum_{\substack{p=1..k \\ q=0..n}} \lambda_{p,q} \hat{Z}_{p,q} \right)(z_{i+1} + \alpha_i z),$$

where we have let $\alpha_i = |y_\varepsilon - z_{i+1}|$. Letting $\tilde{g}_i(z) = \hat{g}_i(z_{i+1} + \alpha_i z)$ and since $\alpha_i \lesssim \sqrt{\frac{\mu_{i-1}}{\mu_i}}$ by definition it is then easily seen that $\tilde{\varphi}_\varepsilon$ satisfies

$$\left| \Delta_{\tilde{g}_i} \tilde{\varphi}_\varepsilon(z) + O(\mu_i \mu_{i-1} |\tilde{\varphi}_\varepsilon(z)|) \right| \lesssim \alpha_i^2 \left(a_i + \varepsilon \mu_i^2 \right) + \left(\frac{\mu_{i+1}}{\mu_i} \right)^{\frac{n-2}{2}} \alpha_i^2 \left(\frac{\mu_{i+1}}{\mu_i} + \alpha_i \right)^{2-n}.$$

Since $\alpha_i \lesssim \sqrt{\frac{\mu_{i-1}}{\mu_i}}$, \tilde{g}_i is C^k -close to the Euclidean metric for any $k \geq 1$. Standard elliptic theory then yields

$$\|\tilde{\varphi}_\varepsilon\|_{C^2(B_0(5/4) \setminus B_0(4/5))} \lesssim \alpha_i^2 \left(a_i + \varepsilon \mu_i^2 \right) + \left(\frac{\mu_{i+1}}{\mu_i} \right)^{\frac{n-2}{2}} \alpha_i^2 \left(\frac{\mu_{i+1}}{\mu_i} + \alpha_i \right)^{2-n}. \quad (3.35)$$

Since $y_\varepsilon \in \hat{B}_i \setminus \hat{B}_{i+1}$, by letting $\tilde{y}_\varepsilon = \exp_{\xi_i}^{g_{\xi_i}}(\mu_i y_\varepsilon)$ there holds $\alpha_i \lesssim \frac{\theta_{i+1}(\tilde{y}_\varepsilon)}{\mu_i}$ and $\frac{\theta_{i+1}(\tilde{y}_\varepsilon)}{\mu_i} \lesssim \alpha_i$. Scaling back (3.35) to the original variables yields, for all $x \in B_i$ with $\theta_{i+1}(x) \leq \mu_i$:

$$\begin{aligned} & |\varphi_\varepsilon(x)| + \theta_{i+1}(x) |\nabla \varphi_\varepsilon(x)| + \theta_{i+1}(x)^2 |\nabla^2 \varphi_\varepsilon(x)| \\ & \lesssim \left(\sum_{p \geq i} \Lambda_p \right) W_i(x) + (a_i + \varepsilon \mu_i^2) \frac{\theta_{i+1}(x)^2}{\mu_i^2} \mu_i^{1-\frac{n}{2}} + \frac{\theta_{i+1}(x)^2}{\mu_i^2} W_{i+1}(x) \mathbb{1}_{\{\theta_{i+1}(x) \leq \mu_i\}} \\ & + \sum_{p \leq i-1} \Lambda_p \mu_p^{1-\frac{n}{2}} \\ & \lesssim (a_i + \sum_{p \geq i} \Lambda_p) W_i(x) + \mu_{i-1}^{1-\frac{n}{2}} \left(\frac{\mu_i}{\theta_i(x)} \right)^2 + \frac{\theta_{i+1}(x)^2}{\mu_i^2} W_{i+1}(x) \mathbb{1}_{\{\theta_{i+1}(x) \leq \mu_i\}} \\ & + \sum_{p \leq i-1} \Lambda_p \mu_p^{1-\frac{n}{2}}, \end{aligned}$$

which proves Claim 3.6 in the case $\theta_{i+1}(x) \leq \mu_i$. We used here that $\mu_i^{1-\frac{n}{2}} \lesssim W_i(x)$ when $x \in B_i$ and $\theta_{i+1}(x) \leq \mu_i$.

Assume then that $|y_\varepsilon - z_{i+1}| \geq 2$. Let, for any $z \in B_0(3/2) \setminus B_0(2/3)$,

$$\tilde{\varphi}_\varepsilon(z) = \left(\hat{\varphi}_\varepsilon - \sum_{\substack{p=1..k \\ q=0..n}} \lambda_{p,q} \hat{Z}_{p,q} \right) (\alpha_i z),$$

where again $\alpha_i = |y_\varepsilon - z_{i+1}|$. Remember that $|z_{i+1}| \leq 1$. Letting $\tilde{g}_i(z) = \hat{g}_i(\alpha_i z)$, and since $1 + |z_{i+1} - \alpha_i z| \gtrsim 1 + \alpha_i \gtrsim 1$ for any $z \in B_0(3/2) \setminus B_0(2/3)$,

it is then easily seen that $\tilde{\varphi}_\varepsilon$ satisfies

$$\begin{aligned} \left| \Delta_{\tilde{g}_i} \tilde{\varphi}_\varepsilon(z) + O(\mu_i \mu_{i-1} |\tilde{\varphi}_\varepsilon(z)|) \right| &\lesssim \frac{a_i}{(1 + \alpha_i)^{n-2}} + \frac{\mu_i^{n-2} \mathbb{1}_{\text{l.c.f.}}}{(1 + \alpha_i)^2} + \frac{\varepsilon \mu_i^2}{(1 + \alpha_i)^{n-4}} \\ &\quad + \left(\frac{\mu_{i+1}}{\mu_i} \right)^{\frac{n-2}{2}} \frac{1}{(1 + \alpha_i)^n}. \end{aligned}$$

Scaling back to the original variables as before proves the Claim when $x \in B_i$ and $\theta_{i+1}(x) \geq \mu_i$, and concludes the proof of Claim 3.6. \square

We now conclude the proof of (3.25).

- A straightforward application of Claim 3.6 yields, together with $(H2_{\ell+1})$ in the case where $\ell \leq k-1$:

$$\begin{aligned} &\int_{\frac{3}{2}\sqrt{\frac{\mu_{\ell-1}}{\mu_\ell}} \leq |y| \leq 2\sqrt{\frac{\mu_{\ell-1}}{\mu_\ell}}} \left(\frac{\mu_\ell}{\mu_{\ell-1}} |\hat{\varphi}_\varepsilon(y)| + \sqrt{\frac{\mu_\ell}{\mu_{\ell-1}}} |\nabla \hat{\varphi}_\varepsilon(y)| \right) |x - y|^{2-n} dy \\ &\lesssim \varepsilon \mu_\ell^2 \sum_{j \geq \ell} a_j + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} \Lambda_j + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1 + |x|)^2} \\ &\quad + \frac{\varepsilon \mu_\ell^2 (1 + \sum_{j \leq \ell-1} a_j)}{(1 + |x|)^{n-4}}. \end{aligned} \tag{3.36}$$

If (M, g) is locally conformally flat, estimates (3.26) – (3.33) and (3.36) together with (3.12) conclude the proof of (3.25).

- It therefore remains to conclude the proof in the non-locally conformally flat case. In that case, using (3.19), $(H1_{\ell+1})$ and $(H2_{\ell+1})$ if $\ell \leq k-1$, and Claim 3.6, we have

$$\begin{aligned} &\int_{\hat{B}_\ell} \left(\mu_\ell^3 |y|^2 |\nabla \hat{\varphi}_\varepsilon(y)| + \mu_\ell^2 |y|^2 |\nabla^2 \hat{\varphi}_\varepsilon(y)| \right) |x - y|^{2-n} dy \lesssim \frac{\mu_\ell^2 (a_\ell + \Lambda_\ell)}{(1 + |x|)^{n-4}} \\ &\quad + \varepsilon \mu_\ell^2 \frac{\sum_{j \leq \ell-1} a_j}{(1 + |x|)^{n-2}} + \left(\sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \right) \hat{W}_{\ell+1}(x) \\ &\quad + \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^2 \hat{W}_{\ell+1}(x) \mathbb{1}_{\{|x - z_{\ell+1}| \leq 1\}} + \frac{\varepsilon \mu_\ell^2}{(1 + |x|)^{n-4}}. \end{aligned} \tag{3.37}$$

Combining (3.26)–(3.34) with (3.37) and (3.12) then concludes the proof of (3.25) in the non-locally conformally flat case. \square

Step 3: Estimation of $|\Pi(\hat{\varphi}_\varepsilon)(x)|$. We prove in this part the following

Claim 3.7. *For all $0 \leq j \leq n$,*

$$\int_{\mathbb{R}^n} \hat{\varphi}_\varepsilon V_j U_0^{2^*-2} dy = O(\varepsilon \mu_\ell^2 \sum_{j \leq \ell-1} a_j) + \begin{cases} O(\varepsilon \mu_\ell^2 a_\ell) & (M, g) \text{ l.c.f.} \\ O(\mu_\ell^2 a_\ell) & \text{otherwise} \end{cases}.$$

As a consequence, for any $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$:

$$|\Pi(\hat{\varphi}_\varepsilon)(x)| \lesssim \frac{\varepsilon \mu_\ell^2 (1 + \sum_{j \leq \ell} a_j)}{(1 + |x|)^{n-4}} + \frac{\mu_\ell^2 a_\ell}{(1 + |x|)^{n-2}} \mathbb{1}_{n.l.c.f.} \quad (3.38)$$

Proof. Let $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$. By definition we have

$$\Pi(\hat{\varphi}_\varepsilon)(x) = \sum_{j=0}^n \frac{1}{\|\nabla V_j\|_{L^2(\mathbb{R}^n)}^2} \left(\int_{\mathbb{R}^n} \hat{\varphi}_\varepsilon V_j U_0^{2^*-2} dy \right) V_j(x),$$

where U_0 is as in (2.16). Since $|V_j(y)| \lesssim U_0(y)$ we only need to prove the first part of the Claim to obtain (3.38). Let $0 \leq j \leq n$. By definition of φ_ε in Proposition 3.1 we have $\varphi_\varepsilon \in K_k^\perp$, which implies that

$$\int_M \varphi_\varepsilon (\Delta_g + c_n S_g + \varepsilon h) Z_{\ell,j} dv_g = 0.$$

By (3.19), (3.21), by (2.19) and (2.20) and since $\hat{Z}_{\ell,j} = V_j$ we have:

$$\begin{aligned} 0 &= \int_M \varphi_\varepsilon (\Delta_g + c_n S_g + \varepsilon h) Z_{\ell,j} dv_g \\ &= \int_{B_\ell} \varphi_\varepsilon (\Delta_g + c_n S_g + \varepsilon h) Z_{\ell,j} dv_g + O\left(\sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j}\right)^{\frac{n-2}{2}} a_j\right) \\ &= \int_{\hat{B}_\ell} \hat{\varphi}_\varepsilon (\Delta_{\hat{g}_\ell}(y) + \mu_\ell^2 c_n S_{\hat{g}_\ell}(y_\ell)) V_j(y) dy + O(\varepsilon \mu_\ell^2 \sum_{j \leq \ell} a_j), \end{aligned} \quad (3.39)$$

where $y_\ell = \exp_{\xi_\ell}^{g_{\xi_\ell}}(\mu_\ell y)$. If (M, g) is locally conformally flat we have

$$(\Delta_{\hat{g}_\ell}(y) + \mu_\ell^2 c_n S_{\hat{g}_\ell}(y_\ell)) V_j(y) = (2^* - 1) U_0^{2^*-2} V_j,$$

so that, with (2.8) and since $\hat{\varphi}_\varepsilon$ is supported in \hat{B}_ℓ ,

$$\int_{\mathbb{R}^n} \hat{\varphi}_\varepsilon U_0^{2^*-2} V_j dy = O\left(\varepsilon \mu_\ell^2 (1 + \sum_{j \leq \ell} a_j)\right).$$

If (M, g) is not locally conformally flat we have

$$(\Delta_{\hat{g}_\ell}(y) + \mu_\ell^2 c_n S_{\hat{g}_\ell}(y_\ell)) V_j(y) = (2^* - 1) U_0^{2^*-2} V_j + O(\mu_\ell^2 (1 + |y|)^{2-n}), \quad (3.40)$$

so that

$$\int_{\hat{B}_\ell} \hat{\varphi}_\varepsilon \left(\Delta_{\hat{g}_\ell}(y) + \mu_\ell^2 c_n S_{\hat{g}_\ell}(y_\ell) \right) V_j(y) dy = \int_{\hat{B}_\ell} \hat{\varphi}_\varepsilon U_0^{2^*-2} V_j dy + O(\mu_\ell^2 a_\ell).$$

Combining with (3.39) concludes the proof of (3.38). \square

Step 4: improvement of (3.25) into an optimal estimate. We prove in this step the following improvement of (3.25):

Claim 3.8. *For any $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$ there holds*

$$\begin{aligned} |\hat{\varphi}_\varepsilon(x)| &\lesssim \varepsilon \mu_\ell^2 \sum_{j \geq \ell} a_j + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} \Lambda_j + \frac{\varepsilon \mu_\ell^2 (1 + \sum_{j \leq \ell} a_j)}{(1 + |x|)^{n-4}} \\ &\quad + \frac{\Lambda_\ell}{(1 + |x|)^{n-2}} + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1 + |x|)^2} \\ &\quad + \sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \hat{W}_j + \frac{\mu_\ell^4 a_\ell}{(1 + |x|)^{n-6}} \mathbf{1}_{n.l.c.f} \\ &\quad + (1 + a_\ell) \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^2 \hat{W}_{\ell+1}(x) \mathbf{1}_{\{|x - z_{\ell+1}| \leq 2\}}. \end{aligned} \quad (3.41)$$

Proof. First, in the locally conformally flat case, (3.41) follows from combining (3.25) with (3.38). We therefore assume that (M, g) is not locally conformally flat. Combining (3.25) and (3.38) we have, for any $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$:

$$\begin{aligned} |\hat{\varphi}_\varepsilon(x)| &\lesssim \varepsilon \mu_\ell^2 \sum_{j \geq \ell} a_j + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} \Lambda_j + \frac{\varepsilon \mu_\ell^2 (1 + \sum_{j \leq \ell} a_j)}{(1 + |x|)^{n-4}} \\ &\quad + \frac{\Lambda_\ell}{(1 + |x|)^{n-2}} + \sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \hat{W}_j + \frac{\mu_\ell^2 a_\ell}{(1 + |x|)^{n-4}} \\ &\quad + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1 + |x|)^2} \\ &\quad + (1 + a_\ell) \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^2 \hat{W}_{\ell+1}(x) \mathbf{1}_{\{|x - z_{\ell+1}| \leq 2\}}. \end{aligned} \quad (3.42)$$

With (3.42) we can now improve (3.38). Using (3.40) together with (3.42) now yields

$$\begin{aligned} &\left| \int_{\hat{B}_\ell} \hat{\varphi}_\varepsilon \left(\Delta_{\hat{g}_\ell}(y) + \mu_\ell^2 c_n S_{\hat{g}_\ell}(y_\ell) \right) V_j(y) dy - \int_{\hat{B}_\ell} \hat{\varphi}_\varepsilon U_0^{2^*-2} V_j dy \right| \\ &\quad \lesssim \varepsilon \mu_\ell^2 \sum_{j=1}^k a_j + \Lambda_\ell + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} \Lambda_j, \end{aligned}$$

where we used (2.8) to claim that in the non-locally conformally flat case we have $\mu_\ell^2 \lesssim \varepsilon$ and thus $\mu_\ell^4 \lesssim \varepsilon \mu_\ell^2$ for all $\ell \geq 1$. The other steps in the proof of (3.38) are left unchanged and we obtain now

$$|\Pi(\hat{\varphi}_\varepsilon)(x)| \lesssim \varepsilon \mu_\ell^2 \sum_{j \geq \ell} a_j + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} \Lambda_j + \frac{\Lambda_\ell + \varepsilon \mu_\ell^2 \sum_{j \leq \ell-1} a_j}{(1+|x|)^{n-2}}. \quad (3.43)$$

Going back to the proof of (3.25) it is easily seen that, when (M, g) is not locally conformally flat, the term $\mu_\ell^2 a_\ell (1+|x|)^{4-n}$ comes from the contribution of (3.37). We are now going to use (3.42) to compute (3.37) again. First, a straightforward adaptation of the proof of Claim 3.6, using now (3.42) to control φ_ε , yields the following improved estimate on the derivatives of φ_ε : for all $x \in C_\ell$,

$$\begin{aligned} & |\varphi_\varepsilon(x)| + \tilde{\theta}_\ell(x) |\nabla \varphi_\varepsilon(x)| + \tilde{\theta}_\ell(x)^2 |\nabla^2 \varphi_\varepsilon(x)| \\ & \lesssim \Lambda_\ell W_\ell(x) + \mu_{\ell-1}^{1-\frac{n}{2}} \left(1 + \sum_{j \geq \ell+1} a_j \right) \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + a_\ell \theta_\ell^2 W_\ell(x) \\ & + (1 + a_\ell) \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbb{1}_{\{\theta_{\ell+1}(x) \leq \mu_\ell\}} + \sum_{p \leq \ell-1} \Lambda_p \mu_p^{1-\frac{n}{2}}, \end{aligned}$$

where C_ℓ is as in (3.17) and where we have again let $\tilde{\theta}_\ell(x) = \min(\theta_{\ell+1}(x), \theta_\ell(x))$. The latter estimate now yields, together with $(H1_{\ell+1})$ when $\ell \leq k-1$:

$$\begin{aligned} & \int_{\hat{B}_\ell} \left(\mu_\ell^3 |y|^2 |\nabla \hat{\varphi}_\varepsilon(y)| + \mu_\ell^2 |y|^2 |\nabla^2 \hat{\varphi}_\varepsilon(y)| \right) |x-y|^{2-n} dy \lesssim \varepsilon \mu_\ell^2 \sum_{j \geq \ell} a_j \\ & + \frac{\Lambda_\ell}{(1+|x|)^{n-2}} + \frac{\varepsilon \mu_\ell^2 (1 + \sum_{j \leq \ell} a_j)}{(1+|x|)^{n-4}} + \left(\sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \right) \hat{W}_{\ell+1}(x) \\ & + \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^2 \hat{W}_{\ell+1}(x) \mathbb{1}_{\{|x - z_{\ell+1}| \leq 1\}} + \frac{\mu_\ell^4 a_\ell}{(1+|x|)^{n-6}} \mathbb{1}_{\text{n.l.c.f.}} \end{aligned} \quad (3.44)$$

Combining (3.26) – (3.34) with (3.43) and (3.44) finally proves (3.41). \square

Step 5: estimation of Λ_ℓ and conclusion. Let $j \in \{0, \dots, n\}$ be fixed. We integrate (3.7) against $Z_{\ell,j}$: using (2.8), (2.21), (2.22), (3.5) and (3.16) we

obtain

$$\begin{aligned}
|\lambda_{\ell,j}| + o(\Lambda_\ell) &= O(\varepsilon\mu_\ell^2) + \int_M \varphi_\varepsilon \left(\Delta_g + c_n S_g + \varepsilon h - f' \left(\sum_{j=1}^k \kappa_j W_j \right) \right) Z_{\ell,j} dv_g \\
&= O(\varepsilon\mu_\ell^2) + O \left(\sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} a_j + \sum_{j \geq \ell+1} \left(\frac{\mu_j}{\mu_\ell} \right)^{\frac{n-2}{2}} a_j \right) \\
&\quad + \int_{B_\ell \setminus B_{\ell+1}} \varphi_\varepsilon \left(\Delta_g + c_n S_g + \varepsilon h - f' \left(\sum_{j=1}^k \kappa_j W_j \right) \right) Z_{\ell,j} dv_g \\
&= O(\varepsilon\mu_\ell^2 (1 + \sum_{j=1}^k a_j)) \\
&\quad + \int_{B_\ell \setminus B_{\ell+1}} \varphi_\varepsilon \left(\Delta_g + c_n S_g + \varepsilon h - (2^* - 1) W_\ell^{2^*-2} \right) Z_{\ell,j} dv_g
\end{aligned}$$

where we used (3.23) and where we also used (2.7) to claim that $\mu_{\ell+1}/\mu_\ell \lesssim \mu_\ell/\mu_{\ell-1}$ for $1 \leq \ell \leq k$. Using (2.19) and (2.20) and controlling $\hat{\varphi}_\varepsilon$ by (3.41) finally yields, after straightforward computations:

$$|\lambda_{\ell,j}| + o(\Lambda_\ell) = O(\varepsilon\mu_\ell^2 (1 + \sum_{j=1}^k a_j)).$$

By (3.20) this yields in turn

$$\Lambda_\ell \lesssim \varepsilon\mu_\ell^2 \left(1 + \sum_{j=1}^k a_j \right), \quad (3.45)$$

which proves that $(H2_\ell)$ defined by $(H2_\ell)$ holds true. By plugging (3.45) into (3.41) we obtain that, for any $x \in \hat{B}_\ell \setminus \hat{B}_{\ell+1}$ and for any $1 \leq \ell \leq k$:

$$\begin{aligned}
|\hat{\varphi}_\varepsilon(x)| &\lesssim \varepsilon\mu_\ell^2 \sum_{j \geq \ell} a_j + \sum_{j \leq \ell-1} \left(\frac{\mu_\ell}{\mu_j} \right)^{\frac{n-2}{2}} \Lambda_j + \frac{\varepsilon\mu_\ell^2 (1 + \sum_{j \leq \ell} a_j)}{(1 + |x|)^{n-4}} \\
&\quad + \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \frac{1}{(1 + |x|)^{n-2}} + \frac{\mu_\ell^4 a_\ell}{(1 + |x|)^{n-6}} \mathbb{1}_{n.l.c.f} \\
&\quad + \sum_{j \geq \ell+1} \mu_{j-1}^2 a_j \hat{W}_j \\
&\quad + (1 + a_\ell) \left(\frac{\mu_{\ell+1}}{\mu_\ell} + |x - z_{\ell+1}| \right)^2 \hat{W}_{\ell+1}(x) \mathbb{1}_{\{|x - z_{\ell+1}| \leq 2\}}.
\end{aligned} \quad (3.46)$$

We can now conclude the proof of Lemma 3.4. If (M, g) is locally conformally flat, $(H1_\ell)$ follows from (3.46) by scaling back in the original variables. In the

non-locally conformally flat case we use (2.8) to claim that $\mu_\ell^2 \lesssim \varepsilon$ and thus

$$\frac{\mu_\ell^4 a_\ell}{(1 + |x|)^{n-6}} \lesssim \varepsilon \mu_\ell^2 a_\ell.$$

Scaling back (3.46) to the original variables and using (2.8) we obtain:

$$\begin{aligned} |\varphi_\varepsilon(x)| &\lesssim \varepsilon \mu_\ell^{3-\frac{n}{2}} \sum_{j \geq \ell} a_j + \sum_{j \leq \ell-1} \mu_j^{1-\frac{n}{2}} \Lambda_j + \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 \\ &\quad + \sum_{j \geq \ell+1} \mu_{j-1}^2 a_j W_j(x) + \varepsilon \left(1 + \sum_{j \leq \ell} a_j \right) \theta_\ell^2(x) W_\ell(x) \\ &\quad + (1 + a_\ell) \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbb{1}_{\{\theta_{\ell+1}(x) \leq \mu_\ell\}}, \end{aligned}$$

which proves that $(H1_\ell)$ holds true. Since in the case $\ell = k$ we did not assume anything, this shows that (H_k) is true. The general case for $1 \leq \ell \leq k$ follows by induction. \square

3.4. Proof of Proposition 3.2, Part 2: end of the proof. The last step of the proof of Proposition 3.2 is an inductive estimation of the a_i .

Claim 3.9. *For any $2 \leq \ell \leq k$ we have*

$$a_\ell \lesssim \left(\frac{\mu_\ell}{\mu_{\ell-1}} + \varepsilon \mu_{\ell-1}^2 \right) \left(1 + \sum_{j=1}^{\ell-1} a_j \right). \quad (3.47)$$

Proof. We prove (3.47) inductively using $(H1_\ell)$ and $(H2_\ell)$. Let $\ell \geq 2$. If $\ell = k$ we do not assume anything, while if $\ell \leq k-1$ we assume that (3.47) is true for any $p \geq \ell+1$. For $x \in M$ let $G_\varepsilon(x, \cdot)$ denote the Green's function for $\Delta_g + c_n S_g + \varepsilon h$ in M . Standard properties of G_ε (Robert [34]) ensure that

$$G_\varepsilon(x, y) \lesssim d_g(x, y)^{2-n} \quad \text{for all } x \neq y.$$

We write a representation formula for φ_ε in M : with (3.7) it writes as

$$\begin{aligned} &\left| \varphi_\varepsilon(x) - \sum_{\substack{i=1..k \\ j=0..n}} \lambda_{i,j} Z_{i,j}(x) \right| \\ &\lesssim \int_M d_g(x, \cdot)^{2-n} |k_\varepsilon| dv_g + \sum_{i=2}^k \int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} |\varphi_\varepsilon| dv_g \\ &\quad + \int_{M \setminus B_{g_{\xi_1}}(\xi_1, r_0)} d_g(x, \cdot)^{2-n} W_1^{2^*-2} |\varphi_\varepsilon| dv_g \\ &\quad + \int_{B_{g_{\xi_1}}(\xi_1, r_0) \setminus B_2} d_g(x, \cdot)^{2-n} W_1^{2^*-2} |\varphi_\varepsilon| dv_g, \end{aligned} \quad (3.48)$$

where r_0 is as in (2.3). Assume now that $x \in C_\ell$, where C_ℓ is defined in (3.17). First, straightforward computations using (2.8), (2.13), (3.5), (A.1) and (A.2) below show that

$$\begin{aligned} \int_M d_g(x, \cdot)^{2-n} |k_\varepsilon| dv_g &\lesssim \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} \\ &\quad + \frac{\mu_\ell}{\mu_{\ell-1}} W_\ell(x) + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbf{1}_{\theta_{\ell+1}(x) \leq \mu_\ell}. \end{aligned}$$

Independently, using the definition of a_1 in (3.19), (A.1) and since $x \in C_\ell$, $\ell \geq 2$, we have

$$\int_{M \setminus B_{g_{\xi_1}}(\xi_1, r_0)} d_g(x, \cdot)^{2-n} W_1^{2^*-2} |\varphi_\varepsilon| dv_g \lesssim a_1 \mu_1^{\frac{n+2}{2}} \lesssim a_1 \varepsilon \mu_1^{3-\frac{n}{2}}.$$

Using (2.13), $(H1_\ell)$, (A.1) and (A.2) we similarly obtain, since $x \in C_\ell$ and $\ell \geq 2$, that:

$$\begin{aligned} &\sum_{i=2}^k \int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} |\varphi_\varepsilon| dv_g + \int_{B_{g_{\xi_1}}(\xi_1, r_0) \setminus B_2} d_g(x, \cdot)^{2-n} W_1^{2^*-2} |\varphi_\varepsilon| dv_g \\ &\lesssim \left(1 + \sum_{j \geq \ell} a_j \right) \left(\mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} + \frac{\mu_\ell}{\mu_{\ell-1}} W_\ell(x) \right. \\ &\quad \left. + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbf{1}_{\theta_{\ell+1}(x) \leq \mu_\ell} \right). \end{aligned}$$

Using (2.13) and $(H2_\ell)$ we also obtain that

$$\left| \sum_{\substack{i=1..k \\ j=0..n}} \lambda_{ij} Z_{i,j}(x) \right| \lesssim \left(1 + \sum_{j=1}^k a_j \right) \left(\frac{\mu_\ell}{\mu_{\ell-1}} W_\ell(x) + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} \right).$$

Remark now that there holds, for any $x \in C_\ell$ and since $\ell \geq 2$:

$$\mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbf{1}_{\theta_{\ell+1}(x) \leq \mu_\ell} \lesssim \frac{\mu_\ell}{\mu_{\ell-1}} W_\ell(x). \quad (3.49)$$

Plugging the latter estimates in (3.48) finally gives

$$|\varphi_\varepsilon(x)| \lesssim \left(1 + \sum_{j=1}^k a_j \right) \left(\frac{\mu_\ell}{\mu_{\ell-1}} W_\ell(x) + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} \right).$$

If $\ell = k$ we do not assume anything, while if $\ell \leq k - 1$ we use in addition (3.47) inductively. Together with the latter estimate we obtain in each case

$$|\varphi_\varepsilon(x)| \lesssim \left(1 + \sum_{j=1}^{\ell} a_j\right) \left(\frac{\mu_\ell}{\mu_{\ell-1}} W_\ell(x) + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}}\right).$$

Let now $x_\ell \in C_\ell$ be a point where a_ℓ is attained (recall that a_ℓ is defined in (3.19)). Then $|\varphi_\varepsilon(x_\ell)| = a_\ell W_\ell(x_\ell)$. Since $W_\ell(x) \gtrsim \mu_{\ell-1}^{1-\frac{n}{2}}$ on C_ℓ the latter estimate gives, together with (2.8),

$$a_\ell \lesssim \left(\frac{\mu_\ell}{\mu_{\ell-1}} + \varepsilon \mu_{\ell-1}^2\right) \left(1 + \sum_{j=1}^{\ell-1} a_j\right) + o(a_\ell),$$

which proves (3.47) for $\ell \geq 2$. □

It remains to estimate a_1 to conclude the proof. This is the content of the following Claim:

Claim 3.10. *We have*

$$a_1 \lesssim \min\left(\frac{\mu_2}{\mu_1}, \varepsilon, \mu_1^2\right).$$

In particular, $a_1 = o(1)$ as $\varepsilon \rightarrow 0$.

Proof. Estimate $(H1_1)$ is for the moment still too rough on $M \setminus B_2$, because of the constant term $\varepsilon \mu_1^{3-\frac{n}{2}}$. As a first thing, we improve the precision of the estimate. Let $x \in M \setminus B_2$. We use again the representation formula (3.48) and we estimate again each term. Since $x \in M \setminus B_2$ we have, with (A.1) and (A.2)

$$\begin{aligned} \int_M d_g(x, \cdot)^{2-n} |k_\varepsilon| dv_g &\lesssim \left\{ \begin{array}{ll} \mu_1^{\frac{n-2}{2}} \left(\frac{\mu_1}{\theta_1(x)}\right)^2 & \text{if } (M, g) \text{ is l.c.f} \\ \varepsilon \theta_1(x)^2 W_1(x) & \text{otherwise} \end{array} \right\} \\ &\quad + \varepsilon \mu_2^2 W_1 + \frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbf{1}_{\theta_2(x) \leq \mu_1}. \end{aligned}$$

By definition of a_1 we also have

$$\int_{M \setminus B_{g_{\xi_1}}(\xi_1, r_0)} d_g(x, \cdot)^{2-n} W_1^{2^*-2} |\varphi_\varepsilon| dv_g \lesssim a_1 \mu_1^{\frac{n+2}{2}}.$$

Using (2.13), $(H1_\ell)$, (3.47), (A.1) and (A.2) we similarly obtain, since $x \in M \setminus B_2$, that:

$$\begin{aligned}
& \sum_{i=2}^k \int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} |\varphi_\varepsilon| dv_g \\
& + \int_{B_{g_{\xi_1}}(\xi_1, r_0) \setminus B_2} d_g(x, \cdot)^{2-n} W_1^{2^*-2} |\varphi_\varepsilon| dv_g \\
& \lesssim (1 + a_1) \left[\varepsilon \mu_1^{3-\frac{n}{2}} \left(\frac{\mu_1}{\theta_1(x)} \right)^2 + \varepsilon \mu_2^2 W_1 + \frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbb{1}_{\theta_2(x) \leq \mu_1} \right].
\end{aligned} \tag{3.50}$$

By $(H2_\ell)$, (3.47) and (2.8) we now have

$$\left| \sum_{\substack{i=1..k \\ j=0..n}} \lambda_{ij} Z_{i,j}(x) \right| \lesssim \varepsilon \mu_1^2 (1 + a_1) W_1(x).$$

Combining the latter estimates into (3.48) then yields

$$\begin{aligned}
|\varphi_\varepsilon(x)| & \lesssim \left\{ \begin{array}{ll} \mu_1^{\frac{n-2}{2}} \left(\frac{\mu_1}{\theta_1(x)} \right)^2 & \text{if } (M, g) \text{ is l.c.f} \\ \varepsilon \theta_1(x)^2 W_1(x) & \text{otherwise} \end{array} \right\} \\
& + (1 + a_1) \left[\varepsilon \mu_1^{3-\frac{n}{2}} \left(\frac{\mu_1}{\theta_1(x)} \right)^2 + \frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbb{1}_{\theta_2(x) \leq \mu_1} \right].
\end{aligned} \tag{3.51}$$

It is easily seen that the limiting term $\varepsilon \mu_1^{3-\frac{n}{2}} \left(\frac{\mu_1}{\theta_1(x)} \right)^2$ in (3.50) is due to the second integral in the left-hand side of (3.50) (the contribution over $B_{g_{\xi_1}}(\xi_1, r_0) \setminus B_2$). We now use (3.51) to improve the precision of this term. After a finite number of iterations using (A.1), (3.50) becomes

$$\begin{aligned}
|\varphi_\varepsilon(x)| & \lesssim (1 + a_1) \left\{ \begin{array}{ll} \mu_1^{\frac{n-2}{2}} \left(\frac{\mu_1}{\theta_1(x)} \right)^2 & \text{if } (M, g) \text{ is l.c.f} \\ \varepsilon \theta_1(x)^2 W_1(x) & \text{otherwise} \end{array} \right\} \\
& + (1 + a_1) \frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbb{1}_{\theta_2(x) \leq \mu_1}.
\end{aligned} \tag{3.52}$$

As before, since $x \in M \setminus B_2$, we have

$$\frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbb{1}_{\theta_2(x) \leq \mu_1} \lesssim \frac{\mu_2}{\mu_1} W_1(x).$$

It remains to apply (3.52) to a point $x_1 \in M \setminus B_2$ where a_1 is attained. We obtain:

$$a_1 \lesssim o(a_1) + \frac{\mu_2}{\mu_1} + \begin{cases} \mu_1^2 & \text{if } (M, g) \text{ is l.c.f.} \\ \varepsilon & \text{otherwise} \end{cases},$$

which concludes the proof of the Claim. \square

End of the proof of Proposition 3.2. We are now ready to conclude the proof of Proposition 3.2. First, Claim 3.10 shows that $a_1 = o(1)$. Using this in (3.52) proves (3.10). Then, using Claim 3.10 together with (3.47) shows inductively that

$$a_\ell \lesssim \frac{\mu_\ell}{\mu_{\ell-1}} + \varepsilon \mu_{\ell-1}^2 \quad (3.53)$$

for all $\ell \geq 2$, so that in particular $a_\ell = o(1)$ for all $1 \leq \ell \leq k$. With $(H2_\ell)$, this proves (3.8). It remains to prove that (3.9) is true. Let $2 \leq \ell \leq k$ be fixed. By (3.53) and (2.8) we have

$$\begin{aligned} \mu_{\ell-1}^{1-\frac{n}{2}} \sum_{j \geq \ell} a_j &\lesssim \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} + \frac{\mu_\ell}{\mu_{\ell-1}} \mu_{\ell-1}^{1-\frac{n}{2}} \\ &\lesssim \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} + \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 \end{aligned}$$

for any $x \in B_\ell \setminus B_{\ell+1}$. Similarly, (3.8) now shows that

$$\sum_{j \leq \ell-1} \mu_j^{1-\frac{n}{2}} \Lambda_j \lesssim \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}}.$$

With (3.53) we also have, for any $p \geq \ell + 1$ and for any $x \in B_\ell \setminus B_{\ell+1}$:

$$\begin{aligned} \mu_{p-1}^2 a_p W_p(x) &\lesssim \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} W_\ell(x) + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbb{1}_{\theta_{\ell+1} \leq \mu_\ell} \\ &\lesssim \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbb{1}_{\theta_{\ell+1} \leq \mu_\ell}. \end{aligned}$$

Finally, since $\ell \geq 2$, we have

$$\varepsilon \theta_\ell^2(x) W_\ell(x) \lesssim \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2.$$

Combining these estimates in $(H1_\ell)$ finally prove (3.9) and concludes the proof of Proposition 3.2. \square

4. THE NONLINEAR PROBLEM

4.1. Solving (1.3) up to kernel elements. We perform in this section a nonlinear fixed-point argument in strong spaces and solve (1.3) up to kernel elements. Let $k \geq 1$ be an integer, $(\kappa_i)_{1 \leq i \leq n} \in \{-1, 1\}^k$ and let $\varepsilon_0 > 0$ be as in Proposition 3.1. Let $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$, where S_k is defined in (2.9). For $0 < \varepsilon \leq \varepsilon_0$ we define a function Ψ_ε in M as follows:

$$\begin{aligned} \bullet \Psi_\varepsilon(x) &= \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\theta_\ell(x)} \right)^2 + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}} + \frac{\theta_{\ell+1}(x)^2}{\mu_\ell^2} W_{\ell+1}(x) \mathbb{1}_{\{\theta_{\ell+1}(x) \leq \mu_\ell\}} \\ &\quad \text{if } x \in B_\ell \setminus B_{\ell+1}, \quad \text{for } \ell \geq 2, \\ \bullet \Psi_\varepsilon(x) &= \left\{ \begin{array}{ll} \frac{\mu_1^{\frac{n+2}{2}}}{\theta_1(x)^2} & \text{if } (M, g) \text{ is l.c.f. around } \xi_1 \\ \frac{\mu_1^{\frac{n+2}{2}}}{\theta_1(x)^{n-4}} & \text{if } (M, g) \text{ is not l.c.f.} \end{array} \right\} + \frac{\theta_2(x)^2}{\mu_1^2} W_2(x) \mathbb{1}_{\{\theta_2(x) \leq \mu_1\}} \\ &\quad \text{if } x \in M \setminus B_2, \end{aligned} \tag{4.1}$$

where W_ℓ is given by (2.11). Let $C > 0$ to be chosen later. We define

$$E_\varepsilon^C = \left\{ \varphi \in C^0(M) \text{ such that } |\varphi(x)| \leq C \Psi_\varepsilon(x) \text{ for all } x \in M \right\} \tag{4.2}$$

endowed with the following norm:

$$\|\varphi\|_* := \left\| \frac{\varphi}{\Psi_\varepsilon} \right\|_{L^\infty(M)} \quad \text{for } \varphi \in E_\varepsilon^C. \tag{4.3}$$

Note that E_ε^C depends on ε and on $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k})$. Since $\Psi_\varepsilon > 0$ in M , it is easily seen that $(E_\varepsilon^C, \|\cdot\|_*)$ is a Banach space. Building on the linear theory of Proposition 3.2 we perform a nonlinear fixed-point procedure in E_ε^C :

Proposition 4.1. *Let $k \geq 1$ be an integer, $(\kappa_i)_{1 \leq i \leq n} \in \{-1, 1\}^k$ and denote $f(u) = (u+)^{2^*-1}$ or $f(u) = |u|^{2^*-2}$. There exist $\varepsilon_1 > 0$ and $C > 0$ such that the following holds. For any $\varepsilon \in (0, \varepsilon_1)$, any $h \in C^0(M)$ and for any $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$, there exists*

$$\varphi_\varepsilon := \varphi_\varepsilon \left((t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \right) \in K_k^\perp$$

which is the unique solution in $K_k^\perp \cap E_\varepsilon^C$ of

$$\Pi_{K_k^\perp} \left(\sum_{i=1}^k \kappa_i W_i + \varphi_\varepsilon - (\Delta_g + c_n S_g + \varepsilon h)^{-1} \left(f \left(\sum_{i=1}^k \kappa_i W_i + \varphi_\varepsilon \right) \right) \right) = 0, \tag{4.4}$$

where E_ε^C is defined in (4.2). In addition,

$$((t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k \mapsto \varphi_\varepsilon \in C^0(M)$$

is continuous for any fixed value of ε .

The main point of Proposition 4.1 is that φ_ε is constructed in the space E_ε^C – and not only in K_k^\perp – and satisfies therefore

$$|\varphi_\varepsilon| \lesssim \Psi_\varepsilon \quad (4.5)$$

in M , where Ψ_ε is given by (4.1). In particular, φ_ε still satisfies the estimates (3.9) and (3.10), which is another way of thinking of the nonlinear problem (4.4) as a perturbation of the linear equation (3.7). Proposition 4.1 therefore provides us with a canonical remainder φ_ε with sharp pointwise estimate that precisely capture the pointwise interactions between the towering bubbling profiles W_i on the whole of M . The information that we obtain in Proposition 4.1 is thus much more precise than a mere energy estimate and will turn out to be crucial to prove Theorem 1.1 in Section 5. Such a precision, of course, comes at the expense of proving sharp pointwise estimates for the linear equation as in Proposition 3.2. Remark that since $\varphi_\varepsilon \in E_\varepsilon^C$ for some C , integrating (4.4) against φ_ε yields

$$\|\varphi_\varepsilon\|_{H^1} = o(1) \quad (4.6)$$

as $\varepsilon \rightarrow 0$, where the H^1 norm is given by (2.14).

Proof. Let $k \geq 1$, $(\kappa_i)_{1 \leq i \leq n} \in \{-1, 1\}^k$, $0 < \varepsilon \leq \varepsilon_0$ and let $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$. In the following proof we will omit, for the sake of clarity, the dependency in ε and $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$ in the quantities $W_i, Z_{i,j}, \dots$ appearing. The proof of Proposition 4.1 follows the standard steps of a nonlinear fixed-point scheme, with the notable exception that the point-fixing is now achieved in the strong space E_ε^C defined by (4.2).

Let $C > 0$ be fixed (to be chosen later). For $\varphi \in E_\varepsilon^C$, as defined in (4.2), we define

$$T(\varphi) = \mathcal{L}_\varepsilon \left(-R_\varepsilon + N(\varphi) \right),$$

where \mathcal{L}_ε is given by (3.4) and where we have let

$$R_\varepsilon = (\triangle_g + c_n S_g + \varepsilon h) \left(\sum_{i=1}^k \kappa_i W_i \right) - f \left(\sum_{i=1}^k \kappa_i W_i \right)$$

and

$$N(\varphi) = f \left(\sum_{i=1}^k \kappa_i W_i + \varphi \right) - f \left(\sum_{i=1}^k \kappa_i W_i \right) - f' \left(\sum_{i=1}^k \kappa_i W_i \right) \varphi.$$

The mapping T is well-defined since $\varphi \in E_\varepsilon^C$ and thus $-R_\varepsilon + N(\varphi) \in C^0(M)$. By (3.4), $T(\varphi)$ solves

$$\begin{aligned} (\Delta_g + c_n S_g + \varepsilon h) T(\varphi) - f' \left(\sum_{i=1}^k \kappa_i W_i \right) T(\varphi) &= -R_\varepsilon + N(\varphi) \\ &+ \sum_{\substack{i=1..k \\ j=0..n}} \lambda_{ij} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j} \end{aligned} \quad (4.7)$$

for some uniquely defined real numbers (λ_{ij}) . We show that T is a contraction on $(E_\varepsilon^C, \|\cdot\|_*)$ for ε small enough, for a suitable fixed value of C .

We first prove that T stabilises E_ε^C . By (4.1) one has

$$\Psi_\varepsilon \lesssim \sum_{i=1}^k W_i,$$

so that for any $\varphi \in E_\varepsilon^C$ we have

$$|N(\varphi)| \lesssim \left(\sum_{i=1}^k W_i \right)^{2^*-3} |\varphi|^2.$$

Let $2 \leq i \leq k$ and $\varphi \in E_\varepsilon^C$. By (4.1) we have, for any $x \in B_i \setminus B_{i+1}$,

$$\begin{aligned} &W_i(x)^{2^*-3} |\varphi(x)|^2 \\ &\lesssim C^2 W_i(x)^{2^*-3} \left(\mu_{i-1}^{2-n} \left(\frac{\mu_i}{\theta_i(x)} \right)^4 + \varepsilon^2 \mu_{i-1}^{6-n} + \left(\frac{\theta_{i+1}(x)}{\mu_i} \right)^4 W_{i+1}^2 \mathbb{1}_{\theta_{i+1}(x) \leq \mu_i} \right) \\ &\lesssim C^2 \left(\frac{\mu_i}{\mu_{i-1}} \right)^2 \mu_{i-1}^{1-\frac{n}{2}} W_i(x)^{2^*-2} + C^2 \varepsilon^2 \mu_{i-1}^4 W_i(x)^{2^*-2} W_{i-1}(x) \\ &+ C^2 \left(\frac{\mu_{i+1}}{\mu_i} \right)^2 W_i(x)^{2^*-2} W_{i+1}(x), \end{aligned} \quad (4.8)$$

where we used (3.49) and (2.8). Similarly for $x \in M \setminus B_2$, we get by (4.1)

$$W_1(x)^{2^*-3} |\varphi(x)|^2 \lesssim C^2 \left(\mu_1^4 \frac{\mu_1^{\frac{n+2}{2}}}{\theta_1(x)^4} + \left(\frac{\mu_2}{\mu_1} \right)^2 W_1(x)^{2^*-2} W_2(x) \right) \quad (4.9)$$

when (M, g) is locally conformally flat around ξ_1 , and

$$W_1(x)^{2^*-3} |\varphi(x)|^2 \lesssim C^2 \left(\mu_1^4 \varepsilon W_1(x) + \left(\frac{\mu_2}{\mu_1} \right)^2 W_1(x)^{2^*-2} W_2(x) \right) \quad (4.10)$$

when (M, g) is not conformally flat. The previous estimates show that there exists a positive constant $D_0 > 0$ such that

$$\frac{N(\varphi)}{D_0 C^2 \left(\mu_1^4 + \frac{\mu_2^2}{\mu_1^2} \right)} \quad (4.11)$$

satisfies (3.5), where C is the constant in (4.2). Independently, by (3.6) we know that there exists a constant $D_1 > 0$ such that R_ε/D_1 also satisfies (3.5). By (4.7) we can then apply Proposition 3.2 which yields a positive constant $C_1 > 0$ independent of ε such that

$$|T(\varphi)(x)| \leq C_1 \left(1 + C^2 \left(\mu_1^4 + \frac{\mu_2^2}{\mu_1^2} \right) \right) \Psi_\varepsilon(x)$$

for any $\varphi \in E_\varepsilon^C$ and $x \in M$, where Ψ_ε is as in (4.1), or in other words

$$\|T(\varphi)\|_* \leq C_1 \left(1 + C^2 \left(\mu_1^4 + \frac{\mu_2^2}{\mu_1^2} \right) \right)$$

where the norm is defined in (4.3). Choose $C = 2C_1$. Since $\mu_1^4 + \frac{\mu_2^2}{\mu_1^2} \rightarrow 0$ as $\varepsilon \rightarrow 0$, for ε small enough we then have $T(\varphi) \in E_\varepsilon^C$.

We now prove that T is a contraction on $(E_\varepsilon^C, \|\cdot\|)$ for this fixed value of C . If $\varphi_1, \varphi_2 \in E_\varepsilon^C$, $T(\varphi_1) - T(\varphi_2)$ satisfies by assumption:

$$\begin{aligned} (\Delta_g + c_n S_g + \varepsilon h)(T(\varphi_1) - T(\varphi_2)) - f' \left(\sum_{i=1}^k \kappa_i W_i \right) (T(\varphi_1) - T(\varphi_2)) \\ = N(\varphi_1) - N(\varphi_2) + \sum_{\substack{i=1..k \\ j=0..n}} \nu_{ij} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j} \end{aligned} \quad (4.12)$$

for some real numbers (ν_{ij}) . By (4.2), (4.3), the mean-value theorem and since $\varphi \in E_\varepsilon^C$ we have

$$\begin{aligned} |N(\varphi_1) - N(\varphi_2)| &\lesssim \left(\sum_{i=1}^k W_i \right)^{2^*-3} (|\varphi_1| + |\varphi_2|) |\varphi_1 - \varphi_2| \\ &\lesssim \left(\sum_{i=1}^k W_i \right)^{2^*-3} \Psi_\varepsilon^2 \|\varphi_1 - \varphi_2\|_* \end{aligned}$$

In (4.8), (4.9) and (4.10) we already showed that

$$\frac{1}{\left(\mu_1^4 + \frac{\mu_2^2}{\mu_1^2} \right)} \left(\sum_{i=1}^k W_i \right)^{2^*-3} \Psi_\varepsilon^2$$

satisfies (3.5) up to some positive constant, and therefore

$$\frac{1}{D_2 \left(\mu_1^4 + \frac{\mu_2^2}{\mu_1^2} \right)} \frac{N(\varphi_1) - N(\varphi_2)}{\|\varphi_1 - \varphi_2\|_*}$$

satisfies (3.5) for some positive constant D_2 independent of ε . With (4.12) we can again apply Proposition 3.2 and we obtain that

$$\|T(\varphi_1) - T(\varphi_2)\|_* \leq C_2 \left(\mu_1^4 + \frac{\mu_2^2}{\mu_1^2} \right) \|\varphi_1 - \varphi_2\|_*$$

for some positive constant C_2 independent of ε . This shows that T is a contraction on E_ε^C for ε small enough. Picard's fixed-point theorem then yields the existence of a unique fixed-point φ_ε of T in E_ε^C which is by (4.7), equivalently, a solution of (4.4).

It remains to see that for a fixed ε the mapping

$$((t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k \mapsto \varphi_\varepsilon \in C^0(M)$$

is continuous. By construction φ_ε satisfies (4.7) with $T(\varphi_\varepsilon) = \varphi_\varepsilon$. By (4.11), $N(\varphi_\varepsilon)$ satisfies (3.5) up to a constant, and therefore Proposition 3.2 applies again: (3.8) in particular shows that the λ_{ij} are uniformly bounded in ε and in the choice of $((t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$. The continuity of $(t_i, z_i)_{1 \leq i \leq \ell} \in S_\ell \mapsto \varphi_{\varepsilon, \ell} \in C^0(M)$ now easily follows from the continuity of the W_i , $1 \leq i \leq k$, from (4.7) and from standard elliptic theory, and by the uniqueness property of $\varphi_{\varepsilon, \ell}$ in $K_k^\perp \cap E_\varepsilon^C$. \square

4.2. Application to *a priori* pointwise estimates. The procedure developed in Propositions 3.2 and 4.1 can be useful when looking for *a priori* estimates. We illustrate this on a toy-model example. Consider for instance $h \in C^0(M)$, a sequence (ε_k) of positive numbers converging to 0 as $k \rightarrow +\infty$ and a sequence $(u_k)_k$ of *positive* functions satisfying

$$\left(\Delta_g + c_n S_g + \varepsilon_k h \right) u_k = u_k^{2^*-1} \quad (4.13)$$

in M . We assume for simplicity that the sequence $(u_k)_k$ blows-up with finite-energy and a zero weak-limit. By Struwe's celebrated compactness result [39] there exists $N \geq 1$ such that

$$\|u_k\|_{H^1} = N K_n^{-n} + o(1)$$

as $k \rightarrow +\infty$, where K_n^{-n} is the energy of the standard bubble in \mathbb{R}^n and where the H^1 norm is given by (2.14). Consider the manifold of the sums of N bubbles in M :

$$\mathcal{M}_N := \left\{ \sum_{i=1}^N W_{\mu_i, \xi_i}, (\mu_i, \xi_i)_{1 \leq i \leq N} \in (\mathbb{R}_+^* \times M)^N \right\},$$

where W_{μ_i, ξ_i} is given by (2.4). It is a $N(n+1)$ -dimensional closed manifold of $H^1(M)$. For any $k \geq 1$, we can thus let $(\mu_{i,k}, \xi_{i,k})_{1 \leq i \leq N}$ be such that

$$W_k := \sum_{i=1}^N W_{\mu_{i,k}, \xi_{i,k}}$$

attains the distance of u_k to \mathcal{M}_N . Letting $\varphi_k = u_k - W_k$ we therefore have

$$\|\varphi_k\|_{H^1} = \inf_{W \in \mathcal{N}} \|u_k - W\|_{H^1}.$$

As a consequence, φ_k is orthogonal to the tangent space to \mathcal{M}_N at $(\mu_{i,k}, \xi_{i,k})_{1 \leq i \leq N}$ which, for k large enough, is given by

$$\mathcal{K}_{N,k} := \text{Span} \left\{ \mathcal{Z}_{i,0}, \mathcal{Z}_{i,j}, \quad 1 \leq i \leq N, 0 \leq j \leq n \right\},$$

where we have let

$$\mathcal{Z}_{i,0} = \frac{2}{n-2} \mu_i \frac{\partial}{\partial \mu_i} W_{\mu_i, \xi_i} \quad \text{and} \quad \mathcal{Z}_{i,j} = -n \mu_i \frac{\partial}{\partial (\xi_i)_j} W_{\mu_i, \xi_i}.$$

Struwe's result asserts in addition that $\|\varphi_k\|_{H^1} \rightarrow 0$ as $k \rightarrow +\infty$. We thus have

$$u_k = \sum_{i=1}^N W_{\mu_{i,k}, \xi_{i,k}} + \varphi_k, \quad \varphi_k \in \mathcal{K}_{N,k}^\perp, \quad \|\varphi_k\|_{H^1(M)} \rightarrow 0 \quad (4.14)$$

for k large enough.

Now, the elements $\mathcal{Z}_{i,0}$ and $\mathcal{Z}_{i,j}$ are respectively equal to $Z_{i,0}$ and $Z_{i,j}$ in (2.15) at first order. A simple adaptation of the results in Robert-Vétois [36] therefore shows that, for $\delta > 0$ small enough fixed there exists, for k large enough, a unique $\psi_k \in \mathcal{K}_{N,k}^\perp$ satisfying $\|\psi_k\|_{H^1} < \delta$ and

$$\Pi_{\mathcal{K}_{N,k}^\perp} \left(\sum_{i=1}^N W_{\mu_{i,k}, \xi_{i,k}} + \psi_k - (\Delta_g + c_n S_g + \varepsilon_k h)^{-1} \left(\left(\sum_{i=1}^N W_{\mu_{i,k}, \xi_{i,k}} + \psi_k \right)_+^{2^*-1} \right) \right) = 0.$$

By (4.14) we thus have $\psi_k = \varphi_k$. Assume now that $(\mu_{i,k}, \xi_{i,k})_{1 \leq i \leq N}$ satisfies (2.8) and (2.10). It is easily seen that the results in Section 3 still apply if we replace $Z_{i,j}$ by $\mathcal{Z}_{i,j}$ and K_k by $\mathcal{K}_{N,k}$, since only the leading-order behavior of $\mathcal{Z}_{i,j}$ comes into play in the proof of Proposition 3.2. We can therefore apply Proposition 4.1 with $\mathcal{K}_{N,k}$ (and, say, $t_i = 1$ for all $1 \leq i \leq N$) which provides us, for k large enough, with a uniquely defined $\tilde{\varphi}_k \in \mathcal{K}_{N,k}^\perp$ which solves the previous equation and satisfies in addition $\|\tilde{\varphi}_k\|_{H^1} = o(1)$ as $k \rightarrow +\infty$ by (4.6). We therefore have $\varphi_k = \psi_k = \tilde{\varphi}_k$ in the end and by (4.2) φ_k satisfies

$$|\varphi_k| \lesssim \Psi_{\varepsilon_k}. \quad (4.15)$$

Proposition 4.1 thus turns the weak H^1 control given by (4.14) into a highly accurate pointwise control. This example is intended as a toy model to match

the analysis developed in this paper. But the method described here is likely to generalize to more general bubbling configurations (with or without a weak limit), to more general potential than $c_n S_g + \varepsilon_k h$ and to sign-changing solutions, provided an analogue of Proposition 3.2 is available.

This approach can be seen as an alternative to the approach in Druet-Hebey-Robert [10] to obtain pointwise blow-up estimates for solutions of critical elliptic equations. Here we reduce the problem to proving sharp pointwise estimates for solutions *of a linear problem*. And the precision of these linear estimates can be expected to only depend on the precision of the *ansatz* chosen for the bubbles (see the discussion surrounding (3.11)). This approach yields an improved precision for symmetry estimates in particular configurations (such as the one described in this paper) and could prove particularly useful in the compactness analysis of highly nonlinear problems, such as critical p -Laplace equations or critical anisotropic equations.

5. EXPANSION OF THE KERNEL COEFFICIENTS AND PROOF OF THEOREM 1.1

In this section we prove Theorem 1.1. Let (M, g) be a locally conformally flat manifold. Let $k \geq 1$ be an integer, ε_1 be given by Proposition 4.1, $h \in C^1(M)$ and $((t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$, where S_k is as in (2.9). Throughout this section we will assume that

$$f(u) = (u_+)^{2^*-1} \quad \text{and} \quad \kappa_i = 1 \quad \forall i \in \{1, \dots, k\}.$$

Let φ_ε be the unique function given by Proposition 4.1. Equation (4.4) together with (4.5) shows the existence of real numbers $(\nu_{i,j})$, $1 \leq i \leq k$ and $0 \leq j \leq n$ such that

$$\begin{aligned} (\Delta_g + c_n S_g + \varepsilon h) \left(\sum_{i=1}^k W_i + \varphi_\varepsilon \right) - \left(\sum_{i=1}^k W_i + \varphi_\varepsilon \right)^{2^*-1} \\ = \sum_{\substack{i=1..k \\ j=0..n}} \nu_{i,j} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j}. \end{aligned} \tag{5.1}$$

The $\nu_{i,j}$ are, by Proposition 4.1, continuous functions of $((t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$. They *a priori* all depend on each variable in S_k . This is a striking difference with the symmetric situation of Morabito-Pistoia-Vaira [26], where an inductive splitting of the remainder term led to a triangular dependency of the coefficients $\nu_{i,j}$. In the present work, we overcome the difficulties due to the non-symmetric background by using the sharp estimates on φ_ε given by (4.5).

We perform an asymptotic expansion of the $\nu_{i,j}$ in (5.1). We state the $\ell = 1$ and $\ell \geq 2$ case separately for clarity but the following two statements are proven together.

Proposition 5.1 (Case $\ell = 1$). *We have:*

$$\|\nabla V_0\|_{L^2(\mathbb{R}^n)}^2 \nu_{1,0}(t_1, \dots, z_k) = \varepsilon^{\frac{n-2}{n-4}} \left(\frac{4D_1}{n-2} h(\xi_1) t_1^2 - 2D_2 t_1^{n-2} m(\xi_1) + \Lambda_{0,1} \right)$$

and, for $1 \leq j \leq n$,

$$\|\nabla V_j\|_{L^2(\mathbb{R}^n)}^2 \nu_{1,j}(t_1, \dots, z_k) = \varepsilon^{\frac{n-1}{n-4}} \left(nD_1 t_1^3 \nabla_j h(\xi_1) - nD_2 t_1^{n-1} \nabla_j m(\xi_1) + \Lambda_{j,1} \right)$$

as $\varepsilon \rightarrow 0$. In these expressions $m(\xi_1)$ denotes the mass of the Green function at ξ_1 as defined in (2.1) and D_1, D_2 are positive constants that only depend on n . Also, V_j is as in (2.17), and $\Lambda_{j,1}$, for $0 \leq j \leq n$, denote continuous functions defined in S_k which converge uniformly to 0 as $\varepsilon \rightarrow 0$.

Proposition 5.2 (Case $2 \leq \ell \leq k$). *For any $2 \leq \ell \leq k$ we have*

$$\|\nabla V_0\|_{L^2(\mathbb{R}^n)}^2 \nu_{\ell,0}(t_1, \dots, z_k) = \varepsilon^{1+2\gamma_\ell} \left(\frac{4D_1}{n-2} h(\xi_1) t_\ell^2 - D_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n-2}{2}} U_0(z_\ell) + \Lambda_{0,\ell} \right)$$

and, for $1 \leq j \leq n$,

$$\|\nabla V_j\|_{L^2(\mathbb{R}^n)}^2 \nu_{\ell,j}(t_1, \dots, z_k) = \varepsilon^{\frac{n}{2}(\gamma_\ell - \gamma_{\ell-1})} \left(-nD_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n}{2}} \partial_j U_0(z_\ell) + \Lambda_{j,\ell} \right)$$

as $\varepsilon \rightarrow 0$, where U_0 is as in (2.16) and γ_ℓ is as in (2.7). Here D_3 is a positive constant, D_1 is the same as in Proposition 5.1 and $\Lambda_{j,\ell}$, for $0 \leq j \leq n$ denote functions defined in S_k which converge uniformly to 0 as $\varepsilon \rightarrow 0$.

In view of Propositions 5.1 and 5.2 and of (2.8) we have to carry on the computations at the order $\varepsilon\mu_1^2$ and $\varepsilon\mu_1^3$ respectively for $\nu_{1,0}$ and $\nu_{1,j}$, $1 \leq j \leq n$, and at the order $\varepsilon\mu_\ell^2$ and $\frac{\mu_\ell}{\mu_{\ell-1}}\varepsilon\mu_\ell^2 = \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n}{2}}$ respectively for $\nu_{\ell,0}$ and $\nu_{\ell,j}$, for $\ell \geq 2$ and $1 \leq j \leq n$. In the following we will let:

$$\Theta_{1,0} = \varepsilon\mu_1^2 \sim \mu_1^{n-2} \quad (5.2)$$

since (M, g) is locally conformally flat and, for any $1 \leq j \leq n$,

$$\Theta_{1,j} = \varepsilon\mu_1^3 \sim \mu_1^{n-1}. \quad (5.3)$$

For $2 \leq \ell \leq k$ we define similarly:

$$\Theta_{\ell,0} = \varepsilon\mu_\ell^2 \quad \text{and, for any } 1 \leq j \leq n, \quad \Theta_{\ell,j} = \frac{\mu_\ell}{\mu_{\ell-1}}\varepsilon\mu_\ell^2 \sim \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n}{2}}. \quad (5.4)$$

For $a, b > 0$, the symbol $a \sim b$ in the previous expressions means that there exists a constant $C > 0$, independent of ε and of the choice of $(t_1, \xi_1, (t_i, z_i)_{2 \leq i \leq k}) \in S_k$, such that

$$\frac{1}{C}a \leq b \leq Ca.$$

Throughout the proof of Propositions 5.1 and 5.2 we will also make use of the following fact:

$$\varepsilon \mu_2^2 \sim \left(\frac{\mu_2}{\mu_1} \right)^{\frac{n-2}{2}} = o(\varepsilon \mu_1^3) \quad (5.5)$$

as $\varepsilon \rightarrow 0$ which, since (M, g) is locally conformally flat, follows easily from (2.8).

Proof of Propositions 5.1 and 5.2. First, φ_ε satisfies (4.5), where Ψ_ε is as in (4.1). As shown in the proof of Proposition 4.1 (see e.g. (4.7) and (4.11)), Proposition 3.2 applies to (5.1) and yields, for any $1 \leq i \leq k$:

$$\sum_{j=0}^n |\nu_{i,j}| \lesssim \varepsilon \mu_i^2. \quad (5.6)$$

As a consequence of (5.6) and of (2.22) we have, for any $\ell \in \{1, \dots, k\}$ and $0 \leq q \leq n$,

$$\int_M \left(\sum_{\substack{i=1..k \\ j=0..n}} \nu_{i,j} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j} \right) Z_{\ell,q} = \|\nabla V_q\|_{L^2(\mathbb{R}^n)}^2 \nu_{\ell,q} + o(\Theta_{\ell,q}). \quad (5.7)$$

Let $1 \leq \ell \leq k$ be fixed. We write (5.1) in the following way:

$$\begin{aligned} & \sum_{\substack{i=1..k \\ j=0..n}} \nu_{i,j} (\Delta_g + c_n S_g + \varepsilon h) Z_{i,j} \\ &= \left(\Delta_g + c_n S_g + \varepsilon h \right) \left(\sum_{i=1}^k W_i \right) - \left(\sum_{i=1}^k W_i \right)^{2^*-1} \\ &+ \left(\Delta_g + c_n S_g + \varepsilon h \right) \varphi_\varepsilon - (2^* - 1) W_\ell^{2^*-2} \varphi_\varepsilon \\ &- (2^* - 1) \left[\left(\sum_{i=1}^k W_i \right)^{2^*-2} - W_\ell^{2^*-2} \right] \varphi_\varepsilon \\ &- \left[\left(\sum_{i=1}^k W_i + \varphi_\varepsilon \right)^{2^*-1} - \left(\sum_{i=1}^k W_i \right)^{2^*-1} - (2^* - 1) \left(\sum_{i=1}^k W_i \right)^{2^*-2} \varphi_\varepsilon \right] \\ &= I_1 + I_2 + I_3 + I_4, \end{aligned} \quad (5.8)$$

where each term I_i , $1 \leq i \leq 4$, is given by the i -th line in the right-hand side of (5.8). We first claim that the sole contribution to $\nu_{\ell,j}$ is given by I_1 :

Lemma 5.3. *For any $1 \leq \ell \leq k$ and any $0 \leq j \leq n$ there holds:*

$$\int_M (I_2 + I_3 + I_4) Z_{\ell,j} dv_g = o(\Theta_{\ell,j}),$$

where $\Theta_{\ell,j}$ is as in (5.2), (5.3) and (5.4).

Proof of Lemma 5.3. We estimate each term separately.

Estimation of I_2 . Let $0 \leq j \leq n$. First, integrating by parts shows that

$$\int_M I_2 Z_{\ell,j} dv_g = \int_M \varphi_{\ell,\varepsilon} \left((\Delta_g + c_n S_g + \varepsilon h) Z_{\ell,j} - (2^* - 1) W_\ell^{2^*-2} Z_{\ell,j} \right) dv_g.$$

Assume first that $j = 0$. By (2.19) and (4.5), and since $|Z_{\ell,0}| \lesssim W_\ell$, we get

$$\begin{aligned} & \left| \int_{B_{\ell+1}} \varphi_{\ell,\varepsilon} \left((\Delta_g + c_n S_g + \varepsilon h) Z_{\ell,j} - (2^* - 1) W_\ell^{2^*-2} Z_{\ell,j} \right) dv_g \right| \\ & \lesssim \mu_\ell^{\frac{n-6}{2}} \int_{B_{\ell+1}} |\varphi_\varepsilon| dv_g \\ & \lesssim \mu_\ell^{\frac{n-6}{2}} \sum_{i=\ell+1}^k \int_{B_i \setminus B_{i+1}} |\varphi_\varepsilon| dv_g = o(\varepsilon \mu_\ell^2). \end{aligned}$$

Similarly, using again (2.19) and (4.5),

$$\begin{aligned} & \left| \int_{M \setminus B_\ell} \varphi_{\ell,\varepsilon} \left((\Delta_g + c_n S_g + \varepsilon h) Z_{\ell,j} - (2^* - 1) W_\ell^{2^*-2} Z_{\ell,j} \right) dv_g \right| \\ & \lesssim \sum_{i=1}^{\ell-1} \int_{B_i \setminus B_{i+1}} \left(\mu_\ell^{\frac{n-2}{2}} + \varepsilon W_\ell \right) |\varphi_\varepsilon| dv_g \\ & = o(\varepsilon \mu_\ell^2), \end{aligned}$$

where this last integral is zero when $\ell = 1$, and

$$\begin{aligned} & \left| \int_{B_\ell \setminus B_{\ell+1}} \varphi_{\ell,\varepsilon} \left((\Delta_g + c_n S_g + \varepsilon h) Z_{\ell,j} - (2^* - 1) W_\ell^{2^*-2} Z_{\ell,j} \right) dv_g \right| \\ & \lesssim \int_{B_\ell \setminus B_{\ell+1}} \left(\mu_\ell^{\frac{n-2}{2}} + \varepsilon W_\ell \right) |\varphi_\varepsilon| dv_g \\ & = o(\varepsilon \mu_\ell^2). \end{aligned}$$

In conclusion we have

$$\int_M I_2 Z_{\ell,0} dv_g = o(\Theta_{\ell,0}), \quad (5.9)$$

for any $1 \leq \ell \leq k$, where $\Theta_{1,0}$ and $\Theta_{\ell,0}$ are given respectively by (5.2) and (5.4).

Assume now that $1 \leq j \leq n$. Integrating by parts and using (2.20) gives

$$\left| \int_M I_2 Z_{\ell,j} dv_g \right| \lesssim \varepsilon \int_M |\varphi_{\ell,\varepsilon}| |Z_{\ell,j}| dv_g + \int_M \mu_\ell^{\frac{n}{2}} \theta_\ell^{2-n} |\varphi_{\ell,\varepsilon}| dv_g.$$

We first have, with (4.5):

$$\begin{aligned} \int_{B_{\ell+1}} (\varepsilon |Z_{\ell,j}| + \mu_\ell^{\frac{n}{2}} \theta_\ell^{2-n}) |\varphi_{\ell,\varepsilon}| dv_g &\lesssim \mu_\ell^{2-\frac{n}{2}} \int_{B_{\ell+1}} |\varphi_\varepsilon| dv_g \\ &= o(\Theta_{\ell,j}) \end{aligned}$$

for all $1 \leq \ell \leq k$, where $\Theta_{\ell,j}$ is given by (5.3) and (5.4). Straightforward computations using (4.5) to estimate the integral over $B_\ell \setminus B_{\ell+1}$ and $M \setminus B_\ell$ similarly give

$$\int_M I_2 Z_{\ell,j} dv_g = o(\Theta_{\ell,j}) \quad (5.10)$$

for any $1 \leq \ell \leq k$ and $1 \leq j \leq n$, where $\Theta_{1,j}$ and $\Theta_{\ell,j}$ are given respectively by (5.3) and (5.4).

Estimation of I_3 . By (4.5) and (5.8) we have

$$|I_3(x)| \lesssim \begin{cases} W_i^{2^*-2} |\phi_\varepsilon| & \text{if } x \in B_i \setminus B_{i+1}, \quad i \neq \ell \\ W_\ell^{2^*-3} (W_{\ell-1} + W_{\ell+1}) |\varphi_\varepsilon| & \text{if } x \in B_\ell \setminus B_{\ell+1} \end{cases},$$

where it is intended that the term $W_{\ell-1}$ vanishes when $\ell = 1$ and the term $W_{\ell+1}$ vanishes when $\ell = k$. So that

$$\begin{aligned} \left| \int_M I_3 Z_{\ell,j} dv_g \right| &\lesssim \sum_{i \neq \ell} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} |\varphi_\varepsilon| |Z_{\ell,j}| dv_g \\ &\quad + \int_{B_\ell \setminus B_{\ell+1}} W_\ell^{2^*-3} (W_{\ell-1} + W_{\ell+1}) |\varphi_\varepsilon| |Z_{\ell,j}| dv_g. \end{aligned}$$

Assume first that $\ell = 1$ and recall that $B_1 = M$. Using (4.5) we obtain

$$\int_{M \setminus B_2} W_1^{2^*-3} W_2 |\varphi_\varepsilon| |Z_{1,j}| dv_g \lesssim o(\mu_1^{n-1}) + \left(\frac{\mu_2}{\mu_1} \right)^{\frac{n+2}{2}} = o(\Theta_{\ell,j})$$

for all $0 \leq j \leq n$, where $\Theta_{1,j}$ is as in (5.2) and (5.3). Similarly we have, using (4.5) and since $|Z_{1,j}| \lesssim W_1$:

$$\begin{aligned} \sum_{i \geq 2} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} |\varphi_\varepsilon| |Z_{1,j}| dv_g &\lesssim \sum_{i \geq 2} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_1 |\varphi_\varepsilon| dv_g \\ &\lesssim \mu_2 \mu_1^{n-2} + \mu_2^{\frac{n+2}{2}} \mu_1^{-\frac{n+2}{2}} + \mu_2^{\frac{n}{2}} \mu_1^{-\frac{n-2}{2}} \\ &= o(\Theta_{1,j}). \end{aligned}$$

In the end we obtain

$$\int_M I_3 Z_{1,j} dv_g = o(\Theta_{1,j}) \quad (5.11)$$

for all $0 \leq j \leq n$.

Assume now that $2 \leq \ell \leq k$. First, straightforward computations using (4.5) show that, for any $0 \leq j \leq n$,

$$\begin{aligned} \int_{B_\ell \setminus B_{\ell+1}} W_\ell^{2^*-3} (W_{\ell-1} + W_{\ell+1}) |\varphi_\varepsilon| |Z_{\ell,j}| dv_g &\lesssim \int_{B_\ell \setminus B_{\ell+1}} W_\ell^{2^*-2} (W_{\ell-1} + W_{\ell+1}) |\varphi_\varepsilon| dv_g \\ &\lesssim \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n+2}{2}} + \varepsilon \mu_\ell^3 + \left(\frac{\mu_{\ell+1}}{\mu_{\ell-1}} \right)^{\frac{n+2}{2}} \\ &= o(\Theta_{\ell,j}). \end{aligned}$$

Similarly, using again (4.5) we have

$$\begin{aligned} \sum_{i \geq \ell+1} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} |\varphi_\varepsilon| |Z_{\ell,j}| dv_g &\lesssim \frac{\mu_{\ell+1}^{\frac{n}{2}}}{\mu_\ell \mu_{\ell-1}^{\frac{n-2}{2}}} + \mu_{\ell+1} \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} \\ &= o(\Theta_{\ell,j}) \end{aligned}$$

for any $0 \leq j \leq n$. Finally, on the one hand, using that $|Z_{\ell,0}| \lesssim W_\ell$ we find

$$\begin{aligned} \sum_{i \leq \ell-1} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} |\varphi_\varepsilon| |Z_{\ell,0}| dv_g &\lesssim \mu_\ell^{\frac{n-2}{2}} \mu_{\ell-2}^{1-\frac{n}{2}} + \mu_\ell^{\frac{n+2}{2}} \mu_{\ell-1}^{-\frac{n+2}{2}} + o \left(\left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \right) \\ &= o(\Theta_{\ell,0}). \end{aligned}$$

On the other hand, using that $|Z_{\ell,j}| \lesssim \mu_\ell^{\frac{n}{2}} \theta_\ell^{1-n}$ for $1 \leq j \leq n$, where θ_ℓ is as in (2.18), we have:

$$\begin{aligned} \sum_{i \leq \ell-1} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} |\varphi_\varepsilon| |Z_{\ell,j}| dv_g &\lesssim \mu_\ell^{\frac{n}{2}} \mu_{\ell-1}^{-1} \mu_{\ell-2}^{1-\frac{n}{2}} + o \left(\left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n}{2}} \right) \\ &= o(\Theta_{\ell,j}). \end{aligned}$$

In the end we thus have

$$\int_M I_3 Z_{\ell,j} dv_g = o(\Theta_{\ell,j}) \quad (5.12)$$

for all $0 \leq j \leq n$.

Estimation of I_4 . We finally estimate the contribution of I_4 . By definition and by (4.5) we have

$$|I_4| \lesssim \left(\sum_{i=1}^k W_i \right)^{2^*-3} |\varphi_\varepsilon|^2 \quad \text{in } M$$

and therefore

$$\left| \int_M I_4 Z_{\ell,j} dv_g \right| \lesssim \sum_{i=1}^k \int_{B_i \setminus B_{i+1}} W_i^{2^*-3} |Z_{\ell,j}| |\varphi_\varepsilon|^2 dv_g.$$

Using again (4.5) and the following estimate on $Z_{\ell,j}$:

$$|Z_{\ell,j}| \lesssim \begin{cases} W_\ell & \text{if } j = 0 \\ \mu_\ell^{\frac{n}{2}} \theta_\ell^{1-n} & \text{if } 1 \leq j \leq n \end{cases}, \quad (5.13)$$

we get by straightforwards computations that mimic those that led to (5.9), (5.10), (5.11) and (5.12) that

$$\int_M I_4 Z_{\ell,j} dv_g = o(\Theta_{\ell,j}), \quad (5.14)$$

for any $1 \leq \ell \leq k$ and $0 \leq j \leq n$, where $\Theta_{\ell,j}$ is as in (5.2), (5.3) and (5.4). Combining (5.9), (5.10), (5.11), (5.12) and (5.14) concludes the proof of Lemma 5.3. \square

Let us now estimate the contribution due to I_1 in (5.8). We first claim that the following holds: for any $1 \leq \ell \leq k$ and $0 \leq j \leq n$,

$$\begin{aligned} \int_M I_1 Z_{\ell,j} dv_g &= \int_M \left[\left(\Delta_g + c_n S_g + \varepsilon h \right) W_\ell - W_\ell^{2^*-1} \right. \\ &\quad \left. - \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] \right] Z_{\ell,j} + o(\Theta_{\ell,j}), \end{aligned} \quad (5.15)$$

where $\Theta_{\ell,j}$ is as in (5.2), (5.3) and (5.4). Indeed, by definition of I_1 in (5.8),

$$I_1 = \sum_{i=1}^k \left[\left(\Delta_g + c_n S_g + \varepsilon h \right) W_i - W_i^{2^*-1} \right] - \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right].$$

By (2.6) and (2.13) and since (M, g) is locally conformally flat we have for any $1 \leq i \leq k$,

$$\left| \left(\Delta_g + c_n S_g + \varepsilon h \right) W_i - W_i^{2^*-1} \right| \lesssim \frac{\mu_i^{\frac{n+2}{2}}}{\theta_i^4} + \varepsilon W_i.$$

Let $i > \ell$. Straightforward computations using (2.8) show that

$$\int_M \frac{\mu_i^{\frac{n+2}{2}}}{\theta_i^4} |Z_{\ell,j}| dv_g + \int_M \varepsilon W_i |Z_{\ell,j}| dv_g \lesssim (\varepsilon + \mu_i^2) \varepsilon \mu_i^2 = o(\Theta_{\ell,j}). \quad (5.16)$$

Similarly, if $i < \ell$,

$$\begin{aligned} &\int_M \frac{\mu_i^{\frac{n+2}{2}}}{\theta_i^4} |Z_{\ell,j}| dv_g + \int_M \varepsilon W_i |Z_{\ell,j}| dv_g \\ &\lesssim \begin{cases} \mu_\ell^{\frac{n-2}{2}} \mu_i^{\frac{n-2}{2}} & j = 0 \\ \mu_\ell^{\frac{n}{2}} \mu_i^{\frac{n-4}{2}} & j \in \{1, \dots, n\} \end{cases} \\ &= o(\Theta_{\ell,j}) \end{aligned} \quad (5.17)$$

again by (2.8). Estimates (5.16) and (5.17) together prove (5.15).

We now prove the following claim:

Lemma 5.4. *For $\ell \geq 2$,*

$$-\int_M \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] Z_{\ell,j} dv_g = -n D_3 \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n}{2}} \partial_j U_0(z_\ell) + o(\Theta_{\ell,j})$$

for some positive constant D_3 (given by (5.18) below), while for $\ell = 1$

$$-\int_M \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] Z_{1,j} dv_g = o(\Theta_{1,j}).$$

Proof. We first assume that $\ell \geq 2$. On $B_{\ell+1}$ we have

$$\left| \left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right| \lesssim W_{\ell+1}^{2^*-2} \left(\sum_{p \geq \ell+2} W_p + W_\ell \right),$$

with the convention that $W_i = 0$ if $i \geq k+1$, so that

$$\begin{aligned} \left| \int_{B_{\ell+1}} \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] Z_{\ell,j} dv_g \right| &\lesssim \left(\frac{\mu_{\ell+2}}{\mu_\ell} \right)^{\frac{n-2}{2}} + \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n}{2}} \\ &= o(\Theta_{\ell,j}), \end{aligned}$$

since by (2.8) we have (for $\ell \geq 2$):

$$\frac{\mu_{\ell+1}}{\mu_\ell} = o\left(\frac{\mu_\ell}{\mu_{\ell-1}} \right).$$

Independently, if $i \leq \ell-1$, we have

$$\left| \left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right| \lesssim W_i^{2^*-2} (W_{i+1} + W_{i-1}) \quad \text{in } B_i \setminus B_{i+1},$$

with the convention that $W_0 = 0$. Using that $|Z_{\ell,0}| \lesssim W_\ell$ and that $|Z_{\ell,j}| \lesssim \mu_\ell^{\frac{n}{2}} \theta_\ell^{1-n}$ then gives that

$$\begin{aligned} \left| \int_{M \setminus B_\ell} \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] Z_{\ell,0} dv_g \right| &\lesssim \sum_{i=1}^{\ell-1} \left(\left(\frac{\mu_\ell}{\mu_i} \right)^{\frac{n}{2}} + \left(\frac{\mu_\ell}{\mu_{i-1}} \right)^{\frac{n-2}{2}} \right) \\ &= o(\Theta_{\ell,0}), \end{aligned}$$

and that

$$\begin{aligned} \left| \int_{M \setminus B_\ell} \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] Z_{\ell,j} dv_g \right| &\lesssim \sum_{i=1}^{\ell-1} \left(\left(\frac{\mu_\ell}{\mu_i} \right)^{\frac{n+1}{2}} + \frac{\mu_\ell^{\frac{n}{2}}}{\mu_{i-1}^{\frac{n-2}{2}} \mu_i} \right) \\ &= o(\Theta_{\ell,j}), \end{aligned}$$

with the convention that the term $\left(\frac{\mu_\ell}{\mu_{i-1}}\right)^{\frac{n-2}{2}}$ vanishes if $i = 1$. It therefore remains to estimate the integral over $B_\ell \setminus B_{\ell+1}$. Using (2.13) we can write that

$$\begin{aligned} & \left| \left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} - (2^* - 1) W_\ell^{2^*-2} \sum_{i=1}^{\ell-1} W_i \right| \\ & \lesssim W_\ell^{2^*-2} \sum_{p \geq \ell+1} W_p + W_\ell^{2^*-3} (W_{\ell+1}^2 + W_{\ell-1}^2), \end{aligned}$$

so that

$$\begin{aligned} & \left| \int_{B_\ell \setminus B_{\ell+1}} \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} - (2^* - 1) W_\ell^{2^*-2} \sum_{i=1}^{\ell-1} W_i \right] Z_{\ell,j} dv_g \right| \\ & \lesssim \left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} + \begin{cases} \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n}{2}} & j = 0 \\ \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n+1}{2}} & j \in \{1, \dots, n\} \end{cases} \\ & = o(\Theta_{\ell,j}). \end{aligned}$$

For the last line we wrote that

$$\left(\frac{\mu_{\ell+1}}{\mu_\ell} \right)^{\frac{n-2}{2}} = \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n-2}{2}} \left(\frac{\mu_{\ell+1} \mu_{\ell-1}}{\mu_\ell^2} \right)^{\frac{n-2}{2}} = o(\Theta_{\ell,j}),$$

and used (2.8), which shows that

$$\left(\frac{\mu_{\ell+1} \mu_{\ell-1}}{\mu_\ell^2} \right)^{\frac{n-2}{2}} = o \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right).$$

We have thus proven that

$$\begin{aligned} & - \int_M \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] Z_{\ell,j} dv_g \\ & = -(2^* - 1) \int_{B_\ell \setminus B_{\ell+1}} W_\ell^{2^*-2} \left(\sum_{i \leq \ell-1} W_i \right) Z_{\ell,j} dv_g + o(\Theta_{\ell,j}). \end{aligned}$$

Let, for any $y \in B_0(\frac{\mu_\ell}{\mu_{\ell-1}})$ and any $i \leq \ell - 1$:

$$\hat{W}_i(y) = \mu_\ell^{\frac{n-2}{2}} W_i(\exp_{\xi_\ell}^{g_{\xi_\ell}}(\mu_\ell y)).$$

As is easily checked, V_j defined in (2.17) satisfies

$$V_j(x) = -n \partial_j U_0(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Let $i \leq \ell - 1$. Using (2.11), changing variables and integrating by parts we have

$$\begin{aligned} (2^* - 1) \int_{B_\ell \setminus B_{\ell+1}} W_\ell^{2^*-2} W_i Z_{\ell,j} &= (2^* - 1) \int_{B_0(\frac{\mu_{\ell-1}}{\mu_\ell})} U_0^{2^*-2} V_j \hat{W}_i dy + o(\Theta_{\ell,j}) \\ &= n \int_{B_0(\frac{\mu_{\ell-1}}{\mu_\ell})} U_0^{2^*-1} \partial_j \hat{W}_i dy + o(\Theta_{\ell,j}). \end{aligned}$$

If $i \leq \ell - 2$ we have, for any $y \in B_0(\frac{\mu_\ell}{\mu_{\ell-1}})$,

$$|\partial_j \hat{W}_i(y)| \lesssim \left(\frac{\mu_\ell}{\mu_i} \right)^{\frac{n}{2}},$$

so that

$$n \int_{B_0(\frac{\mu_{\ell-1}}{\mu_\ell})} U_0^{2^*-1} \partial_j \hat{W}_i dy = o(\Theta_{\ell,j}).$$

If $i = \ell - 1$ we write that

$$|\partial_j \hat{W}_{\ell-1}(y) - \partial_j \hat{W}_{\ell-1}(0)| \lesssim \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n+2}{2}} |y|,$$

so that

$$n \int_{B_0(\frac{\mu_{\ell-1}}{\mu_\ell})} U_0^{2^*-1} \partial_j \hat{W}_{\ell-1} dy = n D_3 \partial_j \hat{W}_{\ell-1}(0) + o(\Theta_{\ell,j}),$$

where we have let

$$D_3 = \int_{\mathbb{R}^n} U_0^{2^*-1} dy. \quad (5.18)$$

It remains to notice that, by definition of ξ_ℓ in (2.10), we have

$$\partial_j \hat{W}_{\ell-1}(0) = \left(\frac{\mu_\ell}{\mu_{\ell-1}} \right)^{\frac{n}{2}} (\partial_j U_0(z_\ell) + o(1)),$$

where U_0 is as in (2.16). This concludes the proof of Lemma 5.4 when $\ell \geq 2$.

It remains to prove the $\ell = 1$ case. We have, with (2.13) and the convention that $W_{k+1} = 0$ and $W_0 = 0$, that

$$\begin{aligned} &\left| - \int_M \left[\left(\sum_{i=1}^k W_i \right)^{2^*-1} - \sum_{i=1}^k W_i^{2^*-1} \right] Z_{1,j} dv_g \right| \\ &\lesssim \sum_{i \geq 1} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} (W_{i-1} + W_{i+1}) |Z_{1,j}| dv_g \\ &\lesssim \sum_{i \geq 2} \left(\frac{\mu_i^{\frac{n}{2}}}{\mu_{i-1} \mu_1^{\frac{n-2}{2}}} + \left(\frac{\mu_{i+1}}{\mu_1} \right)^{\frac{n-2}{2}} \right) + \left(\frac{\mu_2}{\mu_1} \right)^{\frac{n-2}{2}} \\ &= o(\Theta_{1,j}) \end{aligned}$$

for any $0 \leq j \leq n$, by (5.5). This concludes the proof of Lemma 5.4. \square

We are now in position to conclude the proof of Propositions 5.1 and 5.2. By Lemma 5.4 and (5.15) we only need to estimate the first term in the right-hand side of (5.15). We first treat the $j = 0$ case. By definition of W_ℓ in (2.11) and $Z_{\ell,0}$ in (2.15) it is easily seen that there holds

$$\frac{2}{n-2}\mu_\ell \frac{\partial}{\partial \mu_\ell} W_\ell = (n-2)\omega_{n-1}G_{g_\xi}(\xi, x)d_{g_\xi}(\xi, x)^{n-2}Z_{\ell,0} + O(\mu_\ell^{\frac{n-2}{2}})$$

By (2.1) and (2.2) this gives, since (M, g) is locally conformally flat:

$$Z_{\ell,0} = \left(1 + O(d_{g_{\xi_\ell}}(\xi_\ell, x)^{n-2})\right) \frac{2}{n-2}\mu_\ell \frac{\partial}{\partial \mu_\ell} W_\ell + O(\mu_\ell^{\frac{n-2}{2}}).$$

The latter expression then gives

$$\begin{aligned} & \int_M \left[\left(\Delta_g + c_n S_g + \varepsilon h \right) W_\ell - W_\ell^{2^*-1} \right] Z_{\ell,0} dv_g \\ &= \frac{2}{n-2}\mu_\ell \frac{\partial}{\partial \mu_\ell} \left[\frac{1}{2} \int_M |\nabla W_\ell|^2 + (c_n S_g + \varepsilon h) W_\ell^2 dv_g - \frac{1}{2^*} \int_M W_\ell^{2^*} dv_g \right] \\ &+ o(\Theta_{\ell,0}). \end{aligned} \quad (5.19)$$

For any $\ell \geq 1$, The computations in Esposito-Pistoia-Vétois [12] (Lemma 4.1) show that

$$\begin{aligned} & \frac{1}{2} \int_M |\nabla W_\ell|^2 + (c_n S_g + \varepsilon h) W_\ell^2 dv_g - \frac{1}{2^*} \int_M W_\ell^{2^*} dv_g \\ &= D_1 \varepsilon h(\xi_\ell) \mu_\ell^2 - D_2 \mu_\ell^{n-2} m(\xi_\ell) + o(\varepsilon \mu_\ell^2) \end{aligned} \quad (5.20)$$

for some positive constants D_1 and D_2 , where $m(\xi_\ell)$ is the mass as defined in (2.1), and where the $o(\cdot)$ terms are C^1 with respect to μ_ℓ and to ξ_ℓ . Plugging (5.20) into (5.19) yields

$$\begin{aligned} & \int_M \left[\left(\Delta_g + c_n S_g + \varepsilon h \right) W_\ell - W_\ell^{2^*-1} \right] Z_{\ell,0} dv_g \\ &= \frac{4D_1}{n-2} \varepsilon h(\xi_\ell) \mu_\ell^2 - 2D_2 \mu_\ell^{n-2} m(\xi_\ell) + o(\Theta_{\ell,0}). \end{aligned} \quad (5.21)$$

Assume now that $1 \leq j \leq n$. Lemmas 6.2 and 6.3 in [12] similarly show that

$$\begin{aligned} & \int_{B_\ell} \left[\left(\Delta_g + c_n S_g + \varepsilon h \right) W_\ell - W_\ell^{2^*-1} \right] Z_{\ell,j} dv_g \\ &= n\mu_\ell \frac{\partial}{\partial (\xi_\ell)_j} \left[\frac{1}{2} \int_M |\nabla W_\ell|^2 + (c_n S_g + \varepsilon h) W_\ell^2 dv_g - \frac{1}{2^*} \int_M W_\ell^{2^*} dv_g \right] + o(\Theta_{\ell,j}). \end{aligned}$$

With (5.20) this gives:

$$\begin{aligned}
& \int_{B_\ell} \left[(\Delta_g + c_n S_g + \varepsilon h) W_\ell - W_\ell^{2^*-1} \right] Z_{\ell,j} dv_g \\
&= nD_1 \varepsilon \nabla_j h(\xi_\ell) \mu_\ell^3 - nD_2 \mu_\ell^{n-1} \nabla_j m(\xi_\ell) + o(\Theta_{\ell,j}) \\
&= o(\Theta_{\ell,j}).
\end{aligned} \tag{5.22}$$

Coming back to (5.8) and using Lemma 5.3, (5.15), Lemma 5.4, (5.21) and (5.22) concludes the proof of Propositions 5.1 and 5.2. \square

We now conclude the proof of Theorem 1.1:

End of the proof of Theorem 1.1. Recall that (M, g) is a locally conformally flat manifold of dimension $n \geq 7$. Define a function $F : S_k \rightarrow \mathbb{R}^{k(1+n)}$ by:

$$F(t_1, \xi_1, \dots, t_k, z_k) = \left\{ \begin{array}{l} \frac{4D_1}{n-2} h(\xi_1) t_1^2 - 2D_2 t_1^{n-2} m(\xi_1) \\ \left(nD_1 t_1^3 \nabla_j h(\xi_1) - nD_2 t_1^{n-1} \nabla_j m(\xi_1) \right)_{1 \leq j \leq n} \\ \left(\frac{4D_1}{n-2} h(\xi_1) t_\ell^2 - D_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n-2}{2}} U_0(z_\ell) \right) \\ \left(-nD_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n}{2}} \partial_j U_0(z_\ell) \right)_{2 \leq \ell \leq k, 1 \leq j \leq n} \end{array} \right\},$$

where S_k is as in (2.9). Assume that (M, g) is not conformally diffeomorphic to the standard sphere and let ξ_0 be a critical point of the mass function $\xi \mapsto m(\xi) > 0$. Choose now $h \in C^1(M)$ such that $h(\xi_0) = 1$ and $\nabla h(\xi_0) = 0$ and let

$$t_{1,0} = \left(\frac{2D_1}{(n-2)D_2 m(\xi_0)} \right)^{\frac{1}{n-4}}.$$

Assume also that h is chosen so that ξ_0 has non-zero degree as a zero of

$$\xi \mapsto nD_1 t_{1,0}^3 \nabla h(\xi) - nD_2 t_{1,0}^{n-1} \nabla m(\xi).$$

These conditions are obviously met if $h \equiv 1$ and ξ_0 is a *non-degenerate* critical point of the mass function $\xi \mapsto m(\xi)$. Let finally, for $2 \leq \ell \leq k$,

$$t_{\ell,0} = \left(\frac{4D_1}{(n-2)D_3} t_{\ell-1,0}^{\frac{n-2}{2}} \right)^{\frac{2}{n-6}}.$$

It is then easily seen that $(t_{1,0}, \xi_0, (t_{\ell,0}, 0)_{2 \leq \ell \leq k})$ is a zero of F with non-zero degree. In particular, for any ε small enough, by Propositions 5.1 and 5.2, there exists $(t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{\ell,\varepsilon}, z_{\ell,\varepsilon})_{2 \leq \ell \leq k}) \in S_k$ that annihilates the $\nu_{i,j}$ defined

in (5.1). Denote by $\mu_{\ell,\varepsilon}$ and $\xi_{\ell,\varepsilon}$, $1 \leq \ell \leq k$, the parameters associated to $(t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{\ell,\varepsilon}, z_{\ell,\varepsilon})_{2 \leq \ell \leq k}) \in S_k$ as in (2.7) and (2.10). By (5.1), the function

$$\begin{aligned} u_\varepsilon &= \sum_{i=1}^k W_{i,\varepsilon} + \varphi_\varepsilon \\ &= \sum_{i=1}^k W_{\mu_i, \xi_i} + \varphi_\varepsilon \left((t_{1,\varepsilon}, \xi_{1,\varepsilon}, (t_{\ell,\varepsilon}, z_{\ell,\varepsilon})_{2 \leq \ell \leq k}) \right) \end{aligned}$$

solves (1.3). By the pointwise estimates on φ_ε given by Proposition 4.1 it is easily seen that u_ε blows-up with k towering bubbles around ξ_0 . This concludes the proof of Theorem 1.1. \square

6. SIGN-CHANGING BUBBLE TOWERS FOR THE BRÉZIS-NIRENBERG PROBLEM

We prove in this section Theorem 1.2. Let us consider the Brezis-Nirenberg problem

$$\Delta_\xi u - \varepsilon u = |u|^{\frac{4}{n-2}} u \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad (6.1)$$

where Ω is a bounded open domain in \mathbb{R}^n , $n \geq 7$, $\Delta_\xi = -\sum_{i=1}^n \partial_i^2$ and ε is a positive small parameter. For $\mu > 0$ and $\xi \in \Omega$ we introduce the standard bubble

$$U_{\mu,\xi}(x) := \mu^{-\frac{n-2}{2}} U_0 \left(\frac{x - \xi}{\mu} \right), \quad (6.2)$$

where U_0 is given by (2.16), and its projection $P_\Omega U_{\mu,\xi}$ on the space $H_0^1(\Omega)$, i.e. the solution to the Dirichlet boundary problem

$$\Delta_\xi P_\Omega U_{\mu,\xi} = \Delta_\xi U_{\mu,\xi} \text{ in } \Omega, \quad P_\Omega U_{\mu,\xi} = 0 \text{ on } \partial\Omega.$$

It is well known that (see e.g. Proposition 1 in Rey [33]) that

$$P_\Omega U_{\mu,\xi}(x) = U_{\mu,\xi}(x) - c_n \mu^{\frac{n-2}{2}} H(x, \xi) + O\left(\mu^{\frac{n+2}{2}}\right) \quad (6.3)$$

with $c_n := (n(n-2))^{\frac{n-2}{2}} \omega_{n-1}$, and that this expansion holds true uniformly with respect to $x \in \overline{\Omega}$ and ξ in compact sets of Ω . Here $H(\cdot, \xi)$ is the regular part of the Green function G of Δ_ξ in Ω with Dirichlet boundary condition, defined by

$$G(x, y) = \frac{1}{(n-2)\omega_{n-1}} |x - y|^{2-n} - H(x, y) \quad \forall x \neq y \in \Omega. \quad (6.4)$$

For an integer $k \geq 1$, we look for a sign-changing solution to (6.1) having the following form:

$$u(x) = \sum_{\ell=1}^k (-1)^\ell P_\Omega U_{\mu_\ell, \xi_\ell}(x) + \varphi_\varepsilon(x), \quad (6.5)$$

where the concentration parameters satisfy

$$\mu_\ell = t_\ell \epsilon^{\gamma_\ell} \text{ for } t_\ell > 0 \text{ and } \gamma_\ell = \frac{n-2}{2(n-4)} \left(\frac{n-2}{n-6} \right)^{\ell-1} - \frac{1}{2} \quad (6.6)$$

and the concentration points are given by

$$\xi_1 \in \Omega \text{ and } \xi_\ell = \xi_{\ell-1} + \mu_{\ell-1} z_\ell \text{ with } z_\ell \in \mathbb{R}^n, \text{ for any } 2 \leq \ell \leq k \quad (6.7)$$

All the arguments developed in the proof of Theorem 1.1 adapt straightforwardly in this setting with

- the bubbles W_ℓ replaced by the bubbles $P_\Omega U_{\mu_\ell, \xi_\ell}$,
- the elements $Z_{\ell,j}$ in (2.15) replaced by

$$Z_{\Omega, \ell, j} := P_\Omega \left(\mu_\ell^{-\frac{n-2}{2}} V_j \left(\frac{x - \xi_\ell}{\mu_\ell} \right) \right)$$

where V_j is defined in (2.17), and K_k in (2.23) is replaced by

$$K_{\Omega, k} = \text{Span} \left\{ Z_{\Omega, \ell, j}, 1 \leq \ell \leq k, 0 \leq j \leq n \right\}.$$

- and the configuration space S_k defined in (2.9) replaced by the set

$$S_{\Omega, k} = \left\{ (\xi, z_2, \dots, z_k, t_1, \dots, t_k) : d(\xi, \partial\Omega) \geq d, |z_\ell| \leq 1, A^{-1} \leq t_\ell \leq A \right\}$$

for some fixed constants $d > 0$ and $A > 1$.

More precisely, let for $(\xi, z_2, \dots, z_k, t_1, \dots, t_k) \in S_\Omega$ and for $x \in \Omega$:

- $\Psi_{\Omega, \varepsilon}(x) = \mu_{\ell-1}^{1-\frac{n}{2}} \left(\frac{\mu_\ell}{\mu_\ell + |x - \xi_\ell|} \right)^2 + \varepsilon \mu_{\ell-1}^{3-\frac{n}{2}}$
 $+ \frac{(\mu_{\ell+1} + |\xi_{\ell+1} - x|)^2}{\mu_\ell^2} P_\Omega U_{\mu_{\ell+1}, \xi_{\ell+1}}(x) \mathbb{1}_{\{|x - \xi_{\ell+1}| \leq \mu_\ell\}}$
 if $x \in B_\ell \setminus B_{\ell+1}$, for $\ell \geq 2$,
- $\Psi_{\Omega, \varepsilon}(x) = \frac{\mu_1^{\frac{n+2}{2}}}{(\mu_1 + |\xi_1 - x|)^2} + \frac{(\mu_2 + |\xi_2 - x|)^2}{\mu_1^2} P_\Omega U_{\mu_2, \xi_2}(x) \mathbb{1}_{\{|\xi_2 - x| \leq \mu_1\}}$
 if $x \in M \setminus B_2$,

where we have let $B_i = B(\xi_i, \sqrt{\mu_i \mu_{i-1}})$ for $i \geq 2$ and $B_1 = \Omega \setminus B_2$. A careful inspection of the proofs of Proposition 3.2 and Proposition 4.1 shows that the analysis carries on here (since the points ξ_1, \dots, ξ_ℓ live in compact sets in the interior) and yields the following result:

Proposition 6.1. *Let $k \geq 1$ be an integer and let $f(u) = |u|^{2^*-2}u$ for $u \in \mathbb{R}$. There exist $\varepsilon_1 > 0$ and $C > 0$ such that the following holds. For any $\varepsilon \in (0, \varepsilon_1)$ and for any $(\xi, z_2, \dots, z_k, t_1, \dots, t_k) \in S_{\Omega, k}$, there exists*

$$\varphi_\varepsilon := \varphi_\varepsilon(\xi, z_2, \dots, z_k, t_1, \dots, t_k) \in K_{\Omega, k}^\perp$$

which is the unique solution to

$$\Pi_{K_{\Omega,k}^\perp} \left[\sum_{\ell=1}^k (-1)^\ell P_\Omega U_{\mu_\ell, \xi_\ell} + \varphi_\varepsilon - (\Delta_\xi - \varepsilon)^{-1} \left(f \left(\sum_{\ell=1}^k (-1)^\ell P_\Omega U_{\mu_\ell, \xi_\ell} + \varphi_\varepsilon \right) \right) \right] = 0$$

that both belongs to $K_{\Omega,k}^\perp$ and satisfies $|\varphi_\varepsilon| \leq C\Psi_{\Omega,\varepsilon}$. In addition,

$$(\xi, z_2, \dots, z_k, t_1, \dots, t_k) \in S_{\Omega,k} \mapsto \varphi_\varepsilon \in C^0(\overline{\Omega})$$

is continuous for any fixed value of ε .

As in Section 5, this Proposition asserts the existence of $(\nu_{\ell,j})$, $1 \leq \ell \leq k$ and $0 \leq j \leq n$ such that

$$\begin{aligned} (\Delta_\xi - \varepsilon) \left(\sum_{\ell=1}^k (-1)^\ell P_\Omega U_{\mu_\ell, \xi_\ell} + \varphi_\varepsilon \right) - f \left(\sum_{\ell=1}^k (-1)^\ell P_\Omega U_{\mu_\ell, \xi_\ell} + \varphi_\varepsilon \right) \\ = \sum_{\substack{\ell=1..k \\ j=0..n}} \nu_{\ell,j} (\Delta_\xi - \varepsilon) Z_{\Omega,\ell,j} \end{aligned} \quad (6.8)$$

in Ω . Again, the analysis carried on in Section 5 remains true here. By (2.1) and (6.4) the analogue of the mass $m(\xi_1)$ is the opposite of Robin's function $-H(\xi, \xi)$. Taking into account that two consecutive bubbles have different sign so that their interaction is *negative* instead of *positive*, Proposition 5.1 and Proposition 5.1 can be rephrased as follows:

Proposition 6.2.

- (i) If $\ell = 1$ then:

$$\begin{aligned} \|\nabla V_0\|_{L^2(\mathbb{R}^n)}^2 \nu_{1,0}(\xi, z_2, \dots, z_k, t_1, \dots, t_k) \\ = \varepsilon^{1+2\gamma_1} \left(-\frac{4D_1}{n-2} t_1^2 + 2D_2 t_1^{n-2} H(\xi, \xi) + \Lambda_{0,1} \right) \end{aligned}$$

and, for $1 \leq j \leq n$,

$$\|\nabla V_j\|_{L^2(\mathbb{R}^n)}^2 \nu_{1,j}(\xi, z_2, \dots, z_k, t_1, \dots, t_k) = \varepsilon^{1+3\gamma_1} \left(nD_2 t_1^{n-1} \partial_j H(\xi, \xi) + \Lambda_{j,1} \right)$$

as $\varepsilon \rightarrow 0$.

- (ii) If $2 \leq \ell \leq k$

$$\begin{aligned} \|\nabla V_0\|_{L^2(\mathbb{R}^n)}^2 \nu_{\ell,0}(\xi, z_2, \dots, z_k, t_1, \dots, t_k) \\ = \varepsilon^{1+2\gamma_\ell} \left(-\frac{4D_1}{n-2} t_\ell^2 + D_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n-2}{2}} U_0(z_\ell) + \Lambda_{0,\ell} \right) \end{aligned}$$

and, for $1 \leq j \leq n$,

$$\begin{aligned} & \|\nabla V_j\|_{L^2(\mathbb{R}^n)}^2 \nu_{\ell,j}(\xi, z_2, \dots, z_k, t_1, \dots, t_k) \\ &= \varepsilon^{\frac{n}{2}(\gamma_\ell - \gamma_{\ell-1})} \left(nD_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n}{2}} \partial_j U_0(z_\ell) + \Lambda_{j,\ell} \right) \end{aligned}$$

as $\varepsilon \rightarrow 0$.

The functions U_0 is defined in (2.16), the positive constants D_i are the same as in Propositions 5.1 and 5.2 and the $\Lambda_{j,\ell}$'s are continuous functions on $S_{\Omega,k}$ which converge uniformly to 0 as $\varepsilon \rightarrow 0$.

Proof of Theorem 1.2. With Proposition 6.2 the end of the proof of Theorem 1.2 follows the same lines than the one of Theorem 1.1. Let ξ_0 be a non-degenerate interior critical point of the Robin's function $\xi \mapsto H(\xi, \xi)$. Let

$$t_{1,0} = \left(\frac{2D_1}{(n-2)D_2H(\xi_0, \xi_0)} \right)^{\frac{1}{n-4}}$$

and, for $\ell \geq 2$,

$$t_{\ell,0} = \left(\frac{4D_1}{(n-2)D_3} t_{\ell-1,0}^{\frac{n-2}{2}} \right)^{\frac{2}{n-6}}.$$

It is then easily checked that

$$(t_{1,0}, \xi_0, (t_{\ell,0}, 0)_{2 \leq \ell \leq k})$$

is a zero of the system

$$\begin{cases} -\frac{4D_1}{n-2}t_1^2 + 2D_2t_1^{n-2}H(\xi, \xi) = 0 \\ nD_2t_1^{n-1}\partial_j H(\xi, \xi) = 0 \\ \left(\begin{array}{l} -\frac{4D_1}{n-2}t_\ell^2 + D_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n-2}{2}} U_0(z_\ell) \\ nD_3 \left(\frac{t_\ell}{t_{\ell-1}} \right)^{\frac{n}{2}} \partial_j U_0(z_\ell) \end{array} \right)_{2 \leq \ell \leq k} = 0 \end{cases}$$

with non-zero degree. It is then stable under C^0 -perturbations. For any $\varepsilon > 0$ this provides us with $(\xi_\varepsilon, z_{2,\varepsilon}, \dots, z_{k,\varepsilon}, t_{1,\varepsilon}, \dots, t_{k,\varepsilon})$ such that

$$\nu_{\ell,j}(\xi_\varepsilon, z_{2,\varepsilon}, \dots, z_{k,\varepsilon}, t_{1,\varepsilon}, \dots, t_{k,\varepsilon}) = 0$$

for all $1 \leq \ell \leq k$ and $0 \leq j \leq n$. By (6.8), the corresponding

$$\varphi_\varepsilon = \varphi_\varepsilon(\xi_\varepsilon, z_{2,\varepsilon}, \dots, z_{k,\varepsilon}, t_{1,\varepsilon}, \dots, t_{k,\varepsilon})$$

provides us with a solution of (6.1) and concludes the proof of Theorem 1.2. \square

The Robin's function $\xi \mapsto H(\xi, \xi)$ always admits an interior critical point since H is positive and $H(\xi, \xi) \rightarrow +\infty$ as $\xi \rightarrow \partial\Omega$. And for generic domains Ω , all the critical points of the Robin's function are non-degenerate as proved in Micheletti-Pistoia [25]. Finally, here again, if one considers more general equations of the form

$$\Delta_\xi u - \varepsilon h u = |u|^{\frac{4}{n-2}} u \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega$$

for $h \in C^0(\overline{\Omega})$, an analogue of Theorem 1.2 can be proven where towering phenomena occur at *any* critical point of the Robin's function $\xi \mapsto H(\xi, \xi)$ (provided h is suitably chosen). Proposition 6.1 remains indeed true and the details are identical to those developed at the very end of Section 5.

APPENDIX A. TECHNICAL RESULTS

We keep the notations of Section 2. The integer $k \geq 1$ refers to the number of bubbles.

A.1. Proof of (2.21) and (2.22). Let $1 \leq i < j \leq k$ and $0 \leq p, q \leq n$. We first prove (2.21). We write for this, using (2.14), (2.19) and (2.20) and since $|Z_{i,p}| \lesssim W_i$, that

$$\begin{aligned} \langle Z_{i,p}, Z_{i,q} \rangle &= \int_M (\Delta_g + c_n S_g + \varepsilon h) Z_{i,p} Z_{i,q} dv_g \\ &= (2^* - 1) \int_M W_i^{2^*-2} Z_{i,p} Z_{i,q} dv_g + O\left(\int_M W_i^2 dv_g\right) \\ &= (2^* - 1) \int_M W_i^{2^*-2} Z_{i,p} Z_{i,q} dv_g + O(\mu_i^2). \end{aligned}$$

Letting $y = \exp_{\xi_i}^{g_{\xi_i}}(\mu_i x)$ and using the conformal invariance of the conformal laplacian and (2.4) we get:

$$\begin{aligned} \int_M W_i^{2^*-2} Z_{i,p} Z_{i,q} dv_g &= \int_{Bg_{\xi_i}(\xi_i, r_0)} W_i^{2^*-2} Z_{i,p} Z_{i,q} dv_g + O(\mu_i^n) \\ &= \int_{B(0, \frac{r_0}{\mu_i})} U_0^{2^*-2} V_p V_q dx + O(\mu_i^4) \\ &= \int_{\mathbb{R}^n} U_0^{2^*-2} V_p V_q dx + O(\mu_i^4) \\ &= \|\nabla V_p\|_{L^2(\mathbb{R}^n)}^2 \delta_{pq} + O(\mu_i^4), \end{aligned}$$

where r_0 is as in (2.3), which proves (2.21).

We now prove (2.22). With (2.19) and (2.20) we have:

$$\begin{aligned}
& \langle Z_{i,p}, Z_{j,q} \rangle \\
&= \int_M \left[\Delta_g Z_{j,q} + (c_n S_g + \varepsilon h) Z_{j,q} \right] Z_{i,p} dv_g \\
&= (2^* - 1) \int_M W_j^{2^*-2} Z_{j,q} Z_{i,p} dv_g + \int_M \left(\varepsilon Z_{j,q} Z_{i,p} + O(\mu_j W_j |Z_{i,q}|) \right) dv_g \\
&= O \left(\left(\frac{\mu_j}{\mu_i} \right)^{\frac{n-2}{2}} \right),
\end{aligned}$$

where we just used the rough estimate $|Z_{i,p}| \lesssim \mu_i^{1-\frac{n}{2}}$.

A.2. Additional material. We prove two inequalities that were used several times throughout the proof.

Lemma A.1. *Let $i \in \{1, \dots, k\}$, and $x \in M$. We have:*

If $0 \leq p < n - 4$:

$$\begin{aligned}
& \int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} \left(\frac{\mu_i}{\theta_i} \right)^p dv_g \\
& \lesssim \begin{cases} \left(\frac{\mu_i}{\theta_i(x)} \right)^{p+2} & \text{if } x \in B_i \\ \mu_i^{\frac{p+2}{2}} \mu_{i-1}^{\frac{n-4-p}{2}} W_i(x) & \text{if } x \in M \setminus B_i \end{cases} \quad (\text{A.1})
\end{aligned}$$

If $p > n - 4$:

$$\int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} \left(\frac{\mu_i}{\theta_i} \right)^p dv_g \lesssim \left(\frac{\mu_i}{\theta_i(x)} \right)^{n-2} = \mu_i^{\frac{n-2}{2}} W_i(x),$$

where θ_i is as in (2.18).

Proof. Assume first that $d_{g_{\xi_i}}(\xi_i, x) \leq 2\sqrt{\mu_i \mu_{i-1}}$. Then $W_i^{2^*-2} \lesssim \mu_i^2 \theta_i(x)^{-4}$, where θ_i is as in (2.18). If $0 \leq p < n - 4$, the result follows from Giraud's lemma. While if $p > n - 4$ letting $y = \exp_{\xi_i}^{g_{\xi_i}}(\mu_i z)$ in the integral and $\tilde{x} = \frac{1}{\mu_i} \exp_{\xi_i}^{g_{\xi_i}-1}(x)$ yields :

$$\begin{aligned}
& \int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} \left(\frac{\mu_i}{\theta_i} \right)^p dv_g \lesssim \int_{\mathbb{R}^n} |\tilde{x} - z|^{2-n} (1 + |z|)^{-n-2} dz \\
& \lesssim \frac{1}{(1 + |\tilde{x}|)^{n-2}} \lesssim \left(\frac{\mu_i}{\theta_i(x)} \right)^{n-2}.
\end{aligned}$$

Assume then that $d_{g_{\xi_i}}(\xi_i, x) \geq 2\sqrt{\mu_{i-1} \mu_i}$ (note that this only makes sense for $i \geq 2$). Then for any $y \in B_i$ one has $d_g(x, y) \gtrsim d_g(x, \xi_i) \gtrsim \theta_i(x)$. If $p < n - 4$

we then have

$$\begin{aligned}
\int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} \left(\frac{\mu_i}{\theta_i} \right)^p dv_g &\lesssim \theta_i(x)^{2-n} \int_{B_i} W_i^{2^*-2} \left(\frac{\mu_i}{\theta_i} \right)^p \\
&= \theta_i(x)^{2-n} \mu_i^{p+2} (\mu_i \mu_{i-1})^{\frac{n-4-p}{2}} \\
&= \mu_i^{\frac{p+2}{2}} \mu_{i-1}^{\frac{n-4-p}{2}} W_i(x),
\end{aligned}$$

while if $p > n - 4$ we have

$$\begin{aligned}
\int_{B_i \setminus B_{i+1}} d_g(x, \cdot)^{2-n} W_i^{2^*-2} \left(\frac{\mu_i}{\theta_i} \right)^p dv_g &\lesssim \theta_i(x)^{2-n} \int_{B_i} W_i^{2^*-2} \left(\frac{\mu_i}{\theta_i} \right)^p \\
&= \theta_i(x)^{2-n} \mu_i^{n-2} \int_{\mathbb{R}^n} (1 + |y|)^{-p-4} dy \\
&\lesssim \left(\frac{\mu_i}{\theta_i(x)} \right)^{n-2}.
\end{aligned}$$

Note finally that, in the intermediate case $\sqrt{\mu_{i-1}\mu_i} \leq d_{g_{\xi_i}}(\xi_i, x) \leq 2\sqrt{\mu_{i-1}\mu_i}$ we have

$$\left(\frac{\mu_i}{\theta_i(x)} \right)^{p+2} \sim \left(\frac{\mu_i}{\mu_{i-1}} \right)^{\frac{p+2}{2}} \sim \mu_i^{\frac{p+2}{2}} \mu_{i-1}^{\frac{n-4-p}{2}} W_i(x)$$

and this holds true for $0 \leq p \leq n - 4$, so that the result follows. \square

Lemma A.2. *For any $i, j \in \{1, \dots, k\}$, $i < j$, and any $x \in M$, we have*

$$\begin{aligned}
&\int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_j d_g(x, \cdot)^{2-n} dv_g \\
&\lesssim \begin{cases} \left(\frac{\mu_j}{\mu_{i+1}} \right)^{\frac{n-2}{2}} \frac{\mu_{i+1}}{\mu_i} \mu_i^{1-\frac{n}{2}} & \text{if } x \in B_{i+1} \\ \left(\frac{\mu_j}{\mu_i} \right)^{\frac{n-2}{2}} W_i(x) & \text{if } x \in B_i \setminus B_{i+1} \text{ and } \theta_j(x) \geq \mu_i \\ \frac{\theta_j(x)^2}{\mu_i^2} W_j(x) & \text{if } x \in B_i \setminus B_{i+1} \text{ and } \theta_j(x) \leq \mu_i \\ \left(\frac{\mu_j}{\mu_i} \right)^{\frac{n-2}{2}} W_i(x) & \text{if } x \in M \setminus B_i. \end{cases} \quad (\text{A.2})
\end{aligned}$$

Proof. Assume first that $d_{g_{\xi_i}}(\xi_i, x) \geq \sqrt{\mu_i \mu_{i-1}}$, that is $x \in M \setminus B_i$ (this case is empty if $i = 1$). Changing variables in the integral by $y = \exp_{\xi_i}^{g_{\xi_i}}(\mu_i x)$ and using Giraud's lemma yields in this case

$$\int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_j d_g(x, \cdot)^{2-n} dv_g \lesssim \theta_i(x)^{2-n} \mu_j^{\frac{n-2}{2}} = \left(\frac{\mu_j}{\mu_i} \right)^{\frac{n-2}{2}} W_i(x).$$

Assume now that $x \in B_{i+1}$, so in particular $\theta_j(x) \lesssim \sqrt{\mu_{i+1}\mu_i}$ by (2.18). We let the change of variables

$$y = \exp_{\xi_j}^{g_{\xi_j}}(\theta_j(x)z)$$

in the integral. By (2.10) one has

$$\begin{aligned} & \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_j d_g(x, \cdot)^{2-n} dv_g \\ & \lesssim \mu_i^{-2} \mu_j^{\frac{n-2}{2}} \theta_j(x)^{4-n} \\ & \quad \times \int_{B_0 \left(2\sqrt{\frac{\mu_i-1\mu_i}{\theta_j(x)}} \right) \setminus B_0 \left(\frac{1}{2}\sqrt{\frac{\mu_{i+1}\mu_i}{\theta_j(x)}} \right)} |\check{x}_i - z|^{2-n} \left(\frac{\mu_j}{\theta_j(x)} + |z| \right)^{2-n} dz, \end{aligned}$$

where we have let $\check{x}_i = \frac{1}{\theta_j(x)} (\exp_{\xi_j}^{g_{\xi_j}})^{-1}(x)$. As is easily checked, the previous integral is always uniformly integrable whatever the value of $\theta_j(x)$ is, and there holds $|\check{x}_i| \leq 1$ by definition of θ_j . So that in the end

$$\begin{aligned} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_j d_g(x, \cdot)^{2-n} dv_g & \lesssim \mu_i^{-2} \mu_j^{\frac{n-2}{2}} \theta_j(x)^{4-n} \int_{\mathbb{R}^n \setminus B_0 \left(\frac{\sqrt{\mu_{i+1}\mu_i}}{\theta_j(x)} \right)} |z|^{4-2n} dz \\ & \lesssim \mu_i^{-2} \mu_j^{\frac{n-2}{2}} \theta_j(x)^{4-n} \left(\frac{\theta_j(x)}{\sqrt{\mu_{i+1}\mu_i}} \right)^{n-4} \\ & \lesssim \left(\frac{\mu_j}{\mu_{i+1}} \right)^{\frac{n-2}{2}} \frac{\mu_{i+1}}{\mu_i} \mu_i^{1-\frac{n}{2}}. \end{aligned}$$

Assume now that $x \in B_i \setminus B_{i+1}$. The change of variables

$$y = \exp_{\xi_j}^{g_{\xi_j}}(\mu_i z)$$

gives

$$\begin{aligned} & \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_j d_g(x, \cdot)^{2-n} dv_g \\ & \lesssim \mu_j^{\frac{n-2}{2}} \mu_i^{2-n} \int_{\mathbb{R}^n \setminus B_0 \left(\frac{1}{2}\sqrt{\frac{\mu_{i+1}}{\mu_i}} \right)} (1 + |\check{\xi} - y|)^{-4} |y|^{2-n} |\check{x} - y|^{2-n} dy, \end{aligned} \tag{A.3}$$

where we have let $\check{\xi} = \frac{1}{\mu_i} (\exp_{\xi_j}^{g_{\xi_j}})^{-1}(\xi_i)$ and $\check{x} = \frac{1}{\mu_i} (\exp_{\xi_j}^{g_{\xi_j}})^{-1}(x)$. Since $x \notin B_{i+1}$ we have $\theta_j(x) \gtrsim \sqrt{\mu_{i+1}\mu_i}$ and thus $|\check{x}| \gtrsim \frac{\theta_j(x)}{\mu_i}$. Assume first that $|\check{x}| \gtrsim 1$. Then we also have $\theta_j(x) \gtrsim \theta_i(x)$, so that classical integral comparison results yield

$$\int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_j d_g(x, \cdot)^{2-n} dv_g \lesssim \mu_j^{\frac{n-2}{2}} \mu_i^{2-n} (1 + |\check{x}|)^{2-n} \lesssim \left(\frac{\mu_j}{\mu_i} \right)^{\frac{n-2}{2}} W_i(x).$$

Assume finally that $|\check{x}| \lesssim 1$. A standard Giraud-type argument (see e.g. lemma 7.5 in Hebey [13]) yields with (A.3)

$$\begin{aligned} \int_{B_i \setminus B_{i+1}} W_i^{2^*-2} W_j d_g(x, \cdot)^{2-n} dv_g &\lesssim \mu_j^{\frac{n-2}{2}} \mu_i^{2-n} |\check{x}|^{4-n} \\ &\lesssim \frac{\theta_j(x)^2}{\mu_i^2} W_j(x), \end{aligned}$$

which concludes the proof of (A.2). \square

REFERENCES

1. Thierry Aubin, *Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire*, J. Math. Pures Appl. (9) **55** (1976), no. 3, 269–296. MR 431287
2. Mohamed Ben Ayed, Khalil El Mehdi, and Filomena Pacella, *Blow-up and nonexistence of sign changing solutions to the Brezis-Nirenberg problem in dimension three*, Ann. Inst. H. Poincaré Anal. Non Linéaire **23** (2006), no. 4, 567–589. MR 2245756
3. ———, *Blow-up and symmetry of sign-changing solutions to some critical elliptic equations*, J. Differential Equations **230** (2006), no. 2, 771–795. MR 2269943
4. Simon Brendle, *Blow-up phenomena for the Yamabe equation*, J. Amer. Math. Soc. **21** (2008), no. 4, 951–979. MR 2425176 (2009m:53084)
5. Simon Brendle and Fernando C. Marques, *Blow-up phenomena for the Yamabe equation. II*, J. Differential Geom. **81** (2009), no. 2, 225–250. MR 2472174 (2010k:53050)
6. Haïm Brézis and Louis Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), no. 4, 437–477. MR 709644
7. Katiuscia Cerquetti, *A uniqueness result for a semilinear elliptic equation involving the critical Sobolev exponent in symmetric domains*, Asymptot. Anal. **21** (1999), no. 2, 99–115. MR 1723563
8. Olivier Druet, *From one bubble to several bubbles: the low-dimensional case*, J. Differential Geom. **63** (2003), no. 3, 399–473. MR 2015469
9. ———, *Compactness for Yamabe metrics in low dimensions*, Int. Math. Res. Not. (2004), no. 23, 1143–1191. MR 2041549 (2005b:53056)
10. Olivier Druet, Emmanuel Hebey, and Frédéric Robert, *Blow-up theory for elliptic PDEs in Riemannian geometry*, Mathematical Notes, vol. 45, Princeton University Press, Princeton, NJ, 2004. MR 2063399 (2005g:53058)
11. Pierpaolo Esposito, Nassif Ghoussoub, Angela Pistoia, and Giusi Vaira, *Sign-changing solutions for critical equations with hardy potential*, Anal. PDE, To appear.
12. Pierpaolo Esposito, Angela Pistoia, and Jérôme Vétois, *The effect of linear perturbations on the Yamabe problem*, Math. Ann. **358** (2014), no. 1-2, 511–560. MR 3158007
13. Emmanuel Hebey, *Compactness and stability for nonlinear elliptic equations*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2014. MR 3235821
14. Alessandro Iacopetti, *Asymptotic analysis for radial sign-changing solutions of the Brezis-Nirenberg problem*, Ann. Mat. Pura Appl. (4) **194** (2015), no. 6, 1649–1682. MR 3419911
15. Alessandro Iacopetti and Filomena Pacella, *A nonexistence result for sign-changing solutions of the Brezis-Nirenberg problem in low dimensions*, J. Differential Equations **258** (2015), no. 12, 4180–4208. MR 3327552

16. Alessandro Iacopetti and Giusi Vaira, *Sign-changing tower of bubbles for the Brezis-Nirenberg problem*, Commun. Contemp. Math. **18** (2016), no. 1, 1550036, 53. MR 3454626
17. ———, *Sign-changing blowing-up solutions for the Brezis-Nirenberg problem in dimensions four and five*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **18** (2018), no. 1, 1–38. MR 3783782
18. M. A. Khuri, F. C. Marques, and R. M. Schoen, *A compactness theorem for the Yamabe problem*, J. Differential Geom. **81** (2009), no. 1, 143–196. MR 2477893 (2010e:53065)
19. John M. Lee and Thomas H. Parker, *The Yamabe problem*, Bull. Amer. Math. Soc. (N.S.) **17** (1987), no. 1, 37–91. MR 888880 (88f:53001)
20. Yan Yan Li, *Prescribing scalar curvature on S^n and related problems. I*, J. Differential Equations **120** (1995), no. 2, 319–410. MR 1347349
21. Yan Yan Li and Lei Zhang, *Compactness of solutions to the Yamabe problem. III*, J. Funct. Anal. **245** (2007), no. 2, 438–474. MR 2309836 (2008f:53038)
22. Yanyan Li, *Prescribing scalar curvature on S^n and related problems. II. Existence and compactness*, Comm. Pure Appl. Math. **49** (1996), no. 6, 541–597. MR 1383201
23. Yanyan Li and Meijun Zhu, *Yamabe type equations on three-dimensional Riemannian manifolds*, Commun. Contemp. Math. **1** (1999), no. 1, 1–50. MR 1681811 (2000m:53051)
24. Fernando Coda Marques, *A priori estimates for the Yamabe problem in the non-locally conformally flat case*, J. Differential Geom. **71** (2005), no. 2, 315–346. MR 2197144 (2006i:53046)
25. Anna Maria Micheletti and Angela Pistoia, *Non degeneracy of critical points of the Robin function with respect to deformations of the domain*, Potential Anal. **40** (2014), no. 2, 103–116. MR 3152157
26. Filippo Morabito, Angela Pistoia, and Giusi Vaira, *Towering phenomena for the Yamabe equation on symmetric manifolds*, Potential Anal. **47** (2017), no. 1, 53–102. MR 3666799
27. Monica Musso and Angela Pistoia, *Multispoke solutions for a nonlinear elliptic problem involving the critical Sobolev exponent*, Indiana Univ. Math. J. **51** (2002), no. 3, 541–579. MR 1911045
28. ———, *Tower of bubbles for almost critical problems in general domains*, J. Math. Pures Appl. (9) **93** (2010), no. 1, 1–40. MR 2579374
29. Angela Pistoia and Giusi Vaira, *Clustering phenomena for linear perturbation of the Yamabe equation*, Partial differential equations arising from physics and geometry, London Math. Soc. Lecture Note Ser., vol. 450, Cambridge Univ. Press, Cambridge, 2019, pp. 311–331. MR 3930853
30. Bruno Premoselli, *A pointwise finite-dimensional reduction method for a fully coupled system of Einstein-Lichnerowicz type*, Commun. Contemp. Math. **20** (2018), no. 6, 1750076, 72. MR 3848069
31. ———, *A pointwise finite-dimensional reduction method for Einstein-Lichnerowicz-type systems: the six-dimensional case*, Nonlinear Anal. **172** (2018), 200–215. MR 3790374
32. Bruno Premoselli and Pierre-Damien Thizy, *Bubbling above the threshold of the scalar curvature in dimensions four and five*, Calc. Var. Partial Differential Equations **57** (2018), no. 6, Paper No. 147, 39. MR 3858830
33. Olivier Rey, *The role of the Green’s function in a nonlinear elliptic equation involving the critical Sobolev exponent*, J. Funct. Anal. **89** (1990), no. 1, 1–52. MR 1040954
34. Frédéric Robert, *Existence et asymptotiques optimales des fonctions de Green des opérateurs elliptiques d’ordre deux*, <http://www.iecn.un-nancy.fr/frobert/ConstrucGreen.pdf>.

35. Frédéric Robert and Jérôme Vétois, *Examples of non-isolated blow-up for perturbations of the scalar curvature equation on non-locally conformally flat manifolds*, J. Differential Geom. **98** (2014), no. 2, 349–356. MR 3263521
36. ———, *A general theorem for the construction of blowing-up solutions to some elliptic nonlinear equations with Lyapunov-Schmidt's finite-dimensional reduction*, Concentration Compactness and Profile Decomposition (Bangalore, 2011), Trends in Mathematics, Springer, Basel (2014), 85–116.
37. Richard Schoen, *Conformal deformation of a Riemannian metric to constant scalar curvature*, J. Differential Geom. **20** (1984), no. 2, 479–495. MR 788292
38. Richard M. Schoen, *Lecture notes from courses at stanford*, (1988), Written by D. Pollack, preprint.
39. Michael Struwe, *A global compactness result for elliptic boundary value problems involving limiting nonlinearities*, Math. Z. **187** (1984), no. 4, 511–517. MR 760051
40. Pierre-Damien Thizy and Jérôme Vétois, *Positive clusters for smooth perturbations of a critical elliptic equation in dimensions four and five*, J. Funct. Anal. **275** (2018), no. 1, 170–195. MR 3799627
41. Neil S. Trudinger, *Remarks concerning the conformal deformation of Riemannian structures on compact manifolds*, Ann. Scuola Norm. Sup. Pisa (3) **22** (1968), 265–274. MR 0240748
42. Hidehiko Yamabe, *On a deformation of Riemannian structures on compact manifolds*, Osaka Math. J. **12** (1960), 21–37. MR 125546

BRUNO PREMOSELLI, UNIVERSITÉ LIBRE DE BRUXELLES, SERVICE DE GÉOMÉTRIE DIFFÉRENTIELLE CP 218, BOULEVARD DU TRIOMPHE, B-1050 BRUXELLES, BELGIQUE.
E-mail address: bruno.premoselli@ulb.ac.be,